

CIOMS FORM I — ACUTE KIDNEY INJURY SPONTANEOUS REPORT
International Reporting of Adverse Drug Reactions | Case Ref: CIOMS-RENAL-2025-099

I. REACTION INFORMATION

1. PATIENT INITIALS: G.T. (George Taylor)	2. AGE: 67 YRS	3. SEX: MALE	4. WEIGHT: 76 kg	5. COUNTRY: USA
6. DESCRIBE REACTION: 67-year-old male with chronic osteoarthritis initiated Renotril 30 mg PO QD. Baseline Serum Creatinine: 1.0 mg/dL (eGFR 78 mL/min). On day 14 of treatment, patient presented with oliguria (<400 mL/day) and peripheral edema. Laboratory chemistry revealed Serum Creatinine spiked to 4.2 mg/dL (KDIGO Stage 3 Acute Kidney Injury) with BUN 68 mg/dL and potassium 5.6 mEq/L. Renotril permanently withdrawn. Patient hospitalized for 5 days with aggressive IV hydration. Renal function recovered to baseline (Cr 1.1 mg/dL) at 3 weeks post-discharge. Seriousness: [X] Inpatient Hospitalization: YES. Suspect Drug: Renotril 30mg, Lot #RN-2025-412. Reporter: Dr. Keith Miller, MD (Nephrology), Vanderbilt University Medical Center.				