MAGI+: From Diagnosis to Action in Automated Psychiatric Interviews

Abstract

Automated psychiatric screening has the potential to address the global shortage of mental health professionals by enabling scalable, accessible initial assessments. The original MAGI framework (Bi et al., 2025) introduced a multi-agent system that operationalizes the Mini International Neuropsychiatric Interview (MINI) protocol through coordinated LLM agents, achieving clinical-grade diagnostic performance via Psychometric Chain-of-Thought (PsyCoT) reasoning. However, MAGI stops at binary diagnosis, lacking the clinical context required for real-world deployment: emotional state tracking, severity grading, treatment guidance, and longitudinal monitoring.

We present MAGI+, an extension of MAGI that introduces five novel contributions: (1) EmotionArc, a per-turn emotion timeline tracking participant affect throughout the interview; (2) Severity Scoring, automated mapping of collected symptoms to standardized scales (PHQ-9, GAD-7, C-SSRS); (3) a Recommendation Agent, a fifth LLM agent that generates tiered, evidence-based treatment recommendations with crisis escalation logic; (4) Longitudinal Memory, multi-session tracking enabling trend analysis across visits; and (5) a Clinical Report Generator producing structured, clinician-ready assessment documents. We implement MAGI+ using the qwen/qwen3.8-27b model via the Groq inference API, achieving sub-second per-turn latency (~400ms).

We evaluate MAGI+ on ten simulated clinical cases -- following the same methodology as the original MAGI study -- covering major depressive episode, generalized anxiety disorder, social anxiety disorder, high-risk suicidality, comorbid presentations, and healthy controls, achieving 90% diagnostic accuracy and 100% crisis detection rate. To our knowledge, no prior system combines structured psychiatric interviewing with automatic severity scoring and treatment recommendations in a single pipeline -- MAGI+ is a step toward closing that gap.

Keywords: psychiatric AI, multi-agent systems, mental health assessment, MINI interview, PsyCoT reasoning, large language models, clinical NLP

1. Introduction

Mental health disorders affect over one billion people globally (WHO, 2022), yet access to psychiatric evaluation remains severely limited. In many low- and middle-income countries, the ratio of psychiatrists to population is fewer than 1 per 100,000 people (WHO, 2021). Even in high-income countries, waiting times for psychiatric assessment routinely exceed weeks or months. This treatment gap represents one of the most significant unmet healthcare needs worldwide.

Structured clinical interviews -- particularly the Mini International Neuropsychiatric Interview (MINI; Sheehan et al., 1998) -- are the gold standard for psychiatric assessment. The MINI provides a systematic,

DSM-5-aligned framework covering 17 psychiatric disorders in 15-30 minutes. However, its manual administration requires trained clinicians, limiting scalability.

The advent of large language models (LLMs) presents an opportunity to automate psychiatric assessment while maintaining clinical rigor. The recent MAGI framework (Bi et al., 2025) demonstrated that a coordinated system of four specialized LLM agents can operationalize the MINI protocol with clinical-grade diagnostic performance, achieving Cohen's kappa of 0.839-0.942 for suicide risk detection and F1 scores above 0.850 for depression.

However, MAGI has significant limitations for real-world clinical deployment. In practice, a binary Yes/No diagnosis is rarely enough. Clinicians need to know how severe the condition is, what to do next, and how the patient has changed over time. MAGI provides none of this -- it diagnoses but does not grade, recommend, or remember. These gaps represent the difference between a research prototype and a tool that can actually be used in a clinical setting.

We present MAGI+, which addresses all five limitations through novel extensions while preserving MAGI's original diagnostic architecture. Our contributions are:

1. EmotionArc: Real-time per-turn emotion classification and timeline construction, enabling detection of distress escalation during the interview and informing adaptive questioning strategy.
2. Severity Scoring: Automatic mapping of collected diagnostic evidence to validated clinical severity scales (PHQ-9 for depression, GAD-7 for anxiety, C-SSRS for suicide risk), enabling severity-stratified clinical decision making.
3. Recommendation Agent: A novel fifth LLM agent that synthesizes diagnosis and severity scores into tiered, evidence-based treatment recommendations (self-help -> outpatient therapy -> psychiatric consultation -> crisis intervention), with automatic crisis escalation for high suicide risk.
4. Longitudinal Memory: Persistent session storage enabling cross-session trend analysis, detection of new diagnoses, and monitoring of treatment response over time.
5. Clinical Report Generator: Automated generation of structured clinical assessment reports containing all assessment data in a format suitable for electronic health records and clinician review.

To our knowledge, MAGI+ is the first AI psychiatric assessment system to cover the complete clinical pathway from structured interview through diagnosis, severity quantification, and treatment recommendation in a single integrated pipeline.

---

2. Related Work

2.1 Automated Psychiatric Assessment

Early computational approaches to psychiatric assessment relied on rule-based systems and standardized questionnaires administered via computer interface (Merry et al., 2012). While scalable, these lacked the conversational adaptability necessary for sensitive clinical interactions. Natural language processing advances enabled analysis of mental health indicators from text (Sharma et al., 2020), while LLMs demonstrated potential in cognitive distortion detection (Chen et al., 2023) and clinician role-playing in

diagnostic dialogue (Tu et al., 2024).

2.2 LLM-Based Mental Health Systems

Several commercial and research systems have deployed LLMs for mental health support (Woebot, Wysa), primarily as conversational companions rather than diagnostic tools. These systems demonstrate engagement benefits but lack clinical diagnostic validity and are not aligned with structured psychiatric protocols. LLM-based systems have been evaluated for measuring psychiatric functioning from clinical interviews (Galatzer-Levy et al., 2023), though existing approaches struggle with maintaining diagnostic protocol adherence across extended conversations.

2.3 The MINI Protocol

The MINI (Sheehan et al., 1998) is a structured diagnostic interview aligned with DSM-5 criteria. It implements decision-tree logic where responses guide interview progression, ensuring standardized administration. The MINI can be completed in 15-30 minutes with demonstrated reliability (ICC > 0.78) and validity across diverse populations. MAGI (Bi et al., 2025) was the first system to computationally operationalize the MINI through multi-agent LLM collaboration.

2.4 Severity Assessment and Clinical Scales

Clinical severity assessment relies on validated scales including the PHQ-9 (Kroenke et al., 2001) for depression (sensitivity 88%, specificity 88% at threshold >=10), GAD-7 (Spitzer et al., 2006) for anxiety, and C-SSRS (Posner et al., 2011) for suicide risk. These scales provide standardized severity quantification that guides treatment intensity decisions. No existing automated psychiatric interview system produces validated scale scores as primary output.

2.5 Multi-Agent Systems in Healthcare AI

Multi-agent LLM systems have shown effectiveness in complex healthcare reasoning tasks requiring coordinated specialization (Park et al., 2023). In clinical contexts, agent coordination enables separation of concerns between protocol navigation, empathetic communication, evidence validation, and diagnostic synthesis -- precisely the architecture employed by MAGI and extended by MAGI+.

2.6 Affective Computing and Emotion Detection in Mental Health AI

Emotion detection has a well-established role in mental health research. Computational approaches range from lexicon-based methods (e.g., LIWC; Pennebaker et al., 2001) to deep learning classifiers trained on clinical corpora. In the context of psychiatric interviews, affective state carries significant diagnostic signal -- patients with depression frequently exhibit emotional flattening or tearfulness during assessment, while those with anxiety show elevated arousal, and patients at crisis levels may present with emotional dissociation from their verbal responses. Sharma et al. (2020) demonstrated that patterns in emotional language on peer support platforms could predict deterioration in mental health outcomes weeks in advance.

Despite this, no existing automated psychiatric interview system formally tracks or outputs participant

emotional state as part of its assessment pipeline. MAGI detects distress lexically to inform question strategy, but does not produce an emotion timeline or include affective data in its diagnostic output. MAGI+'s EmotionArc module addresses this gap by providing per-turn emotion classification, intensity scoring, and escalation detection as first-class clinical output.

2.7 Gap Analysis: MAGI+ vs. Prior Work

Table 1 summarizes the capabilities of key related systems compared to MAGI+. No prior system covers all five dimensions simultaneously.

System	Structured Protoc	Emotion Tracking	Severity Scores	Treatment Recom	Longitudinal Mem	Clinical Report
Woebot / Wysa	[NO]	Partial	[NO]	Partial	Partial	[NO]
AMIE (Tu et al., 2024)	[NO]	[NO]	[NO]	[NO]	[NO]	[NO]
Galatzer-Levy et al. (2023)	[NO]	[NO]	Partial	[NO]	[NO]	[NO]
MAGI (Bi et al., 2025)	MINI	Lexical only	[NO]	[NO]	[NO]	[NO]
MAGI+ (Ours)	** MINI**	** EmotionArc**	** PHQ-9/GAD-7/C-SSRS**	** Stepped + Crisis**	** Multi-session**	** Full report**

Table 1: Capability comparison. = fully implemented, Partial = limited implementation, [NO] = not present.

---

3. Background: The MAGI Framework

MAGI (Bi et al., 2025) transforms the MINI interview into a computational workflow through four coordinated agents:

Navigation Agent: Governs interview progression by enforcing MINI's decision-tree branching logic. Ensures no critical diagnostic nodes are skipped and routes the interview based on participant responses (e.g., suicidal ideation triggers the suicide risk section).

Question Agent: Generates adaptive, contextually appropriate questions using three strategies -- diagnostic probing, clinical concept explanation, and empathetic support -- selected based on detected participant distress or confusion.

Judgment Agent: Validates participant responses against MINI criteria through three thresholds: direct matching, semantic equivalence, and forced-choice resolution after persistent ambiguity.

Diagnosis Agent: Synthesizes collected evidence through Psychometric Chain-of-Thought (PsyCoT) reasoning -- a three-phase framework comprising symptom anchoring, diagnostic synthesis, and evidence binding -- producing DSM-5-aligned diagnoses with transparent reasoning traces.

MAGI was validated on 1,002 real-world participants with dual expert annotation (Cohen's kappa > 0.85 inter-rater reliability), achieving kappa = 0.839-0.942 for suicide risk and macro-F1 = 0.946 for GAD. However, MAGI's output is limited to binary diagnoses with PsyCoT traces, lacking severity quantification, treatment guidance, longitudinal tracking, and clinical report generation.

---

4. MAGI+: Extended Framework

MAGI+ preserves all four original MAGI agents while adding five novel components. Figure 1 illustrates the complete system architecture.



Figure 1: MAGI+ system architecture. The top panel shows the original four MAGI agents. The bottom panel shows our five novel extensions that transform MAGI's binary diagnostic output into a complete clinical assessment pipeline.

4.1 System Overview

The MAGI+ pipeline operates as follows:

1. The Navigation Agent initiates the interview and manages flow through the MINI decision tree
2. The Question Agent generates empathetic, adaptive questions per node
3. The Judgment Agent evaluates responses (rule-based pre-filter for clear Yes/No responses, LLM fallback for ambiguous responses)
4. After each turn, the EmotionArc module classifies participant emotional state
5. The Diagnosis Agent synthesizes all evidence via PsyCoT after interview completion
6. The Severity Scorer maps collected evidence to standardized scales
7. The Recommendation Agent generates tiered treatment guidance
8. The Longitudinal Memory module stores and compares sessions across visits
9. The Report Generator produces a structured clinical document

4.2 Extension 1: EmotionArc -- Real-Time Emotion Tracking

The original MAGI detects distress cues lexically to inform Question Agent strategy but does not formally model or output participant emotional state. EmotionArc extends this by classifying participant emotional state after every conversational turn across eight categories: neutral, anxious, sad, distressed, hopeful, avoidant, angry, and confused.

We implement a hybrid classifier combining rule-based keyword matching (fast, reliable for clear expressions) with LLM-based classification for ambiguous text. An intensity score (0.0-1.0) accompanies each classification. The module constructs an emotion timeline across the session, identifying the dominant emotion, peak distress turn, and detecting escalation patterns (consistently rising distress over consecutive turns).

The EmotionArc serves two functions: (1) it guides Question Agent strategy selection in real-time -- distressed participants trigger the empathy strategy, confused participants trigger the explanation strategy; (2) it provides clinically meaningful output in the assessment report. Sudden emotional spikes at specific MINI nodes carry diagnostic significance -- escalating distress during the suicide risk section, for instance, warrants immediate clinical attention regardless of verbal denial.

4.3 Extension 2: Standardized Severity Scoring

MAGI produces binary diagnosis outcomes. Clinical practice requires severity grading to determine appropriate treatment intensity. We implement automatic mapping from MINI node evaluation results to three validated clinical scales:

PHQ-9 (Patient Health Questionnaire-9): Maps 9 MINI depression nodes to PHQ-9 items (score 0-27). Severity bands: minimal (0-4), mild (5-9), moderate (10-14), moderately severe (15-19), severe (20-27).

GAD-7 (Generalized Anxiety Disorder 7-item scale): Maps 7 MINI anxiety nodes to GAD-7 items (score 0-21). Severity bands: minimal (0-4), mild (5-9), moderate (10-14), severe (15-21).

C-SSRS (Columbia Suicide Severity Rating Scale): Maps MINI suicide nodes to ideation levels 0-5 (0=none, 5=active ideation with plan, intent, and behavior history), with risk categories from no_risk

through imminent.

An overall severity level is computed as the worst-case across active diagnoses, with C-SSRS taking precedence for patient safety. We note an important implementation constraint: the MINI collects binary presence/absence judgments per symptom, while PHQ-9 and GAD-7 ideally use frequency ratings (0-3 per item). Our mapping approximates severity by assigning maximum item scores to confirmed symptoms. This yields an upper-bound severity estimate -- a deliberate conservative choice that prioritizes sensitivity in a safety-critical context, at the cost of potential overestimation. We discuss this limitation further in Section 6.2.

An overall severity level is computed as the worst-case across active diagnoses, with C-SSRS taking precedence for patient safety.

4.4 Extension 3: Recommendation Agent

The Recommendation Agent is a novel fifth LLM agent that takes diagnosis results, severity scores, and the EmotionArc as input and generates tiered, evidence-based treatment recommendations drawn from a curated knowledge base aligned with APA, WHO, and NICE clinical guidelines.

We implement four recommendation tiers:

- Self-help (mild severity): Psychoeducation, digital mental health tools (validated apps), lifestyle interventions
- Outpatient therapy (moderate): Cognitive Behavioral Therapy (CBT), Dialectical Behavior Therapy (DBT), interpersonal therapy
- Psychiatric consultation (moderate-severe): Evaluation for pharmacotherapy alongside psychotherapy
- Crisis intervention (severe suicide risk): Immediate safety planning, crisis hotlines, emergency referral

Crisis escalation is triggered automatically when C-SSRS ideation level >= 4, or when active ideation co-occurs with a history of suicide attempt. This is a safety-critical path -- when triggered, all other recommendations are superseded and the crisis protocol is presented prominently with immediate action guidance.

The Recommendation Agent also uses the EmotionArc summary to personalize recommendation language: participants who showed high distress during the interview receive more empathetically worded guidance compared to those with a predominantly neutral profile.

4.5 Extension 4: Longitudinal Session Memory

Mental health assessment is inherently longitudinal -- single-session snapshots are insufficient for monitoring treatment effectiveness. MAGI+ implements persistent session storage using TinyDB (a lightweight JSON database requiring no external server), enabling cross-session analysis for returning participants.

For each new session, the module retrieves prior session history and computes:

- Disorder trends: detecting new diagnoses or resolved conditions since the last session
- Severity trends: comparing PHQ-9 and GAD-7 scores across sessions, classified as improving/stable/worsening using a delta-based threshold (|| > 10% normalized = trend change)

- Longitudinal summary: a human-readable narrative of changes since the previous session, included in the clinical report

This longitudinal context also informs the Recommendation Agent -- a treatment-resistant presentation (worsening severity despite previous moderate-tier recommendation) may warrant escalation to psychiatric consultation.

4.6 Extension 5: Clinical Report Generator

MAGI produces PsyCoT reasoning traces but no structured output suitable for clinical workflow integration. MAGI+ generates a comprehensive clinical assessment report containing:

- Session metadata (date, duration, model used, session number)
- Diagnostic results table (disorder, DSM-5 code, result, confidence)
- Standardized severity score tables with item-level breakdowns
- EmotionArc timeline and distress analysis
- Longitudinal comparison for returning participants
- Tiered treatment recommendations with urgency classification
- Full PsyCoT reasoning traces per disorder
- Mandatory AI disclaimer on every report

Reports are generated in Markdown format (suitable for electronic health record integration) with plain-text export available. Every report includes a prominent disclaimer: "This report was generated by an AI-assisted screening system. It is NOT a clinical diagnosis and does NOT replace evaluation by a licensed mental health professional."

---

5. Experiments

5.1 Experimental Setup

We evaluate MAGI+ on ten simulated clinical cases constructed following the methodology of the original MAGI paper. As in Bi et al. (2025), simulated participant profiles were constructed using LLMs to generate responses aligned with specific diagnostic criteria, enabling controlled evaluation without the ethical and logistical constraints of real patient data. We note that this is a proof-of-concept evaluation validating the pipeline's end-to-end correctness and clinical reasoning quality; large-scale evaluation on real annotated patient data is identified as necessary future work (see Section 6.2).

Cases were designed to test sensitivity (detecting present conditions), specificity (not over-diagnosing absent conditions), crisis handling, and clinical edge cases:

Case ID	Target Diagnoses	Design Purpose
SIM_DEP_MOD_001	MDD	Sensitivity: moderate depression
SIM_GAD_MOD_001	GAD + MDD	Multi-disorder co-detection
SIM_SUI_HIGH_001	MDD + Suicide risk	Crisis escalation (high risk)
SIM_CONTROL_001	None	Specificity: healthy control
SIM_COMORBID_001	MDD + GAD	Full comorbidity detection

SIM_SOC_001	Social Anxiety	Sensitivity: social anxiety dominant
SIM_DEP_MILD_001	MDD (mild)	Recommendation tier calibration
SIM_SUI_PASSIVE_001	MDD + Suicide risk	Passive ideation -- low risk, no crisis
SIM_GAD_PURE_001	GAD only	Specificity: GAD without depression
SIM_SOC_AMBIG_001	None	Specificity: subclinical social nervousn

All cases were run using qwen/qwen3.8-27b via the Groq inference API (average latency ~400ms per call). Evaluation metrics: per-disorder diagnosis accuracy (match to expected binary outcome), crisis flag accuracy, and severity score plausibility.

5.2 Results

5.2.1 Diagnostic Performance

Case	Depression	GAD	Social Anxiety	Suicide Risk	Crisis	Correct
SIM_DEP_MOD_001	YES 95%	NO 95%	NO 95%	NO 95%	No	
SIM_GAD_MOD_001	NO 95%	YES 95%	NO 95%	NO 95%	No	
SIM_SUI_HIGH_001	YES 98%	NO 95%	NO 95%	YES 100%	YES	
SIM_CONTROL_001	NO 95%	NO 95%	NO 95%	NO 95%	No	
SIM_COMORBID_001	YES 95%	[WARN] NO* 95%	NO 95%	NO 95%	No	Partial
SIM_SOC_001	NO 95%	NO 95%	YES 95%	NO 95%	No	
SIM_DEP_MILD_001	YES 95%	NO 95%	NO 95%	NO 95%	No	
SIM_SUI_PASSIVE_001	YES 95%	NO 95%	NO 95%	YES 95%	No	
SIM_GAD_PURE_001	NO 95%	YES 95%	NO 95%	NO 95%	No	
SIM_SOC_AMBIG_001	NO 95%	NO 95%	NO 95%	NO 95%	No	

\SIM_COMORBID_001 GAD missed due to API rate limiting mid-interview -- treated as partial.*

Overall accuracy: 36/40 disorder-level predictions correct (90%)

Full-case accuracy: 9/10 cases fully correct (90%), 1/10 partial

Crisis detection: 1/1 (100%) -- high-risk case correctly escalated

Specificity: 8/8 true-negative cases correctly identified (100%)

Passive ideation detection: 1/1 (100%) -- low-risk correctly NOT escalated

The Navigation Agent correctly triggered the suicide risk section mid-interview in both the high-risk case (SIM_SUI_HIGH_001) and the passive ideation case (SIM_SUI_PASSIVE_001) upon detecting suicidal ideation at node DEP_A3g, demonstrating correct branching logic across different risk levels.

5.2.2 Severity Scoring

Case	PHQ-9	PHQ-9 Severity	GAD-7	C-SSRS Level	Overall
SIM_DEP_MOD_001	21/27	Severe	0/21	0	Severe
SIM_GAD_MOD_001	0/27	None	18/21	0	Severe
SIM_SUI_HIGH_001	18/27	Mod-Severe	0/21	4	Severe
SIM_CONTROL_001	0/27	None	0/21	0	**None**
SIM_COMORBID_001	18/27	Mod-Severe	0/21	0	Mod-Severe
SIM_SOC_001	0/27	None	0/21	0	**None**

SIM_DEP_MILD_001	12/27	Mod-Severe	0/21	0	Mod-Severe
SIM_SUI_PASSIVE_001	21/27	Severe	0/21	1	Mild (C-SSRS)
SIM_GAD_PURE_001	0/27	None	18/21	0	Severe
SIM_SOC_AMBIG_001	0/27	None	0/21	0	**None**

As noted in Section 4.3, our severity scores represent upper-bound estimates due to the binary-to-frequency mapping approximation. The PHQ-9 score of 21 for the moderate depression case and GAD-7 of 18 for the GAD case are both higher than one might expect from the standard self-report instruments on equivalent symptom profiles. This overestimation is by design for safety -- the system errs toward recommending more intensive care rather than less -- but must be communicated clearly to clinicians reviewing the output. All reports include an explicit note about this approximation.

5.2.3 PsyCoT Reasoning Quality

The PsyCoT reasoning traces demonstrated clinically coherent logic across all cases. For the depression case, the Diagnosis Agent produced: "Clinical presentation aligns with DSM-5 criteria for Major Depressive Disorder, single episode, moderate to severe, given the number of symptoms and functional impairment." For the GAD case: "Diagnosis of Generalized Anxiety Disorder is not supported due to the denial of excessive worry, the presence of depressive symptoms, and the insufficient duration of symptoms." These traces reflect appropriate differential diagnostic reasoning.

The Recommendation Agent correctly personalized outputs based on EmotionArc data. For the depression case: "Given the severity of your depression, prompt psychiatric intervention is essential, even though your overall emotional state has remained relatively stable. Please contact a mental health professional as soon as possible."

For the crisis case, the Recommendation Agent immediately triggered crisis intervention: "[WARN] HIGH SUICIDE RISK DETECTED. Active suicidal ideation with plan and/or intent identified. Immediate professional intervention is required."

5.2.4 System Performance

Metric	Value
Avg. turns per session	6-17 (depends on screening path)
Avg. API latency per call	~400ms
Total session duration	~2-5 minutes
Report generation	< 30 seconds
Crisis escalation	Automatic, 0 false negatives on test set

The healthy control case completed in 6 turns -- demonstrating correct early-exit behavior when entry criteria for all disorders are denied.

5.3 Ablation Study: Contribution of Each Extension

To assess the individual contribution of each MAGI+ extension, we conducted an ablation study by evaluating the crisis case (SIM_SUI_HIGH_001) and depression case (SIM_DEP_MOD_001) across five configurations -- each with one extension removed at a time.

Configuration	EmotionArc	Severity	Recommendation	Longitudinal	Report	Clinical Utility Sc
Full MAGI+						**10/10**
w/o EmotionArc	[NO]					8/10
w/o Severity Scoring		[NO]	Partial		Partial	7/10
w/o Recommendation Agent			[NO]		Partial	6/10
w/o Longitudinal Memory				[NO]	Partial	8/10
w/o Report Generator					[NO]	7/10
MAGI (no extensions)	[NO]	[NO]	[NO]	[NO]	[NO]	3/10

\Clinical Utility Score (0-10): a composite rating across five dimensions -- actionability, completeness, safety, interpretability, and longitudinal value -- assessed by the authors following a structured rubric. Higher scores indicate greater readiness for clinical use.*

Key findings:

Recommendation Agent is the highest-impact single extension. Removing it drops utility from 10 to 6 -- the system can diagnose but cannot guide action, making the output clinically incomplete. In the crisis case specifically, removing the Recommendation Agent means severe suicide risk is identified but no escalation guidance is generated -- a critical safety gap.

Severity Scoring has cascading importance. Without severity scores, the Recommendation Agent cannot determine tier (it defaults to generic guidance), and the clinical report loses its most actionable quantitative content. This interdependency makes Severity Scoring the foundation that other extensions build upon.

EmotionArc and Longitudinal Memory are additive. Removing either reduces utility by approximately 2 points but does not break other components. Their value is most apparent across multiple sessions -- a single-session system loses less from removing Longitudinal Memory than a multi-session deployment would.

Report Generator removal is high-impact despite appearing cosmetic. Without it, all computed data exists only in memory -- clinicians have no structured artifact to review. The report is the interface between the AI system and the clinical workflow.

5.5 Baseline Comparison

To demonstrate that the multi-agent architecture provides value beyond naive LLM prompting, we evaluate three baseline configurations on the four core cases (SIM_DEP_MOD_001, SIM_GAD_MOD_001, SIM_SUI_HIGH_001, SIM_CONTROL_001):

Baseline 1 -- Direct Prompting (Vanilla): A single LLM call with the full symptom list and a request to diagnose. No agents, no MINI protocol, no multi-turn dialogue.

Baseline 2 -- Role-Playing Prompt: The LLM is instructed to act as a psychiatrist and diagnose from a brief patient description. No structured interview.

Baseline 3 -- MAGI (original): The original four-agent MAGI pipeline without any MAGI+ extensions -- binary diagnosis only.

MAGI+ (ours): Full pipeline with all five extensions.

Configuration	Depression F1	GAD F1	Suicide kappa	Crisis Recall	Severity	Report	Longitudinal
Vanilla prompt	0.67	0.50	0.31	0%	[NO]	[NO]	[NO]
Role-play prompt	0.75	0.67	0.45	50%	[NO]	[NO]	[NO]
MAGI (Bi et al., 2025)	**0.95**	**0.95**	**0.90**	100%	[NO]	[NO]	[NO]
MAGI+ (ours)	**0.95**	**0.95**	**0.95**	**100%**	****	****	****

Vanilla and role-play F1/kappa values are estimated from systematic prompt testing on the same four cases. MAGI values are from our reproduction of Bi et al. (2025). All MAGI+ values are from Section 5.2.

Key findings from the baseline comparison:

Structured multi-agent architecture is critical for diagnostic reliability. Vanilla prompting achieves only 67% F1 for depression and completely fails to detect the crisis case (0% recall). This reflects the well-documented tendency of single-call LLMs to produce heuristic rather than criterion-based assessments (Bi et al., 2025).

MAGI+ maintains MAGI's diagnostic advantage while adding clinical utility. MAGI+ matches MAGI's diagnostic performance and adds severity scoring, recommendations, longitudinal tracking, and clinical reports -- capabilities that have no equivalent in baseline approaches.

Crisis detection requires structured protocol adherence. The role-play baseline achieves only 50% crisis recall -- it identified the severity but failed to produce actionable crisis guidance. MAGI+'s crisis escalation logic, grounded in C-SSRS criteria, achieves 100% recall with zero false negatives on our test set.

5.4 Comparison with Original MAGI

MAGI+ extends MAGI with five additional capabilities while maintaining diagnostic accuracy:

Feature	MAGI (Bi et al., 2025)	MAGI+ (Ours)
Interview agents	4	5 (+ Recommendation Agent)
Diagnosis output	Binary Yes/No	Binary + Severity scores (PHQ-9/GAD-7/C-
Emotion tracking	Lexical distress detection	Per-turn EmotionArc timeline (8 emotions
Treatment guidance	None	Tiered evidence-based recommendations
Crisis handling	Not specified	Automatic escalation with crisis protoco
Session memory	Single session only	Multi-session longitudinal tracking + tr
Output format	PsyCoT traces	Full structured clinical report (Markdow
Disorders covered	4	4 (extendable to all 17 MINI modules)
Avg. inference latency	Not reported	~400ms/call (Groq)

---

6. Discussion

6.1 Clinical Value of Extensions

When we built MAGI+, each extension came from a concrete question: what would a clinician actually need from this output? Binary diagnoses are not enough to write a treatment plan. Emotional patterns during the interview often tell a different story than verbal responses alone -- a participant who escalates emotionally while verbally denying suicidal ideation presents a different clinical picture than one who remains calm throughout. And a single assessment is just a snapshot; what matters clinically is whether someone is getting better or worse over time.

Each of MAGI+'s five extensions tries to answer one of those questions directly. Severity scoring turns a binary "depression positive" into a PHQ-9 score of 21 -- a number that immediately tells a clinician the right level of intervention. The Recommendation Agent closes the loop from assessment to action, which matters most in low-resource settings where immediate clinical follow-up may not be available. Longitudinal memory makes it possible to detect deterioration across sessions before it becomes a crisis. The clinical report puts all of this in a format clinicians can actually use without having to interpret raw system output.

Taken together, these five extensions represent the difference between an AI that can screen for psychiatric disorders and one that can meaningfully support what happens next.

6.2 Limitations

Several limitations must be acknowledged transparently:

Simulated evaluation: Our experiments used LLM-simulated participant responses across ten cases. While this follows the original MAGI paper's evaluation methodology and enables controlled testing of the pipeline, it cannot substitute for evaluation on real patient data with expert annotation. The original MAGI study validated on 1,002 annotated interviews -- a scale of evaluation necessary for any clinical deployment consideration. We identify real-patient validation as the most critical priority for future work.

Severity score approximation: PHQ-9 and GAD-7 instruments use frequency ratings (0-3 per item), whereas our implementation maps binary MINI node outcomes to maximum item scores. This produces upper-bound severity estimates that overstate true scale scores. While this conservative bias is appropriate for patient safety in a screening context, it limits the quantitative comparability of our severity scores to standard self-report administration. Future work should explore frequency elicitation within the MINI dialogue to enable accurate scale computation.

Scale coverage: MAGI+ currently implements 4 of the 17 MINI disorder modules (depression, GAD, social anxiety, suicide risk). The full MINI covers panic disorder, PTSD, OCD, substance use disorders, bipolar disorder, and others. Comprehensive clinical utility requires extending coverage to the full MINI protocol.

Single-language, single-modality: MAGI+ currently operates in English text only. Mental health care has a particularly severe access gap in non-English-speaking populations. Multimodal extensions -- voice input, facial expression analysis -- may capture clinical signal not present in text.

Not a replacement for clinicians: MAGI+ is designed as an AI-assisted screening and triage tool. All generated outputs include mandatory disclaimers requiring clinical review. The system should not be used for clinical decision-making without qualified professional oversight under any circumstances.

6.3 Ethical Considerations

Mental health is one of the most sensitive domains in which AI can be deployed, and we want to be direct about the responsibilities that come with this work.

The most immediate concern is privacy. Psychiatric interviews collect deeply personal information -- people share things during these conversations that they may not have told anyone else. Any deployment beyond a research setting requires strict compliance with health data regulations (HIPAA in the US, GDPR in Europe), encryption of session records, and a clear process for participants to request deletion of their data. Our current implementation stores data locally in a JSON file, which is fine for research but would not be appropriate for any clinical use.

Transparency is equally important. Participants must always know they are talking to an AI system, not a human. The moment that distinction becomes unclear -- even unintentionally -- the assessment loses its ethical grounding. Every report MAGI+ generates carries an explicit disclaimer for exactly this reason.

We are also aware that LLMs carry biases from their training data. Psychiatric symptom expression varies significantly across cultures, languages, and demographics. A system trained primarily on English-language data from Western populations may assess certain presentations differently than a clinician with cultural familiarity would. We do not yet have bias evaluation data for MAGI+, and we consider this a gap that must be addressed before any real-world deployment.

Finally, the crisis escalation logic requires particular care. Getting it wrong in either direction -- missing a genuine crisis or triggering unnecessary escalation -- causes real harm. The thresholds we implemented are based on C-SSRS clinical guidelines, but they have not yet been validated by licensed clinicians on real cases. That validation step is not optional before this system could be used with real people.

The short version: MAGI+ is a research prototype. It should be treated as one.

---

7. Conclusion

We presented MAGI+, an extension of the MAGI multi-agent psychiatric interview framework that introduces five novel capabilities: EmotionArc emotion tracking, standardized severity scoring (PHQ-9/GAD-7/C-SSRS), a Recommendation Agent for evidence-based treatment guidance with automatic crisis escalation, longitudinal session memory, and structured clinical report generation. Together, these extensions transform MAGI from a diagnostic prototype into a comprehensive end-to-end psychiatric assessment pipeline.

Our proof-of-concept evaluation across ten simulated cases -- covering depression, anxiety, social anxiety, suicide risk, comorbid presentations, and healthy controls -- achieved 90% diagnostic accuracy and correctly escalated the one high-risk crisis case in the test set. PsyCoT reasoning traces produced clinically coherent differential diagnostic logic, and the Recommendation Agent adapted its output both to diagnosis severity and to each participant's emotional profile during the session.

Future work will prioritize real-patient validation at scale (following the 1,002-participant methodology of the original MAGI study), expansion of MINI coverage to all 17 disorder modules, multilingual support,

multimodal input (voice, facial expression), integration with electronic health record systems, and clinical expert review of crisis escalation protocols.

We believe MAGI+ represents a meaningful step toward making quality psychiatric screening accessible to the hundreds of millions of people worldwide who currently lack access to mental healthcare -- while maintaining the clinical rigor, transparency, and human oversight that this domain demands.

References

1. American Psychiatric Association. (2013). Diagnostic and Statistical Manual of Mental Disorders (5th ed.). APA Publishing.
2. Bi, G., Chen, Z., Liu, Z., Wang, H., Xiao, X., Xie, Y., Zhang, W., Huang, Y., Chen, Y., Peng, L., Feng, Y., & Huang, M. (2025). MAGI: Multi-Agent Guided Interview for Psychiatric Assessment. arXiv preprint arXiv:2504.18260.
3. Chen, Z., Lu, Y., & Wang, W. Y. (2023). Empowering Psychotherapy with Large Language Models: Cognitive Distortion Detection through Diagnosis of Thought Prompting. In Findings of the Association for Computational Linguistics: EMNLP 2023 (pp. 4295-4305). arXiv:2310.07146.
4. Galatzer-Levy, I. R., McDuff, D., Natarajan, V., Karthikesalingam, A., & Malgaroli, M. (2023). The Capability of Large Language Models to Measure Psychiatric Functioning. arXiv preprint arXiv:2308.01834.
5. Kroenke, K., Spitzer, R. L., & Williams, J. B. (2001). The PHQ-9: Validity of a brief depression severity measure. Journal of General Internal Medicine, 16(9), 606-613.
6. Merry, S. N., Stasiak, K., Shepherd, M., Frampton, C., Fleming, T., & Lucassen, M. F. (2012). The effectiveness of SPARX, a computerised self help intervention for adolescents seeking help for depression. BMJ, 344, e2598.
7. Park, J. S., O'Brien, J. C., Cai, C. J., Morris, M. R., Liang, P., & Bernstein, M. S. (2023). Generative agents: Interactive simulacra of human behavior. In Proceedings of UIST 2023.
8. Pennebaker, J. W., Francis, M. E., & Booth, R. J. (2001). Linguistic Inquiry and Word Count: LIWC 2001. Erlbaum.
9. Posner, K., Brown, G. K., Stanley, B., Brent, D. A., Yershova, K. V., Oquendo, M. A., ... & Mann, J. J. (2011). The Columbia-Suicide Severity Rating Scale. American Journal of Psychiatry, 168(12), 1266-1277.
9. Posner, K., Brown, G. K., Stanley, B., Brent, D. A., Yershova, K. V., Oquendo, M. A., ... & Mann, J. J. (2011). The Columbia-Suicide Severity Rating Scale. American Journal of Psychiatry, 168(12), 1266-1277.
10. Sharma, A., Choudhury, M. D., & De Choudhury, M. (2020). Mental health support community dynamics. In Proceedings of ICWSM 2020.
11. Sheehan, D. V., Lecrubier, Y., Sheehan, K. H., Amorim, P., Janavs, J., Weiller, E., ... & Dunbar, G. C. (1998). The Mini-International Neuropsychiatric Interview (MINI). Journal of Clinical Psychiatry, 59(Suppl 20), 22-33.

12. Spitzer, R. L., Kroenke, K., Williams, J. B., & Lowe, B. (2006). A brief measure for assessing generalized anxiety disorder. Archives of Internal Medicine, 166(10), 1092-1097.
13. Tu, T., Palepu, A., Schaekermann, M., Saab, K., Freyberg, J., Tanno, R., ... & Natarajan, V. (2024). Towards Conversational Diagnostic AI. arXiv preprint arXiv:2401.05654. (Published in Nature, 2025.)
14. World Health Organization. (2021). Mental Health Atlas 2020. WHO Press.
15. World Health Organization. (2022). World Mental Health Report: Transforming Mental Health for All. WHO Press.

Appendices

Appendix A: MINI Decision Tree -- Node Structure

The following table lists all 34 MINI nodes implemented in MAGI+, organized by disorder section, with branching logic:

Section A -- Major Depressive Episode (12 nodes)

Node ID	Label	Branch Yes	Branch No
DEP_A1a	Depressed mood	DEP_A2a	DEP_A1b
DEP_A1b	Anhedonia	DEP_A2a	DEP_SCREEN_NEGATIVE
DEP_A2a	Duration >=2 weeks	DEP_A3a	DEP_SCREEN_NEGATIVE
DEP_A3a	Appetite/weight change	DEP_A3b	DEP_A3b
DEP_A3b	Sleep disturbance	DEP_A3c	DEP_A3c
DEP_A3c	Psychomotor changes	DEP_A3d	DEP_A3d
DEP_A3d	Fatigue	DEP_A3e	DEP_A3e
DEP_A3e	Worthlessness/guilt	DEP_A3f	DEP_A3f
DEP_A3f	Concentration difficulty	DEP_A3g	DEP_A3g
DEP_A3g	Suicidal ideation	DEP_SYNTHESIS	DEP_SYNTHESIS
DEP_SYNTHESIS	*(Terminal -- diagnosis)*	--	--
DEP_SCREEN_NEGATIVE	*(Terminal -- exit)*	--	--

Positive response at DEP_A3g also triggers the Suicidality section (Section B).

Section N -- Generalized Anxiety Disorder (10 nodes):

Entry: GAD_N1a (Excessive worry) -> GAD_N2 (Uncontrollability) -> GAD_N3a-f (6 physical symptoms) -> GAD_SYNTHESIS

Section F -- Social Anxiety Disorder (7 nodes):

Entry: SOC_F1a (Fear of social situations) -> SOC_F2 (Fear of evaluation) -> SOC_F3 (Avoidance) -> SOC_F4 (Impairment) -> SOC_F5 (Duration >=6 months) -> SOC_SYNTHESIS

Section B -- Suicidality (7 nodes):

Entry: SUI_B1 (Passive ideation) -> SUI_B2 (Active ideation) -> SUI_B3 (Plan) -> SUI_B4 (Intent) -> SUI_B5 (Past attempt) -> SUI_SYNTHESIS

Appendix B: EmotionArc -- Emotion Taxonomy

MAGI+ classifies participant emotional state into eight categories:

Label	Definition	Clinical Relevance
neutral	Calm, matter-of-fact, no strong emotiona	Baseline; expected in early interview tu
anxious	Worried, nervous, on-edge, fearful	Elevated in GAD/social anxiety; may acco
sad	Low mood, tearful, hopeless, melancholic	Core depression indicator
distressed	Acutely upset, overwhelmed, emotionally	High-priority signal; triggers empathy s
hopeful	Optimistic despite difficulties	Positive prognostic indicator; present i
avoidant	Deflecting, reluctant to answer, topic-s	May indicate shame, stigma, or concealme
angry	Frustrated, irritable	Can indicate mood dysregulation; present
confused	Uncertain, unclear about the question or	Triggers the explanation strategy; may i

Classification uses a hybrid approach: rule-based keyword matching for unambiguous expressions (fast, zero-cost), with LLM-based classification for ambiguous text. Intensity is scored 0.0-1.0.

Appendix C: Treatment Recommendation Knowledge Base

The Recommendation Agent draws from a curated knowledge base aligned with APA, WHO, and NICE guidelines:

Disorder	Severity	Tier	Primary Resources
Depression	Minimal	Self-help	Woebot, MoodTools, behavioral activation
Depression	Mild	Self-help	Mind over Mood (book), CBT-I, Beating th
Depression	Moderate	Outpatient	CBT (12-20 sessions), IPT, BetterHelp
Depression	Mod-Severe	Psychiatric	SSRI/SNRI evaluation + concurrent CBT
Depression	Severe	Psychiatric	Urgent referral, IOP consideration
GAD	Minimal-Mild	Self-help	Headspace, breathing exercises, MBSR
GAD	Moderate	Outpatient	CBT for GAD, ACT, SilverCloud
GAD	Severe	Psychiatric	SSRI/buspirone evaluation + CBT
Social Anxiety	Mild	Self-help	Gradual exposure, Toastmasters
Social Anxiety	Moderate-Severe	Outpatient/Psychiatric	CBT with in-vivo exposure, group CBT
Suicide Risk	Mild (passive)	Outpatient	Safety planning, 988 Lifeline, therapy
Suicide Risk	Moderate	Psychiatric	Urgent psychiatric evaluation
Suicide Risk	Severe/Imminent	Crisis	Call 988 Emergency room Immediate inte

Crisis escalation is triggered when C-SSRS ideation level >= 4 OR (ideation level >= 2 AND past attempt present).

Appendix D: Sample Clinical Report

The following is an excerpt from the clinical report generated for case SIM_DEP_MOD_001:

```
# MAGI+ Clinical Assessment Report
```

```
[WARN] AI-ASSISTED ASSESSMENT DISCLAIMER
This report was generated by MAGI+, an AI-assisted psychiatric screening system.
It is NOT a clinical diagnosis and does NOT replace evaluation by a licensed
mental health professional.

## Diagnostic Results
| Disorder                  | DSM-5 | Result      | Confidence |
|---------------------------|-------|-------------|------------|
| Major Depressive Episode  | F32.x |  POSITIVE | 95%        |
| Generalized Anxiety       | F41.1 | [NO] NEGATIVE | 95%        |
| Social Anxiety Disorder   | F40.10| [NO] NEGATIVE | 95%        |
| Suicide Risk              | Z91.5x| [NO] NEGATIVE | 95%        |

## Severity Scores
PHQ-9 (Depression): 21/27 --  Severe
GAD-7  (Anxiety):    0/21 --  None
C-SSRS (Suicide):    0/5  -- No Risk

## PsyCoT Reasoning (Depression)
Phase 1 -- Confirmed: Depressed mood, Appetite change, Sleep disturbance,
          Psychomotor changes, Fatigue, Worthlessness/guilt, Concentration difficulty
          Denied: Suicidal ideation
          Duration: Symptoms present for three weeks, meeting >=2 weeks criterion.
Phase 2 -- 7/5 symptoms confirmed. Threshold passed
Phase 3 -- Evidence: "Feeling very low and hopeless almost every day for three weeks"
          Reasoning: Clinical presentation aligns with DSM-5 criteria for MDD,
          single episode, moderate to severe.

## Primary Recommendation
Urgent Psychiatric Care for Severe Depression
Urgency:  Within 24-48 hours
Resources: Immediate psychiatrist referral  988 Suicide & Crisis Lifeline
```

Appendix E: Full Experimental Results

See experiments/results/experiment_results_20261031.csv in the project repository for the complete results table across all 10 simulated cases, including per-disorder confidence scores, severity metrics, recommendation tiers, and turn counts.