# GATE Assignment 2

# Perambuduri Srikaran - AI20BTECH11018

Download all python codes from

https://github.com/srikaran-p/EE3900/tree/main/ GATE Assignment2/codes

Download all latex codes from

https://github.com/srikaran-p/EE3900/tree/main/ GATE Assignment2

## **PROBLEM**

(GATE EC-2006 Q.14) Let  $x(t) \leftrightarrow X(f)$  be Fourier Transform pair. The Fourier Transform of the signal x(5t-3) in terms of X(f) is given as

(A) 
$$\frac{1}{5}e^{-\frac{j6\pi f}{5}}X\left(\frac{f}{5}\right)$$

(B) 
$$\frac{1}{5}e^{\frac{j6\pi f}{5}}X(\frac{f}{5})$$

(B) 
$$\frac{1}{5}e^{\frac{j6\pi f}{5}}X\left(\frac{f}{5}\right)$$
  
(C)  $\frac{1}{5}e^{j6\pi f}X\left(\frac{f}{5}\right)$   
(D)  $\frac{1}{5}e^{j6\pi f}X\left(\frac{f}{5}\right)$ 

(D) 
$$\frac{1}{5}e^{j6\pi f}X\left(\frac{f}{5}\right)$$




Fig. 4: Plot of rect(5t-3)

### SOLUTION

By the time scaling property of Fourier transform,

$$x(\alpha t) \stackrel{\mathcal{F}}{\rightleftharpoons} \frac{1}{|\alpha|} X \left( \frac{f}{|\alpha|} \right)$$
 (0.0.1)

$$x(5t) \stackrel{\mathcal{F}}{\rightleftharpoons} \frac{1}{5}X\left(\frac{f}{5}\right)$$
 (0.0.2)

By the time shifting property of Fourier transform,

$$x(t-t_0) \stackrel{\mathcal{F}}{\rightleftharpoons} e^{-j2\pi f t_0} X(f)$$
 (0.0.3)

$$x\left(5\left(t-\frac{3}{5}\right)\right) \stackrel{\mathcal{F}}{\rightleftharpoons} \frac{1}{5}e^{-\frac{j6\pi f}{5}}X\left(\frac{f}{5}\right)$$
 (0.0.4)

Let x(t) = rect(t) and  $\text{sinc}(t) = \frac{\sin(t)}{t}$ 

$$X(f) = \operatorname{sinc}(f) \tag{0.0.5}$$

For x(t) = rect(5t - 3)

$$\frac{1}{5}e^{-\frac{j6\pi f}{5}}X\left(\frac{f}{5}\right) = \frac{1}{5}e^{-\frac{j6\pi f}{5}}\operatorname{sinc}\left(\frac{f}{5}\right)$$
 (0.0.6)

The correct option is (A).




Fig. 4: Fourier transform Simulation v/s Theoretical