t-Table Cutoff Values

For confidence intervals and two-tailed hypothesis tests, find the risk column of interest, α, at the top of the table; for one-tailed tests, find the risk column at the bottom. The cutoff value is then found at the intersection of the risk column and the *df* row.



Example: For a confidence interval or two-tailed test where $df = 10$ and α is to be .05, the cutoffs are $t = \pm 2.228$.



Example: For an upper-tail H_a with $\alpha = .10$ and $df = 9$, the cutoff is $t = 1.383$. Since all values are given as positive, a minus sign needs to be provided if H_a is lower tail.

Degrees of Freedom, *df*	Two-Tailed α (Probability in Both Tails Combined)					
	.20	.10	.05	.02	.01	.001
1	3.078	6.314	12.706	31.821	63.657	636.619
2	1.886	2.920	4.303	6.965	9.925	31.598
3	1.638	2.353	3.182	4.541	5.841	12.941
4	1.533	2.132	2.776	3.747	4.604	8.610
5	1.476	2.015	2.571	3.365	4.032	6.859
6	1.440	1.943	2.447	3.143	3.707	5.959
7	1.415	1.895	2.365	2.998	3.499	5.405
8	1.397	1.860	2.306	2.896	3.355	5.041
9	1.383	1.833	2.262	2.821	3.250	4.781
10	1.372	1.812	2.228	2.764	3.169	4.587
11	1.363	1.796	2.201	2.718	3.106	4.437
12	1.356	1.782	2.179	2.681	3.055	4.318
13	1.350	1.771	2.160	2.650	3.012	4.221
14	1.345	1.761	2.145	2.624	2.977	4.140
15	1.341	1.753	2.131	2.602	2.947	4.073
16	1.337	1.746	2.120	2.583	2.921	4.015
17	1.333	1.740	2.110	2.567	2.898	3.965
18	1.330	1.734	2.101	2.552	2.878	3.922
19	1.328	1.729	2.093	2.539	2.861	3.883
20	1.325	1.725	2.086	2.528	2.845	3.850
21	1.323	1.721	2.080	2.518	2.831	3.819
22	1.321	1.717	2.074	2.508	2.819	3.792
23	1.319	1.714	2.069	2.500	2.807	3.767
24	1.318	1.711	2.064	2.492	2.797	3.745
25	1.316	1.708	2.060	2.485	2.787	3.725
26	1.315	1.706	2.056	2.479	2.779	3.707
27	1.314	1.703	2.052	2.473	2.771	3.690
28	1.313	1.701	2.048	2.467	2.763	3.674
29	1.311	1.699	2.045	2.462	2.756	3.659
≥ 30	1.282	1.645	1.960	2.326	2.576	3.291
	.10	.05	.025	.01	.005	.0005
	One-Tailed α (Probability in One Tail Only)					

Source: Computed by R. E. Shiffler and A. J. Adams