# Inner Product Spaces

Sriram Vadlamani

February 3, 2020

## Contents

|   | Introduction                               | 3 |
|---|--|---|
|   | 1.1 A bilinear Form                        | 3 |
| 2 | Inner product                              | 3 |
|   | 2.1 Symmetry                               | 3 |
|   | 2.2 Positive definite                      | 3 |
| 3 | Theorems                                   | 3 |
|   | 3.1 Cauchy-Schwartz and Minkowski Theorems | 3 |
| 1 | Orthogonality                              | 4 |
|   | 4.1 Pythagorean Theorem                    | 4 |
|   | 4.2 The orthogonal of a subspace           |   |

#### 1 Introduction

#### 1.1 A bilinear Form

let E be an R-vs and  $\theta: E \times E \to R$ , we say,  $\theta$  is a bilinear form if:  $\forall (u, v, w) \in E^2$  and  $\forall a \in R$ ,

- $\theta(u+v,w) = \theta(u,v) + \theta(u,w)$
- $\theta(au, v) = a\theta(u, v)$

#### Proposition 1

let E be an R-vs of finite dimension 'n'. We have  $B=(e_1,e_2,\,\dots\,,\,e_n)$  the basis of E.

 $\theta: E \times E \to R$  a bilinear form.

Then  $\forall (x,y) \in E^2, \theta(x,y) = X^t \cdot M \cdot Y$ 

Where M is the matrix of the bilinear form defined as:

 $M = \theta(e_i, e_i)$ 

And X and Y are the coordinates of 'x' and 'y'.

## 2 Inner product

A space  $(E, \theta)$  is said to be an inner product space iff

- The bilinear form is 'symmetric'
- The form is positive definite.

#### 2.1 Symmetry

A bilinear form is symmetric iff  $\forall (x,y) \in E^2$  $\theta(x,y) = \theta(y,x)$ 

#### 2.2 Positive definite

A bilinear form is said to be positive definite iff  $\forall x \in E, \theta(x, x) \geq 0$ , and  $\forall x \in E, \theta(x, x) = 0 \Longrightarrow x = 0$ 

#### 3 Theorems

#### 3.1 Cauchy-Schwartz and Minkowski Theorems

Cauchy-Schwartz: Let E be an R-vs and  $\theta: E \to R$  a positive definite and symmetric bilinear form, then

$$\forall (x,y) \in E^2 \mid \theta(x,y) \mid \leq \sqrt{\theta(x,x)} \times \sqrt{\theta(y,y)}$$

**Minkowski's:** Let  $(E, \theta)$  be an inner product space on R, then:  $\forall (x,y) \in E^2, \ \sqrt{\theta(x+y,x+y)} \leq \sqrt{\theta(x,x)} + \sqrt{\theta(y,y)}$ 

## 4 Orthogonality

We call  $N: E \to R$  a norm  $\forall (x,y) \in E^2$  and  $\forall \lambda \in R$  we have:

- $N(x) \geq 0$
- $N(\lambda \cdot x) = \lambda \cdot N(x)$
- $N(x) = 0 \iff x = 0$
- $N(x+y) \leq N(x) + N(y)$

and we say that 'N' is a norm. In geometry, this is what we call a triangular inequality.

**Proposition 2** The norm for any vector  $x \in E$  is  $\sqrt{\theta(x,x)}$ .

#### 4.1 Pythagorean Theorem

for  $(E, \theta)$  an inner product space, and  $\forall (x, y) \in E^2$  such that  $\langle x, y \rangle = 0$ , we have:

$$||x + y||^2 = ||x||^2 + ||y||^2$$

#### 4.2 The orthogonal of a subspace

For  $(E,\theta)$  an inner product space,  $A^{\perp}$  is defined as:  $\{\forall x\in E, \forall y\in A, < x,y>=0\}$  where  $A\subset E$ 

Some important remarks are:

- $A \subset B \Rightarrow B^{\perp} \subset A^{\perp}$
- $A^{\perp} = span(A)^{\perp}$
- $A \cap A^{\perp} \subset \{0\}$