



Food and Agriculture
Organization of the
United Nations



World Health
Organization



REPORT 2021

Pesticide residues in food

Extra Joint FAO/WHO Meeting
on Pesticide Residues



Report 2021

Pesticide Residues in Food

Extra Joint FAO/WHO Meeting on Pesticide Residues

Report of the extra Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Core Assessment Group on Pesticide Residues was held virtually over two sessions from 17 to 21 May and from 7 to 11 June 2021.

Required citation:

FAO and WHO. 2021. *Report 2021 – Pesticide residues in food. Extra Joint FAO/WHO Meeting on Pesticide Residues*. Rome.
<https://doi.org/10.4060/cb6975en>

The designations employed and the presentation of material in this information product do not imply the expression of any opinion whatsoever on the part of the Food and Agriculture Organization of the United Nations (FAO) or the World Health Organization (WHO) concerning the legal or development status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. The mention of specific companies or products of manufacturers, whether or not these have been patented, does not imply that these have been endorsed or recommended by FAO or WHO in preference to others of a similar nature that are not mentioned.

The views expressed in this information product are those of the author(s) and do not necessarily reflect the views or policies of FAO or WHO.

ISBN [FAO] 978-92-5-135028-7

ISBN [WHO] 978-92-4-003769-4 (electronic version)

ISBN [WHO] 978-92-4-003770-0 (print version)

© FAO and WHO, 2021



Some rights reserved. This work is made available under the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 IGO licence (CC BY-NC-SA 3.0 IGO; <https://creativecommons.org/licenses/by-nc-sa/3.0/igo/legalcode>).

Under the terms of this licence, this work may be copied, redistributed and adapted for non-commercial purposes, provided that the work is appropriately cited. In any use of this work, there should be no suggestion that FAO or WHO endorses any specific organization, products or services. The use of the FAO or WHO logo is not permitted. If the work is adapted, then it must be licensed under the same or equivalent Creative Commons licence. If a translation of this work is created, it must include the following disclaimer along with the required citation: "This translation was not created by the Food and Agriculture Organization of the United Nations (FAO) or the World Health Organization (WHO). Neither FAO nor WHO is responsible for the content or accuracy of this translation. The original English edition shall be the authoritative edition.

Disputes arising under the licence that cannot be settled amicably will be resolved by mediation and arbitration as described in Article 8 of the licence except as otherwise provided herein. The applicable mediation rules will be the mediation rules of the World Intellectual Property Organization <http://www.wipo.int/amc/en/mediation/rules> and any arbitration will be conducted in accordance with the Arbitration Rules of the United Nations Commission on International Trade Law (UNCITRAL).

Third-party materials. Users wishing to reuse material from this work that is attributed to a third party, such as tables, figures or images, are responsible for determining whether permission is needed for that reuse and for obtaining permission from the copyright holder. The risk of claims resulting from infringement of any third-party-owned component in the work rests solely with the user.

Sales, rights and licensing. FAO information products are available on the FAO website (www.fao.org/publications) and can be purchased through publications-sales@fao.org. Requests for commercial use should be submitted via: www.fao.org/contact-us/licence-request. Queries regarding rights and licensing should be submitted to: copyright@fao.org.

Contents

List of participants	v
Abbreviations.....	vii
Use of JMPR reports and evaluations by registration authorities	ix
1. Introduction.....	1
1.1. Declaration of interests	1
2. General considerations	3
3. Responses to specific concerns raised by the Codex Committee on Pesticide Residues (CCPR).....	7
4. Dietary exposure assessment for pesticide residues in food.....	9
4.1. Long-term dietary exposure	9
4.2. Acute dietary exposure	10
5. Evaluation of data for acceptable daily intake and acute reference dose for humans, maximum residue levels and supervised trials median residue values.....	11
5.1. ACETAMIPRID (246)	11
5.2. BIXAFEN (262)	13
5.3. CHLORPYRIFOS (017) (R).....	29
5.4. CLOFENTEZINE (156)	31
5.5. CLOTHIANIDIN (238)	35
5.6. CYPERMETHRINS (including alpha- and zeta-cypermethrin) (118).....	47
5.7. CYPRODINIL (207)	49
5.8. DIFENOCONAZOLE (224)	53
5.9. ETHION (034)	59
5.10. ETHIPROLE (304)	61
5.11. FENBUCONAZOLE (197)	65
5.12. FENHEXAMID (215)	67
5.13. FENPICOXAMID (305).....	71
5.14. FLUOPYRAM (243)	79
5.15. IMAZALIL (110)	83
5.16. ISOPROTHIOLANE (299)	87
5.17. ISOXAFLUTOLE (268)	93
5.18. MANDIPROPAMID (231)	95
5.19. METHOPRENE (147)	101
5.20. METHOXYFENOZIDE (209).....	103
5.21. PROTHIOCONAZOLE (232).....	107
5.22. PYDIFLUMETOFEN (309)	115
5.23. QUINOXYFEN (222)	129

5.24.	SPINETORAM (233)	133
5.25.	SPIROMESIFEN (294).....	137
5.26.	SULFOXAFLOL (252)	139
5.27.	TEBUCONAZOLE (189).....	143
5.28.	THIAMETHOXAM (245).....	145
5.29.	TRIFLOXYSTROBIN (213)	155
5.30.	TRINEXAPAC-ETHYL (271).....	165
Annex 1:	Acceptable daily intakes, acute reference doses, recommended maximum residue levels, supervised trials median residue values and other values recorded by the 2021 Extra JMPR Meeting.....	177
Annex 2:	Index of reports and evaluations of pesticides by the JMPR	187
Annex 3:	International estimated daily intakes (IEDIs) of pesticide residues.....	203
Annex 4:	International Estimate of Short-Term Intakes (IESTIs) of pesticide residues.....	395
Annex 5:	Reports and other documents resulting from previous joint meetings of the FAO panel of experts on pesticide residues in food and the environment and the WHO core assessment group on pesticide residues.....	421
Annex 6:	Livestock dietary burdens	431

List of participants

FAO Experts

- Dr Christos Anagnostopoulos**, Laboratory of Pesticide Residues, Benaki Phytopathological Institute, Athens, 14561 Greece
- Dr Michael Doherty**, Office of Pesticide Programs, Health Effects Division, United States Environmental Protection Agency, Arlington, VA 22202, USA.
- Mr Jonathon Giordano**, Office of Pesticide Programs, Health Effects Division, United States, Environmental Protection Agency, Arlington, VA 22202, USA
- Ms Monica Le**, Pest Management Regulatory Agency, Health Canada, Ottawa, Ontario K1A 0K9, Canada
- Dr Mi-Gyung Lee**, Dept. of Food Science & Biotechnology Andong National University, Gyeongsangbuk-do, 36729, Republic of Korea
- Mr David Lunn**, Plants & Pathways Directorate, Ministry for Primary Industries, Wellington, New Zealand.
- Dr Arion Leahigh**, Office of Pesticide Programs, Health Effects Division, United States Environmental Protection Agency, MS 7509C, Washington, DC 20460, USA
- Dr Dugald MacLachlan**, Australian Government Department of Agriculture, Water and the Environment, Canberra, ACT 2601, Australia
- Dr Rebecca Scrivens**, Chemicals Regulation Division, Health & Safety Executive, Mallard House, Kings Pool, 3 Peasholme Green, York, YO1 7PX United Kingdom
- Mr Christian Sieke**, Federal Institute for Risk Assessment, Unit Residues and Analytical Methods, Department Pesticide Safety, Max-Dohrn-Strasse 8–10, 10589 Berlin, Germany (Chairman)
- Ms Monique Thomas**, Pest Management Regulatory Agency, Health Canada, Ottawa, Ontario K1A 0K9, Canada
- Dr Yukiko Yamada**, International Food Safety Consultant, Adjunct Professor, Azabu University; Guest Scholar, National Institute of Health Sciences; Tokyo 1070062, Japan
- Dr Guibiao Ye**, Institute for the Control of Agrochemicals, Ministry of Agriculture and Rural Affairs, Maizidian 22, Chaoyang District, Beijing 100125, People's Republic of China

WHO Experts

- Dr Ian Dewhurst**, Leavening, North Yorkshire YO17 9SA, United Kingdom (Co- Chair)
- Dr Elizabeth Mendez**, Office of Pesticide Programs, Health Effects Division, United States Environmental Protection Agency, Washington DC 20460, United States of America
- Dr Luca Tosti**, Department of Biomedical and Clinical Sciences - University of Milan, University Hospital Luigi Sacco, 20157 Milano, Italy
- Dr Midori Yoshida**, Food Safety Commission of Japan, Tokyo 107-6122, Japan
- Dr Rhian Cope**, Australian Pesticides and Veterinary Medicines Authority, Australia
- Dr Prakashchandra (PV) Shah**, Retired from United States Environmental Protection Agency, United States of America

JMPR Secretariat

Mr Kevin Bodnaruk, West Pymble, NSW 2073, Australia (*FAO Editor*)

Dr Jeevan Khurana, Lyneham, ACT 2602, Australia (*FAO Editor*)

Ms Nora Lune, Department of Food Safety and Zoonoses (FOS), World Health Organization, 1211 Geneva 27, Switzerland (*WHO JMPR Secretariat*)

Mr Soren Madsen, Department of Food Safety and Zoonoses (FOS), World Health Organization, 1211 Geneva 27, Switzerland (*WHO JMPR Secretariat*)

Dr Russell Parry, Shrewsbury SY2 6HZ United Kingdom (*WHO Editor*)

Mr Kim Petersen Department of Food Safety and Zoonoses (FOS), World Health Organization, 1211 Geneva 27, Switzerland (*WHO JMPR Secretariat*)

Dr Xiongwu Qiao, Shanxi Academy of Agricultural Sciences, 81 Longcheng Street, Taiyuan, Shanxi, 030031, China (*CCPR Chairman*)

Ms Ho Ngai Yin, Department of Nutrition and Food Safety World Health Organization, 1211 Geneva 27, Switzerland (*WHO JMPR Secretariat*)

Ms Yong Zhen Yang, Plant Production and Protection Division, Food and Agriculture Organization of the United Nations, Viale delle Terme di Caracalla, 00153 Rome, Italy (*FAO JMPR Secretariat*)

Abbreviations

ADI	acceptable daily intake
AGF	aspirated grain fractions
AR	applied radioactivity
As	as received
ARfD	acute reference dose
BBCH	Biologische Bundesanstalt, Bundessortenamt und Chemische Industrie
bw	body weight
CAC	Codex Alimentarius Commission
CAS	Chemical Abstracts Service
CCPR	Codex Committee on Pesticide Residues
cGAP	critical GAP
DALA	days after last application
DAT	days after treatment
DM	dry matter
DT ₅₀	time required for 50% dissipation of the initial concentration
Dw	dry weight
EFSA	European Food Safety Authority
eq	equivalent(s)
ESI	electrospray ionization
EU	European Union
FAO	Food and Agriculture Organization of the United Nations
GAP	good agricultural practice
GC-ECD	gas chromatography – electron capture detector
GC-FPD	gas chromatography – flame photometric detector
GC-MS	gas chromatography – mass spectrometry
GEMS	Global Environment Monitoring System – Food Contamination Monitoring and Assessment Programme
GI	gastrointestinal tract
GLP	good laboratory practice
HPLC	high performance liquid chromatography
HR	highest residue level in the edible portion of a commodity
HR-P	highest residue level in a processed commodity
IEDI	International Estimated Daily Intake
IESTI	International Estimate of Short-Term Dietary Intake
ILV	independent laboratory validation
ISO	International Organization for Standardization

IUPAC	International Union of Pure and Applied Chemistry
JECFA	Joint FAO/WHO Expert Committee on Food Additives
JMPR	Joint FAO/WHO Meeting on Pesticide Residues
LC-MS/MS	liquid chromatography-tandem mass spectrometry
LOD	limit of detection
LOQ	limit of quantification
MRL	maximum residue limit
NOAEL	no-observed-adverse-effect level
OECD	Organisation for Economic Co-operation and Development
PBI	plant-back interval
PF	processing factor
PHI	pre-harvest interval
Po	post-harvest
ppm	parts per million
QuEChERS	quick, easy, cheap, effective, rugged and safe
QSAR	quantitative structure–activity relationship
RAC	raw agricultural commodity
RSD	relative standard deviation
RTI	re-treatment interval
SC	suspension concentrate
SPE	solid phase extraction
STMR	supervised trials median residue
STMR-P	supervised trials median residue in a processed commodity
TRR	total radioactive residues
TTC	threshold of toxicological concern
UK	United Kingdom
USA	United States of America
WHO	World Health Organization

Use of JMPR reports and evaluations by registration authorities

Most of the summaries and evaluations contained in this report are based on unpublished proprietary data provided for use by JMPR in making its assessments. A registration authority should not grant a registration on the basis of an evaluation unless it has first received authorisation for such use from the owner of the data provided for the JMPR review or has received the data on which the summaries are based, either from the owner of the data or from a second party that has obtained permission from the owner of the data for this purpose.

Pesticide residues in food

Report of the 2021 Extra Joint FAO/WHO Meeting on Pesticide Residues

1. Introduction

An Extra Joint Meeting of the Food and Agriculture Organization of the United Nations (FAO) Panel of Experts on Pesticide Residues in Food and the Environment and the World Health Organization (WHO) Core Assessment Group on Pesticide Residues (JMPR) was held virtually over two sessions from 17 to 21 May and 7 to 11 June.

The meeting was opened by the JMPR Secretariats, Madame YongZhen Yang of FAO and Dr Kim Peterson of the WHO. Madame Yang welcomed and conveyed her appreciation to the panel members for their efforts in participating in preparatory virtual sessions prior the current JMPR Meeting. Dr Peterson also thanked the participants for providing their expertise and for devoting significant time and effort to the important work of the JMPR. He highlighted that holding the current Meeting virtually was an important initiative but that the practice would not be unfamiliar to participants and that despite the challenge of significantly different time zones, was confident the goals of the Meeting would be achieved.

In concluding Madame Yang referenced recent remarks of the FAO Director General Dr Qu Dongyu that the challenges of the pandemic, has meant that new modes of working and innovative systems have had to be created in adapting to this new normal and that working remotely is now part of our daily working life. The current virtual meeting will be an opportunity for the JMPR to identify possibilities of working remotely.

During the meeting, the FAO Panel of Experts on Pesticide Residues in Food was responsible for reviewing residue and analytical aspects of the pesticides under consideration, including data on their metabolism, fate in the environment and use patterns for estimating the maximum levels of residues that might occur as a result of use of the pesticides according to good agricultural practice. The methodologies are described in detail in the FAO Manual on the provision and evaluation of pesticide residue data for the estimation of maximum residue levels in food and feed (2016) hereafter referred to as the FAO Manual. The WHO Core Assessment Group on Pesticide Residues was responsible for reviewing toxicological and related data where necessary and possible.

The Meeting evaluated 29 pesticides for residues with regard to additional uses. The Meeting estimated maximum residue levels and recommended them for use by CCPR and estimated supervised trials median residue (STMR) and highest residue (HR) levels as a basis for estimating dietary exposures.

The Meeting also estimated the dietary exposures (both acute and long-term) of the pesticides reviewed and, on this basis, performed a dietary risk assessment in relation to the relevant ADI and where necessary the ARfD. Cases in which ADIs or ARfDs may be exceeded, if they occur, are clearly indicated in order to facilitate the decision-making process by CCPR.

1.1. Declaration of interests

The Secretariat informed the Meeting that all experts participating in the 2021 Extra JMPR had completed declaration of interest forms and that no conflicts had been identified.

2. General considerations

Feedback on virtual Extra JMPR Meeting

The Covid-19 pandemic forced the cancellation of the JMPR Meeting scheduled for September 2020 in Rome, Italy and precluded an in-person meeting of the 2021 Extra Meeting of the JMPR, scheduled to take place between 12–21 May 2021, also in Rome, Italy.

In order to avoid an increased backlog of compounds scheduled for JMPR review that would have resulted from cancelling the 2021 Extra Meeting, the joint FAO/WHO secretariats considered the use of online video-conferencing. This decision was based on the nature of the work for Extra Meetings, which is less complex than regular annual Meetings and does not address new compound evaluations or periodic re-evaluations.

The joint JMPR Secretariats discussed possible changes to the meeting format with all invited experts and the Codex Secretariat. All experts expressed their willingness and ability to participate in the 2021 Extra Meeting remotely, through video conference, as a one-time measure given the exceptional circumstances of the COVID-19 pandemic. For the purpose of the Extra 2021 Meeting, both the FAO panel and the WHO Core group were represented by a subset of experts and not by the full complement of panel members and temporary advisers normally present for a typical annual Meeting. It must be noted that the scheduled work of the 2021 Extra Meeting was restricted to evaluations of new uses only and although the agenda was expanded to include new uses from the 2020 and 2021 schedules, the overall level of effort and interaction required was significantly less than would be needed for a typical annual JMPR Meeting where new substances and periodic review compounds occupy the majority of the meeting's focus and attention.

Prior to the Meeting, the organizers recognized the challenges associated with the use of an on-line format and the 16-h range in time zones of the participants' locations that needed to be accommodated. Multiple video-conferences were self-organized with consideration to time zone and spread of expertise, resulting in five subgroups of FAO experts commencing activities towards the end of 2020 to facilitate the necessary pre-meeting work, such as identifying potential issues with data and their interpretation, initiating discussions between FAO and WHO experts and preparing documents for discussion at the actual Extra Meeting.

The 2021 Extra Meeting of the JMPR was held online using the Microsoft-Teams platform. The video-conferences were held in two five-day sessions from 17–21 May and from 7–11 June 2021. For a number of practical reasons, video-conferencing with all experts was restricted to 3–4 hour time slots (noon to 16:00, CEST) per day, which allowed less than half the 8–10 hour daily meeting time typical of an in-person JMPR meeting.

The first session focused on a full review of all draft evaluations and appraisals. Modifications to the draft documents were made during the 2-week period prior to the second session. The goal being to have appraisals suitable for sponsor review and plenary discussion during the second session. However, significant review work was still needed during the second session, prior to plenary discussion and ultimately, finalisation and adoption of the report.

The Meeting evaluated 29 compounds and estimated 237 maximum residue levels. The commitment and flexibility of all experts attending the meeting was fully recognized and appreciated by the joint FAO/WHO Secretariats, especially as the scheduled meeting time was well outside of normal working hours for a number of experts.

Experiences and opinions from the experts following participation in the online Meeting are summarised as follows:

- In particular, for sessions with participation of experts from many time zones, discussions involving more people became less feasible. Session timings were limited to a few hours and the efforts taken by individual experts to ensure very early morning or late-night participation and engagement in the Meeting were significant. A 3–4 hour time frame for video

conferencing was considered a maximum for focus and concentration on such technical subject matter.

- During the usual physical meetings, experts are immersed for a period of about two weeks with full attention to the scheduled work of JMPR and face little to no distractions. In contrast, during the virtual meetings some experts attended the video-conferences in parallel to their actual full-time employment. This often resulted in long days (especially for experts from time zones outside CEST) and strongly increased their workloads on-top of their normal daily routine. This also impacts the ability to concentrate on the compounds under JMPR consideration in a global context.
- Exchange of ideas and opportunity to fully utilise individual expertise was more limited in the virtual format and this can have downsides on a personal level as well as in the development of the final conclusions.
- Simple aspects and initial planning are readily considered during virtual meetings.
- For complex, potentially contentious issues a physical meeting is considered optimal. Video-conferences also have technical limitations. Especially for the simultaneous assessment of multiple studies (as required in residues/toxicity assessments), multiple screens or print-outs help in gaining the complete picture. This aspect cannot be reproduced by video-conferencing platforms, because on-screen focus is only on one document or presenter at a time. In addition, informal contacts between experts besides the normal Meeting (e.g., detailed discussions on wordings or translations, seeking clarification with the editor) become more difficult without breaking the progress of the general discussions. The technical possibility for break-out groups was not considered feasible, splitting the Meeting into sub-groups would have resulted in a number of experts being unable to participate in the main discussion.
- It also needs to be mentioned that video-conferencing demands access to good connectivity and IT technology than is required for physical meetings by the experts. Stable and effectively functioning internet connections are a necessity as well as the additional hardware requirements needed to improve the performance during the Meeting (e.g., multiple screens to check documents in parallel and large screens). In addition, not all experts are able (or allowed for IT security reasons) to join video-conferences from their workplaces and need to rely on their personal computing infrastructure. Physical meetings also avoid any additional risk of online security complications.
- The flexibility of online sessions could result in an increased accessibility to the Meeting for experts who may have difficulty travelling and allow contributions by a broader range of individual experts.
- One aspect noted during the Meeting is a lack of experience in changing workflows from physical to remote meetings. In physical JMPR Meetings, working procedures have grown over the years and strongly rely on direct communication and interaction to best support open and independent discussion. A comparable routine still needs to evolve for working in a virtual environment.
- Experts considered that for the moderately simple evaluations considered for this Extra 2021 Meeting, the overall inputs of time, including commenting process outside of the virtual meeting itself, were longer for this virtual meeting than would have been the case to have considered the evaluations through a face-to-face meeting setting.

In summary, the experts noted that an online meeting was not conducive to the essential in-depth robust scientific discussions and, thus, can only partially substitute for face-to-face meetings. Especially for annual JMPR Meetings involving evaluations of new compounds or periodic re-evaluations, for which health-based guidance values and residue definitions need to be established, video-conferencing was not considered a suitable format. The experts noted that the online format is less efficient, more time consuming and does not foster an atmosphere of openness and free interaction between experts on individual level as well as between the two joint groups as in physical

JMPR Meetings. In general, continuation of the online-only meeting format is expected to provide only limited benefits, which overall, are outweighed by counterproductive aspects which would not aid future JMPR decision making.

3. Responses to specific concerns raised by the Codex Committee on Pesticide Residues (CCPR)

No specific concerns raised by the CCPR were considered by the 2021 Extra JMPR Meeting.

4. Dietary exposure assessment for pesticide residues in food

4.1. Long-term dietary exposure

At the present Meeting, an International Estimated Daily Intake (IEDI) was calculated for each compound for which an acceptable daily intake (ADI) was established. The IEDI was calculated by multiplying the median concentrations of residues by the average daily per capita consumption of treated commodities. The concentrations were supervised trials median residues [STMRs] and/or supervised trials median residues in a processed commodity [STMR-Ps]. The per capita food consumption amounts were estimated using the 17 Global Environment Monitoring System – Food Contamination Monitoring and Assessment Programme (GEMS/Food) cluster diets. A detailed description of the method is included in the Environmental Health Criteria 240 (EHC 240) monograph.¹

These IEDIs were expressed as a percentage of the upper bound of the ADIs for a 55 kg or 60 kg person, depending on the region covered by each cluster diet (Table 1). The spreadsheet application is available at http://www.who.int/foodsafety/areas_work/chemical-risks/gems-food/en/.

The detailed calculations of the long-term dietary exposure assessments are given in Annex 3.

Table 1 Summary of long-term dietary exposure assessments (IEDI)

CCPR code	Compound name	ADI (mg/kg bw)	Range of IEDI, as % of the upper bound of the ADI
246	Acetamiprid	0–0.07	0–3
262	Bixafen	0–0.02	4–10
156	Clofentezine	0–0.02	0–4
238	Clothianidin	0–0.1	0–2
207	Cyprodinil	0–0.03	7–70
224	Difenoconazole	0–0.01	10–80
304	Ethiprole	0–0.005	1–6
197	Fenbuconazole	0–0.03	0–3
215	Fenhexamid	0–0.2	0–5
305	Fenpicoxamid	0–0.05	0
243	Fluopyram	0–0.01	10–80
110	Imazalil	0–0.03	3–40
299	Isoprothiolane	0–0.1	0–2
268	Isoxaflutole	0–0.02	0
231	Mandipropamid	0–0.2	0–6
209	Methoxyfenozide	0–0.1	1–7
232	Prothioconazole	-	-
	Prothioconazole-desthio	0–0.01	1–3
309	Pydiflumetofen	0–0.1	2–20
222	Quinoxifen	0–0.2	0–1
147	S-methoprene	0–0.05	10–60
233	Spinetoram	0–0.05	2–20
252	Sulfoxaflor	0–0.05	0–7
189	Tebuconazole	0–0.3	1–9
245	Thiamethoxam	0–0.08	1–7
213	Trifloxystrobin	0–0.04	1–9
271	Trinexapac-ethyl	0–0.3 ^a	0

¹ FAO/WHO (2009). Principles and methods for the risk assessment of chemicals in food. A joint publication of the Food and Agriculture Organization of the United Nations and the World Health Organization. Geneva: World Health Organization (Environmental Health Criteria 240; <https://www.who.int/health-topics/food-safety>).

ADI: acceptable daily intake; bw: body weight; CCPR: Codex Committee on Pesticide Residues; IEDI: international estimated daily intake

^a expressed as trinexapac acid equivalents

There was insufficient information to propose maximum residue levels for chlorpyrifos, cypermethrin, ethion and spiromesifen.

4.2. Acute dietary exposure

At the present Meeting, an International Estimate of Short-Term Intake (IESTI) was calculated for compounds for which an acute reference dose (ARfD) was established. For each relevant food commodity, the highest expected residue (highest residue in the edible portion of a commodity [HR] or highest residue in a processed commodity [HR-P]) and the highest large portion (LP) data for the general population (all ages) and children (6 years and under) were used for the calculation of the IESTI. The LP data are derived from national dietary survey data collated by the WHO. For mixed commodities the STMR is used as the residue level in the IESTI calculation. In the case where a separate ARfD was established for women of childbearing age, the IESTI was calculated for this population group only. A description of the method is included in EHC 240.

These IESTI results are expressed as a percentage of the ARfD (Table 3). The spreadsheet application is available at http://www.fao.org/fileadmin/templates/agphome/documents/Pests_Pesticides/JMPR/IESTI_IEDI_2020Nov.zip.

There was insufficient information to propose maximum residue levels for chlorpyrifos, cypermethrin, ethion and spiromesifen.

The present or previous Meetings agreed that ARfDs for clofentezine, cyprodinil, fenhexamid, fenpicoxamid, isoprothiolane, isoxaflutole, mandipropamid, quinoxifen, S-methoprene, spiromesifen, trifloxystrobin, trinexapac-ethyl were unnecessary. Therefore, for these compounds, an acute dietary exposure assessment was not conducted.

The detailed calculations of acute dietary exposure are given in [Annex 4](#).

Table 3 Summary of acute dietary exposure assessments (IESTI)

CCPR code	Compound name	ARfD (mg/kg bw)	Commodity (maximum % of ARfD)	Exceeding: population, (country)
246	Acetamiprid	0.1	3	
262	Bixafen	0.2	20	
238	Clothianidin	0.6	1	
224	Difenoconazole	0.3	2	
304	Ethiprole	0.005	2	
197	Fenbuconazole	0.2	3	
243	Fluopyram	0.5	0	
110	Imazalil	0.05	50	
209	Methoxyfenozide	0.9	5	
232	Prothioconazole	-	-	
	Prothioconazole-desthio	0.01 WCBA	8	
		1 All except WCBA	0	
309	Pydiflumetofen	0.3	20	
252	Sulfoxaflor	0.3	6	
189	Tebuconazole	0.3	0	
245	Thiamethoxam	1	5	

ARfD: acute reference dose; bw: body weight; CCPR: Codex Committee on Pesticide Residues; IESTI: international estimate of short-term intake; WCBA: women of childbearing age

5. Evaluation of data for acceptable daily intake and acute reference dose for humans, maximum residue levels and supervised trials median residue values

5.1. ACETAMIPRID (246)

RESIDUE AND ANALYTICAL ASPECTS

Acetamiprid is a neonicotinoid insecticide. The compound was evaluated for the first time by the JMPR in 2011 (T,R), where an ADI of 0–0.07 mg/kg bw and an ARfD of 0.1 mg/kg bw was established. Maximum residue levels were recommended by the JMPR in 2011, 2012 (response to concerns raised on several leafy vegetables) and 2015.

The residue definition for acetamiprid in plant commodities for compliance with the MRL and dietary risk assessment is acetamiprid. The residue definition for animal commodities for compliance with the MRL and dietary risk assessment is the sum of acetamiprid and N-desmethyl-acetamiprid, expressed as acetamiprid.

The residue is not fat-soluble.

The 2017 JMPR evaluated acetamiprid for an additional use on pistachio in Iran, but concluded that the provided data did not match the GAP. At the Fiftieth Session of the CCPR it was indicated an alternative GAP would be provided. The current Meeting received a new label to support a maximum residue level recommendation for use of acetamiprid on pistachio, based on field trial data provided in 2017.

Results of supervised residue trials on crops

Pistachio

The critical GAP for acetamiprid on pistachio in Iran is for 3 foliar applications at 0.05 kg ai/ha and a PHI of 28 days. The 2017 Meeting received four trials, conducted in Iran, using an application rate of 3×0.05 kg ai/ha, a re-treatment interval (RTI) of 39–60 days between the first two treatments and 29–32 days between the second and third treatments and a PHI 28–30 days. Information from the Iranian government confirms that these RTI ranges are common practice in the country.

The Meeting decided that the four trials were suitable for maximum residue level estimation. Residue levels (in nutmeat) were ($n = 4$): 0.20, 0.32, 0.34 and 0.49 mg/kg.

The Meeting estimated a maximum residue level of 1 mg/kg and an STMR and HR of 0.33 and 0.51 mg/kg (highest individual value), respectively for pistachio. The Meeting decided to withdraw the maximum residue level for the Group of Tree nuts of 0.06 mg/kg and to recommend a maximum residue level of 0.06 mg/kg for the Group of Tree nuts, except Pistachio.

Residues in animal commodities

Pistachio is not part of the diets listed in the most recent version of the OECD livestock dietary burden calculator. The Meeting therefore did not recalculate the livestock dietary burden and confirms its previous recommendations for animal commodities.

RECOMMENDATIONS

On the basis of the data obtained from supervised trials, the Meeting concluded that the residue levels listed in Annex 1 are suitable for establishing maximum residue limits and for IEDI and IESTI assessments.

Definition of the residue for compliance with the MRL and dietary risk assessment for plant commodities: *Acetamiprid*

Definition of the residue for compliance with the MRL and dietary risk assessment for animal commodities: *Sum of acetamiprid and N-desmethyl-acetamiprid, expressed as acetamiprid.*

The residue is not fat-soluble

DIETARY RISK ASSESSMENT

Long-term dietary exposure

The ADI for acetamiprid is 0–0.07 mg/kg bw. The International Estimated Daily Intakes (IEDIs) for acetamiprid were estimated for the 17 GEMS/Food Consumption Cluster Diets using the STMR or STMR-P values estimated by the JMPR. The results are shown in Annex 3 of the 2021 Extra JMPR Report.

The IEDIs ranged from 0–3% of the maximum ADI. The Meeting concluded that long-term dietary exposure to residues of acetamiprid from uses considered by previous and current JMPR is unlikely to present a public health concern.

Acute dietary exposure

The ARfD for acetamiprid is 0.1 mg/kg bw. The International Estimate of Short-Term Intakes (IESTIs) for acetamiprid were calculated for the food commodities and their processed commodities for which HRs/HR-Ps or STMRs/STMR-Ps were estimated by the present Meeting and for which consumption data were available. The results are shown in Annex 4 of the 2021 Extra JMPR Report.

The IESTI for pistachio varied from 2% of the ARfD for the general population to 3% of the ARfD for children. The Meeting concluded that acute dietary exposure to residues of acetamiprid from use considered by the present Meeting is unlikely to present a public health concern.

5.2. BIXAFEN (262)

RESIDUE AND ANALYTICAL ASPECTS

Bixafen (ISO common name) is a pyrazole-carboxamide fungicide used to control diseases on multiple crops. Bixafen inhibits fungal respiration by binding to mitochondrial respiratory complex II. It was considered for the first time by the 2013 JMPR for toxicology and residues, when an ADI of 0–0.02 mg/kg bw and an ARfD of 0.2 mg/kg bw were established. Bixafen was last reviewed for residues by the 2016 JMPR, where residues in rotational crops were evaluated and recommendations were made for maximum residue levels in plant and animal commodities.

The 2013 JMPR recommended the following residue definition for bixafen:

Definition of the residue for compliance with the MRL for plant commodities: *bixafen*




Definition of the residue for compliance with the MRL for animal commodities and for dietary risk assessment for plant and animal commodities: *sum of bixafen and N-(3',4'-dichloro-5-fluorobiphenyl-2-yl)-3-(difluoromethyl)-1H-pyrazole-4-carboxamide (bixafen-desmethyl), expressed as bixafen*

The residue is fat-soluble.

Bixafen was scheduled at the Fifty-first Session of the CCPR for the evaluation of additional uses by the 2020 JMPR Meeting, which was postponed to the 2021 Extra JMPR. The current Meeting received new information on use patterns for bixafen in pulses, potatoes, cereal grains and oilseed crops, supported by additional plant metabolism studies, field rotational crop studies, analytical methods and recovery data, supervised field trials and studies simulating typical processing conditions.

Table 1 List of compounds and abbreviations

Code Names	Chemical name	Structure	Where found
Bixafen BYF 00587 (AE1698406)	N-(3',4'-dichloro-5-fluorobiphenyl-2-yl)-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide (IUPAC) N-(3',4'-dichloro-5-fluoro[1,1'-biphenyl]-2-yl)-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide (CAS)		Plants Soil Livestock Rats
Bixafen-desmethyl (M21) (BCS-AA10008)	N-(3',4'-dichloro-5-fluorobiphenyl-2-yl)-3-(difluoromethyl)-1H-pyrazole-4-carboxamide		Plants Soil Livestock Rats
Bixafen-pyrazole-4-carboxylic acid (M42) (AE 1954999)	3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxylic acid		Rotational crops Soil Livestock Rats

Code Names	Chemical name	Structure	Where found
Bixafen-pyrazole-4-carboxamide (M43)	3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide		Rotational crops Rats
Bixafen-desmethyl-pyrazole-4-carboxylic acid (tautomer 1 & 2) (M44 & M45) (BCS-AA10651)	3-(difluoromethyl)-1H-pyrazole-4-carboxylic acid □ 5-(difluoromethyl)-1H-pyrazole-4-carboxylic acid		Plants Rotational crops Rats
Bixafen-pyrazolone-4-carboxylic acid (M47)	3-oxo-2,3-dihydro-1H-pyrazole-4-carboxylic acid or 3-hydroxy-1H-pyrazole-4-carboxylic acid		Plants Rotational crops

Plant metabolism

The fate of bixafen in plants was evaluated by the 2013 Meeting following foliar spray application of [pyrazole-5-¹⁴C]-bixafen or [dichlorophenyl-UL-¹⁴C]-bixafen to soya beans or wheat. A detailed assessment of these studies is presented in the 2013 JMPR Report. The current Meeting received additional plant metabolism studies on potatoes and tomato.

The metabolism of both bixafen radiolabels in potatoes was investigated in parallel under semi-enclosed conditions (plant containers, natural sunlight, rain cover) with three foliar applications of 0.24 kg ai/ha each. The treatments were performed at the beginning of flowering (BBCH 61), at fruit development (BBCH 70) and 7 days before harvest (BBCH 97, leaves and stem dead). Potato leaves were harvested at an intermediate growth stage (BBCH 92, beginning of senescence) 28 days after the second application. Mature tubers were sampled 7 days after the final treatment.

Total radioactive residues (TRR) were comparable between both labels, representing 21.8–24.4 mg eq/kg for leaves and 0.002–0.003 mg eq/kg for tubers. All samples were extracted three times with a mixture of acetonitrile/water (4/1; v/v), which recovered ~98% of the TRR from the leaves and 85% of the TRR from the tubers (pyrazole-label only). Post extraction solids amounted 2.0–2.3% of the TRR for leaves (0.445–0.558 mg eq/kg) and 15.2% of the TRR for tubers (< 0.001 mg eq/kg).

The identification of the radioactive residues revealed unchanged bixafen as the predominant residue in potato leaves for both radiolabels (95.6–95.9% TRR). The predominant metabolite in leaves was bixafen-desmethyl, present at 0.6% of the TRR (0.121–0.135 mg eq/kg). No additional

identified metabolites (M42, M43 and M47) or unidentified analytical fractions represented more than 0.4% of the TRR each.

For tubers, only the samples from the [pyrazole-¹⁴C]-label were further investigated. TRR levels were low, allowing only limited identification of the residue. Parent bixafen was the predominant residue in tubers representing 31.4% of the TRR (0.001 mg eq/kg). Identified metabolites were M47 at the LOQ of 0.001 mg eq/kg (25.9% TRR) and the tautomeric mixture of M44 & M45 (12.5% TRR), M43 (9.0% TRR) and M42 (5.9% TRR), all present below 0.001 mg eq/kg.

For tomatoes, the metabolism of bixafen was investigated in a greenhouse by application of [pyrazole-5-¹⁴C]-bixafen or [dichlorophenyl-UL-¹⁴C]-bixafen under identical experimental conditions. The plants received three foliar applications with individual rates of about 0.21–0.23 kg ai/ha each at BBCH 78–80 (development of fruit: fruits have reached typical form and size), 21 days later at BBCH 84–85 (ripening of fruit: 40–50% of fruits show typical fully ripe colour) and further 14 days later at BBCH 87–89 (ripening of fruit: 70–100% of fruits show typical fully ripe colour). Samples of fruits and leaves were collected 3 days after the last treatment (DALT), however only the fruits were further investigated.

The fruits were surface-washed with dichloromethane and subsequently extracted three times with acetonitrile/water (4/1; v/v). TRR levels were 1.75 mg eq/kg for [pyrazole-¹⁴C]-bixafen and 3.19 mg eq/kg for [dichlorophenyl-UL-¹⁴C]-bixafen treated samples. Approximately 93–94% of the radioactivity was recovered in the surface wash and 5.6–6.9% of the TRR in the extract. Post extraction solids amounted to 0.1% of the TRR.

For both labels, in the combined surface washes and extracts, bixafen was the predominant residue representing a total of >99% of the TRR. The only identified metabolite was bixafen-desmethyl, present at 0.1% of the TRR (0.002–0.004 mg eq/kg). In addition, two minor unidentified analytical fractions were found at < 0.1–0.2% of the TRR (0.001–0.005 mg eq/kg).

The Meeting concluded that parent bixafen is the predominant residue in all plant parts (potato leaves and tubers, tomato fruits). The only other metabolite identified was bixafen-desmethyl at minor proportions (<1% TRR). In potato tubers, TRR levels were very low (0.002–0.003 mg eq/kg). Parent bixafen was also the predominant residue (31.4% TRR), however M47 was quantified as a major metabolite by proportion (25.9% TRR), but not based on its concentration (0.001 mg eq/kg). Further metabolites identified in tubers were M44 and M45, M43 and M42, all present below 0.001 mg eq/kg. Metabolites M21 (0.6%; 0.135 mg eq/kg), M43 (0.1%; 0.025 mg eq/kg), M42 (0.1%; 0.023 mg eq/kg) and M47 (< 0.1%; 0.006 mg eq/kg) were also identified in potato leaves.

Environmental fate (rotational crops)

The current Meeting received multiple additional field rotational crop studies, all conducted in Europe with one application of 1.13 kg ai/ha to bare soil. After plant back intervals (PBI) of 30, 120 and 365 days, various crops were planted/seeded and grown to commercial harvest. In parallel, initial residues in soil were measured directly after treatment.

All samples were analysed for bixafen and its desmethyl-metabolite (M21), M43 and M44/45. For strawberries and cabbage, residues were generally below the LOQs of 0.01 mg/kg at each PBI. Also, no residues of bixafen-desmethyl and M43 were quantified in any of the samples. For all other crops investigated, the results for residues of bixafen and M44/45 are summarized in the following Table.

Table 2 Overview of bixafen residues in rotational crops

Crop	Matrix	PBI (days)	Residues (mg eq/kg)	
			Bixafen	M44/45
Leek	Stalks	30	4 × < 0.01	3 × < 0.01, 0.016
		120	4 × < 0.01	3 × < 0.01, 0.034
		365	4 × < 0.01	2 × < 0.01, 0.011, 0.014

Crop	Matrix	PBI	Residues (mg eq/kg)	
		(days)	Bixafen	M44/45
Cabbage	Head	30	$4 \times < 0.01$	$3 \times < 0.01, 0.01$
		120	$4 \times < 0.01$	$4 \times < 0.01$
		365	$4 \times < 0.01$	$4 \times < 0.01$
Courgette	Fruit	30	$4 \times < 0.01$	$3 \times < 0.01, 0.011$
		120	$4 \times < 0.01$	$2 \times < 0.01, 0.015, 0.023$
		365	$4 \times < 0.01$	$3 \times < 0.01, 0.018$
Pea	Seed, dry	30	$3 \times < 0.01, 0.033$	$3 \times < 0.02, 0.037$
		120	$4 \times < 0.01$	$2 \times < 0.02, 0.019, 0.11$
		365	$4 \times < 0.01$	$3 \times < 0.02, 0.061$
	Vines	30	$4 \times < 0.01$	$2 \times < 0.02, 0.030, 0.055$
		120	$4 \times < 0.01$	$2 \times < 0.02, 0.011, 0.082$
		365	$4 \times < 0.01$	$3 \times < 0.02, 0.047$
Potato	Tubers	30	$2 \times < 0.01, 0.018, 0.058$	$< 0.02, 0.012, 0.061, 0.064$
		120	$3 \times < 0.01, 0.019$	$0.016, 0.016, 0.050, 0.060$
		365	$4 \times < 0.01$	$0.012, 0.015, 0.025, 0.038$
Maize	Forage	30	$4 \times < 0.01$	$3 \times < 0.01, 0.030$
		120	$4 \times < 0.01$	$4 \times < 0.01$
		365	$4 \times < 0.01$	$4 \times < 0.01$
	Kernel	30	$4 \times < 0.01$	$3 \times < 0.01, 0.063$
		120	$4 \times < 0.01$	$4 \times < 0.01$
		365	$4 \times < 0.01$	$4 \times < 0.01$
	Stover	30	$4 \times < 0.01$	$3 \times < 0.01, 0.017$
		120	$4 \times < 0.01$	$3 \times < 0.01, 0.025$
		365	$4 \times < 0.01$	$3 \times < 0.01, 0.023$
Oilseed rape	Forage	30	$4 \times < 0.01$	$3 \times < 0.01, 0.015$
		120	$4 \times < 0.01$	$3 \times < 0.01, 0.01$
		365	$4 \times < 0.01$	$2 \times < 0.01, 0.014, 0.042$
	Seed, dry	30	$4 \times < 0.01$	$4 \times < 0.01$
		120	$4 \times < 0.01$	$4 \times < 0.01$
		365	$4 \times < 0.01$	$4 \times < 0.01$

Concerning the uptake of bixafen residues into rotational crops, the 2016 JMPR made a very conservative estimate for the soil plateau concentration equivalent to 0.93 kg ai/ha, which addresses many years of subsequent application at the maximum annual GAP rate (up to 0.25 kg ai/ha and year). This estimation also covers all GAPs evaluated by the current Meeting.

The 2016 JMPR concluded that small findings of bixafen in carrot roots (mean: 0.02 mg/kg, highest residue: 0.028 mg/kg) in field rotational crop studies addressing the estimated soil plateau concentrations were insignificant in view of the overall conservatism involved. However, additional field rotational crop data provided to the current Meeting suggest a potential transfer of parent bixafen into root crops and into pulse crops (not supported with data in 2016).

For peas, representing the pulses crop group, a highest residue of 0.033 mg/kg was found for bixafen in the 30 day crop rotation. The median for the total residue (sum of bixafen and bixafen-desmethyl, expressed as bixafen) was < 0.02 mg/kg.

New data on potato tubers showed quantifiable residues in two out of four field trials at the 30-day PBI and is complimented by three results in carrot roots as evaluated by the 2016 JMPR. The combined range of residues for root and tuber vegetables is: < 0.01 , < 0.01 , 0.014, 0.018, 0.019, 0.028

and 0.058 mg/kg for bixafen and < 0.02, < 0.02, 0.024, 0.028, 0.029, 0.038 and 0.068 mg/kg for the total bixafen residue.

In summary, the Meeting concluded that for root and tuber vegetables and pulse crops, the overall data available suggest a potential uptake of bixafen residues from soil. For all other crops that are potentially rotated, no significant uptake of bixafen residues from soil is expected.

The 2013 JMPR assessed the relevance of the tautomeric mixture M44/M45 against the TTC of a Cramer Class III compound. Based on field rotational crop data provided to the 2016 JMPR and the additional data provided to the current Meeting, the following concentrations of M44/M45 arising were estimated:

Table 3 Overview of M44/M45 residues in rotational crops

Commodity group	Field rotational crop commodity	M44/M45 in mg/kg (highest concentrations per trial from all PBIs)	M44/M45 STMR value in mg/kg
Bulb vegetables	Leek	< 0.01, < 0.01, 0.016, 0.034	0.013
Root and tuber vegetables	Carrot roots (2016 JMPR)	< 0.01, < 0.01, 0.044	
	Potato tuber	0.016, 0.016, 0.061, 0.064	
	Combined	< 0.01, < 0.01, 0.016, 0.016, 0.044, 0.061, 0.064	0.016
Leafy crops and brassica (extrapolated to stalk and stem vegetables)	Lettuce (2016 JMPR)	< 0.01, 0.017, 0.092	
	Cabbage	< 0.01, < 0.01, < 0.01, 0.01	
	Carrot leaves (2016 JMPR)	< 0.01, < 0.01, 0.18	
	Combined	< 0.01, < 0.01, < 0.01, < 0.01, < 0.01, < 0.01, 0.01, 0.017, 0.092, 0.18	0.01
Fruiting vegetables and strawberries (extrapolated to legume vegetables)	Courgettes	< 0.01, < 0.01, 0.015, 0.023	
	Strawberries	< 0.01, < 0.01, < 0.01, < 0.01	
	Combined	< 0.01, < 0.01, < 0.01, < 0.01, < 0.01, < 0.01, 0.015, 0.023	0.01
Pulses	Peas (dry)	< 0.01, < 0.01, 0.037, 0.082	0.0235
Cereal grains	Barley/wheat (2016 JMPR)	< 0.01, < 0.01, < 0.01	
	Maize	< 0.01, < 0.01, < 0.01, 0.063	
	Combined	< 0.01, < 0.01, < 0.01, < 0.01, < 0.01, < 0.01, 0.063	0.01
Oilseeds	Rape seed	< 0.01, < 0.01, < 0.01, < 0.01	0.01

Methods of analysis

The current Meeting received additional analytical methods for the determination of parent bixafen in plant commodities and additional concurrent recovery information for the methods 01012, 01013, 01633/M001 and 00952 evaluated by the 2013 JMPR.

For plant matrices, the new method 01367 (including ILV) was provided, based on the QuEChERS multi-method. Samples are extracted with acetonitrile in the presence of salts added for phase separation and cleaned by SPE. Bixafen is analysed by HPLC-MS/MS (ESI+) and the method was tested in high water, high oil, high protein, high acid and high starch matrices at an LOQ of 0.01 mg/kg.

The methods 01012, 01013 and 01633/M001 for plant commodities and method 00952 for soil were first evaluated by the 2013 JMPR. New recovery data provided to the current Meeting confirmed the previous LOQs of 0.01 mg/kg for bixafen, bixafen-desmethyl (M21), M43 and M44/M45 in all matrices analysed within acceptable ranges.

The Meeting concluded that the presented methods were sufficiently validated and are suitable to measure bixafen, bixafen-desmethyl, M43 and M44/45 in plant commodities and/or soil.

Stability of pesticide residues in stored analytical samples

The current Meeting received additional information on the storage stability of bixafen and bixafen-desmethyl (M21) in plant matrices and of M44/M45 in soil.

In plant matrices, the storage stability of both bixafen and bixafen-desmethyl (M21) was demonstrated at -18 °C in orange fruits and dry bean seeds for at least 24 months.

In two different soil types, M44/M45 was also stable for up to 18 months under freezer storage conditions (-18 °C). At the last sampling interval of 24 months, M44/M45 was also stable in the clay loam investigated but showed significant degradation (< 70% remaining) in the silt loam soil.

The Meeting agreed that the demonstrated storage stability on various representative plant and animal commodities covered the residue sample storage intervals used in the field trials considered by the current Meeting.

Definition of the residue

The current Meeting received new data on the metabolism of bixafen following foliar treatment of potatoes and tomatoes.

Following foliar application to potatoes, bixafen was the predominant residue identified in all matrices. In leaves, unchanged parent bixafen represented 96% of the TRR (20.9–23.3 mg eq/kg). In tubers protected by soil, bixafen was still the major residue (31% TRR), but at low concentrations of 0.001 mg eq/kg. The only metabolite identified in leaves was bixafen desmethyl (M21; 0.6% TRR, 0.12–0.14 mg eq/kg). In tubers, M47 was found at 0.001 mg eq/kg (26% TRR), while M44 and M45, M43 and M42 were identified (5.9–12.5% TRR) at concentrations below 0.001 mg eq/kg.

In tomato fruits collected three days after the final treatment, more than 99% of the TRR was identified as unchanged parent bixafen (1.7–3.2 mg eq/kg). The only identified metabolite was bixafen-desmethyl (M21; 0.1% TRR, up to 0.004 mg eq/kg).

The Meeting confirmed its previous recommendation of bixafen for compliance with the MRL and the sum of bixafen and bixafen-desmethyl (M21) for dietary risk assessment for plant commodities.

Results of supervised residue trials on crops

The Meeting received supervised trial data for applications of bixafen on soya beans, potatoes, barley, maize (including sweet corn), sorghum, wheat, cotton, peanuts and sunflowers, respectively.

Residue values referred to as “total” are the sum of bixafen and M21 (bixafen-desmethyl), expressed as bixafen. Due to the general potential for uptake of residues from soil, LOQ values are always added to calculate the total residue concentration.

Soya beans (dry)

Bixafen is registered in the USA for use on soya beans with a maximum GAP involving two foliar spray applications of 0.077 kg ai/ha each (10-day interval) and a PHI of 21 days.

Supervised field trials conducted in Canada and the USA on soya beans were provided approximating the cGAP use pattern but at higher rates of 0.11 kg ai/ha per application. In absence of supervised field trial data matching the GAP within ±25%, the Meeting decided to scale the available data accordingly with a factor of 0.7 (0.077 kg ai/ha cGAP rate ÷ 0.11 kg ai/ha trial rate).

Unscaled residues of bixafen in soya beans were (n = 21): < 0.01(13), 0.01, 0.011, 0.012, 0.014, 0.014, 0.015, 0.024, 0.029 mg/kg.

Unscaled total residues of bixafen in soya beans were (n = 21): < 0.02(13), 0.020, 0.021, 0.022, 0.024, 0.024, 0.025, 0.034, 0.039 mg/kg.

Scaled (factor 0.7) residues of bixafen in soya beans were (n = 21): < 0.007(13), 0.007, 0.0077, 0.0084, 0.0098, 0.0098, 0.0105, 0.0168, 0.0203 mg/kg.

Scaled (factor 0.7) total residues of bixafen in soya beans were (n = 21): < 0.014(13), 0.014, 0.0147, 0.0154, 0.0168, 0.0168, 0.0175, 0.0238, 0.0273 mg/kg.

In field studies on succeeding crops a highest residue for bixafen in dry pea seeds of 0.033 mg/kg was found. The median total residue was < 0.02 mg/kg. Since soya beans can be both a primary and succeeding crop, the Meeting concluded that additional residues due to uptake via roots need to be taken into account.

The Meeting estimated a maximum residue level of 0.08 mg/kg (by adding the highest residue of 0.033 mg/kg for rotated pulse crops to each individual bixafen result for soya beans) and an STMR value of 0.034 mg/kg (by adding the median total residue of < 0.02 mg/kg for rotated pulse crops to the total median for soya beans) for bixafen in soya beans (dry).

Potatoes

Bixafen is registered in the USA for use on potatoes with a maximum GAP involving three foliar spray applications of 0.08 kg ai/ha each (10-day interval) and a PHI of 14 days.

Supervised field trials conducted in Canada and the USA on potatoes were provided involving four spray applications at <75% GAP rate and a shorter sampling and re-treatment interval of 7 days. The Meeting concluded that these trials do not approximate the cGAP. In addition, due to the number of deviations in the use and sampling intervals, the proportionality principle cannot be applied. No recommendation for the use of bixafen on potatoes can be given.

Barley

The 2016 JMPR assessed the use of bixafen on barley based on a GAP from the United Kingdom and recommended a maximum residue level of 0.4 mg/kg and an STMR value of 0.08 mg/kg for barley grain.

The current Meeting received new information that bixafen is registered in Brazil for the use on barley with a maximum GAP involving four foliar applications of 0.063 kg ai/ha each (15 days interval) and a PHI of 30 days.

Supervised field trials conducted in Brazil on barley were provided approximating the cGAP.

Residues of bixafen in barley grain were (n = 10): 0.11, 0.15, 0.17, 0.21, 0.30, 0.32, 0.53, 0.57, 0.58, 0.77 mg/kg.

Total residues of bixafen in barley grain were (n = 10): 0.12, 0.16, 0.18, 0.22, 0.32, 0.34, 0.54, 0.58, 0.59, 0.78 mg/kg.

The Meeting estimated a maximum residue level of 1.5 mg/kg and an STMR value of 0.33 mg/kg for bixafen in barley grain to replace its previous recommendations.

Maize

Bixafen is registered in the US for the use on maize with a maximum GAP involving two foliar spray applications of 0.077 kg ai/ha each (7 days interval, not after R4 stage/BBCH 83–85) and a PHI of 30 days.

Supervised field trials conducted in Canada and the USA on maize were provided involving two higher applications of 0.11 kg ai/ha at later growth stages, but at longer intervals of 19–36 days.

Residues of bixafen in maize grain were (n = 15): < 0.01(15) mg/kg.

Total residues of bixafen in maize grain were (n = 15): < 0.02(15) mg/kg

The Meeting noted that in these trials, treatment at rates higher than the cGAP and closer to harvest did not result in quantifiable residues in maize grain and the Meeting decided that they could be used to support the cGAP from the USA.

The Meeting estimated a maximum residue level of 0.01(*) mg/kg and an STMR value of 0.02 mg/kg for bixafen in maize grain.

Sweet corn

Bixafen is registered in Brazil for the use on maize (including sweet corn) with a maximum GAP involving two foliar spray applications of 0.063 kg ai/ha each (15 days interval) and a PHI of 15 days.

Supervised field trials conducted in Brazil on sweet corn were provided approximating the critical GAP.

Residues of bixafen in sweet corn cobs without husks were (n = 10): < 0.01(10) mg/kg.

Total residues of bixafen in sweet corn cobs without husks were (n = 10): < 0.02(10) mg/kg

The Meeting estimated a maximum residue level of 0.01(*) mg/kg, an STMR value of 0.02 mg/kg and an HR value of 0.02 mg/kg for bixafen in sweet corn (corn on the cob).

Sorghum

Bixafen is registered in the USA for the use on sorghum with a maximum GAP involving two foliar spray applications of 0.074 kg ai/ha each and a PHI of 30 days.

Supervised field trials conducted in the USA on sorghum were provided approximating the cGAP use pattern but at higher rates of 0.11 kg ai/ha per application. In the absence of supervised field trial data matching the GAP within $\pm 25\%$, the Meeting decided to scale the available data accordingly with a factor of 0.7 (0.074 kg ai/ha cGAP rate $\div$ 0.11 kg ai/ha trial rate).

Unscaled total residues of bixafen in sorghum grain were (n = 9): 0.077, 0.13, 0.17, 0.20, 0.28, 0.35, 0.49, 0.90, 1.9 mg/kg

Scaled (factor 0.7) residues of bixafen in sorghum grain were (n = 9): 0.0126, 0.077, 0.091, 0.098, 0.175, 0.175, 0.336, 0.553, 1.26 mg/kg.

Scaled (factor 0.7) total residues of bixafen in sorghum grain were (n = 9): 0.0539, 0.091, 0.119, 0.14, 0.196, 0.245, 0.343, 0.63, 1.33 mg/kg

The Meeting estimated a maximum residue level of 2 mg/kg and an STMR value of 0.196 mg/kg for bixafen in sorghum.

Wheat, rye and triticale

The 2016 JMPR assessed the use of bixafen on wheat, rye and triticale based on a GAP from the United Kingdom and recommended a maximum residue level of 0.05 mg/kg and an STMR value of 0.02 mg/kg for each of wheat, rye and triticale.

Bixafen is registered in the USA for use on wheat, rye and triticale with a maximum GAP involving two foliar spray applications of 0.075 kg ai/ha each (14 days interval) and a PHI of 30 days.

Supervised field trials on wheat conducted in Canada and the USA were provided, involving two applications of 0.11 kg ai/ha each with intervals of 18–73 days. Although the label recommends an early treatment as conducted in the supervised field trials, the label compliant cGAP allows two late treatments at a 14 days interval with a total rate not exceeding 0.15 kg ai/ha. Consequently, the Meeting concluded that the data do not approximate the cGAP registered in the USA.

In addition, bixafen is registered in Brazil for use on wheat with a maximum GAP involving four foliar spray applications of 0.063 kg ai/ha each (15 days interval) and a PHI of 30 days.

Supervised field trials conducted in Brazil on wheat were provided approximating the cGAP.

Residues of bixafen in wheat grain were (n = 10): 0.02, 0.05, 0.06, 0.06, 0.07, 0.10(3), 0.12, 0.17 mg/kg.

Total residues of bixafen in wheat grain were (n = 10): 0.03, 0.06, 0.07, 0.07, 0.08, 0.11, 0.11, 0.13, 0.14, 0.18 mg/kg.

Based on the Brazilian GAP for wheat only, the Meeting estimated a maximum residue level of 0.3 mg/kg and an STMR value of 0.095 mg/kg for bixafen to replace its previous recommendation on wheat grain.

Cotton

Bixafen is registered in Brazil for the use on cotton with a maximum GAP involving four foliar spray applications of 0.063 kg ai/ha each (15 days interval) and a PHI of 30 days.

Supervised field trials conducted in Brazil on cotton were provided approximating the cGAP.

Residues of bixafen in cotton seeds were (n = 10): < 0.01, < 0.01, 0.01, 0.01, 0.02, 0.02, 0.03, 0.06, 0.06, 0.16 mg/kg.

Total residues of bixafen in cotton seeds were (n = 10): < 0.02, < 0.02, 0.02, 0.02, 0.03, 0.03, 0.04, 0.07, 0.07, 0.17 mg/kg.

The Meeting estimated a maximum residue level of 0.3 mg/kg and an STMR value of 0.03 mg/kg for bixafen in cotton seeds.

Peanuts

Bixafen is registered in the USA for the use on peanuts with a maximum GAP involving two foliar spray applications of 0.10 kg ai/ha each (14-day interval) and a PHI of 14 days.

Supervised field trials on peanuts conducted in the USA were provided, involving four applications of approximately 0.056 kg ai/ha each with intervals of 13–15 days. The Meeting took note of the very slow metabolic degradation of bixafen residues in crops. Combined with low basipetal translocation of bixafen in plants, it was concluded that the splitting into four instead of two treatments had no impact on residues in peanut nutmeats, which are protected by soil. Since the total seasonal rate in the supervised field trials approximates the US GAP, the Meeting considered these trials as suitable to support a recommendation.

Residues of bixafen in peanut nutmeat were (n = 14): < 0.01(14) mg/kg.

Total residues of bixafen in peanut nutmeat seeds were (n = 14): < 0.02(14) mg/kg.

The Meeting estimated a maximum residue level of 0.01 mg/kg and an STMR value of 0.02 mg/kg for bixafen in peanuts.

Sunflower

Bixafen is registered in Brazil for the use on sunflowers with a maximum GAP involving two foliar spray applications of 0.063 kg ai/ha each (15 days interval) and a PHI of 30 days.

Supervised field trials conducted in Brazil on sunflowers were provided approximating the cGAP.

Residues of bixafen in sunflower seeds were (n = 10): < 0.01, < 0.01, 0.01, 0.02, 0.02, 0.05, 0.08, 0.21, 0.56, 1.7 mg/kg.

Total residues of bixafen in sunflower seeds were (n = 10): < 0.02, < 0.02, 0.02, 0.03, 0.03, 0.06, 0.09, 0.22, 0.57, 1.7 mg/kg.

The Meeting estimated a maximum residue level of 3 mg/kg and an STMR value of 0.045 mg/kg for bixafen in sunflower seeds.

Animal Feeds

Maize forage

Bixafen is registered in the US for the use on maize with a maximum GAP involving two foliar spray applications of 0.077 kg ai/ha each (7 days interval) and a PHI of 10 days for forage.

Supervised field trials conducted in Canada and the USA on maize forage were provided, mostly involving a single treatment at 0.11 kg ai/ha and one plot with two applications of 0.11 kg ai/ha, a longer re-treatment interval of 20 days and a longer sampling interval of 30 days.

The Meeting concluded that the available trials do not approximate the registered use in the USA.

Maize fodder

Bixafen is registered in the US for the use on maize with a maximum GAP involving two foliar spray applications of 0.077 kg ai/ha each (7 days interval, not after R4 stage/BBCH 83–85) and a PHI of 30 days for stover.

Supervised field trials conducted in Canada and the USA on maize were provided involving two higher applications of 0.11 kg ai/ha at a longer interval of 20 days. The Meeting noted that bixafen residues in maize forage decline trials decreased during the first 10 days and remained stable afterward. Therefore, the deviation relating to the longer interval between applications in the field trials provided was considered to be of low relevance for the final residue concentration at harvest. In absence of supervised field trial data matching the GAP within $\pm 25\%$, the Meeting decided to scale the available data accordingly with a factor of 0.7 (0.077 kg ai/ha cGAP rate $\div$ 0.11 kg ai/ha trial rate).

Unscaled residues of bixafen in maize stover were (n = 16): 0.32, 0.45, 0.49, 0.76, 1.2, 1.6, 1.7, 1.8, 1.9, 2.0(3), 2.4(3), 3.0 mg/kg.

Unscaled Total residues of bixafen in maize stover were (n = 16): 0.36, 0.53, 0.69, 0.82, 1.3, 1.6, 1.9, 2.0, 2.2(3), 2.4, 2.4, 2.5, 2.8, 3.2 mg/kg

Scaled residues of bixafen in maize stover were (n = 16): 0.224, 0.315, 0.343, 0.532, 0.84, 1.12, 1.19, 1.26, 1.33, 1.4(3), 1.68(3), 2.1 mg/kg.

Scaled Total residues of bixafen in maize stover were (n = 16): 0.252, 0.371, 0.483, 0.574, 0.91, 1.12, 1.33, 1.4, 1.54(3), 1.68, 1.68, 1.75, 1.96, 2.24 mg/kg

The Meeting estimated a maximum residue level of 5 mg/kg (DM, 89% dry-matter content), a median residue of 1.47 mg/kg (as received) and a highest residue of 2.24 mg/kg (as received) for bixafen in maize fodder.

Sorghum forage

Bixafen is registered in the US for the use on sorghum with a maximum GAP involving two foliar spray applications of 0.074 kg ai/ha each (7 days interval) and a PHI of 30 days for forage.

Supervised field trials conducted in the USA on sorghum forage were provided involving a single treatment at 0.11 kg ai/ha with a PHI of approximately 10 days.

The Meeting noted that the cGAP for bixafen also allows two applications on sorghum until the forage growth stage and concluded that the available trials do not approximate the registered use in the USA.

Sorghum fodder

Bixafen is registered in the US for the use on sorghum with a maximum GAP involving two foliar spray applications of 0.074 kg ai/ha each (7 days interval) and a PHI of 30 days.

Supervised field trials conducted in the USA on sorghum were provided involving two higher applications of 0.11 kg ai/ha at longer intervals of 15–22 days. In one decline trial conducted for sorghum forage, bixafen residues decreased significantly between day 10 (0.79 mg/kg) and day 20 (0.092 mg/kg) after the first treatment. The Meeting concluded that the longer interval in the field trials may result in a significant reduction of the residue concentrations at harvest and therefore does not reflect the critical use pattern for sorghum fodder.

Wheat forage

The 2016 JMPR evaluated the use of bixafen on small cereal grains based on a GAP from the United Kingdom and estimated a highest residue of 7.3 mg/kg and a median residue of 3.5 mg/kg for barley and wheat forage (as received) and decided to extrapolate the estimations to oats, rye and triticale forage.

The current Meeting received new GAP information for the use of bixafen on wheat from Brazil, however no residue data on forage were provided.

Bixafen is registered in the USA for the use on wheat, rye and triticale with a maximum GAP involving two foliar spray application of 0.075 kg ai/ha each (14 days interval) and a PHI of 10 days for forage and 20 days for hay.

Supervised field trials on wheat conducted in Canada and the USA were provided, involving single applications of 0.11 kg ai/ha. The cGAP allows two treatments at a 14 days interval with a total rate not exceeding 0.15 kg ai/ha including for forage. Consequently, the Meeting concluded that the data do not approximate the cGAP registered in the USA.

The Meeting confirmed its previous recommendations for bixafen in wheat forage.

Wheat straw

The 2016 JMPR evaluated the use of bixafen on small cereal grains based on a GAP from the United Kingdom and estimated a maximum residue level of 20 mg/kg (dry-weight basis, based on 89% DM content), a highest residue of 11 mg/kg (as received) and a median value of 2.2 mg/kg (as received) for barley and wheat, straw and fodder and decided to extrapolate the estimations to oats, rye and triticale straw and fodder.

The current Meeting received new GAP information for the use of bixafen on wheat from Brazil, however no residue data on straw were provided.

Bixafen is registered in the USA for use on wheat, rye and triticale with a maximum GAP involving two foliar spray applications of 0.075 kg ai/ha each (14 days interval) and a PHI of 30 days.

Supervised field trials on wheat conducted in Canada and the USA were provided, involving two applications of 0.11 kg ai/ha each with intervals of 18–73 days. Although the label recommends an early treatment as conducted in the supervised field trials, the label compliant cGAP allows two late treatments at a 14 day interval with a total rate not exceeding 0.15 kg ai/ha. Consequently, the Meeting concluded that the data do not approximate the cGAP registered in the USA.

The Meeting confirms its previous recommendations for bixafen in wheat straw.

Peanut hay

Bixafen is registered in the USA for use on peanuts with a maximum GAP involving two foliar spray applications of 0.10 kg ai/ha each (14 days interval). However, the label excludes utilisation of forage or hay for animal feed purposes.

Residues in rotational crops

Taking into account new data on field rotational crops, the current Meeting estimated a highest bixafen residue of 0.033 mg/kg for pulses. Median total bixafen residues were 0.02 mg/kg.

Based on these values, the Meeting estimated a maximum residue level of 0.04 mg/kg and an STMR value of 0.02 mg/kg for pulses, except soya beans (dry).

The current Meeting estimated a highest bixafen residue of 0.058 mg/kg for root and tuberous crops. Total bixafen residues were 0.028 mg/kg (median) and 0.068 mg/kg (highest residue).

Based on these values, the Meeting estimated a maximum residue level of 0.06 mg/kg, an STMR value of 0.028 mg/kg and an HR of 0.068 for root and tuber vegetables.

Fate of residues during processing

The fate of bixafen residues has been examined simulating commercial processing of soya beans, potatoes, maize grain, sorghum grain, wheat grain, oilseed rapeseeds and peanuts.

Estimated processing factors for the commodities considered at this Meeting are summarised below. In potatoes, residues in the raw agricultural commodity were below the LOQ. Consequently, no processing factors were derived for this commodity. For processing factors on other commodities, please refer to the evaluation document.

Table 4 Processing factors and processed commodities

Raw commodity	Processed commodity	Bixafen			Total residue		
		Individual processing factors	Mean or best estimate processing factor	Maximum residue level (mg/kg)	Individual processing factors	Mean or best estimate processing factor	STMR-P in mg/kg
Soya beans (dry) (STMR: 0.034 mg/kg, MRL: 0.08 mg/kg)	AGF	310, <u>470</u> , 620	470	-	See bixafen	See bixafen	15.98
	Flour	< 0.09, 0.13	0.11	-			0.00374
	Hulls	2.4, <u>2.8</u> , 7.9	2.8	0.3			0.0952
	Meal	0.11, <u>0.14</u> , < 0.22	0.14	-			0.00476
	Milk	< 0.04, < 0.09	0.065	-			0.00221
	Refined oil (cold press)	1.3, 1.4	1.4	0.15			0.0476
	Refined oil (solvent ext.)	0.35, <u>0.40</u> , 0.50	0.40	-			See cold press
Barley (STMR: 0.33 mg/kg, MRL: 1.5 mg/kg)	Brewer's malt	0.33 ^a , <u>0.8^a</u> , <u>0.92^a</u> , 0.96 ^a	0.86 ^a	-	0.5 ^a , <u>0.91^a</u> , 1 ^a , 1 ^a	0.96 ^a	0.317
	Beer	< 0.04 ^a , < <u>0.05^a</u> , < <u>0.08^a</u> , < 0.33 ^a	< 0.065 ^a	-	0.08 ^a , < <u>0.09^a</u> , < <u>0.13^a</u> , < 0.5 ^a	< 0.11 ^a	0.036
	Pearl barley	0.17 ^a , <u>0.2^a</u> , <u>0.23^a</u> , < 0.33 ^a	0.22 ^a	-	0.19 ^a , <u>0.23^a</u> , <u>0.27^a</u> , < 0.5 ^a	0.25 ^a	0.083
Maize (STMR: 0.02 mg/kg, MRL: 0.01* mg/kg)	AGF	160, <u>160</u> , 180	160	-	See bixafen	See bixafen	3.2
	Bran (dry milling)	2.6, 3.6	3.1	0.03			0.062
	Flour (dry milling)	< 0.63, <u>0.89</u> , 1.5	0.89	-			0.0178

Raw commodity	Processed commodity	Bixafen			Total residue		
		Individual processing factors	Mean or best estimate processing factor	Maximum residue level (mg/kg)	Individual processing factors	Mean or best estimate processing factor	STMR-P in mg/kg
	Germ (dry milling)	See wet milling	See wet milling	-			See wet milling
	Germ (wet milling)	0.63, 1.4	1.0	-			0.02
	Grits (dry milling)	< 0.36, < <u>0.63</u> , < 0.93	< 0.63	-			0.0126
	Meal (dry milling)	< 0.63, <u>0.75</u> , 1.4	0.75	-			0.015
	Refined oil (dry milling)	See wet milling	See wet milling	-			See wet milling
	Refined oil (wet milling)	1.7, <u>1.8</u> , 1.9	1.8	0.02			0.036
	Starch (wet milling)	< 0.36, < <u>0.63</u> , < 0.93	< 0.63	-			0.0126
Sorghum (STMR: 0.02 mg/kg, MRL: 0.01* mg/kg)	AGF (grain sorghum)	26	26	-	26	26	0.52
	Syrup (sweet sorghum)	0.17	0.17	-	0.15	0.15	0.0015
Wheat (STMR: 0.095 mg/kg, MRL: 0.3 mg/kg)	AGF	200	200	-	180	180	17.1
	Bran	0.81, 2.4 ^a , <u>2.6</u> ^a , 2.8 ^a , 3.5 ^a	2.6	0.8	0.88, 2.4 ^a , <u>2.5</u> ^a , 2.7 ^a , 4 ^a	2.5	0.2375
	Flour	0.12, 0.2 ^a , <u>0.2</u> ^a , 0.25 ^a , 0.5 ^a	0.20	-	0.20, 0.33 ^a , <u>0.33</u> ^a , 0.4 ^a , 0.66 ^a	0.33	0.03135
	Germ	0.57, 0.6 ^a , <u>0.75</u> ^a , 1.2 ^a , 1.5 ^a	0.75	-	0.74, 0.8 ^a , <u>0.83</u> ^a , 1.3 ^a , 2.7 ^a	0.83	0.07885
Oilseed rape (STMR: 0.02 mg/kg, MRL: 0.04 mg/kg, based on 2016 JMPR)	Meal (extracted)	< 0.5 ^a , <u>0.72</u> , <u>0.82</u> , 2 ^a	0.77	-	0.66 ^a , <u>0.8</u> , <u>1.2</u> , 1.5 ^a	1.0	0.02
	Oil, crude	0.5 ^a , <u>1</u> ^a , <u>2.4</u> , 2.5	1.7	-	0.66 ^a , <u>1</u> ^a , <u>1.8</u> , 2.0	1.4	0.028
	Oil, refined	< 0.5 ^a , <u>2</u> ^a , <u>2.1</u> , 2.1	2.0	0.08	< 0.66 ^a , <u>1.4</u> ^a , <u>1.5</u> , 1.8	1.4	0.028
Peanut (STMR: 0.02 mg/kg, MRL: 0.01 mg/kg)	Meal	< 0.57	< 0.57	-	See bixafen	See bixafen	0.0114
	Refined oil (edible)	2.2	2.2	0.03			0.044

^a PF estimated by the 2013 JMPR

- no traded commodity or maximum residue level covered by raw agricultural commodity

In addition to the STMR-P values given in the table above, the Meeting estimated maximum residue levels of 0.3 mg/kg for soya bean hulls, 0.15 mg/kg for soya bean refined oil, 0.03 mg/kg for maize bran and 0.02 mg/kg for maize oil.

For wheat bran, the Meeting estimate a maximum residue level of 0.8 mg/kg to replace its previous recommendation of 0.15 mg/kg made by the 2016 JMPR.

For oilseed rape, new processing information were provided to the current Meeting. Based on the combined processing factors including those from the 2016 and 2020 JMPR, the Meeting confirms its previous maximum residue level recommendation of 0.08 mg/kg for bixafen in rapeseed oil, edible.

For peanut oil, edible, the Meeting estimated a maximum residue level of 0.03 mg/kg.

Residues in animal commodities

Farm animal feeding studies

No additional information on farm animals was provided to the current Meeting. Please refer to the 2013 JMPR Report and Evaluation.

Estimated maximum and mean dietary burdens of livestock and animal commodities maximum residue levels

Dietary burdens were calculated for beef cattle, dairy cattle, broilers and laying poultry based on feed items evaluated by the JMPR. The dietary burdens, estimated using the estimated using the most recent version of the OECD livestock dietary burden calculator, are presented in Annex 6 and summarised below.

Table 5 Estimated maximum and mean dietary burdens of farm animals

	Livestock dietary burden for the estimation of maximum residue levels and the dietary exposure, total bixafen, ppm of dry matter diet							
	US-Canada		EU		Australia		Japan	
	max.	mean	max.	Mean	max.	mean	max.	mean
Beef cattle	4.0	2.5	8.6	4.2	29 ^a	14 ^c	0.14	0.14
Dairy cattle	8.4	4.1	8.5	4.1	27 ^b	13 ^d	1.3	0.7
Poultry - broiler	0.18	0.18	0.25	0.21	0.17	0.17	0.16	0.16
Poultry - layer	0.18	0.18	3.2 ^e	1.6 ^f	0.17	0.17	0.14	0.14

^a Highest maximum beef cattle burden suitable for MRL estimates and dietary exposure for mammalian meat

^b Highest maximum dairy cattle burden suitable for MRL estimates for milk

^c Highest mean beef cattle burden suitable for dietary exposure for mammalian meat

^d Highest mean dairy cattle burden suitable for dietary exposure for milk

^e Highest maximum broiler or laying hen burden suitable for MRL estimates and dietary exposure for poultry products and eggs

^f Highest mean broiler or laying hen burden suitable for MRL estimates and dietary exposure for poultry products and eggs

Animal commodities maximum residue levels

In comparison to the estimations made by the 2016 JMPR, the dietary burden remained unchanged for beef and dairy cattle after including the uses considered by the current Meeting and increased insignificantly for laying hens (Max: 3.0 ppm → 3.2 ppm and mean: 1.5 ppm → 1.6 ppm).

Therefore, the Meeting confirms its previous recommendation for bixafen in animal commodities.

RECOMMENDATIONS

On the basis of the data obtained from supervised trials, the Meeting concluded that the residue levels listed in Annex 1 are suitable for establishing maximum residue limits and for IEDI and IESTI assessments.

Definition of the residue for compliance with MRL for plant commodities: *bixafen*

Definition of the residue for compliance with MRL for animal commodities and (for the estimation of dietary intake) for plant and animal commodities: *sum of bixafen and N-(3',4'-dichloro-5-fluorobiphenyl-2-yl)-3-(difluoromethyl)-1H-pyrazole-4-carboxamide (bixafen-desmethyl), expressed as bixafen*

The residue is fat-soluble.

Dietary risk assessment

Long-term dietary exposure

The ADI for bixafen is 0–0.02 mg/kg bw. The International Estimated Daily Intakes (IEDIs) for bixafen were estimated for the 17 GEMS/Food Consumption Cluster Diets using the STMR or STMR-P values estimated by the JMPR. The results are shown in Annex 3 of the 2021 Extra JMPR Report.

The IEDIs ranged from 4–10% of the maximum ADI. The Meeting concluded that long-term dietary exposure to residues of bixafen from uses considered by the JMPR is unlikely to present a public health concern.

Acute dietary exposure

The ARfD for bixafen is 0.2 mg/kg bw. The International Estimate of Short Term Intakes (IESTIs) for bixafen were calculated for the food commodities and their processed commodities for which HRs/HR-Ps or STMRs/STMR-Ps were estimated by the present Meeting and for which consumption data were available. The results are shown in Annex 4 of the 2021 Extra JMPR Report.

The IESTIs for children were 0–20% of the ARfD and for adults 0–20% of the ARfD. The Meeting concluded that the acute dietary exposure to residues of bixafen from the uses considered by the present Meeting is unlikely to present a public health concern.

Assessment of metabolites using the threshold of toxicological concern (TTC) approach

The tautomeric mixture M44/M45 could be assessed using the TTC approach (Cramer Class III threshold of 1.5 µg/kg bw per day) as concluded by the 2013 Meeting.

The current Meeting considered residues of M44/M45 in rotational crops resulting in a dietary exposure estimate of 0.31 µg/kg bw per day.

The Meeting concluded that the estimated dietary exposure to residues of the tautomeric mixture M44/M45 from uses considered by the JMPR is below the TTC for Cramer Class III compounds and is unlikely to present a public health concern. Should further uses be considered in the future, these conclusions may need to be re-evaluated.

5.3. CHLORPYRIFOS (017) (R)

RESIDUE AND ANALYTICAL ASPECTS

Chlorpyrifos is a non-systemic organophosphorus insecticide with contact, stomach and respiratory action. It was first evaluated by the JMPR in 1972 and the last periodic reviews were in 1999 for toxicology and in 2000 for residues. In 2004 and 2006, the JMPR evaluated additional uses.

The 1999 JMPR confirmed an ADI of 0–0.01 mg/kg bw for chlorpyrifos and established an ARfD of 0.1 mg/kg bw. In 2000, the JMPR defined the residue definition as chlorpyrifos for compliance with the MRL and dietary risk assessment for plant and animal commodities. The residue is fat-soluble.

Chlorpyrifos was scheduled at the Fifty-first Session of the CCPR for evaluation of additional uses by the 2020 JMPR, which was postponed to the 2021 Extra JMPR. The Meeting received information on GAP and residue trials on eggplant.

Methods of analysis

Chlorpyrifos residues in eggplant were analysed by a new method using QuEChERS sample preparation method and GC-ECD or GC-FPD. Recovery test results showed recoveries of 80–109%. The LOQ was 0.01 mg/kg. The analytical method used in the eggplant residue trials was considered sufficiently validated.

Stability of residues in stored analytical samples

Residue analysis was performed within 24 hours after sample collection.

Results of supervised residue trials on crops

Fruiting vegetables, other than Cucurbits

Eggplant

Chlorpyrifos is registered for use on eggplant in India with the critical GAP involving an application rate of 0.20 kg ai/ha with no specified PHI (the maximum number of foliar sprays and minimum re-treatment intervals are not specified).

Six decline trials (including a sample collection time of 0 days) were conducted in India during 2014–2015 at a rate of 0.20 kg ai/ha with 2 applications at a 10-day interval. The residue concentrations of chlorpyrifos in eggplant were (n = 6): 0.14, 0.22, 0.67, 0.98, 1.1 and 1.4 mg/kg (highest value of 1.5 mg/kg from replicate plots).

Communication by the sponsor indicated that the local agricultural practice involves re-treatment intervals of 3–6 days. Residues declined with a median half-life of 1.63 days (1st order) and the modelled residue at a spray interval of 3 days and 2 applications differed by more than 25% from the supervised field trials. Therefore, the Meeting concluded that the supervised field trials were conducted at a significantly longer re-treatment interval and cannot be used for the estimation of a maximum residue level.

5.4. CLOFENTEZINE (156)

RESIDUE AND ANALYTICAL ASPECTS

Clofentezine (3,6-bis (2-chlorophenyl)-1,2,4,5-tetrazine) (IUPAC) is an acaricide used for the control of red spider mites on a wide range of crops. It was evaluated for the first time by the 1986 JMPR and re-evaluated for residues several times up to 1992. Clofentezine underwent periodic review of the toxicology in 2005, where an ADI of 0–0.02 mg/kg bw was established and an ARfD was deemed unnecessary. The most recent periodic review of the residues was conducted by the 2007 JMPR.

The 2007 JMPR recommended the following residue definition for clofentezine:

Definition of the residue for compliance with the MRL and dietary risk assessment for plant commodities: *clofentezine*.

Definition of the residue for compliance with the MRL and dietary risk assessment for animal commodities: *sum of clofentezine and all metabolites containing the 2-chlorobenzoyl moiety, expressed as clofentezine*.

The residue is fat-soluble.

Clofentezine was scheduled at the Fifty-first Session of the CCPR for evaluation of additional uses by the 2020 JMPR, which was postponed to the 2021 Extra JMPR. The current Meeting received new information on use patterns for clofentezine in hops, supervised field trials supported by additional plant metabolism studies, analytical methods and storage stability data.

Plant metabolism

The fate of clofentezine in plants was last evaluated by the 2007 JMPR following foliar spray application to lemons, apples, peaches and grapes. A detailed assessment of these studies is presented in the Report of the 2007 JMPR. The current Meeting received an additional plant metabolism study on lettuce.

The metabolism of [tetrazine-3-¹⁴C]-clofentezine in lettuce was investigated under outdoor conditions by the application of one foliar spray at 0.28 kg ai/ha. The treatment was performed at BBCH 39 (rosette development complete). Mature lettuce leaves were sampled 21 days after the treatment.

Total radioactive residues (TRR) were 0.243 mg eq/kg (combustion measurement). The sample was extracted with acetonitrile, followed by a mixture of acetonitrile/water (1/1; v/v), which recovered ~91% of the TRR from the leaves. Post extraction solids amounted 7.4% of the TRR.

The identification of the radioactive residues revealed unchanged clofentezine as the only residue in lettuce, representing 100% of the radioactivity in the extract and 90.2% of the TRR.

Methods of analysis

The current Meeting received additional analytical methods for the determination of parent clofentezine and its metabolite 2-CBN (2-Chlorobenzonitrile) in hops.

Methods 11735 and 14116.6103 both rely on liquid extraction with acetonitrile, followed by solid-phase extraction for clean-up. Parent clofentezine is measured via LC-MS/MS (ESI+), while its metabolite 2-CBN is quantified by GC-MS. Following validation experiments, LOQs of 0.02 mg/kg and 0.05 mg/kg were demonstrated for clofentezine and 2-CBN, respectively.

Storage stability

The current Meeting received information on the storage stability of clofentezine and 2-CBN in dry hops, investigated together with the supervised field trial samples.

In fortified hops samples (homogenized dry cones) clofentezine was quickly degraded. Average percentages remaining compared to initial concentrations (normalized to 100%) at day 0

were 40% (86 days), 16% (182 days) and < 3% (386 days). No concurrent recovery tests have been conducted.




Figure 1 1st order decline of clofentezine residues in dry hops under freezer storage conditions

In addition, incurred residues in two field trial samples were investigated. In the re-analysis at 751–754 days after harvest, 125% and 107% of the first measurement (made 383–386 days after harvest) were recovered. Based on the decline kinetics in the storage stability samples, over 90% of the clofentezine present at harvest would have dissipated by the time the first analysis was made. Without an analysis much closer to harvest, the Meeting could not use the analysis from the field trial samples to ascertain stability of incurred residues during storage.

The Meeting concluded that clofentezine in dry hops cones is not stable during freezer storage. According to the decline observed in stored fortified samples, analysis of the frozen field trial samples should be conducted within one month to recover at least 70% of the initial residue concentration.

For 2-CBN, no decline was observed in fortified samples. The Meeting concluded, that 2-CBN is stable in dry hops cones for at least one year.

Definition of the residue

The current Meeting received new data on the metabolism of clofentezine following foliar treatment of lettuce.

Following foliar application to lettuce, parent clofentezine was the only residue identified and represented > 90% of the TRR.

The Meeting therefore confirmed its previous residue definition for compliance with the MRL and dietary risk assessment for plant commodities.

Results of supervised residue trials on crops

The Meeting received supervised trial data for applications of clofentezine on hops from the USA.

Hops, dry cones

Clofentezine is registered in the USA for the use on hops with a maximum GAP involving one foliar spray of 0.28 kg ai/ha and a PHI of 21 days.

Supervised field trials conducted in the USA on hops were provided approximating the critical GAP. However, several trial samples had been stored longer than one month and degradation of the residue during storage cannot be excluded. Also, two of the trials (11735.19-WA3 & 11735.19-WA4) were conducted in close proximity and on the same dates, these trials were therefore not considered as independent by the Meeting.

Residues of clofentezine in hops, dry cones from independent trials, where analysis was conducted within one month of harvest, were (n = 4): 1.7, 2.0, 2.4 and 2.7 mg/kg.

The Meeting estimated a maximum residue level of 7 mg/kg and an STMR of 2.2 mg/kg for clofentezine in hops, dry cones.

Residues in animal commodities

The additional use on hops does not affect the dietary burden of livestock animals. No re-evaluation of clofentezine residues in animal commodities was conducted.

RECOMMENDATIONS

On the basis of the data obtained from supervised trials, the Meeting concluded that the residue levels listed in Annex 1 are suitable for establishing maximum residue limits and for IEDI assessment.

Definition of the residue for compliance with the MRL and dietary risk assessment for plant commodities: *clofentezine*.

Definition of the residue for compliance with the MRL and dietary risk assessment for animal commodities: *sum of clofentezine and all metabolites containing the 2-chlorobenzoyl moiety, expressed as clofentezine*.

The residue is fat-soluble.

DIETARY RISK ASSESSMENT

Long-term dietary exposure

The ADI for clofentezine is 0–0.02 mg/kg bw. The International Estimated Daily Intakes (IEDIs) for clofentezine were estimated for the 17 GEMS/Food Consumption Cluster Diets using the STMR or STMR-P values estimated by the JMPR. The results are shown in Annex 3 of the 2021 Extra JMPR Report.

The IEDIs ranged from 0–4% of the maximum ADI. The Meeting concluded that long-term dietary exposure to residues of clofentezine from uses considered by the JMPR is unlikely to present a public health concern.

Acute dietary exposure

The 2005 JMPR decided that an ARfD for clofentezine was unnecessary. The Meeting therefore concluded that the acute dietary exposure to residues of clofentezine from the use considered by the Meeting is unlikely to present a public health concern.

5.5. CLOTHIANIDIN (238)

RESIDUE AND ANALYTICAL ASPECTS

Clothianidin (ISO common name) is a broad-spectrum, neonicotinoid insecticide with registered uses on multiple crops. Clothianidin is a major metabolite of thiamethoxam (245). Thiamethoxam was scheduled at the Fifty-first Session of the CCPR for evaluation of additional uses by the 2020 JMPR, which was postponed to the 2021 Extra JMPR. The Meeting also considered residues of clothianidin arising from the uses of thiamethoxam. The current Meeting received information on analytical methods, field trials and processing studies to support estimation of maximum residue levels in commodities of persimmon, barley, rice, sorghum, sweet corn and wheat.

Clothianidin was evaluated for the first time by the 2010 JMPR, which established an acceptable daily intake (ADI) of 0–0.1 mg/kg bw and an acute reference dose (ARfD) of 0.6 mg/kg bw. Clothianidin underwent subsequent evaluations by the JMPR in 2011, 2012 and 2014.

The definition of the residue for compliance with the MRLs and for dietary risk assessment for animal and plant commodities is clothianidin. The residue is not fat-soluble.

Methods of analysis

The Meeting received method validation and concurrent recovery data for use of Method AG-765 (reviewed by the 2010 JMPR) and the related Method GRM.009.02A, an unnamed method for analysis of residues in Japanese persimmon, as well as the QuEChERS multiresidue method. All methods were demonstrated to have adequate performance for recovery of clothianidin, with an LOQ of 0.01 mg/kg (0.02 mg/kg in Japanese persimmon).

Storage stability

No new information on the stability of clothianidin residues during frozen storage was provided. The 2010 JMPR had determined that residues of clothianidin are stable for 1–2 years under frozen (–18 °C) conditions for a large range of commodities, including the high-water and high-starch commodities considered by the current Meeting. The stability of residues in crops under consideration by the present Meeting is considered to be adequately demonstrated for the periods that field trial samples had been stored prior to analysis.

Results of supervised residue trials on crops

The Meeting received data from supervised residue trials generated following application of thiamethoxam to crops and GAP information for thiamethoxam on persimmon, wheat, barley, rice, sorghum (grain and sweet) and sweet corn. In making recommendations for maximum residues levels, the Meeting first determined whether a Codex MRL has been adopted based on the recommendation of previous JMPRs for residues of clothianidin, from the use of clothianidin on the crop. Under the assumption that growers are unlikely to use both thiamethoxam and clothianidin, both at the critical GAP, during any one growing season, the Meeting made a recommendation based on which source (application of clothianidin, *per se*, or residues arising from use of thiamethoxam) resulted in the highest residue estimates.

Japanese Persimmon

The critical thiamethoxam GAP for Japanese persimmon is from the Republic of Korea and consists of three applications, on a 10-day interval, of a 2000-fold dilution of the formulation (equivalent to 5 g thiamethoxam/hL). Applications are made to the point of run-off and there is a 7-day PHI.

Residues of clothianidin arising from use of thiamethoxam in independent trials provided to the current Meeting and approximating the critical GAP were (n = 2): < 0.02 (2) mg/kg.

The 2014 JMPR evaluated residues in Japanese persimmon in trials approximating the critical GAP. Residues of clothianidin arising from use of thiamethoxam in independent trials were (n = 2): < 0.02 (2) mg/kg.

The Meeting decided to combine data from the two sets of trials (n = 4): < 0.02 (4) mg/kg.

The maximum expected residue of clothianidin from the use of thiamethoxam on Japanese persimmon is < 0.02 mg/kg. The Meeting noted that a Codex MRL already exists for residues of clothianidin in Japanese persimmon, via the Group of Pome Fruits, at 0.4 mg/kg (STMR = 0.1 mg/kg, HR = 0.2 mg/kg). The Meeting confirmed its previous recommendation.

Wheat

Labels were provided for registrations of thiamethoxam in Mexico (2 foliar applications at 56 g ai/ha, 7-day RTI, 14-day PHI) and the USA (seed treatment at 52 g ai/100 kg seed + 2 foliar applications at 96 g ai/ha, 5-day RTI, 30-day PHI). Residue decline data indicated that for those two GAPs, the rate is the determining factor in residues at harvest rather than the RTI or the PHI. Therefore, the Meeting decided that the GAP from the USA is the critical GAP.

Trials in the USA approximated the US GAP for application rate and timing, but with a significantly shorter interval between the last application and harvest (14 days). The Meeting decided that the data are not suitable to support recommendations based on the US GAP. The Meeting agreed to consider the GAP from Mexico.

The Meeting determined that the seed treatment application used in the trials was not likely to contribute significantly to residues at harvest and decided that the US trials can be adjusted using proportionality to approximate the GAP from Mexico.

The Meeting noted that the results from the trial in Grand Island, NE (Trial C33-0170) are inconsistent within the residue decline data (< 0.01 mg/kg in the 7, 10, 14 and 17 DALA samples and 5.0 mg/kg in the 21 DALA sample) and in terms of the full set of available field trials (residue range = 0.015 to 0.18 mg/kg). The Meeting agreed not to consider data from that trial in any residue recommendations.

Residues (unscaled) of clothianidin arising from use of thiamethoxam in independent trials were (n = 16): < 0.01 (7), 0.012, 0.017, 0.018, 0.023, 0.026, 0.029, 0.034, 0.13 and 0.15 mg/kg.

When scaled (scaling factors range from 0.57 to 0.59), residues were (n = 16): < 0.01 (7), < 0.01, 0.01, 0.011, 0.013, 0.015, 0.017, 0.019, 0.075 and 0.087 mg/kg.

The Meeting noted that the Codex MRL for clothianidin in wheat grain of 0.02 mg/kg does not accommodate the expected clothianidin residues arising from the use of thiamethoxam considered by the current Meeting. Therefore, the Meeting used the data listed above to estimate a new maximum residue level of 0.15 mg/kg and an STMR of 0.01 mg/kg for wheat grain to replace its previous recommendation.

Based on the registered use including triticale, the Meeting decided to extrapolate the residue estimates to triticale.

Barley

Labels were provided for registrations of thiamethoxam in Mexico (2 foliar applications at 56 g ai/ha, 7-day RTI, 14-day PHI) and the USA (seed treatment at 52 g ai/100 kg seed + 2 foliar applications at 96 g ai/ha, 5-day RTI, 30-day PHI). Residue decline data indicated that for those two GAPs, the rate is the determining factor in residues at harvest rather than the RTI or the PHI. Therefore, the Meeting decided that the GAP from the USA is the critical GAP.

Trials in the USA approximated the US GAP for application rate and timing, but with a significantly shorter interval between the last application and harvest (14 days). The Meeting decided that the data are not suitable to support recommendations based on the US GAP. The Meeting agreed to consider the GAP from Mexico.

The Meeting determined that the seed treatment application used in the trials was not likely to contribute significantly to residues at harvest and decided that the US trials can be adjusted using proportionality to approximate the GAP from Mexico.

Residues (unscaled) of clothianidin arising from use of thiamethoxam in independent trials were (n = 12): < 0.01 (3), 0.015, 0.022, 0.025, 0.028, 0.041, 0.044, 0.049, 0.053 and 0.072 mg/kg.

When scaled (scaling factors range from 0.57 to 0.61), residues were (n = 12): < 0.01 (4), 0.013, 0.014, 0.016, 0.024, 0.026, 0.028, 0.031 and 0.042 mg/kg.

The Meeting noted that the Codex MRL for clothianidin in barley grain of 0.04 mg/kg does not accommodate the expected clothianidin residues arising from the use of thiamethoxam considered by the current Meeting. Therefore, the Meeting used the data listed above to estimate a new maximum residue level of 0.07 mg/kg and an STMR of 0.015 mg/kg for barley grain to replace its previous recommendation.

Based on the registered uses including oats and barley being an example representative commodity of the subgroup that includes oats, the Meeting decided to extrapolate the residue estimates to oats.

Rice

The critical thiamethoxam GAP for rice is from Japan and consists of two applications of a 1000x dilution in up to 1500 L/ha (equivalent to 97.5 g ai/ha) with a PHI of 14 days; a re-treatment interval is not specified.

Residues of clothianidin arising from use of thiamethoxam in husked rice from independent trials approximating the critical GAP were (n = 11): 0.07, 0.09, 0.11 (2), 0.12, 0.14, 0.16 (3), 0.18 and 0.26 mg/kg. Of those, two trials were done in paddy rice with residues of 0.12 and 0.14 mg/kg.

The Meeting noted that the Codex MRL for clothianidin in rice of 0.5 mg/kg (STMR = 0.145 mg/kg) accommodates the expected clothianidin residues arising from the use of thiamethoxam on upland and paddy rice considered by the current Meeting. Therefore, the Meeting confirmed its previous recommendations for residues of clothianidin in rice.

Grain sorghum

The critical thiamethoxam GAP for grain sorghum is from the USA and consists of three applications: a seed treatment at 2.97 g ai/100 kg seed followed by two in-season foliar applications, each at 96 g ai/ha, on a 7-day interval, with a PHI of 14 days.

Trials were conducted in the USA using the in-season applications only. The Meeting determined that residues of thiamethoxam coming from the seed treatment use were not likely to contribute significantly to residues at harvest.

Residues of clothianidin arising from use of thiamethoxam in grain sorghum from independent trials approximating the critical GAP were (n = 11): < 0.01 (4), 0.015, 0.018, 0.029, 0.031, 0.039, 0.046 and 0.078 mg/kg.

The Meeting noted that the Codex MRL for clothianidin in sorghum grain of 0.01 (*) mg/kg does not accommodate the expected clothianidin residues arising from the use of thiamethoxam considered by the current Meeting. Therefore, the Meeting used the data listed above to estimate a new maximum residue level of 0.15 mg/kg and an STMR of 0.018 mg/kg for sorghum grain to replace its previous recommendation.

Sweet corn

The critical thiamethoxam GAP for sweet corn is from the USA and consists of three applications: a seed treatment at 52 g ai/100 kg seed followed by two in-season foliar applications, each at 96 g ai/ha, on a 5-day interval, with a PHI of 1 day.

Residues of clothianidin arising from use of thiamethoxam in sweet corn (kernel + cob with husk removed) from independent trials approximating the critical GAP were (n = 12): < 0.01 (12) mg/kg.

The Meeting noted that a Codex MRL has been adopted for clothianidin residues in sweet corn (corn on the cob) at 0.01 (*) mg/kg (STMR, HR = 0.01 mg/kg). Furthermore, The Meeting noted that the only crop in the Subgroup of sweet corns is sweet corn and that the representative commodity is sweet corn (corn on the cob, kernels plus cob with husk removed) and agreed to estimate a new maximum residue level of 0.01(*) mg/kg, an STMR of 0.01 mg/kg and an HR of 0.01 mg/kg for residues of clothianidin in the Subgroup of sweet corns (Subgroup 020F) and withdrew its previous recommendation for sweet corn (corn on the cob).

Grasses for sugar or syrup production

Sweet sorghum

The critical thiamethoxam GAP for sweet sorghum is from the USA and consists of three applications: a seed treatment at 2.97 g ai/100 kg seed followed by two in-season foliar applications, each at 96 g ai/ha, on a 7-day interval, with a PHI of 14 days.

Trials were conducted using the in-season applications only. The Meeting determined that residues of clothianidin coming from thiamethoxam seed treatment use were not likely to contribute significantly to residues at harvest. Residues of clothianidin arising from use of thiamethoxam in sweet sorghum cane from independent trials approximating the critical GAP were (n = 4): 0.039, 0.062, 0.067 and 0.20 mg/kg.

The Meeting estimated a maximum residue level of 0.4 mg/kg, an HR of 0.2 mg/kg and an STMR of 0.0645 mg/kg for sorgho or sorghum, sweet.

Residues in animal feeds

Forage of cereal grains

Wheat

The critical thiamethoxam GAP for forage is from the USA (seed treatment at 52 g ai/100 kg seed + 2 foliar applications at 96 g ai/ha, 5-day RTI, 7-day PHI).

Residues of clothianidin arising from use of thiamethoxam in forage from trials in the USA approximating the US GAP were (n = 17): < 0.011, < 0.018, 0.017, 0.024 (2), 0.026, 0.043, 0.054, 0.060, 0.072, 0.078, 0.10, 0.13, 0.16, 0.17, 0.19 and 0.21 mg/kg.

The registration in the USA includes barley, oat, rye, triticale and wheat. The residue data (as received) for wheat forage listed above have a median residue of 0.06 mg/kg and a highest residue of 0.21 mg/kg. The Meeting agreed to extrapolate the median and highest residue estimates from wheat forage to forage of barley, oat, rye and triticale.

Sorghum

The critical thiamethoxam GAP for sorghum is from the USA and consists of three applications: a seed treatment at 2.97 g ai/100 kg seed followed by two in-season foliar applications, each at 96 g ai/ha, on a 7-day interval, with a PHI of 14 days.

Trials were conducted using the in-season applications only. The Meeting determined that residues of clothianidin coming from thiamethoxam seed treatment use were not likely to contribute significantly to residues at harvest. Residues of clothianidin arising from use of thiamethoxam in sorghum forage from independent trials approximating the critical GAP were (n = 10): 0.099, 0.10, 0.11, 0.14, 0.17, 0.21, 0.22, 0.26 (2) and 0.29 mg/kg (as received).

The Meeting estimated a median residue of 0.19 mg/kg (as received) and a highest residue of 0.29 mg/kg (as received) in sorghum forage to replace the previous estimates.

Sweet corn

The critical thiamethoxam GAP for sweet corn is from the USA and consists of three applications: a seed treatment at 52 g ai/100 kg seed followed by two in-season foliar applications, each at 96 g ai/ha, on a 7-day interval, with a PHI of 1 day for forages.

Residues of clothianidin arising from use of thiamethoxam in sweet corn forage from independent trials approximating the critical GAP were (n = 12): 0.017, 0.029, 0.037, 0.038, 0.044, 0.078, 0.09 (2), 0.1, 0.12 (2) and 0.22 mg/kg (as received).

The Meeting estimated a median residue of 0.084 mg/kg (as received) and a highest residue of 0.22 mg/kg (as received) in sweet corn forage to replace the previous estimates.

Hay of cereal grains

Wheat

The critical thiamethoxam GAP for hay is from the USA (seed treatment at 52 g ai/100 kg seed + 2 foliar applications at 96 g ai/ha, 5-day RTI, 7-day PHI).

Residues of clothianidin arising from use of thiamethoxam in wheat hay from trials in the USA approximating the US GAP were (n = 17): 0.015, 0.034, 0.058 (2), 0.064, 0.080, 0.091, 0.15, 0.20 (2), 0.23, 0.26, 0.31, 0.40, 0.43, 0.60 and 0.72 mg/kg.

Barley

The critical thiamethoxam GAP for hay is from the USA (seed treatment at 52 g ai/100 kg seed + 2 foliar applications at 96 g ai/ha, 5-day RTI, 7-day PHI).

Residues of clothianidin arising from use of thiamethoxam in barley hay from trials in the USA approximating the US GAP were (n = 12): 0.014, 0.027, 0.048, 0.053, 0.063, 0.067, 0.072, 0.089, 0.096, 0.10, 0.21 and 0.23 mg/kg.

The registration in the USA includes barley, oat, rye, triticale and wheat. The Meeting determined via a Mann-Whitney U Test that the residue data listed above for wheat and barley hay are not statistically different and agreed to combine the data. Residues of clothianidin in wheat and barley hay arising from thiamethoxam treatment approximating the US GAP were (n = 29): 0.014, 0.015, 0.027, 0.034, 0.048, 0.053, 0.058 (2), 0.063, 0.064, 0.067, 0.072, 0.080, 0.089, 0.091, 0.096, 0.10, 0.15, 0.20 (2), 0.21, 0.23 (2), 0.26, 0.31, 0.40, 0.43, 0.60 and 0.72 mg/kg.

Based on the registered uses and on the data for wheat and barley hay and on an OECD standard dry-matter content of 88%, the Meeting estimated a maximum residue level of level of 1 mg/kg (dw), a median residue of 0.09 mg/kg (as received) and a highest residue of 0.72 mg/kg (as received) in hay of wheat, barley, oat, rye and triticale.

Straw of cereal grains

Wheat

Labels were provided for registrations of thiamethoxam in Mexico (2 foliar applications at 56 g ai/ha, 7-day RTI, 14-day PHI) and the USA (seed treatment at 52 g ai/100 kg seed + 2 foliar applications at 96 g ai/ha, 5-day RTI, 30-day PHI).

For wheat straw, the trials in the USA did not match the USA GAP due to the straw being harvested 14 DALA rather than 30 days specified for the PHI. Therefore, the Meeting decided that the data are not suitable to support recommendations based on the US GAP. The Meeting agreed to consider the GAP from Mexico. As noted for wheat grain, the Meeting determined that residues of thiamethoxam coming from the seed treatment use were not likely to contribute significantly to

residues at harvest and that residues from the US trials can be adjusted using proportionality to approximate the GAP from Mexico.

Residues (unscaled) of clothianidin arising from use of thiamethoxam in wheat straw from independent trials were (n = 17): 0.016, 0.021, 0.028, 0.031, 0.071, 0.072, 0.084, 0.087, 0.090, 0.097, 0.10, 0.12, 0.24, 0.26, 0.29, 0.37 and 0.46 mg/kg.

When scaled (scaling factors range from 0.57 to 0.59), residues were (n = 17): < 0.01, 0.012, 0.016, 0.018, 0.041 (2), 0.049, 0.050, 0.052, 0.056, 0.059, 0.070, 0.14, 0.15, 0.17, 0.21 and 0.27 mg/kg.

Barley

Labels were provided for registrations of thiamethoxam in Mexico (2 foliar applications at 56 g ai/ha, 7-day RTI, 14-day PHI) and the USA (seed treatment at 52 g ai/100 kg seed + 2 foliar applications at 96 g ai/ha, 5-day RTI, 30-day PHI).

For barley straw, the trials in the USA did not match the USA GAP due to the straw being harvested 14 DALA rather than 30 days specified for the PHI. Therefore, the Meeting decided that the data are not suitable to support recommendations based on the US GAP. The Meeting agreed to consider the GAP from Mexico. As noted for barley grain, the Meeting determined that residues of thiamethoxam coming from the seed treatment use were not likely to contribute significantly to residues at harvest and that residues from the US trials can be adjusted using proportionality to approximate the GAP from Mexico.

Residues (unscaled) of clothianidin arising from use of thiamethoxam in barley straw from independent trials were (n = 12): < 0.01, 0.022, 0.032, 0.034, 0.041, 0.044, 0.053, 0.059, 0.067, 0.074, 0.085 and 0.11 mg/kg.

When scaled (scaling factors range from 0.57 to 0.59), residues were (n = 12): < 0.01, 0.013, 0.019, 0.020, 0.024, 0.026, 0.030, 0.034, 0.039, 0.042, 0.049 and 0.064 mg/kg.

The registration in Mexico includes barley, oat, triticale and wheat. Noting that residues appear to be similar between the straw data for wheat and barley, the Meeting agreed to explore a common set of residue estimates for these straw commodities. A Mann-Whitney U-Test indicated that the residue distributions from wheat and barley are significantly different. The OECD MRL calculator suggested maximum residue levels of 0.4 mg/kg (median = 0.054 mg/kg) for wheat straw and 0.1 mg/kg for barley (median = 0.028 mg/kg), all on an as-received basis.

The Meeting noted that the maximum residue level, median residue and highest residue estimates in hay are greater than those in straw and agreed to apply those estimates to straw. Furthermore, noting that the straw commodities of different cereal crops are generally indistinguishable from each other when in trade, the Meeting agreed to make recommendations for the straw of cereals included on the Mexico label (barley, oat, triticale, wheat).

Based on OECD standard dry-matter content of 88%, the Meeting estimated a maximum residue level of 1 mg/kg (dw), a median residue of 0.09 mg/kg (as received) and a highest residue of 0.72 mg/kg (as received) in straw and fodder of wheat, barley, oat and triticale.

The above recommendations replace the previous recommendations, each at 0.2 mg/kg, for barley straw and fodder and wheat straw and fodder.

Rice

The critical thiamethoxam GAP for rice is from Japan and consists of two applications of a 1000× dilution in up to 1500 L/ha (equivalent to 97.5 g ai/ha) with a PHI of 14 days; a re-treatment interval is not specified.

Residues of clothianidin arising from use of thiamethoxam in rice straw from independent trials approximating the critical GAP were (n = 11): 0.01 (2), 0.02 (2), 0.03 (3), 0.04, 0.05, 0.10 and 0.13 mg/kg.

Based on OECD dry-matter content of 88%, the Meeting estimated a maximum residue level of 0.2 mg/kg (dw), a median residue of 0.030 mg/kg (as received) and a highest residue of 0.13 mg/kg (as received) for residues of clothianidin in rice straw and fodder, dry.

Sorghum

The critical thiamethoxam GAP for sorghum is from the USA and consists of three applications: a seed treatment at 2.97 g ai/100 kg seed followed by two in-season foliar applications, each at 96 g ai/ha, on a 7-day interval, with a PHI of 14 days.

Trials were conducted using the in-season applications only. The Meeting determined that residues of clothianidin coming from thiamethoxam seed treatment use were not likely to contribute significantly to residues at harvest. Residues of clothianidin arising from use of thiamethoxam in sorghum stover from independent trials approximating the critical GAP were (n = 10): 0.038, 0.045, 0.067, 0.068, 0.077, 0.10, 0.17, 0.18, 0.20 and 0.46 mg/kg (as received).

The Meeting noted that the established Codex MRL for residues of clothianidin in sorghum straw and fodder (dry) of 0.01(*) mg/kg does not accommodate the expected clothianidin residues arising from the use of thiamethoxam considered by the current Meeting. Therefore, the Meeting used the data above to estimate a maximum residue level of 0.8 mg/kg (dw), a median residue of 0.0885 mg/kg (as received) and a highest residue of 0.46 mg/kg (as received) in sorghum straw and fodder, dry to replace the previous estimates.

Sweet corn

The critical thiamethoxam GAP for sweet corn is from the USA and consists of three applications: a seed treatment at 52 g ai/100 kg seed followed by two in-season foliar applications, each at 96 g ai/ha, on a 7-day interval, with a PHI of 26 days.

Residues of clothianidin arising from use of thiamethoxam in sweet corn stover from independent trials approximating the critical GAP were (n = 4): < 0.01 (3) and 0.021 mg/kg (as received).

Based on OECD standard dry-matter content of 83%, the Meeting estimated a maximum residue level of 0.05 mg/kg (dw), a median residue of 0.01 mg/kg (as received) and a highest residue of 0.021 mg/kg (as received) in sweet corn stover.

Fate of residues during processing

The Meeting received data showing the effect of processing wheat grain into aspirated grain fractions, bran, flour, middlings, shorts and germ; rice into hulls, bran and polished rice; grain sorghum into aspirated grain fractions and flour; and sweet sorghum into syrup. Processing factors and residue estimates are summarized below. The Meeting noted the conclusion of the 2010 Meeting that thiamethoxam was “essentially stable during the hydrolysis conditions simulating food processing conditions.” On that basis, the Meeting concluded that clothianidin is not likely to have formed and the processing factors directly reflected the changing concentrations of clothianidin during the processing of cereal grain RACs into processed commodities.

Table 1 Processing factors and residue estimates for clothianidin in wheat, sorghum and sweet sorghum commodities

Raw commodity	Residue in RAC, mg/kg			Processed commodity	Processing Factors		Residue in processed commodity, mg/kg		
	Max	STMR	HR		Individual	Best estimate	Max-P	STMR-P	HR-P
Wheat grain	3	0.01	--	Aspirated fractions	3.8, 10	6.9	--	0.069	--
				Bran	1.8, 1.9	1.85	5.55	0.0185	--
				Flour	0.6, 0.69	0.645	--	0.00645	--
				Middlings	0.78, 0.81	0.795	--	0.00795	--

Raw commodity	Residue in RAC, mg/kg			Processed commodity	Processing Factors		Residue in processed commodity, mg/kg		
	Max	STMR	HR		Individual	Best estimate	Max-P	STMR-P	HR-P
Sorghum grain	0.15	0.018	--	Shorts	0.89, 0.91	0.90	--	0.009	--
				Germ	1.8, 1.8	1.8	5.4	0.018	--
				Aspirated fractions	8.2, 17	12.6	--	0.227	--
				Flour	0.55, 0.72	0.635	--	0.0114	--
Sorghum cane	0.4	0.0645	--	Syrup	0.92	0.92	--	0.059	--

Based on the maximum residue estimates above and the OECD rounding classes, the Meeting estimated maximum residue levels of 6 mg/kg for wheat bran and 6 mg/kg for wheat germ. The Meeting decided to apply the processing factor for wheat bran (1.85) to barley, giving a maximum residue level of 0.15 mg/kg. The estimated STMR-Ps for the processed commodities considered by the Meeting are shown in Table 1. Furthermore, the Meeting decided to apply the processing factors for wheat bran (1.85) and flour (0.645) to barley and triticale, giving STMRs as follows:

Barley bran = 0.028 mg/kg,

Barley flour = 0.0097 mg/kg,

Triticale flour = 0.00645 mg/kg.

For rice commodities, the processing study provided residue analysis for rice grain, rice hulls, rice bran and polished rice, but not for husked rice. The Meeting derived a processing factor for husked rice by mathematically reconstituting the commodity based on residues in bran, residues in polished rice and their respective material balances reported in the processing study. Processing factors describing residues relative to the rice RAC (grain) and from husked rice are shown in Table 2.

Table 2 Processing factors and residue estimates for clothianidin in rice commodities

Raw commodity	Residue in starting commodity, mg/kg			Processed commodity	Processing Factors		Residue in processed commodity, mg/kg		
	Max	STMR	HR		Individual	Best estimate	Max-P	STMR-P	HR-P
Rice grain	See text below			Husked rice	0.38, 0.81	0.595	--	--	--
				Hulls	3.5, 5.0	4.25	--	--	--
				Bran	0.79, 1.4	1.1	--	--	--
				Polished rice	0.32, 0.68	0.5	--	--	--
Husked Rice	0.5 ^a	0.145 ^a		Hulls	6.24, 9.17	7.7	3.8	1.1	--
				Bran	1.81, 2.08	1.95	0.97	0.28	--
				Polished rice	0.833, 0.857	0.845	0.42	0.12	--

^a 2010 JMPR

For rice, the Meeting estimated maximum residue levels of 4 mg/kg for rice hulls, 1 mg/kg for rice bran and 0.5 mg/kg for polished rice. Furthermore, the Meeting used the residue estimates and the processing factor from rice grain to husked rice to derive residue estimates for rice grain. The Meeting estimated a maximum residue level of 0.9 mg/kg and a median residue of 0.3 mg/kg for rice grain.

Residues in animal commodities

The Meeting has added feed items and their associated residues to the dietary burden calculation used by the 2014 JMPR. Dietary burden calculations are provided in Annex 6; the dietary burden estimates are summarized below.

Table 3 Estimated maximum and mean dietary burdens of farm animals

Animal	Dietary burden estimates, ppm							
	Canada-US		European Union		Australia		Japan	
	Maximum	Mean	Maximum	Mean	Maximum	Mean	Maximum	Mean
Beef cattle	0.478	0.239	0.976	0.258	1.184 ^a	0.883 ^c	0.148	0.087
Dairy cattle	0.616	0.378	0.793	0.292	1.143 ^b	0.863 ^d	0.394	0.266
Broiler poultry	0.12	0.115	0.199	0.069	0.239	0.239	0.036	0.036
Laying hen	0.115	0.115	0.313 ^e	0.134	0.239	0.239 ^f	0.08	0.08

^a Highest maximum cattle dietary burden suitable for MRL estimates for mammalian tissues.

^b Highest maximum dairy cattle dietary burden suitable for MRL estimates for milk

^c Highest mean cattle dietary burden suitable for STMR estimates for mammalian tissues

^d Highest mean dairy cattle dietary burden suitable for STMR estimates for milk

^e Highest maximum poultry dietary burden suitable for MRL estimates for poultry tissues and eggs

^f Highest mean poultry dietary burden suitable for STMR estimates for poultry tissues and eggs

Farm animal feeding studies

Cattle

A cattle-feeding study with clothianidin was reviewed by the 2010 JMPR. Animals were dosed at 0.27, 0.8, or 2.6 ppm in the diet. Clothianidin residues were < 0.02 mg/kg in all samples. Based on the results of the feeding study with clothianidin and on the dietary burden, residues of clothianidin from consumption of clothianidin residues in feeds are expected to be < 0.02 mg/kg in all animal commodities.

Clothianidin residues in mammalian commodities may also arise from consumption of thiamethoxam in treated feeds. In a feeding study with thiamethoxam reviewed by the 2010 JMPR, animals were dosed at 2, 6, or 20 ppm in the diet. Clothianidin residues were as follows:

Table 4 Clothianidin residue levels in animal commodities from 2, 6 and 20 ppm dosage groups

Matrix	2 ppm in feed		6 ppm in feed		20 ppm in feed	
	Maximum	Mean	Maximum	Mean	Maximum	Mean
Muscle	< 0.01 mg/kg	< 0.01 mg/kg	< 0.01 mg/kg	< 0.01 mg/kg	< 0.01 mg/kg	< 0.01 mg/kg
Fat	Not analysed				< 0.01	< 0.01
Kidney	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01
Liver	0.049	0.039	0.14	0.12	0.38	0.27
Milk	0.006	< 0.005	0.02	0.014	0.07	0.043

The dietary burdens for thiamethoxam relevant for estimating residues of clothianidin from thiamethoxam in animal commodities (see section on thiamethoxam) and the anticipated residues are shown in Table 5.

Table 5 Anticipated residues of clothianidin from thiamethoxam in animal commodities

Thiamethoxam feeding study	Feed level (ppm) for milk/egg residues	Residues (mg/kg) in milk	Feed level (ppm) for tissue residues	Clothianidin Residues (mg/kg)			
				Muscle	Liver	Kidney	Fat
Maximum residue level beef or dairy cattle							
Feeding study	20	0.043	20	< 0.01	0.38	< 0.01	< 0.01
Dietary burden and high residue	20.6	0.044	20.6	< 0.01	0.39	< 0.01	< 0.01
STMR beef or dairy cattle							
Feeding study	6	0.014	6	< 0.01	0.12	< 0.01	--

Thiamethoxam feeding study	Feed level (ppm) for milk/egg residues	Residues (mg/kg) in milk	Feed level (ppm) for tissue residues	Clothianidin Residues (mg/kg)			
				Muscle	Liver	Kidney	Fat
	20	0.043	20	< 0.01	0.27	< 0.01	< 0.01
Dietary burden and residue estimate	18.8	0.041	18.8	< 0.01	0.257	< 0.01	< 0.01

The Meeting estimated new maximum residue levels for liver at 0.4 mg/kg and milks at 0.05 mg/kg to replace its previous recommendations and confirmed its previous recommendations of 0.02(*) mg/kg for meat (from mammals other than marine mammals); edible offal (mammalian), except liver; and mammalian fat.

Poultry

Residues of clothianidin in poultry commodities may occur following consumption of feed bearing residues of clothianidin as well as consumption of feed bearing residues of thiamethoxam. Poultry-feeding studies with clothianidin or thiamethoxam are not available. In the laying hen clothianidin metabolism study, test animals were dosed at 134 ppm in the feed. In the laying hen thiamethoxam metabolism study, test animals were dosed at either 98 or 112 ppm in the feed. The residues of clothianidin observed in those studies, as well as the estimated residues of clothianidin based on the ratio of the dose from the metabolism study and either the maximum or mean dietary burden for poultry are shown in Table 6.

Table 6 Estimated residues of clothianidin in poultry commodities following consumption of clothianidin or thiamethoxam in the diet

Test substance □	Metabolism study		Maximum		Mean	
	Clothianidin	Thiamethoxam	Clothianidin	Thiamethoxam	Clothianidin	Thiamethoxam
Dose/dietary burden	134 ppm	98 ppm	0.313 ppm	11.8 ppm	0.151 ppm	11.8 ppm
Commodity	Observed clothianidin residue, mg/kg		Estimated clothianidin residue, mg/kg		Estimated clothianidin residue, mg/kg	
Muscle	0.0465	0.0119	0.00011	0.0014	0.000052	0.0014
Fat + Skin	0.00742	0.0277	0.000017	0.0033	0.0000084	0.0033
Liver	0.165	3.071	0.00039	0.37	0.00019	0.37
Eggs	0.142	0.0514	0.00033	0.0062	0.00016	0.0062

The Meeting confirmed its previous recommendations of 0.01(*) mg/kg for poultry meat, poultry fat and eggs and made a new recommendation of 0.4 mg/kg for poultry edible offal, with an HR and STMR both at 0.37 mg/kg, to replace its previous recommendation.

RECOMMENDATIONS

On the basis of the data obtained from supervised trials, the Meeting concluded that the residue levels listed in Annex 1 are suitable for establishing maximum residue limits and for IEDI and IESTI assessments.

Definition of the residue for compliance with the MRL and dietary risk assessment for plant and animal commodities: *clothianidin*.

The residue is not fat-soluble.

DIETARY RISK ASSESSMENT***Long-term dietary exposure***

The ADI for clothianidin is 0–0.1 mg/kg bw. The International Estimated Daily Intakes (IEDIs) for clothianidin were estimated for the 17 GEMS/Food Consumption Cluster Diets using the STMR or STMR-P values estimated by the JMPR. The results are shown in Annex 3 of the 2021 Extra JMPR Report.

The IEDIs ranged from 0–2% of the maximum ADI. The Meeting concluded that long-term dietary exposure to residues of clothianidin from uses considered by the JMPR is unlikely to present a public health concern.

Acute dietary exposure

The ARfD for clothianidin is 0.6 mg/kg bw. The International Estimate of Short-Term Intakes (IESTIs) for clothianidin were calculated for the food commodities and their processed commodities for which HRs/HR-Ps or STMRs/STMR-Ps were estimated by the present Meeting and for which consumption data were available. The results are shown in Annex 4 of the 2021 Extra JMPR Report.

The IESTIs varied from 0–1% of the ARfD for children and for the general population. The Meeting concluded that acute dietary exposure to residues of clothianidin from uses considered by the present Meeting is unlikely to present a public health concern.

5.6. CYPERMETHRINS (including alpha- and zeta-cypermethrin) (118)

RESIDUE AND ANALYTICAL ASPECTS

Cypermethrin is a non-systemic pyrethroid insecticide with contact and stomach action. Cypermethrin was first evaluated by the JMPR in 1979 and periodic reviews were conducted in 2006 for toxicology and in 2008 for residues. Further evaluations for additional uses were conducted in 2009, 2011 and 2019.

For cypermethrins (including alpha- and zeta-cypermethrin), a group ADI of 0–0.02 mg/kg bw and a group ARfD of 0.04 mg/kg bw were established by the 2006 JMPR. In 2008, the JMPR established a residue definition of *cypermethrin (sum of isomers)* for compliance with the MRL and dietary risk assessment for plant and animal commodities. The residue is fat-soluble.

Cypermethrin was scheduled at the Fifty-first Session of the CCPR for evaluation of additional uses by the 2020 JMPR, which was postponed to the 2021 Extra JMPR. The Meeting received information on GAP and residue trials on eggplant.

Methods of analysis

Analysis of cypermethrin in eggplant samples was conducted by a new method using QuEChERS sample preparation and GC-ECD. Recovery test results showed recoveries of 83–105%. The LOQ was 0.05 mg/kg. The analytical method used in the eggplant residue trials was considered sufficiently validated.

Stability of residues in stored analytical samples

Residue analysis was performed within 24 hours after sample collection.

Results of supervised residue trials on crops

Fruiting vegetables, other than Cucurbits

Eggplant

The critical GAP for the use of cypermethrin on eggplants in India is foliar spraying with 0.07 kg ai/ha and a 3-day PHI (the maximum number of sprays and minimum re-treatment intervals are not specified).

Seven residue trials on eggplant (decline trials) were conducted in India during 2015–2017, at a rate of 0.050 kg ai/ha with 2 applications at a 10-day interval and a PHI of 3 days. The residue concentrations of cypermethrin in eggplant were ($n = 7$): < 0.05, < 0.05, 0.090, 0.10, 0.11, 0.18 and 0.57 mg/kg (highest value of 0.64 mg/kg from replicate plots). The trials did not match the application rate of the critical GAP.

Communication by the sponsor indicated that the local agricultural practice involves re-treatment intervals of 3–6 days. Residues declined with a median half-life of 2.35 days (1st order) and the modelled residue at a spray interval of 3 days and 2 applications differed by more than 25% compared to the supervised field trials. The Meeting concluded that the supervised field trials were conducted at a significantly longer re-treatment interval in combination with lower application rates and cannot be used for the estimation of a maximum residue level.

5.7. CYPRODINIL (207)

RESIDUE AND ANALYTICAL ASPECTS

Cyprodinil is a broad-spectrum fungicide belonging to the anilinopyrimidine group used to control a range of pathogens including *Tapesia yallundae*, *Botrytis spp.*, *Alternaria spp.* and *Rhynchosium secalis*. Cyprodinil was first evaluated by the JMPR in 2003 when an ADI of 0–0.03 mg/kg bw was established. The 2003 JMPR determined that an ARfD was unnecessary. The residue definition for both plants and animal commodities, for both compliance with the MRL and dietary risk assessment, is cyprodinil. The residue is fat-soluble.

The Forty-third Codex Alimentarius Commission (2020) approved the new work proposals including the priority list of pesticides for evaluation by the current Meeting. The priority list included cyprodinil for evaluation of additional uses on peas, beans and ginseng. The current Meeting received information on supervised field trials in beans, peas and ginseng, processing studies for ginseng processed commodities as well as information on analytical methods and storage stability.

Methods of analysis

The analytical methods (REM 141.10, CGA 219417 and GRM010.02A) used in the legume and pulse trials received by the Meeting were considered the same as those evaluated by previous meetings (the 2013, 2015 and 2017 JMPR). These methods were validated for the analysis of cyprodinil in beans (whole plant, green with pods and remaining plant) and dry peas with a validated LOQ of 0.01 mg/kg and in dry beans at an LOQ of 0.02 mg/kg.

The current Meeting received method validation data for use of Method JLND2020RS002. The method was demonstrated to have adequate performance for recovery of cyprodinil in ginseng and ginseng processed commodities. An LOQ of 0.01 mg/kg was validated for ginseng and its processed fractions.

Stability of pesticide residues in stored analytical samples

The stability of residues of cyprodinil in stored frozen samples was previously evaluated by the 2003, 2013, 2015 and 2017 JMPR. Stability for at least 2 years has been demonstrated for high-water, high-starch and high-acid commodities and for at least 9 months in high-oil commodities. Samples of legume vegetables were stored frozen for up to ca. 1 year, which is covered by the available data. Although data specific to a high-protein commodity have not been evaluated, the Meeting decided the conclusion could be extrapolated to pulses, which were stored for up to 1 month (dry beans) or up to 8.2 months (dry peas). New storage stability data were provided for cyprodinil in ginseng and its processed fractions. The Meeting concluded that cyprodinil is stable at -18 °C for at least 277 days for ginseng, dried ginseng and red ginseng, which covered the longest storage intervals of field sampling and laboratory analysis in ginseng supervised trials.

Results of supervised residue trials on crops

The Meeting received GAP information for cyprodinil on legume vegetables in Europe (Latvia and Spain), on pulses in Canada and on ginseng and its processed fractions in China. Supervised residues trials were provided for beans (with pods), peas and beans (dried) and ginseng.

Peas (with pods)

The registered GAP in Spain for the use of cyprodinil on protected peas with pods is for two applications at up to 375 g ai/ha with application intervals of 14 days and a minimum PHI of 3 days. The Meeting received information concerning a GAP in Latvia for cyprodinil on unprotected peas with pods involving three applications at up to 375 g ai/ha with application intervals of 10–14 days and a minimum PHI of 14 days.

No residue trials on peas with pods were available to support the estimation of a maximum residue level for peas with pods. The Meeting noted that beans with pods is considered a

representative crop for Codex subgroup 14B (Peas with pods) and that residue trial data on beans could be used to support a recommendation for peas with pods.

In eight supervised residue trials matching the Latvian GAP, residues of cyprodinil were (n = 8): 0.01, 0.03 (2), 0.04 (2), 0.06, 0.13 and 0.25 mg/kg.

In trials approximating the GAP from Spain evaluated by the 2017 JMPR, residues of cyprodinil in beans with pods were (n = 9): 0.34, 0.47, 0.54, 0.58, 0.60, 0.61, 0.75, 0.83 and 1.2 mg/kg.

The Meeting agreed that the GAP in Spain is the critical GAP and decided to extrapolate the recommendation of the 2017 JMPR for the Subgroup of beans with pods to the Subgroup of peas with pods.

The Meeting estimated a maximum residue level of 2 mg/kg and an STMR of 0.60 mg/kg for the Subgroup peas with pods.

Pulses

Beans (dry)

A Codex MRL has been established for residues of cyprodinil in dry beans at 0.2 mg/kg.

The registered GAP in Canada for the use of cyprodinil on dry beans and peas is for three applications at up to 365 g ai/ha with application intervals of 7 days and a minimum PHI of 7 days.

Five supervised residue trials with cyprodinil on beans were conducted in Canada and matching the Canadian GAP. The residues in dried beans were (n = 5): < 0.02, < 0.02, < 0.02, 0.025 and 0.038 mg/kg.

The Meeting noted that the residues in dried beans from the Canadian GAP are accommodated by the existing MRL; however, the Meeting agreed to extrapolate the recommendation made by the 2013 JMPR for beans, dry to the Subgroup of dry beans. The Meeting estimated a maximum residue level of 0.2 mg/kg and an STMR of 0.03 mg/kg for residues of cyprodinil in the Subgroup of dry beans, except soya beans and withdrew its previous recommendation for beans (dry).

Peas (dry)

The critical GAP for the use of cyprodinil on dry peas in Canada consists of three applications at up to 365 g ai/ha with application intervals of 7 days and a PHI of 7 days.

Seven supervised residue trials with cyprodinil on peas were conducted in Canada matching the critical GAP. Residues of cyprodinil in dried peas were (n = 7): 0.037, 0.047, 0.048, 0.054, 0.080, 0.11 and 0.12 mg/kg.

The Meeting estimated a maximum residue level of 0.2 mg/kg and an STMR of 0.054 mg/kg for residues of cyprodinil in the Subgroup of dry peas.

Ginseng

The critical GAP for the use of cyprodinil on ginseng in China consists of two foliar applications at up to 450 g ai/ha with application intervals of 7 days and a minimum PHI of 28 days.

Five supervised residue trials matching the critical GAP were conducted in China. Residues of cyprodinil on ginseng were (n = 5): 0.034, 0.037, 0.045, 0.115 and 0.120 mg/kg.

The Meeting estimated a maximum residue level for cyprodinil in ginseng of 0.3 mg/kg and an STMR of 0.045 mg/kg.

Residues in animal Feeds

Beans and Peas (vines)

The registered GAP in Spain for the use of cyprodinil on protected beans and peas (with pods) is for two applications at up to 375 g ai/ha with application intervals of 14 days and a minimum PHI of 3 days. The Meeting received information concerning a GAP in Latvia for cyprodinil on unprotected beans and peas (with pods) for three applications at up to 375 g ai/ha with application intervals of 10–14 days and a minimum PHI of 14 days.

No trials were available matching the GAP from Spain. Residues of cyprodinil in bean plants from trials matching the Latvian GAP were (n = 8): 0.05 (2), 0.07, 0.08, 0.12, 0.29, 0.32, 0.56 mg/kg. The meeting estimated a median residue of 0.1 mg/kg and highest residue of 0.56 mg/kg for bean vines and pea vines.

Fate of residues during processing

The Meeting received data showing the effect of processing ginseng into dried ginseng and red ginseng.

In the supervised residue trials with cyprodinil on ginseng discussed above, samples of ginseng were processed into dried ginseng and red ginseng. The Meeting selected the higher of the dried or red ginseng residues for making residue estimates. Residues of cyprodinil in processed ginseng were (n = 5): 0.041, 0.062, 0.114, 0.175 and 1.4 mg/kg.

The Meeting estimated a maximum residue level of 3 mg/kg in dried and red ginseng and an STMR-P of 0.114 mg/kg.

Farm animal dietary burdens

The potential dietary burdens of cyprodinil in animal diets were calculated from the highest and median residue levels of animal feeds, estimated using the most recent version of the OECD livestock dietary burden calculator. The maximum and mean dietary burdens were estimated for beef and dairy cattle, laying hens and broiler hens in Australia, Europe, Japan and USA-Canada.

Dietary burdens for cyprodinil are presented in Annex 6 of the 2021 Extra JMPR Report and summarised in Table 2, below, for all uses of cyprodinil on feed items evaluated by this and previous Meetings.

Table 2 Estimated maximum and mean dietary burdens of farm animals for Cyprodinil (ppm)

	USA-Canada		EU		Australia		Japan	
	Max	Mean	Max	Mean	Max	Mean	Max	Mean
Beef cattle	1.084	0.470	15.12	1.934 ^b	6.667	1.476	0.702	0.702
Dairy cattle	2.499	1.030	14.82	1.518	25.32 ^a	1.804 ^c	0.382	0.382
Poultry – broiler	0.690	0.690	1.342	0.672	0.200	0.200	0.079	0.079
Poultry - layer	0.554	0.554	2.265 ^d	0.756 ^e	0.200	0.200	0.160	0.160

^a Highest maximum beef or dairy cattle dietary burden used for mammalian milk and tissue MRL calculations.

^b Highest mean beef or dairy cattle dietary burden used for mammalian tissue STMR estimates.

^c Highest mean dairy cattle dietary burden used for mammalian milk STMR estimates.

^d Highest maximum poultry dietary burden used for poultry tissue and egg MRL calculations.

^e Highest mean poultry dietary burden used for poultry tissue and egg STMR estimates.

Residues in animal commodities

Maximum residue level and STMR estimates for poultry and ruminant commodities from previous meetings were based on maximum / mean dietary burdens of 23.3 / 1.8 ppm in cattle and 4.1 / 0.76 ppm in poultry. The current Meeting concluded that the animal dietary burden had changed insignificantly (or had decreased) and confirmed the previous recommendations.

RECOMMENDATIONS

On the basis of the data obtained from supervised trials, the Meeting concluded that the residue levels listed in Annex 1 are suitable for establishing MRLs and for IEDI assessment.

Definition of the residue for compliance with the MRL and dietary risk assessment for plant and animal commodities: *cyprodinil*.

The residue is fat-soluble.

DIETARY RISK ASSESSMENT***Long-term dietary exposure***

The 2003 JMPR set an ADI of 0–0.03 mg/kg bw for cyprodinil. The International Estimated Daily Intakes (IEDIs) for cyprodinil were estimated for the 17 GEMS/Food Consumption Cluster Diets using the STMR or STMR-P values estimated by the JMPR. The results are shown in Annex 3 of the 2021 Extra JMPR Report.

The IEDIs ranged from 7–70% of the maximum ADI. The Meeting concluded that long-term dietary exposure to residues of cyprodinil from uses considered by the JMPR is unlikely to present a public health concern.

Acute dietary exposure

The 2003 JMPR decided that an ARfD was unnecessary. Therefore, the Meeting concluded that the acute dietary exposure to residues of cyprodinil from the uses considered is unlikely to present a public health concern.

5.8. DIFENOCONAZOLE (224)

RESIDUE AND ANALYTICAL ASPECTS

Difenoconazole (ISO common name) is a broad-spectrum conazole (triazole) fungicide used for the control of diseases in multiple crops. Its fungicidal mode of action is as a sterol demethylation inhibitor. Difenoconazole was evaluated for the first time by JMPR 2007 when an acceptable daily intake (ADI) of 0–0.01 mg/kg bw and an acute reference dose (ARfD) of 0.3 mg/kg bw were established. In 2007, 2010, 2013, 2015 and 2017, the JMPR evaluated the compound for residues and recommended a number of maximum residue levels.






The definition of the residue for both compliance with the MRL and dietary risk assessment for plant commodities is: *difenoconazole* and


The definition of the residue for both compliance with the MRL and dietary risk assessment for animal commodities is: *the sum of difenoconazole and 1-[2-chloro-4-(4-chlorophenoxy)-phenyl]-2-(1,2,4-triazol)-1-yl-ethanol (CGA205375), expressed as difenoconazole.*

The residue is fat-soluble.

Difenoconazole was scheduled at the Fifty-first Session of the CCPR for evaluation of additional uses by the 2020 JMPR, which was postponed to the 2021 Extra JMPR. The current Meeting received storage stability, GAP information, field trials and processing studies to support maximum residue level estimations in commodities of cotton, cranberry, guava and tea. In addition, the current Meeting also received seed treatment and confined rotational crop metabolism studies.

The following compounds are discussed in this document:

Common or code name	Chemical name molecular formula molar mass, g/mol	Structure
Difenoconazole	1-({2-[2-chloro-4-(4-chlorophenoxy)phenyl]-4-methyl-1,3-dioxolan-2-yl} methyl)-1 <i>H</i> -1,2,4-triazole C ₁₉ H ₁₇ Cl ₂ N ₃ O ₃ 406.26	
CGA205375	1-[2-chloro-4-(4-chlorophenoxy)phenyl]-2-(1 <i>H</i> -1,2,4-triazol-1-yl)ethan-1-ol C ₁₆ H ₁₃ Cl ₂ N ₃ O ₂ 350.20	
CGA189138	2-chloro-4-(4-chlorophenoxy)benzoic acid C ₁₃ H ₈ Cl ₂ O ₃ 283.11	
CGA131013 Triazolylalanine	2-amino-3-(1 <i>H</i> -1,2,4-triazol-1-yl)propanoic acid C ₅ H ₈ N ₄ O ₂ 156.14	
CGA142856 Triazolylacetic acid	(1 <i>H</i> -1,2,4-triazol-1-yl)acetic acid C ₄ H ₅ N ₃ O ₂ 127.10	

Common or code name	Chemical name molecular formula molar mass, g/mol	Structure
CGA205369 Triazolylactic acid	2-hydroxy-3-(1 <i>H</i> -1,2,4-triazol-1-yl)propanoic acid $C_5H_7N_3O_2$ 157.13	

Plant metabolism

Wheat (seed treatment)

In a study provided to the current Meeting, wheat grain was treated with either [^{14}C -chlorophenoxy] or [^{14}C -triazolyl]difenoconazole, each at ca. 32 mg ai/100 g seed. Treated seeds were container-grown in a greenhouse and samples of forage (BBCH 14–21), hay (BBCH 56–65) and straw and grain (maturity) were analysed for TRR and extracted to determine the metabolic profile of the residue. Total radioactive residues were about 2- to 3-fold higher from the triazolyl treatment in forage (0.18 vs. 0.097 mg eq/kg), hay (0.11 vs. 0.03 mg eq/kg) and straw (0.31 vs. 0.11 mg eq/kg) samples and approximately 80-fold higher in grain (0.42 vs. 0.005 mg eq/kg). The extractability of residues into acetonitrile/water ranged from 64 to 78% in straw, from 80 to 90% in hay and approximately 90% in forage and grain. In total, approximately 70 to 96% of the unextracted radioactive residues in straw consisted of cellulose, lignin and hemicellulose. Analyses of these components was not undertaken for other commodities.

In terms of specific residues, concentrations of parent difenoconazole were low in all commodities, ranging from not detected to 0.006 mg/kg. Following treatment with chlorophenoxy-labelled compound, the major residue was CGA 205375 (forage: 47% TRR, 0.07 mg eq/kg; hay: 60% TRR, 0.018 mg eq/kg; straw: 12% TRR, 0.013 mg eq/kg), with ca. 85% of the residue in forage and hay occurring as the glycoside-conjugated form (which was not detected in straw). Grain samples from the chlorophenoxy-labelled treatment were not extracted due to low radioactivity. Treatment with triazolyl-labelled difenoconazole resulted, again, in CGA 205375 (free and conjugated) as a major residue in forage, hay and straw (forage: 61% TRR, 0.133 mg eq/kg; hay: 23% TRR, 0.026 mg eq/kg; straw: 11% TRR, 0.03 mg eq/kg), but was not detected in grain; the residue consisted primarily (ca. 85%) in a glycoside-conjugate form in forage and hay. Other major residues observed following treatment with triazolyl-labelled compound were triazolylalanine (CGA131013) (forage: 6.4% TRR, 0.014 mg eq/kg; hay 16% TRR, 0.018 mg eq/kg; straw: 11% TRR, 0.031 mg eq/kg; grain: 27% TRR, 0.11 mg eq/kg) and 1,2,4-triazolylacetic acid (forage: 9.2% TRR, 0.020 mg eq/kg; hay: 30% TRR, 0.034 mg eq/kg; straw: 41% TRR, 0.113 mg eq/kg; grain: 33% TRR, 0.13 mg eq/kg). Unidentified compounds were reported as being $\leq 10\%$ TRR from either treatment in all matrices; however, up to approximately 30% of the radioactivity was not accounted for.

The Meeting noted that metabolism of difenoconazole by plants following foliar (tomato, potato, grape, oilseed rape) and seed-treatment (wheat) applications was evaluated by the 2007 JMPR. Overall, the residues were described as low amounts of difenoconazole and higher amounts of the metabolite CGA 205375. The 2007 Meeting noted that ‘The presence of triazolylalanine and triazolylacetic acid was suggested but not confirmed for extracts of the mature grain (triazole label).’ The results of the current study are generally in good agreement with those from the studies reviewed by the 2007 Meeting. The Meeting decided that the new study does not impact the established residue definitions for difenoconazole.

Environmental fate

Confined rotational crops

In the study provided to the current Meeting, difenoconazole was labelled only in the chlorophenyl moiety. The study was conducted by treating bare soil at a rate of ca. 0.52 kg ai/ha (approximately 0.5X the maximum seasonal rate for most crops). Lettuce, radish, sorghum and wheat were planted into treated soil 30, 60, 120 and 270 days after the application. Total radioactive residues at the 30-day plant-back interval (PBI) ranged from 0.01 mg eq/kg to 0.083 mg eq/kg. In the majority of analysed crop matrices, TRR levels showed little to no decline with increasing plant-back intervals (PBI) of 30–270 days. The exceptions were lettuce and wheat straw, where residues from the 270-day PBI were ca. 2- to 4-fold lower than those at the 30-day PBI. TRRs in wheat grain were very low (0.004–0.006 mg eq/kg) and did not decline with increasing PBI. In most crops, the proportion of unextracted residues (acetonitrile:water; 4:1, v/v) increased with longer PBIs.

In immature and mature lettuce, radish leaves and radish roots, difenoconazole was the principal residue at all PBIs. Difenoconazole was a minor residue in sorghum forage and wheat straw and was not detected in sorghum stover. CGA 205375 and CGA 189138 were minor residues in lettuce and sorghum samples. CGA 205375 was a major residue in radish leaves and roots (generally 15–22% TRR) but occurred at low levels (< 0.007 mg eq/kg). CGA 205375 was also a major residue in wheat straw (ca. 40% TRR, ca. 0.02 mg eq/kg).

The 2007 Meeting also reviewed confined rotational crop studies (phenyl and triazolyl label; lettuce, wheat, maize, sugar beet, mustard, turnips). The TRR from those studies were generally low (< 0.01 mg eq/kg) following treatment with the phenyl label and higher (0.01–0.34 mg eq/kg) from the triazolyl label. The predominant residues were triazolylalanine (10–66% TRR), triazolylacetic acid (0–39% TRR) and triazolylactic acid (0–54% TRR), depending on the matrix.

The 2007 JMPR had noted that the confined rotational crops studies indicated that the water-soluble and mobile metabolites triazolylalanine, triazolylacetic acid and triazolylactic acid would be the predominant residues in rotational crops. Limited field rotational crop studies evaluated by the 2007 JMPR showed absence of residues of triazolylalanine and difenoconazole above their respective LOQs in carrots and spinach as following crops. While the available confined rotational crop studies indicate that difenoconazole may occur in rotational crops, the current Meeting concludes, based on a comparison of residues in confined and field rotational crop studies, that residues of difenoconazole above the LOQ are unlikely to occur.

Methods of analysis

The Meeting received method validation and concurrent recovery data for use of the QuEChERS multiresidue method to analyse residues in guava and concurrent recovery data for use of Method REM 147.08 on cranberry, cotton and tea commodities. Method REM 147.08 was previously reviewed by the 2007, 2010, 2015 and 2017 Meetings. Both methods were demonstrated to have adequate performance for recovery of difenoconazole, with an LOQ of 0.01 mg/kg.

The Meeting also received method validation and concurrent recovery data for Method Meth-160, Revision #2. This method was used for analysis of 1,2,4-triazole, triazolylalanine and triazolylacetic acid in cotton and tea commodities. All three analytes were validated with an LOQ of 0.01 mg/kg in cotton seed and gin by-products. Data were insufficient (too few replicates) to establish a validated LOQ in cotton meal, seed hulls and refined oil; nevertheless, the Meeting noted that recoveries of all three analytes were good ($\geq 89\%$) at the lowest fortification levels (0.01 mg/kg for 1,2,4-triazole and triazolylacetic acid, 0.1 mg/kg for triazolylalanine).

The Meeting decided that the methods used for data collection of metabolite residues are fit for purpose for all analytes.

Stability of pesticide residues in stored analytical samples

Storage stability data were provided for difenoconazole residues in oranges, dried beans, guava and tea. In addition, storage stability data were provided for 1,2,4-triazole; triazolyalanine, triazolyacetic acid and triazollylactic acid residues in tea.

Residues of all analytes were shown to be stable as follows:

Difenoconazole

Oranges: at least 732 days (24 months),

Dried beans: at least 732 days (24 months),

Guava: at least 398 days (13 months),

Dry tea leaves: at least 372 days (12 months),

1,2,4-triazole; triazolyalanine, triazolyacetic acid and triazollylactic acid

Dry tea leaves: at least 407–412 days (13 months).

The Meeting confirms its previous conclusions that residues of difenoconazole are ‘stable at approximately -20 °C for two years in matrices with high oil content (cotton seed, cotton seed meal, cotton seed oil (soya bean only one year)), high water content matrices (potatoes, tomatoes (banana, lettuce only one year)), high starch matrices (wheat grain) and wheat forage and straw’.

Results of supervised residue trials on crops

The Meeting received data from supervised residue trials and GAP information on guava, cranberry, cotton and tea.

Guava

Labels were provided for use of difenoconazole on guava in Egypt and the USA. On a per-hectare basis, the US label appears to be the critical GAP (3 × 128 g ai/ha, 7-day RTI, 0-day PHI versus 3 × 89 g ai/ha, 7-day RTI, 7-day PHI). No field trials matching the US GAP were available; therefore, the Meeting considered the GAP from Egypt.

The Egyptian GAP consists of three applications of 6.25 g ai/hL, using 950–1430 L/ha (equivalent to 59–89 g ai/ha), on a 7-day interval, with a PHI of 7 days.

Residues of difenoconazole in independent trials conducted in Egypt and matching the Egyptian GAP were (n = 6): 0.013, 0.018, 0.033, 0.034, 0.053 and 0.077 mg/kg.

The Meeting estimated a maximum residue level of 0.15 mg/kg, an STMR of 0.0335 mg/kg and an HR of 0.095 mg/kg (from a single sample) for guava.

Cranberry

The critical GAP for cranberry is from the USA and consists of three applications each at 128 g ai/ha, on a 7-day interval, with a PHI of 30 days with a maximum seasonal rate of 381 g ai/ha.

Residues of difenoconazole in independent trials approximating the critical GAP were (n = 4): 0.14, 0.20 (2) and 0.24 mg/kg.

The Meeting estimated a maximum residue level of 0.6 mg/kg, an STMR of 0.2 mg/kg and an HR of 0.26 mg/kg (from a single sample) for cranberry.

Cotton seed

The critical GAP for cotton is from the USA and consists of three applications each at 128 g ai/ha, on a 14-day interval, with a PHI of 45 days with a maximum seasonal rate of 381 g ai/ha.

Residues of difenoconazole in cotton seed from independent trials approximating the critical GAP were (n = 12): < 0.01 (4), 0.012, 0.018, 0.024, 0.036, 0.041, 0.047, 0.14 and 0.30 mg/kg.

The Meeting estimated a maximum residue level of 0.4 mg/kg and an STMR of 0.021 mg/kg for cotton seed.

Tea

The critical GAP for tea is from Japan and consists of two applications of a 2000-fold dilution (5 g ai/100 L) of the formulated product with a 7-day PHI; the minimum re-treatment interval is not specified.

The Meeting considered the three sets of supervised field trials (Japan, 1993, 2016 and Taiwan Province of China, 2006), field trial use patterns matched the critical GAP. Residues of difenoconazole in independent trials were (n = 8): 1.6, 1.9, 2.1, 4.4, 5.3, 7.9, 8.1 and 12 mg/kg in dried tea leaves.

The Meeting estimated a maximum residue level of 20 mg/kg and an STMR of 4.85 mg/kg for tea, green, black (black, fermented and dried).

Residues in animal feeds

Cotton gin by-products

The critical GAP for cotton is from the USA and consists of three applications each at 128 g ai/ha, on a 14-day interval, with a PHI of 45 days with a maximum seasonal rate of 381 g ai/ha.

Residues of difenoconazole in cotton gin by-products from independent trials approximating the critical GAP were (n = 3): 1.2, 0.50 and 5.6 mg/kg.

The Meeting agreed that three trials were not sufficient to make a recommendation for residues of difenoconazole in cotton gin by-products.

Fate of residues during processing

The Meeting received data showing the effect of processing cotton seed into meal, hulls and refined oil. In addition, the Meeting received data on residues in tea infusion created from treated dried tea leaves. Processing factors and residue estimates are summarized below.

Table 1 Processing factors and residue estimates for difenoconazole

Raw commodity	Residue in RAC, mg/kg			Processed commodity	Processing Factors		Residue in processed commodity, mg/kg		
	Max	STMR	HR		Individual	Best estimate	Max-P	STMR-P	HR-P
Cotton seed	0.4	0.021	--	Meal	< 0.066, 0.089	0.089	--	0.0019	--
				Hulls	< 0.066, 0.43	0.43	--	0.0090	--
				RBD oil ^a	< 0.066, < 0.088	< 0.066	--	< 0.0014	--
Tea	20	4.85	--	Tea infusion	0.0010 (3), 0.0011, 0.0012, 0.0015 (2), 0.0016, 0.0017 (2), 0.0019, < 0.017	0.0015	--	0.0072	

^a Refined Bleached Deodorized

Residues of difenoconazole did not concentrate in any of the processed commodities considered by the Meeting; therefore, the Meeting did not estimate maximum residues for those commodities. Residues for estimating animal dietary burden and for dietary risk assessment are provided in the table above.

Residues in animal commodities

Of the uses considered by the Meeting, only cotton meal, cotton hulls and cotton gin by-products are significant animal feed items. The Meeting was not able to make residue recommendations for cotton gin by-products. A comparison of the most recent burden estimates used for residue estimation (2013 JMPR) with those estimated by this meeting using current practices are shown below.

Table 2 Dietary burden (ppm) comparison from the 2013 JMPR and the current Meeting

Animal	Maximum		Mean	
	2013 JMPR	Current	2013 JMPR	Current
Beef cattle	17.9	17.9	15.3	15.0
Dairy cattle	14.9	14.7	12.4	12.1
Poultry (broiler and layer)	1.89	1.20	1.11	0.80

As the dietary burdens were essentially unchanged or are lower than those from the previous Meeting, the Meeting confirmed its previous recommendations for residues in animal commodities.

RECOMMENDATIONS

On the basis of the data obtained from supervised trials, the Meeting concluded that the residue levels listed in Annex 1 are suitable for establishing maximum residue limits and for IEDI and IESTI assessments.

Definition of the residue for compliance with the MRL and dietary risk assessment for plant commodities: *difenoconazole*.

Definition of the residue for compliance with the MRL and dietary risk assessment for animal commodities: *sum of difenoconazole and 1-[2-chloro-4-(4-chloro-phenoxy)-phenyl]-2-(1,2,4-triazol)-1-yl-ethanol (CGA205375), expressed as difenoconazole*

The residue is fat-soluble.

DIETARY RISK ASSESSMENT

Long-term dietary exposure

The ADI for difenoconazole is 0–0.01 mg/kg bw. The International Estimated Daily Intakes (IEDIs) for difenoconazole were estimated for the 17 GEMS/Food Consumption Cluster Diets using the STMR or STMR-P values estimated by the JMPR. The results are shown in Annex 3 of the 2021 Extra JMPR Report.

The IEDIs ranged from 10–80% of the maximum ADI. The Meeting concluded that long-term dietary exposure to residues of difenoconazole from uses considered by the JMPR is unlikely to present a public health concern.

Acute dietary exposure

The ARfD for difenoconazole is 0.3 mg/kg bw. The International Estimate of Short-Term Intakes (IESTIs) for difenoconazole were calculated for the food commodities and their processed commodities for which HRs/HR-Ps or STMRs/STMR-Ps were estimated by the present Meeting and for which consumption data were available. The results are shown in Annex 4 of the 2021 Extra JMPR Report.

The IESTIs varied from 0–2% of the ARfD for children and 0–2% of the ARfD for the general population. The Meeting concluded that acute dietary exposure to residues of difenoconazole from uses considered by the present Meeting is unlikely to present a public health concern.

5.9. ETHION (034)

TOXICOLOGY

Ethion was reviewed by the 2021 Extra Joint FAO/WHO Meeting on Pesticide Residues (JMPR) for additional uses. The FAO panel sought the advice of the WHO panel as to whether an estimate for acute dietary exposure needed to be performed for ethion.

The toxicity of ethion was previously evaluated by JMPR in 1968, 1969, 1970, 1972, 1975, 1982, 1983, 1986 and 1990. No new information on the toxicology of ethion had been provided to the 2021 Extra Meeting and no copies of original study reports were available. The 1990 JMPR established an ADI for ethion of 0–0.002 mg/kg bw. The ADI was based on the NOAEL of 0.2 mg/kg bw per day for decreased fetal body weight and delayed ossification in a rat developmental toxicity study, to which a safety factor of 100 was applied. No assessment for the establishment of an ARfD had been performed by a previous JMPR.

The 2021 Extra Meeting noted that JECFA had reviewed ethion in 2017 for veterinary uses (JECFA, 2018²) and established an ARfD of 0.02 mg/kg bw, based on the NOAEL of 0.15 mg/kg bw per day for erythrocyte acetylcholinesterase inhibition in a human volunteer study

The 2021 Extra Meeting noted that JECFA had identified a potential concern for exposures in pregnant women and had recommended that repeat exposures in high-percentile pregnant consumers or a suitable surrogate population should be addressed.

In the absence of any information provided on ethion and noting that the evaluation procedures used by JECFA and JMPR to establish HBGVs are very similar, the Meeting proposed that the JECFA 2017 conclusions on dietary exposures are used pending a periodic review of ethion by JMPR.

A toxicological addendum was not prepared.

RESIDUE AND ANALYTICAL ASPECTS

Ethion is an organophosphate insecticide and acaricide with non-systemic and contact action. It was first evaluated by the JMPR in 1968 and most recently for toxicology in 1990 and residues in 1994. The 1990 JMPR established an ADI of 0–0.002 mg/kg bw and the 1994 JMPR defined the residue as ethion (fat-soluble).

Currently, ethion is included in the Codex Alimentarius Commission (CAC) list of pesticides whose MRLs (CXLs) or Guideline levels (GLs) have been deleted by the CAC and for which no MRLs have been proposed. The CXLs for ethion exist only for spices based on monitoring data.

Ethion was scheduled at the Fifty-first Session of the CCPR for evaluation of additional uses by the 2020 JMPR, which was postponed to the 2021 Extra JMPR. The Meeting received information on GAP and residue trials on chili pepper.

Methods of analysis

Analysis of ethion in chili peppers was conducted by a new method using QuEChERS sample preparation and GC-FPD or GC-ECD. Recovery test results showed recoveries of 74–111%. The LOQ value was 0.05 mg/kg. The analytical method used in the chili pepper residue trials was considered sufficiently validated.

² JECFA, (2018), Joint FAO/WHO Expert Meeting on Food Additives, 85th Meeting 2017, *Residues evaluation of certain veterinary drugs*, FAO JECFA monographs 21, FAO, Rome 2018

Stability of pesticide residues in stored analytical samples

Residue analysis was performed within 24 hours after sample collection.

Results of supervised residue trials on crops***Fruiting vegetables, other than Cucurbits******Chili pepper***

The critical GAP for the use of ethion on chili pepper in India involves foliar sprays at 1.0 kg ai/ha and a 5-day PHI (the maximum number of sprays and minimum re-treatment intervals are not specified).

Six residue trials (decline) on chili pepper were conducted in India during 2014–2016, at a rate of 0.50 kg ai/ha with 2 applications at a 10-day interval and a PHI of 5 days. The residue concentrations of ethion in chili pepper were (n = 6): 0.96, 0.96, 1.1, 1.6, 1.9 and 2.6 mg/kg (highest value of 3.0 mg/kg from replicate plots). The trials did not match the application rate of the critical GAP.

Communication by the sponsor indicated that the local agricultural practice involves re-treatment intervals of 2–4 days. Residues declined with a median half-life of 4.1 days (1st order) and the modelled residue at a spray interval of 2 days and 2 applications differed by more than 25% compared to the supervised field trials. The Meeting concluded that the supervised field trials were conducted at a significantly longer re-treatment interval in combination with lower application rates and cannot be used for the estimation of a maximum residue level.

5.10. ETHIPROLE (304)

RESIDUE AND ANALYTICAL ASPECTS

Ethiprole is a non-systemic phenylpyrazole insecticide. It was first evaluated by the 2018 JMPR for toxicology and residues. The 2018 JMPR established an ADI of 0–0.005 mg/kg bw, an ARfD of 0.005 mg/kg bw and recommended the following residue definitions:

Definition of the residue for compliance with the MRL for plant commodities: Ethiprole

Definition of the residue for dietary risk assessment for plant commodities: Sum of ethiprole, 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-4-(ethylsulfinyl)-1H-pyrazole-3-carboxamide (ethiprole amide) and 5-amino-1-(2,6-dichloro-4-(trifluoromethylphenyl)-4-ethylsulfonylpyrazole-3-carbonitrile (ethiprole-sulfone), expressed as parent equivalents.

Definition of the residue for compliance with the MRL and dietary risk assessment for animal commodities: Sum of ethiprole and 5-amino-1-(2,6-dichloro-4-(trifluoromethylphenyl)-4-ethylsulfonylpyrazole-3-carbonitrile (ethiprole-sulfone), expressed as parent equivalents.

The residue is fat-soluble.

On the basis of the above residue definitions, the 2018 JMPR estimated maximum residue levels for rice, coffee and related commodities as well as commodities of animal origin.

The Forty-third Codex Alimentarius Commission (2020) approved the new work proposals including the priority list of pesticides for evaluation by the current Meeting. The priority list included ethiprole for evaluation of an additional use on soya beans. The present Meeting received new validation information on an analytical method, supervised residue trials and processing studies on soya beans.

Methods of analysis

The 2018 JMPR evaluated three LC-MS/MS methods for the determination of ethiprole, ethiprole-sulfone and ethiprole-amide in supervised trials, including Method 01128. The current Meeting received new information on the validation of Method 01128 used in the supervised trials on soya beans. The Method was validated with an LOQ of 0.002 mg/kg for the three analytes in soya beans and their processed commodities except aspirated grain fractions. In aspirated grain fractions, recoveries were only tested at a fortification level of 100 mg/kg for each analyte with the resulting procedural recoveries and relative standard deviations within the acceptable range.

Results of supervised residue trials on crops

The current Meeting received information on supervised trials using foliar spray of ethiprole conducted in support of estimating maximum residue level for soya bean (dry).

For the calculation of the sum of ethiprole, ethiprole-sulfone and ethiprole-amide, expressed as parent equivalents (total residues), the Meeting used the approach agreed at the 2018 JMPR. When the concentration of any analyte was below the LOQ, it was regarded to be at the LOQ; and when the concentrations of all three analytes were below the LOQ, the total residue is expressed as below the sum of the LOQ values (< 0.006 mg/kg).

Pulses

Soya bean (dry)

Critical GAP in Brazil for soya bean allows two foliar applications at a maximum rate of 0.2 kg ai/ha with a re-treatment interval of 7 days and PHI of 14 days.

Eight field trials were conducted on soya beans in Brazil in 2018–2019. In the trials conducted in 2019, duplicate samples were taken at each DALA point.

Ethiprole residues from the trials matching the critical GAP were in rank order (n = 8): < 0.002 (4), 0.003, 0.004, 0.007 and 0.028 mg/kg.

The Meeting estimated a maximum residue level of 0.05 mg/kg for soya bean (dry).

Corresponding total residues from the trials were in rank order (n = 8): < 0.006 (4), 0.007, 0.008, 0.012 and 0.038 mg/kg.

The Meeting estimated an STMR of 0.0065 mg/kg for soya bean (dry).

Fate of residues during processing

Processing

The effects of processing on the concentrations of ethiprole residues were evaluated by the 2018 JMPR for rice and coffee beans.

The current Meeting received information on the processing of soya beans to processed commodities, relevant to the current evaluation.

The calculated processing factors for these commodities together with calculated STMR-Ps are summarized below. Since the residue concentrations in the second trial were close to the LOQ or lower than LOQ, the Meeting selected processing factors calculated using the first trial (first value for commodity) as best estimates.

Table 1 Processing Factors and STMR-P/HR-P

Commodity	Ethiprole		Total ethiprole residues		
	Individual processing factors	Best estimate	Individual processing factors	Best estimate	STMR/STMR-P
Seed (RAC)	(0.047, 0.004 mg/kg)	-	(0.055, 0.008 mg/kg)	-	0.0065
Flour	0.26, 0.50	0.26	0.29, 0.75	0.29	0.002
Soya milk	< 0.04, 0.50	< 0.04	< 0.11, < 0.75	< 0.11	0.0007
Refined oil		< 0.04		0.24	0.002
Cold-pressed	< 0.04, 0.50	< 0.04	0.24, < 0.75	0.24	0.002
Solvent-extracted	< 0.04, 0.50	< 0.04	0.11, < 0.75	0.11	0.0007
Meal	0.21, 0.50	0.21	0.25, 0.75	0.25	0.002
Aspirated grain fractions ^a	755, 400	755	765, 323	765	4.97
Hulls	7.0, 4.5	7.0	6.9, 3.1	6.9	0.045

^a Recoveries were only validated for the three analytes at a fortification level of 100 mg/kg in aspirated grain fractions, while the concentrations of the analytes were 0.054–35.5 mg eq/kg.

Based on the maximum residue level of 0.05 mg/kg for soya bean (dry) and the processing factor of 7.0 for ethiprole in soya bean hulls, the Meeting estimated a maximum residue level of 0.4 mg/kg for soya bean hulls.

Residues in animal commodities

The 2018 JMPR calculated the maximum and mean dietary burdens using the STMR/STMR-Ps estimated for rice bran, rice grain and rice hulls on the basis of the OECD Animal Feeding Table. The current Meeting evaluated the supervised residue trials and processing studies on soya beans. Among those commodities for which data were available, soya beans, soya aspirated grain fractions, soya bean meal, soya bean hulls may be fed to livestock. The newly calculated dietary burdens showed minor increase of 0.01 ppm which corresponded to a less than 1% increase compared to the previously calculated highest dietary burdens for cattle and poultry. The current Meeting concluded that it was unnecessary to review the recommendations for animal commodities made by the 2018 JMPR.

RECOMMENDATIONS

On the basis of the data from supervised trials, the Meeting concluded that the residue levels listed in Annex 1 are suitable for establishing maximum residue limits and for IEDI and IESTI assessment.

Definition of the residue for compliance with the MRL for plant commodities: *Ethiprole*

Definition of the residue for dietary risk assessment for plant commodities: *Sum of ethiprole, 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-4-(ethylsulfinyl)-1H-pyrazole-3-carboxamide (ethiprole-amide) and 5-amino-1-(2,6-dichloro-4-(trifluoromethylphenyl)-4-ethylsulfonylpyrazole-3-carbonitrile (ethiprole-sulfone), expressed as parent equivalents.*

Definition of the residue for compliance with the MRL and dietary risk assessment for animal commodities: *Sum of ethiprole and 5-amino-1-(2,6-dichloro-4-(trifluoromethylphenyl)-4-ethylsulfonylpyrazole-3-carbonitrile (ethiprole-sulfone), expressed as parent equivalents.*

The residue is fat-soluble.

DIETARY RISK ASSESSMENT

Long-term dietary exposure

The ADI for ethiprole is 0–0.005 mg/kg bw. The International Daily Intakes (IEDIs) for ethiprole were estimated for the 17 GEMS/Food Consumption Cluster Diets using the STMR and STMR-P values estimated by JMPR. The results are shown in Annex 3 of the 2021 Extra JMPR Report.

The IEDIs ranged from 1–6% of the maximum ADI. The Meeting concluded that long-term dietary exposure to residues of ethiprole from uses considered by JMPR is unlikely to present a public health concern.

Acute dietary exposure

The ARfD for ethiprole is 0.005 mg/kg bw. The international Estimate of Short-Term Intakes (IESTIs) for ethiprole were calculated for soya bean (dry) and the processed commodities for which STMR/STMR-Ps were estimated by the present Meeting and for which consumption data were available. The results are shown in Annex 4 of the 2021 Extra JMPR report.

The IESTIs varied from 0–2% and 0–1% of the ARfD for children and general population, respectively. The Meeting concluded that acute dietary exposure to residues of ethiprole from uses considered by the present Meeting is unlikely to present a public health concern.

5.11. FENBUCONAZOLE (197)

RESIDUE AND ANALYTICAL ASPECTS

Fenbuconazole, whose IUPAC name is 4-(4-chlorophenyl)-2-phenyl- 2-(1*H*-1,2,4-triazol-1-ylmethyl) butyronitrile, is a triazole fungicide intended for agricultural and horticultural use for the control of a variety of fungal infections of crops.

Fenbuconazole was first evaluated by the 1997 JMPR and an ADI of 0–0.03 mg/kg bw was established. The residue definition is fenbuconazole, both for compliance with the MRL and dietary risk assessment for plant and animal commodities. The residue is not fat-soluble.

The 2012 JMPR established an ARfD of 0.2 mg/kg bw. The compound was also evaluated by the JMPR in 2009 and 2013 for additional uses.

At the Fifty-first Session of CCPR (2019), fenbuconazole was scheduled for evaluation of an additional use on tea by the 2020 JMPR, which was postponed to the 2021 Extra JMPR. The current Meeting received new information on use patterns, an additional analytical method, supervised field trials and a processing study on tea.

Methods of analysis

The Meeting received information on a new analytical method for fenbuconazole.

In the new method based on LC-MS/MS, fenbuconazole was extracted from dry tea leaves with acetonitrile and cleaned-up before analysis. Tea infusion (filtrate of soaking tea leaves in boiling water) was directly analysed by the same LC-MS/MS method. Recovery data support an LOQ of 0.01 mg/kg. The Meeting agreed that the method is suitable for analysing fenbuconazole in dried tea leaves and tea infusion.

Stability of pesticide residues in stored analytical samples

Stability studies of fenbuconazole in tea leaves (dry) were available. The Meeting concluded that fenbuconazole in tea leaves stored at ≤ -20 °C was stable for at least 3 months. All the field trial residue samples were analysed within this period.

Results of supervised residue trials on crops

Tea, green, black (black, fermented and dried)

The critical GAP for tea in Japan is two applications before the first pick and two applications between subsequent picks at 4.4 g ai/hL with a PHI of 7 days. Re-treatment interval between the two applications is not specified on the label. However, the Meeting noted that an interval of 7 days was common in practice in Japan.

In trials matching the GAP, residues of fenbuconazole in tea leaves from the first pick were (n = 4): 2.4, 4.0, 4.2 and 14 mg/kg.

For the second pick, following two further applications, residues were (n = 3): 3.5, 4.5 and 11 mg/kg after 7 DALA.

The Meeting considered that residues after the first pick and before the second set of sprays would not contribute significantly to the final residue taking into account growth dilution during the interval of at least 50 days between the second treatment (before the first pick) and the third treatment (after the first pick).

The Meeting agreed that the trials were sufficiently independent even though the samples were taken at the first pick and 2nd pick from the same plot and decided to use the residue data from both picks for estimation of maximum residue levels. The combined data set of residues of fenbuconazole in tea leaves (from both first and second picks) were (n = 7): 2.4, 3.5, 4.0, 4.2, 4.5, 11 and 14 mg/kg.

The Meeting estimated a maximum residue level and STMR value of 30 mg/kg and 4.2 mg/kg, respectively, for fenbuconazole in tea, green, black (black, fermented and dried).

Fate of residues during processing

The Meeting received information on the fate of fenbuconazole residues when tea infusion was produced from tea leaves. An estimated processing factor and the derived STMR-P are summarized in the table below.

Table 1 Processing Factors and STMR-P/HR-P

RAC	Processed commodity	Calculated processing factors ^a	PF (mean or best estimate)	RAC STMR (mg/kg)	STMR-P (mg/kg)
Tea leaves	Tea infusion	0.0035, 0.0037, 0.0041, 0.0044, 0.0046, 0.0046, < 0.005	0.0044	4.2	0.0018

^a Each value represents a separate trial. The factor is the ratio of the residue in processed commodity divided by the residue in the RAC.

RECOMMENDATIONS

On the basis of the data from supervised trials the Meeting concluded that the residue levels listed in Annex 1 are suitable for establishing maximum residue limits and for IEDI and IESTI assessment.

Definition of the residue for compliance with the MRL and dietary risk assessment for plant and animal commodities: fenbuconazole.

The residue is not fat-soluble.

DIETARY RISK ASSESSMENT

Long-term dietary exposure

The ADI for fenbuconazole is 0–0.03 mg/kg bw. The International Estimated Daily Intakes (IEDIs) for fenbuconazole were estimated for the 17 GEMS/Food Consumption Cluster Diets using the STMR or STMR-P values estimated by the JMPR. The results are shown in Annex 3 of the 2021 Extra JMPR Report.

The IEDIs ranged from 0–3% of the maximum ADI for fenbuconazole. The Meeting concluded that long-term dietary exposure to residues of fenbuconazole from uses considered by the JMPR is unlikely to present a public health concern.

Acute dietary exposure

The ARfD for fenbuconazole is 0.2 mg/kg bw. The International Estimate of Short-Term Intakes (IESTIs) for fenbuconazole were calculated for tea leaves and infusion for which STMRs/STMR-Ps were estimated by the present Meeting and for which consumption data were available. The results are shown in Annex 4 of the 2021 Extra JMPR Report.

The IESTIs from tea varied from 2–3% of the ARfD for children and 0–3% of the ARfD for the general population. The Meeting concluded that acute dietary exposure to residues of fenbuconazole from uses considered by the present Meeting is unlikely to present a public health concern.

5.12. FENHEXAMID (215)

RESIDUE AND ANALYTICAL ASPECTS

Fenhexamid is a hydroxylaniline protective fungicide whose mode of action is by the inhibition of germ tube elongation, mycelial growth and spore germination. Fenhexamid was evaluated for the first time by the 2005 JMPR where an ADI of 0–0.2 mg/kg bw was established and it was concluded that an ARfD was unnecessary. The 2005 JMPR also concluded that the residue definition for plant and animal commodities for compliance with the MRL and dietary risk assessment was parent fenhexamid.

Fenhexamid was scheduled by the Fifty-first Session of the CCPR for the evaluation of additional uses by the 2020 JMPR, which was postponed to the 2021 Extra JMPR. The current Meeting received new information on analytical methods, storage stability and on the use patterns for fenhexamid in pear, ginseng, asparagus, spring onion and bulb vegetables supported by supervised field trials.

Methods of analysis

The Meeting received descriptions and validation data for the analytical methods (specific method codes were not available) in plant matrices. Plant matrices (high water content) are extracted with acetone and cleaned up by solid phase extraction. The residues are detected with LC-MS and generally achieved an LOQ of 0.02 mg/kg for pear, asparagus and onions. The recoveries were in the range of 77–102% with repeatability below 14%.

For ginseng, the Meeting noted that the performance of the analytical method was poor (53–84%) and the mean procedural recoveries were not acceptable (58–58.3%).

Stability of pesticide residues in stored analytical samples

Information provided to the 2005 JMPR demonstrated that fenhexamid residues were stable for 12 months in high-water commodities (peaches, cherries, plums, tomatoes). The current Meeting received additional residue stability data for pear, ginseng, asparagus and green onions from samples stored concurrently with residue trial samples. Stability was demonstrated for at least 7.7, 7.2 and 17.7 months for pear, asparagus and green onion, respectively, covering the storage period of samples in the residue trials.

In ginseng, 60% of the residue was found after 4.4 months, however the procedural recoveries were not acceptable (mean recovery of the fresh fortification samples was 62.5%) thus the meeting could not conclude on an acceptable storage stability period for ginseng. **Error! Reference source not found.**

Results of supervised residue trials on crops

The Meeting received supervised residue trial data for foliar applications of fenhexamid on pears, ginseng, asparagus, spring and bulb onion.

Pear

Fenhexamid is registered in the USA for use as a post-harvest dip or drench application on pears at a maximum rate of 0.34 kg ai in 378.5 L of water to 90.7 tn. of fruit (equivalent to 0.09 kg ai/hL or 3.7 g ai/1 tn. fruit). The GAP does not allow a combination of a post-harvest and pre-harvest treatment.

In five trials matching the GAP for post-harvest dip, residues of fenhexamid were (n = 5): 0.9, 1.8, 2.6, 2.9 and 2.9 mg/kg. In five additional trials matching the GAP for post-harvest drench, residues of fenhexamid were (n = 5): 1.1, 1.3, 1.7, 2.05 and 2.2 mg/kg.

Two further trials were carried out with one post-harvest spray application of 0.34 kg ai in 378.5 L of water in the packing line, however only one of the trials matched the GAP for post-harvest use with residues of fenhexamid at 3.25 mg/kg.

The combined data were (n = 11): 0.9, 1.1, 1.3, 1.7, 1.8, 2.05, 2.2, 2.6, 2.9, 2.9 and 3.25 mg/kg.

The Meeting estimated a maximum residue level of 6 Po mg/kg (based on the mean + 4×SD) and an STMR of 2.05 mg/kg for pears based on the combined dataset.

Ginseng

The critical GAP for fenhexamid on ginseng in the USA is for 4 applications up to BBCH 99 at 840 g ai/ha (interval 7–14 days) and 0 days PHI. The meeting received five trials according to the critical GAP from which 4 were considered independent. The Meeting noted that the performance of the analytical method was poor (53–84%) and the mean procedural recoveries were not acceptable (58–58.3%), thus the meeting was unable to recommend a maximum residue level for ginseng roots.

Asparagus

The critical GAP for fenhexamid on asparagus in the USA is for 4 applications to the ferns up to BBCH 95 at 840 g ai/ha (interval 7–14 days) and 180 days PHI. The meeting received 2 independent, overdosed trials (at 4190–4260 g ai/ha). The residues in spears were (n = 2): < 0.02 and < 0.02 mg/kg. Since in both trials residues were below the LOQ and the applications were made to ferns prior to senescence, about 6 months before spear emergence, the limited number of trials was considered acceptable.

The Meeting estimated a maximum residue level of 0.02 mg/kg and an STMR of 0.02 mg/kg for asparagus.

Subgroup of Bulb Onions

The critical GAP for fenhexamid on bulb onion (Onion, bulb, subgroup 3-07A) in the USA is for 4 applications up to BBCH 48 at 840 g ai/ha (7-day intervals up to the day of harvest) and 0 days PHI. The meeting received seven independent trials matching the critical GAP. The residues in bulb onions were (n = 7): < 0.02(2), 0.034, 0.055, 0.069, 0.33 and 1.4 mg/kg.

The Meeting estimated a maximum residue level of 3 mg/kg and an STMR of 0.055 mg/kg for the Subgroup of Bulb Onions.

Subgroup of Green Onions

The critical GAP for fenhexamid on spring onion (Onion, green, subgroup 3-07B) in the USA is for 4 applications up to BBCH 40 at 840 g ai/ha (7-day intervals up to the day of harvest) and a 0-day PHI. The meeting received 3 independent trials according to the critical GAP. The residues in green onions were (n = 3): 2.55, 4.35 and 11.7 mg/kg.

The Meeting concluded that the data, were not sufficient to give a recommendation for Green onions as a minor crop.

RECOMMENDATIONS

On the basis of the data from supervised trials, the Meeting concluded that the residue levels listed in Annex 1 are suitable for establishing maximum residue limits and for IEDI assessment.

The residue definition for compliance with the MRL and dietary risk assessment for plant and animal commodities is *fenhexamid*.

The residue is fat-soluble

DIETARY RISK ASSESSMENT***Long-term dietary exposure***

The ADI for fenhexamid is 0–0.2 mg/kg bw. The International Estimated Daily Intakes (IEDIs) for fenhexamid were estimated for the 17 GEMS/Food Consumption Cluster Diets using the STMR or STMR-P values estimated by the JMPR. The results are shown in Annex 3 of the 2021 Extra JMPR Report.

The IEDIs ranged from 0–5% of the maximum ADI. The Meeting concluded that long-term dietary exposure to residues of fenhexamid from uses considered by the JMPR is unlikely to present a public health concern.

Acute dietary exposure

The 2005 JMPR decided that an ARfD for fenhexamid was unnecessary. The Meeting therefore concluded that the acute dietary exposure to residues of fenhexamid from the uses considered is unlikely to present a public health concern.

5.13. FENPICOXAMID (305)

TOXICOLOGY

Fenpicoxamid is the ISO-approved common name for [2-[[[(3*R*,7*R*,8*R*,9*S*)-7-benzyl-9-methyl-8-(2-methylpropanoyloxy)-2,6-dioxo-1,5-dioxonan-3-yl]carbamoyl]-4-methoxypyridin-3-yl]oxymethyl 2-methylpropanoate, with the CAS number 517875-34-2. It was evaluated by the JMPR in 2018, when an acceptable daily intake (ADI) of 0–0.05 mg/kg bw was established and it was concluded that an acute reference dose (ARfD) was unnecessary (Annex 5, reference 147).

No new information on the toxicology of fenpicoxamid was provided to the 2021 Extra Meeting.

Fenpicoxamid was reviewed by the 2021 Extra JMPR for additional uses. The FAO panel sought the advice of the WHO panel regarding the toxicity of two metabolites, X12326349 and X696872.

Toxicological data on metabolites

The metabolites X12326349 and X696872 were evaluated following the scheme for metabolites developed by JMPR (WHO, 2015³).

No toxicity data were provided on the metabolites, but predictive modelling of oral absorption was made available by the sponsor (Damoradski, 2021⁴).

The oral absorption of fenpicoxamid is 5–20% (JMPR, 2018). The modelling of the oral absorption of X12326349 and X696872 indicated that, at low doses relevant to dietary risk assessment, their oral absorption would be of the same order as that of fenpicoxamid (*ibid*).

Urinary levels of X12326349, following the oral administration of fenpicoxamid to rats, were 2–7% of the administered dose, equating to approximately 30–40% of the absorbed dose (JMPR, 2018).

Urinary levels of X696872 following the oral administration of fenpicoxamid to rats were 1.3–7% of the administered dose, equating to approximately 30% of the absorbed dose (JMPR, 2018).

Toxicological evaluation

As greater than 10% of the absorbed dose of fenpicoxamid was metabolized to each of X12326349 and X696872, the toxicity of X12326349 and X696872 can be considered adequately tested by the in-situ production of these metabolites during toxicity studies performed with fenpicoxamid alone. The ADI of fenpicoxamid also applies to the metabolites.

A toxicological addendum was not prepared.

³ WHO (2015). Pesticide residues in food. Guidance document for WHO monographers and reviewers. WHO Core Assessment Group on Pesticide Residues. WHO, Geneva, 2005. January 2015

⁴ Damoradski J. (2021) Fenpicoxamid metabolites Fa rat 2 mg–60 mg. Unpublished report. Provided to JMPR by Corteva Agriscience, Indianapolis USA.







RESIDUE AND ANALYTICAL ASPECTS



Fenpicoxamid is a picolinamide fungicide. The mode of action is by the inhibition of mitochondrial complex III to disrupt spore germination and germ tube elongation. Fenpicoxamid was evaluated for the first time by the 2018 JMPR where an ADI of 0–0.05 mg/kg bw was established and it was concluded that an ARfD was unnecessary. The residue definition for plant commodities for compliance with the MRL and dietary risk assessment was parent fenpicoxamid. The 2018 JMPR did not recommend residue definitions for animal commodities (compliance with the MRL and dietary risk assessment) but concluded that if future uses of fenpicoxamid resulted in an increase in the dietary exposure to the animal metabolite X12326349 and the hydrolysis products X12314005, X12019520, X12335723 and X12264475, reconsideration of the residue definitions may become necessary.

It was scheduled at the Fifty-first Session of the CCPR for the evaluation of additional uses by the 2020 JMPR, which was postponed to the 2021 Extra JMPR. The current Meeting received new information on use patterns for fenpicoxamid in wheat, similar grains and pseudo cereals without husks supported by additional analytical methods and recovery data, storage stability studies, supervised field trials, feeding studies and studies simulating typical processing conditions.

The abbreviations and structures used for the major metabolites discussed in this appraisal are shown in Table 1.

Table 1 List of identified metabolites for fenpicoxamid discussed by the current Meeting

Name	Chemical name	Structure
fenpicoxamid	(3 <i>S</i> ,6 <i>S</i> ,7 <i>R</i> ,8 <i>R</i>)-8-benzyl-3-{3-[(isobutyryloxy) methoxy]-4-methoxypyridine-2-carboxamido}-6-methyl-4,9-dioxo-1,5-dioxonan-7-yl isobutyrate	
X12326349	2-benzyl-2,5-dideoxy-4-O-[<i>N</i> -(3-hydroxy-O-4-methoxypyridine-2-carbonyl)-L-seryl]-L-arabinonic acid	
X12019520	(5 <i>S</i>)-3-benzyl-5-methylfuran-2(5 <i>H</i>)-one	
X12314005	(2 <i>S</i> ,3 <i>R</i> ,4 <i>R</i>)-4-benzyl-2-methyl-5-oxotetrahydrofuran-3-yl 2-methylpropanoate	
X12335723	<i>N</i> -(4-methoxy-3-[(2-methylpropanoyl)oxy]methoxy)pyridine-2-carbonyl)-L-serine	
X12264475	<i>N</i> -(3-hydroxy-4-methoxypyridine-2-carbonyl)-DL-serine	

Name	Chemical name	Structure
X642188	(3S,6S,7R,8R)-8-benzyl-3-[[[(3-hydroxy-4-methoxypyridin-2-yl)carbonyl]amino]-6-methyl-4,9-dioxo-1,5-dioxonan-7-yl 2-methylpropanoate	
X696872	N-[(3S,7R,8R,9S)-7-benzyl-8-hydroxy-9-methyl-2,6-dioxo-1,5-dioxonan-3-yl]-3-hydroxy-4-methoxypyridine-2-carboxamide	

Methods of analysis

A LC-MS/MS based analytical method (120615) for the determination of fenpicoxamid and its plant metabolite X642188 in wheat grain, straw and whole plant with an LOQ of 0.01 mg/kg for each analyte was evaluated by the 2018 JMPR. The current Meeting received new procedural recovery data for a similar method involving the same principles acetonitrile/water/phosphoric acid extraction and LC-MS/MS determination of fenpicoxamid and its metabolites (X642188, X12335723, X12019520, X12314005, X12264475) in wheat grain, wheat bran, refined flour and wholemeal bread supporting the LOQ of 0.01 mg/kg.

A similar method was evaluated by the 2018 JMPR for the analysis of fenpicoxamid and its animal metabolites X642188 and X12326349 in animal commodities with an LOQ of 0.01 mg/kg for each analyte.

The Meeting concluded that both methods were demonstrated to have adequate performance for recovery of fenpicoxamid and its metabolites, with an LOQ of 0.01 mg/kg (for each compound).

Stability of pesticide residues in stored analytical samples

The storage stability of fenpicoxamid and its metabolite X642188 was investigated by the 2018 JMPR. Additional storage stability studies for dry and animal commodities were provided to the current meeting. The Meeting concluded that fenpicoxamid and X642188 are stable for at least 24 months in wheat grain, straw, forage, bran, flour, germ and bread when stored frozen at -18 °C. All samples of animal commodities were analysed within 38 days of sampling. This period is covered by the available storage stability study on animal commodities.

Definition of the residue

The 2018 JMPR established the current residue definition for fenpicoxamid in plant commodities.

For plant commodities, the 2018 JMPR concluded that fenpicoxamid is a suitable marker for enforcement of maximum residue levels and for dietary risk assessment for plant commodities. However, it was also recognised that in processed commodities, the degradates X12314005, X12016520, X12335723 and X12264475 may become relevant for commodities subject to high-temperature conditions. The new uses evaluated by the current Meeting relate to cereal grains, which are mostly subject to thermal treatment before consumption. In four processing trials involving field treatment according to the critical GAP, all of these degradates were analysed in raw and processed wheat commodities. Quantifiable residues above the LOQ of 0.01 mg/kg were only found for X12264475 in dried gluten. In absence of specific toxicological data, X12264475 was characterized as a Cramer Class III compound by the 2018 Meeting. Based on the very low X12264475 residue levels in dried gluten (up to 0.014 mg/kg) and the negligible relevance of the commodity in the daily diet, the current Meeting concluded that the dietary exposure will not be significant or reach the TTC for a Cramer Class III compound.

The Meeting confirmed its previous residue definition for fenpicoxamid in plant matrices. Should the Meeting evaluate future uses involving commodities subject to high-temperature conditions, these compounds may need to be re-considered for assessing dietary risk.

For animal commodities, the 2018 JMPR did not recommend a residue definition. The compounds X12326349, 13495S-3S, X12264475 were found as major metabolites in **ruminants** while the co-eluting metabolites X119634422/MW208 and the metabolites X696872, X12264475 and X129300 represented the predominant residue in **poultry**. Since total radioactive residues in both species were very low, the 2018 Meeting concluded that dietary burdens up to 20 ppm for ruminants and 10 ppm for laying hens will not lead to residues above 0.01 mg/kg in animal commodities.

The current Meeting received a new farm animal feeding study on dairy cows with dose rates of 4.5, 13.4 and 67.3 ppm. While residues of parent fenpicoxamid and X642188 remained below the LOQ in all matrices and dose groups, X12326349 was detected (LOD 0.003 mg/kg) in the 4.5 ppm dose group and quantified at concentrations of 0.011 mg/kg and 0.047 mg/kg in tissues in the 13.4 ppm and 67.3 ppm dose groups, respectively. In the highest dose group quantifiable residues were also found in kidney (0.061 mg/kg) and perirenal fat (0.012 mg/kg). In view of the maximum dietary burden for beef cattle of 13.6 ppm arising from the new use on cereals, residues near or above 0.01 mg/kg have to be expected in ruminant liver and kidney. The current Meeting reviewed the available data and recognized that urinary levels of X12326349 represented 30–40% of the absorbed dose. Accordingly, it can be considered as having been addressed by studies with the parent and as such HBGVs for fenpicoxamid also apply to X12326349.

Residues of X12326349 were primarily found in liver and kidney. Only in the highest dose group were residues quantified in perirenal fat (0.012 mg/kg), while residues in muscle remained below the LOQ (0.003 mg/kg). However, the data does not indicate a strong tendency to partition into fatty compartments.

Since X12326349 was the only residue found in livestock animal feed studies and an analytical method capable of measuring the analyte at LOQs of 0.01 mg/kg is available, the current Meeting concluded that it represents a suitable residue definition both for setting of maximum residue levels and for the estimation of the dietary exposure.

The definition of the residue for compliance with the MRL and dietary risk assessment for milk and tissues of mammals (other than marine mammals): *2-benzyl-2,5-dideoxy-4-O-[[3-hydroxy-4-methoxypyridin-2-yl]carbonyl]-L-seryl]-L-arabinoic acid (X12326349), expressed as fenpicoxamid.*

The residue is not fat-soluble.

The current Meeting confirms its previous decision not to recommend a residue definition for poultry commodities (muscle, fat, offals and eggs). If future uses of fenpicoxamid result in an increase in the dietary exposure to the animal metabolite X12326349 and the hydrolysis products X12314005, X12019520, X12335723 and X12264475 in poultry, reconsideration of the residue definitions may become necessary.

Results of supervised residue trials on crops

The Meeting received new GAP information and supervised residue trial data for the foliar application of fenpicoxamid on wheat and related cereal grains.

Rye, triticale and wheat

The critical GAP for fenpicoxamid on rye, triticale and wheat (including durum and spelt) is similar in several European countries. As a representative, the GAP in Belgium is for 2 foliar applications up to BBCH 69 (end of flowering) at 100 g ai/ha with a 14-day re-treatment interval. The PHI is limited by the growth stage at the last application. The meeting received 36 supervised field trials matching the critical GAP, from which 18 were considered independent.

In wheat grain, residues of fenpicoxamid were (n = 18): 0.010 (2), 0.011, 0.012 (3), 0.014, 0.016, 0.020, 0.023, 0.024, 0.030, 0.034, 0.036, 0.04 (2), 0.069 and 0.095 mg/kg.

The Meeting estimated a maximum residue level of 0.15 mg/kg and an STMR of 0.0215 mg/kg for wheat. Noting that the GAP also applied to rye and triticale the Meeting decided to extrapolate the maximum residue level to rye and triticale.

Residues in animal feeds

Forage of cereal straw

Based on a New Zealand GAP for wheat of 2 foliar applications up to BBCH 69 (end of flowering) at 100 g ai/ha (interval 14 days) with a PHI of 28 days for forage. The Meeting received eight independent trials that reported residues of fenpicoxamid in wheat forage, residues were (n = 8): 0.31, 0.32, 0.66, 0.77, 0.83, 1.3, 1.6, 2.2 mg/kg.

The Meeting estimated a median and highest residue of 0.80 mg/kg and 2.2 mg/kg, respectively, for wheat forage.

Fodder and straw

The critical GAP for fenpicoxamid on rye, triticale and wheat (including durum and spelt) is similar in several European countries. As a representative, the GAP in Belgium is for 2 foliar applications up to BBCH 69 (end of flowering) at 100 g ai/ha (interval 14 days). This GAP only allows for feeding of wheat straw and fodder. The PHI is covered by the growth stage at the last application. The meeting received 36 supervised field trials according to the critical GAP, from which 18 were considered as independent.

In wheat straw residues of fenpicoxamid were (n = 18): 0.84, 1.2, 1.8, 2.4, 2.5, 2.7, 2.7, 3.5, 4.7, 4.7, 5.3, 6.3, 6.3, 6.8, 8.7, 9.9, 10.6 and 12 mg/kg.

The Meeting estimated a maximum residue level of 30 mg/kg (dry weight basis) and a median residue of 4.7 mg/kg (as received) and a highest residue of 12 mg/kg (as received) for wheat straw and fodder, dry.

Fate of residues during processing

The Meeting received new processing studies on wheat. Processed commodities were derived using simulated commercial practices. The resulting processing factors and STMR-P estimates for fenpicoxamid are summarized in Table 2 below.

Table 2 Estimation of 1processing factors for wheat processed fractions from field-incurred residues from foliar treatment with fenpicoxamid

Raw commodity	Processed commodity	fenpicoxamid			
		Individual processing factors	Median or best estimate processing factor	STMR or STMR-P (mg/kg)	Maximum residue level (mg/kg)
Wheat grain (STMR: 0.0215 mg/kg)	Bran	0.5, 0.56, 0.57, 0.61, 0.78, <u>0.86</u> , <u>0.88</u> , 0.91, 0.93, 1.24, 1.25, 1.7 4	0.87	0.019	-
	Refined flour (type 550)	0.15, 0.2, 0.22, 0.25, 0.26, 0.28, 0.29, 0.31, 0.34, 0.35, 0.36, 0.44	0.29	0.0062	-
	White bread	0.07, 0.09, 0.11, < 0.13, 0.13, 0.13, 0.16, < 0.18, < 0.2, < 0.24, < 0.29, < 0.29	0.15	0.0032	-

Raw commodity	Processed commodity	fenpicoxamid			
		Individual processing factors	Median or best estimate processing factor	STMR or STMR-P (mg/kg)	Maximum residue level (mg/kg)
	Wholemeal flour	0.26, 0.27, 0.36, 0.38, 0.38, 0.4, 0.41, 0.44, 0.5, 0.57, 0.6, 0.71	0.41	0.0088	-
	Wholemeal bread	0.13, 0.14, < 0.16, 0.18, 0.18, 0.2, 0.2, < 0.2, 0.23, 0.26, < 0.29, < 0.29	0.20	0.0043	-
	Wheat germ	0.08, 0.11, < 0.13, 0.15, < 0.16, 0.16, < 0.18, < 0.2, < 0.24, 0.26, < 0.29, < 0.29	0.17	0.0036	-
	Wheat starch	< 0.03, < 0.03, < 0.04, 0.04, < 0.05, <u>< 0.13</u> , <u>< 0.16</u> , < 0.18, < 0.2, < 0.24, < 0.29, < 0.29	0.15	0.0032	-
	Wheat gluten	0.15, < 0.16, 0.17, 0.18, < 0.2, < 0.24, < 0.29, < 0.29, 0.29, 0.31, 0.32, 0.52	0.27	0.0058	-

Residues in animal commodities

The 2018 JMPR did not recommend residue definitions for animal commodities as they were not necessary for the uses considered by that meeting. Commodities considered by the current meeting can be used as feed items, thus dietary burdens were calculated for beef cattle, dairy cattle, broilers and laying poultry based on the feed items (cereal forage and grain) evaluated by the current meeting. The dietary burdens, estimated using the most recent version of the OECD livestock dietary burden calculator, are presented in Annex 6 of the 2021 Extra JMPR Report and summarised in Table 3 below.

Table 3 Estimated maximum and mean dietary burdens of farm animals

	Livestock dietary burden for the estimation of maximum residue levels and the dietary exposure, ppm of dry matter diet							
	US-Canada		EU		Australia		Japan	
	max.	mean	max.	Mean	max.	mean	max.	mean
Beef cattle	1.4	0.54	2.7	1.1	12.7 ^a	4.9 ^b	0.009	0.009
Dairy cattle	2.2	0.86	2.7	1.1	6.3 ^c	2.4 ^d	0	0
Poultry - broiler	0.025	0.025	0.018	0.018	0.013	0.013	0	0
Poultry - layer	0.025	0.025	1.4 ^e	0.54 ^f	0.009	0.009	0	0

^a Suitable for estimation of maximum residue levels in meat.

^b Suitable for estimation of median residue levels in meat

^c Suitable for estimation of maximum residue levels in milk

^d Suitable for estimation of median residue levels in milk

^e Suitable for estimation of maximum residue levels in poultry meat and eggs

^f Suitable for estimation of median residue levels in poultry meat and eggs

The fenpicoxamid dietary burden reached a maximum level of 12.7 ppm diet in beef cattle, 6.3 ppm diet in dairy cattle and 1.4 mg/kg diet in poultry. The mean dietary burdens were 4.9 mg/kg in beef cattle, 2.4 mg/kg diet in dairy cattle and 0.54 mg/kg in poultry.

Residue in animal commodities**Farm animal feeding studies****Cattle**

A new lactating dairy cow feeding study was provided with dosing at three levels equivalent to 4.5, 13.44 and 67.31 ppm in the diet. Residues of fenpicoxamid and X642188 were < 0.01 mg/kg in all matrices. Residues of X12326349 were < 0.01 mg/kg in milk, subcutaneous/mesenteric fat and muscle, but were quantifiable in liver (0.011–0.047 mg/kg), kidney (0.061 mg/kg) and perirenal fat (0.012 mg/kg).

Table 4 Maximum residue level, STMR and HR in cattle commodities

	Feed level (ppm DM) for milk residues	Residues (mg/kg) in milk ^a	Feed level (ppm DM) for tissue residues	Residues ^a (mg/kg)			
				Muscle	Liver	Kidney	Fat
Maximum residue level and HR in beef or dairy cattle							
Feeding study	4.5	< 0.013	4.5	< 0.013	< 0.013	< 0.013	< 0.013
	13.4	< 0.013	13.4	< 0.013	0.015	< 0.013	< 0.013
MRL estimate ^b	6.3	0.015* ^c	12.7	0.015* ^c	0.02	0.015 ^c	0.015 ^c
STMR in beef or dairy cattle							
Feeding study	4.5	< 0.013	4.5	< 0.013	< 0.013	< 0.013	< 0.013
	13.4	< 0.013	13.4	< 0.013	0.015	< 0.013	< 0.013
Dietary burden and STMR estimate ^c	2.4	0 ^b	4.9	0 ^b	0.013	0.013	0.013

^a Residues of X12326349 expressed as fenpicoxamid calculated using a molecular weight conversion factor of 1.33. X12326349 in fenpicoxamid equivalents = X12326349 mg/kg × (614.66/462.2)

^b Residues are below LOQ at an overdosed rate and therefore residues at the dietary burden are essentially zero

^c MRL estimate based on limit of quantification (i.e. calculated value of < 0.013 mg/kg rounded up to the next MRL class of 0.015 mg/kg)

Since the residue definition is X12326349 expressed as fenpicoxamid, a conversion factor of 1.33 was used based on the molecular weight (X12326349 in fenpicoxamid equivalents = X12326349 mg/kg × (614.66/462.2)).

The Meeting estimated maximum residue levels of 0.02 mg/kg for edible offal, mammalian, 0.015 mg/kg for fat and 0.015(*) mg/kg for milk and meat.

The Meeting estimated STMRs of 0.013 mg/kg for liver, kidney and fat and STMRs of 0 mg/kg for muscle and milk.

Poultry

For poultry, in the 2018 JMPR a metabolism study in laying hens was conducted at a dose level of *ca.* 10 ppm DM which covers the currently calculated dietary burden (1.4 ppm). In this study, TRR were very low in eggs (0.001–0.007 mg/kg) and low in tissues (0.005–0.153 mg/kg). Only liver samples were taken for identification and fenpicoxamid was not detected (< 0.003 mg/kg). Other metabolites were determined to be below the limit of quantification (< 0.01 mg/kg). Therefore, fenpicoxamid and metabolites are expected to be below the LOQ (< 0.01 mg/kg) in all poultry products when fenpicoxamid is applied according to the intended GAP.

The current Meeting confirms its previous decision not to recommend a residue definition for poultry commodities (muscle, fat, offal and eggs). If future uses of fenpicoxamid result in an increase of the dietary exposure of the animal metabolite X12326349 and the products X12314005, X12019520, X12335723 and X12264475 in poultry, reconsideration of the residue definitions may become necessary.

RECOMMENDATIONS

On the basis of the data from supervised trials, the Meeting concluded that the residue levels listed in Annex 1 are suitable for establishing maximum residue limits and for IEDI assessment.

The residue definition for compliance with the MRL and dietary risk assessment for plant commodities is *fenpicoxamid*.

The residue definition for compliance with the MRL and dietary risk assessment for milk and tissues of mammals (other than marine mammals): is *2-benzyl-2,5-dideoxy-4-O-[N- [(3-hydroxy-4-methoxypyridin- 2-yl)carbonyl]-L-seryl]-L-arabinoic acid (X12326349), expressed as fenpicoxamid*.

The residue is not fat-soluble.

DIETARY RISK ASSESSMENT

Long-term dietary exposure

The ADI for fenpicoxamid is 0–0.05 mg/kg bw. The International Estimated Daily Intakes (IEDIs) for fenpicoxamid were estimated for the 17 GEMS/Food Consumption Cluster Diets using the STMR or STMR-P values estimated by the JMPR. The results are shown in Annex 3 of the 2021 Extra JMPR Report.

The IEDIs were 0% of the maximum ADI. The Meeting concluded that long-term dietary exposure to residues of fenpicoxamid from uses considered by the JMPR is unlikely to present a public health concern.

Acute dietary exposure

The 2018 JMPR decided that an ARfD for fenpicoxamid was unnecessary. The Meeting therefore concluded that the acute dietary exposure to residues of fenpicoxamid from the uses considered is unlikely to present a public health concern.

5.14. FLUOPYRAM (243)

RESIDUE AND ANALYTICAL ASPECTS

Fluopyram, a pyridylethylamide broad spectrum fungicide, was first evaluated by the JMPR for toxicology and residues in 2010 and subsequently in 2012, 2014, 2015 and 2017 for residues. The 2010 JMPR established an ADI of 0–0.01 mg/kg bw and an ARfD of 0.5 mg/kg bw.

The 2010 JMPR recommended the following residue definitions:

Definition of the residue for compliance with the MRL and dietary risk assessment for plant commodities: *Fluopyram*

Definition of the residue for compliance with the MRL for animal commodities: *Sum of fluopyram and 2-(trifluoromethyl)benzamide, expressed as fluopyram.*

Definition of the residue for dietary risk assessment for animal commodities: *Sum of fluopyram, 2-(trifluoromethyl)benzamide and the combined residues of N-{(E)-2-[3-chloro-5-(trifluoromethyl)pyridin-2-yl]ethenyl}-2-trifluoromethyl)benzamide and N-{(Z)-2-[3-chloro-5-(trifluoromethyl)pyridin-2-yl]ethenyl}-2-trifluoromethyl)benzamide, all expressed as fluopyram.*

The residue is not fat-soluble.

The Forty-third Codex Alimentarius Commission (2020) approved the new work proposals including the priority list of pesticides for evaluation by the current Meeting. The priority list included fluopyram for the evaluation of an additional use on coffee plants.

The current Meeting received new information on GAP, validation of analytical methods, supervised residue trials on coffee and processing studies on coffee beans.

Methods of analysis

The current Meeting received new information on the validation of Method 00984 and Method 00984-M001 used in supervised trials on coffee. These LC-MS/MS methods were evaluated by the JMPR in 2010 and later for determining fluopyram and its three metabolites in many other matrices.

The methods were validated with LOQs of 0.01 mg/kg for fluopyram and (2-trifluoromethyl)benzamide in green coffee beans, roasted coffee beans and instant coffee. The mean recoveries and RSDs were within the acceptable range in these commodities.

Results of supervised residue trials on crops

The current Meeting received information on supervised trials on coffee plants using drench spray application of fluopyram.

Seed for beverages and sweets

Coffee beans

Critical GAP in Brazil for coffee plants allows one strip or drench application to soil at a maximum rate of 0.5 kg ai/ha per season and a PHI of 90 days.

GAP in Guatemala and Nicaragua for coffee plants allows one application in nursery plants (0.075 g ai/plant) and one application to the ground in mature plantations at the same rate as in Brazil with a PHI of 120 days. However, since it normally takes more than three years for coffee plants to become harvestable, the Meeting considered that the application in nursery plants would not lead to residues in harvested coffee beans and that the GAP in Brazil with a shorter PHI was more critical than that in Guatemala and Nicaragua. In the trials conducted in Central America, samples were collected at DAT that were shorter than the PHI specified in the GAP in Guatemala and Nicaragua.

A total of ten field trials were conducted on coffee plants in coffee growing regions in Brazil (5) in 2015 and in Colombia (2), Costa Rica (1), Guatemala (1) and Mexico (1) in 2016. Duplicate

samples were taken in these trials. The harvested coffee cherries were processed using the dry method in the trials in Brazil and using the wet method in the trials in Central America.

Fluopyram residues from trials in Brazil according to the GAP in Brazil were in rank order (n = 5): < 0.01 (4) and 0.012 mg/kg.

Fluopyram residues from trials in Central America according to the GAP in Brazil were in rank order (n = 5): < 0.01 (5) mg/kg.

While the green coffee bean samples were prepared by two different methods (dry method and wet method) in Brazil and Central America, the Meeting concluded that since the residue levels were low, the residue populations could be considered together.

After combining the data, fluopyram residues were in rank order (n = 10): < 0.01 (9) and 0.012 mg/kg. The Meeting estimated a maximum residue level of 0.015 mg/kg for coffee beans. The Meeting estimated an STMR of 0.01 mg/kg.

Fate of residues during processing

The current Meeting received information on the processing of green coffee beans to roasted coffee beans and freeze-dried instant coffee. For obtaining the raw material green beans, fluopyram was applied at an exaggerated rate (5N of the maximum rate specified in GAP in Brazil) in the field. Fluopyram concentrations in all the samples of green beans (RAC) and roasted coffee and instant coffee were < 0.01 mg/kg. Therefore, processing factors could not be calculated.

As fluopyram was detected slightly above the LOQ in one sample of green beans from the trial conducted in accordance with GAP in Brazil and was below the LOQ in all the samples in the processing studies using an exaggerated 5N rate in the field phase, the Meeting agreed to use 0.01 mg/kg as the STMR-P for roasted beans and instant coffee for dietary exposure assessment.

Residues in animal commodities

As coffee beans or their by-products/processed commodities are not included in the OECD Animal Feeding Table, the current Meeting concluded that it was unnecessary to re-calculate animal dietary burdens or review the recommendations for animal commodities made by the 2017 Jmpr.

RECOMMENDATIONS

On the basis of the data from supervised trials, the Meeting concluded that the residue levels listed in Annex 1 are suitable for establishing maximum residue limits and for IEDI and IESTI assessment.

Definition of the residue for compliance with the MRL and dietary risk assessment for plant commodities: *Fluopyram*

Definition of the residue for compliance with the MRL for animal commodities: *Sum of fluopyram and 2-(trifluoromethyl)benzamide, expressed as fluopyram.*

Definition of the residue for dietary risk assessment for animal commodities: *Sum of fluopyram, 2-(trifluoromethyl)benzamide and the combined residues of N-[(E)-2-[3-chloro-5-(trifluoromethyl)pyridin-2-yl]ethenyl]-2-trifluoromethylbenzamide and N-[(Z)-2-[3-chloro-5-(trifluoromethyl)pyridin-2-yl]ethenyl]-2-trifluoromethylbenzamide, all expressed as fluopyram.*

The residue is not fat-soluble.

DIETARY RISK ASSESSMENT***Long-term dietary exposure***

The ADI for fluopyram is 0–0.01 mg/kg bw. The International Daily Intakes (IEDIs) for fluopyram were estimated for the 17 GEMS/Food Consumption Cluster Diets using the STMR and STMR-P values estimated by JMPR. The results are shown in Annex 3 of the 2021 Extra JMPR Report.

The IEDIs ranged from 10–80% of the maximum ADI. The Meeting concluded that long-term dietary exposure to residues of fluopyram from uses considered by the JMPR is unlikely to present a public health concern.

Acute dietary exposure

The ARfD for fluopyram is 0.5 mg/kg bw. The international Estimate of Short-Term Intakes (IESTIs) for fluopyram were calculated for coffee beans and the processed commodities for which STMR/STMR-Ps were estimated by the present Meeting and for which consumption data were available. The results are shown in Annex 4 of the 2021 Extra JMPR report.

The IESTIs were 0% of the ARfD for children and the general population. The Meeting concluded that acute dietary exposure to residues of fluopyram from uses considered by the present Meeting is unlikely to present a public health concern.

5.15. **IMAZALIL (110)**

RESIDUE AND ANALYTICAL ASPECTS

Imazalil, an imidazole fungicide with protective, curative and anti-sporulant activity, was first evaluated by the JMPR in 1977 and was re-evaluated under the periodic review programme for toxicology and residues by the 2018 JMPR. FAO Specifications for imazalil (technical material and related formulations) were published in 2001.

The 2018 JMPR confirmed the ADI of 0–0.03 mg/kg bw (established by the 1991 JMPR) and the ARfD of 0.05 mg/kg bw (established by the 2005 JMPR).

Residue definitions established by the 2018 JMPR are:

Definition of the residue for compliance with the MRL for plant and animal commodities:
imazalil

Definition of the residue for dietary risk assessment for plant commodities: *free and conjugated imazalil*

Definition of the residue for dietary risk assessment animal commodities: *sum of imazalil and the metabolite R061000 ((RS)-3-[2-(2,4-dichlorophenyl)-2-(2,3-dihydroxypropoxy)ethyl]imidazolidine-2,4-dione (+)-1-[2-(2,4-dichlorophenyl)-2-[(2,3-dihydroxypropyl)oxy]ethyl]-2,5-imidazolidinedione), expressed as imazalil equivalents.*

The residue is not fat-soluble.

The 2018 JMPR also concluded that while conjugated imazalil is likely to contribute to the dietary exposure for foliar-treated crops, it is not relevant for post-harvest and seed treatment uses.

The Forty-third Codex Alimentarius Commission (2020) approved the new work proposals including the priority list of pesticides for evaluation by the current Meeting. The priority list included imazalil for evaluation of additional uses by the 2021 Extra JMPR. The Meeting received new US labels (confirming the citrus GAP information reported by the 2018 JMPR) and new supporting residue information for mandarins and grapefruits.

Methods of analysis

The Meeting received additional validation information on the analytical method (based on the QuEChERS extraction and LC-MS/MS Method evaluated by the 2018 JMPR) used for measuring free imazalil and its alcohol metabolite (R014821) in the new supervised residue trials on mandarins and grapefruits.

The Meeting concluded that for mandarins and grapefruit, the method used in the new residue trials was sufficiently validated with an LOQ of 0.01 mg/kg and is suitable to measure imazalil and metabolite R014821 in citrus commodities.

Results of supervised residue trials on crops

New supervised trials from Europe on mandarins and grapefruit were available to supplement those reported by the 2018 JMPR. The demonstrated stability of residues in frozen samples (at least 12 months) covered the storage intervals in the trials considered by the Meeting.

The Meeting also received product labels from the USA that confirmed the GAP reported by the 2018 JMPR.

Citrus fruits

The 2018 JMPR recommended maximum residue levels of 15 mg/kg for the Subgroup of Lemons and Limes and 8 mg/kg for the Subgroup of Oranges, Sweet, Sour but concluded there were insufficient data to recommend maximum residue levels for the sub-group of Mandarins and the sub-group of Pummelo and Grapefruits.

The current Meeting received new supervised residue trials on mandarins and grapefruit (involving combined post-harvest dip plus wax spray treatments) to supplement the residue trial information reviewed by the 2018 JMPR.

The 2018 JMPR identified the critical GAP in the USA for citrus fruits as being the combination of 2 post-harvest applications, dip or drench at 0.075 kg ai/hL followed by a wax application of 0.2 kg ai/hL, for a total application rate of 0.275 kg ai/hL and a minimum post-treatment interval of 0 days.

Mandarins

In four trials matching the critical GAP for citrus fruits, residues of imazalil in mandarins (whole fruit) were: 2.1, 2.6, 3.2 and 3.8 mg/kg and residues in flesh were: < 0.01, 0.01, 0.041 and 0.077 mg/kg.

Grapefruit

In four trials matching the critical GAP for citrus fruits, residues of imazalil in grapefruit (whole fruit) were: 2.6, 3.1, 3.2 and 4.7 mg/kg and residues in flesh were: < 0.01 (4) mg/kg in flesh. The median whole fruit residue is 3.15 mg/kg.

Orange

The Meeting noted that in the eight orange trials evaluated by the 2018 JMPR and matching the US GAP for citrus, residues of imazalil in oranges (whole fruit) were: 2.5, 2.7, 3.0, 3.4, 3.6, 4.2, 4.4 and 4.8 mg/kg and residues in flesh were: 0.06, 0.06, 0.07, 0.07, 0.11, 0.13, 0.21 and 0.26 mg/kg.

Lemon

The Meeting noted that in the nine lemon trials evaluated by the 2018 JMPR and matching the US GAP for citrus, imazalil residues in lemons (whole fruit) were: 3.0, 3.1, 3.3, 4.1, 5.0, 5.7, 6.4, 7.8 and 9.7 mg/kg and residues in flesh were: 0.05, 0.05, 0.11, 0.12, 0.18, 0.2, 0.23, 0.27 and 0.36 mg/kg. The highest single residue value in whole fruit was 9.9 mg/kg.

Since the US GAP is for the citrus fruits group, the median residues of mandarins, grapefruit, oranges and lemons are within a 5-fold difference and a statistical analysis (Kruskal-Wallis) shows that the four data sets are not significantly different, the Meeting agreed to combine the datasets to estimate a maximum residue level for citrus fruit.

The combined dataset of residues in citrus (whole fruit) is (n = 25): 2.1, 2.5, 2.6, 2.6, 2.7, 3.0, 3.0, 3.1, 3.1, 3.2, 3.2, 3.3, 3.4, 3.6, 3.8, 4.1, 4.2, 4.4, 4.7, 4.8, 5.0, 5.7, 6.4, 7.8 and 9.7 mg/kg and imazalil residues in flesh are: < 0.01 (5), 0.01, 0.041, 0.05 (2), 0.06 (2), 0.07, 0.07, 0.077, 0.11 (2), 0.12, 0.13, 0.18, 0.2, 0.21, 0.23, 0.26, 0.27 and 0.36 mg/kg. The median whole fruit residue is 3.4 mg/kg.

The Meeting estimated a maximum residue level of 15 mg/kg (based on the mean residue + 4SD) for Citrus fruits and withdrew the previous maximum residue level recommendations of 15 mg/kg for the Subgroup of Lemons and Limes and 8 mg/kg for the Subgroup of Oranges, Sweet, Sour.

The Meeting also estimated an STMR of 0.07 mg/kg and an HR of 0.36 mg/kg for Citrus fruits (flesh) except kumquats and an STMR of 3.4 mg/kg and an HR of 9.9 mg/kg (highest individual residue) for kumquats (whole fruit).

Fate of residues during processing

The 2018 JMPR reviewed information on the fate of imazalil during processing and the estimated processing factors for citrus commodities are listed in Table 1.

STMR-Ps and HR-Ps were calculated for mandarin and grapefruit processed commodities using the higher of the orange or lemon processing factors estimated by the 2018 JMPR.

Residues in processed commodities

Table 1 Citrus commodity processing factors (estimated by the 2018 JMPR) and calculated imazalil STMR-Ps and HR-Ps for citrus commodities based on the highest of the orange or lemon processing factors

Processed commodity	Mean or median processing factors		STMR-P (mg/kg) [STMR _{RAC} × PF]	HR-P (mg/kg) [HR _{RAC} × PF]
	Orange	Lemon	Citrus [median: 3.4]	Citrus [Hi Res: 9.9]
Juice	0.1	0.05	0.34	-
Chopped fresh peel	0.39	0.36	1.3	-
Cold pressed oil	28.6	3.5	97	-
Marmalade	0.27		0.92	-
Canned	0.03		0.1	-
Dry pomace (dried pulp)	4.5		15	

The Meeting applied the orange dry pomace processing factor (4.5) to the estimated maximum residue level for citrus fruits (15 mg/kg) to estimate a maximum residue level of 70 mg/kg (dw) for citrus pulp, dry.

The Meeting applied the orange oil processing factor (28.6) to the estimated maximum residue level for citrus fruits (15 mg/kg) to estimate a maximum residue level of 500 mg/kg for citrus oil, edible.

Residues in animal commodities

Farm animal dietary burden

Dietary burden calculations for beef cattle and dairy cattle and poultry were calculated by the 2018 JMPR, using the OECD diets listed in Appendix IX of the 2016 Edition of the FAO Manual. Maximum dietary burdens were 28.9 ppm (beef cattle), 23.4 ppm (dairy cattle) and 2.3 ppm (poultry) and mean dietary burdens were 25.3 ppm (beef cattle), 19.8 ppm (dairy cattle) and 1.1 ppm (poultry).

The Meeting recalculated these dietary burdens to account for the higher residues in dry citrus pulp and while the dietary burdens based on the Australian, Japanese and US-Canadian (poultry only) diets increased albeit insignificantly, the highest maximum and mean dietary burdens (based on the EU diet) remained unchanged.

Residue in animal commodities

Since the revised maximum and mean dietary burdens for beef and dairy cattle and for poultry broilers and layers were the same as those calculated by the 2018 JMPR, the Meeting agreed that the maximum residue levels, STMRs and HRs recommended by the 2018 JMPR for animal commodities did not need to be revised.

RECOMMENDATIONS

On the basis of the data obtained from supervised trials, the Meeting concluded that the residue levels listed in Annex 1 are suitable for establishing maximum residue limits and for IEDI and IESTI assessments.

Definition of the residue for compliance with the MRL for plant and animal commodities: *imazalil*.

Definition of the residue for dietary risk assessment for plant commodities: *free and conjugated imazalil*.

Definition of the residue for dietary risk assessment for animal commodities: *the sum of imazalil and the metabolite R061000 ((RS)-3-[2-(2,4-dichlorophenyl)-2-(2,3-dihydroxypropoxy)ethyl] imidazolidine-2,4-dione (+)-1-[2-(2,4-dichlorophenyl)-2-[(2,3-dihydroxypropyl)oxy]ethyl]-2,5-imidazolidinedione), expressed as imazalil equivalents.*

The residue is not fat-soluble.

DIETARY RISK ASSESSMENT

Long-term dietary exposure

The ADI for imazalil is 0–0.03 mg/kg bw. The International Estimated Daily Intakes (IEDIs) for imazalil were estimated for the 17 GEMS/Food Consumption Cluster Diets using the STMR or STMR-P values estimated by the JMPR. The results are shown in Annex 3 of the 2021 Extra JMPR Report.

The IEDIs ranged from 3–40% of the maximum ADI. The Meeting concluded that long-term dietary exposure to residues of imazalil from uses considered by the JMPR is unlikely to present a public health concern.

Acute dietary exposure

The ARfD for imazalil is 0.05 mg/kg bw. The International Estimate of Short-Term Intakes (IESTIs) for imazalil were calculated for the food commodities and their processed commodities for which HRs/HR-Ps or STMRs/STMR-Ps were estimated by the present Meeting and for which consumption data were available. The results are shown in Annex 4 of the 2021 Extra JMPR Report.

The IESTIs varied from 0–40% of the ARfD for children and 0–50% of the ARfD for the general population. The Meeting concluded that acute dietary exposure to residues of imazalil from uses considered by the present Meeting is unlikely to present a public health concern.

5.16. ISOPROTHIOLANE (299)

RESIDUE AND ANALYTICAL ASPECTS

Isoprothiolane is a fungicide belonging to the family of dicarboxylic acids, which act by inhibition of phospholipid biosynthesis. It was considered for the first time for toxicology and residues by the 2017 JMPR. An ADI of 0–0.1 mg/kg bw was established and an ARfD was considered to be unnecessary. The 2017 JMPR decided on the following residue definitions based on a metabolism study on paddy rice and confined rotational crops and recommended maximum residue levels for husked rice, rice straw and for animal commodities.

The residue definition for compliance with the MRL for plant commodities is *isoprothiolane*.

The residue definition for dietary risk assessment for rice is *isoprothiolane*.


The residue definition for dietary risk assessment for plant commodities other than rice is the sum of *isoprothiolane*, *diisopropyl-4-hydroxy-1,3-dithiolan-2-ylidenemalonate (M-3)*; *free and conjugated and 1-hydroxypropan-2-yl propan-2-yl 1,3-dithiolan-2-ylidenemalonate (M-5)*; *free and conjugated, expressed as isoprothiolane*.

The residue definition for compliance with the MRL and dietary risk assessment for animal commodities is the sum of *isoprothiolane* and *2-(1,3-dithiolan-2-ylidene)-3-oxo-3-(propan-2-yloxy)propanoic acid (M-2)*, expressed as *isoprothiolane*.

The residue is not fat-soluble.

Isoprothiolane was scheduled at the Fifty-first Session of the CCPR for consideration of additional uses by the 2020 JMPR, which was postponed to the 2021 Extra JMPR.




The Meeting received information on metabolism in apples and grapes, physical and chemical properties, analytical methodology, storage stability, a registered use pattern for bananas and supervised residue trials on bananas.



Isoprothiolane (MW 290.4 g/mol)

Table 1 The following abbreviations are used for Isoprothiolane and metabolites/degradates discussed

Name	Chemical name	Structure
Isoprothiolane (parent)	Diisopropyl 1,3-dithiolan-2-ylidenemalonate	
Isoprothiolane monosulfoxide (M-1)	Diisopropyl 1-oxo-1,3-dithiolan-2-ylidenemalonate	
Isoprothiolane monoester (M-2)	Monoisopropyl 1,3-dithiolan-2-ylidenemalonate	

Name	Chemical name	Structure
4-hydroxy isoprothiolane (M-3)	Diisopropyl 4- hydroxy-1,3- dithiolan-2-ylidenemalonate	
Didehydro isoprothiolane (M-4)	Diisopropyl 1,3- dithiol-2-ylidenemalonate	
Hydroxyl-Isopropyl (M-5)	1-Hydroxypropan-2-yl isopropyl 1,3-dithiolan-2-ylidenemalonate	

Plant metabolism

The Meeting received plant metabolism studies on apples and grapes following applications of [^{14}C]-isoprothiolane.

Apples - Direct application to fruit and leaves

In a metabolism study conducted on apples, ^{14}C -isoprothiolane (radiolabelled in dithiolane ring) was applied as a solution in methanol directly to apple fruit and leaves at 3.14 μg ai/fruit or 1.57 μg ai/leaf. The application rate was equivalent to ca. 0.6 kg ai/ha. The fruits and leaves were collected 7 and 14 days after treatment (DAT).

Samples were surface washed with methanol (MeOH), then sequentially extracted with MeOH followed by MeOH/water (4:1 v/v). Extraction efficiencies were 92.1–98.8%.

In fruit, isoprothiolane was the predominant residue, making up 49.3% TRR in the 7DAT samples and 26.6% TRR in the 14DAT samples. The 4-hydroxy-isoprothiolane (M-3) metabolite (up to 2.1% TRR), together with its glucose conjugates, totalled 9% TRR (7DAT) and 14.1% TRR (14DAT). Other characterised metabolites observed in fruit were isoprothiolane-monosulfoxide (M-1) and didehydro-isoprothiolane (M-4), present at 4.1–4.2% TRR and 4.9–6.6% TRR respectively.

In leaves, isoprothiolane was also the predominant residue, making up 53.9% TRR in the 7DAT samples and 40.3% TRR in the 14DAT samples. The 4-hydroxy-isoprothiolane (M-3) metabolite, together with its glucose conjugates, totalled 13.1% TRR (7DAT) and 17.3% TRR (14DAT). Other characterised metabolites observed in leaves were isoprothiolane-monosulfoxide (M-1) and didehydro-isoprothiolane (M-4), present at up to 9% TRR and 3.8% TRR, respectively. The isoprothiolane monoester (M-2) was only detected (0.1% TRR) in the 14DAT leaves.

Apples - Soil application

In a metabolism study conducted on apples, the soil around pot-grown dwarf apple trees was treated once with ^{14}C -isoprothiolane (radiolabelled in dithiolane ring) using a methanol solution of isoprothiolane of 2.27 mg/mL (equivalent to 360 g ai/tree). Samples of apple fruits and leaves were collected at 7, 28 and 61 days after treatment (DAT). Samples were sequentially extracted with MeOH and MeOH/water (4:1 v/v) with extraction efficiencies of 95% (fruit) and 65–72% (leaves).

In fruit, residues were detected only in the 61 DAT samples and since the TRR was < 0.01 mg eq/kg, no metabolite identification was performed.

In leaves at 61 DAT, isoprothiolane (parent) was detected only at low levels (0.7% TRR), along with the isoprothiolane-monoester (M-2) and isoprothiolane-monosulfoxide (M-1) metabolites, present at 1.2% TRR and 3% TRR, respectively. The major metabolite fraction at 65.5% TRR was at the origin on the thin layer chromatograms (TLC), part of which was characterised by β -glucosidase treatment as the glucoside of isoprothiolane-monoester (M-2) at 13.8% TRR. The remainder of the TLC origin material comprised multiple unidentified components (each <10% TRR).

Grapes - Foliar application

In the metabolism study conducted on grapes, three foliar applications of [^{14}C]- isoprothiolane were applied to grapevines at 1.46 kg ai/ha followed by a fourth application at 0.4 kg ai/ha. The applications were made at pre harvest intervals (PHIs) of 70, 42, 21 and 0 days. Samples of grapes were collected 20 days after the 3rd application (20DAA3) and 0 days after the 4th application (0 DAA4), at a crop growth stage of BBCH 89. Samples were rinsed twice with acetonitrile and stored in a freezer until processing. The acetonitrile rinses were filtered prior to analysis. The total ^{14}C content of the acetonitrile rinses was measured by Liquid Scintillation Counting (LSC). Sequential extraction with acetonitrile; acetonitrile:water (1:1 v/v); acetonitrile:0.2N HCl (1:1 v/v) and acetonitrile:0.2N NH_4OH (1:1 v/v) was able to extract 89.2–90.6% TRR (2.73–3.43 mg eq/kg).

Isoprothiolane was the major radioactive component identified, representing 47–50.7% TRR in grapes. Isoprothiolane-monosulfoxide (M-1) was also an abundant component representing between 11.4–19.5% TRR. Three different glycoside conjugates of 4-hydroxy-isoprothiolane (M-3) were present, ranging individually from 2.3–7.6% TRR. The total free and conjugated 4-hydroxy-isoprothiolane represented 13.2–14% TRR.

Methods of analysis

Analytical methods have been reported and validated for the analysis of isoprothiolane in bananas.

The data generation method involved a number of extractions with acetonitrile: water (1:1, v/v) and acetonitrile. Conjugates were cleaved with β -glucosidase enzyme and 0.1M sodium acetate buffer solution (pH 5) with a 24-hour incubation period. The residues were quantified by LC-MS/MS. The LOQs for isoprothiolane, isoprothiolane monosulfoxide (M-1) and 4-hydroxy isoprothiolane (M-3) were demonstrated to be 0.01 mg/kg for each analyte in banana (whole fruit), banana (flesh) and banana (peel).

For MRL compliance, a multi-residue method is available for analysis of isoprothiolane on bananas based on QuEChERS extraction/clean-up and HPLC-MS/MS detections. The LOQ for isoprothiolane was determined to be 0.01 mg/kg in banana (whole fruit).

The Meeting concluded that the above methods were sufficiently validated and are suitable for measuring residues of isoprothiolane and/or its -monosulfoxide and its free and conjugated -4-hydroxy metabolites in bananas.

Stability of pesticide residues in stored analytical samples

Residues of isoprothiolane were observed to be stable for at least 14 months in banana whole fruit and for 6 months in banana peel and flesh under freezer storage conditions. Residues of 4-hydroxy-isoprothiolane and isoprothiolane monosulfoxide were observed to be stable for at least 17 months under freezer storage conditions in whole fruit, flesh and peel. These durations covered the longest period of storage for all samples obtained from supervised residue trials.

Definition of the residue

The 2017 JMPR considered a metabolism study conducted on paddy rice, in which isoprothiolane (parent) was the predominant residue, accounting for 16–76% TRR in rice grain, hulls and stems/leaves. The 2017 JMPR also considered a confined rotational study conducted on lettuce, radish and wheat. Isoprothiolane was observed in most matrices at the 30-day plant-back interval, however, the metabolites 4-hydroxy-isoprothiolane (M-3) and hydroxyl-isopropyl (M-5) generally made up the majority of the residue. Isoprothiolane was not observed in wheat grain.

Based on these studies, for plant commodities, the 2017 JMPR established a residue definition of isoprothiolane (parent only) for MRL-compliance and also for dietary risk assessment for rice. For other plant commodities, the established residue definition for dietary risk assessment is the sum of isoprothiolane and metabolites M-3 and M-5 (expressed as parent compound).

Plant commodities

In the grape metabolism study, isoprothiolane (parent) was the major radioactive component identified representing 47–50.7% TRR. Isoprothiolane monosulfoxide (M-1) was also a relatively abundant component representing between 11.4–19.5% TRR. Three different glycoside conjugates of 4-hydroxy-isoprothiolane were present, individually ranging from 2.3–7.6% TRR. The total residues of 4-hydroxy-isoprothiolane (M-3) and its conjugates represented 13.2–14% TRR. No hydroxyl-isopropyl (M-5) was detected in the study.

In the metabolism study on apples involving direct application to leaves and fruit, isoprothiolane (parent) was the most abundant component in fruit representing 26.6–49.3% TRR. In fruit, the 4-hydroxy-isoprothiolane (M-3) and its conjugates accounted for 9–14.1% TRR, with other identified metabolites being the monosulfoxide derivative (M-1), present at about 4.1–4.2% TRR and the didehydro derivative (M-4), accounting for 4.9–6.6% TRR.

In the supervised bananas field trials, isoprothiolane was detected at residue levels of < 0.01–0.59 mg/kg. Residues of 4-hydroxy isoprothiolane (M-3) were detected in whole fruit in 7 of the 9 independent trials at < 0.01–0.02 mg/kg. In the three trials where peel and flesh were analysed, no quantifiable residues of 4-hydroxy isoprothiolane were detected in the flesh. Quantifiable residues of isoprothiolane monosulfoxide (M-1) were not observed in whole fruit and residues of hydroxyl-isopropyl (M-5) were not analysed.

Based on the available metabolism studies, the Meeting noted that isoprothiolane was the predominant residue in fruit matrices. The Meeting confirmed its previous recommendation of isoprothiolane as a suitable residue definition for MRL-compliance in plant matrices.

For the purpose of dietary exposure estimation, parent isoprothiolane and 4-hydroxy isoprothiolane (free and conjugated) represent a significant part of the residue in fruit crops. The only additional major metabolite found in fruit was isoprothiolane-monosulfoxide (M-1) in grapes (up to 19.5% TRR) and in apples (up to 4.2% TRR), but it was not found in the flesh of bananas in the supervised field trials.

The Meeting agreed that the current residue definition for dietary risk assessment also applies to banana but noted that should additional uses be considered in future this residue definition will need to be revisited.

Results of supervised residue trials on crops

The Meeting received GAP information and residue trials on bananas.

Bananas

The cGAP for bananas in Guatemala, is for up to 12 foliar applications at 400 g ai/ha, with a re-treatment interval of 21 days and a PHI of 0 days. Twelve residue trials, matching cGAP, from Colombia, Costa Rica, Ecuador, Guatemala and Honduras were available to the Meeting. The Meeting noted that while all trials matched the cGAP, not all trials were independent. Residues were observed to be higher from the unbagged trials and these results were used for maximum residue level estimation.

In nine independent trials matching the cGAP the residues in banana (whole fruit) were (n = 9): 0.065, 0.09, 0.13, 0.16, 0.17, 0.28, 0.31, 0.47 and 0.56 mg/kg.

The Meeting estimated a maximum residue level of 1 mg/kg for isoprothiolane in bananas.

Based on the information available the Meeting agreed that the -hydroxyl-isopropyl metabolite (M-5) could be excluded from the STMR calculations in banana as this metabolite has only been reported in rotational crops.

For dietary risk assessment, in the above trials, total residues (i.e., parent plus the free and conjugated M-3 metabolite) in whole fruit were (n = 9): 0.075, 0.1, 0.14, 0.17, 0.18, 0.29, 0.32, 0.48

and 0.57 mg/kg and in three of these trials, total residues in flesh were ($n = 3$): < 0.02 , < 0.02 and 0.02 mg/kg.

In the one trial where quantifiable residues (0.02 mg/kg) were found in the flesh total residues in whole fruit were 0.465 mg/kg (a flesh:whole fruit ratio of 0.043). The Meeting agreed to apply this ratio to the median total residues in whole fruit (0.18 mg/kg) to estimate an STMR of 0.0077 mg/kg for bananas.

Residues in animal commodities

The 2017 JMPR considered residues in animal commodities and recommended maximum residue levels. As bananas are not considered to be a livestock feed item, the livestock dietary burden remains unchanged. No amendments to the previously considered STMRs and maximum residue levels for animal commodities are required.

RECOMMENDATIONS

On the basis of the data obtained from supervised trials, the Meeting concluded that the residue levels listed in Annex 1 are suitable for establishing maximum residue limits and for IEDI assessment.

The residue definition for compliance with the MRL for plant commodities is *isoprothiolane*.

The residue definition for dietary risk assessment for rice is *isoprothiolane*.

The residue definition for dietary risk assessment for plant commodities other than rice is the sum of *isoprothiolane*, *diisopropyl-4-hydroxy-1,3-dithiolan-2-ylidenemalonate (M-3)*; *free and conjugated* and *1-hydroxypropan-2-yl propan-2-yl 1,3-dithiolan-2-ylidenemalonate (M-5)*; *free and conjugated*, expressed as *isoprothiolane*.

The residue definition for compliance with the MRL and dietary risk assessment for animal commodities is the sum of *isoprothiolane* and *2-(1,3-dithiolan-2-ylidene)-3-oxo-3-(propan-2-yl oxy)propanoic acid (M-2)*, expressed as *isoprothiolane*.

The residue is not fat-soluble.

DIETARY RISK ASSESSMENT

Long-term dietary exposure

The ADI for isoprothiolane is 0–0.1 mg/kg bw. The International Estimated Daily Intakes (IEDIs) for isoprothiolane were estimated for the 17 GEMS/Food Consumption Cluster Diets using the STMR or STMR-P values estimated by the JMPR. The results are shown in Annex 3 of the 2021 Extra JMPR Report.

The IEDIs ranged from 0–2% of the maximum ADI. The Meeting concluded that long-term dietary exposure to residues of isoprothiolane from uses considered by the JMPR is unlikely to present a public health concern.

Acute dietary exposure

The 2017 JMPR decided that an ARfD is unnecessary. The Meeting concluded that the acute dietary exposure to residues of isoprothiolane from the uses considered is unlikely to present a public health concern.

5.17. ISOXAFLUTOLE (268)

RESIDUE AND ANALYTICAL ASPECTS

Isoxaflutole is a herbicide that was first evaluated by the 2013 JMPR, when an ADI of 0–0.02 mg/kg bw was established and an ARfD was considered unnecessary.

The residue definition for isoxaflutole for compliance with the MRL for plant and animal commodities and for dietary risk assessment for plant commodities, including tolerant soya beans, is the *sum of isoxaflutole and isoxaflutole diketonitrile, expressed as isoxaflutole*.

The residue definition for isoxaflutole for dietary risk assessment for animal commodities is the *sum of isoxaflutole, isoxaflutole diketonitrile, RPA 205834 (2-aminomethylene-1-cyclopropyl-3-(2-mesyl-4-trifluoromethylphenyl)-propane-1,3-dione) and RPA 207048 (1-cyclopropyl-2-hydroxymethylene-3-(2-mesyl-4-trifluoromethylphenyl)-propane-1,3-dione), including their conjugates, expressed as isoxaflutole*.

The residue is not fat-soluble.

Isoxaflutole was scheduled at the Fifty-first Session of the CCPR (2019) for evaluation of additional uses by the 2020 JMPR, which was postponed to the 2021 Extra JMPR. The Meeting received supervised residue trials on soya bean that had been previously evaluated by the 2013 JMPR and new GAP information on isoxaflutole in/on glyphosate/HPPD tolerant soya bean.

Results of supervised residue trials on crops

Soya bean (dry)

Isoxaflutole is registered for use in the USA as a post-emergence application to glyphosate tolerant and isoxaflutole-tolerant soya bean. The critical GAP is a single application of 0.09 kg ai/ha up to bud not including first bloom growth stage (BBCH 59). The PHI is at least 70 days. Livestock are not allowed to graze or feed forage or hay from treated soya bean.

A total of 77 residue trials were conducted with isoxaflutole in/on glyphosate/HPPD tolerant soya bean in Canada and the USA, where the compound was applied at various BBCH stages and harvested at different days after the application.

The Meeting agreed to consider residue data from all trials approximating the post-emergence (BBCH 59) timing that were conducted with the PHI of at least 70 days.

The residues are (n = 24): < 0.02 (15), 0.02, 0.021, 0.022, 0.023, 0.025, 0.026, 0.028, 0.030 and 0.037 mg/kg.

The Meeting estimated a maximum residue level of 0.04 mg/kg and an STMR of 0.02 mg/kg for isoxaflutole in soya beans.

Fate of residues during processing

Processing factors estimated for soya beans by the 2013 JMPR are 1.2 for meal, 0.79 for hulls, < 0.4 for RBD oil (Refined, bleached and deodorized) and milk and 6 for aspirated grain fractions.

Based on an STMR of 0.02 mg/kg for soya beans, the Meeting estimated STMR-Ps of 0.008 mg/kg for soya bean refined oil and milk and median residues of 0.024 mg/kg for soya bean meal, 0.016 mg/kg for soya bean hulls and 0.12 mg/kg for soya bean aspirated grain fractions.

Residues in animal commodities

Including the median residues of soya bean processed commodities and the STMR for soya bean (dry) did not significantly impact the dietary burden of isoxaflutole estimated by the 2013 JMPR.

The Meeting confirmed its previous recommendations for isoxaflutole in animal commodities.

RECOMMENDATIONS

On the basis of the data obtained from supervised trials, the Meeting concluded that the residue levels listed in Annex 1 are suitable for establishing maximum residue limits and for IEDI assessment.

Definition of the residue for isoxaflutole for compliance with the MRL for plant and animal commodities and for dietary risk assessment for plant commodities, including tolerant soya beans, is the *sum of isoxaflutole and isoxaflutole diketonitrile, expressed as isoxaflutole*.

The residue definition for isoxaflutole for dietary risk assessment for animal commodities is the *sum of isoxaflutole, isoxaflutole diketonitrile, RPA 205834 (2-aminomethylene-1-cyclopropyl-3-(2-mesyl-4-trifluoromethylphenyl)-propane-1,3-dione) and RPA 207048 (1-cyclopropyl-2-hydroxymethylene-3-(2-mesyl-4-trifluoromethylphenyl)-propane-1,3-dione), including their conjugates, expressed as isoxaflutole*.

The residue is not fat-soluble.

DIETARY RISK ASSESSMENT

Long-term dietary exposure

The ADI for isoxaflutole is 0–0.02 mg/kg bw. The International Estimated Daily Intakes (IEDIs) for isoxaflutole were estimated for the 17 GEMS/Food Consumption Cluster diets using the STMR or STMR-P values estimated by the JMPR. The results are shown in Annex 3 of the 2021 Extra JMPR Report.

The IEDIs accounted for 0% of the maximum ADI. The Meeting concluded that the long-term dietary exposure to residues of isoxaflutole from uses considered by the JMPR is unlikely to present a public health concern

Acute dietary exposure

The 2013 JMPR decided that an ARfD for isoxaflutole is unnecessary. The Meeting therefore concluded that the acute dietary exposure to residues of isoxaflutole from the uses considered is unlikely to present a public health concern.

5.18. MANDIPROPAMID (231)

RESIDUE AND ANALYTICAL ASPECTS

Mandipropamid is a fungicide in the mandelamide class used for the control of foliar oomycete pathogens in a range of crops, including *Plasmopara viticola* in grapes, *Phytophthora infestans* in potatoes and tomatoes and *Pseudoperonospora cubensis* in cucurbits. Mandipropamid was first evaluated by the JMPR in 2008 when an ADI of 0–0.2 mg/kg bw was established and maximum residue levels were recommended for various crops. An ARfD was considered unnecessary. Additional uses were evaluated by the 2013 and 2018 JMPR. The JMPR agreed on the following residue definition for mandipropamid:

Definition of the residue for compliance with the MRL and dietary risk assessment for plant and animal commodities: *mandipropamid*.

The residue is not fat-soluble.

The Forty-third Codex Alimentarius Commission (2020) approved the new work proposals including the priority list of pesticides for evaluation by the current Meeting. The priority list included mandipropamid for evaluation of an additional use on citrus fruits.

The current Meeting received information on field trials and processing studies to support the estimation of new maximum residue levels in citrus fruit and citrus processed commodities.

Methods of analysis

The current Meeting received method validation and concurrent recovery data for use of Method WO 9.210 v.1 and v.2, both based on the validated reference method RAM 415/02. Both methods were demonstrated to have adequate performance for recovery of mandipropamid in citrus fruits and citrus processed commodities. An LOQ of 0.01 mg/kg was established for both methods for all citrus raw agricultural commodities and processed fractions.

The 2008 JMPR concluded that the multi-residue method DFG S19 is suitable for monitoring residues of mandipropamid in plant commodities, with an LOQ of 0.01 mg/kg.

Stability of pesticide residues in stored analytical samples

Storage stability data were previously provided for mandipropamid in various commodities. The 2018 JMPR concluded that mandipropamid is stable at -20 °C for at least 24 months in tomatoes (fruit and paste), grapes (fruit and juice), potatoes (tubers and granules/flakes), lettuce, cucumbers, wheat (forage, grain and straw) and soya beans (beans, hulls, meal and oil); and for at least 12 months in green beans (beans, pods with seeds and beans, whole plants with pods). The storage stability data are adequate to support the supervised field trials and processing study currently under evaluation.

Results of supervised residue trials on crops

The Meeting received GAP information for mandipropamid on citrus fruits in the USA and supervised residue trials for mandipropamid on lemons, mandarins, oranges and grapefruit.

Citrus Fruit

For all fruit, samples for whole fruit analysis were quartered in the field and samples for peel and pulp residue distribution were left whole. Comparison of whole fruit residues between quartered samples and samples left whole indicate that quartering in the field does not affect the stability of mandipropamid residue levels in citrus fruit.

The registered GAP in the USA for citrus fruits allows two foliar applications at 146 g ai/ha (seasonal application up to 291 g ai/ha) with a minimum re-treatment interval (RTI) of 30 days and a 0-day pre-harvest interval (PHI).

Lemon

Five field trials on lemons were conducted in the USA matching the USA GAP. Residues of mandipropamid were (n = 5): 0.085, 0.095, 0.098, 0.19 and 0.23 mg/kg.

Mandarin

Two field trials were conducted in the USA matching the USA GAP. Residues of mandipropamid were (n = 2): 0.096 and 0.17 mg/kg.

Orange

Eight independent field trials on oranges were conducted in the USA. Six trials matched the USA GAP. At two trials, both applications were made at ~40% of the GAP application rate (57–58 g ai/ha per application). Residues following applications matching the USA GAP were (n = 6): 0.050, 0.081, 0.088, 0.11, 0.15 and 0.19 mg/kg. Residues following application at ~40% of the GAP application rate were (n = 2): 0.034 and 0.047 mg/kg.

Residues scaled to the USA GAP (scaling factors of 0.98–2.5x) were (n = 8): 0.049, 0.081, 0.085, 0.087, 0.11, 0.12, 0.15 and 0.19 mg/kg.

Grapefruit

Six field trials on grapefruit were conducted in the USA. Five trials matched the USA GAP. At one trial, both applications were made at ~40% of the GAP application rate (58 g ai/ha per application). Residues following applications matching the USA GAP were (n = 5): 0.043, 0.050, 0.055, 0.057 and 0.079 mg/kg. The residue following application at ~40% of the GAP application rate was 0.034 mg/kg.

Residues scaled to the USA GAP (scaling factors of 0.97–2.5) were (n = 6): 0.043, 0.050, 0.053, 0.057, 0.077 and 0.085 mg/kg.

The median whole fruit residues for oranges, lemons and grapefruits were within 5x of one another. However, the Kruskal-Wallis test showed that the residue values were from different populations. The Meeting noted that the USA GAP covers all citrus fruit and estimated the following maximum residue levels and median values for citrus subgroups:

Subgroup of Lemons and Limes: Maximum residue level of 0.5 mg/kg. Median value of 0.098 mg/kg.

Subgroup of Oranges, Sweet, Sour: Maximum residue level of 0.4 mg/kg. Median value of 0.099 mg/kg.

Subgroup of Pummelo and Grapefruits: Maximum residue level of 0.2 mg/kg. Median value of 0.055 mg/kg.

Subgroup of Mandarins: An insufficient number of trials on mandarin were available to estimate an STMR or maximum residue level. However, the Meeting noted that residues in lemons were shown to be similar to or greater than residues in mandarins. The Meeting decided to extrapolate the residues from lemon and estimated a maximum residue level of 0.5 mg/kg and a median value of 0.098 mg/kg.

At one lemon trial, one mandarin trial, two orange trials and two grapefruit trials, additional fruit samples were separated into peel and flesh fractions. For all citrus fruit commodities (n = 6), residues in the flesh were below the LOQ (< 0.01 mg/kg). The Meeting concluded that the STMR for the flesh of all citrus fruit is 0.01 mg/kg.

Fate of residues during processing

The Meeting received an orange processing study. The processed fractions included orange dried pulp, orange juice and orange oil. The Meeting decided to extrapolate processing factors from orange to all citrus commodities. Processing factors and residue estimates are summarized below.

Table 1 Residues of mandipropamid in citrus processed commodities

Processed Commodity	Processing Factor	Residue (RAC)		Residue (Processed Commodity)	
		Maximum Residue Level	Median	Max-P (mg/kg)	STMR-P ^a (mg/kg)
Orange Juice	< 0.043	0.4	0.099	--	< 0.0043
Orange Oil	45			--	4.5
Orange Dried Pulp	2.9			--	0.29
Lemon/Mandarin Juice ^a	< 0.043	0.5	0.098	--	< 0.0042
Lemon/Mandarin Oil ^b	45			22.5	--
Lemon/Mandarin Dried ^c Pulp	2.9			1.45	--
Grapefruit Juice ^a	< 0.043	0.2	0.055	--	< 0.0024
Grapefruit Oil ^b	45			--	--
Grapefruit Dried Pulp ^c	2.9			--	--

^a Lemon/mandarin and grapefruit juice processing factor extrapolated from orange processing study

^b Lemon/mandarin and grapefruit oil processing factor extrapolated from orange processing study

^c Lemon/mandarin and grapefruit dried pulp processing factor extrapolated from orange processing study

Citrus dried pulp: Based on the maximum residue level of 0.5 mg/kg for lemons and median value of 0.099 mg/kg for oranges, the Meeting estimated a maximum residue level of 1.5 mg/kg and a median value of 0.29 mg/kg.

Citrus oil: Based on the maximum residue level of 0.5 mg/kg for lemons and median value of 0.099 mg/kg for oranges, the Meeting estimated a maximum residue level of 30 mg/kg and an STMR of 4.5 mg/kg.

Residues in animal commodities

Farm animal dietary burdens

Dietary burdens were calculated for beef cattle, dairy cattle, broilers and laying poultry based on feed items evaluated by the JMPR. The dietary burdens were estimated using the most recent version of the OECD livestock dietary burden calculator. Dietary burdens for mandipropamid are presented in Annex 6 and summarised in Table 2, below. In addition to citrus dried pulp, kale leaves were added to the previous dietary burdens as Codex has an established MRL for the leafy vegetable Crop Group.

The Australian beef cattle maximum and mean dietary burdens of 17.51 and 9.07 ppm, respectively and dairy cattle maximum and mean dietary burdens of 49.21 and 23.77 ppm, respectively, were refined to exclude bean forage and kale. Mandipropamid is not registered for use on beans in Australia and Australia does not import forages. The labelled use for kale in Australia prohibits feeding to livestock and Australia does not import kale as a feed item. The refined Australian beef cattle maximum and mean dietary burdens are 1.62 and 1.61 ppm, respectively and the refined Australian dairy cattle maximum and mean dietary burdens are 0 ppm.

Table 2 Livestock dietary burdens for mandipropamid (ppm)

	USA-Canada		EU		Australia		Japan	
	Max	Mean	Max	Mean	Max	Mean	Max	Mean
Beef cattle	0.21	0.13	15.58	7.70	1.62	1.61	--	--
Dairy cattle	0.070	0.043	20.84 ^a	10.15 ^b	--	--	--	--

	USA-Canada		EU		Australia		Japan	
	Max	Mean	Max	Mean	Max	Mean	Max	Mean
Poultry – broiler	--	--	0.041	0.014	--	--	--	--
Poultry - layer	--	--	1.97 ^c	1.20 ^d	--	--	--	--

^a Highest maximum beef or dairy cattle dietary burden used for mammalian milk and tissue maximum residue level calculations.

^b Highest mean beef or dairy cattle dietary burden used for mammalian milk and tissue STMR estimates.

^c Highest maximum poultry dietary burden used for poultry tissue and egg maximum residue level calculations.

^d Highest mean poultry dietary burden used for poultry tissue and egg STMR estimates.

Animal commodity maximum residue levels

The highest maximum and mean estimated dietary burdens for cattle are 20.84 and 10.15 ppm, respectively. No animal feeding studies on ruminants are available, therefore the lactating goat metabolism study provided to the 2008 JMPR was used to estimate residues of mandipropamid in mammalian commodities.

Lactating goats were fed for seven days with [¹⁴C] mandipropamid equivalent to 30 ppm in the diet for the methoxyphenyl label and 27 or 45 ppm in the diet for the chlorophenyl label. The methoxyphenyl label produced higher residues of mandipropamid in milk and all tissues and therefore was used as the basis for anticipated residue calculations. Anticipated residues of mandipropamid in tissues are presented in Table 3.

Table 3 Anticipated Residues in Ruminant Milk and Tissues

	Feed level (ppm)	Estimated Residue (mg/kg)				
		Milk	Muscle	Liver	Kidney	Fat
Metabolism study	30	0.0008	ND	0.0065	ND	0.019
Maximum dietary burden and estimated residues	20.84	< 0.001 ^a	NA	0.0045	NA	0.013
Mean dietary burden and estimated residue	10.15	< 0.001 ^a	NA	0.0022	NA	0.0064

ND = Not detected. NA = Not applicable.

^a The STMR anticipated residue of < 0.001 mg/kg in milk is considered negligible and a value of 0 was used in the dietary assessment.

Applying the maximum cattle dietary burden, the Meeting estimated a maximum residue level of 0.02 mg/kg for mandipropamid in mammalian fat (excluding milk fats) to replace its previous recommendation of 0.01(*) mg/kg. The Meeting confirmed its previous maximum residue level estimations of 0.01(*) mg/kg for all other mammalian commodities.

Applying the mean cattle dietary burden, the Meeting concluded that the previous STMR of 0 mg/kg for mammalian meat and milk are appropriate. The Meeting estimated new STMRs of 0.0064 mg/kg for mammalian fat (excluding milk fats) and 0.0022 mg/kg for mammalian edible offal.

The maximum and mean estimated dietary burdens for poultry are unchanged from those estimated by the 2018 JMPR. Therefore, the Meeting confirmed its previous recommendations for poultry commodities.

RECOMMENDATIONS

On the basis of the data obtained from supervised trials, the Meeting concluded that the residue levels listed in Annex 1 are suitable for establishing MRLs and for IEDI assessment.

Definition of the residue for compliance with the MRL and dietary risk assessment for plant and animal commodities: *mandipropamid*.

The residue is not fat-soluble.

DIETARY RISK ASSESSMENT

Long-term dietary exposure

The ADI for mandipropamid is 0–0.2 mg/kg bw. The International Estimated Daily Intakes (IEDIs) for mandipropamid were estimated for the 17 GEMS/Food Consumption Cluster Diets using the STMR or STMR-P values estimated by the JMPR. The results are shown in Annex 3 of the 2021 Extra JMPR Report.

The IEDIs ranged from 0–6% of the maximum ADI. The Meeting concluded that long-term dietary exposure to residues of mandipropamid from uses considered by the JMPR is unlikely to present a public health concern.

Acute dietary exposure

The 2008 JMPR decided that an ARfD for mandipropamid was unnecessary. Therefore, the Meeting concluded that the acute dietary exposure to residues of mandipropamid from the uses considered is unlikely to present a public health concern.

5.19. METHOPRENE (147)

RESIDUE AND ANALYTICAL ASPECTS

Methoprene is an insect growth regulator that was first evaluated by the JMPR in 1984, it received a periodic review in 2001 for toxicology and in 2005 for residues and was evaluated for additional uses in 2016 and 2019 (extra Meeting).

S-methoprene is the biologically active enantiomer in the racemic methoprene and constitutes 50% of methoprene.

An ADI was established at 0–0.09 mg/kg bw for racemic methoprene and at 0–0.05 mg/kg bw for S-methoprene; an ARfD was considered unnecessary.

The residue definition for compliance with the MRL and dietary risk assessment for plant and animal commodities is Methoprene. The residue is fat-soluble.

S-Methoprene was scheduled at the Fifty-first Session of the CCPR (2019) for evaluation by the 2020 JMPR, which was postponed to the 2021 Extra JMPR.

The Meeting received post-harvest supervised residue trials on soya beans for S-methoprene and GAP information.

Results of supervised residue trials on crops

Soya bean (dry) - post harvest use

The critical GAP for S-methoprene in the USA for protection of stored food, grain, animal feed and seeds used for oil is 2.4 g ai/tonne with no withholding period specified. Five trials were conducted in the USA with a single application at 2.4 g ai/tonne and samples collected after treatment gave methoprene residues in rank order (n = 5): 2.1, 2.4, 2.4, 2.5 and 2.6 mg/kg.

As in the trials, where S-methoprene was applied separately to different lots simulating commercial application practice, the results reflected a high recovery of applied methoprene (88 to 108% of the 2.4 g ai/tonne applied in all the trials), the Meeting decided that the application rate determined the level of residue expected at the zero-day withholding period of the GAP and that residues of up to about 2.4 mg/kg can be anticipated.

The Meeting estimated a maximum residue level of 3 (Po) mg/kg and an STMR of 2.4 mg/kg for soya bean (dry).

Residues in animal commodities

Soya bean seed can be fed to livestock. The 2019 Extra JMPR evaluated residues of methoprene in cereal grains, oilseeds and peanuts. Estimation by the present Meeting, now including soya bean (dry), does not increase the previously estimated (2019) maximum dietary burdens of 13.46 ppm in the diet of cattle and 10.62 ppm for poultry. The Meeting confirmed its previous conclusions for animal commodities.

RECOMMENDATIONS

On the basis of the data obtained from supervised trials, the Meeting concluded that the residue levels listed in Annex 1 are suitable for establishing maximum residue limits and for IEDI assessment.

Definition of the residue for compliance with the MRL and dietary risk assessment for plant commodities: methoprene

Definition of the residue for compliance with the MRL and dietary risk assessment for animal commodities: methoprene

The residue is fat-soluble.

DIETARY RISK ASSESSMENT***Long-term dietary exposure***

The ADI for S-methoprene is 0–0.05 mg/kg bw. The International Estimated Daily Intakes (IEDIs) for methoprene were estimated for the 17 GEMS/Food Consumption Cluster Diets using the STMR or STMR-P values estimated by the JMPR. The results are shown in Annex 3 of the 2021 Extra JMPR Report.

Assuming the residues are S-methoprene, the IEDIs ranged from 10-60% of the maximum ADI. The Meeting concluded that long-term dietary exposure to residues of methoprene from uses considered by the JMPR is unlikely to present a public health concern.

Acute dietary exposure

The 2001 JMPR decided that an ARfD for methoprene was unnecessary. The Meeting therefore concluded that the acute dietary exposure to residues of methoprene from the uses considered is unlikely to present a public health concern.

5.20. METHOXYFENOZIDE (209)

RESIDUE AND ANALYTICAL ASPECTS

Methoxyfenozide is an insecticide that mimics moulting hormone of Lepidopterous larvae. The JMPR evaluated it first in 2003 for toxicology and residues and in 2006, 2009 and 2012 for additional uses. The 2003 JMPR established an ADI of 0–0.1 mg/kg bw and an ARfD of 0.9 mg/kg bw. The residue definition is methoxyfenozide for compliance with the MRL and dietary risk assessment for plant and animal commodities. The residue is fat-soluble in its distribution between meat muscle and fat, but not in its distribution in milk.

The Forty-third Codex Alimentarius Commission (2020) approved the new work proposals including the priority list of pesticides for evaluation by the current Meeting. The priority list included methoxyfenozide for evaluation of additional uses. The Meeting received information on residue trials on basil, coffee bean, rice, sugar cane and tea.

Methods of analysis

Methods used for analysis of methoxyfenozide were based on Method 02.25 in the basil and rice trials and Method 02.24 in the coffee and sugar cane trials. The two methods, using determination by LC-MS/MS, were considered by the 2012 JMPR as validated. In recovery tests for methoxyfenozide residues in basil, coffee, rice and sugar cane, recovery results were satisfactory. The LOQ values were 0.01 mg/kg in fresh basil, coffee (green bean, roasted bean and soluble coffee) and sugar cane (stalk, brown sugar and molasses) and 0.05 mg/kg in dried basil and rice (grain, hulls, bran and polished rice).

In the tea trials conducted in 2016–2018, analytical methods for tea leaves and infusion involved extraction with acetonitrile (omitting extraction for infusion samples), clean-up steps by column chromatography and determination by LC-MS/MS. In recovery tests for methoxyfenozide in dried leaves and infusion, recovery results were satisfactory and the LOQ values were 0.01 mg/kg in both matrices. For the 1998 trials, a different analytical method based on HPLC-UV was used and the Meeting estimated an LOQ of 0.8 mg/kg, based on the lowest fortification level with acceptable recoveries.

Stability of pesticide residues in stored analytical samples

Storage stability tests, covering the actual storage days of samples (covering >90% in basil samples) were conducted for basil (fresh, dry), rice (hulls, bran, polished rice) and dried tea leaves, demonstrating that methoxyfenozide residues were stable. For coffee bean and sugar cane samples, the maximum storage interval was 268 days and 200 days, respectively. The previous JMPR (2003) considered that residues were stable for at least for 6–12 months in various plant commodities, when stored frozen. In addition, the 2012 JMPR considered that residues were stable for at least 21 months in spring onion and 15 months in dry pea. Based on the available information, the Meeting considered that methoxyfenozide residues were stable for the sample storage intervals used in the residue field trials considered.

Results of supervised residue trials on crops

Cereal grains – rice

The critical GAP on rice in the USA is two applications at 0.18 kg ai/ha with a minimum re-treatment interval of 10 days and a PHI of 14 days. The Meeting received eight trials (three independent trials) conducted in California, USA, at rates of 0.21–0.22 kg ai/ha with two applications 8–11 days apart and harvest 13 days after the last application. The residues in rice were ($n = 3$): 8.3, 9.1 and 12 mg/kg.

The Meeting considered the number of trials insufficient for estimating a maximum residue level for rice.

Grasses for sugar or syrup – sugar cane

The critical GAP on sugar cane in Brazil is two applications at 0.06 kg ai/ha and a PHI of 60 days (the re-treatment interval is specified as, determined by re-infestation). Eight trials were conducted in Brazil, matching the critical GAP. The residues were (n = 8): < 0.01 (7) and 0.01 mg/kg.

The Meeting estimated a maximum residue level of 0.015 mg/kg, an STMR of 0.01 mg/kg and an HR of 0.01 mg/kg for methoxyfenozide in sugar cane.

Seeds for beverages and sweets – Coffee bean

The critical GAP on coffee in Brazil is two applications at 0.09 kg ai/ha with a minimum re-treatment interval of 30 days and a PHI of 7 days. Six trials were conducted in Brazil, matching the critical GAP. The residues in coffee bean (green, dry) were (n = 6): < 0.01, 0.01, 0.02, 0.035, 0.045 and 0.061 mg/kg.

The Meeting estimated a maximum residue level of 0.15 mg/kg and an STMR of 0.0275 mg/kg for methoxyfenozide in coffee bean.

Herbs – basil

The critical GAP on herbs in the USA is an application rate of 0.28 kg ai/ha (1.1 kg ai/ha/year) with a minimum re-treatment interval of 10 days and a PHI of 1 day. Five trials on basil were conducted in the USA, matching the critical GAP. The residues in fresh basil were (n = 5): 9.9, 15, 19, 21 and 43 mg/kg (highest analytical value 47 mg/kg). The residues in dried basil were (n = 5): 60, 92, 100, 110 and 190 mg/kg (highest analytical value 194 mg/kg).

The Meeting estimated a maximum residue level of 80 mg/kg, an STMR of 19 mg/kg and an HR of 47 mg/kg for methoxyfenozide in basil, leaves. In addition, the Meeting estimated a maximum residue level of 400 mg/kg, an STMR of 100 mg/kg and an HR of 194 mg/kg for methoxyfenozide in basil, dry.

Teas – green tea

The critical GAP on tea in Japan is two applications with a dilution rate of 4,000 times (equivalent to 5 g/hL) and a PHI of 7 days. Six trials (2016–2018) on tea were conducted in Japan, matching the critical GAP. The residues in dried tea leaves were (n = 6): 9.1, 18, 28, 29, 29 and 35 mg/kg.

The Meeting estimated a maximum residue level of 80 mg/kg and an STMR of 28.5 mg/kg for methoxyfenozide in tea, green, black (black, fermented and dried).

Residues in animal feeds

A maximum residue level on rice as a food commodity could not be estimated, therefore, the Meeting did not estimate a maximum residue level on rice straw.

Fate of residues during processing

The Meeting received information on the fate of methoxyfenozide during processing of coffee bean, rice, sugar cane and tea. Processing factors for rice, sugar cane and green tea were estimated and are shown in the table below. For coffee bean, the residue concentration in the raw agricultural commodity was below the limit of quantification, therefore, the processing factors could not be estimated.

Table 1 Processing factors for rice, sugar cane and green tea

Raw agricultural commodity [STMR] (mg/kg)	Processed commodity	Individual processing factors	Mean or best estimate processing factor	STMR-P = STMR _{RAC} ×Pf (mg/kg)
Rice grain	Rice, hull	4.5	4.5	
	Rice, bran	1.0	1.0	
	Polished rice	0.013	0.013	
Sugar cane, stalk [0.01]	Molasses	7	7	0.07
	Brown sugar	6	6	0.06
Dried tea leaves [28.5]	Tea infusion	0.005 (5), 0.006	0.005	0.14

For rice, the maximum residue level on the raw agricultural commodity was not estimated in this Meeting, therefore, the STMR-Ps were not estimated.

A maximum residue level for sugar cane molasses was estimated as 0.1 mg/kg based on the maximum residue level for sugar cane and the estimated processing factor (0.015 mg/kg×Pf, 7).

Animal commodity maximum residue levels

The Meeting noted that of the commodities considered at the current Meeting, only sugar cane molasses is a relevant animal feed item. However, as the inclusion of sugar cane molasses does not significantly increase the animal dietary burden, the Meeting confirmed its previous recommendations for animal commodities.

RECOMMENDATIONS

On the basis of the data obtained from supervised trials, the Meeting concluded that residue levels listed in Annex 1 are suitable for establishing maximum residue limits and for IEDI and IESTI assessments.

Definition of the residue for compliance with the MRL for plant and animal commodities:
methoxyfenozide

Definition of the residue for dietary risk assessment for plant and animal commodities:
methoxyfenozide

The residue is fat-soluble in its distribution between meat muscle and fat, but not in its distribution in milk.

DIETARY RISK ASSESSMENT

Long-term dietary exposure

The ADI for methoxyfenozide is 0–0.1 mg/kg bw. The International Estimated Daily Intakes (IEDIs) for methoxyfenozide were estimated for the 17 GEMS/Food Consumption Cluster Diets using the STMR or STMR-P values estimated by the JMPR. The results are shown in Annex 3 of the 2021 Extra JMPR Report.

The IEDIs ranged from 1–7% of the maximum ADI. The Meeting concluded that long-term dietary exposure to residues of methoxyfenozide from uses considered by the JMPR is unlikely to present a public health concern.

Acute dietary exposure

The ARfD for methoxyfenozide is 0.9 mg/kg bw. The International Estimate of Short Term Intakes (IESTIs) for methoxyfenozide were calculated for the food commodities for which HRs/HR-Ps or STMRs/STMR-Ps were estimated by the present Meeting and for which consumption data were available. The results are shown in Annex 4 of the 2021 Extra JMPR Report.

The IESTIs varied from 0–4% of the ARfD for children and 0–5% of the ARfD for the general population. The Meeting concluded that acute dietary exposure to residues of methoxyfenozide from uses considered by the present Meeting is unlikely to present a public health concern.

5.21. PROTHIOCONAZOLE (232)

RESIDUE AND ANALYTICAL ASPECTS

Prothioconazole is a broad-spectrum systemic fungicide. It was first evaluated by the JMPR in 2008 (T, R). The latest residue review was done in 2017 (R). It was scheduled at the Fifty-first Session of the CCPR for the evaluation of additional uses by the 2020 JMPR, which was postponed to the 2021 Extra JMPR.

The toxicological review established an ADI of 0–0.05 mg/kg bw and an ARfD of 0.8 mg/kg bw for women of child-bearing age for the parent prothioconazole; an ADI of 0–0.01 mg/kg bw and an ARfD of 0.01 mg/kg bw for women of child-bearing age; and 1 mg/kg bw for general population for prothioconazole-desthio. The residue definition for plant commodities for enforcement and dietary risk assessment is prothioconazole-desthio. The residue definition for animal commodities for enforcement is prothioconazole-desthio and for dietary risk assessment the sum of prothioconazole-desthio, prothioconazole-desthio-3-hydroxy, prothioconazole-desthio-4-hydroxy and their conjugates, expressed as prothioconazole-desthio. The residue covered by the residue definition for compliance is considered not fat-soluble.

For the current Meeting, data were provided on residues in oilseed rape, linseed (flax) and sunflower together with processing studies of rape and sunflower seed as well as currently registered use patterns.

Methods of analysis

The Meeting received additional validation information on analytical methods previously evaluated by the 2008, 2009 and 2014 JMPRs for prothioconazole and metabolites in oilseeds (rape, linseed, sunflower). The Meeting concluded that the presented methods were sufficiently validated and are suitable to measure prothioconazole, prothioconazole-desthio as well as the metabolites prothioconazole-desthio- α -hydroxy (α -OH), prothioconazole-desthio-3-hydroxy (3-OH), prothioconazole-desthio-4-hydroxy (4-OH), prothioconazole-desthio-5-hydroxy (5-OH) and prothioconazole-desthio-6-hydroxy (6-OH) as well as the triazole metabolites 1,2,4-triazole (1,2,4-T), triazole acetic acid (TAA), triazole lactic acid (TLA) and triazole alanine (TA). The LOQs are 0.01 mg/kg individually for prothioconazole and metabolites in the oilseeds (rape, linseed, sunflower) as well as in processed commodities. The LOQs for the triazole metabolites 1,2,4-triazole (1,2,4-T), triazole acetic acid (TAA), triazole lactic acid (TLA) and triazole alanine (TA) in the same commodities were also 0.01 mg/kg for each compound. One method measured prothioconazole, prothioconazole-desthio and prothioconazole sulfonic acid in sunflower seed and processed commodities with LOQs of 0.01 mg/kg for each compound.

Stability of pesticide residues in stored analytical samples

A storage stability study was provided to cover the short-term increase in temperature that occurred during shipment for rape samples in one trial prior to frozen storage. Residues of prothioconazole were stable in rape seed and rape green material when stored at 5 °C for five days. The Meeting considered the short-term increase in temperature of samples during shipment in one trial to have no impact on the residues measured.

The demonstrated stability intervals for prothioconazole and metabolites in frozen storage of at least 36 months for rape seed and 24 months for rape pods and plant material reported by the 2008 and 2014 JMPR encompass the duration of storage in the residue trials evaluated. An additional storage stability study for prothioconazole, prothioconazole-desthio and prothioconazole sulfonic acid on sunflower seed demonstrated stability for at least 521 days.

The periods of demonstrated stability adequately cover the frozen storage intervals of samples in supervised trials considered by the current Meeting.

Results of supervised residue trials on crops

Supervised trials were available for the use of prothioconazole on a number of oilseed crops; rape, flax (linseed) and sunflower.

Rape

The critical GAP for prothioconazole on rape in Hungary is two foliar applications at 125 g ai/ha at a minimum interval of 14 days and a PHI of 28 days.

In thirty-two trials approximating GAP residues were (n = 32): < 0.01, 0.01, 0.01, 0.02, 0.02, 0.02, 0.025, 0.026, 0.03, 0.03, 0.03, 0.03, 0.03, 0.03, 0.033, 0.038, 0.04, 0.04, 0.043, 0.05, 0.05, 0.054, 0.055, 0.056, 0.06, 0.064, 0.081, 0.083, 0.09, 0.11, 0.13, 0.14 mg/kg.

The Meeting estimated a maximum residue level of 0.2 mg/kg and an STMR of 0.039 mg/kg for prothioconazole in rape seed.

Linseed (flax)

The Meeting received supervised residue trials on flax. In Greece, the cGAP is two foliar applications at 175 g ai/ha, with a minimum re-treatment interval of 14 days and a PHI of 56 days. No trials matching cGAP in Greece were made available to the Meeting. However, as rapeseed is the representative commodity for the subgroup “Small seed oilseeds”, the Meeting agreed to utilise trials on rapeseed matching the GAP for flax to estimate residues and to extrapolate the results to flax seed.

In four trials approximating cGAP in Greece, residues in rape seed were:

2 × 150 g ai/ha: < 0.01, < 0.01, < 0.01 and 0.02 mg/kg.

In a further seven trials at lower application rates residues were:

2 × 125 g ai/ha: < 0.01, < 0.01 and < 0.01 mg/kg.

2 × 100 g ai/ha: < 0.01, < 0.01, < 0.01 and < 0.01 mg/kg.

The Meeting agreed to use proportionality to scale the residues.

Rate (g ai/ha)	Residue (mg/kg)	Proportionality scaling factor	Scaled residues
2×100	< 0.01 (4)	1.75	< 0.0175 (4)
2×125	< 0.01 (3)	1.4	< 0.014 (3)
2×150	< 0.01 (3), 0.02	1.17	< 0.0117 (3), 0.0234

After scaling, residues approximating cGAP are (n = 11): < 0.0117 (3), < 0.014 (3), < 0.0175 (4) and 0.0234 mg/kg.

The Meeting estimated a maximum residue level of 0.03 mg/kg and an STMR of 0.014 mg/kg for prothioconazole in linseed.

Sunflower

The Meeting received supervised residue trials on sunflower. In Hungary, cGAP is two foliar applications at 125 g ai/ha at an interval of 14 days with a PHI of 28 days.

In eighteen trials on sunflower matching cGAP residues at harvest were (n = 18): < 0.01, < 0.01, < 0.01, < 0.01, < 0.01, < 0.01, < 0.01, < 0.01, 0.01, 0.01, 0.01, 0.017, 0.017, 0.02, 0.035, 0.043 and 0.13 mg/kg.

The critical GAP for prothioconazole on sunflower in Brazil is two foliar applications at 87.5 g ai/ha at a minimum interval of 15 days and a PHI of 30 days.

In ten trials from Brazil approximating GAP residues of prothioconazole-desthio were (n = 10): < 0.009, < 0.009, < 0.009, < 0.009, < 0.009, < 0.009, 0.009, 0.036, 0.063 and 0.32 mg/kg.

The dataset from Brazil leads to the higher maximum residue level and the Meeting decided to utilise this dataset to make a recommendation for a maximum residue level of 0.5 mg/kg and an STMR of 0.009 mg/kg for prothioconazole in sunflower seed.

Residues in animal feeds

Rape forage (green) = green material prior to BBCH 60

Rape forage corresponds to green material prior to BBCH 60. No samples of plant material were analysed at growth stages before BBCH 60. The data are not suitable to estimate residues in rape forage.

Fate of residues during processing

The Meeting received information on the fate of prothioconazole residues during processing of rape and sunflower seed. Residues of prothioconazole metabolites were measured using method 00979 (desthio, alpha-OH, 3-OH, 4-OH, 5-OH, 6-OH). As prothioconazole residues may be converted to prothioconazole-desthio, the processing factor for prothioconazole-desthio is calculated as (prothioconazole-desthio in the processed commodity)/(sum of prothioconazole and prothioconazole-desthio in the raw commodity). To calculate processing factors, residues of prothioconazole should be measured in the raw commodity or evidence is required that prothioconazole residues are not detected. In previous trials on rape seed assessed by the 2008 and 2014 JMPR, prothioconazole was converted to a mixture of prothioconazole sulfonic acid and prothioconazole-desthio with residues reported for both the sulfonic acid and prothioconazole-desthio and for the sum of prothioconazole sulfonic acid and prothioconazole-desthio which is referred to as “total prothioconazole”. In these trials it is not possible to determine whether prothioconazole is present at levels above the LOD and it is therefore not appropriate to use the processing studies provided to derive processing factors. However, the processing factors reported by the 2008 JMPR for rape seed for total prothioconazole are relevant as the sum of prothioconazole and prothioconazole-desthio was measured and are < 0.7 for both rape seed meal and refined oil.

Table 1 Processing factors and median residue values for prothioconazole-desthio used for estimation of maximum residue levels.

Processed commodity	Raw commodity [STMR/ median residue]	Processing factor	STMR-P (mg/kg)	Median residue-P (mg/kg)
Rape meal	0.039	< 0.7	0.0273	0.04641
Rape oil – refined		< 0.7	0.0273	
Sunflower meal	0.009	0.57	0.00513	0.008721
Sunflower crude oil		1.07	0.00963	

STMR-P = for plant commodity risk assessment residue definition – prothioconazole-desthio

Median residue-P = for compounds relevant for the livestock dietary burden for the livestock residue definition for risk assessment

Conversion factor for median residue for use in livestock burden calculations = 1.7

The Meeting applied the rape seed refined oil processing factor (0.7) to the estimated maximum residue level for rape seed (0.2 mg/kg) to estimate a maximum residue level of 0.15 mg/kg for rape seed oil, edible.

The Meeting considered the sunflower crude oil processing factor (1.07) and the estimated maximum residue level for sunflower seed (0.5 mg/kg) to estimate a maximum residue level of 0.5 mg/kg for sunflower seed oil, edible.

Residues in animal commodities

Farm animal dietary burden

The residue definition for risk assessment for livestock commodities differs from that for compliance for plant and livestock commodities. To estimate inputs for dietary intake calculations it is necessary to account for the transfer of compounds relevant to the livestock commodity residue definition from plants to livestock tissues. Metabolism data reported in the 2008 JMPR were used to estimate conversion factors to convert prothioconazole-desthio residues to those that may be converted on feeding to livestock to components in the risk assessment residue definition (prothioconazole, prothioconazole sulfonic acid M02, prothioconazole-desthio-3-hydroxy M14 and prothioconazole-desthio-4-hydroxy M15 and their conjugates M21, M22, M54, M55). As foliar application leads to the highest residues, in considering conversion factors, the average of values available following foliar application were used, i.e. data for seed treatment were not considered relevant. Conversion factors for wheat were extrapolated to all forage and fodders except peanut hay while factors for sugar beet tops and sugar beet roots were extrapolated to potatoes.

The following conversion factors were obtained:

Cereal forage	1.6
Cereal hay	2.1
Cereal straw	2.6
Grains/seeds	1.7
Peanut hay	2.4
Sugar beet tops	1.6
Sugar beet roots/potato tubers	1.0

The feed commodities listed in the Table below were used in estimating livestock dietary burdens.

Some processed and forage commodities do not appear in the Recommendations Table (because no maximum residue level is needed), but they are used in estimating livestock dietary burdens. Those commodities are included in the list below.

Table 2 Potential livestock feed items and residues of compounds in feed relevant to livestock risk assessment

Codex classification	Prothioconazole-desthio		Conversion factor ^A	Residues relevant to livestock risk assessment dietary burden	
	Median residue (mg/kg)	Highest residue (mg/kg)		Median residue (mg/kg)	Highest residue (mg/kg)
Barley grain	0.035		1.7	0.0595	
Barley forage	1.2	5.4	1.6	1.92	8.64
Cotton seed	0.052		1.7	0.0884	
Cotton gin by-products	1.1	1.8	2.6	2.86	4.68
Cotton seed meal	0.005		1.7	0.0085	
Cotton seed hulls	0.008		1.7	0.0136	
Fodder of cereal grains (dry)	1.5	4.8	2.1	3.15	10.08
Linseed/flax seed	0.014		1.7	0.0238	
Maize	0.018		1.7	0.0306	
Maize forage/silage	2.15	4.08	1.6	3.44	6.528
Maize fodder	3.48	6.7	2.1	7.308	14.07
Oat grain	0.01		1.7	0.017	
Peanut fodder (hay)	4.08	11.6	2.4	9.792	27.84
Potato culls	0.01		1	0.01	

Codex classification	Prothioconazole-desthio		Conversion factor ^A	Residues relevant to livestock risk assessment dietary burden	
	Median residue (mg/kg)	Highest residue (mg/kg)		Median residue (mg/kg)	Highest residue (mg/kg)
Pulses (grain)	0.05		1.7	0.085	
Rape seed	0.039		1.7	0.0663	
Rape seed meal	0.0273		1.7	0.04641	
Rye grain	0.01		1.7	0.017	
Soya bean (dry)	0.05		1.7	0.085	
Soya bean aspirated grain fractions	3.75		1.7	6.375	
Soya bean hulls	0.025		1.7	0.0425	
Straw and fodder (dry) of cereals	0.65	1.9	2.6	1.69	4.94
Sugar beet	0.05		1	0.05	
Sugar beet tops	1.5	3.9	1.6	2.4	6.24
Sweet corn fodder	3.48	6.7	2.1	7.308	14.07
Triticale grain	0.01		1.7	0.017	
Wheat grain	0.02		1.7	0.034	
Wheat bran	0.048		1.7	0.0816	
Wheat milled by-products	5		1.7	8.5	

^a While the Meeting noted conversion factors for processed commodities might be different to those of the RAC, the Meeting agreed, in the absence of better information, to also use the conversion factors derived for the RAC for livestock feed commodities derived from processing.

Dietary burdens were calculated for beef cattle, dairy cattle, broilers and laying poultry based on feed items evaluated by the JMPR by the current Meeting. The dietary burdens were estimated using the most recent version of the OECD livestock dietary burden calculator, are presented in Annex 6 and summarised below.

Table 3 Estimated maximum and mean dietary burdens of farm animals (prothioconazole-desthio)

	Animal dietary burden: prothioconazole-desthio, ppm of dry matter diet							
	US-Canada		EU		Australia		Japan	
	max	mean	Max	mean	max	mean	max	mean
Beef cattle	4.1	3.4	13.9	5.7	15.8	5.5	3.1	3.1
Dairy cattle	10.0	4.8	14.6	5.8	15.8a b	5.5	7.7	5.2
Broilers	2.9	2.9	1.2	1.2	1.2	1.2	0.3	0.3
Layers	2.9c d	2.9	2.1	1.5	1.2	1.2	1.7	1.7

^a Highest maximum beef or dairy cattle dietary burden suitable for MRL estimates for mammalian tissues

^b Highest maximum dairy cattle dietary burden suitable for MRL estimates for mammalian milk

^c Highest maximum broiler or laying hen dietary burden suitable for MRL estimates for poultry tissues

^d Highest maximum laying hen dietary burden suitable for MRL estimates for eggs.

Table 4 Estimated maximum and mean dietary burdens of farm animals (residues relevant to estimation of livestock residues according to the risk assessment definition)

	Animal dietary burden: relevant to risk assessment, ppm of dry matter diet							
	US-Canada		EU		Australia		Japan	
	max	mean	Max	mean	max	mean	Max	mean
Beef cattle	7.1	5.7	22.2	9.3	31.2	10.8	5.3	5.3
Dairy cattle	17.8	9.4	23.3	9.5	31.2 ^a	10.8 ^b	12.5	8.7
Broilers	4.9	4.9	2.0	2.0	2.0	2.0	0.5	0.5
Layers	4.9 ^c	4.9 ^d	3.6	2.6	2.0	2.0	2.9	2.9

^a Highest maximum beef or dairy cattle dietary burden suitable for HR estimates for mammalian tissues and milk

^b Highest mean beef or dairy cattle dietary burden suitable for STMR estimates for mammalian tissues and milk.

^c Highest maximum broiler or laying hen dietary burden suitable for HR estimates for poultry tissues and eggs

^d Highest mean broiler or laying hen dietary burden suitable for STMR estimates for poultry tissues and eggs.

Animal commodity maximum residue levels

Cattle

The calculations used to estimate highest total residues for use in estimating maximum residue levels, STMR and HR values are shown below.

Table 5 Anticipated residues of prothioconazole-desthio in mammalian commodities

	Feed Level (ppm) for milk residues	Prothioconazole-desthio residues (mg eq/kg) in milk	Feed Level (ppm) for tissue residues	Prothioconazole-desthio residues (mg eq/kg)			
				Muscle	Liver	Kidney	Fat
Highest residue for maximum residue level estimation (beef or dairy cattle)							
Feeding Study	29 ^a	< 0.004	5.1 29 ^a	< 0.01	0.03 0.18	0.01 0.03	0.01
Dietary burden and estimate of highest residue	15.8	< 0.0022	15.8	< 0.005	0.129	0.023	0.0054

^a The nominal feeding levels reported by the 2008 JMPR were 4, 25 and 100 ppm, however, the actual feed levels were 5.1, 29 and 125 ppm. The actual feed levels are used here.

	Feed Level (ppm) for milk residues	Total residues (mg eq/kg) in milk	Feed Level (ppm) for tissue residues	Total residues ^A (mg eq/kg)			
				Muscle	Liver	Kidney	Fat
HR Determination (beef or dairy cattle)							
Feeding Study			29 ^b 125 ^b	0.03	0.26 1.6	0.17 1.1	0.02 0.14
Dietary burden and estimate of highest residue			31.2	0.007	0.291	0.191	0.023
STMR Determination (beef or dairy cattle)							
Feeding Study	125 ^{b c}	0.011	5.5 ^b 29 ^b	< 0.01	0.04 0.22	0.023 0.14	0.013
Dietary burden and estimate of highest residue	10.8	0.001	10.8	< 0.004	0.077	0.049	0.005

No residues detected above LOQ for the 5.1 and 29 ppm dose groups, mean residues were 0.011 mg/kg for the 125 ppm dose group.

^a Total residues = sum of prothioconazole, prothioconazole-desthio, prothioconazole-desthio-3-hydroxy, prothioconazole-desthio-4-hydroxy and their conjugates expressed as prothioconazole-desthio.

^b The nominal feeding levels reported by the 2008 JMPR were 4, 25 and 100 ppm, however, the actual feed levels were 5.1, 29 and 125 ppm. The actual feed levels are used here.

^c Residues were <LOQ in milk at the 5.1 and 29 ppm feed levels. As residues in milk were detected at the 125 ppm feed level, the Meeting decided to utilise the results from the 125 ppm feed level to estimate residues in milk.

The current Meeting has estimated livestock dietary burdens based on the different residue definitions for compliance and risk assessment, including accounting for the different compounds that need to be included in the estimation of the livestock dietary burdens. Additionally, the OECD

livestock dietary burden calculator has been updated several times since 2009. The Meeting has taken the opportunity to make revised recommendations with the newly developed livestock dietary burdens and most recent version of the OECD livestock dietary burden calculator.

The Meeting estimated the following maximum residue levels: mammalian fat (except milk fat) 0.01 mg/kg and edible offal 0.15 mg/kg to replace its previous recommendations and confirmed its previous estimations for milk (0.004* mg/kg) and meat (mammalian except marine mammals) (0.01 mg/kg). The Meeting estimated the following HRs: muscle 0.007 mg/kg; fat 0.023 mg/kg; liver 0.291 mg/kg and kidney 0.191 mg/kg and STMRs: muscle 0.004 mg/kg; fat 0.005 mg/kg; liver 0.077 mg/kg, kidney 0.049 mg/kg and milk 0.001 mg/kg.

For poultry, the Meeting noted that the 2008 JMPR had concluded that the poultry feeding study designs did not reflect the residue composition in feed and that the results could not be used for estimating maximum residue limits or STMR values. The 2017 JMPR noted in a metabolism study on laying hens with the parent prothioconazole conducted at an exaggerated rate of 171 ppm feed and slaughter at 5 hours after the last dose, residues of prothioconazole-desthio were 0.167, 0.007, 0.006 and 0.13 mg eq/kg in liver, eggs, muscle and fat, respectively. Total radioactive residues (TRR) were 4.017, 0.036, 0.089 and 0.45 mg/kg for liver, eggs, muscle and fat, respectively. Assuming residues are proportional to the level in feed, residues of prothioconazole-desthio following exposure to feed at 2.9 ppm would be 0.0028, 0.0001, 0.0001 and 0.0022 mg/kg in liver, eggs, muscle and fat, below the LOQs of 0.005 mg/kg for eggs and 0.01 mg/kg for tissues. Total residues on feeding at 4.9 ppm would be 0.118, 0.001, 0.0026 and 0.0129 mg/kg residue in liver, eggs, muscle and fat, respectively.

The Meeting estimated the following maximum residue levels for poultry commodities: poultry fat 0.01 mg/kg and poultry edible offal 0.01 mg/kg to replace its previous recommendations and confirmed its previous estimations for eggs (0.005(*) mg/kg) and poultry meat (0.01(*) mg/kg). The Meeting estimated the following HRs: poultry meat 0.0026 mg/kg; poultry fat 0.0129 mg/kg; liver 0.118 mg/kg and eggs 0.001 mg/kg and STMRs at the same levels.

RECOMMENDATIONS

On the basis of the data obtained from supervised trials, the Meeting concluded that the residue levels listed in Annex 1 are suitable for establishing maximum residue limits and for IEDI and IESTI assessments.

Definition of the residue for compliance with the MRL and dietary risk assessment for plant commodities: *prothioconazole-desthio*.

Definition of the residue for compliance with the MRL for animal commodities: *prothioconazole-desthio*; and

Definition of the residue for dietary risk assessment for animal commodities: the *sum of prothioconazole-desthio, prothioconazole-desthio-3-hydroxy, prothioconazole-desthio-4-hydroxy and their conjugates expressed as prothioconazole-desthio*.

The residue is not fat-soluble.

DIETARY RISK ASSESSMENT

Long-term dietary exposure

The ADI for prothioconazole-desthio is 0–0.01 mg/kg bw. The International Estimated Daily Intakes (IEDIs) for prothioconazole-desthio were estimated for the 17 GEMS/Food Consumption Cluster Diets using the STMR or STMR-P values estimated by the JMPR. The results are shown in Annex 3 of the 2021 Extra JMPR Report.

The IEDIs ranged from 1–3% of the maximum ADI. The Meeting concluded that long-term dietary exposure to residues of prothioconazole from uses considered by the JMPR is unlikely to present a public health concern.

Acute dietary exposure

The ARfD for prothioconazole-desthio is 0.01 mg/kg bw for women of child-bearing age and 1 mg/kg bw for the general population. The International Estimate of Short-Term Intakes (IESTIs) for prothioconazole-desthio were calculated for the food commodities and their processed commodities for which HRs/HR-Ps or STMRs/STMR-Ps were estimated by the present Meeting and for which consumption data were available. The results are shown in Annex 4 of the 2021 Extra JMPR Report.

The IESTIs varied from 0–8% of the ARfD for women of child-bearing age and 0% of the ARfD for children and the general population. The Meeting concluded that acute dietary exposure to residues of prothioconazole from uses considered by the present Meeting is unlikely to present a public health concern.

5.22. PYDIFLUMETOFEN (309)

RESIDUE AND ANALYTICAL ASPECTS

Pydiflumetofen is a broad-spectrum fungicide belonging to the carboxamide group. It acts through inhibition of succinate dehydrogenase in complex II of fungal mitochondrial respiration.

Pydiflumetofen was first evaluated for toxicology and residues by the JMPR in 2018. An ADI of 0–0.1 mg/kg bw and an ARfD of 0.3 mg/kg bw were established. The residue definition for compliance with the MRL for plant and animal commodities and dietary risk assessment for plant commodities is pydiflumetofen. The residue definition for dietary risk assessment for animal commodities other than mammalian liver and kidney is the sum of pydiflumetofen and 2,4,6-TCP (2,4,6-Trichlorophenol) and its conjugates, expressed as pydiflumetofen and for dietary risk assessment for mammalian liver and kidney is the sum of pydiflumetofen, 2,4,6-trichlorophenol (2, 4, 6-TCP) and its conjugates and SYN547897 and its conjugates, expressed as pydiflumetofen. The residue is fat-soluble.

The 2018 JMPR noted that pydiflumetofen residues are very persistent in soil (up to 2380 days DT₅₀) and may be taken up by rotational crops.

At the Fifty-first Session of the CCPR, pydiflumetofen was scheduled for evaluation by the 2020 JMPR for several new uses, which was postponed to the 2021 Extra JMPR.

The Meeting received information from the manufacturer on use patterns, supervised residue trials on lemons, oranges, grapefruits, apples, pears, peaches, cherries, plums, blueberries, strawberries, bulb onions, green onions, cabbage, cauliflower, broccoli, beans with pods, peas with pods, beans without pods, peas without pods, sugar beet, carrots, radish, sorghum, almonds, pecans, sunflower and cottonseed and processing studies.

Methods of analysis

The LC-MS/MS analytical method (GRM061.03A) used for analysis of residues of pydiflumetofen in plant commodities, with n LOQ of 0.01 mg/kg, was reviewed by the 2018 JMPR. All samples collected from the supervised residue trials provided to the current Meeting were analysed using the same method. The methods are fit for purpose based on the recoveries from the concurrent method validation.

Stability of pesticide residues in stored analytical samples

The stability of residues of pydiflumetofen during frozen storage was evaluated by the 2018 JMPR. Pydiflumetofen was determined to be stable when stored frozen for at least 23 months at -20 °C in commodities representative of the high water, high acid, high starch, high protein and high oil commodity groups. The periods of demonstrated stability adequately covered the frozen storage intervals of the samples in the supervised residue trials on crops considered by the current Meeting.

Results of supervised residue trials on crops

For maximum residue level estimation of pydiflumetofen residues in primary crop commodities, the addition of residues arising from direct treatment in combination with root uptake of pydiflumetofen from previous years, as determined at the 2019 Meeting, must be taken into account. The crop groups for plant food and feed established in the Codex Classification of Foods and Animal Feeds were used to give recommendations on the overall residue levels of pydiflumetofen expected in these commodities. As the seasonal application rates for the primary crop uses considered by the current Meeting are equivalent to or less than those considered at the 2019 Meeting, new scaling of the rotational crop residues is not required.

The Meeting noted that the use of statistical methods for the estimation of maximum residue levels is not possible when considering potential carryover of residues in succeeding crops.

The Meeting recognised that for permanent crops and crops cultivated in/on culture soil/medium and water, the contribution from carryover is not needed, as those crops are not expected to be subject to significant uptake of pydiflumetofen from the soil.

Citrus Fruits

The critical GAP for citrus fruits is from the USA; 4×85 g ai/ha, 7-day RTI, 0-day PHI. The Meeting received supervised residue trials conducted on lemon, mandarin, orange and grapefruit in the USA matching the critical GAP.

Pydiflumetofen residues in whole lemons in ranked order were (n = 6): 0.02, 0.14, 0.25, 0.29, 0.38 and 0.46 mg/kg.

Pydiflumetofen residues in whole mandarins in ranked order were (n = 4): 0.17, 0.24, 0.28 and 0.56 mg/kg.

Pydiflumetofen residues in whole oranges in ranked order were (n = 10): 0.11, 0.16, 0.17, 0.18, 0.21, 0.23, 0.31, 0.36, 0.40 and 0.68 mg/kg.

Pydiflumetofen residues in whole grapefruits in ranked order were (n = 7): 0.09, 0.12 (2), 0.13, 0.15, 0.16 and 0.58 mg/kg.

The Meeting noted that the GAP covers the group of citrus fruits and that median residues of lemons, mandarins, oranges and grapefruits are within a 5-fold difference. The Kruskal-Wallis H-test also determined that the datasets were from the same population. Therefore, the Meeting decided to combine the four datasets of lemons, mandarins, oranges and grapefruits.

Combined pydiflumetofen residues in lemons, mandarins, oranges and grapefruits were (n = 27): 0.02, 0.09, 0.11, 0.12 (2), 0.13, 0.14, 0.15, 0.16 (2), 0.17 (2), 0.18, 0.21, 0.23, 0.24, 0.25, 0.28, 0.29, 0.31, 0.36, 0.38, 0.40, 0.46, 0.56, 0.58 and 0.68 mg/kg (highest single value of 0.76 mg/kg).

Citrus fruits are cultivated as permanent crops and are not expected to be subject to significant uptake of pydiflumetofen from the soil. The Meeting estimated a maximum residue level of 0.9 mg/kg for the Group of citrus fruits.

For dietary risk assessment, the Meeting calculated a ratio of residues in flesh (0.042 mg/kg) to whole orange (0.179 mg/kg) and applied this ratio of 0.24 to all samples of citrus fruits, where combined pydiflumetofen residues were (n = 27): 0.01, 0.02, 0.03 (5), 0.04 (6), 0.06 (4), 0.07 (3), 0.09 (2), 0.10, 0.11, 0.13, 0.14 and 0.16 mg/kg, from which an HR of 0.16 and STMR of 0.05 mg/kg were estimated for citrus flesh. The Meeting also estimated an HR of 0.76 mg/kg and an STMR of 0.21 mg/kg for kumquats. An HR of 0.76 mg/kg and a median residue of 0.21 mg/kg were estimated for whole citrus fruits and used for estimation of HR-Ps and STMR-Ps for processed commodities.

Pome Fruits

The critical GAP for pome fruits is from the USA; 4×50 g ai/ha, 7-day RTI, 30-day PHI. The Meeting received supervised residue trials conducted on apple and pear in Canada and the USA matching the critical GAP.

Pydiflumetofen residues in apples in ranked order were (n = 14): 0.05 (2), 0.06 (5), 0.07, 0.08 (2), 0.09, 0.10 (2) and 0.11 mg/kg (highest value of 0.13 mg/kg from replicate samples).

Pydiflumetofen residues in pears in ranked order were (n = 11): 0.02 (3), 0.04, 0.05 (3), 0.07, 0.08, 0.09 and 0.12 mg/kg (highest value of 0.13 mg/kg from replicate samples).

The Meeting noted that the GAP covers the group of pome fruits except persimmon, Japanese and that median residues in apples and pears are within a 5-fold difference. The Mann-Whitney U-test also determined that the datasets were from the same population. Therefore, the Meeting decided to combine the two datasets of apples and pears.

Combined pydiflumetofen residues in apples and pears were (n = 25): 0.02 (3), 0.04, 0.05 (5), 0.06 (5), 0.07 (2), 0.08 (3), 0.09 (2), 0.10 (2), 0.11 and 0.12 mg/kg.

Pome fruits are cultivated as permanent crops and are not expected to be subject to a potential uptake of pydiflumetofen from the soil. The Meeting estimated a maximum residue level of 0.2 mg/kg, an HR of 0.13 mg/kg and an STMR of 0.06 mg/kg for the Group of pome fruits, except persimmon, Japanese.

Stone fruits

The critical GAP for stone fruits is from the USA; 4×75 g ai/ha, 7-day RTI, 0-day PHI. The Meeting received supervised residue trials on peaches, cherries and plums conducted in Canada and the USA matching the critical GAP.

Peaches

Pydiflumetofen residues in peaches in ranked order were (n = 12): 0.09, 0.16, 0.19, 0.20 (2), 0.21 (2), 0.22 (2), 0.25, 0.29 and 0.78 mg/kg (highest value of 0.80 mg/kg from replicate samples).

Residue levels in the field trials from Canada and the USA were reported as fruit without stone. At the 2017 Meeting, it was concluded that the weight ratio of flesh to whole fruit for peaches ranged between 0.85 and 0.96 and that correcting the residue levels for this weight/weight ratio would lead to the same maximum residue level.

Peaches are cultivated as permanent crops and are not expected to be subject to a potential uptake of pydiflumetofen from the soil. The Meeting estimated a maximum residue level of 1 mg/kg, an HR of 0.80 mg/kg and an STMR value of 0.21 mg/kg for the Subgroup of peaches.

Cherries

Pydiflumetofen residues in cherries in ranked order were (n = 14): 0.14, 0.18, 0.18, 0.21, 0.24, 0.26, 0.37, 0.42, 0.44, 0.53, 0.61, 0.72, 0.90 and 1.6 mg/kg (highest value of 1.7 mg/kg from replicate samples).

Residue levels in the field trials from Canada and the USA were reported as fruit without stone. At the 2017 Meeting, it was concluded that, based on the cherry trials, the contribution of the pit to the weight of the whole fruit is approximately 10%. Correcting the residue levels using this weight/weight ratio would lead to the same maximum residue level.

Cherries are cultivated as permanent crops and are not expected to be subject to a potential uptake of pydiflumetofen from the soil. The Meeting estimated a maximum residue level of 2.0 mg/kg, an HR of 1.7 mg/kg and an STMR of 0.395 mg/kg for the Subgroup of cherries.

Plums

Pydiflumetofen residues in plums in ranked order were (n = 9): 0.06, 0.11, 0.12, 0.13, 0.15, 0.16, 0.21, 0.32 and 0.35 mg/kg (highest value of 0.37 mg/kg from replicate samples).

Residue levels in the field trials from Canada and the USA were reported as fruit without stone. At the 2017 Meeting, it was concluded that, the weight ratio of flesh to whole fruit for plums ranged between 0.86 and 0.97 and that correcting the residue levels for this weight/weight ratio would lead to the same maximum residue level.

Plums are cultivated as permanent crops and are not expected to be subject to a potential uptake of pydiflumetofen from the soil. The Meeting estimated a maximum residue level of 0.6 mg/kg, an HR of 0.37 mg/kg and an STMR value of 0.15 mg/kg for the Subgroup of plums.

Bushberries

The critical GAP for the bushberry crop subgroup is from the USA; 2×150 g ai/ha, 7-day RTI, 0-day PHI. The Meeting received supervised residue trials on highbush blueberries conducted in Canada and the USA and approximating the critical GAP.

Pydiflumetofen residues in highbush blueberries in ranked order were (n = 10): 0.40, 0.66 (2), 0.69, 0.86, 0.90, 0.96, 1.4, 1.9 and 3.6 mg/kg (highest value of 3.9 mg/kg from replicate samples).

Bushberries are cultivated as permanent crops and are not expected to be subject to a potential uptake of pydiflumetofen from the soil. Noting that blueberries is the representative crop of the subgroup bush berries, the Meeting estimated a maximum residue level of 5 mg/kg, an HR of 3.9 mg/kg and an STMR value of 0.88 mg/kg for the Subgroup of bush berries and extrapolated these values to elderberries.

Low growing berries

In field studies on succeeding crops, reviewed by the 2019 Meeting, the scaled highest residue in fruits (strawberries) was < 0.02 mg/kg.

The current Meeting received supervised residue trials conducted in Canada and the USA on strawberries matching the critical GAP from the USA, for the low growing berry, except cranberry, subgroup; 2×150 g ai/ha, 7-day RTI, 0-day PHI.

Pydiflumetofen residues in strawberries in ranked order were (n = 10): 0.08, 0.09, 0.10, 0.16, 0.18, 0.19, 0.30, 0.44, 0.46 and 0.56 mg/kg (highest value of 0.62 mg/kg from replicate samples).

The Meeting concluded that residues from uptake of pydiflumetofen via the roots are insignificant in comparison to residue levels following direct treatment. The Meeting estimated a maximum residue level of 1 mg/kg, an HR of 0.62 mg/kg and an STMR of 0.185 mg/kg for the Subgroup of low growing berries, except cranberries.

Bulb vegetables

The current Meeting received supervised residue trials conducted in the USA on bulb vegetables matching the critical GAP from the USA for bulb vegetables; 3×125 g ai/ha, 7-day RTI, 7-day PHI.

Bulb onions

Pydiflumetofen residues in bulb onions in ranked order were (n = 8): < 0.01 (2), 0.01, 0.05 (2), 0.06 (2) and 0.12 mg/kg (highest value of 0.13 mg/kg from replicate samples).

Bulb onions may be subject to crop rotation, however, no residue data on suitable representative rotational crops of bulb or stem vegetables were provided to the 2019 Meeting. Based on the consistent range of residues across rotational crop commodities and that the edible portion of bulb onion is below ground, the Meeting agreed to use residues from a root crop to estimate the uptake of pydiflumetofen via the roots. Therefore, the Meeting decided to extrapolate the mean, median and highest residues in the succeeding crop of radish roots of 0.02, 0.02 and 0.07 mg/kg, respectively, to bulb onions.

The Meeting estimated a maximum residue level of 0.3 mg/kg, an HR of 0.20 mg/kg (scaled highest residue found in succeeding crop (radish root) of 0.07 mg/kg added to the highest residue in bulb onions of 0.13 mg/kg obtained from supervised field trials) and an STMR of 0.07 mg/kg (scaled mean residue found in succeeding crop (radish root) of 0.02 mg/kg added to the median residue in bulb onion of 0.05 mg/kg obtained from supervised field trials) for the Subgroup of bulb onions.

Green onions

Pydiflumetofen residues in green onions in ranked order were (n = 4): 0.28, 0.30, 0.36 and 1.1 mg/kg (highest value of 1.3 mg/kg from replicate samples).

Green onions may be subject to crop rotation, however, no residue data on suitable representative rotational crops of bulb or stem vegetables were provided to the 2019 Meeting. Based on the consistent range of residues across rotational crop commodities and that the edible portion of green onion includes the aerial portion of the plant, the Meeting agreed to use residues from a leafy crop to estimate the uptake of pydiflumetofen via the roots. Therefore, the Meeting decided to extrapolate the mean, median and highest residues in the succeeding crop of mature spinach of 0.03, 0.02 and 0.09 mg/kg, respectively, to green onions.

The Meeting estimated a maximum residue level of 1.5 mg/kg, an HR of 1.39 mg/kg (scaled highest residue found in succeeding crop (mature spinach) of 0.09 mg/kg added to the highest residue in green onions of 1.3 mg/kg obtained from supervised field trials) and an STMR of 0.36 mg/kg (scaled mean residue found in succeeding crop (mature spinach) of 0.03 mg/kg added to the median residue in bulb onion of 0.33 mg/kg obtained from supervised field trials) for the Subgroup of green onions.

Brassica vegetables

In field studies on succeeding crops, reviewed by the 2019 Meeting, the scaled mean, median and highest residues in leafy vegetables and Brassicas (mature spinach) were 0.03, 0.02 and 0.09 mg/kg, respectively. Using this data, the Meeting recommended a maximum residue level of 0.1 mg/kg for the Group of Brassica vegetables (except Brassica leafy vegetables).

The current Meeting received supervised residue trials conducted in the USA on Brassica vegetables matching the critical GAP for Brassica vegetables from the USA; 3×125 g ai/ha, 7-day RTI, 0-day PHI.

Flowerhead Brassicas

Pydiflumetofen residues in broccoli and cauliflower in ranked order were (n = 8): 0.04, 0.05, 0.31, 0.36, 0.42, 0.67, 0.94 and 1.4 mg/kg (highest value of 1.5 mg/kg from replicate samples).

The Meeting estimated a maximum residue level of 3.0 mg/kg, an HR of 1.5 mg/kg and an STMR of 0.39 mg/kg for the Subgroup of flowerhead Brassicas.

Head Brassicas

Pydiflumetofen residues in cabbage with wrapper leaves in ranked order were (n = 6): 0.30, 0.33, 0.36, 0.88, 0.95 and 0.99 mg/kg.

Pydiflumetofen residues in cabbage without wrapper leaves in ranked order were (n = 6): < 0.01, 0.02, 0.03, 0.04, 0.05 and 0.12 mg/kg (highest value of 0.13 mg/kg from replicate samples).

The Meeting concluded that for estimating a maximum residue level and for calculating the animal dietary burden, residues from uptake of pydiflumetofen via the roots of head Brassicas are insignificant in comparison to residue levels following direct treatment. However, for dietary risk assessment, the Meeting agreed to take into account the residues from rotational crops since the contribution is more significant in comparison to residues resulting from direct treatment in cabbage without wrapper leaves.

The Meeting estimated a maximum residue level of 2.0 mg/kg, an HR of 0.22 mg/kg (scaled highest residue found in succeeding crop (spinach) 0.09 mg/kg added to the highest residue in cabbage without wrapper leaves of 0.13 mg/kg obtained from supervised field trials) and an STMR of 0.065 mg/kg (scaled mean residue found in succeeding crop (spinach) 0.03 mg/kg added to the median residue in cabbage without wrapper leaves of 0.035 mg/kg obtained from supervised field trials) for the Subgroup of head Brassicas.

The Meeting estimated a highest residue and a median residue for animal dietary burden calculation of 0.99 and 0.62 mg/kg, respectively, for cabbage leaves.

Stem Brassicas

Although pydiflumetofen is not registered for use on the Subgroup of stem Brassicas, these crops may still be subject to crop rotation and therefore contain pydiflumetofen residues after uptake via the roots. The Meeting estimated a maximum residue level of 0.1 mg/kg for the Subgroup of stem Brassicas, an HR of 0.09 mg/kg and an STMR of 0.02 mg/kg and withdrew its previous recommendation for the Group of Brassica vegetables (except Brassica leafy vegetables).

Legume vegetables

In field studies on succeeding crops, reviewed by the 2019 Meeting, the scaled mean, median and highest residues in beans fresh seeds were each < 0.02 mg/kg. Using this data, the Meeting recommended a maximum residue level of 0.02 mg/kg for the Group of legume vegetables.

The current Meeting received a critical GAP for legume vegetables from the USA; 2×200 g ai/ha, 14-day RTI, 14-day PHI. The legume vegetable field trials from Canada and the USA (2×200 g ai/ha, 7-day RTI, 14-day PHI) differed from the critical GAP with regard to the RTI. Comparison of the application scenarios of the trials with the critical GAP, showed that the expected residues are within 20% of the critical GAP. Therefore, the Meeting concluded that the supervised residue trials could be used for estimation of the maximum residue level.

Beans with pods

In trials approximating the critical GAP, pydiflumetofen residues in beans with pods in ranked order were (n = 10): 0.01 (2), 0.02 (2), 0.03, 0.06, 0.08, 0.14, 0.26 and 0.43 mg/kg (highest value of 0.47 mg/kg from replicate samples).

The Meeting concluded that residues from uptake of pydiflumetofen via the roots are insignificant in comparison to residue levels following direct treatment.

The Meeting estimated a maximum residue level of 0.7 mg/kg, an HR of 0.47 mg/kg and an STMR of 0.045 mg/kg for the Subgroup of beans with pods.

Peas with pods

In trials approximating the critical GAP, pydiflumetofen residues in peas with pods in ranked order were (n = 7): 0.01 (2), 0.05, 0.12, 0.16 and 0.64 (2) mg/kg (highest value of 0.84 mg/kg from replicate samples).

The Meeting concluded that residues from uptake of pydiflumetofen via the roots are insignificant in comparison to residue levels following direct treatment.

The Meeting estimated a maximum residue level of 1.5 mg/kg, an HR of 0.84 mg/kg and an STMR of 0.12 mg/kg for the Subgroup of peas with pods.

Succulent beans without pods

In trials approximating the critical GAP, pydiflumetofen residues in beans without pods in ranked order were (n = 9): 0.010 (2), 0.011, 0.013(2), 0.017, 0.018, 0.034 and 0.065 mg/kg (highest value of 0.079 mg/kg from replicate samples).

The Meeting concluded that residues in beans without pods may be influenced significantly by uptake of pydiflumetofen from the soil. For the estimation of a maximum residue level, HR and STMR, all scaled residues were < 0.02 mg/kg in beans fresh seeds from the succeeding crop field trials. Therefore, a value of 0.02 mg/kg was added to the median residue obtained from supervised field trials on beans without pods of 0.013 mg/kg for an overall STMR for beans without pods of 0.033 mg/kg. For the estimation of a maximum residue level, the value of 0.02 mg/kg was added to the highest residue of 0.079 mg/kg found in supervised field trials, resulting in an overall highest residue in succulent beans without pods of 0.099 mg/kg.

The Meeting estimated a maximum residue level of 0.15 mg/kg, an HR of 0.099 mg/kg and an STMR of 0.033 mg/kg for the Subgroup of succulent beans without pods.

Succulent peas without pods

In trials approximating the critical GAP, pydiflumetofen residues in peas without pods in ranked order were (n = 10): < 0.01 (5), 0.011, 0.012, 0.013 (2) and 0.018 mg/kg (highest value of 0.022 mg/kg from replicate samples).

The Meeting concluded that residues in peas without pods may be influenced significantly by uptake of pydiflumetofen from the soil. For the estimation of a maximum residue level, HR and STMR, all scaled residues were < 0.02 mg/kg in beans fresh seeds from the succeeding crop field trials. Therefore, a value of 0.02 mg/kg was added to the median residue obtained from supervised field trials on peas without pods of 0.011 mg/kg for an overall STMR for peas without pods of 0.031 mg/kg. For the estimation of a maximum residue level, the value of 0.02 mg/kg was added to the highest residue of 0.022 mg/kg found in supervised field trials, resulting in an overall highest residue in succulent peas without pods of 0.042 mg/kg.

The Meeting estimated a maximum residue level of 0.05 mg/kg, an HR of 0.042 mg/kg and an STMR of 0.031 mg/kg for the Subgroup of succulent peas without pods.

Underground immature beans and peas

Although pydiflumetofen is not registered for use on the Subgroup of underground immature beans and peas, these crops may still be subject to crop rotation and therefore contain pydiflumetofen residues after uptake via the roots. The Meeting estimated a maximum residue level of 0.02 mg/kg, an HR of 0.02 mg/kg and an STMR of 0.02 mg/kg for the Subgroup of underground immature beans and peas and withdrew its previous recommendation for the Group of legume vegetables.

Root vegetables

In field studies on succeeding crops, reviewed by the 2019 Meeting, the scaled mean, median and highest residues in root and tuber vegetables (radish) were 0.02, 0.02 and 0.07 mg/kg. Using this data, the previous Meeting estimated a maximum residue level of 0.1 mg/kg for the Subgroup of root vegetables.

The current Meeting received supervised residue trials conducted in Canada and the USA on root vegetables matching the critical GAP for root vegetables from the USA; 4×75 g ai/ha, 7-day RTI, 7-day PHI.

Pydiflumetofen residues in sugar beets in ranked order were (n = 17): 0.02 (2), 0.04 (2), 0.05, 0.06 (5), 0.08, 0.09, 0.10 (2), 0.12, 0.13 and 0.14 mg/kg (highest value of 0.15 mg/kg from replicate samples).

Pydiflumetofen residues in carrots in ranked order were (n = 6): 0.02, 0.03, 0.06, 0.07, 0.08 and 0.10 mg/kg (highest value of 0.11 mg/kg from replicate samples).

Pydiflumetofen residues in radish roots in ranked order were (n = 5): 0.01, 0.02, 0.14 and 0.17 (2) mg/kg (highest value of 0.18 mg/kg from replicate samples).

The Meeting noted that the GAP covers the Subgroup of root vegetables and that median residues of sugar beet roots, carrots and radish roots are within a 5-fold difference. The Kruskal Wallis H-test also determined that the datasets were from the same population. Therefore, the Meeting decided to combine the three datasets of sugar beet roots, carrots and radish roots.

Combined pydiflumetofen residues in sugar beet roots, carrots and radish roots were (n = 28): 0.01, 0.02 (4), 0.03, 0.04 (2), 0.05, 0.06 (6), 0.07, 0.08 (2), 0.09, 0.10 (3), 0.12, 0.13, 0.14 (2) and 0.17 (2) mg/kg.

The Meeting concluded that residues in sugar beet roots, carrots and radish roots may be influenced significantly by uptake of pydiflumetofen from the soil. The Meeting decided to add the

scaled mean residue found in field studies on succeeding crops (radish) of 0.02 mg/kg to the median residue obtained from supervised field trials on sugar beet roots, carrots and radish roots of 0.06 mg/kg for an overall STMR for sugar beet roots, carrots and radish roots of 0.08 mg/kg. For the estimation of a maximum residue level, the highest residue of 0.07 mg/kg found in radish roots from the succeeding crop field trials was added to the highest residue of 0.18 mg/kg found in supervised field trials, resulting in an overall highest residue in root vegetables of 0.25 mg/kg.

The Meeting estimated a maximum residue level of 0.3 mg/kg, an HR value of 0.25 mg/kg and an STMR value of 0.08 mg/kg for the Subgroup of root vegetables to replace its previous recommendation.

Sorghum

In field studies on succeeding crops, reviewed by the 2019 Meeting, the scaled mean, median and highest residues in cereals (wheat) were < 0.03, < 0.03, < 0.03 mg/kg, respectively. Using this data, the Meeting recommended a maximum residue level of 0.03 mg/kg for the Subgroup of sorghum grain and millet.

The current Meeting received supervised residue trials conducted in the USA on sorghum matching the critical GAP for sorghum from the USA; 2×126 g ai/ha, 7-day RTI, 21-day PHI.

Pydiflumetofen residues in sorghum grain in ranked order were (n = 12): 0.11 (3), 0.30, 0.44, 0.45, 0.58, 0.64 (2), 0.70, 1.1 and 1.8 mg/kg.

The Meeting concluded that residues from uptake of pydiflumetofen via the roots are insignificant in comparison to residue levels following direct treatment.

The Meeting estimated a maximum residue level of 3 mg/kg and an STMR of 0.515 mg/kg for grain sorghum. As the registered use is only for grain sorghum, the Meeting estimated a maximum residue level of 0.03 mg/kg and STMR of 0.03 mg/kg for the Subgroup of sorghum grain and millet, except grain sorghum and withdrew its previous recommendation for the Subgroup of sorghum grain and millet.

Tree nuts

The critical GAP for tree nuts is from the USA; 3×100 g ai/ha, 7-day RTI, 14-day PHI. The Meeting received supervised residue trials conducted in the USA on almond and pecan approximating the critical GAP.

Pydiflumetofen residues in almonds in ranked order were (n = 5): < 0.01 (3) and 0.03 (2) mg/kg.

Pydiflumetofen residues in pecans in ranked order were (n = 5): < 0.01 (4) and 0.02 mg/kg.

The Meeting noted that the GAP covers the group of tree nuts and that median residues of almonds and pecans are within a 5-fold difference. Therefore, the Meeting decided to combine the two datasets of almonds and pecans.

Combined pydiflumetofen residues in almonds and pecans were (n = 10): ≤ 0.01 (7), 0.02 and 0.03 (2) mg/kg.

Tree nuts are cultivated as permanent crops and are not expected to be subject to a potential uptake of pydiflumetofen from the soil. The Meeting estimated a maximum residue level of 0.05 mg/kg, an HR of 0.03 mg/kg and an STMR of 0.01 mg/kg for the Group of tree nuts.

Subgroups of sunflower seeds and cotton seeds

In field studies on succeeding crops, reviewed by the 2019 Meeting, the scaled mean, median and highest residues in wheat straw (worse case) were 0.10, 0.08 and 0.28 mg/kg, respectively. Using this data, the Meeting recommended a maximum residue level of 0.3 mg/kg for the Subgroups of sunflower seeds and cottonseed.

Sunflower seeds

The current Meeting received supervised residue trials conducted in Canada and the USA on sunflower seed matching the critical GAP for the sunflower seed subgroup from the USA; 2×200 g ai/ha, 14-day RTI, 30-day PHI.

Pydiflumetofen residues in sunflower seeds in ranked order were (n = 11): 0.02 (3), 0.05, 0.07 (2), 0.08, 0.12, 0.13 (2) and 0.37 mg/kg (highest value of 0.41 mg/kg from replicate samples).

The Meeting concluded that residues in sunflower seeds may be influenced by uptake of pydiflumetofen from the soil. Following a re-assessment of the field studies on succeeding crops, the Meeting concluded that the soya bean seed data was most relevant to oilseeds, where all scaled residues were < 0.02 mg/kg in soya bean seeds. Therefore, for the estimation of a maximum residue level, the value of 0.02 mg/kg was added to the highest residue of 0.41 mg/kg found in supervised field trials, resulting in an overall highest residue in sunflower seeds of 0.43 mg/kg. For the estimation of the STMR, a value of 0.02 mg/kg was added to the median residue of 0.07 mg/kg for an overall STMR for sunflower seeds of 0.09 mg/kg.

Noting that the GAP covers the sunflower seed subgroup, the Meeting estimated a maximum residue level of 0.5 mg/kg and an STMR of 0.09 mg/kg for the Subgroup of sunflower seeds to replace its previous recommendation.

Cotton seeds

The critical GAP for cotton seed is from the USA; 3×75 g ai/ha, 10-day RTI, 45-day PHI. None of the cotton seed supervised residue trials were conducted according to the US GAP.

Although the Meeting could not estimate a maximum residue level on cotton seeds to account for primary use of pydiflumetofen, cotton seeds may still be subject to crop rotation and therefore contain pydiflumetofen residues after uptake via the roots. Similar to sunflower seeds, the Meeting concluded that the secondary crop field studies on soya bean seeds were most relevant to oilseeds, where all scaled pydiflumetofen residues were < 0.02 mg/kg. Therefore, the Meeting estimated a maximum residue level of 0.02 mg/kg and an STMR of 0.02 mg/kg for cottonseed to replace its previous recommendation.

Residues in animal feeds

Leaves of root vegetables

In field studies on succeeding crops, reviewed by the 2019 JMPR, the scaled mean, median and highest residues in radish tops were 0.02, 0.02, 0.05 mg/kg, respectively. Using this data, the 2019 JMPR recommended a maximum residue level of 0.07 mg/kg for the Subgroup of Leaves of root and tuber vegetables, except leaves of tuber vegetables.

The US GAP prohibits the feeding of radish tops to livestock, therefore, only sugar beet tops were considered as animal feed. In trials matching the critical GAP, pydiflumetofen residues in sugar beet tops in ranked order were (n = 17); 0.76, 0.77, 0.92, 0.97, 1.0 (2), 1.4 (2), 1.5 (2), 1.6, 1.7, 3.4, 3.6, 5.3, 5.7 and 6.3 mg/kg (highest value of 8.0 mg/kg from replicate samples).

The Meeting concluded that residues from uptake of pydiflumetofen via the roots are insignificant in comparison to residue levels following direct treatment.

The Meeting estimated a maximum residue level of 40 mg/kg (dw; using the OECD standard dry matter content of 23%), a highest residue of 8.0 mg/kg and a median residue of 1.5 mg/kg for sugar beet tops (as received) and withdraws its previous recommendation for the subgroup.

Sorghum forage

In field studies on succeeding crops, reviewed by the 2019 Meeting, the scaled mean, median and highest residues in cereal forages (wheat forage) were 0.03, 0.03, 0.03 mg/kg, respectively.

In trials matching the critical GAP, pydiflumetofen residues in sorghum forage in ranked order were (n = 12): 0.05, 0.09 (2), 0.14 (2), 0.18, 0.28, 0.45 (2), 0.53, 0.61 and 0.70 mg/kg (highest value of 0.73 mg/kg from replicate samples).

The Meeting concluded that residues from uptake of pydiflumetofen via the roots are insignificant in comparison to residue levels following direct treatment.

The Meeting estimated a median residue value of 0.23 mg/kg and a highest residue value of 0.73 mg/kg for sorghum forage (as received).

Sorghum stover

In field studies on succeeding crops, reviewed by the 2019 Meeting, the scaled mean, median and highest residues in cereal straw (wheat straw) were 0.10, 0.08, 0.28 mg/kg, respectively. Using this data, the 2019 Meeting recommended a maximum residue level of 0.3 mg/kg (dry weight) for sorghum straw and fodder, dry.

In trials matching the critical GAP, pydiflumetofen residues in sorghum stover (dry weight) in ranked order were (n = 12): 0.07, 0.15, 0.19, 0.21, 0.25, 0.38, 0.62, 0.66, 1.2, 3.4, 5.0 and 6.0 mg/kg (highest value of 7.6 mg/kg from replicate samples).

The Meeting concluded that residues from uptake of pydiflumetofen via the roots are insignificant in comparison to residue levels following direct treatment.

The Meeting estimated a maximum residue level of 10 mg/kg (dry weight), a highest residue value of 7.6 mg/kg and a median residue value of 0.50 mg/kg for sorghum straw and fodder, dry to replace its previous recommendation.

Almond hulls

In trials matching the critical GAP, pydiflumetofen residues in almond hulls in ranked order were (n = 5): 0.37, 1.4, 1.6, 3.6 and 3.9 mg/kg.

The Meeting estimated a maximum residue level of 10 mg/kg (dw; using the OECD standard dry matter content of 90%) and a median residue value of 1.6 mg/kg (as received) for almond hulls.

Cotton gin by-products

None of the cotton seed supervised residue trials were conducted according to the critical GAP from the USA; 3×75 g ai/ha, 10-day RTI, 45-day PHI.

Therefore, the Meeting did not estimate a maximum residue level for cotton gin by-products.

Fate of residues during processing

Processing data on oranges, apples, plums, sugar beets, grain sorghum, almonds, sunflower and cotton were provided. All data relevant for an estimation of maximum residue levels in processed commodities or for dietary exposure calculations are summarized in the following table.

Table 1 Processing Factors and STMR-P/HR-P

RAC [STMR/HR]	Processed commodity	Individual processing factors	Mean or best estimate processing factor	STMR-P = $STMR_{RAC} \times PF$ (mg/kg)	HR-P = $HR_{RAC} \times PF$ (mg/kg)
Citrus [0.21 mg/kg / 0.76 mg/kg]	Peel	1.78, 1.94	1.86	0.39	1.4
	Juice	< 0.02, < 0.02	< 0.02	0.004	
	Oil	33.7, 52.7	43.2	9.1	
	Dried pulp	1.45	1.45	0.30	
Apple [0.06 mg/kg / 0.13 mg/kg]	Canned apple	0.03	0.03	0.002	
	Apple juice	0.06	0.06	0.004	
	Apple sauce	0.06	0.06	0.004	

RAC [STMR/HR]	Processed commodity	Individual processing factors	Mean or best estimate processing factor	STMR-P = $STMR_{RAC} \times PF$ (mg/kg)	HR-P = $HR_{RAC} \times PF$ (mg/kg)
	Dried apple	0.41	0.41	0.02	0.05
	Apple wet pomace	3.77	3.77	0.22	
Pear [0.05 mg/kg / 0.13 mg/kg]	Canned fruit	0.09	0.09	0.004	
	Dried fruit	0.62	0.62	0.03	0.08
	Juice	0.11	0.11	0.006	
	Wet pomace	3.0	3.0	0.15	
Plum [0.15 mg/kg / 0.37 mg/kg]	Juice	0.01, 0.02	0.02	0.003	
	Puree	0.15, 0.27	0.21	0.03	
	Prune, dried	1.8, 2.8	2.3	0.34	0.85
Sugar beet [0.06 mg/kg / 0.15 mg/kg]	Refined sugar	< 0.07, 0.07	0.07	0.004	
	Molasses	0.50, 0.68	0.59	0.04	
	Dried pulp	1.8, 6.0	3.9	0.24	
Grain sorghum [0.515 mg/kg / 1.84 mg/kg]	Flour	0.82, 0.91	0.87	0.45	
	Aspirated grain fraction	61, 63	62	31.9	
Almond [0.01 mg/kg / 0.03 mg/kg]	Roasted	0.37	0.37	0.004	0.01
	Oil	0.34	0.34	0.003	
Sunflower [0.09 mg/kg / 0.43 mg/kg]	Oil (refined)	0.02, 0.09	0.06	0.005	
	Meal	< 0.02, 0.06	0.04	0.004	
Cotton [0.02 mg/kg / 0.02 mg/kg]	Oil (refined)	0.02, 0.04	0.03	0.0006	
	Hulls	0.12, 0.13	0.13	0.003	
	Meal	< 0.01, < 0.01	< 0.01	0.0002	

Using the estimated maximum residue level of 0.9 mg/kg for the group of citrus fruits and applying the processing factors of 43.2 for citrus oil and 1.45 for citrus pulp, dry, the Meeting estimated maximum residue levels of 40 mg/kg for citrus oil and 1.5 mg/kg for citrus pulp, dry.

Using the estimated maximum residue level of 0.6 mg/kg for the subgroup of plums and applying the processing factor of 2.3 for dried prune plums, the Meeting estimated a maximum residue level of 1.5 mg/kg for plums, dried (prunes).

Residues in animal commodities

Farm animal feeding studies

Farm animal feeding studies (lactating dairy cow and laying hen) are reported in the evaluation of the 2018 JMPR.

Farm animal dietary burden

The dietary burdens for beef cattle, dairy cattle, broilers and laying poultry determined at the 2019 Meeting were updated with the feed items listed in the table below.

Table 2 Potential feed items

Codex Classification	Commodity	Median residue (-P) mg/kg	Highest residue (-P) mg/kg
Orange	Dried pulp	0.30	
Apple	Wet pomace	0.22	
Brassica Head Vegetables	Cabbage head with wrapper	0.62	1.2

Codex Classification	Commodity	Median residue (-P) mg/kg	Highest residue (-P) mg/kg
	leaves		
Root vegetables	Carrot culls, swede roots, turnip roots	0.08	0.24
Sugar beet	Molasses	0.04	
	Dried pulp	0.24	
Leaves of root vegetables	Sugar beet tops	1.5	8.0
Grain sorghum	Grain	0.515	
	Aspirated grain fractions	31.9	
	Forage	0.23	0.73
	Stover	0.50	7.6
Almonds	Hulls	1.6	
Sunflower	Meal	0.004	

The dietary burdens, estimated using the most recent version of the OECD livestock dietary burden calculator, are presented in Annex 6 and summarized below.

Table 3 Estimated maximum and mean dietary burdens of farm animals

	Animal Dietary Burden: Pydiflumetofen, ppm of dry matter diet							
	US-Canada		EU		Australia		Japan	
	Max	Mean	Max	Mean	Max	Mean	Max	Mean
Beef cattle	12	5.6	31	9.0	45 ^a	19 ^b	1.4	1.0
Dairy cattle	28	9.5	31	10	45 ^c	13 ^d	13	4.7
Poultry - broiler	0.56	0.56	0.76	0.59	0.51	0.51	0.42	0.42
Poultry - layer	0.56	0.56	8.7 ^e	3.0 ^f	0.51	0.51	0.46	0.46

^a Highest maximum beef cattle dietary burden suitable for maximum residue level estimates for mammalian tissues

^b Highest mean beef cattle dietary burden suitable for STMR estimates for mammalian tissues

^c Highest maximum dairy cattle dietary burden suitable for MRL estimate for milk

^d Highest mean dairy cattle dietary burden suitable for STMR estimate for milk

^e Highest maximum poultry dietary burden suitable for MRL estimates for poultry tissues and eggs

^f Highest mean poultry dietary burden suitable for STMR estimates for poultry tissues and eggs

Animal commodity maximum residue levels

Cattle residues in tissues and milk at the expected dietary burden for dairy cattle are shown in the Table below.

Table 4 Anticipated residues of pydiflumetofen in mammalian commodities

	Feed level (ppm) for milk residues	Total residues in milk (mg/kg)	Feed level (ppm) for tissue residues	Total residues ^a (mg/kg)			
				Muscle	Liver	Kidney	Fat
MRL Determination							
Feeding study	45	< 0.01	45	< 0.01	0.05	< 0.01	0.05
Dietary burden	45	< 0.01	45	< 0.01	0.05	< 0.01	0.05
HR Determination							
Feeding study			45	< 0.02	0.44	0.30	0.07
Dietary burden			45	< 0.02	0.44	0.30	0.07
STMR Determination							
Feeding study	15	< 0.02	15	< 0.02	0.06	0.07	0.02
			45	< 0.02	0.29	0.25	0.06
Dietary burden	13	< 0.02	19	< 0.02	0.09	0.09	0.02

^a For MRL determination: pydiflumetofen; For HR and STMR determination for milk, muscle and fat: sum of pydiflumetofen and 2,4,6-TCP and its conjugates, expressed as pydiflumetofen; For HR and STMR determination for liver and kidney: sum of pydiflumetofen, 2,4,6-TCP and its conjugates and SYN547897 and its conjugates, expressed as pydiflumetofen

Based on pydiflumetofen residues in milk and cattle tissues, the Meeting confirmed the maximum residue levels estimated by the 2019 Meeting (0.01 (*) mg/kg in milk, 0.1 mg/kg in mammalian meat (in the fat), mammalian edible offal and mammalian fat).

Based on the highest estimated total residues of pydiflumetofen and 2,4,6-TCP expressed as pydiflumetofen in muscle and fat, the Meeting estimated HR values of 0.02 mg/kg in mammalian meat and 0.07 mg/kg in mammalian fat.

Based on the highest estimated total residues of pydiflumetofen, 2,4,6-TCP and SYN547897 expressed as pydiflumetofen in liver and kidney, the Meeting estimated an HR value of 0.44 mg/kg in liver and 0.30 mg/kg in kidney.

Based on the mean estimated total residues of pydiflumetofen and 2,4,6-TCP expressed as pydiflumetofen in milk, muscle and fat, the Meeting estimated STMR values of 0.02 mg/kg in milk, 0.02 mg/kg in mammalian meat and 0.02 mg/kg in mammalian fat.

Based on the mean estimated total residues of pydiflumetofen, 2,4,6-TCP and SYN547897 expressed as pydiflumetofen in liver and kidney, the Meeting estimated STMR values of 0.09 mg/kg for each.

Table 5 Anticipated residues of pydiflumetofen in poultry commodities

	Feed level (ppm) for egg residues	Total residues in egg (mg/kg)	Feed level (ppm) for tissue residues	Total residues (mg/kg)		
				Muscle	Liver	Fat
MRL Determination - Pydiflumetofen						
Feeding study	9	0.011	9	< 0.01	< 0.01	< 0.01
Dietary burden	8.7	0.011	8.7	< 0.01	< 0.01	< 0.01
HR Determination - Sum of pydiflumetofen and 2,4,6-TCP and its conjugates, expressed as pydiflumetofen						
Feeding study	9	0.023	9	< 0.02	< 0.02	< 0.02
Dietary burden	8.7	0.023	8.7	< 0.02	< 0.02	< 0.02
STMR Determination - Sum of pydiflumetofen and 2,4,6-TCP and its conjugates, expressed as pydiflumetofen						
Feeding study	3	< 0.02	3	< 0.02	< 0.02	< 0.02
Dietary burden	3.0	< 0.02	3.0	< 0.02	< 0.02	< 0.02

Based on pydiflumetofen residues in eggs and poultry tissues, the Meeting confirmed the maximum residue levels estimated by the 2019 Meeting (0.02 mg/kg in eggs, 0.01 (*) mg/kg in poultry meat, poultry edible offal of and poultry fat).

Based on the highest estimated total residues of pydiflumetofen and 2,4,6-TCP expressed as pydiflumetofen in eggs, muscle, liver and fat, the Meeting estimated HR values of 0.023 mg/kg in eggs and 0.02 mg/kg in poultry meat, poultry, edible offal of and poultry fat.

Based on the mean estimated total residues of pydiflumetofen and 2,4,6-TCP expressed as pydiflumetofen in eggs, muscle, liver and fat, the Meeting estimated STMR values of 0.02 mg/kg in eggs, poultry meat, poultry, edible offal of and poultry fat.

RECOMMENDATIONS

On the basis of the data obtained from supervised trials, the Meeting concluded that the residue levels listed in Annex 1 are suitable for establishing maximum residue limits and for IEDI and IESTI assessments.

Definition of the residue for compliance with the MRL for plant and animal commodities and dietary risk assessment for plant commodities: pydiflumetofen.

Definition of the residue for dietary risk assessment for animal commodities other than mammalian liver and kidney: Sum of pydiflumetofen and 2,4,6-TCP and its conjugates, expressed as pydiflumetofen

Definition of the residue for dietary risk assessment for mammalian liver and kidney: Sum of pydiflumetofen, 2,4,6-TCP and its conjugates and SYN547897 and its conjugates, expressed as pydiflumetofen.

The residue is fat-soluble.

DIETARY RISK ASSESSMENT

Long-term dietary exposure

The ADI for pydiflumetofen is 0–0.1 mg/kg bw. The International Estimated Daily Intakes (IEDIs) for pydiflumetofen were estimated for the 17 GEMS/Food Consumption Cluster Diets using the STMR or STMR-P values estimated by the JMPR. The results are shown in Annex 3 of the 2021 Extra JMPR Report.

The IEDIs ranged from 2–20% of the maximum ADI. The Meeting concluded that long-term dietary exposure to residues of pydiflumetofen from uses considered by the JMPR is unlikely to present a public health concern.

Acute dietary exposure

The ARfD for pydiflumetofen is 0.3 mg/kg bw. The International Estimate of Short-Term Intakes (IESTIs) for pydiflumetofen were calculated for the food commodities and their processed commodities for which HRs/HR-Ps or STMRs/STMR-Ps were estimated by the present Meeting and for which consumption data were available. The results are shown in Annex 4 of the 2021 Extra JMPR Report.

The IESTIs were 0–20% of the ARfD for the general population and 0–20% of the ARfD for children. The Meeting concluded that acute dietary exposure to residues of pydiflumetofen from uses considered by the present Meeting are unlikely to present a public health concern.

5.23. QUINOXYFEN (222)

RESIDUE AND ANALYTICAL ASPECTS

Quinoxifen is a fungicide used for protection against powdery mildew diseases on a variety of crops.

Quinoxifen was first evaluated for toxicology and residues by JMPR in 2006. An ADI of 0–0.2 mg/kg bw was established while an ARfD was unnecessary. The residue definition for compliance with the MRL and dietary risk assessment for plant and animal commodities is quinoxifen. The residue is fat-soluble.

The Forty-third Codex Alimentarius Commission (2020) approved the new work proposals including the priority list of pesticides for evaluation by the current Meeting. The priority list included quinoxifen for evaluation of additional uses by the 2021 Extra JMPR.

The Meeting received information from the manufacturer on use patterns, supervised residue trials on stone fruits, melons, winter squash, tomato and globe artichokes and processing studies.

Methods of analysis

The GC-MSD analytical method ERC 95.26 and the LC-MS/MS analytical method ERC 95.26.S1 used for analysis of residues of quinoxifen in plant commodities, with an LOQ of 0.01 mg/kg, were reviewed by the 2006 JMPR. All of the samples collected in the supervised residue trials and processing studies provided to the current Meeting were analysed using the same methods. The methods are fit for purpose based on the provided results from concurrent recovery samples.

Stability of pesticide residues in stored analytical samples

The stability of residues of quinoxifen during frozen storage was evaluated by the 2006 JMPR. Quinoxifen was determined to be stable when stored frozen for at least 320 days at -15 °C in commodities representative of the high-water content commodity group. Together with the concurrent storage stability data, the Meeting concluded that quinoxifen is stable in the test crops and processed commodities under frozen storage conditions.

Results of supervised residue trials on crops

Supervised trials were available for the use of quinoxifen on cherries, plums, peaches, melons, winter squash, tomato and globe artichoke.

Stone fruits

Cherries

The Meeting received a new GAP for cherries from the USA allowing 4–5 foliar applications at a rate of 128 g ai/ha with a minimum 7-day RTI and a 7-day PHI and seasonal application rates not exceeding 585 g ai/ha.

In the supervised residue trials from the USA on cherries, previously reviewed by the 2006 JMPR and approximating the critical GAP, quinoxifen residues in cherries in ranked order were (n = 7): 0.05, 0.08, 0.12, 0.14 (2), 0.15 and 0.27 mg/kg.

Residues in the field trials were reported as fruit without stone. At the 2017 JMPR, it was concluded that, based on the trials on cherries, the contribution of the pit to the weight of the whole fruit was approximately 10%. Correcting the residue levels using this weight/weight ratio would lead to the same maximum residue level.

Noting that the USA GAP is for the cherries group, the Meeting estimated a maximum residue level of 0.5 mg/kg and an STMR of 0.14 mg/kg for the Subgroup of cherries, except choke cherries and withdrew its previous recommendation.

Stone fruits except cherries

The critical GAP for stone fruits except cherries in the USA includes apricot, chickasaw plum, damson plum, Japanese plum, nectarine, peach, plum, plumcot, prune (fresh). The critical GAP allows 4 foliar spray applications at 128 g ai/ha, a 10-day RTI and a 7-day PHI, with a seasonal rate not exceeding 512 g ai/ha. The Meeting received trials conducted in the USA approximating the critical GAP with the exception of the RTIs which were 6–8 days.

Supervised field trials were on average conducted at rates 15% greater than the critical GAP and the RTIs were significantly shorter. Therefore, without crop-specific residue decline data to estimate half-lives, the Meeting concluded that the overall impact of these parameters on the residues was >25% and that there were insufficient data to estimate a maximum residue level.

Fruiting vegetables – Cucurbits

Melons

The 2006 JMPR recommended a maximum residue level of 0.1 mg/kg for melons. The Meeting considered if the previously evaluated data would support extrapolation to the subgroup of melons.

The critical GAP for the melons subgroup (including citron melon, muskmelon (cantaloupe, casaba, crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon, true cantaloupe, watermelon) is from the USA which allows 4 foliar spray applications at 110 g ai/ha, a 10-day RTI and a 3-day PHI, with a seasonal rate not exceeding 438 g ai/ha.

The Meeting received the same trials conducted in Canada and the USA as those reviewed by the 2006 Meeting, where 4–5 foliar spray applications were made at 139–169 g ai/ha, 6–8-day RTIs, seasonal application rates of 582–748 g ai/ha and 2–4-day PHIs. The 2006 Meeting also reviewed trials conducted in Europe involving 3 foliar spray applications at 0.005–0.0075 kg ai/hL and PHIs of 7–8-days, however, these trials do not match the USA GAP and were not considered by the current Meeting.

As supervised field trials were conducted at rates 33–71% greater than the critical GAP and the RTIs were shorter, the Meeting concluded that the overall impact of these parameters on the residues was > 25%. Therefore, using the latest available tools and following the most current assessment practices, the Meeting decided that a maximum residue level could not be extended to the subgroup.

Winter squashes

The critical GAP for the winter squash, gourds and pumpkins is from the USA which allows 4 foliar spray applications at 146 g ai/ha, 10-day RTI and a 3-day PHI, with a seasonal rate not exceeding 583 g ai/ha. The Meeting received trials from the USA on winter squashes approximating the critical GAP with the exception of the RTIs, the majority of which ranged from 6–8 days.

While supervised field trials were conducted according to the critical GAP in most respects, the RTIs were significantly shorter. Without crop-specific residue decline data to estimate half-lives, the Meeting concluded that the overall impact of these parameters on the residues was > 25% and that there were insufficient data to estimate a maximum residue level for the subgroup.

Tomato

The critical GAP for tomatoes is from the USA which allows 4 foliar spray applications at 110 g ai/ha, 10-day RTIs and a 3-day PHI, with a seasonal rate not exceeding 438 g ai/ha. The Meeting received trials from the USA on tomatoes where 4 foliar spray applications were made at 143–162 g ai/ha, 6–8-day RTIs, seasonal application rates of 574–733 g ai/ha and 2–4-day PHIs.

As supervised field trials were conducted at rates 31–67% greater than the critical GAP and the RTIs were significantly shorter, the Meeting concluded that the overall impact of these parameters

on the residues was >25%. Therefore, the Meeting decided that a maximum residue level could not be estimated.

Globe artichokes

The critical GAP for globe artichokes is from the USA where the number of spray applications at 110 g ai/ha is unspecified, however the seasonal rate must not exceed 583 g ai/ha, the RTI is 10 days and the PHI is 0 days. The Meeting received trials from the USA on globe artichokes where 4 foliar spray applications were made at 145–151 g ai/ha, 6–7-day RTIs, seasonal application rates of 583–600 g ai/ha and a 0-day PHI.

The number of trials were insufficient and the use pattern did not match the critical GAP with respect to the RTIs (6–7 days vs 10 days). Therefore, the Meeting could not recommend a maximum residue level.

Residues in processed commodities

At the current Meeting, processing studies were reviewed for plums and tomato. As the Meeting could not recommend maximum residue levels for plums and tomatoes, residues in processed commodities were not estimated.

Residues in animal commodities

Farm animal dietary burden

Dietary burdens were calculated for beef cattle, dairy cattle, broilers and laying poultry at the 2006 JMPR. The dietary burdens, were estimated using the diets listed in Appendix IX of the FAO Manual, included barley (grain and straw) and sugar beet leaves. None of the crops assessed at the current Meeting are fed to livestock, therefore no updates to the livestock dietary burdens were warranted. The Meeting confirms its previous recommendations for residues in animal commodities made by the 2006 JMPR.

RECOMMENDATIONS

On the basis of the data obtained from supervised trials, the Meeting concluded that the residue levels listed in Annex 1 are suitable for establishing maximum residue limits and for IEDI assessment.

Definition of the residue for compliance with the MRL and dietary risk assessment for plant and animal commodities: *quinoxifen*.

The residue is fat-soluble.

DIETARY RISK ASSESSMENT

Long-term dietary exposure

The ADI for quinoxifen is 0–0.2 mg/kg bw. The International Estimated Daily Intakes (IEDIs) for quinoxifen were estimated for the 17 GEMS/Food Consumption Cluster Diets using the STMR or STMR-P values estimated by the JMPR. The results are shown in Annex 3 of the 2021 Extra JMPR Report.

The IEDIs ranged from 0–1% of the maximum ADI. The Meeting concluded that long-term dietary exposure to residues of quinoxifen from uses considered by the JMPR is unlikely to present a public health concern.

Acute dietary exposure

The 2006 JMPR decided that an ARfD for quinoxifen was unnecessary. The Meeting therefore concluded that the acute dietary exposure to residues of quinoxifen from the uses considered is unlikely to present a public health concern.

5.24. SPINETORAM (233)

RESIDUE AND ANALYTICAL ASPECTS

Spinetoram is a spinosyn insecticide obtained by chemical modification of a fermentation product of *Saccharopolyspora spinose*. It consists of two closely related active ingredients, spinetoram-J and spinetoram-L, present in approximately a three to one ratio.

It was first evaluated by the 2008 JMPR (T, R), which established an ADI of 0–0.05 mg/kg bw and an ARfD was unnecessary. The 2008 JMPR also concluded the residue definition of spinetoram for plant and animal commodities:

Definition of the residue for compliance with the MRL for plant and animal commodities: *Spinetoram*.

Definition of the residue for dietary risk assessment for plant and animal commodities: *Spinetoram and N-demethyl and N-formyl metabolites of the major spinetoram component*.

The residue is fat-soluble.

Spinetoram was subsequently reviewed by 2012 and 2017 JMPR for additional uses. Codex MRLs have been established for a wide range of crops and products of animal origin.

The Forty-third Codex Alimentarius Commission (2020) approved the new work proposals including the priority list of pesticides for evaluation by the current Meeting. The priority list included spinetoram for evaluation of additional uses on pitaya (dragon fruit) and tea.

The current Meeting received GAP information and supervised residue trials on pitaya (dragon fruit) and tea.

Methods of analysis

The Meeting received descriptions and validation data for analytical methods for residues of spinetoram J and L in dried tea leaves and tea infusion along with the supervised trials.

The dry tea was first swollen by water, extracted with methanol/water (8:2, v/v), purified using polymer-based cartridges and an NH₂ cartridge column and quantified by LC/MS/MS. Tea infusions were extracted with acetonitrile/water (1:1, v/v), purified on an SPE column and analysed by LC-MS/MS. The LOQ of 0.01 mg/kg was successfully validated for spinetoram-J and spinetoram-L.

The methods are considered suitable for analysis of spinetoram J and L in dry green tea and tea infusions.

GRM 05.03 and GRM 05.04, already reviewed by the 2008 JMPR, were satisfactorily validated to determine spinetoram-J and spinetoram-L and the metabolites N-demethyl-spinetoram-J and N-formyl-spinetoram-J in pitaya, with an LOQ of 0.01 mg/kg.

Stability of pesticide residues in stored analytical samples

The Meeting received frozen storage stability data for spinetoram and its N-demethyl and N-formyl metabolites in pitaya and dry green tea included in supervised trials. The Meeting concluded that spinetoram and its N-demethyl-spinetoram-J and N-formyl-spinetoram-J metabolites were stable for at least 11 months (342 days) in pitaya and spinetoram was stable for at least 5 months (166 days) in dry green tea samples stored frozen at -20 °C.

The demonstrated storage stability intervals covered the duration of storage in the residue trials evaluated by the current Meeting, except for one sample in the pitaya trials stored for 362 days, which is less than 10% beyond the covered period of 342 days and was considered unlikely to have a significant impact on the residues.

Results of supervised residue trials on crops

The Meeting received information on supervised trials of spinetoram on pitaya (dragon fruits) and tea.

Pitaya

The critical GAP for spinetoram on pitaya in the USA is 4 applications at 123 g ai/ha with an interval of 4 days and a PHI of 1 day.

Four trials were conducted on pitaya in the USA with four applications at rates approximating 110 g ai/ha and re-treatment intervals of 3–6 days and a PHI of 1 day. The residues of spinetoram (J and L) in fruits were (n = 4): 0.015, 0.044, 0.047 and 0.20 mg/kg. The residues of spinetoram-J, spinetoram-L, N-demethyl-spinetoram-J and N-formyl spinetoram-J were (n = 4): 0.071, 0.088, 0.095 and 0.64 mg/kg.

The Meeting estimated a maximum residue level of 0.5 mg/kg and an STMR of 0.0915 mg/kg for spinetoram in pitaya.

Tea, Green, Black (black, fermented and dried)

The critical GAP for spinetoram on tea in Japan is 1 application at 4.68 g ai/hL (2500 times dilution with spray volume of 4000 L/ha) and a PHI of 1 day.

Eight trials were conducted on tea in Japan matching the critical GAP. The residues of spinetoram (J and L) in dry green tea were (n = 8): 6.1, 12.5, 16, 20(2), 31, 32 and 40 mg/kg. The residues of N-demethyl-spinetoram-J and N-formyl spinetoram-J were not analysed.

The Meeting estimated a maximum residue level of 70 mg/kg for spinetoram in tea, green, black (black, fermented and dried).

In an apple metabolism study evaluated by the 2008 JMPR (one foliar application at a rate of 1.8 kg ai/ha), the residues of spinetoram-J in leaves at 1 day after application were 78.6% TRR, while the residues of N-demethyl and N-formyl spinetoram-J were 12.8% TRR, the ratio of the two metabolites to spinetoram-J was 0.16.

The Meeting noted that tea infusions represent a small part of the diet and their contribution to the overall exposure is negligible. In trials on leek, spring onion and spinach reported previously by the JMPR (2012 and 2017) the ratio of N-demethyl-spinetoram-J to spinetoram J ranged from 0.07 to 0.63 and for N-formyl-spinetoram J to spinetoram J from 0.31 to 4.5. If the metabolites are assumed to be present at 1× and 5× the level of spinetoram J, a conservative assumption, the STMR would be $20 + 16.2 + 5 \times 16.2 = 118$ mg/kg. The latter could be used in estimating exposure in the absence of measured residues.

The Meeting estimated an STMR of 118 mg/kg for spinetoram in tea, green, black (black, fermented and dried).

Fate of residues during processing

The Meeting received processing studies for tea infusion. However, as not all the metabolites included in the residue definition for risk assessment were measured, the data could not be used to estimate processing factors for tea infusion.

RECOMMENDATIONS

On the basis of the data obtained from supervised trials, the Meeting concluded that the residue levels listed in Annex 1 are suitable for establishing maximum residue limits and for IEDI assessments.

Definition of the residue for compliance with the MRL for plant and animal commodities: *Spinetoram*.

Definition of the residue for dietary risk assessment for plant and animal commodities: *Spinetoram and N-demethyl and N-formyl metabolites of the major spinetoram component*.

The residue is fat-soluble.

DIETARY RISK ASSESSMENT

Long-term dietary exposure

The ADI for spinetoram is 0–0.05 mg/kg bw. The International Estimated Daily Intakes (IEDIs) for spinetoram were estimated for the 17 GEMS/Food Consumption Cluster Diets using the STMR or STMR-P values estimated by the JMPR. The results are shown in Annex 3 of the 2021 Extra JMPR Report. The IEDIs ranged from 2–20% of maximum ADI of 0.05 mg/kg bw. The Meeting concluded that the long-term dietary exposure to residues of spinetoram from uses considered by the JMPR is unlikely to present a public health concern.

Acute dietary exposure

The 2008 JMPR decided that an ARfD for spinetoram was unnecessary. The Meeting therefore concluded that the acute dietary exposure to residues of spinetoram from uses considered by the current Meeting is unlikely to present a public health concern.

5.25. SPIROMESIFEN (294)

RESIDUE AND ANALYTICAL ASPECTS

Spiromesifen is a contact insecticide-acaricide belonging to the titronic acid class of compounds. The mode of action is inhibition of lipid biosynthesis, especially triglycerides and free fatty acids.

Spiromesifen was first evaluated by the 2016 JMPR where an ADI of 0–0.03 mg/kg bw was established and an ARfD was determined to be unnecessary. The residue definition for compliance with the MRL for plant and animal commodities and for dietary risk assessment for animal commodities is sum of spiromesifen and spiromesifen-enol, expressed as spiromesifen. For dietary risk assessment for plant commodities, the residue definition is sum of spiromesifen, spiromesifen-enol and 4-hydroxymethyl-spiromesifen-enol (free and conjugated), expressed as spiromesifen. The residue is fat-soluble.

Spiromesifen was scheduled at the Fifty-first Session of the CCPR for the evaluation of additional uses by the 2020 JMPR, which was postponed to the 2021 Extra JMPR. The Meeting received information on GAP and residue trials on chili pepper and okra.

Methods of analysis

Spiromesifen residues (parent only) in green chili pepper and okra were analysed based on the method previously evaluated by the JMPR. The method involved QuEChERS sample preparation method and determination of spiromesifen by LC-MS/MS or GC-MS. Recovery tests for the method validation showed mean recoveries of 70–100% in chili pepper and 83–114% in okra. The LOQ values (LC-MS/MS and GC-MS) for spiromesifen were 0.05 mg/kg in chili pepper and okra. The analytical method used in the chili pepper and okra residue trials was considered sufficiently validated.

Stability of pesticide residues in stored analytical samples

Residue analysis was performed within 24 hours after sample collection.

Results of supervised residue trials on crops

Fruiting vegetables, other than Cucurbits

Chili pepper

The critical GAP for the use of spiromesifen on chili pepper in India involves foliar sprays at 0.096 kg ai/ha and a 7-day PHI (the maximum number of sprays and minimum re-treatment intervals are not specified).

Six trials on field-grown chili peppers (decline trials) were conducted in India during 2015–2017 at a rate of 0.096 kg ai/ha with 2 applications at a 10-day interval and a PHI of 7 days. The residue concentrations of spiromesifen in green chili peppers were (n = 6): < 0.05, 0.060, 0.060, 0.22, 0.36 and 0.59 mg/kg.

Communication by the sponsor indicated that the local agricultural practice involves re-treatment intervals of 2–4 days. Residues declined with a median half-life of 4.07 days (1st order) and the modelled residue at a spray interval of 2 days and 2 applications differed by more than 25% compared with the supervised residue field trials. The Meeting concluded that the supervised field trials were conducted at a significantly longer re-treatment interval and cannot be used for the estimation of a maximum residue level.

Okra

The critical GAP for the use of spiromesifen on okra in India involves foliar sprays at 0.12 kg ai/ha and a 3-day PHI (the maximum number of sprays and minimum re-treatment intervals are not specified).

Seven trials on field-grown okra (decline trials) were conducted in India during 2015–2018 at a rate of 0.096–0.12 kg ai/ha with 2 applications at a 10-day interval and a PHI of 3 days. The residue concentrations of spiromesifen in okra were ($n = 7$): < 0.05 (3), 0.14, 0.15, 0.46 and 0.77 mg/kg.

Communication by the sponsor indicated that the local agricultural practice involves re-treatment intervals of 2–4 days. Residues declined with a median half-life of 2.41 days (1st order) and the modelled residue at a spray interval of 2 days and 2 applications differed by more than 25% compared with the supervised residue field trials. The Meeting concluded that the supervised field trials were conducted at a significantly longer re-treatment interval and cannot be used for the estimation of a maximum residue level.

5.26. **SULFOXAFLOR (252)**

RESIDUE AND ANALYTICAL ASPECTS

Sulfoxaflor (ISO common name) is a broad-spectrum, sulfoximine insecticide with registered uses on multiple crops. It was evaluated for the first time by the 2011 JMPR, which established an acceptable daily intake (ADI) of 0–0.05 mg/kg bw and an acute reference dose (ARfD) of 0.3 mg/kg bw. Sulfoxaflor underwent subsequent evaluations by the JMPR in 2014 and 2016.

The definition of the residue for compliance with the MRL and dietary risk assessment for plant and animal commodities is sulfoxaflor. The residue is not fat-soluble.

Sulfoxaflor was scheduled at the Fifty-first Session of the CCPR for evaluation of additional uses by the 2020 JMPR, which was postponed to the 2021 Extra JMPR. The current Meeting received information on analytical methods, field trials and processing studies to support estimation of new maximum residue levels in commodities of avocado, mango, bush berries, cane berries, globe artichoke, asparagus, sunflower and coffee.

Methods of analysis

The Meeting received method validation and concurrent recovery data for use of Method 091116S1 (reviewed by the 2011 JMPR) and the QuEChERS multiresidue method. All methods were demonstrated to have adequate performance for recovery of sulfoxaflor, with an LOQ of 0.01 mg/kg based on recoveries from method-validation and concurrent-recovery samples.

Stability of pesticide residues in stored analytical samples

Storage stability data were provided for blueberry, cane berry, artichoke, asparagus and sunflower commodities. Little, if any, dissipation during frozen storage was observed in any of the commodities tested. The Meeting concluded that residues of sulfoxaflor are stable as follows:

- Blueberry: at least 2.1 years,
- Cane berry: at least 1.5 years,
- Artichoke: at least 2 years,
- Asparagus: at least 10 months,
- Sunflower seed: at least 2 years,
- Sunflower meal: at least 22 months and
- Sunflower refined oil: at least 23 months.

Results of supervised residue trials on crops

The Meeting received data from supervised residue trials and GAP information on avocado, mango, bush berries, cane berries, artichoke, asparagus, sunflower and coffee.

Avocado

The critical GAP for avocado is from the USA. The label provides for up to four applications, each at 96 g ai/ha, on a 7-day interval, with a 7-day PHI. The label also specifies an annual limit of 298 g ai/ha; thus, the critical GAP is, essentially, three applications at the maximum rate.

Residues of sulfoxaflor in avocado (whole fruit) from independent trials approximating the critical GAP were (n = 5): < 0.013, 0.019, 0.049, 0.050 and 0.066 mg/kg.

The Meeting estimated a maximum residue level of 0.15 mg/kg for sulfoxaflor in avocado.

Residues of sulfoxaflor in avocado (flesh) from independent trials approximating the critical GAP were (n = 5): < 0.01, < 0.012, 0.011, 0.025 and 0.034 mg/kg.

The Meeting estimated an STMR of 0.011 mg/kg and an HR of 0.036 mg/kg (from a single sample) for residues of sulfoxaflor in avocado flesh.

Mango

The critical GAP for mango is from Kenya and consists of three applications, on a 14-day interval, each at 96 g ai/ha, with a 3-day PHI.

Field trials from Ghana, Kenya, Senegal, Tanzania and Uganda were conducted in accordance with the cGAP for application rate, re-treatment interval and PHI; however, two applications were made rather than three. Residue decline data for mango indicate a half-life of sulfoxaflor of approximately 3 days. Based on the half-life, the Meeting decided that a first application (31 days before harvest) would not contribute significantly to residues at harvest. Therefore, the Meeting determined that the trials sufficiently approximate the cGAP and are suitable for making residue recommendations.

Residues of sulfoxaflor in mango from independent trials approximating the cGAP were (n = 6): 0.026, 0.036, 0.039, 0.054, 0.10 and 0.11 mg/kg.

The Meeting estimated a maximum residue level of 0.3 mg/kg for sulfoxaflor in mango. The median residue and highest residue were 0.047 and 0.133 (from a single whole fruit sample) mg/kg, respectively.

Samples from two exaggerated rate trials were separated into peel and flesh. In comparison to whole fruit, residues in flesh were lower by a factor of 0.46 ($0.028 \text{ mg/kg} \div 0.061 \text{ mg/kg}$) and < 0.048 ($< 0.01 \text{ mg/kg} \div 0.21 \text{ mg/kg}$). Applying the factor of 0.46 derived from finite residues to the median and highest whole fruit residues gives an STMR of 0.022 mg/kg and an HR of 0.061 mg/kg in mango flesh for dietary risk assessment.

Subgroup of Bush berries

Blueberry

The critical GAP for blueberry is from the USA. The label provides for up to four applications, each at 96 g ai/ha, on a 7-day interval, with a 1-day PHI. The label also specifies an annual limit of 298 g ai/ha; thus, the critical GAP is, essentially, three applications at the maximum rate.

Residues of sulfoxaflor in blueberry from independent trials approximating the critical GAP were (n = 11): 0.16, 0.17, 0.27, 0.30, 0.31, 0.39 (2), 0.42 (2), 0.61 and 1.3 mg/kg.

Noting that the registered use is for the Bush berry subgroup, the Meeting estimated a maximum residue level of 2 mg/kg, an STMR of 0.39 mg/kg and an HR of 1.4 mg/kg (from a single sample) for sulfoxaflor in the Subgroup of Bush berries. Furthermore, as the US crop subgroup contains elderberry, the Meeting decided to extrapolate the recommendation to elderberry.

Subgroup of Cane berries

The critical GAP for cane berry is from the USA. The label provides for up to four applications, each at 96 g ai/ha, on a 7-day interval, with a 1-day PHI. The label also specifies an annual limit of 298 g ai/ha; thus, the critical GAP is, essentially, three applications at the maximum rate.

Residues of sulfoxaflor in blackberries and raspberries from independent trials approximating the critical GAP were (n = 7): 0.24, 0.28, 0.40, 0.44, 0.49, 0.54 and 0.74 mg/kg.

Noting that the registered use is for the Cane berry Subgroup, the Meeting estimated a maximum residue level of 1.5 mg/kg, an STMR of 0.44 mg/kg and an HR of 0.78 mg/kg (from a single sample) for sulfoxaflor in the Subgroup of Cane berries.

Globe artichoke

The critical GAP for globe artichoke is from the USA. The label provides for up to four applications, each at 96 g ai/ha, on a 14-day interval, with a 3-day PHI. The label also specifies an annual limit of 298 g ai/ha; thus, the critical GAP is, essentially, three applications at the maximum rate.

Residues of sulfoxaflor in globe artichoke from independent trials approximating the critical GAP were (n = 3): 0.22, 0.23 and 0.26 mg/kg.

The Meeting agreed that three trials in globe artichoke were not sufficient to estimate a maximum residue level.

Asparagus

The critical GAP for asparagus is from the USA. The label provides for up to four applications, each at 96 g ai/ha, on a 14-day interval. Application is to the ferns after harvest; as such, no PHI is specified on the label. The label specifies an annual limit of 298 g ai/ha; thus, the critical GAP is, essentially, three applications at the maximum rate.

Residues of sulfoxaflor in asparagus spears from independent trials approximating the critical GAP (harvested at least 4 months after the last treatment to the ferns) were (n = 6): ≤ 0.01 (5) and 0.011 mg/kg.

The Meeting estimated a maximum residue level of 0.015 mg/kg, an STMR of 0.01 mg/kg and an HR of 0.011 mg/kg for sulfoxaflor in asparagus.

Sunflower seeds

The critical GAP for sunflower is from the USA and consists of two applications, on a 7-day interval, each at 96 g ai/ha, with a 14-day PHI.

Residues of sulfoxaflor in sunflower seed from independent trials approximating the critical GAP were (n = 4): < 0.01, 0.013, 0.018 and 0.15 mg/kg.

The Meeting agreed that four trials in sunflower seed are not sufficient for estimating a maximum residue level.

Coffee

The critical GAP for coffee is from Vietnam and consists of application of a 0.125 g/L dilution of the product sprayed thoroughly to the whole plant. The number of treatments, the re-treatment interval and the spray volume are not specified and the PHI is 3 days.

Residues of sulfoxaflor in coffee berries (dried beans with shell) from independent trials approximating the critical GAP were (n = 10): < 0.01, 0.01 (3), 0.03, 0.04, 0.05, 0.06, 0.09 and 0.15 mg/kg.

The Meeting estimated a maximum residue level of 0.3 mg/kg and an STMR of 0.035 mg/kg.

Fate of residues during processing

The Meeting received data showing the effect of processing sunflower seed into meal and refined oil and coffee into roasted beans and instant coffee. Processing factors and residue estimates are summarized below. The Meeting could not estimate a maximum residue level for sunflower seed and did not make residue estimates for processed sunflower seed commodities.

The estimated STMR-Ps for the processed coffee commodities considered by the Meeting are shown in Table 1.

Table 1 Processing factors and residue estimates for sulfoxaflor

Raw commodity	Residue in RAC, mg/kg			Processed commodity	Processing Factors		Residue in processed commodity, mg/kg		
	Max	STMR	HR		Individual	Best estimate	Max-P	STMR-P	HR-P
Coffee (dried beans with shell)	0.3	0.035	--	Roasted beans	0.45, 0.50	0.475	--	0.0167	--
				Instant coffee	2.9, 1.9	2.4	--	0.084	--

Residues in animal commodities

The recommendations made by the current Meeting did not include animal feed items; therefore, the Meeting confirmed is previous recommendations for residues in animal commodities.

RECOMMENDATIONS

On the basis of the data obtained from supervised trials, the Meeting concluded that the residue levels listed in Annex 1 are suitable for establishing maximum residue limits and for IEDI and IESTI assessments.

Definition of the residue for compliance with the MRL and dietary risk assessment for plant and animal commodities: *sulfoxaflor*.

The residue is not fat-soluble.

DIETARY RISK ASSESSMENT

Long-term dietary exposure

The ADI for sulfoxaflor is 0–0.05 mg/kg bw. The International Estimated Daily Intakes (IEDIs) for sulfoxaflor were estimated for the 17 GEMS/Food Consumption Cluster Diets using the STMR or STMR-P values estimated by the JMPR. The results are shown in Annex 3 of the 2021 Extra JMPR Report.

The IEDIs ranged from 0–7% of the maximum ADI. The Meeting concluded that long-term dietary exposure to residues of sulfoxaflor from uses considered by the JMPR is unlikely to present a public health concern.

Acute dietary exposure

The ARfD for sulfoxaflor is 0.3 mg/kg bw. The International Estimate of Short-Term Intakes (IESTIs) for sulfoxaflor were calculated for the food commodities and their processed commodities for which HRs/HR-Ps or STMRs/STMR-Ps were estimated by the present Meeting and for which consumption data were available. The results are shown in Annex 4 of the 2021 Extra JMPR Report.

The IESTIs varied from 0–5% of the ARfD for children and 0–6% of the ARfD for the general population. The Meeting concluded that acute dietary exposure to residues of sulfoxaflor from uses considered by the present Meeting is unlikely to present a public health concern.

5.27. **TEBUCONAZOLE (189)**

RESIDUE AND ANALYTICAL ASPECTS

Tebuconazole is a triazole broad spectrum fungicide, acting by disrupting membrane function and inhibiting sterol biosynthesis. It is a systemic fungicide (primarily translocated acropetally), with protectant and curative activity against wide range of diseases including rusts, smut, bunt, powdery mildew, leaf spots and blights.

Tebuconazole was evaluated for toxicology and residues by the JMPR in 1994 and periodic reviews were conducted by the 2010 JMPR (toxicology) and the 2011 JMPR (residues). FAO Specifications for tebuconazole (technical material and related formulations) were published in 2000.

An ADI of 0–0.03 mg/kg bw and an ARfD of 0.3 mg/kg bw were established by the 2010 JMPR.

The definition of the residue for compliance with the MRL and dietary risk assessment for plant and animal commodities is: *tebuconazole*. The residue is not fat-soluble.

Tebuconazole was scheduled at the Fifty-first Session of the CCPR for evaluation of additional uses by the 2020 JMPR, which was postponed to the 2021 Extra JMPR. The Meeting received new (revised) GAP information and new supporting residue information for coffee.

Results of supervised residue trials on crops

New supervised trials from Brazil were available for the use of tebuconazole on coffee to supplement those reported by the 2008 JMPR. The analytical method used in these trials was reviewed by the 2011 JMPR and additional method validation studies for coffee bean were considered by the Meeting. The demonstrated stability of residues in frozen samples (at least 30 months) covered the storage intervals in the trials considered by the Meeting.

A new product label was available from Brazil and the Meeting noted that a number of authorised tebuconazole formulations with label claims for use on coffee were listed on the MAPA AgroFit website.

Coffee bean

In 2011 the JMPR identified the critical GAP for tebuconazole on coffee as three foliar applications of 0.25 kg ai/ha, with a PHI of 30 days and estimated a maximum residue level of 0.1 mg/kg for tebuconazole in coffee beans.

The current Meeting received new Brazilian GAP information and supporting residue trials. Noting that the critical GAP identified by the 2011 JMPR was no longer authorised, the Meeting identified a new critical GAP of 3 foliar applications of 0.2 kg ai/ha, with a 21-day re-treatment interval and a PHI of 30 days.

In trials conducted in Brazil and matching the new critical GAP, residues of tebuconazole in coffee beans were (n = 7): 0.03 (3), 0.04, 0.05, 0.09 and 0.2 mg/kg.

The Meeting estimated a maximum residue level of 0.4 mg/kg and an STMR of 0.04 mg/kg for tebuconazole in coffee beans to replace the previous maximum residue level recommendation of 0.1 mg/kg.

Fate of residues during processing

The 2011 JMPR reviewed information on the fate of tebuconazole during processing of coffee beans.

The Meeting confirmed the 2011 JMPR decision that one processing study was not sufficient to recommend a maximum residue level for coffee, roasted.

Table 1 Processing factors estimated by the 2011 JMPR for dietary exposure estimation of tebuconazole residues in coffee commodities

Raw commodity [STMR]	Processed commodity	Individual processing factors	Mean or best estimate processing factor	STMR-P [$\text{STMR}_{\text{RAC}} \times \text{PF}$] (mg/kg)	Median residue [$\text{STMR}_{\text{RAC}} \times \text{PF}$] (mg/kg)
Coffee bean [0.04 mg/kg]	Roasted beans	2	2	0.08	
	Freeze dried (instant) coffee	0.8	0.8	0.032	

RECOMMENDATIONS

On the basis of the data obtained from supervised trials, the Meeting concluded that the residue levels listed in Annex 1 are suitable for establishing maximum residue limits and for IEDI and IESTI assessments.

Definition of the residue for compliance with the MRL and dietary risk assessment for plant and animal commodities: *Tebuconazole*

The residue is not fat-soluble.

DIETARY RISK ASSESSMENT

Long-term dietary exposure

The ADI for tebuconazole is 0–0.3 mg/kg bw. The International Estimated Daily Intakes (IEDIs) for tebuconazole were estimated for the 17 GEMS/Food Consumption Cluster Diets using the STMR or STMR-P values estimated by the JMPR. The results are shown in Annex 3 of the 2021 Extra JMPR Report.

The IEDIs ranged from 1–9% of the maximum ADI. The Meeting concluded that long-term dietary exposure to residues of tebuconazole from uses considered by the JMPR is unlikely to present a public health concern.

Acute dietary exposure

The ARfD for tebuconazole is 0.3 mg/kg bw. The International Estimate of Short-Term Intakes (IESTIs) for tebuconazole were calculated for the food commodities and their processed commodities for which HRs/HR-Ps or STMRs/STMR-Ps were estimated by the present Meeting and for which consumption data were available. The results are shown in Annex 4 of the 2021 Extra JMPR Report.

The IESTIs were 0% of the ARfD for children and for the general population. The Meeting concluded that acute dietary exposure to residues of tebuconazole from uses considered by the present Meeting is unlikely to present a public health concern.

5.28. THIAMETHOXAM (245)

RESIDUE AND ANALYTICAL ASPECTS

Thiamethoxam (ISO common name) is a broad-spectrum, neonicotinoid insecticide with registered uses on multiple crops. It was evaluated for the first time by the 2010 JMPR, which established an acceptable daily intake (ADI) of 0–0.08 mg/kg bw and an acute reference dose (ARfD) of 1 mg/kg bw. Thiamethoxam underwent subsequent evaluations by the JMPR in 2011, 2012 and 2014.

The definition of the residue for compliance with the MRL for animal and plant commodities is thiamethoxam. For dietary risk assessment, the residue definitions are thiamethoxam and clothianidin (a.k.a. CGA322704), assessed separately, for plant and animal commodities except poultry and the sum of thiamethoxam, CGA 265307 (N-(2-chlorothiazol-5-ylmethyl)-N'-nitroguanidine) and MU3 (amino-([(2-chlorothiazol-5-ylmethyl)-amino]-methylene)-hydrazide), expressed as thiamethoxam, along with clothianidin (assessed separately) for poultry commodities. The residue is not fat-soluble.

Thiamethoxam was scheduled at the Fifty-first Session of the CCPR for evaluation of additional uses by the 2020 JMPR, which was postponed to the 2021 Extra JMPR. The current Meeting received information on analytical methods, field trials and processing studies to support estimation of maximum residue levels in persimmon, barley, rice, sorghum, sweet corn and wheat.

Clothianidin (238) is a metabolite of thiamethoxam and a registered active ingredient. In addition to considering residues of thiamethoxam, the Meeting also considered residues of clothianidin arising from the uses of thiamethoxam (see section 5.5 Clothianidin (238)).

Methods of analysis

The Meeting received method validation and concurrent recovery data for use of Method AG-765 (reviewed by the 2010 JMPR) and the related Method GRM.009.02A, an unnamed method for analysis of residues in Japanese persimmon, as well as the QuEChERS multiresidue method. All methods were demonstrated to have adequate performance for recovery of thiamethoxam, with an LOQ of 0.01 mg/kg (0.02 mg/kg in Japanese persimmon).

Stability of pesticide residues in stored analytical samples

No new information on the stability of thiamethoxam residues during frozen storage was provided. The 2010 JMPR had determined that residues of thiamethoxam are stable for 1–2 years under frozen (–18 °C) conditions for a large range of commodities, including the high-water and high-starch commodities considered by the current Meeting. The stability of residues in the crops under consideration by the current Meeting is considered to be adequately demonstrated for the periods that field trial samples were stored prior to analysis.

Results of supervised residue trials on crops

The Meeting received GAP information and data from supervised residue trials on persimmon, wheat, barley, rice, sorghum (grain and sweet) and sweet corn.

Persimmon, Japanese

The critical GAP for Japanese persimmon is from the Republic of Korea and consists of three applications, on a 10-day interval, of a 2000-fold dilution of the formulation (equivalent to 5 g ai/hL). Applications are made to the point of run-off and there is a 7-day PHI.

Residues of thiamethoxam in independent trials provided to the current Meeting approximating the critical GAP were (n = 2): 0.092 and 0.26 mg/kg.

The 2014 Meeting evaluated residues in Japanese persimmon in trials approximating the critical GAP. Residues of thiamethoxam in independent trials were (n = 2): 0.14 and 0.19 mg/kg.

The Meeting decided to combine the data from the two sets of trials (n = 4): 0.092, 0.14, 0.19 and 0.26 mg/kg.

The Meeting estimated a maximum residue level of 0.6 mg/kg, an STMR of 0.165 mg/kg and an HR of 0.26 mg/kg for Japanese persimmon.

Wheat

Labels were provided for registrations in Mexico (2 foliar applications at 56 g ai/ha, 7-day RTI, 14-day PHI) and the USA (seed treatment at 52 g ai/100 kg seed + 2 foliar applications at 96 g ai/ha, 5-day RTI, 30-day PHI). Residue decline data indicated that for those two GAPs, the rate is the determining factor in residues at harvest rather than the RTI or the PHI. Therefore, the Meeting decided that the GAP from the USA is the critical GAP.

Trials in the USA approximated the US GAP for application rate and timing, but with a significantly shorter interval between the last application and harvest (14 days). The Meeting decided that the data are not suitable to support recommendations based on the US GAP. The Meeting agreed to consider the GAP from Mexico.

The Meeting determined that the seed treatment application used in the trials was not likely to contribute significantly to residues at harvest and decided that the US trials can be adjusted using proportionality to approximate the GAP from Mexico.

The Meeting noted that the results from the trial in Grand Island, NE (Trial C33-0170) are inconsistent within the residue decline data (< 0.01 mg/kg in the 7, 10, 14 and 17 DALA samples and 5.0 mg/kg in the 21 DALA sample) and also in terms of the full set of available field trials (residue range = 0.015 to 0.18 mg/kg). The Meeting agreed not to consider data from that trial in any residue recommendations.

Residues (unscaled) of thiamethoxam in independent trials were (n = 16): 0.015, 0.021, 0.024, 0.032, 0.038, 0.041, 0.047, 0.052, 0.059, 0.063, 0.070, 0.091, 0.10, 0.13 (2) and 0.18 mg/kg.

When scaled (scaling factors range from 0.57 to 0.59), residues were (n = 16): < 0.01, 0.012, 0.014, 0.019, 0.022, 0.024, 0.027, 0.030, 0.034, 0.037, 0.041, 0.053, 0.058, 0.074, 0.076 and 0.10 mg/kg.

The Meeting estimated a maximum residue level of 0.15 mg/kg and an STMR of 0.032 mg/kg for wheat grain. Based on the registered use including triticale, the Meeting decided to extrapolate the residue estimates to triticale.

Barley

Labels were provided for registrations in Mexico (2 foliar applications at 56 g ai/ha, 7-day RTI, 14-day PHI) and the USA (seed treatment at 52 g ai/100 kg seed + 2 foliar applications at 96 g ai/ha, 5-day RTI, 30-day PHI). Residue decline data indicated that for those two GAPs, the rate is the determining factor in residues at harvest rather than the RTI or the PHI. Therefore, the Meeting decided that the GAP from the USA is the critical GAP.

Trials in the USA approximated the US GAP for application rate and timing, but with a significantly shorter interval between the last application and harvest (14 days). The Meeting decided that the data are not suitable to support recommendations based on the US GAP. The Meeting agreed to consider the GAP from Mexico.

The Meeting determined that the seed treatment application used in the trials was not likely to contribute significantly to residues at harvest and decided that the US trials can be adjusted using proportionality to approximate the GAP from Mexico.

Residues (unscaled) of thiamethoxam in independent trials were (n = 12): 0.030, 0.042, 0.067, 0.12 (2), 0.16, 0.22, 0.27 (2), 0.31, 0.34 and 0.53 mg/kg.

When scaled (scaling factors range from 0.57 to 0.61), residues were (n = 12): 0.017, 0.024, 0.041, 0.070 (2), 0.093, 0.13, 0.15, 0.16, 0.18, 0.20 and 0.31 mg/kg.

The Meeting estimated a maximum residue level of 0.5 mg/kg and an STMR of 0.112 mg/kg for barley grain. Based on the registered uses including oat and barley being an example representative commodity of the subgroup that includes barley and oat, the Meeting decided to extrapolate the residue estimates to oat.

Rice

The critical GAP for rice is from Japan and consists of two applications of a 1000x dilution in up to 1500 L/ha (equivalent to 97.5 g ai/ha) with a PHI of 14 days; a re-treatment interval is not specified.

Residues of thiamethoxam in husked rice from independent trials conducted in the USA and approximating the critical GAP were (n = 11): 0.92, 1.3, 1.5 (2), 1.6, 1.7, 1.8, 2.0 (3) and 2.3 mg/kg. Of those, two trials were done in paddy rice with residues of 1.7 and 2.0 mg/kg.

Since the residue populations of husked paddy rice and husked upland rice are similar, the Meeting estimated a maximum residue level of 5 mg/kg and an STMR of 1.7 mg/kg for husked rice.

Grain sorghum

The critical GAP for grain sorghum is from the USA and consists of three applications: a seed treatment at 2.97 g ai/100 kg seed followed by two in-season foliar applications, each at 96 g ai/ha, on a 7-day interval, with a PHI of 14 days.

Trials were conducted in the USA using the in-season applications only. The Meeting determined that residues of thiamethoxam coming from the seed treatment use were not likely to contribute significantly to residues at harvest. Residues of thiamethoxam in grain sorghum from independent trials approximating the critical GAP were (n = 11): 0.013, 0.021, 0.044, 0.046, 0.049, 0.079, 0.13, 0.14, 0.15, 0.26 and 0.35 mg/kg.

The Meeting estimated a maximum residue level of 0.6 mg/kg and an STMR of 0.079 mg/kg for grain sorghum.

Sweet corn

The critical GAP for sweet corn is from the USA and consists of three applications: a seed treatment at 52 g ai/100 kg seed followed by two in-season foliar applications, each at 96 g ai/ha, on a 5-day interval, with a PHI of 1 day.

Residues of thiamethoxam in sweet corn (kernel + cob with husk removed) from independent trials conducted in the USA and approximating the critical GAP were (n = 12): < 0.01 (12) mg/kg.

The Meeting noted that the only crop in the Subgroup of sweet corns is sweet corn and that the representative commodity is sweet corn (corn on the cob, kernels plus cob with husk removed) and agreed to estimate a maximum residue level of 0.01(*) mg/kg, an STMR of 0.01 mg/kg and an HR of 0.01 mg/kg for residues of thiamethoxam in the Subgroup of sweet corns (Subgroup 020F).

Grasses for sugar or syrup production

Sweet sorghum

The critical GAP for sweet sorghum is from the USA and consists of three applications: a seed treatment at 2.97 g ai/100 kg seed followed by two in-season foliar applications, each at 96 g ai/ha, on a 7-day interval, with a PHI of 14 days.

Trials were conducted in the USA using the in-season applications only. The Meeting determined that residues of thiamethoxam coming from the seed treatment use were not likely to contribute significantly to residues at harvest. Residues of thiamethoxam in sweet sorghum cane from independent trials approximating the critical GAP were (n = 4): < 0.01, < 0.01, 0.056 and 0.24 mg/kg.

The Meeting estimated a maximum residue level of 0.6 mg/kg, an STMR of 0.033 mg/kg and an HR of 0.24 mg/kg for sorgho or sorghum, sweet.

Residues in animal feeds**Forages of cereal grains****Wheat**

The critical GAP for forage is from the USA (seed treatment at 52 g ai/100 kg seed + 2 foliar applications at 96 g ai/ha, 5-day RTI, 7-day PHI).

Residues in forage from trials in the USA approximating the US GAP were (n = 17): < 0.015, 0.011, 0.023, 0.03, 0.032, 0.036, 0.072, 0.079, 0.12, 0.18, 0.33 (2), 0.46, 0.73, 1.1 (2) and 2.0 mg/kg.

The registration in the USA includes barley, oat, rye, triticale and wheat. Based on the residue data (as received) for wheat forage listed above, the Meeting estimated a median residue of 0.12 mg/kg and a highest residue of 2 mg/kg for wheat forage. The Meeting agreed to extrapolate the median and highest residue estimates from wheat forage to forage of barley, oat, rye and triticale.

Hay of cereal grains**Wheat**

The critical GAP for hay is from the USA (seed treatment at 52 g ai/100 kg seed + 2 foliar applications at 96 g ai/ha, 5-day RTI, 7-day PHI).

Residues in wheat hay from trials in the USA approximating the US GAP were (n = 17): 0.014, 0.015, 0.034, 0.036, 0.049, 0.054, 0.058, 0.16, 0.20, 0.32, 0.41, 0.52, 0.91, 1.5, 2.0, 2.1 and 6.7 mg/kg.

Barley

The critical GAP for hay is from the USA (seed treatment at 52 g ai/100 kg seed + 2 foliar applications at 96 g ai/ha, 5-day RTI, 7-day PHI).

Residues in barley hay from trials in the USA approximating the US GAP were (n = 12): 0.023, 0.058, 0.083, 0.12, 0.17, 0.19, 0.25, 0.34, 0.39, 0.42, 0.46 and 0.52 mg/kg.

The registration in the USA includes barley, oat, rye, triticale and wheat. The Meeting determined via a Mann-Whitney U Test that the residue data listed above for wheat and barley hay are not statistically different and agreed to combine the data. Residues of thiamethoxam in wheat and barley hay following treatment approximating the US GAP were (n = 29): 0.014, 0.015, 0.023, 0.034, 0.036, 0.049, 0.054, 0.058, 0.058, 0.083, 0.12, 0.16, 0.17, 0.19, 0.20, 0.25, 0.32, 0.34, 0.39, 0.41, 0.42, 0.46, 0.52, 0.52, 0.91, 1.5, 2.0, 2.1 and 6.7 mg/kg.

Based on the registered uses and on the data for wheat and barley hay and on an OECD standard dry-matter content of 88%, the Meeting estimated a maximum residue level of 8 mg/kg (dw), a median residue of 0.2 mg/kg (as received) and a highest residue of 6.7 mg/kg (as received) in hay of wheat, barley, oat, rye and triticale.

Straw of cereal grains**Wheat**

Labels were provided for registrations in Mexico (2 foliar applications at 56 g ai/ha, 7-day RTI, 14-day PHI) and the USA (seed treatment at 52 g ai/100 kg seed + 2 foliar applications at 96 g ai/ha, 5-day RTI, 30-day PHI).

For wheat straw, the trials in the USA did not match the USA GAP due to the straw being harvested 14 DALA rather than the 30 days specified for the PHI. Therefore, the Meeting decided that the data are not suitable to support recommendations based on the USA GAP. The Meeting agreed to consider the GAP from Mexico. As noted for wheat grain, the Meeting determined that residues of

thiamethoxam coming from the seed treatment use were not likely to contribute significantly to residues at harvest and that residues from the USA trials can be adjusted using proportionality to approximate the GAP from Mexico.

Residues (unscaled) of thiamethoxam in wheat straw from independent trials were (n = 17): 0.14, 0.20, 0.21, 0.25, 0.43, 0.49, 0.75, 0.84, 0.92, 1.6, 1.7 (3), 1.9, 2.4, 3.3 and 3.9 mg/kg.

When scaled (scaling factors range from 0.57 to 0.59), residues in wheat straw were (n = 17): 0.081, 0.12, 0.12, 0.15, 0.25, 0.28, 0.43, 0.49, 0.53, 0.93, 0.98, 0.99, 1.0, 1.1, 1.4, 1.9 and 2.2 mg/kg.

Barley

Labels were provided for registrations in Mexico (2 foliar applications at 56 g ai/ha, 7-day RTI, 14-day PHI) and the USA (seed treatment at 52 g ai/100 kg seed + 2 foliar applications at 96 g ai/ha, 5-day RTI, 30-day PHI).

For barley straw, the trials in the USA did not match the USA GAP due to the straw being harvested 14 DALA rather than the 30 days specified for the PHI. Therefore, the Meeting decided that the data are not suitable to support recommendations based on the US GAP. The Meeting agreed to consider the GAP from Mexico. As noted for barley grain, the Meeting determined that residues of thiamethoxam coming from the seed treatment use were not likely to contribute significantly to residues at harvest and that residues from the US trials can be adjusted using proportionality to approximate the GAP from Mexico.

Residues (unscaled) of thiamethoxam in barley straw from independent trials were (n = 12): 0.014, 0.016, 0.073, 0.078, 0.24, 0.25, 0.25, 0.26, 0.49, 0.53, 0.54, 2.3 mg/kg.

When scaled (scaling factors range from 0.57 to 0.59), residues in barley straw were (n = 12): < 0.01 (2), 0.042, 0.046, 0.14 (2), 0.15, 0.16, 0.28, 0.31 (2) and 1.3 mg/kg.

The registration in Mexico includes barley, oat, triticale and wheat. Noting that residues appear to be similar between the straw data for wheat and barley, the Meeting agreed to explore a common set of residue estimates for these straw commodities. A Mann-Whitney U-Test indicated that the residue distributions from wheat and barley are significantly different. The OECD MRL calculator suggested maximum residue levels of 4 mg/kg (median = 0.53 mg/kg) for wheat straw and 2 mg/kg for barley (median = 0.145 mg/kg), all on an as-received basis.

The Meeting noted that the maximum residue level and highest residue estimates in hay are greater than those in straw and agreed to apply those estimates to straw and to use the more conservative median residue from wheat for estimating dietary burdens. Furthermore, noting that the straw commodities of different cereal crops are generally indistinguishable from each other when in trade, the Meeting agreed to make recommendations for the straw of cereals included on the Mexico label (barley, oat, triticale, wheat).

Based on OECD standard dry-matter content of 88%, the Meeting estimated a maximum residue level of 8 mg/kg (dw), a median residue of 0.53 mg/kg (as received) and a highest residue of 6.7 mg/kg (as received) for residues of thiamethoxam in straw and fodder, dry of wheat, barley, oat and triticale.

The above recommendations replace the previous recommendations, each at 2 mg/kg, for barley straw and fodder, dry and wheat straw and fodder, dry.

Rice

The critical GAP for rice is from Japan and consists of two applications of a 1000x dilution in up to 1500 L/ha (equivalent to 97.5 g ai/ha) with a PHI of 14 days; a re-treatment interval is not specified.

Residues of thiamethoxam in rice straw from independent trials approximating the critical GAP were (n = 11): 0.2 (2), 0.3, 0.36(2), 0.47, 0.51, 0.54, 0.55 and 1.0 (2) mg/kg.

Based on OECD dry-matter content of 88%, the Meeting estimated a maximum residue level of 3 mg/kg (dw), a median residue of 0.47 mg/kg (as received) and a highest residue of 1 mg/kg (as received) for residues of thiamethoxam in rice straw and fodder, dry.

Sorghum

The critical GAP for sorghum is from the USA and consists of three applications: a seed treatment at 2.97 g ai/100 kg seed followed by two in-season foliar applications, each at 96 g ai/ha, on a 7-day interval, with a PHI of 14 days.

Trials were conducted in the USA using the in-season applications only. The Meeting determined that residues of thiamethoxam coming from the seed treatment use were not likely to contribute significantly to residues at harvest.

Residues of thiamethoxam in sorghum forage from independent trials approximating the critical GAP were (n = 10): < 0.01, 0.016, 0.018, 0.028, 0.036, 0.083, 0.12, 0.15, 0.17 and 0.30 mg/kg (as received).

The Meeting estimated a median residue of 0.06 mg/kg (as received) and a highest residue of 0.3 mg/kg (as received) in sorghum forage.

Residues of thiamethoxam in sorghum stover from independent trials approximating the critical GAP were (n = 10): 0.075, 0.076, 0.077, 0.087, 0.12, 0.15, 0.16, 0.19, 0.35 and 0.46 mg/kg (as received).

Based on OECD standard dry-matter content of 88%, the Meeting estimated a maximum residue level of 0.8 mg/kg (dw), a median residue of 0.14 mg/kg (as received) and a highest residue of 0.49 mg/kg (as received) in sorghum straw and fodder (dry).

Sweet corn

The critical GAP for sweet corn is from the USA and consists of three applications: a seed treatment at 52 g ai/100 kg seed followed by two in-season foliar applications, each at 96 g ai/ha, on a 5-day interval, with a PHI of 1 day for forage and 26 days for stover.

Residues of thiamethoxam in sweet corn forage from independent trials approximating the critical GAP were (n = 12): 0.39, 0.73, 1.0, 1.3, 1.4, 1.4, 1.6, 1.6, 1.6, 1.9, 1.9 and 2.8 mg/kg (as received).

The Meeting estimated a median residue of 1.5 mg/kg (as) and a highest residue of 2.8 mg/kg (as received) in sweet corn forage.

Residues of thiamethoxam in sweet corn stover from independent trials approximating the critical GAP were (n = 4): 0.010, 0.011, 0.022 and 0.099 mg/kg (as received).

Based on OECD standard dry-matter content of 83%, the Meeting estimated a maximum residue level of 0.25 mg/kg (dw), a median residue of 0.0165 mg/kg (as received) and a highest residue of 0.099 mg/kg (as received) in sweet corn stover.

Fate of residues during processing

The Meeting received data showing the effect of processing wheat grain into aspirated grain fractions, bran, flour, middlings, shorts and germ; rice into hulls, bran and polished rice; grain sorghum into aspirated grain fractions and flour; and sweet sorghum into syrup. Processing factors and residue estimates are summarized below.

Table 1 Processing factors and residue estimates for thiamethoxam in wheat, sorghum and sweet sorghum commodities

Raw commodity	Residue in RAC, mg/kg			Processed commodity	Processing Factors		Residue in processed commodity, mg/kg		
	Max	STMR	HR		Individual	Best estimate	Max-P	STMR-P	HR-P
Wheat grain	0.15	0.032	--	Aspirated fractions	9.2, 20	14.6	--	0.467	--
				Bran	2.1, 2.7	2.4	0.36	0.0768	--
				Flour	0.10, 0.11	0.105	--	0.00336	--
				Middlings	0.30, 0.43	0.365	--	0.0117	--
				Shorts	0.39, 0.50	0.445	--	0.0142	--
				Germ	1.4, 1.5	1.45	0.218	0.0464	--
Sorghum grain	0.6	0.079	--	Aspirated fractions	11, 12	11.5	--	0.908	--
				Flour	0.73, 0.75	0.74	--	0.0548	--
Sorghum cane	0.6	0.033	--	Syrup	0.81	0.81	--	0.0267	--

Based on the maximum residue estimates above and the OECD rounding classes, the Meeting estimated maximum residue levels of 0.4 mg/kg for wheat bran and 0.3 mg/kg for wheat germ. The Meeting decided to apply the processing factor for wheat bran (2.4) to barley bran, giving a maximum residue level of 1.5 mg/kg. The estimated STMR-Ps for the processed commodities considered by the Meeting are shown in Table 1. Furthermore, the Meeting decided to apply the processing factors for wheat bran (2.4) and flour (0.105) to barley and triticale, giving STMR-Ps as follows:

Barley bran = 0.269 mg/kg,

Barley flour = 0.0118 mg/kg,

Triticale flour = 0.00336 mg/kg.

For rice commodities, the processing study provided residue analysis for rice grain, rice hulls, rice bran and polished rice, but not for husked rice. The Meeting derived a processing factor for husked rice by mathematically reconstituting the commodity based on residues in bran, residues in polished rice and their respective material balances reported in the processing study. Conversely, the residue trials in rice only reported results for husked rice. Processing factors describing residues relative to the rice RAC (grain) and from husked rice are shown in Table 2.

Table 2 Processing factors and residue estimates for thiamethoxam in rice commodities

Commodity	Residue in starting commodity, mg/kg			Processed commodity	Processing Factors		Residue in processed commodity, mg/kg		
	Max	STMR	HR		Individual	Best estimate	Max-P	STMR-P	HR-P
Rice grain	See text below			Husked rice	0.069, 0.14	0.104	No field trial data were provided for rice grain		
				Hulls	4.6, 6.4	5.5			
				Bran	0.39, 0.75	0.57			
				Polished rice	0.032, 0.051	0.0415			
Husked Rice	5 ^a	1.7 ^a		Hulls	66, 45	56	280	95.2	--
				Bran	5.6, 5.4	5.5	27.5	9.35	--
				Polished rice	0.46, 0.37	0.41	2.07	0.704	--

^a From field trials

For rice, the Meeting estimated maximum residue levels of 300 mg/kg for rice hulls, 30 mg/kg for rice bran and 3 mg/kg for polished rice. Furthermore, the Meeting used the residue estimates and the processing factor from rice grain to husked rice to derive residue estimates for rice grain. The Meeting estimated a maximum residue level of 50 mg/kg and a median residue of 17 mg/kg for rice grain.

Residues in animal commodities

The Meeting has added feed items and their associated residues to the dietary burden calculation used by the 2014 JMPR. Dietary burden calculations are provided in Annex 6; the dietary burden estimates are summarized below.

Table 3 Estimated maximum and mean dietary burdens of farm animals

Animal	Dietary burden estimates, ppm							
	Canada-US		European Union		Australia		Japan	
	Maximum	Mean	Maximum	Mean	Maximum	Mean	Maximum	Mean
Beef cattle	6.89	5.55	7.16	1.63	18.2	17.3	2.72	2.40
Dairy cattle	9.73	6.87	9.45	4.16	20.6 ^{a b}	18.8 ^c	1.83	1.26
Broiler poultry	5.0	5.0	1.39	1.16	11.8	11.8	0.59	0.59
Laying hen	5.0	5.0	2.62	1.10	11.8 ^d	11.8 ^e	2.13	2.13

^a Highest maximum cattle dietary burden suitable for MRL estimates for mammalian tissues.

^b Highest maximum dairy cattle dietary burden suitable for MRL estimates for milk

^c Highest mean cattle dietary burden suitable for STMR estimates for mammalian tissues and milk

^d Highest maximum poultry dietary burden suitable for MRL estimates for poultry tissues and eggs

^e Highest mean poultry dietary burden suitable for STMR estimates for poultry tissues and eggs

Farm animal feeding studies

Cattle

A cattle-feeding study with thiamethoxam was reviewed by the 2010 JMPR. Animals were dosed at 2, 6, or 20 ppm in the diet. Thiamethoxam residues were as follows:

Matrix	2 ppm in feed		6 ppm in feed		20 ppm in feed	
	Maximum	Mean	Maximum	Mean	Maximum	Mean
Muscle	< 0.01 mg/kg	< 0.01 mg/kg	0.01 mg/kg	< 0.01 mg/kg	0.06 mg/kg	0.035 mg/kg
Fat	Not analysed		Not analysed		< 0.01	< 0.01
Kidney	< 0.01	< 0.01	< 0.01	< 0.01	0.04	0.0266
Liver	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01	< 0.01
Milk	0.01	0.0073	0.05	0.036	0.19	0.124

Interpolation between the 6-ppm and 20-ppm feeding levels gives the anticipated residues in mammalian commodities shown below.

Thiamethoxam feeding study	Feed level (ppm) for milk/egg residues	Residues (mg/kg) in milk	Feed level (ppm) for tissue residues	Residues (mg/kg)			
				Muscle	Liver	Kidney	Fat
MRL beef or dairy cattle							
Feeding study	20	0.124	20	0.06	< 0.01	0.04	< 0.01
Dietary burden and high residue	20.6	0.128	20.6	0.062	< 0.01	0.041	< 0.01
STMR beef or dairy cattle							
Feeding study	6	0.036	6	< 0.01	< 0.01	< 0.01	< 0.01
	20	0.124	20	0.035	< 0.01	0.0266	< 0.01
Dietary burden and residue estimate	18.8	0.096	18.8	0.033	< 0.01	0.025	< 0.01

The Meeting estimated new maximum residue levels, replacing the previous recommendations for meat, milks and edible offal as follows: Meat (from mammals other than marine mammals) at 0.07 mg/kg, Milks at 0.15 mg/kg, Edible offal (mammalian) at 0.05 mg/kg and Mammalian fats (except milk fats) at 0.01(*) mg/kg.

Poultry

A poultry-feeding study for thiamethoxam is not available. In the laying hen metabolism study reviewed by the 2010 JMPR, test animals were dosed at 98 or 112 ppm thiamethoxam in the feed.

To evaluate residues relevant for compliance with the MRL and for dietary risk assessment, the Meeting used the highest maximum and mean dietary burden for poultry (both at 11.8 ppm) and the 98-ppm dose level to estimate residues in poultry commodities. Concentrations of thiamethoxam (residue definition for compliance with the MRL) were up to 0.195 mg/kg in muscle, 0.0435 mg/kg in fat+skin, 0.0184 mg/kg in liver and 0.033 mg/kg in eggs. The resulting estimated residues are 0.023 mg/kg in muscle, 0.0052 mg/kg in fat, 0.0022 mg/kg in liver and 0.004 mg/kg in eggs.

The Meeting confirmed its previous recommendations of 0.01(*) mg/kg for poultry edible offal and eggs, estimated a new maximum residue level of 0.01(*) mg/kg for poultry fat and estimated a new maximum residue level of 0.03 mg/kg for poultry meat to replace its previous recommendation.

In the metabolism study concentrations of thiamethoxam + CGA 265307 + MU3, expressed as thiamethoxam (residue definition for dietary assessment), were up to 0.534 mg/kg in muscle, 0.276 mg/kg in fat+skin, 2.96 mg/kg in liver and 0.232 mg/kg in eggs. Estimated HRs and STMRS are 0.064 mg/kg in muscle, 0.033 mg/kg in fat, 0.36 mg/kg in liver and 0.028 mg/kg in eggs.

RECOMMENDATIONS

On the basis of the data obtained from supervised trials, the Meeting concluded that the residue levels listed in Annex 1 are suitable for establishing maximum residue limits and for IEDI and IESTI assessments.

Definition of the residue for compliance with the MRL and dietary risk assessment for plant commodities: *thiamethoxam*.

Definition of the residue for compliance with the MRL and dietary risk assessment for animal commodities (except poultry): *thiamethoxam and clothianidin* (considered separately).

Definition of the residue for dietary risk assessment for poultry: *sum of thiamethoxam, CGA 265307 and MU3, expressed as thiamethoxam and clothianidin* (clothianidin considered separately).

The residue is not fat-soluble.

DIETARY RISK ASSESSMENT

Long-term dietary exposure

The ADI for thiamethoxam is 0–0.08 mg/kg bw. The International Estimated Daily Intakes (IEDIs) for thiamethoxam were estimated for the 17 GEMS/Food Consumption Cluster Diets using the STMR or STMR-P values estimated by the JMPR. The results are shown in Annex 3 of the 2021 Extra JMPR Report.

The IEDIs ranged from 1–7% of the maximum ADI. The Meeting concluded that long-term dietary exposure to residues of thiamethoxam from uses considered by the JMPR is unlikely to present a public health concern.

Acute dietary exposure

The ARfD for thiamethoxam is 1.0 mg/kg bw. The International Estimate of Short-Term Intakes (IESTIs) for thiamethoxam were calculated for the food commodities and their processed

commodities for which HRs/HR-Ps or STMRs/STMR-Ps were estimated by the present Meeting and for which consumption data were available. The results are shown in Annex 4 of the 2021 Extra JMPR Report.

The IESTIs varied from 0–5% of the ARfD for children and 0–3% of the ARfD for the general population. The Meeting concluded that acute dietary exposure to residues of thiamethoxam from uses considered by the present Meeting is unlikely to present a public health concern.

5.29. TRIFLOXYSTROBIN (213)

RESIDUE AND ANALYTICAL ASPECTS

Trifloxystrobin is a strobilurin broad-spectrum contact fungicide that was first evaluated for toxicology and residues by the JMPR in 2004. The Meeting derived an ADI of 0–0.04 mg/kg bw and concluded that an ARfD is unnecessary. The residue definition for compliance with the MRL for plant commodities is trifloxystrobin *per se* and for compliance with the MRL for animal commodities as well as for dietary risk assessment for both plant and animal commodities is the sum of trifloxystrobin and (E,E)-methoxyimino-{2-[1-(3-trifluoromethyl-phenyl) ethylidene-aminooxymethyl]-phenyl}acetic acid = CGA321113 (expressed as trifloxystrobin equivalents). The residue is fat-soluble.

Trifloxystrobin was scheduled at the Fifty-first Session of the CCPR for the evaluation of additional uses by the 2020 JMPR, which was postponed to the 2021 Extra JMPR, for citrus fruits, cane berries, bush berries, leaf lettuce, legume vegetables, dry beans and peas, tree nuts, linseed and coffee.

Methods of analysis

Methods 01013, 01313, 00765 and 200177, which were previously evaluated by the 2004, 2012 and 2015 JMPR, were used for the analysis of trifloxystrobin and CGA321113 in citrus fruits, tree nuts, head lettuce, coffee beans, fresh beans and dry peas. The current Meeting received study reports and method validation data for method 01313/M001 and QuEChERS multiresidue method 01207 which were used for the analysis of trifloxystrobin and CGA321113 in raspberries, currants, lettuce, peas and beans. The current Meeting also received concurrent recovery data for analytical methods GIR/MET/TRIFLOXY/01, TF-003-P16-01 and 00742/E001 for the analysis of trifloxystrobin and CGA321113 in raspberries, currants, flax and beans with pods. In general, the new methods considered by the current Meeting involved extraction with acetonitrile/water and final determination was achieved using LC-MS/MS. Among all available methods, the validated LOQ for trifloxystrobin and CGA321113 ranged from 0.005–0.02 mg/kg for each analyte. Mean recoveries were, with a few exceptions, within the acceptable range of 70–120% with a RSD of < 20%.

The Meeting concluded that for the commodities considered by the Meeting, the methods used in the new residue trials were sufficiently validated and suitable to measure trifloxystrobin and CGA321113 in plant commodities.

Stability of pesticide residues in stored analytical samples

The stability of residues in samples on frozen storage was evaluated by the 2004 JMPR for a range of commodities. Samples in the trials considered by the current Meeting were stored frozen for periods less than the period of stability demonstrated in studies supplied to the 2004 JMPR and were satisfactory.

Results of supervised residue trials on crops

The Meeting received information on supervised field trials on citrus (orange, lemon, grapefruit), raspberry, currant, leaf lettuce, beans (fresh), peas (fresh and dried), tree nuts (almond and pecan), linseed and coffee.

Residues for maximum residue level estimation are expressed in mg trifloxystrobin/kg. Residues for dietary risk assessment include parent trifloxystrobin and metabolite CGA321113. The sum of trifloxystrobin and CGA321113 was calculated and expressed as trifloxystrobin on the basis of the relative molecular masses. A conversion factor of 1.036 is required to express CGA321113 as trifloxystrobin. As CGA321113 does not generally constitute a significant proportion of the residue in crops, when the levels of trifloxystrobin or CGA321113 were below the LOQ, their sum was calculated as in the examples provided by the 2004 JMPR and copied below.

Trifloxystrobin (mg/kg)	CGA321113 (mg/kg)	Total (expressed as trifloxystrobin) (mg/kg)
< 0.02	< 0.02	< 0.02
< 0.02	0.03	0.05
0.10	< 0.02	0.10
0.92	0.16	1.1

Citrus Fruits

Citrus fruits were previously evaluated by the 2004 JMPR where a maximum residue level of 0.5 mg/kg and an STMR of 0.095 mg/kg were estimated based on the GAP from the USA, comprising 4 foliar applications at 140 g ai/ha with a PHI of 30 days. This recommendation was based on residue trials from the USA on oranges, lemons and grapefruits (4 × 170 g ai/ha, 26–32-day PHIs).

The current Meeting received a new critical GAP for the USA involving a shorter PHI of 7 days and a minimum re-treatment interval of 7 days. No trials provided to the Meeting matched this GAP with respect to the RTI. The RTI used in the citrus supervised residue trials provided to the Meeting (i.e. 21-days) is significantly longer than the 7-day RTI from the critical GAP. Residue decline data provided and available to the Meeting for citrus fruits indicated that the trials underestimate residues expected in citrus fruits by more than 25% when treated according to the critical GAP.

As the Meeting received a new critical GAP that was not supported by the available field trials and no suitable GAP from another country was provided, following current assessment practices the Meeting withdrew its previous recommendation for citrus fruits.

Cane berries

Raspberries

The critical GAP for protected cane berries in the Netherlands and field grown raspberries and blackberries in Austria and Portugal is 2 foliar applications (SC formulation) × 200 g ai/ha, 7-day RTI and 3-day PHI.

The Meeting received supervised residue trials conducted in Europe on raspberries (protected and unprotected) matching the critical GAP.

For estimation of maximum residue levels, residue levels of trifloxystrobin (parent only) in ranked order were:

Raspberry (protected) (n = 10): 0.086, 0.17, 0.25, 0.33, 0.48, 0.51, 0.52, 0.57, 0.98 and 1.4 mg/kg.

Raspberry (unprotected) (n = 8): 0.098, 0.20, 0.35, 0.36, 0.64, 0.70, 1.0 and 1.5 mg/kg.

For estimation of STMRs, total residue levels of trifloxystrobin and CGA321113 in ranked order were:

Raspberry (protected) (n = 10): 0.10, 0.18, 0.29, 0.38, 0.53, 0.59, 0.60, 0.62, 1.1 and 1.4 mg/kg.

Raspberry (unprotected) (n = 8): 0.12, 0.23, 0.37, 0.39, 0.71, 0.80, 1.1 and 1.5 mg/kg.

The Mann-Whitney U-test determined that the protected and unprotected raspberry datasets were not statistically different. Therefore, the Meeting decided to combine the datasets of protected and unprotected raspberries.

Trifloxystrobin (parent only) residues in raspberries (protected and unprotected) in ranked order were (n = 18): 0.086, 0.098, 0.17, 0.20, 0.25, 0.33, 0.35, 0.36, 0.48, 0.51, 0.52, 0.57, 0.64, 0.70, 0.98, 1.0, 1.4 and 1.5 mg/kg.

Total residue levels of trifloxystrobin and CGA321113 (protected and unprotected) in raspberries in ranked order were (n = 18): 0.10, 0.12, 0.18, 0.23, 0.29, 0.37, 0.38, 0.39, 0.53, 0.59, 0.60, 0.62, 0.71, 0.80, 1.1 (2), 1.4 and 1.5 mg/kg.

Noting that raspberry is a representative commodity for the cane berries subgroup and that the Netherlands GAP includes all commodities in this subgroup, the Meeting estimated a maximum residue level of 3 mg/kg and an STMR value of 0.56 mg/kg for the subgroup of cane berries.

Bush berries

Currants

The critical GAP for protected berries (including currants, gooseberries, blueberries, bilberries, foxberries and rose hips) from the Netherlands and field-grown currants, gooseberries and blueberries from Austria and Spain is 2 foliar applications (SC formulation) × 200 g ai/ha, 7-day RTI and 7-day PHI.

The Meeting received supervised residue trials conducted in Europe on currants (protected and unprotected) matching the critical GAP.

For estimation of maximum residue levels, residue levels of trifloxystrobin (parent only) in ranked order were:

Currants (protected) (n = 8): 0.15, 0.27, 0.31 (2), 0.35, 0.36, 0.51 and 0.77 mg/kg.

Currants (unprotected) (n = 8): 0.13, 0.14, 0.22, 0.29, 0.46, 0.64, 1.2 and 2.1 mg/kg.

For estimation of STMRs, total residue levels of trifloxystrobin and CGA321113 in ranked order were:

Currants (protected) (n = 8): 0.15, 0.27, 0.31 (2), 0.35, 0.36, 0.51 and 0.77 mg/kg.

Currants (unprotected) (n = 8): 0.13, 0.14, 0.22, 0.29, 0.46, 0.65, 1.2 and 2.1 mg/kg.

The Mann-Whitney U-test determined that the protected and unprotected currants datasets were not statistically different. Therefore, the Meeting decided to combine the datasets of protected and unprotected currants.

Trifloxystrobin (parent only) residues in currants (protected and unprotected) in ranked order were (n = 16): 0.13, 0.14, 0.15, 0.22, 0.27, 0.29, 0.31 (2), 0.35, 0.36, 0.46, 0.51, 0.64, 0.77, 1.2 and 2.1 mg/kg.

Total residue levels of trifloxystrobin and CGA321113 in currants in ranked order were (n = 16): 0.13, 0.14, 0.15, 0.22, 0.27, 0.29, 0.31 (2), 0.35, 0.36, 0.46, 0.51, 0.65, 0.77, 1.2 and 2.1 mg/kg.

Noting that currants is a representative commodity for the bush berries subgroup and that the Netherlands GAP includes currants, gooseberries, blueberries, bilberries, foxberries (red bilberries) and rose hips, the Meeting estimated a maximum residue level of 3 mg/kg and an STMR value of 0.33 mg/kg for currants, blueberries, gooseberries, billberries, red bilberries and rose hips.

Lettuce, leaf

The critical GAP for lettuce is from Austria on protected lettuce; 2 foliar applications (SC formulation) at 200 g ai/ha, 7-day RTI, with a 7-day PHI. The Meeting received supervised residue trials conducted in Europe on protected leaf lettuce matching the critical GAP.

For estimation of maximum residue levels, residue levels of trifloxystrobin (parent only) in ranked order were:

Lettuce, leaf (protected) (n = 13): 0.85, 1.2, 1.3, 1.9, 2.6, 3.1, 3.2, 3.5, 3.8, 4.5, 6.2, 8.6 and 9.9 mg/kg.

For estimation of STMRs, total residue levels of trifloxystrobin and CGA321113 in ranked order were:

Lettuce, leaf (protected) (n = 13): 0.90, 1.3 (2), 1.9, 2.8, 3.2, 3.3, 3.5, 3.8, 4.7, 6.7, 8.7 and 10 mg/kg.

The Meeting estimated a maximum residue level of 15 mg/kg and STMR of 3.3 mg/kg for leaf lettuce. The Meeting noted that the GAP in Austria also covers protected lambs lettuce (corn salad). As leaf lettuce is a representative leafy greens commodity, the Meeting agreed to extrapolate the maximum residue level of 15 mg/kg and an STMR of 3.3 mg/kg to corn salad based on the dataset of protected leaf lettuce.

Legume vegetables

Subgroup of beans with pods:

The critical GAP for kidney beans (common bean (poroto)) is from Austria; 2 foliar applications (SC formulation) at 200 g ai/ha, with a 7-day RTI and 7-day PHI.

The Meeting received supervised residue trials conducted in Europe on beans with pods matching the critical GAP.

Trifloxystrobin (parent only) residues in beans with pods in ranked order were (n = 13): 0.02, 0.03 (2), 0.038, 0.045, 0.063, 0.079, 0.09, 0.12, 0.16, 0.21 (2) and 0.34 mg/kg.

Total residue levels of trifloxystrobin and CGA321113 in beans with pods in ranked order were (n = 13): 0.03 (2), 0.04, 0.051, 0.059, 0.063, 0.09, 0.099, 0.12, 0.19, 0.23, 0.26 and 0.35 mg/kg.

The Meeting considered the data would support a maximum residue level of 0.5 mg/kg and an STMR of 0.09 mg/kg for common bean (poroto).

The Meeting noted a GAP for the subgroup of beans with pods from the Netherlands; 2 foliar applications (SC formulation) × 200 g ai/ha, 14-day RTI and 14-day PHI.

The Meeting also received supervised residue trials conducted in Europe on beans with pods approximating the Netherlands GAP but with RTIs of 7-days. Although the residue trials have a shorter RTI compared to the Netherlands GAP, based on residue decline data for beans with pods the Meeting estimated that residues from the field trials would be within 25% of the Netherlands GAP.

Trifloxystrobin (parent only) residues in beans with pods in ranked order were (n = 10): < 0.01 (2), 0.012, 0.02 (2), 0.023, 0.046, 0.05, 0.09 and 0.34 mg/kg.

Total residue levels of trifloxystrobin and CGA321113 in beans with pods in ranked order were (n = 10): < 0.01 (2), 0.012, 0.02 (2), 0.023, 0.05, 0.064, 0.09 and 0.35 mg/kg.

The Meeting considered the data would support a maximum residue level of 0.5 mg/kg and an STMR of 0.022 mg/kg based on the Netherlands GAP.

Noting that the above estimates are the same the Meeting decided to recommend a maximum residue level of 0.5 mg/kg and an STMR of 0.09 mg/kg for the subgroup of beans with pods.

Subgroup of peas with pods

Based on the available residue data, the Meeting concluded that the critical GAP for peas with pods is from France; 1 foliar application using an SC formulation, at 200 g ai/ha, with a PHI of 7 days.

The Meeting received supervised residue trials conducted in Europe on peas with pods, where 2 foliar spray applications (SC formulation) were made at a nominal rate of 200 g ai/ha per application and a 7-day RTI. In 13 trials, peas with pods were harvested at a “-0” day PHI (i.e. just before the last application; 7 days after the first application) which matches the critical GAP from France. As the level of CGA321113 was below the LOQ in all samples, the data populations for enforcement and risk assessment purposes are identical.

Trifloxystrobin (and total trifloxystrobin and CGA321113) residues in peas with pods in ranked order were (n = 13): 0.021, 0.035, 0.037, 0.046, 0.054, 0.063, 0.073, 0.081, 0.19, 0.20, 0.24, 0.30 and 0.86 mg/kg.

The Meeting estimated a maximum residue level of 1.5 mg/kg and an STMR value of 0.073 mg/kg for the subgroup of peas with pods.

Subgroup of succulent beans without pods

The critical GAP for succulent kidney beans without pods is from Austria; 2 foliar applications (SC formulation) at 200 g ai/ha, with a 7-day RTI and a 7-day PHI.

The Meeting received two supervised residue trials conducted in Europe on beans without pods that differed from the critical GAP with regards to the pre-harvest interval. As there were no trials conducted in accordance with the critical GAP and there was an insufficient number of trials to support any other GAP provided for succulent beans without pods, the Meeting did not estimate a maximum residue level or STMR for the subgroup of succulent beans without pods.

Subgroup of succulent peas without pods

The critical GAPs for succulent peas without pods are from Austria and the Netherlands; 2 foliar applications (SC formulation) at 200 g ai/ha, with a 14-day RTI and a 14-day PHI.

The Meeting received supervised residue trials conducted in Europe on peas without pods that differed from the critical GAP with regards to the re-treatment interval (i.e. 7-day RTIs in the trials versus the 14-day RTI from the critical GAP). The residue decline data for succulent peas without pods were insufficient to determine whether the residues in the trials approximated residues according to the critical GAPs.

As there were no trials conducted in accordance with the critical GAP or any other GAP provided for succulent peas without pods, the Meeting did not estimate a maximum residue level or STMR for the subgroup of succulent peas without pods.

Pulses

Subgroup of dry beans and Subgroup of dry peas

The critical GAP for the subgroup of dry beans and subgroup of dry peas is from Austria and the Netherlands; 2 foliar applications (SC formulation) at 200 g ai/ha, with a 14-day RTI and a 14-day PHI.

The Meeting received supervised residue trials conducted in Europe on dry peas, where 2 foliar applications (SC formulation) were made at a nominal rate of 200 g ai/ha per application and 7-day RTI. Dry pea seeds were harvested at PHIs of 14, 20–22 and 28–48 days. Only three trials were provided in which peas without pods were harvested at a 14-day PHI which are insufficient to support a recommendation based on the Austria and the Netherlands GAPs.

As none of the European dry pea trials reflected the critical GAP or any other GAP provided for dry beans or peas, the Meeting could not estimate a maximum residue level or STMR for the subgroup of dry beans or the subgroup of dry peas.

Tree nuts

The critical GAP for tree nuts is from the USA; 4 foliar applications (SC formulation) at 140 g ai/ha, with a 7-day RTI and a 14-day PHI. The Meeting received supervised residue trials conducted in the USA on almonds and pecans where 3 foliar spray applications (SC formulation) were made at a nominal rate of 160 g ai/ha (first application) and 200 g ai/ha (second and third applications), 7-day RTI and a 14-day PHI.

As none of the USA tree nut trials reflected the critical GAP, with regard to application rate and number of applications and the proportionality approach could not be considered, the Meeting was unable to estimate a maximum residue level or STMR for the tree nuts group.

Linseed

The critical GAP for linseed (flax) is from Canada; 1 foliar application (SC formulation) at 132 g ai/ha and a 36-day PHI. The Meeting received supervised residue trials conducted in Canada on flax matching the critical GAP.

Trifloxystrobin (parent only) residues in flax seed in ranked order were (n = 11): < 0.01 (4), 0.013, 0.015, 0.023, 0.04, 0.042, 0.054 and 0.28 mg/kg.

Total residue levels of trifloxystrobin and CGA321113 in flax seed in ranked order were (n = 11): < 0.01 (4), 0.013, 0.015, 0.023, 0.04, 0.042, 0.055 and 0.29 mg/kg.

The Meeting estimated a maximum residue level of 0.4 mg/kg and an STMR of 0.015 mg/kg for linseed.

Coffee beans

The critical GAP for coffee is from Brazil; 3 foliar applications (SC formulation) at 100 g ai/ha, with a 21-day RTI and a 30-day PHI with 0.25% v/v methylated soybean oil. The Meeting received supervised residue trials conducted in Brazil on coffee matching the critical GAP.

Trifloxystrobin (parent only) residues in coffee beans in ranked order were (n = 7): < 0.01 (6) and 0.01 mg/kg.

Total residue levels of trifloxystrobin and CGA321113 in dry coffee beans in ranked order were (n = 7): ≤ 0.01 (4), 0.01, 0.02 and 0.03 mg/kg.

The Meeting estimated a maximum residue level of 0.015 mg/kg and an STMR value of 0.01 mg/kg for coffee beans.

Residues in animal feeds

Bean forage

The critical GAP for beans is from Austria; 2 foliar applications (SC formulation) at 200 g ai/ha, with a 7 day RTI and 7-day PHI. The Meeting received three supervised residue trials conducted in Europe on bean green material (bean forage).

The Meeting concluded that there are an insufficient number of trials conducted at the critical GAP or any other GAP provided for beans to estimate a median or highest residue for bean forage.

Pea vines

The critical GAPs for peas are from Austria and the Netherlands: 2 foliar applications (SC formulation) at 200 g ai/ha, with a 14-day RTI and a 14-day PHI. The Meeting received supervised residue trials conducted Europe on pea rest of plant and green material (pea vines) where 2 foliar spray applications (SC formulation) were made at a nominal rate of 200 g ai/ha, with a 7-day RTI and a 14-day PHI.

Although the residue trials have a shorter RTI compared to the critical GAPs, based on the residue decline data for pea vines the Meeting estimated that residues from the field trials would be within 25% of the critical GAP.

Trifloxystrobin (parent only) residues in pea vines in ranked order were (n = 17): 0.25, 0.26, 0.9, 0.91, 0.99, 1.0, 1.1, 1.5, 2.2, 2.8, 3.7, 4.5, 5.3, 5.4, 5.6, 5.8 and 11 mg/kg.

Total residue levels of trifloxystrobin and CGA321113 in pea vines in ranked order were (n = 17): 0.27, 0.28, 0.95, 1.0 (3), 1.1, 1.7, 2.3, 2.9, 3.9, 4.7, 5.4, 5.5, 5.6, 6.1 and 11 mg/kg.

The Meeting estimated a highest residue of 11 mg/kg and a median value of 2.3 mg/kg for pea vines.

Pea hay

The Meeting received GAP information for pea hay from Austria, France and the Netherlands. None of the available trials matched these GAPs. The Meeting could not estimate a maximum residue level, highest residue, or median value for pea hay.

Almond hulls

None of the almond supervised residue trials were conducted according to the US GAP; 4 × 140 g ai/ha, an RTI of 7–21 days and a 14-day PHI. The Meeting was unable to estimate a maximum residue level or median value for almond hulls.

Fate of residues in processing

Processing data on beans with pods, linseed and coffee were provided. All data relevant for an estimation of maximum residue levels in processed commodities or for dietary exposure calculations are summarized in the following table.

Table 1 Processing factors and residue estimates for trifloxystrobin and CGA321113

Raw commodity	Residue in RAC, mg/kg		Processed commodity	Processing Factors		Residue in processed commodity, mg/kg	
	Max	STMR		Trifloxystrobin [mean or best estimate]	Total ^a [mean or best estimate]	Max-P	STMR-P
Linseed	0.4	0.015	Meal	< 0.15, 0.14 [< 0.15]	1.2, 0.33 [0.77]	--	0.012
			Oil, crude (pressed)	0.82, 0.83 [0.83]	0.89, 0.74 [0.82]	--	--
			Oil, refined (solvent extracted)	0.60, 0.98 [0.79]	0.66, 0.88 [0.77]	--	0.012
Coffee, green beans	0.01*	0.01	Instant	- ^b	0.15	--	0.0015
			Roasted	- ^b	0.15	--	0.0015

^a Processing factor for the sum of trifloxystrobin and CGA321113

^b No processing factors for instant or roasted coffee could be derived for residues of trifloxystrobin since no quantifiable residues were observed in the RAC (green beans), despite the use of exaggerated application rates (approximately 2.5× the maximum label rate).

RESIDUES IN ANIMAL COMMODITIES

Farm animal feeding studies

Farm animal feeding studies (lactating dairy cow and laying hen) are reported in the evaluation of the 2004 JMPR.

Farm animal dietary burden

The Meeting has added feed items (pea vines and linseed meal) and their associated residues to the dietary burden calculations used by the 2015 JMPR. Cotton meal, cotton hulls and cotton gin-by-products from the 2017 JMPR were also included. Dietary burden calculations are provided in Annex 6; the dietary burden estimates are summarized below.

The Australian maximum beef and dairy dietary burdens exceeded the maximum dose in the feeding study. The Meeting refined the dietary burden to exclude pea vines from the Australian

livestock diets as trifloxystrobin is not registered for use on peas in Australia and forages are generally not traded internationally.

Table 2 Estimated maximum and mean dietary burdens of farm animals

Animal	Dietary burden estimates, ppm							
	Canada-US		European Union		Australia		Japan	
	Maximum	Mean	Maximum	Mean	Maximum	Mean	Maximum	Mean
Beef cattle	1.54	0.51	15.59 ^a	3.74 ^c	8.22	5.00	4.52	0.84
Dairy cattle	6.38	1.68	15.26 ^b	3.66 ^d	8.19	4.11	2.10	0.42
Broiler poultry	0.10	0.10	0.07	0.07	0.15	0.15	0.03	0.03
Laying hen	0.10	0.10	4.98 ^e	1.09 ^f	0.15	0.15	0.07	0.07

^a Highest maximum beef cattle dietary burden suitable for MRL estimates for mammalian tissues.

^b Highest maximum dairy cattle dietary burden suitable for MRL estimates for milk.

^c Highest mean cattle dietary burden suitable for STMR estimates for mammalian tissues.

^d Highest mean dairy cattle dietary burden suitable for STMR estimates for milk.

^e Highest maximum poultry dietary burden suitable for MRL estimates for poultry tissues and eggs.

^f Highest mean poultry dietary burden suitable for STMR estimates for poultry tissues and eggs.

Animal commodity maximum residue levels

Based on the estimated dietary burden and the results of the animal feeding studies evaluated by the 2004 JMPR, the calculations used to estimate highest total residues for use in estimating maximum residue levels and STMR values are shown below.

Table 3 Anticipated residues in mammalian commodities

Trifloxystrobin feeding study	Feed level (ppm) for milk residues	Total Residues (mg/kg) in milk ^a	Feed level (ppm) for tissue residues	Total Residues ^a (mg/kg)			
				Muscle	Liver	Kidney	Fat
Maximum residue level beef or dairy cattle							
Feeding study ^b	5.90	-	5.90	-	< 0.04	< 0.04	0.04
	21.00	< 0.02	21.00	< 0.04	0.11	0.04	0.08
Dietary burden and high residue	15.26	< 0.012	15.59	< 0.026	0.085	< 0.04	0.066
STMR beef or dairy cattle							
Feeding study ^c	2.00	-	2.00	-	< 0.04	< 0.04	< 0.04
	5.90	-	5.90	-	< 0.04	< 0.04	< 0.04
	21.00	< 0.02	21.00	< 0.04	0.07	0.04	0.06
Dietary burden and residue estimate	3.66	< 0.005	3.74	< 0.01	< 0.04	< 0.04	< 0.04

^a Total residues = trifloxystrobin and CGA321113.

^b highest residues for tissues and mean residues for milk

^c mean residues for tissues and mean residues for milk

The Meeting confirmed its previous recommendation of a maximum residue level of 0.02* mg/kg for milks.

The Meeting estimated maximum residue levels of 0.07 mg/kg for meat, based on fat (from mammals other than marine mammals) and mammalian fats (except milk fats) and 0.09 mg/kg for edible offal (mammalian). The Meeting estimated STMRs of 0.01 mg/kg for meat (muscle), 0.04 mg/kg for mammalian fat, 0.04 mg/kg for liver, 0.04 mg/kg for kidney and 0.005 mg/kg in milks. These recommendations are intended to replace all previous recommendations for all ruminant matrices.

In the feeding study in laying hens, no trifloxystrobin or CGA321113 residues (total residue, < 0.04 mg/kg) were detected in eggs, tissues or organs from hens at the highest feeding level of 15 ppm. As the maximum and mean dietary burdens of 4.98 mg/kg and 1.09 mg/kg, respectively, were much lower, the Meeting estimated a maximum residue level of 0.04(*) mg/kg for poultry fats,

confirmed its previous recommendations of maximum residue levels of 0.04(*) mg/kg for eggs, poultry meat (fat) and edible offal of poultry and recommended that the STMR values should be 0.0046 mg/kg in eggs, poultry meat, edible offal and fat.

RECOMMENDATION

On the basis of the data obtained from supervised residue trials the Meeting concluded that the residue levels listed in Annex 1 are suitable for establishing maximum residue limits and for IEDI assessment.

Definition of residue for compliance with the MRL for plant commodities: *trifloxystrobin*

Definition of residue for compliance with the MRL for animal commodities and dietary risk assessment for plant and animal commodities : *sum of trifloxystrobin and (E,E)-methoxyimino-{2-[1-(3-trifluoromethyl-phenyl) ethylidene-aminooxymethyl]-phenyl}acetic acid = CGA321113) (expressed as trifloxystrobin equivalents).*

The residue is fat-soluble.

DIETARY RISK ASSESSMENT

Long-term dietary exposure

The ADI for trifloxystrobin is 0–0.04 mg/kg bw. The International Estimated Daily Intakes (IEDIs) for trifloxystrobin were estimated for the 17 GEMS/Food Consumption Cluster Diets using the STMR or STMR-P values estimated by the JMPR. The results are shown in Annex 3 of the 2021 Extra JMPR Report.

The IEDIs ranged from 1–9% of the maximum ADI. The Meeting concluded that long-term dietary exposure to residues of trifloxystrobin from uses considered by the JMPR is unlikely to present a public health concern.

Acute dietary exposure

The 2004 JMPR decided that an ARfD for trifloxystrobin was unnecessary. The Meeting therefore concluded that the acute dietary exposure to residues of trifloxystrobin resulting from uses that have been considered by the JMPR is unlikely to present a public health concern.

5.30. **TRINEXAPAC-ETHYL (271)**

TOXICOLOGY

Trinexapac-ethyl is the ISO-approved common name for 4-(cyclopropyl- α -hydroxymethylene)-3,5-dioxo-cyclohexanecarboxylic acid ethyl ester (IUPAC), with CAS No. 95266-40-3. Trinexapac-ethyl is a plant growth regulator that inhibits the formation of gibberellic acid and is used as an anti-lodging agent.

Trinexapac-ethyl was previously evaluated by JMPR in 2013 when an acceptable daily intake (ADI) of 0–0.03 mg/kg bw was established and it was concluded that an acute reference dose (ARfD) was unnecessary (Annex 5, reference 128).

Toxicological data on metabolites

Metabolites CGA113745, CGA224439 and CGA313458 were found variously in animal metabolism studies (but not in rat studies), plants and in high temperature processed commodities.

A 90-day toxicity study via gavage for CGA22439 was provided to JMPR 2014 (Annex 5, reference 133) because of the formation of a photolytic degradate of aminocyclopyrachlor. No other toxicological studies are available to JMPR to assess the toxicological relevance of these three metabolites, therefore evaluation of potential toxicity is based on QSAR and expert judgement using JMPR's metabolite assessment scheme included in the 2015 guidance document for WHO monographers (JMPR (WHO, 2015⁵)). No specific data were provided and the Meeting performed a QSAR analysis to provide a prediction of the potential toxicity of these metabolites in relation to the parent compound.

Metabolites CGA113745 (3,5-dioxocyclohexane-1-carboxylic acid) and CGA313458 ((2RS)-2-(4-cyclopropyl-2,4-dioxobutyl) butanedioic acid).

The potential toxicity of metabolites CGA113745 and CGA313458 was evaluated using Derek Nexus (an expert system for prediction of toxicity) implemented in Nexus (v. 2.3.0) and Knowledge Base Derek KB 2020 1.0. The software compares the structures with known toxicity end points. When a structural alert is identified the programme assigns a probability to the expression of toxicity by the compound. The parent compound trinexapac-ethyl was used as a reference chemical for prediction of alerts for the metabolites. The results of the Derek predictions are shown in Table 1. Predictions for metabolites CGA113745 and CGA313458 are qualitatively similar to the parent compound. It did not give any alerts for genotoxicity. Both compounds fall in Cramer class III.

Metabolite CGA224439 (cyclopropanecarboxylic acid; CPCA)




Derek in silico predictions for metabolite CGA224439 are shown in Table 1. The metabolite CGA224439 predictions are qualitatively different to those of the parent compound. An alert was triggered for plausible carcinogenicity in mammals due to the irritating properties of the material in the gastrointestinal (GI) tract. Derek also predicted hepatotoxicity as plausible, due to alert based on short chain fatty acid. There was an alert for GI tract irritation as plausible based on short chain alkyl carboxylic acid. There was also an equivocal alert for mitochondrial dysfunction based on short chain fatty acid. The metabolite is inactive in in vitro mutagenicity tests on bacteria and it is not expected to be a skin sensitizer in mammals. CPCA falls in Cramer class II.

In a 90-day toxicity study in rats, which tested CPCA doses of 0, 2, 10, 30 and 60 mg/kg bw per day administered by gavage, the NOAEL was 10 mg/kg bw per day, based on increased aspartate transaminase (AST; females only), increased total bile acids (females only), decreased globulin with an associated decrease in total protein (males only) and adverse microscopic findings in the heart, liver (females only) and thymus (females only) at 30 mg/kg bw per day (Annex

⁵ WHO (2015). Pesticide residues in food. Guidance document for WHO monographers and reviewers. WHO Core Assessment Group on Pesticide Residues. WHO, Geneva, 2005. January 2015

5, reference 133). The NOAEL for trinexapac-ethyl in the 90-day dietary study in rats was 500 ppm, equal to 34 mg/kg bw per day (Annex 5, reference 128). Therefore CGA224439 appears to be more toxic in rats than the parent trinexapac-ethyl following repeated oral dosing.

Table 1 Derek predictions for trinexapac-ethyl and its metabolites (CGA13745, CGA313458 and CGA 224439)

Name/s	Chemical structure	Derek alerts					
		Geno-toxicity	Skin sensitization	Carcino-genicity	Hepato-toxicity	Irritation (GI tract)	Mitochondrial disfunction
Trinexapac-ethyl		No alert	+ ^a	No alert	No alert	No alert	No alert
CGA13745 3,5-Dioxocyclohexane-1-carboxylic acid		No alert	+ ^a	No alert	No alert	No alert	No alert
CGA313458 (2 <i>RS</i>)-2-(4-Cyclopropyl-2,4-dioxobutyl)butanedioic acid		No alert	+ ^a	No alert	No alert	No alert	No alert
CGA224439 Cyclopropanecarboxylic acid		No alert	No alert	+ ^b	+ ^c	+ ^c	= ^c

Key: ~ indicates inactive; = indicates equivocal; + indicates plausible; ++ indicates probable

^a Plausible alert based on 1,3-diketone

^b Plausible alert based on short chain alkyl carboxylic acid and GI irritation

^c Alert based on short chain alkyl carboxylic acid/short chain fatty acid

TOXOLOGICAL EVALUATION

Metabolites CGA113745 and CGA3134548

In the absence of compound-specific data and as per JMPR guidance (WHO, 2015b), the meeting concluded that the reference values for CGA113745 and CGA3134548 could not be set, therefore it is appropriate to compare the exposure estimate to the non-genotoxic TTC value for Cramer class III (1.5 µg/kg bw per day).

A toxicological addendum was not prepared

Metabolite CGA224439

In the absence of compound-specific data and as per JMPR guidance (2015b), the meeting concluded that the reference values for CGA224439 could not be set, therefore it is appropriate to compare the exposure estimate to the non-genotoxic TTC value for Cramer class II (9 µg/kg bw per day).

A toxicological addendum was not prepared

RESIDUE AND ANALYTICAL ASPECTS

Trinexapac-ethyl is a synthetic plant growth regulator used for growth management of crops. It was first evaluated by JMPR in 2013 (T, R). An ADI of 0–0.3 mg/kg bw, expressed as trinexapac acid equivalents, was established in 2013 JMPR and the Meeting concluded that an ARfD for trinexapac-ethyl was unnecessary.

The definition of the residue for compliance with the MRL for plant and animal commodities and for dietary risk assessment for animal commodities is: *Trinexapac acid*.







The definition of the residue for dietary risk assessment for plant commodities is: *Trinexapac acid and its conjugates, expressed as trinexapac acid*.

The residue is not fat-soluble.

It was scheduled at the Fifty-first Session of the CCPR for the evaluation of additional uses by the 2020 JMPR, which was postponed to the 2021 Extra JMPR. The Meeting received information on the metabolism in wheat and oilseed rape, methods of residue analysis, freezer storage stability, the GAP, supervised residue trials on rice, as well as industrial processing studies.

Table 1 The structures of trinexapac-ethyl and its key metabolites discussed are shown below

Compound	Structure
Trinexapac-ethyl	
Trinexapac acid CGA179500	
Tricarballic acid CGA275537	
Trans aconitic acid CGA312753	
Mono-ethyl ester of trans aconitic acid	
CGA313458	

Compound	Structure
CGA329773	
CGA224439	
CGA113745	
Cyclodion acid	
3-carboxyl-7-cyclopropyl-5,7-dioxoheptanoic acid	
CGA300405	

Plant Metabolism

Oilseed rape

Oilseed rape plants were treated at a rate of 393.8 g ai/ha with [cyclohexadione-1,2,6-¹⁴C]-trinexapac-ethyl formulated as a microemulsion at BBCH 55 by foliar spray application. Plants were collected (fully ripe) between 67 and 91 days after application (BBCH 89). Only the seeds were analysed. Total radioactive residue in the seeds was 0.39 mg eq/kg. The extraction of ¹⁴C residues in seed (twice with acetonitrile:water:hexane 16:4:10 (v/v/v), followed by twice with acetonitrile:water 50:50 (v/v) at room temperature) was 67% TRR (0.27 mg eq/kg). No parent trinexapac-ethyl was detected in seeds. Trinexapac acid was identified as the major metabolite, accounting for 22% TRR, 0.086 mg eq/kg (19% TRR free and 2.8% TRR conjugated). The tricarballic acid was the minor metabolite detected

(1.0% TRR; 0.004 mg eq/kg, exclusively in the conjugated form). A major part of the radioactivity was identified as oleic acid (22.7% TRR; 0.09 mg eq/kg, detected following saponification of radioactivity associated with hexane-extracted endogenous oils). The individual levels of other unassigned radiocomponents from extracted residues never exceeded 6.7% TRR, (0.027 mg eq/kg). The characterisation of the unextracted radioactive residues was conducted, however, no identification was achieved due to low residue levels released.

Spring wheat

Spring wheat plants were treated once at rate of 211 g ai/ha with [cyclohexadione-¹⁴C]-trinexapac-ethyl as a microemulsion at BBCH 37 by foliar spray application. Samples were collected at forage growth stage (BBCH 43), at hay growth stage (BBCH 77) and at maturity (BBCH 89).

The total radioactive residues (TRR) for harvested commodities were 1.8 mg eq/kg in forage, 2.0 mg eq/kg in hay, 1.4 mg eq/kg in straw and 1.4 mg eq/kg in grain. Good extractability with acetonitrile/water (three times with 4:1 (v/v) and once with 1:1 (v/v) at room temperature) was achieved in forage, grain and hay ($\geq 84\%$ TRR) with lower extractability in straw (65% TRR). Metabolism of parent trinexapac-ethyl was extensive, the parent was only detected in forage with a low residue level of 0.3% TRR, 0.006 mg eq/kg. The principal metabolites identified in grain were trinexapac acid (40% TRR, 0.58 mg eq/kg; 28% TRR free and 12% TRR conjugated) in grain, the minor metabolites identified in grain were tricarballic acid ethyl ester metabolite CGA300405 (0.8% TRR; 0.012 mg eq/kg), tricarballic acid metabolite CGA275537 (2.0% TRR; 0.03 mg eq/kg) and a hydroxylated metabolite of trinexapac acid (12% TRR, 0.18 mg eq/kg).

The principal metabolites identified in forage, hay and straw were trinexapac acid (5.5–23% TRR; 0.075–0.45 mg eq/kg), the tricarballic acid ethyl ester metabolite (CGA300405, 8.0–21% TRR, 0.13–0.37 mg eq/kg) and tricarballic acid (CGA275537, 7.8–10% TRR, 0.11–0.14 mg eq/kg). Formation of citric acid and subsequent incorporation of small ¹⁴C containing moieties into the broader pool of natural biosynthetic products was observed (characterised by the presence of ¹⁴C-glucose in hay, straw and grain). All metabolites with the exception of CGA329773 and citric acid were found in their free and conjugated forms.

The characterisation of the unextracted radioactive residues in the hay, grain and straw samples was accomplished with a clean fractionation technique, which released additional small amounts of the same metabolites as observed in the extracted fractions. CGA300405 was identified as the highest residue in all samples analysed (0.3–0.6% TRR; 0.004–0.013 mg eq/kg). Acid hydrolysis of a hemicellulose fraction (12% TRR; 0.16 mg eq/kg) derived from straw showed the majority of this fraction to comprise [¹⁴C]-glucose, demonstrating extensive incorporation into natural components.

The Meeting concluded that the metabolism profile of trinexapac-ethyl in rapeseed and spring wheat was comparable to that previously evaluated by the 2013 JMPR and the Meeting confirmed the current residue definition for plant commodities.

Methods of analysis

The Meeting received information on the description and validation of analytical methods for trinexapac acid in plant matrices, as well as a multi-residue method for trinexapac acid in plant and animal matrices.

The method GRM020.05A for free trinexapac acid involved extraction of the samples with methanol/water/phosphate buffer (pH7, v/v/v 30:56:14), clean-up with SPE and quantification with HPLC-MS/MS (monitoring for quantitation $m/z=223 \rightarrow 135$ and for confirmation $m/z = 223 \rightarrow 179$ or $m/z = 223 \rightarrow 83$). An LOQ of 0.01 mg/kg was validated for free trinexapac acid in barley, tomato, apple and sunflower seed.

The method GRM020.09A for free and conjugated trinexapac acid involved sequential extraction with acetonitrile/water (80/20 v/v) and acetonitrile/water (50/50 v/v) for free trinexapac acid and hydrolysis under 0.05M sodium hydroxide at room temperature overnight (minimum 12 hours) for conjugated trinexapac acid, with quantification by HPLC-MS/MS (quantitation

$m/z=223 \rightarrow 135$, confirmatory $m/z = 223 \rightarrow 83$ or $m/z = 223 \rightarrow 125$). An LOQ of 0.01 mg/kg in cereal grain and 0.05 mg/kg in cereal straw was validated for trinexapac acid.

The extraction and hydrolysis were shown to be satisfactory by reference to the metabolism study on spring wheat. The Meeting concluded that the presented methods are suitable to measure either free or, free and conjugated trinexapac acid in plant commodities.

The multi-residues method of QuEChERS was newly validated for determination of free trinexapac acid in plant and animal matrices, with an LOQ of 0.01 mg/kg.

Stability of pesticide residues in stored analytical samples

The Meeting received additional information on storage stability of cyclopropane carboxylic acid in cereal grain and its processed commodities.

Storage stability studies showed that residues of cyclopropane carboxylic acid is stable for at least 12 months at $<-18\text{ }^{\circ}\text{C}$ for high starch commodities and processed commodities from cereal grain (flour, bran, bread and beer).

The samples of rice grain and straw in supervised trials available to the Meeting were stored deep frozen for a maximum of 12 months (365 days) and were covered by storage stability evaluated in 2013 JMPR.

Definition of the residue

The current Meeting received new plant metabolism data indicating that the metabolism profile of trinexapac-ethyl in rapeseed and spring wheat was comparable to that previously evaluated by the 2013 JMPR and the Meeting confirmed the current residue definition for plant commodities.

The current Meeting also received new information on the nature of residues during processing following high temperature hydrolysis. In combination with the information considered by the 2013 JMPR, it was confirmed that trinexapac acid shows limited stability under conditions representative of pasteurisation, baking, brewing, boiling and sterilisation. Major degradates were CGA113745 (10–12% AR), CGA313458 (16–21% AR) and cyclopropane carboxylic acid (CGA224439, 16% AR).

CGA113745 was also found in goat liver, kidney and fat up to 16% of the TRR. The residue ratio of CGA113745 to trinexapac acid was up to 0.25.

CGA313458 was also found in plant matrices. The residue ratios compared to trinexapac acid were 0.3 in rice grain and 0.03 in rape seeds.

The information on the toxicity for all three metabolites was limited and the Meeting decided to assess them using the thresholds of toxicological concern (TTC) approach in the following classes:

CGA113745 and CGA313458	Cramer Class III compounds
CGA224439	Cramer Class II compound

Results of supervised residue trials on crops

Supervised trials were available for the use of trinexapac-ethyl on rice. Product labels were available from the USA.

Rice

The critical GAP for trinexapac ethyl on rice in the USA is one application at 50 g ai/ha (up to panicle formation, panicle 1–2mm, BBCH32) with a PHI of 50 days.

In six trials conducted on rice in the USA approximating critical GAP, the residues of trinexapac acid (free and conjugated) in grains were ($n = 6$): 0.017, 0.023, 0.054, 0.073, 0.18 and 0.23 mg/kg. The free trinexapac acid was not separately analysed.

In four trials conducted in USA with longer days after last application (DALA, 85-100 days after application), the residues of trinexapac acid (free and conjugated) in grain were (n = 4): 0.035, 0.050, 0.074 and 0.10 mg/kg. The free trinexapac acid was not separately analysed.

Considering the ratios of % TRR for free compared to free and conjugated (total) trinexapac acid in rice grain from three metabolism studies, that were 0.86 (10% free/11.6% total), 0.82 (23% free/28.1% total) and 0.92 (20.2% free/24% total), the conjugated residues contributed less than 20% of the total. The Meeting agreed to recommend the maximum residue level based on the residues of total trinexapac acid.

Noting all trials were conducted at growth stage of BBCH 30–32 and the residue decline studies showed that longer sampling intervals had little or no impact on the residues of trinexapac acid in rice grain, the Meeting decided to use the combined data to make estimation. The combined data set was (n = 10): 0.017, 0.023, 0.035, 0.050, 0.054, 0.073, 0.074, 0.10, 0.18 and 0.23 mg/kg.

The Meeting estimated a maximum residue level of 0.5 mg/kg and an STMR of 0.064 mg/kg for trinexapac-ethyl in rice grain.

Rye

The critical GAP for trinexapac-ethyl on rye in the USA is one application at 123 g ai/ha with a PHI of 45 days. No data reflecting the critical GAP on rye were available.

In trials evaluated by the 2013 JMPR, the residues of free trinexapac acid in wheat grain were (n = 18): 0.10, 0.25, 0.32, 0.34, 0.35, 0.46, 0.49, 0.55, 0.55, 0.57, 0.77, 0.88, 0.91, 0.98, 0.99, 1.0, 1.4 and 2.0 mg/kg. The residues of free and conjugated trinexapac acid in wheat and barley grain were (n = 30): 0.03, 0.07, 0.08, 0.15, 0.27, 0.31, 0.32, 0.40, 0.44, 0.45, 0.47, 0.50, 0.52, 0.53, 0.53, 0.60, 0.72, 0.76, 0.77, 0.78, 0.82, 0.83, 0.85, 0.99, 1.0, 1.0, 1.1, 1.2, 1.6 and 3.3 mg/kg.

The current Meeting recognized that rye and wheat (spring wheat, winter wheat and durum wheat) have the same GAP and show comparable residues after early treatment. Application was before flowering. Therefore, the current Meeting agreed to extrapolate the maximum residue level of 3 mg/kg estimated for wheat grain to rye grain and to extrapolate the STMR of 0.57 mg/kg from wheat and barley to rye grain.

Residues in animal feeds

Rice straw

The Meeting received supervised residue trials on rice. The critical GAP for trinexapac-ethyl on rice in the USA is one application at 50 g ai/ha (up to panicle formation (panicle 1–2mm, BBCH32) with a PHI of 50 days.

In six trials conducted on rice in the USA approximating critical GAP, the residues of trinexapac acid (free and conjugated) in straw were (n = 6): < 0.01(2), 0.012, 0.016, 0.036 and 0.045 mg/kg. The free trinexapac acid was not separately analysed.

In four trials conducted in the USA with longer days after last application (DALA, 85–100 days after application, more than 25%), the residues of trinexapac acid (free and conjugated) in straw were (n = 4): < 0.01, 0.01, 0.017 and 0.02 mg/kg. The free trinexapac acid was not separately analysed.

Considering the ratios of % TRR for free compared to total trinexapac acid in rice straw in three metabolism studies on rice, that were 0.86 (4.2% free/4.9% total), 0.97 (6.7% free/6.9% total) and 0.80 (6.7% free/8.4% total), the conjugated residues of trinexapac acid contributed less than 20% of the total. The Meeting agreed to recommend the maximum residue level based on the residues of total trinexapac acid.

Noting that all trials were conducted at a growth stage of BBCH 30–32 and the residue decline studies showed that longer sampling intervals had little or no impact on the residues of

trinexapac acid in rice straw, the Meeting decided to use the combined data to make a maximum residue level estimation. The combined data set was (n = 10): <0.01(3), 0.01, 0.012, 0.016, 0.017, 0.02, 0.036 and 0.045 mg/kg.

The Meeting estimated a maximum residue level of 0.08 mg/kg (dw), a median residue of 0.014 mg/kg (as received) and a highest residue of 0.045 mg/kg (as received) for trinexapac-ethyl in rice straw.

Rye

The critical GAP for trinexapac-ethyl on rye in the USA is one application at 123 g ai/ha with a PHI of 45 days. No data reflecting the critical GAP on rye were available.

Rye forage

In trials evaluated by the 2013 JMPR, the residues of total trinexapac acid in wheat forage collected 30 days after application (dry weight) at the GAP application rate were (n = 18): 0.08, 0.12, 0.16, 0.24, 0.28, 0.32, 0.32, 0.36, 0.40, 0.40, 0.48, 0.68, 0.68, 0.88, 0.92, 1.32, 1.52 and 3.76 mg/kg.

The current Meeting recognized that rye and wheat (spring wheat, winter wheat and durum wheat) have the same GAP and show comparable residues after early treatment. Application was before flowering. The current Meeting agreed to extrapolate the median residue of 0.40 mg/kg and highest residue of 3.76 mg/kg for trinexapac acid in wheat forage to rye forage (dry).

Rye hay and straw

In the trials evaluated by the 2013 JMPR, the residues of free trinexapac acid in wheat hay collected 30 days after application (dry weight) at the GAP application rate were (n = 18): <0.06(3), 0.06, 0.07, 0.08, 0.08, 0.09, 0.09, 0.10, 0.13, 0.13, 0.15, 0.17, 0.20, 0.22, 0.22 and 0.85 mg/kg. The residues of total trinexapac acid in barley and wheat hay collected 30 days after application (dry weight) according to the GAP were (n = 29): <0.011, 0.03, 0.03, 0.05, 0.05, 0.06, 0.07, 0.10, 0.11, 0.13, 0.13, 0.15, 0.16, 0.17, 0.19, 0.20, 0.22, 0.27, 0.28, 0.34, 0.35, 0.38, 0.45, 0.47, 0.55, 0.57, 0.67, 0.89 and 1.34 mg/kg.

The current Meeting noted that rye and wheat (spring wheat, winter wheat and durum wheat) have the same GAP and normally show comparable residues after early treatment.

The current Meeting agreed to extrapolate the maximum residue level of 0.9 mg/kg for trinexapac-ethyl in wheat straw and fodder (dry) to rye straw and fodder (dry) and to extrapolate the median residue of 0.19 mg/kg and highest residue of 1.34 mg/kg for trinexapac-ethyl for wheat straw and fodder (dry) and barley straw and fodder (dry) to rye straw and fodder (dry).

Fate of residues during processing

The Meeting received two new studies on high temperature hydrolysis of residues of trinexapac acid

One study showed that trinexapac acid (hydroxymethylene-C¹⁴ labelled) underwent limited degradation under conditions representative of pasteurisation (pH 4, 90 °C, 20 minutes) and sterilisation (pH 6, 120 °C, 20 minutes) and degraded at significant amounts to cyclopropane carboxylic acid (16%) and CGA313458 (18%) under conditions representative of baking, brewing and boiling (pH 5, 100 °C, 60 minutes). Another study showed that trinexapac acid (1,6-cyclohexyl and hydroxymethylene-C¹⁴ labelled) is hydrolytically stable under conditions representative of pasteurisation, baking, brewing, boiling and sterilisation and two metabolites, cyclodion acid and 3-carboxyl-7-cyclopropyl-5,7-dioxoheptanoic acid increased from 1.7 to 3.4% and 2 to 3.5%, respectively, under condition of baking, brewing, boiling (pH 5, 100 °C, 60 minutes).

The 2013 JMPR concluded that trinexapac acid (cyclohexyl C¹⁴ labelled) undergoes limited degradation during the simulation of pasteurisation (pH 4, 90 °C, 25 minutes), baking, boiling and brewing (pH 5, 100 °C, 60 minutes) and sterilisation (pH 6, 120 °C, 20 minutes). Trinexapac acid was

the major component of the radioactive material at the end of hydrolysis reactions (51–59%) and the major degradates were CGA113745 (10–12%) and CGA313458 (16–21%).

The Meeting concluded that trinexapac acid might undergo limited hydrolysis under high temperature conditions, resulting in > 60% of *trinexapac acid* remaining at the end of each treatment.

The Meeting also received processing studies for barley, rice and wheat. Estimated processing factors for the commodities considered at this Meeting are summarised below.

Table 2 Processing factors for estimation of STMR-Ps (free and conjugated)

	Processed commodity	Median or best estimate processing factor	STMR-P = STMR _{RAC} × PF (mg/kg)
Barley (STMR 0.57 mg/kg)	Pot Barley	0.20	0.11
	Pearled Barley	0.52	0.30
	Barley Bran	1.1	0.63
	Barley Flour	0.43	0.25
	Brewing Malt	0.49	0.28
	Beer	0.06	0.03
Wheat (STMR 0.57 mg/kg)	White Flour	0.4	0.23
	Bran	1.2	0.68
	Whole meal Flour	0.72	0.41
	Wholemeal Bread	0.62	0.35
	Germ	0.66	0.38
	Dry Gluten	0.24	0.14
	Dry Starch	0.08	0.046
	Gluten Feed Meal	0.18	0.10 (median residue)
Rice (STMR 0.064 mg/kg)	Rice (polished)	1.3	0.08
	Rice hulls	0.41	0.03
	Rice bran	5.1	0.33

Table 3 Processing factors for estimation of maximum residue levels in processed commodities (free acid)

	Processed commodity	Mean or best estimate processing factor	Maximum residue level for processed commodity (mg/kg)
Barley (maximum residue level 3 mg/kg)	Barley Bran	1.3	3.9
Rice (maximum residue level 0.5 mg/kg)	Rice (polished)	1.3	0.65
	Rice bran	5.1	2.55
Wheat (maximum residue level 3 mg/kg)	Wheat bran	1.6	4.8

Based on the processing factor of 1.3 for barley bran and the barley grain maximum residue level of 3 mg/kg, the calculated highest residues in barley bran are 3.9 mg/kg. The Meeting estimated a maximum residue level for trinexapac-ethyl in barley bran of 4 mg/kg, to replace the previous recommendation of 6 mg/kg.

Based on the processing factors of 1.3 for polished rice and 5.1 for rice bran and the rice grain maximum residue level of 0.5 mg/kg, the calculated highest residues in polished rice and rice bran are

0.65 and 2.55 mg/kg, respectively. The Meeting estimated a maximum residue level for trinexapac-ethyl in polished rice of 0.7 mg/kg, in rice bran of 3 mg/kg.

Based on the processing factor of 1.6 for wheat bran and the wheat grain maximum residue level of 3 mg/kg, the calculated highest residues in wheat bran are 4.8 mg/kg. The Meeting estimated a maximum residue level for trinexapac-ethyl in wheat bran of 5 mg/kg, to replace the previous recommendation of 8 mg/kg.

Residues in animal commodities

The current Meeting calculated dietary burdens based on residue estimates from previous meeting with updates to reflect feed commodities addressed by the current Meeting. Potential feed items include: wheat, barley, oat and triticale grain, straw, forage, hay and silage, wheat milled by-products (bran), wheat aspirated grain fraction, barley bran fractions, sugarcane molasses and bagasse and rape forage and rape seed meal, rice grain and straw, rye grain and straw. The dietary burdens estimated using the most recent version of the OECD livestock dietary burden calculator, are presented in Annex 6 and summarised below.

Table 4 Estimated maximum and mean dietary burdens of farm animals

	Animal dietary burden: Trinexapac acid, ppm of dry matter diet							
	US-Canada		EU		Australia		Japan	
	max	Mean	max	Mean	max	Mean	max	Mean
Beef cattle	1.14	0.741	1.743	0.726	3.969	0.752	0.791	0.791 (0.97) ^a
Dairy cattle	1.702	0.682	1.743	0.674	4.052 (3.76) ^a	0.729 (0.86) ^a	0.861	0.693
Poultry broiler	0.778	0.78 (0.94) ^a	0.635	0.635	0.698	0.6981	0.11	0.11
Poultry layer	0.778	0.778 (0.94) ^a	1.011 (1.08) ^a	0.7	0.697	0.697	0.273	0.273

^a values in parenthesis from the 2013 evaluation

Despite the additional feed items, the Meeting noted that the dietary burdens changed within a range of -18% to 8% compared with those estimated by the 2013 JMPR to derive residue estimates for animal commodities. The Meeting confirmed its previous recommendations for animal commodities.

RECOMMENDATIONS

On the basis of the data obtained from supervised trials, the Meeting concluded that the residue levels listed in Annex 1 are suitable for establishing maximum residue limits and for IEDI assessments.

The definition of the residue for compliance with the MRL for plant and animal commodities and for dietary risk assessment for animal commodities: *Trinexapac acid*.

The definition of the residue for dietary risk assessment for plant commodities: *Trinexapac acid and its conjugates, expressed as trinexapac acid*.

The residue is not fat-soluble.

DIETARY RISK ASSESSMENT

Long-term dietary exposure

The ADI for trinexapac-ethyl is 0–0.3 mg/kg bw (expressed as trinexapac acid equivalents). The International Estimated Daily Intakes (IEDIs) for trinexapac-ethyl were estimated for the 17 GEMS/Food Consumption Cluster Diets using the STMR or STMR-P values estimated by the JMPR. The results are shown in Annex 3 of the 2021 Extra JMPR Report.

The IEDIs were 0% of the maximum ADI. The Meeting concluded that long-term dietary exposure to residues of trinexapac-ethyl from uses considered by the JMPR is unlikely to present a public health concern.

Acute dietary exposure

The 2013 JMPR decided that an ARfD for trinexapac-ethyl was unnecessary. The Meeting therefore concluded that the acute dietary exposure to residues of trinexapac-ethyl from the uses considered is unlikely to present a public health concern.

Assessment of metabolites using the threshold of toxicological concern (TTC) approach

The metabolite CGA113745 could be assessed using the TTC approach (Cramer Class III threshold of 1.5 µg/kg bw per day).

The present Meeting noted that the ratio of residues for CGA113745 compared to trinexapac acid in animal commodities was up to 0.25. In addition, trinexapac acid could be converted into CGA113745 during high temperature hydrolysis by up to 12% of the AR. The Meeting applied a conservative ratio of residues of 0.37 (0.25+0.12) to the trinexapac acid IEDI to derive an upper limit estimate of dietary exposure to CGA113745.

Based on the highest IEDI for trinexapac acid of 1.5 µg/kg bw per day and the overall ratio of 0.37 for formation of CGA113745, an upper long-term dietary exposure of 0.56 µg/kg bw per day was estimated.

The Meeting concluded that the estimated dietary exposure to residues of CGA113745 from uses considered by the JMPR is below the TTC for Cramer Class III compounds and is unlikely to present a public health concern. Should further uses be considered in the future, these conclusions may need to be re-evaluated.

The metabolite CGA313458 could be assessed using the TTC approach (Cramer Class III threshold of 1.5 µg/kg bw per day).

The present Meeting noted that the ratio of residues for CGA313458 compared to trinexapac acid in plant commodities was up to 0.3 (rice grain). In addition, trinexapac acid could be converted during high temperature hydrolysis into CGA313458 by up to 21%. The Meeting applied a conservative ratio of residues of 0.51 (0.3+0.21) to the trinexapac acid IEDI to derive an upper limit estimate of dietary exposure to CGA313458.

Based on the highest IEDI for trinexapac acid of 1.5 µg/kg bw per day and the overall ratio of 0.51 for the formation of CGA313458, a long-term dietary exposure of 0.77 µg/kg bw per day was estimated.

The Meeting concluded that the estimated dietary exposure to residues of CGA313458 from uses considered by the JMPR is below the TTC for Cramer Class III compounds and is unlikely to present a public health concern. Should further uses be considered in the future, these conclusions may need to be re-evaluated.

The metabolite CGA224439 could be assessed using the TTC approach (Cramer Class II threshold of 9 µg/kg bw per day).

The present Meeting noted that CGA224439 is only formed during high temperature hydrolysis in limited amounts (up to 16% AR) and applied a conservative ratio of residues of 0.16 to the trinexapac acid IEDI to derive an upper limit estimate of the dietary exposure to CGA224439.

Based on the highest IEDI for trinexapac acid of 1.5 µg/kg bw per day and the ratio of 0.16 for the formation of CGA224439, a long-term dietary exposure of 0.24 µg/kg bw per day was estimated.

The Meeting concluded that the estimated dietary exposure to residues of CGA224439 from uses considered by the JMPR is below the TTC for Cramer Class II compounds and is unlikely to present a public health concern. Should further uses be considered in the future, these conclusions may need to be re-evaluated.

Annex 1: Acceptable daily intakes, acute reference doses, recommended maximum residue levels, supervised trials median residue values and other values recorded by the 2021 Extra JMPR Meeting.

Pesticide (Codex reference number)	CCN	Commodity	Recommended Maximum residue level (mg/kg)		STMR or STMR-P mg/kg	HR or HR-P mg/kg
			New	Previous		
Acetamiprid (246) ADI: 0–0.07 mg/kg bw ARfD: 0.1 mg/kg bw	TN 0085	Tree nuts, Group of	W	0.06	0.01	0.05
	TN 0085	Tree nuts, Group of, except Pistachio nut	0.06		0.01	0.05
	TN 0675	Pistachio nut	1		0.33	0.51
	Definition of the residue for compliance with the MRL and for dietary risk assessment for plant commodities: <i>Acetamiprid</i> . Definition of the residue for compliance with the MRL and for dietary risk assessment for animal commodities: <i>Sum of acetamiprid and N-desmethyl-acetamiprid, expressed as acetamiprid</i> . The residue is not fat-soluble					
Bixafen (262) ADI: 0–0.02 mg/kg bw ARfD: 0.2 mg/kg bw	GC 0640	Barley	1.5	0.4	0.33	
	SO 0691	Cottonseed	0.3		0.03	
	AS 0645	Maize fodder (dry)	5 (dw)		Median: 1.47 (as)	Highest: 2.24 (as)
	GC 0645	Maize	0.01(*)		0.02	
	AS 3490	Maize bran, unprocessed	0.03		0.062	
	OR 0645	Maize oil, Edible	0.02		0.036	
	SO 0697	Peanut	0.01		0.02	
	OR 0697	Peanut oil, Edible	0.03		0.044	
	VD 0070	Pulses, Group of (except Soya bean (dry))	0.04 ^a		0.02 ^a	
	VR 0075	Root and tuber vegetables, Group of	0.06 ^a		0.028 ^a	0.068 ^a
	GC 0651	Sorghum Grain	2		0.196	
	VD 0541	Soya bean (dry)	0.08		0.034	
	AB 0541	Soya bean hulls	0.3		0.0952	
	OR 0541	Soya bean oil, Refined	0.15		0.0476	
	SO 0702	Sunflower seed	3		0.045	
	GC 0447	Sweet corn (corn on the cob) (kernels plus cob with husk removed)	0.01(*)		0.02	0.02
	CF 0654	Wheat bran, Processed	0.8	0.15	0.2375	
	GC 0654	Wheat	0.3	0.05	0.095	
		Beer			0.036	
		Brewer's malt			0.317	
		Pearl barley			0.083	
		Soya bean, flour			0.00374	
		Soya bean milk			0.00221	
	CF 1211	Wheat flour			0.03135	
	CF 1210	Wheat germ			0.07885	
^a based on rotational crops Definition of the residue for compliance with the MRL for plant commodities: <i>Bixafen</i> . Definition of the residue for compliance with the MRL for animal commodities and dietary risk assessment for plant and animal commodities: <i>Sum of bixafen and N-(3',4'-dichloro-5-fluorobiphenyl-2-yl)-3-(difluoromethyl)-1H-pyrazole-4-carboxamide (bixafen-desmethyl), expressed as bixafen</i> . The residue is fat-soluble.						
Clofentezine (156) ADI: 0–0.02 mg/kg bw ARfD: Unnecessary	DH 1100	Hops, Dry	7		2.2	
Definition of the residue for compliance with the MRL and dietary risk assessment for plant commodities: <i>Clofentezine</i> . Definition of the residue for compliance with the MRL and dietary risk assessment for animal commodities: <i>Sum of clofentezine and all metabolites containing the 2-chlorobenzoyl moiety, expressed as clofentezine</i> . The residue is fat-soluble.						
Clothianidin (238) ADI: 0–0.1 mg/kg bw ARfD: 0.6 mg/kg bw	CF 0640	Barley bran, Processed	0.15 c,T		0.028 c,T	
		Barley hay	1 (dw)		Median: 0.09 (as)	Highest: 0.72 (as)
	AS 0640	Barley straw and fodder, Dry	1 (dw) c,T	0.2	Median: 0.09 (as) c,T	Highest: 0.72 (as) c,T
	GC 0640	Barley	0.07 c,T	0.04	0.015 c,T	
	PE 0112	Eggs	0.01(*)	0.01(*)	0.0062	0.0062

Pesticide (Codex reference number)	CCN	Commodity	Recommended Maximum residue level (mg/kg)		STMR or STMR-P mg/kg	HR or HR-P mg/kg
			New	Previous		
	MO 0099	Liver of cattle, goats, pigs & sheep	0.4	0.2	0.257	0.39
	ML 0106	Milks	0.05	0.02	0.041	
	AS 0647	Oat straw and fodder, Dry	1 (dw) c,T		Median: 0.09 (as) c,T	Highest: 0.72 (as) c,T
	GC 0647	Oats	0.07 c,T		0.015 c,T	
		Oat hay	1 (dw)		Median: 0.09 (as)	Highest: 0.72 (as)
	PO 0111	Poultry, edible offal of	0.4	0.1	0.37	0.37
	PF 0111	Poultry fats	0.01(*)	0.01(*)	0.0033	0.0033
	PM 0110	Poultry meat	0.01(*)	0.01(*)	0.0014	0.0014
	CM 1206	Rice bran, Unprocessed	1 C,t		0.28 C,t	
	CM 1207	Rice hulls	4 C,t		1.1 C,t	
	AS 0649	Rice straw and fodder, Dry	0.2 (dw) C,t		Median: 0.03 (as) C,t	Highest: 0.13 (as) C,t
	GC 0649	Rice	0.9 C,t		0.3 C,t	
	CM 0649	Rice, husked	0.5 C,t	0.5	0.145	
	CM 1205	Rice, Polished	0.5 C,t		0.12 C,t	
		Rye hay	1 (dw)		Median: 0.09 (as)	Highest: 0.72 (as)
	AS 0651	Sorghum straw and fodder, Dry	0.8 (dw) c,T	0.01	Median: 0.0885 (as) c,T	Highest: 0.46 (as) c,T
	GC 0651	Sorghum Grain	0.15 c,T	0.01(*)	0.018 c,T	
	GS 0658	Sorgo or Sorghum, Sweet	0.4 T		0.0645 T	0.2 T
	GC 2090	Sweet Corns, Subgroup of	0.01(*) C,t		0.01 C,t	0.01 C,t
	GC 0447	Sweet corn (Corn-on-the-cob) (kernels plus cob with husk removed)	W	0.01(*)		
	AS 0447	Sweet corn fodder	0.05 (dw) C,t		Median: 0.01 (as) c,T	Highest: 0.021 (as) c,T
	GC 0653	Triticale	0.15 c,T		0.01 c,T	
		Triticale hay	1 (dw)		Median: 0.09 (as)	Highest: 0.72 (as)
	AS 0653	Triticale straw and fodder, Dry	1 (dw) c,T		Median: 0.09 (as) c,T	Highest: 0.72 (as) c,T
	CF 0654	Wheat bran, Processed	6 c,T		0.0185 c,T	
	CF 1210	Wheat germ	6 c,T	0.02(*)	0.018 c,T	
	AS 0654	Wheat straw and fodder, Dry	1 (dw) c,T	0.2	Median: 0.09 (as) c,T	Highest: 0.72 (as) c,T
	GC 0654	Wheat	0.15 c,T	0.02	0.01 c,T	
		Wheat hay	1 (dw)		Median: 0.09 (as)	Highest: 0.72 (as)
		Wheat and triticale flour			0.00645	
		Barley flour			0.0097	
		Sorghum (grain) flour			0.0114	
		Sorghum (sweet) syrup			0.059	
T = based on thiamethoxam use only; c,T or C,t = combined clothianidin and thiamethoxam use						
Definition of the residue for compliance with the MRL and for dietary risk assessment for plant and animal commodities: <i>Clothianidin</i> .						
The residue is not fat-soluble.						
Cyprodinil (207) ADI: 0–0.03 mg/kg bw ARfD: Unnecessary	VR 0604	Ginseng	0.3		0.045	
	DV 0604	Ginseng, dried including red ginseng	3		0.114	
	VP 2061	Peas with pods, Subgroup of	2		0.60	
	VD 0071	Beans (dry)	W	0.2		
	VD 2065	Dry beans, Subgroup of (except soya beans)	0.2		0.03	
	VD 2066	Dry peas, Subgroup of	0.2		0.054	
Definition of the residue for compliance with the MRL and dietary risk assessment for plant and animal commodities: <i>Cyprodinil</i> .						
The residue is fat-soluble						
Difenoconazole (224) ADI: 0–0.01 mg/kg bw ARfD: 0.3 mg/kg bw	FT 0336	Guava	0.15		0.0335	0.095
	FB 0265	Cranberry	0.6		0.2	0.26
	SO 0691	Cottonseed	0.4		0.021	
	DT 1114	Tea, green, black (black, fermented and dried)	20		4.85	
	OR 0691	Cotton seed oil, edible			0.0014	
		Tea infusion			0.0072	

Pesticide (Codex reference number)	CCN	Commodity	Recommended Maximum residue level (mg/kg)		STMR or STMR-P mg/kg	HR or HR-P mg/kg
			New	Previous		
Definition of the residue for compliance with the MRL and dietary risk assessment for plant commodities: <i>Difenoconazole</i> .						
Definition of the residue for compliance with the MRL and dietary risk assessment for animal commodities: <i>Sum of difenoconazole and 1-[2-chloro-4-(4-chloro-phenoxy)-phenyl]-2-(1,2,4-triazol)-1-yl-ethanol</i> , expressed as <i>difenoconazole</i> .						
The residue is fat-soluble.						
Ethion (034)						
ADI: 0.002 mg/kg bw						
ARfD: 0.02 mg/kg bw						
Definition of the residue for compliance with the MRL and dietary risk assessment for plant and animal commodities: <i>Ethion</i> .						
The residue is fat-soluble.						
Ethiprole (304)	VD 0541	Soya bean (dry)	0.05		0.0065	
ADI: 0–0.005 mg/kg bw	AB 0541	Soya bean hulls	0.4		0.045	
ARfD: 0.005 mg/kg bw	OR 0541	Soya bean oil, refined			0.002	
		Soya bean flour			0.002	
		Soya milk			0.0007	
Definition of the residue for compliance with the MRL for plant commodities: <i>Ethiprole</i> .						
Definition of the residue for dietary risk assessment for plant commodities: <i>Sum of ethiprole, 5-amino-1-[2,6-dichloro-4-(trifluoromethyl)phenyl]-4-(ethylsulfinyl)-1H-pyrazole-3-carboxamide (ethiprole-amide) and 5-amino-1-(2,6-dichloro-4-trifluoromethylphenyl)-4-ethylsulfonylpyrazole-3-carbonitrile (ethiprole-sulfone)</i> , expressed as parent equivalents.						
Definition of the residue for compliance with the MRL and dietary risk assessment for animal commodities: <i>Sum of ethiprole and 5-amino-1-(2,6-dichloro-4-trifluoromethylphenyl)-4-ethylsulfonylpyrazole-3-carbonitrile (ethiprole-sulfone)</i> , expressed as parent equivalents.						
The residue is fat-soluble						
Fenbuconazole (197)	DT 1114	Tea, green, black (black, fermented and dried)	30		4.2	
ADI: 0–0.03 mg/kg bw		Tea infusion			0.0018	
ARfD: 0.2 mg/kg bw						
Definition of the residue for compliance with the MRL and dietary risk assessment for plant and animal commodities: <i>Fenbuconazole</i> .						
The residue is not fat-soluble.						
Fenhexamid (215)	VS 0621	Asparagus	0.02		0.02	
ADI: 0–0.2 mg/kg bw	VA 2031	Bulb onions, Subgroup of	3		0.055	
ARfD: Unnecessary	FP 0230	Pear	6 Po		2.05	
Definition of the residue for compliance with the MRL and dietary risk assessment in plant and animal commodities: <i>Fenhexamid</i> .						
The residue is fat-soluble						
Fenpicoxamid (305)	MO 0105	Edible offal (mammalian)	0.02		0.013	
ADI: 0–0.05 mg/kg bw	MF 0100	Mammalian fats (except milk fats)	0.015		0.013	
ARfD: Unnecessary	MM 0095	Meat (from mammals other than marine mammals)	0.015(*)		0	
	ML 0106	Milks	0.015(*)		0	
	GC 0650	Rye	0.15		0.0215	
	GC 0653	Triticale	0.15		0.0215	
	AS 0654	Wheat straw and fodder, dry	30 (dw)		Median: 4.7 (as)	Highest: 12 (as)
	GC 0654	Wheat	0.15		0.0215	
	GC 0653	Triticale flour (white and wholemeal)			0.0088 ^a	
	CF 1210	Wheat germ			0.0036	
	CP 1212	Wheat wholemeal bread			0.0043	
	CP 1211	Wheat white bread			0.0032	
	CF 1211	Wheat white flour			0.0062	
	CF 1250	Rye flour (white and wholemeal)			0.0088 ^a	
	-	Wheat, bulgur			0.0088 ^a	
	-	Wheat starch			0.0032	
	-	Wheat gluten			0.0058	
^a PF for wheat wholemeal flour was used.						
Definition of the residue for compliance with the MRL and dietary risk assessment for plant commodities: <i>Fenpicoxamid</i> .						
Definition of the residue for compliance with the MRL and dietary risk assessment for milk and tissues of mammals (other than marine mammals): <i>2-benzyl-2,5-dideoxy-4-O-[N-1-(3-hydroxy-4-methoxypyridin-2-yl)carbonyl]-L-seryl]-L-arabinoic acid (X12326349)</i> , expressed as <i>fenpicoxamid</i> .						

Pesticide (Codex reference number)	CCN	Commodity	Recommended Maximum residue level (mg/kg)		STMR or STMR-P mg/kg	HR or HR-P mg/kg
			New	Previous		
as fenpicoxamid.						
The residue is not fat-soluble						
Fluopyram (243) ADI: 0–0.01 mg/kg bw ARfD: 0.5 mg/kg bw	SB 0716	Coffee beans	0.015		0.01	
	SM 0716	Coffee beans, roasted			0.01	
		Instant coffee			0.01	
Definition of the residue for compliance with the MRL and dietary risk assessment for plant commodities: Fluopyram.						
Definition of the residue for compliance with the MRL for animal commodities: Sum of fluopyram and 2(trifluoromethyl)benzamide, expressed as fluopyram.						
Definition of the residue for dietary risk assessment for animal commodities: Sum of fluopyram, 2(trifluoromethyl)benzamide and the combined residues of the E-olefine and Z-olefine isomers of fluopyram, all expressed as fluopyram.						
Although fluopyram (parent compound) is fat-soluble, the 2-(trifluoromethyl)benzamide metabolite (the major component of the residue) is not fat-soluble.						
Imazalil (110) ADI: 0–0.03 mg/kg bw ARfD: 0.05 mg/kg bw	FC 0001	Citrus Fruit, Group of	15 Po		0.07 (except kumquats) 3.4 (kumquats)	0.36 (except kumquats) 9.9 (kumquats)
	FC 0002	Lemons and limes, Subgroup of	W	15 Po		
	FC 0004	Oranges, Sweet, Sour, Subgroup of	W	8 Po		
	AB 0001	Citrus pulp, dry	70 (dw) Po		15 (dw)	
	OR 0001	Citrus oil, edible	500 Po		97	
	JF 0001	Citrus juice			0.34	
		Citrus canned			0.1	
		Citrus marmalade			0.92	
		Citrus peel (chopped)			1.3	
Definition of the residue for compliance with the MRL for plant and animal commodities: imazalil.						
Definition of the residue for dietary risk assessment for plant commodities: free and conjugated imazalil.						
Definition of the residue for dietary risk assessment for animal commodities: sum of imazalil and the metabolite R061000 ((RS)-3-[2-(2,4-dichlorophenyl)-2-(2,3-dihydroxypropoxy)ethyl]imidazolidine- 2,4-dione (+)-1-[2-(2,4-dichlorophenyl)-2-[(2,3-dihydroxypropyl)oxy]ethyl]-2,5-imidazolidinedione), expressed as imazalil equivalents.						
The residue is not fat-soluble						
Isoprothiolane (299) ADI: 0–0.1 mg/kg bw ARfD: Unnecessary	FI 0327	Banana	1		0.0077	
Definition of the residue for compliance with the MRL for plant commodities: Isoprothiolane.						
Definition of the residue for dietary risk assessment for rice: Isoprothiolane.						
Definition of the residue for dietary risk assessment for plants other than rice: Sum of isoprothiolane, diisopropyl-4-hydroxy-1,3-dithiolan-2-ylidenemalonate (M-3); free and conjugated and 1-hydroxypropan-2-yl propan-2-yl 1,3-dithiolan-2-ylidenemalonate (M-5); free and conjugated, expressed as isoprothiolane.						
Definition of the residue for compliance with the MRL and dietary risk assessment for animal commodities: sum of isoprothiolane and 2-(1,3-dithiolan-2-ylidene)-3-oxo-3-(propan-2-yloxy)propanoic acid (M-2), expressed as isoprothiolane.						
The residue is not fat-soluble						
Isoxaflutole (268) ADI: 0–0.02 mg/kg bw ARfD: Unnecessary	VD 0541	Soya bean (dry)	0.04		0.02	
	OR 0541	Soya bean oil, refined			0.008	
		Soya bean milk			0.008	
Definition of the residue for compliance with the MRL and dietary risk assessment for plant commodities: Sum of isoxaflutole and isoxaflutole diketonitrile, expressed as isoxaflutole.						
Definition of the residue for compliance with the MRL for animal commodities: Sum of isoxaflutole and isoxaflutole diketonitrile, expressed as isoxaflutole.						
Definition of the residue for dietary risk assessment for animal commodities: sum of isoxaflutole, isoxaflutole diketonitrile, RPA 205834 (2-aminomethylene-1-cyclopropyl-3-(2-mesyl-4-trifluoromethylphenyl)-propane-1,3-dione) and RPA 207048 (1-cyclopropyl-2-hydroxymethylene-3-(2-mesyl-4-trifluoromethylphenyl)-propane-1,3-dione), including their conjugates, expressed as isoxaflutole.						
The residue not fat-soluble.						
Mandipropamid (231) ADI: 0–0.2 mg/kg bw	OR 0001	Citrus oil, edible	30		4.5	
	AB 0001	Citrus pulp, dry	1.5		Median: 0.29	

Pesticide (Codex reference number)	CCN	Commodity	Recommended Maximum residue level (mg/kg)		STMR or STMR-P mg/kg	HR or HR-P mg/kg
			New	Previous		
ARfD: Unnecessary	MO 0105	Edible offal (mammalian)	0.01(*)	0.01(*)	0.0022	
	FC 0002	Lemons and Limes, Subgroup of	0.5		0.01	
	MF 0100	Mammalian fats (except milk fats)	0.02	0.01(*)	0.0064	
	FC 0003	Mandarins, Subgroup of	0.5		0.01	
	FC 0004	Oranges, Sweet, Sour (including Orange-like hybrids), Subgroup of	0.4		0.01	
	FC 0005	Pummelo and Grapefruits (including Shaddock-like hybrids, among other Grapefruit), Subgroup of	0.2		0.01	
		Citrus flesh (excluding kumquat commodities)			0.01	
	JF 0203	Grapefruit juice			0.0024	
	FC 0303	Kumquat, raw (including juice)			0.098	
		Lemon/lime/mandarin juice			0.0042	
	JF 0004	Orange juice			0.0043	
Definition of the residue for compliance with the MRL and dietary risk assessment for plant and animal commodities: <i>Mandipropamid</i> . The residue is not fat-soluble.						
Methoprene (147) ADI: 0-0.09 mg/kg bw for the R,S racemate; 0- 0.05 mg/kg bw for S-methoprene ARfD: Unnecessary	VD 0541	Soya bean (dry)	3 Po		2.4	
Definition of the residue for compliance with the MRL and dietary risk assessment for plant and animal commodities: <i>Methoprene</i> . The residue is fat-soluble						
Methoxyfenozide (209) ADI: 0-0.1 mg/kg bw ARfD: 0.9 mg/kg bw	DH 0722	Basil, dry	400		100	194
	HH 0722	Basil, leaves	80		19	47
	SB 0716	Coffee bean	0.15		0.0275	
	GS 0659	Sugar cane	0.015		0.01	0.01
	DM 0659	Sugar cane molasses	0.1		0.07	
	DT 1114	Tea, Green, Black (black, fermented and dried)	80		28.5	
		Brown sugar			0.06	
		Tea infusion			0.14	
Definition of the residue for compliance with the MRL and dietary risk assessment for plant and animal commodities: <i>Methoxyfenozide</i> . The residue is fat-soluble, but is not classified as fat-soluble with respect to its distribution in milk.						
Prothioconazole (232) ADI: 0-0.05 mg/kg bw ARfD: 0.8 mg/kg bw (for women of child-bearing age) ARfD: Unnecessary (general population) Prothioconazole-desthio: ADI: 0-0.01 mg/kg bw ARfD: 0.01 mg/kg bw (woman of child-bearing age) ARfD: 1 mg/kg bw (general population)	MO 0105	Edible offal (Mammalian)	0.15	0.3	Liver: 0.077 Kidney: 0.049	Liver: 0.291 Kidney: 0.191
	PE 0112	Eggs	0.005(*)	0.005(*)	0.001	0.001
	SO 0693	Linseed	0.03		0.014	
	MF 0100	Mammalian fats (except milk fats)	0.01	0.02	0.005	0.023
	MM 0095	Meat (from mammals other than marine mammals)	0.01	0.01	Muscle: 0.004 Fat: 0.005	Muscle: 0.007 Fat: 0.023
	ML 0106	Milks	0.004(*)	0.004(*)	0.001	
	PO 0111	Poultry, Edible offal of	0.01	0.1	0.118	0.118
	PF 0111	Poultry, fats	0.01	0.01(*)	0.0129	0.0129
	PM 0110	Poultry meat	0.01(*)	0.01(*)	Muscle: 0.0026 Fat: 0.0129	Muscle: 0.0026 Fat: 0.0129
	SO 0495	Rape seed	0.2	0.1	0.039	
	OR 0495	Rape seed oil, Edible	0.15		0.0273	
	SO 2091	Sunflower seeds, Subgroup of	0.5		0.009	
	OC 0702	Sunflower seed oil, crude	0.5		0.0096	

Pesticide (Codex reference number)	CCN	Commodity	Recommended Maximum residue level (mg/kg)		STMR or STMR-P mg/kg	HR or HR-P mg/kg
			New	Previous		
Definition of the residue for compliance with the MRL and dietary risk assessment for plant commodities: <i>Prothioconazole-desthio</i> .						
Definition of the residue for compliance with the MRL for animal commodities: <i>Prothioconazole-desthio</i> .						
Definition of the residue for dietary risk assessment for animal commodities: <i>Sum of prothioconazole-desthio, prothioconazole-desthio-3-hydroxy, prothioconazole-desthio-4-hydroxy and their conjugates expressed as prothioconazole-desthio</i> .						
The residue is not fat-soluble						
Pydiflumetofen (309) ADI: 0–0.1 mg/kg bw ARfD: 0.3 mg/kg bw	AM 0660	Almond hulls	10 (dw)		Median: 1.6 (as)	
	VP 2060	Beans with pods, Subgroup of	0.7		0.045	0.47
	VB 0040	Brassica vegetables (except Brassica leafy vegetables), Group of	W	0.1		
	VA 2031	Bulb onions, Subgroup of	0.3		0.07	0.20
	FB 2006	Bush berries, Subgroup of	5		0.88	3.9
	FS 0013	Cherries, Subgroup of	2		0.395	1.7
	FC 0001	Citrus Fruit, Group of	0.9		0.05 (except kumquats) 0.21 (kumquats)	0.16 (except kumquats) 0.76 (kumquats)
	OR 0001	Citrus oil, edible	40		9.1	
	AB 0001	Citrus pulp, dry	1.5		0.30	
	SO 0691	Cottonseed	0.02 ^a	0.3	0.02 ^a	
	MO 0105	Edible offal (mammalian)	0.1	0.1	Liver: 0.09 Kidney: 0.09	Liver: 0.44 Kidney: 0.30
	PE 0112	Eggs	0.02	0.02	0.02	0.023
	FB 0267	Elderberries	5		0.88	3.9
	VB 0042	Flowerhead Brassicas, Subgroup of	3		0.39	1.5
	VA 2032	Green onions, Subgroup of	1.5		0.36	1.39
	VB 2036	Head Brassicas, Subgroup of	2		0.065	0.22
	VL 2052	Leaves of root and tuber vegetables, Subgroup of (except leaves of tuber vegetables)	W	0.07		
	VP 0060	Legume vegetables, Group of	W	0.02		
	FB 2009	Low growing berries, Subgroup of (except cranberries)	1		0.185	0.62
	MF 0100	Mammalian fats (except milk fats)	0.1	0.1	0.02	0.07
	MM 0095	Meat (from mammals other than marine mammals)	0.1 (fat)	0.1 (fat)	Muscle: 0.02 Fat: 0.02	Muscle: 0.02 Fat: 0.07
	ML 0106	Milks	0.01(*)	0.01(*)	0.02	
	FS 2001	Peaches, Subgroup of	1		0.21	0.80
	VP 2061	Peas with pods, Subgroup of	1.5		0.12	0.84
	DF 0014	Prunes	1.5		0.34	0.85
	FS 0014	Plums, Subgroup of	0.6		0.15	0.37
	FP 0009	Pome fruit, Group of, except Persimmon, Japanese	0.2		0.06	0.13
	PO 0111	Poultry, Edible offal of	0.01(*)	0.01(*)	0.02 (liver)	0.02 (liver)
	PF 0111	Poultry fats	0.01(*)	0.01(*)	0.02	0.02
	PM 0110	Poultry meat	0.01(*)	0.01(*)	0.02	0.02
	VR 2070	Root vegetables, Subgroup of	0.3	0.1	0.08	0.25
	GC 2089	Sorghum Grain and Millet, Subgroup of	W	0.03		
	GC 2089	Sorghum Grain and Millet, Subgroup of, except grain sorghum	0.03		0.03	
	GC 0651	Sorghum Grain	3		0.515	
	AS 0651	Sorghum straw and fodder, dry	10 (dw)	0.3 (dw)	Median: 0.50 (dw)	Highest: 7.6 (dw)
	VB 2016	Stem Brassicas, Subgroup of	0.1 ^a		0.02 ^a	0.09 ^a
	VP 2062	Succulent beans without pods, Subgroup of	0.15		0.033	0.099
	VP 2063	Succulent peas without pods, Subgroup of	0.05		0.031	0.042
	AV 0596	Sugar beet leaves or tops	40 (dw)		Median: 1.50 (as)	Highest: 8.0 (as)
	SO 2091	Sunflower seeds, Subgroup of	0.5	0.3	0.09	
	TN 0085	Tree nuts, Group of	0.05		0.01	0.03
	VP 2064	Underground immature beans and peas, Subgroup of	0.02 ^a		0.02 ^a	0.02 ^a
	JF 0226	Apple juice			0.004	
		Apple sauce			0.004	
		Apple, canned			0.002	

Pesticide (Codex reference number)	CCN	Commodity	Recommended Maximum residue level (mg/kg)		STMR or STMR-P mg/kg	HR or HR-P mg/kg
			New	Previous		
	DF 0226	Apples, dried			0.02	0.05
	JF 0001	Citrus juice			0.004	
		Citrus peel			0.39	1.4
	OR 0691	Cotton seed oil (refined)			0.0006 ^a	
		Pear, canned			0.004	
		Pear, dried			0.03	0.08
		Pear juice			0.006	
		Sugar beet refined sugar			0.004	
		Sorghum flour			0.45	
	OR 0702	Sunflower oil, edible			0.005	
^a based on rotational crops						
Definition of the residue for compliance with the MRL and dietary risk assessment for plant commodities: <i>Pydiflumetofen</i> .						
Definition of the residue for compliance with the MRL for animal commodities: <i>Pydiflumetofen</i> .						
Definition of the residue for dietary risk assessment for animal commodities other than mammalian liver and kidney: <i>Sum of pydiflumetofen and 2,4,6-trichlorophenol (2,4,6-TCP) and its conjugates, expressed as pydiflumetofen</i> .						
Definition of the residue for dietary risk assessment for mammalian liver and kidney: <i>Sum of pydiflumetofen, 2,4,6-trichlorophenol (2,4,6-TCP) and its conjugates and 3-(difluoromethyl)-N-methoxy-1-methyl-N-[1-methyl-2-(2,4,6-trichloro-3-hydroxy-phenyl) ethyl]pyrazole-4-carboxamide (SYN547897) and its conjugates, expressed as pydiflumetofen</i> .						
The residue is fat-soluble						
Quinoxifen (222) ADI: 0–0.2 mg/kg bw ARfD: Unnecessary	FS 0013	Cherries	W	0.4		
	FS 0013	Cherries, Subgroup of (except Choke cherries)	0.5		0.14	
Definition of the residue for compliance with the MRL and dietary risk assessment: <i>Quinoxifen</i> .						
The residue is fat-soluble.						
Spinetoram (233) ADI: 0–0.05 mg/kg bw ARfD: Unnecessary	FI 2540	Pitaya	0.5		0.0915	
	DT 1114	Tea, Green, Black (black, fermented and dried)	70		118	
Definition of the residue for compliance with the MRL for plant and animal commodities: <i>Spinetoram</i> .						
Definition of the residue for dietary risk assessment for plant and animal commodities: <i>Spinetoram and N-demethyl and N-formyl metabolites of the major spinetoram component</i> .						
The residue is fat-soluble.						
Sulfoxaflor (252) ADI: 0–0.05 mg/kg bw ARfD: 0.3 mg/kg bw	VS 0621	Asparagus	0.015		0.01	0.011
	FI 0326	Avocado	0.15		0.011	0.036
	FB 2006	Bush berries, Subgroup of	2		0.39	1.4
	FB 2005	Cane berries, Subgroup of	1.5		0.44	0.78
	SB 0716	Coffee beans	0.3		0.035	
	FB 0267	Elderberry	2		0.39	1.4
	FI 0345	Mango	0.3		0.022	0.061
		Coffee (instant)			0.084	
	SM 0716	Coffee beans, roasted			0.0167	
Definition of the residue for compliance with the MRL and dietary risk assessment for plant and animal commodities: <i>Sulfoxaflor</i> .						
The residue is not fat-soluble						
Tebuconazole (189) ADI: 0–0.03 mg/kg bw ARfD: 0.3 mg/kg bw	SB 0716	Coffee beans	0.4	0.1	0.04	
	SM 0716	Coffee beans, roasted			0.08	
		Freeze dried (instant) coffee			0.032	
Definition of the residue for compliance with the MRL and dietary risk assessment for plant and animal commodities: <i>Tebuconazole</i> .						
Residue is not fat-soluble						
Thiamethoxam (245) ADI: 0–0.08 mg/kg bw ARfD: 1 mg/kg bw	CF 0640	Barley bran	1.5		0.269	
		Barley hay	8 (dw)		0.2 (as)	6.7 (as)
	AS 0640	Barley straw and fodder, dry	8 (dw)	2	0.53 (as)	6.7 (as)
	GC 0640	Barley	0.5		0.112	
	MO 0105	Edible offal (mammalian)	0.05	0.01(*)	0.025	0.041
	PE 0112	Eggs	0.01(*)	0.01(*)	0.028	0.028
	MF 0100	Mammalian fats (except milk fats)	0.01(*)		0.01	0.01
	MM 0095	Meat (from mammals other than marine mammals)	0.07	0.02	0.033	0.062

Pesticide (Codex reference number)	CCN	Commodity	Recommended Maximum residue level (mg/kg)		STMR or STMR-P mg/kg	HR or HR-P mg/kg
			New	Previous		
	ML 0106	Milks	0.15	0.05	0.096	
		Oat hay	8 (dw)		0.2 (as)	6.7 (as)
	AS 0647	Oat straw and fodder, dry	8 (dw)		0.53 (as)	6.7 (as)
	GC 0647	Oats	0.5		0.112	
	FP 0307	Persimmon, Japanese	0.6		0.165	0.26
	PF 0111	Poultry fats	0.01(*)		0.033	0.033
	PM 0110	Poultry meat	0.03	0.01(*)	0.064	0.064
	PO 0111	Poultry, edible offal of	0.01(*)	0.01(*)	0.36	0.36
	GC 0649	Rice	50		17	
	CM 1206	Rice bran	30		9.35	
	CM 1207	Rice hulls	300		95.2	
	AS 0649	Rice straw and fodder, dry	3 (dw)		0.47 (as)	1 (as)
	CM 0649	Rice, husked	5		1.7	
	CM 1205	Rice, polished	3		0.704	
		Rye hay	8 (dw)		0.2 (as)	6.7 (as)
	AS 0651	Sorghum straw and fodder (dry)	0.8 (dw)		0.14 (as)	0.49 (as)
	GC 0651	Sorghum grain	0.6		0.079	
	GS 0658	Sorgo or sorghum, sweet	0.6		0.033	0.24
	GC 2090	Sweet corns, Subgroup of	0.01(*)		0.01	0.01
	AS 0447	Sweet corn fodder	0.25 (dw)		0.0165 (as)	0.099 (as)
	GC 0653	Triticale	0.15		0.032	
		Triticale hay	8 (dw)		0.2 (as)	6.7 (as)
	AS 0653	Triticale straw and fodder, dry	8 (dw)		0.53 (as)	6.7 (as)
	GC 0654	Wheat	0.15		0.032	
	CF 0654	Wheat bran	0.4		0.0768	
	CF 1210	Wheat germ	0.3		0.0464	
		Wheat hay	8 (dw)		0.2 (as)	6.7 (as)
	AS 0654	Wheat straw and fodder, dry	8 (dw)	2	0.53 (as)	6.7 (as)
		Barley flour			0.0118	
		Sorghum (grain) flour			0.0548	
		Sorghum (sweet) syrup			0.0267	
		Triticale flour			0.00336	
		Wheat and triticale flour			0.00336	
Definition of the residue for compliance with the MRL for animal and plant commodities: <i>Thiamethoxam</i> .						
Definition of the residue for dietary risk assessment for plant and animal commodities (except poultry): <i>Thiamethoxam</i> and <i>clothianidin</i> (considered separately).						
Definition of the residue for dietary risk assessment for poultry: <i>Sum of thiamethoxam, CGA 265307 and MU3, expressed as thiamethoxam; and clothianidin</i> (clothianidin to be considered separately from thiamethoxam).						
The residue is not fat-soluble						
Trifloxystrobin (213) ADI: 0–0.04 mg/kg bw ARfD: Unnecessary	VP 2060	Beans with pods, Subgroup of	0.5		0.09	
	FB 0261	Bilberry	3		0.33	
	FB 0263	Bilberry, red	3		0.33	
	FB 0020	Blueberries	3		0.33	
	FB 2005	Cane berries, Subgroup of	3		0.56	
	FC 0001	Citrus fruits	W	0.5		
	AB 0001	Citrus pulp, dry	W	1		
	SB 0716	Coffee beans	0.015		0.01	
	FB 0021	Currant, black, red, white	3		0.33	
	VL 0470	Corn salad	15		3.3	
	MO 0105	Edible offal (mammalian)	0.09		Kidney 0.04 Liver 0.04	
	PE 0112	Eggs	0.04(*)	0.04(*)	0.0046	
	FB 0268	Gooseberry	3		0.33	
	MO 0098	Kidney of cattle, goats, pig and sheep	W	0.04(*)		
	VL 0483	Lettuce, leaf	15		3.3	
	SO 0693	Linseed	0.4		0.015	
	MO 0099	Liver of cattle, goats, pigs & sheep	W	0.05		
	MF 0100	Mammalian fats (except milk fats)	0.07		0.04	
	MM 0095	Meat (from mammals other than marine mammals)	0.07 (fat)	0.05 (fat)	Fat 0.04 Muscle 0.007	
	ML 0106	Milks	0.02(*)	0.02(*)	0.003	
	VP 2061	Peas with pods, Subgroup of	1.5		0.073	
	PF 0111	Poultry fats	0.04(*)		0.0046	

Pesticide (Codex reference number)	CCN	Commodity	Recommended Maximum residue level (mg/kg)		STMR or STMR-P mg/kg	HR or HR-P mg/kg
			New	Previous		
	PM 0110	Poultry meat	0.04(*) (fat)	0.04(*) (fat)	0.0046	
	PO 0111	Poultry, edible offal of	0.04(*)	0.04(*)	0.0046	
	FB 0273	Rose hips	3		0.33	
		Coffee, instant			0.0015	
	SM 0716	Coffee beans, roasted			0.0015	
		Linseed oil, refined			0.012	
Definition of the residue for compliance with the MRL for plant commodities: <i>Trifloxystrobin</i> .						
Definition of the residue for dietary risk assessment for plant commodities: <i>Sum of trifloxystrobin and [(E,E)-methoxyimino-{2-[1-(3-trifluoromethylphenyl)ethylideneaminooxymethyl]phenyl}acetic acid]</i> (CGA 321113), expressed as <i>trifloxystrobin</i> .						
Definition of the residue for compliance with the MRL dietary risk assessment for animal commodities: <i>Sum of trifloxystrobin and [(E,E)-methoxyimino-{2-[1-(3-trifluoromethylphenyl)ethylideneamino-oxymethyl]phenyl}acetic acid]</i> (CGA 321113), expressed as <i>trifloxystrobin</i> .						
The residue is fat-soluble						
Trinexapac-ethyl (271) ADI: 0–0.3 mg/kg bw ARfD: Unnecessary	CF 0640	Barley bran, processed	4	6	0.63	
	GC 0649	Rice	0.5		0.064	
	CM 1205	Rice polished	0.7		0.08	
		Rice bran	3		0.33	
	AS 0649	Rice straw and fodder, dry	0.08 (dw)		Median: 0.014 (as)	Highest: 0.045 (as)
	GC 0650	Rye	3		0.57	
	AS 0650	Rye straw and fodder, dry	0.9 (dw)		Median: 0.19 (dw)	Highest: 1.34 (dw)
	CM 0654	Wheat bran, unprocessed	5	8	0.68	
		Barley flour			0.25	
		Barley, pearled			0.30	
		Barley malt			0.28	
		Barley beer			0.03	
		Wheat, white flour			0.23	
	CF 1210	Wheat germ			0.38	
		Wheat gluten			0.14	
		Wheat starch			0.046	
		Wheat whole meal bread			0.35	
	Definition of the residue for compliance with the MRL for plant and animal commodities and for dietary risk assessment for animal commodities: <i>Trinexapac acid</i> .					
Definition of the residue for dietary risk assessment for plant commodities: <i>Trinexapac acid and its conjugates, expressed as trinexapac acid</i> .						
The residue is not fat-soluble.						

Annex 2: Index of reports and evaluations of pesticides by the JMPR

Numbers in parentheses after the names of pesticides are Codex classification numbers. The abbreviations used are:

T, evaluation of toxicology

R, evaluation of residue and analytical aspects

E, evaluation of effects on the environment

Abamectin (177)	1992 (T,R), 1994 (T,R), 1995 (T), 1997 (T,R), 2000 (R), 2015 (R), 2017 (T), 2018 (R)
Acephate (095)	1976 (T,R), 1979 (R), 1981 (R), 1982 (T), 1984 (T,R), 1987 (T), 1988 (T), 1990 (T,R), 1991 (corr. to 1990 R evaluation), 1994 (R), 1996 (R), 2002 (T), 2003 (R), 2004 (corr. to 2003 report), 2005 (T), 2006 (R), 2011 (R)
Acetamiprid (246)	2011 (T, R), 2012 (R), 2015 (R), 2017 (R), 2021 (R)
Acetochlor (280)	2015 (T, R), 2019 (T, R)
Acibenzolar-S-methyl (288)	2016 (T, R)
Acrylonitrile	1965 (T, R)
Aldicarb (117)	1979 (T, R), 1982 (T,R), 1985 (R), 1988 (R), 1990 (R), 1991 (corr. to 1990 evaluation), 1992 (T), 1993 (R), 1994 (R), 1996 (R), 2001 (R), 2002 (R), 2006 (R)
Aldrin (001)	1965 (T), 1966 (T, R), 1967 (R), 1974 (R), 1975 (R), 1977 (T), 1990 (R), 1992 (R)
Allethrin	1965 (T, R)
Ametoctradin (253)	2012 (T, R)
Aminocarb (134)	1978 (T, R), 1979 (T, R)
Aminocyclopyrachlor (272)	2014 (T, R)
Aminomethylphosphonic acid (AMPA, 198)	1997 (T, R)
Aminopyralid (220)	2006 (T, R), 2007 (T,R)
Amitraz (122)	1980 (T, R), 1983 (R), 1984 (T,R), 1985 (R), 1986 (R), 1989 (R), 1990 (T,R), 1991 (R & corr. to 1990 R evaluation), 1998 (T)
Amitrole (079)	1974 (T, R), 1977 (T), 1993 (T, R), 1997 (T), 1998 (R)
Anilazine (163)	1989 (T, R), 1992 (R)
Atrazine	2007 (T)
Azinphos-ethyl (068)	1973 (T, R), 1983 (R)
Azinphos-methyl (002)	1965 (T), 1968 (T, R), 1972 (R), 1973 (T), 1974 (R), 1991 (T, R), 1992 (corr. to 1991 report), 1993 (R), 1995 (R), 2007 (T)

Azocyclotin (129)	1979 (R), 1981 (T), 1982 (R), 1983 (R), 1985 (R), 1989 (T,R), 1991 (R), 1994 (T), 2005 (T,R)
Azoxystrobin (229)	2008 (T,R), 2011 (R), 2012 (R), 2013 (R), 2017 (R), 2019 (R)
Benalaxyl (155)	1986 (R), 1987 (T), 1988 (R), 1992 (R), 1993 (R), 2005 (T), 2009 (R)
Bendiocarb (137)	1982 (T,R), 1984 (T,R), 1989 (R), 1990 (R)
Benomyl (069)	1973 (T,R), 1975 (T,R), 1978 (T,R), 1983 (T,R), 1988 (R), 1990 (R), 1994 (R), 1995 (T,E), 1998 (R)
Bentazone (172)	1991 (T,R), 1992 (corr. to 1991 report, Annex I), 1994 (R), 1995 (R), 1998 (T,R), 1999 (corr. to 1998 report), 2004 (T), 2012 (T), 2013 (R), 2016 (T), 2018 (R)
Benzovindiflupyr (261)	2013 (T), 2014 (R), 2016 (R)
BHC (technical-grade)	1965 (T), 1968 (T,R), 1973 (T,R) (see also Lindane)
Bicyclopyrone (295)	2017 (T, R)
Bifenazate (219)	2006 (T,R), 2008 (R), 2010 (R)
Bifenthrin (178)	1992 (T,R), 1995 (R), 1996 (R), 1997 (R), 2009 (T), 2010 (R), 2015 (R)
Binapacryl (003)	1969 (T,R), 1974 (R), 1982 (T), 1984 (R), 1985 (T,R)
Bioresmethrin (093)	1975 (R), 1976 (T,R), 1991 (T,R)
Biphenyl	See Diphenyl
Bitertanol (144)	1983 (T), 1984 (R), 1986 (R), 1987 (T), 1988 (R), 1989 (R), 1991 (R), 1998 (T), 1999 (R), 2002 (R)
Bixafen (262)	2013 (T,R), 2016 (R), 2021 (R)
Boscalid (221)	2006 (T,R), 2008 (R), 2010 (R), 2019 (T, R)
Bromide ion (047)	1968 (R), 1969 (T, R), 1971 (R), 1979 (R), 1981 (R), 1983 (R), 1988 (T, R), 1989 (R), 1992 (R)
Bromomethane (052)	1965 (T, R), 1966 (T,R), 1967 (R), 1968 (T,R), 1971 (R), 1979 (R), 1985 (R), 1992 (R)
Bromophos (004)	1972 (T,R), 1975 (R), 1977 (T,R), 1982 (R), 1984 (R), 1985 (R)
Bromophos-ethyl (005)	1972 (T,R), 1975 (T,R), 1977 (R)
Bromopropylate (070)	1973 (T,R), 1993 (T,R)
Butocarboxim (139)	1983 (R), 1984 (T), 1985 (T), 1986 (R)
Buprofezin (173)	1991 (T,R), 1995 (R), 1996 (corr. to 1995 report.), 1999 (R), 2008 (T,R), 2009 (R), 2012 (R), 2014 (R), 2016 (R)
sec-Butylamine (089)	1975 (T,R), 1977 (R), 1978 (T,R), 1979 (R), 1980 (R), 1981 (T), 1984 (T,R: withdrawal of temporary ADI, but no evaluation)
Cadusafos (174)	1991 (T,R), 1992 (R), 1992 (R), 2009 (R), 2010 (R)

Campheclor (071)	1968 (T,R), 1973 (T,R)
Captafol (006)	1969 (T,R), 1973 (T,R), 1974 (R), 1976 (R), 1977 (T,R), 1982 (T), 1985 (T,R), 1986 (corr. to 1985 report), 1990 (R), 1999 (ARfD)
Captan (007)	1965 (T), 1969 (T,R), 1973 (T), 1974 (R), 1977 (T,R), 1978 (T,R), 1980 (R), 1982 (T), 1984 (T,R), 1986 (R), 1987 (R and corr. to 1986 R evaluation), 1990 (T,R), 1991 (corr. to 1990 R evaluation), 1994 (R), 1995 (T), 1997 (R), 2000 (R), 2004 (T), 2007 (T), 2017 (R)
Carbaryl (008)	1965 (T), 1966 (T,R), 1967 (T,R), 1968 (R), 1969 (T,R), 1970 (R), 1973 (T,R), 1975 (R), 1976 (R), 1977 (R), 1979 (R), 1984 (R), 1996 (T), 2001 (T), 2002 (R), 2007 (R)
Carbendazim (072)	1973 (T,R), 1976 (R), 1977 (T), 1978 (R), 1983 (T,R), 1985 (T,R), 1987 (R), 1988 (R), 1990 (R), 1994 (R), 1995 (T,E), 1998 (T,R), 2003 (R), 2005 (T), 2012 (R)
Carbofuran (096)	1976 (T,R), 1979 (T,R), 1980 (T), 1982 (T), 1991 (R), 1993 (R), 1996 (T), 1997 (R), 1999 (corr. to 1997 report), 2002 (T,R), 2003 (R) (See also carbosulfan), 2004 (R), 2008 (T), 2009 (R)
Carbon disulfide (009)	1965 (T,R), 1967 (R), 1968 (R), 1971 (R), 1985 (R)
Carbon tetrachloride (010)	1965 (T,R), 1967 (R), 1968 (T,R), 1971 (R), 1979 (R), 1985 (R)
Carbophenothion (011)	1972 (T,R), 1976 (T,R), 1977 (T,R), 1979 (T,R), 1980 (T,R), 1983 (R)
Carbosulfan (145)	1984 (T,R), 1986 (T), 1991 (R), 1992 (corr. to 1991 report), 1993 (R), 1997 (R), 1999 (R), 2002 (R), 2003 (T,R), 2004 (R, corr. to 2003 report)
Cartap (097)	1976 (T,R), 1978 (T,R), 1995 (T,R)
Chinomethionat (080)	1968 (T,R) (as oxythioquinox), 1974 (T,R), 1977 (T,R), 1981 (T,R), 1983 (R), 1984 (T,R), 1987 (T)
Chlorantraniliprole (230)	2008 (T,R), 2010 (R), 2013 (R), 2014 (R), 2016 (R), 2019 (R)
Chlorbenside	1965 (T)
Chlordane (012)	1965 (T), 1967 (T,R), 1969 (R), 1970 (T,R), 1972 (R), 1974 (R), 1977 (T,R), 1982 (T), 1984 (T,R), 1986 (T)
Chlordimeform (013)	1971 (T,R), 1975 (T,R), 1977 (T), 1978 (T,R), 1979 (T), 1980 (T), 1985 (T), 1986 (R), 1987 (T)
Chlorfenapyr (254)	2013 (T), 2018 (T,R)
Chlorfenson	1965 (T)
Chlorfenvinphos (014)	1971 (T,R), 1984 (R), 1994 (T), 1996 (R)

Chlormequat (015)	1970 (T,R), 1972 (T,R), 1976 (R), 1985 (R), 1994 (T,R), 1997 (T), 1999 (ARfD), 2000 (R), 2017 (T, R)
Chlorobenzilate (016)	1965 (T), 1968 (T,R), 1972 (R), 1975 (R), 1977 (R), 1980 (T)
Chloropicrin	1965 (T,R)
Chloropropylate	1968 (T,R), 1972 (R)
Chlorothalonil (081)	1974 (T,R), 1977 (T,R), 1978 (R), 1979 (T,R), 1981 (T,R), 1983 (T,R), 1984 (corr. to 1983 report and T evaluation), 1985 (T,R), 1987 (T), 1988 (R), 1990 (T,R), 1991 (corr. to 1990 evaluation), 1992 (T), 1993 (R), 1997 (R), 2009 (T), 2010 (R), 2012 (R), 2015 (R), 2019 (T, R)
Chlorpropham (201)	1965 (T), 2000 (T), 2001 (R), 2005 (T), 2008 (R)
Chlorpyrifos (017)	1972 (T,R), 1974 (R), 1975 (R), 1977 (T,R), 1981 (R), 1982 (T,R), 1983 (R), 1989 (R), 1995 (R), 1999 (T), 2000 (R), 2004 (R), 2006 (R), 2021 (R)
Chlorpyrifos-methyl (090)	1975 (T,R), 1976 (R, Annex I only), 1979 (R), 1990 (R), 1991 (T,R), 1992 (T and corr. to 1991 report), 1993 (R), 1994 (R), 2001 (T), 2009 (R)
Chlorthion	1965 (T)
Clethodim (187)	1994 (T,R), 1997 (R), 1999 (R), 2002 (R)
Clofentezine (156)	1986 (T,R), 1987 (R), 1989 (R), 1990 (R), 1992 (R), 2005 (T), 2007 (R), 2021 (R)
Clothianidin (238)	2010 (T,R), 2011 (R), 2014 (R), 2021 (R)
Coumaphos (018)	1968 (T,R), 1972 (R), 1975 (R), 1978 (R), 1980 (T,R), 1983 (R), 1987 (T), 1990 (T,R)
Crufomate (019)	1968 (T,R), 1972 (R)
Cyanophenfos (091)	1975 (T,R), 1978 (T: ADI extended, but no evaluation), 1980 (T), 1982 (R), 1983 (T)
Cyantraniliprole (263)	2013 (T,R), 2015 (R), 2018 (R)
Cyazofamid (281)	2015 (T, R), 2018 (R)
Cyclaniliprole (296)	2017 (T, R)
Cycloxydim (179)	1992 (T,R), 1993 (R), 2009 (T), 2012 (R)
Cyflumetofen (273)	2014 (T,R)
Cyfluthrin (157)	1986 (R), 1987 (T and corr. to 1986 report), 1989 (R), 1990 (R), 1992 (R), 2006 (T), 2007 (R)
Cyhalothrin (including lambda-cyhalothrin(146)	1984 (T,R), 1986 (R), 1988 (R), 2007 (T), 2008 (R), 2015 (R), 2018 (T)
Cyhexatin (067)	1970 (T,R), 1973 (T,R), 1974 (R), 1975 (R), 1977 (T), 1978 (T,R), 1980 (T), 1981 (T), 1982 (R), 1983 (R), 1985 (R), 1988 (T), 1989 (T), 1991 (T,R), 1992 (R), 1994 (T), 2005 (T,R)

Cypermethrin (118)	1979 (T,R), 1981 (T,R), 1982 (R), 1983 (R), 1984 (R), 1985 (R), 1986 (R), 1987 (corr. to 1986 evaluation), 1988 (R), 1990 (R), 2006 (T), 2008 (R), 2009 (R), 2011 (R), 2021 (R)
Cyproconazole (239)	2010 (T,R), 2013 (R)
Cyprodinil (207)	2003 (T,R), 2004 (corr. to 2003 report), 2013 (R), 2015 (R), 2017 (R), 2018 (R), 2019 (T, R), 2021 (R)
Cyromazine (169)	1990 (T,R), 1991 (corr. to 1990 R evaluation), 1992 (R), 2006 (T), 2007 (R), 2012 (R)
2,4-D (020)	1970 (T,R), 1971 (T,R), 1974 (T,R), 1975 (T,R), 1980 (R), 1985 (R), 1986 (R), 1987 (corr. to 1986 report, Annex I), 1996 (T), 1997 (E), 1998 (R), 2001 (R), 2017 (R)
Daminozide (104)	1977 (T,R), 1983 (T), 1989 (T,R), 1991 (T)
DDT (021)	1965 (T), 1966 (T,R), 1967 (T,R), 1968 (T,R), 1969 (T,R), 1978 (R), 1979 (T), 1980 (T), 1983 (T), 1984 (T), 1993 (R), 1994 (R), 1996 (R)
Deltamethrin (135)	1980 (T,R), 1981 (T,R), 1982 (T,R), 1984 (R), 1985 (R), 1986 (R), 1987 (R), 1988 (R), 1990 (R), 1992 (R), 2000 (T), 2002 (R), 2016 (R)
Demeton (092)	1965 (T), 1967 (R), 1975 (R), 1982 (T)
Demeton-S-methyl (073)	1973 (T,R), 1979 (R), 1982 (T), 1984 (T,R), 1989 (T,R), 1992 (R), 1998 (R)
Demeton-S-methylsulfon (164)	1973 (T,R), 1982 (T), 1984 (T,R), 1989 (T,R), 1992 (R)
Dialifos (098)	1976 (T,R), 1982 (T), 1985 (R)
Diazinon (022)	1965 (T), 1966 (T), 1967 (R), 1968 (T,R), 1970 (T,R), 1975 (R), 1979 (R), 1993 (T,R), 1994 (R), 1996 (R), 1999 (R), 2001 (T), 2006 (T,R), 2016 (T)
1,2-Dibromoethane (023)	1965 (T,R), 1966 (T,R), 1967 (R), 1968 (R), 1971 (R), 1979 (R), 1985 (R)
Dicamba (240)	2010 (T,R), 2011 (R), 2012 (R), 2013 (R), 2019 (T, R)
Dichlobenil (274)	2014 (T,R)
Dicloran (083)	2003 (R)
Dichlorfluanid (082)	1969 (T,R), 1974 (T,R), 1977 (T,R), 1979 (T,R), 1981 (R), 1982 (R), 1983 (T,R), 1985 (R)
1,2-Dichloroethane (024)	1965 (T,R), 1967 (R), 1971 (R), 1979 (R), 1985 (R)
Dichlorvos (025)	1965 (T,R), 1966 (T,R), 1967 (T,R), 1969 (R), 1970 (T,R), 1974 (R), 1977 (T), 1993 (T,R), 2011 (T), 2012 (R)
Dicloran (083)	1974 (T,R), 1977 (T,R), 1998 (T,R)
Dicofol (026)	1968 (T,R), 1970 (R), 1974 (R), 1992 (T,R), 1994 (R), 2011 (T), 2012 (R)

Dieldrin (001)	1965 (T), 1966 (T,R), 1967 (T,R), 1968 (R), 1969 (R), 1970 (T,R), 1974 (R), 1975 (R), 1977 (T), 1990 (R), 1992 (R)
Difenoconazole (224)	2007 (T,R), 2010 (R), 2013 (R), 2015 (R), 2017 (R), 2021 (R)
Diflubenzuron (130)	1981 (T,R), 1983 (R), 1984 (T,R), 1985 (T,R), 1988 (R), 2001 (T), 2002 (R), 2011 (R)
Dimethenamid-P (214)	2005 (T,R)
Dimethipin (151)	1985 (T,R), 1987 (T,R), 1988 (T,R), 1999 (T), 2001 (R), 2004 (T)
Dimethoate (027)	1965 (T), 1966 (T), 1967 (T,R), 1970 (R), 1973 (R in evaluation of formothion), 1977 (R), 1978 (R), 1983 (R) 1984 (T,R), 1986 (R), 1987 (T,R), 1988 (R), 1990 (R), 1991 (corr. to 1990 evaluation), 1994 (R), 1996 (T), 1998 (R), 2003 (T,R), 2004 (corr. to 2003 report), 2006 (R), 2008 (R)
Dimethomorph (225)	2007 (T,R), 2014 (R), 2016 (R)
Dimethrin	1965 (T)
Dinocap (087)	1969 (T,R), 1974 (T,R), 1989 (T,R), 1992 (R), 1998 (R), 1999 (R), 2000 (T), 2001 (R)
Dinotefuran (255)	2012 (T,R)
Dioxathion (028)	1968 (T,R), 1972 (R)
Diphenyl (029)	1966 (T,R), 1967 (T)
Diphenylamine (030)	1969 (T,R), 1976 (T,R), 1979 (R), 1982 (T), 1984 (T,R), 1998 (T), 2001 (R), 2003 (R), 2008 (R)
Diquat (031)	1970 (T,R), 1972 (T,R), 1976 (R), 1977 (T,R), 1978 (R), 1994 (R), 2013 (T,R), 2018 (R)
Disulfoton (074)	1973 (T,R), 1975 (T,R), 1979 (R), 1981 (R), 1984 (R), 1991 (T,R), 1992 (corr. to 1991 report, Annex I), 1994 (R), 1996 (T), 1998 (R), 2006 (R)
Dithianon (180)	1992 (T,R), 1995 (R), 1996 (corr. to 1995 report), 2010 (T), 2013 (T,R)
Dithiocarbamates (105)	1965 (T), 1967 (T,R), 1970 (T,R), 1983 (R propineb, thiram), 1984 (R propineb), 1985 (R), 1987 (T thiram), 1988 (R thiram), 1990 (R), 1991 (corr. to 1990 evaluation), 1992 (T thiram), 1993 (T,R), 1995 (R), 1996 (T,R ferbam, ziram; R thiram), 2004 (R), 2012 (R), 2014 (R)
4,6-Dinitro- <i>ortho</i> -cresol (DNOC)	1965 (T)
Dodine (084)	1974 (T,R), 1976 (T,R), 1977 (R), 2000 (T), 2003 (R), 2004 (corr. to 2003 report)
Edifenphos (099)	1976 (T,R), 1979 (T,R), 1981 (T,R)
Emamectin benzoate (247)	2011 (T,R), 2014 (R)

Endosulfan (032)	1965 (T), 1967 (T,R), 1968 (T,R), 1971 (R), 1974 (R), 1975 (R), 1982 (T), 1985 (T,R), 1989 (T,R), 1993 (R), 1998 (T), 2006 (R), 2010 (R)
Endrin (033)	1965 (T), 1970 (T,R), 1974 (R), 1975 (R), 1990 (R), 1992 (R)
Esfenvalerate (204)	2002 (T,R)
Ethephon (106)	1977 (T,R), 1978 (T,R), 1983 (R), 1985 (R), 1993 (T), 1994 (R), 1995 (T), 1997 (T), 2002 (T), 2015 (T, R)
Ethiofencarb (107)	1977 (T,R), 1978 (R), 1981 (R), 1982 (T,R), 1983 (R)
Ethion (034)	1968 (T,R), 1969 (R), 1970 (R), 1972 (T,R), 1975 (R), 1982 (T), 1983 (R), 1985 (T), 1986 (T), 1989 (T), 1990 (T), 1994 (R), 2021 (T, R)
Ethiprole (304)	2018 (T, R), 2021 (R)
Ethoprophos (149)	1983 (T), 1984 (R), 1987 (T), 1999 (T), 2004 (R)
Ethoxyquin (035)	1969 (T,R), 1998 (T), 1999 (R), 2005 (T), 2008 (R)
Ethylene dibromide	See 1,2-Dibromoethane
Ethylene dichloride	See 1,2-Dichloroethane
Ethylene oxide	1965 (T,R), 1968 (T,R), 1971 (R)
Ethylenethiourea (ETU) (108)	1974 (R), 1977 (T,R), 1986 (T,R), 1987 (R), 1988 (T,R), 1990 (R), 1993 (T,R)
Etofenprox (184)	1993 (T,R), 2011 (T,R)
Etoxazole (241)	2010 (T,R), 2011 (R)
Etrimfos (123)	1980 (T,R), 1982 (T,R), 1986 (T,R), 1987 (R), 1988 (R), 1989 (R), 1990 (R)
Famoxadone (208)	2003 (T,R)
Fenamidone (264)	2013 (T), 2014 (T,R)
Fenamiphos (085)	1974 (T,R), 1977 (R), 1978 (R), 1980 (R), 1985 (T), 1987 (T), 1997 (T), 1999 (R), 2002 (T), 2006 (R)
Fenarimol (192)	1995 (T,R,E), 1996 (R and corr. to 1995 report)
Fenazaquin (297)	2017 (T, R), 2019 (R)
Fenbuconazole (197)	1997 (T,R), 2009 (R), 2012 (T), 2013 (R), 2021 (R)
Fenbutatin oxide (109)	1977 (T,R), 1979 (R), 1992 (T), 1993 (R)
Fenchlorfos (036)	1968 (T,R), 1972 (R), 1983 (R)
Fenhexamid (215)	2005 (T,R), 2021 (R)
Fenitrothion (037)	1969 (T,R), 1974 (T,R), 1976 (R), 1977 (T,R), 1979 (R), 1982 (T), 1983 (R), 1984 (T,R), 1986 (T,R), 1987 (R and corr. to 1986 R evaluation), 1988 (T), 1989 (R), 2000 (T), 2003 (R), 2004 (R, corr. to 2003 report), 2007 (T,R)
Fenpicoxamid (305)	2018 (T,R), 2021 (T, R)
Fenpropathrin (185)	1993 (T,R), 2006 (R), 2012 (T), 2014 (R)

Fenpropimorph (188)	1994 (T), 1995 (R), 1999 (R), 2001 (T), 2004 (T), 2016 (T), 2017 (T, R)
Fenpyrazamine (298)	2017 (R, T)
Fenpyroximate (193)	1995 (T,R), 1996 (corr. to 1995 report), 1999 (R), 2004 (T), 2007 (T), 2010 (R), 2013 (R), 2017 (T, R), 2018 (R)
Fensulfothion (038)	1972 (T,R), 1982 (T), 1983 (R)
Fenthion (039)	1971 (T,R), 1975 (T,R), 1977 (R), 1978 (T,R), 1979 (T), 1980 (T), 1983 (R), 1989 (R), 1995 (T,R,E), 1996 (corr. to 1995 report), 1997 (T), 2000 (R)
Fentin compounds (040)	1965 (T), 1970 (T,R), 1972 (R), 1986 (R), 1991 (T,R), 1993 (R), 1994 (R)
Fenvalerate (119)	1979 (T,R), 1981 (T,R), 1982 (T), 1984 (T,R), 1985 (R), 1986 (T,R), 1987 (R and corr. to 1986 report), 1988 (R), 1990 (R), 1991 (corr. to 1990 R evaluation), 2012 (T,R)
Ferbam	See Dithiocarbamates, 1965 (T), 1967 (T,R), 1996 (T,R)
Fipronil (202)	1997 (T), 2000 (T), 2001 (R), 2016 (R)
Fipronil-desulfinyl	1997 (T)
Flonicamid (282)	2015 (T,R), 2016 (R), 2017 (R), 2019 (R)
Fluazifop-P-butyl	2016 (T,R)
Flubendiamide (242)	2010 (T,R)
Flucythrinate (152)	1985 (T,R), 1987 (R), 1988 (R), 1989 (R), 1990 (R), 1993 (R)
Fludioxonil (211)	2004 (T,R), 2006 (R), 2010 (R), 2012 (R), 2013 (R), 2018 (R)
Fluensulfone (265)	2013 (T), 2014 (T,R), 2016 (T,R), 2017 (R)
Flufenoxuron (275)	2014 (T,R)
Flumethrin (195)	1996 (T,R)
Fluazinam (306)	2018 (T,R)
Fluopicolide (235)	2009 (T,R), 2014 (R)
Fluopyram (243)	2010 (T,R), 2012 (R), 2014 (R), 2015 (R), 2017 (R), 2021 (R)
Flupyradifurone (285)	2015 (T), 2016 (R), 2017 (R), 2019 (R)
Flusilazole (165)	1989 (T,R), 1990 (R), 1991 (R), 1993 (R), 1995 (T), 2007 (T,R)
Flutolanil (205)	2002 (T,R), 2013 (R)
Flutriafol (248)	2011 (T,R), 2015 (R)
Fluxapyroxad (256)	2012 (T,R), 2015 (R), 2018 (T,R)

Folpet (041)	1969 (T,R), 1973 (T), 1974 (R), 1982 (T), 1984 (T,R), 1986 (T), 1987 (R), 1990 (T,R), 1991 (corr. to 1990 R evaluation), 1993 (T,R), 1994 (R), 1995 (T), 1997 (R), 1998 (R), 1999 (R), 2002 (T), 2004 (T), 2007 (T)
Formothion (042)	1969 (T,R), 1972 (R), 1973 (T,R), 1978 (R), 1998 (R)
Fosetyl Aluminium (302)	2017 (T, R), 2019 (R)
Glufosinate-ammonium (175)	1991 (T,R), 1992 (corr. to 1991 report, Annex I), 1994 (R), 1998 (R), 1999 (T,R), 2012 (T,R), 2014 (R)
Glyphosate (158)	1986 (T,R), 1987 (R and corr. to 1986 report), 1988 (R), 1994 (R), 1997 (T,R), 2004 (T), 2005 (R), 2011 (T,R), 2013 (R), 2016 (T), 2019 (R)
Guazatine (114)	1978 (T,R), 1980 (R), 1997 (T,R)
Haloxypop (194)	1995 (T,R), 1996 (R and corr. to 1995 report), 2001 (R), 2006 (T), 2009 (R)
Heptachlor (043)	1965 (T), 1966 (T,R), 1967 (R), 1968 (R), 1969 (R), 1970 (T,R), 1974 (R), 1975 (R), 1977 (R), 1987 (R), 1991 (T,R), 1992 (corr. to 1991 report, Annex I), 1993 (R), 1994 (R)
Hexachlorobenzene (044)	1969 (T,R), 1973 (T,R), 1974 (T,R), 1978 (T), 1985 (R)
Hexaconazole (170)	1990 (T,R), 1991 (R and corr. to 1990 R evaluation), 1993 (R)
Hexythiazox (176)	1991 (T,R), 1994 (R), 1998 (R), 2008 (T), 2009 (R)
Hydrogen cyanide (045)	1965 (T,R)
Hydrogen phosphide (046)	1965 (T,R), 1966 (T,R), 1967 (R), 1969 (R), 1971 (R)
Imazalil (110)	1977 (T,R), 1980 (T,R), 1984 (T,R), 1985 (T,R), 1986 (T), 1988 (R), 1989 (R), 1991 (T), 1994 (R), 2000 (T), 2001 (T), 2005 (T), 2018 (T,R), 2021 (R)
Imazamox (276)	2014 (T,R), 2017 (R)
Imazapic (266)	2013 (T,R), 2015 (R)
Imazapyr (267)	2013 (T,R), 2015 (R), 2017 (R)
Imazethapyr (289)	2016 (T,R)
Imidacloprid (206)	2001 (T), 2002 (R), 2006 (R), 2008 (R), 2012 (R), 2015 (R), 2017 (R)
Indoxacarb (216)	2005 (T,R), 2007 (R), 2009 (R), 2012 (R), 2013 (R)
Iprodione (111)	1977 (T,R), 1980 (R), 1992 (T), 1994 (R), 1995 (T), 2001 (R)
Isofenphos (131)	1981 (T,R), 1982 (T,R), 1984 (R), 1985 (R), 1986 (T,R), 1988 (R), 1992 (R)
Isofetamid (290)	2016 (T,R), 2018 (R)
Isoprothiolane (299)	2017 (T, R), 2021 (R)
Isopyrazam (249)	2011 (T,R), 2017 (R)

Isoxaflutole (268)	2013 (T,R), 2021 (R)
Kresoxim-methyl (199)	1998 (T,R), 2001 (R), 2018 (T,R)
Lead arsenate	1965 (T), 1968 (T,R)
Leptophos (088)	1974 (T,R), 1975 (T,R), 1978 (T,R)
Lindane (048)	1965 (T), 1966 (T,R), 1967 (R), 1968 (R), 1969 (R), 1970 (T,R, published as Annex VI to 1971 evaluations), 1973 (T,R), 1974 (R), 1975 (R), 1977 (T,R), 1978 (R), 1979 (R), 1989 (T,R), 1997 (T), 2002 (T), 2003 (R), 2004 (corr. to 2003 report), 2015 (R)
Lufenuron (286)	2015 (T, R), 2018 (R)
Malathion (049)	1965 (T), 1966 (T,R), 1967 (corr. to 1966 R evaluation), 1968 (R), 1969 (R), 1970 (R), 1973 (R), 1975 (R), 1977 (R), 1984 (R), 1997 (T), 1999 (R), 2000 (R), 2003 (T), 2004 (R), 2005 (R), 2008 (R), 2013 (R), 2016 (T)
Maleic hydrazide (102)	1976 (T,R), 1977 (T,R), 1980 (T), 1984 (T,R), 1996 (T), 1998 (R)
Mancozeb (050)	1967 (T,R), 1970 (T,R), 1974 (R), 1977 (R), 1980 (T,R), 1993 (T,R)
Mandestrobin (307)	2018 (T,R)
Mandipropamid (231)	2008 (T,R), 2013 (R), 2018 (R), 2021 (R)
Maneb	See Dithiocarbamates, 1965 (T), 1967 (T,R), 1987 (T), 1993 (T,R)
MCPA (257)	2012 (T,R)
Mecarbam (124)	1980 (T,R), 1983 (T,R), 1985 (T,R), 1986 (T,R), 1987 (R)
Meptyldinocap (244)	2010 (T, R)
Mesotrione (277)	2014 (T, R), 2019 (T, R)
Metaflumizone (236)	2009 (T, R), 2019 (T, R)
Metalaxyl (138)	1982 (T, R), 1984 (R), 1985 (R), 1986 (R), 1987 (R), 1989 (R), 1990 (R), 1992 (R), 1995 (R)
Metalaxyl –M (212)	2002 (T), 2004 (R)
Methacrifos (125)	1980 (T,R), 1982 (T), 1986 (T), 1988 (T), 1990 (T,R), 1992 (R)
Methamidophos (100)	1976 (T,R), 1979 (R), 1981 (R), 1982 (T,R), 1984 (R), 1985 (T), 1989 (R), 1990 (T,R), 1994 (R), 1996 (R), 1997 (R), 2002 (T), 2003 (R), 2004 (R, corr. to 2003 report)
Methidathion (051)	1972 (T,R), 1975 (T,R), 1979 (R), 1992 (T,R), 1994 (R), 1997 (T)
Methiocarb (132)	1981 (T,R), 1983 (T,R), 1984 (T), 1985 (T), 1986 (R), 1987 (T,R), 1988 (R), 1998 (T), 1999 (R), 2005 (R)

Methomyl (094)	1975 (R), 1976 (R), 1977 (R), 1978 (R), 1986 (T,R), 1987 (R), 1988 (R), 1989 (T,R), 1990 (R), 1991 (R), 2001 (T,R), 2004 (R), 2008 (R)
Methoprene (147)	1984 (T,R), 1986 (R), 1987 (T and corr. to 1986 report), 1988 (R), 1989 (R), 2001 (T), 2005 (R), 2016 (R), 2019 (R), 2021 (R)
Methoxychlor	1965 (T), 1977 (T)
Methoxyfenozide (209)	2003 (T,R), 2004 (corr. to 2003 report), 2006 (R), 2009 (R), 2012 (R), 2021 (R)
Methyl bromide (052)	See Bromomethane
Metrafenone (278)	2014 (T,R), 2016 (R)
Metiram (186)	1993 (T), 1995 (R)
Mevinphos (053)	1965 (T), 1972 (T,R), 1996 (T), 1997 (E,R), 2000 (R)
MGK (264)	1967 (T,R)
Monocrotophos (054)	1972 (T,R), 1975 (T,R), 1991 (T,R), 1993 (T), 1994 (R)
Myclobutanil (181)	1992 (T,R), 1997 (R), 1998 (R), (2001 (R)), 2014 (T,R)
Nabam	See Dithiocarbamates, 1965 (T), 1976 (T,R)
Natamycin (300)	2017 (T, R)
Nitrofen (140)	1983 (T,R)
Norflurazon (308)	2018 (T,R)
Novaluron (217)	2005 (T,R), 2010 (R)
Omethoate (055)	1971 (T,R), 1975 (T,R), 1978 (T,R), 1979 (T), 1981 (T,R), 1984 (R), 1985 (T), 1986 (R), 1987 (R), 1988 (R), 1990 (R), 1998 (R)
Organomercury compounds	1965 (T), 1966 (T,R), 1967 (T,R)
Oxamyl (126)	1980 (T,R), 1983 (R), 1984 (T), 1985 (T,R), 1986 (R), 2002 (T,R), 2017 (T, R)
Oxathiapiprolin (291)	2016 (T,R), 2018 (R)
Oxydemeton-methyl (166)	1965 (T, as demeton-S-methyl sulfoxide), 1967 (T), 1968 (R), 1973 (T,R), 1982 (T), 1984 (T,R), 1989 (T,R), 1992 (R), 1998 (R), 1999 (corr. to 1992 report), 2002 (T), 2004 (R)
Oxythioquinox	See Chinomethionat
Paclobutrazol (161)	1988 (T,R), 1989 (R)
Paraquat (057)	1970 (T,R), 1972 (T,R), 1976 (T,R), 1978 (R), 1981 (R), 1982 (T), 1985 (T), 1986 (T), 2003 (T), 2004 (R), 2009 (R)
Parathion (058)	1965 (T), 1967 (T,R), 1969 (R), 1970 (R), 1984 (R), 1991 (R), 1995 (T,R), 1997 (R), 2000 (R)
Parathion-methyl (059)	1965 (T), 1968 (T, R), 1972 (R), 1975 (T,R), 1978 (T,R), 1979 (T), 1980 (T), 1982 (T),

	1984 (T,R), 1991 (R), 1992 (R), 1994 (R), 1995 (T), 2000 (R), 2003 (R)
Penconazole (182)	1992 (T, R), 1995 (R), 2015 (T), 2016 (R)
Pendimethalin (292)	2016 (T, R), 2019 (R)
Penthiopyrad (253)	2011 (T), 2012 (R), 2013 (R)
Permethrin (120)	1979 (T, R), 1980 (R), 1981 (T,R), 1982 (R), 1983 (R), 1984 (R), 1985 (R), 1986 (T,R), 1987 (T), 1988 (R), 1989 (R), 1991 (R), 1992 (corr. to 1991 report), 1999 (T)
2-Phenylphenol (056)	1969 (T,R), 1975 (R), 1983 (T), 1985 (T,R), 1989 (T), 1990 (T,R), 1999 (T,R), 2002 (R)
Phenothrin (127)	1979 (R), 1980 (T,R), 1982 (T), 1984 (T), 1987 (R), 1988 (T,R)
Phenthoate (128)	1980 (T,R), 1981 (R), 1984 (T)
Phorate (112)	1977 (T,R), 1982 (T), 1983 (T), 1984 (R), 1985 (T), 1990 (R), 1991 (R), 1992 (R), 1993 (T), 1994 (T), 1996 (T), 2004 (T), 2005 (R), 2012 (R), 2014 (R)
Phosalone (060)	1972 (T,R), 1975 (R), 1976 (R), 1993 (T), 1994 (R), 1997 (T), 1999 (R), 2001 (T)
Phosmet (103)	1976 (R), 1977 (corr. to 1976 R evaluation), 1978 (T,R), 1979 (T,R), 1981 (R), 1984 (R), 1985 (R), 1986 (R), 1987 (R and corr. to 1986 R evaluation), 1988 (R), 1994 (T), 1997 (R), 1998 (T), 2002 (R), 2003 (R), 2007 (R)
Phosphine	See Hydrogen phosphide
Phosphamidon (061)	1965 (T), 1966 (T), 1968 (T,R), 1969 (R), 1972 (R), 1974 (R), 1982 (T), 1985 (T), 1986 (T)
Phosphonic acid (301)	2017 (T, R)
Phoxim (141)	1982 (T), 1983 (R), 1984 (T,R), 1986 (R), 1987 (R), 1988 (R)
Picoxystrobin (258)	2012 (T,R), 2013 (R), 2016 (R), 2017 (R)
Pinoxaden (293)	2016 (T,R)
Piperonyl butoxide (062)	1965 (T,R), 1966 (T,R), 1967 (R), 1969 (R), 1972 (T,R), 1992 (T,R), 1995 (T), 2001 (R), 2002 (R)
Pirimicarb (101)	1976 (T,R), 1978 (T,R), 1979 (R), 1981 (T,R), 1982 (T), 1985 (R), 2004 (T), 2006 (R)
Pirimiphos-methyl (086)	1974 (T,R), 1976 (T,R), 1977 (R), 1979 (R), 1983 (R), 1985 (R), 1992 (T), 1994 (R), 2003 (R), 2004 (R, corr. to 2003 report), 2006 (T)
Prochloraz (142)	1983 (T,R), 1985 (R), 1987 (R), 1988 (R), 1989 (R), 1990 (R), 1991 (corr. to 1990 report, Annex I and R evaluation), 1992 (R), 2001 (T), 2004 (R), 2009 (R)
Procymidone(136)	1981 (R), 1982 (T), 1989 (T,R), 1990 (R), 1991 (corr. to 1990 Annex I), 1993 (R), 1998 (R), 2007 (T)

Profenofos (171)	1990 (T,R), 1992 (R), 1994 (R), 1995 (R), 2007 (T), 2008 (R), 2011 (R), 2018 (R)
Propamocarb (148)	1984 (T,R), 1986 (T,R), 1987 (R), 2005 (T), 2006 (R), 2014 (R), 2018 (R)
Propargite (113)	1977 (T,R), 1978 (R), 1979 (R), 1980 (T,R), 1982 (T,R), 1999 (T), 2002 (R), 2006 (R)
Propham (183)	1965 (T), 1992 (T,R)
Propiconazole (160)	1987 (T, R), 1991 (R), 1994 (R), 2004 (T), 2006 (R), 2007 (R), 2013 (R), 2014 (R), 2015 (R), 2017 (R), 2018 (R)
Propineb	1977 (T, R), 1980 (T), 1983 (T), 1984 (R), 1985 (T, R), 1993 (T, R), 2004 (R)
Propoxur (075)	1973 (T, R), 1977 (R), 1981 (R), 1983 (R), 1989 (T), 1991 (R), 1996 (R)
Propylene oxide (250)	2011 (T, R), 2017 (T, R)
Propylenethiourea (PTU, 150)	1993 (T, R), 1994 (R), 1999 (T)
Prothioconazole (232)	2008 (T, R), 2009 (R), 2014 (R), 2017 (R), 2021 (R)
Pydiflumetofen (309)	2018 (T, R), 2021 (R)
Pymetrozine (279)	2014 (T, R)
Pyraclostrobin (210)	2003 (T), 2004 (R), 2006 (R), 2011 (R), 2012 (R), 2014 (R), 2018 (T, R)
Pyrazophos (153)	1985 (T, R), 1987 (R), 1992 (T,R), 1993 (R)
Pyrethrins (063)	1965 (T), 1966 (T, R), 1967 (R), 1968 (R), 1969 (R), 1970 (T), 1972 (T,R), 1974 (R), 1999 (T), 2000 (R), 2003 (T,R), 2005 (R)
Pyrimethanil (226)	2007 (T, R), 2013 (R)
Pyriofenone (310)	2018 (T, R)
Pyriproxyfen (200)	1999 (R, T), 2000 (R), 2001 (T), 2018 (R)
Quinclorac (287)	2015 (T, R), 2017 (R)
Quinoxifen (223)	2006 (T, R), 2021 (R)
Quintozene (064)	1969 (T, R), 1973 (T,R), 1974 (R), 1975 (T,R), 1976 (Annex I, corr. to 1975 R evaluation), 1977 (T,R), 1995 (T,R), 1998 (R)
Saflufenacil (251)	2011 (T, R), 2016 (R), 2017 (R)
Sedaxane (259)	2012 (T, R), 2014 (R)
Spices	2004 (R), 2005 (R), 2007 (R), 2010 (R), 2015 (R)
Spinetoram (233)	2008 (T, R), 2012 (R), 2017 (R), 2021 (R)
Spinosad (203)	2001 (T, R), 2004 (R), 2008 (R), 2011 (R)
Spirodiclofen (237)	2009 (T, R)
Spiromesifen (294)	2016 (T, R), 2021 (R)

Spirotetramat (234)	2008 (T, R), 2011 (R), 2012 (R), 2013 (R), 2015 (R), 2019 (R)
Sulfoxaflor (252)	2011 (T, R), 2013 (R), 2014 (R), 2016 (R), 2018 (R), 2021 (R)
Sulfuryl fluoride (218)	2005 (T, R)
2,4,5-T (121)	1970 (T,R), 1979 (T,R), 1981 (T)
Tebuconazole (189)	1994 (T, R), 1996 (corr. to Annex II of 1995 report), 1997 (R), 2008 (R), 2010 (T), 2011 (R), 2015 (R), 2017 (R), 2019 (R), 2021 (R)
Tebufenozide (196)	1996 (T, R), 1997 (R), 1999 (R), 2001 (T, R), 2003 (T)
Tecnazine (115)	1974 (T, R), 1978 (T, R), 1981 (R), 1983 (T), 1987 (R), 1989 (R), 1994 (T, R)
Teflubenzuron (190)	1994 (T), 1996 (R), 2016 (T, R)
Temephos	2006 (T)
Terbufos (167)	1989 (T, R), 1990 (T,R), 2003 (T), 2005 (R)
Thiabendazole (065)	1970 (T, R), 1971 (R), 1972 (R), 1975 (R), 1977 (T,R), 1979 (R), 1981 (R), 1997 (R), 2000 (R), 2006 (T,R), 2019 (T, R)
Thiacloprid (223)	2006 (T, R)
Thiamethoxam (245)	2010 (T, R), 2011 (R), 2012 (R), 2014 (R)
Thiodicarb (154)	1985 (T, R), 1986 (T), 1987 (R), 1988 (R), 2000 (T), 2001 (R)
Thiometon (076)	1969 (T, R), 1973 (T,R), 1976 (R), 1979 (T,R), 1988 (R)
Thiophanate-methyl (077)	1973 (T, R), 1975 (T, R), 1977 (T), 1978 (R), 1988 (R), 2002 (R), 1990 (R), 1994 (R), 1995 (T,E), 1998 (T,R), 2006 (T), 2017 (T)
Thiram (105)	See Dithiocarbamates, 1965 (T), 1967 (T, R), 1970 (T,R), 1974 (T), 1977 (T), 1983 (R), 1984 (R), 1985 (T,R), 1987 (T), 1988 (R), 1989 (R), 1992 (T), 1996 (R)
Tioxazafen (211)	2018 (T, R)
Tolclofos-methyl (191)	1994 (T, R), 1996 (corr. to Annex II of 1995 report)
Tolfenpyrad (269)	2013 (T), 2016 (R)
Tolyfluanid (162)	1988 (T, R), 1990 (R), 1991 (corr. to 1990 report), 2002 (T, R), 2003 (R)
Toxaphene	See Camphechlor
Triadimefon (133)	1979 (R), 1981 (T, R), 1983 (T,R), 1984 (R), 1985 (T,R), 1986 (R), 1987 (R and corr. to 1986 R evaluation), 1988 (R), 1989 (R), 1992 (R), 1995 (R), 2004 (T), 2007 (R)

Triadimenol (168)	1989 (T, R), 1992 (R), 1995 (R), 2004 (T), 2007 (R), 2014 (R)
Triazolylalanine	1989 (T, R)
Triazophos (143)	1982 (T), 1983 (R), 1984 (corr. to 1983 report, Annex I), 1986 (T, R), 1990 (R), 1991 (T and corr. to 1990 R evaluation), 1992 (R), 1993 (T, R), 2002 (T), 2007 (R), 2010 (R), 2013 (R)
Trichlorfon (066)	1971 (T,R), 1975 (T,R), 1978 (T,R), 1987 (R)
Trichloronat	1971 (T,R)
Trichloroethylene	1968 (R)
Tricyclohexyltin hydroxide	See Cyhexatin
Trifloxystrobin (213)	2004 (T, R), 2012 (R), 2015 (R), 2017 (R), 2021 (R)
Triflumezopyrim (303)	2017 (T, R)
Triflumizole (270)	2013 (T, R)
Triforine (116)	1977 (T), 1978 (T, R), 1997 (T), 2004 (R), 2014 (T,R)
Trinexapac-ethyl (271)	2013 (T,R), 2021(T, R)
Triphenyltin compounds	See Fentin compounds
Vamidothion (078)	1973 (T, R), 1982 (T), 1985 (T, R), 1987 (R), 1988 (T), 1990 (R), 1992 (R)
Vinclozolin (159)	1986 (T, R), 1987 (R and corr. to 1986 report and R evaluation), 1988 (T, R), 1989 (R), 1990 (R), 1992 (R), 1995 (T)
Zineb (105)	See Dithiocarbamates, 1965 (T), 1967 (T, R), 1993 (T)
Ziram (105)	See Dithiocarbamates, 1965 (T), 1967 (T, R), 1996 (T,R)
Zoxamide (227)	2007 (T, R), 2009 (R)

Annex 3

Annex 3: International estimated daily intakes (IEDIs) of pesticide residues

ACETAMIPRID (246)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.07 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
FC 0001	Group of Citrus fruit, raw (incl kumquat commodities)	RAC	0.25	32.25	8.06	11.67	2.92	16.70	4.18	76.01	19.00	33.90	8.48	92.97	23.24
JF 0001	Group of Citrus fruit, juice	PP	0.03	1.30	0.04	2.37	0.07	0.22	0.01	13.88	0.42	0.75	0.02	2.63	0.08
FP 0009	Group of Pome fruits, raw (incl apple cider, excl apple juice)	RAC	0.225	19.35	4.35	34.06	7.66	17.87	4.02	25.74	5.79	7.69	1.73	56.85	12.79
JF 0226	Apple juice, single strength (incl. concentrated)	PP	0.2	0.32	0.06	3.07	0.61	0.07	0.01	5.00	1.00	0.29	0.06	5.57	1.11
FS 0013	Subgroup of Cherries, raw	RAC	0.45	0.92	0.41	9.15	4.12	0.01	0.00	0.61	0.27	0.06	0.03	6.64	2.99
FS 0014	Subgroup of Plums, raw	RAC	0.04	2.40	0.10	8.60	0.34	0.06	0.00	2.52	0.10	0.58	0.02	4.16	0.17
DF 0014	Plums, dried (prunes)	PP	0.12	0.09	0.01	0.06	0.01	0.01	0.00	0.18	0.02	0.04	0.00	0.06	0.01
FS 2001	Subgroup of peaches, raw (incl dried apricots)	RAC	0.2	8.01	1.60	5.87	1.17	0.18	0.04	8.19	1.64	1.64	0.33	22.46	4.49
FB 2005	Subgroup of Caneberries, raw	RAC	0.64	0.42	0.27	1.05	0.67	0.01	0.01	0.02	0.01	0.02	0.01	1.24	0.79
FB 2006	Subgroup of Bush berries, raw (including processed)	RAC	0.64	0.53	0.34	1.31	0.84	0.40	0.26	1.66	1.06	0.01	0.01	0.99	0.63
FB 2007	Subgroup of Large shrub/tree berries, raw (including processed)	RAC	0.64	0.62	0.40	0.33	0.21	0.34	0.22	1.42	0.91	0.01	0.01	1.51	0.97
FB 0269	Grapes, raw (incl must, incl wine, excl dried, excl juice)	RAC	0.085	13.94	1.18	26.46	2.25	2.79	0.24	18.58	1.58	8.54	0.73	59.95	5.10
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	0.08	0.51	0.04	0.51	0.04	0.01	0.00	1.27	0.10	0.12	0.01	2.07	0.17
JF 0269	Grape juice (from wine grapes)	PP	0.13	0.14	0.02	0.29	0.04	0.05	0.01	0.30	0.04	0.24	0.03	0.05	0.01
FB 0265	Cranberry, raw	RAC	0.64	0.02	0.01	0.01	0.01	NC	-	0.03	0.02	0.01	0.01	0.01	0.01
FB 0275	Strawberry, raw	RAC	0.1	0.70	0.07	2.01	0.20	0.04	0.00	1.36	0.14	0.37	0.04	2.53	0.25
VA 2031	Subgroup of bulb onions	RAC	0.01	31.65	0.32	43.28	0.43	3.68	0.04	38.48	0.38	20.46	0.20	47.29	0.47
VA 2032	Subgroup of Green Onions	RAC	0.38	2.64	1.00	3.09	1.17	1.05	0.40	2.89	1.10	0.61	0.23	5.24	1.99
VB 0042	Subgroup of Flowerhead Brassica, raw	RAC	0.02	2.54	0.05	0.49	0.01	0.01	0.00	3.57	0.07	7.79	0.16	3.12	0.06
VB 2036	Subgroup of Head Brassicas, raw	RAC	0.02	3.81	0.08	38.89	0.78	0.77	0.02	6.23	0.12	4.27	0.09	14.28	0.29
VC 0424	Cucumber, raw	RAC	0.057	8.01	0.46	30.66	1.75	1.45	0.08	19.84	1.13	0.27	0.02	34.92	1.99
VC 0425	Gherkin, raw	RAC	0.05	1.73	0.09	6.64	0.33	0.31	0.02	4.29	0.21	0.29	0.01	7.56	0.38
VC 0431	Squash, Summer (Courgette, Marrow, Zucchini, Zucchini), raw	RAC	0.05	0.78	0.04	2.06	0.10	0.30	0.02	1.61	0.08	2.25	0.11	2.36	0.12

ACETAMIPRID (246)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.07 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
VC 2040	Subgroup of Melons, Pumpkins and Winter squashes	RAC	0.05	42.62	2.13	46.85	2.34	4.21	0.21	67.02	3.35	12.84	0.64	110.47	5.52
VO 0448	Tomato, raw (incl juice, incl canned, excl paste)	RAC	0.04	42.41	1.70	76.50	3.06	10.69	0.43	85.07	3.40	24.98	1.00	203.44	8.14
-	Tomato, paste (i.e. concentrated tomato sauce/puree)	PP	0.09	2.34	0.21	1.33	0.12	1.57	0.14	4.24	0.38	0.34	0.03	2.83	0.25
VO 0051	Subgroup of peppers, raw (incl dried sweet peppers, excl dried chili peppers), incl okra	RAC	0.04	10.45	0.42	13.74	0.55	13.81	0.55	14.53	0.58	15.25	0.61	27.93	1.12
-	Peppers, sweet, dried	PP	0.4	0.42	0.17	0.53	0.21	0.84	0.34	0.50	0.20	0.95	0.38	0.37	0.15
VO 2046	Subgroup of eggplants	RAC	0.04	5.58	0.22	4.31	0.17	0.89	0.04	9.31	0.37	13.64	0.55	20.12	0.80
VL 0460	Amaranth leaves, raw (i.e. Bledo)	RAC	0.64	1.09	0.70	1.94	1.24	1.20	0.77	2.91	1.86	NC	-	1.41	0.90
VL 0464	Chard, raw (i.e. Beet leaves; Silver beet)	RAC	0.64	0.40	0.26	0.70	0.45	0.44	0.28	1.06	0.68	4.66	2.98	0.51	0.33
VL 0465	Chervil, raw	RAC	0.64	0.19	0.12	0.34	0.22	0.21	0.13	0.52	0.33	NC	-	0.25	0.16
VL 2752	Chrysanthemum, edible leaved	RAC	0.64	0.02	0.01	0.04	0.03	0.02	0.01	0.06	0.04	NC	-	0.03	0.02
VL 0470	Corn salad (Lambs lettuce)	RAC	0.64	0.64	0.41	1.13	0.72	0.70	0.45	1.70	1.09	NC	-	0.82	0.52
VL 0474	Dandelion, raw (i.e. Laiteron, Pissenlit)	RAC	0.64	0.13	0.08	0.23	0.15	0.14	0.09	0.34	0.22	1.44	0.92	0.16	0.10
VL 0476	Endive, raw (i.e. scarole)	RAC	0.64	0.03	0.02	0.02	0.01	0.01	0.01	0.40	0.26	0.05	0.03	0.39	0.25
VL 0483	Lettuce, leaf, raw	RAC	0.64	0.53	0.34	0.36	0.23	0.16	0.10	6.21	3.97	1.90	1.22	6.05	3.87
VL 2765	Perilla leaves (Sesame leaves)	RAC	0.64	0.15	0.10	0.27	0.17	0.17	0.11	0.40	0.26	NC	-	0.19	0.12
VL 0492	Purslane, raw	RAC	0.64	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	NC	-	0.01	0.01
VL 0502	Spinach, raw	RAC	0.51	0.74	0.38	0.22	0.11	0.02	0.01	0.91	0.46	0.04	0.02	2.92	1.49
VL 0503	Spinach, Indian, raw (i.e. vine spinach)	RAC	0.51	0.01	0.01	0.02	0.01	0.02	0.01	0.04	0.02	0.16	0.08	0.02	0.01
VL 0485	Mustard greens, raw (i.e. Indian mustard, Amsoi, mustard cabbage)	RAC	2	0.03	0.06	0.31	0.62	0.01	0.02	0.05	0.10	0.47	0.94	0.11	0.22
VP 2060	Subgroup of beans with pods (all commodities within this group)	RAC	0.01	0.68	0.01	NC	-	NC	-	0.39	0.00	0.22	0.00	0.49	0.00
VP 0062	Beans without pods: (Phaseolus spp.) (succulent seeds), raw	RAC	0.03	1.56	0.05	0.60	0.02	0.49	0.01	1.18	0.04	0.90	0.03	7.79	0.23
VP 2063	Subgroup of succulent peas without pods	RAC	0.03	1.97	0.06	0.51	0.02	0.02	0.00	0.79	0.02	3.68	0.11	3.80	0.11
VS 0624	Celery	RAC	0.3	2.14	0.64	3.79	1.14	2.35	0.71	5.69	1.71	0.02	0.01	2.75	0.83
VS 0621	Asparagus, raw	RAC	0.26	0.01	0.00	0.01	0.00	0.01	0.00	0.03	0.01	0.07	0.02	0.21	0.05
GC 2090	Subgroup of Sweet Corns	RAC	0.01	0.14	0.00	0.94	0.01	5.70	0.06	2.61	0.03	1.94	0.02	0.22	0.00

Annex 3

ACETAMIPRID (246)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.07 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
TN 0085	Group of Tree nuts raw, excl coconut commodities	RAC	0.01	2.33	0.02	2.07	0.02	0.39	0.00	3.75	0.04	0.93	0.01	6.38	0.06
TN 0660	Almonds, nutmeat	RAC	0.01	1.38	0.01	0.08	0.00	0.01	0.00	1.00	0.01	0.06	0.00	0.81	0.01
TN 0662	Brazil nuts, nutmeat	RAC	0.01	0.01	0.00	0.01	0.00	0.03	0.00	0.02	0.00	0.02	0.00	0.01	0.00
TN 0295	Cashew nuts, nutmeat	RAC	0.01	0.01	0.00	0.02	0.00	0.24	0.00	0.47	0.00	0.32	0.00	0.05	0.00
TN 0664	Chestnut, raw	RAC	0.01	0.03	0.00	0.02	0.00	0.01	0.00	0.31	0.00	0.09	0.00	0.67	0.01
TN 0665	Coconut, nutmeat (incl. copra, incl desiccated, incl oil)	RAC	0.01	1.73	0.02	1.20	0.01	6.63	0.07	10.18	0.10	13.07	0.13	2.98	0.03
TN 0666	Hazelnuts, nutmeat	RAC	0.01	0.03	0.00	0.13	0.00	0.01	0.00	0.11	0.00	0.01	0.00	1.11	0.01
TN 0669	Macadamia nuts, nutmeat (i.e. Queensland nuts)	RAC	0.01	0.01	0.00	0.01	0.00	0.01	0.00	0.04	0.00	NC	-	0.03	0.00
TN 0672	Pecan, nutmeat	RAC	0.01	0.05	0.00	0.05	0.00	0.02	0.00	0.14	0.00	0.09	0.00	0.13	0.00
TN 0673	Pine nut, nutmeat (i.e. pignolia nuts)	RAC	0.01	0.18	0.00	0.18	0.00	0.08	0.00	0.49	0.00	0.25	0.00	0.43	0.00
TN 0675	Pistachio nut, nutmeat	RAC	0.33	0.41	0.14	0.07	0.02	0.01	0.00	0.85	0.28	0.02	0.01	1.08	0.36
TN 0678	Walnut, nutmeat	RAC	0.01	0.23	0.00	1.49	0.01	0.01	0.00	0.33	0.00	0.07	0.00	2.06	0.02
SO 0691	Cotton seed, raw (incl oil)	RAC	0.09	20.53	1.85	9.80	0.88	6.42	0.58	4.73	0.43	7.14	0.64	18.68	1.68
OR 0691	Cotton seed oil, edible	PP	0.004	3.22	0.01	1.54	0.01	1.01	0.00	0.74	0.00	1.12	0.00	2.93	0.01
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.022	24.96	0.55	57.95	1.27	16.70	0.37	38.38	0.84	26.46	0.58	29.00	0.64
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.017	6.24	0.11	14.49	0.25	4.18	0.07	9.60	0.16	6.62	0.11	7.25	0.12
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.017	3.29	0.06	6.14	0.10	0.82	0.01	1.57	0.03	2.23	0.04	1.07	0.02
MO 0105	Edible offal (mammalian), raw	RAC	0.11	4.79	0.53	9.68	1.06	2.97	0.33	5.49	0.60	3.84	0.42	5.03	0.55
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.019	289.65	5.50	485.88	9.23	26.92	0.51	239.03	4.54	199.91	3.80	180.53	3.43
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0	13.17	0.00	26.78	0.00	7.24	0.00	116.71	0.00	22.54	0.00	32.09	0.00
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0	1.46	0.00	2.98	0.00	0.80	0.00	12.97	0.00	2.50	0.00	3.57	0.00
PF 0111	Poultry fat, raw (incl rendered)	RAC	0	0.10	0.00	0.10	0.00	NC	-	0.10	0.00	0.10	0.00	0.10	0.00
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.01	0.12	0.00	0.12	0.00	0.11	0.00	5.37	0.05	0.24	0.00	0.10	0.00
PE 0112	Eggs, raw, (incl dried)	RAC	0	7.84	0.00	23.08	0.00	2.88	0.00	14.89	0.00	9.81	0.00	14.83	0.00
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-

ACETAMIPRID (246)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.07 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
	Total intake (ug/person)=				35.9		50.2		16.0		61.7		27.7		90.3
	Bodyweight per region (kg bw) =				60		60		60		60		60		60
	ADI (ug/person)=				4200		4200		4200		4200		4200		4200
	%ADI=				0.9%		1.2%		0.4%		1.5%		0.7%		2.1%
	Rounded %ADI=				1%		1%		0%		1%		1%		2%

ACETAMIPRID (246)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.07 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
FC 0001	Group of Citrus fruit, raw (incl kumquat commodities)	RAC	0.25	38.66	9.67	54.93	13.73	26.36	6.59	51.46	12.87	51.06	12.77	466.36	116.59
JF 0001	Group of Citrus fruit, juice	PP	0.03	36.84	1.11	3.75	0.11	0.30	0.01	21.62	0.65	21.82	0.65	46.67	1.40
FP 0009	Group of Pome fruits, raw (incl apple cider, excl apple juice)	RAC	0.225	51.09	11.50	65.40	14.72	42.71	9.61	45.29	10.19	62.51	14.06	7.74	1.74
JF 0226	Apple juice, single strength (incl. concentrated)	PP	0.2	14.88	2.98	11.98	2.40	0.15	0.03	9.98	2.00	30.32	6.06	3.47	0.69
FS 0013	Subgroup of Cherries, raw	RAC	0.45	1.40	0.63	4.21	1.89	0.04	0.02	2.93	1.32	1.50	0.68	NC	-
FS 0014	Subgroup of Plums, raw	RAC	0.04	3.75	0.15	3.33	0.13	5.94	0.24	2.64	0.11	2.50	0.10	0.06	0.00
DF 0014	Plums, dried (prunes)	PP	0.12	0.61	0.07	0.35	0.04	0.05	0.01	0.35	0.04	0.49	0.06	0.13	0.02
FS 2001	Subgroup of peaches, raw (incl dried apricots)	RAC	0.2	13.03	2.61	16.29	3.26	8.29	1.66	12.95	2.59	5.35	1.07	0.04	0.01
FB 2005	Subgroup of Caneberries, raw	RAC	0.64	0.56	0.36	1.43	0.92	0.14	0.09	1.23	0.79	1.14	0.73	0.01	0.01
FB 2006	Subgroup of Bush berries, raw (including processed)	RAC	0.64	1.31	0.84	5.50	3.52	0.01	0.01	2.57	1.64	0.82	0.52	2.15	1.38
FB 2007	Subgroup of Large shrub/tree berries, raw (including processed)	RAC	0.64	8.26	5.29	0.14	0.09	0.07	0.04	0.13	0.08	0.19	0.12	1.87	1.20
FB 0269	Grapes, raw (incl must, incl wine, excl dried, excl juice)	RAC	0.085	128.64	10.93	97.04	8.25	7.74	0.66	43.94	3.73	88.68	7.54	8.80	0.75
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	0.08	3.09	0.25	1.51	0.12	0.03	0.00	1.38	0.11	4.26	0.34	0.42	0.03
JF 0269	Grape juice (from wine grapes)	PP	0.13	0.56	0.07	1.96	0.25	0.02	0.00	2.24	0.29	2.27	0.30	0.34	0.04
FB 0265	Cranberry, raw	RAC	0.64	0.06	0.04	0.01	0.01	0.01	0.01	1.22	0.78	0.11	0.07	NC	-
FB 0275	Strawberry, raw	RAC	0.1	4.49	0.45	5.66	0.57	0.02	0.00	6.63	0.66	5.75	0.58	0.05	0.01

Annex 3

ACETAMIPRID (246)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.07 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
VA 2031	Subgroup of bulb onions	RAC	0.01	20.67	0.21	31.32	0.31	37.52	0.38	35.08	0.35	11.77	0.12	13.74	0.14
VA 2032	Subgroup of Green Onions	RAC	0.38	5.57	2.12	5.15	1.96	1.77	0.67	4.28	1.63	17.34	6.59	6.48	2.46
VB 0042	Subgroup of Flowerhead Brassica, raw	RAC	0.02	9.50	0.19	6.77	0.14	NC	-	3.21	0.06	9.36	0.19	0.75	0.02
VB 2036	Subgroup of Head Brassicas, raw	RAC	0.02	11.20	0.22	29.79	0.60	25.06	0.50	34.72	0.69	8.73	0.17	15.64	0.31
VC 0424	Cucumber, raw	RAC	0.057	6.72	0.38	11.03	0.63	32.10	1.83	15.10	0.86	4.05	0.23	9.57	0.55
VC 0425	Gherkin, raw	RAC	0.05	0.41	0.02	5.89	0.29	NC	-	0.06	0.00	0.37	0.02	2.07	0.10
VC 0431	Squash, Summer (Courgette, Marrow, Zucchetti, Zucchini), raw	RAC	0.05	NC	-	NC	-	5.48	0.27	NC	-	NC	-	1.03	0.05
VC 2040	Subgroup of Melons, Pumpkins and Winter squashes	RAC	0.05	20.68	1.03	25.00	1.25	85.72	4.29	34.31	1.72	11.54	0.58	23.32	1.17
VO 0448	Tomato, raw (incl juice, incl canned, excl paste)	RAC	0.04	44.88	1.80	55.49	2.22	35.44	1.42	75.65	3.03	27.00	1.08	9.61	0.38
-	Tomato, paste (i.e. concentrated tomato sauce/puree)	PP	0.09	4.96	0.45	3.20	0.29	0.15	0.01	1.61	0.14	6.88	0.62	0.52	0.05
VO 0051	Subgroup of peppers, raw (incl dried sweet peppers, excl dried chili peppers), incl okra	RAC	0.04	6.39	0.26	15.53	0.62	19.13	0.77	10.53	0.42	8.29	0.33	5.25	0.21
-	Peppers, sweet, dried	PP	0.4	0.11	0.04	0.21	0.08	0.36	0.14	0.21	0.08	0.25	0.10	0.15	0.06
VO 2046	Subgroup of eggplants	RAC	0.04	1.01	0.04	1.69	0.07	21.37	0.85	3.00	0.12	1.40	0.06	NC	-
VL 0460	Amaranth leaves, raw (i.e. Bledo)	RAC	0.64	NC	-	NC	-	47.45	30.37	NC	-	NC	-	2.07	1.32
VL 0464	Chard, raw (i.e. Beet leaves; Silver beet)	RAC	0.64	NC	-	NC	-	NC	-	NC	-	NC	-	0.75	0.48
VL 0465	Chervil, raw	RAC	0.64	NC	-	NC	-	8.39	5.37	NC	-	NC	-	0.37	0.24
VL 2752	Chrysanthemum, edible leaved	RAC	0.64	NC	-	NC	-	NC	-	0.32	0.20	NC	-	0.04	0.03
VL 0470	Corn salad (Lambs lettuce)	RAC	0.64	1.41	0.90	4.28	2.74	NC	-	0.03	0.02	5.11	3.27	1.20	0.77
VL 0474	Dandelion, raw (i.e. Laiteron, Pissenlit)	RAC	0.64	0.05	0.03	0.01	0.01	NC	-	NC	-	0.01	0.01	0.24	0.15
VL 0476	Endive, raw (i.e. scarole)	RAC	0.64	0.21	0.13	0.93	0.60	NC	-	0.30	0.19	2.14	1.37	0.14	0.09
VL 0483	Lettuce, leaf, raw	RAC	0.64	14.50	9.28	11.76	7.53	13.14	8.41	19.50	12.48	4.81	3.08	2.23	1.43
VL 2765	Perilla leaves (Sesame leaves)	RAC	0.64	NC	-	NC	-	NC	-	2.23	1.43	NC	-	0.29	0.19
VL 0492	Purslane, raw	RAC	0.64	0.01	0.01	NC	-	NC	-	NC	-	NC	-	0.01	0.01
VL 0502	Spinach, raw	RAC	0.51	2.20	1.12	1.76	0.90	13.38	6.82	2.94	1.50	5.53	2.82	0.02	0.01
VL 0503	Spinach, Indian, raw (i.e. vine spinach)	RAC	0.51	NC	-	NC	-	NC	-	NC	-	NC	-	0.03	0.02
VL 0485	Mustard greens, raw (i.e. Indian mustard, Amsoi,	RAC	2	NC	-	NC	-	NC	-	NC	-	NC	-	0.13	0.26

ACETAMIPRID (246)															
International Estimated Daily Intake (IEDI)				ADI = 0 - 0.07 mg/kg bw											
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
	mustard cabbage)														
VP 2060	Subgroup of beans with pods (all commodities within this group)	RAC	0.01	5.07	0.05	0.83	0.01	0.17	0.00	3.70	0.04	NC	-	NC	-
VP 0062	Beans without pods: (Phaseolus spp.) (succulent seeds), raw	RAC	0.03	2.21	0.07	5.25	0.16	4.17	0.13	1.61	0.05	16.95	0.51	0.17	0.01
VP 2063	Subgroup of succulent peas without pods	RAC	0.03	10.72	0.32	1.99	0.06	2.72	0.08	4.26	0.13	4.23	0.13	NC	-
VS 0624	Celery	RAC	0.3	7.68	2.30	2.85	0.86	NC	-	3.34	1.00	16.83	5.05	4.04	1.21
VS 0621	Asparagus, raw	RAC	0.26	0.84	0.22	2.08	0.54	7.11	1.85	1.01	0.26	1.69	0.44	0.04	0.01
GC 2090	Subgroup of Sweet Corns	RAC	0.01	11.43	0.11	3.71	0.04	0.74	0.01	13.63	0.14	3.07	0.03	1.50	0.02
TN 0085	Group of Tree nuts raw, excl coconut commodities	RAC	0.01	4.38	0.04	6.21	0.06	1.94	0.02	3.76	0.04	7.65	0.08	4.01	0.04
TN 0660	Almonds, nutmeat	RAC	0.01	0.81	0.01	2.21	0.02	0.03	0.00	1.02	0.01	1.47	0.01	NC	-
TN 0662	Brazil nuts, nutmeat	RAC	0.01	0.12	0.00	0.05	0.00	0.01	0.00	0.05	0.00	0.13	0.00	NC	-
TN 0295	Cashew nuts, nutmeat	RAC	0.01	0.59	0.01	0.23	0.00	0.18	0.00	0.52	0.01	1.75	0.02	2.78	0.03
TN 0664	Chestnut, raw	RAC	0.01	0.34	0.00	0.21	0.00	1.14	0.01	0.52	0.01	0.09	0.00	NC	-
TN 0665	Coconut, nutmeat (incl. copra, incl desiccated, incl oil)	RAC	0.01	4.13	0.04	2.73	0.03	13.15	0.13	5.85	0.06	6.92	0.07	22.24	0.22
TN 0666	Hazelnuts, nutmeat	RAC	0.01	0.45	0.00	1.12	0.01	0.02	0.00	0.34	0.00	1.63	0.02	NC	-
TN 0669	Macadamia nuts, nutmeat (i.e. Queensland nuts)	RAC	0.01	NC	-	0.40	0.00	NC	-	NC	-	NC	-	0.07	0.00
TN 0672	Pecan, nutmeat	RAC	0.01	0.38	0.00	NC	-	NC	-	0.27	0.00	NC	-	0.26	0.00
TN 0673	Pine nut, nutmeat (i.e. pignolia nuts)	RAC	0.01	0.99	0.01	0.66	0.01	0.22	0.00	0.27	0.00	1.89	0.02	0.89	0.01
TN 0675	Pistachio nut, nutmeat	RAC	0.33	0.35	0.12	0.48	0.16	0.07	0.02	0.39	0.13	0.23	0.08	0.02	0.01
TN 0678	Walnut, nutmeat	RAC	0.01	0.34	0.00	0.84	0.01	0.28	0.00	0.39	0.00	0.45	0.00	NC	-
SO 0691	Cotton seed, raw (incl oil)	RAC	0.09	10.71	0.96	4.23	0.38	7.19	0.65	7.54	0.68	5.66	0.51	2.38	0.21
OR 0691	Cotton seed oil, edible	PP	0.004	1.68	0.01	0.66	0.00	1.13	0.00	1.18	0.00	0.89	0.00	0.37	0.00
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.022	112.02	2.46	120.71	2.66	63.46	1.40	88.99	1.96	96.24	2.12	41.02	0.90
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.017	28.01	0.48	30.18	0.51	15.86	0.27	22.25	0.38	24.06	0.41	10.25	0.17
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.017	6.44	0.11	15.51	0.26	3.79	0.06	8.29	0.14	18.44	0.31	8.00	0.14
MO	Edible offal (mammalian), raw	RAC	0.11	15.17	1.67	5.19	0.57	6.30	0.69	6.78	0.75	3.32	0.37	3.17	0.35

Annex 3

ACETAMIPRID (246)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.07 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
0105															
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.019	388.92	7.39	335.88	6.38	49.15	0.93	331.25	6.29	468.56	8.90	245.45	4.66
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0	66.38	0.00	48.47	0.00	21.58	0.00	78.41	0.00	48.04	0.00	76.01	0.00
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0	7.38	0.00	5.39	0.00	2.40	0.00	8.71	0.00	5.34	0.00	8.45	0.00
PF 0111	Poultry fat, raw (incl rendered)	RAC	0	0.10	0.00	0.10	0.00	NC	-	0.10	0.00	0.71	0.00	NC	-
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.01	0.33	0.00	0.72	0.01	0.27	0.00	0.35	0.00	0.80	0.01	NC	-
PE 0112	Eggs, raw, (incl dried)	RAC	0	25.84	0.00	29.53	0.00	28.05	0.00	33.19	0.00	36.44	0.00	8.89	0.00
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-
Total intake (ug/person)=				81.6		83.0		87.3		74.9		85.4		142.3	
Bodyweight per region (kg bw) =				60		60		55		60		60		60	
ADI (ug/person)=				4200		4200		3850		4200		4200		4200	
%ADI=				1.9%		2.0%		2.3%		1.8%		2.0%		3.4%	
Rounded %ADI=				2%		2%		2%		2%		2%		3%	

ACETAMIPRID (246)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.07 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
FC 0001	Group of Citrus fruit, raw (incl kumquat commodities)	RAC	0.25	20.93	5.23	2.35	0.59	30.71	7.68	0.15	0.04	4.45	1.11
JF 0001	Group of Citrus fruit, juice	PP	0.03	0.11	0.00	0.29	0.01	13.55	0.41	0.14	0.00	0.33	0.01
FP 0009	Group of Pome fruits, raw (incl apple cider, excl apple juice)	RAC	0.225	68.85	15.49	10.93	2.46	70.82	15.93	189.78	42.70	19.56	4.40
JF 0226	Apple juice, single strength (incl. concentrated)	PP	0.2	0.03	0.01	0.10	0.02	7.19	1.44	0.03	0.01	NC	-
FS 0013	Subgroup of Cherries, raw	RAC	0.45	0.01	0.00	0.01	0.00	5.96	2.68	0.01	0.00	NC	-
FS 0014	Subgroup of Plums, raw	RAC	0.04	0.07	0.00	0.01	0.00	15.56	0.62	0.01	0.00	NC	-
DF 0014	Plums, dried (prunes)	PP	0.12	0.01	0.00	0.01	0.00	0.37	0.04	0.01	0.00	NC	-
FS 2001	Subgroup of peaches, raw (incl dried apricots)	RAC	0.2	0.02	0.00	0.01	0.00	10.76	2.15	0.01	0.00	NC	-

ACETAMIPRID (246)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.07 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
FB 2005	Subgroup of Caneberries, raw	RAC	0.64	0.01	0.01	7.30	4.67	2.29	1.47	0.01	0.01	NC	-
FB 2006	Subgroup of Bush berries, raw (including processed)	RAC	0.64	0.82	0.52	4.05	2.59	5.94	3.80	0.43	0.28	2.66	1.70
FB 2007	Subgroup of Large shrub/tree berries, raw (including processed)	RAC	0.64	0.71	0.45	7.32	4.68	NC	-	0.38	0.24	2.32	1.48
FB 0269	Grapes, raw (incl must, incl wine, excl dried, excl juice)	RAC	0.085	0.57	0.05	0.69	0.06	98.34	8.36	0.73	0.06	44.12	3.75
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	0.08	0.01	0.00	0.13	0.01	1.06	0.08	0.01	0.00	0.03	0.00
JF 0269	Grape juice (from wine grapes)	PP	0.13	0.01	0.00	0.01	0.00	0.41	0.05	0.01	0.00	NC	-
FB 0265	Cranberry, raw	RAC	0.64	NC	-	NC	-	0.03	0.02	NC	-	NC	-
FB 0275	Strawberry, raw	RAC	0.1	0.01	0.00	0.01	0.00	3.35	0.34	0.01	0.00	0.01	0.00
VA 2031	Subgroup of bulb onions	RAC	0.01	9.83	0.10	22.30	0.22	34.69	0.35	9.65	0.10	2.39	0.02
VA 2032	Subgroup of Green Onions	RAC	0.38	1.45	0.55	1.50	0.57	1.42	0.54	0.01	0.00	6.30	2.39
VB 0042	Subgroup of Flowerhead Brassica, raw	RAC	0.02	0.02	0.00	0.02	0.00	4.86	0.10	0.01	0.00	NC	-
VB 2036	Subgroup of Head Brassicas, raw	RAC	0.02	5.31	0.11	4.16	0.08	52.05	1.04	0.01	0.00	NC	-
VC 0424	Cucumber, raw	RAC	0.057	0.68	0.04	1.81	0.10	10.40	0.59	0.01	0.00	0.04	0.00
VC 0425	Gherkin, raw	RAC	0.05	0.15	0.01	0.39	0.02	3.15	0.16	0.01	0.00	0.01	0.00
VC 0431	Squash, Summer (Courgette, Marrow, Zucchini, Zucchini), raw	RAC	0.05	0.09	0.00	1.01	0.05	NC	-	1.91	0.10	NC	-
VC 2040	Subgroup of Melons, Pumpkins and Winter squashes	RAC	0.05	5.04	0.25	6.54	0.33	38.26	1.91	11.70	0.59	NC	-
VO 0448	Tomato, raw (incl juice, incl canned, excl paste)	RAC	0.04	13.17	0.53	4.92	0.20	62.69	2.51	1.04	0.04	0.11	0.00
-	Tomato, paste (i.e. concentrated tomato sauce/puree)	PP	0.09	0.58	0.05	0.22	0.02	2.21	0.20	0.24	0.02	3.10	0.28
VO 0051	Subgroup of peppers, raw (incl dried sweet peppers, excl dried chilipeppers), incl okra	RAC	0.04	15.20	0.61	14.23	0.57	25.14	1.01	0.91	0.04	NC	-
-	Peppers, sweet, dried	PP	0.4	0.58	0.23	1.27	0.51	1.21	0.48	0.12	0.05	NC	-
VO 2046	Subgroup of eggplants	RAC	0.04	1.31	0.05	8.26	0.33	3.95	0.16	0.01	0.00	NC	-
VL 0460	Amaranth leaves, raw (i.e. Bledo)	RAC	0.64	1.87	1.20	1.35	0.86	NC	-	1.27	0.81	2.53	1.62
VL 0464	Chard, raw (i.e. Beet leaves; Silver beet)	RAC	0.64	0.68	0.44	0.49	0.31	NC	-	0.46	0.29	0.92	0.59
VL 0465	Chervil, raw	RAC	0.64	0.33	0.21	0.24	0.15	NC	-	0.22	0.14	0.45	0.29
VL 2752	Chrysanthemum, edible leaved	RAC	0.64	0.04	0.03	0.03	0.02	NC	-	0.03	0.02	0.05	0.03

Annex 3

ACETAMIPRID (246)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.07 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
VL 0470	Corn salad (Lambs lettuce)	RAC	0.64	1.09	0.70	0.79	0.51	NC	-	0.74	0.47	1.47	0.94
VL 0474	Dandelion, raw (i.e. Laiteron, Pissenlit)	RAC	0.64	0.22	0.14	0.16	0.10	NC	-	0.15	0.10	0.29	0.19
VL 0476	Endive, raw (i.e. scarole)	RAC	0.64	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	NC	-
VL 0483	Lettuce, leaf, raw	RAC	0.64	0.29	0.19	0.03	0.02	6.71	4.29	0.01	0.01	NC	-
VL 2765	Perilla leaves (Sesame leaves)	RAC	0.64	0.26	0.17	0.19	0.12	NC	-	0.18	0.12	0.35	0.22
VL 0492	Purslane, raw	RAC	0.64	0.01	0.01	0.01	0.01	NC	-	0.01	0.01	0.01	0.01
VL 0502	Spinach, raw	RAC	0.51	0.17	0.09	0.01	0.01	0.81	0.41	0.01	0.01	NC	-
VL 0503	Spinach, Indian, raw (i.e. vine spinach)	RAC	0.51	0.02	0.01	0.02	0.01	NC	-	0.02	0.01	0.03	0.02
VL 0485	Mustard greens, raw (i.e. Indian mustard, Amsoi, mustard cabbage)	RAC	2	0.04	0.08	0.03	0.06	NC	-	0.01	0.02	NC	-
VP 2060	Subgroup of beans with pods (all commodities within this group)	RAC	0.01	NC	-	NC	-	NC	-	NC	-	NC	-
VP 0062	Beans without pods: (Phaseolus spp.) (succulent seeds), raw	RAC	0.03	0.30	0.01	3.13	0.09	4.11	0.12	0.01	0.00	NC	-
VP 2063	Subgroup of succulent peas without pods	RAC	0.03	0.21	0.01	0.02	0.00	5.51	0.17	0.02	0.00	NC	-
VS 0624	Celery	RAC	0.3	3.66	1.10	2.65	0.80	4.84	1.45	2.47	0.74	4.94	1.48
VS 0621	Asparagus, raw	RAC	0.26	0.01	0.00	0.01	0.00	0.17	0.04	0.01	0.00	NC	-
GC 2090	Subgroup of Sweet Corns	RAC	0.01	3.63	0.04	20.50	0.21	8.78	0.09	0.02	0.00	0.17	0.00
TN 0085	Group of Tree nuts raw, excl coconut commodities	RAC	0.01	1.61	0.02	1.16	0.01	3.31	0.03	0.02	0.00	0.07	0.00
TN 0660	Almonds, nutmeat	RAC	0.01	0.01	0.00	0.01	0.00	0.61	0.01	0.01	0.00	NC	-
TN 0662	Brazil nuts, nutmeat	RAC	0.01	0.01	0.00	0.01	0.00	0.02	0.00	0.01	0.00	NC	-
TN 0295	Cashew nuts, nutmeat	RAC	0.01	0.91	0.01	0.14	0.00	0.11	0.00	0.01	0.00	NC	-
TN 0664	Chestnut, raw	RAC	0.01	0.01	0.00	0.01	0.00	0.75	0.01	0.01	0.00	NC	-
TN 0665	Coconut, nutmeat (incl. copra, incl desiccated, incl oil)	RAC	0.01	2.77	0.03	134.37	1.34	2.81	0.03	0.70	0.01	317.67	3.18
TN 0666	Hazelnuts, nutmeat	RAC	0.01	0.01	0.00	0.01	0.00	0.21	0.00	0.01	0.00	NC	-
TN 0669	Macadamia nuts, nutmeat (i.e. Queensland nuts)	RAC	0.01	0.04	0.00	0.05	0.00	NC	-	0.01	0.00	0.01	0.00
TN 0672	Pecan, nutmeat	RAC	0.01	0.15	0.00	0.22	0.00	0.31	0.00	0.01	0.00	0.01	0.00
TN 0673	Pine nut, nutmeat (i.e. pignolia nuts)	RAC	0.01	0.51	0.01	0.74	0.01	0.36	0.00	0.01	0.00	0.05	0.00
TN 0675	Pistachio nut, nutmeat	RAC	0.33	0.01	0.00	0.01	0.00	0.15	0.05	0.01	0.00	NC	-

ACETAMIPRID (246)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.07 mg/kg bw						
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person								
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake	
TN 0678	Walnut, nutmeat	RAC	0.01	0.01	0.00	0.01	0.00	0.81	0.01	0.01	0.00	NC	-	
SO 0691	Cotton seed, raw (incl oil)	RAC	0.09	8.14	0.73	0.32	0.03	2.84	0.26	2.69	0.24	0.97	0.09	
OR 0691	Cotton seed oil, edible	PP	0.004	1.28	0.01	0.05	0.00	0.45	0.00	0.42	0.00	0.15	0.00	
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.022	23.34	0.51	40.71	0.90	97.15	2.14	18.06	0.40	57.71	1.27	
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.017	5.84	0.10	10.18	0.17	24.29	0.41	4.52	0.08	14.43	0.25	
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.017	1.05	0.02	1.14	0.02	18.69	0.32	0.94	0.02	3.12	0.05	
MO 0105	Edible offal (mammalian), raw	RAC	0.11	4.64	0.51	1.97	0.22	10.01	1.10	3.27	0.36	3.98	0.44	
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.019	108.75	2.07	70.31	1.34	436.11	8.29	61.55	1.17	79.09	1.50	
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0	3.53	0.00	10.83	0.00	51.36	0.00	4.53	0.00	50.00	0.00	
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0	0.39	0.00	1.20	0.00	5.71	0.00	0.50	0.00	5.56	0.00	
PF 0111	Poultry fat, raw (incl rendered)	RAC	0	NC	-	NC	-	0.32	0.00	NC	-	NC	-	
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.01	0.10	0.00	0.70	0.01	0.97	0.01	0.10	0.00	NC	-	
PE 0112	Eggs, raw, (incl dried)	RAC	0	3.84	0.00	4.41	0.00	27.25	0.00	1.13	0.00	7.39	0.00	
-	-	-		-	-	-	-	-	-	-	-	-	-	
Total intake (ug/person)=					32.7		25.4		73.4		49.3		27.3	
Bodyweight per region (kg bw) =					60		60		60		60		60	
ADI (ug/person)=					4200		4200		4200		4200		4200	
%ADI=					0.8%		0.6%		1.7%		1.2%		0.7%	
Rounded %ADI=					1%		1%		2%		1%		1%	

Annex 3

BIXAFEN (262)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.02 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
VD 0071	Beans, dry, raw (<i>Phaseolus</i> spp)	RAC	0.02	2.39	0.05	1.61	0.03	10.47	0.21	1.84	0.04	12.90	0.26	7.44	0.15
VD 0523	Broad bean, dry, raw (incl horse-bean, field bean) (<i>Vicia faba</i>)	RAC	0.02	1.27	0.03	0.01	0.00	0.12	0.00	2.49	0.05	0.23	0.00	5.54	0.11
VD 0527	Cowpea, dry, raw (<i>Vigna sinensis</i> , <i>Dolichos sinensis</i>)	RAC	0.02	0.05	0.00	NC	-	1.74	0.03	0.01	0.00	0.01	0.00	0.07	0.00
VD 0541	Soya bean, dry, raw (<i>Glycine soja</i>)	RAC	0.034	0.58	0.02	0.05	0.00	0.37	0.01	0.03	0.00	1.65	0.06	0.30	0.01
-	Soya paste (i.e. miso)	PP	0.034	NC	-	NC	-	NC	-	NC	-	NC	-	NC	-
-	Soya curd (i.e. tofu)	PP	0.034	NC	-	NC	-	NC	-	NC	-	NC	-	NC	-
OR 0541	Soya oil, refined	PP	0.0476	12.99	0.62	10.43	0.50	3.63	0.17	13.10	0.62	10.70	0.51	13.10	0.62
-	Soya sauce	PP	0.034	0.01	0.00	0.02	0.00	0.01	0.00	0.34	0.01	0.03	0.00	0.01	0.00
-	Soya flour	PP	0.00374	0.05	0.00	0.86	0.00	0.02	0.00	1.02	0.00	0.01	0.00	0.15	0.00
VD 2066	Subgroup of dry peas, raw	RAC	0.02	9.09	0.18	3.35	0.07	1.06	0.02	9.48	0.19	15.11	0.30	10.58	0.21
VD 2067	Subgroup of dry underground pulses, raw	RAC	0.02	NC	-	NC	-	0.20	0.00	NC	-	NC	-	NC	-
VR 0075	Group of Root and tuber vegetables, raw (incl processed)	RAC	0.028	87.83	2.46	374.04	10.47	668.92	18.73	121.64	3.41	94.20	2.64	247.11	6.92
GC 0650	Rye, raw (incl flour)	RAC	0.02	0.13	0.00	19.38	0.39	0.10	0.00	0.12	0.00	0.03	0.00	2.15	0.04
GC 0653	Triticale, raw (incl flour)	RAC	0.02	NC	-	NC	-	NC	-	0.01	0.00	0.39	0.01	NC	-
GC 0654	Wheat, raw (incl meslin)	RAC	0.095	0.01	0.00	1.12	0.11	NC	-	0.01	0.00	0.56	0.05	NC	-
-	Wheat, bulgur	PP	0.095	NC	-	NC	-	NC	-	0.03	0.00	NC	-	NC	-
CF 1210	Wheat, germ	PP	0.07885	NC	-	NC	-	0.01	0.00	0.01	0.00	0.14	0.01	0.01	0.00
CP 1212	Wheat, wholemeal bread	PP	0.095	0.01	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.03	0.00	0.01	0.00
CP 1211	Wheat, white bread	PP	0.095	0.25	0.02	0.63	0.06	0.12	0.01	0.43	0.04	1.39	0.13	0.22	0.02
-	Wheat, Fermented Beverages (Korean jakju and takju)	PP	0.095	NC	-	NC	-	NC	-	NC	-	NC	-	NC	-
CF 1211	Wheat, white flour (incl white flour products: starch, gluten, macaroni, pastry)	PP	0.03135	301.24	9.44	268.64	8.42	30.21	0.95	222.51	6.98	134.73	4.22	343.12	10.76
GC 0640	Barley, raw (incl malt extract, incl flour & grits, excl pot&pearled, excl beer, excl malt)	RAC	0.33	7.91	2.61	0.64	0.21	0.15	0.05	0.18	0.06	1.21	0.40	0.41	0.14
-	Barley, pot&pearled	PP	0.083	7.12	0.59	7.34	0.61	0.02	0.00	0.03	0.00	0.67	0.06	0.20	0.02
-	Barley beer	PP	0.036	4.87	0.18	93.78	3.38	24.28	0.87	12.76	0.46	39.28	1.41	18.15	0.65
-	Barley Malt	PP	0.317	0.09	0.03	1.04	0.33	0.18	0.06	0.33	0.10	0.04	0.01	0.10	0.03

BIXAFEN (262)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.02 mg/kg bw								
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day			Intake as ug/person/day									
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake	
GC 0647	Oats, raw (incl rolled)	RAC	0.08	0.05	0.00	7.05	0.56	0.10	0.01	1.71	0.14	0.96	0.08	0.04	0.00	
GC 0651	Sorghum, raw (incl flour, incl beer) (i.e. Chicken corn, Dari seed, Durra, Feterita)	RAC	0.196	4.34	0.85	0.01	0.00	16.25	3.19	15.82	3.10	10.97	2.15	2.92	0.57	
GC 0645	Maize, raw (incl glucose & dextrose & isoglucose, incl beer, excl flour, excl oil, excl germ, excl starch)	RAC	0.02	0.84	0.02	0.24	0.00	1.56	0.03	0.46	0.01	2.21	0.04	13.13	0.26	
CF 1255	Maize, flour (white flour and wholemeal flour)	PP	0.0178	22.72	0.40	35.61	0.63	87.27	1.55	34.92	0.62	46.71	0.83	49.12	0.87	
-	Maize, germ	PP	0.02	0.01	0.00	NC	-	0.01	0.00	0.01	0.00	0.22	0.00	NC	-	
-	Maize starch	PP	0.0126	0.08	0.00	NC	-	0.01	0.00	2.29	0.03	0.08	0.00	0.11	0.00	
OR 0645	Maize oil	PP	0.036	0.96	0.03	0.85	0.03	0.29	0.01	5.42	0.20	0.42	0.02	2.10	0.08	
GC 0447	Sweet corn on the cob, raw (incl frozen kernels, incl canned kernels)	RAC	0.02	0.14	0.00	0.94	0.02	5.70	0.11	2.61	0.05	1.94	0.04	0.22	0.00	
SO 0495	Rape seed, raw	RAC	0.02	0.02	0.00	NC	-	NC	-	0.01	0.00	0.75	0.02	0.01	0.00	
OR 0495	Rape seed oil, edible	PP	0.03	0.35	0.01	0.44	0.01	0.19	0.01	0.97	0.03	3.28	0.10	0.77	0.02	
SO 2091	Subgroup of Sunflower seeds, raw (incl processed)	RAC	0.045	7.43	0.33	36.06	1.62	1.15	0.05	8.77	0.39	5.74	0.26	13.63	0.61	
SO 0691	Cotton seed, raw (incl oil)	RAC	0.03	20.53	0.62	9.80	0.29	6.42	0.19	4.73	0.14	7.14	0.21	18.68	0.56	
SO 0697	Peanuts, nutmeat, raw (incl roasted, incl butter, excl oil)	RAC	0.02	0.46	0.01	1.21	0.02	6.64	0.13	2.71	0.05	1.26	0.03	1.84	0.04	
OR 0697	Peanut oil, edible	PP	0.044	0.36	0.02	0.01	0.00	2.57	0.11	0.07	0.00	2.29	0.10	0.36	0.02	
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.5	24.96	12.48	57.95	28.98	16.70	8.35	38.38	19.19	26.46	13.23	29.00	14.50	
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.21	6.24	1.31	14.49	3.04	4.18	0.88	9.60	2.02	6.62	1.39	7.25	1.52	
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.5	3.29	1.65	6.14	3.07	0.82	0.41	1.57	0.79	2.23	1.12	1.07	0.54	
MO 0105	Edible offal (mammalian), raw	RAC	1.7	4.79	8.14	9.68	16.46	2.97	5.05	5.49	9.33	3.84	6.53	5.03	8.55	
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.082	289.65	23.75	485.88	39.84	26.92	2.21	239.03	19.60	199.91	16.39	180.53	14.80	
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0	13.17	0.00	26.78	0.00	7.24	0.00	116.71	0.00	22.54	0.00	32.09	0.00	
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0.02	1.46	0.03	2.98	0.06	0.80	0.02	12.97	0.26	2.50	0.05	3.57	0.07	
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.02	0.10	0.00	0.10	0.00	NC	-	0.10	0.00	0.10	0.00	0.10	0.00	
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.02	0.12	0.00	0.12	0.00	0.11	0.00	5.37	0.11	0.24	0.00	0.10	0.00	

Annex 3

BIXAFEN (262)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.02 mg/kg bw								
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day			Intake as ug/person/day									
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake	
PE 0112	Eggs, raw, (incl dried)	RAC	0.02	7.84	0.16	23.08	0.46	2.88	0.06	14.89	0.30	9.81	0.20	14.83	0.30	
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-	
Total intake (ug/person)=				66.1			119.7		43.5		68.3		52.9		63.0	
Bodyweight per region (kg bw) =				60			60		60		60		60		60	
ADI (ug/person)=				1200			1200		1200		1200		1200		1200	
%ADI=				5.5%			10.0%		3.6%		5.7%		4.4%		5.3%	
Rounded %ADI=				6%			10%		4%		6%		4%		5%	

BIXAFEN (262)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.02 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day			Intake as ug/person/day								
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
VD 0071	Beans, dry, raw (Phaseolus spp)	RAC	0.02	1.51	0.03	1.50	0.03	1.90	0.04	5.11	0.10	1.36	0.03	23.43	0.47
VD 0523	Broad bean, dry, raw (incl horse-bean, field bean) (Vicia faba)	RAC	0.02	0.02	0.00	0.01	0.00	1.16	0.02	0.40	0.01	NC	-	0.06	0.00
VD 0527	Cowpea, dry, raw (Vigna sinensis, Dolichos sinensis)	RAC	0.02	NC	-	NC	-	0.16	0.00	0.01	0.00	NC	-	NC	-
VD 0541	Soya bean, dry, raw (Glycine soja)	RAC	0.034	0.02	0.00	0.33	0.01	6.64	0.23	3.94	0.13	NC	-	5.78	0.20
-	Soya paste (i.e. miso)	PP	0.034	NC	-	NC	-	NC	-	1.87	0.06	NC	-	NC	-
-	Soya curd (i.e. tofu)	PP	0.034	NC	-	NC	-	0.68	0.02	0.87	0.03	NC	-	NC	-
OR 0541	Soya oil, refined	PP	0.0476	19.06	0.91	21.06	1.00	5.94	0.28	33.78	1.61	40.05	1.91	13.39	0.64
-	Soya sauce	PP	0.034	0.45	0.02	0.29	0.01	2.93	0.10	4.35	0.15	0.09	0.00	0.70	0.02
-	Soya flour	PP	0.00374	0.22	0.00	0.27	0.00	0.29	0.00	0.17	0.00	NC	-	NC	-
VD 2066	Subgroup of dry peas, raw	RAC	0.02	5.01	0.10	3.76	0.08	1.82	0.04	3.44	0.07	3.49	0.07	5.15	0.10
VD 2067	Subgroup of dry underground pulses, raw	RAC	0.02	NC	-	NC	-	NC	-	NC	-	NC	-	NC	-
VR 0075	Group of Root and tuber vegetables, raw (incl processed)	RAC	0.028	290.31	8.13	300.35	8.41	214.25	6.00	242.72	6.80	348.67	9.76	137.52	3.85
GC 0650	Rye, raw (incl flour)	RAC	0.02	3.21	0.06	35.38	0.71	0.21	0.00	6.50	0.13	1.49	0.03	NC	-

BIXAFEN (262)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.02 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
GC 0653	Triticale, raw (incl flour)	RAC	0.02	0.01	0.00	0.17	0.00	0.29	0.01	0.01	0.00	NC	-	NC	-
GC 0654	Wheat, raw (incl meslin)	RAC	0.095	NC	-	NC	-	NC	-	0.01	0.00	NC	-	NC	-
-	Wheat, bulgur	PP	0.095	NC	-	NC	-	0.01	0.00	NC	-	NC	-	NC	-
CF 1210	Wheat, germ	PP	0.07885	0.97	0.08	0.10	0.01	0.03	0.00	0.01	0.00	NC	-	0.04	0.00
CP 1212	Wheat, wholemeal bread	PP	0.095	0.03	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.05	0.00	0.02	0.00
CP 1211	Wheat, white bread	PP	0.095	1.30	0.12	0.46	0.04	0.06	0.01	0.22	0.02	2.44	0.23	0.77	0.07
-	Wheat, Fermented Beverages (Korean jakju and takju)	PP	0.095	NC	-	NC	-	NC	-	4.36	0.41	NC	-	NC	-
CF 1211	Wheat, white flour (incl white flour products: starch, gluten, macaroni, pastry)	PP	0.03135	198.08	6.21	193.03	6.05	106.24	3.33	185.09	5.80	168.67	5.29	131.59	4.13
GC 0640	Barley, raw (incl malt extract, incl flour & grits, excl pot&pearled, excl beer, excl malt)	RAC	0.33	0.82	0.27	0.21	0.07	0.09	0.03	1.53	0.50	1.58	0.52	0.63	0.21
-	Barley, pot&pearled	PP	0.083	0.57	0.05	2.56	0.21	0.33	0.03	0.56	0.05	0.36	0.03	NC	-
-	Barley beer	PP	0.036	180.21	6.49	259.46	9.34	45.91	1.65	172.36	6.20	234.42	8.44	65.30	2.35
-	Barley Malt	PP	0.317	0.19	0.06	NC	-	0.04	0.01	0.08	0.03	NC	-	2.14	0.68
GC 0647	Oats, raw (incl rolled)	RAC	0.08	7.50	0.60	6.26	0.50	0.15	0.01	4.87	0.39	3.16	0.25	2.98	0.24
GC 0651	Sorghum, raw (incl flour, incl beer) (i.e. Chicken corn, Dari seed, Durra, Feterita)	RAC	0.196	NC	-	NC	-	1.44	0.28	1.15	0.23	NC	-	7.12	1.40
GC 0645	Maize, raw (incl glucose & dextrose & isoglucose, incl beer, excl flour, excl oil, excl germ, excl starch)	RAC	0.02	0.10	0.00	9.93	0.20	1.40	0.03	10.26	0.21	0.33	0.01	0.04	0.00
CF 1255	Maize, flour (white flour and wholemeal flour)	PP	0.0178	14.27	0.25	12.86	0.23	19.71	0.35	12.55	0.22	4.21	0.07	52.30	0.93
-	Maize, germ	PP	0.02	0.01	0.00	NC	-	NC	-	0.01	0.00	NC	-	0.01	0.00
-	Maize starch	PP	0.0126	NC	-	NC	-	0.19	0.00	7.13	0.09	NC	-	NC	-
OR 0645	Maize oil	PP	0.036	0.90	0.03	0.47	0.02	0.15	0.01	3.01	0.11	1.86	0.07	0.36	0.01
GC 0447	Sweet corn on the cob, raw (incl frozen kernels, incl canned kernels)	RAC	0.02	11.43	0.23	3.71	0.07	0.74	0.01	13.63	0.27	3.07	0.06	1.50	0.03
SO 0495	Rape seed, raw	RAC	0.02	NC	-	NC	-	0.01	0.00	NC	-	NC	-	NC	-
OR 0495	Rape seed oil, edible	PP	0.03	12.52	0.38	7.63	0.23	3.00	0.09	6.01	0.18	NC	-	NC	-
SO 2091	Subgroup of Sunflower seeds, raw (incl processed)	RAC	0.045	23.43	1.05	29.34	1.32	1.24	0.06	14.00	0.63	6.48	0.29	6.91	0.31
SO 0691	Cotton seed, raw (incl oil)	RAC	0.03	10.71	0.32	4.23	0.13	7.19	0.22	7.54	0.23	5.66	0.17	2.38	0.07

Annex 3

BIXAFEN (262)															
International Estimated Daily Intake (IEDI)															
ADI = 0 - 0.02 mg/kg bw															
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
SO 0697	Peanuts, nutmeat, raw (incl roasted, incl butter, excl oil)	RAC	0.02	3.26	0.07	2.22	0.04	5.38	0.11	4.85	0.10	1.54	0.03	1.82	0.04
OR 0697	Peanut oil, edible	PP	0.044	1.02	0.04	0.23	0.01	1.81	0.08	0.42	0.02	5.23	0.23	0.01	0.00
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.5	112.02	56.01	120.71	60.36	63.46	31.73	88.99	44.50	96.24	48.12	41.02	20.51
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.21	28.01	5.88	30.18	6.34	15.86	3.33	22.25	4.67	24.06	5.05	10.25	2.15
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.5	6.44	3.22	15.51	7.76	3.79	1.90	8.29	4.15	18.44	9.22	8.00	4.00
MO 0105	Edible offal (mammalian), raw	RAC	1.7	15.17	25.79	5.19	8.82	6.30	10.71	6.78	11.53	3.32	5.64	3.17	5.39
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.082	388.92	31.89	335.88	27.54	49.15	4.03	331.25	27.16	468.56	38.42	245.45	20.13
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0	66.38	0.00	48.47	0.00	21.58	0.00	78.41	0.00	48.04	0.00	76.01	0.00
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0.02	7.38	0.15	5.39	0.11	2.40	0.05	8.71	0.17	5.34	0.11	8.45	0.17
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.02	0.10	0.00	0.10	0.00	NC	-	0.10	0.00	0.71	0.01	NC	-
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.02	0.33	0.01	0.72	0.01	0.27	0.01	0.35	0.01	0.80	0.02	NC	-
PE 0112	Eggs, raw, (incl dried)	RAC	0.02	25.84	0.52	29.53	0.59	28.05	0.56	33.19	0.66	36.44	0.73	8.89	0.18
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-
Total intake (ug/person)=				149.0		140.3		65.4		117.6		134.8		68.3	
Bodyweight per region (kg bw) =				60		60		55		60		60		60	
ADI (ug/person)=				1200		1200		1100		1200		1200		1200	
%ADI=				12.4%		11.7%		5.9%		9.8%		11.2%		5.7%	
Rounded %ADI=				10%		10%		6%		10%		10%		6%	

BIXAFEN (262)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.02 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
VD 0071	Beans, dry, raw (Phaseolus spp)	RAC	0.02	7.11	0.14	2.33	0.05	3.76	0.08	44.70	0.89	3.27	0.07
VD 0523	Broad bean, dry, raw (incl horse-bean, field bean) (Vicia faba)	RAC	0.02	3.70	0.07	0.03	0.00	0.17	0.00	0.01	0.00	NC	-
VD 0527	Cowpea, dry, raw (Vigna sinensis, Dolichos sinensis)	RAC	0.02	12.77	0.26	0.99	0.02	0.01	0.00	4.33	0.09	NC	-
VD 0541	Soya bean, dry, raw (Glycine soja)	RAC	0.034	2.76	0.09	0.07	0.00	0.33	0.01	3.16	0.11	NC	-
-	Soya paste (i.e. miso)	PP	0.034	NC	-	NC	-	NC	-	NC	-	NC	-
-	Soya curd (i.e. tofu)	PP	0.034	NC	-	NC	-	NC	-	NC	-	NC	-
OR 0541	Soya oil, refined	PP	0.0476	2.32	0.11	2.54	0.12	18.70	0.89	2.51	0.12	6.29	0.30
-	Soya sauce	PP	0.034	0.01	0.00	0.13	0.00	0.17	0.01	0.01	0.00	0.56	0.02
-	Soya flour	PP	0.00374	0.11	0.00	0.08	0.00	0.07	0.00	0.01	0.00	0.03	0.00
VD 2066	Subgroup of dry peas, raw	RAC	0.02	4.43	0.09	11.36	0.23	4.22	0.08	9.36	0.19	1.21	0.02
VD 2067	Subgroup of dry underground pulses, raw	RAC	0.02	0.20	0.00	NC	-	NC	-	NC	-	NC	-
VR 0075	Group of Root and tuber vegetables, raw (incl processed)	RAC	0.028	282.25	7.90	232.11	6.50	281.91	7.89	620.21	17.37	459.96	12.88
GC 0650	Rye, raw (incl flour)	RAC	0.02	0.03	0.00	0.01	0.00	13.95	0.28	0.01	0.00	0.88	0.02
GC 0653	Triticale, raw (incl flour)	RAC	0.02	0.01	0.00	NC	-	NC	-	NC	-	NC	-
GC 0654	Wheat, raw (incl meslin)	RAC	0.095	NC	-	NC	-	NC	-	NC	-	0.97	0.09
-	Wheat, bulgur	PP	0.095	0.01	0.00	NC	-	NC	-	NC	-	NC	-
CF 1210	Wheat, germ	PP	0.07885	0.04	0.00	0.01	0.00	0.01	0.00	0.01	0.00	NC	-
CP 1212	Wheat, wholemeal bread	PP	0.095	0.01	0.00	0.01	0.00	0.03	0.00	0.01	0.00	0.01	0.00
CP 1211	Wheat, white bread	PP	0.095	0.43	0.04	0.41	0.04	1.56	0.15	0.11	0.01	0.07	0.01
-	Wheat, Fermented Beverages (Korean jakju and takju)	PP	0.095	NC	-	NC	-	NC	-	NC	-	NC	-
CF 1211	Wheat, white flour (incl white flour products: starch, gluten, macaroni, pastry)	PP	0.03135	44.78	1.40	86.96	2.73	214.05	6.71	20.31	0.64	103.60	3.25
GC 0640	Barley, raw (incl malt extract, incl flour & grits, excl pot&pearled, excl beer, excl malt)	RAC	0.33	0.09	0.03	0.01	0.00	0.80	0.26	0.01	0.00	0.11	0.04
-	Barley, pot&pearled	PP	0.083	5.46	0.45	0.01	0.00	1.44	0.12	0.01	0.00	NC	-
-	Barley beer	PP	0.036	16.25	0.59	11.36	0.41	225.21	8.11	19.49	0.70	52.17	1.88
-	Barley Malt	PP	0.317	0.01	0.00	0.11	0.03	0.67	0.21	0.01	0.00	4.61	1.46

Annex 3

BIXAFEN (262)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.02 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
GC 0647	Oats, raw (incl rolled)	RAC	0.08	0.37	0.03	0.07	0.01	2.79	0.22	0.10	0.01	NC	-
GC 0651	Sorghum, raw (incl flour, incl beer) (i.e. Chicken corn, Dari seed, Durra, Feterita)	RAC	0.196	89.16	17.48	2.02	0.40	NC	-	35.38	6.93	NC	-
GC 0645	Maize, raw (incl glucose & dextrose & isoglucose, incl beer, excl flour, excl oil, excl germ, excl starch)	RAC	0.02	0.54	0.01	0.51	0.01	3.26	0.07	7.96	0.16	NC	-
CF 1255	Maize, flour (white flour and wholemeal flour)	PP	0.0178	94.34	1.68	8.09	0.14	28.03	0.50	55.94	1.00	28.07	0.50
-	Maize, germ	PP	0.02	0.01	0.00	NC	-	NC	-	NC	-	NC	-
-	Maize starch	PP	0.0126	0.02	0.00	0.01	0.00	NC	-	NC	-	NC	-
OR 0645	Maize oil	PP	0.036	0.33	0.01	0.07	0.00	0.81	0.03	0.01	0.00	NC	-
GC 0447	Sweet corn on the cob, raw (incl frozen kernels, incl canned kernels)	RAC	0.02	3.63	0.07	20.50	0.41	8.78	0.18	0.02	0.00	0.17	0.00
SO 0495	Rape seed, raw	RAC	0.02	NC	-	0.01	0.00	NC	-	NC	-	NC	-
OR 0495	Rape seed oil, edible	PP	0.03	0.07	0.00	0.03	0.00	4.62	0.14	0.03	0.00	NC	-
SO 2091	Subgroup of Sunflower seeds, raw (incl processed)	RAC	0.045	0.99	0.04	0.22	0.01	32.01	1.44	12.12	0.55	0.48	0.02
SO 0691	Cotton seed, raw (incl oil)	RAC	0.03	8.14	0.24	0.32	0.01	2.84	0.09	2.69	0.08	0.97	0.03
SO 0697	Peanuts, nutmeat, raw (incl roasted, incl butter, excl oil)	RAC	0.02	7.14	0.14	0.45	0.01	1.87	0.04	6.22	0.12	0.53	0.01
OR 0697	Peanut oil, edible	PP	0.044	5.02	0.22	0.05	0.00	0.17	0.01	0.29	0.01	NC	-
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.5	23.34	11.67	40.71	20.36	97.15	48.58	18.06	9.03	57.71	28.86
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.21	5.84	1.23	10.18	2.14	24.29	5.10	4.52	0.95	14.43	3.03
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.5	1.05	0.53	1.14	0.57	18.69	9.35	0.94	0.47	3.12	1.56
MO 0105	Edible offal (mammalian), raw	RAC	1.7	4.64	7.89	1.97	3.35	10.01	17.02	3.27	5.56	3.98	6.77
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.082	108.75	8.92	70.31	5.77	436.11	35.76	61.55	5.05	79.09	6.49
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0	3.53	0.00	10.83	0.00	51.36	0.00	4.53	0.00	50.00	0.00
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0.02	0.39	0.01	1.20	0.02	5.71	0.11	0.50	0.01	5.56	0.11
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.02	NC	-	NC	-	0.32	0.01	NC	-	NC	-
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.02	0.10	0.00	0.70	0.01	0.97	0.02	0.10	0.00	NC	-
PE 0112	Eggs, raw, (incl dried)	RAC	0.02	3.84	0.08	4.41	0.09	27.25	0.55	1.13	0.02	7.39	0.15
-	-	-		-	-	-	-	-	-	-	-	-	-

BIXAFEN (262)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.02 mg/kg bw					
Codex Code	Commodity description	Expr as	STM R mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
	Total intake (ug/person)=				61.4		43.4		144.0		50.1		67.5
	Bodyweight per region (kg bw) =				60		60		60		60		60
	ADI (ug/person)=				1200		1200		1200		1200		1200
	%ADI=				5.1%		3.6%		12.0%		4.2%		5.6%
	Rounded %ADI=				5%		4%		10%		4%		6%

Annex 3

CLOFENTIZINE (156)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.02 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
FC 0002	Subgroup of Lemons and limes, raw (incl lemon juice) (incl kumquat commodities)	RAC	0.02	4.82	0.10	2.45	0.05	3.93	0.08	25.44	0.51	8.74	0.17	16.23	0.32
FC 0003	Subgroup of Mandarins, raw (incl mandarin juice)	RAC	0.02	6.18	0.12	3.66	0.07	0.25	0.01	6.82	0.14	3.49	0.07	19.38	0.39
FC 0004	Subgroup of Oranges, sweet, sour, raw	RAC	0.02	20.66	0.41	5.23	0.10	11.90	0.24	37.90	0.76	21.16	0.42	56.46	1.13
JF 0004	Subgroup of Oranges, juice (single strength, incl. concentrated)	PP	0.014	1.27	0.02	2.20	0.03	0.09	0.00	11.81	0.17	0.46	0.01	1.69	0.02
FC 0005	Subgroup of Pummelo and grapefruits, raw (incl grapefruit juice)	RAC	0.02	0.66	0.01	0.69	0.01	0.96	0.02	10.20	0.20	1.25	0.03	2.97	0.06
FP 0009	Group of Pome fruits, raw (incl apple cider, excl apple juice)	RAC	0.05	19.35	0.97	34.06	1.70	17.87	0.89	25.74	1.29	7.69	0.38	56.85	2.84
JF 0226	Apple juice, single strength (incl. concentrated)	PP	0.0055	0.32	0.00	3.07	0.02	0.07	0.00	5.00	0.03	0.29	0.00	5.57	0.03
FS 0012	Group of Stone fruits, raw (incl dried plums, incl dried apricots)	RAC	0.11	11.60	1.28	23.79	2.62	0.25	0.03	11.84	1.30	2.41	0.27	33.44	3.68
FB 0021	Currants, Black, Red, White, raw	RAC	0.04	0.02	0.00	0.74	0.03	0.01	0.00	0.03	0.00	0.01	0.00	0.01	0.00
FB 0269	Grapes, raw (incl must, excl dried, excl juice, excl wine)	RAC	0.25	13.02	3.26	9.25	2.31	0.03	0.01	16.91	4.23	3.70	0.93	54.44	13.61
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	0.28	0.51	0.14	0.51	0.14	0.01	0.00	1.27	0.36	0.12	0.03	2.07	0.58
JF 0269	Grape juice (from wine grapes)	PP	0	0.14	0.00	0.29	0.00	0.05	0.00	0.30	0.00	0.24	0.00	0.05	0.00
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.011	0.67	0.01	12.53	0.14	2.01	0.02	1.21	0.01	3.53	0.04	4.01	0.04
FB 1236	Wine grapes, raw (incl must, juice, wine)	RAC	0.011	14.11	0.16	26.83	0.30	2.85	0.03	18.95	0.21	8.84	0.10	60.01	0.66
FB 0275	Strawberry, raw	RAC	0.72	0.70	0.50	2.01	1.45	0.04	0.03	1.36	0.98	0.37	0.27	2.53	1.82
VC 0424	Cucumber, raw	RAC	0.125	8.01	1.00	30.66	3.83	1.45	0.18	19.84	2.48	0.27	0.03	34.92	4.37
VC 0046	Melons, except watermelon, raw (Cantaloupe)	RAC	0	8.90	0.00	8.64	0.00	0.80	0.00	17.90	0.00	2.80	0.00	29.17	0.00
VO 0448	Tomato, raw (incl juice, incl paste, incl canned)	RAC	0.09	51.75	4.66	81.80	7.36	16.99	1.53	102.02	9.18	26.32	2.37	214.77	19.33
TN 0085	Group of Tree nuts, raw (incl processed)	RAC	0.05	4.06	0.20	3.27	0.16	7.01	0.35	13.93	0.70	14.01	0.70	9.36	0.47
DH 1100	Hops, dry	RAC	2.2	0.01	0.02	0.04	0.09	0.01	0.02	0.01	0.02	NC	-	0.01	0.02
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat)	RAC	0	31.20	0.00	72.44	0.00	20.88	0.00	47.98	0.00	33.08	0.00	36.25	0.00
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0	3.29	0.00	6.14	0.00	0.82	0.00	1.57	0.00	2.23	0.00	1.07	0.00
MO 0105	Edible offal (mammalian), raw	RAC	0.05	4.79	0.24	9.68	0.48	2.97	0.15	5.49	0.27	3.84	0.19	5.03	0.25
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0	289.65	0.00	485.88	0.00	26.92	0.00	239.03	0.00	199.91	0.00	180.53	0.00

CLOFENTIZINE (156)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.02 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
PM 0110	Poultry meat, raw (incl prepared)	RAC	0	14.63	0.00	29.76	0.00	8.04	0.00	129.68	0.00	25.04	0.00	35.66	0.00
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0	0.12	0.00	0.12	0.00	0.11	0.00	5.37	0.00	0.24	0.00	0.10	0.00
PE 0112	Eggs, raw, (incl dried)	RAC	0	7.84	0.00	23.08	0.00	2.88	0.00	14.89	0.00	9.81	0.00	14.83	0.00
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-
Total intake (ug/person)=				13.1		20.9		3.6		22.8		6.0		49.6	
Bodyweight per region (kg bw) =				60		60		60		60		60		60	
ADI (ug/person)=				1200		1200		1200		1200		1200		1200	
%ADI=				1.1%		1.7%		0.3%		1.9%		0.5%		4.1%	
Rounded %ADI=				1%		2%		0%		2%		1%		4%	

CLOFENTIZINE (156)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.02 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
FC 0002	Subgroup of Lemons and limes, raw (incl lemon juice) (incl kumquat commodities)	RAC	0.02	10.12	0.20	15.69	0.31	2.88	0.06	12.30	0.25	22.32	0.45	6.59	0.13
FC 0003	Subgroup of Mandarins, raw (incl mandarin juice)	RAC	0.02	12.42	0.25	14.99	0.30	16.08	0.32	10.78	0.22	9.94	0.20	NC	-
FC 0004	Subgroup of Oranges, sweet, sour, raw	RAC	0.02	15.68	0.31	24.00	0.48	6.80	0.14	29.09	0.58	15.39	0.31	160.47	3.21
JF 0004	Subgroup of Oranges, juice (single strength, incl. concentrated)	PP	0.014	33.31	0.47	1.78	0.02	0.28	0.00	18.97	0.27	14.01	0.20	13.36	0.19
FC 0005	Subgroup of Pummelo and grapefruits, raw (incl grapefruit juice)	RAC	0.02	8.21	0.16	4.60	0.09	0.64	0.01	5.85	0.12	19.98	0.40	368.86	7.38
FP 0009	Group of Pome fruits, raw (incl apple cider, excl apple juice)	RAC	0.05	51.09	2.55	65.40	3.27	42.71	2.14	45.29	2.26	62.51	3.13	7.74	0.39
JF 0226	Apple juice, single strength (incl. concentrated)	PP	0.0055	14.88	0.08	11.98	0.07	0.15	0.00	9.98	0.05	30.32	0.17	3.47	0.02
FS 0012	Group of Stone fruits, raw (incl dried plums, incl dried apricots)	RAC	0.11	19.98	2.20	24.87	2.74	14.41	1.59	19.54	2.15	10.78	1.19	0.50	0.06
FB 0021	Currants, Black, Red, White, raw	RAC	0.04	0.48	0.02	4.23	0.17	NC	-	1.51	0.06	0.49	0.02	NC	-
FB 0269	Grapes, raw (incl must, excl dried, excl juice, excl wine)	RAC	0.25	6.48	1.62	11.31	2.83	5.21	1.30	9.50	2.38	4.66	1.17	0.78	0.20
DF 0269	Grapes, dried (= currants, raisins and sultanas)	PP	0.28	3.09	0.87	1.51	0.42	0.03	0.01	1.38	0.39	4.26	1.19	0.42	0.12

Annex 3

CLOFENTIZINE (156)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.02 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
	(from table-grapes)														
JF 0269	Grape juice (from wine grapes)	PP	0	0.56	0.00	1.96	0.00	0.02	0.00	2.24	0.00	2.27	0.00	0.34	0.00
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.011	88.93	0.98	62.41	0.69	1.84	0.02	25.07	0.28	61.17	0.67	5.84	0.06
FB 1236	Wine grapes, raw (incl must, juice, wine)	RAC	0.011	129.34	1.42	99.46	1.09	7.76	0.09	46.71	0.51	91.48	1.01	9.23	0.10
FB 0275	Strawberry, raw	RAC	0.72	4.49	3.23	5.66	4.08	0.02	0.01	6.63	4.77	5.75	4.14	0.05	0.04
VC 0424	Cucumber, raw	RAC	0.125	6.72	0.84	11.03	1.38	32.10	4.01	15.10	1.89	4.05	0.51	9.57	1.20
VC 0046	Melons, except watermelon, raw (Cantaloupe)	RAC	0	9.20	0.00	11.95	0.00	14.63	0.00	8.99	0.00	7.86	0.00	2.46	0.00
VO 0448	Tomato, raw (incl juice, incl paste, incl canned)	RAC	0.09	64.74	5.83	68.31	6.15	36.05	3.24	82.09	7.39	54.50	4.91	11.69	1.05
TN 0085	Group of Tree nuts, raw (incl processed)	RAC	0.05	8.52	0.43	8.94	0.45	15.09	0.75	9.60	0.48	14.57	0.73	26.26	1.31
DH 1100	Hops, dry	RAC	2.2	NC	-	NC	-	0.02	0.04	0.02	0.04	NC	-	NC	-
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat)	RAC	0	140.03	0.00	150.89	0.00	79.32	0.00	111.24	0.00	120.30	0.00	51.27	0.00
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0	6.44	0.00	15.51	0.00	3.79	0.00	8.29	0.00	18.44	0.00	8.00	0.00
MO 0105	Edible offal (mammalian), raw	RAC	0.05	15.17	0.76	5.19	0.26	6.30	0.32	6.78	0.34	3.32	0.17	3.17	0.16
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0	388.92	0.00	335.88	0.00	49.15	0.00	331.25	0.00	468.56	0.00	245.45	0.00
PM 0110	Poultry meat, raw (incl prepared)	RAC	0	73.76	0.00	53.86	0.00	23.98	0.00	87.12	0.00	53.38	0.00	84.45	0.00
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0	0.33	0.00	0.72	0.00	0.27	0.00	0.35	0.00	0.80	0.00	NC	-
PE 0112	Eggs, raw, (incl dried)	RAC	0	25.84	0.00	29.53	0.00	28.05	0.00	33.19	0.00	36.44	0.00	8.89	0.00
Total intake (ug/person)=				22.2		24.8		14.1		24.4		20.5		15.6	
Bodyweight per region (kg bw) =				60		60		55		60		60		60	
ADI (ug/person)=				1200		1200		1100		1200		1200		1200	
%ADI=				1.9%		2.1%		1.3%		2.0%		1.7%		1.3%	
Rounded %ADI=				2%		2%		1%		2%		2%		1%	

CLOFENTIZINE (156)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.02 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
FC 0002	Subgroup of Lemons and limes, raw (incl lemon juice) (incl kumquat commodities)	RAC	0.02	18.97	0.38	0.97	0.02	6.23	0.12	0.09	0.00	3.35	0.07
FC 0003	Subgroup of Mandarins, raw (incl mandarin juice)	RAC	0.02	0.16	0.00	0.27	0.01	9.06	0.18	0.01	0.00	0.02	0.00
FC 0004	Subgroup of Oranges, sweet, sour, raw	RAC	0.02	1.18	0.02	1.11	0.02	14.28	0.29	0.05	0.00	1.08	0.02
JF 0004	Subgroup of Oranges, juice (single strength, incl. concentrated)	PP	0.014	0.08	0.00	0.26	0.00	12.61	0.18	0.14	0.00	0.33	0.00
FC 0005	Subgroup of Pummelo and grapefruits, raw (incl grapefruit juice)	RAC	0.02	0.68	0.01	0.05	0.00	3.21	0.06	0.01	0.00	NC	-
FP 0009	Group of Pome fruits, raw (incl apple cider, excl apple juice)	RAC	0.05	68.85	3.44	10.93	0.55	70.82	3.54	189.78	9.49	19.56	0.98
JF 0226	Apple juice, single strength (incl. concentrated)	PP	0.0055	0.03	0.00	0.10	0.00	7.19	0.04	0.03	0.00	NC	-
FS 0012	Group of Stone fruits, raw (incl dried plums, incl dried apricots)	RAC	0.11	0.09	0.01	0.03	0.00	33.36	3.67	0.01	0.00	NC	-
FB 0021	Currants, Black, Red, White, raw	RAC	0.04	0.01	0.00	NC	-	0.74	0.03	NC	-	NC	-
FB 0269	Grapes, raw (incl must, excl dried, excl juice, excl wine)	RAC	0.25	0.14	0.04	0.36	0.09	15.33	3.83	0.01	0.00	0.28	0.07
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	0.28	0.01	0.00	0.13	0.04	1.06	0.30	0.01	0.00	0.03	0.01
JF 0269	Grape juice (from wine grapes)	PP	0	0.01	0.00	0.01	0.00	0.41	0.00	0.01	0.00	NC	-
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.011	0.31	0.00	0.23	0.00	60.43	0.66	0.52	0.01	31.91	0.35
FB 1236	Wine grapes, raw (incl must, juice, wine)	RAC	0.011	0.58	0.01	0.70	0.01	98.85	1.09	0.73	0.01	44.12	0.49
FB 0275	Strawberry, raw	RAC	0.72	0.01	0.01	0.01	0.01	3.35	2.41	0.01	0.01	0.01	0.01
VC 0424	Cucumber, raw	RAC	0.125	0.68	0.09	1.81	0.23	10.40	1.30	0.01	0.00	0.04	0.01
VC 0046	Melons, except watermelon, raw (Cantaloupe)	RAC	0	0.19	0.00	0.10	0.00	4.98	0.00	0.01	0.00	NC	-
VO 0448	Tomato, raw (incl juice, incl paste, incl canned)	RAC	0.09	15.50	1.40	5.78	0.52	71.52	6.44	2.00	0.18	12.50	1.13
TN 0085	Group of Tree nuts, raw (incl processed)	RAC	0.05	4.39	0.22	135.53	6.78	6.11	0.31	0.72	0.04	317.74	15.89
DH 1100	Hops, dry	RAC	2.2	NC	-	NC	-	0.04	0.09	NC	-	NC	-
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat)	RAC	0	29.18	0.00	50.89	0.00	121.44	0.00	22.58	0.00	72.14	0.00
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0	1.05	0.00	1.14	0.00	18.69	0.00	0.94	0.00	3.12	0.00
MO 0105	Edible offal (mammalian), raw	RAC	0.05	4.64	0.23	1.97	0.10	10.01	0.50	3.27	0.16	3.98	0.20
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0	108.75	0.00	70.31	0.00	436.11	0.00	61.55	0.00	79.09	0.00

Annex 3

CLOFENTIZINE (156)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.02 mg/kg bw						
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person								
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake	
PM 0110	Poultry meat, raw (incl prepared)	RAC	0	3.92	0.00	12.03	0.00	57.07	0.00	5.03	0.00	55.56	0.00	
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0	0.10	0.00	0.70	0.00	0.97	0.00	0.10	0.00	NC	-	
PE 0112	Eggs, raw, (incl dried)	RAC	0	3.84	0.00	4.41	0.00	27.25	0.00	1.13	0.00	7.39	0.00	
Total intake (ug/person)=					5.9		8.4		25.0		9.9		19.2	
Bodyweight per region (kg bw) =					60		60		60		60		60	
ADI (ug/person)=					1200		1200		1200		1200		1200	
%ADI=					0.5%		0.7%		2.1%		0.8%		1.6%	
Rounded %ADI=					0%		1%		2%		1%		2%	

CLOTHIANIDIN (238)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw								
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day			Intake as ug/person/day									
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake	
FC 0001	Group of Citrus fruit, raw (incl citrus fruit juice, incl kumquat commodities)	RAC	0.02	34.91	0.70	16.51	0.33	17.23	0.34	104.48	2.09	35.57	0.71	98.49	1.97	
FP 0009	Group of Pome fruits, raw (incl. apple juice, incl apple cider)	RAC	0.1	19.79	1.98	38.25	3.83	17.96	1.80	32.56	3.26	8.08	0.81	64.45	6.45	
FS 0012	Group of Stone fruits, raw (incl dried apricots, excl dried plums)	RAC	0.04	11.33	0.45	23.62	0.94	0.24	0.01	11.32	0.45	2.28	0.09	33.26	1.33	
DF 0014	Plums, dried (prunes)	PP	0.07	0.09	0.01	0.06	0.00	0.01	0.00	0.18	0.01	0.04	0.00	0.06	0.00	
FB 2005	Subgroup of Caneberries, raw	RAC	0.01	0.42	0.00	1.05	0.01	0.01	0.00	0.02	0.00	0.02	0.00	1.24	0.01	
FB 2006	Subgroup of Bush berries, raw (including processed)	RAC	0.01	0.53	0.01	1.31	0.01	0.40	0.00	1.66	0.02	0.01	0.00	0.99	0.01	
FB 2007	Subgroup of Large shrub/tree berries, raw (including processed)	RAC	0.01	0.62	0.01	0.33	0.00	0.34	0.00	1.42	0.01	0.01	0.00	1.51	0.02	
FB 0269	Grapes, raw (incl must, incl wine, excl dried, excl juice)	RAC	0.12	13.94	1.67	26.46	3.18	2.79	0.33	18.58	2.23	8.54	1.02	59.95	7.19	
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	0.31	0.51	0.16	0.51	0.16	0.01	0.00	1.27	0.39	0.12	0.04	2.07	0.64	
JF 0269	Grape juice (from wine grapes)	PP	0.18	0.14	0.03	0.29	0.05	0.05	0.01	0.30	0.05	0.24	0.04	0.05	0.01	
FI 0326	Avocado, raw	RAC	0.01	0.13	0.00	0.03	0.00	2.05	0.02	2.54	0.03	2.34	0.02	0.12	0.00	
FI 0327	Banana, raw (incl plantains) (incl dried)	RAC	0.02	5.23	0.10	6.94	0.14	99.45	1.99	32.47	0.65	48.30	0.97	24.70	0.49	
FI 0345	Mango, raw (incl canned mango, incl mango juice)	RAC	0.02	10.48	0.21	0.01	0.00	7.24	0.14	6.87	0.14	19.98	0.40	6.25	0.13	
FI 0350	Papaya, raw	RAC	0	0.35	0.00	0.01	0.00	3.05	0.00	0.80	0.00	7.28	0.00	1.00	0.00	
FI 0353	Pineapple, raw (incl canned pineapple, incl pineapple juice, incl dried pineapple)	RAC	0	0.61	0.00	1.56	0.00	7.89	0.00	9.36	0.00	8.76	0.00	1.30	0.00	
VB 0040	Group of Brassica vegetables (excl Brassica leafy vegetables), raw	RAC	0.015	6.43	0.10	40.26	0.60	0.80	0.01	9.94	0.15	12.07	0.18	17.73	0.27	
VC 0045	Group of Fruiting vegetables, cucurbits, raw	RAC	0.02	53.14	1.06	86.21	1.72	6.28	0.13	92.76	1.86	15.64	0.31	155.30	3.11	
VO 0050	Group of Fruiting vegetables other than cucurbits, raw, (incl processed commodities, excl dried chilli peppers)	RAC	0.02	67.79	1.36	99.85	2.00	31.70	0.63	125.86	2.52	55.22	1.10	262.82	5.26	
VL 0053	Group of Leafy vegetables, raw	RAC	0.52	8.47	4.40	22.36	11.63	7.74	4.02	25.51	13.27	45.77	23.80	21.22	11.03	
VP 0061	Beans with pods (Phaseolus spp): (immature pods + succulent seeds)	RAC	0.07	0.68	0.05	NC	-	NC	-	0.39	0.03	0.22	0.02	0.49	0.03	
014B	Peas with pods	-	0.01	-	-	-	-	-	-	-	-	-	-	-	-	
VP 0062	Beans without pods: (Phaseolus spp.) (succulent seeds), raw	RAC	0.01	1.56	0.02	0.60	0.01	0.49	0.00	1.18	0.01	0.90	0.01	7.79	0.08	
VP 0523	Broad bean without pods (succulent seeds)	RAC	0.01	3.51	0.04	0.43	0.00	0.01	0.00	0.60	0.01	0.29	0.00	0.78	0.01	

Annex 3

CLOTHIANIDIN (238)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
	(Vicia spp), raw														
VP 0064	Peas without pods (Pisum spp) (succulent seeds)	RAC	0.01	1.97	0.02	0.51	0.01	0.02	0.00	0.79	0.01	3.68	0.04	3.80	0.04
VD 0070	Group of Pulses, raw (incl processed)	RAC	0.02	87.29	1.75	64.04	1.28	37.15	0.74	89.82	1.80	91.02	1.82	98.20	1.96
VD 0520	Bambara groundnut (dry) (Vigna subterranea), raw	RAC	0.01	NC	-	NC	-	0.20	0.00	NC	-	NC	-	NC	-
VR 0075	Group of Root and tuber vegetables, raw (incl processed)	RAC	0.02	87.83	1.76	374.04	7.48	668.92	13.38	121.64	2.43	94.20	1.88	247.11	4.94
VS 0623	Cardoon	RAC	0.01	0.24	0.00	0.43	0.00	0.27	0.00	0.64	0.01	NC	-	0.31	0.00
VS 0624	Celery	RAC	0.01	2.14	0.02	3.79	0.04	2.35	0.02	5.69	0.06	0.02	0.00	2.75	0.03
VS 0627	Rhubarb	RAC	0.012	0.73	0.01	1.30	0.02	0.80	0.01	1.95	0.02	NC	-	0.94	0.01
VS 0621	Asparagus, raw	RAC	0.01	0.01	0.00	0.01	0.00	0.01	0.00	0.03	0.00	0.07	0.00	0.21	0.00
VS 0622	Bamboo shoots, raw	RAC	0.01	1.72	0.02	3.05	0.03	1.89	0.02	4.59	0.05	NC	-	2.21	0.02
VS 0620	Artichoke globe, raw	RAC	0.024	0.69	0.02	0.01	0.00	0.01	0.00	0.32	0.01	0.26	0.01	1.21	0.03
VS 0626	Palm hearts, raw	RAC	0.01	0.39	0.00	0.70	0.01	0.43	0.00	1.05	0.01	2.27	0.02	0.51	0.01
GC 0650	Rye, raw	RAC	0.01	NC	-	NC	-	0.02	0.00	0.01	0.00	0.01	0.00	0.01	0.00
CF 1250	Rye, flour (white flour and wholemeal flour)	PP	0.00645	0.11	0.00	15.51	0.10	0.06	0.00	0.10	0.00	0.03	0.00	1.72	0.01
GC 0653	Triticale, raw	RAC	0.01	NC	-	NC	-	NC	-	0.01	0.00	NC	-	NC	-
GC 0653	Triticale, flour (white flour and wholemeal flour)	PP	0.00645	NC	-	NC	-	NC	-	NC	-	0.31	0.00	NC	-
GC 0654	Wheat, raw (incl bulgur, incl fermented beverages, excl germ, excl wholemeal bread, excl white flour products, excl white bread)	RAC	0.01	0.01	0.00	1.12	0.01	NC	-	0.03	0.00	0.56	0.01	NC	-
CF 1210	Wheat, germ	PP	0.018	NC	-	NC	-	0.01	0.00	0.01	0.00	0.14	0.00	0.01	0.00
CP 1212	Wheat, wholemeal bread	PP	0.01	0.01	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.03	0.00	0.01	0.00
CP 1211	Wheat, white bread	PP	0.01	0.25	0.00	0.63	0.01	0.12	0.00	0.43	0.00	1.39	0.01	0.22	0.00
CF 1211	Wheat, white flour (incl white flour products: starch, gluten, macaroni, pastry)	PP	0.00645	301.24	1.94	268.64	1.73	30.21	0.19	222.51	1.44	134.73	0.87	343.12	2.21
GC 0640	Barley, raw (incl malt extract, incl beer, incl malt, excl pot&pearled, excl flour & grits)	RAC	0.015	3.55	0.05	19.31	0.29	4.98	0.07	3.02	0.05	7.85	0.12	3.98	0.06
-	Barley, flour (white flour and wholemeal flour)	PP	0.0097	2.93	0.03	0.30	0.00	0.02	0.00	0.01	0.00	0.48	0.00	0.01	0.00
GC 0647	Oats, raw (incl rolled)	RAC	0.015	0.05	0.00	7.05	0.11	0.10	0.00	1.71	0.03	0.96	0.01	0.04	0.00
CM 0649 (GC 0649)	Rice, husked, dry (incl flour, incl oil, incl beverages, incl starch, excl polished)	REP	0.145	1.26	0.18	1.58	0.23	31.05	4.50	5.43	0.79	0.90	0.13	2.18	0.32

[illegible]

Annex 3

CLOTHIANIDIN (238)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
	Total intake (ug/person)=				28.6		45.5		36.4		52.5		58.2		62.5
	Bodyweight per region (kg bw) =				60		60		60		60		60		60
	ADI (ug/person)=				6000		6000		6000		6000		6000		6000
	%ADI=				0.5%		0.8%		0.6%		0.9%		1.0%		1.0%
	Rounded %ADI=				0%		1%		1%		1%		1%		1%

CLOTHIANIDIN (238)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
FC 0001	Group of Citrus fruit, raw (incl citrus fruit juice, incl kumquat commodities)	RAC	0.02	114.42	2.29	62.91	1.26	26.97	0.54	96.72	1.93	96.22	1.92	563.19	11.26
FP 0009	Group of Pome fruits, raw (incl. apple juice, incl apple cider)	RAC	0.1	71.38	7.14	81.73	8.17	42.91	4.29	58.89	5.89	103.85	10.39	12.48	1.25
FS 0012	Group of Stone fruits, raw (incl dried apricots, excl dried plums)	RAC	0.04	18.18	0.73	23.83	0.95	14.27	0.57	18.52	0.74	9.35	0.37	0.11	0.00
DF 0014	Plums, dried (prunes)	PP	0.07	0.61	0.04	0.35	0.02	0.05	0.00	0.35	0.02	0.49	0.03	0.13	0.01
FB 2005	Subgroup of Caneberries, raw	RAC	0.01	0.56	0.01	1.43	0.01	0.14	0.00	1.23	0.01	1.14	0.01	0.01	0.00
FB 2006	Subgroup of Bush berries, raw (including processed)	RAC	0.01	1.31	0.01	5.50	0.06	0.01	0.00	2.57	0.03	0.82	0.01	2.15	0.02
FB 2007	Subgroup of Large shrub/tree berries, raw (including processed)	RAC	0.01	8.26	0.08	0.14	0.00	0.07	0.00	0.13	0.00	0.19	0.00	1.87	0.02
FB 0269	Grapes, raw (incl must, incl wine, excl dried, excl juice)	RAC	0.12	128.64	15.44	97.04	11.64	7.74	0.93	43.94	5.27	88.68	10.64	8.80	1.06
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	0.31	3.09	0.96	1.51	0.47	0.03	0.01	1.38	0.43	4.26	1.32	0.42	0.13
JF 0269	Grape juice (from wine grapes)	PP	0.18	0.56	0.10	1.96	0.35	0.02	0.00	2.24	0.40	2.27	0.41	0.34	0.06
FI 0326	Avocado, raw	RAC	0.01	2.65	0.03	0.87	0.01	0.46	0.00	1.64	0.02	1.30	0.01	0.96	0.01
FI 0327	Banana, raw (incl plantains) (incl dried)	RAC	0.02	25.76	0.52	23.65	0.47	23.83	0.48	24.37	0.49	19.43	0.39	101.55	2.03
FI 0345	Mango, raw (incl canned mango, incl mango juice)	RAC	0.02	1.80	0.04	0.63	0.01	10.05	0.20	1.07	0.02	3.52	0.07	16.44	0.33
FI 0350	Papaya, raw	RAC	0	0.31	0.00	0.18	0.00	1.50	0.00	0.51	0.00	0.54	0.00	1.08	0.00

CLOTHIANIDIN (238)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
FI 0353	Pineapple, raw (incl canned pineapple, incl pineapple juice, incl dried pineapple)	RAC	0	13.13	0.00	11.13	0.00	6.94	0.00	14.36	0.00	36.74	0.00	18.81	0.00
VB 0040	Group of Brassica vegetables (excl Brassica leafy vegetables), raw	RAC	0.015	20.71	0.31	39.81	0.60	25.06	0.38	37.93	0.57	18.12	0.27	16.74	0.25
VC 0045	Group of Fruiting vegetables, cucurbits, raw	RAC	0.02	27.81	0.56	41.93	0.84	123.30	2.47	49.47	0.99	15.95	0.32	35.99	0.72
VO 0050	Group of Fruiting vegetables other than cucurbits, raw, (incl processed commodities, excl dried chilli peppers)	RAC	0.02	72.14	1.44	85.53	1.71	76.55	1.53	95.63	1.91	64.19	1.28	16.94	0.34
VL 0053	Group of Leafy vegetables, raw	RAC	0.52	18.83	9.79	21.85	11.36	121.23	63.04	43.09	22.41	18.18	9.45	18.32	9.53
VP 0061	Beans with pods (Phaseolus spp): (immature pods + succulent seeds)	RAC	0.07	5.07	0.35	0.83	0.06	0.17	0.01	3.70	0.26	NC	-	NC	-
014B	Peas with pods	-	0.01	-	-	-	-	-	-	-	-	-	-	-	-
VP 0062	Beans without pods: (Phaseolus spp.) (succulent seeds), raw	RAC	0.01	2.21	0.02	5.25	0.05	4.17	0.04	1.61	0.02	16.95	0.17	0.17	0.00
VP 0523	Broad bean without pods (succulent seeds) (Vicia spp), raw	RAC	0.01	0.22	0.00	0.84	0.01	0.15	0.00	0.48	0.00	2.04	0.02	NC	-
VP 0064	Peas without pods (Pisum spp) (succulent seeds)	RAC	0.01	10.72	0.11	1.99	0.02	2.72	0.03	4.26	0.04	4.23	0.04	NC	-
VD 0070	Group of Pulses, raw (incl processed)	RAC	0.02	112.88	2.26	123.05	2.46	47.73	0.95	204.75	4.10	227.52	4.55	110.05	2.20
VD 0520	Bambara groundnut (dry) (Vigna subterranea), raw	RAC	0.01	NC	-	NC	-	NC	-	NC	-	NC	-	NC	-
VR 0075	Group of Root and tuber vegetables, raw (incl processed)	RAC	0.02	290.31	5.81	300.35	6.01	214.25	4.29	242.72	4.85	348.67	6.97	137.52	2.75
VS 0623	Cardoon	RAC	0.01	0.02	0.00	3.49	0.03	NC	-	0.02	0.00	NC	-	0.46	0.00
VS 0624	Celery	RAC	0.01	7.68	0.08	2.85	0.03	NC	-	3.34	0.03	16.83	0.17	4.04	0.04
VS 0627	Rhubarb	RAC	0.012	1.61	0.02	2.23	0.03	NC	-	0.52	0.01	7.63	0.09	1.39	0.02
VS 0621	Asparagus, raw	RAC	0.01	0.84	0.01	2.08	0.02	7.11	0.07	1.01	0.01	1.69	0.02	0.04	0.00
VS 0622	Bamboo shoots, raw	RAC	0.01	0.92	0.01	0.55	0.01	61.79	0.62	NC	-	1.72	0.02	3.26	0.03
VS 0620	Artichoke globe, raw	RAC	0.024	0.98	0.02	3.65	0.09	0.07	0.00	1.67	0.04	0.26	0.01	NC	-
VS 0626	Palm hearts, raw	RAC	0.01	0.51	0.01	0.73	0.01	3.54	0.04	0.01	0.00	0.66	0.01	0.75	0.01
GC 0650	Rye, raw	RAC	0.01	0.01	0.00	NC	-	0.06	0.00	0.01	0.00	NC	-	NC	-
CF 1250	Rye, flour (white flour and wholemeal flour)	PP	0.00645	2.57	0.02	28.31	0.18	0.12	0.00	5.20	0.03	1.20	0.01	NC	-
GC 0653	Triticale, raw	RAC	0.01	NC	-	NC	-	0.01	0.00	0.01	0.00	NC	-	NC	-
GC 0653	Triticale, flour (white flour and wholemeal flour)	PP	0.00645	0.01	0.00	0.14	0.00	0.23	0.00	NC	-	NC	-	NC	-

Annex 3

CLOTHIANIDIN (238)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
GC 0654	Wheat, raw (incl bulgur, incl fermented beverages, excl germ, excl wholemeal bread, excl white flour products, excl white bread)	RAC	0.01	NC	-	NC	-	0.02	0.00	0.83	0.01	NC	-	NC	-
CF 1210	Wheat, germ	PP	0.018	0.97	0.02	0.10	0.00	0.03	0.00	0.01	0.00	NC	-	0.04	0.00
CP 1212	Wheat, wholemeal bread	PP	0.01	0.03	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.05	0.00	0.02	0.00
CP 1211	Wheat, white bread	PP	0.01	1.30	0.01	0.46	0.00	0.06	0.00	0.22	0.00	2.44	0.02	0.77	0.01
CF 1211	Wheat, white flour (incl white flour products: starch, gluten, macaroni, pastry)	PP	0.00645	198.08	1.28	193.03	1.25	106.24	0.69	185.09	1.19	168.67	1.09	131.59	0.85
GC 0640	Barley, raw (incl malt extract, incl beer, incl malt, excl pot&pearled, excl flour & grits)	RAC	0.015	35.17	0.53	49.45	0.74	8.86	0.13	34.31	0.51	44.87	0.67	15.82	0.24
-	Barley, flour (white flour and wholemeal flour)	PP	0.0097	0.08	0.00	0.03	0.00	0.01	0.00	0.05	0.00	0.68	0.01	0.05	0.00
GC 0647	Oats, raw (incl rolled)	RAC	0.015	7.50	0.11	6.26	0.09	0.15	0.00	4.87	0.07	3.16	0.05	2.98	0.04
CM 0649 (GC 0649)	Rice, husked, dry (incl flour, incl oil, incl beverages, incl starch, excl polished)	REP	0.145	3.70	0.54	2.11	0.31	1.51	0.22	1.75	0.25	0.29	0.04	5.12	0.74
CM 1205	Rice polished, dry	PP	0.12	13.38	1.61	10.80	1.30	262.08	31.45	57.16	6.86	12.83	1.54	62.78	7.53
-	Rice bran oil	PP	0.28	NC	-	NC	-	0.15	0.04	0.10	0.03	NC	-	NC	-
GC 0651	Sorghum, raw (i.e. Chicken corn, Dari seed, Durra, Feterita)	RAC	0.018	NC	-	NC	-	0.01	0.00	1.15	0.02	NC	-	7.12	0.13
-	Sorghum, flour (white flour and wholemeal flour)	PP	0.0114	NC	-	NC	-	1.29	0.01	0.01	0.00	NC	-	NC	-
GC 2090	Subgroup of Sweet Corns	RAC	0.01	11.43	0.11	3.71	0.04	0.74	0.01	13.63	0.14	3.07	0.03	1.50	0.02
GS 0659	Sugar cane, raw (incl sugar, incl molasses)	RAC	0.03	92.24	2.77	95.72	2.87	28.47	0.85	77.39	2.32	117.73	3.53	103.90	3.12
TN 0672	Pecan, nutmeat	RAC	0.01	0.38	0.00	NC	-	NC	-	0.27	0.00	NC	-	0.26	0.00
SO 0091	Subgroup of Oilseeds, raw (incl processed)	RAC	0.02	78.69	1.57	62.08	1.24	39.56	0.79	52.51	1.05	37.72	0.75	16.21	0.32
SB 0715	Cacao bean, raw (incl roasted, incl powder, incl butter, incl paste, incl nes products)	RAC	0.02	7.54	0.15	5.59	0.11	0.29	0.01	4.14	0.08	1.27	0.03	5.29	0.11
SB 0716	Coffee bean raw (incl roasted, incl instant coffee, incl substitutes)	RAC	0.015	10.90	0.16	12.44	0.19	0.77	0.01	9.48	0.14	22.07	0.33	8.15	0.12
HH 0738	Mint, raw	RAC	0.11	NC	-	NC	-	NC	-	NC	-	NC	-	NC	-
HS 0444	Peppers, chili, dried	PP	0.2	0.11	0.02	0.21	0.04	0.36	0.07	0.21	0.04	0.25	0.05	0.15	0.03
DH 1100	Hops, dry	RAC	0.026	NC	-	NC	-	0.02	0.00	0.02	0.00	NC	-	NC	-
DT 1114	Tea, green or black, fermented and dried	RAC	0.12	2.71	0.33	0.82	0.10	1.14	0.14	1.59	0.19	1.82	0.22	0.53	0.06

CLOTHIANIDIN (238)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw													
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day			Intake as ug/person/day														
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake						
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat)	RAC	0.02	140.03	2.80	150.89	3.02	79.32	1.59	111.24	2.22	120.30	2.41	51.27	1.03						
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.02	6.44	0.13	15.51	0.31	3.79	0.08	8.29	0.17	18.44	0.37	8.00	0.16						
MO 0105	Edible offal (mammalian), raw	RAC	0.035	15.17	0.53	5.19	0.18	6.30	0.22	6.78	0.24	3.32	0.12	3.17	0.11						
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.004	388.92	1.56	335.88	1.34	49.15	0.20	331.25	1.33	468.56	1.87	245.45	0.98						
PM 0110	Poultry meat, raw (incl prepared)	RAC	0.0014	73.76	0.10	53.86	0.08	23.98	0.03	87.12	0.12	53.38	0.07	84.45	0.12						
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0.0014	66.38	0.09	48.47	0.07	21.58	0.03	78.41	0.11	48.04	0.07	76.01	0.11						
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0.0033	7.38	0.02	5.39	0.02	2.40	0.01	8.71	0.03	5.34	0.02	8.45	0.03						
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.0033	0.10	0.00	0.10	0.00	NC	-	0.10	0.00	0.71	0.00	NC	-						
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.37	0.33	0.12	0.72	0.27	0.27	0.10	0.35	0.13	0.80	0.30	NC	-						
PE 0112	Eggs, raw, (incl dried)	RAC	0.0062	25.84	0.16	29.53	0.18	28.05	0.17	33.19	0.21	36.44	0.23	8.89	0.06						
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-						
Total intake (ug/person)=				62.9			60.7			117.3			68.0			62.8			48.0		
Bodyweight per region (kg bw) =				60			60			55			60			60			60		
ADI (ug/person)=				6000			6000			5500			6000			6000			6000		
%ADI=				1.0%			1.0%			2.1%			1.1%			1.0%			0.8%		
Rounded %ADI=				1%			1%			2%			1%			1%			1%		

Annex 3

CLOTHIANIDIN (238)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
FC 0001	Group of Citrus fruit, raw (incl citrus fruit juice, incl kumquat commodities)	RAC	0.02	21.16	0.42	2.94	0.06	58.52	1.17	0.44	0.01	5.13	0.10
FP 0009	Group of Pome fruits, raw (incl. apple juice, incl apple cider)	RAC	0.1	68.89	6.89	11.06	1.11	80.62	8.06	189.82	18.98	19.56	1.96
FS 0012	Group of Stone fruits, raw (incl dried apricots, excl dried plums)	RAC	0.04	0.09	0.00	0.02	0.00	32.27	1.29	0.01	0.00	NC	-
DF 0014	Plums, dried (prunes)	PP	0.07	0.01	0.00	0.01	0.00	0.37	0.03	0.01	0.00	NC	-
FB 2005	Subgroup of Caneberries, raw	RAC	0.01	0.01	0.00	7.30	0.07	2.29	0.02	0.01	0.00	NC	-
FB 2006	Subgroup of Bush berries, raw (including processed)	RAC	0.01	0.82	0.01	4.05	0.04	5.94	0.06	0.43	0.00	2.66	0.03
FB 2007	Subgroup of Large shrub/tree berries, raw (including processed)	RAC	0.01	0.71	0.01	7.32	0.07	NC	-	0.38	0.00	2.32	0.02
FB 0269	Grapes, raw (incl must, incl wine, excl dried, excl juice)	RAC	0.12	0.57	0.07	0.69	0.08	98.34	11.80	0.73	0.09	44.12	5.29
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	0.31	0.01	0.00	0.13	0.04	1.06	0.33	0.01	0.00	0.03	0.01
JF 0269	Grape juice (from wine grapes)	PP	0.18	0.01	0.00	0.01	0.00	0.41	0.07	0.01	0.00	NC	-
FI 0326	Avocado, raw	RAC	0.01	1.12	0.01	0.01	0.00	0.84	0.01	0.01	0.00	6.60	0.07
FI 0327	Banana, raw (incl plantains) (incl dried)	RAC	0.02	44.80	0.90	118.17	2.36	25.25	0.51	454.49	9.09	310.23	6.20
FI 0345	Mango, raw (incl canned mango, incl mango juice)	RAC	0.02	12.25	0.25	6.83	0.14	0.76	0.02	0.01	0.00	20.12	0.40
FI 0350	Papaya, raw	RAC	0	6.47	0.00	0.25	0.00	0.19	0.00	0.01	0.00	26.42	0.00
FI 0353	Pineapple, raw (incl canned pineapple, incl pineapple juice, incl dried pineapple)	RAC	0	8.51	0.00	6.27	0.00	6.89	0.00	0.18	0.00	24.94	0.00
VB 0040	Group of Brassica vegetables (excl Brassica leafy vegetables), raw	RAC	0.015	5.46	0.08	4.28	0.06	58.72	0.88	0.02	0.00	NC	-
VC 0045	Group of Fruiting vegetables, cucurbits, raw	RAC	0.02	5.96	0.12	9.74	0.19	51.82	1.04	13.61	0.27	0.05	0.00
VO 0050	Group of Fruiting vegetables other than cucurbits, raw, (incl processed commodities, excl dried chilli peppers)	RAC	0.02	32.01	0.64	28.27	0.57	100.61	2.01	2.91	0.06	12.50	0.25
VL 0053	Group of Leafy vegetables, raw	RAC	0.52	12.42	6.46	8.75	4.55	7.53	3.92	7.07	3.68	14.11	7.34
VP 0061	Beans with pods (Phaseolus spp): (immature pods + succulent seeds)	RAC	0.07	NC	-	NC	-	NC	-	NC	-	NC	-
014B	Peas with pods	-	0.01	-	-	-	-	-	-	-	-	-	-
VP 0062	Beans without pods: (Phaseolus spp.) (succulent seeds), raw	RAC	0.01	0.30	0.00	3.13	0.03	4.11	0.04	0.01	0.00	NC	-

CLOTHIANIDIN (238)		International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
VP 0523	Broad bean without pods (succulent seeds) (Vicia spp), raw	RAC	0.01	0.07	0.00	0.01	0.00	0.76	0.01	NC	-	NC	-
VP 0064	Peas without pods (Pisum spp) (succulent seeds)	RAC	0.01	0.21	0.00	0.02	0.00	5.51	0.06	0.02	0.00	NC	-
VD 0070	Group of Pulses, raw (incl processed)	RAC	0.02	46.57	0.93	30.77	0.62	112.53	2.25	75.53	1.51	43.68	0.87
VD 0520	Bambara groundnut (dry) (Vigna subterranea), raw	RAC	0.01	0.20	0.00	NC	-	NC	-	NC	-	NC	-
VR 0075	Group of Root and tuber vegetables, raw (incl processed)	RAC	0.02	282.25	5.65	232.11	4.64	281.91	5.64	620.21	12.40	459.96	9.20
VS 0623	Cardoon	RAC	0.01	0.41	0.00	0.30	0.00	NC	-	0.28	0.00	0.56	0.01
VS 0624	Celery	RAC	0.01	3.66	0.04	2.65	0.03	4.84	0.05	2.47	0.02	4.94	0.05
VS 0627	Rhubarb	RAC	0.012	1.26	0.02	0.91	0.01	0.96	0.01	0.85	0.01	1.70	0.02
VS 0621	Asparagus, raw	RAC	0.01	0.01	0.00	0.01	0.00	0.17	0.00	0.01	0.00	NC	-
VS 0622	Bamboo shoots, raw	RAC	0.01	2.95	0.03	2.13	0.02	1.52	0.02	2.00	0.02	3.99	0.04
VS 0620	Artichoke globe, raw	RAC	0.024	0.01	0.00	NC	-	0.08	0.00	0.01	0.00	NC	-
VS 0626	Palm hearts, raw	RAC	0.01	0.67	0.01	0.49	0.00	NC	-	0.46	0.00	0.91	0.01
GC 0650	Rye, raw	RAC	0.01	0.01	0.00	NC	-	NC	-	0.01	0.00	NC	-
CF 1250	Rye, flour (white flour and wholemeal flour)	PP	0.00645	0.02	0.00	0.01	0.00	11.16	0.07	0.01	0.00	0.70	0.00
GC 0653	Triticale, raw	RAC	0.01	0.01	0.00	NC	-	NC	-	NC	-	NC	-
GC 0653	Triticale, flour (white flour and wholemeal flour)	PP	0.00645	NC	-	NC	-	NC	-	NC	-	NC	-
GC 0654	Wheat, raw (incl bulgur, incl fermented beverages, excl germ, excl wholemeal bread, excl white flour products, excl white bread)	RAC	0.01	0.01	0.00	NC	-	NC	-	NC	-	0.97	0.01
CF 1210	Wheat, germ	PP	0.018	0.04	0.00	0.01	0.00	0.01	0.00	0.01	0.00	NC	-
CP 1212	Wheat, wholemeal bread	PP	0.01	0.01	0.00	0.01	0.00	0.03	0.00	0.01	0.00	0.01	0.00
CP 1211	Wheat, white bread	PP	0.01	0.43	0.00	0.41	0.00	1.56	0.02	0.11	0.00	0.07	0.00
CF 1211	Wheat, white flour (incl white flour products: starch, gluten, macaroni, pastry)	PP	0.00645	44.78	0.29	86.96	0.56	214.05	1.38	20.31	0.13	103.60	0.67
GC 0640	Barley, raw (incl malt extract, incl beer, incl malt, excl pot&pearled, excl flour & grits)	RAC	0.015	3.15	0.05	2.31	0.03	43.92	0.66	3.72	0.06	16.26	0.24
-	Barley, flour (white flour and wholemeal flour)	PP	0.0097	0.02	0.00	NC	-	0.32	0.00	0.01	0.00	NC	-
GC 0647	Oats, raw (incl rolled)	RAC	0.015	0.37	0.01	0.07	0.00	2.79	0.04	0.10	0.00	NC	-
CM 0649 (GC 0649)	Rice, husked, dry (incl flour, incl oil, incl beverages, incl starch, excl polished)	REP	0.145	13.58	1.97	4.29	0.62	2.17	0.31	0.01	0.00	8.84	1.28

Annex 3

CLOTHIANIDIN (238)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
CM 1205	Rice polished, dry	PP	0.12	30.20	3.62	218.34	26.20	12.77	1.53	15.24	1.83	51.35	6.16
-	Rice bran oil	PP	0.28	NC	-	0.60	0.17	NC	-	NC	-	NC	-
GC 0651	Sorghum, raw (i.e. Chicken corn, Dari seed, Durra, Feterita)	RAC	0.018	NC	-	NC	-	NC	-	NC	-	NC	-
-	Sorghum, flour (white flour and wholemeal flour)	PP	0.0114	75.99	0.87	1.82	0.02	NC	-	19.82	0.23	NC	-
GC 2090	Subgroup of Sweet Corns	RAC	0.01	3.63	0.04	20.50	0.21	8.78	0.09	0.02	0.00	0.17	0.00
GS 0659	Sugar cane, raw (incl sugar, incl molasses)	RAC	0.03	33.75	1.01	106.29	3.19	78.09	2.34	29.09	0.87	45.70	1.37
TN 0672	Pecan, nutmeat	RAC	0.01	0.15	0.00	0.22	0.00	0.31	0.00	0.01	0.00	0.01	0.00
SO 0091	Subgroup of Oilseeds, raw (incl processed)	RAC	0.02	94.19	1.88	14.81	0.30	55.34	1.11	32.27	0.65	55.44	1.11
SB 0715	Cacao bean, raw (incl roasted, incl powder, incl butter, incl paste, incl nes products)	RAC	0.02	0.11	0.00	0.89	0.02	6.28	0.13	0.17	0.00	2.31	0.05
SB 0716	Coffee bean raw (incl roasted, incl instant coffee, incl substitutes)	RAC	0.015	0.95	0.01	1.32	0.02	11.64	0.17	2.96	0.04	14.73	0.22
HH 0738	Mint, raw	RAC	0.11	NC	-	NC	-	NC	-	NC	-	NC	-
HS 0444	Peppers, chili, dried	PP	0.2	0.58	0.12	1.27	0.25	1.21	0.24	0.12	0.02	NC	-
DH 1100	Hops, dry	RAC	0.026	NC	-	NC	-	0.04	0.00	NC	-	NC	-
DT 1114	Tea, green or black, fermented and dried	RAC	0.12	0.53	0.06	5.25	0.63	0.63	0.08	0.56	0.07	0.82	0.10
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat)	RAC	0.02	29.18	0.58	50.89	1.02	121.44	2.43	22.58	0.45	72.14	1.44
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.02	1.05	0.02	1.14	0.02	18.69	0.37	0.94	0.02	3.12	0.06
MO 0105	Edible offal (mammalian), raw	RAC	0.035	4.64	0.16	1.97	0.07	10.01	0.35	3.27	0.11	3.98	0.14
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.004	108.75	0.44	70.31	0.28	436.11	1.74	61.55	0.25	79.09	0.32
PM 0110	Poultry meat, raw (incl prepared)	RAC	0.0014	3.92	0.01	12.03	0.02	57.07	0.08	5.03	0.01	55.56	0.08
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0.0014	3.53	0.00	10.83	0.02	51.36	0.07	4.53	0.01	50.00	0.07
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0.0033	0.39	0.00	1.20	0.00	5.71	0.02	0.50	0.00	5.56	0.02
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.0033	NC	-	NC	-	0.32	0.00	NC	-	NC	-
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.37	0.10	0.04	0.70	0.26	0.97	0.36	0.10	0.04	NC	-
PE 0112	Eggs, raw, (incl dried)	RAC	0.0062	3.84	0.02	4.41	0.03	27.25	0.17	1.13	0.01	7.39	0.05
-	-	-		-	-	-	-	-	-	-	-	-	-
Total intake (ug/person)=				33.7		48.6		53.1		51.0		45.3	

CLOTHIANIDIN (238)			International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw						
Codex Code	Commodity description	Expr as	STM mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
	Bodyweight per region (kg bw) =				60		60		60		60		60
	ADI (ug/person)=				6000		6000		6000		6000		6000
	%ADI=				0.6%		0.8%		0.9%		0.8%		0.8%
	Rounded %ADI=				1%		1%		1%		1%		1%

Annex 3

CYPRODINIL (207)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
FP 0009	Group of Pome fruits, raw (incl apple cider, excl apple juice)	RAC	0.48	19.35	9.29	34.06	16.35	17.87	8.58	25.74	12.36	7.69	3.69	56.85	27.29
JF 0226	Apple juice, single strength (incl. concentrated)	PP	0.015	0.32	0.00	3.07	0.05	0.07	0.00	5.00	0.08	0.29	0.00	5.57	0.08
FS 0012	Group of Stone fruits, raw (incl dried apricots, excl dried plums)	RAC	0.68	11.33	7.70	23.62	16.06	0.24	0.16	11.32	7.70	2.28	1.55	33.26	22.62
DF 0014	Plums, dried (prunes)	PP	1.2	0.09	0.11	0.06	0.07	0.01	0.01	0.18	0.22	0.04	0.05	0.06	0.07
FB 2005	Subgroup of Caneberries, raw	RAC	2.2	0.42	0.92	1.05	2.31	0.01	0.02	0.02	0.04	0.02	0.04	1.24	2.73
FB 2006	Subgroup of Bush berries, raw (including processed)	RAC	2.2	0.53	1.17	1.31	2.88	0.40	0.88	1.66	3.65	0.01	0.02	0.99	2.18
FB 2007	Subgroup of Large shrub/tree berries, raw (including processed)	RAC	2.2	0.62	1.36	0.33	0.73	0.34	0.75	1.42	3.12	0.01	0.02	1.51	3.32
FB 0269	Grapes, raw (incl must, excl dried, excl juice, excl wine)	RAC	0.79	13.02	10.29	9.25	7.31	0.03	0.02	16.91	13.36	3.70	2.92	54.44	43.01
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	1.7	0.51	0.87	0.51	0.87	0.01	0.02	1.27	2.16	0.12	0.20	2.07	3.52
JF 0269	Grape juice (from wine grapes)	PP	0.12	0.14	0.02	0.29	0.03	0.05	0.01	0.30	0.04	0.24	0.03	0.05	0.01
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.062	0.67	0.04	12.53	0.78	2.01	0.12	1.21	0.08	3.53	0.22	4.01	0.25
FB 2009	Subgroup of Low growing berries, raw	RAC	2.2	0.71	1.56	2.02	4.44	0.04	0.09	1.39	3.06	0.37	0.81	2.53	5.57
FT 0336	Guava, raw	RAC	0.485	0.47	0.23	0.01	0.00	0.48	0.23	0.49	0.24	4.42	2.14	0.06	0.03
FI 0326	Avocado, raw	RAC	0.265	0.13	0.03	0.03	0.01	2.05	0.54	2.54	0.67	2.34	0.62	0.12	0.03
FI 0355	Pomegranate, raw, (incl processed)	RAC	3.3	3.40	11.22	2.10	6.93	2.65	8.75	10.89	35.94	NC	-	6.67	22.01
-	Onions, dry, raw	RAC	0.065	29.36	1.91	37.50	2.44	3.56	0.23	34.78	2.26	18.81	1.22	43.38	2.82
VA 2032	Subgroup of Green Onions	RAC	0.065	2.64	0.17	3.09	0.20	1.05	0.07	2.89	0.19	0.61	0.04	5.24	0.34
VB 0042	Subgroup of Flowerhead Brassica, raw	RAC	0.27	2.54	0.69	0.49	0.13	0.01	0.00	3.57	0.96	7.79	2.10	3.12	0.84
VB 0041	Cabbages, head, raw	RAC	0.03	2.73	0.08	27.92	0.84	0.55	0.02	4.47	0.13	4.27	0.13	10.25	0.31
VC 0045	Group of Fruiting vegetables, cucurbits, raw	RAC	0.09	53.14	4.78	86.21	7.76	6.28	0.57	92.76	8.35	15.64	1.41	155.30	13.98
VO 0448	Tomato, raw (incl canned, excl juice, excl paste)	RAC	0.24	42.04	10.09	76.13	18.27	10.69	2.57	84.59	20.30	24.92	5.98	203.27	48.78
-	Tomato, paste (i.e. concentrated tomato sauce/puree)	PP	0.48	2.34	1.12	1.33	0.64	1.57	0.75	4.24	2.04	0.34	0.16	2.83	1.36
JF 0448	Tomato, juice (single strength, incl concentrated)	PP	0.036	0.29	0.01	0.29	0.01	0.01	0.00	0.38	0.01	0.05	0.00	0.14	0.01
VO 0051	Subgroup of peppers, raw (incl dried sweet peppers, excl dried chilipeppers), incl okra	RAC	0.24	10.45	2.51	13.74	3.30	13.81	3.31	14.53	3.49	15.25	3.66	27.93	6.70
VO 2046	Subgroup of eggplants	RAC	0.24	5.58	1.34	4.31	1.03	0.89	0.21	9.31	2.23	13.64	3.27	20.12	4.83

CYPRODINIL (207)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
VL 2050	Subgroup of Leafy greens	RAC	11	3.93	43.23	5.28	58.08	3.07	33.77	14.53	159.83	8.25	90.75	12.75	140.25
VL 0054	Subgroup of Leaves of Brassicaceae, raw	RAC	0.37	2.63	0.97	9.27	3.43	1.86	0.69	5.82	2.15	19.53	7.23	4.90	1.81
VL 2052	Subgroup of Leaves of Root and Tuber Vegetables	RAC	11	0.18	1.98	0.31	3.41	0.84	9.24	0.47	5.17	2.06	22.66	0.23	2.53
VL 2053	Subgroup of Leaves of trees, shrubs and vines	RAC	11	0.39	4.29	0.69	7.59	0.43	4.73	1.04	11.44	4.57	50.27	0.50	5.50
VL 2054	Subgroup of Leafy aquatic vegetables	RAC	11	1.21	13.31	2.15	23.65	1.33	14.63	3.24	35.64	11.36	124.96	1.56	17.16
VL 2832	Witloof chicory (sprouts)	RAC	11	0.03	0.33	0.02	0.22	0.01	0.11	0.36	3.96	0.06	0.66	0.35	3.85
VP 2060	Subgroup of beans with pods (all commodities within this group)	RAC	0.6	0.68	0.41	NC	-	NC	-	0.39	0.23	0.22	0.13	0.49	0.29
VP 0062	Beans without pods: (Phaseolus spp.) (succulent seeds), raw	RAC	0.02	1.56	0.03	0.60	0.01	0.49	0.01	1.18	0.02	0.90	0.02	7.79	0.16
VD 0071	Beans, dry, raw (Phaseolus spp)	RAC	0.03	2.39	0.07	1.61	0.05	10.47	0.31	1.84	0.06	12.90	0.39	7.44	0.22
VD 0523	Broad bean, dry, raw (incl horse-bean, field bean) (Vicia faba)	RAC	0.03	1.27	0.04	0.01	0.00	0.12	0.00	2.49	0.07	0.23	0.01	5.54	0.17
VD 0527	Cowpea, dry, raw (Vigna sinensis, Dolichos sinensis)	RAC	0.03	0.05	0.00	NC	-	1.74	0.05	0.01	0.00	0.01	0.00	0.07	0.00
VD 0541	Soya bean, dry, raw (incl paste, incl curd, incl oil, incl sauce)	RAC	0.085	72.79	6.19	59.05	5.02	20.55	1.75	74.20	6.31	61.12	5.20	73.24	6.23
-	Beans (dry) NES: including inter alia lablab or hyacinth bean (Dolichos spp.); jack or sword bean (Canavalia spp.); winged bean (Psophocarpus tetragonolobus); guar bean (Cyamopsis tetragonoloba); velvet bean (Stizolobium spp.); yam bean (Pachyrhizus erosus)	RAC	0.03	1.70	0.05	0.01	0.00	3.00	0.09	1.80	0.05	1.64	0.05	1.33	0.04
VD 2066	Subgroup of dry peas, raw	RAC	0.054	9.09	0.49	3.35	0.18	1.06	0.06	9.48	0.51	15.11	0.82	10.58	0.57
VR 0577	Carrots, raw	RAC	0.195	9.51	1.85	30.78	6.00	0.37	0.07	8.75	1.71	2.80	0.55	6.10	1.19
VR 0588	Parsnip, raw	RAC	0.09	0.59	0.05	1.05	0.09	0.65	0.06	1.58	0.14	NC	-	0.76	0.07
VR 0494	Radish roots, raw	RAC	0.01	2.31	0.02	4.09	0.04	2.53	0.03	6.15	0.06	5.88	0.06	2.97	0.03
VR 0589	Potato, raw (incl flour, incl frozen, incl starch, incl tapioca)	RAC	0.01	59.74	0.60	316.14	3.16	9.78	0.10	60.26	0.60	54.12	0.54	119.82	1.20
VS 0624	Celery	RAC	8.45	2.14	18.08	3.79	32.03	2.35	19.86	5.69	48.08	0.02	0.17	2.75	23.24
VS 0620	Artichoke globe, raw	RAC	1.2	0.69	0.83	0.01	0.01	0.01	0.01	0.32	0.38	0.26	0.31	1.21	1.45
GC 0654	Wheat, raw (incl bulgur, incl fermented beverages, incl white flour products, incl white bread, excl germ, excl wholemeal bread)	RAC	0.07	381.15	26.68	341.54	23.91	38.34	2.68	281.87	19.73	172.65	12.09	434.06	30.38

Annex 3

CYPRODINIL (207)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
CP 1212	Wheat, wholemeal bread	PP	0.036	0.01	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.03	0.00	0.01	0.00
CF 1211	Wheat, white flour	PP	0.019	299.27	5.69	263.32	5.00	27.93	0.53	214.18	4.07	133.47	2.54	340.03	6.46
GC 0640	Barley, raw (incl malt extract, incl pot&pearled, incl flour & grits, incl malt, excl beer)	RAC	0.58	18.98	11.01	13.35	7.74	0.42	0.24	0.67	0.39	2.30	1.33	0.86	0.50
-	Barley beer	PP	0.0058	4.87	0.03	93.78	0.54	24.28	0.14	12.76	0.07	39.28	0.23	18.15	0.11
TN 0660	Almonds, nutmeat	RAC	0.02	1.38	0.03	0.08	0.00	0.01	0.00	1.00	0.02	0.06	0.00	0.81	0.02
TN 0662	Brazil nuts, nutmeat	RAC	0.01	0.01	0.00	0.01	0.00	0.03	0.00	0.02	0.00	0.02	0.00	0.01	0.00
TN 0295	Cashew nuts, nutmeat	RAC	0.01	0.01	0.00	0.02	0.00	0.24	0.00	0.47	0.00	0.32	0.00	0.05	0.00
TN 0664	Chestnut, raw	RAC	0.01	0.03	0.00	0.02	0.00	0.01	0.00	0.31	0.00	0.09	0.00	0.67	0.01
TN 0665	Coconut, nutmeat (incl. copra, incl desiccated, incl oil)	RAC	0.01	1.73	0.02	1.20	0.01	6.63	0.07	10.18	0.10	13.07	0.13	2.98	0.03
TN 0666	Hazelnuts, nutmeat	RAC	0.01	0.03	0.00	0.13	0.00	0.01	0.00	0.11	0.00	0.01	0.00	1.11	0.01
TN 0669	Macadamia nuts, nutmeat (i.e. Queensland nuts)	RAC	0.01	0.01	0.00	0.01	0.00	0.01	0.00	0.04	0.00	NC	-	0.03	0.00
TN 0672	Pecan, nutmeat	RAC	0.01	0.05	0.00	0.05	0.00	0.02	0.00	0.14	0.00	0.09	0.00	0.13	0.00
TN 0673	Pine nut, nutmeat (i.e. pignolia nuts)	RAC	0.01	0.18	0.00	0.18	0.00	0.08	0.00	0.49	0.00	0.25	0.00	0.43	0.00
TN 0678	Walnut, nutmeat	RAC	0.01	0.23	0.00	1.49	0.01	0.01	0.00	0.33	0.00	0.07	0.00	2.06	0.02
SO 0495	Rape seed, raw (incl oil)	RAC	0.02	0.93	0.02	1.16	0.02	0.49	0.01	2.53	0.05	9.32	0.19	2.02	0.04
HH 0092	Group of Herbs, raw (incl dried)	RAC	5.05	1.69	8.53	1.91	9.65	1.18	5.96	3.35	16.92	0.55	2.78	1.64	8.28
HS 0444	Peppers, chili, dried	PP	2	0.42	0.84	0.53	1.06	0.84	1.68	0.50	1.00	0.95	1.90	0.37	0.74
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat)	RAC	0	31.20	0.00	72.44	0.00	20.88	0.00	47.98	0.00	33.08	0.00	36.25	0.00
MO 0105	Edible offal (mammalian), raw	RAC	0	4.79	0.00	9.68	0.00	2.97	0.00	5.49	0.00	3.84	0.00	5.03	0.00
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0	289.65	0.00	485.88	0.00	26.92	0.00	239.03	0.00	199.91	0.00	180.53	0.00
PM 0110	Poultry meat, raw (incl prepared)	RAC	0	14.63	0.00	29.76	0.00	8.04	0.00	129.68	0.00	25.04	0.00	35.66	0.00
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0	0.12	0.00	0.12	0.00	0.11	0.00	5.37	0.00	0.24	0.00	0.10	0.00
PE 0112	Eggs, raw, (incl dried)	RAC	0	7.84	0.00	23.08	0.00	2.88	0.00	14.89	0.00	9.81	0.00	14.83	0.00
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-
Total intake (ug/person)=				213.2		284.4		124.8		441.5		356.3		465.2	
Bodyweight per region (kg bw) =				60		60		60		60		60		60	

CYPRODINIL (207)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day			Intake as ug/person/day								
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
	ADI (ug/person)=				1800		1800		1800		1800		1800		1800
	%ADI=				11.8%		15.8%		6.9%		24.5%		19.8%		25.8%
	Rounded %ADI=				10%		20%		7%		20%		20%		30%

CYPRODINIL (207)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
FP 0009	Group of Pome fruits, raw (incl apple cider, excl apple juice)	RAC	0.48	51.09	24.52	65.40	31.39	42.71	20.50	45.29	21.74	62.51	30.00	7.74	3.72
JF 0226	Apple juice, single strength (incl. concentrated)	PP	0.015	14.88	0.22	11.98	0.18	0.15	0.00	9.98	0.15	30.32	0.45	3.47	0.05
FS 0012	Group of Stone fruits, raw (incl dried apricots, excl dried plums)	RAC	0.68	18.18	12.36	23.83	16.20	14.27	9.70	18.52	12.59	9.35	6.36	0.11	0.07
DF 0014	Plums, dried (prunes)	PP	1.2	0.61	0.73	0.35	0.42	0.05	0.06	0.35	0.42	0.49	0.59	0.13	0.16
FB 2005	Subgroup of Caneberries, raw	RAC	2.2	0.56	1.23	1.43	3.15	0.14	0.31	1.23	2.71	1.14	2.51	0.01	0.02
FB 2006	Subgroup of Bush berries, raw (including processed)	RAC	2.2	1.31	2.88	5.50	12.10	0.01	0.02	2.57	5.65	0.82	1.80	2.15	4.73
FB 2007	Subgroup of Large shrub/tree berries, raw (including processed)	RAC	2.2	8.26	18.17	0.14	0.31	0.07	0.15	0.13	0.29	0.19	0.42	1.87	4.11
FB 0269	Grapes, raw (incl must, excl dried, excl juice, excl wine)	RAC	0.79	6.48	5.12	11.31	8.93	5.21	4.12	9.50	7.51	4.66	3.68	0.78	0.62
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	1.7	3.09	5.25	1.51	2.57	0.03	0.05	1.38	2.35	4.26	7.24	0.42	0.71
JF 0269	Grape juice (from wine grapes)	PP	0.12	0.56	0.07	1.96	0.24	0.02	0.00	2.24	0.27	2.27	0.27	0.34	0.04
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.062	88.93	5.51	62.41	3.87	1.84	0.11	25.07	1.55	61.17	3.79	5.84	0.36
FB 2009	Subgroup of Low growing berries, raw	RAC	2.2	4.55	10.01	5.66	12.45	0.02	0.04	7.85	17.27	5.86	12.89	0.05	0.11
FT 0336	Guava, raw	RAC	0.485	0.01	0.00	NC	-	0.42	0.20	NC	-	NC	-	NC	-
FI 0326	Avocado, raw	RAC	0.265	2.65	0.70	0.87	0.23	0.46	0.12	1.64	0.43	1.30	0.34	0.96	0.25
FI 0355	Pomegranate, raw, (incl processed)	RAC	3.3	7.91	26.10	9.72	32.08	7.67	25.31	5.26	17.36	9.04	29.83	14.43	47.62
-	Onions, dry, raw	RAC	0.065	19.69	1.28	29.83	1.94	24.64	1.60	31.35	2.04	9.72	0.63	12.59	0.82

Annex 3

CYPRODINIL (207)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
VA 2032	Subgroup of Green Onions	RAC	0.065	5.57	0.36	5.15	0.33	1.77	0.12	4.28	0.28	17.34	1.13	6.48	0.42
VB 0042	Subgroup of Flowerhead Brassica, raw	RAC	0.27	9.50	2.57	6.77	1.83	NC	-	3.21	0.87	9.36	2.53	0.75	0.20
VB 0041	Cabbages, head, raw	RAC	0.03	8.97	0.27	27.12	0.81	1.44	0.04	24.96	0.75	4.55	0.14	11.23	0.34
VC 0045	Group of Fruiting vegetables, cucurbits, raw	RAC	0.09	27.81	2.50	41.93	3.77	123.30	11.10	49.47	4.45	15.95	1.44	35.99	3.24
VO 0448	Tomato, raw (incl canned, excl juice, excl paste)	RAC	0.24	43.88	10.53	55.41	13.30	35.38	8.49	74.88	17.97	26.50	6.36	9.51	2.28
-	Tomato, paste (i.e. concentrated tomato sauce/puree)	PP	0.48	4.96	2.38	3.20	1.54	0.15	0.07	1.61	0.77	6.88	3.30	0.52	0.25
JF 0448	Tomato, juice (single strength, incl concentrated)	PP	0.036	0.80	0.03	0.07	0.00	0.05	0.00	0.61	0.02	0.40	0.01	0.08	0.00
VO 0051	Subgroup of peppers, raw (incl dried sweet peppers, excl dried chilipeppers), incl okra	RAC	0.24	6.39	1.53	15.53	3.73	19.13	4.59	10.53	2.53	8.29	1.99	5.25	1.26
VO 2046	Subgroup of eggplants	RAC	0.24	1.01	0.24	1.69	0.41	21.37	5.13	3.00	0.72	1.40	0.34	NC	-
VL 2050	Subgroup of Leafy greens	RAC	11	18.38	202.18	18.73	206.03	82.36	905.96	25.32	278.52	17.60	193.60	7.37	81.07
VL 0054	Subgroup of Leaves of Brassicaceae, raw	RAC	0.37	0.10	0.04	NC	-	26.78	9.91	5.00	1.85	0.58	0.21	5.68	2.10
VL 2052	Subgroup of Leaves of Root and Tuber Vegetables	RAC	11	NC	-	NC	-	NC	-	NC	-	NC	-	0.33	3.63
VL 2053	Subgroup of Leaves of trees, shrubs and vines	RAC	11	NC	-	NC	-	NC	-	NC	-	NC	-	0.74	8.14
VL 2054	Subgroup of Leafy aquatic vegetables	RAC	11	0.35	3.85	3.13	34.43	3.74	41.14	1.79	19.69	NC	-	2.30	25.30
VL 2832	Witloof chicory (sprouts)	RAC	11	1.50	16.50	0.95	10.45	NC	-	1.84	20.24	0.65	7.15	0.13	1.43
VP 2060	Subgroup of beans with pods (all commodities within this group)	RAC	0.6	5.07	3.04	0.83	0.50	0.17	0.10	3.70	2.22	NC	-	NC	-
VP 0062	Beans without pods: (Phaseolus spp.) (succulent seeds), raw	RAC	0.02	2.21	0.04	5.25	0.11	4.17	0.08	1.61	0.03	16.95	0.34	0.17	0.00
VD 0071	Beans, dry, raw (Phaseolus spp)	RAC	0.03	1.51	0.05	1.50	0.05	1.90	0.06	5.11	0.15	1.36	0.04	23.43	0.70
VD 0523	Broad bean, dry, raw (incl horse-bean, field bean) (Vicia faba)	RAC	0.03	0.02	0.00	0.01	0.00	1.16	0.03	0.40	0.01	NC	-	0.06	0.00
VD 0527	Cowpea, dry, raw (Vigna sinensis, Dolichos sinensis)	RAC	0.03	NC	-	NC	-	0.16	0.00	0.01	0.00	NC	-	NC	-
VD 0541	Soya bean, dry, raw (incl paste, incl curd, incl oil, incl sauce)	RAC	0.085	106.33	9.04	117.78	10.01	42.12	3.58	195.70	16.63	222.52	18.91	80.47	6.84
-	Beans (dry) NES: including inter alia lablab or hyacinth bean (Dolichos spp.); jack or sword bean (Canavalia spp.); winged bean (Psophocarpus tetragonolobus); guar bean (Cyamopsis tetragonoloba); velvet bean (Stizolobium spp.); yam bean (Pachyrrhizus erosus)	RAC	0.03	0.01	0.00	NC	-	0.57	0.02	0.11	0.00	0.16	0.00	0.94	0.03

CYPRODINIL (207)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
VD 2066	Subgroup of dry peas, raw	RAC	0.054	5.01	0.27	3.76	0.20	1.82	0.10	3.44	0.19	3.49	0.19	5.15	0.28
VR 0577	Carrots, raw	RAC	0.195	26.26	5.12	27.13	5.29	10.07	1.96	16.49	3.22	44.69	8.71	8.75	1.71
VR 0588	Parsnip, raw	RAC	0.09	4.42	0.40	0.06	0.01	NC	-	NC	-	NC	-	1.12	0.10
VR 0494	Radish roots, raw	RAC	0.01	3.83	0.04	11.99	0.12	NC	-	5.26	0.05	2.19	0.02	4.37	0.04
VR 0589	Potato, raw (incl flour, incl frozen, incl starch, incl tapioca)	RAC	0.01	225.03	2.25	234.24	2.34	71.48	0.71	177.55	1.78	234.55	2.35	37.71	0.38
VS 0624	Celery	RAC	8.45	7.68	64.90	2.85	24.08	NC	-	3.34	28.22	16.83	142.21	4.04	34.14
VS 0620	Artichoke globe, raw	RAC	1.2	0.98	1.18	3.65	4.38	0.07	0.08	1.67	2.00	0.26	0.31	NC	-
GC 0654	Wheat, raw (incl bulgur, incl fermented beverages, incl white flour products, incl white bread, excl germ, excl wholemeal bread)	RAC	0.07	252.06	17.64	244.62	17.12	134.41	9.41	235.10	16.46	216.33	15.14	167.34	11.71
CP 1212	Wheat, wholemeal bread	PP	0.036	0.03	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.05	0.00	0.02	0.00
CF 1211	Wheat, white flour	PP	0.019	182.77	3.47	187.54	3.56	103.82	1.97	180.42	3.43	164.00	3.12	118.84	2.26
GC 0640	Barley, raw (incl malt extract, incl pot&pearled, incl flour & grits, incl malt, excl beer)	RAC	0.58	1.94	1.13	4.15	2.41	0.66	0.38	2.50	1.45	2.14	1.24	3.52	2.04
-	Barley beer	PP	0.0058	180.21	1.05	259.46	1.50	45.91	0.27	172.36	1.00	234.42	1.36	65.30	0.38
TN 0660	Almonds, nutmeat	RAC	0.02	0.81	0.02	2.21	0.04	0.03	0.00	1.02	0.02	1.47	0.03	NC	-
TN 0662	Brazil nuts, nutmeat	RAC	0.01	0.12	0.00	0.05	0.00	0.01	0.00	0.05	0.00	0.13	0.00	NC	-
TN 0295	Cashew nuts, nutmeat	RAC	0.01	0.59	0.01	0.23	0.00	0.18	0.00	0.52	0.01	1.75	0.02	2.78	0.03
TN 0664	Chestnut, raw	RAC	0.01	0.34	0.00	0.21	0.00	1.14	0.01	0.52	0.01	0.09	0.00	NC	-
TN 0665	Coconut, nutmeat (incl. copra, incl desiccated, incl oil)	RAC	0.01	4.13	0.04	2.73	0.03	13.15	0.13	5.85	0.06	6.92	0.07	22.24	0.22
TN 0666	Hazelnuts, nutmeat	RAC	0.01	0.45	0.00	1.12	0.01	0.02	0.00	0.34	0.00	1.63	0.02	NC	-
TN 0669	Macadamia nuts, nutmeat (i.e. Queensland nuts)	RAC	0.01	NC	-	0.40	0.00	NC	-	NC	-	NC	-	0.07	0.00
TN 0672	Pecan, nutmeat	RAC	0.01	0.38	0.00	NC	-	NC	-	0.27	0.00	NC	-	0.26	0.00
TN 0673	Pine nut, nutmeat (i.e. pignolia nuts)	RAC	0.01	0.99	0.01	0.66	0.01	0.22	0.00	0.27	0.00	1.89	0.02	0.89	0.01
TN 0678	Walnut, nutmeat	RAC	0.01	0.34	0.00	0.84	0.01	0.28	0.00	0.39	0.00	0.45	0.00	NC	-
SO 0495	Rape seed, raw (incl oil)	RAC	0.02	32.68	0.65	19.91	0.40	7.83	0.16	15.69	0.31	NC	-	NC	-
HH 0092	Group of Herbs, raw (incl dried)	RAC	5.05	2.61	13.18	2.31	11.67	8.89	44.89	3.92	19.80	1.16	5.86	2.06	10.40
HS 0444	Peppers, chili, dried	PP	2	0.11	0.22	0.21	0.42	0.36	0.72	0.21	0.42	0.25	0.50	0.15	0.30

Annex 3

CYPRODINIL (207)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat)	RAC	0	140.03	0.00	150.89	0.00	79.32	0.00	111.24	0.00	120.30	0.00	51.27	0.00
MO 0105	Edible offal (mammalian), raw	RAC	0	15.17	0.00	5.19	0.00	6.30	0.00	6.78	0.00	3.32	0.00	3.17	0.00
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0	388.92	0.00	335.88	0.00	49.15	0.00	331.25	0.00	468.56	0.00	245.45	0.00
PM 0110	Poultry meat, raw (incl prepared)	RAC	0	73.76	0.00	53.86	0.00	23.98	0.00	87.12	0.00	53.38	0.00	84.45	0.00
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0	0.33	0.00	0.72	0.00	0.27	0.00	0.35	0.00	0.80	0.00	NC	-
PE 0112	Eggs, raw, (incl dried)	RAC	0	25.84	0.00	29.53	0.00	28.05	0.00	33.19	0.00	36.44	0.00	8.89	0.00
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-
Total intake (ug/person)=				480.9		487.0		1113.6		538.5		519.5		264.6	
Bodyweight per region (kg bw) =				60		60		55		60		60		60	
ADI (ug/person)=				1800		1800		1650		1800		1800		1800	
%ADI=				26.7%		27.1%		67.5%		29.9%		28.9%		14.7%	
Rounded %ADI=				30%		30%		70%		30%		30%		10%	

CYPRODINIL (207)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
FP 0009	Group of Pome fruits, raw (incl apple cider, excl apple juice)	RAC	0.48	68.85	33.05	10.93	5.25	70.82	33.99	189.78	91.09	19.56	9.39
JF 0226	Apple juice, single strength (incl. concentrated)	PP	0.015	0.03	0.00	0.10	0.00	7.19	0.11	0.03	0.00	NC	-
FS 0012	Group of Stone fruits, raw (incl dried apricots, excl dried plums)	RAC	0.68	0.09	0.06	0.02	0.01	32.27	21.94	0.01	0.01	NC	-
DF 0014	Plums, dried (prunes)	PP	1.2	0.01	0.01	0.01	0.01	0.37	0.44	0.01	0.01	NC	-
FB 2005	Subgroup of Caneberries, raw	RAC	2.2	0.01	0.02	7.30	16.06	2.29	5.04	0.01	0.02	NC	-
FB 2006	Subgroup of Bush berries, raw (including processed)	RAC	2.2	0.82	1.80	4.05	8.91	5.94	13.07	0.43	0.95	2.66	5.85
FB 2007	Subgroup of Large shrub/tree berries, raw (including processed)	RAC	2.2	0.71	1.56	7.32	16.10	NC	-	0.38	0.84	2.32	5.10
FB 0269	Grapes, raw (incl must, excl dried, excl juice, excl wine)	RAC	0.79	0.14	0.11	0.36	0.28	15.33	12.11	0.01	0.01	0.28	0.22
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	1.7	0.01	0.02	0.13	0.22	1.06	1.80	0.01	0.02	0.03	0.05
JF 0269	Grape juice (from wine grapes)	PP	0.12	0.01	0.00	0.01	0.00	0.41	0.05	0.01	0.00	NC	-
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.062	0.31	0.02	0.23	0.01	60.43	3.75	0.52	0.03	31.91	1.98
FB 2009	Subgroup of Low growing berries, raw	RAC	2.2	0.01	0.02	0.01	0.02	3.37	7.41	0.01	0.02	0.01	0.02
FT 0336	Guava, raw	RAC	0.485	0.10	0.05	0.08	0.04	NC	-	0.14	0.07	3.11	1.51
FI 0326	Avocado, raw	RAC	0.265	1.12	0.30	0.01	0.00	0.84	0.22	0.01	0.00	6.60	1.75
FI 0355	Pomegranate, raw, (incl processed)	RAC	3.3	5.49	18.12	27.17	89.66	NC	-	2.89	9.54	17.87	58.97
-	Onions, dry, raw	RAC	0.065	9.01	0.59	20.24	1.32	30.90	2.01	9.61	0.62	2.11	0.14
VA 2032	Subgroup of Green Onions	RAC	0.065	1.45	0.09	1.50	0.10	1.42	0.09	0.01	0.00	6.30	0.41
VB 0042	Subgroup of Flowerhead Brassica, raw	RAC	0.27	0.02	0.01	0.02	0.01	4.86	1.31	0.01	0.00	NC	-
VB 0041	Cabbages, head, raw	RAC	0.03	3.82	0.11	2.99	0.09	49.16	1.47	0.01	0.00	NC	-
VC 0045	Group of Fruiting vegetables, cucurbits, raw	RAC	0.09	5.96	0.54	9.74	0.88	51.82	4.66	13.61	1.22	0.05	0.00
VO 0448	Tomato, raw (incl canned, excl juice, excl paste)	RAC	0.24	13.10	3.14	4.90	1.18	62.16	14.92	1.04	0.25	0.09	0.02
-	Tomato, paste (i.e. concentrated tomato sauce/puree)	PP	0.48	0.58	0.28	0.22	0.11	2.21	1.06	0.24	0.12	3.10	1.49
JF 0448	Tomato, juice (single strength, incl concentrated)	PP	0.036	0.05	0.00	0.01	0.00	0.42	0.02	0.01	0.00	0.02	0.00
VO 0051	Subgroup of peppers, raw (incl dried sweet peppers, excl dried chilipeppers), incl okra	RAC	0.24	15.20	3.65	14.23	3.42	25.14	6.03	0.91	0.22	NC	-

Annex 3

CYPRODINIL (207)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day			Intake = daily intake: ug/person						
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
VO 2046	Subgroup of eggplants	RAC	0.24	1.31	0.31	8.26	1.98	3.95	0.95	0.01	0.00	NC	-
VL 2050	Subgroup of Leafy greens	RAC	11	4.99	54.89	3.29	36.19	7.53	82.83	3.05	33.55	6.09	66.99
VL 0054	Subgroup of Leaves of Brassicaceae, raw	RAC	0.37	3.58	1.32	2.64	0.98	NC	-	1.83	0.68	3.65	1.35
VL 2052	Subgroup of Leaves of Root and Tuber Vegetables	RAC	11	0.30	3.30	0.22	2.42	NC	-	0.20	2.20	0.41	4.51
VL 2053	Subgroup of Leaves of trees, shrubs and vines	RAC	11	0.67	7.37	0.48	5.28	NC	-	0.45	4.95	0.90	9.90
VL 2054	Subgroup of Leafy aquatic vegetables	RAC	11	2.08	22.88	1.50	16.50	0.01	0.11	1.41	15.51	2.81	30.91
VL 2832	Witloof chicory (sprouts)	RAC	11	0.02	0.22	0.01	0.11	0.01	0.11	0.01	0.11	NC	-
VP 2060	Subgroup of beans with pods (all commodities within this group)	RAC	0.6	NC	-	NC	-	NC	-	NC	-	NC	-
VP 0062	Beans without pods: (Phaseolus spp.) (succulent seeds), raw	RAC	0.02	0.30	0.01	3.13	0.06	4.11	0.08	0.01	0.00	NC	-
VD 0071	Beans, dry, raw (Phaseolus spp)	RAC	0.03	7.11	0.21	2.33	0.07	3.76	0.11	44.70	1.34	3.27	0.10
VD 0523	Broad bean, dry, raw (incl horse-bean, field bean) (Vicia faba)	RAC	0.03	3.70	0.11	0.03	0.00	0.17	0.01	0.01	0.00	NC	-
VD 0527	Cowpea, dry, raw (Vigna sinensis, Dolichos sinensis)	RAC	0.03	12.77	0.38	0.99	0.03	0.01	0.00	4.33	0.13	NC	-
VD 0541	Soya bean, dry, raw (incl paste, incl curd, incl oil, incl sauce)	RAC	0.085	15.80	1.34	14.29	1.21	104.36	8.87	17.11	1.45	35.20	2.99
-	Beans (dry) NES: including inter alia lablab or hyacinth bean (Dolichos spp.); jack or sword bean (Canavalia spp.); winged bean (Psophocarpus tetragonolobus); guar bean (Cyamopsis tetragonoloba); velvet bean (Stizolobium spp.); yam bean (Pachyrrhizus erosus)	RAC	0.03	2.54	0.08	1.77	0.05	0.03	0.00	0.03	0.00	3.99	0.12
VD 2066	Subgroup of dry peas, raw	RAC	0.054	4.43	0.24	11.36	0.61	4.22	0.23	9.36	0.51	1.21	0.07
VR 0577	Carrots, raw	RAC	0.195	2.07	0.40	3.00	0.59	25.29	4.93	0.05	0.01	NC	-
VR 0588	Parsnip, raw	RAC	0.09	1.02	0.09	0.74	0.07	3.50	0.32	0.69	0.06	1.37	0.12
VR 0494	Radish roots, raw	RAC	0.01	3.96	0.04	2.86	0.03	3.30	0.03	2.67	0.03	5.34	0.05
VR 0589	Potato, raw (incl flour, incl frozen, incl starch, incl tapioca)	RAC	0.01	23.96	0.24	13.56	0.14	213.41	2.13	104.35	1.04	8.56	0.09
VS 0624	Celery	RAC	8.45	3.66	30.93	2.65	22.39	4.84	40.90	2.47	20.87	4.94	41.74
VS 0620	Artichoke globe, raw	RAC	1.2	0.01	0.01	NC	-	0.08	0.10	0.01	0.01	NC	-
GC 0654	Wheat, raw (incl bulgur, incl fermented beverages, incl white flour products, incl white bread, excl germ, excl wholemeal bread)	RAC	0.07	57.15	4.00	110.46	7.73	272.58	19.08	25.81	1.81	132.04	9.24

Total intake (ug/person)=	208.3	254.2	316.7	196.3	277.9
Bodyweight per region (kg bw) =	60	60	60	60	60

Annex 3

CYPRODINIL (207)		International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw							
Codex Code	Commodity description	Expr as	STM R mg/kg	Diets: g/person/day			Intake = daily intake: ug/person						
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
	ADI (ug/person)=				1800		1800		1800		1800		1800
	%ADI=				11.6%		14.1%		17.6%		10.9%		15.4%
	Rounded %ADI=				10%		10%		20%		10%		20%

DIFENOCONAZOLE (224)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.01 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
FC 0001	Group of Citrus fruit, raw (incl kumquat commodities)	RAC	0.16	32.25	5.16	11.67	1.87	16.70	2.67	76.01	12.16	33.90	5.42	92.97	14.88
JF 0001	Group of Citrus fruit, juice	PP	0.002	1.30	0.00	2.37	0.00	0.22	0.00	13.88	0.03	0.75	0.00	2.63	0.01
FP 0009	Group of Pome fruits, raw (incl. apple juice, incl apple cider)	RAC	0.86	19.79	17.02	38.25	32.90	17.96	15.45	32.56	28.00	8.08	6.95	64.45	55.43
FS 0013	Subgroup of Cherries, raw	RAC	0.04	0.92	0.04	9.15	0.37	0.01	0.00	0.61	0.02	0.06	0.00	6.64	0.27
FS 0014	Subgroup of Plums, raw (incl dried plums)	RAC	0.04	2.67	0.11	8.77	0.35	0.07	0.00	3.03	0.12	0.70	0.03	4.34	0.17
FS 2001	Subgroup of peaches, raw (incl dried apricots)	RAC	0.15	8.01	1.20	5.87	0.88	0.18	0.03	8.19	1.23	1.64	0.25	22.46	3.37
FB 0020	Blueberries, raw	RAC	1	0.01	0.01	0.03	0.03	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
FB 0269	Grapes, raw (incl must, excl dried, excl juice, excl wine)	RAC	0.52	13.02	6.77	9.25	4.81	0.03	0.02	16.91	8.79	3.70	1.92	54.44	28.31
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	1.1	0.51	0.56	0.51	0.56	0.01	0.01	1.27	1.40	0.12	0.13	2.07	2.28
JF 0269	Grape juice (from wine grapes)	PP	0.24	0.14	0.03	0.29	0.07	0.05	0.01	0.30	0.07	0.24	0.06	0.05	0.01
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.094	0.67	0.06	12.53	1.18	2.01	0.19	1.21	0.11	3.53	0.33	4.01	0.38
FB 0265	Cranberry, raw	RAC	0.2	0.02	0.00	0.01	0.00	NC	-	0.03	0.01	0.01	0.00	0.01	0.00
FB 0275	Strawberry, raw	RAC	0.42	0.70	0.29	2.01	0.84	0.04	0.02	1.36	0.57	0.37	0.16	2.53	1.06
FT 0305	Table olives, raw (incl preserved)	RAC	0.465	0.70	0.33	0.32	0.15	0.01	0.00	1.53	0.71	0.17	0.08	1.85	0.86
FT 0336	Guava, raw	RAC	0.0335	0.47	0.02	0.01	0.00	0.48	0.02	0.49	0.02	4.42	0.15	0.06	0.00
FI 0326	Avocado, raw	RAC	0.05	0.13	0.01	0.03	0.00	2.05	0.10	2.54	0.13	2.34	0.12	0.12	0.01
FI 0327	Banana, raw (incl plantains) (incl dried)	RAC	0.02	5.23	0.10	6.94	0.14	99.45	1.99	32.47	0.65	48.30	0.97	24.70	0.49
FI 0345	Mango, raw (incl canned mango, incl mango juice)	RAC	0.03	10.48	0.31	0.01	0.00	7.24	0.22	6.87	0.21	19.98	0.60	6.25	0.19
FI 0350	Papaya, raw	RAC	0.065	0.35	0.02	0.01	0.00	3.05	0.20	0.80	0.05	7.28	0.47	1.00	0.07
FI 2540	Pitaya, raw (i.e dragon fruit or pitahaya)	RAC	0.034	0.01	0.00	0.01	0.00	0.01	0.00	0.01	0.00	NC	-	0.01	0.00
FI 0351	Passion fruit, raw	RAC	0.01	0.58	0.01	0.01	0.00	0.59	0.01	0.60	0.01	0.18	0.00	0.08	0.00
-	Onions, dry, raw	RAC	0.015	29.36	0.44	37.50	0.56	3.56	0.05	34.78	0.52	18.81	0.28	43.38	0.65
VA 0384	Leek, raw	RAC	0.08	0.18	0.01	1.59	0.13	0.03	0.00	0.28	0.02	0.01	0.00	3.21	0.26
-	Onions, green, raw	RAC	2.8	2.45	6.86	1.49	4.17	1.02	2.86	2.60	7.28	0.60	1.68	2.03	5.68
VB 0040	Group of Brassica vegetables (excl Brassica leafy vegetables), raw	RAC	0.35	6.43	2.25	40.26	14.09	0.80	0.28	9.94	3.48	12.07	4.22	17.73	6.21

Annex 3

DIFENOCONAZOLE (224)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.01 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
VC 0424	Cucumber, raw	RAC	0.04	8.01	0.32	30.66	1.23	1.45	0.06	19.84	0.79	0.27	0.01	34.92	1.40
VC 0425	Gherkin, raw	RAC	0.04	1.73	0.07	6.64	0.27	0.31	0.01	4.29	0.17	0.29	0.01	7.56	0.30
VC 0431	Squash, Summer (Courgette, Marrow, Zucchini, Zucchini), raw	RAC	0.04	0.78	0.03	2.06	0.08	0.30	0.01	1.61	0.06	2.25	0.09	2.36	0.09
VC 0046	Melons, except watermelon, raw (Cantaloupe)	RAC	0.14	8.90	1.25	8.64	1.21	0.80	0.11	17.90	2.51	2.80	0.39	29.17	4.08
VC 0432	Watermelon, raw	RAC	0.01	28.96	0.29	25.65	0.26	1.56	0.02	39.26	0.39	4.94	0.05	66.90	0.67
VO 0448	Tomato, raw	RAC	0.1	41.73	4.17	75.65	7.57	10.66	1.07	82.87	8.29	24.75	2.48	200.93	20.09
-	Tomato, canned (& peeled)	PP	0.01	0.20	0.00	0.31	0.00	0.02	0.00	1.11	0.01	0.11	0.00	1.50	0.02
-	Tomato, paste (i.e. concentrated tomato sauce/puree)	PP	0.22	2.34	0.51	1.33	0.29	1.57	0.35	4.24	0.93	0.34	0.07	2.83	0.62
JF 0448	Tomato, juice (single strength, incl concentrated)	PP	0.031	0.29	0.01	0.29	0.01	0.01	0.00	0.38	0.01	0.05	0.00	0.14	0.00
VO 0442	Okra, raw (i.e. Lady's Finger, Gombo)	RAC	0.14	1.97	0.28	NC	-	3.68	0.52	3.24	0.45	5.72	0.80	1.57	0.22
VO 0444	Peppers, chili, raw	RAC	0.24	3.99	0.96	7.30	1.75	2.93	0.70	5.62	1.35	NC	-	17.44	4.19
VO 0445	Peppers, sweet, raw (incl dried)	RAC	0.14	4.49	0.63	6.44	0.90	7.21	1.01	5.68	0.80	9.52	1.33	8.92	1.25
VO 0440	Egg plant, raw (i.e. aubergine)	RAC	0.14	5.58	0.78	4.31	0.60	0.89	0.12	9.31	1.30	13.64	1.91	20.12	2.82
VL 0483	Lettuce, leaf, raw	RAC	0.41	0.53	0.22	0.36	0.15	0.16	0.07	6.21	2.55	1.90	0.78	6.05	2.48
VP 0060	Group of Legume vegetables, raw	RAC	0.07	7.73	0.54	1.53	0.11	0.51	0.04	2.95	0.21	5.08	0.36	12.86	0.90
VD 0071	Beans, dry, raw (Phaseolus spp)	RAC	0.011	2.39	0.03	1.61	0.02	10.47	0.12	1.84	0.02	12.90	0.14	7.44	0.08
VD 0523	Broad bean, dry, raw (incl horse-bean, field bean) (Vicia faba)	RAC	0.011	1.27	0.01	0.01	0.00	0.12	0.00	2.49	0.03	0.23	0.00	5.54	0.06
VD 0541	Soya bean, dry, raw (incl flour, incl paste, incl curd, incl sauce, excl oil)	RAC	0.01	0.63	0.01	1.09	0.01	0.40	0.00	1.40	0.01	1.68	0.02	0.48	0.00
OR 0541	Soya oil, refined	PP	0.08	12.99	1.04	10.43	0.83	3.63	0.29	13.10	1.05	10.70	0.86	13.10	1.05
-	Beans (dry) NES: including inter alia lablab or hyacinth bean (Dolichos spp.); jack or sword bean (Canavalia spp.); winged bean (Psophocarpus tetragonolobus); guar bean (Cyamopsis tetragonoloba); velvet bean (Stizolobium spp.); yam bean (Pachyrhizus erosus)	RAC	0.028	1.70	0.05	0.01	0.00	3.00	0.08	1.80	0.05	1.64	0.05	1.33	0.04
VD 0533	Lentil (dry) (Lens spp), raw	RAC	0.028	2.12	0.06	0.01	0.00	0.03	0.00	3.21	0.09	1.60	0.04	4.90	0.14
VD 0537	Pigeon pea (dry) (Cajanus spp), raw	RAC	0.028	NC	-	NC	-	0.10	0.00	0.07	0.00	3.38	0.09	NC	-

DIFENOCONAZOLE (224)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.01 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
VR 0577	Carrots, raw	RAC	0.05	9.51	0.48	30.78	1.54	0.37	0.02	8.75	0.44	2.80	0.14	6.10	0.31
VR 0578	Celeriac, raw	RAC	0.12	1.70	0.20	3.01	0.36	1.87	0.22	4.53	0.54	NC	-	2.19	0.26
VR 0596	Sugar beet, raw (incl sugar)	RAC	0.02	0.13	0.00	NC	-	0.08	0.00	0.66	0.01	0.47	0.01	88.94	1.78
VR 0589	Potato, raw (incl flour, incl frozen, incl starch, incl tapioca)	RAC	1.2	59.74	71.69	316.14	379.37	9.78	11.74	60.26	72.31	54.12	64.94	119.82	143.78
VS 0624	Celery	RAC	0.14	2.14	0.30	3.79	0.53	2.35	0.33	5.69	0.80	0.02	0.00	2.75	0.39
VS 0621	Asparagus, raw	RAC	0.02	0.01	0.00	0.01	0.00	0.01	0.00	0.03	0.00	0.07	0.00	0.21	0.00
VS 0620	Artichoke globe, raw	RAC	0.51	0.69	0.35	0.01	0.01	0.01	0.01	0.32	0.16	0.26	0.13	1.21	0.62
CM 0649 (GC 0649)	Rice, husked, dry (incl flour, incl oil, incl beverages, incl starch, excl polished)	REP	1.1	1.26	1.39	1.58	1.74	31.05	34.16	5.43	5.97	0.90	0.99	2.18	2.40
CM 1205	Rice polished, dry	PP	0.0086	34.21	0.29	10.39	0.09	41.72	0.36	82.38	0.71	150.24	1.29	70.47	0.61
GC 0447	Sweet corn on the cob, raw (incl frozen kernels, incl canned kernels)	RAC	0.01	0.14	0.00	0.94	0.01	5.70	0.06	2.61	0.03	1.94	0.02	0.22	0.00
TN 0085	Group of Tree nuts, raw (incl processed)	RAC	0.01	4.06	0.04	3.27	0.03	7.01	0.07	13.93	0.14	14.01	0.14	9.36	0.09
SO 0495	Rape seed, raw	RAC	0.03	0.02	0.00	NC	-	NC	-	0.01	0.00	0.75	0.02	0.01	0.00
OR 0495	Rape seed oil, edible	PP	0.002	0.35	0.00	0.44	0.00	0.19	0.00	0.97	0.00	3.28	0.01	0.77	0.00
SO 0702	Sunflower seed, raw (incl oil)	RAC	0.01	7.40	0.07	35.86	0.36	1.15	0.01	8.76	0.09	5.45	0.05	13.62	0.14
OR 0691	Cotton seed oil, edible	PP	0.0014	3.22	0.00	1.54	0.00	1.01	0.00	0.74	0.00	1.12	0.00	2.93	0.00
SO 0697	Peanuts, nutmeat, raw (incl roasted, incl oil, incl butter)	RAC	0	1.30	0.00	1.23	0.00	12.62	0.00	2.87	0.00	6.59	0.00	2.67	0.00
SO 0305	Olives for oil production, raw	RAC	0.465	1.47	0.68	0.67	0.31	NC	-	1.26	0.59	0.04	0.02	7.63	3.55
-	Olive oil (virgin and residue oil)	PP	0.7	2.17	1.52	0.13	0.09	0.05	0.04	1.32	0.92	0.10	0.07	2.76	1.93
SB 0716	Coffee bean raw (incl roasted, incl instant coffee, incl substitutes)	RAC	0.01	1.36	0.01	3.59	0.04	1.44	0.01	5.18	0.05	2.02	0.02	1.70	0.02
HS 0444	Peppers, chili, dried	PP	1.1	0.42	0.46	0.53	0.58	0.84	0.92	0.50	0.55	0.95	1.05	0.37	0.41
DT 1114	Tea, green or black, fermented and dried, (including concentrates)	RAC	4.85	2.28	11.06	1.98	9.60	0.46	2.23	2.43	11.79	1.29	6.26	3.04	14.74
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.047	24.96	1.17	57.95	2.72	16.70	0.79	38.38	1.80	26.46	1.24	29.00	1.36
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.14	6.24	0.87	14.49	2.03	4.18	0.58	9.60	1.34	6.62	0.93	7.25	1.02
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.14	3.29	0.46	6.14	0.86	0.82	0.11	1.57	0.22	2.23	0.31	1.07	0.15

Annex 3

DIFENOCONAZOLE (224)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.01 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
MO 0105	Edible offal (mammalian), raw	RAC	0.71	4.79	3.40	9.68	6.87	2.97	2.11	5.49	3.90	3.84	2.73	5.03	3.57
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.011	289.65	3.19	485.88	5.34	26.92	0.30	239.03	2.63	199.91	2.20	180.53	1.99
PM 0110	Poultry meat, raw (incl prepared)	RAC	0.0002	14.63	0.00	29.76	0.01	8.04	0.00	129.68	0.03	25.04	0.01	35.66	0.01
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.0002	0.10	0.00	0.10	0.00	NC	-	0.10	0.00	0.10	0.00	0.10	0.00
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.0002	0.12	0.00	0.12	0.00	0.11	0.00	5.37	0.00	0.24	0.00	0.10	0.00
PE 0112	Eggs, raw, (incl dried)	RAC	0.011	7.84	0.09	23.08	0.25	2.88	0.03	14.89	0.16	9.81	0.11	14.83	0.16
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

Total intake (ug/person)=	150.6	491.1	82.8	191.9	116.0	340.4
Bodyweight per region (kg bw) =	60	60	60	60	60	60
ADI (ug/person)=	600	600	600	600	600	600
%ADI=	25.1%	81.9%	13.8%	32.0%	19.3%	56.7%
Rounded %ADI=	30%	80%	10%	30%	20%	60%

DIFENOCONAZOLE (224)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.01 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
FC 0001	Group of Citrus fruit, raw (incl kumquat commodities)	RAC	0.16	38.66	6.19	54.93	8.79	26.36	4.22	51.46	8.23	51.06	8.17	466.36	74.62
JF 0001	Group of Citrus fruit, juice	PP	0.002	36.84	0.07	3.75	0.01	0.30	0.00	21.62	0.04	21.82	0.04	46.67	0.09
FP 0009	Group of Pome fruits, raw (incl. apple juice, incl apple cider)	RAC	0.86	71.38	61.39	81.73	70.29	42.91	36.90	58.89	50.65	103.85	89.31	12.48	10.73
FS 0013	Subgroup of Cherries, raw	RAC	0.04	1.40	0.06	4.21	0.17	0.04	0.00	2.93	0.12	1.50	0.06	NC	-
FS 0014	Subgroup of Plums, raw (incl dried plums)	RAC	0.04	5.55	0.22	4.37	0.17	6.08	0.24	3.66	0.15	3.93	0.16	0.46	0.02
FS 2001	Subgroup of peaches, raw (incl dried apricots)	RAC	0.15	13.03	1.95	16.29	2.44	8.29	1.24	12.95	1.94	5.35	0.80	0.04	0.01
FB 0020	Blueberries, raw	RAC	1	0.04	0.04	0.23	0.23	0.01	0.01	0.83	0.83	0.33	0.33	NC	-
FB 0269	Grapes, raw (incl must, excl dried, excl juice, excl wine)	RAC	0.52	6.48	3.37	11.31	5.88	5.21	2.71	9.50	4.94	4.66	2.42	0.78	0.41

DIFENOCONAZOLE (224)															
International Estimated Daily Intake (IEDI)				ADI = 0 - 0.01 mg/kg bw											
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	1.1	3.09	3.40	1.51	1.66	0.03	0.03	1.38	1.52	4.26	4.69	0.42	0.46
JF 0269	Grape juice (from wine grapes)	PP	0.24	0.56	0.13	1.96	0.47	0.02	0.00	2.24	0.54	2.27	0.54	0.34	0.08
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.094	88.93	8.36	62.41	5.87	1.84	0.17	25.07	2.36	61.17	5.75	5.84	0.55
FB 0265	Cranberry, raw	RAC	0.2	0.06	0.01	0.01	0.00	0.01	0.00	1.22	0.24	0.11	0.02	NC	-
FB 0275	Strawberry, raw	RAC	0.42	4.49	1.89	5.66	2.38	0.02	0.01	6.63	2.78	5.75	2.42	0.05	0.02
FT 0305	Table olives, raw (incl preserved)	RAC	0.465	2.00	0.93	2.48	1.15	0.01	0.00	1.21	0.56	1.64	0.76	0.27	0.13
FT 0336	Guava, raw	RAC	0.0335	0.01	0.00	NC	-	0.42	0.01	NC	-	NC	-	NC	-
FI 0326	Avocado, raw	RAC	0.05	2.65	0.13	0.87	0.04	0.46	0.02	1.64	0.08	1.30	0.07	0.96	0.05
FI 0327	Banana, raw (incl plantains) (incl dried)	RAC	0.02	25.76	0.52	23.65	0.47	23.83	0.48	24.37	0.49	19.43	0.39	101.55	2.03
FI 0345	Mango, raw (incl canned mango, incl mango juice)	RAC	0.03	1.80	0.05	0.63	0.02	10.05	0.30	1.07	0.03	3.52	0.11	16.44	0.49
FI 0350	Papaya, raw	RAC	0.065	0.31	0.02	0.18	0.01	1.50	0.10	0.51	0.03	0.54	0.04	1.08	0.07
FI 2540	Pitaya, raw (i.e dragon fruit or pitahaya)	RAC	0.034	NC	-	NC	-	0.08	0.00	NC	-	NC	-	NC	-
FI 0351	Passion fruit, raw	RAC	0.01	0.01	0.00	0.01	0.00	NC	-	NC	-	0.02	0.00	NC	-
-	Onions, dry, raw	RAC	0.015	19.69	0.30	29.83	0.45	24.64	0.37	31.35	0.47	9.72	0.15	12.59	0.19
VA 0384	Leek, raw	RAC	0.08	4.01	0.32	4.41	0.35	0.72	0.06	0.54	0.04	16.41	1.31	0.03	0.00
-	Onions, green, raw	RAC	2.8	1.55	4.34	0.74	2.07	1.05	2.94	3.74	10.47	0.94	2.63	6.45	18.06
VB 0040	Group of Brassica vegetables (excl Brassica leafy vegetables), raw	RAC	0.35	20.71	7.25	39.81	13.93	25.06	8.77	37.93	13.28	18.12	6.34	16.74	5.86
VC 0424	Cucumber, raw	RAC	0.04	6.72	0.27	11.03	0.44	32.10	1.28	15.10	0.60	4.05	0.16	9.57	0.38
VC 0425	Gherkin, raw	RAC	0.04	0.41	0.02	5.89	0.24	NC	-	0.06	0.00	0.37	0.01	2.07	0.08
VC 0431	Squash, Summer (Courgette, Marrow, Zucchetti, Zucchini), raw	RAC	0.04	NC	-	NC	-	5.48	0.22	NC	-	NC	-	1.03	0.04
VC 0046	Melons, except watermelon, raw (Cantaloupe)	RAC	0.14	9.20	1.29	11.95	1.67	14.63	2.05	8.99	1.26	7.86	1.10	2.46	0.34
VC 0432	Watermelon, raw	RAC	0.01	4.60	0.05	9.82	0.10	68.50	0.69	13.19	0.13	1.99	0.02	14.56	0.15
VO 0448	Tomato, raw	RAC	0.1	32.13	3.21	51.27	5.13	34.92	3.49	73.37	7.34	15.15	1.52	8.88	0.89
-	Tomato, canned (& peeled)	PP	0.01	7.57	0.08	2.66	0.03	0.30	0.00	0.97	0.01	7.31	0.07	0.41	0.00
-	Tomato, paste (i.e. concentrated tomato sauce/puree)	PP	0.22	4.96	1.09	3.20	0.70	0.15	0.03	1.61	0.35	6.88	1.51	0.52	0.11

Annex 3

DIFENOCONAZOLE (224)															
International Estimated Daily Intake (IEDI)				ADI = 0 - 0.01 mg/kg bw											
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
JF 0448	Tomato, juice (single strength, incl concentrated)	PP	0.031	0.80	0.02	0.07	0.00	0.05	0.00	0.61	0.02	0.40	0.01	0.08	0.00
VO 0442	Okra, raw (i.e. Lady's Finger, Gombo)	RAC	0.14	NC	-	NC	-	0.04	0.01	0.17	0.02	NC	-	0.72	0.10
VO 0444	Peppers, chili, raw	RAC	0.24	5.57	1.34	14.00	3.36	8.25	1.98	5.77	1.38	6.44	1.55	2.53	0.61
VO 0445	Peppers, sweet, raw (incl dried)	RAC	0.14	0.82	0.11	1.53	0.21	10.85	1.52	4.59	0.64	1.84	0.26	2.00	0.28
VO 0440	Egg plant, raw (i.e. aubergine)	RAC	0.14	1.01	0.14	1.69	0.24	21.37	2.99	3.00	0.42	1.40	0.20	NC	-
VL 0483	Lettuce, leaf, raw	RAC	0.41	14.50	5.95	11.76	4.82	13.14	5.39	19.50	8.00	4.81	1.97	2.23	0.91
VP 0060	Group of Legume vegetables, raw	RAC	0.07	18.21	1.27	8.91	0.62	7.22	0.51	10.04	0.70	23.22	1.63	0.17	0.01
VD 0071	Beans, dry, raw (Phaseolus spp)	RAC	0.011	1.51	0.02	1.50	0.02	1.90	0.02	5.11	0.06	1.36	0.01	23.43	0.26
VD 0523	Broad bean, dry, raw (incl horse-bean, field bean) (Vicia faba)	RAC	0.011	0.02	0.00	0.01	0.00	1.16	0.01	0.40	0.00	NC	-	0.06	0.00
VD 0541	Soya bean, dry, raw (incl flour, incl paste, incl curd, incl sauce, excl oil)	RAC	0.01	0.47	0.00	0.77	0.01	9.12	0.09	8.05	0.08	0.04	0.00	6.06	0.06
OR 0541	Soya oil, refined	PP	0.08	19.06	1.52	21.06	1.68	5.94	0.48	33.78	2.70	40.05	3.20	13.39	1.07
-	Beans (dry) NES: including inter alia lablab or hyacinth bean (Dolichos spp.); jack or sword bean (Canavalia spp.); winged bean (Psophocarpus tetragonolobus); guar bean (Cyamopsis tetragonoloba); velvet bean (Stizolobium spp.); yam bean (Pachyrhizus erosus)	RAC	0.028	0.01	0.00	NC	-	0.57	0.02	0.11	0.00	0.16	0.00	0.94	0.03
VD 0533	Lentil (dry) (Lens spp), raw	RAC	0.028	0.95	0.03	1.18	0.03	0.40	0.01	0.96	0.03	0.71	0.02	1.28	0.04
VD 0537	Pigeon pea (dry) (Cajanus spp), raw	RAC	0.028	NC	-	NC	-	0.20	0.01	NC	-	NC	-	NC	-
VR 0577	Carrots, raw	RAC	0.05	26.26	1.31	27.13	1.36	10.07	0.50	16.49	0.82	44.69	2.23	8.75	0.44
VR 0578	Celeriac, raw	RAC	0.12	2.97	0.36	1.79	0.21	NC	-	0.06	0.01	16.91	2.03	3.22	0.39
VR 0596	Sugar beet, raw (incl sugar)	RAC	0.02	0.01	0.00	NC	-	0.01	0.00	0.01	0.00	NC	-	NC	-
VR 0589	Potato, raw (incl flour, incl frozen, incl starch, incl tapioca)	RAC	1.2	225.03	270.04	234.24	281.09	71.48	85.78	177.55	213.06	234.55	281.46	37.71	45.25
VS 0624	Celery	RAC	0.14	7.68	1.08	2.85	0.40	NC	-	3.34	0.47	16.83	2.36	4.04	0.57
VS 0621	Asparagus, raw	RAC	0.02	0.84	0.02	2.08	0.04	7.11	0.14	1.01	0.02	1.69	0.03	0.04	0.00
VS 0620	Artichoke globe, raw	RAC	0.51	0.98	0.50	3.65	1.86	0.07	0.04	1.67	0.85	0.26	0.13	NC	-
CM 0649 (GC 0649)	Rice, husked, dry (incl flour, incl oil, incl beverages, incl starch, excl polished)	REP	1.1	3.70	4.07	2.11	2.32	1.51	1.66	1.75	1.93	0.29	0.32	5.12	5.63

DIFENOCONAZOLE (224)															
International Estimated Daily Intake (IEDI)				ADI = 0 - 0.01 mg/kg bw											
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
CM 1205	Rice polished, dry	PP	0.0086	13.38	0.12	10.80	0.09	262.08	2.25	57.16	0.49	12.83	0.11	62.78	0.54
GC 0447	Sweet corn on the cob, raw (incl frozen kernels, incl canned kernels)	RAC	0.01	11.43	0.11	3.71	0.04	0.74	0.01	13.63	0.14	3.07	0.03	1.50	0.02
TN 0085	Group of Tree nuts, raw (incl processed)	RAC	0.01	8.52	0.09	8.94	0.09	15.09	0.15	9.60	0.10	14.57	0.15	26.26	0.26
SO 0495	Rape seed, raw	RAC	0.03	NC	-	NC	-	0.01	0.00	NC	-	NC	-	NC	-
OR 0495	Rape seed oil, edible	PP	0.002	12.52	0.03	7.63	0.02	3.00	0.01	6.01	0.01	NC	-	NC	-
SO 0702	Sunflower seed, raw (incl oil)	RAC	0.01	23.40	0.23	29.33	0.29	1.24	0.01	13.85	0.14	6.48	0.06	6.91	0.07
OR 0691	Cotton seed oil, edible	PP	0.0014	1.68	0.00	0.66	0.00	1.13	0.00	1.18	0.00	0.89	0.00	0.37	0.00
SO 0697	Peanuts, nutmeat, raw (incl roasted, incl oil, incl butter)	RAC	0	5.63	0.00	2.75	0.00	9.58	0.00	5.82	0.00	13.71	0.00	1.84	0.00
SO 0305	Olives for oil production, raw	RAC	0.465	0.35	0.16	0.01	0.00	0.01	0.00	0.57	0.27	0.06	0.03	NC	-
-	Olive oil (virgin and residue oil)	PP	0.7	3.40	2.38	9.49	6.64	0.02	0.01	4.28	3.00	2.74	1.92	0.48	0.34
SB 0716	Coffee bean raw (incl roasted, incl instant coffee, incl substitutes)	RAC	0.01	10.90	0.11	12.44	0.12	0.77	0.01	9.48	0.09	22.07	0.22	8.15	0.08
HS 0444	Peppers, chili, dried	PP	1.1	0.11	0.12	0.21	0.23	0.36	0.40	0.21	0.23	0.25	0.28	0.15	0.17
DT 1114	Tea, green or black, fermented and dried, (including concentrates)	RAC	4.85	2.91	14.11	1.73	8.39	1.14	5.53	1.85	8.97	2.29	11.11	0.74	3.59
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.047	112.02	5.27	120.71	5.67	63.46	2.98	88.99	4.18	96.24	4.52	41.02	1.93
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.14	28.01	3.92	30.18	4.22	15.86	2.22	22.25	3.11	24.06	3.37	10.25	1.44
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.14	6.44	0.90	15.51	2.17	3.79	0.53	8.29	1.16	18.44	2.58	8.00	1.12
MO 0105	Edible offal (mammalian), raw	RAC	0.71	15.17	10.77	5.19	3.68	6.30	4.47	6.78	4.81	3.32	2.36	3.17	2.25
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.011	388.92	4.28	335.88	3.69	49.15	0.54	331.25	3.64	468.56	5.15	245.45	2.70
PM 0110	Poultry meat, raw (incl prepared)	RAC	0.0002	73.76	0.01	53.86	0.01	23.98	0.00	87.12	0.02	53.38	0.01	84.45	0.02
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.0002	0.10	0.00	0.10	0.00	NC	-	0.10	0.00	0.71	0.00	NC	-
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.0002	0.33	0.00	0.72	0.00	0.27	0.00	0.35	0.00	0.80	0.00	NC	-
PE 0112	Eggs, raw, (incl dried)	RAC	0.011	25.84	0.28	29.53	0.32	28.05	0.31	33.19	0.37	36.44	0.40	8.89	0.10
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-
Total intake (ug/person)=				437.6		459.2		187.0		371.4		460.6		186.1	

Annex 3

DIFENOCONAZOLE (224)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.01 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day			Intake as ug/person/day								
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
	Bodyweight per region (kg bw) =				60		60		55		60		60		60
	ADI (ug/person)=				600		600		550		600		600		600
	%ADI=				72.9%		76.5%		34.0%		61.9%		76.8%		31.0%
	Rounded %ADI=				70%		80%		30%		60%		80%		30%

DIFENOCONAZOLE (224)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.01 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day			Intake = daily intake: ug/person						
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
FC 0001	Group of Citrus fruit, raw (incl kumquat commodities)	RAC	0.16	20.93	3.35	2.35	0.38	30.71	4.91	0.15	0.02	4.45	0.71
JF 0001	Group of Citrus fruit, juice	PP	0.002	0.11	0.00	0.29	0.00	13.55	0.03	0.14	0.00	0.33	0.00
FP 0009	Group of Pome fruits, raw (incl. apple juice, incl apple cider)	RAC	0.86	68.89	59.25	11.06	9.51	80.62	69.33	189.82	163.25	19.56	16.82
FS 0013	Subgroup of Cherries, raw	RAC	0.04	0.01	0.00	0.01	0.00	5.96	0.24	0.01	0.00	NC	
FS 0014	Subgroup of Plums, raw (incl dried plums)	RAC	0.04	0.07	0.00	0.02	0.00	16.65	0.67	0.01	0.00	NC	
FS 2001	Subgroup of peaches, raw (incl dried apricots)	RAC	0.15	0.02	0.00	0.01	0.00	10.76	1.61	0.01	0.00	NC	
FB 0020	Blueberries, raw	RAC	1	NC	-	NC	-	0.20	0.20	NC	-	NC	
FB 0269	Grapes, raw (incl must, excl dried, excl juice, excl wine)	RAC	0.52	0.14	0.07	0.36	0.19	15.33	7.97	0.01	0.01	0.28	0.15
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	1.1	0.01	0.01	0.13	0.14	1.06	1.17	0.01	0.01	0.03	0.03
JF 0269	Grape juice (from wine grapes)	PP	0.24	0.01	0.00	0.01	0.00	0.41	0.10	0.01	0.00	NC	
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.094	0.31	0.03	0.23	0.02	60.43	5.68	0.52	0.05	31.91	3.00
FB 0265	Cranberry, raw	RAC	0.2	NC	-	NC	-	0.03	0.01	NC	-	NC	
FB 0275	Strawberry, raw	RAC	0.42	0.01	0.00	0.01	0.00	3.35	1.41	0.01	0.00	0.01	0.00
FT 0305	Table olives, raw (incl preserved)	RAC	0.465	0.01	0.00	0.01	0.00	1.75	0.81	0.01	0.00	0.24	0.11
FT 0336	Guava, raw	RAC	0.0335	0.10	0.00	0.08	0.00	NC	-	0.14	0.00	3.11	0.10
FI 0326	Avocado, raw	RAC	0.05	1.12	0.06	0.01	0.00	0.84	0.04	0.01	0.00	6.60	0.33

DIFENOCONAZOLE (224)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.01 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
FI 0327	Banana, raw (incl plantains) (incl dried)	RAC	0.02	44.80	0.90	118.17	2.36	25.25	0.51	454.49	9.09	310.23	6.20
FI 0345	Mango, raw (incl canned mango, incl mango juice)	RAC	0.03	12.25	0.37	6.83	0.20	0.76	0.02	0.01	0.00	20.12	0.60
FI 0350	Papaya, raw	RAC	0.065	6.47	0.42	0.25	0.02	0.19	0.01	0.01	0.00	26.42	1.72
FI 2540	Pitaya, raw (i.e dragon fruit or pitahaya)	RAC	0.034	0.01	0.00	0.01	0.00	NC	-	0.01	0.00	0.04	0.00
FI 0351	Passion fruit, raw	RAC	0.01	0.12	0.00	0.10	0.00	0.01	0.00	0.18	0.00	3.81	0.04
-	Onions, dry, raw	RAC	0.015	9.01	0.14	20.24	0.30	30.90	0.46	9.61	0.14	2.11	0.03
VA 0384	Leek, raw	RAC	0.08	0.02	0.00	1.44	0.12	1.22	0.10	0.01	0.00	NC	-
-	Onions, green, raw	RAC	2.8	1.43	4.00	0.05	0.14	0.20	0.56	NC	-	6.30	17.64
VB 0040	Group of Brassica vegetables (excl Brassica leafy vegetables), raw	RAC	0.35	5.46	1.91	4.28	1.50	58.72	20.55	0.02	0.01	NC	-
VC 0424	Cucumber, raw	RAC	0.04	0.68	0.03	1.81	0.07	10.40	0.42	0.01	0.00	0.04	0.00
VC 0425	Gherkin, raw	RAC	0.04	0.15	0.01	0.39	0.02	3.15	0.13	0.01	0.00	0.01	0.00
VC 0431	Squash, Summer (Courgette, Marrow, Zucchini, Zucchini), raw	RAC	0.04	0.09	0.00	1.01	0.04	NC	-	1.91	0.08	NC	-
VC 0046	Melons, except watermelon, raw (Cantaloupe)	RAC	0.14	0.19	0.03	0.10	0.01	4.98	0.70	0.01	0.00	NC	-
VC 0432	Watermelon, raw	RAC	0.01	4.29	0.04	0.30	0.00	28.70	0.29	0.01	0.00	NC	-
VO 0448	Tomato, raw	RAC	0.1	12.99	1.30	4.79	0.48	58.40	5.84	0.92	0.09	0.09	0.01
-	Tomato, canned (& peeled)	PP	0.01	0.07	0.00	0.08	0.00	2.42	0.02	0.07	0.00	NC	-
-	Tomato, paste (i.e. concentrated tomato sauce/puree)	PP	0.22	0.58	0.13	0.22	0.05	2.21	0.49	0.24	0.05	3.10	0.68
JF 0448	Tomato, juice (single strength, incl concentrated)	PP	0.031	0.05	0.00	0.01	0.00	0.42	0.01	0.01	0.00	0.02	0.00
VO 0442	Okra, raw (i.e. Lady's Finger, Gombo)	RAC	0.14	6.23	0.87	0.10	0.01	NC	-	NC	-	NC	-
VO 0444	Peppers, chili, raw	RAC	0.24	3.47	0.83	3.56	0.85	16.30	3.91	0.01	0.00	NC	-
VO 0445	Peppers, sweet, raw (incl dried)	RAC	0.14	5.49	0.77	10.57	1.48	8.84	1.24	0.91	0.13	NC	-
VO 0440	Egg plant, raw (i.e. aubergine)	RAC	0.14	1.31	0.18	8.26	1.16	3.95	0.55	0.01	0.00	NC	-
VL 0483	Lettuce, leaf, raw	RAC	0.41	0.29	0.12	0.03	0.01	6.71	2.75	0.01	0.00	NC	-
VP 0060	Group of Legume vegetables, raw	RAC	0.07	0.58	0.04	3.16	0.22	10.38	0.73	0.04	0.00	NC	-
VD 0071	Beans, dry, raw (Phaseolus spp)	RAC	0.011	7.11	0.08	2.33	0.03	3.76	0.04	44.70	0.49	3.27	0.04
VD 0523	Broad bean, dry, raw (incl horse-bean, field bean) (Vicia faba)	RAC	0.011	3.70	0.04	0.03	0.00	0.17	0.00	0.01	0.00	NC	-

Annex 3

DIFENOCONAZOLE (224)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.01 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
VD 0541	Soya bean, dry, raw (incl flour, incl paste, incl curd, incl sauce, excl oil)	RAC	0.01	2.89	0.03	0.21	0.00	0.48	0.00	3.16	0.03	0.26	0.00
OR 0541	Soya oil, refined	PP	0.08	2.32	0.19	2.54	0.20	18.70	1.50	2.51	0.20	6.29	0.50
-	Beans (dry) NES: including inter alia lablab or hyacinth bean (Dolichos spp.); jack or sword bean (Canavalia spp.); winged bean (Psophocarpus tetragonolobus); guar bean (Cyamopsis tetragonoloba); velvet bean (Stizolobium spp.); yam bean (Pachyrrhizus erosus)	RAC	0.028	2.54	0.07	1.77	0.05	0.03	0.00	0.03	0.00	3.99	0.11
VD 0533	Lentil (dry) (Lens spp), raw	RAC	0.028	0.67	0.02	7.26	0.20	0.37	0.01	0.08	0.00	NC	-
VD 0537	Pigeon pea (dry) (Cajanus spp), raw	RAC	0.028	1.14	0.03	0.03	0.00	NC	-	5.53	0.15	NC	-
VR 0577	Carrots, raw	RAC	0.05	2.07	0.10	3.00	0.15	25.29	1.26	0.05	0.00	NC	-
VR 0578	Celeriac, raw	RAC	0.12	2.91	0.35	2.10	0.25	7.59	0.91	1.97	0.24	3.93	0.47
VR 0596	Sugar beet, raw (incl sugar)	RAC	0.02	3.93	0.08	1.68	0.03	NC	-	NC	-	36.12	0.72
VR 0589	Potato, raw (incl flour, incl frozen, incl starch, incl tapioca)	RAC	1.2	23.96	28.75	13.56	16.27	213.41	256.09	104.35	125.22	8.56	10.27
VS 0624	Celery	RAC	0.14	3.66	0.51	2.65	0.37	4.84	0.68	2.47	0.35	4.94	0.69
VS 0621	Asparagus, raw	RAC	0.02	0.01	0.00	0.01	0.00	0.17	0.00	0.01	0.00	NC	-
VS 0620	Artichoke globe, raw	RAC	0.51	0.01	0.01	NC	-	0.08	0.04	0.01	0.01	NC	-
CM 0649 (GC 0649)	Rice, husked, dry (incl flour, incl oil, incl beverages, incl starch, excl polished)	REP	1.1	13.58	14.94	4.29	4.72	2.17	2.39	0.01	0.01	8.84	9.72
CM 1205	Rice polished, dry	PP	0.0086	30.20	0.26	218.34	1.88	12.77	0.11	15.24	0.13	51.35	0.44
GC 0447	Sweet corn on the cob, raw (incl frozen kernels, incl canned kernels)	RAC	0.01	3.63	0.04	20.50	0.21	8.78	0.09	0.02	0.00	0.17	0.00
TN 0085	Group of Tree nuts, raw (incl processed)	RAC	0.01	4.39	0.04	135.53	1.36	6.11	0.06	0.72	0.01	317.74	3.18
SO 0495	Rape seed, raw	RAC	0.03	NC	-	0.01	0.00	NC	-	NC	-	NC	-
OR 0495	Rape seed oil, edible	PP	0.002	0.07	0.00	0.03	0.00	4.62	0.01	0.03	0.00	NC	-
SO 0702	Sunflower seed, raw (incl oil)	RAC	0.01	0.94	0.01	0.22	0.00	32.01	0.32	12.12	0.12	0.48	0.00
OR 0691	Cotton seed oil, edible	PP	0.0014	1.28	0.00	0.05	0.00	0.45	0.00	0.42	0.00	0.15	0.00
SO 0697	Peanuts, nutmeat, raw (incl roasted, incl oil, incl butter)	RAC	0	18.82	0.00	0.57	0.00	2.28	0.00	6.90	0.00	0.53	0.00
SO 0305	Olives for oil production, raw	RAC	0.465	NC	-	NC	-	0.02	0.01	NC	-	NC	-
-	Olive oil (virgin and residue oil)	PP	0.7	0.03	0.02	0.02	0.01	2.14	1.50	0.01	0.01	0.10	0.07

DIFENOCONAZOLE (224)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.01 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
SB 0716	Coffee bean raw (incl roasted, incl instant coffee, incl substitutes)	RAC	0.01	0.95	0.01	1.32	0.01	11.64	0.12	2.96	0.03	14.73	0.15
HS 0444	Peppers, chili, dried	PP	1.1	0.58	0.64	1.27	1.40	1.21	1.33	0.12	0.13	NC	-
DT 1114	Tea, green or black, fermented and dried, (including concentrates)	RAC	4.85	0.53	2.57	5.25	25.46	0.86	4.17	0.56	2.72	0.88	4.27
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.047	23.34	1.10	40.71	1.91	97.15	4.57	18.06	0.85	57.71	2.71
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.14	5.84	0.82	10.18	1.42	24.29	3.40	4.52	0.63	14.43	2.02
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.14	1.05	0.15	1.14	0.16	18.69	2.62	0.94	0.13	3.12	0.44
MO 0105	Edible offal (mammalian), raw	RAC	0.71	4.64	3.29	1.97	1.40	10.01	7.11	3.27	2.32	3.98	2.83
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.011	108.75	1.20	70.31	0.77	436.11	4.80	61.55	0.68	79.09	0.87
PM 0110	Poultry meat, raw (incl prepared)	RAC	0.0002	3.92	0.00	12.03	0.00	57.07	0.01	5.03	0.00	55.56	0.01
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.0002	NC	-	NC	-	0.32	0.00	NC	-	NC	-
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.0002	0.10	0.00	0.70	0.00	0.97	0.00	0.10	0.00	NC	-
PE 0112	Eggs, raw, (incl dried)	RAC	0.011	3.84	0.04	4.41	0.05	27.25	0.30	1.13	0.01	7.39	0.08
-	-	-		-	-	-	-	-	-	-	-	-	-
Total intake (ug/person)=				130.3		77.6		426.9		307.4		87.8	
Bodyweight per region (kg bw) =				60		60		60		60		60	
ADI (ug/person)=				600		600		600		600		600	
%ADI=				21.7%		12.9%		71.2%		51.2%		14.6%	
Rounded %ADI=				20%		10%		70%		50%		10%	

ETHIPROLE (304)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.005 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
VD 0541	Soya bean, dry, raw (incl flour, incl paste, incl curd, incl sauce, excl oil)	RAC	0.0065	0.63	0.00	1.09	0.01	0.40	0.00	1.40	0.01	1.68	0.01	0.48	0.00
OR 0541	Soya oil, refined	PP	0.002	12.99	0.03	10.43	0.02	3.63	0.01	13.10	0.03	10.70	0.02	13.10	0.03
CM 0649 (GC 0649)	Rice, husked, dry (incl flour, incl oil, incl beverages, incl starch, excl polished)	REP	0.14	1.26	0.18	1.58	0.22	31.05	4.35	5.43	0.76	0.90	0.13	2.18	0.31

Annex 3

ETHIPROLE (304)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.005 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
CM 1205	Rice polished, dry	PP	0.04	34.21	1.37	10.39	0.42	41.72	1.67	82.38	3.30	150.24	6.01	70.47	2.82
SB 0716	Coffee bean, raw (i.e. green coffee)	RAC	0.0245	0.96	0.02	0.16	0.00	0.91	0.02	0.27	0.01	1.37	0.03	0.46	0.01
SM 0716	Coffee bean, roasted	PP	0.044	0.19	0.01	0.91	0.04	0.16	0.01	2.50	0.11	0.39	0.02	0.40	0.02
-	Coffee bean, instant coffee (incl essences and concentrates)	PP	0.048	0.07	0.00	0.94	0.05	0.07	0.00	0.70	0.03	0.07	0.00	0.29	0.01
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.021	24.96	0.52	57.95	1.22	16.70	0.35	38.38	0.81	26.46	0.56	29.00	0.61
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.094	6.24	0.59	14.49	1.36	4.18	0.39	9.60	0.90	6.62	0.62	7.25	0.68
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.094	3.29	0.31	6.14	0.58	0.82	0.08	1.57	0.15	2.23	0.21	1.07	0.10
MO 0105	Edible offal (mammalian), raw	RAC	0.076	4.79	0.36	9.68	0.74	2.97	0.23	5.49	0.42	3.84	0.29	5.03	0.38
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.011	289.65	3.19	485.88	5.34	26.92	0.30	239.03	2.63	199.91	2.20	180.53	1.99
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0.02	13.17	0.26	26.78	0.54	7.24	0.14	116.71	2.33	22.54	0.45	32.09	0.64
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0.037	1.46	0.05	2.98	0.11	0.80	0.03	12.97	0.48	2.50	0.09	3.57	0.13
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.037	0.10	0.00	0.10	0.00	NC	-	0.10	0.00	0.10	0.00	0.10	0.00
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.031	0.12	0.00	0.12	0.00	0.11	0.00	5.37	0.17	0.24	0.01	0.10	0.00
PE 0112	Eggs, raw, (incl dried)	RAC	0.03	7.84	0.24	23.08	0.69	2.88	0.09	14.89	0.45	9.81	0.29	14.83	0.44
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-
Total intake (ug/person)=				7.1		11.3		7.7		12.6		10.9		8.2	
Bodyweight per region (kg bw) =				60		60		60		60		60		60	
ADI (ug/person)=				300		300		300		300		300		300	
%ADI=				2.4%		3.8%		2.6%		4.2%		3.6%		2.7%	
Rounded %ADI=				2%		4%		3%		4%		4%		3%	

ETHIPROLE (304)															
International Estimated Daily Intake (IEDI)				ADI = 0 - 0.005 mg/kg bw											
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
VD 0541	Soya bean, dry, raw (incl flour, incl paste, incl curd, incl sauce, excl oil)	RAC	0.0065	0.47	0.00	0.77	0.01	9.12	0.06	8.05	0.05	0.04	0.00	6.06	0.04
OR 0541	Soya oil, refined	PP	0.002	19.06	0.04	21.06	0.04	5.94	0.01	33.78	0.07	40.05	0.08	13.39	0.03
CM 0649 (GC 0649)	Rice, husked, dry (incl flour, incl oil, incl beverages, incl starch, excl polished)	REP	0.14	3.70	0.52	2.11	0.30	1.51	0.21	1.75	0.25	0.29	0.04	5.12	0.72
CM 1205	Rice polished, dry	PP	0.04	13.38	0.54	10.80	0.43	262.08	10.48	57.16	2.29	12.83	0.51	62.78	2.51
SB 0716	Coffee bean, raw (i.e. green coffee)	RAC	0.0245	0.60	0.01	NC	-	0.62	0.02	1.71	0.04	NC	-	3.51	0.09
SM 0716	Coffee bean, roasted	PP	0.044	7.02	0.31	9.75	0.43	0.02	0.00	5.09	0.22	13.38	0.59	0.77	0.03
-	Coffee bean, instant coffee (incl essences and concentrates)	PP	0.048	0.75	0.04	0.30	0.01	0.04	0.00	0.67	0.03	2.43	0.12	1.43	0.07
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.021	112.02	2.35	120.71	2.53	63.46	1.33	88.99	1.87	96.24	2.02	41.02	0.86
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.094	28.01	2.63	30.18	2.84	15.86	1.49	22.25	2.09	24.06	2.26	10.25	0.96
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.094	6.44	0.61	15.51	1.46	3.79	0.36	8.29	0.78	18.44	1.73	8.00	0.75
MO 0105	Edible offal (mammalian), raw	RAC	0.076	15.17	1.15	5.19	0.39	6.30	0.48	6.78	0.52	3.32	0.25	3.17	0.24
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.011	388.92	4.28	335.88	3.69	49.15	0.54	331.25	3.64	468.56	5.15	245.45	2.70
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0.02	66.38	1.33	48.47	0.97	21.58	0.43	78.41	1.57	48.04	0.96	76.01	1.52
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0.037	7.38	0.27	5.39	0.20	2.40	0.09	8.71	0.32	5.34	0.20	8.45	0.31
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.037	0.10	0.00	0.10	0.00	NC	-	0.10	0.00	0.71	0.03	NC	-
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.031	0.33	0.01	0.72	0.02	0.27	0.01	0.35	0.01	0.80	0.02	NC	-
PE 0112	Eggs, raw, (incl dried)	RAC	0.03	25.84	0.78	29.53	0.89	28.05	0.84	33.19	1.00	36.44	1.09	8.89	0.27
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-
Total intake (ug/person)=				14.9		14.2		16.4		14.7		15.1		11.1	
Bodyweight per region (kg bw) =				60		60		55		60		60		60	
ADI (ug/person)=				300		300		275		300		300		300	
%ADI=				5.0%		4.7%		5.9%		4.9%		5.0%		3.7%	
Rounded %ADI=				5%		5%		6%		5%		5%		4%	

Annex 3

ETHIPROLE (304)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.005 mg/kg bw						
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person								
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake	
VD 0541	Soya bean, dry, raw (incl flour, incl paste, incl curd, incl sauce, excl oil)	RAC	0.0065	2.89	0.02	0.21	0.00	0.48	0.00	3.16	0.02	0.26	0.00	
OR 0541	Soya oil, refined	PP	0.002	2.32	0.00	2.54	0.01	18.70	0.04	2.51	0.01	6.29	0.01	
CM 0649 (GC 0649)	Rice, husked, dry (incl flour, incl oil, incl beverages, incl starch, excl polished)	REP	0.14	13.58	1.90	4.29	0.60	2.17	0.30	0.01	0.00	8.84	1.24	
CM 1205	Rice polished, dry	PP	0.04	30.20	1.21	218.34	8.73	12.77	0.51	15.24	0.61	51.35	2.05	
SB 0716	Coffee bean, raw (i.e. green coffee)	RAC	0.0245	0.83	0.02	0.69	0.02	1.09	0.03	2.91	0.07	0.82	0.02	
SM 0716	Coffee bean, roasted	PP	0.044	0.02	0.00	0.41	0.02	7.50	0.33	0.01	0.00	0.06	0.00	
-	Coffee bean, instant coffee (incl essences and concentrates)	PP	0.048	0.03	0.00	0.05	0.00	0.60	0.03	0.01	0.00	5.53	0.27	
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.021	23.34	0.49	40.71	0.85	97.15	2.04	18.06	0.38	57.71	1.21	
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.094	5.84	0.55	10.18	0.96	24.29	2.28	4.52	0.42	14.43	1.36	
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.094	1.05	0.10	1.14	0.11	18.69	1.76	0.94	0.09	3.12	0.29	
MO 0105	Edible offal (mammalian), raw	RAC	0.076	4.64	0.35	1.97	0.15	10.01	0.76	3.27	0.25	3.98	0.30	
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.011	108.75	1.20	70.31	0.77	436.11	4.80	61.55	0.68	79.09	0.87	
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0.02	3.53	0.07	10.83	0.22	51.36	1.03	4.53	0.09	50.00	1.00	
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0.037	0.39	0.01	1.20	0.04	5.71	0.21	0.50	0.02	5.56	0.21	
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.037	NC	-	NC	-	0.32	0.01	NC	-	NC	-	
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.031	0.10	0.00	0.70	0.02	0.97	0.03	0.10	0.00	NC	-	
PE 0112	Eggs, raw, (incl dried)	RAC	0.03	3.84	0.12	4.41	0.13	27.25	0.82	1.13	0.03	7.39	0.22	
-	-	-		-	-	-	-	-	-	-	-	-	-	
Total intake (ug/person)=					6.0		12.6		15.0		2.7		9.1	
Bodyweight per region (kg bw) =					60		60		60		60		60	
ADI (ug/person)=					300		300		300		300		300	
%ADI=					2.0%		4.2%		5.0%		0.9%		3.0%	
Rounded %ADI=					2%		4%		5%		1%		3%	

FENBUCONAZOLE (197)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
FC 0002	Subgroup of Lemons and limes, raw (incl lemon juice) (incl kumquat commodities)	RAC	0.018	4.82	0.09	2.45	0.04	3.93	0.07	25.44	0.46	8.74	0.16	16.23	0.29
FC 0003	Subgroup of Mandarins, raw (incl mandarin juice)	RAC	0.01	6.18	0.06	3.66	0.04	0.25	0.00	6.82	0.07	3.49	0.03	19.38	0.19
FC 0004	Subgroup of Oranges, sweet, sour, raw (incl orange juice)	RAC	0.01	23.26	0.23	9.71	0.10	12.09	0.12	62.02	0.62	22.09	0.22	59.91	0.60
FC 0005	Subgroup of Pummelo and grapefruits, raw (incl grapefruit juice)	RAC	0.01	0.66	0.01	0.69	0.01	0.96	0.01	10.20	0.10	1.25	0.01	2.97	0.03
FP 0009	Group of Pome fruits, raw (incl. apple juice, incl apple cider)	RAC	0.12	19.79	2.37	38.25	4.59	17.96	2.16	32.56	3.91	8.08	0.97	64.45	7.73
FS 0013	Subgroup of Cherries, raw	RAC	0.36	0.92	0.33	9.15	3.29	0.01	0.00	0.61	0.22	0.06	0.02	6.64	2.39
FS 0014	Subgroup of Plums, raw (incl dried plums)	RAC	0.08	2.67	0.21	8.77	0.70	0.07	0.01	3.03	0.24	0.70	0.06	4.34	0.35
FS 0240	Apricot, raw (incl dried)	RAC	0.25	5.15	1.29	3.66	0.92	0.03	0.01	2.25	0.56	0.17	0.04	6.80	1.70
-	Peaches and nectarines, raw	RAC	0.25	2.87	0.72	2.21	0.55	0.15	0.04	5.94	1.49	1.47	0.37	15.66	3.92
FB 0020	Blueberries, raw	RAC	0.06	0.01	0.00	0.03	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.01	0.00
FB 0269	Grapes, raw (incl must, incl dried, excl juice, excl wine)	RAC	0.3	15.15	4.55	11.38	3.41	0.05	0.02	22.18	6.65	4.19	1.26	63.07	18.92
JF 0269	Grape juice (from wine grapes)	PP	0.03	0.14	0.00	0.29	0.01	0.05	0.00	0.30	0.01	0.24	0.01	0.05	0.00
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.018	0.67	0.01	12.53	0.23	2.01	0.04	1.21	0.02	3.53	0.06	4.01	0.07
FB 0265	Cranberry, raw	RAC	0.13	0.02	0.00	0.01	0.00	NC	-	0.03	0.00	0.01	0.00	0.01	0.00
FI 0327	Banana, raw (incl plantains) (incl dried)	RAC	0.01	5.23	0.05	6.94	0.07	99.45	0.99	32.47	0.32	48.30	0.48	24.70	0.25
VC 0424	Cucumber, raw	RAC	0.02	8.01	0.16	30.66	0.61	1.45	0.03	19.84	0.40	0.27	0.01	34.92	0.70
VC 0431	Squash, Summer (Courgette, Marrow, Zucchini, Zucchini), raw	RAC	0.02	0.78	0.02	2.06	0.04	0.30	0.01	1.61	0.03	2.25	0.05	2.36	0.05
VC 0046	Melons, except watermelon, raw (Cantaloupe)	RAC	0.0025	8.90	0.02	8.64	0.02	0.80	0.00	17.90	0.04	2.80	0.01	29.17	0.07
VO 0051	Subgroup of peppers, raw (incl dried sweet peppers, excl dried chilipeppers), excl okra	RAC	0.15	8.48	1.27	13.74	2.06	10.13	1.52	11.29	1.69	9.52	1.43	26.36	3.95
GC 0650	Rye, raw (incl flour)	RAC	0.02	0.13	0.00	19.38	0.39	0.10	0.00	0.12	0.00	0.03	0.00	2.15	0.04
GC 0654	Wheat, raw (incl bulgur, incl fermented beverages, incl germ, incl wholemeal bread, excl white flour products, excl white bread)	RAC	0.02	0.01	0.00	1.13	0.02	0.01	0.00	0.05	0.00	0.74	0.01	0.01	0.00
CP 1211	Wheat, white bread	PP	0.0092	0.25	0.00	0.63	0.01	0.12	0.00	0.43	0.00	1.39	0.01	0.22	0.00
CF 1211	Wheat, white flour	PP	0.005	299.27	1.50	263.32	1.32	27.93	0.14	214.18	1.07	133.47	0.67	340.03	1.70

Annex 3

FENBUCONAZOLE (197)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw								
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day										
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake	
GC 0640	Barley, raw (incl malt extract, incl pot&pearled, incl flour & grits, incl beer, incl malt)	RAC	0.03	19.91	0.60	31.16	0.93	5.04	0.15	3.10	0.09	9.77	0.29	4.31	0.13	
TN 0085	Group of Tree nuts, raw (incl processed)	RAC	0	4.06	0.00	3.27	0.00	7.01	0.00	13.93	0.00	14.01	0.00	9.36	0.00	
SO 0495	Rape seed, raw (incl oil)	RAC	0.05	0.93	0.05	1.16	0.06	0.49	0.02	2.53	0.13	9.32	0.47	2.02	0.10	
SO 0702	Sunflower seed, raw (incl oil)	RAC	0.02	7.40	0.15	35.86	0.72	1.15	0.02	8.76	0.18	5.45	0.11	13.62	0.27	
SO 0697	Peanuts, nutmeat, raw (incl roasted, incl butter, excl oil)	RAC	0.03	0.46	0.01	1.21	0.04	6.64	0.20	2.71	0.08	1.26	0.04	1.84	0.06	
OR 0697	Peanut oil, edible	PP	0.04	0.36	0.01	0.01	0.00	2.57	0.10	0.07	0.00	2.29	0.09	0.36	0.01	
HS 0444	Peppers, chili, dried	PP	1.5	0.42	0.63	0.53	0.80	0.84	1.26	0.50	0.75	0.95	1.43	0.37	0.56	
DT 1114	Tea, green or black, fermented and dried	RAC	4.2	2.28	9.58	1.92	8.06	0.46	1.93	2.40	10.08	1.29	5.42	3.04	12.77	
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat)	RAC	0.003	31.20	0.09	72.44	0.22	20.88	0.06	47.98	0.14	33.08	0.10	36.25	0.11	
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0	3.29	0.00	6.14	0.00	0.82	0.00	1.57	0.00	2.23	0.00	1.07	0.00	
MO 0105	Edible offal (mammalian), raw	RAC	0.02	4.79	0.10	9.68	0.19	2.97	0.06	5.49	0.11	3.84	0.08	5.03	0.10	
PM 0110	Poultry meat, raw (incl prepared)	RAC	0	14.63	0.00	29.76	0.00	8.04	0.00	129.68	0.00	25.04	0.00	35.66	0.00	
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0	0.12	0.00	0.12	0.00	0.11	0.00	5.37	0.00	0.24	0.00	0.10	0.00	
PE 0112	Eggs, raw, (incl dried)	RAC	0	7.84	0.00	23.08	0.00	2.88	0.00	14.89	0.00	9.81	0.00	14.83	0.00	
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-	
Total intake (ug/person)=					24.1		29.4		9.0		29.5		13.9		57.1	
Bodyweight per region (kg bw) =					60		60		60		60		60		60	
ADI (ug/person)=					1800		1800		1800		1800		1800		1800	
%ADI=					1.3%		1.6%		0.5%		1.6%		0.8%		3.2%	
Rounded %ADI=					1%		2%		0%		2%		1%		3%	

FENBUCONAZOLE (197)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
FC 0002	Subgroup of Lemons and limes, raw (incl lemon juice) (incl kumquat commodities)	RAC	0.018	10.12	0.18	15.69	0.28	2.88	0.05	12.30	0.22	22.32	0.40	6.59	0.12
FC 0003	Subgroup of Mandarins, raw (incl mandarin juice)	RAC	0.01	12.42	0.12	14.99	0.15	16.08	0.16	10.78	0.11	9.94	0.10	NC	-
FC 0004	Subgroup of Oranges, sweet, sour, raw (incl orange juice)	RAC	0.01	83.66	0.84	27.64	0.28	7.37	0.07	67.80	0.68	43.97	0.44	187.74	1.88
FC 0005	Subgroup of Pummelo and grapefruits, raw (incl grapefruit juice)	RAC	0.01	8.21	0.08	4.60	0.05	0.64	0.01	5.85	0.06	19.98	0.20	368.86	3.69
FP 0009	Group of Pome fruits, raw (incl. apple juice, incl apple cider)	RAC	0.12	71.38	8.57	81.73	9.81	42.91	5.15	58.89	7.07	103.85	12.46	12.48	1.50
FS 0013	Subgroup of Cherries, raw	RAC	0.36	1.40	0.50	4.21	1.52	0.04	0.01	2.93	1.05	1.50	0.54	NC	-
FS 0014	Subgroup of Plums, raw (incl dried plums)	RAC	0.08	5.55	0.44	4.37	0.35	6.08	0.49	3.66	0.29	3.93	0.31	0.46	0.04
FS 0240	Apricot, raw (incl dried)	RAC	0.25	4.27	1.07	3.31	0.83	0.06	0.02	2.86	0.72	1.71	0.43	NC	-
-	Peaches and nectarines, raw	RAC	0.25	8.76	2.19	12.98	3.25	8.23	2.06	10.09	2.52	3.64	0.91	0.04	0.01
FB 0020	Blueberries, raw	RAC	0.06	0.04	0.00	0.23	0.01	0.01	0.00	0.83	0.05	0.33	0.02	NC	-
FB 0269	Grapes, raw (incl must, incl dried, excl juice, excl wine)	RAC	0.3	19.37	5.81	17.62	5.29	5.33	1.60	15.24	4.57	22.40	6.72	2.51	0.75
JF 0269	Grape juice (from wine grapes)	PP	0.03	0.56	0.02	1.96	0.06	0.02	0.00	2.24	0.07	2.27	0.07	0.34	0.01
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.018	88.93	1.60	62.41	1.12	1.84	0.03	25.07	0.45	61.17	1.10	5.84	0.11
FB 0265	Cranberry, raw	RAC	0.13	0.06	0.01	0.01	0.00	0.01	0.00	1.22	0.16	0.11	0.01	NC	-
FI 0327	Banana, raw (incl plantains) (incl dried)	RAC	0.01	25.76	0.26	23.65	0.24	23.83	0.24	24.37	0.24	19.43	0.19	101.55	1.02
VC 0424	Cucumber, raw	RAC	0.02	6.72	0.13	11.03	0.22	32.10	0.64	15.10	0.30	4.05	0.08	9.57	0.19
VC 0431	Squash, Summer (Courgette, Marrow, Zucchini, Zucchini), raw	RAC	0.02	NC	-	NC	-	5.48	0.11	NC	-	NC	-	1.03	0.02
VC 0046	Melons, except watermelon, raw (Cantaloupe)	RAC	0.0025	9.20	0.02	11.95	0.03	14.63	0.04	8.99	0.02	7.86	0.02	2.46	0.01
VO 0051	Subgroup of peppers, raw (incl dried sweet peppers, excl dried chilepeppers), excl okra	RAC	0.15	6.39	0.96	15.53	2.33	19.09	2.86	10.36	1.55	8.29	1.24	4.53	0.68
GC 0650	Rye, raw (incl flour)	RAC	0.02	3.21	0.06	35.38	0.71	0.21	0.00	6.50	0.13	1.49	0.03	NC	-
GC 0654	Wheat, raw (incl bulgur, incl fermented beverages, incl germ, incl wholemeal bread, excl white flour products, excl white bread)	RAC	0.02	1.00	0.02	0.11	0.00	0.05	0.00	0.84	0.02	0.06	0.00	0.06	0.00
CP 1211	Wheat, white bread	PP	0.0092	1.30	0.01	0.46	0.00	0.06	0.00	0.22	0.00	2.44	0.02	0.77	0.01
CF 1211	Wheat, white flour	PP	0.005	182.77	0.91	187.54	0.94	103.82	0.52	180.42	0.90	164.00	0.82	118.84	0.59

Annex 3

FENBUCONAZOLE (197)															
International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw											
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
GC 0640	Barley, raw (incl malt extract, incl pot&pearled, incl flour & grits, incl beer, incl malt)	RAC	0.03	36.18	1.09	53.45	1.60	9.39	0.28	35.25	1.06	46.68	1.40	15.92	0.48
TN 0085	Group of Tree nuts, raw (incl processed)	RAC	0	8.52	0.00	8.94	0.00	15.09	0.00	9.60	0.00	14.57	0.00	26.26	0.00
SO 0495	Rape seed, raw (incl oil)	RAC	0.05	32.68	1.63	19.91	1.00	7.83	0.39	15.69	0.78	NC	-	NC	-
SO 0702	Sunflower seed, raw (incl oil)	RAC	0.02	23.40	0.47	29.33	0.59	1.24	0.02	13.85	0.28	6.48	0.13	6.91	0.14
SO 0697	Peanuts, nutmeat, raw (incl roasted, incl butter, excl oil)	RAC	0.03	3.26	0.10	2.22	0.07	5.38	0.16	4.85	0.15	1.54	0.05	1.82	0.05
OR 0697	Peanut oil, edible	PP	0.04	1.02	0.04	0.23	0.01	1.81	0.07	0.42	0.02	5.23	0.21	0.01	0.00
HS 0444	Peppers, chili, dried	PP	1.5	0.11	0.17	0.21	0.32	0.36	0.54	0.21	0.32	0.25	0.38	0.15	0.23
DT 1114	Tea, green or black, fermented and dried	RAC	4.2	2.71	11.38	0.82	3.44	1.14	4.79	1.59	6.68	1.82	7.64	0.53	2.23
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat)	RAC	0.003	140.03	0.42	150.89	0.45	79.32	0.24	111.24	0.33	120.30	0.36	51.27	0.15
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0	6.44	0.00	15.51	0.00	3.79	0.00	8.29	0.00	18.44	0.00	8.00	0.00
MO 0105	Edible offal (mammalian), raw	RAC	0.02	15.17	0.30	5.19	0.10	6.30	0.13	6.78	0.14	3.32	0.07	3.17	0.06
PM 0110	Poultry meat, raw (incl prepared)	RAC	0	73.76	0.00	53.86	0.00	23.98	0.00	87.12	0.00	53.38	0.00	84.45	0.00
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0	0.33	0.00	0.72	0.00	0.27	0.00	0.35	0.00	0.80	0.00	NC	-
PE 0112	Eggs, raw, (incl dried)	RAC	0	25.84	0.00	29.53	0.00	28.05	0.00	33.19	0.00	36.44	0.00	8.89	0.00
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-
Total intake (ug/person)=				39.4		35.0		20.7		30.9		36.4		14.0	
Bodyweight per region (kg bw) =				60		60		55		60		60		60	
ADI (ug/person)=				1800		1800		1650		1800		1800		1800	
%ADI=				2.2%		1.9%		1.3%		1.7%		2.0%		0.8%	
Rounded %ADI=				2%		2%		1%		2%		2%		1%	

FENBUCONAZOLE (197)													
International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw									
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
FC 0002	Subgroup of Lemons and limes, raw (incl lemon juice) (incl kumquat commodities)	RAC	0.018	18.97	0.34	0.97	0.02	6.23	0.11	0.09	0.00	3.35	0.06
FC 0003	Subgroup of Mandarins, raw (incl mandarin juice)	RAC	0.01	0.16	0.00	0.27	0.00	9.06	0.09	0.01	0.00	0.02	0.00
FC 0004	Subgroup of Oranges, sweet, sour, raw (incl orange juice)	RAC	0.01	1.34	0.01	1.65	0.02	40.03	0.40	0.33	0.00	1.76	0.02
FC 0005	Subgroup of Pummelo and grapefruits, raw (incl grapefruit juice)	RAC	0.01	0.68	0.01	0.05	0.00	3.21	0.03	0.01	0.00	NC	-
FP 0009	Group of Pome fruits, raw (incl. apple juice, incl apple cider)	RAC	0.12	68.89	8.27	11.06	1.33	80.62	9.67	189.82	22.78	19.56	2.35
FS 0013	Subgroup of Cherries, raw	RAC	0.36	0.01	0.00	0.01	0.00	5.96	2.15	0.01	0.00	NC	-
FS 0014	Subgroup of Plums, raw (incl dried plums)	RAC	0.08	0.07	0.01	0.02	0.00	16.65	1.33	0.01	0.00	NC	-
FS 0240	Apricot, raw (incl dried)	RAC	0.25	0.01	0.00	0.01	0.00	3.29	0.82	0.01	0.00	NC	-
-	Peaches and nectarines, raw	RAC	0.25	0.02	0.01	0.01	0.00	7.47	1.87	0.01	0.00	NC	-
FB 0020	Blueberries, raw	RAC	0.06	NC	-	NC	-	0.20	0.01	NC	-	NC	-
FB 0269	Grapes, raw (incl must, incl dried, excl juice, excl wine)	RAC	0.3	0.16	0.05	0.92	0.28	19.73	5.92	0.02	0.01	0.40	0.12
JF 0269	Grape juice (from wine grapes)	PP	0.03	0.01	0.00	0.01	0.00	0.41	0.01	0.01	0.00	NC	-
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.018	0.31	0.01	0.23	0.00	60.43	1.09	0.52	0.01	31.91	0.57
FB 0265	Cranberry, raw	RAC	0.13	NC	-	NC	-	0.03	0.00	NC	-	NC	-
FI 0327	Banana, raw (incl plantains) (incl dried)	RAC	0.01	44.80	0.45	118.17	1.18	25.25	0.25	454.49	4.54	310.23	3.10
VC 0424	Cucumber, raw	RAC	0.02	0.68	0.01	1.81	0.04	10.40	0.21	0.01	0.00	0.04	0.00
VC 0431	Squash, Summer (Courgette, Marrow, Zucchini, Zucchini), raw	RAC	0.02	0.09	0.00	1.01	0.02	NC	-	1.91	0.04	NC	-
VC 0046	Melons, except watermelon, raw (Cantaloupe)	RAC	0.0025	0.19	0.00	0.10	0.00	4.98	0.01	0.01	0.00	NC	-
VO 0051	Subgroup of peppers, raw (incl dried sweet peppers, excl dried chili peppers), excl okra	RAC	0.15	8.97	1.35	14.13	2.12	25.14	3.77	0.91	0.14	NC	-
GC 0650	Rye, raw (incl flour)	RAC	0.02	0.03	0.00	0.01	0.00	13.95	0.28	0.01	0.00	0.88	0.02
GC 0654	Wheat, raw (incl bulgur, incl fermented beverages, incl germ, incl wholemeal bread, excl white flour products, excl white bread)	RAC	0.02	0.05	0.00	0.01	0.00	0.05	0.00	0.01	0.00	0.97	0.02
CP 1211	Wheat, white bread	PP	0.0092	0.43	0.00	0.41	0.00	1.56	0.01	0.11	0.00	0.07	0.00
CF 1211	Wheat, white flour	PP	0.005	43.75	0.22	85.81	0.43	206.68	1.03	19.38	0.10	92.92	0.46
GC 0640	Barley, raw (incl malt extract, incl pot&pearled, incl flour & grits, incl beer, incl malt)	RAC	0.03	11.58	0.35	2.33	0.07	46.71	1.40	3.72	0.11	16.26	0.49

Annex 3

FENBUCONAZOLE (197)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw						
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person								
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake	
TN 0085	Group of Tree nuts, raw (incl processed)	RAC	0	4.39	0.00	135.53	0.00	6.11	0.00	0.72	0.00	317.74	0.00	
SO 0495	Rape seed, raw (incl oil)	RAC	0.05	0.19	0.01	0.07	0.00	12.07	0.60	0.08	0.00	NC	-	
SO 0702	Sunflower seed, raw (incl oil)	RAC	0.02	0.94	0.02	0.22	0.00	32.01	0.64	12.12	0.24	0.48	0.01	
SO 0697	Peanuts, nutmeat, raw (incl roasted, incl butter, excl oil)	RAC	0.03	7.14	0.21	0.45	0.01	1.87	0.06	6.22	0.19	0.53	0.02	
OR 0697	Peanut oil, edible	PP	0.04	5.02	0.20	0.05	0.00	0.17	0.01	0.29	0.01	NC	-	
HS 0444	Peppers, chili, dried	PP	1.5	0.58	0.87	1.27	1.91	1.21	1.82	0.12	0.18	NC	-	
DT 1114	Tea, green or black, fermented and dried	RAC	4.2	0.53	2.23	5.25	22.05	0.63	2.65	0.56	2.35	0.82	3.44	
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat)	RAC	0.003	29.18	0.09	50.89	0.15	121.44	0.36	22.58	0.07	72.14	0.22	
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0	1.05	0.00	1.14	0.00	18.69	0.00	0.94	0.00	3.12	0.00	
MO 0105	Edible offal (mammalian), raw	RAC	0.02	4.64	0.09	1.97	0.04	10.01	0.20	3.27	0.07	3.98	0.08	
PM 0110	Poultry meat, raw (incl prepared)	RAC	0	3.92	0.00	12.03	0.00	57.07	0.00	5.03	0.00	55.56	0.00	
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0	0.10	0.00	0.70	0.00	0.97	0.00	0.10	0.00	NC	-	
PE 0112	Eggs, raw, (incl dried)	RAC	0	3.84	0.00	4.41	0.00	27.25	0.00	1.13	0.00	7.39	0.00	
-	-	-		-	-	-	-	-	-	-	-	-	-	
Total intake (ug/person)=					14.8		29.7		36.8		30.8		11.0	
Bodyweight per region (kg bw) =					60		60		60		60		60	
ADI (ug/person)=					1800		1800		1800		1800		1800	
%ADI=					0.8%		1.6%		2.0%		1.7%		0.6%	
Rounded %ADI=					1%		2%		2%		2%		1%	

FENHEXAMID (215)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.2 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
FP 0230	Pear, raw	RAC	2.05	2.16	4.43	6.24	12.79	0.05	0.10	4.07	8.34	1.16	2.38	5.34	10.95
FS 0013	Subgroup of Cherries, raw	RAC	1.35	0.92	1.24	9.15	12.35	0.01	0.01	0.61	0.82	0.06	0.08	6.64	8.96
FS 0014	Subgroup of Plums, raw (incl dried plums)	RAC	0.31	2.67	0.83	8.77	2.72	0.07	0.02	3.03	0.94	0.70	0.22	4.34	1.35
FS 0240	Apricot, raw (incl dried)	RAC	3.85	5.15	19.83	3.66	14.09	0.03	0.12	2.25	8.66	0.17	0.65	6.80	26.18
-	Peaches and nectarines, raw	RAC	3.85	2.87	11.05	2.21	8.51	0.15	0.58	5.94	22.87	1.47	5.66	15.66	60.29
FB 0264	Blackberries, raw	RAC	2	0.35	0.70	0.11	0.22	0.01	0.02	0.02	0.04	0.01	0.02	1.23	2.46
FB 0266	Dewberries, incl boysen- & loganberry, raw	RAC	2	0.01	0.02	0.01	0.02	0.01	0.02	0.01	0.02	0.01	0.02	0.01	0.02
FB 0272	Raspberries, red, black, raw	RAC	2	0.07	0.14	0.93	1.86	0.01	0.02	0.01	0.02	0.01	0.02	0.01	0.02
FB 0020	Blueberries, raw	RAC	1.65	0.01	0.02	0.03	0.05	0.01	0.02	0.01	0.02	0.01	0.02	0.01	0.02
FB 0021	Currants, Black, Red, White, raw	RAC	1.65	0.02	0.03	0.74	1.22	0.01	0.02	0.03	0.05	0.01	0.02	0.01	0.02
FB 0268	Gooseberry, raw	RAC	1.65	0.01	0.02	0.24	0.40	NC	-	0.01	0.02	0.01	0.02	NC	-
FB 0267	Elderberries, raw (incl processed)	RAC	1.65	0.44	0.73	0.27	0.45	0.34	0.56	1.41	2.33	NC	-	0.87	1.44
FB 0269	Grapes, raw (i.e. table grapes)	RAC	4.3	12.68	54.52	9.12	39.22	0.03	0.13	16.88	72.58	3.70	15.91	54.42	234.01
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	8	0.51	4.08	0.51	4.08	0.01	0.08	1.27	10.16	0.12	0.96	2.07	16.56
-	Graps must (from wine-grapes)	PP	1.8	0.33	0.59	0.13	0.23	0.01	0.02	0.02	0.04	0.01	0.02	0.02	0.04
FB 0275	Strawberry, raw	RAC	3.3	0.70	2.31	2.01	6.63	0.04	0.13	1.36	4.49	0.37	1.22	2.53	8.35
FI 0341	Kiwifruit, raw	RAC	6.3	0.03	0.19	0.36	2.27	0.01	0.06	1.17	7.37	0.06	0.38	0.69	4.35
VA 2031	Subgroup of bulb onions	RAC	0.055	31.65	1.74	43.28	2.38	3.68	0.20	38.48	2.12	20.46	1.13	47.29	2.60
VC 0424	Cucumber, raw	RAC	0.185	8.01	1.48	30.66	5.67	1.45	0.27	19.84	3.67	0.27	0.05	34.92	6.46
VC 0425	Gherkin, raw	RAC	0.185	1.73	0.32	6.64	1.23	0.31	0.06	4.29	0.79	0.29	0.05	7.56	1.40
VC 0431	Squash, Summer (Courgette, Marrow, Zucchini, Zucchini), raw	RAC	0.185	0.78	0.14	2.06	0.38	0.30	0.06	1.61	0.30	2.25	0.42	2.36	0.44
VO 0448	Tomato, raw	RAC	0.395	41.73	16.48	75.65	29.88	10.66	4.21	82.87	32.73	24.75	9.78	200.93	79.37
-	Tomato, paste (i.e. concentrated tomato sauce/puree)	PP	2.05	2.34	4.80	1.33	2.73	1.57	3.22	4.24	8.69	0.34	0.70	2.83	5.80
JF 0448	Tomato, juice (single strength, incl concentrated)	PP	0.13	0.29	0.04	0.29	0.04	0.01	0.00	0.38	0.05	0.05	0.01	0.14	0.02
VO 0445	Peppers, sweet, raw	RAC	0.71	1.43	1.02	2.61	1.85	1.05	0.75	2.01	1.43	2.59	1.84	6.24	4.43
VO 0440	Egg plant, raw (i.e. aubergine)	RAC	0.71	5.58	3.96	4.31	3.06	0.89	0.63	9.31	6.61	13.64	9.68	20.12	14.29

Annex 3

FENHEXAMID (215)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.2 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
VL 0483	Lettuce, leaf, raw	RAC	11.5	0.53	6.10	0.36	4.14	0.16	1.84	6.21	71.42	1.90	21.85	6.05	69.58
VS 0621	Asparagus, raw	RAC	0.02	0.01	0.00	0.01	0.00	0.01	0.00	0.03	0.00	0.07	0.00	0.21	0.00
TN 0660	Almonds, nutmeat	RAC	0.02	1.38	0.03	0.08	0.00	0.01	0.00	1.00	0.02	0.06	0.00	0.81	0.02
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat)	RAC	0	31.20	0.00	72.44	0.00	20.88	0.00	47.98	0.00	33.08	0.00	36.25	0.00
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 80% as muscle	RAC	0	24.96	0.00	57.95	0.00	16.70	0.00	38.38	0.00	26.46	0.00	29.00	0.00
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0	6.24	0.00	14.49	0.00	4.18	0.00	9.60	0.00	6.62	0.00	7.25	0.00
MO 0105	Edible offal (mammalian), raw	RAC	0	4.79	0.00	9.68	0.00	2.97	0.00	5.49	0.00	3.84	0.00	5.03	0.00
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0	289.65	0.00	485.88	0.00	26.92	0.00	239.03	0.00	199.91	0.00	180.53	0.00
PM 0110	Poultry meat, raw (incl prepared)	RAC	0	14.63	0.00	29.76	0.00	8.04	0.00	129.68	0.00	25.04	0.00	35.66	0.00
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0	13.17	0.00	26.78	0.00	7.24	0.00	116.71	0.00	22.54	0.00	32.09	0.00
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0	1.46	0.00	2.98	0.00	0.80	0.00	12.97	0.00	2.50	0.00	3.57	0.00
PF 0111	Poultry fat, raw (incl rendered)	RAC	0	0.10	0.00	0.10	0.00	NC	-	0.10	0.00	0.10	0.00	0.10	0.00
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0	0.12	0.00	0.12	0.00	0.11	0.00	5.37	0.00	0.24	0.00	0.10	0.00
PE 0112	Eggs, raw, (incl dried)	RAC	0	7.84	0.00	23.08	0.00	2.88	0.00	14.89	0.00	9.81	0.00	14.83	0.00
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-
Total intake (ug/person)=				136.8		158.5		13.1		266.6		73.1		559.4	
Bodyweight per region (kg bw) =				60		60		60		60		60		60	
ADI (ug/person)=				12000		12000		12000		12000		12000		12000	
%ADI=				1.1%		1.3%		0.1%		2.2%		0.6%		4.7%	
Rounded %ADI=				1%		1%		0%		2%		1%		5%	

FENHEXAMID (215)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.2 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
FP 0230	Pear, raw	RAC	2.05	8.79	18.02	8.44	17.30	12.37	25.36	9.60	19.68	10.27	21.05	0.23	0.47
FS 0013	Subgroup of Cherries, raw	RAC	1.35	1.40	1.89	4.21	5.68	0.04	0.05	2.93	3.96	1.50	2.03	NC	-
FS 0014	Subgroup of Plums, raw (incl dried plums)	RAC	0.31	5.55	1.72	4.37	1.35	6.08	1.88	3.66	1.13	3.93	1.22	0.46	0.14
FS 0240	Apricot, raw (incl dried)	RAC	3.85	4.27	16.44	3.31	12.74	0.06	0.23	2.86	11.01	1.71	6.58	NC	-
-	Peaches and nectarines, raw	RAC	3.85	8.76	33.73	12.98	49.97	8.23	31.69	10.09	38.85	3.64	14.01	0.04	0.15
FB 0264	Blackberries, raw	RAC	2	0.09	0.18	0.52	1.04	0.14	0.28	0.24	0.48	NC	-	0.01	0.02
FB 0266	Dewberries, incl boysen- & loganberry, raw	RAC	2	0.01	0.02	NC	-	0.01	0.02	0.01	0.02	NC	-	0.01	0.02
FB 0272	Raspberries, red, black, raw	RAC	2	0.47	0.94	0.91	1.82	0.01	0.02	0.99	1.98	1.14	2.28	NC	-
FB 0020	Blueberries, raw	RAC	1.65	0.04	0.07	0.23	0.38	0.01	0.02	0.83	1.37	0.33	0.54	NC	-
FB 0021	Currants, Black, Red, White, raw	RAC	1.65	0.48	0.79	4.23	6.98	NC	-	1.51	2.49	0.49	0.81	NC	-
FB 0268	Gooseberry, raw	RAC	1.65	0.03	0.05	1.04	1.72	0.01	0.02	0.23	0.38	NC	-	NC	-
FB 0267	Elderberries, raw (incl processed)	RAC	1.65	8.20	13.53	0.14	0.23	NC	-	NC	-	NC	-	1.87	3.09
FB 0269	Grapes, raw (i.e. table grapes)	RAC	4.3	6.33	27.22	11.22	48.25	5.21	22.40	9.38	40.33	4.55	19.57	0.78	3.35
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	8	3.09	24.72	1.51	12.08	0.03	0.24	1.38	11.04	4.26	34.08	0.42	3.36
-	Graps must (from wine-grapes)	PP	1.8	0.16	0.29	0.09	0.16	0.01	0.02	0.12	0.22	0.11	0.20	NC	-
FB 0275	Strawberry, raw	RAC	3.3	4.49	14.82	5.66	18.68	0.02	0.07	6.63	21.88	5.75	18.98	0.05	0.17
FI 0341	Kiwifruit, raw	RAC	6.3	2.46	15.50	3.62	22.81	0.04	0.25	1.48	9.32	7.43	46.81	0.03	0.19
VA 2031	Subgroup of bulb onions	RAC	0.055	20.67	1.14	31.32	1.72	37.52	2.06	35.08	1.93	11.77	0.65	13.74	0.76
VC 0424	Cucumber, raw	RAC	0.185	6.72	1.24	11.03	2.04	32.10	5.94	15.10	2.79	4.05	0.75	9.57	1.77
VC 0425	Gherkin, raw	RAC	0.185	0.41	0.08	5.89	1.09	NC	-	0.06	0.01	0.37	0.07	2.07	0.38
VC 0431	Squash, Summer (Courgette, Marrow, Zucchini, Zucchini), raw	RAC	0.185	NC	-	NC	-	5.48	1.01	NC	-	NC	-	1.03	0.19
VO 0448	Tomato, raw	RAC	0.395	32.13	12.69	51.27	20.25	34.92	13.79	73.37	28.98	15.15	5.98	8.88	3.51
-	Tomato, paste (i.e. concentrated tomato sauce/puree)	PP	2.05	4.96	10.17	3.20	6.56	0.15	0.31	1.61	3.30	6.88	14.10	0.52	1.07
JF 0448	Tomato, juice (single strength, incl concentrated)	PP	0.13	0.80	0.10	0.07	0.01	0.05	0.01	0.61	0.08	0.40	0.05	0.08	0.01
VO 0445	Peppers, sweet, raw	RAC	0.71	NC	-	NC	-	8.25	5.86	3.03	2.15	NC	-	0.91	0.65
VO 0440	Egg plant, raw (i.e. aubergine)	RAC	0.71	1.01	0.72	1.69	1.20	21.37	15.17	3.00	2.13	1.40	0.99	NC	-

Annex 3

FENHEXAMID (215)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.2 mg/kg bw												
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day			Intake as ug/person/day													
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake					
VL 0483	Lettuce, leaf, raw	RAC	11.5	14.50	166.75	11.76	135.24	13.14	151.11	19.50	224.25	4.81	55.32	2.23	25.65					
VS 0621	Asparagus, raw	RAC	0.02	0.84	0.02	2.08	0.04	7.11	0.14	1.01	0.02	1.69	0.03	0.04	0.00					
TN 0660	Almonds, nutmeat	RAC	0.02	0.81	0.02	2.21	0.04	0.03	0.00	1.02	0.02	1.47	0.03	NC	-					
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat)	RAC	0	140.03	0.00	150.89	0.00	79.32	0.00	111.24	0.00	120.30	0.00	51.27	0.00					
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0	112.02	0.00	120.71	0.00	63.46	0.00	88.99	0.00	96.24	0.00	41.02	0.00					
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0	28.01	0.00	30.18	0.00	15.86	0.00	22.25	0.00	24.06	0.00	10.25	0.00					
MO 0105	Edible offal (mammalian), raw	RAC	0	15.17	0.00	5.19	0.00	6.30	0.00	6.78	0.00	3.32	0.00	3.17	0.00					
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0	388.92	0.00	335.88	0.00	49.15	0.00	331.25	0.00	468.56	0.00	245.45	0.00					
PM 0110	Poultry meat, raw (incl prepared)	RAC	0	73.76	0.00	53.86	0.00	23.98	0.00	87.12	0.00	53.38	0.00	84.45	0.00					
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0	66.38	0.00	48.47	0.00	21.58	0.00	78.41	0.00	48.04	0.00	76.01	0.00					
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0	7.38	0.00	5.39	0.00	2.40	0.00	8.71	0.00	5.34	0.00	8.45	0.00					
PF 0111	Poultry fat, raw (incl rendered)	RAC	0	0.10	0.00	0.10	0.00	NC	-	0.10	0.00	0.71	0.00	NC	-					
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0	0.33	0.00	0.72	0.00	0.27	0.00	0.35	0.00	0.80	0.00	NC	-					
PE 0112	Eggs, raw, (incl dried)	RAC	0	25.84	0.00	29.53	0.00	28.05	0.00	33.19	0.00	36.44	0.00	8.89	0.00					
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-					
Total intake (ug/person)=				362.8			369.4			278.0			429.8			246.1		44.9		
Bodyweight per region (kg bw) =				60			60			55			60			60				
ADI (ug/person)=				12000			12000			11000			12000			12000				
%ADI=				3.0%			3.1%			2.5%			3.6%			2.1%			0.4%	
Rounded %ADI=				3%			3%			3%			4%			2%			0%	

FENHEXAMID (215)		International Estimated Daily Intake (IEDI)				ADI = 0 - 0.2 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
FP 0230	Pear, raw	RAC	2.05	0.07	0.14	0.14	0.29	9.45	19.37	0.01	0.02	0.14	0.29
FS 0013	Subgroup of Cherries, raw	RAC	1.35	0.01	0.01	0.01	0.01	5.96	8.05	0.01	0.01	NC	-
FS 0014	Subgroup of Plums, raw (incl dried plums)	RAC	0.31	0.07	0.02	0.02	0.01	16.65	5.16	0.01	0.00	NC	-
FS 0240	Apricot, raw (incl dried)	RAC	3.85	0.01	0.04	0.01	0.04	3.29	12.67	0.01	0.04	NC	-
-	Peaches and nectarines, raw	RAC	3.85	0.02	0.08	0.01	0.04	7.47	28.76	0.01	0.04	NC	-
FB 0264	Blackberries, raw	RAC	2	0.01	0.02	7.29	14.58	0.25	0.50	0.01	0.02	NC	-
FB 0266	Dewberries, incl boysen- & loganberry, raw	RAC	2	0.01	0.02	0.01	0.02	NC	-	0.01	0.02	NC	-
FB 0272	Raspberries, red, black, raw	RAC	2	0.01	0.02	0.01	0.02	2.04	4.08	0.01	0.02	NC	-
FB 0020	Blueberries, raw	RAC	1.65	NC	-	NC	-	0.20	0.33	NC	-	NC	-
FB 0021	Currants, Black, Red, White, raw	RAC	1.65	0.01	0.02	NC	-	0.74	1.22	NC	-	NC	-
FB 0268	Gooseberry, raw	RAC	1.65	NC	-	NC	-	0.12	0.20	NC	-	NC	-
FB 0267	Elderberries, raw (incl processed)	RAC	1.65	0.71	1.17	3.52	5.81	NC	-	0.38	0.63	2.32	3.83
FB 0269	Grapes, raw (i.e. table grapes)	RAC	4.3	0.14	0.60	0.36	1.55	15.22	65.45	0.01	0.04	0.09	0.39
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	8	0.01	0.08	0.13	1.04	1.06	8.48	0.01	0.08	0.03	0.24
-	Graps must (from wine-grapes)	PP	1.8	0.01	0.02	0.01	0.02	0.11	0.20	0.01	0.02	0.19	0.34
FB 0275	Strawberry, raw	RAC	3.3	0.01	0.03	0.01	0.03	3.35	11.06	0.01	0.03	0.01	0.03
FI 0341	Kiwifruit, raw	RAC	6.3	0.01	0.06	0.01	0.06	2.00	12.60	0.01	0.06	NC	-
VA 2031	Subgroup of bulb onions	RAC	0.055	9.83	0.54	22.30	1.23	34.69	1.91	9.65	0.53	2.39	0.13
VC 0424	Cucumber, raw	RAC	0.185	0.68	0.13	1.81	0.33	10.40	1.92	0.01	0.00	0.04	0.01
VC 0425	Gherkin, raw	RAC	0.185	0.15	0.03	0.39	0.07	3.15	0.58	0.01	0.00	0.01	0.00
VC 0431	Squash, Summer (Courgette, Marrow, Zucchini, Zucchini), raw	RAC	0.185	0.09	0.02	1.01	0.19	NC	-	1.91	0.35	NC	-
VO 0448	Tomato, raw	RAC	0.395	12.99	5.13	4.79	1.89	58.40	23.07	0.92	0.36	0.09	0.04
-	Tomato, paste (i.e. concentrated tomato sauce/puree)	PP	2.05	0.58	1.19	0.22	0.45	2.21	4.53	0.24	0.49	3.10	6.36
JF 0448	Tomato, juice (single strength, incl concentrated)	PP	0.13	0.05	0.01	0.01	0.00	0.42	0.05	0.01	0.00	0.02	0.00
VO 0445	Peppers, sweet, raw	RAC	0.71	1.24	0.88	1.27	0.90	NC	-	0.01	0.01	NC	-
VO 0440	Egg plant, raw (i.e. aubergine)	RAC	0.71	1.31	0.93	8.26	5.86	3.95	2.80	0.01	0.01	NC	-

Annex 3

FENHEXAMID (215)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.2 mg/kg bw						
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person								
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake	
VL 0483	Lettuce, leaf, raw	RAC	11.5	0.29	3.34	0.03	0.35	6.71	77.17	0.01	0.12	NC	-	
VS 0621	Asparagus, raw	RAC	0.02	0.01	0.00	0.01	0.00	0.17	0.00	0.01	0.00	NC	-	
TN 0660	Almonds, nutmeat	RAC	0.02	0.01	0.00	0.01	0.00	0.61	0.01	0.01	0.00	NC	-	
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat)	RAC	0	29.18	0.00	50.89	0.00	121.44	0.00	22.58	0.00	72.14	0.00	
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0	23.34	0.00	40.71	0.00	97.15	0.00	18.06	0.00	57.71	0.00	
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0	5.84	0.00	10.18	0.00	24.29	0.00	4.52	0.00	14.43	0.00	
MO 0105	Edible offal (mammalian), raw	RAC	0	4.64	0.00	1.97	0.00	10.01	0.00	3.27	0.00	3.98	0.00	
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0	108.75	0.00	70.31	0.00	436.11	0.00	61.55	0.00	79.09	0.00	
PM 0110	Poultry meat, raw (incl prepared)	RAC	0	3.92	0.00	12.03	0.00	57.07	0.00	5.03	0.00	55.56	0.00	
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0	3.53	0.00	10.83	0.00	51.36	0.00	4.53	0.00	50.00	0.00	
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0	0.39	0.00	1.20	0.00	5.71	0.00	0.50	0.00	5.56	0.00	
PF 0111	Poultry fat, raw (incl rendered)	RAC	0	NC	-	NC	-	0.32	0.00	NC	-	NC	-	
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0	0.10	0.00	0.70	0.00	0.97	0.00	0.10	0.00	NC	-	
PE 0112	Eggs, raw, (incl dried)	RAC	0	3.84	0.00	4.41	0.00	27.25	0.00	1.13	0.00	7.39	0.00	
-	-	-		-	-	-	-	-	-	-	-	-	-	
Total intake (ug/person)=					14.5		34.8		290.2		2.9		11.7	
Bodyweight per region (kg bw) =					60		60		60		60		60	
ADI (ug/person)=					12000		12000		12000		12000		12000	
%ADI=					0.1%		0.3%		2.4%		0.0%		0.1%	
Rounded %ADI=					0%		0%		2%		0%		0%	

FENPICOXAMID (305)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.05 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
FI 0327	Banana, raw (incl plantains) (incl dried)	RAC	0.01	5.23	0.05	6.94	0.07	99.45	0.99	32.47	0.32	48.30	0.48	24.70	0.25
GC 0650	Rye, raw	RAC	0.0215	NC	-	NC	-	0.02	0.00	0.01	0.00	0.01	0.00	0.01	0.00
CF 1250	Rye, flour (white flour and wholemeal flour)	PP	0.0088	0.11	0.00	15.51	0.14	0.06	0.00	0.10	0.00	0.03	0.00	1.72	0.02
GC 0653	Triticale, raw	RAC	0.0215	NC	-	NC	-	NC	-	0.01	0.00	NC	-	NC	-
GC 0653	Triticale, flour (white flour and wholemeal flour)	PP	0.0088	NC	-	NC	-	NC	-	NC	-	0.31	0.00	NC	-
GC 0654	Wheat, raw (incl meslin)	RAC	0.0215	0.01	0.00	1.12	0.02	NC	-	0.01	0.00	0.56	0.01	NC	-
-	Wheat, bulgur	PP	0.0088	NC	-	NC	-	NC	-	0.03	0.00	NC	-	NC	-
CF 1210	Wheat, germ	PP	0.0099	NC	-	NC	-	0.01	0.00	0.01	0.00	0.14	0.00	0.01	0.00
CP 1212	Wheat, wholemeal bread	PP	0.0043	0.01	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.03	0.00	0.01	0.00
CP 1211	Wheat, white bread	PP	0.0032	0.25	0.00	0.63	0.00	0.12	0.00	0.43	0.00	1.39	0.00	0.22	0.00
CF 1211	Wheat, white flour	PP	0.0062	299.27	1.86	263.32	1.63	27.93	0.17	214.18	1.33	133.47	0.83	340.03	2.11
-	Wheat, starch	PP	0.0032	0.02	0.00	NC	-	0.01	0.00	0.05	0.00	0.13	0.00	0.01	0.00
-	Wheat, gluten	PP	0.0058	0.01	0.00	0.01	0.00	0.01	0.00	0.27	0.00	0.01	0.00	0.03	0.00
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0	24.96	0.00	57.95	0.00	16.70	0.00	38.38	0.00	26.46	0.00	29.00	0.00
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.013	6.24	0.08	14.49	0.19	4.18	0.05	9.60	0.12	6.62	0.09	7.25	0.09
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.013	3.29	0.04	6.14	0.08	0.82	0.01	1.57	0.02	2.23	0.03	1.07	0.01
MO 0105	Edible offal (mammalian), raw	RAC	0.013	4.79	0.06	9.68	0.13	2.97	0.04	5.49	0.07	3.84	0.05	5.03	0.07

Total intake (ug/person)=

2.1

2.3

1.3

1.9

1.5

2.5

Bodyweight per region (kg bw) =

60

60

60

60

60

60

ADI (ug/person)=

3000

3000

3000

3000

3000

3000

%ADI=

0.1%

0.1%

0.0%

0.1%

0.0%

0.1%

Rounded %ADI=

0%

0%

0%

0%

0%

0%

Annex 3

FENPICOXAMID (305)

International Estimated Daily Intake (IEDI)

ADI = 0 - 0.05 mg/kg bw

Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
FI 0327	Banana, raw (incl plantains) (incl dried)	RAC	0.01	25.76	0.26	23.65	0.24	23.83	0.24	24.37	0.24	19.43	0.19	101.55	1.02
GC 0650	Rye, raw	RAC	0.0215	0.01	0.00	NC	-	0.06	0.00	0.01	0.00	NC	-	NC	-
CF 1250	Rye, flour (white flour and wholemeal flour)	PP	0.0088	2.57	0.02	28.31	0.25	0.12	0.00	5.20	0.05	1.20	0.01	NC	-
GC 0653	Triticale, raw	RAC	0.0215	NC	-	NC	-	0.01	0.00	0.01	0.00	NC	-	NC	-
GC 0653	Triticale, flour (white flour and wholemeal flour)	PP	0.0088	0.01	0.00	0.14	0.00	0.23	0.00	NC	-	NC	-	NC	-
GC 0654	Wheat, raw (incl meslin)	RAC	0.0215	NC	-	NC	-	NC	-	0.01	0.00	NC	-	NC	-
-	Wheat, bulgur	PP	0.0088	NC	-	NC	-	0.01	0.00	NC	-	NC	-	NC	-
CF 1210	Wheat, germ	PP	0.0099	0.97	0.01	0.10	0.00	0.03	0.00	0.01	0.00	NC	-	0.04	0.00
CP 1212	Wheat, wholemeal bread	PP	0.0043	0.03	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.05	0.00	0.02	0.00
CP 1211	Wheat, white bread	PP	0.0032	1.30	0.00	0.46	0.00	0.06	0.00	0.22	0.00	2.44	0.01	0.77	0.00
CF 1211	Wheat, white flour	PP	0.0062	182.77	1.13	187.54	1.16	103.82	0.64	180.42	1.12	164.00	1.02	118.84	0.74
-	Wheat, starch	PP	0.0032	NC	-	NC	-	0.01	0.00	0.31	0.00	NC	-	NC	-
-	Wheat, gluten	PP	0.0058	0.68	0.00	NC	-	0.01	0.00	0.01	0.00	NC	-	NC	-
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0	112.02	0.00	120.71	0.00	63.46	0.00	88.99	0.00	96.24	0.00	41.02	0.00
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.013	28.01	0.36	30.18	0.39	15.86	0.21	22.25	0.29	24.06	0.31	10.25	0.13
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.013	6.44	0.08	15.51	0.20	3.79	0.05	8.29	0.11	18.44	0.24	8.00	0.10
MO 0105	Edible offal (mammalian), raw	RAC	0.013	15.17	0.20	5.19	0.07	6.30	0.08	6.78	0.09	3.32	0.04	3.17	0.04

Total intake (ug/person)=

2.1

2.3

1.2

1.9

1.8

2.0

Bodyweight per region (kg bw) =

60

60

55

60

60

60

ADI (ug/person)=

3000

3000

2750

3000

3000

3000

%ADI=

0.1%

0.1%

0.0%

0.1%

0.1%

0.1%

Rounded %ADI=

0%

0%

0%

0%

0%

0%

FENPICOXAMID (305)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.05 mg/kg bw						
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person								
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake	
FI 0327	Banana, raw (incl plantains) (incl dried)	RAC	0.01	44.80	0.45	118.17	1.18	25.25	0.25	454.49	4.54	310.23	3.10	
GC 0650	Rye, raw	RAC	0.0215	0.01	0.00	NC	-	NC	-	0.01	0.00	NC	-	
CF 1250	Rye, flour (white flour and wholemeal flour)	PP	0.0088	0.02	0.00	0.01	0.00	11.16	0.10	0.01	0.00	0.70	0.01	
GC 0653	Triticale, raw	RAC	0.0215	0.01	0.00	NC	-	NC	-	NC	-	NC	-	
GC 0653	Triticale, flour (white flour and wholemeal flour)	PP	0.0088	NC	-	NC	-	NC	-	NC	-	NC	-	
GC 0654	Wheat, raw (incl meslin)	RAC	0.0215	NC	-	NC	-	NC	-	NC	-	0.97	0.02	
-	Wheat, bulgur	PP	0.0088	0.01	0.00	NC	-	NC	-	NC	-	NC	-	
CF 1210	Wheat, germ	PP	0.0099	0.04	0.00	0.01	0.00	0.01	0.00	0.01	0.00	NC	-	
CP 1212	Wheat, wholemeal bread	PP	0.0043	0.01	0.00	0.01	0.00	0.03	0.00	0.01	0.00	0.01	0.00	
CP 1211	Wheat, white bread	PP	0.0032	0.43	0.00	0.41	0.00	1.56	0.00	0.11	0.00	0.07	0.00	
CF 1211	Wheat, white flour	PP	0.0062	43.75	0.27	85.81	0.53	206.68	1.28	19.38	0.12	92.92	0.58	
-	Wheat, starch	PP	0.0032	0.01	0.00	0.02	0.00	NC	-	NC	-	NC	-	
-	Wheat, gluten	PP	0.0058	0.01	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.19	0.00	
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0	23.34	0.00	40.71	0.00	97.15	0.00	18.06	0.00	57.71	0.00	
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.013	5.84	0.08	10.18	0.13	24.29	0.32	4.52	0.06	14.43	0.19	
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.013	1.05	0.01	1.14	0.01	18.69	0.24	0.94	0.01	3.12	0.04	
MO 0105	Edible offal (mammalian), raw	RAC	0.013	4.64	0.06	1.97	0.03	10.01	0.13	3.27	0.04	3.98	0.05	
-	-	-		-	-	-	-	-	-	-	-	-	-	
Total intake (ug/person)=					0.9		1.9		2.3		4.8		4.0	
Bodyweight per region (kg bw) =					60		60		60		60		60	
ADI (ug/person)=					3000		3000		3000		3000		3000	
%ADI=					0.0%		0.1%		0.1%		0.2%		0.1%	
Rounded %ADI=					0%		0%		0%		0%		0%	

Annex 3

FLUOPYRAM (243)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.01 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
FC 0002	Subgroup of Lemons and limes, raw (incl lemon juice) (incl kumquat commodities)	RAC	0.325	4.82	1.57	2.45	0.80	3.93	1.28	25.44	8.27	8.74	2.84	16.23	5.27
FC 0003	Subgroup of Mandarins, raw (incl mandarin juice)	RAC	0.15	6.18	0.93	3.66	0.55	0.25	0.04	6.82	1.02	3.49	0.52	19.38	2.91
FC 0004	Subgroup of Oranges, sweet, sour, raw (incl orange juice)	RAC	0.15	23.26	3.49	9.71	1.46	12.09	1.81	62.02	9.30	22.09	3.31	59.91	8.99
FC 0005	Subgroup of Pummelo and grapefruits, raw (incl grapefruit juice)	RAC	0.14	0.66	0.09	0.69	0.10	0.96	0.13	10.20	1.43	1.25	0.18	2.97	0.42
FP 0009	Group of Pome fruits, raw (incl apple cider, excl apple juice)	RAC	0.135	19.35	2.61	34.06	4.60	17.87	2.41	25.74	3.47	7.69	1.04	56.85	7.67
JF 0226	Apple juice, single strength (incl. concentrated)	PP	0.01	0.32	0.00	3.07	0.03	0.07	0.00	5.00	0.05	0.29	0.00	5.57	0.06
FS 0013	Subgroup of Cherries, raw	RAC	0.57	0.92	0.52	9.15	5.22	0.01	0.01	0.61	0.35	0.06	0.03	6.64	3.78
FS 0014	Subgroup of Plums, raw (incl dried plums)	RAC	0.13	2.67	0.35	8.77	1.14	0.07	0.01	3.03	0.39	0.70	0.09	4.34	0.56
FS 2001	Subgroup of peaches, raw (incl dried apricots)	RAC	0.22	8.01	1.76	5.87	1.29	0.18	0.04	8.19	1.80	1.64	0.36	22.46	4.94
FB 2005	Subgroup of Caneberries, raw	RAC	0.83	0.42	0.35	1.05	0.87	0.01	0.01	0.02	0.02	0.02	0.02	1.24	1.03
FB 2006	Subgroup of Bush berries, raw (including processed)	RAC	1.15	0.53	0.61	1.31	1.51	0.40	0.46	1.66	1.91	0.01	0.01	0.99	1.14
FB 0269	Grapes, raw (incl must, excl dried, excl juice, excl wine)	RAC	0.58	13.02	7.55	9.25	5.37	0.03	0.02	16.91	9.81	3.70	2.15	54.44	31.58
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	1.68	0.51	0.86	0.51	0.86	0.01	0.02	1.27	2.13	0.12	0.20	2.07	3.48
JF 0269	Grape juice (from wine grapes)	PP	0.012	0.14	0.00	0.29	0.00	0.05	0.00	0.30	0.00	0.24	0.00	0.05	0.00
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.1	0.67	0.07	12.53	1.25	2.01	0.20	1.21	0.12	3.53	0.35	4.01	0.40
FB 0275	Strawberry, raw	RAC	0.025	0.70	0.02	2.01	0.05	0.04	0.00	1.36	0.03	0.37	0.01	2.53	0.06
FI 0327	Banana, raw (incl plantains) (incl dried)	RAC	0.175	5.23	0.92	6.94	1.21	99.45	17.40	32.47	5.68	48.30	8.45	24.70	4.32
FI 0345	Mango, raw (incl canned mango, incl mango juice)	RAC	0.02	10.48	0.21	0.01	0.00	7.24	0.14	6.87	0.14	19.98	0.40	6.25	0.13
VA 0381	Garlic, raw	RAC	0.01	2.29	0.02	5.78	0.06	0.11	0.00	3.69	0.04	1.65	0.02	3.91	0.04
-	Onions, dry, raw	RAC	0.01	29.36	0.29	37.50	0.38	3.56	0.04	34.78	0.35	18.81	0.19	43.38	0.43
VA 0384	Leek, raw	RAC	0.01	0.18	0.00	1.59	0.02	0.03	0.00	0.28	0.00	0.01	0.00	3.21	0.03
-	Onions, green, raw	RAC	5.1	2.45	12.50	1.49	7.60	1.02	5.20	2.60	13.26	0.60	3.06	2.03	10.35
VB 0400	Broccoli, raw	RAC	0.05	0.88	0.04	0.17	0.01	0.01	0.00	1.25	0.06	3.00	0.15	1.09	0.05
VB 0404	Cauliflower, raw	RAC	0.01	1.65	0.02	0.32	0.00	0.01	0.00	2.33	0.02	4.79	0.05	2.03	0.02

FLUOPYRAM (243)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.01 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
VB 0402	Brussels sprouts, raw	RAC	0.06	0.63	0.04	6.41	0.38	0.13	0.01	1.03	0.06	NC	-	2.35	0.14
VB 0041	Cabbages, head, raw	RAC	0.01	2.73	0.03	27.92	0.28	0.55	0.01	4.47	0.04	4.27	0.04	10.25	0.10
VC 0424	Cucumber, raw	RAC	0.11	8.01	0.88	30.66	3.37	1.45	0.16	19.84	2.18	0.27	0.03	34.92	3.84
VO 0448	Tomato, raw	RAC	0.11	41.73	4.59	75.65	8.32	10.66	1.17	82.87	9.12	24.75	2.72	200.93	22.10
-	Tomato, canned (& peeled)	PP	0.023	0.20	0.00	0.31	0.01	0.02	0.00	1.11	0.03	0.11	0.00	1.50	0.03
-	Tomato, paste (i.e. concentrated tomato sauce/puree)	PP	0.08	2.34	0.19	1.33	0.11	1.57	0.13	4.24	0.34	0.34	0.03	2.83	0.23
JF 0448	Tomato, juice (single strength, incl concentrated)	PP	0.04	0.29	0.01	0.29	0.01	0.01	0.00	0.38	0.02	0.05	0.00	0.14	0.01
VO 0051	Subgroup of peppers, raw (incl dried sweet peppers, excl dried chilipeppers), excl okra	RAC	0.14	8.48	1.19	13.74	1.92	10.13	1.42	11.29	1.58	9.52	1.33	26.36	3.69
VO 2046	Subgroup of eggplants	RAC	0.11	5.58	0.61	4.31	0.47	0.89	0.10	9.31	1.02	13.64	1.50	20.12	2.21
VL 0483	Lettuce, leaf, raw	RAC	2.2	0.53	1.17	0.36	0.79	0.16	0.35	6.21	13.66	1.90	4.18	6.05	13.31
VL 2832	Witloof chicory (sprouts)	RAC	0.02	0.03	0.00	0.02	0.00	0.01	0.00	0.36	0.01	0.06	0.00	0.35	0.01
VP 0061	Beans with pods (Phaseolus spp): (immature pods + succulent seeds)	RAC	0.2	0.68	0.14	NC	-	NC	-	0.39	0.08	0.22	0.04	0.49	0.10
VP 2062	Subgroup of succulent beans without pods (all commodities within this group)	RAC	0.03	5.07	0.15	1.02	0.03	0.49	0.01	1.78	0.05	1.19	0.04	8.57	0.26
VP 2063	Subgroup of succulent peas without pods	RAC	0.03	1.97	0.06	0.51	0.02	0.02	0.00	0.79	0.02	3.68	0.11	3.80	0.11
VD 0071	Beans, dry, raw (Phaseolus spp)	RAC	0.015	2.39	0.04	1.61	0.02	10.47	0.16	1.84	0.03	12.90	0.19	7.44	0.11
VD 0523	Broad bean, dry, raw (incl horse-bean, field bean) (Vicia faba)	RAC	0.015	1.27	0.02	0.01	0.00	0.12	0.00	2.49	0.04	0.23	0.00	5.54	0.08
VD 0527	Cowpea, dry, raw (Vigna sinensis, Dolichos sinensis)	RAC	0.015	0.05	0.00	NC	-	1.74	0.03	0.01	0.00	0.01	0.00	0.07	0.00
VD 0541	Soya bean, dry, raw (incl flour, incl paste, incl curd, incl sauce, excl oil)	RAC	0.0205	0.63	0.01	1.09	0.02	0.40	0.01	1.40	0.03	1.68	0.03	0.48	0.01
OR 0541	Soya oil, refined	PP	0.00041	12.99	0.01	10.43	0.00	3.63	0.00	13.10	0.01	10.70	0.00	13.10	0.01
-	Beans (dry) NES: including inter alia lablab or hyacinth bean (Dolichos spp.); jack or sword bean (Canavalia spp.); winged bean (Psophocarpus tetragonolobus); guar bean (Cyamopsis tetragonoloba); velvet bean (Stizolobium spp.); yam bean (Pachyrrhizus erosus)	RAC	0.015	1.70	0.03	0.01	0.00	3.00	0.05	1.80	0.03	1.64	0.02	1.33	0.02
VD 2066	Subgroup of dry peas, raw	RAC	0.058	9.09	0.53	3.35	0.19	1.06	0.06	9.48	0.55	15.11	0.88	10.58	0.61

Annex 3

FLUOPYRAM (243)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.01 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
VR 0577	Carrots, raw	RAC	0.09	9.51	0.86	30.78	2.77	0.37	0.03	8.75	0.79	2.80	0.25	6.10	0.55
VR 0596	Sugar beet, raw (incl sugar)	RAC	0.01	0.13	0.00	NC	-	0.08	0.00	0.66	0.01	0.47	0.00	88.94	0.89
VR 0589	Potato, raw (incl flour, incl frozen, incl starch, incl tapioca)	RAC	0.021	59.74	1.25	316.14	6.64	9.78	0.21	60.26	1.27	54.12	1.14	119.82	2.52
VS 0621	Asparagus, raw	RAC	0	0.01	0.00	0.01	0.00	0.01	0.00	0.03	0.00	0.07	0.00	0.21	0.00
VS 0620	Artichoke globe, raw	RAC	0.13	0.69	0.09	0.01	0.00	0.01	0.00	0.32	0.04	0.26	0.03	1.21	0.16
GC 0650	Rye, raw (incl flour)	RAC	0.19	0.13	0.02	19.38	3.68	0.10	0.02	0.12	0.02	0.03	0.01	2.15	0.41
GC 0653	Triticale, raw (incl flour)	RAC	0.19	NC	-	NC	-	NC	-	0.01	0.00	0.39	0.07	NC	-
GC 0654	Wheat, raw (incl bulgur, incl fermented beverages, incl germ, incl wholemeal bread, excl white flour products, excl white bread)	RAC	0.19	0.01	0.00	1.13	0.21	0.01	0.00	0.05	0.01	0.74	0.14	0.01	0.00
CF 1211	Wheat, white flour (incl white flour products: starch, gluten, macaroni, pastry)	PP	0.023	301.24	6.93	268.64	6.18	30.21	0.69	222.51	5.12	134.73	3.10	343.12	7.89
GC 0640	Barley, raw (incl malt extract, incl pot&pearled, incl flour & grits, incl beer, incl malt)	RAC	0.017	19.91	0.34	31.16	0.53	5.04	0.09	3.10	0.05	9.77	0.17	4.31	0.07
GC 0647	Oats, raw (incl rolled)	RAC	0.017	0.05	0.00	7.05	0.12	0.10	0.00	1.71	0.03	0.96	0.02	0.04	0.00
CM 0649 (GC 0649)	Rice, husked, dry (incl flour, incl oil, incl beverages, incl starch, excl polished)	REP	0.178	1.26	0.22	1.58	0.28	31.05	5.53	5.43	0.97	0.90	0.16	2.18	0.39
CM 1205	Rice polished, dry	PP	0.0676	34.21	2.31	10.39	0.70	41.72	2.82	82.38	5.57	150.24	10.16	70.47	4.76
GC 0645	Maize, raw (incl glucose & dextrose & isoglucose, incl beer, incl germ, excl flour, excl oil, excl starch)	RAC	0.01	0.84	0.01	0.24	0.00	1.56	0.02	0.46	0.00	2.44	0.02	13.13	0.13
CF 1255	Maize, flour (white flour and wholemeal flour)	PP	0.0085	22.72	0.19	35.61	0.30	87.27	0.74	34.92	0.30	46.71	0.40	49.12	0.42
OR 0645	Maize oil	PP	0.0058	0.96	0.01	0.85	0.00	0.29	0.00	5.42	0.03	0.42	0.00	2.10	0.01
GC 2090	Subgroup of Sweet Corns	RAC	0.01	0.14	0.00	0.94	0.01	5.70	0.06	2.61	0.03	1.94	0.02	0.22	0.00
TN 0085	Group of Tree nuts, raw (incl processed)	RAC	0.01	4.06	0.04	3.27	0.03	7.01	0.07	13.93	0.14	14.01	0.14	9.36	0.09
SO 0495	Rape seed, raw	RAC	0.33	0.02	0.01	NC	-	NC	-	0.01	0.00	0.75	0.25	0.01	0.00
OR 0495	Rape seed oil, edible	PP	0.23	0.35	0.08	0.44	0.10	0.19	0.04	0.97	0.22	3.28	0.75	0.77	0.18
SO 0702	Sunflower seed, raw	RAC	0.0066	0.09	0.00	0.33	0.00	0.09	0.00	0.24	0.00	0.02	0.00	0.01	0.00
OR 0702	Sunflower seed oil, edible	PP	0.00066	2.97	0.00	14.42	0.01	0.43	0.00	3.46	0.00	2.20	0.00	5.53	0.00
SO 0691	Cotton seed, raw (incl oil)	RAC	0.0585	20.53	1.20	9.80	0.57	6.42	0.38	4.73	0.28	7.14	0.42	18.68	1.09

FLUOPYRAM (243)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.01 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
SO 0697	Peanuts, nutmeat, raw (incl roasted, excl oil, excl butter)	RAC	0.033	0.46	0.02	1.21	0.04	6.64	0.22	2.52	0.08	1.25	0.04	1.83	0.06
OR 0697	Peanut oil, edible	PP	0.00033	0.36	0.00	0.01	0.00	2.57	0.00	0.07	0.00	2.29	0.00	0.36	0.00
-	Peanut butter	PP	0.0073	0.01	0.00	0.01	0.00	0.01	0.00	0.19	0.00	0.01	0.00	0.01	0.00
SB 0716	Coffee bean raw (incl roasted, incl instant coffee, incl substitutes)	RAC	0.01	1.36	0.01	3.59	0.04	1.44	0.01	5.18	0.05	2.02	0.02	1.70	0.02
HH 0722	Basil leaves, raw (incl dried)	RAC	19	0.14	2.66	0.26	4.94	0.16	3.04	0.38	7.22	NC	-	0.19	3.61
HS 0444	Peppers, chili, dried	PP	1.4	0.42	0.59	0.53	0.74	0.84	1.18	0.50	0.70	0.95	1.33	0.37	0.52
DH 1100	Hops, dry	RAC	10.35	0.01	0.10	0.04	0.41	0.01	0.10	0.01	0.10	NC	-	0.01	0.10
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.51	24.96	12.73	57.95	29.56	16.70	8.52	38.38	19.58	26.46	13.50	29.00	14.79
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.67	6.24	4.18	14.49	9.71	4.18	2.80	9.60	6.43	6.62	4.43	7.25	4.86
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.67	3.29	2.20	6.14	4.11	0.82	0.55	1.57	1.05	2.23	1.49	1.07	0.72
MO 0105	Edible offal (mammalian), raw	RAC	3.8	4.79	18.20	9.68	36.78	2.97	11.29	5.49	20.86	3.84	14.59	5.03	19.11
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.48	289.65	139.03	485.88	233.22	26.92	12.92	239.03	114.73	199.91	95.96	180.53	86.65
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0.19	13.17	2.50	26.78	5.09	7.24	1.37	116.71	22.18	22.54	4.28	32.09	6.10
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0.28	1.46	0.41	2.98	0.83	0.80	0.23	12.97	3.63	2.50	0.70	3.57	1.00
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.28	0.10	0.03	0.10	0.03	NC	-	0.10	0.03	0.10	0.03	0.10	0.03
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.88	0.12	0.11	0.12	0.11	0.11	0.10	5.37	4.73	0.24	0.21	0.10	0.09
PE 0112	Eggs, raw, (incl dried)	RAC	0.46	7.84	3.61	23.08	10.62	2.88	1.32	14.89	6.85	9.81	4.51	14.83	6.82
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-
Total intake (ug/person)=				244.4		408.7		87.2		311.7		193.0		298.7	
Bodyweight per region (kg bw) =				60		60		60		60		60		60	
ADI (ug/person)=				600		600		600		600		600		600	
%ADI=				40.7%		68.1%		14.5%		51.9%		32.2%		49.8%	
Rounded %ADI=				40%		70%		10%		50%		30%		50%	

Annex 3

FLUOPYRAM (243)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.01 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
FC 0002	Subgroup of Lemons and limes, raw (incl lemon juice) (incl kumquat commodities)	RAC	0.325	10.12	3.29	15.69	5.10	2.88	0.94	12.30	4.00	22.32	7.25	6.59	2.14
FC 0003	Subgroup of Mandarins, raw (incl mandarin juice)	RAC	0.15	12.42	1.86	14.99	2.25	16.08	2.41	10.78	1.62	9.94	1.49	NC	-
FC 0004	Subgroup of Oranges, sweet, sour, raw (incl orange juice)	RAC	0.15	83.66	12.55	27.64	4.15	7.37	1.11	67.80	10.17	43.97	6.60	187.74	28.16
FC 0005	Subgroup of Pummelo and grapefruits, raw (incl grapefruit juice)	RAC	0.14	8.21	1.15	4.60	0.64	0.64	0.09	5.85	0.82	19.98	2.80	368.86	51.64
FP 0009	Group of Pome fruits, raw (incl apple cider, excl apple juice)	RAC	0.135	51.09	6.90	65.40	8.83	42.71	5.77	45.29	6.11	62.51	8.44	7.74	1.04
JF 0226	Apple juice, single strength (incl. concentrated)	PP	0.01	14.88	0.15	11.98	0.12	0.15	0.00	9.98	0.10	30.32	0.30	3.47	0.03
FS 0013	Subgroup of Cherries, raw	RAC	0.57	1.40	0.80	4.21	2.40	0.04	0.02	2.93	1.67	1.50	0.86	NC	-
FS 0014	Subgroup of Plums, raw (incl dried plums)	RAC	0.13	5.55	0.72	4.37	0.57	6.08	0.79	3.66	0.48	3.93	0.51	0.46	0.06
FS 2001	Subgroup of peaches, raw (incl dried apricots)	RAC	0.22	13.03	2.87	16.29	3.58	8.29	1.82	12.95	2.85	5.35	1.18	0.04	0.01
FB 2005	Subgroup of Caneberries, raw	RAC	0.83	0.56	0.46	1.43	1.19	0.14	0.12	1.23	1.02	1.14	0.95	0.01	0.01
FB 2006	Subgroup of Bush berries, raw (including processed)	RAC	1.15	1.31	1.51	5.50	6.33	0.01	0.01	2.57	2.96	0.82	0.94	2.15	2.47
FB 0269	Grapes, raw (incl must, excl dried, excl juice, excl wine)	RAC	0.58	6.48	3.76	11.31	6.56	5.21	3.02	9.50	5.51	4.66	2.70	0.78	0.45
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	1.68	3.09	5.19	1.51	2.54	0.03	0.05	1.38	2.32	4.26	7.16	0.42	0.71
JF 0269	Grape juice (from wine grapes)	PP	0.012	0.56	0.01	1.96	0.02	0.02	0.00	2.24	0.03	2.27	0.03	0.34	0.00
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.1	88.93	8.89	62.41	6.24	1.84	0.18	25.07	2.51	61.17	6.12	5.84	0.58
FB 0275	Strawberry, raw	RAC	0.025	4.49	0.11	5.66	0.14	0.02	0.00	6.63	0.17	5.75	0.14	0.05	0.00
FI 0327	Banana, raw (incl plantains) (incl dried)	RAC	0.175	25.76	4.51	23.65	4.14	23.83	4.17	24.37	4.26	19.43	3.40	101.55	17.77
FI 0345	Mango, raw (incl canned mango, incl mango juice)	RAC	0.02	1.80	0.04	0.63	0.01	10.05	0.20	1.07	0.02	3.52	0.07	16.44	0.33
VA 0381	Garlic, raw	RAC	0.01	0.98	0.01	1.49	0.01	12.88	0.13	3.74	0.04	2.05	0.02	1.14	0.01
-	Onions, dry, raw	RAC	0.01	19.69	0.20	29.83	0.30	24.64	0.25	31.35	0.31	9.72	0.10	12.59	0.13
VA 0384	Leek, raw	RAC	0.01	4.01	0.04	4.41	0.04	0.72	0.01	0.54	0.01	16.41	0.16	0.03	0.00
-	Onions, green, raw	RAC	5.1	1.55	7.91	0.74	3.77	1.05	5.36	3.74	19.07	0.94	4.79	6.45	32.90
VB 0400	Broccoli, raw	RAC	0.05	4.24	0.21	1.76	0.09	NC	-	0.51	0.03	3.79	0.19	0.26	0.01
VB 0404	Cauliflower, raw	RAC	0.01	5.27	0.05	5.01	0.05	NC	-	2.70	0.03	5.57	0.06	0.49	0.00
VB 0402	Brussels sprouts, raw	RAC	0.06	2.24	0.13	2.67	0.16	6.23	0.37	0.32	0.02	4.19	0.25	2.58	0.15

FLUOPYRAM (243)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.01 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
VB 0041	Cabbages, head, raw	RAC	0.01	8.97	0.09	27.12	0.27	1.44	0.01	24.96	0.25	4.55	0.05	11.23	0.11
VC 0424	Cucumber, raw	RAC	0.11	6.72	0.74	11.03	1.21	32.10	3.53	15.10	1.66	4.05	0.45	9.57	1.05
VO 0448	Tomato, raw	RAC	0.11	32.13	3.53	51.27	5.64	34.92	3.84	73.37	8.07	15.15	1.67	8.88	0.98
-	Tomato, canned (& peeled)	PP	0.023	7.57	0.17	2.66	0.06	0.30	0.01	0.97	0.02	7.31	0.17	0.41	0.01
-	Tomato, paste (i.e. concentrated tomato sauce/puree)	PP	0.08	4.96	0.40	3.20	0.26	0.15	0.01	1.61	0.13	6.88	0.55	0.52	0.04
JF 0448	Tomato, juice (single strength, incl concentrated)	PP	0.04	0.80	0.03	0.07	0.00	0.05	0.00	0.61	0.02	0.40	0.02	0.08	0.00
VO 0051	Subgroup of peppers, raw (incl dried sweet peppers, excl dried chilipeppers), excl okra	RAC	0.14	6.39	0.89	15.53	2.17	19.09	2.67	10.36	1.45	8.29	1.16	4.53	0.63
VO 2046	Subgroup of eggplants	RAC	0.11	1.01	0.11	1.69	0.19	21.37	2.35	3.00	0.33	1.40	0.15	NC	-
VL 0483	Lettuce, leaf, raw	RAC	2.2	14.50	31.90	11.76	25.87	13.14	28.91	19.50	42.90	4.81	10.58	2.23	4.91
VL 2832	Witloof chicory (sprouts)	RAC	0.02	1.50	0.03	0.95	0.02	NC	-	1.84	0.04	0.65	0.01	0.13	0.00
VP 0061	Beans with pods (Phaseolus spp): (immature pods + succulent seeds)	RAC	0.2	5.07	1.01	0.83	0.17	0.17	0.03	3.70	0.74	NC	-	NC	-
VP 2062	Subgroup of succulent beans without pods (all commodities within this group)	RAC	0.03	2.42	0.07	6.09	0.18	4.33	0.13	2.09	0.06	18.99	0.57	0.17	0.01
VP 2063	Subgroup of succulent peas without pods	RAC	0.03	10.72	0.32	1.99	0.06	2.72	0.08	4.26	0.13	4.23	0.13	NC	-
VD 0071	Beans, dry, raw (Phaseolus spp)	RAC	0.015	1.51	0.02	1.50	0.02	1.90	0.03	5.11	0.08	1.36	0.02	23.43	0.35
VD 0523	Broad bean, dry, raw (incl horse-bean, field bean) (Vicia faba)	RAC	0.015	0.02	0.00	0.01	0.00	1.16	0.02	0.40	0.01	NC	-	0.06	0.00
VD 0527	Cowpea, dry, raw (Vigna sinensis, Dolichos sinensis)	RAC	0.015	NC	-	NC	-	0.16	0.00	0.01	0.00	NC	-	NC	-
VD 0541	Soya bean, dry, raw (incl flour, incl paste, incl curd, incl sauce, excl oil)	RAC	0.0205	0.47	0.01	0.77	0.02	9.12	0.19	8.05	0.17	0.04	0.00	6.06	0.12
OR 0541	Soya oil, refined	PP	0.00041	19.06	0.01	21.06	0.01	5.94	0.00	33.78	0.01	40.05	0.02	13.39	0.01
-	Beans (dry) NES: including inter alia lablab or hyacinth bean (Dolichos spp.); jack or sword bean (Canavalia spp.); winged bean (Psophocarpus tetragonolobus); guar bean (Cyamopsis tetragonoloba); velvet bean (Stizolobium spp.); yam bean (Pachyrrhizus erosus)	RAC	0.015	0.01	0.00	NC	-	0.57	0.01	0.11	0.00	0.16	0.00	0.94	0.01
VD 2066	Subgroup of dry peas, raw	RAC	0.058	5.01	0.29	3.76	0.22	1.82	0.11	3.44	0.20	3.49	0.20	5.15	0.30
VR 0577	Carrots, raw	RAC	0.09	26.26	2.36	27.13	2.44	10.07	0.91	16.49	1.48	44.69	4.02	8.75	0.79
VR 0596	Sugar beet, raw (incl sugar)	RAC	0.01	0.01	0.00	NC	-	0.01	0.00	0.01	0.00	NC	-	NC	-

Annex 3

FLUOPYRAM (243)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.01 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
VR 0589	Potato, raw (incl flour, incl frozen, incl starch, incl tapioca)	RAC	0.021	225.03	4.73	234.24	4.92	71.48	1.50	177.55	3.73	234.55	4.93	37.71	0.79
VS 0621	Asparagus, raw	RAC	0	0.84	0.00	2.08	0.00	7.11	0.00	1.01	0.00	1.69	0.00	0.04	0.00
VS 0620	Artichoke globe, raw	RAC	0.13	0.98	0.13	3.65	0.47	0.07	0.01	1.67	0.22	0.26	0.03	NC	-
GC 0650	Rye, raw (incl flour)	RAC	0.19	3.21	0.61	35.38	6.72	0.21	0.04	6.50	1.24	1.49	0.28	NC	-
GC 0653	Triticale, raw (incl flour)	RAC	0.19	0.01	0.00	0.17	0.03	0.29	0.06	0.01	0.00	NC	-	NC	-
GC 0654	Wheat, raw (incl bulgur, incl fermented beverages, incl germ, incl wholemeal bread, excl white flour products, excl white bread)	RAC	0.19	1.00	0.19	0.11	0.02	0.05	0.01	0.84	0.16	0.06	0.01	0.06	0.01
CF 1211	Wheat, white flour (incl white flour products: starch, gluten, macaroni, pastry)	PP	0.023	198.08	4.56	193.03	4.44	106.24	2.44	185.09	4.26	168.67	3.88	131.59	3.03
GC 0640	Barley, raw (incl malt extract, incl pot&pearled, incl flour & grits, incl beer, incl malt)	RAC	0.017	36.18	0.62	53.45	0.91	9.39	0.16	35.25	0.60	46.68	0.79	15.92	0.27
GC 0647	Oats, raw (incl rolled)	RAC	0.017	7.50	0.13	6.26	0.11	0.15	0.00	4.87	0.08	3.16	0.05	2.98	0.05
CM 0649 (GC 0649)	Rice, husked, dry (incl flour, incl oil, incl beverages, incl starch, excl polished)	REP	0.178	3.70	0.66	2.11	0.38	1.51	0.27	1.75	0.31	0.29	0.05	5.12	0.91
CM 1205	Rice polished, dry	PP	0.0676	13.38	0.90	10.80	0.73	262.08	17.72	57.16	3.86	12.83	0.87	62.78	4.24
GC 0645	Maize, raw (incl glucose & dextrose & isoglucose, incl beer, incl germ, excl flour, excl oil, excl starch)	RAC	0.01	0.10	0.00	9.93	0.10	1.40	0.01	10.26	0.10	0.33	0.00	0.05	0.00
CF 1255	Maize, flour (white flour and wholemeal flour)	PP	0.0085	14.27	0.12	12.86	0.11	19.71	0.17	12.55	0.11	4.21	0.04	52.30	0.44
OR 0645	Maize oil	PP	0.0058	0.90	0.01	0.47	0.00	0.15	0.00	3.01	0.02	1.86	0.01	0.36	0.00
GC 2090	Subgroup of Sweet Corns	RAC	0.01	11.43	0.11	3.71	0.04	0.74	0.01	13.63	0.14	3.07	0.03	1.50	0.02
TN 0085	Group of Tree nuts, raw (incl processed)	RAC	0.01	8.52	0.09	8.94	0.09	15.09	0.15	9.60	0.10	14.57	0.15	26.26	0.26
SO 0495	Rape seed, raw	RAC	0.33	NC	-	NC	-	0.01	0.00	NC	-	NC	-	NC	-
OR 0495	Rape seed oil, edible	PP	0.23	12.52	2.88	7.63	1.75	3.00	0.69	6.01	1.38	NC	-	NC	-
SO 0702	Sunflower seed, raw	RAC	0.0066	0.01	0.00	1.32	0.01	0.03	0.00	1.17	0.01	NC	-	0.02	0.00
OR 0702	Sunflower seed oil, edible	PP	0.00066	9.50	0.01	11.37	0.01	0.49	0.00	5.15	0.00	2.63	0.00	2.80	0.00
SO 0691	Cotton seed, raw (incl oil)	RAC	0.0585	10.71	0.63	4.23	0.25	7.19	0.42	7.54	0.44	5.66	0.33	2.38	0.14
SO 0697	Peanuts, nutmeat, raw (incl roasted, excl oil, excl butter)	RAC	0.033	3.19	0.11	2.19	0.07	5.36	0.18	4.82	0.16	1.40	0.05	1.06	0.03

FLUOPYRAM (243)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.01 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
OR 0697	Peanut oil, edible	PP	0.00033	1.02	0.00	0.23	0.00	1.81	0.00	0.42	0.00	5.23	0.00	0.01	0.00
-	Peanut butter	PP	0.0073	0.07	0.00	0.04	0.00	0.01	0.00	0.03	0.00	0.15	0.00	0.75	0.01
SB 0716	Coffee bean raw (incl roasted, incl instant coffee, incl substitutes)	RAC	0.01	10.90	0.11	12.44	0.12	0.77	0.01	9.48	0.09	22.07	0.22	8.15	0.08
HH 0722	Basil leaves, raw (incl dried)	RAC	19	0.52	9.88	0.05	0.95	3.23	61.37	0.18	3.42	0.12	2.28	0.27	5.13
HS 0444	Peppers, chili, dried	PP	1.4	0.11	0.15	0.21	0.29	0.36	0.50	0.21	0.29	0.25	0.35	0.15	0.21
DH 1100	Hops, dry	RAC	10.35	NC	-	NC	-	0.02	0.21	0.02	0.21	NC	-	NC	-
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.51	112.02	57.13	120.71	61.56	63.46	32.36	88.99	45.39	96.24	49.08	41.02	20.92
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.67	28.01	18.76	30.18	20.22	15.86	10.63	22.25	14.91	24.06	16.12	10.25	6.87
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.67	6.44	4.31	15.51	10.39	3.79	2.54	8.29	5.55	18.44	12.35	8.00	5.36
MO 0105	Edible offal (mammalian), raw	RAC	3.8	15.17	57.65	5.19	19.72	6.30	23.94	6.78	25.76	3.32	12.62	3.17	12.05
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.48	388.92	186.68	335.88	161.22	49.15	23.59	331.25	159.00	468.56	224.91	245.45	117.82
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0.19	66.38	12.61	48.47	9.21	21.58	4.10	78.41	14.90	48.04	9.13	76.01	14.44
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0.28	7.38	2.07	5.39	1.51	2.40	0.67	8.71	2.44	5.34	1.49	8.45	2.36
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.28	0.10	0.03	0.10	0.03	NC	-	0.10	0.03	0.71	0.20	NC	-
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.88	0.33	0.29	0.72	0.63	0.27	0.24	0.35	0.31	0.80	0.70	NC	-
PE 0112	Eggs, raw, (incl dried)	RAC	0.46	25.84	11.89	29.53	13.58	28.05	12.90	33.19	15.27	36.44	16.76	8.89	4.09
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-
Total intake (ug/person)=				483.3		418.7		266.6		428.3		434.0		347.5	
Bodyweight per region (kg bw) =				60		60		55		60		60		60	
ADI (ug/person)=				600		600		550		600		600		600	
%ADI=				80.6%		69.8%		48.5%		71.4%		72.3%		57.9%	
Rounded %ADI=				80%		70%		50%		70%		70%		60%	

FLUOPYRAM (243)

International Estimated Daily Intake (IEDI)

ADI = 0 - 0.01 mg/kg bw

Annex 3

Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
FC 0002	Subgroup of Lemons and limes, raw (incl lemon juice) (incl kumquat commodities)	RAC	0.325	18.97	6.17	0.97	0.32	6.23	2.02	0.09	0.03	3.35	1.09
FC 0003	Subgroup of Mandarins, raw (incl mandarin juice)	RAC	0.15	0.16	0.02	0.27	0.04	9.06	1.36	0.01	0.00	0.02	0.00
FC 0004	Subgroup of Oranges, sweet, sour, raw (incl orange juice)	RAC	0.15	1.34	0.20	1.65	0.25	40.03	6.00	0.33	0.05	1.76	0.26
FC 0005	Subgroup of Pummelo and grapefruits, raw (incl grapefruit juice)	RAC	0.14	0.68	0.10	0.05	0.01	3.21	0.45	0.01	0.00	NC	-
FP 0009	Group of Pome fruits, raw (incl apple cider, excl apple juice)	RAC	0.135	68.85	9.29	10.93	1.48	70.82	9.56	189.78	25.62	19.56	2.64
JF 0226	Apple juice, single strength (incl. concentrated)	PP	0.01	0.03	0.00	0.10	0.00	7.19	0.07	0.03	0.00	NC	-
FS 0013	Subgroup of Cherries, raw	RAC	0.57	0.01	0.01	0.01	0.01	5.96	3.40	0.01	0.01	NC	-
FS 0014	Subgroup of Plums, raw (incl dried plums)	RAC	0.13	0.07	0.01	0.02	0.00	16.65	2.16	0.01	0.00	NC	-
FS 2001	Subgroup of peaches, raw (incl dried apricots)	RAC	0.22	0.02	0.00	0.01	0.00	10.76	2.37	0.01	0.00	NC	-
FB 2005	Subgroup of Caneberries, raw	RAC	0.83	0.01	0.01	7.30	6.06	2.29	1.90	0.01	0.01	NC	-
FB 2006	Subgroup of Bush berries, raw (including processed)	RAC	1.15	0.82	0.94	4.05	4.66	5.94	6.83	0.43	0.49	2.66	3.06
FB 0269	Grapes, raw (incl must, excl dried, excl juice, excl wine)	RAC	0.58	0.14	0.08	0.36	0.21	15.33	8.89	0.01	0.01	0.28	0.16
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	1.68	0.01	0.02	0.13	0.22	1.06	1.78	0.01	0.02	0.03	0.05
JF 0269	Grape juice (from wine grapes)	PP	0.012	0.01	0.00	0.01	0.00	0.41	0.00	0.01	0.00	NC	-
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.1	0.31	0.03	0.23	0.02	60.43	6.04	0.52	0.05	31.91	3.19
FB 0275	Strawberry, raw	RAC	0.025	0.01	0.00	0.01	0.00	3.35	0.08	0.01	0.00	0.01	0.00
FI 0327	Banana, raw (incl plantains) (incl dried)	RAC	0.175	44.80	7.84	118.17	20.68	25.25	4.42	454.49	79.54	310.23	54.29
FI 0345	Mango, raw (incl canned mango, incl mango juice)	RAC	0.02	12.25	0.25	6.83	0.14	0.76	0.02	0.01	0.00	20.12	0.40
VA 0381	Garlic, raw	RAC	0.01	0.82	0.01	2.06	0.02	3.79	0.04	0.03	0.00	0.29	0.00
-	Onions, dry, raw	RAC	0.01	9.01	0.09	20.24	0.20	30.90	0.31	9.61	0.10	2.11	0.02
VA 0384	Leek, raw	RAC	0.01	0.02	0.00	1.44	0.01	1.22	0.01	0.01	0.00	NC	-
-	Onions, green, raw	RAC	5.1	1.43	7.29	0.05	0.26	0.20	1.02	NC	-	6.30	32.13
VB 0400	Broccoli, raw	RAC	0.05	0.01	0.00	0.01	0.00	2.13	0.11	0.01	0.00	NC	-
VB 0404	Cauliflower, raw	RAC	0.01	0.01	0.00	0.01	0.00	2.73	0.03	0.01	0.00	NC	-
VB 0402	Brussels sprouts, raw	RAC	0.06	0.88	0.05	0.69	0.04	2.89	0.17	0.01	0.00	NC	-
VB 0041	Cabbages, head, raw	RAC	0.01	3.82	0.04	2.99	0.03	49.16	0.49	0.01	0.00	NC	-

FLUOPYRAM (243)

International Estimated Daily Intake (IEDI)

ADI = 0 - 0.01 mg/kg bw

Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
VC 0424	Cucumber, raw	RAC	0.11	0.68	0.07	1.81	0.20	10.40	1.14	0.01	0.00	0.04	0.00
VO 0448	Tomato, raw	RAC	0.11	12.99	1.43	4.79	0.53	58.40	6.42	0.92	0.10	0.09	0.01
-	Tomato, canned (& peeled)	PP	0.023	0.07	0.00	0.08	0.00	2.42	0.06	0.07	0.00	NC	-
-	Tomato, paste (i.e. concentrated tomato sauce/puree)	PP	0.08	0.58	0.05	0.22	0.02	2.21	0.18	0.24	0.02	3.10	0.25
JF 0448	Tomato, juice (single strength, incl concentrated)	PP	0.04	0.05	0.00	0.01	0.00	0.42	0.02	0.01	0.00	0.02	0.00
VO 0051	Subgroup of peppers, raw (incl dried sweet peppers, excl dried chili peppers), excl okra	RAC	0.14	8.97	1.26	14.13	1.98	25.14	3.52	0.91	0.13	NC	-
VO 2046	Subgroup of eggplants	RAC	0.11	1.31	0.14	8.26	0.91	3.95	0.43	0.01	0.00	NC	-
VL 0483	Lettuce, leaf, raw	RAC	2.2	0.29	0.64	0.03	0.07	6.71	14.76	0.01	0.02	NC	-
VL 2832	Witloof chicory (sprouts)	RAC	0.02	0.02	0.00	0.01	0.00	0.01	0.00	0.01	0.00	NC	-
VP 0061	Beans with pods (Phaseolus spp): (immature pods + succulent seeds)	RAC	0.2	NC	-	NC	-	NC	-	NC	-	NC	-
VP 2062	Subgroup of succulent beans without pods (all commodities within this group)	RAC	0.03	0.37	0.01	3.14	0.09	4.88	0.15	0.01	0.00	NC	-
VP 2063	Subgroup of succulent peas without pods	RAC	0.03	0.21	0.01	0.02	0.00	5.51	0.17	0.02	0.00	NC	-
VD 0071	Beans, dry, raw (Phaseolus spp)	RAC	0.015	7.11	0.11	2.33	0.03	3.76	0.06	44.70	0.67	3.27	0.05
VD 0523	Broad bean, dry, raw (incl horse-bean, field bean) (Vicia faba)	RAC	0.015	3.70	0.06	0.03	0.00	0.17	0.00	0.01	0.00	NC	-
VD 0527	Cowpea, dry, raw (Vigna sinensis, Dolichos sinensis)	RAC	0.015	12.77	0.19	0.99	0.01	0.01	0.00	4.33	0.06	NC	-
VD 0541	Soya bean, dry, raw (incl flour, incl paste, incl curd, incl sauce, excl oil)	RAC	0.0205	2.89	0.06	0.21	0.00	0.48	0.01	3.16	0.06	0.26	0.01
OR 0541	Soya oil, refined	PP	0.00041	2.32	0.00	2.54	0.00	18.70	0.01	2.51	0.00	6.29	0.00
-	Beans (dry) NES: including inter alia lablab or hyacinth bean (Dolichos spp.); jack or sword bean (Canavalia spp.); winged bean (Psophocarpus tetragonolobus); guar bean (Cyamopsis tetragonoloba); velvet bean (Stizolobium spp.); yam bean (Pachyrrhizus erosus)	RAC	0.015	2.54	0.04	1.77	0.03	0.03	0.00	0.03	0.00	3.99	0.06
VD 2066	Subgroup of dry peas, raw	RAC	0.058	4.43	0.26	11.36	0.66	4.22	0.24	9.36	0.54	1.21	0.07
VR 0577	Carrots, raw	RAC	0.09	2.07	0.19	3.00	0.27	25.29	2.28	0.05	0.00	NC	-
VR 0596	Sugar beet, raw (incl sugar)	RAC	0.01	3.93	0.04	1.68	0.02	NC	-	NC	-	36.12	0.36
VR 0589	Potato, raw (incl flour, incl frozen, incl starch, incl tapioca)	RAC	0.021	23.96	0.50	13.56	0.28	213.41	4.48	104.35	2.19	8.56	0.18

Annex 3

FLUOPYRAM (243)

International Estimated Daily Intake (IEDI)

ADI = 0 - 0.01 mg/kg bw

Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
VS 0621	Asparagus, raw	RAC	0	0.01	0.00	0.01	0.00	0.17	0.00	0.01	0.00	NC	-
VS 0620	Artichoke globe, raw	RAC	0.13	0.01	0.00	NC	-	0.08	0.01	0.01	0.00	NC	-
GC 0650	Rye, raw (incl flour)	RAC	0.19	0.03	0.01	0.01	0.00	13.95	2.65	0.01	0.00	0.88	0.17
GC 0653	Triticale, raw (incl flour)	RAC	0.19	0.01	0.00	NC	-	NC	-	NC	-	NC	-
GC 0654	Wheat, raw (incl bulgur, incl fermented beverages, incl germ, incl wholemeal bread, excl white flour products, excl white bread)	RAC	0.19	0.05	0.01	0.01	0.00	0.05	0.01	0.01	0.00	0.97	0.18
CF 1211	Wheat, white flour (incl white flour products: starch, gluten, macaroni, pastry)	PP	0.023	44.78	1.03	86.96	2.00	214.05	4.92	20.31	0.47	103.60	2.38
GC 0640	Barley, raw (incl malt extract, incl pot&pearled, incl flour & grits, incl beer, incl malt)	RAC	0.017	11.58	0.20	2.33	0.04	46.71	0.79	3.72	0.06	16.26	0.28
GC 0647	Oats, raw (incl rolled)	RAC	0.017	0.37	0.01	0.07	0.00	2.79	0.05	0.10	0.00	NC	-
CM 0649 (GC 0649)	Rice, husked, dry (incl flour, incl oil, incl beverages, incl starch, excl polished)	REP	0.178	13.58	2.42	4.29	0.76	2.17	0.39	0.01	0.00	8.84	1.57
CM 1205	Rice polished, dry	PP	0.0676	30.20	2.04	218.34	14.76	12.77	0.86	15.24	1.03	51.35	3.47
GC 0645	Maize, raw (incl glucose & dextrose & isoglucose, incl beer, incl germ, excl flour, excl oil, excl starch)	RAC	0.01	0.55	0.01	0.51	0.01	3.26	0.03	7.96	0.08	NC	-
CF 1255	Maize, flour (white flour and wholemeal flour)	PP	0.0085	94.34	0.80	8.09	0.07	28.03	0.24	55.94	0.48	28.07	0.24
OR 0645	Maize oil	PP	0.0058	0.33	0.00	0.07	0.00	0.81	0.00	0.01	0.00	NC	-
GC 2090	Subgroup of Sweet Corns	RAC	0.01	3.63	0.04	20.50	0.21	8.78	0.09	0.02	0.00	0.17	0.00
TN 0085	Group of Tree nuts, raw (incl processed)	RAC	0.01	4.39	0.04	135.53	1.36	6.11	0.06	0.72	0.01	317.74	3.18
SO 0495	Rape seed, raw	RAC	0.33	NC	-	0.01	0.00	NC	-	NC	-	NC	-
OR 0495	Rape seed oil, edible	PP	0.23	0.07	0.02	0.03	0.01	4.62	1.06	0.03	0.01	NC	-
SO 0702	Sunflower seed, raw	RAC	0.0066	0.02	0.00	0.01	0.00	0.03	0.00	2.23	0.01	NC	-
OR 0702	Sunflower seed oil, edible	PP	0.00066	0.37	0.00	0.09	0.00	12.98	0.01	4.01	0.00	0.20	0.00
SO 0691	Cotton seed, raw (incl oil)	RAC	0.0585	8.14	0.48	0.32	0.02	2.84	0.17	2.69	0.16	0.97	0.06
SO 0697	Peanuts, nutmeat, raw (incl roasted, excl oil, excl butter)	RAC	0.033	7.14	0.24	0.42	0.01	1.83	0.06	6.22	0.21	0.53	0.02
OR 0697	Peanut oil, edible	PP	0.00033	5.02	0.00	0.05	0.00	0.17	0.00	0.29	0.00	NC	-
-	Peanut butter	PP	0.0073	0.01	0.00	0.03	0.00	0.05	0.00	NC	-	NC	-
SB 0716	Coffee bean raw (incl roasted, incl instant coffee, incl substitutes)	RAC	0.01	0.95	0.01	1.32	0.01	11.64	0.12	2.96	0.03	14.73	0.15

FLUOPYRAM (243)

International Estimated Daily Intake (IEDI)

ADI = 0 - 0.01 mg/kg bw

Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
HH 0722	Basil leaves, raw (incl dried)	RAC	19	0.25	4.75	0.18	3.42	0.13	2.47	0.17	3.23	0.33	6.27
HS 0444	Peppers, chili, dried	PP	1.4	0.58	0.81	1.27	1.78	1.21	1.69	0.12	0.17	NC	-
DH 1100	Hops, dry	RAC	10.35	NC	-	NC	-	0.04	0.41	NC	-	NC	-
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.51	23.34	11.91	40.71	20.76	97.15	49.55	18.06	9.21	57.71	29.43
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.67	5.84	3.91	10.18	6.82	24.29	16.27	4.52	3.03	14.43	9.67
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.67	1.05	0.70	1.14	0.76	18.69	12.52	0.94	0.63	3.12	2.09
MO 0105	Edible offal (mammalian), raw	RAC	3.8	4.64	17.63	1.97	7.49	10.01	38.04	3.27	12.43	3.98	15.12
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.48	108.75	52.20	70.31	33.75	436.11	209.33	61.55	29.54	79.09	37.96
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0.19	3.53	0.67	10.83	2.06	51.36	9.76	4.53	0.86	50.00	9.50
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0.28	0.39	0.11	1.20	0.34	5.71	1.60	0.50	0.14	5.56	1.56
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.28	NC	-	NC	-	0.32	0.09	NC	-	NC	-
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.88	0.10	0.09	0.70	0.62	0.97	0.85	0.10	0.09	NC	-
PE 0112	Eggs, raw, (incl dried)	RAC	0.46	3.84	1.77	4.41	2.03	27.25	12.54	1.13	0.52	7.39	3.40
-	-	-		-	-	-	-	-	-	-	-	-	-

Total intake (ug/person)=

139.4

138.8

460.1

172.1

225.0

Bodyweight per region (kg bw) =

60

60

60

60

60

ADI (ug/person)=

600

600

600

600

600

%ADI=

23.2%

23.1%

76.7%

28.7%

37.5%

Rounded %ADI=

20%

20%

80%

30%

40%

Annex 3

IMAZALIL (110)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
FC 0001	Group of Citrus fruit, raw (incl citrus fruit juice, excl kumquat commodities)	RAC	0.07	32.55	2.28	16.24	1.14	14.04	0.98	90.04	6.30	33.92	2.37	96.78	6.77
JF 0001	Group of Citrus fruit, juice	PP	0.34	1.30	0.44	2.37	0.81	0.22	0.07	13.88	4.72	0.75	0.26	2.63	0.89
FC 0303	Kumquats, raw (incl juice)	RAC	3.4	2.36	8.02	0.27	0.92	3.19	10.85	14.44	49.10	1.66	5.64	1.71	5.81
FI 0327	Banana, raw (incl plantains) (incl dried)	RAC	0.05	5.23	0.26	6.94	0.35	99.45	4.97	32.47	1.62	48.30	2.42	24.70	1.24
VO 0448	Tomato, raw (incl juice, incl paste, incl canned)	RAC	0.13	51.75	6.73	81.80	10.63	16.99	2.21	102.02	13.26	26.32	3.42	214.77	27.92
VR 0589	Potato, raw (incl flour, incl frozen, incl starch, incl tapioca)	RAC	2.2	59.74	131.43	316.14	695.51	9.78	21.52	60.26	132.57	54.12	119.06	119.82	263.60
GC 0653	Triticale, raw (incl flour)	RAC	0	NC	-	NC	-	NC	-	0.01	0.00	0.39	0.00	NC	-
GC 0654	Wheat, raw (incl bulgur, incl fermented beverages, incl germ, incl wholemeal bread, incl white flour products, incl white bread)	RAC	0	381.15	0.00	341.55	0.00	38.35	0.00	281.89	0.00	172.83	0.00	434.07	0.00
GC 0640	Barley, raw (incl malt extract, incl pot&pearled, incl flour & grits, incl beer, incl malt)	RAC	0	19.91	0.00	31.16	0.00	5.04	0.00	3.10	0.00	9.77	0.00	4.31	0.00
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.4	24.96	9.98	57.95	23.18	16.70	6.68	38.38	15.35	26.46	10.59	29.00	11.60
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.4	6.24	2.50	14.49	5.80	4.18	1.67	9.60	3.84	6.62	2.65	7.25	2.90
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.04	3.29	0.13	6.14	0.25	0.82	0.03	1.57	0.06	2.23	0.09	1.07	0.04
MO 0105	Edible offal (mammalian), raw	RAC	0.42	4.79	2.01	9.68	4.07	2.97	1.25	5.49	2.31	3.84	1.61	5.03	2.11
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.04	289.65	11.59	485.88	19.44	26.92	1.08	239.03	9.56	199.91	8.00	180.53	7.22
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0.04	13.17	0.53	26.78	1.07	7.24	0.29	116.71	4.67	22.54	0.90	32.09	1.28
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0.04	1.46	0.06	2.98	0.12	0.80	0.03	12.97	0.52	2.50	0.10	3.57	0.14
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.04	0.10	0.00	0.10	0.00	NC	-	0.10	0.00	0.10	0.00	0.10	0.00
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.04	0.12	0.00	0.12	0.00	0.11	0.00	5.37	0.21	0.24	0.01	0.10	0.00
PE 0112	Eggs, raw, (incl dried)	RAC	0.02	7.84	0.16	23.08	0.46	2.88	0.06	14.89	0.30	9.81	0.20	14.83	0.30
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-
Total intake (ug/person)=				176.1		763.7		51.7		244.4		157.3		331.8	
Bodyweight per region (kg bw) =				60		60		60		60		60		60	
ADI (ug/person)=				1800		1800		1800		1800		1800		1800	
%ADI=				9.8%		42.4%		2.9%		13.6%		8.7%		18.4%	

IMAZALIL (110)			International Estimated Daily Intake (IEDI)								ADI = 0 - 0.03 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day			Intake as ug/person/day									
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake	
	Rounded %ADI=				10%		40%		3%		10%		9%		20%	

IMAZALIL (110)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
FC 0001	Group of Citrus fruit, raw (incl citrus fruit juice, excl kumquat commodities)	RAC	0.07	109.75	7.68	57.06	3.99	25.01	1.75	95.27	6.67	79.17	5.54	561.83	39.33
JF 0001	Group of Citrus fruit, juice	PP	0.34	36.84	12.53	3.75	1.28	0.30	0.10	21.62	7.35	21.82	7.42	46.67	15.87
FC 0303	Kumquats, raw (incl juice)	RAC	3.4	4.67	15.88	5.86	19.92	1.96	6.66	1.45	4.93	17.05	57.97	1.37	4.66
FI 0327	Banana, raw (incl plantains) (incl dried)	RAC	0.05	25.76	1.29	23.65	1.18	23.83	1.19	24.37	1.22	19.43	0.97	101.55	5.08
VO 0448	Tomato, raw (incl juice, incl paste, incl canned)	RAC	0.13	64.74	8.42	68.31	8.88	36.05	4.69	82.09	10.67	54.50	7.09	11.69	1.52
VR 0589	Potato, raw (incl flour, incl frozen, incl starch, incl tapioca)	RAC	2.2	225.03	495.07	234.24	515.33	71.48	157.26	177.55	390.61	234.55	516.01	37.71	82.96
GC 0653	Triticale, raw (incl flour)	RAC	0	0.01	0.00	0.17	0.00	0.29	0.00	0.01	0.00	NC	-	NC	-
GC 0654	Wheat, raw (incl bulgur, incl fermented beverages, incl germ, incl wholemeal bread, incl white flour products, incl white bread)	RAC	0	253.07	0.00	244.73	0.00	134.44	0.00	235.10	0.00	216.39	0.00	167.40	0.00
GC 0640	Barley, raw (incl malt extract, incl pot&pearled, incl flour & grits, incl beer, incl malt)	RAC	0	36.18	0.00	53.45	0.00	9.39	0.00	35.25	0.00	46.68	0.00	15.92	0.00
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.4	112.02	44.81	120.71	48.28	63.46	25.38	88.99	35.60	96.24	38.50	41.02	16.41
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.4	28.01	11.20	30.18	12.07	15.86	6.35	22.25	8.90	24.06	9.62	10.25	4.10
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.04	6.44	0.26	15.51	0.62	3.79	0.15	8.29	0.33	18.44	0.74	8.00	0.32
MO 0105	Edible offal (mammalian), raw	RAC	0.42	15.17	6.37	5.19	2.18	6.30	2.65	6.78	2.85	3.32	1.39	3.17	1.33
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.04	388.92	15.56	335.88	13.44	49.15	1.97	331.25	13.25	468.56	18.74	245.45	9.82
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0.04	66.38	2.66	48.47	1.94	21.58	0.86	78.41	3.14	48.04	1.92	76.01	3.04
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0.04	7.38	0.30	5.39	0.22	2.40	0.10	8.71	0.35	5.34	0.21	8.45	0.34
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.04	0.10	0.00	0.10	0.00	NC	-	0.10	0.00	0.71	0.03	NC	-

Annex 3

IMAZALIL (110)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.04	0.33	0.01	0.72	0.03	0.27	0.01	0.35	0.01	0.80	0.03	NC	-
PE 0112	Eggs, raw, (incl dried)	RAC	0.02	25.84	0.52	29.53	0.59	28.05	0.56	33.19	0.66	36.44	0.73	8.89	0.18
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Total intake (ug/person)=				622.5		630.0		209.7		486.5		666.9		184.9	
Bodyweight per region (kg bw) =				60		60		55		60		60		60	
ADI (ug/person)=				1800		1800		1650		1800		1800		1800	
%ADI=				34.6%		35.0%		12.7%		27.0%		37.1%		10.3%	
Rounded %ADI=				30%		30%		10%		30%		40%		10%	

IMAZALIL (110)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day				Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake		
FC 0001	Group of Citrus fruit, raw (incl citrus fruit juice, excl kumquat commodities)	RAC	0.07	2.80	0.20	2.71	0.19	56.74	3.97	0.36	0.03	1.78	0.12		
JF 0001	Group of Citrus fruit, juice	PP	0.34	0.11	0.04	0.29	0.10	13.55	4.61	0.14	0.05	0.33	0.11		
FC 0303	Kumquats, raw (incl juice)	RAC	3.4	18.35	62.39	0.23	0.78	1.78	6.05	0.08	0.27	3.35	11.39		
FI 0327	Banana, raw (incl plantains) (incl dried)	RAC	0.05	44.80	2.24	118.17	5.91	25.25	1.26	454.49	22.72	310.23	15.51		
VO 0448	Tomato, raw (incl juice, incl paste, incl canned)	RAC	0.13	15.50	2.02	5.78	0.75	71.52	9.30	2.00	0.26	12.50	1.63		
VR 0589	Potato, raw (incl flour, incl frozen, incl starch, incl tapioca)	RAC	2.2	23.96	52.71	13.56	29.83	213.41	469.50	104.35	229.57	8.56	18.83		
GC 0653	Triticale, raw (incl flour)	RAC	0	0.01	0.00	NC	-	NC	-	NC	-	NC	-		
GC 0654	Wheat, raw (incl bulgur, incl fermented beverages, incl germ, incl wholemeal bread, incl white flour products, incl white bread)	RAC	0	57.20	0.00	110.47	0.00	272.62	0.00	25.82	0.00	132.04	0.00		
GC 0640	Barley, raw (incl malt extract, incl pot&pearled, incl flour & grits, incl beer, incl malt)	RAC	0	11.58	0.00	2.33	0.00	46.71	0.00	3.72	0.00	16.26	0.00		
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.4	23.34	9.34	40.71	16.28	97.15	38.86	18.06	7.23	57.71	23.08		
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.4	5.84	2.33	10.18	4.07	24.29	9.72	4.52	1.81	14.43	5.77		

IMAZALIL (110)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw						
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person								
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake	
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.04	1.05	0.04	1.14	0.05	18.69	0.75	0.94	0.04	3.12	0.12	
MO 0105	Edible offal (mammalian), raw	RAC	0.42	4.64	1.95	1.97	0.83	10.01	4.20	3.27	1.37	3.98	1.67	
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.04	108.75	4.35	70.31	2.81	436.11	17.44	61.55	2.46	79.09	3.16	
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0.04	3.53	0.14	10.83	0.43	51.36	2.05	4.53	0.18	50.00	2.00	
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0.04	0.39	0.02	1.20	0.05	5.71	0.23	0.50	0.02	5.56	0.22	
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.04	NC	-	NC	-	0.32	0.01	NC	-	NC	-	
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.04	0.10	0.00	0.70	0.03	0.97	0.04	0.10	0.00	NC	-	
PE 0112	Eggs, raw, (incl dried)	RAC	0.02	3.84	0.08	4.41	0.09	27.25	0.55	1.13	0.02	7.39	0.15	
-	-	-		-	-	-	-	-	-	-	-	-	-	
Total intake (ug/person)=					137.8		62.2		568.5		266.0		83.8	
Bodyweight per region (kg bw) =					60		60		60		60		60	
ADI (ug/person)=					1800		1800		1800		1800		1800	
%ADI=					7.7%		3.5%		31.6%		14.8%		4.7%	
Rounded %ADI=					8%		3%		30%		10%		5%	

Annex 3

ISOPROTHIOLANE (299)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
FI 0327	Banana, raw (incl plantains) (incl dried)	RAC	0.0077	5.23	0.04	6.94	0.05	99.45	0.77	32.47	0.25	48.30	0.37	24.70	0.19
CM 0649 (GC 0649)	Rice, husked, dry (incl flour, incl oil, incl beverages, incl starch, excl polished)	REP	1.54	1.26	1.94	1.58	2.43	31.05	47.82	5.43	8.36	0.90	1.39	2.18	3.36
CM 1205	Rice polished, dry	PP	0.38	34.21	13.00	10.39	3.95	41.72	15.85	82.38	31.30	150.24	57.09	70.47	26.78
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat)	RAC	0.01	31.20	0.31	72.44	0.72	20.88	0.21	47.98	0.48	33.08	0.33	36.25	0.36
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.01	24.96	0.25	57.95	0.58	16.70	0.17	38.38	0.38	26.46	0.26	29.00	0.29
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.01	6.24	0.06	14.49	0.14	4.18	0.04	9.60	0.10	6.62	0.07	7.25	0.07
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.01	3.29	0.03	6.14	0.06	0.82	0.01	1.57	0.02	2.23	0.02	1.07	0.01
MO 0105	Edible offal (mammalian), raw	RAC	0.01	4.79	0.05	9.68	0.10	2.97	0.03	5.49	0.05	3.84	0.04	5.03	0.05
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.01	289.65	2.90	485.88	4.86	26.92	0.27	239.03	2.39	199.91	2.00	180.53	1.81
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-
Total intake (ug/person)=				18.6		12.9		65.2		43.3		61.6		32.9	
Bodyweight per region (kg bw) =				60		60		60		60		60		60	
ADI (ug/person)=				6000		6000		6000		6000		6000		6000	
%ADI=				0.3%		0.2%		1.1%		0.7%		1.0%		0.5%	
Rounded %ADI=				0%		0%		1%		1%		1%		1%	

ISOPROTHIOLANE (299)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
FI 0327	Banana, raw (incl plantains) (incl dried)	RAC	0.0077	25.76	0.20	23.65	0.18	23.83	0.18	24.37	0.19	19.43	0.15	101.55	0.78
CM 0649 (GC 0649)	Rice, husked, dry (incl flour, incl oil, incl beverages, incl starch, excl polished)	REP	1.54	3.70	5.70	2.11	3.25	1.51	2.33	1.75	2.70	0.29	0.45	5.12	7.88
CM 1205	Rice polished, dry	PP	0.38	13.38	5.08	10.80	4.10	262.08	99.59	57.16	21.72	12.83	4.88	62.78	23.86
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat)	RAC	0.01	140.03	1.40	150.89	1.51	79.32	0.79	111.24	1.11	120.30	1.20	51.27	0.51
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.01	112.02	1.12	120.71	1.21	63.46	0.63	88.99	0.89	96.24	0.96	41.02	0.41
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.01	28.01	0.28	30.18	0.30	15.86	0.16	22.25	0.22	24.06	0.24	10.25	0.10
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.01	6.44	0.06	15.51	0.16	3.79	0.04	8.29	0.08	18.44	0.18	8.00	0.08
MO 0105	Edible offal (mammalian), raw	RAC	0.01	15.17	0.15	5.19	0.05	6.30	0.06	6.78	0.07	3.32	0.03	3.17	0.03
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.01	388.92	3.89	335.88	3.36	49.15	0.49	331.25	3.31	468.56	4.69	245.45	2.45
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-
Total intake (ug/person)=				17.9		14.1		104.3		30.3		12.8		36.1	
Bodyweight per region (kg bw) =				60		60		55		60		60		60	
ADI (ug/person)=				6000		6000		5500		6000		6000		6000	
%ADI=				0.3%		0.2%		1.9%		0.5%		0.2%		0.6%	
Rounded %ADI=				0%		0%		2%		1%		0%		1%	

Annex 3

ISOPROTHIOLANE (299)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw						
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person								
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake	
FI 0327	Banana, raw (incl plantains) (incl dried)	RAC	0.0077	44.80	0.34	118.17	0.91	25.25	0.19	454.49	3.50	310.23	2.39	
CM 0649 (GC 0649)	Rice, husked, dry (incl flour, incl oil, incl beverages, incl starch, excl polished)	REP	1.54	13.58	20.91	4.29	6.61	2.17	3.34	0.01	0.02	8.84	13.61	
CM 1205	Rice polished, dry	PP	0.38	30.20	11.48	218.34	82.97	12.77	4.85	15.24	5.79	51.35	19.51	
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat)	RAC	0.01	29.18	0.29	50.89	0.51	121.44	1.21	22.58	0.23	72.14	0.72	
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.01	23.34	0.23	40.71	0.41	97.15	0.97	18.06	0.18	57.71	0.58	
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.01	5.84	0.06	10.18	0.10	24.29	0.24	4.52	0.05	14.43	0.14	
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.01	1.05	0.01	1.14	0.01	18.69	0.19	0.94	0.01	3.12	0.03	
MO 0105	Edible offal (mammalian), raw	RAC	0.01	4.64	0.05	1.97	0.02	10.01	0.10	3.27	0.03	3.98	0.04	
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.01	108.75	1.09	70.31	0.70	436.11	4.36	61.55	0.62	79.09	0.79	
-	-	-		-	-	-	-	-	-	-	-	-	-	
Total intake (ug/person)=					34.5		92.2		15.5		10.4		37.8	
Bodyweight per region (kg bw) =					60		60		60		60		60	
ADI (ug/person)=					6000		6000		6000		6000		6000	
%ADI=					0.6%		1.5%		0.3%		0.2%		0.6%	
Rounded %ADI=					1%		2%		0%		0%		1%	

[illegible]

Annex 3

ISOXAFLUTOLE (268)

International Estimated Daily Intake (IEDI)

ADI = 0 - 0.02 mg/kg bw

Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
VD 0541	Soya bean, dry, raw (incl flour, incl curd, excl oil, excl paste, excl sauce)	RAC	0.02	0.28	0.01	0.65	0.01	7.93	0.16	5.36	0.11	NC	-	5.78	0.12
-	Soya paste (i.e. miso)	PP	0.02	NC	-	NC	-	NC	-	1.87	0.04	NC	-	NC	-
OR 0541	Soya oil, refined	PP	0.008	19.06	0.15	21.06	0.17	5.94	0.05	33.78	0.27	40.05	0.32	13.39	0.11
-	Soya sauce	PP	0.02	0.45	0.01	0.29	0.01	2.93	0.06	4.35	0.09	0.10	0.00	0.70	0.01
VD 0524	Chick-pea (dry) (Cicer spp), raw	RAC	0	0.27	0.00	1.33	0.00	0.32	0.00	0.15	0.00	0.10	0.00	0.10	0.00
GC 0645	Maize, raw (incl glucose & dextrose & isoglucose, incl flour, incl oil, incl beer, incl germ, incl starch)	RAC	0.02	18.51	0.37	26.18	0.52	26.04	0.52	39.99	0.80	7.36	0.15	64.58	1.29
GC 0447	Sweet corn on the cob, raw (incl frozen kernels, incl canned kernels)	RAC	0	11.43	0.00	3.71	0.00	0.74	0.00	13.63	0.00	3.07	0.00	1.50	0.00
GS 0659	Sugar cane, raw (incl sugar, incl molasses)	RAC	0	92.24	0.00	95.72	0.00	28.47	0.00	77.39	0.00	117.73	0.00	103.90	0.00
SO 0698	Poppy seed, raw (incl oil)	RAC	0	0.10	0.00	0.25	0.00	0.10	0.00	0.10	0.00	NC	-	NC	-
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat)	RAC	0	140.03	0.00	150.89	0.00	79.32	0.00	111.24	0.00	120.30	0.00	51.27	0.00
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0	6.44	0.00	15.51	0.00	3.79	0.00	8.29	0.00	18.44	0.00	8.00	0.00
MO 0105	Edible offal (mammalian), raw	RAC	0.2	15.17	3.03	5.19	1.04	6.30	1.26	6.78	1.36	3.32	0.66	3.17	0.63
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0	388.92	0.00	335.88	0.00	49.15	0.00	331.25	0.00	468.56	0.00	245.45	0.00
PM 0110	Poultry meat, raw (incl prepared)	RAC	0	73.76	0.00	53.86	0.00	23.98	0.00	87.12	0.00	53.38	0.00	84.45	0.00
PF 0111	Poultry fat, raw (incl rendered)	RAC	0	0.10	0.00	0.10	0.00	NC	-	0.10	0.00	0.71	0.00	NC	-
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.1	0.33	0.03	0.72	0.07	0.27	0.03	0.35	0.04	0.80	0.08	NC	-
PE 0112	Eggs, raw, (incl dried)	RAC	0	25.84	0.00	29.53	0.00	28.05	0.00	33.19	0.00	36.44	0.00	8.89	0.00

Total intake (ug/person)=

3.6

1.8

2.1

2.7

1.2

2.2

Bodyweight per region (kg bw) =

60

60

55

60

60

60

ADI (ug/person)=

1200

1200

1100

1200

1200

1200

%ADI=

0.3%

0.2%

0.2%

0.2%

0.1%

0.2%

Rounded %ADI=

0%

0%

0%

0%

0%

0%

ISOXAFLUTOLE (268)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.02 mg/kg bw						
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person								
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake	
VD 0541	Soya bean, dry, raw (incl flour, incl curd, excl oil, excl paste, excl sauce)	RAC	0.02	2.89	0.06	0.16	0.00	0.42	0.01	3.16	0.06	0.10	0.00	
-	Soya paste (i.e. miso)	PP	0.02	NC	-	NC	-	NC	-	NC	-	NC	-	
OR 0541	Soya oil, refined	PP	0.008	2.32	0.02	2.54	0.02	18.70	0.15	2.51	0.02	6.29	0.05	
-	Soya sauce	PP	0.02	0.10	0.00	0.13	0.00	0.17	0.00	0.10	0.00	0.56	0.01	
VD 0524	Chick-pea (dry) (Cicer spp), raw	RAC	0	1.09	0.00	1.56	0.00	0.33	0.00	0.18	0.00	0.47	0.00	
GC 0645	Maize, raw (incl glucose & dextrose & isoglucose, incl flour, incl oil, incl beer, incl germ, incl starch)	RAC	0.02	116.66	2.33	10.52	0.21	38.46	0.77	76.60	1.53	34.44	0.69	
GC 0447	Sweet corn on the cob, raw (incl frozen kernels, incl canned kernels)	RAC	0	3.63	0.00	20.50	0.00	8.78	0.00	0.10	0.00	0.17	0.00	
GS 0659	Sugar cane, raw (incl sugar, incl molasses)	RAC	0	33.75	0.00	106.29	0.00	78.09	0.00	29.09	0.00	45.70	0.00	
SO 0698	Poppy seed, raw (incl oil)	RAC	0	0.10	0.00	0.10	0.00	0.11	0.00	NC	-	NC	-	
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat)	RAC	0	29.18	0.00	50.89	0.00	121.44	0.00	22.58	0.00	72.14	0.00	
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0	1.05	0.00	1.14	0.00	18.69	0.00	0.94	0.00	3.12	0.00	
MO 0105	Edible offal (mammalian), raw	RAC	0.2	4.64	0.93	1.97	0.39	10.01	2.00	3.27	0.65	3.98	0.80	
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0	108.75	0.00	70.31	0.00	436.11	0.00	61.55	0.00	79.09	0.00	
PM 0110	Poultry meat, raw (incl prepared)	RAC	0	3.92	0.00	12.03	0.00	57.07	0.00	5.03	0.00	55.56	0.00	
PF 0111	Poultry fat, raw (incl rendered)	RAC	0	NC	-	NC	-	0.32	0.00	NC	-	NC	-	
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.1	0.10	0.01	0.70	0.07	0.97	0.10	0.10	0.01	NC	-	
PE 0112	Eggs, raw, (incl dried)	RAC	0	3.84	0.00	4.41	0.00	27.25	0.00	1.13	0.00	7.39	0.00	
Total intake (ug/person)=					3.3		0.7		3.0		2.3		1.5	
Bodyweight per region (kg bw) =					60		60		60		60		60	
ADI (ug/person)=					1200		1200		1200		1200		1200	
%ADI=					0.3%		0.1%		0.3%		0.2%		0.1%	
Rounded %ADI=					0%		0%		0%		0%		0%	

Annex 3

MANDIPROPAMID (231)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.2 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
FC 0002	Subgroup of Lemons and limes, raw (excl kumquat commodities)	RAC	0.01	2.42	0.02	2.15	0.02	0.43	0.00	10.74	0.11	6.59	0.07	14.06	0.14
FC 0303	Kumquats, raw (incl juice)	RAC	0.098	2.36	0.23	0.27	0.03	3.19	0.31	14.44	1.42	1.66	0.16	1.71	0.17
-	Lemon, juice (single strength, incl. concentrated)	PP	0.0042	0.01	0.00	0.01	0.00	0.11	0.00	0.09	0.00	0.18	0.00	0.17	0.00
FC 0003	Subgroup of Mandarins, raw	RAC	0.01	6.18	0.06	3.66	0.04	0.25	0.00	6.82	0.07	3.49	0.03	19.38	0.19
-	Subgroup of Mandarins, juice	PP	0.0042	NC	-	NC	-	NC	-	NC	-	NC	-	NC	-
FC 0004	Subgroup of Oranges, sweet, sour, raw	RAC	0.01	20.66	0.21	5.23	0.05	11.90	0.12	37.90	0.38	21.16	0.21	56.46	0.56
JF 0004	Subgroup of Oranges, juice (single strength, incl. concentrated)	PP	0.0043	1.27	0.01	2.20	0.01	0.09	0.00	11.81	0.05	0.46	0.00	1.69	0.01
FC 0005	Subgroup of Pummelo and grapefruits, raw	RAC	0.01	0.64	0.01	0.35	0.00	0.93	0.01	6.10	0.06	1.01	0.01	1.36	0.01
JF 0203	Grapefruits, juice (single strength, incl. concentrated)	PP	0.0024	0.01	0.00	0.16	0.00	0.02	0.00	1.97	0.00	0.12	0.00	0.77	0.00
FB 0269	Grapes, raw (incl must, excl dried, excl juice, excl wine)	RAC	0.43	13.02	5.60	9.25	3.98	0.03	0.01	16.91	7.27	3.70	1.59	54.44	23.41
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	1.68	0.51	0.86	0.51	0.86	0.01	0.02	1.27	2.13	0.12	0.20	2.07	3.48
JF 0269	Grape juice (from wine grapes)	PP	0.14	0.14	0.02	0.29	0.04	0.05	0.01	0.30	0.04	0.24	0.03	0.05	0.01
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.366	0.67	0.25	12.53	4.59	2.01	0.74	1.21	0.44	3.53	1.29	4.01	1.47
-	Onions, dry, raw	RAC	0.01	29.36	0.29	37.50	0.38	3.56	0.04	34.78	0.35	18.81	0.19	43.38	0.43
-	Onions, green, raw	RAC	0.48	2.45	1.18	1.49	0.72	1.02	0.49	2.60	1.25	0.60	0.29	2.03	0.97
VB 0400	Broccoli, raw	RAC	0.435	0.88	0.38	0.17	0.07	0.01	0.00	1.25	0.54	3.00	1.31	1.09	0.47
VB 0041	Cabbages, head, raw	RAC	0.01	2.73	0.03	27.92	0.28	0.55	0.01	4.47	0.04	4.27	0.04	10.25	0.10
VC 0424	Cucumber, raw	RAC	0.02	8.01	0.16	30.66	0.61	1.45	0.03	19.84	0.40	0.27	0.01	34.92	0.70
VC 0431	Squash, Summer (Courgette, Marrow, Zucchini, Zucchini), raw	RAC	0.04	0.78	0.03	2.06	0.08	0.30	0.01	1.61	0.06	2.25	0.09	2.36	0.09
VC 0046	Melons, except watermelon, raw (Cantaloupe)	RAC	0.115	8.90	1.02	8.64	0.99	0.80	0.09	17.90	2.06	2.80	0.32	29.17	3.35
VO 0448	Tomato, raw (incl paste, excl juice, excl canned)	RAC	0.06	51.07	3.06	80.96	4.86	16.96	1.02	99.83	5.99	26.09	1.57	212.26	12.74
-	Tomato, canned (& peeled)	PP	0.022	0.20	0.00	0.31	0.01	0.02	0.00	1.11	0.02	0.11	0.00	1.50	0.03
JF 0448	Tomato, juice (single strength, incl concentrated)	PP	0.059	0.29	0.02	0.29	0.02	0.01	0.00	0.38	0.02	0.05	0.00	0.14	0.01
VO 0444	Peppers, chili, raw	RAC	0.84	3.99	3.35	7.30	6.13	2.93	2.46	5.62	4.72	NC	-	17.44	14.65
VO 0445	Peppers, sweet, raw (incl dried)	RAC	0.12	4.49	0.54	6.44	0.77	7.21	0.87	5.68	0.68	9.52	1.14	8.92	1.07

Total intake (ug/person)=	73.2	174.0	60.4	190.3	271.3	195.4
Bodyweight per region (kg bw) =	60	60	60	60	60	60

Annex 3

MANDIPROPAMID (231)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.2 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day			Intake as ug/person/day								
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
	ADI (ug/person)=				12000		12000		12000		12000		12000		12000
	%ADI=				0.6%		1.4%		0.5%		1.6%		2.3%		1.6%
	Rounded %ADI=				1%		1%		1%		2%		2%		2%

MANDIPROPAMID (231)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.2 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day			Intake as ug/person/day								
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
FC 0002	Subgroup of Lemons and limes, raw (excl kumquat commodities)	RAC	0.01	3.78	0.04	8.84	0.09	0.92	0.01	6.71	0.07	4.09	0.04	4.57	0.05
FC 0303	Kumquats, raw (incl juice)	RAC	0.098	4.67	0.46	5.86	0.57	1.96	0.19	1.45	0.14	17.05	1.67	1.37	0.13
-	Lemon, juice (single strength, incl. concentrated)	PP	0.0042	0.60	0.00	0.36	0.00	0.01	0.00	1.49	0.01	0.43	0.00	0.24	0.00
FC 0003	Subgroup of Mandarins, raw	RAC	0.01	12.34	0.12	14.99	0.15	16.08	0.16	10.76	0.11	9.94	0.10	NC	-
-	Subgroup of Mandarins, juice	PP	0.0042	0.04	0.00	NC	-	0.01	0.00	0.01	0.00	NC	-	NC	-
FC 0004	Subgroup of Oranges, sweet, sour, raw	RAC	0.01	15.68	0.16	24.00	0.24	6.80	0.07	29.09	0.29	15.39	0.15	160.47	1.60
JF 0004	Subgroup of Oranges, juice (single strength, incl. concentrated)	PP	0.0043	33.31	0.14	1.78	0.01	0.28	0.00	18.97	0.08	14.01	0.06	13.36	0.06
FC 0005	Subgroup of Pummelo and grapefruits, raw	RAC	0.01	2.19	0.02	1.24	0.01	0.60	0.01	3.44	0.03	4.60	0.05	299.96	3.00
JF 0203	Grapefruits, juice (single strength, incl. concentrated)	PP	0.0024	2.89	0.01	1.61	0.00	0.02	0.00	1.15	0.00	7.39	0.02	33.07	0.08
FB 0269	Grapes, raw (incl must, excl dried, excl juice, excl wine)	RAC	0.43	6.48	2.79	11.31	4.86	5.21	2.24	9.50	4.09	4.66	2.00	0.78	0.34
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	1.68	3.09	5.19	1.51	2.54	0.03	0.05	1.38	2.32	4.26	7.16	0.42	0.71
JF 0269	Grape juice (from wine grapes)	PP	0.14	0.56	0.08	1.96	0.27	0.02	0.00	2.24	0.31	2.27	0.32	0.34	0.05
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.366	88.93	32.55	62.41	22.84	1.84	0.67	25.07	9.18	61.17	22.39	5.84	2.14
-	Onions, dry, raw	RAC	0.01	19.69	0.20	29.83	0.30	24.64	0.25	31.35	0.31	9.72	0.10	12.59	0.13
-	Onions, green, raw	RAC	0.48	1.55	0.74	0.74	0.36	1.05	0.50	3.74	1.80	0.94	0.45	6.45	3.10
VB 0400	Broccoli, raw	RAC	0.435	4.24	1.84	1.76	0.77	NC	-	0.51	0.22	3.79	1.65	0.26	0.11
VB 0041	Cabbages, head, raw	RAC	0.01	8.97	0.09	27.12	0.27	1.44	0.01	24.96	0.25	4.55	0.05	11.23	0.11

MANDIPROPAMID (231)															
International Estimated Daily Intake (IEDI)				ADI = 0 - 0.2 mg/kg bw											
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
VC 0424	Cucumber, raw	RAC	0.02	6.72	0.13	11.03	0.22	32.10	0.64	15.10	0.30	4.05	0.08	9.57	0.19
VC 0431	Squash, Summer (Courgette, Marrow, Zucchini), raw	RAC	0.04	NC	-	NC	-	5.48	0.22	NC	-	NC	-	1.03	0.04
VC 0046	Melons, except watermelon, raw (Cantaloupe)	RAC	0.115	9.20	1.06	11.95	1.37	14.63	1.68	8.99	1.03	7.86	0.90	2.46	0.28
VO 0448	Tomato, raw (incl paste, excl juice, excl canned)	RAC	0.06	51.98	3.12	64.09	3.85	35.52	2.13	79.82	4.79	42.65	2.56	10.96	0.66
-	Tomato, canned (& peeled)	PP	0.022	7.57	0.17	2.66	0.06	0.30	0.01	0.97	0.02	7.31	0.16	0.41	0.01
JF 0448	Tomato, juice (single strength, incl concentrated)	PP	0.059	0.80	0.05	0.07	0.00	0.05	0.00	0.61	0.04	0.40	0.02	0.08	0.00
VO 0444	Peppers, chili, raw	RAC	0.84	5.57	4.68	14.00	11.76	8.25	6.93	5.77	4.85	6.44	5.41	2.53	2.13
VO 0445	Peppers, sweet, raw (incl dried)	RAC	0.12	0.82	0.10	1.53	0.18	10.85	1.30	4.59	0.55	1.84	0.22	2.00	0.24
VL 0053	Group of Leafy vegetables, raw	RAC	5.65	18.83	106.39	21.85	123.45	121.23	684.95	43.09	243.46	18.18	102.72	18.32	103.51
VP 2060	Subgroup of beans with pods (all commodities within this group)	RAC	0.22	5.07	1.12	0.83	0.18	0.17	0.04	3.70	0.81	NC	-	NC	-
VR 0589	Potato, raw (incl flour, incl frozen, incl starch, incl tapioca)	RAC	0.0185	225.03	4.16	234.24	4.33	71.48	1.32	177.55	3.28	234.55	4.34	37.71	0.70
VS 0624	Celery	RAC	2.7	7.68	20.74	2.85	7.70	NC	-	3.34	9.02	16.83	45.44	4.04	10.91
-	Barley beer	PP	0.057	180.21	10.27	259.46	14.79	45.91	2.62	172.36	9.82	234.42	13.36	65.30	3.72
-	Millet beer	PP	0.057	NC	-	NC	-	0.02	0.00	NC	-	NC	-	NC	-
-	Sorghum beer	PP	0.057	NC	-	NC	-	0.01	0.00	NC	-	NC	-	NC	-
-	Maize beer	PP	0.057	NC	-	NC	-	NC	-	1.99	0.11	NC	-	NC	-
SB 0715	Cacao bean, raw (incl roasted)	RAC	0.01	NC	-	NC	-	0.01	0.00	0.26	0.00	NC	-	1.41	0.01
-	cacao paste	PP	0.01	NC	-	NC	-	0.01	0.00	0.06	0.00	NC	-	NC	-
DM 0715	cacao powder	PP	0.005	2.78	0.01	1.82	0.01	0.20	0.00	1.66	0.01	0.04	0.00	0.74	0.00
DM 1215	cacao butter	PP	0.005	0.98	0.00	0.59	0.00	0.01	0.00	0.06	0.00	1.05	0.01	NC	-
-	cacao products NES	PP	0.01	3.79	0.04	3.18	0.03	0.09	0.00	2.10	0.02	0.17	0.00	3.13	0.03
HS 0444	Peppers, chili, dried	PP	0.84	0.11	0.09	0.21	0.18	0.36	0.30	0.21	0.18	0.25	0.21	0.15	0.13
DH 1100	Hops, dry	RAC	28.5	NC	-	NC	-	0.02	0.57	0.02	0.57	NC	-	NC	-
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat)	RAC	0	140.03	0.00	150.89	0.00	79.32	0.00	111.24	0.00	120.30	0.00	51.27	0.00
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.0064	6.44	0.04	15.51	0.10	3.79	0.02	8.29	0.05	18.44	0.12	8.00	0.05

Annex 3

MANDIPROPAMID (231)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.2 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
MO 0105	Edible offal (mammalian), raw	RAC	0.0022	15.17	0.03	5.19	0.01	6.30	0.01	6.78	0.01	3.32	0.01	3.17	0.01
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0	388.92	0.00	335.88	0.00	49.15	0.00	331.25	0.00	468.56	0.00	245.45	0.00
PM 0110	Poultry meat, raw (incl prepared)	RAC	0	73.76	0.00	53.86	0.00	23.98	0.00	87.12	0.00	53.38	0.00	84.45	0.00
PF 0111	Poultry fat, raw (incl rendered)	RAC	0	0.10	0.00	0.10	0.00	NC	-	0.10	0.00	0.71	0.00	NC	-
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0	0.33	0.00	0.72	0.00	0.27	0.00	0.35	0.00	0.80	0.00	NC	-
PE 0112	Eggs, raw, (incl dried)	RAC	0	25.84	0.00	29.53	0.00	28.05	0.00	33.19	0.00	36.44	0.00	8.89	0.00
Total intake (ug/person)=				196.6		201.5		706.9		298.1		211.8		134.2	
Bodyweight per region (kg bw) =				60		60		55		60		60		60	
ADI (ug/person)=				12000		12000		11000		12000		12000		12000	
%ADI=				1.6%		1.7%		6.4%		2.5%		1.8%		1.1%	
Rounded %ADI=				2%		2%		6%		2%		2%		1%	

MANDIPROPAMID (231)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.2 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
FC 0002	Subgroup of Lemons and limes, raw (excl kumquat commodities)	RAC	0.01	0.61	0.01	0.73	0.01	4.01	0.04	0.01	0.00	NC	-
FC 0303	Kumquats, raw (incl juice)	RAC	0.098	18.35	1.80	0.23	0.02	1.78	0.17	0.08	0.01	3.35	0.33
-	Lemon, juice (single strength, incl. concentrated)	PP	0.0042	0.01	0.00	0.01	0.00	0.16	0.00	0.01	0.00	NC	-
FC 0003	Subgroup of Mandarins, raw	RAC	0.01	0.16	0.00	0.27	0.00	9.06	0.09	0.01	0.00	0.02	0.00
-	Subgroup of Mandarins, juice	PP	0.0042	0.01	0.00	NC	-	NC	-	NC	-	NC	-
FC 0004	Subgroup of Oranges, sweet, sour, raw	RAC	0.01	1.18	0.01	1.11	0.01	14.28	0.14	0.05	0.00	1.08	0.01
JF 0004	Subgroup of Oranges, juice (single strength, incl. concentrated)	PP	0.0043	0.08	0.00	0.26	0.00	12.61	0.05	0.14	0.00	0.33	0.00
FC 0005	Subgroup of Pummelo and grapefruits, raw	RAC	0.01	0.63	0.01	0.01	0.00	1.58	0.02	0.01	0.00	NC	-
JF 0203	Grapefruits, juice (single strength, incl. concentrated)	PP	0.0024	0.03	0.00	0.02	0.00	0.78	0.00	0.01	0.00	NC	-

MANDIPROPAMID (231)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.2 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
FB 0269	Grapes, raw (incl must, excl dried, excl juice, excl wine)	RAC	0.43	0.14	0.06	0.36	0.15	15.33	6.59	0.01	0.00	0.28	0.12
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	1.68	0.01	0.02	0.13	0.22	1.06	1.78	0.01	0.02	0.03	0.05
JF 0269	Grape juice (from wine grapes)	PP	0.14	0.01	0.00	0.01	0.00	0.41	0.06	0.01	0.00	NC	-
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.366	0.31	0.11	0.23	0.08	60.43	22.12	0.52	0.19	31.91	11.68
-	Onions, dry, raw	RAC	0.01	9.01	0.09	20.24	0.20	30.90	0.31	9.61	0.10	2.11	0.02
-	Onions, green, raw	RAC	0.48	1.43	0.69	0.05	0.02	0.20	0.10	NC	-	6.30	3.02
VB 0400	Broccoli, raw	RAC	0.435	0.01	0.00	0.01	0.00	2.13	0.93	0.01	0.00	NC	-
VB 0041	Cabbages, head, raw	RAC	0.01	3.82	0.04	2.99	0.03	49.16	0.49	0.01	0.00	NC	-
VC 0424	Cucumber, raw	RAC	0.02	0.68	0.01	1.81	0.04	10.40	0.21	0.01	0.00	0.04	0.00
VC 0431	Squash, Summer (Courgette, Marrow, Zucchini, Zucchini), raw	RAC	0.04	0.09	0.00	1.01	0.04	NC	-	1.91	0.08	NC	-
VC 0046	Melons, except watermelon, raw (Cantaloupe)	RAC	0.115	0.19	0.02	0.10	0.01	4.98	0.57	0.01	0.00	NC	-
VO 0448	Tomato, raw (incl paste, excl juice, excl canned)	RAC	0.06	15.33	0.92	5.65	0.34	67.23	4.03	1.88	0.11	12.48	0.75
-	Tomato, canned (& peeled)	PP	0.022	0.07	0.00	0.08	0.00	2.42	0.05	0.07	0.00	NC	-
JF 0448	Tomato, juice (single strength, incl concentrated)	PP	0.059	0.05	0.00	0.01	0.00	0.42	0.02	0.01	0.00	0.02	0.00
VO 0444	Peppers, chili, raw	RAC	0.84	3.47	2.91	3.56	2.99	16.30	13.69	0.01	0.01	NC	-
VO 0445	Peppers, sweet, raw (incl dried)	RAC	0.12	5.49	0.66	10.57	1.27	8.84	1.06	0.91	0.11	NC	-
VL 0053	Group of Leafy vegetables, raw	RAC	5.65	12.42	70.17	8.75	49.44	7.53	42.54	7.07	39.95	14.11	79.72
VP 2060	Subgroup of beans with pods (all commodities within this group)	RAC	0.22	NC	-	NC	-	NC	-	NC	-	NC	-
VR 0589	Potato, raw (incl flour, incl frozen, incl starch, incl tapioca)	RAC	0.0185	23.96	0.44	13.56	0.25	213.41	3.95	104.35	1.93	8.56	0.16
VS 0624	Celery	RAC	2.7	3.66	9.88	2.65	7.16	4.84	13.07	2.47	6.67	4.94	13.34
-	Barley beer	PP	0.057	16.25	0.93	11.36	0.65	225.21	12.84	19.49	1.11	52.17	2.97
-	Millet beer	PP	0.057	5.80	0.33	NC	-	NC	-	22.96	1.31	NC	-
-	Sorghum beer	PP	0.057	24.90	1.42	NC	-	NC	-	70.33	4.01	NC	-
-	Maize beer	PP	0.057	1.03	0.06	NC	-	NC	-	40.94	2.33	NC	-
SB 0715	Cacao bean, raw (incl roasted)	RAC	0.01	0.01	0.00	0.53	0.01	0.33	0.00	0.08	0.00	NC	-

Annex 3

MANDIPROPAMID (231)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.2 mg/kg bw						
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person								
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake	
-	cacao paste	PP	0.01	0.01	0.00	0.01	0.00	NC	-	0.01	0.00	NC	-	
DM 0715	cacao powder	PP	0.005	0.05	0.00	0.20	0.00	1.17	0.01	0.01	0.00	1.80	0.01	
DM 1215	cacao butter	PP	0.005	0.01	0.00	0.01	0.00	0.38	0.00	0.01	0.00	NC	-	
-	cacao products NES	PP	0.01	0.04	0.00	0.17	0.00	4.41	0.04	0.08	0.00	0.51	0.01	
HS 0444	Peppers, chili, dried	PP	0.84	0.58	0.49	1.27	1.07	1.21	1.02	0.12	0.10	NC	-	
DH 1100	Hops, dry	RAC	28.5	NC	-	NC	-	0.04	1.14	NC	-	NC	-	
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat)	RAC	0	29.18	0.00	50.89	0.00	121.44	0.00	22.58	0.00	72.14	0.00	
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.0064	1.05	0.01	1.14	0.01	18.69	0.12	0.94	0.01	3.12	0.02	
MO 0105	Edible offal (mammalian), raw	RAC	0.0022	4.64	0.01	1.97	0.00	10.01	0.02	3.27	0.01	3.98	0.01	
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0	108.75	0.00	70.31	0.00	436.11	0.00	61.55	0.00	79.09	0.00	
PM 0110	Poultry meat, raw (incl prepared)	RAC	0	3.92	0.00	12.03	0.00	57.07	0.00	5.03	0.00	55.56	0.00	
PF 0111	Poultry fat, raw (incl rendered)	RAC	0	NC	-	NC	-	0.32	0.00	NC	-	NC	-	
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0	0.10	0.00	0.70	0.00	0.97	0.00	0.10	0.00	NC	-	
PE 0112	Eggs, raw, (incl dried)	RAC	0	3.84	0.00	4.41	0.00	27.25	0.00	1.13	0.00	7.39	0.00	
Total intake (ug/person)=					91.1		64.0		127.3		58.1		112.2	
Bodyweight per region (kg bw) =					60		60		60		60		60	
ADI (ug/person)=					12000		12000		12000		12000		12000	
%ADI=					0.8%		0.5%		1.1%		0.5%		0.9%	
Rounded %ADI=					1%		1%		1%		0%		1%	

METHOXYFENOZIDE (209)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
FC 0001	Group of Citrus fruit, raw (excl kumquat commodities)	RAC	0.28	29.89	8.37	11.40	3.19	13.51	3.78	61.57	17.24	32.24	9.03	91.26	25.55
JF 0001	Group of Citrus fruit, juice	PP	0.062	1.30	0.08	2.37	0.15	0.22	0.01	13.88	0.86	0.75	0.05	2.63	0.16
FP 0009	Group of Pome fruits, raw (incl apple cider, excl apple juice)	RAC	0.43	19.35	8.32	34.06	14.65	17.87	7.68	25.74	11.07	7.69	3.31	56.85	24.45
JF 0226	Apple juice, single strength (incl. concentrated)	PP	0.13	0.32	0.04	3.07	0.40	0.07	0.01	5.00	0.65	0.29	0.04	5.57	0.72
FS 0012	Group of Stone fruits, raw (incl dried apricots, excl dried plums)	RAC	0.44	11.33	4.99	23.62	10.39	0.24	0.11	11.32	4.98	2.28	1.00	33.26	14.63
DF 0014	Plums, dried (prunes)	PP	0.44	0.09	0.04	0.06	0.03	0.01	0.00	0.18	0.08	0.04	0.02	0.06	0.03
FB 0020	Blueberries, raw	RAC	1.25	0.01	0.01	0.03	0.04	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
FB 0269	Grapes, raw (incl must, excl dried, excl juice, excl wine)	RAC	0.33	13.02	4.30	9.25	3.05	0.03	0.01	16.91	5.58	3.70	1.22	54.44	17.97
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	0.73	0.51	0.37	0.51	0.37	0.01	0.01	1.27	0.93	0.12	0.09	2.07	1.51
JF 0269	Grape juice (from wine grapes)	PP	0.1	0.14	0.01	0.29	0.03	0.05	0.01	0.30	0.03	0.24	0.02	0.05	0.01
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.13	0.67	0.09	12.53	1.63	2.01	0.26	1.21	0.16	3.53	0.46	4.01	0.52
FB 0265	Cranberry, raw	RAC	0.1	0.02	0.00	0.01	0.00	NC	-	0.03	0.00	0.01	0.00	0.01	0.00
FB 0275	Strawberry, raw	RAC	0.24	0.70	0.17	2.01	0.48	0.04	0.01	1.36	0.33	0.37	0.09	2.53	0.61
FI 0326	Avocado, raw	RAC	0.13	0.13	0.02	0.03	0.00	2.05	0.27	2.54	0.33	2.34	0.30	0.12	0.02
FI 0350	Papaya, raw	RAC	0.31	0.35	0.11	0.01	0.00	3.05	0.95	0.80	0.25	7.28	2.26	1.00	0.31
VB 0400	Broccoli, raw	RAC	0.94	0.88	0.83	0.17	0.16	0.01	0.01	1.25	1.18	3.00	2.82	1.09	1.02
VB 0041	Cabbages, head, raw	RAC	0.93	2.73	2.54	27.92	25.97	0.55	0.51	4.47	4.16	4.27	3.97	10.25	9.53
VC 2039	Subgroup of Cucumbers and Squashes, raw	RAC	0.091	10.52	0.96	39.36	3.58	2.07	0.19	25.74	2.34	2.80	0.25	44.83	4.08
VC 0046	Melons, except watermelon, raw (Cantaloupe)	RAC	0.091	8.90	0.81	8.64	0.79	0.80	0.07	17.90	1.63	2.80	0.25	29.17	2.65
VC 0429	Pumpkins, raw	RAC	0.091	4.76	0.43	12.56	1.14	1.85	0.17	9.86	0.90	5.11	0.47	14.39	1.31
VO 0448	Tomato, raw	RAC	0.2	41.73	8.35	75.65	15.13	10.66	2.13	82.87	16.57	24.75	4.95	200.93	40.19
-	Tomato, canned (& peeled)	PP	0.06	0.20	0.01	0.31	0.02	0.02	0.00	1.11	0.07	0.11	0.01	1.50	0.09
-	Tomato, paste (i.e. concentrated tomato sauce/puree)	PP	0.44	2.34	1.03	1.33	0.59	1.57	0.69	4.24	1.87	0.34	0.15	2.83	1.25
JF 0448	Tomato, juice (single strength, incl concentrated)	PP	0.06	0.29	0.02	0.29	0.02	0.01	0.00	0.38	0.02	0.05	0.00	0.14	0.01
VO 0444	Peppers, chili, raw	RAC	0.16	3.99	0.64	7.30	1.17	2.93	0.47	5.62	0.90	NC	-	17.44	2.79

Annex 3

METHOXYFENOZIDE (209)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
VO 0445	Peppers, sweet, raw	RAC	0.16	1.43	0.23	2.61	0.42	1.05	0.17	2.01	0.32	2.59	0.41	6.24	1.00
-	Peppers, sweet, dried	PP	1.6	0.42	0.67	0.53	0.85	0.84	1.34	0.50	0.80	0.95	1.52	0.37	0.59
VL 0483	Lettuce, leaf, raw	RAC	12	0.53	6.36	0.36	4.32	0.16	1.92	6.21	74.52	1.90	22.80	6.05	72.60
VL 0485	Mustard greens, raw (i.e. Indian mustard, Amsoi, mustard cabbage)	RAC	16	0.03	0.48	0.31	4.96	0.01	0.16	0.05	0.80	0.47	7.52	0.11	1.76
VL 0494	Radish leaves, raw	RAC	0.75	0.26	0.20	0.45	0.34	0.28	0.21	0.68	0.51	NC	-	0.33	0.25
VP 0061	Beans with pods (Phaseolus spp): (immature pods + succulent seeds)	RAC	0.065	0.68	0.04	NC	-	NC	-	0.39	0.03	0.22	0.01	0.49	0.03
014B	Peas with pods	-	0.1	-	-	-	-	-	-	-	-	-	-	-	-
VP 0062	Beans without pods: (Phaseolus spp.) (succulent seeds), raw	RAC	0.05	1.56	0.08	0.60	0.03	0.49	0.02	1.18	0.06	0.90	0.05	7.79	0.39
VP 0064	Peas without pods (Pisum spp) (succulent seeds)	RAC	0.05	1.97	0.10	0.51	0.03	0.02	0.00	0.79	0.04	3.68	0.18	3.80	0.19
VD 0071	Beans, dry, raw (Phaseolus spp)	RAC	0.05	2.39	0.12	1.61	0.08	10.47	0.52	1.84	0.09	12.90	0.65	7.44	0.37
VD 0527	Cowpea, dry, raw (Vigna sinensis, Dolichos sinensis)	RAC	0.56	0.05	0.03	NC	-	1.74	0.97	0.01	0.01	0.01	0.01	0.07	0.04
VD 0072	Peas (dry) (Pisum spp), raw	RAC	0.17	1.62	0.28	3.22	0.55	0.92	0.16	1.50	0.26	2.90	0.49	0.17	0.03
VR 0577	Carrots, raw	RAC	0.13	9.51	1.24	30.78	4.00	0.37	0.05	8.75	1.14	2.80	0.36	6.10	0.79
VR 0494	Radish roots, raw	RAC	0.08	2.31	0.18	4.09	0.33	2.53	0.20	6.15	0.49	5.88	0.47	2.97	0.24
VR 0596	Sugar beet, raw (incl sugar)	RAC	0.11	0.13	0.01	NC	-	0.08	0.01	0.66	0.07	0.47	0.05	88.94	9.78
VR 0508	Sweet potato, raw (incl dried)	RAC	0.01	0.18	0.00	0.18	0.00	42.16	0.42	1.61	0.02	3.06	0.03	6.67	0.07
VS 0624	Celery	RAC	3.4	2.14	7.28	3.79	12.89	2.35	7.99	5.69	19.35	0.02	0.07	2.75	9.35
GC 0645	Maize, raw (incl glucose & dextrose & isoglucose, incl flour, incl oil, incl beer, incl germ, incl starch)	RAC	0.02	29.81	0.60	44.77	0.90	108.95	2.18	52.37	1.05	60.28	1.21	75.69	1.51
GC 2090	Subgroup of Sweet Corns	RAC	0.02	0.14	0.00	0.94	0.02	5.70	0.11	2.61	0.05	1.94	0.04	0.22	0.00
GS 0659	Sugar cane, raw	RAC	0.01	38.16	0.38	NC	-	12.58	0.13	0.34	0.00	17.79	0.18	42.78	0.43
-	Sugar cane, molasses	PP	0.07	NC	-	NC	-	NC	-	NC	-	0.01	0.00	NC	-
-	Sugar cane, sugar (incl non-centrifugal sugar, incl refined sugar and maltose)	PP	0.06	61.52	3.69	86.27	5.18	18.80	1.13	80.02	4.80	66.39	3.98	56.32	3.38
TN 0085	Group of Tree nuts, raw (incl processed)	RAC	0.012	4.06	0.05	3.27	0.04	7.01	0.08	13.93	0.17	14.01	0.17	9.36	0.11
OR 0691	Cotton seed oil, edible	PP	0.15	3.22	0.48	1.54	0.23	1.01	0.15	0.74	0.11	1.12	0.17	2.93	0.44
SO 0697	Peanuts, nutmeat, raw (incl roasted, incl oil, incl	RAC	0.01	1.30	0.01	1.23	0.01	12.62	0.13	2.87	0.03	6.59	0.07	2.67	0.03

METHOXYFENOZIDE (209)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw																			
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day																			
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake												
	butter)																										
SO 0697	Peanuts, nutmeat, raw (incl roasted, incl butter, excl oil)	RAC	0.01	0.46	0.00	1.21	0.01	6.64	0.07	2.71	0.03	1.26	0.01	1.84	0.02												
OR 0697	Peanut oil, edible	PP	0.029	0.36	0.01	0.01	0.00	2.57	0.07	0.07	0.00	2.29	0.07	0.36	0.01												
SB 0716	Coffee bean raw (incl roasted, incl instant coffee, incl substitutes)	RAC	0.028	1.36	0.04	3.59	0.10	1.44	0.04	5.18	0.15	2.02	0.06	1.70	0.05												
HH 0722	Basil leaves, raw (incl dried)	RAC	19	0.14	2.66	0.26	4.94	0.16	3.04	0.38	7.22	NC	-	0.19	3.61												
HS 0444	Peppers, chili, dried	PP	1.6	0.42	0.67	0.53	0.85	0.84	1.34	0.50	0.80	0.95	1.52	0.37	0.59												
DT 1114	Tea, green or black, fermented and dried, (including concentrates)	RAC	29	2.28	66.12	1.98	57.42	0.46	13.34	2.43	70.47	1.29	37.41	3.04	88.16												
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.003	24.96	0.07	57.95	0.17	16.70	0.05	38.38	0.12	26.46	0.08	29.00	0.09												
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.036	6.24	0.22	14.49	0.52	4.18	0.15	9.60	0.35	6.62	0.24	7.25	0.26												
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.036	3.29	0.12	6.14	0.22	0.82	0.03	1.57	0.06	2.23	0.08	1.07	0.04												
MO 0105	Edible offal (mammalian), raw	RAC	0.025	4.79	0.12	9.68	0.24	2.97	0.07	5.49	0.14	3.84	0.10	5.03	0.13												
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.03	289.65	8.69	485.88	14.58	26.92	0.81	239.03	7.17	199.91	6.00	180.53	5.42												
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0	13.17	0.00	26.78	0.00	7.24	0.00	116.71	0.00	22.54	0.00	32.09	0.00												
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0	1.46	0.00	2.98	0.00	0.80	0.00	12.97	0.00	2.50	0.00	3.57	0.00												
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0	0.12	0.00	0.12	0.00	0.11	0.00	5.37	0.00	0.24	0.00	0.10	0.00												
PE 0112	Eggs, raw, (incl dried)	RAC	0	7.84	0.00	23.08	0.00	2.88	0.00	14.89	0.00	9.81	0.00	14.83	0.00												
Total intake (ug/person)=				143.8				197.2				54.4				263.3				116.8				351.2			
Bodyweight per region (kg bw) =				60				60				60				60				60							
ADI (ug/person)=				6000				6000				6000				6000				6000							
%ADI=				2.4%				3.3%				0.9%				4.4%				1.9%				5.9%			
Rounded %ADI=				2%				3%				1%				4%				2%				6%			

Annex 3

METHOXYFENOZIDE (209)															
International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw											
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
FC 0001	Group of Citrus fruit, raw (excl kumquat commodities)	RAC	0.28	33.99	9.52	49.07	13.74	24.40	6.83	50.01	14.00	34.01	9.52	464.99	130.20
JF 0001	Group of Citrus fruit, juice	PP	0.062	36.84	2.28	3.75	0.23	0.30	0.02	21.62	1.34	21.82	1.35	46.67	2.89
FP 0009	Group of Pome fruits, raw (incl apple cider, excl apple juice)	RAC	0.43	51.09	21.97	65.40	28.12	42.71	18.37	45.29	19.47	62.51	26.88	7.74	3.33
JF 0226	Apple juice, single strength (incl. concentrated)	PP	0.13	14.88	1.93	11.98	1.56	0.15	0.02	9.98	1.30	30.32	3.94	3.47	0.45
FS 0012	Group of Stone fruits, raw (incl dried apricots, excl dried plums)	RAC	0.44	18.18	8.00	23.83	10.49	14.27	6.28	18.52	8.15	9.35	4.11	0.11	0.05
DF 0014	Plums, dried (prunes)	PP	0.44	0.61	0.27	0.35	0.15	0.05	0.02	0.35	0.15	0.49	0.22	0.13	0.06
FB 0020	Blueberries, raw	RAC	1.25	0.04	0.05	0.23	0.29	0.01	0.01	0.83	1.04	0.33	0.41	NC	-
FB 0269	Grapes, raw (incl must, excl dried, excl juice, excl wine)	RAC	0.33	6.48	2.14	11.31	3.73	5.21	1.72	9.50	3.14	4.66	1.54	0.78	0.26
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	0.73	3.09	2.26	1.51	1.10	0.03	0.02	1.38	1.01	4.26	3.11	0.42	0.31
JF 0269	Grape juice (from wine grapes)	PP	0.1	0.56	0.06	1.96	0.20	0.02	0.00	2.24	0.22	2.27	0.23	0.34	0.03
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.13	88.93	11.56	62.41	8.11	1.84	0.24	25.07	3.26	61.17	7.95	5.84	0.76
FB 0265	Cranberry, raw	RAC	0.1	0.06	0.01	0.01	0.00	0.01	0.00	1.22	0.12	0.11	0.01	NC	-
FB 0275	Strawberry, raw	RAC	0.24	4.49	1.08	5.66	1.36	0.02	0.00	6.63	1.59	5.75	1.38	0.05	0.01
FI 0326	Avocado, raw	RAC	0.13	2.65	0.34	0.87	0.11	0.46	0.06	1.64	0.21	1.30	0.17	0.96	0.12
FI 0350	Papaya, raw	RAC	0.31	0.31	0.10	0.18	0.06	1.50	0.47	0.51	0.16	0.54	0.17	1.08	0.33
VB 0400	Broccoli, raw	RAC	0.94	4.24	3.99	1.76	1.65	NC	-	0.51	0.48	3.79	3.56	0.26	0.24
VB 0041	Cabbages, head, raw	RAC	0.93	8.97	8.34	27.12	25.22	1.44	1.34	24.96	23.21	4.55	4.23	11.23	10.44
VC 2039	Subgroup of Cucumbers and Squashes, raw	RAC	0.091	7.14	0.65	16.92	1.54	37.58	3.42	15.16	1.38	4.42	0.40	12.67	1.15
VC 0046	Melons, except watermelon, raw (Cantaloupe)	RAC	0.091	9.20	0.84	11.95	1.09	14.63	1.33	8.99	0.82	7.86	0.72	2.46	0.22
VC 0429	Pumpkins, raw	RAC	0.091	6.88	0.63	3.23	0.29	2.59	0.24	12.12	1.10	1.68	0.15	6.30	0.57
VO 0448	Tomato, raw	RAC	0.2	32.13	6.43	51.27	10.25	34.92	6.98	73.37	14.67	15.15	3.03	8.88	1.78
-	Tomato, canned (& peeled)	PP	0.06	7.57	0.45	2.66	0.16	0.30	0.02	0.97	0.06	7.31	0.44	0.41	0.02
-	Tomato, paste (i.e. concentrated tomato sauce/puree)	PP	0.44	4.96	2.18	3.20	1.41	0.15	0.07	1.61	0.71	6.88	3.03	0.52	0.23
JF 0448	Tomato, juice (single strength, incl concentrated)	PP	0.06	0.80	0.05	0.07	0.00	0.05	0.00	0.61	0.04	0.40	0.02	0.08	0.00
VO 0444	Peppers, chili, raw	RAC	0.16	5.57	0.89	14.00	2.24	8.25	1.32	5.77	0.92	6.44	1.03	2.53	0.40

METHOXYFENOZIDE (209)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
VO 0445	Peppers, sweet, raw	RAC	0.16	NC	-	NC	-	8.25	1.32	3.03	0.48	NC	-	0.91	0.15
-	Peppers, sweet, dried	PP	1.6	0.11	0.18	0.21	0.34	0.36	0.58	0.21	0.34	0.25	0.40	0.15	0.24
VL 0483	Lettuce, leaf, raw	RAC	12	14.50	174.00	11.76	141.12	13.14	157.68	19.50	234.00	4.81	57.72	2.23	26.76
VL 0485	Mustard greens, raw (i.e. Indian mustard, Amsoi, mustard cabbage)	RAC	16	NC	-	NC	-	NC	-	NC	-	NC	-	0.13	2.08
VL 0494	Radish leaves, raw	RAC	0.75	NC	-	NC	-	NC	-	3.78	2.84	NC	-	0.48	0.36
VP 0061	Beans with pods (Phaseolus spp): (immature pods + succulent seeds)	RAC	0.065	5.07	0.33	0.83	0.05	0.17	0.01	3.70	0.24	NC	-	NC	-
014B	Peas with pods	-	0.1	-	-	-	-	-	-	-	-	-	-	-	-
VP 0062	Beans without pods: (Phaseolus spp.) (succulent seeds), raw	RAC	0.05	2.21	0.11	5.25	0.26	4.17	0.21	1.61	0.08	16.95	0.85	0.17	0.01
VP 0064	Peas without pods (Pisum spp) (succulent seeds)	RAC	0.05	10.72	0.54	1.99	0.10	2.72	0.14	4.26	0.21	4.23	0.21	NC	-
VD 0071	Beans, dry, raw (Phaseolus spp)	RAC	0.05	1.51	0.08	1.50	0.08	1.90	0.10	5.11	0.26	1.36	0.07	23.43	1.17
VD 0527	Cowpea, dry, raw (Vigna sinensis, Dolichos sinensis)	RAC	0.56	NC	-	NC	-	0.16	0.09	0.01	0.01	NC	-	NC	-
VD 0072	Peas (dry) (Pisum spp), raw	RAC	0.17	3.80	0.65	1.25	0.21	0.90	0.15	2.33	0.40	2.70	0.46	3.83	0.65
VR 0577	Carrots, raw	RAC	0.13	26.26	3.41	27.13	3.53	10.07	1.31	16.49	2.14	44.69	5.81	8.75	1.14
VR 0494	Radish roots, raw	RAC	0.08	3.83	0.31	11.99	0.96	NC	-	5.26	0.42	2.19	0.18	4.37	0.35
VR 0596	Sugar beet, raw (incl sugar)	RAC	0.11	0.01	0.00	NC	-	0.01	0.00	0.01	0.00	NC	-	NC	-
VR 0508	Sweet potato, raw (incl dried)	RAC	0.01	0.93	0.01	0.32	0.00	64.65	0.65	5.37	0.05	0.30	0.00	3.13	0.03
VS 0624	Celery	RAC	3.4	7.68	26.11	2.85	9.69	NC	-	3.34	11.36	16.83	57.22	4.04	13.74
GC 0645	Maize, raw (incl glucose & dextrose & isoglucose, incl flour, incl oil, incl beer, incl germ, incl starch)	RAC	0.02	18.51	0.37	26.18	0.52	26.04	0.52	39.99	0.80	7.36	0.15	64.58	1.29
GC 2090	Subgroup of Sweet Corns	RAC	0.02	11.43	0.23	3.71	0.07	0.74	0.01	13.63	0.27	3.07	0.06	1.50	0.03
GS 0659	Sugar cane, raw	RAC	0.01	NC	-	NC	-	4.27	0.04	0.01	0.00	NC	-	3.24	0.03
-	Sugar cane, molasses	PP	0.07	NC	-	NC	-	0.08	0.01	NC	-	NC	-	NC	-
-	Sugar cane, sugar (incl non-centrifugal sugar, incl refined sugar and maltose)	PP	0.06	92.24	5.53	95.72	5.74	24.12	1.45	77.39	4.64	117.73	7.06	100.67	6.04
TN 0085	Group of Tree nuts, raw (incl processed)	RAC	0.012	8.52	0.10	8.94	0.11	15.09	0.18	9.60	0.12	14.57	0.17	26.26	0.32
OR 0691	Cotton seed oil, edible	PP	0.15	1.68	0.25	0.66	0.10	1.13	0.17	1.18	0.18	0.89	0.13	0.37	0.06
SO 0697	Peanuts, nutmeat, raw (incl roasted, incl oil, incl	RAC	0.01	5.63	0.06	2.75	0.03	9.58	0.10	5.82	0.06	13.71	0.14	1.84	0.02

Annex 3

METHOXYFENOZIDE (209)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
	butter)														
SO 0697	Peanuts, nutmeat, raw (incl roasted, incl butter, excl oil)	RAC	0.01	3.26	0.03	2.22	0.02	5.38	0.05	4.85	0.05	1.54	0.02	1.82	0.02
OR 0697	Peanut oil, edible	PP	0.029	1.02	0.03	0.23	0.01	1.81	0.05	0.42	0.01	5.23	0.15	0.01	0.00
SB 0716	Coffee bean raw (incl roasted, incl instant coffee, incl substitutes)	RAC	0.028	10.90	0.31	12.44	0.35	0.77	0.02	9.48	0.27	22.07	0.62	8.15	0.23
HH 0722	Basil leaves, raw (incl dried)	RAC	19	0.52	9.88	0.05	0.95	3.23	61.37	0.18	3.42	0.12	2.28	0.27	5.13
HS 0444	Peppers, chili, dried	PP	1.6	0.11	0.18	0.21	0.34	0.36	0.58	0.21	0.34	0.25	0.40	0.15	0.24
DT 1114	Tea, green or black, fermented and dried, (including concentrates)	RAC	29	2.91	84.39	1.73	50.17	1.14	33.06	1.85	53.65	2.29	66.41	0.74	21.46
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.003	112.02	0.34	120.71	0.36	63.46	0.19	88.99	0.27	96.24	0.29	41.02	0.12
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.036	28.01	1.01	30.18	1.09	15.86	0.57	22.25	0.80	24.06	0.87	10.25	0.37
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.036	6.44	0.23	15.51	0.56	3.79	0.14	8.29	0.30	18.44	0.66	8.00	0.29
MO 0105	Edible offal (mammalian), raw	RAC	0.025	15.17	0.38	5.19	0.13	6.30	0.16	6.78	0.17	3.32	0.08	3.17	0.08
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.03	388.92	11.67	335.88	10.08	49.15	1.47	331.25	9.94	468.56	14.06	245.45	7.36
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0	66.38	0.00	48.47	0.00	21.58	0.00	78.41	0.00	48.04	0.00	76.01	0.00
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0	7.38	0.00	5.39	0.00	2.40	0.00	8.71	0.00	5.34	0.00	8.45	0.00
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0	0.33	0.00	0.72	0.00	0.27	0.00	0.35	0.00	0.80	0.00	NC	-
PE 0112	Eggs, raw, (incl dried)	RAC	0	25.84	0.00	29.53	0.00	28.05	0.00	33.19	0.00	36.44	0.00	8.89	0.00
Total intake (ug/person)=				406.7		340.1		311.1		426.7		294.1		243.6	
Bodyweight per region (kg bw) =				60		60		55		60		60		60	
ADI (ug/person)=				6000		6000		5500		6000		6000		6000	
%ADI=				6.8%		5.7%		5.7%		7.1%		4.9%		4.1%	
Rounded %ADI=				7%		6%		6%		7%		5%		4%	

METHOXYFENOZIDE (209)		International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw							
Codex Code	Commodity description	Expras	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
FC 0001	Group of Citrus fruit, raw (excl kumquat commodities)	RAC	0.28	2.57	0.72	2.12	0.59	28.93	8.10	0.07	0.02	1.10	0.31
JF 0001	Group of Citrus fruit, juice	PP	0.062	0.11	0.01	0.29	0.02	13.55	0.84	0.14	0.01	0.33	0.02
FP 0009	Group of Pome fruits, raw (incl apple cider, excl apple juice)	RAC	0.43	68.85	29.61	10.93	4.70	70.82	30.45	189.78	81.61	19.56	8.41
JF 0226	Apple juice, single strength (incl. concentrated)	PP	0.13	0.03	0.00	0.10	0.01	7.19	0.93	0.03	0.00	NC	-
FS 0012	Group of Stone fruits, raw (incl dried apricots, excl dried plums)	RAC	0.44	0.09	0.04	0.02	0.01	32.27	14.20	0.01	0.00	NC	-
DF 0014	Plums, dried (prunes)	PP	0.44	0.01	0.00	0.01	0.00	0.37	0.16	0.01	0.00	NC	-
FB 0020	Blueberries, raw	RAC	1.25	NC	-	NC	-	0.20	0.25	NC	-	NC	-
FB 0269	Grapes, raw (incl must, excl dried, excl juice, excl wine)	RAC	0.33	0.14	0.05	0.36	0.12	15.33	5.06	0.01	0.00	0.28	0.09
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	0.73	0.01	0.01	0.13	0.09	1.06	0.77	0.01	0.01	0.03	0.02
JF 0269	Grape juice (from wine grapes)	PP	0.1	0.01	0.00	0.01	0.00	0.41	0.04	0.01	0.00	NC	-
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.13	0.31	0.04	0.23	0.03	60.43	7.86	0.52	0.07	31.91	4.15
FB 0265	Cranberry, raw	RAC	0.1	NC	-	NC	-	0.03	0.00	NC	-	NC	-
FB 0275	Strawberry, raw	RAC	0.24	0.01	0.00	0.01	0.00	3.35	0.80	0.01	0.00	0.01	0.00
FI 0326	Avocado, raw	RAC	0.13	1.12	0.15	0.01	0.00	0.84	0.11	0.01	0.00	6.60	0.86
FI 0350	Papaya, raw	RAC	0.31	6.47	2.01	0.25	0.08	0.19	0.06	0.01	0.00	26.42	8.19
VB 0400	Broccoli, raw	RAC	0.94	0.01	0.01	0.01	0.01	2.13	2.00	0.01	0.01	NC	-
VB 0041	Cabbages, head, raw	RAC	0.93	3.82	3.55	2.99	2.78	49.16	45.72	0.01	0.01	NC	-
VC 2039	Subgroup of Cucumbers and Squashes, raw	RAC	0.091	0.92	0.08	3.20	0.29	13.55	1.23	1.91	0.17	0.05	0.00
VC 0046	Melons, except watermelon, raw (Cantaloupe)	RAC	0.091	0.19	0.02	0.10	0.01	4.98	0.45	0.01	0.00	NC	-
VC 0429	Pumpkins, raw	RAC	0.091	0.56	0.05	6.14	0.56	4.59	0.42	11.70	1.06	NC	-
VO 0448	Tomato, raw	RAC	0.2	12.99	2.60	4.79	0.96	58.40	11.68	0.92	0.18	0.09	0.02
-	Tomato, canned (& peeled)	PP	0.06	0.07	0.00	0.08	0.00	2.42	0.15	0.07	0.00	NC	-
-	Tomato, paste (i.e. concentrated tomato sauce/puree)	PP	0.44	0.58	0.26	0.22	0.10	2.21	0.97	0.24	0.11	3.10	1.36
JF 0448	Tomato, juice (single strength, incl concentrated)	PP	0.06	0.05	0.00	0.01	0.00	0.42	0.03	0.01	0.00	0.02	0.00
VO 0444	Peppers, chili, raw	RAC	0.16	3.47	0.56	3.56	0.57	16.30	2.61	0.01	0.00	NC	-

Annex 3

METHOXYFENOZIDE (209)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
VO 0445	Peppers, sweet, raw	RAC	0.16	1.24	0.20	1.27	0.20	NC	-	0.01	0.00	NC	-
-	Peppers, sweet, dried	PP	1.6	0.58	0.93	1.27	2.03	1.21	1.94	0.12	0.19	NC	-
VL 0483	Lettuce, leaf, raw	RAC	12	0.29	3.48	0.03	0.36	6.71	80.52	0.01	0.12	NC	-
VL 0485	Mustard greens, raw (i.e. Indian mustard, Amsoi, mustard cabbage)	RAC	16	0.04	0.64	0.03	0.48	NC	-	0.01	0.16	NC	-
VL 0494	Radish leaves, raw	RAC	0.75	0.44	0.33	0.32	0.24	NC	-	0.30	0.23	0.59	0.44
VP 0061	Beans with pods (Phaseolus spp): (immature pods + succulent seeds)	RAC	0.065	NC	-	NC	-	NC	-	NC	-	NC	-
014B	Peas with pods	-	0.1	-	-	-	-	-	-	-	-	-	-
VP 0062	Beans without pods: (Phaseolus spp.) (succulent seeds), raw	RAC	0.05	0.30	0.02	3.13	0.16	4.11	0.21	0.01	0.00	NC	-
VP 0064	Peas without pods (Pisum spp) (succulent seeds)	RAC	0.05	0.21	0.01	0.02	0.00	5.51	0.28	0.02	0.00	NC	-
VD 0071	Beans, dry, raw (Phaseolus spp)	RAC	0.05	7.11	0.36	2.33	0.12	3.76	0.19	44.70	2.24	3.27	0.16
VD 0527	Cowpea, dry, raw (Vigna sinensis, Dolichos sinensis)	RAC	0.56	12.77	7.15	0.99	0.55	0.01	0.01	4.33	2.42	NC	-
VD 0072	Peas (dry) (Pisum spp), raw	RAC	0.17	1.53	0.26	2.52	0.43	3.52	0.60	3.56	0.61	0.74	0.13
VR 0577	Carrots, raw	RAC	0.13	2.07	0.27	3.00	0.39	25.29	3.29	0.05	0.01	NC	-
VR 0494	Radish roots, raw	RAC	0.08	3.96	0.32	2.86	0.23	3.30	0.26	2.67	0.21	5.34	0.43
VR 0596	Sugar beet, raw (incl sugar)	RAC	0.11	3.93	0.43	1.68	0.18	NC	-	NC	-	36.12	3.97
VR 0508	Sweet potato, raw (incl dried)	RAC	0.01	28.83	0.29	61.55	0.62	0.15	0.00	221.94	2.22	NC	-
VS 0624	Celery	RAC	3.4	3.66	12.44	2.65	9.01	4.84	16.46	2.47	8.40	4.94	16.80
GC 0645	Maize, raw (incl glucose & dextrose & isoglucose, incl flour, incl oil, incl beer, incl germ, incl starch)	RAC	0.02	116.66	2.33	10.52	0.21	38.46	0.77	76.60	1.53	34.44	0.69
GC 2090	Subgroup of Sweet Corns	RAC	0.02	3.63	0.07	20.50	0.41	8.78	0.18	0.02	0.00	0.17	0.00
GS 0659	Sugar cane, raw	RAC	0.01	5.62	0.06	50.91	0.51	NC	-	11.04	0.11	0.10	0.00
-	Sugar cane, molasses	PP	0.07	NC	-	NC	-	NC	-	NC	-	NC	-
-	Sugar cane, sugar (incl non-centrifugal sugar, incl refined sugar and maltose)	PP	0.06	28.13	1.69	55.38	3.32	78.09	4.69	18.04	1.08	45.60	2.74
TN 0085	Group of Tree nuts, raw (incl processed)	RAC	0.012	4.39	0.05	135.53	1.63	6.11	0.07	0.72	0.01	317.74	3.81
OR 0691	Cotton seed oil, edible	PP	0.15	1.28	0.19	0.05	0.01	0.45	0.07	0.42	0.06	0.15	0.02
SO 0697	Peanuts, nutmeat, raw (incl roasted, incl oil, incl butter)	RAC	0.01	18.82	0.19	0.57	0.01	2.28	0.02	6.90	0.07	0.53	0.01

METHOXYFENOZIDE (209)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
SO 0697	Peanuts, nutmeat, raw (incl roasted, incl butter, excl oil)	RAC	0.01	7.14	0.07	0.45	0.00	1.87	0.02	6.22	0.06	0.53	0.01
OR 0697	Peanut oil, edible	PP	0.029	5.02	0.15	0.05	0.00	0.17	0.00	0.29	0.01	NC	-
SB 0716	Coffee bean raw (incl roasted, incl instant coffee, incl substitutes)	RAC	0.028	0.95	0.03	1.32	0.04	11.64	0.33	2.96	0.08	14.73	0.41
HH 0722	Basil leaves, raw (incl dried)	RAC	19	0.25	4.75	0.18	3.42	0.13	2.47	0.17	3.23	0.33	6.27
HS 0444	Peppers, chili, dried	PP	1.6	0.58	0.93	1.27	2.03	1.21	1.94	0.12	0.19	NC	-
DT 1114	Tea, green or black, fermented and dried, (including concentrates)	RAC	29	0.53	15.37	5.25	152.25	0.86	24.94	0.56	16.24	0.88	25.52
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.003	23.34	0.07	40.71	0.12	97.15	0.29	18.06	0.05	57.71	0.17
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.036	5.84	0.21	10.18	0.37	24.29	0.87	4.52	0.16	14.43	0.52
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.036	1.05	0.04	1.14	0.04	18.69	0.67	0.94	0.03	3.12	0.11
MO 0105	Edible offal (mammalian), raw	RAC	0.025	4.64	0.12	1.97	0.05	10.01	0.25	3.27	0.08	3.98	0.10
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.03	108.75	3.26	70.31	2.11	436.11	13.08	61.55	1.85	79.09	2.37
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0	3.53	0.00	10.83	0.00	51.36	0.00	4.53	0.00	50.00	0.00
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0	0.39	0.00	1.20	0.00	5.71	0.00	0.50	0.00	5.56	0.00
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0	0.10	0.00	0.70	0.00	0.97	0.00	0.10	0.00	NC	-
PE 0112	Eggs, raw, (incl dried)	RAC	0	3.84	0.00	4.41	0.00	27.25	0.00	1.13	0.00	7.39	0.00
Total intake (ug/person)=				96.4		192.5		289.3		124.9		88.1	
Bodyweight per region (kg bw) =				60		60		60		60		60	
ADI (ug/person)=				6000		6000		6000		6000		6000	
%ADI=				1.6%		3.2%		4.8%		2.1%		1.5%	
Rounded %ADI=				2%		3%		5%		2%		1%	

Annex 3

PROTHIOCONAZOLE (232)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.01 mg/kg bw				ADI is for prothioconazole-desthio			
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day			Intake as ug/person/day								
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
FB 2006	Subgroup of Bush berries, raw (including processed)	RAC	0.52	0.53	0.28	1.31	0.68	0.40	0.21	1.66	0.86	0.01	0.01	0.99	0.51
FB 0265	Cranberry, raw	RAC	0.025	0.02	0.00	0.01	0.00	NC	-	0.03	0.00	0.01	0.00	0.01	0.00
VC 0045	Group of Fruiting vegetables, cucurbits, raw	RAC	0.045	53.14	2.39	86.21	3.88	6.28	0.28	92.76	4.17	15.64	0.70	155.30	6.99
VD 0523	Broad bean, dry, raw (incl horse-bean, field bean) (Vicia faba)	RAC	0.05	1.27	0.06	0.01	0.00	0.12	0.01	2.49	0.12	0.23	0.01	5.54	0.28
VD 0527	Cowpea, dry, raw (Vigna sinensis, Dolichos sinensis)	RAC	0.05	0.05	0.00	NC	-	1.74	0.09	0.01	0.00	0.01	0.00	0.07	0.00
VD 0541	Soya bean, dry, raw (incl paste, incl curd, incl oil, incl sauce)	RAC	0.009	72.79	0.66	59.05	0.53	20.55	0.18	74.20	0.67	61.12	0.55	73.24	0.66
-	Beans (dry) NES: including inter alia lablab or hyacinth bean (Dolichos spp.); jack or sword bean (Canavalia spp.); winged bean (Psophocarpus tetragonolobus); guar bean (Cyamopsis tetragonoloba); velvet bean (Stizolobium spp.); yam bean (Pachyrrhizus erosus)	RAC	0.05	1.70	0.09	0.01	0.00	3.00	0.15	1.80	0.09	1.64	0.08	1.33	0.07
VD 2066	Subgroup of dry peas, raw	RAC	0.05	9.09	0.45	3.35	0.17	1.06	0.05	9.48	0.47	15.11	0.76	10.58	0.53
VR 0596	Sugar beet, raw (incl sugar)	RAC	0.05	0.13	0.01	NC	-	0.08	0.00	0.66	0.03	0.47	0.02	88.94	4.45
VR 0589	Potato, raw (incl flour, incl frozen, incl starch, incl tapioca)	RAC	0.01	59.74	0.60	316.14	3.16	9.78	0.10	60.26	0.60	54.12	0.54	119.82	1.20
GC 0650	Rye, raw (incl flour)	RAC	0.01	0.13	0.00	19.38	0.19	0.10	0.00	0.12	0.00	0.03	0.00	2.15	0.02
GC 0653	Triticale, raw (incl flour)	RAC	0.01	NC	-	NC	-	NC	-	0.01	0.00	0.39	0.00	NC	-
GC 0654	Wheat, raw (incl bulgur, incl fermented beverages, excl germ, excl wholemeal bread, excl white flour products, excl white bread)	RAC	0.01	0.01	0.00	1.12	0.01	NC	-	0.03	0.00	0.56	0.01	NC	-
CF 1210	Wheat, germ	PP	0.04	NC	-	NC	-	0.01	0.00	0.01	0.00	0.14	0.01	0.01	0.00
CP 1212	Wheat, wholemeal bread	PP	0.01	0.01	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.03	0.00	0.01	0.00
CP 1211	Wheat, white bread	PP	0.01	0.25	0.00	0.63	0.01	0.12	0.00	0.43	0.00	1.39	0.01	0.22	0.00
CF 1211	Wheat, white flour (incl white flour products: starch, gluten, macaroni, pastry)	PP	0.01	301.24	3.01	268.64	2.69	30.21	0.30	222.51	2.23	134.73	1.35	343.12	3.43
GC 0640	Barley, raw (incl malt extract, incl pot&pearled, incl flour & grits, incl beer, incl malt)	RAC	0.035	19.91	0.70	31.16	1.09	5.04	0.18	3.10	0.11	9.77	0.34	4.31	0.15
GC 0647	Oats, raw (incl rolled)	RAC	0.01	0.05	0.00	7.05	0.07	0.10	0.00	1.71	0.02	0.96	0.01	0.04	0.00
GC 0645	Maize, raw (incl glucose & dextrose & isoglucose, incl starch, excl flour, excl oil,	RAC	0.01	0.97	0.01	0.24	0.00	0.70	0.01	4.09	0.04	2.33	0.02	13.31	0.13

PROTHIOCONAZOLE (232)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.01 mg/kg bw				ADI is for prothioconazole-desthio			
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day			Intake as ug/person/day								
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
	excl beer, excl germ)														
CF 1255	Maize, flour (white flour and wholemeal flour)	PP	0.0057	22.72	0.13	35.61	0.20	87.27	0.50	34.92	0.20	46.71	0.27	49.12	0.28
-	Maize, germ	PP	0.01	0.01	0.00	NC	-	0.01	0.00	0.01	0.00	0.22	0.00	NC	-
-	Maize starch	PP	0.0028	0.08	0.00	NC	-	0.01	0.00	2.29	0.01	0.08	0.00	0.11	0.00
-	Maize beer	PP	0.01	NC	-	NC	-	4.61	0.05	NC	-	NC	-	NC	-
OR 0645	Maize oil	PP	0.0028	0.96	0.00	0.85	0.00	0.29	0.00	5.42	0.02	0.42	0.00	2.10	0.01
GC 0447	Sweet corn on the cob, raw (incl frozen kernels, incl canned kernels)	RAC	0.018	0.14	0.00	0.94	0.02	5.70	0.10	2.61	0.05	1.94	0.03	0.22	0.00
SO 0693	Linseed, raw (incl oil)	RAC	0.014	0.02	0.00	NC	-	NC	-	0.01	0.00	0.13	0.00	NC	-
SO 0495	Rape seed, raw	RAC	0.039	0.02	0.00	NC	-	NC	-	0.01	0.00	0.75	0.03	0.01	0.00
OR 0495	Rape seed oil, edible	PP	0.0273	0.35	0.01	0.44	0.01	0.19	0.01	0.97	0.03	3.28	0.09	0.77	0.02
SO 2091	Subgroup of Sunflower seeds, raw	RAC	0.009	0.09	0.00	0.33	0.00	0.09	0.00	0.24	0.00	0.02	0.00	0.01	0.00
OR 0702	Sunflower seed oil, edible	PP	0.00963	2.97	0.03	14.42	0.14	0.43	0.00	3.46	0.03	2.20	0.02	5.53	0.05
SO 0691	Cotton seed, raw (incl oil)	RAC	0.052	20.53	1.07	9.80	0.51	6.42	0.33	4.73	0.25	7.14	0.37	18.68	0.97
OR 0691	Cotton seed oil, edible	PP	0.01	3.22	0.03	1.54	0.02	1.01	0.01	0.74	0.01	1.12	0.01	2.93	0.03
SO 0697	Peanuts, nutmeat, raw (incl roasted, incl oil, incl butter)	RAC	0.01	1.30	0.01	1.23	0.01	12.62	0.13	2.87	0.03	6.59	0.07	2.67	0.03
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.004	24.96	0.10	57.95	0.23	16.70	0.07	38.38	0.15	26.46	0.11	29.00	0.12
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.005	6.24	0.03	14.49	0.07	4.18	0.02	9.60	0.05	6.62	0.03	7.25	0.04
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.005	3.29	0.02	6.14	0.03	0.82	0.00	1.57	0.01	2.23	0.01	1.07	0.01
MO 0105	Edible offal (mammalian), raw	RAC	0.077	4.79	0.37	9.68	0.75	2.97	0.23	5.49	0.42	3.84	0.30	5.03	0.39
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.001	289.65	0.29	485.88	0.49	26.92	0.03	239.03	0.24	199.91	0.20	180.53	0.18
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0.00255	13.17	0.03	26.78	0.07	7.24	0.02	116.71	0.30	22.54	0.06	32.09	0.08
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0.0129	1.46	0.02	2.98	0.04	0.80	0.01	12.97	0.17	2.50	0.03	3.57	0.05
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.0129	0.10	0.00	0.10	0.00	NC	-	0.10	0.00	0.10	0.00	0.10	0.00
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.118	0.12	0.01	0.12	0.01	0.11	0.01	5.37	0.63	0.24	0.03	0.10	0.01

Annex 3

PROTHIOCONAZOLE (232)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.01 mg/kg bw				ADI is for prothioconazole-desthio				
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day			Intake as ug/person/day									
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake	
PE 0112	Eggs, raw, (incl dried)	RAC	0.00103	7.84	0.01	23.08	0.02	2.88	0.00	14.89	0.02	9.81	0.01	14.83	0.02	
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-	
Total intake (ug/person)=				10.4			15.1		3.1		12.0		6.1		20.7	
Bodyweight per region (kg bw) =				60			60		60		60		60		60	
ADI (ug/person)=				600			600		600		600		600		600	
%ADI=				1.7%			2.5%		0.5%		2.0%		1.0%		3.4%	
Rounded %ADI=				2%			3%		1%		2%		1%		3%	

PROTHIOCONAZOLE (232)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.01 mg/kg bw				ADI is for prothioconazole-desthio			
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day			Intake as ug/person/day								
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
FB 2006	Subgroup of Bush berries, raw (including processed)	RAC	0.52	1.31	0.68	5.50	2.86	0.01	0.01	2.57	1.34	0.82	0.43	2.15	1.12
FB 0265	Cranberry, raw	RAC	0.025	0.06	0.00	0.01	0.00	0.01	0.00	1.22	0.03	0.11	0.00	NC	-
VC 0045	Group of Fruiting vegetables, cucurbits, raw	RAC	0.045	27.81	1.25	41.93	1.89	123.30	5.55	49.47	2.23	15.95	0.72	35.99	1.62
VD 0523	Broad bean, dry, raw (incl horse-bean, field bean) (Vicia faba)	RAC	0.05	0.02	0.00	0.01	0.00	1.16	0.06	0.40	0.02	NC	-	0.06	0.00
VD 0527	Cowpea, dry, raw (Vigna sinensis, Dolichos sinensis)	RAC	0.05	NC	-	NC	-	0.16	0.01	0.01	0.00	NC	-	NC	-
VD 0541	Soya bean, dry, raw (incl paste, incl curd, incl oil, incl sauce)	RAC	0.009	106.33	0.96	117.78	1.06	42.12	0.38	195.70	1.76	222.52	2.00	80.47	0.72
-	Beans (dry) NES: including inter alia lablab or hyacinth bean (Dolichos spp.); jack or sword bean (Canavalia spp.); winged bean (Psophocarpus tetragonolobus); guar bean (Cyamopsis tetragonoloba); velvet bean (Stizolobium spp.); yam bean (Pachyrrhizus erosus)	RAC	0.05	0.01	0.00	NC	-	0.57	0.03	0.11	0.01	0.16	0.01	0.94	0.05
VD 2066	Subgroup of dry peas, raw	RAC	0.05	5.01	0.25	3.76	0.19	1.82	0.09	3.44	0.17	3.49	0.17	5.15	0.26
VR 0596	Sugar beet, raw (incl sugar)	RAC	0.05	0.01	0.00	NC	-	0.01	0.00	0.01	0.00	NC	-	NC	-

PROTHIOCONAZOLE (232)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.01 mg/kg bw				ADI is for prothioconazole-desthio			
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day			Intake as ug/person/day								
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
VR 0589	Potato, raw (incl flour, incl frozen, incl starch, incl tapioca)	RAC	0.01	225.03	2.25	234.24	2.34	71.48	0.71	177.55	1.78	234.55	2.35	37.71	0.38
GC 0650	Rye, raw (incl flour)	RAC	0.01	3.21	0.03	35.38	0.35	0.21	0.00	6.50	0.07	1.49	0.01	NC	-
GC 0653	Triticale, raw (incl flour)	RAC	0.01	0.01	0.00	0.17	0.00	0.29	0.00	0.01	0.00	NC	-	NC	-
GC 0654	Wheat, raw (incl bulgur, incl fermented beverages, excl germ, excl wholemeal bread, excl white flour products, excl white bread)	RAC	0.01	NC	-	NC	-	0.02	0.00	0.83	0.01	NC	-	NC	-
CF 1210	Wheat, germ	PP	0.04	0.97	0.04	0.10	0.00	0.03	0.00	0.01	0.00	NC	-	0.04	0.00
CP 1212	Wheat, wholemeal bread	PP	0.01	0.03	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.05	0.00	0.02	0.00
CP 1211	Wheat, white bread	PP	0.01	1.30	0.01	0.46	0.00	0.06	0.00	0.22	0.00	2.44	0.02	0.77	0.01
CF 1211	Wheat, white flour (incl white flour products: starch, gluten, macaroni, pastry)	PP	0.01	198.08	1.98	193.03	1.93	106.24	1.06	185.09	1.85	168.67	1.69	131.59	1.32
GC 0640	Barley, raw (incl malt extract, incl pot&pearled, incl flour & grits, incl beer, incl malt)	RAC	0.035	36.18	1.27	53.45	1.87	9.39	0.33	35.25	1.23	46.68	1.63	15.92	0.56
GC 0647	Oats, raw (incl rolled)	RAC	0.01	7.50	0.08	6.26	0.06	0.15	0.00	4.87	0.05	3.16	0.03	2.98	0.03
GC 0645	Maize, raw (incl glucose & dextrose & isoglucose, incl starch, excl flour, excl oil, excl beer, excl germ)	RAC	0.01	0.10	0.00	9.93	0.10	1.71	0.02	21.20	0.21	0.33	0.00	0.04	0.00
CF 1255	Maize, flour (white flour and wholemeal flour)	PP	0.0057	14.27	0.08	12.86	0.07	19.71	0.11	12.55	0.07	4.21	0.02	52.30	0.30
-	Maize, germ	PP	0.01	0.01	0.00	NC	-	NC	-	0.01	0.00	NC	-	0.01	0.00
-	Maize starch	PP	0.0028	NC	-	NC	-	0.19	0.00	7.13	0.02	NC	-	NC	-
-	Maize beer	PP	0.01	NC	-	NC	-	NC	-	1.99	0.02	NC	-	NC	-
OR 0645	Maize oil	PP	0.0028	0.90	0.00	0.47	0.00	0.15	0.00	3.01	0.01	1.86	0.01	0.36	0.00
GC 0447	Sweet corn on the cob, raw (incl frozen kernels, incl canned kernels)	RAC	0.018	11.43	0.21	3.71	0.07	0.74	0.01	13.63	0.25	3.07	0.06	1.50	0.03
SO 0693	Linseed, raw (incl oil)	RAC	0.014	NC	-	NC	-	0.02	0.00	0.01	0.00	NC	-	NC	-
SO 0495	Rape seed, raw	RAC	0.039	NC	-	NC	-	0.01	0.00	NC	-	NC	-	NC	-
OR 0495	Rape seed oil, edible	PP	0.0273	12.52	0.34	7.63	0.21	3.00	0.08	6.01	0.16	NC	-	NC	-
SO 2091	Subgroup of Sunflower seeds, raw	RAC	0.009	0.01	0.00	1.32	0.01	0.03	0.00	1.18	0.01	NC	-	0.02	0.00
OR	Sunflower seed oil, edible	PP	0.00963	9.50	0.09	11.37	0.11	0.49	0.00	5.15	0.05	2.63	0.03	2.80	0.03

Annex 3

PROTHIOCONAZOLE (232)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.01 mg/kg bw				ADI is for prothioconazole-desthio									
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day			Intake as ug/person/day														
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake						
0702																					
SO 0691	Cotton seed, raw (incl oil)	RAC	0.052	10.71	0.56	4.23	0.22	7.19	0.37	7.54	0.39	5.66	0.29	2.38	0.12						
OR 0691	Cotton seed oil, edible	PP	0.01	1.68	0.02	0.66	0.01	1.13	0.01	1.18	0.01	0.89	0.01	0.37	0.00						
SO 0697	Peanuts, nutmeat, raw (incl roasted, incl oil, incl butter)	RAC	0.01	5.63	0.06	2.75	0.03	9.58	0.10	5.82	0.06	13.71	0.14	1.84	0.02						
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.004	112.02	0.45	120.71	0.48	63.46	0.25	88.99	0.36	96.24	0.38	41.02	0.16						
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.005	28.01	0.14	30.18	0.15	15.86	0.08	22.25	0.11	24.06	0.12	10.25	0.05						
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.005	6.44	0.03	15.51	0.08	3.79	0.02	8.29	0.04	18.44	0.09	8.00	0.04						
MO 0105	Edible offal (mammalian), raw	RAC	0.077	15.17	1.17	5.19	0.40	6.30	0.49	6.78	0.52	3.32	0.26	3.17	0.24						
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.001	388.92	0.39	335.88	0.34	49.15	0.05	331.25	0.33	468.56	0.47	245.45	0.25						
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0.00255	66.38	0.17	48.47	0.12	21.58	0.06	78.41	0.20	48.04	0.12	76.01	0.19						
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0.0129	7.38	0.10	5.39	0.07	2.40	0.03	8.71	0.11	5.34	0.07	8.45	0.11						
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.0129	0.10	0.00	0.10	0.00	NC	-	0.10	0.00	0.71	0.01	NC	-						
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.118	0.33	0.04	0.72	0.08	0.27	0.03	0.35	0.04	0.80	0.09	NC	-						
PE 0112	Eggs, raw, (incl dried)	RAC	0.00103	25.84	0.03	29.53	0.03	28.05	0.03	33.19	0.03	36.44	0.04	8.89	0.01						
Total intake (ug/person)=					12.7			15.1			10.0			13.6			11.3		7.6		
Bodyweight per region (kg bw) =					60			60			55			60			60			60	
ADI (ug/person)=					600			600			550			600			600			600	
%ADI=					2.1%			2.5%			1.8%			2.3%			1.9%			1.3%	
Rounded %ADI=					2%			3%			2%			2%			2%			1%	

PROTHIOCONAZOLE (232)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.01 mg/kg bw		ADI is for prothioconazole-desthio			
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake		G16 diet	G16 intake	G17 diet
FB 2006	Subgroup of Bush berries, raw (including processed)	RAC	0.52	0.82	0.43	4.05	2.11	5.94	3.09	0.43	0.22	2.66	1.38
FB 0265	Cranberry, raw	RAC	0.025	NC	-	NC	-	0.03	0.00	NC	-	NC	-
VC 0045	Group of Fruiting vegetables, cucurbits, raw	RAC	0.045	5.96	0.27	9.74	0.44	51.82	2.33	13.61	0.61	0.05	0.00
VD 0523	Broad bean, dry, raw (incl horse-bean, field bean) (Vicia faba)	RAC	0.05	3.70	0.19	0.03	0.00	0.17	0.01	0.01	0.00	NC	-
VD 0527	Cowpea, dry, raw (Vigna sinensis, Dolichos sinensis)	RAC	0.05	12.77	0.64	0.99	0.05	0.01	0.00	4.33	0.22	NC	-
VD 0541	Soya bean, dry, raw (incl paste, incl curd, incl oil, incl sauce)	RAC	0.009	15.80	0.14	14.29	0.13	104.36	0.94	17.11	0.15	35.20	0.32
-	Beans (dry) NES: including inter alia lablab or hyacinth bean (Dolichos spp.); jack or sword bean (Canavalia spp.); winged bean (Psophocarpus tetragonolobus); guar bean (Cyamopsis tetragonoloba); velvet bean (Stizolobium spp.); yam bean (Pachyrrhizus erosus)	RAC	0.05	2.54	0.13	1.77	0.09	0.03	0.00	0.03	0.00	3.99	0.20
VD 2066	Subgroup of dry peas, raw	RAC	0.05	4.43	0.22	11.36	0.57	4.22	0.21	9.36	0.47	1.21	0.06
VR 0596	Sugar beet, raw (incl sugar)	RAC	0.05	3.93	0.20	1.68	0.08	NC	-	NC	-	36.12	1.81
VR 0589	Potato, raw (incl flour, incl frozen, incl starch, incl tapioca)	RAC	0.01	23.96	0.24	13.56	0.14	213.41	2.13	104.35	1.04	8.56	0.09
GC 0650	Rye, raw (incl flour)	RAC	0.01	0.03	0.00	0.01	0.00	13.95	0.14	0.01	0.00	0.88	0.01
GC 0653	Triticale, raw (incl flour)	RAC	0.01	0.01	0.00	NC	-	NC	-	NC	-	NC	-
GC 0654	Wheat, raw (incl bulgur, incl fermented beverages, excl germ, excl wholemeal bread, excl white flour products, excl white bread)	RAC	0.01	0.01	0.00	NC	-	NC	-	NC	-	0.97	0.01
CF 1210	Wheat, germ	PP	0.04	0.04	0.00	0.01	0.00	0.01	0.00	0.01	0.00	NC	-
CP 1212	Wheat, wholemeal bread	PP	0.01	0.01	0.00	0.01	0.00	0.03	0.00	0.01	0.00	0.01	0.00
CP 1211	Wheat, white bread	PP	0.01	0.43	0.00	0.41	0.00	1.56	0.02	0.11	0.00	0.07	0.00
CF 1211	Wheat, white flour (incl white flour products: starch, gluten, macaroni, pastry)	PP	0.01	44.78	0.45	86.96	0.87	214.05	2.14	20.31	0.20	103.60	1.04
GC 0640	Barley, raw (incl malt extract, incl pot&pearled, incl flour & grits, incl beer, incl malt)	RAC	0.035	11.58	0.41	2.33	0.08	46.71	1.63	3.72	0.13	16.26	0.57
GC 0647	Oats, raw (incl rolled)	RAC	0.01	0.37	0.00	0.07	0.00	2.79	0.03	0.10	0.00	NC	-
GC 0645	Maize, raw (incl glucose & dextrose & isoglucose, incl starch, excl flour, excl oil, excl beer, excl germ)	RAC	0.01	0.37	0.00	0.52	0.01	3.26	0.03	0.18	0.00	NC	-

Annex 3

PROTHIOCONAZOLE (232)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.01 mg/kg bw		ADI is for prothioconazole-desthio			
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake		G16 diet	G16 intake	G17 diet
CF 1255	Maize, flour (white flour and wholemeal flour)	PP	0.0057	94.34	0.54	8.09	0.05	28.03	0.16	55.94	0.32	28.07	0.16
-	Maize, germ	PP	0.01	0.01	0.00	NC	-	NC	-	NC	-	NC	-
-	Maize starch	PP	0.0028	0.02	0.00	0.01	0.00	NC	-	NC	-	NC	-
-	Maize beer	PP	0.01	1.03	0.01	NC	-	NC	-	40.94	0.41	NC	-
OR 0645	Maize oil	PP	0.0028	0.33	0.00	0.07	0.00	0.81	0.00	0.01	0.00	NC	-
GC 0447	Sweet corn on the cob, raw (incl frozen kernels, incl canned kernels)	RAC	0.018	3.63	0.07	20.50	0.37	8.78	0.16	0.02	0.00	0.17	0.00
SO 0693	Linseed, raw (incl oil)	RAC	0.014	0.07	0.00	NC	-	0.03	0.00	NC	-	NC	-
SO 0495	Rape seed, raw	RAC	0.039	NC	-	0.01	0.00	NC	-	NC	-	NC	-
OR 0495	Rape seed oil, edible	PP	0.0273	0.07	0.00	0.03	0.00	4.62	0.13	0.03	0.00	NC	-
SO 2091	Subgroup of Sunflower seeds, raw	RAC	0.009	0.06	0.00	0.01	0.00	0.03	0.00	2.23	0.02	NC	-
OR 0702	Sunflower seed oil, edible	PP	0.00963	0.37	0.00	0.09	0.00	12.98	0.12	4.01	0.04	0.20	0.00
SO 0691	Cotton seed, raw (incl oil)	RAC	0.052	8.14	0.42	0.32	0.02	2.84	0.15	2.69	0.14	0.97	0.05
OR 0691	Cotton seed oil, edible	PP	0.01	1.28	0.01	0.05	0.00	0.45	0.00	0.42	0.00	0.15	0.00
SO 0697	Peanuts, nutmeat, raw (incl roasted, incl oil, incl butter)	RAC	0.01	18.82	0.19	0.57	0.01	2.28	0.02	6.90	0.07	0.53	0.01
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.004	23.34	0.09	40.71	0.16	97.15	0.39	18.06	0.07	57.71	0.23
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.005	5.84	0.03	10.18	0.05	24.29	0.12	4.52	0.02	14.43	0.07
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.005	1.05	0.01	1.14	0.01	18.69	0.09	0.94	0.00	3.12	0.02
MO 0105	Edible offal (mammalian), raw	RAC	0.077	4.64	0.36	1.97	0.15	10.01	0.77	3.27	0.25	3.98	0.31
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.001	108.75	0.11	70.31	0.07	436.11	0.44	61.55	0.06	79.09	0.08
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0.00255	3.53	0.01	10.83	0.03	51.36	0.13	4.53	0.01	50.00	0.13
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0.0129	0.39	0.01	1.20	0.02	5.71	0.07	0.50	0.01	5.56	0.07
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.0129	NC	-	NC	-	0.32	0.00	NC	-	NC	-
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.118	0.10	0.01	0.70	0.08	0.97	0.11	0.10	0.01	NC	-
PE 0112	Eggs, raw, (incl dried)	RAC	0.00103	3.84	0.00	4.41	0.00	27.25	0.03	1.13	0.00	7.39	0.01
-	-	-		-	-	-	-	-	-	-	-	-	-
Total intake (ug/person)=				5.2		5.6		15.6		4.5		6.6	

PROTHIOCONAZOLE (232)				International Estimated Daily Intake (IEDI)			ADI = 0 - 0.01 mg/kg bw			ADI is for prothioconazole-desthio			
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
	Bodyweight per region (kg bw) =				60		60		60		60		60
	ADI (ug/person)=				600		600		600		600		600
	%ADI=				0.9%		0.9%		2.6%		0.8%		1.1%
	Rounded %ADI=				1%		1%		3%		1%		1%

Annex 3

PYDIFLUMETOFEN (309)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day			Intake as ug/person/day								
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
FC 0001	Group of Citrus fruit, raw (incl kumquat commodities)	RAC	0.21	32.25	6.77	11.67	2.45	16.70	3.51	76.01	15.96	33.90	7.12	92.97	19.52
JF 0001	Group of Citrus fruit, juice	PP	0.004	1.30	0.01	2.37	0.01	0.22	0.00	13.88	0.06	0.75	0.00	2.63	0.01
-	Lemon, juice (single strength, incl. concentrated)	PP	0.004	0.01	0.00	0.01	0.00	0.11	0.00	0.09	0.00	0.18	0.00	0.17	0.00
-	Subgroup of Mandarins, juice	PP	0.004	NC	-	NC	-	NC	-	NC	-	NC	-	NC	-
JF 0004	Subgroup of Oranges, juice (single strength, incl. concentrated)	PP	0.004	1.27	0.01	2.20	0.01	0.09	0.00	11.81	0.05	0.46	0.00	1.69	0.01
JF 0203	Grapefruits, juice (single strength, incl. concentrated)	PP	0.004	0.01	0.00	0.16	0.00	0.02	0.00	1.97	0.01	0.12	0.00	0.77	0.00
FP 0009	Group of Pomefruits, raw	RAC	0.06	19.24	1.15	33.89	2.03	3.34	0.20	25.53	1.53	7.59	0.46	56.76	3.41
JF 0226	Apple juice, single strength (incl. concentrated)	PP	0.004	0.32	0.00	3.07	0.01	0.07	0.00	5.00	0.02	0.29	0.00	5.57	0.02
-	Apple cider (i.e. fermented apple juice)	PP	0.004	0.07	0.00	0.12	0.00	10.66	0.04	0.15	0.00	0.07	0.00	0.07	0.00
FS 0013	Subgroup of Cherries, raw	RAC	0.395	0.92	0.36	9.15	3.61	0.01	0.00	0.61	0.24	0.06	0.02	6.64	2.62
FS 0014	Subgroup of Plums, raw	RAC	0.15	2.40	0.36	8.60	1.29	0.06	0.01	2.52	0.38	0.58	0.09	4.16	0.62
DF 0014	Plums, dried (prunes)	PP	0.34	0.09	0.03	0.06	0.02	0.01	0.00	0.18	0.06	0.04	0.01	0.06	0.02
FS 2001	Subgroup of peaches, raw (incl dried apricots)	RAC	0.21	8.01	1.68	5.87	1.23	0.18	0.04	8.19	1.72	1.64	0.34	22.46	4.72
FB 2006	Subgroup of Bush berries, raw (including processed)	RAC	0.88	0.53	0.47	1.31	1.15	0.40	0.35	1.66	1.46	0.01	0.01	0.99	0.87
FB 0267	Elderberries, raw (incl processed)	RAC	0.88	0.44	0.39	0.27	0.24	0.34	0.30	1.41	1.24	NC	-	0.87	0.77
FB 0269	Grapes, raw (i.e. table grapes)	RAC	0.29	12.68	3.68	9.12	2.64	0.03	0.01	16.88	4.90	3.70	1.07	54.42	15.78
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	0.71	0.51	0.36	0.51	0.36	0.01	0.01	1.27	0.90	0.12	0.09	2.07	1.47
JF 0269	Grape juice (from wine grapes)	PP	0.017	0.14	0.00	0.29	0.00	0.05	0.00	0.30	0.01	0.24	0.00	0.05	0.00
-	Graps must (from wine-grapes)	PP	0.31	0.33	0.10	0.13	0.04	0.01	0.00	0.02	0.01	0.01	0.00	0.02	0.01
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.091	0.67	0.06	12.53	1.14	2.01	0.18	1.21	0.11	3.53	0.32	4.01	0.36
FB 0275	Strawberry, raw	RAC	0.185	0.70	0.13	2.01	0.37	0.04	0.01	1.36	0.25	0.37	0.07	2.53	0.47
VA 2031	Subgroup of bulb onions	RAC	0.07	31.65	2.22	43.28	3.03	3.68	0.26	38.48	2.69	20.46	1.43	47.29	3.31
VA 2032	Subgroup of Green Onions	RAC	0.36	2.64	0.95	3.09	1.11	1.05	0.38	2.89	1.04	0.61	0.22	5.24	1.89
VB 0042	Subgroup of Flowerhead Brassica, raw	RAC	0.39	2.54	0.99	0.49	0.19	0.01	0.00	3.57	1.39	7.79	3.04	3.12	1.22
VB 2036	Subgroup of Head Brassicas, raw	RAC	0.065	3.81	0.25	38.89	2.53	0.77	0.05	6.23	0.40	4.27	0.28	14.28	0.93

PYDIFLUMETOFEN (309)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
VB 2016	Subgroup of Stem Brassicas, raw	RAC	0.02	0.09	0.00	0.89	0.02	0.02	0.00	0.14	0.00	NC	-	0.33	0.01
VC 0045	Group of Fruiting vegetables, cucurbits, raw	RAC	0.12	53.14	6.38	86.21	10.35	6.28	0.75	92.76	11.13	15.64	1.88	155.30	18.64
VO 0448	Tomato, raw	RAC	0.11	41.73	4.59	75.65	8.32	10.66	1.17	82.87	9.12	24.75	2.72	200.93	22.10
-	Tomato, canned (& peeled)	PP	0.005	0.20	0.00	0.31	0.00	0.02	0.00	1.11	0.01	0.11	0.00	1.50	0.01
-	Tomato, paste (i.e. concentrated tomato sauce/puree)	PP	0.075	2.34	0.18	1.33	0.10	1.57	0.12	4.24	0.32	0.34	0.03	2.83	0.21
JF 0448	Tomato, juice (single strength, incl concentrated)	PP	0.005	0.29	0.00	0.29	0.00	0.01	0.00	0.38	0.00	0.05	0.00	0.14	0.00
VO 0051	Subgroup of peppers, raw (Capsicum spp. Only), excl okra	RAC	0.11	5.42	0.60	9.91	1.09	3.97	0.44	7.63	0.84	2.59	0.28	23.68	2.60
VO 0442	Okra, raw (i.e. Lady's Finger, Gombo)	RAC	0.02	1.97	0.04	NC	-	3.68	0.07	3.24	0.06	5.72	0.11	1.57	0.03
VO 2046	Subgroup of eggplants	RAC	0.11	5.58	0.61	4.31	0.47	0.89	0.10	9.31	1.02	13.64	1.50	20.12	2.21
VL 2050	Subgroup of Leafy greens	RAC	12.5	3.93	49.13	5.28	66.00	3.07	38.38	14.53	181.63	8.25	103.13	12.75	159.38
VL 0054	Subgroup of Leaves of Brassicaceae, raw	RAC	0.02	2.63	0.05	9.27	0.19	1.86	0.04	5.82	0.12	19.53	0.39	4.90	0.10
VL 2052	Subgroup of Leaves of Root and Tuber Vegetables	RAC	0.02	0.18	0.00	0.31	0.01	0.84	0.02	0.47	0.01	2.06	0.04	0.23	0.00
VP 2060	Subgroup of beans with pods (all commodities within this group)	RAC	0.045	0.68	0.03	NC	-	NC	-	0.39	0.02	0.22	0.01	0.49	0.02
014B	Peas with pods	-	0.12	-	-	-	-	-	-	-	-	-	-	-	-
VP 2062	Subgroup of succulent beans without pods (all commodities within this group)	RAC	0.033	5.07	0.17	1.02	0.03	0.49	0.02	1.78	0.06	1.19	0.04	8.57	0.28
VP 2063	Subgroup of succulent peas without pods	RAC	0.031	1.97	0.06	0.51	0.02	0.02	0.00	0.79	0.02	3.68	0.11	3.80	0.12
014E	Underground immature beans and peas	-	0.02	-	-	-	-	-	-	-	-	-	-	-	-
VD 0071	Beans, dry, raw (Phaseolus spp)	RAC	0.028	2.39	0.07	1.61	0.05	10.47	0.29	1.84	0.05	12.90	0.36	7.44	0.21
VD 0523	Broad bean, dry, raw (incl horse-bean, field bean) (Vicia faba)	RAC	0.028	1.27	0.04	0.01	0.00	0.12	0.00	2.49	0.07	0.23	0.01	5.54	0.16
VD 0527	Cowpea, dry, raw (Vigna sinensis, Dolichos sinensis)	RAC	0.028	0.05	0.00	NC	-	1.74	0.05	0.01	0.00	0.01	0.00	0.07	0.00
VD 0541	Soya bean, dry, raw (Glycine soja)	RAC	0.028	0.58	0.02	0.05	0.00	0.37	0.01	0.03	0.00	1.65	0.05	0.30	0.01
-	Soya paste (i.e. miso)	PP	0.004	NC	-	NC	-	NC	-	NC	-	NC	-	NC	-
-	Soya curd (i.e. tofu)	PP	0.004	NC	-	NC	-	NC	-	NC	-	NC	-	NC	-
OR 0541	Soya oil, refined	PP	0.005	12.99	0.06	10.43	0.05	3.63	0.02	13.10	0.07	10.70	0.05	13.10	0.07
-	Soya sauce	PP	0.002	0.01	0.00	0.02	0.00	0.01	0.00	0.34	0.00	0.03	0.00	0.01	0.00

Annex 3

PYDIFLUMETOFEN (309)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
-	Soya flour	PP	0.002	0.05	0.00	0.86	0.00	0.02	0.00	1.02	0.00	0.01	0.00	0.15	0.00
-	Beans (dry) NES: including inter alia lablab or hyacinth bean (Dolichos spp.); jack or sword bean (Canavalia spp.); winged bean (Psophocarpus tetragonolobus); guar bean (Cyamopsis tetragonoloba); velvet bean (Stizolobium spp.); yam bean (Pachyrhizus erosus)	RAC	0.028	1.70	0.05	0.01	0.00	3.00	0.08	1.80	0.05	1.64	0.05	1.33	0.04
VD 2066	Subgroup of dry peas, raw	RAC	0.028	9.09	0.25	3.35	0.09	1.06	0.03	9.48	0.27	15.11	0.42	10.58	0.30
VR 2070	Subgroup of Root vegetables, raw	RAC	0.08	24.72	1.98	57.71	4.62	17.01	1.36	49.58	3.97	9.33	0.75	114.41	9.15
-	Sugar beet, sugar	PP	0.004	0.02	0.00	NC	-	0.01	0.00	0.09	0.00	0.07	0.00	12.63	0.05
VR 0573	Arrowroot, raw	RAC	0.03	1.53	0.05	0.01	0.00	0.93	0.03	1.33	0.04	0.47	0.01	0.02	0.00
VR 0463	Cassava raw (incl starch, incl tapioca, incl flour) (i.e. Manioc)	RAC	0.03	0.08	0.00	0.01	0.00	482.56	14.48	0.99	0.03	25.75	0.77	3.29	0.10
VR 0585	Jerusalem artichoke, raw (i.e. Topinambur)	RAC	0.03	1.57	0.05	0.01	0.00	0.96	0.03	1.36	0.04	0.48	0.01	0.02	0.00
VR 0589	Potato, raw (incl frozen, incl tapioca, excl flour, excl starch)	RAC	0.03	59.36	1.78	315.64	9.47	9.38	0.28	55.58	1.67	53.71	1.61	119.55	3.59
-	Potato, flour	PP	0.014	0.05	0.00	0.10	0.00	0.09	0.00	0.88	0.01	0.09	0.00	0.06	0.00
-	Potato, starch	PP	0.014	0.03	0.00	0.01	0.00	0.01	0.00	0.15	0.00	0.01	0.00	0.01	0.00
VR 0508	Sweet potato, raw (incl dried)	RAC	0.03	0.18	0.01	0.18	0.01	42.16	1.26	1.61	0.05	3.06	0.09	6.67	0.20
VR 0504	Tannia, raw (i.e. Tanier, Yautia)	RAC	0.03	NC	-	NC	-	NC	-	0.01	0.00	0.26	0.01	1.27	0.04
VR 0505	Taro, raw (i.e. Dasheen, Eddoe)	RAC	0.03	0.01	0.00	NC	-	25.12	0.75	0.04	0.00	0.01	0.00	0.97	0.03
VR 0600	Yams, raw (incl dried)	RAC	0.03	0.02	0.00	NC	-	90.40	2.71	6.45	0.19	0.74	0.02	0.65	0.02
VS 2080	Subgroup of stems and petioles	RAC	4.4	3.11	13.68	5.52	24.29	3.42	15.05	8.29	36.48	0.02	0.09	4.00	17.60
GC 0648	Quinoa, raw	RAC	0.063	NC	-	NC	-	NC	-	NC	-	0.07	0.00	NC	-
GC 0650	Rye, raw (incl flour)	RAC	0.063	0.13	0.01	19.38	1.22	0.10	0.01	0.12	0.01	0.03	0.00	2.15	0.14
GC 0653	Triticale, raw (incl flour)	RAC	0.063	NC	-	NC	-	NC	-	0.01	0.00	0.39	0.02	NC	-
GC 0654	Wheat, raw (incl bulgur, incl fermented beverages, excl germ, excl wholemeal bread, excl white flour products, excl white bread)	RAC	0.063	0.01	0.00	1.12	0.07	NC	-	0.03	0.00	0.56	0.04	NC	-
CF 1210	Wheat, germ	PP	0.091	NC	-	NC	-	0.01	0.00	0.01	0.00	0.14	0.01	0.01	0.00
CP 1212	Wheat, wholemeal bread	PP	0.027	0.01	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.03	0.00	0.01	0.00

PYDIFLUMETOFEN (309)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
CP 1211	Wheat, white bread	PP	0.02	0.25	0.01	0.63	0.01	0.12	0.00	0.43	0.01	1.39	0.03	0.22	0.00
CF 1211	Wheat, white flour	PP	0.02	299.27	5.99	263.32	5.27	27.93	0.56	214.18	4.28	133.47	2.67	340.03	6.80
-	Wheat, starch	PP	0.002	0.02	0.00	NC	-	0.01	0.00	0.05	0.00	0.13	0.00	0.01	0.00
-	Wheat, gluten	PP	0.11	0.01	0.00	0.01	0.00	0.01	0.00	0.27	0.03	0.01	0.00	0.03	0.00
-	Wheat, macaroni, dry	PP	0.02	0.72	0.01	2.20	0.04	1.22	0.02	3.99	0.08	0.53	0.01	1.66	0.03
-	Wheat, pastry, baked	PP	0.02	1.21	0.02	3.13	0.06	1.05	0.02	4.02	0.08	0.60	0.01	1.40	0.03
GC 0640	Barley, raw (incl malt extract, incl beer, incl malt, excl pot&pearled, excl flour & grits)	RAC	0.23	3.55	0.82	19.31	4.44	4.98	1.15	3.02	0.69	7.85	1.81	3.98	0.92
-	Barley, pot&pearled	PP	0.01	7.12	0.07	7.34	0.07	0.02	0.00	0.03	0.00	0.67	0.01	0.20	0.00
-	Barley, flour (white flour and wholemeal flour)	PP	0.053	2.93	0.16	0.30	0.02	0.02	0.00	0.01	0.00	0.48	0.03	0.01	0.00
GC 0641	Buckwheat, raw (incl flour)	RAC	0.23	NC	-	0.40	0.09	0.01	0.00	0.01	0.00	0.07	0.02	0.09	0.02
GC 0647	Oats, raw	RAC	0.23	0.01	0.00	NC	-	0.01	0.00	0.45	0.10	0.01	0.00	0.01	0.00
GC 0647	Oats, rolled (i.e. oatmeal dry)	PP	0.003	0.03	0.00	3.88	0.01	0.05	0.00	0.69	0.00	0.53	0.00	0.02	0.00
GC 2088	Subgroup of rice cereals	REP	0.03	45.40	1.36	14.99	0.45	84.88	2.55	111.73	3.35	194.75	5.84	93.12	2.79
GC 0646	Millet, raw (incl flour, incl beer)	RAC	0.03	1.46	0.04	2.32	0.07	5.84	0.18	0.89	0.03	16.17	0.49	0.01	0.00
GC 0651	Sorghum, raw (incl beer, excl flour)(i.e. Chicken corn, Dari seed, Durra, Feterita)	RAC	0.515	NC	-	0.01	0.01	3.34	1.72	0.01	0.01	NC	-	NC	-
-	Sorghum, flour (white flour and wholemeal flour)	PP	0.45	3.91	1.76	NC	-	11.62	5.23	14.24	6.41	9.87	4.44	2.62	1.18
GC 0645	Maize, raw (incl glucose & dextrose & isoglucose, incl beer, excl flour, excl oil, excl germ, excl starch)	RAC	0.03	0.84	0.03	0.24	0.01	1.56	0.05	0.46	0.01	2.21	0.07	13.13	0.39
CF 1255	Maize, flour (white flour and wholemeal flour)	PP	0.048	22.72	1.09	35.61	1.71	87.27	4.19	34.92	1.68	46.71	2.24	49.12	2.36
-	Maize, germ	PP	0.063	0.01	0.00	NC	-	0.01	0.00	0.01	0.00	0.22	0.01	NC	-
-	Maize starch	PP	0.013	0.08	0.00	NC	-	0.01	0.00	2.29	0.03	0.08	0.00	0.11	0.00
OR 0645	Maize oil	PP	0.057	0.96	0.05	0.85	0.05	0.29	0.02	5.42	0.31	0.42	0.02	2.10	0.12
-	Cereals, NES, raw (including processed) : canagua, quihuicha, Job's tears and wild rice	RAC	0.03	2.04	0.06	2.99	0.09	1.86	0.06	19.17	0.58	3.33	0.10	1.66	0.05
GC 2090	Subgroup of Sweet Corns	RAC	0.03	0.14	0.00	0.94	0.03	5.70	0.17	2.61	0.08	1.94	0.06	0.22	0.01
TN 0085	Group of Tree nuts, raw (incl processed)	RAC	0.01	4.06	0.04	3.27	0.03	7.01	0.07	13.93	0.14	14.01	0.14	9.36	0.09
SO 0090	Subgroup of Mustard seeds, raw (incl flour, incl oil)	RAC	0.0945	0.02	0.00	0.05	0.00	0.01	0.00	0.31	0.03	0.03	0.00	0.04	0.00

Annex 3

PYDIFLUMETOFEN (309)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
SO 0693	Linseed, raw (incl oil)	RAC	0.0945	0.02	0.00	NC	-	NC	-	0.01	0.00	0.13	0.01	NC	-
SO 0698	Poppy seed, raw (incl oil)	RAC	0.0945	0.01	0.00	0.01	0.00	0.01	0.00	0.03	0.00	0.01	0.00	0.01	0.00
SO 0495	Rape seed, raw	RAC	0.0945	0.02	0.00	NC	-	NC	-	0.01	0.00	0.75	0.07	0.01	0.00
OR 0495	Rape seed oil, edible	PP	0.035	0.35	0.01	0.44	0.02	0.19	0.01	0.97	0.03	3.28	0.11	0.77	0.03
SO 0700	Sesame seed, raw (incl oil)	RAC	0.0945	1.22	0.12	0.01	0.00	0.54	0.05	4.23	0.40	0.82	0.08	2.77	0.26
SO 2091	Subgroup of Sunflower seeds, raw (incl processed)	RAC	0.09	7.43	0.67	36.06	3.25	1.15	0.10	8.77	0.79	5.74	0.52	13.63	1.23
OR 0702	Sunflower seed oil, edible	PP	0.005	2.97	0.01	14.42	0.07	0.43	0.00	3.46	0.02	2.20	0.01	5.53	0.03
OR 0691	Cotton seed oil, edible	PP	0.0006	3.22	0.00	1.54	0.00	1.01	0.00	0.74	0.00	1.12	0.00	2.93	0.00
SO 0697	Peanuts, nutmeat, raw (incl roasted, incl butter, excl oil)	RAC	0.03	0.46	0.01	1.21	0.04	6.64	0.20	2.71	0.08	1.26	0.04	1.84	0.06
OR 0697	Peanut oil, edible	PP	0.072	0.36	0.03	0.01	0.00	2.57	0.19	0.07	0.01	2.29	0.16	0.36	0.03
HS 0444	Peppers, chili, dried	PP	1.1	0.42	0.46	0.53	0.58	0.84	0.92	0.50	0.55	0.95	1.05	0.37	0.41
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.02	24.96	0.50	57.95	1.16	16.70	0.33	38.38	0.77	26.46	0.53	29.00	0.58
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.02	6.24	0.12	14.49	0.29	4.18	0.08	9.60	0.19	6.62	0.13	7.25	0.15
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.02	3.29	0.07	6.14	0.12	0.82	0.02	1.57	0.03	2.23	0.04	1.07	0.02
MO 0105	Edible offal (mammalian), raw	RAC	0.09	4.79	0.43	9.68	0.87	2.97	0.27	5.49	0.49	3.84	0.35	5.03	0.45
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.02	289.65	5.79	485.88	9.72	26.92	0.54	239.03	4.78	199.91	4.00	180.53	3.61
PM 0110	Poultry meat, raw (incl prepared)	RAC	0.02	14.63	0.29	29.76	0.60	8.04	0.16	129.68	2.59	25.04	0.50	35.66	0.71
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.02	0.10	0.00	0.10	0.00	NC	-	0.10	0.00	0.10	0.00	0.10	0.00
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.02	0.12	0.00	0.12	0.00	0.11	0.00	5.37	0.11	0.24	0.00	0.10	0.00
PE 0112	Eggs, raw, (incl dried)	RAC	0.02	7.84	0.16	23.08	0.46	2.88	0.06	14.89	0.30	9.81	0.20	14.83	0.30
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-
Total intake (ug/person)=				120.1		179.6		101.8		310.9		155.0		318.1	
Bodyweight per region (kg bw) =				60		60		60		60		60		60	
ADI (ug/person)=				6000		6000		6000		6000		6000		6000	
%ADI=				2.0%		3.0%		1.7%		5.2%		2.6%		5.3%	

PYDIFLUMETOFEN (309)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
	Rounded %ADI=				2%		3%		2%		5%		3%		5%

PYDIFLUMETOFEN (309)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
FC 0001	Group of Citrus fruit, raw (incl kumquat commodities)	RAC	0.21	38.66	8.12	54.93	11.54	26.36	5.54	51.46	10.81	51.06	10.72	466.36	97.94
JF 0001	Group of Citrus fruit, juice	PP	0.004	36.84	0.15	3.75	0.02	0.30	0.00	21.62	0.09	21.82	0.09	46.67	0.19
-	Lemon, juice (single strength, incl. concentrated)	PP	0.004	0.60	0.00	0.36	0.00	0.01	0.00	1.49	0.01	0.43	0.00	0.24	0.00
-	Subgroup of Mandarins, juice	PP	0.004	0.04	0.00	NC	-	0.01	0.00	0.01	0.00	NC	-	NC	-
JF 0004	Subgroup of Oranges, juice (single strength, incl. concentrated)	PP	0.004	33.31	0.13	1.78	0.01	0.28	0.00	18.97	0.08	14.01	0.06	13.36	0.05
JF 0203	Grapefruits, juice (single strength, incl. concentrated)	PP	0.004	2.89	0.01	1.61	0.01	0.02	0.00	1.15	0.00	7.39	0.03	33.07	0.13
FP 0009	Group of Pomefruits, raw	RAC	0.06	37.39	2.24	58.13	3.49	37.64	2.26	44.80	2.69	62.17	3.73	6.47	0.39
JF 0226	Apple juice, single strength (incl. concentrated)	PP	0.004	14.88	0.06	11.98	0.05	0.15	0.00	9.98	0.04	30.32	0.12	3.47	0.01
-	Apple cider (i.e. fermented apple juice)	PP	0.004	10.05	0.04	5.34	0.02	3.72	0.01	0.36	0.00	0.25	0.00	0.93	0.00
FS 0013	Subgroup of Cherries, raw	RAC	0.395	1.40	0.55	4.21	1.66	0.04	0.02	2.93	1.16	1.50	0.59	NC	-
FS 0014	Subgroup of Plums, raw	RAC	0.15	3.75	0.56	3.33	0.50	5.94	0.89	2.64	0.40	2.50	0.38	0.06	0.01
DF 0014	Plums, dried (prunes)	PP	0.34	0.61	0.21	0.35	0.12	0.05	0.02	0.35	0.12	0.49	0.17	0.13	0.04
FS 2001	Subgroup of peaches, raw (incl dried apricots)	RAC	0.21	13.03	2.74	16.29	3.42	8.29	1.74	12.95	2.72	5.35	1.12	0.04	0.01
FB 2006	Subgroup of Bush berries, raw (including processed)	RAC	0.88	1.31	1.15	5.50	4.84	0.01	0.01	2.57	2.26	0.82	0.72	2.15	1.89
FB 0267	Elderberries, raw (incl processed)	RAC	0.88	8.20	7.22	0.14	0.12	NC	-	NC	-	NC	-	1.87	1.65
FB 0269	Grapes, raw (i.e. table grapes)	RAC	0.29	6.33	1.84	11.22	3.25	5.21	1.51	9.38	2.72	4.55	1.32	0.78	0.23
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	0.71	3.09	2.19	1.51	1.07	0.03	0.02	1.38	0.98	4.26	3.02	0.42	0.30
JF 0269	Grape juice (from wine grapes)	PP	0.017	0.56	0.01	1.96	0.03	0.02	0.00	2.24	0.04	2.27	0.04	0.34	0.01
-	Graps must (from wine-grapes)	PP	0.31	0.16	0.05	0.09	0.03	0.01	0.00	0.12	0.04	0.11	0.03	NC	-

Annex 3

PYDIFLUMETOFEN (309)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.091	88.93	8.09	62.41	5.68	1.84	0.17	25.07	2.28	61.17	5.57	5.84	0.53
FB 0275	Strawberry, raw	RAC	0.185	4.49	0.83	5.66	1.05	0.02	0.00	6.63	1.23	5.75	1.06	0.05	0.01
VA 2031	Subgroup of bulb onions	RAC	0.07	20.67	1.45	31.32	2.19	37.52	2.63	35.08	2.46	11.77	0.82	13.74	0.96
VA 2032	Subgroup of Green Onions	RAC	0.36	5.57	2.01	5.15	1.85	1.77	0.64	4.28	1.54	17.34	6.24	6.48	2.33
VB 0042	Subgroup of Flowerhead Brassica, raw	RAC	0.39	9.50	3.71	6.77	2.64	NC	-	3.21	1.25	9.36	3.65	0.75	0.29
VB 2036	Subgroup of Head Brassicas, raw	RAC	0.065	11.20	0.73	29.79	1.94	25.06	1.63	34.72	2.26	8.73	0.57	15.64	1.02
VB 2016	Subgroup of Stem Brassicas, raw	RAC	0.02	NC	-	3.25	0.07	NC	-	NC	-	0.02	0.00	0.36	0.01
VC 0045	Group of Fruiting vegetables, cucurbits, raw	RAC	0.12	27.81	3.34	41.93	5.03	123.30	14.80	49.47	5.94	15.95	1.91	35.99	4.32
VO 0448	Tomato, raw	RAC	0.11	32.13	3.53	51.27	5.64	34.92	3.84	73.37	8.07	15.15	1.67	8.88	0.98
-	Tomato, canned (& peeled)	PP	0.005	7.57	0.04	2.66	0.01	0.30	0.00	0.97	0.00	7.31	0.04	0.41	0.00
-	Tomato, paste (i.e. concentrated tomato sauce/puree)	PP	0.075	4.96	0.37	3.20	0.24	0.15	0.01	1.61	0.12	6.88	0.52	0.52	0.04
JF 0448	Tomato, juice (single strength, incl concentrated)	PP	0.005	0.80	0.00	0.07	0.00	0.05	0.00	0.61	0.00	0.40	0.00	0.08	0.00
VO 0051	Subgroup of peppers, raw (Capsicum spp. Only), excl okra	RAC	0.11	5.57	0.61	14.00	1.54	16.50	1.82	8.80	0.97	6.44	0.71	3.44	0.38
VO 0442	Okra, raw (i.e. Lady's Finger, Gombo)	RAC	0.02	NC	-	NC	-	0.04	0.00	0.17	0.00	NC	-	0.72	0.01
VO 2046	Subgroup of eggplants	RAC	0.11	1.01	0.11	1.69	0.19	21.37	2.35	3.00	0.33	1.40	0.15	NC	-
VL 2050	Subgroup of Leafy greens	RAC	12.5	18.38	229.75	18.73	234.13	82.36	1029.50	25.32	316.50	17.60	220.00	7.37	92.13
VL 0054	Subgroup of Leaves of Brassicaceae, raw	RAC	0.02	0.10	0.00	NC	-	26.78	0.54	5.00	0.10	0.58	0.01	5.68	0.11
VL 2052	Subgroup of Leaves of Root and Tuber Vegetables	RAC	0.02	NC	-	NC	-	NC	-	NC	-	NC	-	0.33	0.01
VP 2060	Subgroup of beans with pods (all commodities within this group)	RAC	0.045	5.07	0.23	0.83	0.04	0.17	0.01	3.70	0.17	NC	-	NC	-
014B	Peas with pods	-	0.12	-	-	-	-	-	-	-	-	-	-	-	-
VP 2062	Subgroup of succulent beans without pods (all commodities within this group)	RAC	0.033	2.42	0.08	6.09	0.20	4.33	0.14	2.09	0.07	18.99	0.63	0.17	0.01
VP 2063	Subgroup of succulent peas without pods	RAC	0.031	10.72	0.33	1.99	0.06	2.72	0.08	4.26	0.13	4.23	0.13	NC	-
014E	Underground immature beans and peas	-	0.02	-	-	-	-	-	-	-	-	-	-	-	-
VD 0071	Beans, dry, raw (Phaseolus spp)	RAC	0.028	1.51	0.04	1.50	0.04	1.90	0.05	5.11	0.14	1.36	0.04	23.43	0.66

PYDIFLUMETOFEN (309)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
VD 0523	Broad bean, dry, raw (incl horse-bean, field bean) (Vicia faba)	RAC	0.028	0.02	0.00	0.01	0.00	1.16	0.03	0.40	0.01	NC	-	0.06	0.00
VD 0527	Cowpea, dry, raw (Vigna sinensis, Dolichos sinensis)	RAC	0.028	NC	-	NC	-	0.16	0.00	0.01	0.00	NC	-	NC	-
VD 0541	Soya bean, dry, raw (Glycine soja)	RAC	0.028	0.02	0.00	0.33	0.01	6.64	0.19	3.94	0.11	NC	-	5.78	0.16
-	Soya paste (i.e. miso)	PP	0.004	NC	-	NC	-	NC	-	1.87	0.01	NC	-	NC	-
-	Soya curd (i.e. tofu)	PP	0.004	NC	-	NC	-	0.68	0.00	0.87	0.00	NC	-	NC	-
OR 0541	Soya oil, refined	PP	0.005	19.06	0.10	21.06	0.11	5.94	0.03	33.78	0.17	40.05	0.20	13.39	0.07
-	Soya sauce	PP	0.002	0.45	0.00	0.29	0.00	2.93	0.01	4.35	0.01	0.09	0.00	0.70	0.00
-	Soya flour	PP	0.002	0.22	0.00	0.27	0.00	0.29	0.00	0.17	0.00	NC	-	NC	-
-	Beans (dry) NES: including inter alia lablab or hyacinth bean (Dolichos spp.); jack or sword bean (Canavalia spp.); winged bean (Psophocarpus tetragonolobus); guar bean (Cyamopsis tetragonoloba); velvet bean (Stizolobium spp.); yam bean (Pachyrrhizus erosus)	RAC	0.028	0.01	0.00	NC	-	0.57	0.02	0.11	0.00	0.16	0.00	0.94	0.03
VD 2066	Subgroup of dry peas, raw	RAC	0.028	5.01	0.14	3.76	0.11	1.82	0.05	3.44	0.10	3.49	0.10	5.15	0.14
VR 2070	Subgroup of Root vegetables, raw	RAC	0.08	64.22	5.14	65.78	5.26	49.73	3.98	57.68	4.61	113.82	9.11	37.27	2.98
-	Sugar beet, sugar	PP	0.004	0.01	0.00	NC	-	0.01	0.00	NC	-	NC	-	NC	-
VR 0573	Arrowroot, raw	RAC	0.03	0.02	0.00	0.01	0.00	2.05	0.06	0.21	0.01	NC	-	0.76	0.02
VR 0463	Cassava raw (incl starch, incl tapioca, incl flour) (i.e. Manioc)	RAC	0.03	0.01	0.00	NC	-	20.96	0.63	0.14	0.00	NC	-	9.62	0.29
VR 0585	Jerusalem artichoke, raw (i.e. Topinambur)	RAC	0.03	0.11	0.00	0.01	0.00	NC	-	0.22	0.01	NC	-	0.78	0.02
VR 0589	Potato, raw (incl frozen, incl tapioca, excl flour, excl starch)	RAC	0.03	221.34	6.64	224.14	6.72	70.42	2.11	172.24	5.17	227.39	6.82	37.41	1.12
-	Potato, flour	PP	0.014	0.81	0.01	0.48	0.01	0.19	0.00	0.25	0.00	1.57	0.02	0.07	0.00
-	Potato, starch	PP	0.014	NC	-	1.74	0.02	0.05	0.00	0.92	0.01	NC	-	NC	-
VR 0508	Sweet potato, raw (incl dried)	RAC	0.03	0.93	0.03	0.32	0.01	64.65	1.94	5.37	0.16	0.30	0.01	3.13	0.09
VR 0504	Tannia, raw (i.e. Tanier, Yautia)	RAC	0.03	NC	-	NC	-	NC	-	0.01	0.00	NC	-	10.74	0.32
VR 0505	Taro, raw (i.e. Dasheen, Eddoe)	RAC	0.03	NC	-	NC	-	1.93	0.06	0.84	0.03	NC	-	19.94	0.60
VR 0600	Yams, raw (incl dried)	RAC	0.03	NC	-	NC	-	0.03	0.00	0.71	0.02	NC	-	17.57	0.53

Annex 3

PYDIFLUMETOFEN (309)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
VS 2080	Subgroup of stems and petioles	RAC	4.4	9.31	40.96	8.57	37.71	NC	-	3.88	17.07	24.46	107.62	5.89	25.92
GC 0648	Quinoa, raw	RAC	0.063	NC	-	NC	-	NC	-	NC	-	NC	-	NC	-
GC 0650	Rye, raw (incl flour)	RAC	0.063	3.21	0.20	35.38	2.23	0.21	0.01	6.50	0.41	1.49	0.09	NC	-
GC 0653	Triticale, raw (incl flour)	RAC	0.063	0.01	0.00	0.17	0.01	0.29	0.02	0.01	0.00	NC	-	NC	-
GC 0654	Wheat, raw (incl bulgur, incl fermented beverages, excl germ, excl wholemeal bread, excl white flour products, excl white bread)	RAC	0.063	NC	-	NC	-	0.02	0.00	0.83	0.05	NC	-	NC	-
CF 1210	Wheat, germ	PP	0.091	0.97	0.09	0.10	0.01	0.03	0.00	0.01	0.00	NC	-	0.04	0.00
CP 1212	Wheat, wholemeal bread	PP	0.027	0.03	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.05	0.00	0.02	0.00
CP 1211	Wheat, white bread	PP	0.02	1.30	0.03	0.46	0.01	0.06	0.00	0.22	0.00	2.44	0.05	0.77	0.02
CF 1211	Wheat, white flour	PP	0.02	182.77	3.66	187.54	3.75	103.82	2.08	180.42	3.61	164.00	3.28	118.84	2.38
-	Wheat, starch	PP	0.002	NC	-	NC	-	0.01	0.00	0.31	0.00	NC	-	NC	-
-	Wheat, gluten	PP	0.11	0.68	0.07	NC	-	0.01	0.00	0.01	0.00	NC	-	NC	-
-	Wheat, macaroni, dry	PP	0.02	6.71	0.13	4.98	0.10	2.12	0.04	1.90	0.04	2.89	0.06	4.12	0.08
-	Wheat, pastry, baked	PP	0.02	7.93	0.16	0.51	0.01	0.29	0.01	2.44	0.05	1.78	0.04	8.64	0.17
GC 0640	Barley, raw (incl malt extract, incl beer, incl malt, excl pot&pearled, excl flour & grits)	RAC	0.23	35.17	8.09	49.45	11.37	8.86	2.04	34.31	7.89	44.87	10.32	15.82	3.64
-	Barley, pot&pearled	PP	0.01	0.57	0.01	2.56	0.03	0.33	0.00	0.56	0.01	0.36	0.00	NC	-
-	Barley, flour (white flour and wholemeal flour)	PP	0.053	0.08	0.00	0.03	0.00	0.01	0.00	0.05	0.00	0.68	0.04	0.05	0.00
GC 0641	Buckwheat, raw (incl flour)	RAC	0.23	0.01	0.00	0.79	0.18	0.18	0.04	0.35	0.08	NC	-	NC	-
GC 0647	Oats, raw	RAC	0.23	NC	-	NC	-	0.01	0.00	0.01	0.00	NC	-	0.23	0.05
GC 0647	Oats, rolled (i.e. oatmeal dry)	PP	0.003	4.12	0.01	3.44	0.01	0.08	0.00	2.67	0.01	1.74	0.01	1.51	0.00
GC 2088	Subgroup of rice cereals	REP	0.03	20.96	0.63	16.04	0.48	339.67	10.19	75.51	2.27	16.86	0.51	86.13	2.58
GC 0646	Millet, raw (incl flour, incl beer)	RAC	0.03	0.03	0.00	0.16	0.00	1.75	0.05	0.69	0.02	NC	-	NC	-
GC 0651	Sorghum, raw (incl beer, excl flour)(i.e. Chicken corn, Dari seed, Durra, Feterita)	RAC	0.515	NC	-	NC	-	0.01	0.01	1.15	0.59	NC	-	7.12	3.67
-	Sorghum, flour (white flour and wholemeal flour)	PP	0.45	NC	-	NC	-	1.29	0.58	0.01	0.00	NC	-	NC	-
GC 0645	Maize, raw (incl glucose & dextrose & isoglucose, incl beer, excl flour, excl oil, excl germ, excl starch)	RAC	0.03	0.10	0.00	9.93	0.30	1.40	0.04	10.26	0.31	0.33	0.01	0.04	0.00
CF 1255	Maize, flour (white flour and wholemeal flour)	PP	0.048	14.27	0.68	12.86	0.62	19.71	0.95	12.55	0.60	4.21	0.20	52.30	2.51

PYDIFLUMETOFEN (309)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
-	Maize, germ	PP	0.063	0.01	0.00	NC	-	NC	-	0.01	0.00	NC	-	0.01	0.00
-	Maize starch	PP	0.013	NC	-	NC	-	0.19	0.00	7.13	0.09	NC	-	NC	-
OR 0645	Maize oil	PP	0.057	0.90	0.05	0.47	0.03	0.15	0.01	3.01	0.17	1.86	0.11	0.36	0.02
-	Cereals, NES, raw (including processed) : canagua, quihuicha, Job's tears and wild rice	RAC	0.03	6.17	0.19	3.01	0.09	0.76	0.02	3.30	0.10	3.38	0.10	15.84	0.48
GC 2090	Subgroup of Sweet Corns	RAC	0.03	11.43	0.34	3.71	0.11	0.74	0.02	13.63	0.41	3.07	0.09	1.50	0.05
TN 0085	Group of Tree nuts, raw (incl processed)	RAC	0.01	8.52	0.09	8.94	0.09	15.09	0.15	9.60	0.10	14.57	0.15	26.26	0.26
SO 0090	Subgroup of Mustard seeds, raw (incl flour, incl oil)	RAC	0.0945	0.30	0.03	0.48	0.05	0.33	0.03	0.63	0.06	1.03	0.10	0.40	0.04
SO 0693	Linseed, raw (incl oil)	RAC	0.0945	NC	-	NC	-	0.02	0.00	0.01	0.00	NC	-	NC	-
SO 0698	Poppy seed, raw (incl oil)	RAC	0.0945	0.02	0.00	0.25	0.02	0.01	0.00	0.02	0.00	NC	-	NC	-
SO 0495	Rape seed, raw	RAC	0.0945	NC	-	NC	-	0.01	0.00	NC	-	NC	-	NC	-
OR 0495	Rape seed oil, edible	PP	0.035	12.52	0.44	7.63	0.27	3.00	0.11	6.01	0.21	NC	-	NC	-
SO 0700	Sesame seed, raw (incl oil)	RAC	0.0945	0.61	0.06	0.09	0.01	1.53	0.14	0.85	0.08	0.08	0.01	0.14	0.01
SO 2091	Subgroup of Sunflower seeds, raw (incl processed)	RAC	0.09	23.43	2.11	29.34	2.64	1.24	0.11	14.00	1.26	6.48	0.58	6.91	0.62
OR 0702	Sunflower seed oil, edible	PP	0.005	9.50	0.05	11.37	0.06	0.49	0.00	5.15	0.03	2.63	0.01	2.80	0.01
OR 0691	Cotton seed oil, edible	PP	0.0006	1.68	0.00	0.66	0.00	1.13	0.00	1.18	0.00	0.89	0.00	0.37	0.00
SO 0697	Peanuts, nutmeat, raw (incl roasted, incl butter, excl oil)	RAC	0.03	3.26	0.10	2.22	0.07	5.38	0.16	4.85	0.15	1.54	0.05	1.82	0.05
OR 0697	Peanut oil, edible	PP	0.072	1.02	0.07	0.23	0.02	1.81	0.13	0.42	0.03	5.23	0.38	0.01	0.00
HS 0444	Peppers, chili, dried	PP	1.1	0.11	0.12	0.21	0.23	0.36	0.40	0.21	0.23	0.25	0.28	0.15	0.17
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.02	112.02	2.24	120.71	2.41	63.46	1.27	88.99	1.78	96.24	1.92	41.02	0.82
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.02	28.01	0.56	30.18	0.60	15.86	0.32	22.25	0.44	24.06	0.48	10.25	0.21
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.02	6.44	0.13	15.51	0.31	3.79	0.08	8.29	0.17	18.44	0.37	8.00	0.16
MO 0105	Edible offal (mammalian), raw	RAC	0.09	15.17	1.37	5.19	0.47	6.30	0.57	6.78	0.61	3.32	0.30	3.17	0.29
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.02	388.92	7.78	335.88	6.72	49.15	0.98	331.25	6.63	468.56	9.37	245.45	4.91
PM	Poultry meat, raw (incl prepared)	RAC	0.02	73.76	1.48	53.86	1.08	23.98	0.48	87.12	1.74	53.38	1.07	84.45	1.69

Annex 3

PYDIFLUMETOFEN (309)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
0110															
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.02	0.10	0.00	0.10	0.00	NC	-	0.10	0.00	0.71	0.01	NC	-
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.02	0.33	0.01	0.72	0.01	0.27	0.01	0.35	0.01	0.80	0.02	NC	-
PE 0112	Eggs, raw, (incl dried)	RAC	0.02	25.84	0.52	29.53	0.59	28.05	0.56	33.19	0.66	36.44	0.73	8.89	0.18
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-
Total intake (ug/person)=				367.0		377.6		1101.0		426.1		420.5		264.0	
Bodyweight per region (kg bw) =				60		60		55		60		60		60	
ADI (ug/person)=				6000		6000		5500		6000		6000		6000	
%ADI=				6.1%		6.3%		20.0%		7.1%		7.0%		4.4%	
Rounded %ADI=				6%		6%		20%		7%		7%		4%	

PYDIFLUMETOFEN (309)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day				Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake		
FC 0001	Group of Citrus fruit, raw (incl kumquat commodities)	RAC	0.21	20.93	4.40	2.35	0.49	30.71	6.45	0.15	0.03	4.45	0.93		
JF 0001	Group of Citrus fruit, juice	PP	0.004	0.11	0.00	0.29	0.00	13.55	0.05	0.14	0.00	0.33	0.00		
-	Lemon, juice (single strength, incl. concentrated)	PP	0.004	0.01	0.00	0.01	0.00	0.16	0.00	0.01	0.00	NC	-		
-	Subgroup of Mandarins, juice	PP	0.004	0.01	0.00	NC	-	NC	-	NC	-	NC	-		
JF 0004	Subgroup of Oranges, juice (single strength, incl. concentrated)	PP	0.004	0.08	0.00	0.26	0.00	12.61	0.05	0.14	0.00	0.33	0.00		
JF 0203	Grapefruits, juice (single strength, incl. concentrated)	PP	0.004	0.03	0.00	0.02	0.00	0.78	0.00	0.01	0.00	NC	-		
FP 0009	Group of Pomefruits, raw	RAC	0.06	2.39	0.14	10.93	0.66	69.47	4.17	1.59	0.10	19.56	1.17		
JF 0226	Apple juice, single strength (incl. concentrated)	PP	0.004	0.03	0.00	0.10	0.00	7.19	0.03	0.03	0.00	NC	-		
-	Apple cider (i.e. fermented apple juice)	PP	0.004	48.75	0.20	0.01	0.00	0.99	0.00	138.03	0.55	NC	-		
FS 0013	Subgroup of Cherries, raw	RAC	0.395	0.01	0.00	0.01	0.00	5.96	2.35	0.01	0.00	NC	-		

PYDIFLUMETOFEN (309)		International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
FS 0014	Subgroup of Plums, raw	RAC	0.15	0.07	0.01	0.01	0.00	15.56	2.33	0.01	0.00	NC	-
DF 0014	Plums, dried (prunes)	PP	0.34	0.01	0.00	0.01	0.00	0.37	0.13	0.01	0.00	NC	-
FS 2001	Subgroup of peaches, raw (incl dried apricots)	RAC	0.21	0.02	0.00	0.01	0.00	10.76	2.26	0.01	0.00	NC	-
FB 2006	Subgroup of Bush berries, raw (including processed)	RAC	0.88	0.82	0.72	4.05	3.56	5.94	5.23	0.43	0.38	2.66	2.34
FB 0267	Elderberries, raw (incl processed)	RAC	0.88	0.71	0.62	3.52	3.10	NC	-	0.38	0.33	2.32	2.04
FB 0269	Grapes, raw (i.e. table grapes)	RAC	0.29	0.14	0.04	0.36	0.10	15.22	4.41	0.01	0.00	0.09	0.03
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	0.71	0.01	0.01	0.13	0.09	1.06	0.75	0.01	0.01	0.03	0.02
JF 0269	Grape juice (from wine grapes)	PP	0.017	0.01	0.00	0.01	0.00	0.41	0.01	0.01	0.00	NC	-
-	Graps must (from wine-grapes)	PP	0.31	0.01	0.00	0.01	0.00	0.11	0.03	0.01	0.00	0.19	0.06
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.091	0.31	0.03	0.23	0.02	60.43	5.50	0.52	0.05	31.91	2.90
FB 0275	Strawberry, raw	RAC	0.185	0.01	0.00	0.01	0.00	3.35	0.62	0.01	0.00	0.01	0.00
VA 2031	Subgroup of bulb onions	RAC	0.07	9.83	0.69	22.30	1.56	34.69	2.43	9.65	0.68	2.39	0.17
VA 2032	Subgroup of Green Onions	RAC	0.36	1.45	0.52	1.50	0.54	1.42	0.51	0.01	0.00	6.30	2.27
VB 0042	Subgroup of Flowerhead Brassica, raw	RAC	0.39	0.02	0.01	0.02	0.01	4.86	1.90	0.01	0.00	NC	-
VB 2036	Subgroup of Head Brassicas, raw	RAC	0.065	5.31	0.35	4.16	0.27	52.05	3.38	0.01	0.00	NC	-
VB 2016	Subgroup of Stem Brassicas, raw	RAC	0.02	0.12	0.00	0.10	0.00	1.81	0.04	0.01	0.00	NC	-
VC 0045	Group of Fruiting vegetables, cucurbits, raw	RAC	0.12	5.96	0.72	9.74	1.17	51.82	6.22	13.61	1.63	0.05	0.01
VO 0448	Tomato, raw	RAC	0.11	12.99	1.43	4.79	0.53	58.40	6.42	0.92	0.10	0.09	0.01
-	Tomato, canned (& peeled)	PP	0.005	0.07	0.00	0.08	0.00	2.42	0.01	0.07	0.00	NC	-
-	Tomato, paste (i.e. concentrated tomato sauce/puree)	PP	0.075	0.58	0.04	0.22	0.02	2.21	0.17	0.24	0.02	3.10	0.23
JF 0448	Tomato, juice (single strength, incl concentrated)	PP	0.005	0.05	0.00	0.01	0.00	0.42	0.00	0.01	0.00	0.02	0.00
VO 0051	Subgroup of peppers, raw (Capsicum spp. Only), excl okra	RAC	0.11	4.72	0.52	4.83	0.53	16.30	1.79	0.01	0.00	NC	-
VO 0442	Okra, raw (i.e. Lady's Finger, Gombo)	RAC	0.02	6.23	0.12	0.10	0.00	NC	-	NC	-	NC	-
VO 2046	Subgroup of eggplants	RAC	0.11	1.31	0.14	8.26	0.91	3.95	0.43	0.01	0.00	NC	-
VL 2050	Subgroup of Leafy greens	RAC	12.5	4.99	62.38	3.29	41.13	7.53	94.13	3.05	38.13	6.09	76.13
VL 0054	Subgroup of Leaves of Brassicaceae, raw	RAC	0.02	3.58	0.07	2.64	0.05	NC	-	1.83	0.04	3.65	0.07

Annex 3

PYDIFLUMETOFEN (309)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
VL 2052	Subgroup of Leaves of Root and Tuber Vegetables	RAC	0.02	0.30	0.01	0.22	0.00	NC	-	0.20	0.00	0.41	0.01
VP 2060	Subgroup of beans with pods (all commodities within this group)	RAC	0.045	NC	-	NC	-	NC	-	NC	-	NC	-
014B	Peas with pods	-	0.12	-	-	-	-	-	-	-	-	-	-
VP 2062	Subgroup of succulent beans without pods (all commodities within this group)	RAC	0.033	0.37	0.01	3.14	0.10	4.88	0.16	0.01	0.00	NC	-
VP 2063	Subgroup of succulent peas without pods	RAC	0.031	0.21	0.01	0.02	0.00	5.51	0.17	0.02	0.00	NC	-
014E	Underground immature beans and peas	-	0.02	-	-	-	-	-	-	-	-	-	-
VD 0071	Beans, dry, raw (Phaseolus spp)	RAC	0.028	7.11	0.20	2.33	0.07	3.76	0.11	44.70	1.25	3.27	0.09
VD 0523	Broad bean, dry, raw (incl horse-bean, field bean) (Vicia faba)	RAC	0.028	3.70	0.10	0.03	0.00	0.17	0.00	0.01	0.00	NC	-
VD 0527	Cowpea, dry, raw (Vigna sinensis, Dolichos sinensis)	RAC	0.028	12.77	0.36	0.99	0.03	0.01	0.00	4.33	0.12	NC	-
VD 0541	Soya bean, dry, raw (Glycine soja)	RAC	0.028	2.76	0.08	0.07	0.00	0.33	0.01	3.16	0.09	NC	-
-	Soya paste (i.e. miso)	PP	0.004	NC	-	NC	-	NC	-	NC	-	NC	-
-	Soya curd (i.e. tofu)	PP	0.004	NC	-	NC	-	NC	-	NC	-	NC	-
OR 0541	Soya oil, refined	PP	0.005	2.32	0.01	2.54	0.01	18.70	0.09	2.51	0.01	6.29	0.03
-	Soya sauce	PP	0.002	0.01	0.00	0.13	0.00	0.17	0.00	0.01	0.00	0.56	0.00
-	Soya flour	PP	0.002	0.11	0.00	0.08	0.00	0.07	0.00	0.01	0.00	0.03	0.00
-	Beans (dry) NES: including inter alia lablab or hyacinth bean (Dolichos spp.); jack or sword bean (Canavalia spp.); winged bean (Psophocarpus tetragonolobus); guar bean (Cyamopsis tetragonoloba); velvet bean (Stizolobium spp.); yam bean (Pachyrrhizus erosus)	RAC	0.028	2.54	0.07	1.77	0.05	0.03	0.00	0.03	0.00	3.99	0.11
VD 2066	Subgroup of dry peas, raw	RAC	0.028	4.43	0.12	11.36	0.32	4.22	0.12	9.36	0.26	1.21	0.03
VR 2070	Subgroup of Root vegetables, raw	RAC	0.08	31.84	2.55	23.38	1.87	68.28	5.46	17.52	1.40	71.01	5.68
-	Sugar beet, sugar	PP	0.004	0.56	0.00	0.24	0.00	NC	-	NC	-	5.13	0.02
VR 0573	Arrowroot, raw	RAC	0.03	13.83	0.41	18.24	0.55	0.01	0.00	0.05	0.00	19.60	0.59
VR 0463	Cassava raw (incl starch, incl tapioca, incl flour) (i.e. Manioc)	RAC	0.03	91.92	2.76	34.12	1.02	NC	-	259.92	7.80	45.48	1.36
VR 0585	Jerusalem artichoke, raw (i.e. Topinambur)	RAC	0.03	14.22	0.43	18.75	0.56	0.01	0.00	0.06	0.00	20.14	0.60
VR 0589	Potato, raw (incl frozen, incl tapioca, excl flour, excl	RAC	0.03	23.72	0.71	12.61	0.38	211.06	6.33	101.89	3.06	8.53	0.26

PYDIFLUMETOFEN (309)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
	starch)												
-	Potato, flour	PP	0.014	0.05	0.00	0.20	0.00	0.52	0.01	0.54	0.01	0.01	0.00
-	Potato, starch	PP	0.014	0.01	0.00	0.01	0.00	NC	-	NC	-	NC	-
VR 0508	Sweet potato, raw (incl dried)	RAC	0.03	28.83	0.86	61.55	1.85	0.15	0.00	221.94	6.66	NC	-
VR 0504	Tannia, raw (i.e. Tanier, Yautia)	RAC	0.03	NC	-	NC	-	0.01	0.00	NC	-	NC	-
VR 0505	Taro, raw (i.e. Dasheen, Eddoe)	RAC	0.03	6.71	0.20	31.91	0.96	NC	-	10.73	0.32	264.31	7.93
VR 0600	Yams, raw (incl dried)	RAC	0.03	70.93	2.13	30.62	0.92	0.07	0.00	5.65	0.17	30.85	0.93
VS 2080	Subgroup of stems and petioles	RAC	4.4	5.33	23.45	3.85	16.94	5.80	25.52	3.60	15.84	7.20	31.68
GC 0648	Quinoa, raw	RAC	0.063	NC	-	NC	-	NC	-	NC	-	NC	-
GC 0650	Rye, raw (incl flour)	RAC	0.063	0.03	0.00	0.01	0.00	13.95	0.88	0.01	0.00	0.88	0.06
GC 0653	Triticale, raw (incl flour)	RAC	0.063	0.01	0.00	NC	-	NC	-	NC	-	NC	-
GC 0654	Wheat, raw (incl bulgur, incl fermented beverages, excl germ, excl wholemeal bread, excl white flour products, excl white bread)	RAC	0.063	0.01	0.00	NC	-	NC	-	NC	-	0.97	0.06
CF 1210	Wheat, germ	PP	0.091	0.04	0.00	0.01	0.00	0.01	0.00	0.01	0.00	NC	-
CP 1212	Wheat, wholemeal bread	PP	0.027	0.01	0.00	0.01	0.00	0.03	0.00	0.01	0.00	0.01	0.00
CP 1211	Wheat, white bread	PP	0.02	0.43	0.01	0.41	0.01	1.56	0.03	0.11	0.00	0.07	0.00
CF 1211	Wheat, white flour	PP	0.02	43.75	0.88	85.81	1.72	206.68	4.13	19.38	0.39	92.92	1.86
-	Wheat, starch	PP	0.002	0.01	0.00	0.02	0.00	NC	-	NC	-	NC	-
-	Wheat, gluten	PP	0.11	0.01	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.19	0.02
-	Wheat, macaroni, dry	PP	0.02	0.52	0.01	0.63	0.01	2.99	0.06	0.26	0.01	5.18	0.10
-	Wheat, pastry, baked	PP	0.02	0.51	0.01	0.51	0.01	4.36	0.09	0.67	0.01	5.32	0.11
GC 0640	Barley, raw (incl malt extract, incl beer, incl malt, excl pot&pearled, excl flour & grits)	RAC	0.23	3.15	0.72	2.31	0.53	43.92	10.10	3.72	0.86	16.26	3.74
-	Barley, pot&pearled	PP	0.01	5.46	0.05	0.01	0.00	1.44	0.01	0.01	0.00	NC	-
-	Barley, flour (white flour and wholemeal flour)	PP	0.053	0.02	0.00	NC	-	0.32	0.02	0.01	0.00	NC	-
GC 0641	Buckwheat, raw (incl flour)	RAC	0.23	0.04	0.01	2.82	0.65	0.01	0.00	0.01	0.00	NC	-
GC 0647	Oats, raw	RAC	0.23	0.01	0.00	0.02	0.00	NC	-	0.09	0.02	NC	-
GC 0647	Oats, rolled (i.e. oatmeal dry)	PP	0.003	0.20	0.00	0.03	0.00	1.54	0.00	0.01	0.00	NC	-

Annex 3

PYDIFLUMETOFEN (309)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
GC 2088	Subgroup of rice cereals	REP	0.03	52.55	1.58	286.02	8.58	18.64	0.56	19.67	0.59	75.09	2.25
GC 0646	Millet, raw (incl flour, incl beer)	RAC	0.03	61.13	1.83	0.78	0.02	NC	-	33.55	1.01	NC	-
GC 0651	Sorghum, raw (incl beer, excl flour)(i.e. Chicken corn, Dari seed, Durra, Feterita)	RAC	0.515	4.73	2.44	NC	-	NC	-	13.36	6.88	NC	-
-	Sorghum, flour (white flour and wholemeal flour)	PP	0.45	75.99	34.20	1.82	0.82	NC	-	19.82	8.92	NC	-
GC 0645	Maize, raw (incl glucose & dextrose & isoglucose, incl beer, excl flour, excl oil, excl germ, excl starch)	RAC	0.03	0.54	0.02	0.51	0.02	3.26	0.10	7.96	0.24	NC	-
CF 1255	Maize, flour (white flour and wholemeal flour)	PP	0.048	94.34	4.53	8.09	0.39	28.03	1.35	55.94	2.69	28.07	1.35
-	Maize, germ	PP	0.063	0.01	0.00	NC	-	NC	-	NC	-	NC	-
-	Maize starch	PP	0.013	0.02	0.00	0.01	0.00	NC	-	NC	-	NC	-
OR 0645	Maize oil	PP	0.057	0.33	0.02	0.07	0.00	0.81	0.05	0.01	0.00	NC	-
-	Cereals, NES, raw (including processed) : canagua, quihuicha, Job's tears and wild rice	RAC	0.03	17.71	0.53	2.00	0.06	9.61	0.29	0.45	0.01	4.55	0.14
GC 2090	Subgroup of Sweet Corns	RAC	0.03	3.63	0.11	20.50	0.62	8.78	0.26	0.02	0.00	0.17	0.01
TN 0085	Group of Tree nuts, raw (incl processed)	RAC	0.01	4.39	0.04	135.53	1.36	6.11	0.06	0.72	0.01	317.74	3.18
SO 0090	Subgroup of Mustard seeds, raw (incl flour, incl oil)	RAC	0.0945	0.04	0.00	0.19	0.02	0.32	0.03	0.06	0.01	0.01	0.00
SO 0693	Linseed, raw (incl oil)	RAC	0.0945	0.07	0.01	NC	-	0.03	0.00	NC	-	NC	-
SO 0698	Poppy seed, raw (incl oil)	RAC	0.0945	0.01	0.00	0.01	0.00	0.11	0.01	NC	-	NC	-
SO 0495	Rape seed, raw	RAC	0.0945	NC	-	0.01	0.00	NC	-	NC	-	NC	-
OR 0495	Rape seed oil, edible	PP	0.035	0.07	0.00	0.03	0.00	4.62	0.16	0.03	0.00	NC	-
SO 0700	Sesame seed, raw (incl oil)	RAC	0.0945	2.34	0.22	0.66	0.06	0.26	0.02	9.84	0.93	NC	-
SO 2091	Subgroup of Sunflower seeds, raw (incl processed)	RAC	0.09	0.99	0.09	0.22	0.02	32.01	2.88	12.12	1.09	0.48	0.04
OR 0702	Sunflower seed oil, edible	PP	0.005	0.37	0.00	0.09	0.00	12.98	0.06	4.01	0.02	0.20	0.00
OR 0691	Cotton seed oil, edible	PP	0.0006	1.28	0.00	0.05	0.00	0.45	0.00	0.42	0.00	0.15	0.00
SO 0697	Peanuts, nutmeat, raw (incl roasted, incl butter, excl oil)	RAC	0.03	7.14	0.21	0.45	0.01	1.87	0.06	6.22	0.19	0.53	0.02
OR 0697	Peanut oil, edible	PP	0.072	5.02	0.36	0.05	0.00	0.17	0.01	0.29	0.02	NC	-
HS 0444	Peppers, chili, dried	PP	1.1	0.58	0.64	1.27	1.40	1.21	1.33	0.12	0.13	NC	-
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.02	23.34	0.47	40.71	0.81	97.15	1.94	18.06	0.36	57.71	1.15

PYDIFLUMETOFEN (309)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.1 mg/kg bw					
Codex Code	Commodity description	Expr as	STM R mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.02	5.84	0.12	10.18	0.20	24.29	0.49	4.52	0.09	14.43	0.29
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.02	1.05	0.02	1.14	0.02	18.69	0.37	0.94	0.02	3.12	0.06
MO 0105	Edible offal (mammalian), raw	RAC	0.09	4.64	0.42	1.97	0.18	10.01	0.90	3.27	0.29	3.98	0.36
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.02	108.75	2.18	70.31	1.41	436.11	8.72	61.55	1.23	79.09	1.58
PM 0110	Poultry meat, raw (incl prepared)	RAC	0.02	3.92	0.08	12.03	0.24	57.07	1.14	5.03	0.10	55.56	1.11
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.02	NC	-	NC	-	0.32	0.01	NC	-	NC	-
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.02	0.10	0.00	0.70	0.01	0.97	0.02	0.10	0.00	NC	-
PE 0112	Eggs, raw, (incl dried)	RAC	0.02	3.84	0.08	4.41	0.09	27.25	0.55	1.13	0.02	7.39	0.15
-	-	-		-	-	-	-	-	-	-	-	-	-
Total intake (ug/person)=				159.5		99.7		226.5		105.2		155.3	
Bodyweight per region (kg bw) =				60		60		60		60		60	
ADI (ug/person)=				6000		6000		6000		6000		6000	
%ADI=				2.7%		1.7%		3.8%		1.8%		2.6%	
Rounded %ADI=				3%		2%		4%		2%		3%	

Annex 3

QUINOXYFEN (222)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.2 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
FS 0013	Subgroup of Cherries, raw	RAC	0.14	0.92	0.13	9.15	1.28	0.01	0.00	0.61	0.09	0.06	0.01	6.64	0.93
FB 0021	Currants, Black, Red, White, raw	RAC	0.2	0.02	0.00	0.74	0.15	0.01	0.00	0.03	0.01	0.01	0.00	0.01	0.00
FB 0269	Grapes, raw (i.e. table grapes)	RAC	0.13	12.68	1.65	9.12	1.19	0.03	0.00	16.88	2.19	3.70	0.48	54.42	7.07
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	0.099	0.51	0.05	0.51	0.05	0.01	0.00	1.27	0.13	0.12	0.01	2.07	0.20
JF 0269	Grape juice (from wine grapes)	PP	0.009	0.14	0.00	0.29	0.00	0.05	0.00	0.30	0.00	0.24	0.00	0.05	0.00
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.015	0.67	0.01	12.53	0.19	2.01	0.03	1.21	0.02	3.53	0.05	4.01	0.06
FB 0275	Strawberry, raw	RAC	0.32	0.70	0.22	2.01	0.64	0.04	0.01	1.36	0.44	0.37	0.12	2.53	0.81
VO 0445	Peppers, sweet, raw	RAC	0.15	1.43	0.21	2.61	0.39	1.05	0.16	2.01	0.30	2.59	0.39	6.24	0.94
-	Peppers, sweet, dried	PP	1.5	0.42	0.63	0.53	0.80	0.84	1.26	0.50	0.75	0.95	1.43	0.37	0.56
VL 0483	Lettuce, leaf, raw	RAC	3.8	0.53	2.01	0.36	1.37	0.16	0.61	6.21	23.60	1.90	7.22	6.05	22.99
VR 0596	Sugar beet, raw (incl sugar)	RAC	0.01	0.13	0.00	NC	-	0.08	0.00	0.66	0.01	0.47	0.00	88.94	0.89
GC 0654	Wheat, raw (incl bulgur, incl fermented beverages, incl germ, incl wholemeal bread, incl white flour products, incl white bread)	RAC	0.01	381.15	3.81	341.55	3.42	38.35	0.38	281.89	2.82	172.83	1.73	434.07	4.34
GC 0640	Barley, raw (incl malt extract, incl pot&pearled, incl flour & grits, incl malt, excl beer)	RAC	0.01	18.98	0.19	13.35	0.13	0.42	0.00	0.67	0.01	2.30	0.02	0.86	0.01
-	Barley beer	PP	0.001	4.87	0.00	93.78	0.09	24.28	0.02	12.76	0.01	39.28	0.04	18.15	0.02
DH 1100	Hops, dry	RAC	0.22	0.01	0.00	0.04	0.01	0.01	0.00	0.01	0.00	NC	-	0.01	0.00
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.002	24.96	0.05	57.95	0.12	16.70	0.03	38.38	0.08	26.46	0.05	29.00	0.06
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.01	6.24	0.06	14.49	0.14	4.18	0.04	9.60	0.10	6.62	0.07	7.25	0.07
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.01	3.29	0.03	6.14	0.06	0.82	0.01	1.57	0.02	2.23	0.02	1.07	0.01
MO 0105	Edible offal (mammalian), raw	RAC	0.01	4.79	0.05	9.68	0.10	2.97	0.03	5.49	0.05	3.84	0.04	5.03	0.05
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.02	289.65	5.79	485.88	9.72	26.92	0.54	239.03	4.78	199.91	4.00	180.53	3.61
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0	13.17	0.00	26.78	0.00	7.24	0.00	116.71	0.00	22.54	0.00	32.09	0.00
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0.013	1.46	0.02	2.98	0.04	0.80	0.01	12.97	0.17	2.50	0.03	3.57	0.05
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.013	0.10	0.00	0.10	0.00	NC	-	0.10	0.00	0.10	0.00	0.10	0.00
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.009	0.12	0.00	0.12	0.00	0.11	0.00	5.37	0.05	0.24	0.00	0.10	0.00

QUINOXYFEN (222)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.2 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
PE 0112	Eggs, raw, (incl dried)	RAC	0.003	7.84	0.02	23.08	0.07	2.88	0.01	14.89	0.04	9.81	0.03	14.83	0.04
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-
Total intake (ug/person)=				15.0		20.0		3.2		35.7		15.7		42.7	
Bodyweight per region (kg bw) =				60		60		60		60		60		60	
ADI (ug/person)=				12000		12000		12000		12000		12000		12000	
%ADI=				0.1%		0.2%		0.0%		0.3%		0.1%		0.4%	
Rounded %ADI=				0%		0%		0%		0%		0%		0%	

QUINOXYFEN (222)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.2 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
FS 0013	Subgroup of Cherries, raw	RAC	0.14	1.40	0.20	4.21	0.59	0.04	0.01	2.93	0.41	1.50	0.21	NC	-
FB 0021	Currants, Black, Red, White, raw	RAC	0.2	0.48	0.10	4.23	0.85	NC	-	1.51	0.30	0.49	0.10	NC	-
FB 0269	Grapes, raw (i.e. table grapes)	RAC	0.13	6.33	0.82	11.22	1.46	5.21	0.68	9.38	1.22	4.55	0.59	0.78	0.10
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	0.099	3.09	0.31	1.51	0.15	0.03	0.00	1.38	0.14	4.26	0.42	0.42	0.04
JF 0269	Grape juice (from wine grapes)	PP	0.009	0.56	0.01	1.96	0.02	0.02	0.00	2.24	0.02	2.27	0.02	0.34	0.00
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.015	88.93	1.33	62.41	0.94	1.84	0.03	25.07	0.38	61.17	0.92	5.84	0.09
FB 0275	Strawberry, raw	RAC	0.32	4.49	1.44	5.66	1.81	0.02	0.01	6.63	2.12	5.75	1.84	0.05	0.02
VO 0445	Peppers, sweet, raw	RAC	0.15	NC	-	NC	-	8.25	1.24	3.03	0.45	NC	-	0.91	0.14
-	Peppers, sweet, dried	PP	1.5	0.11	0.17	0.21	0.32	0.36	0.54	0.21	0.32	0.25	0.38	0.15	0.23
VL 0483	Lettuce, leaf, raw	RAC	3.8	14.50	55.10	11.76	44.69	13.14	49.93	19.50	74.10	4.81	18.28	2.23	8.47
VR 0596	Sugar beet, raw (incl sugar)	RAC	0.01	0.01	0.00	NC	-	0.01	0.00	0.01	0.00	NC	-	NC	-
GC 0654	Wheat, raw (incl bulgur, incl fermented beverages, incl germ, incl wholemeal bread, incl white flour products, incl white bread)	RAC	0.01	253.07	2.53	244.73	2.45	134.44	1.34	235.10	2.35	216.39	2.16	167.40	1.67
GC 0640	Barley, raw (incl malt extract, incl pot&pearled, incl flour & grits, incl malt, excl beer)	RAC	0.01	1.94	0.02	4.15	0.04	0.66	0.01	2.50	0.03	2.14	0.02	3.52	0.04

Annex 3

QUINOXYFEN (222)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.2 mg/kg bw								
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day										
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake	
-	Barley beer	PP	0.001	180.21	0.18	259.46	0.26	45.91	0.05	172.36	0.17	234.42	0.23	65.30	0.07	
DH 1100	Hops, dry	RAC	0.22	NC	-	NC	-	0.02	0.00	0.02	0.00	NC	-	NC	-	
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.002	112.02	0.22	120.71	0.24	63.46	0.13	88.99	0.18	96.24	0.19	41.02	0.08	
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.01	28.01	0.28	30.18	0.30	15.86	0.16	22.25	0.22	24.06	0.24	10.25	0.10	
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.01	6.44	0.06	15.51	0.16	3.79	0.04	8.29	0.08	18.44	0.18	8.00	0.08	
MO 0105	Edible offal (mammalian), raw	RAC	0.01	15.17	0.15	5.19	0.05	6.30	0.06	6.78	0.07	3.32	0.03	3.17	0.03	
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.02	388.92	7.78	335.88	6.72	49.15	0.98	331.25	6.63	468.56	9.37	245.45	4.91	
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0	66.38	0.00	48.47	0.00	21.58	0.00	78.41	0.00	48.04	0.00	76.01	0.00	
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0.013	7.38	0.10	5.39	0.07	2.40	0.03	8.71	0.11	5.34	0.07	8.45	0.11	
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.013	0.10	0.00	0.10	0.00	NC	-	0.10	0.00	0.71	0.01	NC	-	
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.009	0.33	0.00	0.72	0.01	0.27	0.00	0.35	0.00	0.80	0.01	NC	-	
PE 0112	Eggs, raw, (incl dried)	RAC	0.003	25.84	0.08	29.53	0.09	28.05	0.08	33.19	0.10	36.44	0.11	8.89	0.03	
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-	
Total intake (ug/person)=					70.9		61.2		55.3		89.4		35.4		16.2	
Bodyweight per region (kg bw) =					60		60		55		60		60		60	
ADI (ug/person)=					12000		12000		11000		12000		12000		12000	
%ADI=					0.6%		0.5%		0.5%		0.7%		0.3%		0.1%	
Rounded %ADI=					1%		1%		1%		1%		0%		0%	

QUINOXYFEN (222)													
International Estimated Daily Intake (IEDI)				ADI = 0 - 0.2 mg/kg bw									
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
FS 0013	Subgroup of Cherries, raw	RAC	0.14	0.01	0.00	0.01	0.00	5.96	0.83	0.01	0.00	NC	-
FB 0021	Currants, Black, Red, White, raw	RAC	0.2	0.01	0.00	NC	-	0.74	0.15	NC	-	NC	-
FB 0269	Grapes, raw (i.e. table grapes)	RAC	0.13	0.14	0.02	0.36	0.05	15.22	1.98	0.01	0.00	0.09	0.01
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	0.099	0.01	0.00	0.13	0.01	1.06	0.10	0.01	0.00	0.03	0.00
JF 0269	Grape juice (from wine grapes)	PP	0.009	0.01	0.00	0.01	0.00	0.41	0.00	0.01	0.00	NC	-
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.015	0.31	0.00	0.23	0.00	60.43	0.91	0.52	0.01	31.91	0.48
FB 0275	Strawberry, raw	RAC	0.32	0.01	0.00	0.01	0.00	3.35	1.07	0.01	0.00	0.01	0.00
VO 0445	Peppers, sweet, raw	RAC	0.15	1.24	0.19	1.27	0.19	NC	-	0.01	0.00	NC	-
-	Peppers, sweet, dried	PP	1.5	0.58	0.87	1.27	1.91	1.21	1.82	0.12	0.18	NC	-
VL 0483	Lettuce, leaf, raw	RAC	3.8	0.29	1.10	0.03	0.11	6.71	25.50	0.01	0.04	NC	-
VR 0596	Sugar beet, raw (incl sugar)	RAC	0.01	3.93	0.04	1.68	0.02	NC	-	NC	-	36.12	0.36
GC 0654	Wheat, raw (incl bulgur, incl fermented beverages, incl germ, incl wholemeal bread, incl white flour products, incl white bread)	RAC	0.01	57.20	0.57	110.47	1.10	272.62	2.73	25.82	0.26	132.04	1.32
GC 0640	Barley, raw (incl malt extract, incl pot&pearled, incl flour & grits, incl malt, excl beer)	RAC	0.01	8.50	0.09	0.17	0.00	3.92	0.04	0.02	0.00	6.34	0.06
-	Barley beer	PP	0.001	16.25	0.02	11.36	0.01	225.21	0.23	19.49	0.02	52.17	0.05
DH 1100	Hops, dry	RAC	0.22	NC	-	NC	-	0.04	0.01	NC	-	NC	-
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.002	23.34	0.05	40.71	0.08	97.15	0.19	18.06	0.04	57.71	0.12
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.01	5.84	0.06	10.18	0.10	24.29	0.24	4.52	0.05	14.43	0.14
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.01	1.05	0.01	1.14	0.01	18.69	0.19	0.94	0.01	3.12	0.03
MO 0105	Edible offal (mammalian), raw	RAC	0.01	4.64	0.05	1.97	0.02	10.01	0.10	3.27	0.03	3.98	0.04
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.02	108.75	2.18	70.31	1.41	436.11	8.72	61.55	1.23	79.09	1.58
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0	3.53	0.00	10.83	0.00	51.36	0.00	4.53	0.00	50.00	0.00
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0.013	0.39	0.01	1.20	0.02	5.71	0.07	0.50	0.01	5.56	0.07
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.013	NC	-	NC	-	0.32	0.00	NC	-	NC	-
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.009	0.10	0.00	0.70	0.01	0.97	0.01	0.10	0.00	NC	-
PE 0112	Eggs, raw, (incl dried)	RAC	0.003	3.84	0.01	4.41	0.01	27.25	0.08	1.13	0.00	7.39	0.02

Annex 3

QUINOXYFEN (222)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.2 mg/kg bw						
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person								
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake	
-	-	-		-	-	-	-	-	-	-	-	-	-	
Total intake (ug/person)=					5.3		5.1		45.0		1.9		4.3	
Bodyweight per region (kg bw) =					60		60		60		60		60	
ADI (ug/person)=					12000		12000		12000		12000		12000	
%ADI=					0.0%		0.0%		0.4%		0.0%		0.0%	
Rounded %ADI=					0%		0%		0%		0%		0%	

METHOPRENE (147)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.05 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
VD 0541	Soya bean, dry, raw (incl paste, incl curd, incl oil, incl sauce)	RAC	2.4	72.79	174.70	59.05	141.72	20.55	49.32	74.20	178.08	61.12	146.69	73.24	175.78
GC 0648	Quinoa, raw	RAC	4.85	NC	-	NC	-	NC	-	NC	-	0.07	0.34	NC	-
GC 0650	Rye, raw (incl flour)	RAC	4.85	0.13	0.63	19.38	93.99	0.10	0.49	0.12	0.58	0.03	0.15	2.15	10.43
GC 0653	Triticale, raw (incl flour)	RAC	4.85	NC	-	NC	-	NC	-	0.01	0.05	0.39	1.89	NC	-
GC 0654	Wheat, raw (incl meslin)	RAC	4.85	0.01	0.05	1.12	5.43	NC	-	0.01	0.05	0.56	2.72	NC	-
-	Wheat, bulgur	PP	4.85	NC	-	NC	-	NC	-	0.03	0.15	NC	-	NC	-
CF 1210	Wheat, germ	PP	23.3	NC	-	NC	-	0.01	0.23	0.01	0.23	0.14	3.26	0.01	0.23
CP 1212	Wheat, wholemeal bread	PP	4.51	0.01	0.05	0.01	0.05	0.01	0.05	0.01	0.05	0.03	0.14	0.01	0.05
CP 1211	Wheat, white bread	PP	1.72	0.25	0.43	0.63	1.08	0.12	0.21	0.43	0.74	1.39	2.39	0.22	0.38
-	Wheat, Fermented Beverages (Korean jakju and takju)	PP	4.85	NC	-	NC	-	NC	-	NC	-	NC	-	NC	-
CF 1211	Wheat, white flour (incl white flour products: starch, gluten, macaroni, pastry)	PP	1.72	301.24	518.13	268.64	462.06	30.21	51.96	222.51	382.72	134.73	231.74	343.12	590.17
GC 0640	Barley, raw (incl malt extract, incl pot&pearled, incl flour & grits, incl beer, incl malt)	RAC	4.85	19.91	96.56	31.16	151.13	5.04	24.44	3.10	15.04	9.77	47.38	4.31	20.90
GC 0641	Buckwheat, raw (incl flour)	RAC	4.85	NC	-	0.40	1.94	0.01	0.05	0.01	0.05	0.07	0.34	0.09	0.44
GC 0647	Oats, raw (incl rolled)	RAC	4.85	0.05	0.24	7.05	34.19	0.10	0.49	1.71	8.29	0.96	4.66	0.04	0.19
CM 0649 (GC 0649)	Rice, husked, dry (incl flour, incl oil, incl beverages, incl starch, excl polished)	REP	1.07	1.26	1.35	1.58	1.69	31.05	33.22	5.43	5.81	0.90	0.96	2.18	2.33
CM 1205	Rice polished, dry	PP	0.1	34.21	3.42	10.39	1.04	41.72	4.17	82.38	8.24	150.24	15.02	70.47	7.05
GC 0646	Millet, raw (incl flour, incl beer)	RAC	4.85	1.46	7.08	2.32	11.25	5.84	28.32	0.89	4.32	16.17	78.42	0.01	0.05
GC 0651	Sorghum, raw (incl flour, incl beer) (i.e. Chicken corn, Dari seed, Durra, Feterita)	RAC	4.85	4.34	21.05	0.01	0.05	16.25	78.81	15.82	76.73	10.97	53.20	2.92	14.16
GC 0645	Maize, raw (incl glucose & dextrose & isoglucose, incl flour, incl beer, incl germ, incl starch, excl oil)	RAC	4.85	28.85	139.92	43.93	213.06	108.66	527.00	46.94	227.66	59.87	290.37	73.58	356.86
OR 0645	Maize oil	PP	0	0.96	0.00	0.85	0.00	0.29	0.00	5.42	0.00	0.42	0.00	2.10	0.00
-	Cereals, NES, raw (including processed) : canagua, quihuicha, Job's tears and wild rice	RAC	4.85	2.04	9.89	2.99	14.50	1.86	9.02	19.17	92.97	3.33	16.15	1.66	8.05
SO 2090	Subgroup of Small seed oilseeds, raw (incl processed)	RAC	2	2.19	4.38	1.22	2.44	1.05	2.10	7.10	14.20	10.29	20.58	4.83	9.66
SO 2091	Subgroup of Sunflower seeds, raw (incl processed)	RAC	2	7.43	14.86	36.06	72.12	1.15	2.30	8.77	17.54	5.74	11.48	13.63	27.26

Annex 3

METHOPRENE (147)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.05 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
SO 0691	Cotton seed, raw (incl oil)	RAC	2	20.53	41.06	9.80	19.60	6.42	12.84	4.73	9.46	7.14	14.28	18.68	37.36
-	Castor bean, raw (incl oil)	RAC	2	NC	-	0.07	0.14	NC	-	NC	-	NC	-	0.01	0.02
SO 3155	Melonseed	RAC	2	0.06	0.12	NC	-	1.08	2.16	0.38	0.76	0.01	0.02	0.25	0.50
SO 0696	Palm kernels, raw (incl oil)	RAC	2	5.81	11.62	3.77	7.54	20.07	40.14	24.53	49.06	5.94	11.88	8.99	17.98
SO 0697	Peanuts, nutmeat, raw (incl roasted, incl oil, incl butter)	RAC	5	1.30	6.50	1.23	6.15	12.62	63.10	2.87	14.35	6.59	32.95	2.67	13.35
SO 0701	Shea nut (karite nuts), nutmeat, raw (incl butter)	RAC	2	NC	-	NC	-	0.34	0.68	NC	-	NC	-	NC	-
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.007	24.96	0.17	57.95	0.41	16.70	0.12	38.38	0.27	26.46	0.19	29.00	0.20
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.092	6.24	0.57	14.49	1.33	4.18	0.38	9.60	0.88	6.62	0.61	7.25	0.67
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.092	3.29	0.30	6.14	0.56	0.82	0.08	1.57	0.14	2.23	0.21	1.07	0.10
MO 0105	Edible offal (mammalian), raw	RAC	0.014	4.79	0.07	9.68	0.14	2.97	0.04	5.49	0.08	3.84	0.05	5.03	0.07
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.044	289.65	12.74	485.88	21.38	26.92	1.18	239.03	10.52	199.91	8.80	180.53	7.94
PM 0110	Poultry meat, raw (incl prepared)	RAC	0.007	14.63	0.10	29.76	0.21	8.04	0.06	129.68	0.91	25.04	0.18	35.66	0.25
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.007	0.12	0.00	0.12	0.00	0.11	0.00	5.37	0.04	0.24	0.00	0.10	0.00
PE 0112	Eggs, raw, (incl dried)	RAC	0.006	7.84	0.05	23.08	0.14	2.88	0.02	14.89	0.09	9.81	0.06	14.83	0.09
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-
Total intake (ug/person)=				1066.1		1265.3		933.0		1120.0		997.1		1302.5	
Bodyweight per region (kg bw) =				60		60		60		60		60		60	
ADI (ug/person)=				3000		3000		3000		3000		3000		3000	
%ADI=				35.5%		42.2%		31.1%		37.3%		33.2%		43.4%	
Rounded %ADI=				40%		40%		30%		40%		30%		40%	

METHOPRENE (147)															
International Estimated Daily Intake (IEDI)				ADI = 0 - 0.05 mg/kg bw											
Codex Code	Commodity description	Expr as	STM ^R mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
VD 0541	Soya bean, dry, raw (incl paste, incl curd, incl oil, incl sauce)	RAC	2.4	106.33	255.19	117.78	282.67	42.12	101.09	195.70	469.68	222.52	534.05	80.47	193.13
GC 0648	Quinoa, raw	RAC	4.85	NC	-	NC	-	NC	-	NC	-	NC	-	NC	-
GC 0650	Rye, raw (incl flour)	RAC	4.85	3.21	15.57	35.38	171.59	0.21	1.02	6.50	31.53	1.49	7.23	NC	-
GC 0653	Triticale, raw (incl flour)	RAC	4.85	0.01	0.05	0.17	0.82	0.29	1.41	0.01	0.05	NC	-	NC	-
GC 0654	Wheat, raw (incl meslin)	RAC	4.85	NC	-	NC	-	NC	-	0.01	0.05	NC	-	NC	-
-	Wheat, bulgur	PP	4.85	NC	-	NC	-	0.01	0.05	NC	-	NC	-	NC	-
CF 1210	Wheat, germ	PP	23.3	0.97	22.60	0.10	2.33	0.03	0.70	0.01	0.23	NC	-	0.04	0.93
CP 1212	Wheat, wholemeal bread	PP	4.51	0.03	0.14	0.01	0.05	0.01	0.05	0.01	0.05	0.05	0.23	0.02	0.09
CP 1211	Wheat, white bread	PP	1.72	1.30	2.24	0.46	0.79	0.06	0.10	0.22	0.38	2.44	4.20	0.77	1.32
-	Wheat, Fermented Beverages (Korean jakju and takju)	PP	4.85	NC	-	NC	-	NC	-	4.36	21.15	NC	-	NC	-
CF 1211	Wheat, white flour (incl white flour products: starch, gluten, macaroni, pastry)	PP	1.72	198.08	340.70	193.03	332.01	106.24	182.73	185.09	318.35	168.67	290.11	131.59	226.33
GC 0640	Barley, raw (incl malt extract, incl pot&pearled, incl flour & grits, incl beer, incl malt)	RAC	4.85	36.18	175.47	53.45	259.23	9.39	45.54	35.25	170.96	46.68	226.40	15.92	77.21
GC 0641	Buckwheat, raw (incl flour)	RAC	4.85	0.01	0.05	0.79	3.83	0.18	0.87	0.35	1.70	NC	-	NC	-
GC 0647	Oats, raw (incl rolled)	RAC	4.85	7.50	36.38	6.26	30.36	0.15	0.73	4.87	23.62	3.16	15.33	2.98	14.45
CM 0649 (GC 0649)	Rice, husked, dry (incl flour, incl oil, incl beverages, incl starch, excl polished)	REP	1.07	3.70	3.96	2.11	2.26	1.51	1.62	1.75	1.87	0.29	0.31	5.12	5.48
CM 1205	Rice polished, dry	PP	0.1	13.38	1.34	10.80	1.08	262.08	26.21	57.16	5.72	12.83	1.28	62.78	6.28
GC 0646	Millet, raw (incl flour, incl beer)	RAC	4.85	0.03	0.15	0.16	0.78	1.75	8.49	0.69	3.35	NC	-	NC	-
GC 0651	Sorghum, raw (incl flour, incl beer) (i.e. Chicken corn, Dari seed, Durra, Feterita)	RAC	4.85	NC	-	NC	-	1.44	6.98	1.15	5.58	NC	-	7.12	34.53
GC 0645	Maize, raw (incl glucose & dextrose & isoglucose, incl flour, incl beer, incl germ, incl starch, excl oil)	RAC	4.85	17.61	85.41	25.71	124.69	25.89	125.57	36.98	179.35	5.49	26.63	64.23	311.52
OR 0645	Maize oil	PP	0	0.90	0.00	0.47	0.00	0.15	0.00	3.01	0.00	1.86	0.00	0.36	0.00
-	Cereals, NES, raw (including processed) : canagua, quihuicha, Job's tears and wild rice	RAC	4.85	6.17	29.92	3.01	14.60	0.76	3.69	3.30	16.01	3.38	16.39	15.84	76.82
SO 2090	Subgroup of Small seed oilseeds, raw (incl processed)	RAC	2	33.60	67.20	20.72	41.44	9.72	19.44	17.21	34.42	1.11	2.22	0.54	1.08

Annex 3

METHOPRENE (147)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.05 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
SO 2091	Subgroup of Sunflower seeds, raw (incl processed)	RAC	2	23.43	46.86	29.34	58.68	1.24	2.48	14.00	28.00	6.48	12.96	6.91	13.82
SO 0691	Cotton seed, raw (incl oil)	RAC	2	10.71	21.42	4.23	8.46	7.19	14.38	7.54	15.08	5.66	11.32	2.38	4.76
-	Castor bean, raw (incl oil)	RAC	2	NC	-	NC	-	0.01	0.02	NC	-	NC	-	NC	-
SO 3155	Melonseed	RAC	2	NC	-	NC	-	0.01	0.02	NC	-	NC	-	NC	-
SO 0696	Palm kernels, raw (incl oil)	RAC	2	5.33	10.66	5.04	10.08	11.83	23.66	7.94	15.88	10.77	21.54	4.53	9.06
SO 0697	Peanuts, nutmeat, raw (incl roasted, incl oil, incl butter)	RAC	5	5.63	28.15	2.75	13.75	9.58	47.90	5.82	29.10	13.71	68.55	1.84	9.20
SO 0701	Shea nut (karite nuts), nutmeat, raw (incl butter)	RAC	2	NC	-	NC	-	NC	-	NC	-	NC	-	NC	-
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.007	112.02	0.78	120.71	0.84	63.46	0.44	88.99	0.62	96.24	0.67	41.02	0.29
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.092	28.01	2.58	30.18	2.78	15.86	1.46	22.25	2.05	24.06	2.21	10.25	0.94
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.092	6.44	0.59	15.51	1.43	3.79	0.35	8.29	0.76	18.44	1.70	8.00	0.74
MO 0105	Edible offal (mammalian), raw	RAC	0.014	15.17	0.21	5.19	0.07	6.30	0.09	6.78	0.09	3.32	0.05	3.17	0.04
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.044	388.92	17.11	335.88	14.78	49.15	2.16	331.25	14.58	468.56	20.62	245.45	10.80
PM 0110	Poultry meat, raw (incl prepared)	RAC	0.007	73.76	0.52	53.86	0.38	23.98	0.17	87.12	0.61	53.38	0.37	84.45	0.59
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.007	0.33	0.00	0.72	0.01	0.27	0.00	0.35	0.00	0.80	0.01	NC	-
PE 0112	Eggs, raw, (incl dried)	RAC	0.006	25.84	0.16	29.53	0.18	28.05	0.17	33.19	0.20	36.44	0.22	8.89	0.05
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-
Total intake (ug/person)=				1165.4		1380.0		619.6		1391.0		1264.6		999.5	
Bodyweight per region (kg bw) =				60		60		55		60		60		60	
ADI (ug/person)=				3000		3000		2750		3000		3000		3000	
%ADI=				38.8%		46.0%		22.5%		46.4%		42.2%		33.3%	
Rounded %ADI=				40%		50%		20%		50%		40%		30%	

METHOPRENE (147)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.05 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
VD 0541	Soya bean, dry, raw (incl paste, incl curd, incl oil, incl sauce)	RAC	2.4	15.80	37.92	14.29	34.30	104.36	250.46	17.11	41.06	35.20	84.48
GC 0648	Quinoa, raw	RAC	4.85	NC	-	NC	-	NC	-	NC	-	NC	-
GC 0650	Rye, raw (incl flour)	RAC	4.85	0.03	0.15	0.01	0.05	13.95	67.66	0.01	0.05	0.88	4.27
GC 0653	Triticale, raw (incl flour)	RAC	4.85	0.01	0.05	NC	-	NC	-	NC	-	NC	-
GC 0654	Wheat, raw (incl meslin)	RAC	4.85	NC	-	NC	-	NC	-	NC	-	0.97	4.70
-	Wheat, bulgur	PP	4.85	0.01	0.05	NC	-	NC	-	NC	-	NC	-
CF 1210	Wheat, germ	PP	23.3	0.04	0.93	0.01	0.23	0.01	0.23	0.01	0.23	NC	-
CP 1212	Wheat, wholemeal bread	PP	4.51	0.01	0.05	0.01	0.05	0.03	0.14	0.01	0.05	0.01	0.05
CP 1211	Wheat, white bread	PP	1.72	0.43	0.74	0.41	0.71	1.56	2.68	0.11	0.19	0.07	0.12
-	Wheat, Fermented Beverages (Korean jakju and takju)	PP	4.85	NC	-	NC	-	NC	-	NC	-	NC	-
CF 1211	Wheat, white flour (incl white flour products: starch, gluten, macaroni, pastry)	PP	1.72	44.78	77.02	86.96	149.57	214.05	368.17	20.31	34.93	103.60	178.19
GC 0640	Barley, raw (incl malt extract, incl pot&pearled, incl flour & grits, incl beer, incl malt)	RAC	4.85	11.58	56.16	2.33	11.30	46.71	226.54	3.72	18.04	16.26	78.86
GC 0641	Buckwheat, raw (incl flour)	RAC	4.85	0.04	0.19	2.82	13.68	0.01	0.05	0.01	0.05	NC	-
GC 0647	Oats, raw (incl rolled)	RAC	4.85	0.37	1.79	0.07	0.34	2.79	13.53	0.10	0.49	NC	-
CM 0649 (GC 0649)	Rice, husked, dry (incl flour, incl oil, incl beverages, incl starch, excl polished)	REP	1.07	13.58	14.53	4.29	4.59	2.17	2.32	0.01	0.01	8.84	9.46
CM 1205	Rice polished, dry	PP	0.1	30.20	3.02	218.34	21.83	12.77	1.28	15.24	1.52	51.35	5.14
GC 0646	Millet, raw (incl flour, incl beer)	RAC	4.85	61.13	296.48	0.78	3.78	NC	-	33.55	162.72	NC	-
GC 0651	Sorghum, raw (incl flour, incl beer) (i.e. Chicken corn, Dari seed, Durra, Feterita)	RAC	4.85	89.16	432.43	2.02	9.80	NC	-	35.38	171.59	NC	-
GC 0645	Maize, raw (incl glucose & dextrose & isoglucose, incl flour, incl beer, incl germ, incl starch, excl oil)	RAC	4.85	116.33	564.20	10.45	50.68	37.65	182.60	76.60	371.51	34.44	167.03
OR 0645	Maize oil	PP	0	0.33	0.00	0.07	0.00	0.81	0.00	0.01	0.00	NC	-
-	Cereals, NES, raw (including processed) : canagua, quihuicha, Job's tears and wild rice	RAC	4.85	17.71	85.89	2.00	9.70	9.61	46.61	0.45	2.18	4.55	22.07
SO 2090	Subgroup of Small seed oilseeds, raw (incl processed)	RAC	2	2.63	5.26	0.93	1.86	12.79	25.58	9.98	19.96	0.01	0.02
SO 2091	Subgroup of Sunflower seeds, raw (incl processed)	RAC	2	0.99	1.98	0.22	0.44	32.01	64.02	12.12	24.24	0.48	0.96

Annex 3

METHOPRENE (147)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.05 mg/kg bw						
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person								
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake	
SO 0691	Cotton seed, raw (incl oil)	RAC	2	8.14	16.28	0.32	0.64	2.84	5.68	2.69	5.38	0.97	1.94	
-	Castor bean, raw (incl oil)	RAC	2	NC	-	NC	-	NC	-	NC	-	NC	-	
SO 3155	Melonseed	RAC	2	1.81	3.62	NC	-	0.01	0.02	NC	-	NC	-	
SO 0696	Palm kernels, raw (incl oil)	RAC	2	60.84	121.68	12.77	25.54	5.41	10.82	0.57	1.14	53.45	106.90	
SO 0697	Peanuts, nutmeat, raw (incl roasted, incl oil, incl butter)	RAC	5	18.82	94.10	0.57	2.85	2.28	11.40	6.90	34.50	0.53	2.65	
SO 0701	Shea nut (karite nuts), nutmeat, raw (incl butter)	RAC	2	0.95	1.90	NC	-	NC	-	NC	-	NC	-	
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.007	23.34	0.16	40.71	0.28	97.15	0.68	18.06	0.13	57.71	0.40	
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.092	5.84	0.54	10.18	0.94	24.29	2.23	4.52	0.42	14.43	1.33	
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.092	1.05	0.10	1.14	0.10	18.69	1.72	0.94	0.09	3.12	0.29	
MO 0105	Edible offal (mammalian), raw	RAC	0.014	4.64	0.06	1.97	0.03	10.01	0.14	3.27	0.05	3.98	0.06	
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.044	108.75	4.79	70.31	3.09	436.11	19.19	61.55	2.71	79.09	3.48	
PM 0110	Poultry meat, raw (incl prepared)	RAC	0.007	3.92	0.03	12.03	0.08	57.07	0.40	5.03	0.04	55.56	0.39	
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.007	0.10	0.00	0.70	0.00	0.97	0.01	0.10	0.00	NC	-	
PE 0112	Eggs, raw, (incl dried)	RAC	0.006	3.84	0.02	4.41	0.03	27.25	0.16	1.13	0.01	7.39	0.04	
-	-	-		-	-	-	-	-	-	-	-	-	-	
Total intake (ug/person)=					1822.1		346.5		1304.3		893.3		672.8	
Bodyweight per region (kg bw) =					60		60		60		60		60	
ADI (ug/person)=					3000		3000		3000		3000		3000	
%ADI=					60.7%		11.5%		43.5%		29.8%		22.4%	
Rounded %ADI=					60%		10%		40%		30%		20%	

SPINETORAM (233)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.05 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
FC 0003	Subgroup of Mandarins, raw	RAC	0.0605	6.18	0.37	3.66	0.22	0.25	0.02	6.82	0.41	3.49	0.21	19.38	1.17
FC 0004	Subgroup of Oranges, sweet, sour, raw	RAC	0.0435	20.66	0.90	5.23	0.23	11.90	0.52	37.90	1.65	21.16	0.92	56.46	2.46
JF 0004	Subgroup of Oranges, juice (single strength, incl. concentrated)	PP	0.003	1.27	0.00	2.20	0.01	0.09	0.00	11.81	0.04	0.46	0.00	1.69	0.01
FP 0009	Group of Pome fruits, raw (incl. apple juice, incl apple cider)	RAC	0.025	19.79	0.49	38.25	0.96	17.96	0.45	32.56	0.81	8.08	0.20	64.45	1.61
JF 0226	Apple juice, single strength (incl. concentrated)	PP	0.011	0.32	0.00	3.07	0.03	0.07	0.00	5.00	0.06	0.29	0.00	5.57	0.06
FS 0013	Subgroup of Cherries, raw	RAC	0.0205	0.92	0.02	9.15	0.19	0.01	0.00	0.61	0.01	0.06	0.00	6.64	0.14
FS 0014	Subgroup of Plums, raw (incl dried plums)	RAC	0.02	2.67	0.05	8.77	0.18	0.07	0.00	3.03	0.06	0.70	0.01	4.34	0.09
FS 0240	Apricot, raw (incl dried)	RAC	0.0485	5.15	0.25	3.66	0.18	0.03	0.00	2.25	0.11	0.17	0.01	6.80	0.33
-	Peaches and nectarines, raw	RAC	0.055	2.87	0.16	2.21	0.12	0.15	0.01	5.94	0.33	1.47	0.08	15.66	0.86
FB 0272	Raspberries, red, black, raw	RAC	0.42	0.07	0.03	0.93	0.39	0.01	0.00	0.01	0.00	0.01	0.00	0.01	0.00
FB 0020	Blueberries, raw	RAC	0.12	0.01	0.00	0.03	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.01	0.00
FB 0021	Currants, Black, Red, White, raw	RAC	0.18	0.02	0.00	0.74	0.13	0.01	0.00	0.03	0.01	0.01	0.00	0.01	0.00
FB 0269	Grapes, raw (incl must, incl dried, incl juice, incl wine)	RAC	0.074	16.25	1.20	28.96	2.14	2.87	0.21	24.22	1.79	9.33	0.69	68.64	5.08
FB 0275	Strawberry, raw	RAC	0.026	0.70	0.02	2.01	0.05	0.04	0.00	1.36	0.04	0.37	0.01	2.53	0.07
FT 0305	Table olives, raw (incl preserved)	RAC	0.02	0.70	0.01	0.32	0.01	0.01	0.00	1.53	0.03	0.17	0.00	1.85	0.04
FI 0343	Litchi, raw (incl processed)	RAC	0.02	2.32	0.05	1.43	0.03	1.81	0.04	7.42	0.15	NC	-	4.54	0.09
FI 0326	Avocado, raw	RAC	0.02	0.13	0.00	0.03	0.00	2.05	0.04	2.54	0.05	2.34	0.05	0.12	0.00
FI 0345	Mango, raw (incl canned mango, incl mango juice)	RAC	0.02	10.48	0.21	0.01	0.00	7.24	0.14	6.87	0.14	19.98	0.40	6.25	0.13
FI 2540	Pitaya, raw (i.e dragon fruit or pitahaya)	RAC	0.0915	0.01	0.00	0.01	0.00	0.01	0.00	0.01	0.00	NC	-	0.01	0.00
FI 0351	Passion fruit, raw	RAC	0.12	0.58	0.07	0.01	0.00	0.59	0.07	0.60	0.07	0.18	0.02	0.08	0.01
VA 2031	Subgroup of bulb onions	RAC	0.01	31.65	0.32	43.28	0.43	3.68	0.04	38.48	0.38	20.46	0.20	47.29	0.47
VA 0384	Leek, raw	RAC	0.026	0.18	0.00	1.59	0.04	0.03	0.00	0.28	0.01	0.01	0.00	3.21	0.08
-	Onions, green, raw	RAC	0.33	2.45	0.81	1.49	0.49	1.02	0.34	2.60	0.86	0.60	0.20	2.03	0.67
VB 0040	Group of Brassica vegetables (excl Brassica leafy vegetables), raw	RAC	0.05	6.43	0.32	40.26	2.01	0.80	0.04	9.94	0.50	12.07	0.60	17.73	0.89
VC 2039	Subgroup of Cucumbers and Squashes, raw	RAC	0.02	10.52	0.21	39.36	0.79	2.07	0.04	25.74	0.51	2.80	0.06	44.83	0.90

Annex 3

SPINETORAM (233)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.05 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
VC 0046	Melons, except watermelon, raw (Cantaloupe)	RAC	0.02	8.90	0.18	8.64	0.17	0.80	0.02	17.90	0.36	2.80	0.06	29.17	0.58
VO 0448	Tomato, raw (incl juice, incl paste, incl canned)	RAC	0.02	51.75	1.04	81.80	1.64	16.99	0.34	102.02	2.04	26.32	0.53	214.77	4.30
VO 0051	Subgroup of peppers, raw (incl dried sweet peppers, excl dried chilipeppers), incl okra	RAC	0.026	10.45	0.27	13.74	0.36	13.81	0.36	14.53	0.38	15.25	0.40	27.93	0.73
VL 0483	Lettuce, leaf, raw	RAC	0.895	0.53	0.47	0.36	0.32	0.16	0.14	6.21	5.56	1.90	1.70	6.05	5.41
VL 0502	Spinach, raw	RAC	1.6	0.74	1.18	0.22	0.35	0.02	0.03	0.91	1.46	0.04	0.06	2.92	4.67
VP 0061	Beans with pods (Phaseolus spp): (immature pods + succulent seeds)	RAC	0.024	0.68	0.02	NC	-	NC	-	0.39	0.01	0.22	0.01	0.49	0.01
VD 0541	Soya bean, dry, raw (incl paste, incl curd, incl oil, incl sauce)	RAC	0.02	72.79	1.46	59.05	1.18	20.55	0.41	74.20	1.48	61.12	1.22	73.24	1.46
VR 0596	Sugar beet, raw (incl sugar)	RAC	0.02	0.13	0.00	NC	-	0.08	0.00	0.66	0.01	0.47	0.01	88.94	1.78
VR 0589	Potato, raw (incl flour, incl frozen, incl starch, incl tapioca)	RAC	0.02	59.74	1.19	316.14	6.32	9.78	0.20	60.26	1.21	54.12	1.08	119.82	2.40
VS 0624	Celery	RAC	0.3	2.14	0.64	3.79	1.14	2.35	0.71	5.69	1.71	0.02	0.01	2.75	0.83
CM 0649 (GC 0649)	Rice, husked, dry (incl polished, incl flour, incl starch, incl oil, incl beverages)	REP	0.04	45.40	1.82	14.99	0.60	84.88	3.40	111.73	4.47	194.75	7.79	93.12	3.72
GC 0645	Maize, raw (incl glucose & dextrose & isoglucose, incl flour, incl oil, incl beer, incl germ, incl starch)	RAC	0.02	29.81	0.60	44.77	0.90	108.95	2.18	52.37	1.05	60.28	1.21	75.69	1.51
GC 0447	Sweet corn on the cob, raw (incl frozen kernels, incl canned kernels)	RAC	0.02	0.14	0.00	0.94	0.02	5.70	0.11	2.61	0.05	1.94	0.04	0.22	0.00
TN 0085	Group of Tree nuts, raw (incl processed)	RAC	0.02	4.06	0.08	3.27	0.07	7.01	0.14	13.93	0.28	14.01	0.28	9.36	0.19
SO 0691	Cotton seed, raw (incl oil)	RAC	0	20.53	0.00	9.80	0.00	6.42	0.00	4.73	0.00	7.14	0.00	18.68	0.00
HS 0444	Peppers, chili, dried	PP	0.26	0.42	0.11	0.53	0.14	0.84	0.22	0.50	0.13	0.95	0.25	0.37	0.10
DT 1114	Tea, green or black, fermented and dried, (including concentrates)	RAC	118	2.28	269.04	1.98	233.64	0.46	54.28	2.43	286.74	1.29	152.22	3.04	358.72
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.005	24.96	0.12	57.95	0.29	16.70	0.08	38.38	0.19	26.46	0.13	29.00	0.15
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.025	6.24	0.16	14.49	0.36	4.18	0.10	9.60	0.24	6.62	0.17	7.25	0.18
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.025	3.29	0.08	6.14	0.15	0.82	0.02	1.57	0.04	2.23	0.06	1.07	0.03
MO 0105	Edible offal (mammalian), raw	RAC	0.005	4.79	0.02	9.68	0.05	2.97	0.01	5.49	0.03	3.84	0.02	5.03	0.03
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.003	289.65	0.87	485.88	1.46	26.92	0.08	239.03	0.72	199.91	0.60	180.53	0.54
PM 0110	Poultry meat, raw (incl prepared) - 90% as	RAC	0.0002	13.17	0.00	26.78	0.01	7.24	0.00	116.71	0.02	22.54	0.00	32.09	0.01

SPINETORAM (233)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.05 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
	muscle														
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0.004	1.46	0.01	2.98	0.01	0.80	0.00	12.97	0.05	2.50	0.01	3.57	0.01
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.004	0.10	0.00	0.10	0.00	NC	-	0.10	0.00	0.10	0.00	0.10	0.00
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.0004	0.12	0.00	0.12	0.00	0.11	0.00	5.37	0.00	0.24	0.00	0.10	0.00
PE 0112	Eggs, raw, (incl dried)	RAC	0.0004	7.84	0.00	23.08	0.01	2.88	0.00	14.89	0.01	9.81	0.00	14.83	0.01
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-
Total intake (ug/person)=				284.9		257.8		64.8		316.2		171.5		402.5	
Bodyweight per region (kg bw) =				60		60		60		60		60		60	
ADI (ug/person)=				3000		3000		3000		3000		3000		3000	
%ADI=				9.5%		8.6%		2.2%		10.5%		5.7%		13.4%	
Rounded %ADI=				9%		9%		2%		10%		6%		10%	

SPINETORAM (233)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.05 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
FC 0003	Subgroup of Mandarins, raw	RAC	0.0605	12.34	0.75	14.99	0.91	16.08	0.97	10.76	0.65	9.94	0.60	NC	-
FC 0004	Subgroup of Oranges, sweet, sour, raw	RAC	0.0435	15.68	0.68	24.00	1.04	6.80	0.30	29.09	1.27	15.39	0.67	160.47	6.98
JF 0004	Subgroup of Oranges, juice (single strength, incl. concentrated)	PP	0.003	33.31	0.10	1.78	0.01	0.28	0.00	18.97	0.06	14.01	0.04	13.36	0.04
FP 0009	Group of Pome fruits, raw (incl. apple juice, incl apple cider)	RAC	0.025	71.38	1.78	81.73	2.04	42.91	1.07	58.89	1.47	103.85	2.60	12.48	0.31
JF 0226	Apple juice, single strength (incl. concentrated)	PP	0.011	14.88	0.16	11.98	0.13	0.15	0.00	9.98	0.11	30.32	0.33	3.47	0.04
FS 0013	Subgroup of Cherries, raw	RAC	0.0205	1.40	0.03	4.21	0.09	0.04	0.00	2.93	0.06	1.50	0.03	NC	-
FS 0014	Subgroup of Plums, raw (incl dried plums)	RAC	0.02	5.55	0.11	4.37	0.09	6.08	0.12	3.66	0.07	3.93	0.08	0.46	0.01
FS 0240	Apricot, raw (incl dried)	RAC	0.0485	4.27	0.21	3.31	0.16	0.06	0.00	2.86	0.14	1.71	0.08	NC	-
-	Peaches and nectarines, raw	RAC	0.055	8.76	0.48	12.98	0.71	8.23	0.45	10.09	0.55	3.64	0.20	0.04	0.00
FB 0272	Raspberries, red, black, raw	RAC	0.42	0.47	0.20	0.91	0.38	0.01	0.00	0.99	0.42	1.14	0.48	NC	-

Annex 3

SPINETORAM (233)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.05 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
FB 0020	Blueberries, raw	RAC	0.12	0.04	0.00	0.23	0.03	0.01	0.00	0.83	0.10	0.33	0.04	NC	-
FB 0021	Currants, Black, Red, White, raw	RAC	0.18	0.48	0.09	4.23	0.76	NC	-	1.51	0.27	0.49	0.09	NC	-
FB 0269	Grapes, raw (incl must, incl dried, incl juice, incl wine)	RAC	0.074	142.23	10.53	105.77	7.83	7.87	0.58	52.44	3.88	109.22	8.08	10.96	0.81
FB 0275	Strawberry, raw	RAC	0.026	4.49	0.12	5.66	0.15	0.02	0.00	6.63	0.17	5.75	0.15	0.05	0.00
FT 0305	Table olives, raw (incl preserved)	RAC	0.02	2.00	0.04	2.48	0.05	0.01	0.00	1.21	0.02	1.64	0.03	0.27	0.01
FI 0343	Litchi, raw (incl processed)	RAC	0.02	8.00	0.16	3.70	0.07	2.91	0.06	0.05	0.00	11.86	0.24	9.83	0.20
FI 0326	Avocado, raw	RAC	0.02	2.65	0.05	0.87	0.02	0.46	0.01	1.64	0.03	1.30	0.03	0.96	0.02
FI 0345	Mango, raw (incl canned mango, incl mango juice)	RAC	0.02	1.80	0.04	0.63	0.01	10.05	0.20	1.07	0.02	3.52	0.07	16.44	0.33
FI 2540	Pitaya, raw (i.e dragon fruit or pitahaya)	RAC	0.0915	NC	-	NC	-	0.08	0.01	NC	-	NC	-	NC	-
FI 0351	Passion fruit, raw	RAC	0.12	0.01	0.00	0.01	0.00	NC	-	NC	-	0.02	0.00	NC	-
VA 2031	Subgroup of bulb onions	RAC	0.01	20.67	0.21	31.32	0.31	37.52	0.38	35.08	0.35	11.77	0.12	13.74	0.14
VA 0384	Leek, raw	RAC	0.026	4.01	0.10	4.41	0.11	0.72	0.02	0.54	0.01	16.41	0.43	0.03	0.00
-	Onions, green, raw	RAC	0.33	1.55	0.51	0.74	0.24	1.05	0.35	3.74	1.23	0.94	0.31	6.45	2.13
VB 0040	Group of Brassica vegetables (excl Brassica leafy vegetables), raw	RAC	0.05	20.71	1.04	39.81	1.99	25.06	1.25	37.93	1.90	18.12	0.91	16.74	0.84
VC 2039	Subgroup of Cucumbers and Squashes, raw	RAC	0.02	7.14	0.14	16.92	0.34	37.58	0.75	15.16	0.30	4.42	0.09	12.67	0.25
VC 0046	Melons, except watermelon, raw (Cantaloupe)	RAC	0.02	9.20	0.18	11.95	0.24	14.63	0.29	8.99	0.18	7.86	0.16	2.46	0.05
VO 0448	Tomato, raw (incl juice, incl paste, incl canned)	RAC	0.02	64.74	1.29	68.31	1.37	36.05	0.72	82.09	1.64	54.50	1.09	11.69	0.23
VO 0051	Subgroup of peppers, raw (incl dried sweet peppers, excl dried chilipeppers), incl okra	RAC	0.026	6.39	0.17	15.53	0.40	19.13	0.50	10.53	0.27	8.29	0.22	5.25	0.14
VL 0483	Lettuce, leaf, raw	RAC	0.895	14.50	12.98	11.76	10.53	13.14	11.76	19.50	17.45	4.81	4.30	2.23	2.00
VL 0502	Spinach, raw	RAC	1.6	2.20	3.52	1.76	2.82	13.38	21.41	2.94	4.70	5.53	8.85	0.02	0.03
VP 0061	Beans with pods (Phaseolus spp): (immature pods + succulent seeds)	RAC	0.024	5.07	0.12	0.83	0.02	0.17	0.00	3.70	0.09	NC	-	NC	-
VD 0541	Soya bean, dry, raw (incl paste, incl curd, incl oil, incl sauce)	RAC	0.02	106.33	2.13	117.78	2.36	42.12	0.84	195.70	3.91	222.52	4.45	80.47	1.61
VR 0596	Sugar beet, raw (incl sugar)	RAC	0.02	0.01	0.00	NC	-	0.01	0.00	0.01	0.00	NC	-	NC	-
VR 0589	Potato, raw (incl flour, incl frozen, incl starch, incl tapioca)	RAC	0.02	225.03	4.50	234.24	4.68	71.48	1.43	177.55	3.55	234.55	4.69	37.71	0.75
VS 0624	Celery	RAC	0.3	7.68	2.30	2.85	0.86	NC	-	3.34	1.00	16.83	5.05	4.04	1.21

SPINETORAM (233)															
International Estimated Daily Intake (IEDI)				ADI = 0 - 0.05 mg/kg bw											
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
CM 0649 (GC 0649)	Rice, husked, dry (incl polished, incl flour, incl starch, incl oil, incl beverages)	REP	0.04	20.96	0.84	16.04	0.64	339.67	13.59	75.51	3.02	16.86	0.67	86.13	3.45
GC 0645	Maize, raw (incl glucose & dextrose & isoglucose, incl flour, incl oil, incl beer, incl germ, incl starch)	RAC	0.02	18.51	0.37	26.18	0.52	26.04	0.52	39.99	0.80	7.36	0.15	64.58	1.29
GC 0447	Sweet corn on the cob, raw (incl frozen kernels, incl canned kernels)	RAC	0.02	11.43	0.23	3.71	0.07	0.74	0.01	13.63	0.27	3.07	0.06	1.50	0.03
TN 0085	Group of Tree nuts, raw (incl processed)	RAC	0.02	8.52	0.17	8.94	0.18	15.09	0.30	9.60	0.19	14.57	0.29	26.26	0.53
SO 0691	Cotton seed, raw (incl oil)	RAC	0	10.71	0.00	4.23	0.00	7.19	0.00	7.54	0.00	5.66	0.00	2.38	0.00
HS 0444	Peppers, chili, dried	PP	0.26	0.11	0.03	0.21	0.05	0.36	0.09	0.21	0.05	0.25	0.07	0.15	0.04
DT 1114	Tea, green or black, fermented and dried, (including concentrates)	RAC	118	2.91	343.38	1.73	204.14	1.14	134.52	1.85	218.30	2.29	270.22	0.74	87.32
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.005	112.02	0.56	120.71	0.60	63.46	0.32	88.99	0.44	96.24	0.48	41.02	0.21
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.025	28.01	0.70	30.18	0.75	15.86	0.40	22.25	0.56	24.06	0.60	10.25	0.26
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.025	6.44	0.16	15.51	0.39	3.79	0.09	8.29	0.21	18.44	0.46	8.00	0.20
MO 0105	Edible offal (mammalian), raw	RAC	0.005	15.17	0.08	5.19	0.03	6.30	0.03	6.78	0.03	3.32	0.02	3.17	0.02
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.003	388.92	1.17	335.88	1.01	49.15	0.15	331.25	0.99	468.56	1.41	245.45	0.74
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0.0002	66.38	0.01	48.47	0.01	21.58	0.00	78.41	0.02	48.04	0.01	76.01	0.02
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0.004	7.38	0.03	5.39	0.02	2.40	0.01	8.71	0.03	5.34	0.02	8.45	0.03
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.004	0.10	0.00	0.10	0.00	NC	-	0.10	0.00	0.71	0.00	NC	-
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.0004	0.33	0.00	0.72	0.00	0.27	0.00	0.35	0.00	0.80	0.00	NC	-
PE 0112	Eggs, raw, (incl dried)	RAC	0.0004	25.84	0.01	29.53	0.01	28.05	0.01	33.19	0.01	36.44	0.01	8.89	0.00
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-
Total intake (ug/person)=				392.5		249.2		193.5		270.8		319.0		112.2	
Bodyweight per region (kg bw) =				60		60		55		60		60		60	
ADI (ug/person)=				3000		3000		2750		3000		3000		3000	
%ADI=				13.1%		8.3%		7.0%		9.0%		10.6%		3.7%	

Annex 3

SPINETORAM (233)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.05 mg/kg bw								
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day			Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake	
Rounded %ADI=				10%			8%		7%		9%		10%		4%	

SPINETORAM (233)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.05 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
FC 0003	Subgroup of Mandarins, raw	RAC	0.0605	0.16	0.01	0.27	0.02	9.06	0.55	0.01	0.00	0.02	0.00
FC 0004	Subgroup of Oranges, sweet, sour, raw	RAC	0.0435	1.18	0.05	1.11	0.05	14.28	0.62	0.05	0.00	1.08	0.05
JF 0004	Subgroup of Oranges, juice (single strength, incl. concentrated)	PP	0.003	0.08	0.00	0.26	0.00	12.61	0.04	0.14	0.00	0.33	0.00
FP 0009	Group of Pome fruits, raw (incl. apple juice, incl apple cider)	RAC	0.025	68.89	1.72	11.06	0.28	80.62	2.02	189.82	4.75	19.56	0.49
JF 0226	Apple juice, single strength (incl. concentrated)	PP	0.011	0.03	0.00	0.10	0.00	7.19	0.08	0.03	0.00	NC	-
FS 0013	Subgroup of Cherries, raw	RAC	0.0205	0.01	0.00	0.01	0.00	5.96	0.12	0.01	0.00	NC	-
FS 0014	Subgroup of Plums, raw (incl dried plums)	RAC	0.02	0.07	0.00	0.02	0.00	16.65	0.33	0.01	0.00	NC	-
FS 0240	Apricot, raw (incl dried)	RAC	0.0485	0.01	0.00	0.01	0.00	3.29	0.16	0.01	0.00	NC	-
-	Peaches and nectarines, raw	RAC	0.055	0.02	0.00	0.01	0.00	7.47	0.41	0.01	0.00	NC	-
FB 0272	Raspberries, red, black, raw	RAC	0.42	0.01	0.00	0.01	0.00	2.04	0.86	0.01	0.00	NC	-
FB 0020	Blueberries, raw	RAC	0.12	NC	-	NC	-	0.20	0.02	NC	-	NC	-
FB 0021	Currants, Black, Red, White, raw	RAC	0.18	0.01	0.00	NC	-	0.74	0.13	NC	-	NC	-
FB 0269	Grapes, raw (incl must, incl dried, incl juice, incl wine)	RAC	0.074	0.60	0.04	1.26	0.09	103.25	7.64	0.74	0.05	44.23	3.27
FB 0275	Strawberry, raw	RAC	0.026	0.01	0.00	0.01	0.00	3.35	0.09	0.01	0.00	0.01	0.00
FT 0305	Table olives, raw (incl preserved)	RAC	0.02	0.01	0.00	0.01	0.00	1.75	0.04	0.01	0.00	0.24	0.00
FI 0343	Litchi, raw (incl processed)	RAC	0.02	3.74	0.07	18.51	0.37	4.87	0.10	1.97	0.04	12.17	0.24
FI 0326	Avocado, raw	RAC	0.02	1.12	0.02	0.01	0.00	0.84	0.02	0.01	0.00	6.60	0.13
FI 0345	Mango, raw (incl canned mango, incl mango juice)	RAC	0.02	12.25	0.25	6.83	0.14	0.76	0.02	0.01	0.00	20.12	0.40
FI 2540	Pitaya, raw (i.e dragon fruit or pitahaya)	RAC	0.0915	0.01	0.00	0.01	0.00	NC	-	0.01	0.00	0.04	0.00

SPINETORAM (233)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.05 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
FI 0351	Passion fruit, raw	RAC	0.12	0.12	0.01	0.10	0.01	0.01	0.00	0.18	0.02	3.81	0.46
VA 2031	Subgroup of bulb onions	RAC	0.01	9.83	0.10	22.30	0.22	34.69	0.35	9.65	0.10	2.39	0.02
VA 0384	Leek, raw	RAC	0.026	0.02	0.00	1.44	0.04	1.22	0.03	0.01	0.00	NC	-
-	Onions, green, raw	RAC	0.33	1.43	0.47	0.05	0.02	0.20	0.07	NC	-	6.30	2.08
VB 0040	Group of Brassica vegetables (excl Brassica leafy vegetables), raw	RAC	0.05	5.46	0.27	4.28	0.21	58.72	2.94	0.02	0.00	NC	-
VC 2039	Subgroup of Cucumbers and Squashes, raw	RAC	0.02	0.92	0.02	3.20	0.06	13.55	0.27	1.91	0.04	0.05	0.00
VC 0046	Melons, except watermelon, raw (Cantaloupe)	RAC	0.02	0.19	0.00	0.10	0.00	4.98	0.10	0.01	0.00	NC	-
VO 0448	Tomato, raw (incl juice, incl paste, incl canned)	RAC	0.02	15.50	0.31	5.78	0.12	71.52	1.43	2.00	0.04	12.50	0.25
VO 0051	Subgroup of peppers, raw (incl dried sweet peppers, excl dried chilipeppers), incl okra	RAC	0.026	15.20	0.40	14.23	0.37	25.14	0.65	0.91	0.02	NC	-
VL 0483	Lettuce, leaf, raw	RAC	0.895	0.29	0.26	0.03	0.03	6.71	6.01	0.01	0.01	NC	-
VL 0502	Spinach, raw	RAC	1.6	0.17	0.27	0.01	0.02	0.81	1.30	0.01	0.02	NC	-
VP 0061	Beans with pods (Phaseolus spp): (immature pods + succulent seeds)	RAC	0.024	NC	-	NC	-	NC	-	NC	-	NC	-
VD 0541	Soya bean, dry, raw (incl paste, incl curd, incl oil, incl sauce)	RAC	0.02	15.80	0.32	14.29	0.29	104.36	2.09	17.11	0.34	35.20	0.70
VR 0596	Sugar beet, raw (incl sugar)	RAC	0.02	3.93	0.08	1.68	0.03	NC	-	NC	-	36.12	0.72
VR 0589	Potato, raw (incl flour, incl frozen, incl starch, incl tapioca)	RAC	0.02	23.96	0.48	13.56	0.27	213.41	4.27	104.35	2.09	8.56	0.17
VS 0624	Celery	RAC	0.3	3.66	1.10	2.65	0.80	4.84	1.45	2.47	0.74	4.94	1.48
CM 0649 (GC 0649)	Rice, husked, dry (incl polished, incl flour, incl starch, incl oil, incl beverages)	REP	0.04	52.55	2.10	286.02	11.44	18.64	0.75	19.67	0.79	75.09	3.00
GC 0645	Maize, raw (incl glucose & dextrose & isoglucose, incl flour, incl oil, incl beer, incl germ, incl starch)	RAC	0.02	116.66	2.33	10.52	0.21	38.46	0.77	76.60	1.53	34.44	0.69
GC 0447	Sweet corn on the cob, raw (incl frozen kernels, incl canned kernels)	RAC	0.02	3.63	0.07	20.50	0.41	8.78	0.18	0.02	0.00	0.17	0.00
TN 0085	Group of Tree nuts, raw (incl processed)	RAC	0.02	4.39	0.09	135.53	2.71	6.11	0.12	0.72	0.01	317.74	6.35
SO 0691	Cotton seed, raw (incl oil)	RAC	0	8.14	0.00	0.32	0.00	2.84	0.00	2.69	0.00	0.97	0.00
HS 0444	Peppers, chili, dried	PP	0.26	0.58	0.15	1.27	0.33	1.21	0.31	0.12	0.03	NC	-
DT 1114	Tea, green or black, fermented and dried, (including concentrates)	RAC	118	0.53	62.54	5.25	619.50	0.86	101.48	0.56	66.08	0.88	103.84
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.005	23.34	0.12	40.71	0.20	97.15	0.49	18.06	0.09	57.71	0.29

Annex 3

SPINETORAM (233)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.05 mg/kg bw						
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person								
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake	
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.025	5.84	0.15	10.18	0.25	24.29	0.61	4.52	0.11	14.43	0.36	
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.025	1.05	0.03	1.14	0.03	18.69	0.47	0.94	0.02	3.12	0.08	
MO 0105	Edible offal (mammalian), raw	RAC	0.005	4.64	0.02	1.97	0.01	10.01	0.05	3.27	0.02	3.98	0.02	
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.003	108.75	0.33	70.31	0.21	436.11	1.31	61.55	0.18	79.09	0.24	
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0.0002	3.53	0.00	10.83	0.00	51.36	0.01	4.53	0.00	50.00	0.01	
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0.004	0.39	0.00	1.20	0.00	5.71	0.02	0.50	0.00	5.56	0.02	
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.004	NC	-	NC	-	0.32	0.00	NC	-	NC	-	
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.0004	0.10	0.00	0.70	0.00	0.97	0.00	0.10	0.00	NC	-	
PE 0112	Eggs, raw, (incl dried)	RAC	0.0004	3.84	0.00	4.41	0.00	27.25	0.01	1.13	0.00	7.39	0.00	
-	-	-		-	-	-	-	-	-	-	-	-	-	
Total intake (ug/person)=					74.2		638.8		140.8		77.1		125.4	
Bodyweight per region (kg bw) =					60		60		60		60		60	
ADI (ug/person)=					3000		3000		3000		3000		3000	
%ADI=					2.5%		21.3%		4.7%		2.6%		4.2%	
Rounded %ADI=					2%		20%		5%		3%		4%	

SULFOXAFLOR (252)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.05 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
FC 0002	Subgroup of Lemons and limes, raw (incl lemon juice) (incl kumquat commodities)	RAC	0.038	4.82	0.18	2.45	0.09	3.93	0.15	25.44	0.97	8.74	0.33	16.23	0.62
FC 0003	Subgroup of Mandarins, raw (incl mandarin juice)	RAC	0.26	6.18	1.61	3.66	0.95	0.25	0.07	6.82	1.77	3.49	0.91	19.38	5.04
FC 0004	Subgroup of Oranges, sweet, sour, raw	RAC	0.26	20.66	5.37	5.23	1.36	11.90	3.09	37.90	9.85	21.16	5.50	56.46	14.68
JF 0004	Subgroup of Oranges, juice (single strength, incl. concentrated)	PP	0.036	1.27	0.05	2.20	0.08	0.09	0.00	11.81	0.43	0.46	0.02	1.69	0.06
FC 0005	Subgroup of Pummelo and grapefruits, raw (incl grapefruit juice)	RAC	0.0125	0.66	0.01	0.69	0.01	0.96	0.01	10.20	0.13	1.25	0.02	2.97	0.04
FP 0009	Group of Pome fruits, raw (incl apple cider, excl apple juice)	RAC	0.067	19.35	1.30	34.06	2.28	17.87	1.20	25.74	1.72	7.69	0.52	56.85	3.81
JF 0226	Apple juice, single strength (incl. concentrated)	PP	0.067	0.32	0.02	3.07	0.21	0.07	0.00	5.00	0.34	0.29	0.02	5.57	0.37
FS 0013	Subgroup of Cherries, raw	RAC	0.34	0.92	0.31	9.15	3.11	0.01	0.00	0.61	0.21	0.06	0.02	6.64	2.26
FS 0014	Subgroup of Plums, raw (incl dried plums)	RAC	0.038	2.67	0.10	8.77	0.33	0.07	0.00	3.03	0.12	0.70	0.03	4.34	0.16
FS 2001	Subgroup of peaches, raw (incl dried apricots)	RAC	0.061	8.01	0.49	5.87	0.36	0.18	0.01	8.19	0.50	1.64	0.10	22.46	1.37
FB 2005	Subgroup of Caneberries, raw	RAC	0.44	0.42	0.18	1.05	0.46	0.01	0.00	0.02	0.01	0.02	0.01	1.24	0.55
FB 2006	Subgroup of Bush berries, raw (including processed)	RAC	0.39	0.53	0.21	1.31	0.51	0.40	0.16	1.66	0.65	0.01	0.00	0.99	0.39
FB 0267	Elderberries, raw (incl processed)	RAC	0.39	0.44	0.17	0.27	0.11	0.34	0.13	1.41	0.55	NC	-	0.87	0.34
FB 0269	Grapes, raw (incl must, incl juice, excl dried, excl wine)	RAC	0.14	13.19	1.85	9.61	1.35	0.09	0.01	17.28	2.42	4.00	0.56	54.50	7.63
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	0.49	0.51	0.25	0.51	0.25	0.01	0.00	1.27	0.62	0.12	0.06	2.07	1.01
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.098	0.67	0.07	12.53	1.23	2.01	0.20	1.21	0.12	3.53	0.35	4.01	0.39
FB 0275	Strawberry, raw	RAC	0.19	0.70	0.13	2.01	0.38	0.04	0.01	1.36	0.26	0.37	0.07	2.53	0.48
FI 0326	Avocado, raw	RAC	0.011	0.13	0.00	0.03	0.00	2.05	0.02	2.54	0.03	2.34	0.03	0.12	0.00
FI 0345	Mango, raw (incl canned mango, incl mango juice)	RAC	0.022	10.48	0.23	0.01	0.00	7.24	0.16	6.87	0.15	19.98	0.44	6.25	0.14
VA 0381	Garlic, raw	RAC	0.01	2.29	0.02	5.78	0.06	0.11	0.00	3.69	0.04	1.65	0.02	3.91	0.04
-	Onions, dry, raw	RAC	0	29.36	0.00	37.50	0.00	3.56	0.00	34.78	0.00	18.81	0.00	43.38	0.00
-	Onions, green, raw	RAC	0.11	2.45	0.27	1.49	0.16	1.02	0.11	2.60	0.29	0.60	0.07	2.03	0.22
VB 0400	Broccoli, raw	RAC	0.074	0.88	0.07	0.17	0.01	0.01	0.00	1.25	0.09	3.00	0.22	1.09	0.08
VB 0404	Cauliflower, raw	RAC	0.012	1.65	0.02	0.32	0.00	0.01	0.00	2.33	0.03	4.79	0.06	2.03	0.02

Annex 3

SULFOXAFLO (252)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.05 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
VB 0041	Cabbages, head, raw	RAC	0.099	2.73	0.27	27.92	2.76	0.55	0.05	4.47	0.44	4.27	0.42	10.25	1.01
VB 0467	Chinese cabbage, type pe-tsai, raw	RAC	1.2	0.45	0.54	4.56	5.47	0.09	0.11	0.73	0.88	NC	-	1.67	2.00
VC 0045	Group of Fruiting vegetables, cucurbits, raw	RAC	0.029	53.14	1.54	86.21	2.50	6.28	0.18	92.76	2.69	15.64	0.45	155.30	4.50
VO 0448	Tomato, raw (incl canned, excl juice, excl paste)	RAC	0.11	42.04	4.62	76.13	8.37	10.69	1.18	84.59	9.30	24.92	2.74	203.27	22.36
-	Tomato, paste (i.e. concentrated tomato sauce/puree)	PP	0.23	2.34	0.54	1.33	0.31	1.57	0.36	4.24	0.98	0.34	0.08	2.83	0.65
JF 0448	Tomato, juice (single strength, incl concentrated)	PP	0.052	0.29	0.02	0.29	0.02	0.01	0.00	0.38	0.02	0.05	0.00	0.14	0.01
VO 0442	Okra, raw (i.e. Lady's Finger, Gombo)	RAC	0.11	1.97	0.22	NC	-	3.68	0.40	3.24	0.36	5.72	0.63	1.57	0.17
VO 0444	Peppers, chili, raw	RAC	0.11	3.99	0.44	7.30	0.80	2.93	0.32	5.62	0.62	NC	-	17.44	1.92
VO 0445	Peppers, sweet, raw (incl dried)	RAC	0.11	4.49	0.49	6.44	0.71	7.21	0.79	5.68	0.62	9.52	1.05	8.92	0.98
VO 0440	Egg plant, raw (i.e. aubergine)	RAC	0.11	5.58	0.61	4.31	0.47	0.89	0.10	9.31	1.02	13.64	1.50	20.12	2.21
VL 2050	Subgroup of Leafy greens	RAC	1.2	3.93	4.72	5.28	6.34	3.07	3.68	14.53	17.44	8.25	9.90	12.75	15.30
VL 0054	Subgroup of Leaves of Brassicaceae, raw	RAC	1.2	2.63	3.16	9.27	11.12	1.86	2.23	5.82	6.98	19.53	23.44	4.90	5.88
VL 2052	Subgroup of Leaves of Root and Tuber Vegetables	RAC	1.2	0.18	0.22	0.31	0.37	0.84	1.01	0.47	0.56	2.06	2.47	0.23	0.28
VL 2053	Subgroup of Leaves of trees, shrubs and vines	RAC	1.2	0.39	0.47	0.69	0.83	0.43	0.52	1.04	1.25	4.57	5.48	0.50	0.60
VL 2054	Subgroup of Leafy aquatic vegetables	RAC	1.2	1.21	1.45	2.15	2.58	1.33	1.60	3.24	3.89	11.36	13.63	1.56	1.87
VD 0071	Beans, dry, raw (Phaseolus spp)	RAC	0.075	2.39	0.18	1.61	0.12	10.47	0.79	1.84	0.14	12.90	0.97	7.44	0.56
VD 0541	Soya bean, dry, raw (incl flour, incl paste, incl curd, incl sauce, excl oil)	RAC	0.011	0.63	0.01	1.09	0.01	0.40	0.00	1.40	0.02	1.68	0.02	0.48	0.01
OR 0541	Soya oil, refined	PP	0.0033	12.99	0.04	10.43	0.03	3.63	0.01	13.10	0.04	10.70	0.04	13.10	0.04
VR 0574	Beetroot, raw	RAC	0.01	3.42	0.03	6.06	0.06	3.75	0.04	9.11	0.09	NC	-	4.39	0.04
VR 0575	Burdock, greater or edible, raw	RAC	0.01	0.03	0.00	0.06	0.00	0.04	0.00	0.09	0.00	NC	-	0.04	0.00
VR 0577	Carrots, raw	RAC	0.01	9.51	0.10	30.78	0.31	0.37	0.00	8.75	0.09	2.80	0.03	6.10	0.06
VR 0578	Celeriac, raw	RAC	0.01	1.70	0.02	3.01	0.03	1.87	0.02	4.53	0.05	NC	-	2.19	0.02
VR 0469	Chicory, roots, raw	RAC	0.01	0.01	0.00	0.20	0.00	0.01	0.00	0.01	0.00	0.02	0.00	0.01	0.00
VR 0583	Horseradish, raw	RAC	0.01	0.51	0.01	0.91	0.01	0.56	0.01	1.37	0.01	NC	-	0.66	0.01
VR 0587	Parsley turnip-rooted, raw	RAC	0.01	0.32	0.00	0.57	0.01	0.35	0.00	0.85	0.01	NC	-	0.41	0.00
VR 0588	Parsnip, raw	RAC	0.01	0.59	0.01	1.05	0.01	0.65	0.01	1.58	0.02	NC	-	0.76	0.01

SULFOXAFLOR (252)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.05 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
VR 0494	Radish roots, raw	RAC	0.01	2.31	0.02	4.09	0.04	2.53	0.03	6.15	0.06	5.88	0.06	2.97	0.03
VR 0591	Japanese radish, raw (i.e. Chinese radish, Daikon)	RAC	0.01	1.90	0.02	3.36	0.03	2.08	0.02	5.06	0.05	NC	-	2.44	0.02
VR 0498	Salsify, raw (i.e. Oysterplant)	RAC	0.01	0.21	0.00	0.37	0.00	0.23	0.00	0.55	0.01	NC	-	0.27	0.00
VR 0596	Sugar beet, raw	RAC	0.014	NC	-	NC	-	NC	-	NC	-	0.01	0.00	NC	-
-	Sugar beet, sugar	PP	0.025	0.02	0.00	NC	-	0.01	0.00	0.09	0.00	0.07	0.00	12.63	0.32
VR 0497	Swede, raw (i.e. Rutabaga)	RAC	0.01	1.58	0.02	2.80	0.03	1.74	0.02	4.21	0.04	NC	-	2.03	0.02
VR 0506	Turnip, garden, raw	RAC	0.01	2.50	0.03	4.44	0.04	2.75	0.03	6.67	0.07	0.14	0.00	3.22	0.03
VR 0573	Arrowroot, raw	RAC	0.01	1.53	0.02	0.01	0.00	0.93	0.01	1.33	0.01	0.47	0.00	0.02	0.00
VR 0463	Cassava raw (incl starch, incl tapioca, incl flour) (i.e. Manioc)	RAC	0.01	0.08	0.00	0.01	0.00	482.56	4.83	0.99	0.01	25.75	0.26	3.29	0.03
VR 0585	Jerusalem artichoke, raw (i.e. Topinambur)	RAC	0.01	1.57	0.02	0.01	0.00	0.96	0.01	1.36	0.01	0.48	0.00	0.02	0.00
VR 0589	Potato, raw (incl flour, incl frozen, incl starch, incl tapioca)	RAC	0.01	59.74	0.60	316.14	3.16	9.78	0.10	60.26	0.60	54.12	0.54	119.82	1.20
VR 0508	Sweet potato, raw (incl dried)	RAC	0.01	0.18	0.00	0.18	0.00	42.16	0.42	1.61	0.02	3.06	0.03	6.67	0.07
VR 0504	Tannia, raw (i.e. Tanier, Yautia)	RAC	0.01	NC	-	NC	-	NC	-	0.01	0.00	0.26	0.00	1.27	0.01
VR 0505	Taro, raw (i.e. Dasheen, Eddoe)	RAC	0.01	0.01	0.00	NC	-	25.12	0.25	0.04	0.00	0.01	0.00	0.97	0.01
VR 0600	Yams, raw (incl dried)	RAC	0.01	0.02	0.00	NC	-	90.40	0.90	6.45	0.06	0.74	0.01	0.65	0.01
VR 2072	Subgroup of Aquatic root and tuber vegetables	RAC	0.01	NC	-	NC	-	NC	-	NC	-	NC	-	NC	-
VS 0624	Celery	RAC	0.19	2.14	0.41	3.79	0.72	2.35	0.45	5.69	1.08	0.02	0.00	2.75	0.52
VS 0621	Asparagus, raw	RAC	0.01	0.01	0.00	0.01	0.00	0.01	0.00	0.03	0.00	0.07	0.00	0.21	0.00
GC 0653	Triticale, raw (incl flour)	RAC	0.025	NC	-	NC	-	NC	-	0.01	0.00	0.39	0.01	NC	-
GC 0654	Wheat, raw (incl bulgur, incl fermented beverages, incl germ, incl wholemeal bread, incl white flour products, incl white bread)	RAC	0.025	381.15	9.53	341.55	8.54	38.35	0.96	281.89	7.05	172.83	4.32	434.07	10.85
GC 0640	Barley, raw (incl malt extract, incl malt, excl pot&pearled, excl flour & grits,excl beer)	RAC	0.063	2.62	0.17	1.50	0.09	0.37	0.02	0.60	0.04	0.38	0.02	0.53	0.03
-	Barley, pot&pearled	PP	0.044	7.12	0.31	7.34	0.32	0.02	0.00	0.03	0.00	0.67	0.03	0.20	0.01
-	Barley beer	PP	0.013	4.87	0.06	93.78	1.22	24.28	0.32	12.76	0.17	39.28	0.51	18.15	0.24
SO 0495	Rape seed, raw	RAC	0.045	0.02	0.00	NC	-	NC	-	0.01	0.00	0.75	0.03	0.01	0.00
OR 0495	Rape seed oil, edible	PP	0.014	0.35	0.00	0.44	0.01	0.19	0.00	0.97	0.01	3.28	0.05	0.77	0.01

Annex 3

SULFOXAFLOR (252)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.05 mg/kg bw								
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day										
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake	
OR 0691	Cotton seed oil, edible	PP	0.002	3.22	0.01	1.54	0.00	1.01	0.00	0.74	0.00	1.12	0.00	2.93	0.01	
SB 0716	Coffee bean, raw (i.e. green coffee)	RAC	0.035	0.96	0.03	0.16	0.01	0.91	0.03	0.27	0.01	1.37	0.05	0.46	0.02	
SM 0716	Coffee bean, roasted	PP	0.0167	0.19	0.00	0.91	0.02	0.16	0.00	2.50	0.04	0.39	0.01	0.40	0.01	
-	Coffee bean, instant coffee (incl essences and concentrates)	PP	0.102	0.07	0.01	0.94	0.10	0.07	0.01	0.70	0.07	0.07	0.01	0.29	0.03	
HS 0444	Peppers, chili, dried	PP	1.1	0.42	0.46	0.53	0.58	0.84	0.92	0.50	0.55	0.95	1.05	0.37	0.41	
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.045	24.96	1.12	57.95	2.61	16.70	0.75	38.38	1.73	26.46	1.19	29.00	1.31	
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.03	6.24	0.19	14.49	0.43	4.18	0.13	9.60	0.29	6.62	0.20	7.25	0.22	
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.03	3.29	0.10	6.14	0.18	0.82	0.02	1.57	0.05	2.23	0.07	1.07	0.03	
MO 0105	Edible offal (mammalian), raw	RAC	0.13	4.79	0.62	9.68	1.26	2.97	0.39	5.49	0.71	3.84	0.50	5.03	0.65	
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.05	289.65	14.48	485.88	24.29	26.92	1.35	239.03	11.95	199.91	10.00	180.53	9.03	
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0.015	13.17	0.20	26.78	0.40	7.24	0.11	116.71	1.75	22.54	0.34	32.09	0.48	
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0.005	1.46	0.01	2.98	0.01	0.80	0.00	12.97	0.06	2.50	0.01	3.57	0.02	
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.005	0.10	0.00	0.10	0.00	NC	-	0.10	0.00	0.10	0.00	0.10	0.00	
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.046	0.12	0.01	0.12	0.01	0.11	0.01	5.37	0.25	0.24	0.01	0.10	0.00	
PE 0112	Eggs, raw, (incl dried)	RAC	0.013	7.84	0.10	23.08	0.30	2.88	0.04	14.89	0.19	9.81	0.13	14.83	0.19	
Total intake (ug/person)=					61.1		101.0		30.9		95.7		91.6		126.0	
Bodyweight per region (kg bw) =					60		60		60		60		60		60	
ADI (ug/person)=					3000		3000		3000		3000		3000		3000	
%ADI=					2.0%		3.4%		1.0%		3.2%		3.1%		4.2%	
Rounded %ADI=					2%		3%		1%		3%		3%		4%	

SULFOXAFLOL (252)		International Estimated Daily Intake (IEDI)				ADI = 0 - 0.05 mg/kg bw									
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
FC 0002	Subgroup of Lemons and limes, raw (incl lemon juice) (incl kumquat commodities)	RAC	0.038	10.12	0.38	15.69	0.60	2.88	0.11	12.30	0.47	22.32	0.85	6.59	0.25
FC 0003	Subgroup of Mandarins, raw (incl mandarin juice)	RAC	0.26	12.42	3.23	14.99	3.90	16.08	4.18	10.78	2.80	9.94	2.58	NC	-
FC 0004	Subgroup of Oranges, sweet, sour, raw	RAC	0.26	15.68	4.08	24.00	6.24	6.80	1.77	29.09	7.56	15.39	4.00	160.47	41.72
JF 0004	Subgroup of Oranges, juice (single strength, incl. concentrated)	PP	0.036	33.31	1.20	1.78	0.06	0.28	0.01	18.97	0.68	14.01	0.50	13.36	0.48
FC 0005	Subgroup of Pummelo and grapefruits, raw (incl grapefruit juice)	RAC	0.0125	8.21	0.10	4.60	0.06	0.64	0.01	5.85	0.07	19.98	0.25	368.86	4.61
FP 0009	Group of Pome fruits, raw (incl apple cider, excl apple juice)	RAC	0.067	51.09	3.42	65.40	4.38	42.71	2.86	45.29	3.03	62.51	4.19	7.74	0.52
JF 0226	Apple juice, single strength (incl. concentrated)	PP	0.067	14.88	1.00	11.98	0.80	0.15	0.01	9.98	0.67	30.32	2.03	3.47	0.23
FS 0013	Subgroup of Cherries, raw	RAC	0.34	1.40	0.48	4.21	1.43	0.04	0.01	2.93	1.00	1.50	0.51	NC	-
FS 0014	Subgroup of Plums, raw (incl dried plums)	RAC	0.038	5.55	0.21	4.37	0.17	6.08	0.23	3.66	0.14	3.93	0.15	0.46	0.02
FS 2001	Subgroup of peaches, raw (incl dried apricots)	RAC	0.061	13.03	0.79	16.29	0.99	8.29	0.51	12.95	0.79	5.35	0.33	0.04	0.00
FB 2005	Subgroup of Caneberries, raw	RAC	0.44	0.56	0.25	1.43	0.63	0.14	0.06	1.23	0.54	1.14	0.50	0.01	0.00
FB 2006	Subgroup of Bush berries, raw (including processed)	RAC	0.39	1.31	0.51	5.50	2.15	0.01	0.00	2.57	1.00	0.82	0.32	2.15	0.84
FB 0267	Elderberries, raw (incl processed)	RAC	0.39	8.20	3.20	0.14	0.05	NC	-	NC	-	NC	-	1.87	0.73
FB 0269	Grapes, raw (incl must, incl juice, excl dried, excl wine)	RAC	0.14	7.18	1.01	13.73	1.92	5.24	0.73	12.27	1.72	7.46	1.04	1.21	0.17
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	0.49	3.09	1.51	1.51	0.74	0.03	0.01	1.38	0.68	4.26	2.09	0.42	0.21
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.098	88.93	8.72	62.41	6.12	1.84	0.18	25.07	2.46	61.17	5.99	5.84	0.57
FB 0275	Strawberry, raw	RAC	0.19	4.49	0.85	5.66	1.08	0.02	0.00	6.63	1.26	5.75	1.09	0.05	0.01
FI 0326	Avocado, raw	RAC	0.011	2.65	0.03	0.87	0.01	0.46	0.01	1.64	0.02	1.30	0.01	0.96	0.01
FI 0345	Mango, raw (incl canned mango, incl mango juice)	RAC	0.022	1.80	0.04	0.63	0.01	10.05	0.22	1.07	0.02	3.52	0.08	16.44	0.36
VA 0381	Garlic, raw	RAC	0.01	0.98	0.01	1.49	0.01	12.88	0.13	3.74	0.04	2.05	0.02	1.14	0.01
-	Onions, dry, raw	RAC	0	19.69	0.00	29.83	0.00	24.64	0.00	31.35	0.00	9.72	0.00	12.59	0.00
-	Onions, green, raw	RAC	0.11	1.55	0.17	0.74	0.08	1.05	0.12	3.74	0.41	0.94	0.10	6.45	0.71
VB 0400	Broccoli, raw	RAC	0.074	4.24	0.31	1.76	0.13	NC	-	0.51	0.04	3.79	0.28	0.26	0.02
VB 0404	Cauliflower, raw	RAC	0.012	5.27	0.06	5.01	0.06	NC	-	2.70	0.03	5.57	0.07	0.49	0.01
VB 0041	Cabbages, head, raw	RAC	0.099	8.97	0.89	27.12	2.68	1.44	0.14	24.96	2.47	4.55	0.45	11.23	1.11

Annex 3

SULFOXAFLOL (252)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.05 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
VB 0467	Chinese cabbage, type pe-tsai, raw	RAC	1.2	NC	-	NC	-	17.39	20.87	9.44	11.33	NC	-	1.83	2.20
VC 0045	Group of Fruiting vegetables, cucurbits, raw	RAC	0.029	27.81	0.81	41.93	1.22	123.30	3.58	49.47	1.43	15.95	0.46	35.99	1.04
VO 0448	Tomato, raw (incl canned, excl juice, excl paste)	RAC	0.11	43.88	4.83	55.41	6.10	35.38	3.89	74.88	8.24	26.50	2.92	9.51	1.05
-	Tomato, paste (i.e. concentrated tomato sauce/puree)	PP	0.23	4.96	1.14	3.20	0.74	0.15	0.03	1.61	0.37	6.88	1.58	0.52	0.12
JF 0448	Tomato, juice (single strength, incl concentrated)	PP	0.052	0.80	0.04	0.07	0.00	0.05	0.00	0.61	0.03	0.40	0.02	0.08	0.00
VO 0442	Okra, raw (i.e. Lady's Finger, Gombo)	RAC	0.11	NC	-	NC	-	0.04	0.00	0.17	0.02	NC	-	0.72	0.08
VO 0444	Peppers, chili, raw	RAC	0.11	5.57	0.61	14.00	1.54	8.25	0.91	5.77	0.63	6.44	0.71	2.53	0.28
VO 0445	Peppers, sweet, raw (incl dried)	RAC	0.11	0.82	0.09	1.53	0.17	10.85	1.19	4.59	0.50	1.84	0.20	2.00	0.22
VO 0440	Egg plant, raw (i.e. aubergine)	RAC	0.11	1.01	0.11	1.69	0.19	21.37	2.35	3.00	0.33	1.40	0.15	NC	-
VL 2050	Subgroup of Leafy greens	RAC	1.2	18.38	22.06	18.73	22.48	82.36	98.83	25.32	30.38	17.60	21.12	7.37	8.84
VL 0054	Subgroup of Leaves of Brassicaceae, raw	RAC	1.2	0.10	0.12	NC	-	26.78	32.14	5.00	6.00	0.58	0.70	5.68	6.82
VL 2052	Subgroup of Leaves of Root and Tuber Vegetables	RAC	1.2	NC	-	NC	-	NC	-	NC	-	NC	-	0.33	0.40
VL 2053	Subgroup of Leaves of trees, shrubs and vines	RAC	1.2	NC	-	NC	-	NC	-	NC	-	NC	-	0.74	0.89
VL 2054	Subgroup of Leafy aquatic vegetables	RAC	1.2	0.35	0.42	3.13	3.76	3.74	4.49	1.79	2.15	NC	-	2.30	2.76
VD 0071	Beans, dry, raw (Phaseolus spp)	RAC	0.075	1.51	0.11	1.50	0.11	1.90	0.14	5.11	0.38	1.36	0.10	23.43	1.76
VD 0541	Soya bean, dry, raw (incl flour, incl paste, incl curd, incl sauce, excl oil)	RAC	0.011	0.47	0.01	0.77	0.01	9.12	0.10	8.05	0.09	0.04	0.00	6.06	0.07
OR 0541	Soya oil, refined	PP	0.0033	19.06	0.06	21.06	0.07	5.94	0.02	33.78	0.11	40.05	0.13	13.39	0.04
VR 0574	Beetroot, raw	RAC	0.01	9.91	0.10	6.34	0.06	NC	-	9.65	0.10	19.11	0.19	6.47	0.06
VR 0575	Burdock, greater or edible, raw	RAC	0.01	NC	-	NC	-	NC	-	0.48	0.00	NC	-	0.06	0.00
VR 0577	Carrots, raw	RAC	0.01	26.26	0.26	27.13	0.27	10.07	0.10	16.49	0.16	44.69	0.45	8.75	0.09
VR 0578	Celeriac, raw	RAC	0.01	2.97	0.03	1.79	0.02	NC	-	0.06	0.00	16.91	0.17	3.22	0.03
VR 0469	Chicory, roots, raw	RAC	0.01	0.01	0.00	0.51	0.01	0.01	0.00	0.01	0.00	21.12	0.21	NC	-
VR 0583	Horseradish, raw	RAC	0.01	0.01	0.00	0.42	0.00	13.01	0.13	0.26	0.00	2.70	0.03	0.97	0.01
VR 0587	Parsley turnip-rooted, raw	RAC	0.01	NC	-	NC	-	NC	-	NC	-	NC	-	0.61	0.01
VR 0588	Parsnip, raw	RAC	0.01	4.42	0.04	0.06	0.00	NC	-	NC	-	NC	-	1.12	0.01
VR 0494	Radish roots, raw	RAC	0.01	3.83	0.04	11.99	0.12	NC	-	5.26	0.05	2.19	0.02	4.37	0.04

SULFOXAFLO (252)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.05 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
VR 0591	Japanese radish, raw (i.e. Chinese radish, Daikon)	RAC	0.01	NC	-	NC	-	26.64	0.27	18.92	0.19	NC	-	3.59	0.04
VR 0498	Salsify, raw (i.e. Oysterplant)	RAC	0.01	1.02	0.01	0.52	0.01	NC	-	NC	-	2.08	0.02	0.39	0.00
VR 0596	Sugar beet, raw	RAC	0.014	0.01	0.00	NC	-	0.01	0.00	0.01	0.00	NC	-	NC	-
-	Sugar beet, sugar	PP	0.025	0.01	0.00	NC	-	0.01	0.00	NC	-	NC	-	NC	-
VR 0497	Swede, raw (i.e. Rutabaga)	RAC	0.01	10.01	0.10	1.66	0.02	NC	-	NC	-	3.06	0.03	2.99	0.03
VR 0506	Turnip, garden, raw	RAC	0.01	5.78	0.06	15.35	0.15	NC	-	6.54	0.07	1.95	0.02	4.73	0.05
VR 0573	Arrowroot, raw	RAC	0.01	0.02	0.00	0.01	0.00	2.05	0.02	0.21	0.00	NC	-	0.76	0.01
VR 0463	Cassava raw (incl starch, incl tapioca, incl flour) (i.e. Manioc)	RAC	0.01	0.01	0.00	NC	-	20.96	0.21	0.14	0.00	NC	-	9.62	0.10
VR 0585	Jerusalem artichoke, raw (i.e. Topinambur)	RAC	0.01	0.11	0.00	0.01	0.00	NC	-	0.22	0.00	NC	-	0.78	0.01
VR 0589	Potato, raw (incl flour, incl frozen, incl starch, incl tapioca)	RAC	0.01	225.03	2.25	234.24	2.34	71.48	0.71	177.55	1.78	234.55	2.35	37.71	0.38
VR 0508	Sweet potato, raw (incl dried)	RAC	0.01	0.93	0.01	0.32	0.00	64.65	0.65	5.37	0.05	0.30	0.00	3.13	0.03
VR 0504	Tannia, raw (i.e. Tanier, Yautia)	RAC	0.01	NC	-	NC	-	NC	-	0.01	0.00	NC	-	10.74	0.11
VR 0505	Taro, raw (i.e. Dasheen, Eddoe)	RAC	0.01	NC	-	NC	-	1.93	0.02	0.84	0.01	NC	-	19.94	0.20
VR 0600	Yams, raw (incl dried)	RAC	0.01	NC	-	NC	-	0.03	0.00	0.71	0.01	NC	-	17.57	0.18
VR 2072	Subgroup of Aquatic root and tuber vegetables	RAC	0.01	NC	-	NC	-	3.42	0.03	NC	-	NC	-	NC	-
VS 0624	Celery	RAC	0.19	7.68	1.46	2.85	0.54	NC	-	3.34	0.63	16.83	3.20	4.04	0.77
VS 0621	Asparagus, raw	RAC	0.01	0.84	0.01	2.08	0.02	7.11	0.07	1.01	0.01	1.69	0.02	0.04	0.00
GC 0653	Triticale, raw (incl flour)	RAC	0.025	0.01	0.00	0.17	0.00	0.29	0.01	0.01	0.00	NC	-	NC	-
GC 0654	Wheat, raw (incl bulgur, incl fermented beverages, incl germ, incl wholemeal bread, incl white flour products, incl white bread)	RAC	0.025	253.07	6.33	244.73	6.12	134.44	3.36	235.10	5.88	216.39	5.41	167.40	4.19
GC 0640	Barley, raw (incl malt extract, incl malt, excl pot&pearled, excl flour & grits,excl beer)	RAC	0.063	0.93	0.06	0.15	0.01	0.14	0.01	1.56	0.10	0.33	0.02	3.42	0.22
-	Barley, pot&pearled	PP	0.044	0.57	0.03	2.56	0.11	0.33	0.01	0.56	0.02	0.36	0.02	NC	-
-	Barley beer	PP	0.013	180.21	2.34	259.46	3.37	45.91	0.60	172.36	2.24	234.42	3.05	65.30	0.85
SO 0495	Rape seed, raw	RAC	0.045	NC	-	NC	-	0.01	0.00	NC	-	NC	-	NC	-
OR 0495	Rape seed oil, edible	PP	0.014	12.52	0.18	7.63	0.11	3.00	0.04	6.01	0.08	NC	-	NC	-
OR 0691	Cotton seed oil, edible	PP	0.002	1.68	0.00	0.66	0.00	1.13	0.00	1.18	0.00	0.89	0.00	0.37	0.00

Annex 3

SULFOXAFLOL (252)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.05 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
SB 0716	Coffee bean, raw (i.e. green coffee)	RAC	0.035	0.60	0.02	NC	-	0.62	0.02	1.71	0.06	NC	-	3.51	0.12
SM 0716	Coffee bean, roasted	PP	0.0167	7.02	0.12	9.75	0.16	0.02	0.00	5.09	0.09	13.38	0.22	0.77	0.01
-	Coffee bean, instant coffee (incl essences and concentrates)	PP	0.102	0.75	0.08	0.30	0.03	0.04	0.00	0.67	0.07	2.43	0.25	1.43	0.15
HS 0444	Peppers, chili, dried	PP	1.1	0.11	0.12	0.21	0.23	0.36	0.40	0.21	0.23	0.25	0.28	0.15	0.17
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.045	112.02	5.04	120.71	5.43	63.46	2.86	88.99	4.00	96.24	4.33	41.02	1.85
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.03	28.01	0.84	30.18	0.91	15.86	0.48	22.25	0.67	24.06	0.72	10.25	0.31
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.03	6.44	0.19	15.51	0.47	3.79	0.11	8.29	0.25	18.44	0.55	8.00	0.24
MO 0105	Edible offal (mammalian), raw	RAC	0.13	15.17	1.97	5.19	0.67	6.30	0.82	6.78	0.88	3.32	0.43	3.17	0.41
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.05	388.92	19.45	335.88	16.79	49.15	2.46	331.25	16.56	468.56	23.43	245.45	12.27
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0.015	66.38	1.00	48.47	0.73	21.58	0.32	78.41	1.18	48.04	0.72	76.01	1.14
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0.005	7.38	0.04	5.39	0.03	2.40	0.01	8.71	0.04	5.34	0.03	8.45	0.04
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.005	0.10	0.00	0.10	0.00	NC	-	0.10	0.00	0.71	0.00	NC	-
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.046	0.33	0.02	0.72	0.03	0.27	0.01	0.35	0.02	0.80	0.04	NC	-
PE 0112	Eggs, raw, (incl dried)	RAC	0.013	25.84	0.34	29.53	0.38	28.05	0.36	33.19	0.43	36.44	0.47	8.89	0.12
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-
Total intake (ug/person)=				105.4		109.8		194.0		125.8		102.2		103.4	
Bodyweight per region (kg bw) =				60		60		55		60		60		60	
ADI (ug/person)=				3000		3000		2750		3000		3000		3000	
%ADI=				3.5%		3.7%		7.1%		4.2%		3.4%		3.4%	
Rounded %ADI=				4%		4%		7%		4%		3%		3%	

SULFOXAFLOR (252)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.05 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day			Intake = daily intake: ug/person						
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
FC 0002	Subgroup of Lemons and limes, raw (incl lemon juice) (incl kumquat commodities)	RAC	0.038	18.97	0.72	0.97	0.04	6.23	0.24	0.09	0.00	3.35	0.13
FC 0003	Subgroup of Mandarins, raw (incl mandarin juice)	RAC	0.26	0.16	0.04	0.27	0.07	9.06	2.36	0.01	0.00	0.02	0.01
FC 0004	Subgroup of Oranges, sweet, sour, raw	RAC	0.26	1.18	0.31	1.11	0.29	14.28	3.71	0.05	0.01	1.08	0.28
JF 0004	Subgroup of Oranges, juice (single strength, incl. concentrated)	PP	0.036	0.08	0.00	0.26	0.01	12.61	0.45	0.14	0.01	0.33	0.01
FC 0005	Subgroup of Pummelo and grapefruits, raw (incl grapefruit juice)	RAC	0.0125	0.68	0.01	0.05	0.00	3.21	0.04	0.01	0.00	NC	-
FP 0009	Group of Pome fruits, raw (incl apple cider, excl apple juice)	RAC	0.067	68.85	4.61	10.93	0.73	70.82	4.74	189.78	12.72	19.56	1.31
JF 0226	Apple juice, single strength (incl. concentrated)	PP	0.067	0.03	0.00	0.10	0.01	7.19	0.48	0.03	0.00	NC	-
FS 0013	Subgroup of Cherries, raw	RAC	0.34	0.01	0.00	0.01	0.00	5.96	2.03	0.01	0.00	NC	-
FS 0014	Subgroup of Plums, raw (incl dried plums)	RAC	0.038	0.07	0.00	0.02	0.00	16.65	0.63	0.01	0.00	NC	-
FS 2001	Subgroup of peaches, raw (incl dried apricots)	RAC	0.061	0.02	0.00	0.01	0.00	10.76	0.66	0.01	0.00	NC	-
FB 2005	Subgroup of Caneberries, raw	RAC	0.44	0.01	0.00	7.30	3.21	2.29	1.01	0.01	0.00	NC	-
FB 2006	Subgroup of Bush berries, raw (including processed)	RAC	0.39	0.82	0.32	4.05	1.58	5.94	2.32	0.43	0.17	2.66	1.04
FB 0267	Elderberries, raw (incl processed)	RAC	0.39	0.71	0.28	3.52	1.37	NC	-	0.38	0.15	2.32	0.90
FB 0269	Grapes, raw (incl must, incl juice, excl dried, excl wine)	RAC	0.14	0.15	0.02	0.38	0.05	15.84	2.22	0.01	0.00	0.28	0.04
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	0.49	0.01	0.00	0.13	0.06	1.06	0.52	0.01	0.00	0.03	0.01
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.098	0.31	0.03	0.23	0.02	60.43	5.92	0.52	0.05	31.91	3.13
FB 0275	Strawberry, raw	RAC	0.19	0.01	0.00	0.01	0.00	3.35	0.64	0.01	0.00	0.01	0.00
FI 0326	Avocado, raw	RAC	0.011	1.12	0.01	0.01	0.00	0.84	0.01	0.01	0.00	6.60	0.07
FI 0345	Mango, raw (incl canned mango, incl mango juice)	RAC	0.022	12.25	0.27	6.83	0.15	0.76	0.02	0.01	0.00	20.12	0.44
VA 0381	Garlic, raw	RAC	0.01	0.82	0.01	2.06	0.02	3.79	0.04	0.03	0.00	0.29	0.00
-	Onions, dry, raw	RAC	0	9.01	0.00	20.24	0.00	30.90	0.00	9.61	0.00	2.11	0.00
-	Onions, green, raw	RAC	0.11	1.43	0.16	0.05	0.01	0.20	0.02	NC	-	6.30	0.69
VB 0400	Broccoli, raw	RAC	0.074	0.01	0.00	0.01	0.00	2.13	0.16	0.01	0.00	NC	-
VB 0404	Cauliflower, raw	RAC	0.012	0.01	0.00	0.01	0.00	2.73	0.03	0.01	0.00	NC	-
VB 0041	Cabbages, head, raw	RAC	0.099	3.82	0.38	2.99	0.30	49.16	4.87	0.01	0.00	NC	-

Annex 3

SULFOXAFLOL (252)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.05 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
VB 0467	Chinese cabbage, type pe-tsai, raw	RAC	1.2	0.62	0.74	0.49	0.59	NC	-	0.01	0.01	NC	-
VC 0045	Group of Fruiting vegetables, cucurbits, raw	RAC	0.029	5.96	0.17	9.74	0.28	51.82	1.50	13.61	0.39	0.05	0.00
VO 0448	Tomato, raw (incl canned, excl juice, excl paste)	RAC	0.11	13.10	1.44	4.90	0.54	62.16	6.84	1.04	0.11	0.09	0.01
-	Tomato, paste (i.e. concentrated tomato sauce/puree)	PP	0.23	0.58	0.13	0.22	0.05	2.21	0.51	0.24	0.06	3.10	0.71
JF 0448	Tomato, juice (single strength, incl concentrated)	PP	0.052	0.05	0.00	0.01	0.00	0.42	0.02	0.01	0.00	0.02	0.00
VO 0442	Okra, raw (i.e. Lady's Finger, Gombo)	RAC	0.11	6.23	0.69	0.10	0.01	NC	-	NC	-	NC	-
VO 0444	Peppers, chili, raw	RAC	0.11	3.47	0.38	3.56	0.39	16.30	1.79	0.01	0.00	NC	-
VO 0445	Peppers, sweet, raw (incl dried)	RAC	0.11	5.49	0.60	10.57	1.16	8.84	0.97	0.91	0.10	NC	-
VO 0440	Egg plant, raw (i.e. aubergine)	RAC	0.11	1.31	0.14	8.26	0.91	3.95	0.43	0.01	0.00	NC	-
VL 2050	Subgroup of Leafy greens	RAC	1.2	4.99	5.99	3.29	3.95	7.53	9.04	3.05	3.66	6.09	7.31
VL 0054	Subgroup of Leaves of Brassicaceae, raw	RAC	1.2	3.58	4.30	2.64	3.17	NC	-	1.83	2.20	3.65	4.38
VL 2052	Subgroup of Leaves of Root and Tuber Vegetables	RAC	1.2	0.30	0.36	0.22	0.26	NC	-	0.20	0.24	0.41	0.49
VL 2053	Subgroup of Leaves of trees, shrubs and vines	RAC	1.2	0.67	0.80	0.48	0.58	NC	-	0.45	0.54	0.90	1.08
VL 2054	Subgroup of Leafy aquatic vegetables	RAC	1.2	2.08	2.50	1.50	1.80	0.01	0.01	1.41	1.69	2.81	3.37
VD 0071	Beans, dry, raw (Phaseolus spp)	RAC	0.075	7.11	0.53	2.33	0.17	3.76	0.28	44.70	3.35	3.27	0.25
VD 0541	Soya bean, dry, raw (incl flour, incl paste, incl curd, incl sauce, excl oil)	RAC	0.011	2.89	0.03	0.21	0.00	0.48	0.01	3.16	0.03	0.26	0.00
OR 0541	Soya oil, refined	PP	0.0033	2.32	0.01	2.54	0.01	18.70	0.06	2.51	0.01	6.29	0.02
VR 0574	Beetroot, raw	RAC	0.01	5.86	0.06	4.23	0.04	9.46	0.09	3.96	0.04	7.91	0.08
VR 0575	Burdock, greater or edible, raw	RAC	0.01	0.06	0.00	0.04	0.00	NC	-	0.04	0.00	0.08	0.00
VR 0577	Carrots, raw	RAC	0.01	2.07	0.02	3.00	0.03	25.29	0.25	0.05	0.00	NC	-
VR 0578	Celeriac, raw	RAC	0.01	2.91	0.03	2.10	0.02	7.59	0.08	1.97	0.02	3.93	0.04
VR 0469	Chicory, roots, raw	RAC	0.01	0.01	0.00	0.03	0.00	0.10	0.00	NC	-	NC	-
VR 0583	Horseradish, raw	RAC	0.01	0.88	0.01	0.63	0.01	0.54	0.01	0.59	0.01	1.19	0.01
VR 0587	Parsley turnip-rooted, raw	RAC	0.01	0.55	0.01	0.40	0.00	4.29	0.04	0.37	0.00	0.74	0.01
VR 0588	Parsnip, raw	RAC	0.01	1.02	0.01	0.74	0.01	3.50	0.04	0.69	0.01	1.37	0.01
VR 0494	Radish roots, raw	RAC	0.01	3.96	0.04	2.86	0.03	3.30	0.03	2.67	0.03	5.34	0.05
VR 0591	Japanese radish, raw (i.e. Chinese radish, Daikon)	RAC	0.01	3.25	0.03	2.35	0.02	NC	-	2.20	0.02	4.39	0.04

SULFOXAFLOL (252)		International Estimated Daily Intake (IEDI)				ADI = 0 - 0.05 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
VR 0498	Salsify, raw (i.e. Oysterplant)	RAC	0.01	0.36	0.00	0.26	0.00	NC	-	0.24	0.00	0.48	0.00
VR 0596	Sugar beet, raw	RAC	0.014	0.01	0.00	NC	-	NC	-	NC	-	NC	-
-	Sugar beet, sugar	PP	0.025	0.56	0.01	0.24	0.01	NC	-	NC	-	5.13	0.13
VR 0497	Swede, raw (i.e. Rutabaga)	RAC	0.01	2.71	0.03	1.96	0.02	7.80	0.08	1.83	0.02	3.66	0.04
VR 0506	Turnip, garden, raw	RAC	0.01	4.29	0.04	3.10	0.03	6.41	0.06	2.90	0.03	5.79	0.06
VR 0573	Arrowroot, raw	RAC	0.01	13.83	0.14	18.24	0.18	0.01	0.00	0.05	0.00	19.60	0.20
VR 0463	Cassava raw (incl starch, incl tapioca, incl flour) (i.e. Manioc)	RAC	0.01	91.92	0.92	34.12	0.34	NC	-	259.92	2.60	45.48	0.45
VR 0585	Jerusalem artichoke, raw (i.e. Topinambur)	RAC	0.01	14.22	0.14	18.75	0.19	0.01	0.00	0.06	0.00	20.14	0.20
VR 0589	Potato, raw (incl flour, incl frozen, incl starch, incl tapioca)	RAC	0.01	23.96	0.24	13.56	0.14	213.41	2.13	104.35	1.04	8.56	0.09
VR 0508	Sweet potato, raw (incl dried)	RAC	0.01	28.83	0.29	61.55	0.62	0.15	0.00	221.94	2.22	NC	-
VR 0504	Tannia, raw (i.e. Tanier, Yautia)	RAC	0.01	NC	-	NC	-	0.01	0.00	NC	-	NC	-
VR 0505	Taro, raw (i.e. Dasheen, Eddoe)	RAC	0.01	6.71	0.07	31.91	0.32	NC	-	10.73	0.11	264.31	2.64
VR 0600	Yams, raw (incl dried)	RAC	0.01	70.93	0.71	30.62	0.31	0.07	0.00	5.65	0.06	30.85	0.31
VR 2072	Subgroup of Aquatic root and tuber vegetables	RAC	0.01	NC	-	NC	-	NC	-	NC	-	NC	-
VS 0624	Celery	RAC	0.19	3.66	0.70	2.65	0.50	4.84	0.92	2.47	0.47	4.94	0.94
VS 0621	Asparagus, raw	RAC	0.01	0.01	0.00	0.01	0.00	0.17	0.00	0.01	0.00	NC	-
GC 0653	Triticale, raw (incl flour)	RAC	0.025	0.01	0.00	NC	-	NC	-	NC	-	NC	-
GC 0654	Wheat, raw (incl bulgur, incl fermented beverages, incl germ, incl wholemeal bread, incl white flour products, incl white bread)	RAC	0.025	57.20	1.43	110.47	2.76	272.62	6.82	25.82	0.65	132.04	3.30
GC 0640	Barley, raw (incl malt extract, incl malt, excl pot&pearled, excl flour & grits, excl beer)	RAC	0.063	0.06	0.00	0.15	0.01	1.13	0.07	0.01	0.00	6.34	0.40
-	Barley, pot&pearled	PP	0.044	5.46	0.24	0.01	0.00	1.44	0.06	0.01	0.00	NC	-
-	Barley beer	PP	0.013	16.25	0.21	11.36	0.15	225.21	2.93	19.49	0.25	52.17	0.68
SO 0495	Rape seed, raw	RAC	0.045	NC	-	0.01	0.00	NC	-	NC	-	NC	-
OR 0495	Rape seed oil, edible	PP	0.014	0.07	0.00	0.03	0.00	4.62	0.06	0.03	0.00	NC	-
OR 0691	Cotton seed oil, edible	PP	0.002	1.28	0.00	0.05	0.00	0.45	0.00	0.42	0.00	0.15	0.00
SB 0716	Coffee bean, raw (i.e. green coffee)	RAC	0.035	0.83	0.03	0.69	0.02	1.09	0.04	2.91	0.10	0.82	0.03

Annex 3

SULFOXAFLOR (252)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.05 mg/kg bw						
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person								
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake	
SM 0716	Coffee bean, roasted	PP	0.0167	0.02	0.00	0.41	0.01	7.50	0.13	0.01	0.00	0.06	0.00	
-	Coffee bean, instant coffee (incl essences and concentrates)	PP	0.102	0.03	0.00	0.05	0.01	0.60	0.06	0.01	0.00	5.53	0.56	
HS 0444	Peppers, chili, dried	PP	1.1	0.58	0.64	1.27	1.40	1.21	1.33	0.12	0.13	NC	-	
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) -80% as muscle	RAC	0.045	23.34	1.05	40.71	1.83	97.15	4.37	18.06	0.81	57.71	2.60	
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat) - 20% as fat	RAC	0.03	5.84	0.18	10.18	0.31	24.29	0.73	4.52	0.14	14.43	0.43	
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.03	1.05	0.03	1.14	0.03	18.69	0.56	0.94	0.03	3.12	0.09	
MO 0105	Edible offal (mammalian), raw	RAC	0.13	4.64	0.60	1.97	0.26	10.01	1.30	3.27	0.43	3.98	0.52	
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.05	108.75	5.44	70.31	3.52	436.11	21.81	61.55	3.08	79.09	3.95	
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0.015	3.53	0.05	10.83	0.16	51.36	0.77	4.53	0.07	50.00	0.75	
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0.005	0.39	0.00	1.20	0.01	5.71	0.03	0.50	0.00	5.56	0.03	
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.005	NC	-	NC	-	0.32	0.00	NC	-	NC	-	
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.046	0.10	0.00	0.70	0.03	0.97	0.04	0.10	0.00	NC	-	
PE 0112	Eggs, raw, (incl dried)	RAC	0.013	3.84	0.05	4.41	0.06	27.25	0.35	1.13	0.01	7.39	0.10	
Total intake (ug/person)=					39.5		35.2		99.8		37.9		44.4	
Bodyweight per region (kg bw) =					60		60		60		60		60	
ADI (ug/person)=					3000		3000		3000		3000		3000	
%ADI=					1.3%		1.2%		3.3%		1.3%		1.5%	
Rounded %ADI=					1%		1%		3%		1%		1%	

TEBUCONAZOLE (184)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
FP 0226	Apple, raw	RAC	0.275	13.39	3.68	26.46	7.28	0.52	0.14	16.07	4.42	6.37	1.75	47.79	13.14
JF 0226	Apple juice, single strength (incl. concentrated)	PP	0.063	0.32	0.02	3.07	0.19	0.07	0.00	5.00	0.32	0.29	0.02	5.57	0.35
FP 0230	Pear, raw	RAC	0.275	2.16	0.59	6.24	1.72	0.05	0.01	4.07	1.12	1.16	0.32	5.34	1.47
FS 0013	Subgroup of Cherries, raw	RAC	0.86	0.92	0.79	9.15	7.87	0.01	0.01	0.61	0.52	0.06	0.05	6.64	5.71
FS 0014	Subgroup of Plums, raw	RAC	0.08	2.40	0.19	8.60	0.69	0.06	0.00	2.52	0.20	0.58	0.05	4.16	0.33
DF 0014	Plums, dried (prunes)	PP	0.232	0.09	0.02	0.06	0.01	0.01	0.00	0.18	0.04	0.04	0.01	0.06	0.01
FS 0240	Apricot, raw (incl dried)	RAC	0.46	5.15	2.37	3.66	1.68	0.03	0.01	2.25	1.04	0.17	0.08	6.80	3.13
-	Peaches and nectarines, raw	RAC	0.46	2.87	1.32	2.21	1.02	0.15	0.07	5.94	2.73	1.47	0.68	15.66	7.20
FB 0267	Elderberries, raw (incl processed)	RAC	0.345	0.44	0.15	0.27	0.09	0.34	0.12	1.41	0.49	NC	-	0.87	0.30
FB 0269	Grapes, raw (incl juice, excl must, excl dried, excl wine)	RAC	0.72	12.86	9.26	9.49	6.83	0.09	0.06	17.25	12.42	3.99	2.87	54.48	39.23
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	0.86	0.51	0.44	0.51	0.44	0.01	0.01	1.27	1.09	0.12	0.10	2.07	1.78
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.2	0.67	0.13	12.53	2.51	2.01	0.40	1.21	0.24	3.53	0.71	4.01	0.80
FT 0305	Table olives, raw	RAC	0	0.08	0.00	0.16	0.00	NC	-	0.24	0.00	0.01	0.00	0.65	0.00
FI 0327	Banana, raw (incl plantains) (incl dried)	RAC	0.01	5.23	0.05	6.94	0.07	99.45	0.99	32.47	0.32	48.30	0.48	24.70	0.25
FI 0345	Mango, raw (incl canned mango, incl mango juice)	RAC	0.05	10.48	0.52	0.01	0.00	7.24	0.36	6.87	0.34	19.98	1.00	6.25	0.31
FI 0350	Papaya, raw	RAC	0.18	0.35	0.06	0.01	0.00	3.05	0.55	0.80	0.14	7.28	1.31	1.00	0.18
FI 0351	Passion fruit, raw	RAC	0.1	0.58	0.06	0.01	0.00	0.59	0.06	0.60	0.06	0.18	0.02	0.08	0.01
VA 0381	Garlic, raw	RAC	0.02	2.29	0.05	5.78	0.12	0.11	0.00	3.69	0.07	1.65	0.03	3.91	0.08
-	Onions, dry, raw	RAC	0.055	29.36	1.61	37.50	2.06	3.56	0.20	34.78	1.91	18.81	1.03	43.38	2.39
VA 0384	Leek, raw	RAC	0.195	0.18	0.04	1.59	0.31	0.03	0.01	0.28	0.05	0.01	0.00	3.21	0.63
-	Onions, green, raw	RAC	0.1	2.45	0.25	1.49	0.15	1.02	0.10	2.60	0.26	0.60	0.06	2.03	0.20
VB 0400	Broccoli, raw	RAC	0.015	0.88	0.01	0.17	0.00	0.01	0.00	1.25	0.02	3.00	0.05	1.09	0.02
VB 0404	Cauliflower, raw	RAC	0.05	1.65	0.08	0.32	0.02	0.01	0.00	2.33	0.12	4.79	0.24	2.03	0.10
VB 0402	Brussels sprouts, raw	RAC	0.095	0.63	0.06	6.41	0.61	0.13	0.01	1.03	0.10	NC	-	2.35	0.22
VB 0041	Cabbages, head, raw	RAC	0.05	2.73	0.14	27.92	1.40	0.55	0.03	4.47	0.22	4.27	0.21	10.25	0.51
VC 0424	Cucumber, raw	RAC	0.05	8.01	0.40	30.66	1.53	1.45	0.07	19.84	0.99	0.27	0.01	34.92	1.75

Annex 3

TEBUCONAZOLE (184)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
VC 0431	Squash, Summer (Courgette, Marrow, Zucchini, Zucchini), raw	RAC	0.05	0.78	0.04	2.06	0.10	0.30	0.02	1.61	0.08	2.25	0.11	2.36	0.12
VC 0046	Melons, except watermelon, raw (Cantaloupe)	RAC	0.02	8.90	0.18	8.64	0.17	0.80	0.02	17.90	0.36	2.80	0.06	29.17	0.58
VO 0448	Tomato, raw (incl canned, excl juice, excl paste)	RAC	0.15	42.04	6.31	76.13	11.42	10.69	1.60	84.59	12.69	24.92	3.74	203.27	30.49
-	Tomato, paste (i.e. concentrated tomato sauce/puree)	PP	0.48	2.34	1.12	1.33	0.64	1.57	0.75	4.24	2.04	0.34	0.16	2.83	1.36
JF 0448	Tomato, juice (single strength, incl concentrated)	PP	0.082	0.29	0.02	0.29	0.02	0.01	0.00	0.38	0.03	0.05	0.00	0.14	0.01
VO 0445	Peppers, sweet, raw (incl dried)	RAC	0.185	4.49	0.83	6.44	1.19	7.21	1.33	5.68	1.05	9.52	1.76	8.92	1.65
VO 0440	Egg plant, raw (i.e. aubergine)	RAC	0.04	5.58	0.22	4.31	0.17	0.89	0.04	9.31	0.37	13.64	0.55	20.12	0.80
VP 2060	Subgroup of beans with pods (all commodities within this group)	RAC	0.315	0.68	0.21	NC	-	NC	-	0.39	0.12	0.22	0.07	0.49	0.15
VD 0071	Beans, dry, raw (Phaseolus spp)	RAC	0.05	2.39	0.12	1.61	0.08	10.47	0.52	1.84	0.09	12.90	0.65	7.44	0.37
VD 0541	Soya bean, dry, raw (incl flour, incl paste, incl curd, incl sauce, excl oil)	RAC	0.02	0.63	0.01	1.09	0.02	0.40	0.01	1.40	0.03	1.68	0.03	0.48	0.01
OR 0541	Soya oil, refined	PP	0.001	12.99	0.01	10.43	0.01	3.63	0.00	13.10	0.01	10.70	0.01	13.10	0.01
VR 0577	Carrots, raw	RAC	0.11	9.51	1.05	30.78	3.39	0.37	0.04	8.75	0.96	2.80	0.31	6.10	0.67
VS 0621	Asparagus, raw	RAC	0.02	0.01	0.00	0.01	0.00	0.01	0.00	0.03	0.00	0.07	0.00	0.21	0.00
VS 0620	Artichoke globe, raw	RAC	0.145	0.69	0.10	0.01	0.00	0.01	0.00	0.32	0.05	0.26	0.04	1.21	0.18
GC 0650	Rye, raw (incl flour)	RAC	0.05	0.13	0.01	19.38	0.97	0.10	0.01	0.12	0.01	0.03	0.00	2.15	0.11
GC 0653	Triticale, raw (incl flour)	RAC	0.05	NC	-	NC	-	NC	-	0.01	0.00	0.39	0.02	NC	-
GC 0654	Wheat, raw (incl bulgur, incl fermented beverages, incl germ, incl wholemeal bread, incl white flour products, incl white bread)	RAC	0.05	381.15	19.06	341.55	17.08	38.35	1.92	281.89	14.09	172.83	8.64	434.07	21.70
GC 0640	Barley, raw (incl malt extract, incl flour & grits, incl malt, excl pot&pearled, excl beer)	RAC	0.085	8.03	0.68	2.05	0.17	0.39	0.03	0.62	0.05	1.27	0.11	0.56	0.05
-	Barley, pot&pearled	PP	0.085	7.12	0.61	7.34	0.62	0.02	0.00	0.03	0.00	0.67	0.06	0.20	0.02
-	Barley beer	PP	0.002	4.87	0.01	93.78	0.19	24.28	0.05	12.76	0.03	39.28	0.08	18.15	0.04
GC 0647	Oats, raw	RAC	0.085	0.01	0.00	NC	-	0.01	0.00	0.45	0.04	0.01	0.00	0.01	0.00
020C	Rice cereals	-	0.085	-	-	-	-	-	-	-	-	-	-	-	-
CM 0649 (GC 0649)	Rice, husked, dry (incl polished, incl flour, incl starch, incl oil, incl beverages)	REP	0.275	45.40	12.49	14.99	4.12	84.88	23.34	111.73	30.73	194.75	53.56	93.12	25.61
GC 0447	Sweet corn on the cob, raw (incl frozen kernels, incl canned kernels)	RAC	0.06	0.14	0.01	0.94	0.06	5.70	0.34	2.61	0.16	1.94	0.12	0.22	0.01

TEBUCONAZOLE (184)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
TN 0085	Group of Tree nuts, raw (incl processed)	RAC	0	4.06	0.00	3.27	0.00	7.01	0.00	13.93	0.00	14.01	0.00	9.36	0.00
SO 0495	Rape seed, raw	RAC	0.1	0.02	0.00	NC	-	NC	-	0.01	0.00	0.75	0.08	0.01	0.00
OR 0495	Rape seed oil, edible	PP	0.11	0.35	0.04	0.44	0.05	0.19	0.02	0.97	0.11	3.28	0.36	0.77	0.08
SO 2091	Subgroup of Sunflower seeds, raw (incl processed)	RAC	0.04	7.43	0.30	36.06	1.44	1.15	0.05	8.77	0.35	5.74	0.23	13.63	0.55
SO 0691	Cotton seed, raw (incl oil)	RAC	0.05	20.53	1.03	9.80	0.49	6.42	0.32	4.73	0.24	7.14	0.36	18.68	0.93
SO 0697	Peanuts, nutmeat, raw (incl roasted, incl butter, excl oil)	RAC	0.035	0.46	0.02	1.21	0.04	6.64	0.23	2.71	0.09	1.26	0.04	1.84	0.06
OR 0697	Peanut oil, edible	PP	0	0.36	0.00	0.01	0.00	2.57	0.00	0.07	0.00	2.29	0.00	0.36	0.00
SB 0716	Coffee bean, raw (i.e. green coffee)	RAC	0.04	0.96	0.04	0.16	0.01	0.91	0.04	0.27	0.01	1.37	0.05	0.46	0.02
SM 0716	Coffee bean, roasted	PP	0.08	0.19	0.02	0.91	0.07	0.16	0.01	2.50	0.20	0.39	0.03	0.40	0.03
-	Coffee bean, instant coffee (incl essences and concentrates)	PP	0.032	0.07	0.00	0.94	0.03	0.07	0.00	0.70	0.02	0.07	0.00	0.29	0.01
HS 0444	Peppers, chili, dried	PP	1.85	0.42	0.78	0.53	0.98	0.84	1.55	0.50	0.93	0.95	1.76	0.37	0.68
DH 1100	Hops, dry	RAC	9.65	0.01	0.10	0.04	0.39	0.01	0.10	0.01	0.10	NC	-	0.01	0.10
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat)	RAC	0	31.20	0.00	72.44	0.00	20.88	0.00	47.98	0.00	33.08	0.00	36.25	0.00
MO 0105	Edible offal (mammalian), raw	RAC	0.06	4.79	0.29	9.68	0.58	2.97	0.18	5.49	0.33	3.84	0.23	5.03	0.30
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0	289.65	0.00	485.88	0.00	26.92	0.00	239.03	0.00	199.91	0.00	180.53	0.00
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0	13.17	0.00	26.78	0.00	7.24	0.00	116.71	0.00	22.54	0.00	32.09	0.00
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0	1.46	0.00	2.98	0.00	0.80	0.00	12.97	0.00	2.50	0.00	3.57	0.00
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.05	0.12	0.01	0.12	0.01	0.11	0.01	5.37	0.27	0.24	0.01	0.10	0.01
PE 0112	Eggs, raw, (incl dried)	RAC	0	7.84	0.00	23.08	0.00	2.88	0.00	14.89	0.00	9.81	0.00	14.83	0.00
Total intake (ug/person)=				68.0				81.1				35.8			
Bodyweight per region (kg bw) =				60				60				60			
ADI (ug/person)=				1800				1800				1800			
%ADI=				3.8%				4.5%				2.0%			
Rounded %ADI=				4%				5%				2%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			
												5%			

Annex 3

TEBUCONAZOLE (184)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
FP 0226	Apple, raw	RAC	0.275	27.44	7.55	49.21	13.53	21.57	5.93	31.09	8.55	51.60	14.19	1.77	0.49
JF 0226	Apple juice, single strength (incl. concentrated)	PP	0.063	14.88	0.94	11.98	0.75	0.15	0.01	9.98	0.63	30.32	1.91	3.47	0.22
FP 0230	Pear, raw	RAC	0.275	8.79	2.42	8.44	2.32	12.37	3.40	9.60	2.64	10.27	2.82	0.23	0.06
FS 0013	Subgroup of Cherries, raw	RAC	0.86	1.40	1.20	4.21	3.62	0.04	0.03	2.93	2.52	1.50	1.29	NC	-
FS 0014	Subgroup of Plums, raw	RAC	0.08	3.75	0.30	3.33	0.27	5.94	0.48	2.64	0.21	2.50	0.20	0.06	0.00
DF 0014	Plums, dried (prunes)	PP	0.232	0.61	0.14	0.35	0.08	0.05	0.01	0.35	0.08	0.49	0.11	0.13	0.03
FS 0240	Apricot, raw (incl dried)	RAC	0.46	4.27	1.96	3.31	1.52	0.06	0.03	2.86	1.32	1.71	0.79	NC	-
-	Peaches and nectarines, raw	RAC	0.46	8.76	4.03	12.98	5.97	8.23	3.79	10.09	4.64	3.64	1.67	0.04	0.02
FB 0267	Elderberries, raw (incl processed)	RAC	0.345	8.20	2.83	0.14	0.05	NC	-	NC	-	NC	-	1.87	0.65
FB 0269	Grapes, raw (incl juice, excl must, excl dried, excl wine)	RAC	0.72	7.03	5.06	13.65	9.83	5.23	3.77	12.15	8.75	7.35	5.29	1.21	0.87
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	0.86	3.09	2.66	1.51	1.30	0.03	0.03	1.38	1.19	4.26	3.66	0.42	0.36
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.2	88.93	17.79	62.41	12.48	1.84	0.37	25.07	5.01	61.17	12.23	5.84	1.17
FT 0305	Table olives, raw	RAC	0	0.04	0.00	0.01	0.00	0.01	0.00	0.03	0.00	0.01	0.00	NC	-
FI 0327	Banana, raw (incl plantains) (incl dried)	RAC	0.01	25.76	0.26	23.65	0.24	23.83	0.24	24.37	0.24	19.43	0.19	101.55	1.02
FI 0345	Mango, raw (incl canned mango, incl mango juice)	RAC	0.05	1.80	0.09	0.63	0.03	10.05	0.50	1.07	0.05	3.52	0.18	16.44	0.82
FI 0350	Papaya, raw	RAC	0.18	0.31	0.06	0.18	0.03	1.50	0.27	0.51	0.09	0.54	0.10	1.08	0.19
FI 0351	Passion fruit, raw	RAC	0.1	0.01	0.00	0.01	0.00	NC	-	NC	-	0.02	0.00	NC	-
VA 0381	Garlic, raw	RAC	0.02	0.98	0.02	1.49	0.03	12.88	0.26	3.74	0.07	2.05	0.04	1.14	0.02
-	Onions, dry, raw	RAC	0.055	19.69	1.08	29.83	1.64	24.64	1.36	31.35	1.72	9.72	0.53	12.59	0.69
VA 0384	Leek, raw	RAC	0.195	4.01	0.78	4.41	0.86	0.72	0.14	0.54	0.11	16.41	3.20	0.03	0.01
-	Onions, green, raw	RAC	0.1	1.55	0.16	0.74	0.07	1.05	0.11	3.74	0.37	0.94	0.09	6.45	0.65
VB 0400	Broccoli, raw	RAC	0.015	4.24	0.06	1.76	0.03	NC	-	0.51	0.01	3.79	0.06	0.26	0.00
VB 0404	Cauliflower, raw	RAC	0.05	5.27	0.26	5.01	0.25	NC	-	2.70	0.14	5.57	0.28	0.49	0.02
VB 0402	Brussels sprouts, raw	RAC	0.095	2.24	0.21	2.67	0.25	6.23	0.59	0.32	0.03	4.19	0.40	2.58	0.25
VB 0041	Cabbages, head, raw	RAC	0.05	8.97	0.45	27.12	1.36	1.44	0.07	24.96	1.25	4.55	0.23	11.23	0.56

TEBUCONAZOLE (184)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
VC 0424	Cucumber, raw	RAC	0.05	6.72	0.34	11.03	0.55	32.10	1.61	15.10	0.76	4.05	0.20	9.57	0.48
VC 0431	Squash, Summer (Courgette, Marrow, Zucchini, Zucchini), raw	RAC	0.05	NC	-	NC	-	5.48	0.27	NC	-	NC	-	1.03	0.05
VC 0046	Melons, except watermelon, raw (Cantaloupe)	RAC	0.02	9.20	0.18	11.95	0.24	14.63	0.29	8.99	0.18	7.86	0.16	2.46	0.05
VO 0448	Tomato, raw (incl canned, excl juice, excl paste)	RAC	0.15	43.88	6.58	55.41	8.31	35.38	5.31	74.88	11.23	26.50	3.98	9.51	1.43
-	Tomato, paste (i.e. concentrated tomato sauce/puree)	PP	0.48	4.96	2.38	3.20	1.54	0.15	0.07	1.61	0.77	6.88	3.30	0.52	0.25
JF 0448	Tomato, juice (single strength, incl concentrated)	PP	0.082	0.80	0.07	0.07	0.01	0.05	0.00	0.61	0.05	0.40	0.03	0.08	0.01
VO 0445	Peppers, sweet, raw (incl dried)	RAC	0.185	0.82	0.15	1.53	0.28	10.85	2.01	4.59	0.85	1.84	0.34	2.00	0.37
VO 0440	Egg plant, raw (i.e. aubergine)	RAC	0.04	1.01	0.04	1.69	0.07	21.37	0.85	3.00	0.12	1.40	0.06	NC	-
VP 2060	Subgroup of beans with pods (all commodities within this group)	RAC	0.315	5.07	1.60	0.83	0.26	0.17	0.05	3.70	1.17	NC	-	NC	-
VD 0071	Beans, dry, raw (Phaseolus spp)	RAC	0.05	1.51	0.08	1.50	0.08	1.90	0.10	5.11	0.26	1.36	0.07	23.43	1.17
VD 0541	Soya bean, dry, raw (incl flour, incl paste, incl curd, incl sauce, excl oil)	RAC	0.02	0.47	0.01	0.77	0.02	9.12	0.18	8.05	0.16	0.04	0.00	6.06	0.12
OR 0541	Soya oil, refined	PP	0.001	19.06	0.02	21.06	0.02	5.94	0.01	33.78	0.03	40.05	0.04	13.39	0.01
VR 0577	Carrots, raw	RAC	0.11	26.26	2.89	27.13	2.98	10.07	1.11	16.49	1.81	44.69	4.92	8.75	0.96
VS 0621	Asparagus, raw	RAC	0.02	0.84	0.02	2.08	0.04	7.11	0.14	1.01	0.02	1.69	0.03	0.04	0.00
VS 0620	Artichoke globe, raw	RAC	0.145	0.98	0.14	3.65	0.53	0.07	0.01	1.67	0.24	0.26	0.04	NC	-
GC 0650	Rye, raw (incl flour)	RAC	0.05	3.21	0.16	35.38	1.77	0.21	0.01	6.50	0.33	1.49	0.07	NC	-
GC 0653	Triticale, raw (incl flour)	RAC	0.05	0.01	0.00	0.17	0.01	0.29	0.01	0.01	0.00	NC	-	NC	-
GC 0654	Wheat, raw (incl bulgur, incl fermented beverages, incl germ, incl wholemeal bread, incl white flour products, incl white bread)	RAC	0.05	253.07	12.65	244.73	12.24	134.44	6.72	235.10	11.76	216.39	10.82	167.40	8.37
GC 0640	Barley, raw (incl malt extract, incl flour & grits, incl malt, excl pot&pearled, excl beer)	RAC	0.085	1.07	0.09	0.21	0.02	0.15	0.01	1.64	0.14	1.58	0.13	3.52	0.30
-	Barley, pot&pearled	PP	0.085	0.57	0.05	2.56	0.22	0.33	0.03	0.56	0.05	0.36	0.03	NC	-
-	Barley beer	PP	0.002	180.21	0.36	259.46	0.52	45.91	0.09	172.36	0.34	234.42	0.47	65.30	0.13
GC 0647	Oats, raw	RAC	0.085	NC	-	NC	-	0.01	0.00	0.01	0.00	NC	-	0.23	0.02
020C	Rice cereals	-	0.085	-	-	-	-	-	-	-	-	-	-	-	-
CM 0649 (GC	Rice, husked, dry (incl polished, incl flour, incl starch, incl oil, incl beverages)	REP	0.275	20.96	5.76	16.04	4.41	339.67	93.41	75.51	20.77	16.86	4.64	86.13	23.69

Annex 3

TEBUCONAZOLE (184)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw								
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day										
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake	
0649)																
GC 0447	Sweet corn on the cob, raw (incl frozen kernels, incl canned kernels)	RAC	0.06	11.43	0.69	3.71	0.22	0.74	0.04	13.63	0.82	3.07	0.18	1.50	0.09	
TN 0085	Group of Tree nuts, raw (incl processed)	RAC	0	8.52	0.00	8.94	0.00	15.09	0.00	9.60	0.00	14.57	0.00	26.26	0.00	
SO 0495	Rape seed, raw	RAC	0.1	NC	-	NC	-	0.01	0.00	NC	-	NC	-	NC	-	
OR 0495	Rape seed oil, edible	PP	0.11	12.52	1.38	7.63	0.84	3.00	0.33	6.01	0.66	NC	-	NC	-	
SO 2091	Subgroup of Sunflower seeds, raw (incl processed)	RAC	0.04	23.43	0.94	29.34	1.17	1.24	0.05	14.00	0.56	6.48	0.26	6.91	0.28	
SO 0691	Cotton seed, raw (incl oil)	RAC	0.05	10.71	0.54	4.23	0.21	7.19	0.36	7.54	0.38	5.66	0.28	2.38	0.12	
SO 0697	Peanuts, nutmeat, raw (incl roasted, incl butter, excl oil)	RAC	0.035	3.26	0.11	2.22	0.08	5.38	0.19	4.85	0.17	1.54	0.05	1.82	0.06	
OR 0697	Peanut oil, edible	PP	0	1.02	0.00	0.23	0.00	1.81	0.00	0.42	0.00	5.23	0.00	0.01	0.00	
SB 0716	Coffee bean, raw (i.e. green coffee)	RAC	0.04	0.60	0.02	NC	-	0.62	0.02	1.71	0.07	NC	-	3.51	0.14	
SM 0716	Coffee bean, roasted	PP	0.08	7.02	0.56	9.75	0.78	0.02	0.00	5.09	0.41	13.38	1.07	0.77	0.06	
-	Coffee bean, instant coffee (incl essences and concentrates)	PP	0.032	0.75	0.02	0.30	0.01	0.04	0.00	0.67	0.02	2.43	0.08	1.43	0.05	
HS 0444	Peppers, chili, dried	PP	1.85	0.11	0.20	0.21	0.39	0.36	0.67	0.21	0.39	0.25	0.46	0.15	0.28	
DH 1100	Hops, dry	RAC	9.65	NC	-	NC	-	0.02	0.19	0.02	0.19	NC	-	NC	-	
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat)	RAC	0	140.03	0.00	150.89	0.00	79.32	0.00	111.24	0.00	120.30	0.00	51.27	0.00	
MO 0105	Edible offal (mammalian), raw	RAC	0.06	15.17	0.91	5.19	0.31	6.30	0.38	6.78	0.41	3.32	0.20	3.17	0.19	
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0	388.92	0.00	335.88	0.00	49.15	0.00	331.25	0.00	468.56	0.00	245.45	0.00	
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0	66.38	0.00	48.47	0.00	21.58	0.00	78.41	0.00	48.04	0.00	76.01	0.00	
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0	7.38	0.00	5.39	0.00	2.40	0.00	8.71	0.00	5.34	0.00	8.45	0.00	
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.05	0.33	0.02	0.72	0.04	0.27	0.01	0.35	0.02	0.80	0.04	NC	-	
PE 0112	Eggs, raw, (incl dried)	RAC	0	25.84	0.00	29.53	0.00	28.05	0.00	33.19	0.00	36.44	0.00	8.89	0.00	
Total intake (ug/person)=					89.3		94.7		135.9		94.7		81.4		46.8	
Bodyweight per region (kg bw) =					60		60		55		60		60		60	
ADI (ug/person)=					1800		1800		1650		1800		1800		1800	
%ADI=					5.0%		5.3%		8.2%		5.3%		4.5%		2.6%	

TEBUCONAZOLE (184)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw								
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day			Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake	
Rounded %ADI=				5%			5%		8%		5%		5%		3%	

TEBUCONAZOLE (184)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw					
Codex Code		Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person						
					G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet
FP 0226	Apple, raw	RAC	0.275	0.21	0.06	2.05	0.56	54.48	14.98	0.10	0.03	1.38	0.38
JF 0226	Apple juice, single strength (incl. concentrated)	PP	0.063	0.03	0.00	0.10	0.01	7.19	0.45	0.03	0.00	NC	-
FP 0230	Pear, raw	RAC	0.275	0.07	0.02	0.14	0.04	9.45	2.60	0.01	0.00	0.14	0.04
FS 0013	Subgroup of Cherries, raw	RAC	0.86	0.01	0.01	0.01	0.01	5.96	5.13	0.01	0.01	NC	-
FS 0014	Subgroup of Plums, raw	RAC	0.08	0.07	0.01	0.01	0.00	15.56	1.24	0.01	0.00	NC	-
DF 0014	Plums, dried (prunes)	PP	0.232	0.01	0.00	0.01	0.00	0.37	0.09	0.01	0.00	NC	-
FS 0240	Apricot, raw (incl dried)	RAC	0.46	0.01	0.00	0.01	0.00	3.29	1.51	0.01	0.00	NC	-
-	Peaches and nectarines, raw	RAC	0.46	0.02	0.01	0.01	0.00	7.47	3.44	0.01	0.00	NC	-
FB 0267	Elderberries, raw (incl processed)	RAC	0.345	0.71	0.24	3.52	1.21	NC	-	0.38	0.13	2.32	0.80
FB 0269	Grapes, raw (incl juice, excl must, excl dried, excl wine)	RAC	0.72	0.15	0.11	0.38	0.27	15.73	11.33	0.01	0.01	0.09	0.06
DF 0269	Grapes, dried (= currants, raisins and sultanas) (from table-grapes)	PP	0.86	0.01	0.01	0.13	0.11	1.06	0.91	0.01	0.01	0.03	0.03
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.2	0.31	0.06	0.23	0.05	60.43	12.09	0.52	0.10	31.91	6.38
FT 0305	Table olives, raw	RAC	0	NC	-	NC	-	0.01	0.00	NC	-	NC	-
FI 0327	Banana, raw (incl plantains) (incl dried)	RAC	0.01	44.80	0.45	118.17	1.18	25.25	0.25	454.49	4.54	310.23	3.10
FI 0345	Mango, raw (incl canned mango, incl mango juice)	RAC	0.05	12.25	0.61	6.83	0.34	0.76	0.04	0.01	0.00	20.12	1.01
FI 0350	Papaya, raw	RAC	0.18	6.47	1.16	0.25	0.05	0.19	0.03	0.01	0.00	26.42	4.76
FI 0351	Passion fruit, raw	RAC	0.1	0.12	0.01	0.10	0.01	0.01	0.00	0.18	0.02	3.81	0.38
VA 0381	Garlic, raw	RAC	0.02	0.82	0.02	2.06	0.04	3.79	0.08	0.03	0.00	0.29	0.01
-	Onions, dry, raw	RAC	0.055	9.01	0.50	20.24	1.11	30.90	1.70	9.61	0.53	2.11	0.12
VA 0384	Leek, raw	RAC	0.195	0.02	0.00	1.44	0.28	1.22	0.24	0.01	0.00	NC	-

Annex 3

TEBUCONAZOLE (184)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
-	Onions, green, raw	RAC	0.1	1.43	0.14	0.05	0.01	0.20	0.02	NC	-	6.30	0.63
VB 0400	Broccoli, raw	RAC	0.015	0.01	0.00	0.01	0.00	2.13	0.03	0.01	0.00	NC	-
VB 0404	Cauliflower, raw	RAC	0.05	0.01	0.00	0.01	0.00	2.73	0.14	0.01	0.00	NC	-
VB 0402	Brussels sprouts, raw	RAC	0.095	0.88	0.08	0.69	0.07	2.89	0.27	0.01	0.00	NC	-
VB 0041	Cabbages, head, raw	RAC	0.05	3.82	0.19	2.99	0.15	49.16	2.46	0.01	0.00	NC	-
VC 0424	Cucumber, raw	RAC	0.05	0.68	0.03	1.81	0.09	10.40	0.52	0.01	0.00	0.04	0.00
VC 0431	Squash, Summer (Courgette, Marrow, Zucchini, Zucchini), raw	RAC	0.05	0.09	0.00	1.01	0.05	NC	-	1.91	0.10	NC	-
VC 0046	Melons, except watermelon, raw (Cantaloupe)	RAC	0.02	0.19	0.00	0.10	0.00	4.98	0.10	0.01	0.00	NC	-
VO 0448	Tomato, raw (incl canned, excl juice, excl paste)	RAC	0.15	13.10	1.97	4.90	0.74	62.16	9.32	1.04	0.16	0.09	0.01
-	Tomato, paste (i.e. concentrated tomato sauce/puree)	PP	0.48	0.58	0.28	0.22	0.11	2.21	1.06	0.24	0.12	3.10	1.49
JF 0448	Tomato, juice (single strength, incl concentrated)	PP	0.082	0.05	0.00	0.01	0.00	0.42	0.03	0.01	0.00	0.02	0.00
VO 0445	Peppers, sweet, raw (incl dried)	RAC	0.185	5.49	1.02	10.57	1.96	8.84	1.64	0.91	0.17	NC	-
VO 0440	Egg plant, raw (i.e. aubergine)	RAC	0.04	1.31	0.05	8.26	0.33	3.95	0.16	0.01	0.00	NC	-
VP 2060	Subgroup of beans with pods (all commodities within this group)	RAC	0.315	NC	-	NC	-	NC	-	NC	-	NC	-
VD 0071	Beans, dry, raw (Phaseolus spp)	RAC	0.05	7.11	0.36	2.33	0.12	3.76	0.19	44.70	2.24	3.27	0.16
VD 0541	Soya bean, dry, raw (incl flour, incl paste, incl curd, incl sauce, excl oil)	RAC	0.02	2.89	0.06	0.21	0.00	0.48	0.01	3.16	0.06	0.26	0.01
OR 0541	Soya oil, refined	PP	0.001	2.32	0.00	2.54	0.00	18.70	0.02	2.51	0.00	6.29	0.01
VR 0577	Carrots, raw	RAC	0.11	2.07	0.23	3.00	0.33	25.29	2.78	0.05	0.01	NC	-
VS 0621	Asparagus, raw	RAC	0.02	0.01	0.00	0.01	0.00	0.17	0.00	0.01	0.00	NC	-
VS 0620	Artichoke globe, raw	RAC	0.145	0.01	0.00	NC	-	0.08	0.01	0.01	0.00	NC	-
GC 0650	Rye, raw (incl flour)	RAC	0.05	0.03	0.00	0.01	0.00	13.95	0.70	0.01	0.00	0.88	0.04
GC 0653	Triticale, raw (incl flour)	RAC	0.05	0.01	0.00	NC	-	NC	-	NC	-	NC	-
GC 0654	Wheat, raw (incl bulgur, incl fermented beverages, incl germ, incl wholemeal bread, incl white flour products, incl white bread)	RAC	0.05	57.20	2.86	110.47	5.52	272.62	13.63	25.82	1.29	132.04	6.60
GC 0640	Barley, raw (incl malt extract, incl flour & grits, incl malt, excl pot&pearled, excl beer)	RAC	0.085	0.10	0.01	0.15	0.01	1.71	0.15	0.02	0.00	6.34	0.54
-	Barley, pot&pearled	PP	0.085	5.46	0.46	0.01	0.00	1.44	0.12	0.01	0.00	NC	-

TEBUCONAZOLE (184)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
-	Barley beer	PP	0.002	16.25	0.03	11.36	0.02	225.21	0.45	19.49	0.04	52.17	0.10
GC 0647	Oats, raw	RAC	0.085	0.01	0.00	0.02	0.00	NC	-	0.09	0.01	NC	-
020C	Rice cereals	-	0.085	-	-	-	-	-	-	-	-	-	-
CM 0649 (GC 0649)	Rice, husked, dry (incl polished, incl flour, incl starch, incl oil, incl beverages)	REP	0.275	52.55	14.45	286.02	78.66	18.64	5.13	19.67	5.41	75.09	20.65
GC 0447	Sweet corn on the cob, raw (incl frozen kernels, incl canned kernels)	RAC	0.06	3.63	0.22	20.50	1.23	8.78	0.53	0.02	0.00	0.17	0.01
TN 0085	Group of Tree nuts, raw (incl processed)	RAC	0	4.39	0.00	135.53	0.00	6.11	0.00	0.72	0.00	317.74	0.00
SO 0495	Rape seed, raw	RAC	0.1	NC	-	0.01	0.00	NC	-	NC	-	NC	-
OR 0495	Rape seed oil, edible	PP	0.11	0.07	0.01	0.03	0.00	4.62	0.51	0.03	0.00	NC	-
SO 2091	Subgroup of Sunflower seeds, raw (incl processed)	RAC	0.04	0.99	0.04	0.22	0.01	32.01	1.28	12.12	0.48	0.48	0.02
SO 0691	Cotton seed, raw (incl oil)	RAC	0.05	8.14	0.41	0.32	0.02	2.84	0.14	2.69	0.13	0.97	0.05
SO 0697	Peanuts, nutmeat, raw (incl roasted, incl butter, excl oil)	RAC	0.035	7.14	0.25	0.45	0.02	1.87	0.07	6.22	0.22	0.53	0.02
OR 0697	Peanut oil, edible	PP	0	5.02	0.00	0.05	0.00	0.17	0.00	0.29	0.00	NC	-
SB 0716	Coffee bean, raw (i.e. green coffee)	RAC	0.04	0.83	0.03	0.69	0.03	1.09	0.04	2.91	0.12	0.82	0.03
SM 0716	Coffee bean, roasted	PP	0.08	0.02	0.00	0.41	0.03	7.50	0.60	0.01	0.00	0.06	0.00
-	Coffee bean, instant coffee (incl essences and concentrates)	PP	0.032	0.03	0.00	0.05	0.00	0.60	0.02	0.01	0.00	5.53	0.18
HS 0444	Peppers, chili, dried	PP	1.85	0.58	1.07	1.27	2.35	1.21	2.24	0.12	0.22	NC	-
DH 1100	Hops, dry	RAC	9.65	NC	-	NC	-	0.04	0.39	NC	-	NC	-
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat)	RAC	0	29.18	0.00	50.89	0.00	121.44	0.00	22.58	0.00	72.14	0.00
MO 0105	Edible offal (mammalian), raw	RAC	0.06	4.64	0.28	1.97	0.12	10.01	0.60	3.27	0.20	3.98	0.24
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0	108.75	0.00	70.31	0.00	436.11	0.00	61.55	0.00	79.09	0.00
PM 0110	Poultry meat, raw (incl prepared) - 90% as muscle	RAC	0	3.53	0.00	10.83	0.00	51.36	0.00	4.53	0.00	50.00	0.00
PM 0110	Poultry meat, raw (incl prepared) - 10% as fat	RAC	0	0.39	0.00	1.20	0.00	5.71	0.00	0.50	0.00	5.56	0.00
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.05	0.10	0.01	0.70	0.04	0.97	0.05	0.10	0.01	NC	-
PE 0112	Eggs, raw, (incl dried)	RAC	0	3.84	0.00	4.41	0.00	27.25	0.00	1.13	0.00	7.39	0.00
-	-	-		-	-	-	-	-	-	-	-	-	-
Total intake (ug/person)=				27.8		97.3		101.5		16.4		47.9	

Annex 3

TEBUCONAZOLE (184)			International Estimated Daily Intake (IEDI)				ADI = 0 - 0.03 mg/kg bw						
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
	Bodyweight per region (kg bw) =				60		60		60		60		60
	ADI (ug/person)=				1800		1800		1800		1800		1800
	%ADI=				1.5%		5.4%		5.6%		0.9%		2.7%
	Rounded %ADI=				2%		5%		6%		1%		3%

THIAMETHOXAM (245)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.08 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day			Intake as ug/person/day								
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
FC 0002	Subgroup of Lemons and limes, raw (incl lemon juice) (incl kumquat commodities)	RAC	0.028	4.82	0.13	2.45	0.07	3.93	0.11	25.44	0.71	8.74	0.24	16.23	0.45
FC 0003	Subgroup of Mandarins, raw (incl mandarin juice)	RAC	0.028	6.18	0.17	3.66	0.10	0.25	0.01	6.82	0.19	3.49	0.10	19.38	0.54
FC 0004	Subgroup of Oranges, sweet, sour, raw	RAC	0.028	20.66	0.58	5.23	0.15	11.90	0.33	37.90	1.06	21.16	0.59	56.46	1.58
JF 0004	Subgroup of Oranges, juice (single strength, incl. concentrated)	PP	0.007	1.27	0.01	2.20	0.02	0.09	0.00	11.81	0.08	0.46	0.00	1.69	0.01
FC 0005	Subgroup of Pummelo and grapefruits, raw (incl grapefruit juice)	RAC	0.028	0.66	0.02	0.69	0.02	0.96	0.03	10.20	0.29	1.25	0.04	2.97	0.08
FP 0226	Apple, raw (incl cider, excl juice)	RAC	0.07	13.49	0.94	26.63	1.86	15.05	1.05	16.28	1.14	6.47	0.45	47.88	3.35
JF 0226	Apple juice, single strength (incl. concentrated)	PP	0.065	0.32	0.02	3.07	0.20	0.07	0.00	5.00	0.33	0.29	0.02	5.57	0.36
FP 0228	Loquat, raw (incl processed) (i.e. Japanese medlar)	RAC	0.07	0.59	0.04	0.36	0.03	0.46	0.03	1.88	0.13	NC	-	1.15	0.08
FP 0229	Medlar, raw (incl processed)	RAC	0.07	0.47	0.03	0.29	0.02	0.36	0.03	1.49	0.10	NC	-	0.92	0.06
FP 0230	Pear, raw	RAC	0.07	2.16	0.15	6.24	0.44	0.05	0.00	4.07	0.28	1.16	0.08	5.34	0.37
FP 0307	Persimmon, Japanese, raw (i.e. Kaki fruit)	RAC	0.165	1.91	0.32	0.01	0.00	1.94	0.32	1.96	0.32	NC	-	0.25	0.04
FP 0231	Quince, raw	RAC	0.07	0.73	0.05	0.54	0.04	0.01	0.00	0.07	0.00	0.06	0.00	1.31	0.09
FS 0012	Group of Stone fruits, raw (incl dried apricots, excl dried plums)	RAC	0.195	11.33	2.21	23.62	4.61	0.24	0.05	11.32	2.21	2.28	0.44	33.26	6.49
DF 0014	Plums, dried (prunes)	PP	0.16	0.09	0.01	0.06	0.01	0.01	0.00	0.18	0.03	0.04	0.01	0.06	0.01
FB 2005	Subgroup of Caneberries, raw	RAC	0.055	0.42	0.02	1.05	0.06	0.01	0.00	0.02	0.00	0.02	0.00	1.24	0.07
FB 2006	Subgroup of Bush berries, raw (including processed)	RAC	0.055	0.53	0.03	1.31	0.07	0.40	0.02	1.66	0.09	0.01	0.00	0.99	0.05
FB 2007	Subgroup of Large shrub/tree berries, raw (including processed)	RAC	0.055	0.62	0.03	0.33	0.02	0.34	0.02	1.42	0.08	0.01	0.00	1.51	0.08
FB 0269	Grapes, raw (incl must, incl dried, incl juice, excl wine)	RAC	0.055	15.33	0.84	11.75	0.65	0.11	0.01	22.55	1.24	4.49	0.25	63.13	3.47
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.055	0.67	0.04	12.53	0.69	2.01	0.11	1.21	0.07	3.53	0.19	4.01	0.22
FB 2009	Subgroup of Low growing berries, raw	RAC	0.055	0.71	0.04	2.02	0.11	0.04	0.00	1.39	0.08	0.37	0.02	2.53	0.14
FI 0326	Avocado, raw	RAC	0.08	0.13	0.01	0.03	0.00	2.05	0.16	2.54	0.20	2.34	0.19	0.12	0.01
FI 0327	Banana, raw (incl plantains) (incl dried)	RAC	0.02	5.23	0.10	6.94	0.14	99.45	1.99	32.47	0.65	48.30	0.97	24.70	0.49
FI 0345	Mango, raw (incl canned mango, incl mango juice)	RAC	0.03	10.48	0.31	0.01	0.00	7.24	0.22	6.87	0.21	19.98	0.60	6.25	0.19
FI 0353	Pineapple, raw (incl canned pineapple, incl pineapple juice, incl dried pineapple)	RAC	0.01	0.61	0.01	1.56	0.02	7.89	0.08	9.36	0.09	8.76	0.09	1.30	0.01

Annex 3

THIAMETHOXAM (245)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.08 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day			Intake as ug/person/day								
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
VB 0040	Group of Brassica vegetables (excl Brassica leafy vegetables), raw	RAC	0.53	6.43	3.41	40.26	21.34	0.80	0.42	9.94	5.27	12.07	6.40	17.73	9.40
VC 0045	Group of Fruiting vegetables, cucurbits, raw	RAC	0.105	53.14	5.58	86.21	9.05	6.28	0.66	92.76	9.74	15.64	1.64	155.30	16.31
VO 0448	Tomato, raw (incl canned, excl juice, excl paste)	RAC	0.08	42.04	3.36	76.13	6.09	10.69	0.86	84.59	6.77	24.92	1.99	203.27	16.26
-	Tomato, paste (i.e. concentrated tomato sauce/puree)	PP	0.24	2.34	0.56	1.33	0.32	1.57	0.38	4.24	1.02	0.34	0.08	2.83	0.68
JF 0448	Tomato, juice (single strength, incl concentrated)	PP	0.054	0.29	0.02	0.29	0.02	0.01	0.00	0.38	0.02	0.05	0.00	0.14	0.01
VO 0442	Okra, raw (i.e. Lady's Finger, Gombo)	RAC	0.08	1.97	0.16	NC	-	3.68	0.29	3.24	0.26	5.72	0.46	1.57	0.13
VO 0444	Peppers, chili, raw	RAC	0.08	3.99	0.32	7.30	0.58	2.93	0.23	5.62	0.45	NC	-	17.44	1.40
VO 0445	Peppers, sweet, raw (incl dried)	RAC	0.08	4.49	0.36	6.44	0.52	7.21	0.58	5.68	0.45	9.52	0.76	8.92	0.71
VO 0440	Egg plant, raw (i.e. aubergine)	RAC	0.08	5.58	0.45	4.31	0.34	0.89	0.07	9.31	0.74	13.64	1.09	20.12	1.61
VL 2050	Subgroup of Leafy greens	RAC	0.54	3.93	2.12	5.28	2.85	3.07	1.66	14.53	7.85	8.25	4.46	12.75	6.89
VP 0061	Beans with pods (Phaseolus spp): (immature pods + succulent seeds)	RAC	0.08	0.68	0.05	NC	-	NC	-	0.39	0.03	0.22	0.02	0.49	0.04
014B	Peas with pods	-	0.01	-	-	-	-	-	-	-	-	-	-	-	-
VP 0062	Beans without pods: (Phaseolus spp.) (succulent seeds), raw	RAC	0.01	1.56	0.02	0.60	0.01	0.49	0.00	1.18	0.01	0.90	0.01	7.79	0.08
VP 0523	Broad bean without pods (succulent seeds) (Vicia spp), raw	RAC	0.01	3.51	0.04	0.43	0.00	0.01	0.00	0.60	0.01	0.29	0.00	0.78	0.01
VP 0064	Peas without pods (Pisum spp) (succulent seeds)	RAC	0.01	1.97	0.02	0.51	0.01	0.02	0.00	0.79	0.01	3.68	0.04	3.80	0.04
VD 0070	Group of Pulses, raw (incl processed)	RAC	0.02	87.29	1.75	64.04	1.28	37.15	0.74	89.82	1.80	91.02	1.82	98.20	1.96
VD 0520	Bambara groundnut (dry) (Vigna subterranea), raw	RAC	0.01	NC	-	NC	-	0.20	0.00	NC	-	NC	-	NC	-
VR 0075	Group of Root and tuber vegetables, raw (incl processed)	RAC	0.01	87.83	0.88	374.04	3.74	668.92	6.69	121.64	1.22	94.20	0.94	247.11	2.47
VS 0620	Artichoke globe, raw	RAC	0.23	0.69	0.16	0.01	0.00	0.01	0.00	0.32	0.07	0.26	0.06	1.21	0.28
VF 2084	Group of edible fungi (cultivated & wild), raw (incl canned, incl dried)	RAC	0.08	0.09	0.01	0.56	0.04	0.02	0.00	2.65	0.21	0.11	0.01	0.51	0.04
GC 0653	Triticale, raw	RAC	0.032	NC	-	NC	-	NC	-	0.01	0.00	NC	-	NC	-
GC 0654	Wheat, raw (incl meslin)	RAC	0.032	0.01	0.00	1.12	0.04	NC	-	0.01	0.00	0.56	0.02	NC	-
-	Wheat, bulgur	PP	0.14	NC	-	NC	-	NC	-	0.03	0.00	NC	-	NC	-
CF 1210	Wheat, germ	PP	0.0493	NC	-	NC	-	0.01	0.00	0.01	0.00	0.14	0.01	0.01	0.00
CP 1212	Wheat, wholemeal bread	PP	0.014	0.01	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.03	0.00	0.01	0.00

THIAMETHOXAM (245)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.08 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
CP 1211	Wheat, white bread	PP	0.014	0.25	0.00	0.63	0.01	0.12	0.00	0.43	0.01	1.39	0.02	0.22	0.00
-	Wheat, Fermented Beverages (Korean jakju and takju)	PP	0.02	NC	-	NC	-	NC	-	NC	-	NC	-	NC	-
CF 1211	Wheat, white flour (incl white flour products: starch, gluten, macaroni, pastry)	PP	0.00336	301.24	1.01	268.64	0.90	30.21	0.10	222.51	0.75	134.73	0.45	343.12	1.15
GC 0640	Barley, raw (incl malt extract, incl pot&pearled, incl beer, incl malt, excl flour & grits)	RAC	0.112	14.50	1.62	30.61	3.43	5.01	0.56	3.08	0.34	8.88	0.99	4.29	0.48
-	Barley, pot&pearled	PP	0.03	7.12	0.21	7.34	0.22	0.02	0.00	0.03	0.00	0.67	0.02	0.20	0.01
-	Barley, flour (white flour and wholemeal flour)	PP	0.01	2.93	0.03	0.30	0.00	0.02	0.00	0.01	0.00	0.48	0.00	0.01	0.00
GC 0647	Oats, raw (incl rolled)	RAC	0.112	0.05	0.01	7.05	0.79	0.10	0.01	1.71	0.19	0.96	0.11	0.04	0.00
CM 0649 (GC 0649)	Rice, husked, dry (incl flour, incl oil, incl beverages, incl starch, excl polished)	REP	1.7	1.26	2.14	1.58	2.69	31.05	52.79	5.43	9.23	0.90	1.53	2.18	3.71
CM 1205	Rice polished, dry	PP	0.704	34.21	24.08	10.39	7.31	41.72	29.37	82.38	58.00	150.24	105.77	70.47	49.61
GC 0651	Sorghum, raw (incl beer, excl flour)(i.e. Chicken corn, Dari seed, Durra, Feterita)	RAC	0.079	NC	-	0.01	0.00	3.34	0.26	0.01	0.00	NC	-	NC	-
-	Sorghum, flour (white flour and wholemeal flour)	PP	0.0548	3.91	0.21	NC	-	11.62	0.64	14.24	0.78	9.87	0.54	2.62	0.14
GC 0645	Maize, raw (incl glucose & dextrose & isoglucose, incl flour, incl oil, incl beer, incl germ, incl starch)	RAC	0.02	29.81	0.60	44.77	0.90	108.95	2.18	52.37	1.05	60.28	1.21	75.69	1.51
GC 0447	Sweet corn on the cob, raw (incl frozen kernels, incl canned kernels)	RAC	0.01	0.14	0.00	0.94	0.01	5.70	0.06	2.61	0.03	1.94	0.02	0.22	0.00
GC 0447	Sweet corn on the cob, raw (i.e kernels plus cob without husks)	RAC	0.01	0.08	0.00	0.09	0.00	5.67	0.06	0.05	0.00	1.77	0.02	NC	-
-	Sugar crops NES, raw (incl sugar, syrup and others): sugar maple, sweet sorghum, sugar palm	RAC	0.033	1.30	0.04	2.72	0.09	0.92	0.03	3.26	0.11	0.71	0.02	0.90	0.03
TN 0672	Pecan, nutmeat	RAC	0.01	0.05	0.00	0.05	0.00	0.02	0.00	0.14	0.00	0.09	0.00	0.13	0.00
SO 0091	Subgroup of Oilseeds, raw	RAC	0.02	1.34	0.03	1.35	0.03	8.56	0.17	3.76	0.08	2.22	0.04	3.38	0.07
SO 0090	Subgroup of Mustard seeds, raw (incl flour, incl oil)	RAC	0.02	0.02	0.00	0.05	0.00	0.01	0.00	0.31	0.01	0.03	0.00	0.04	0.00
SO 0693	Linseed, raw (incl oil)	RAC	0.02	0.02	0.00	NC	-	NC	-	0.01	0.00	0.13	0.00	NC	-
SO 0698	Poppy seed, raw (incl oil)	RAC	0.02	0.01	0.00	0.01	0.00	0.01	0.00	0.03	0.00	0.01	0.00	0.01	0.00
SO 0495	Rape seed, raw (incl oil)	RAC	0.02	0.93	0.02	1.16	0.02	0.49	0.01	2.53	0.05	9.32	0.19	2.02	0.04
SO 0700	Sesame seed, raw (incl oil)	RAC	0.02	1.22	0.02	0.01	0.00	0.54	0.01	4.23	0.08	0.82	0.02	2.77	0.06
SO 0699	Safflower seed, raw (incl oil)	RAC	0.02	0.03	0.00	0.20	0.00	0.01	0.00	0.01	0.00	0.29	0.01	0.01	0.00

Annex 3

THIAMETHOXAM (245)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.08 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day			Intake as ug/person/day								
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
OR 0691	Cotton seed oil, edible	PP	0.0004	3.22	0.00	1.54	0.00	1.01	0.00	0.74	0.00	1.12	0.00	2.93	0.00
-	Castor bean, raw (incl oil)	RAC	0.02	NC	-	0.07	0.00	NC	-	NC	-	NC	-	0.01	0.00
SO 0696	Palm kernels, raw (incl oil)	RAC	0.02	5.81	0.12	3.77	0.08	20.07	0.40	24.53	0.49	5.94	0.12	8.99	0.18
SO 0697	Peanuts, nutmeat, raw (incl roasted, incl oil, incl butter)	RAC	0.02	1.30	0.03	1.23	0.02	12.62	0.25	2.87	0.06	6.59	0.13	2.67	0.05
SO 0701	Shea nut (karite nuts), nutmeat, raw (incl butter)	RAC	0.02	NC	-	NC	-	0.34	0.01	NC	-	NC	-	NC	-
SO 0305	Olives for oil production, raw (incl oil)	RAC	0.02	12.61	0.25	1.35	0.03	0.27	0.01	8.04	0.16	0.58	0.01	21.80	0.44
SO 0696	Palm fruit (African Oil Palm), raw (incl oil)	RAC	0.02	28.87	0.58	1.09	0.02	53.08	1.06	80.61	1.61	24.20	0.48	17.72	0.35
SB 0715	Cacao bean, raw (incl roasted, incl powder, incl butter, incl paste, incl nes products)	RAC	0.02	0.72	0.01	4.20	0.08	0.60	0.01	4.21	0.08	0.42	0.01	0.78	0.02
SB 0716	Coffee bean, raw (i.e. green coffee)	RAC	0.035	0.96	0.03	0.16	0.01	0.91	0.03	0.27	0.01	1.37	0.05	0.46	0.02
SM 0716	Coffee bean, roasted	PP	0.0049	0.19	0.00	0.91	0.00	0.16	0.00	2.50	0.01	0.39	0.00	0.40	0.00
-	Coffee bean, instant coffee (incl essences and concentrates)	PP	0.035	0.07	0.00	0.94	0.03	0.07	0.00	0.70	0.02	0.07	0.00	0.29	0.01
-	Coffee bean, substitutes, containing coffee	PP	0.035	0.01	0.00	0.01	0.00	0.16	0.01	0.17	0.01	0.02	0.00	0.03	0.00
HH 0738	Mint, raw	RAC	0.34	0.50	0.17	0.01	0.00	NC	-	NC	-	NC	-	NC	-
HS 0444	Peppers, chili, dried	PP	0.8	0.42	0.34	0.53	0.42	0.84	0.67	0.50	0.40	0.95	0.76	0.37	0.30
DH 1100	Hops, dry	RAC	0.028	0.01	0.00	0.04	0.00	0.01	0.00	0.01	0.00	NC	-	0.01	0.00
DT 1114	Tea, green or black, fermented and dried, (including concentrates)	RAC	4.1	2.28	9.35	1.98	8.12	0.46	1.89	2.43	9.96	1.29	5.29	3.04	12.46
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat)	RAC	0.062	31.20	1.93	72.44	4.49	20.88	1.29	47.98	2.97	33.08	2.05	36.25	2.25
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.01	3.29	0.03	6.14	0.06	0.82	0.01	1.57	0.02	2.23	0.02	1.07	0.01
MO 0105	Edible offal (mammalian), raw	RAC	0.041	4.79	0.20	9.68	0.40	2.97	0.12	5.49	0.23	3.84	0.16	5.03	0.21
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.096	289.65	27.81	485.88	46.64	26.92	2.58	239.03	22.95	199.91	19.19	180.53	17.33
PM 0110	Poultry meat, raw (incl prepared)	RAC	0.064	14.63	0.94	29.76	1.90	8.04	0.51	129.68	8.30	25.04	1.60	35.66	2.28
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.033	0.10	0.00	0.10	0.00	NC	-	0.10	0.00	0.10	0.00	0.10	0.00
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.36	0.12	0.04	0.12	0.04	0.11	0.04	5.37	1.93	0.24	0.09	0.10	0.04
PE 0112	Eggs, raw, (incl dried)	RAC	0.028	7.84	0.22	23.08	0.65	2.88	0.08	14.89	0.42	9.81	0.27	14.83	0.42
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-

THIAMETHOXAM (245)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.08 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day			Intake as ug/person/day								
				G01 diet	G01 intake	G02 diet	G02 intake	G03 diet	G03 intake	G04 diet	G04 intake	G05 diet	G05 intake	G06 diet	G06 intake
	Total intake (ug/person)=				97.5		135.0		112.4		165.9		165.3		169.8
	Bodyweight per region (kg bw) =				60		60		60		60		60		60
	ADI (ug/person)=				4800		4800		4800		4800		4800		4800
	%ADI=				2.0%		2.8%		2.3%		3.5%		3.4%		3.5%
	Rounded %ADI=				2%		3%		2%		3%		3%		4%

THIAMETHOXAM (245)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.08 mg/kg bw							
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
FC 0002	Subgroup of Lemons and limes, raw (incl lemon juice) (incl kumquat commodities)	RAC	0.028	10.12	0.28	15.69	0.44	2.88	0.08	12.30	0.34	22.32	0.62	6.59	0.18
FC 0003	Subgroup of Mandarins, raw (incl mandarin juice)	RAC	0.028	12.42	0.35	14.99	0.42	16.08	0.45	10.78	0.30	9.94	0.28	NC	-
FC 0004	Subgroup of Oranges, sweet, sour, raw	RAC	0.028	15.68	0.44	24.00	0.67	6.80	0.19	29.09	0.81	15.39	0.43	160.47	4.49
JF 0004	Subgroup of Oranges, juice (single strength, incl. concentrated)	PP	0.007	33.31	0.23	1.78	0.01	0.28	0.00	18.97	0.13	14.01	0.10	13.36	0.09
FC 0005	Subgroup of Pummelo and grapefruits, raw (incl grapefruit juice)	RAC	0.028	8.21	0.23	4.60	0.13	0.64	0.02	5.85	0.16	19.98	0.56	368.86	10.33
FP 0226	Apple, raw (incl cider, excl juice)	RAC	0.07	41.14	2.88	56.49	3.95	26.64	1.86	31.58	2.21	51.94	3.64	3.05	0.21
JF 0226	Apple juice, single strength (incl. concentrated)	PP	0.065	14.88	0.97	11.98	0.78	0.15	0.01	9.98	0.65	30.32	1.97	3.47	0.23
FP 0228	Loquat, raw (incl processed) (i.e. Japanese medlar)	RAC	0.07	0.96	0.07	NC	-	NC	-	3.92	0.27	NC	-	2.49	0.17
FP 0229	Medlar, raw (incl processed)	RAC	0.07	NC	-	NC	-	NC	-	NC	-	NC	-	1.98	0.14
FP 0230	Pear, raw	RAC	0.07	8.79	0.62	8.44	0.59	12.37	0.87	9.60	0.67	10.27	0.72	0.23	0.02
FP 0307	Persimmon, Japanese, raw (i.e. Kaki fruit)	RAC	0.165	0.01	0.00	0.30	0.05	3.59	0.59	0.15	0.02	0.02	0.00	NC	-
FP 0231	Quince, raw	RAC	0.07	0.19	0.01	0.18	0.01	0.11	0.01	0.04	0.00	0.28	0.02	NC	-
FS 0012	Group of Stone fruits, raw (incl dried apricots, excl dried plums)	RAC	0.195	18.18	3.55	23.83	4.65	14.27	2.78	18.52	3.61	9.35	1.82	0.11	0.02
DF 0014	Plums, dried (prunes)	PP	0.16	0.61	0.10	0.35	0.06	0.05	0.01	0.35	0.06	0.49	0.08	0.13	0.02
FB 2005	Subgroup of Caneberries, raw	RAC	0.055	0.56	0.03	1.43	0.08	0.14	0.01	1.23	0.07	1.14	0.06	0.01	0.00

Annex 3

THIAMETHOXAM (245)															
International Estimated Daily Intake (IEDI)				ADI = 0 - 0.08 mg/kg bw											
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
FB 2006	Subgroup of Bush berries, raw (including processed)	RAC	0.055	1.31	0.07	5.50	0.30	0.01	0.00	2.57	0.14	0.82	0.05	2.15	0.12
FB 2007	Subgroup of Large shrub/tree berries, raw (including processed)	RAC	0.055	8.26	0.45	0.14	0.01	0.07	0.00	0.13	0.01	0.19	0.01	1.87	0.10
FB 0269	Grapes, raw (incl must, incl dried, incl juice, excl wine)	RAC	0.055	20.07	1.10	20.04	1.10	5.35	0.29	18.01	0.99	25.20	1.39	2.94	0.16
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.055	88.93	4.89	62.41	3.43	1.84	0.10	25.07	1.38	61.17	3.36	5.84	0.32
FB 2009	Subgroup of Low growing berries, raw	RAC	0.055	4.55	0.25	5.66	0.31	0.02	0.00	7.85	0.43	5.86	0.32	0.05	0.00
FI 0326	Avocado, raw	RAC	0.08	2.65	0.21	0.87	0.07	0.46	0.04	1.64	0.13	1.30	0.10	0.96	0.08
FI 0327	Banana, raw (incl plantains) (incl dried)	RAC	0.02	25.76	0.52	23.65	0.47	23.83	0.48	24.37	0.49	19.43	0.39	101.55	2.03
FI 0345	Mango, raw (incl canned mango, incl mango juice)	RAC	0.03	1.80	0.05	0.63	0.02	10.05	0.30	1.07	0.03	3.52	0.11	16.44	0.49
FI 0353	Pineapple, raw (incl canned pineapple, incl pineapple juice, incl dried pineapple)	RAC	0.01	13.13	0.13	11.13	0.11	6.94	0.07	14.36	0.14	36.74	0.37	18.81	0.19
VB 0040	Group of Brassica vegetables (excl Brassica leafy vegetables), raw	RAC	0.53	20.71	10.98	39.81	21.10	25.06	13.28	37.93	20.10	18.12	9.60	16.74	8.87
VC 0045	Group of Fruiting vegetables, cucurbits, raw	RAC	0.105	27.81	2.92	41.93	4.40	123.30	12.95	49.47	5.19	15.95	1.67	35.99	3.78
VO 0448	Tomato, raw (incl canned, excl juice, excl paste)	RAC	0.08	43.88	3.51	55.41	4.43	35.38	2.83	74.88	5.99	26.50	2.12	9.51	0.76
-	Tomato, paste (i.e. concentrated tomato sauce/puree)	PP	0.24	4.96	1.19	3.20	0.77	0.15	0.04	1.61	0.39	6.88	1.65	0.52	0.12
JF 0448	Tomato, juice (single strength, incl concentrated)	PP	0.054	0.80	0.04	0.07	0.00	0.05	0.00	0.61	0.03	0.40	0.02	0.08	0.00
VO 0442	Okra, raw (i.e. Lady's Finger, Gombo)	RAC	0.08	NC	-	NC	-	0.04	0.00	0.17	0.01	NC	-	0.72	0.06
VO 0444	Peppers, chili, raw	RAC	0.08	5.57	0.45	14.00	1.12	8.25	0.66	5.77	0.46	6.44	0.52	2.53	0.20
VO 0445	Peppers, sweet, raw (incl dried)	RAC	0.08	0.82	0.07	1.53	0.12	10.85	0.87	4.59	0.37	1.84	0.15	2.00	0.16
VO 0440	Egg plant, raw (i.e. aubergine)	RAC	0.08	1.01	0.08	1.69	0.14	21.37	1.71	3.00	0.24	1.40	0.11	NC	-
VL 2050	Subgroup of Leafy greens	RAC	0.54	18.38	9.93	18.73	10.11	82.36	44.47	25.32	13.67	17.60	9.50	7.37	3.98
VP 0061	Beans with pods (Phaseolus spp): (immature pods + succulent seeds)	RAC	0.08	5.07	0.41	0.83	0.07	0.17	0.01	3.70	0.30	NC	-	NC	-
014B	Peas with pods	-	0.01	-	-	-	-	-	-	-	-	-	-	-	-
VP 0062	Beans without pods: (Phaseolus spp.) (succulent seeds), raw	RAC	0.01	2.21	0.02	5.25	0.05	4.17	0.04	1.61	0.02	16.95	0.17	0.17	0.00
VP 0523	Broad bean without pods (succulent seeds) (Vicia spp), raw	RAC	0.01	0.22	0.00	0.84	0.01	0.15	0.00	0.48	0.00	2.04	0.02	NC	-
VP 0064	Peas without pods (Pisum spp) (succulent seeds)	RAC	0.01	10.72	0.11	1.99	0.02	2.72	0.03	4.26	0.04	4.23	0.04	NC	-

THIAMETHOXAM (245)		International Estimated Daily Intake (IEDI)				ADI = 0 - 0.08 mg/kg bw									
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day		Intake as ug/person/day									
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
VD 0070	Group of Pulses, raw (incl processed)	RAC	0.02	112.88	2.26	123.05	2.46	47.73	0.95	204.75	4.10	227.52	4.55	110.05	2.20
VD 0520	Bambara groundnut (dry) (Vigna subterranea), raw	RAC	0.01	NC	-	NC	-	NC	-	NC	-	NC	-	NC	-
VR 0075	Group of Root and tuber vegetables, raw (incl processed)	RAC	0.01	290.31	2.90	300.35	3.00	214.25	2.14	242.72	2.43	348.67	3.49	137.52	1.38
VS 0620	Artichoke globe, raw	RAC	0.23	0.98	0.23	3.65	0.84	0.07	0.02	1.67	0.38	0.26	0.06	NC	-
VF 2084	Group of edible fungi (cultivated & wild), raw (incl canned, incl dried)	RAC	0.08	7.31	0.58	5.92	0.47	1.26	0.10	3.73	0.30	14.85	1.19	0.57	0.05
GC 0653	Triticale, raw	RAC	0.032	NC	-	NC	-	0.01	0.00	0.01	0.00	NC	-	NC	-
GC 0654	Wheat, raw (incl meslin)	RAC	0.032	NC	-	NC	-	NC	-	0.01	0.00	NC	-	NC	-
-	Wheat, bulgur	PP	0.14	NC	-	NC	-	0.01	0.00	NC	-	NC	-	NC	-
CF 1210	Wheat, germ	PP	0.0493	0.97	0.05	0.10	0.00	0.03	0.00	0.01	0.00	NC	-	0.04	0.00
CP 1212	Wheat, wholemeal bread	PP	0.014	0.03	0.00	0.01	0.00	0.01	0.00	0.01	0.00	0.05	0.00	0.02	0.00
CP 1211	Wheat, white bread	PP	0.014	1.30	0.02	0.46	0.01	0.06	0.00	0.22	0.00	2.44	0.03	0.77	0.01
-	Wheat, Fermented Beverages (Korean jakju and takju)	PP	0.02	NC	-	NC	-	NC	-	4.36	0.09	NC	-	NC	-
CF 1211	Wheat, white flour (incl white flour products: starch, gluten, macaroni, pastry)	PP	0.00336	198.08	0.67	193.03	0.65	106.24	0.36	185.09	0.62	168.67	0.57	131.59	0.44
GC 0640	Barley, raw (incl malt extract, incl pot&pearled, incl beer, incl malt, excl flour & grits)	RAC	0.112	36.04	4.04	53.39	5.98	9.38	1.05	35.17	3.94	45.43	5.09	15.82	1.77
-	Barley, pot&pearled	PP	0.03	0.57	0.02	2.56	0.08	0.33	0.01	0.56	0.02	0.36	0.01	NC	-
-	Barley, flour (white flour and wholemeal flour)	PP	0.01	0.08	0.00	0.03	0.00	0.01	0.00	0.05	0.00	0.68	0.01	0.05	0.00
GC 0647	Oats, raw (incl rolled)	RAC	0.112	7.50	0.84	6.26	0.70	0.15	0.02	4.87	0.55	3.16	0.35	2.98	0.33
CM 0649 (GC 0649)	Rice, husked, dry (incl flour, incl oil, incl beverages, incl starch, excl polished)	REP	1.7	3.70	6.29	2.11	3.59	1.51	2.57	1.75	2.98	0.29	0.49	5.12	8.70
CM 1205	Rice polished, dry	PP	0.704	13.38	9.42	10.80	7.60	262.08	184.50	57.16	40.24	12.83	9.03	62.78	44.20
GC 0651	Sorghum, raw (incl beer, excl flour)(i.e. Chicken corn, Dari seed, Durra, Feterita)	RAC	0.079	NC	-	NC	-	0.01	0.00	1.15	0.09	NC	-	7.12	0.56
-	Sorghum, flour (white flour and wholemeal flour)	PP	0.0548	NC	-	NC	-	1.29	0.07	0.01	0.00	NC	-	NC	-
GC 0645	Maize, raw (incl glucose & dextrose & isoglucose, incl flour, incl oil, incl beer, incl germ, incl starch)	RAC	0.02	18.51	0.37	26.18	0.52	26.04	0.52	39.99	0.80	7.36	0.15	64.58	1.29

Annex 3

THIAMETHOXAM (245)															
International Estimated Daily Intake (IEDI)				ADI = 0 - 0.08 mg/kg bw											
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
GC 0447	Sweet corn on the cob, raw (incl frozen kernels, incl canned kernels)	RAC	0.01	11.43	0.11	3.71	0.04	0.74	0.01	13.63	0.14	3.07	0.03	1.50	0.02
GC 0447	Sweet corn on the cob, raw (i.e kernels plus cob without husks)	RAC	0.01	4.94	0.05	0.30	0.00	0.58	0.01	5.33	0.05	0.07	0.00	NC	-
-	Sugar crops NES, raw (incl sugar, syrup and others): sugar maple, sweet sorghum, sugar palm	RAC	0.033	4.87	0.16	2.50	0.08	0.89	0.03	40.03	1.32	1.05	0.03	2.83	0.09
TN 0672	Pecan, nutmeat	RAC	0.01	0.38	0.00	NC	-	NC	-	0.27	0.00	NC	-	0.26	0.00
SO 0091	Subgroup of Oilseeds, raw	RAC	0.02	2.58	0.05	4.10	0.08	5.77	0.12	6.63	0.13	1.81	0.04	0.64	0.01
SO 0090	Subgroup of Mustard seeds, raw (incl flour, incl oil)	RAC	0.02	0.30	0.01	0.48	0.01	0.33	0.01	0.63	0.01	1.03	0.02	0.40	0.01
SO 0693	Linseed, raw (incl oil)	RAC	0.02	NC	-	NC	-	0.02	0.00	0.01	0.00	NC	-	NC	-
SO 0698	Poppy seed, raw (incl oil)	RAC	0.02	0.02	0.00	0.25	0.01	0.01	0.00	0.02	0.00	NC	-	NC	-
SO 0495	Rape seed, raw (incl oil)	RAC	0.02	32.68	0.65	19.91	0.40	7.83	0.16	15.69	0.31	NC	-	NC	-
SO 0700	Sesame seed, raw (incl oil)	RAC	0.02	0.61	0.01	0.09	0.00	1.53	0.03	0.85	0.02	0.08	0.00	0.14	0.00
SO 0699	Safflower seed, raw (incl oil)	RAC	0.02	0.02	0.00	0.01	0.00	0.01	0.00	0.16	0.00	NC	-	NC	-
OR 0691	Cotton seed oil, edible	PP	0.0004	1.68	0.00	0.66	0.00	1.13	0.00	1.18	0.00	0.89	0.00	0.37	0.00
-	Castor bean, raw (incl oil)	RAC	0.02	NC	-	NC	-	0.01	0.00	NC	-	NC	-	NC	-
SO 0696	Palm kernels, raw (incl oil)	RAC	0.02	5.33	0.11	5.04	0.10	11.83	0.24	7.94	0.16	10.77	0.22	4.53	0.09
SO 0697	Peanuts, nutmeat, raw (incl roasted, incl oil, incl butter)	RAC	0.02	5.63	0.11	2.75	0.06	9.58	0.19	5.82	0.12	13.71	0.27	1.84	0.04
SO 0701	Shea nut (karite nuts), nutmeat, raw (incl butter)	RAC	0.02	NC	-	NC	-	NC	-	NC	-	NC	-	NC	-
SO 0305	Olives for oil production, raw (incl oil)	RAC	0.02	17.78	0.36	48.67	0.97	0.10	0.00	22.50	0.45	14.09	0.28	2.46	0.05
SO 0696	Palm fruit (African Oil Palm), raw (incl oil)	RAC	0.02	12.11	0.24	1.38	0.03	24.43	0.49	6.52	0.13	14.27	0.29	1.35	0.03
SB 0715	Cacao bean, raw (incl roasted, incl powder, incl butter, incl paste, incl nes products)	RAC	0.02	7.54	0.15	5.59	0.11	0.29	0.01	4.14	0.08	1.27	0.03	5.29	0.11
SB 0716	Coffee bean, raw (i.e. green coffee)	RAC	0.035	0.60	0.02	NC	-	0.62	0.02	1.71	0.06	NC	-	3.51	0.12
SM 0716	Coffee bean, roasted	PP	0.0049	7.02	0.03	9.75	0.05	0.02	0.00	5.09	0.02	13.38	0.07	0.77	0.00
-	Coffee bean, instant coffee (incl essences and concentrates)	PP	0.035	0.75	0.03	0.30	0.01	0.04	0.00	0.67	0.02	2.43	0.09	1.43	0.05
-	Coffee bean, substitutes, containing coffee	PP	0.035	0.08	0.00	0.09	0.00	0.02	0.00	0.02	0.00	0.07	0.00	0.15	0.01
HH 0738	Mint, raw	RAC	0.34	NC	-	NC	-	NC	-	NC	-	NC	-	NC	-
HS 0444	Peppers, chili, dried	PP	0.8	0.11	0.09	0.21	0.17	0.36	0.29	0.21	0.17	0.25	0.20	0.15	0.12

THIAMETHOXAM (245)															
International Estimated Daily Intake (IEDI)				ADI = 0 - 0.08 mg/kg bw											
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets as g/person/day				Intake as ug/person/day							
				G07 diet	G07 intake	G08 diet	G08 intake	G09 diet	G09 intake	G10 diet	G10 intake	G11 diet	G11 intake	G12 diet	G12 intake
DH 1100	Hops, dry	RAC	0.028	NC	-	NC	-	0.02	0.00	0.02	0.00	NC	-	NC	-
DT 1114	Tea, green or black, fermented and dried, (including concentrates)	RAC	4.1	2.91	11.93	1.73	7.09	1.14	4.67	1.85	7.59	2.29	9.39	0.74	3.03
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat)	RAC	0.062	140.03	8.68	150.89	9.36	79.32	4.92	111.24	6.90	120.30	7.46	51.27	3.18
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.01	6.44	0.06	15.51	0.16	3.79	0.04	8.29	0.08	18.44	0.18	8.00	0.08
MO 0105	Edible offal (mammalian), raw	RAC	0.041	15.17	0.62	5.19	0.21	6.30	0.26	6.78	0.28	3.32	0.14	3.17	0.13
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.096	388.92	37.34	335.88	32.24	49.15	4.72	331.25	31.80	468.56	44.98	245.45	23.56
PM 0110	Poultry meat, raw (incl prepared)	RAC	0.064	73.76	4.72	53.86	3.45	23.98	1.53	87.12	5.58	53.38	3.42	84.45	5.40
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.033	0.10	0.00	0.10	0.00	NC	-	0.10	0.00	0.71	0.02	NC	-
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.36	0.33	0.12	0.72	0.26	0.27	0.10	0.35	0.13	0.80	0.29	NC	-
PE 0112	Eggs, raw, (incl dried)	RAC	0.028	25.84	0.72	29.53	0.83	28.05	0.79	33.19	0.93	36.44	1.02	8.89	0.25
-	-	-		-	-	-	-	-	-	-	-	-	-	-	-
Total intake (ug/person)=				141.2				141.6				296.0			
Bodyweight per region (kg bw) =				60				60				55			
ADI (ug/person)=				4800				4800				4400			
%ADI=				2.9%				3.0%				6.7%			
Rounded %ADI=				3%				3%				7%			

Annex 3

THIAMETHOXAM (245)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.08 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
FC 0002	Subgroup of Lemons and limes, raw (incl lemon juice) (incl kumquat commodities)	RAC	0.028	18.97	0.53	0.97	0.03	6.23	0.17	0.09	0.00	3.35	0.09
FC 0003	Subgroup of Mandarins, raw (incl mandarin juice)	RAC	0.028	0.16	0.00	0.27	0.01	9.06	0.25	0.01	0.00	0.02	0.00
FC 0004	Subgroup of Oranges, sweet, sour, raw	RAC	0.028	1.18	0.03	1.11	0.03	14.28	0.40	0.05	0.00	1.08	0.03
JF 0004	Subgroup of Oranges, juice (single strength, incl. concentrated)	PP	0.007	0.08	0.00	0.26	0.00	12.61	0.09	0.14	0.00	0.33	0.00
FC 0005	Subgroup of Pummelo and grapefruits, raw (incl grapefruit juice)	RAC	0.028	0.68	0.02	0.05	0.00	3.21	0.09	0.01	0.00	NC	-
FP 0226	Apple, raw (incl cider, excl juice)	RAC	0.07	66.67	4.67	2.06	0.14	55.83	3.91	188.29	13.18	1.38	0.10
JF 0226	Apple juice, single strength (incl. concentrated)	PP	0.065	0.03	0.00	0.10	0.01	7.19	0.47	0.03	0.00	NC	-
FP 0228	Loquat, raw (incl processed) (i.e. Japanese medlar)	RAC	0.07	0.94	0.07	4.68	0.33	NC	-	0.50	0.04	3.08	0.22
FP 0229	Medlar, raw (incl processed)	RAC	0.07	0.75	0.05	3.73	0.26	4.87	0.34	0.40	0.03	2.45	0.17
FP 0230	Pear, raw	RAC	0.07	0.07	0.00	0.14	0.01	9.45	0.66	0.01	0.00	0.14	0.01
FP 0307	Persimmon, Japanese, raw (i.e. Kaki fruit)	RAC	0.165	0.41	0.07	0.32	0.05	0.02	0.00	0.58	0.10	12.51	2.06
FP 0231	Quince, raw	RAC	0.07	NC	-	NC	-	0.65	0.05	NC	-	NC	-
FS 0012	Group of Stone fruits, raw (incl dried apricots, excl dried plums)	RAC	0.195	0.09	0.02	0.02	0.00	32.27	6.29	0.01	0.00	NC	-
DF 0014	Plums, dried (prunes)	PP	0.16	0.01	0.00	0.01	0.00	0.37	0.06	0.01	0.00	NC	-
FB 2005	Subgroup of Caneberries, raw	RAC	0.055	0.01	0.00	7.30	0.40	2.29	0.13	0.01	0.00	NC	-
FB 2006	Subgroup of Bush berries, raw (including processed)	RAC	0.055	0.82	0.05	4.05	0.22	5.94	0.33	0.43	0.02	2.66	0.15
FB 2007	Subgroup of Large shrub/tree berries, raw (including processed)	RAC	0.055	0.71	0.04	7.32	0.40	NC	-	0.38	0.02	2.32	0.13
FB 0269	Grapes, raw (incl must, incl dried, incl juice, excl wine)	RAC	0.055	0.17	0.01	0.94	0.05	20.24	1.11	0.03	0.00	0.40	0.02
-	Grape wine (incl vermouths) (from wine-grapes)	PP	0.055	0.31	0.02	0.23	0.01	60.43	3.32	0.52	0.03	31.91	1.76
FB 2009	Subgroup of Low growing berries, raw	RAC	0.055	0.01	0.00	0.01	0.00	3.37	0.19	0.01	0.00	0.01	0.00
FI 0326	Avocado, raw	RAC	0.08	1.12	0.09	0.01	0.00	0.84	0.07	0.01	0.00	6.60	0.53
FI 0327	Banana, raw (incl plantains) (incl dried)	RAC	0.02	44.80	0.90	118.17	2.36	25.25	0.51	454.49	9.09	310.23	6.20
FI 0345	Mango, raw (incl canned mango, incl mango juice)	RAC	0.03	12.25	0.37	6.83	0.20	0.76	0.02	0.01	0.00	20.12	0.60
FI 0353	Pineapple, raw (incl canned pineapple, incl pineapple juice, incl dried pineapple)	RAC	0.01	8.51	0.09	6.27	0.06	6.89	0.07	0.18	0.00	24.94	0.25
VB 0040	Group of Brassica vegetables (excl Brassica leafy	RAC	0.53	5.46	2.89	4.28	2.27	58.72	31.12	0.02	0.01	NC	-

THIAMETHOXAM (245)													
International Estimated Daily Intake (IEDI)				ADI = 0 - 0.08 mg/kg bw									
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
	vegetables), raw												
VC 0045	Group of Fruiting vegetables, cucurbits, raw	RAC	0.105	5.96	0.63	9.74	1.02	51.82	5.44	13.61	1.43	0.05	0.01
VO 0448	Tomato, raw (incl canned, excl juice, excl paste)	RAC	0.08	13.10	1.05	4.90	0.39	62.16	4.97	1.04	0.08	0.09	0.01
-	Tomato, paste (i.e. concentrated tomato sauce/puree)	PP	0.24	0.58	0.14	0.22	0.05	2.21	0.53	0.24	0.06	3.10	0.74
JF 0448	Tomato, juice (single strength, incl concentrated)	PP	0.054	0.05	0.00	0.01	0.00	0.42	0.02	0.01	0.00	0.02	0.00
VO 0442	Okra, raw (i.e. Lady's Finger, Gombo)	RAC	0.08	6.23	0.50	0.10	0.01	NC	-	NC	-	NC	-
VO 0444	Peppers, chili, raw	RAC	0.08	3.47	0.28	3.56	0.28	16.30	1.30	0.01	0.00	NC	-
VO 0445	Peppers, sweet, raw (incl dried)	RAC	0.08	5.49	0.44	10.57	0.85	8.84	0.71	0.91	0.07	NC	-
VO 0440	Egg plant, raw (i.e. aubergine)	RAC	0.08	1.31	0.10	8.26	0.66	3.95	0.32	0.01	0.00	NC	-
VL 2050	Subgroup of Leafy greens	RAC	0.54	4.99	2.69	3.29	1.78	7.53	4.07	3.05	1.65	6.09	3.29
VP 0061	Beans with pods (Phaseolus spp): (immature pods + succulent seeds)	RAC	0.08	NC	-	NC	-	NC	-	NC	-	NC	-
014B	Peas with pods	-	0.01	-	-	-	-	-	-	-	-	-	-
VP 0062	Beans without pods: (Phaseolus spp.) (succulent seeds), raw	RAC	0.01	0.30	0.00	3.13	0.03	4.11	0.04	0.01	0.00	NC	-
VP 0523	Broad bean without pods (succulent seeds) (Vicia spp), raw	RAC	0.01	0.07	0.00	0.01	0.00	0.76	0.01	NC	-	NC	-
VP 0064	Peas without pods (Pisum spp) (succulent seeds)	RAC	0.01	0.21	0.00	0.02	0.00	5.51	0.06	0.02	0.00	NC	-
VD 0070	Group of Pulses, raw (incl processed)	RAC	0.02	46.57	0.93	30.77	0.62	112.53	2.25	75.53	1.51	43.68	0.87
VD 0520	Bambara groundnut (dry) (Vigna subterranea), raw	RAC	0.01	0.20	0.00	NC	-	NC	-	NC	-	NC	-
VR 0075	Group of Root and tuber vegetables, raw (incl processed)	RAC	0.01	282.25	2.82	232.11	2.32	281.91	2.82	620.21	6.20	459.96	4.60
VS 0620	Artichoke globe, raw	RAC	0.23	0.01	0.00	NC	-	0.08	0.02	0.01	0.00	NC	-
VF 2084	Group of edible fungi (cultivated & wild), raw (incl canned, incl dried)	RAC	0.08	0.02	0.00	0.04	0.00	3.73	0.30	0.01	0.00	NC	-
GC 0653	Triticale, raw	RAC	0.032	0.01	0.00	NC	-	NC	-	NC	-	NC	-
GC 0654	Wheat, raw (incl meslin)	RAC	0.032	NC	-	NC	-	NC	-	NC	-	0.97	0.03
-	Wheat, bulgur	PP	0.14	0.01	0.00	NC	-	NC	-	NC	-	NC	-
CF 1210	Wheat, germ	PP	0.0493	0.04	0.00	0.01	0.00	0.01	0.00	0.01	0.00	NC	-
CP 1212	Wheat, wholemeal bread	PP	0.014	0.01	0.00	0.01	0.00	0.03	0.00	0.01	0.00	0.01	0.00

Annex 3

THIAMETHOXAM (245)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.08 mg/kg bw					
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person							
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
CP 1211	Wheat, white bread	PP	0.014	0.43	0.01	0.41	0.01	1.56	0.02	0.11	0.00	0.07	0.00
-	Wheat, Fermented Beverages (Korean jakju and takju)	PP	0.02	NC	-	NC	-	NC	-	NC	-	NC	-
CF 1211	Wheat, white flour (incl white flour products: starch, gluten, macaroni, pastry)	PP	0.00336	44.78	0.15	86.96	0.29	214.05	0.72	20.31	0.07	103.60	0.35
GC 0640	Barley, raw (incl malt extract, incl pot&pearled, incl beer, incl malt, excl flour & grits)	RAC	0.112	11.54	1.29	2.33	0.26	46.13	5.17	3.72	0.42	16.26	1.82
-	Barley, pot&pearled	PP	0.03	5.46	0.16	0.01	0.00	1.44	0.04	0.01	0.00	NC	-
-	Barley, flour (white flour and wholemeal flour)	PP	0.01	0.02	0.00	NC	-	0.32	0.00	0.01	0.00	NC	-
GC 0647	Oats, raw (incl rolled)	RAC	0.112	0.37	0.04	0.07	0.01	2.79	0.31	0.10	0.01	NC	-
CM 0649 (GC 0649)	Rice, husked, dry (incl flour, incl oil, incl beverages, incl starch, excl polished)	REP	1.7	13.58	23.09	4.29	7.29	2.17	3.69	0.01	0.02	8.84	15.03
CM 1205	Rice polished, dry	PP	0.704	30.20	21.26	218.34	153.71	12.77	8.99	15.24	10.73	51.35	36.15
GC 0651	Sorghum, raw (incl beer, excl flour)(i.e. Chicken corn, Dari seed, Durra, Feterita)	RAC	0.079	4.73	0.37	NC	-	NC	-	13.36	1.06	NC	-
-	Sorghum, flour (white flour and wholemeal flour)	PP	0.0548	75.99	4.16	1.82	0.10	NC	-	19.82	1.09	NC	-
GC 0645	Maize, raw (incl glucose & dextrose & isoglucose, incl flour, incl oil, incl beer, incl germ, incl starch)	RAC	0.02	116.66	2.33	10.52	0.21	38.46	0.77	76.60	1.53	34.44	0.69
GC 0447	Sweet corn on the cob, raw (incl frozen kernels, incl canned kernels)	RAC	0.01	3.63	0.04	20.50	0.21	8.78	0.09	0.02	0.00	0.17	0.00
GC 0447	Sweet corn on the cob, raw (i.e kernels plus cob without husks)	RAC	0.01	3.61	0.04	20.45	0.20	6.00	0.06	0.01	0.00	0.17	0.00
-	Sugar crops NES, raw (incl sugar, syrup and others): sugar maple, sweet sorghum, sugar palm	RAC	0.033	0.49	0.02	0.63	0.02	4.52	0.15	0.40	0.01	5.87	0.19
TN 0672	Pecan, nutmeat	RAC	0.01	0.15	0.00	0.22	0.00	0.31	0.00	0.01	0.00	0.01	0.00
SO 0091	Subgroup of Oilseeds, raw	RAC	0.02	9.97	0.20	0.84	0.02	1.73	0.03	13.57	0.27	0.53	0.01
SO 0090	Subgroup of Mustard seeds, raw (incl flour, incl oil)	RAC	0.02	0.04	0.00	0.19	0.00	0.32	0.01	0.06	0.00	0.01	0.00
SO 0693	Linseed, raw (incl oil)	RAC	0.02	0.07	0.00	NC	-	0.03	0.00	NC	-	NC	-
SO 0698	Poppy seed, raw (incl oil)	RAC	0.02	0.01	0.00	0.01	0.00	0.11	0.00	NC	-	NC	-
SO 0495	Rape seed, raw (incl oil)	RAC	0.02	0.19	0.00	0.07	0.00	12.07	0.24	0.08	0.00	NC	-
SO 0700	Sesame seed, raw (incl oil)	RAC	0.02	2.34	0.05	0.66	0.01	0.26	0.01	9.84	0.20	NC	-
SO 0699	Safflower seed, raw (incl oil)	RAC	0.02	0.05	0.00	NC	-	NC	-	NC	-	NC	-
OR 0691	Cotton seed oil, edible	PP	0.0004	1.28	0.00	0.05	0.00	0.45	0.00	0.42	0.00	0.15	0.00

THIAMETHOXAM (245)				International Estimated Daily Intake (IEDI)				ADI = 0 - 0.08 mg/kg bw						
Codex Code	Commodity description	Expr as	STMR mg/kg	Diets: g/person/day		Intake = daily intake: ug/person								
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake	
-	Castor bean, raw (incl oil)	RAC	0.02	NC	-	NC	-	NC	-	NC	-	NC	-	
SO 0696	Palm kernels, raw (incl oil)	RAC	0.02	60.84	1.22	12.77	0.26	5.41	0.11	0.57	0.01	53.45	1.07	
SO 0697	Peanuts, nutmeat, raw (incl roasted, incl oil, incl butter)	RAC	0.02	18.82	0.38	0.57	0.01	2.28	0.05	6.90	0.14	0.53	0.01	
SO 0701	Shea nut (karite nuts), nutmeat, raw (incl butter)	RAC	0.02	0.95	0.02	NC	-	NC	-	NC	-	NC	-	
SO 0305	Olives for oil production, raw (incl oil)	RAC	0.02	0.18	0.00	0.11	0.00	11.00	0.22	0.06	0.00	0.49	0.01	
SO 0696	Palm fruit (African Oil Palm), raw (incl oil)	RAC	0.02	36.35	0.73	7.16	0.14	2.99	0.06	22.89	0.46	28.38	0.57	
SB 0715	Cacao bean, raw (incl roasted, incl powder, incl butter, incl paste, incl nes products)	RAC	0.02	0.11	0.00	0.89	0.02	6.28	0.13	0.17	0.00	2.31	0.05	
SB 0716	Coffee bean, raw (i.e. green coffee)	RAC	0.035	0.83	0.03	0.69	0.02	1.09	0.04	2.91	0.10	0.82	0.03	
SM 0716	Coffee bean, roasted	PP	0.0049	0.02	0.00	0.41	0.00	7.50	0.04	0.01	0.00	0.06	0.00	
-	Coffee bean, instant coffee (incl essences and concentrates)	PP	0.035	0.03	0.00	0.05	0.00	0.60	0.02	0.01	0.00	5.53	0.19	
-	Coffee bean, substitutes, containing coffee	PP	0.035	0.01	0.00	0.03	0.00	0.13	0.00	0.01	0.00	NC	-	
HH 0738	Mint, raw	RAC	0.34	NC	-	NC	-	NC	-	NC	-	NC	-	
HS 0444	Peppers, chili, dried	PP	0.8	0.58	0.46	1.27	1.02	1.21	0.97	0.12	0.10	NC	-	
DH 1100	Hops, dry	RAC	0.028	NC	-	NC	-	0.04	0.00	NC	-	NC	-	
DT 1114	Tea, green or black, fermented and dried, (including concentrates)	RAC	4.1	0.53	2.17	5.25	21.53	0.86	3.53	0.56	2.30	0.88	3.61	
MM 0095	MEAT FROM MAMMALS other than marine mammals, raw (incl prepared meat)	RAC	0.062	29.18	1.81	50.89	3.16	121.44	7.53	22.58	1.40	72.14	4.47	
MF 0100	Mammalian fats, raw, excl milk fats (incl rendered fats)	RAC	0.01	1.05	0.01	1.14	0.01	18.69	0.19	0.94	0.01	3.12	0.03	
MO 0105	Edible offal (mammalian), raw	RAC	0.041	4.64	0.19	1.97	0.08	10.01	0.41	3.27	0.13	3.98	0.16	
ML 0106	Milks, raw or skimmed (incl dairy products)	RAC	0.096	108.75	10.44	70.31	6.75	436.11	41.87	61.55	5.91	79.09	7.59	
PM 0110	Poultry meat, raw (incl prepared)	RAC	0.064	3.92	0.25	12.03	0.77	57.07	3.65	5.03	0.32	55.56	3.56	
PF 0111	Poultry fat, raw (incl rendered)	RAC	0.033	NC	-	NC	-	0.32	0.01	NC	-	NC	-	
PO 0111	Poultry edible offal, raw (incl prepared)	RAC	0.36	0.10	0.04	0.70	0.25	0.97	0.35	0.10	0.04	NC	-	
PE 0112	Eggs, raw, (incl dried)	RAC	0.028	3.84	0.11	4.41	0.12	27.25	0.76	1.13	0.03	7.39	0.21	
-	-	-		-	-	-	-	-	-	-	-	-	-	
Total intake (ug/person)=					90.6		211.4		152.7		59.9		97.7	

Annex 3

THIAMETHOXAM (245)													
International Estimated Daily Intake (IEDI)				ADI = 0 - 0.08 mg/kg bw									
Codex Code	Commodity description	Expr as	STM R mg/kg	Diets: g/person/day			Intake = daily intake: ug/person						
				G13 diet	G13 intake	G14 diet	G14 intake	G15 diet	G15 intake	G16 diet	G16 intake	G17 diet	G17 intake
	Bodyweight per region (kg bw) =				60		60		60		60		60
	ADI (ug/person)=				4800		4800		4800		4800		4800
	%ADI=				1.9%		4.4%		3.2%		1.2%		2.0%
	Rounded %ADI=				2%		4%		3%		1%		2%

Annex 4

Annex 4: International Estimate of Short-Term Intakes (IESTIs) of pesticide residues

ACETAMIPRID (246)											IESTI					
Acute RfD= 0.1 mg/kg bw (100 µg/kg bw)											Maximum %ARfD:					
											all		gen pop		child	
Codex Code	Commodity	Processing	STMR or STMR-P mg/kg	HR or HR-P mg/kg	DCF	Country	Population group	n	Large portion, g/person	Unit weight, edible portion, g	Variability factor	Case	IESTI µg/kg bw/day	% acute RfD rounded	% acute RfD rounded	% acute RfD rounded
TN 0675	Pistachio nut (all commodities)	highest utilisation: Total	0.33	0.51	1.000	PRIMO-IE	child	P97.5	115.86	<25	NR	1	0.01 - 2.95	0% - 3%	0% - 2%	1% - 3%

BIXAFEN (262)											IESTI					
Acute RfD= 0.2 mg/kg bw (200 µg/kg bw)											Maximum %ARfD:					
											all		gen pop		child	
Codex Code	Commodity	Processing	STMR or STMR-P mg/kg	HR or HR-P mg/kg	DCF	Country	Population group	n	Large portion, g/person	Unit weight, edible portion, g	Variability factor	Case	IESTI µg/kg bw/day	% acute RfD rounded	% acute RfD rounded	% acute RfD rounded
VD 0071	Beans (dry) (Phaseolus spp) (all commodities)	highest utilisation: Total	0.02		1.000	PRIMO-UK	infant	P97.5	159.00	<25	NR	3	0.04 - 0.37	0% - 0%	0% - 0%	0% - 0%
VD 0523	Broad bean (dry) (Vicia spp)	cooked/boiled	0.02		0.400	CN	Gen pop, > 1 yrs	737	1190.24	<25	NR	3	0.179	0%	0%	0%
VD 0531	Lablab bean (dry) (Lablab spp)	cooked/boiled	0.02		0.400	CN	Gen pop, > 1 yrs	1219	972.42	<25	NR	3	0.146	0%	0%	-
VD 0541	Soya bean (dry) (Glycine spp) (all commodities)	highest utilisation: Oil (refined)	0.00221 - 0.0476		1.000	US	Child, < 6 yrs	6511	37.75	NR	NR	3	0.02 - 0.12	0% - 0%	0% - 0%	0% - 0%
VD 0072	Peas (dry) (Pisum spp)	cooked/boiled	0.02		0.400	CN	Gen pop, > 1 yrs	268	1673.82	<25	NR	3	0.252	0%	0%	0%
VD 0524	Chick-pea (dry) (Cicer spp)	canned/preserved	0.02		0.400	PRIMO-NL	child	P100	328.80	<25	NR	3	0.143	0%	0%	0%
VD 0533	Lentil (dry) (Lens spp) (all commodities)	highest utilisation: Total	0.02		1.000	PRIMO-UK	Child, 11-14 yrs	P97.5	321.50	<25	NR	3	0.04 - 0.13	0% - 0%	0% - 0%	0% - 0%
VD 0537	Pigeon pea (dry) (Cajanus spp)	Total	0.02		1.000	AU	Gen pop, > 2 yrs	129	95.83	<25	NR	3	0.029	0%	0%	0%
VR 0574	Beetroot (all commodities)	highest utilisation: Total	0.028	0.068	1.000	AU	Child, 2-6 yrs	53	314.08	135	3	2a	0.02 - 2.09	0% - 1%	0% - 0%	0% - 1%

BIXAFEN (262)										IESTI						
Acute RfD= 0.2 mg/kg bw (200 µg/kg bw)										Maximum %ARfD:				20%	20%	20%
													all	gen pop	child	
Codex Code	Commodity	Processing	STMR or STMR-P mg/kg	HR or HR-P mg/kg	DCF	Country	Population group	n	Large portion, g/person	Unit weight, edible portion, g	Variability factor	Case	IESTI µg/kg bw/day	% acute RfD rounded	% acute RfD rounded	% acute RfD rounded
VR 0575	Burdock, greater or edible	Total	0.028	0.068	1.000	JP	Child, 1-6 yrs	122	35.70	68	3	2b	0.444	0%	0%	0%
VR 0577	Carrot (all commodities)	highest utilisation: raw with skin	0.028	0.068	1.000	CN	Child, 1-6 yrs	400	234.68	300	3	2b	0 - 2.97	0% - 1%	0% - 1%	0% - 1%
VR 0578	Celeriac (Turnip rooted celery) (all commodities)	highest utilisation: Total	0.028	0.068	1.000	PRIMO-BE	toddler	P100	196.90	749	3	2b	0.01 - 2.26	0% - 1%	0% - 0%	0% - 1%
VR 0469	Chicory, roots (all commodities)	highest utilisation: Total	0.028	0.068	1.000	AU	Gen pop, > 2 yrs	175	26.16	48	3	2b	0.02 - 0.08	0% - 0%	0% - 0%	0% - 0%
VR 0583	Horseradish (all commodities)	highest utilisation: Total	0.028	0.068	1.000	PRIMO-DE	Gen pop	P97.5	79.50	220	3	2b	0 - 0.21	0% - 0%	0% - 0%	0% - 0%
VR 0587	Parsley, turnip-rooted (Hamburg roots) (all commodities)	highest utilisation: Total	0.028	0.068	1.000	PRIMO-NL	Gen pop	P97.5	96.60	140	3	2b	0.2 - 0.3	0% - 0%	0% - 0%	0% - 0%
VR 0588	Parsnip (all commodities)	highest utilisation: cooked/boiled (without skin)	0.028	0.068	1.000	PRIMO-NL	child	E	133.30	227	3	2b	0.16 - 1.48	0% - 1%	0% - 0%	1% - 1%
VR 0494	Radish (all commodities)	highest utilisation: Total	0.028	0.068	1.000	PRIMO-NL	child	E	64.40	172	3	2b	0 - 0.71	0% - 0%	0% - 0%	0% - 0%
VR 0590	Radish, black (all commodities)	highest utilisation: Total	0.028	0.068	1.000	PRIMO-NL	child	E	64.40	172	3	2b	0 - 0.71	0% - 0%	0% - 0%	0% - 0%
VR 0591	Radish, Japanese (Chinese radish, Daikon) (all commodities)	highest utilisation: raw without skin	0.028	0.068	1.000	CN	Child, 1-6 yrs	1187	293.37	1000	3	2b	0 - 3.71	0% - 2%	0% - 1%	0% - 2%
VR 0498	Salsify (Oyster plant) (all commodities)	highest utilisation: Total	0.028	0.068	1.000	PRIMO-BE	toddler	P100	99.90	75	3	2a	0 - 0.96	0% - 0%	0% - 0%	0% - 0%
VR 0596	Sugar beet (all commodities)	highest utilisation: composite foods; unspecified ind processed	0.028	0.068	1.000	NL	Child, 2-6 yrs	2554	168.93	NR	NR	3	0.06 - 0.26	0% - 0%	0% - 0%	0% - 0%
VR 0497	Swede (Rutabaga) (all commodities)	highest utilisation: Total	0.028	0.068	1.000	PRIMO-UK	infant	P97.5	90.00	500	3	2b	0 - 2.11	0% - 1%	0% - 1%	0% - 1%
VR 0506	Turnip, garden (all commodities)	highest utilisation: Total	0.028	0.068	1.000	US	Gen pop, 0-85 yrs	53	1533.22	105	3	2a	0.2 - 1.68	0% - 1%	0% - 1%	0% - 1%
VR 0573	Arrowroot (all commodities)	highest utilisation: starch	0.028	0.068	1.000	PRIMO-NL	child	E	12.40	NR	NR	3	0 - 0.02	0% - 0%	0% - 0%	0% - 0%
VR 0463	Cassava (Manioc) (all commodities)	highest utilisation: cooked/boiled (without	0.028	0.068	1.000	PRIMO-NL	Gen pop	E	250.00	356	3	2b	0.02 - 0.78	0% - 0%	0% - 0%	0% - 0%

Annex 4

BIXAFEN (262)										IESTI						
Acute RfD= 0.2 mg/kg bw (200 µg/kg bw)										Maximum %ARfD:			20%	20%	20%	
													all	gen pop	child	
Codex Code	Commodity	Processing	STMR or STMR-P mg/kg	HR or HR-P mg/kg	DCF	Country	Population group	n	Large portion, g/person	Unit weight, edible portion, g	Variability factor	Case	IESTI µg/kg bw/day	% acute RfD rounded	% acute RfD rounded	% acute RfD rounded
		peel)														
VR 0585	Jerusalem artichoke (i.e. Topinambur) (all commodities)	highest utilisation: cooked/boiled (without peel)	0.028	0.068	1.000	PRIMO-NL	child	E	133.30	56	3	2a	0.37 - 0.91	0% - 0%	0% - 0%	0% - 0%
VR 0589	Potato (all commodities)	highest utilisation: Total	0.028	0.068	1.000	PRIMO-UK	infant	P97.5	191.10	216	3	2b	0.02 - 4.48	0% - 2%	0% - 1%	0% - 2%
VR 0508	Sweet potato (all commodities)	highest utilisation: Total	0.028	0.068	1.000	CA	Child, <6 yrs	91	358.61	546	3	2b	0.2 - 5.73	0% - 3%	0% - 1%	1% - 3%
VR 0504	Tannia (Tanier, Yautia) (all commodities)	highest utilisation: cooked/boiled (without peel)	0.028	0.068	1.000	NL	Gen pop, > 1 yrs	E	249.97	170	3	2a	0.02 - 0.61	0% - 0%	0% - 0%	0% - 0%
VR 0505	Taro (Dasheen, Eddoe) (all commodities)	highest utilisation: cooked/boiled (without peel)	0.028	0.068	1.000	CN	Child, 1-6 yrs	199	384.18	668	3	2b	0.5 - 4.86	0% - 2%	0% - 1%	0% - 2%
VR 0600	Yams (all commodities)	highest utilisation: Total	0.028	0.068	1.000	PRIMO-UK	adult	P97.5	693.70	365	3	2a	0.49 - 1.27	0% - 1%	0% - 1%	1% - 1%
VR 3001	Chinese water chestnut (all commodities)	highest utilisation: cooked/boiled without peel	0.028	0.068	1.000	TH	Child, 3-6 yrs	169	134.24	<25	NR	1	0.06 - 0.53	0% - 0%	0% - 0%	0% - 0%
VR 3002	Lotus tuber	Total	0.028	0.068	1.000	CN	Child, 1-6 yrs	200	256.38	120	3	2a	2.092	1%	0%	1%
GC 0653	Triticale	Total	0.02		1.000	DE	Gen pop, 14-80 yrs	27100	394.70	<25	NR	3	0.103	0%	0%	0%
GC 0654	Wheat (all commodities)	highest utilisation: flour (cereals)	0.03135 - 0.2375		1.000	CN	Child, 1-6 yrs	3556	415.87	NR	NR	3	0.05 - 0.81	0% - 0%	0% - 0%	0% - 0%
GC 0640	Barley (all commodities)	highest utilisation: beer	0.036 - 0.33		0.190	CA	Gen pop, all ages	2514	21271.20	NR	NR	3	0.12 - 1.85	0% - 1%	0% - 1%	0% - 0%
GC 0651	Sorghum grain (Chicken corn, Dari seed, Durra, Feterita) (all commodities)	highest utilisation: cooked/boiled	0.196		0.400	CN	Gen pop, > 1 yrs	356	1348.67	<25	NR	3	0.31 - 1.99	0% - 1%	0% - 1%	0% - 0%
GC 0645	Maize (corn) (all commodities)	highest utilisation: Flour (cereals)	0.0126 - 0.062		1.000	CN	Child, 1-6 yrs	213	361.24	NR	NR	3	0.01 - 0.4	0% - 0%	0% - 0%	0% - 0%
GC 0447	Sweet corn (corn-on-the-cob) (kernels plus cob with husks removed) (all commodities)	highest utilisation: cooked/boiled (corn-on-the-cob)	0.02	0.02	1.000	TH	Child, 3-6 yrs	1383	196.99	191	3	2a	0.02 - 0.68	0% - 0%	0% - 0%	0% - 0%

BIXAFEN (262)										IESTI						
Acute RfD= 0.2 mg/kg bw (200 µg/kg bw)										Maximum %ARfD:						
										20% 20% 20%						
										all gen pop child						
Codex Code	Commodity	Processing	STMR or STMR-P mg/kg	HR or HR-P mg/kg	DCF	Country	Population group	n	Large portion, g/person	Unit weight, edible portion, g	Variability factor	Case	IESTI µg/kg bw/day	% acute RfD rounded	% acute RfD rounded	% acute RfD rounded
SO 0495	Rape seed (Canola) (all commodities)	highest utilisation: Oil (refined)	0.02 - 0.03		1.000	CA	Child, <6 yrs	1127	26.46	NR	NR	3	0.05 - 0.05	0% - 0%	0% - 0%	0% - 0%
SO 0702	Sunflower seed (all commodities)	highest utilisation: Total	0.045		1.000	CA	women, 15-49 yrs	121	296.25	<25	NR	3	0.02 - 0.21	0% - 0%	0% - 0%	0% - 0%
SO 0691	Cotton seed (all commodities)	highest utilisation: Oil (refined)	0.03		1.000	US	Child, < 6 yrs	6354	3.13	NR	NR	3	0.01 - 0.01	0% - 0%	0% - 0%	0% - 0%
SO 0697	Peanut, shelled (groundnut) (all commodities)	highest utilisation: raw incl roasted	0.02 - 0.044		1.000	CN	Child, 1-6 yrs	290	163.07	<25	NR	3	0.01 - 0.2	0% - 0%	0% - 0%	0% - 0%
MM 0095	Meat from mammals other than marine mammals	Total	NA	NA	1.000	CN	Child, 1-6 yrs	302	264.84	NR	NR	1	NA	7%	5%	7%
MM 0095	Meat from mammals other than marine mammals: 20% as fat	Total		1.3	1.000	CN	Child, 1-6 yrs	302	52.97	NR	NR	1	4.268	2%	2%	2%
MM 0095	Meat from mammals other than marine mammals: 80% as muscle	Total		0.71	1.000	CN	Child, 1-6 yrs	302	211.87	NR	NR	1	9.323	5%	3%	5%
MF 0100	Mammalian fats (except milk fats)	Total		1.3	1.000	PRIMO-FR	adult	P97.5	134.79	NR	NR	1	2.639	1%	1%	1%
MO 1280	Cattle, kidney	Total		0.93	1.000	PRIMO-UK	toddler	P97.5	54.60	NR	NR	1	3.502	2%	1%	2%
MO 1281	Cattle, liver	Total		3.9	1.000	PRIMO-UK	infant	P97.5	70.20	NR	NR	1	31.469	20%	10%	20%
MO 0814	Goat, edible offal of (includes kidney and liver)	Total		3.9	1.000	NL	Gen pop, > 1 yrs	0	NC	NR	NR	1	NC	NC	NC	NC
MO 0816	Horse, edible offal of (includes kidney and liver)	Total		3.9	1.000	NL	Gen pop, > 1 yrs	0	NC	NR	NR	1	NC	NC	NC	NC
MO 1284	Pig kidney	Total		0.93	1.000	CN	Gen pop, > 1 yrs	205	224.78	NR	NR	1	3.927	2%	2%	1%
MO 1285	Pig liver	Total		3.9	1.000	BR	Gen pop, > 10 yrs	20	505.00	NR	NR	1	30.507	20%	20%	5%
MO 1288	Sheep kidney	Total		0.93	1.000	AU	gen pop, > 2 yrs	112	87.63	NR	NR	1	1.216	1%	1%	-
MO 1289	Sheep liver	Total		3.9	1.000	AU	Child, 2-16 yrs	2	169.85	NR	NR	1	17.432	9%	5%	9%

Annex 4

BIXAFEN (262)										IESTI						
Acute RfD= 0.2 mg/kg bw (200 µg/kg bw)										Maximum %ARfD:			20%	20%	20%	
												all	gen pop	child		
Codex Code	Commodity	Processing	STMR or STMR-P mg/kg	HR or HR-P mg/kg	DCF	Country	Population group	n	Large portion, g/person	Unit weight, edible portion, g	Variability factor	Case	IESTI µg/kg bw/day	% acute RfD rounded	% acute RfD rounded	% acute RfD rounded
ML 0106	Milks	Total	0.082		1.000	PRIMO-UK	infant	P97.5	1080.70	NR	NR	3	10.186	5%	2%	5%
FM 0812	Cattle milk fat	Total	2.05		1.000	BR	Gen pop, > 10 yrs	441	150.00	NR	NR	3	4.763	2%	2%	2%
PM 0110	Poultry meat	Total	NA	NA	1.000	CN	Child, 1-6 yrs	175	347.00	NR	NR	1	NA	0%	0%	0%
PM 0110	Poultry meat: 10% as fat	Total		0.04	1.000	CN	Child, 1-6 yrs	175	34.70	NR	NR	1	0.086	0%	0%	0%
PM 0110	Poultry meat: 90% as muscle	Total		0	1.000	CN	Child, 1-6 yrs	175	312.30	NR	NR	1	0.000	0%	0%	0%
PF 0111	Poultry, fats	Total		0.04	1.000	CA	Child, <6 yrs	66	49.38	NR	NR	1	0.116	0%	0%	0%
PO 0111	Poultry, edible offal (includes kidney, liver and skin)	Total		0.03	1.000	CN	Gen pop, > 1 yrs	421	345.63	NR	NR	1	0.195	0%	0%	0%
PE 0112	Eggs	Total		0.047	1.000	PRIMO-UK	infant	P97.5	108.00	NR	NR	1	0.583	0%	0%	0%

CLOTHIANIDIN (238)										IESTI							
Acute RfD= 0.6 mg/kg bw (600 µg/kg bw)										Maximum %ARfD:				1%	1%	1%	
														all	gen pop	child	
Codex Code	Commodity	Processing	STMR or STMR-P mg/kg	HR or HR-P mg/kg	DCF	Country	Population group	n	Large portion, g/person	Unit weight, edible portion, g	Variability factor	Case	IESTI µg/kg bw/day	% acute RfD rounded	% acute RfD rounded	% acute RfD rounded	
FP 0307	Persimmon, Japanese (i.e. Kaki fruit) (all commodities)	highest utilisation: raw with peel (incl consumption without peel)	0	0.2	1.000	TH	Child, 3-6 yrs	20	264.88	228	3	2a	7.62 - 8.42	1% - 1%	0% - 1%	1% - 1%	
GC 0650	Rye (all commodities)	highest utilisation: flakes	0.01	0	1.000	CA	Child, <6 yrs	1909	539.23	NR	NR	3	0.01 - 0.34	0% - 0%	0% - 0%	0% - 0%	
GC 0653	Triticale	Total	0.01		1.000	DE	Gen pop, 14-80 yrs	27100	394.70	<25	NR	3	0.052	0%	0%	0%	
GC 0654	Wheat (all commodities)	highest utilisation: flakes	0.01	0	1.000	CA	Child, <6 yrs	1909	539.23	NR	NR	3	0.01 - 0.34	0% - 0%	0% - 0%	0% - 0%	
GC 0640	Barley (all commodities)	highest utilisation: beer	0.015	0	0.190	CA	Gen pop, all ages	2514	21271.20	NR	NR	3	0 - 0.77	0% - 0%	0% - 0%	0% - 0%	
GC 0647	Oats (all commodities)	highest utilisation: flakes (rolled oats)	0.015	0	1.000	CA	Child, <6 yrs	1909	539.23	NR	NR	3	0.01 - 0.51	0% - 0%	0% - 0%	0% - 0%	
GC 0649	Rice (all commodities)	highest utilisation: Total	0.0725 - 0.159	0	1.000	CA	Child, <6 yrs	666	461.40	<25	NR	3	0.01 - 4.38	0% - 1%	0% - 0%	0% - 1%	
GC 0651	Sorghum grain (Chicken corn, Dari seed, Durra, Feterita) (all commodities)	highest utilisation: cooked/boiled	0.0095 - 0.015	0	0.400	CN	Gen pop, > 1 yrs	356	1348.67	<25	NR	3	0.01 - 0.15	0% - 0%	0% - 0%	0% - 0%	
GC 3081	Baby corn (all commodities)	highest utilisation: canned/preserved (imm cobs)	0	0.01	1.000	NL	Child, 2-6 yrs	E	75.00	12	NR	1	0.04 - 0.04	0% - 0%	0% - 0%	0% - 0%	
GC 0447	Sweet corn (corn-on-the-cob) (kernels plus cob with husks removed) (all commodities)	highest utilisation: cooked/boiled (corn-on-the-cob)	0.01	0.01	1.000	TH	Child, 3-6 yrs	1383	196.99	191	3	2a	0.01 - 0.34	0% - 0%	0% - 0%	0% - 0%	
GC 1275	Sweet corn (whole kernel without cob or husk) (all commodities)	highest utilisation: canned/preserved (kernels)	0.01	0.01	1.000	CA	Child, <6 yrs	289	153.76	NR	NR	3	0.01 - 0.1	0% - 0%	0% - 0%	0% - 0%	
GS 0658	Sorgho or Sorghum, sweet	Total	0.062		1.000	CN	Child, 1-6 yrs	249	131.48	<25	NR	1	ND	-	-	-	
MM 0095	Meat from mammals other than marine mammals	Total		0.02	1.000	CN	Child, 1-6 yrs	302	264.84	NR	NR	1	0.328	0%	0%	0%	
MF 0100	Mammalian fats (except milk fats)	Total		0.02	1.000	PRIMO-FR	adult	P97.5	134.79	NR	NR	1	0.041	0%	0%	0%	

Annex 4

CLOTHIANIDIN (238)										IESTI						
Acute RfD= 0.6 mg/kg bw (600 µg/kg bw)										Maximum %ARfD:			1%	1%	1%	
												all	gen pop	child		
Codex Code	Commodity	Processing	STMR or STMR-P mg/kg	HR or HR-P mg/kg	DCF	Country	Population group	n	Large portion, g/person	Unit weight, edible portion, g	Variability factor	Case	IESTI µg/kg bw/day	% acute RfD rounded	% acute RfD rounded	% acute RfD rounded
MO 0105	Edible offal (mammalian)	Total		0.351	1.000	ZA	Gen pop, > 10 yrs	-	523.58	NR	NR	1	3.299	1%	1%	0%
ML 0106	Milks	Total	0.042		1.000	PRIMO-UK	infant	P97.5	1080.70	NR	NR	3	5.217	1%	0%	1%
PM 0110	Poultry meat	Total		0.0014	1.000	CN	Child, 1-6 yrs	175	347.00	NR	NR	1	0.030	0%	0%	0%
PM 0110	Poultry meat: 10% as fat	Total		0.0033	1.000	CN	Child, 1-6 yrs	175	34.70	NR	NR	1	0.007	0%	0%	0%
PM 0110	Poultry meat: 90% as muscle	Total		0.0014	1.000	CN	Child, 1-6 yrs	175	312.30	NR	NR	1	0.027	0%	0%	0%
PF 0111	Poultry, fats	Total		0.0033	1.000	CA	Child, <6 yrs	66	49.38	NR	NR	1	0.010	0%	0%	0%
PO 0111	Poultry, edible offal (includes kidney, liver and skin)	Total		0.37	1.000	CN	Gen pop, > 1 yrs	421	345.63	NR	NR	1	2.403	0%	0%	0%
PE 0112	Eggs	Total		0.0062	1.000	PRIMO-UK	infant	P97.5	108.00	NR	NR	1	0.077	0%	0%	0%

DIFENOCONAZOLE (224)											IESTI					
Acute RfD= 0.3 mg/kg bw (300 µg/kg bw)											Maximum %ARfD:			2%	2%	2%
														all	gen pop	child
Codex Code	Commodity	Processing	STMR or STMR-P mg/kg	HR or HR-P mg/kg	DCF	Country	Population group	n	Large portion, g/person	Unit weight, edible portion, g	Variability factor	Case	IESTI µg/kg bw/day	% acute RfD rounded	% acute RfD rounded	% acute RfD rounded
FB 0265	Cranberry (all commodities)	highest utilisation: Total	0.2	0.26	1.000	AU	Child, 2-16 yrs	103	279.66	<25	NR	1	0 - 1.91	0% - 1%	0% - 0%	0% - 1%
FT 0336	Guava (all commodities)	highest utilisation: raw with peel	0	0.095	1.000	BR	Gen pop, > 10 yrs	398	850.00	170	3	2a	0.95 - 1.75	0% - 1%	0% - 1%	1% - 1%
FT 0337	Guava	canned/preserved	0.0335		1.000	NL	Child, 2-6 yrs	E	30.89	NR	NR	3	0.056	0%	0%	0%
FT 0338	Guava	juice (pasteurised)	0.0335		1.000	BR	Gen pop, > 10 yrs	1210	720.00	NR	NR	3	0.374	0%	0%	0%
FT 0339	Guava	canned babyfood	0.0335		1.000	NL	toddler, 8-20 m	43	7.76	NR	NR	3	0.025	0%	NC	0%
FT 0340	Guava	composite foods; unspecified ind processed	0.0335		1.000	NL	Child, 2-6 yrs	103	1.69	NR	NR	3	0.003	0%	0%	0%
SO 0691	Cotton seed (all commodities)	highest utilisation: Oil (refined)	0.0014 - 0.021	0	1.000	US	Child, < 6 yrs	6354	3.13	NR	NR	3	0 - 0	0% - 0%	0% - 0%	0% - 0%
DT 1114	Tea, green, black (black, fermented and dried) (all commodities)	highest utilisation: Total	0.0072 - 4.85	0	1.000	PRIMO-IE	child	P97.5	30.60	<25	NR	3	0.01 - 7.42	0% - 2%	0% - 2%	2% - 2%

Annex 4

ETHIPROLE (304)											IESTI					
Acute RfD= 0.005 mg/kg bw (5 µg/kg bw)											Maximum %ARfD:					
											2%		1%		2%	
											all		gen pop		child	
Codex Code	Commodity	Processing	STMR or STMR-P mg/kg	HR or HR-P mg/kg	DCF	Country	Population group	n	Large portion, g/person	Unit weight, edible portion, g	Variability factor	Case	IESTI µg/kg bw/day	% acute RfD rounded	% acute RfD rounded	% acute RfD rounded
VD 0541	Soya bean (dry) (Glycine spp) (all commodities)	highest utilisation: Total	0.0007 - 0.0065		1.000	CN	Child, 1-6 yrs	179	239.05	<25	NR	3	0 - 0.1	0% - 2%	0% - 1%	0% - 2%

FENBUCONAZOLE (197)											IESTI					
Acute RfD= 0.2 mg/kg bw (200 µg/kg bw)											Maximum %ARfD:					
											3%		3%		3%	
											all		gen pop		child	
Codex Code	Commodity	Processing	STMR or STMR-P mg/kg	HR or HR-P mg/kg	DCF	Country	Population group	n	Large portion, g/person	Unit weight, edible portion, g	Variability factor	Case	IESTI µg/kg bw/day	% acute RfD rounded	% acute RfD rounded	% acute RfD rounded
DT 1114	Tea, green, black (black, fermented and dried) (all commodities)	highest utilisation: Total	0.0018 - 4.2		1.000	PRIMO-IE	child	P97.5	NR				0 - 0	0% - 0%	0% - 3%	2% - 3%

FLUOPYRAM (243)											IESTI					
Acute RfD= 0.5 mg/kg bw (500 µg/kg bw)											Maximum %ARfD:			0%	0%	0%
														all	gen pop	child
Codex Code	Commodity	Processing	STMR or STMR-P mg/kg	HR or HR-P mg/kg	DCF	Country Population n			Large portion, g/person	Unit weight, edible portion, g	Variability factor	Case	IESTI µg/kg bw/day	% acute RfD rounded	% acute RfD rounded	% acute RfD rounded
SM 0716	Coffee bean (all commodities)	highest utilisation: extract (beverage)	0.01	0	0.180	CA	women, 15-49 yrs	2666	2088.65	NR	NR	3	0 - 0.05	0% - 0%	0% - 0%	0% - 0%

IMAZALIL (110)										IESTI						
Acute RfD= 0.05 mg/kg bw (50 µg/kg bw)										Maximum %ARfD:				50%	50%	40%
													all	gen pop	child	
Codex Code	Commodity	Processing	STMR or STMR-P mg/kg	HR or HR-P mg/kg	DCF	Country	Population group	n	Large portion, g/person	Unit weight, edible portion, g	Variability factor	Case	IESTI µg/kg bw/day	% acute RfD rounded	% acute RfD rounded	% acute RfD rounded
FC 0303	Kumquats (all commodities)	highest utilisation: Total	0	9.9	1.000	JP	Gen pop, > 1 yrs	135	120.00	<25	NR	1	1.8 - 23.67	4% - 50%	4% - 50%	40% - 40%
FC 0204	Lemon (all commodities)	highest utilisation: Total	0.07 - 0.92	0.36	1.000	PRIMO-DE	child	P95	125.50	71	3	2a	0.04 - 5.98	0% - 10%	0% - 4%	0% - 10%
FC 0205	Lime (all commodities)	highest utilisation: Total	0.07 - 0.34	0.36	1.000	AU	Gen pop, > 2 yrs	579	259.21	49	3	2a	0.01 - 1.92	0% - 4%	0% - 4%	0% - 2%
FC 0003	Subgroup of Mandarins (incl mandarin-like hybrids) (all commodities)	highest utilisation: raw, without peel	0.07 - 0.92	0.36	1.000	CN	Child, 1-6 yrs	151	586.75	124	3	2a	0.01 - 18.64	0% - 40%	0% - 20%	0% - 40%
FC 0004	Subgroup of Oranges, sweet, sour (incl orange-like hybrids) (all commodities)	highest utilisation: Total	0.07 - 0.92	0.36	1.000	AU	Child, 2-6 yrs	1735	800.83	156	3	2a	0.05 - 21.08	0% - 40%	0% - 20%	0% - 40%
FC 0005	Subgroup of Pummelo and Grapefruits (incl Shaddock-like hybrids, among others Grapefruit) (all commodities)	highest utilisation: Total	0.07 - 0.34	0.36	1.000	PRIMO-DE	child	P90	253.56	270	3	2b	0.01 - 16.96	0% - 30%	0% - 20%	0% - 30%
HS 3381	Lemon, peel	Oil (refined)	97		1.000	NL	Gen pop, > 1 yrs	0	NC	NR	NR	3	NC	NC	NC	NC

Annex 4

IMAZALIL (110)										IESTI						
Acute RfD= 0.05 mg/kg bw (50 µg/kg bw)										Maximum %ARfD:						
										all			gen pop		child	
Codex Code	Commodity	Processing	STMR or STMR-P mg/kg	HR or HR-P mg/kg	DCF	Country	Population group	n	Large portion, g/person	Unit weight, edible portion, g	Variability factor	Case	IESTI µg/kg bw/day	% acute RfD rounded	% acute RfD rounded	% acute RfD rounded
HS 3382	Orange, peel	Oil (refined)	97		1.000	NL	Gen pop, > 1 yrs	0	NC	NR	NR	3	NC	NC	NC	NC

METHOXYFENOZIDE (209)										IESTI						
Acute RfD= 0.9 mg/kg bw (900 µg/kg bw)										Maximum %ARfD:						
										all			gen pop		child	
Codex Code	Commodity	Processing	STMR or STMR-P mg/kg	HR or HR-P mg/kg	DCF	Country	Population group	n	Large portion, g/person	Unit weight, edible portion, g	Variability factor	Case	IESTI µg/kg bw/day	% acute RfD rounded	% acute RfD rounded	% acute RfD rounded
GS 0659	Sugar cane (all commodities)	highest utilisation: sugar	0.01 - 0.07	0.01	1.000	PRIMO-NL	child	E	168.90	NR	NR	3	0.02 - 0.55	0% - 0%	0% - 0%	0% - 0%
SM 0716	Coffee bean	raw incl roasted	0.028		1.000	NL	Gen pop, > 1 yrs	0	NC	0	NR	3	NC	NC	NC	NC
HH 0722	Basil, leaves (all commodities)	highest utilisation: raw	19 - 100	47 - 194	1.000	DE	Child, 2-4 yrs	138	11.10	<25	NR	1	5.38 - 32.3	1% - 4%	0% - 1%	1% - 4%
DT 1114	Tea, green, black (black, fermented and dried) (all commodities)	highest utilisation: raw = dried	0.14 - 28.5		1.000	CN	Gen pop, > 1 yrs	679	75.88	<25	NR	3	5.72 - 40.63	1% - 5%	1% - 5%	3% - 3%

PROTHIOCONAZOLE (232) - Acute dietary exposure for the general population and children													IEST I			
Acute RfD= 1 mg/kg bw (1000 µg/kg bw)													Maximum %ARfD:			
ARfD is for prothioconazole-desthio													all gen pop child			
Codex Code	Commodity	Processing	STMR or STMR-P mg/kg	HR or HR-P mg/kg	DCF P mg/kg	Country	Population group	n	Large portion, g/person	Unit weight, edible portion, g	Variability factor	Case	IESTI µg/kg bw/day	% acute RfD rounded	% acute RfD rounded	% acute RfD rounded
SO 0693	Linseed (Flax-seed)	Total	0.014		1.000	CA	Gen pop, all ages	291	81.08	<25	NR	3	0.016	0%	0%	0%
SO 0495	Rape seed (Canola) (all commodities)	highest utilisation: Oil (refined)	0.0273 - 0.039		1.000	CA	Child, <6 yrs	1127	26.46	NR	NR	3	0.03 - 0.04	0% - 0%	0% - 0%	0% - 0%
SO 0699	Safflower seed (all commodities)	highest utilisation: Total	0.009		1.000	PRIM O-DE	child	P95	49.74	<25	NR	3	0 - 0.03	0% - 0%	0% - 0%	0% - 0%
SO 0702	Sunflower seed (all commodities)	highest utilisation: Oil (refined)	0.009 - 0.00963		1.000	PRIM O-NL	child	P97.5	21.60	NR	NR	3	0 - 0.01	0% - 0%	0% - 0%	0% - 0%
MM 0095	Meat from mammals other than marine mammals	Total	NA	NA	1.000	CN	Child, 1-6 yrs	302	264.84	NR	NR	1	NA	0%	0%	0%
MM 0095	Meat from mammals other than marine mammals: 20% as fat	Total		0.023	1.000	CN	Child, 1-6 yrs	302	52.97	NR	NR	1	0.076	0%	0%	0%
MM 0095	Meat from mammals other than marine mammals: 80% as muscle	Total		0.007	1.000	CN	Child, 1-6 yrs	302	211.87	NR	NR	1	0.092	0%	0%	0%
MF 0100	Mammalian fats (except milk fats)	Total		0.023	1.000	PRIM O-FR	adult	P97.5	134.79	NR	NR	1	0.047	0%	0%	0%
MO 0105	Edible offal (mammalian)	Total		0.291	1.000	ZA	Gen pop, > 10 yrs	-	523.58	NR	NR	1	2.735	0%	0%	0%
ML 0106	Milks	Total	0.001		1.000	PRIM O-UK	infant	P97.5	1080.70	NR	NR	3	0.124	0%	0%	0%
PM 0110	Poultry meat	Total	NA	NA	1.000	CN	Child, 1-6 yrs	175	347.00	NR	NR	1	NA	0%	0%	0%
PM 0110	Poultry meat: 10% as fat	Total		0.0129	1.000	CN	Child, 1-6 yrs	175	34.70	NR	NR	1	0.028	0%	0%	0%
PM 0110	Poultry meat: 90% as muscle	Total		0.00255	1.000	CN	Child, 1-6 yrs	175	312.30	NR	NR	1	0.049	0%	0%	0%
PF 0111	Poultry, fats	Total		0.0129	1.000	CA	Child, <6 yrs	66	49.38	NR	NR	1	0.037	0%	0%	0%
PO 0111	Poultry, edible offal (includes kidney, liver and skin)	Total		0.118	1.000	CN	Gen pop, > 1 yrs	421	345.63	NR	NR	1	0.766	0%	0%	0%
PE 0112	Eggs	Total		0.00103	1.000	PRIM O-UK	infant	P97.5	108.00	NR	NR	1	0.013	0%	0%	0%

Annex 4

Acute dietary exposure for women of child bearing age

PROTHIOCONAZOLE (232)														
Acute RfD= 0.01 mg/kg bw (10 µg/kg bw)														
Maximum %ARfD: 8%														
ARfD is for prothioconazole-destho														
women														
Codex Code	Commodity	Processing	STMR or STMR-P mg/kg	HR or HR-P mg/kg	DCF	Country	Population group	n	Large portion, g/person	Unit weight, edible portion, g	Varia-bility factor	Case	IESTI µg/kg bw/day	% acute RfD rounded
SO 0693	Linseed (Flax-seed)	Total	0.014		1.000	CA	women, 15-49 yrs	38	64.29	<25	NR	3	0.012	0%
SO 0495	Rape seed (Canola) (all commodities)	highest utilisation: Oil (refined)	0.0273 - 0.039		1.000	CA	women, 15-49 yrs	2612	44.97	NR	NR	3	0.02 - 0.02	0% - 0%
SO 0699	Safflower seed (all commodities)	highest utilisation: Total	0.009		1.000	PRIM O-DE	women, 14-50 yrs	P97.5	28.07	<25	NR	3	0 - 0	0% - 0%
SO 0702	Sunflower seed (all commodities)	highest utilisation: raw	0.009 - 0.00963		1.000	NL	gen pop, > 1 yrs	369	39.41	<25	NR	3	0 - 0.01	0% - 0%
MM 0095	Meat from mammals other than marine mammals	Total	NA	NA	1.000	DE	Women, 14-50 yrs	25	521.10	NR	NR	1	NA	1%
MM 0095	Meat from mammals other than marine mammals: 20% as fat	Total		0.023	1.000	DE	Women, 14-50 yrs	25	104.22	NR	NR	1	0.036	0%
MM 0095	Meat from mammals other than marine mammals: 80% as muscle	Total		0.007	1.000	DE	Women, 14-50 yrs	25	416.88	NR	NR	1	0.043	0%
MF 0100	Mammalian fats (except milk fats)	Total		0.023	1.000	US	women, 13-49 yrs	6730	45.60	NR	NR	1	0.014	0%
MO 0105	Edible offal (mammalian)	Total		0.291	1.000	PRIM O-DE	women, 14-50 yrs	P100	188.92	NR	NR	1	0.815	8%
ML 0106	Milks	Total	0.001		1.000	DE	Women, 14-50 yrs	6554	1276.50	NR	NR	3	0.019	0%
PM 0110	Poultry meat	Total	NA	NA	1.000	CA	women, 15-49 yrs	2127	384.47	NR	NR	1	NA	0%
PM 0110	Poultry meat: 10% as fat	Total		0.0129	1.000	CA	women, 15-49 yrs	2127	38.45	NR	NR	1	0.007	0%
PM 0110	Poultry meat: 90% as muscle	Total		0.00255	1.000	CA	women, 15-49 yrs	2127	346.02	NR	NR	1	0.013	0%
PF 0111	Poultry, fats	Total		0.0129	1.000	CA	women, 15-49 yrs	195	78.63	NR	NR	1	0.015	0%
PO 0111	Poultry, edible offal (includes kidney, liver and skin)	Total		0.118	1.000	CN	Gen pop, > 1 yrs	421	345.63	NR	NR	1	0.766	8%

PROTHIOCONAZOLE (232) Acute RfD= 0.01 mg/kg bw (10 µg/kg bw) ARfD is for prothioconazole-desthio														
IESTI												Maximum %ARfD:		
												8%		
												women		
Codex Code	Commodity	Processing	STMR or STMR-P mg/kg	HR or HR-P mg/kg	DCF	Country	Population group	n	Large portion, g/person	Unit weight, edible portion, g	Varia-bility factor	Case	IESTI µg/kg bw/day	% acute RfD rounded
PE 0112	Eggs	Total		0.00103	1.000	CA	women, 15-49 yrs	3395	136.90	NR	NR	1	0.002	0%

Annex 4

PYDIFLUMETOFEN (309)														
Acute RfD= 0.3 mg/kg bw (300 µg/kg bw)														
IESTI														
Maximum %ARfD:														
Codex Code	Commodity	Processing	STMR or STMR-P mg/kg	HR or HR-P mg/kg	DCF	Country	Population group	n	Large portion, g/person	Unit weight, edible portion, g	Variability factor	Case	IESTI µg/kg bw/day	all
														% acute RfD rounded
														20%
														% acute RfD rounded
														20%
														% acute RfD rounded
														20%
														% acute RfD rounded
FC 0303	Kumquats (all commodities)	highest utilisation: Total	0	0.76	1.000	JP	Gen pop, > 1 yrs	135	120.00	<25	NR	1	0.14 - 1.82	0% - 1%
FC 0204	Lemon (all commodities)	highest utilisation: Total	0.004 - 0.21	0.76	1.000	PRIMO-DE	child	P95	125.50	71	3	2a	0.01 - 12.62	0% - 4%
FC 0205	Lime (all commodities)	highest utilisation: Total	0.004 - 0.21	0.76	1.000	AU	Gen pop, > 2 yrs	579	259.21	49	3	2a	0.01 - 4.05	0% - 1%
FC 0003	Subgroup of Mandarins (incl mandarin-like hybrids) (all commodities)	highest utilisation: raw, without peel	0.004 - 0.21	0.76	1.000	CN	Child, 1-6 yrs	151	586.75	124	3	2a	0.04 - 39.35	0% - 10%
FC 0004	Subgroup of Oranges, sweet, sour (incl orange-like hybrids) (all commodities)	highest utilisation: Total	0.004 - 0.21	0.76	1.000	AU	Child, 2-6 yrs	1735	800.83	156	3	2a	0.14 - 44.5	0% - 10%
FC 0005	Subgroup of Pummelo and Grapefruits (incl Shaddock-like hybrids, among others Grapefruit) (all commodities)	highest utilisation: Total	0.004 - 0.21	0.76	1.000	PRIMO-DE	child	P90	253.56	270	3	2b	0.04 - 35.8	0% - 10%
FP 0226	Apple (all commodities)	highest utilisation: raw with peel (incl consumption without peel)	0.002 - 0.06	0.046 - 0.13	1.000	CN	Child, 1-6 yrs	1314	403.39	255	3	2a	0.04 - 7.36	0% - 2%
FP 2220	Azarole (Mediterranean medlar) (all commodities)	highest utilisation: juice (pasteurised)	0.06	0.13	1.000	PRIMO-DE	child	P97.5	89.63	NR	NR	3	0.12 - 0.33	0% - 0%
FP 0227	Crab-apple (all commodities)	highest utilisation: raw with peel	0	0.13	1.000	CN	Gen pop, > 1 yrs	204	488.33	<25	NR	1	1.19 - 1.19	0% - 0%
FP 0228	Loquat (Japanese medlar) (all commodities)	highest utilisation: raw without peel	0	0.13	1.000	JP	Gen pop, > 1 yrs	113	326.40	49	3	2a	0.23 - 1.02	0% - 0%
FP 0229	Medlar (all commodities)	highest utilisation: Total	0	0.13	1.000	PRIMO-ES	child	P97.5	116.99	60	3	2a	0.89 - 0.89	0% - 0%
FP 0230	Pear (all commodities)	highest utilisation: Total	0.004 - 0.06	0.08 - 0.13	1.000	CA	Child, <6 yrs	175	498.28	255	3	2a	0.02 - 9	0% - 3%

PYDIFLUMETOFEN (309)										IESTI						
Acute RfD= 0.3 mg/kg bw (300 µg/kg bw)										Maximum %ARfD:				20%	20%	20%
Codex Code	Commodity	Processing	STMR or STMR-P mg/kg	HR or HR-P mg/kg	DCF	Country	Population group	n	Large portion, g/person	Unit weight, edible portion, g	Variability factor	Case	IESTI µg/kg bw/day	% acute RfD rounded	% acute RfD rounded	% acute RfD rounded
FP 0231	Quince (all commodities)	highest utilisation: Total	0.06	0.13	1.000	PRIMO-ES	child	P97.5	169.60	301	3	2b	0 - 1.92	0% - 1%	0% - 0%	0% - 1%
FS 0013	Subgroup of Cherries (all commodities)	highest utilisation: Total	0.395	1.7	1.000	PRIMO-DK	child	P97.5	269.00	<25	NR	1	0.16 - 20.79	0% - 7%	0% - 6%	0% - 7%
FS 0014	Subgroup of Plums (all commodities)	highest utilisation: dried (prunes)	0.15 - 0.34	0.37 - 0.85	1.000	AU	Child, 2-6 yrs	13	447.59	10	NR	1	0.07 - 20.02	0% - 7%	0% - 2%	0% - 7%
FS 0240	Apricot (all commodities)	highest utilisation: Total	0.21	0.8	1.000	PRIMO-DE	child	P95	264.86	50	3	2a	0.09 - 18.07	0% - 6%	0% - 5%	0% - 6%
FS 2237	Japanese apricot (ume)	Total		0.8	1.000	JP	Child, 1-6 yrs	25	25.50	<25	NR	1	1.127	0%	0%	0%
FS 0245	Nectarine (all commodities)	highest utilisation: raw with peel (incl consumption without peel)	0.21	0.8	1.000	NL	toddler, 8-20 m	6	183.60	131	3	2a	0.09 - 34.94	0% - 10%	0% - 3%	0% - 10%
FS 0247	Peach (all commodities)	highest utilisation: raw with peel (incl consumption without peel)	0.21	0.8	1.000	JP	Child, 1-6 yrs	76	306.00	255	3	2a	0.09 - 42.12	0% - 10%	0% - 5%	0% - 10%
FB 0020	Blueberries (all commodities)	highest utilisation: Total	0.88	3.9	1.000	CA	Child, <6 yrs	189	176.21	<25	NR	1	0.13 - 44.64	0% - 10%	0% - 10%	0% - 10%
FB 0021	Currants, black, red, white (all commodities)	highest utilisation: Total	0.88	3.9	1.000	AU	Gen pop, > 2 yrs	322	797.60	<25	NR	1	0.57 - 46.43	0% - 20%	0% - 20%	0% - 10%
FB 0268	Gooseberry (all commodities)	highest utilisation: Total	0.88	3.9	1.000	PRIMO-DE	child	P100	94.96	<25	NR	1	0.17 - 22.93	0% - 8%	0% - 7%	0% - 8%
FB 0273	Rose hips (all commodities)	highest utilisation: Total	0.88	3.9	1.000	PRIMO-FI	women	P97.5	156.60	<25	NR	1	0.56 - 8.58	0% - 3%	0% - 3%	0% - 1%
FB 0267	Elderberries (all commodities)	highest utilisation: Total	0.88	3.9	1.000	CN	Gen pop, > 1 yrs	136	420.22	29	3	2a	0.6 - 35.04	0% - 10%	0% - 10%	0% - 5%
FB 0275	Strawberry (all commodities)	highest utilisation: Raw with skin	0.185	0.62	1.000	NL	toddler, 8-20 m	52	166.73	18	NR	1	0.1 - 10.13	0% - 3%	0% - 2%	0% - 3%
VA 0381	Garlic (all commodities)	highest utilisation: raw without skin	0.07	0.2	1.000	CN	Child, 1-6 yrs	290	174.44	62	3	2a	0 - 3.71	0% - 1%	0% - 1%	0% - 1%
VA 0385	Onion, bulb (all commodities)	highest utilisation: raw without skin	0.07	0.2	1.000	JP	Child, 1-6 yrs	748	102.00	244	3	2b	0.03 - 3.73	0% - 1%	0% - 1%	0% - 1%
VA 0386	Onion, Chinese (all commodities)	highest utilisation: raw	0	0.2	1.000	CN	Child, 1-6 yrs	196	136.53	130	3	2a	0.82 - 4.91	0% - 2%	0% - 1%	0% - 2%
VA 0388	Shallot (all commodities)	highest utilisation: raw without skin	0.07	0.2	1.000	CN	Child, 1-6 yrs	480	115.81	51	3	2a	0.06 - 2.71	0% - 1%	0% - 0%	0% - 1%

Annex 4

PYDIFLUMETOFEN (309)										IESTI						
Acute RfD= 0.3 mg/kg bw (300 µg/kg bw)										Maximum %ARfD:				20%	20%	20%
Codex Code	Commodity	Processing	STMR or STMR-P mg/kg	HR or HR-P mg/kg	DCF	Country	Population group	n	Large portion, g/person	Unit weight, edible portion, g	Variability factor	Case	IESTI µg/kg bw/day	% acute RfD rounded	% acute RfD rounded	% acute RfD rounded
VA 2605	Chives (all commodities)	highest utilisation: Total	0	1.39	1.000	PRIMO-CZ	Child, child, 7-10 yrs	P97.5	26.49	<25	NR	1	0.26 - 1.14	0% - 0%	0% - 0%	0% - 0%
VA 0384	Leek (all commodities)	highest utilisation: cooked/boiled	0.36	1.39	1.000	PRIMO-NL	toddler	E	100.70	142	3	2b	0.04 - 41.17	0% - 10%	0% - 6%	0% - 10%
VA 0387	Onion, Welsh (Japanese bunching onion, Multiplying onion) (all commodities)	highest utilisation: Total	0	1.39	1.000	PRIMO-UK	child, 4-6 yrs	P97.5	93.60	38	3	2a	8.76 - 11.5	3% - 4%	1% - 2%	3% - 4%
VA 0389	Spring onion (all commodities)	highest utilisation: Total	0	1.39	1.000	PRIMO-UK	child, 4-6 yrs	P97.5	93.60	38	3	2a	4.3 - 11.5	1% - 4%	1% - 1%	1% - 4%
VB 0400	Broccoli (all commodities)	highest utilisation: cooked/boiled	0.39	1.5	1.000	PRIMO-NL	toddler	P97.5	160.70	286	3	2b	0.71 - 70.9	0% - 20%	0% - 8%	0% - 20%
VB 0404	Cauliflower (all commodities)	highest utilisation: cooked/boiled	0.39	1.5	1.000	PRIMO-NL	toddler	P97.5	142.00	749	3	2b	0.06 - 62.65	0% - 20%	0% - 10%	0% - 20%
VB 0402	Brussels sprouts (all commodities)	highest utilisation: cooked/boiled	0.065	0.22	1.000	PRIMO-NL	toddler	P90	103.80	<25	NR	1	0 - 2.24	0% - 1%	0% - 1%	0% - 1%
VB 0041	Cabbages, head (all commodities)	highest utilisation: raw	0.065	0.22	1.000	CN	Child, 1-6 yrs	287	255.54	1403	3	2b	0.05 - 10.45	0% - 3%	0% - 2%	0% - 3%
VB 0467	Chinese cabbage (type Pe-tsai) (all commodities)	highest utilisation: Total	0.065	0.22	1.000	CN	Child, 1-6 yrs	2788	336.16	2162	3	2b	0.02 - 13.75	0% - 5%	0% - 3%	0% - 5%
VB 0405	Kohlrabi (all commodities)	highest utilisation: Total	0	0.09	1.000	PRIMO-DE	child	P95	167.96	265	3	2b	0.28 - 2.81	0% - 1%	0% - 0%	1% - 1%
VP 0061	Beans with pods (Phaseolus spp): (immature pods + succulent seeds) (all commodities)	highest utilisation: Total	0.045	0.47	1.000	CA	Child, <6 yrs	261	203.31	<25	NR	1	0.07 - 6.35	0% - 2%	0% - 1%	0% - 2%
VP 0522	Broad bean with pods (immature pods + succulent seeds) (Vicia spp) (all commodities)	highest utilisation: Total	0	0.47	1.000	US	Child, < 6 yrs	221	93.96	9	NR	1	1.3 - 3.05	0% - 1%	0% - 0%	0% - 1%
VP 0542	Sword bean with pods (immature pods + succulent seeds) (Canavalia spp)	highest utilisation: cooked/boiled	0	0.47	1.000	CN	Gen pop, > 1 yrs	891	316.83	<25	NR	1	2.8 - 2.8	1% - 1%	1% - 1%	0% - 0%

PYDIFLUMETOFEN (309)										IESTI						
Acute RfD= 0.3 mg/kg bw (300 µg/kg bw)										Maximum %ARfD:				20%	20%	20%
Codex Code	Commodity	Processing	STMR or STMR-P mg/kg	HR or HR-P mg/kg	DCF	Country	Population group	n	Large portion, g/person	Unit weight, edible portion, g	Variability factor	Case	IESTI µg/kg bw/day	all % acute RfD rounded	gen pop % acute RfD rounded	child % acute RfD rounded
	(all commodities)															
VP 0063	Peas with pods (Pisum spp) immature pods + succulent seeds) (all commodities)	highest utilisation: cooked/boiled	0.12	0.84	1.000	CN	Child, 1-6 yrs	1056	290.21	6	NR	1	4.12 - 15.11	1% - 5%	1% - 3%	1% - 5%
VP 0553	Lentil with pods (immature pods + succulent seeds) (Lens spp) (all commodities)	highest utilisation: cooked/boiled	0	0.84	1.000	CN	Child, 1-6 yrs	371	345.76	<25	NR	1	4.83 - 18	2% - 6%	1% - 3%	2% - 6%
VP 0062	Beans without pods: (Phaseolus spp.) (succulent seeds) (all commodities)	highest utilisation: Total	0.033	0.099	1.000	PRIMO-IE	child	P97.5	157.79	<25	NR	1	0 - 0.78	0% - 0%	0% - 0%	0% - 0%
VP 0523	Broad bean without pods (succulent seeds) (Vicia spp) (all commodities)	highest utilisation: frozen	0.033	0.099	1.000	NL	Child, 2-6 yrs	E	100.00	6	NR	1	0 - 0.54	0% - 0%	0% - 0%	0% - 0%
VP 0064	Peas without pods (Pisum spp) (succulent seeds) (all commodities)	highest utilisation: Total	0.031	0.042	1.000	PRIMO-UK	infant	P97.5	71.30	<25	NR	1	0.03 - 0.34	0% - 0%	0% - 0%	0% - 0%
VP 0520	Bambara groundnut without pods (succulent seeds) (Vigna subterranea)	Total		0.02	1.000	AU	Gen pop, > 2 yrs	22	186.07	<25	NR	1	0.056	0%	0%	-
VR 0574	Beetroot (all commodities)	highest utilisation: Total	0.08	0.25	1.000	AU	Child, 2-6 yrs	53	314.08	135	3	2a	0.05 - 7.7	0% - 3%	0% - 1%	0% - 3%
VR 0575	Burdock, greater or edible	Total		0.25	1.000	JP	Child, 1-6 yrs	122	35.70	68	3	2b	1.633	1%	0%	1%
VR 0577	Carrot (all commodities)	highest utilisation: raw with skin	0.08	0.25	1.000	CN	Child, 1-6 yrs	400	234.68	300	3	2b	0.01 - 10.91	0% - 4%	0% - 1%	0% - 4%
VR 0578	Celeriac (Turnip rooted celery) (all commodities)	highest utilisation: Total	0.08	0.25	1.000	PRIMO-BE	toddler	P100	196.90	749	3	2b	0.01 - 8.3	0% - 3%	0% - 1%	0% - 3%
VR 0469	Chicory, roots (all commodities)	highest utilisation: Total	0.08	0.25	1.000	AU	Gen pop, > 2 yrs	175	26.16	48	3	2b	0.06 - 0.29	0% - 0%	0% - 0%	0% - 0%

Annex 4

PYDIFLUMETOFEN (309)										IESTI						
Acute RfD= 0.3 mg/kg bw (300 µg/kg bw)										Maximum %ARfD:				20%	20%	20%
Codex Code	Commodity	Processing	STMR or STMR-P mg/kg	HR or HR-P mg/kg	DCF	Country	Population group	n	Large portion, g/person	Unit weight, edible portion, g	Variability factor	Case	IESTI µg/kg bw/day	all % acute RfD rounded	gen pop % acute RfD rounded	child % acute RfD rounded
VR 0583	Horseradish (all commodities)	highest utilisation: Total	0	0.25	1.000	PRIMO-DE	Gen pop	P97.5	79.50	220	3	2b	0.01 - 0.78	0% - 0%	0% - 0%	0% - 0%
VR 0587	Parsley, turnip-rooted (Hamburg roots) (all commodities)	highest utilisation: Total	0.08	0.25	1.000	PRIMO-NL	Gen pop	P97.5	96.60	140	3	2b	0.56 - 1.1	0% - 0%	0% - 0%	0% - 0%
VR 0588	Parsnip (all commodities)	highest utilisation: cooked/boiled (without skin)	0.08	0.25	1.000	PRIMO-NL	child	E	133.30	227	3	2b	0.57 - 5.43	0% - 2%	0% - 1%	1% - 2%
VR 0494	Radish (all commodities)	highest utilisation: Total	0.08	0.25	1.000	PRIMO-NL	child	E	64.40	172	3	2b	0.01 - 2.63	0% - 1%	0% - 0%	0% - 1%
VR 0590	Radish, black (all commodities)	highest utilisation: Total	0.08	0.25	1.000	PRIMO-NL	child	E	64.40	172	3	2b	0.01 - 2.63	0% - 1%	0% - 0%	0% - 1%
VR 0591	Radish, Japanese (Chinese radish, Daikon) (all commodities)	highest utilisation: raw without skin	0.08	0.25	1.000	CN	Child, 1-6 yrs	1187	293.37	1000	3	2b	0.01 - 13.64	0% - 5%	0% - 2%	0% - 5%
VR 0498	Salsify (Oyster plant) (all commodities)	highest utilisation: Total	0.08	0.25	1.000	PRIMO-BE	toddler	P100	99.90	75	3	2a	0 - 3.52	0% - 1%	0% - 0%	0% - 1%
VR 0596	Sugar beet (all commodities)	highest utilisation: composite foods; unspecified ind processed	0.004 - 0.08	0	1.000	NL	Child, 2-6 yrs	2554	168.93	NR	NR	3	0.04 - 0.73	0% - 0%	0% - 0%	0% - 0%
VR 0497	Swede (Rutabaga) (all commodities)	highest utilisation: Total	0.08	0.25	1.000	PRIMO-UK	infant	P97.5	90.00	500	3	2b	0 - 7.76	0% - 3%	0% - 2%	0% - 3%
VR 0506	Turnip, garden (all commodities)	highest utilisation: Total	0.08	0.25	1.000	US	Gen pop, 0-85 yrs	53	1533.22	105	3	2a	0.59 - 6.16	0% - 2%	0% - 2%	0% - 1%
GC 0644	Job's tears (all commodities)	highest utilisation: cooked/boiled	0.03	0.03	0.300	TH	Child, 3-6 yrs	134	85.50	<25	NR	3	0.05 - 0.05	0% - 0%	0% - 0%	0% - 0%
GC 0646	Millet (common millet, proso millet) (all commodities)	highest utilisation: Total	0.03	0	1.000	CN	Child, 1-6 yrs	826	219.53	<25	NR	3	0.02 - 0.41	0% - 0%	0% - 0%	0% - 0%
GC 0651	Sorghum grain (Chicken corn, Dari seed, Durra, Feterita) (all commodities)	highest utilisation: cooked/boiled	0.515	0	0.400	CN	Gen pop, > 1 yrs	356	1348.67	<25	NR	3	0.81 - 5.22	0% - 2%	0% - 2%	0% - 1%
TN 0660	Almonds (all commodities)	highest utilisation: Total	0.003 - 0.01	0.01 - 0.03	1.000	CA	Child, <6 yrs	62	63.32	<25	NR	1	0 - 0.12	0% - 0%	0% - 0%	0% - 0%
TN 0662	Brazil nut (all commodities)	highest utilisation: Total	0	0.03	1.000	PRIMO-UK	child, 4-6 yrs	P97.5	17.80	<25	NR	1	0.02 - 0.03	0% - 0%	0% - 0%	0% - 0%

PYDIFLUMETOFEN (309)															
Acute RfD= 0.3 mg/kg bw (300 µg/kg bw)															
IESTI															
Maximum %ARfD:															
Codex Code	Commodity	Processing	STMR or STMR-P mg/kg	HR or HR-P mg/kg	DCF	Country	Population group	n	Large portion, g/person	Unit weight, edible portion, g	Variability factor	Case	IESTI µg/kg bw/day	all	20%
														% acute RfD rounded	% acute RfD rounded
TN 0295	Cashew nut (all commodities)	highest utilisation: raw incl roasted	0.01	0.03	1.000	TH	child, 3-6 yrs	374	98.84	<25	NR	1	0.08 - 0.17	0% - 0%	0% - 0%
TN 0664	Chestnut (all commodities)	highest utilisation: Total	0	0.03	1.000	CN	Gen pop, > 1 yrs	807	475.25	<25	NR	1	0.07 - 0.27	0% - 0%	0% - 0%
TN 0665	Coconut (all commodities)	highest utilisation: raw (i.e. nutmeat)	0.01	0.03	1.000	TH	child, 3-6 yrs	826	423.40	383	3	2a	0.01 - 2.09	0% - 1%	0% - 1%
TN 0666	Hazelnut (all commodities)	highest utilisation: Total	0.01	0.03	1.000	PRIMO-IE	child	P97.5	65.42	<25	NR	1	0.01 - 0.1	0% - 0%	0% - 0%
TN 0669	Macadamia nut (all commodities)	highest utilisation: Total	0.01	0.03	1.000	PRIMO-DE	women, 14-50 yrs	P100	141.69	<25	NR	1	0 - 0.06	0% - 0%	0% - 0%
TN 0672	Pecan (all commodities)	highest utilisation: Total	0.01	0.03	1.000	PRIMO-DE	child	P100	44.41	<25	NR	1	0.02 - 0.08	0% - 0%	0% - 0%
TN 0673	Pine nut (all commodities)	highest utilisation: Total	0	0.03	1.000	BR	Gen pop, > 10 yrs	47	200.00	<25	NR	1	0.02 - 0.09	0% - 0%	0% - 0%
TN 0675	Pistachio nut (all commodities)	highest utilisation: Total	0.01	0.03	1.000	PRIMO-IE	child	P97.5	115.86	<25	NR	1	0 - 0.17	0% - 0%	0% - 0%
TN 0678	Walnut (all commodities)	highest utilisation: Total	0.01	0.03	1.000	PRIMO-BE	toddler	P100	60.00	<25	NR	1	0 - 0.1	0% - 0%	0% - 0%
SO 0702	Sunflower seed (all commodities)	highest utilisation: Total	0.005 - 0.09	0	1.000	CA	women, 15-49 yrs	121	296.25	<25	NR	3	0.01 - 0.42	0% - 0%	0% - 0%
SO 0691	Cotton seed (all commodities)	highest utilisation: Oil (refined)	0.0006 - 0.02	0.02	1.000	US	Child, < 6 yrs	6354	3.13	NR	NR	3	0 - 0	0% - 0%	0% - 0%
HS 3381	Lemon, peel (all commodities)	highest utilisation: raw peel (C. limon Burm.f.)	0.39 - 9.1	1.4	1.000	AU	Child, 2-6 yrs	602	1.15	67	3	2b	0.25 - 0.25	0% - 0%	0% - 0%
HS 3382	Orange, peel (all commodities)	highest utilisation: raw peel (C. sinensis Osbeck)	0.39 - 9.1	1.4	1.000	AU	Gen pop, > 2 yrs	698	15.10	49	3	2b	0.95 - 0.95	0% - 0%	0% - 0%
MM 0095	Meat from mammals other than marine mammals	Total	NA	NA	1.000	CN	Child, 1-6 yrs	302	264.84	NR	NR	1	NA	0%	0%
MM 0095	Meat from mammals other than marine mammals: 20% as fat	Total		0.07	1.000	CN	Child, 1-6 yrs	302	52.97	NR	NR	1	0.230	0%	0%
MM 0095	Meat from mammals other than marine mammals: 80% as muscle	Total		0.02	1.000	CN	Child, 1-6 yrs	302	211.87	NR	NR	1	0.263	0%	0%

Annex 4

PYDIFLUMETOFEN (309)										IESTI						
Acute RfD= 0.3 mg/kg bw (300 µg/kg bw)										Maximum %ARfD:						
Codex Code	Commodity	Processing	STMR or STMR-P mg/kg	HR or HR-P mg/kg	DCF	Country	Population group	n	Large portion, g/person	Unit weight, edible portion, g	Variability factor	Case	IESTI µg/kg bw/day	20% all % acute RfD rounded	20% gen pop % acute RfD rounded	20% child % acute RfD rounded
MF 0100	Mammalian fats (except milk fats)	Total		0.07	1.000	PRIMO-FR	adult	P97.5	134.79	NR	NR	1	0.142	0%	0%	0%
MO 0105	Edible offal (mammalian)	Total		0.44	1.000	ZA	Gen pop, > 10 yrs	-	523.58	NR	NR	1	4.136	1%	1%	1%
ML 0106	Milks	Total	0.02		1.000	PRIMO-UK	infant	P97.5	1080.70	NR	NR	3	2.484	1%	0%	1%
PM 0110	Poultry meat	Total	NA	NA	1.000	CN	Child, 1-6 yrs	175	347.00	NR	NR	1	NA	0%	0%	0%
PM 0110	Poultry meat: 10% as fat	Total		0.02	1.000	CN	Child, 1-6 yrs	175	34.70	NR	NR	1	0.043	0%	0%	0%
PM 0110	Poultry meat: 90% as muscle	Total		0.02	1.000	CN	Child, 1-6 yrs	175	312.30	NR	NR	1	0.387	0%	0%	0%
PF 0111	Poultry, fats	Total		0.02	1.000	CA	Child, <6 yrs	66	49.38	NR	NR	1	0.058	0%	0%	0%
PO 0111	Poultry, edible offal (includes kidney, liver and skin)	Total		0.02	1.000	CN	Gen pop, > 1 yrs	421	345.63	NR	NR	1	0.130	0%	0%	0%
PE 0112	Eggs	Total		0.023	1.000	PRIMO-UK	infant	P97.5	108.00	NR	NR	1	0.286	0%	0%	0%

SULFOXAFLOL (252)										IESTI						
Acute RfD= 0.3 mg/kg bw (300 µg/kg bw)										Maximum %ARfD:						
										6%		6%		5%		
										all		gen pop		child		
Codex Code	Commodity	Processing	STMR or STMR-P mg/kg	HR or HR-P mg/kg	DCF	Country	Population group	n	Large portion, g/person	Unit weight, edible portion, g	Variability factor	Case	IESTI µg/kg bw/day	% acute RfD rounded	% acute RfD rounded	% acute RfD rounded
FB 0264	Blackberries (all commodities)	highest utilisation: Total	0.44	0.78	1.000	PRIMO-UK	toddler	P97.5	155.40	<25	NR	1	0.08 - 8.36	0% - 3%	0% - 2%	0% - 3%
FB 0266	Dewberries (incl Boysenberry, Loganberry) (all commodities)	highest utilisation: Total	0	0.78	1.000	PRIMO-UK	toddler	P97.5	25.50	<25	NR	1	1.37 - 1.37	0% - 0%	0% - 0%	0% - 0%
FB 0272	Raspberries, red, black (all commodities)	highest utilisation: Total	0.44	0.78	1.000	PRIMO-IE	child	P97.5	184.76	<25	NR	1	0.26 - 7.21	0% - 2%	0% - 1%	0% - 2%
FB 0020	Blueberries (all commodities)	highest utilisation: Total	0.39	1.4	1.000	CA	Child, <6 yrs	189	176.21	<25	NR	1	0.06 - 16.02	0% - 5%	0% - 4%	0% - 5%
FB 0021	Currants, black, red, white (all commodities)	highest utilisation: Total	0.39	1.4	1.000	AU	Gen pop, > 2 yrs	322	797.60	<25	NR	1	0.25 - 16.67	0% - 6%	0% - 6%	0% - 4%
FB 0268	Gooseberry (all commodities)	highest utilisation: Total	0.39	1.4	1.000	PRIMO-DE	child	P100	94.96	<25	NR	1	0.07 - 8.23	0% - 3%	0% - 2%	0% - 3%
FB 0273	Rose hips (all commodities)	highest utilisation: Total	0.39	1.4	1.000	PRIMO-FI	women	P97.5	156.60	<25	NR	1	0.25 - 3.08	0% - 1%	0% - 1%	0% - 1%
FB 0267	Elderberries (all commodities)	highest utilisation: Total	0.39	1.4	1.000	CN	Gen pop, > 1 yrs	136	420.22	29	3	2a	0.27 - 12.58	0% - 4%	0% - 4%	0% - 2%
FI 0326	Avocado (all commodities)	highest utilisation: Total	0	0.036	1.000	AU	Child, 2-6 yrs	182	229.90	171	3	2a	0.55 - 1.08	0% - 0%	0% - 0%	0% - 0%
FI 0345	Mango (all commodities)	highest utilisation: raw without peel	0.022	0.061	1.000	NL	toddler, 8-20 m	11	160.43	289	3	2b	0.01 - 2.88	0% - 1%	0% - 0%	0% - 1%
VS 0621	Asparagus (all commodities)	highest utilisation: Total	0.01	0.011	1.000	US	Child, < 6 yrs	23	279.99	9	NR	1	0.11 - 0.21	0% - 0%	0% - 0%	0% - 0%
SB 0716	Coffee bean (all commodities)	highest utilisation: extract (beverage)	0.0167 - 0.035	0	0.180	CA	women, 15-49 yrs	2666	2088.65	NR	NR	3	0 - 0.19	0% - 0%	0% - 0%	0% - 0%

Annex 4

TEBUCONAZOLE (189)											IESTI					
Acute RfD= 0.3 mg/kg bw (300 µg/kg bw)											Maximum %ARfD:					
											0%		0%		0%	
											all		gen pop		child	
Codex Code	Commodity	Processing	STMR or STMR-P mg/kg	HR or HR-P mg/kg	DCF	Country Population n			Large portion, g/person	Unit weight, edible portion, g	Variability factor	Case	IESTI µg/kg bw/day	% acute RfD rounded	% acute RfD rounded	% acute RfD rounded
SB 0716	Coffee bean (all commodities)	highest utilisation: extract (beverage)	0.04 - 0.08		0.180	CA	women, 15-49 yrs	2666	2088.65	NR	NR	3	0 - 0.22	0% - 0%	0% - 0%	0% - 0%

THIAMETHOXAM (245)										IESTI						
Acute RfD= 1 mg/kg bw (1000 µg/kg bw)										Maximum %ARfD:			5%	3%	5%	
													all	gen pop	child	
Codex Code	Commodity	Processing	STMR or STMR-P mg/kg	HR or HR-P mg/kg	DCF	Country	Population group	n	Large portion, g/person	Unit weight, edible portion, g	Variability factor	Case	IESTI µg/kg bw/day	% acute RfD rounded	% acute RfD rounded	% acute RfD rounded
FP 0307	Persimmon, Japanese (i.e. Kaki fruit) (all commodities)	highest utilisation: raw with peel (incl consumption without peel)		0.27	1.000	TH	Child, 3-6 yrs	20	264.88	228	3	2a	10.28 - 11.37	1% - 1%	0% - 1%	1% - 1%
GC 0653	Triticale	Total	0.032		1.000	DE	Gen pop, 14-80 yrs	27100	394.70	<25	NR	3	0.165	0%	0%	0%
GC 0654	Wheat (all commodities)	highest utilisation: flakes	0.00336 - 0.0768		1.000	CA	Child, <6 yrs	1909	539.23	NR	NR	3	0.03 - 1.1	0% - 0%	0% - 0%	0% - 0%
GC 0640	Barley (all commodities)	highest utilisation: beer	0.112 - 0.269		0.190	CA	Gen pop, all ages	2514	21271.20	NR	NR	3	0.01 - 5.75	0% - 1%	0% - 1%	0% - 0%
GC 0647	Oats (all commodities)	highest utilisation: flakes (rolled oats)	0.112		1.000	CA	Child, <6 yrs	1909	539.23	NR	NR	3	0.06 - 3.84	0% - 0%	0% - 0%	0% - 0%
GC 0649	Rice (all commodities)	highest utilisation: Total	0.704 - 9.35		1.000	CA	Child, <6 yrs	666	461.40	<25	NR	3	0.1 - 51.34	0% - 5%	0% - 3%	0% - 5%
GC 0651	Sorghum grain (Chicken corn, Dari seed, Durra, Feterita) (all commodities)	highest utilisation: cooked/boiled	0.0548 - 0.079		0.400	CN	Gen pop, > 1 yrs	356	1348.67	<25	NR	3	0.09 - 0.8	0% - 0%	0% - 0%	0% - 0%
GC 3081	Baby corn (all commodities)	highest utilisation: canned/preserved (imm cobs)		0.01	1.000	NL	Child, 2-6 yrs	E	75.00	12	NR	1	0.04 - 0.04	0% - 0%	0% - 0%	0% - 0%
GC 0447	Sweet corn (corn-on-the-cob) (kernels plus cob with husks removed) (all commodities)	highest utilisation: cooked/boiled (corn-on-the-cob)	0.01	0.01	1.000	TH	Child, 3-6 yrs	1383	196.99	191	3	2a	0.01 - 0.34	0% - 0%	0% - 0%	0% - 0%
GC 1275	Sweet corn (whole kernel without cob or husk) (all commodities)	highest utilisation: canned/preserved (kernels)	0.01	0.01	1.000	CA	Child, <6 yrs	289	153.76	NR	NR	3	0.01 - 0.1	0% - 0%	0% - 0%	0% - 0%
GS 0658	Sorgho or Sorghum, sweet	Total		0.24	1.000	CN	Child, 1-6 yrs	249	131.48	<25	NR	1	1.956	0%	0%	0%
MM 0095	Meat from mammals other than marine mammals	Total	NA	NA	1.000	CN	Child, 1-6 yrs	302	264.84	NR	NR	1	NA	0%	0%	0%
MM 0095	Meat from mammals other than marine mammals: 20% as fat	Total		0.01	1.000	CN	Child, 1-6 yrs	302	52.97	NR	NR	1	0.033	0%	0%	0%
MM 0095	Meat from mammals	Total		0.062	1.000	CN	Child, 1-6	302	211.87	NR	NR	1	0.814	0%	0%	0%

Annex 4

THIAMETHOXAM (245)										IESTI						
Acute RfD= 1 mg/kg bw (1000 µg/kg bw)										Maximum %ARfD:						
										all gen pop child						
Codex Code	Commodity	Processing	STMR or STMR-P mg/kg	HR or HR-P mg/kg	DCF	Country	Population group	n	Large portion, g/person	Unit weight, edible portion, g	Variability factor	Case	IESTI µg/kg bw/day	% acute RfD rounded	% acute RfD rounded	% acute RfD rounded
	other than marine mammals: 80% as muscle						yrs									
MF 0100	Mammalian fats (except milk fats)	Total		0.01	1.000	PRIMO-FR	adult	P97.5	134.79	NR	NR	1	0.020	0%	0%	0%
MO 0105	Edible offal (mammalian)	Total		0.041	1.000	ZA	Gen pop, > 10 yrs	-	523.58	NR	NR	1	0.385	0%	0%	0%
ML 0106	Milks	Total	0.096		1.000	PRIMO-UK	infant	P97.5	1080.70	NR	NR	3	11.925	1%	0%	1%
PM 0110	Poultry meat	Total	NA	NA	1.000	CN	Child, 1-6 yrs	175	347.00	NR	NR	1	NA	0%	0%	0%
PM 0110	Poultry meat: 10% as fat	Total		0.033	1.000	CN	Child, 1-6 yrs	175	34.70	NR	NR	1	0.071	0%	0%	0%
PM 0110	Poultry meat: 90% as muscle	Total		0.064	1.000	CN	Child, 1-6 yrs	175	312.30	NR	NR	1	1.239	0%	0%	0%
PF 0111	Poultry, fats	Total		0.033	1.000	CA	Child, <6 yrs	66	49.38	NR	NR	1	0.096	0%	0%	0%
PO 0111	Poultry, edible offal (includes kidney, liver and skin)	Total		0.36	1.000	CN	Gen pop, > 1 yrs	421	345.63	NR	NR	1	2.338	0%	0%	0%
PE 0112	Eggs	Total		0.028	1.000	PRIMO-UK	infant	P97.5	108.00	NR	NR	1	0.348	0%	0%	0%

Annex 5: Reports and other documents resulting from previous joint meetings of the FAO panel of experts on pesticide residues in food and the environment and the WHO core assessment group on pesticide residues.

1. Principles governing consumer safety in relation to pesticide residues. Report of a meeting of a WHO Expert Committee on Pesticide Residues held jointly with the FAO Panel of Experts on the Use of Pesticides in Agriculture. FAO Plant Production and Protection Division Report, No. PL/1961/11; WHO Technical Report Series, No. 240, 1962.
2. Evaluation of the toxicity of pesticide residues in food. Report of a Joint Meeting of the FAO Committee on Pesticides in Agriculture and the WHO Expert Committee on Pesticide Residues. FAO Meeting Report, No. PL/1963/13; WHO/Food Add./23, 1964.
3. Evaluation of the toxicity of pesticide residues in food. Report of the Second Joint Meeting of the FAO Committee on Pesticides in Agriculture and the WHO Expert Committee on Pesticide Residues. FAO Meeting Report, No. PL/1965/10; WHO/Food Add./26.65, 1965.
4. Evaluation of the toxicity of pesticide residues in food. FAO Meeting Report, No. PL/1965/10/1; WHO/Food Add./27.65, 1965.
5. Evaluation of the hazards to consumers resulting from the use of fumigants in the protection of food. FAO Meeting Report, No. PL/1965/10/2; WHO/Food Add./28.65, 1965.
6. Pesticide residues in food. Joint report of the FAO Working Party on Pesticide Residues and the WHO Expert Committee on Pesticide Residues. FAO Agricultural Studies, No. 73; WHO Technical Report Series, No. 370, 1967.
7. Evaluation of some pesticide residues in food. FAO/PL:CP/15; WHO/Food Add./67.32, 1967.
8. Pesticide residues. Report of the 1967 Joint Meeting of the FAO Working Party and the WHO Expert Committee. FAO Meeting Report, No. PL:1967/M/11; WHO Technical Report Series, No. 391, 1968.
9. 1967 Evaluations of some pesticide residues in food. FAO/PL:1967/M/11/1; WHO/Food Add./68.30, 1968.
10. Pesticide residues in food. Report of the 1968 Joint Meeting of the FAO Working Party of Experts on Pesticide Residues and the WHO Expert Committee on Pesticide Residues. FAO Agricultural Studies, No. 78; WHO Technical Report Series, No. 417, 1968.
11. 1968 Evaluations of some pesticide residues in food. FAO/PL:1968/M/9/1; WHO/Food Add./69.35, 1969.
12. Pesticide residues in food. Report of the 1969 Joint Meeting of the FAO Working Party of Experts on Pesticide Residues and the WHO Expert Group on Pesticide Residues. FAO Agricultural Studies, No. 84; WHO Technical Report Series, No. 458, 1970.
13. 1969 Evaluations of some pesticide residues in food. FAO/PL:1969/M/17/1; WHO/Food Add./70.38, 1970.
14. Pesticide residues in food. Report of the 1970 Joint Meeting of the FAO Working Party of Experts on Pesticide Residues and the WHO Expert Committee on Pesticide Residues. FAO Agricultural Studies, No. 87; WHO Technical Report Series, No. 4574, 1971.
15. 1970 Evaluations of some pesticide residues in food. AGP:1970/M/12/1; WHO/Food Add./71.42, 1971.
16. Pesticide residues in food. Report of the 1971 Joint Meeting of the FAO Working Party of Experts on Pesticide Residues and the WHO Expert Committee on Pesticide Residues. FAO Agricultural Studies, No. 88; WHO Technical Report Series, No. 502, 1972.
17. 1971 Evaluations of some pesticide residues in food. AGP:1971/M/9/1; WHO Pesticide Residue Series, No. 1, 1972.

18. Pesticide residues in food. Report of the 1972 Joint Meeting of the FAO Working Party of Experts on Pesticide Residues and the WHO Expert Committee on Pesticide Residues. FAO Agricultural Studies, No. 90; WHO Technical Report Series, No. 525, 1973.
19. 1972 Evaluations of some pesticide residues in food. AGP:1972/M/9/1; WHO Pesticide Residue Series, No. 2, 1973.
20. Pesticide residues in food. Report of the 1973 Joint Meeting of the FAO Working Party of Experts on Pesticide Residues and the WHO Expert Committee on Pesticide Residues. FAO Agricultural Studies, No. 92; WHO Technical Report Series, No. 545, 1974.
21. 1973 Evaluations of some pesticide residues in food. FAO/AGP/1973/M/9/1; WHO Pesticide Residue Series, No. 3, 1974.
22. Pesticide residues in food. Report of the 1974 Joint Meeting of the FAO Working Party of Experts on Pesticide Residues and the WHO Expert Committee on Pesticide Residues. FAO Agricultural Studies, No. 97; WHO Technical Report Series, No. 574, 1975.
23. 1974 Evaluations of some pesticide residues in food. FAO/AGP/1974/M/11; WHO Pesticide Residue Series, No. 4, 1975.
24. Pesticide residues in food. Report of the 1975 Joint Meeting of the FAO Working Party of Experts on Pesticide Residues and the WHO Expert Committee on Pesticide Residues. FAO Plant Production and Protection Series, No. 1; WHO Technical Report Series, No. 592, 1976.
25. 1975 Evaluations of some pesticide residues in food. AGP:1975/M/13; WHO Pesticide Residue Series, No. 5, 1976.
26. Pesticide residues in food. Report of the 1976 Joint Meeting of the FAO Panel of Experts on Pesticide Residues and the Environment and the WHO Expert Group on Pesticide Residues. FAO Food and Nutrition Series, No. 9; FAO Plant Production and Protection Series, No. 8; WHO Technical Report Series, No. 612, 1977.
27. 1976 Evaluations of some pesticide residues in food. AGP:1976/M/14, 1977.
28. Pesticide residues in food – 1977. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues and Environment and the WHO Expert Group on Pesticide Residues. FAO Plant Production and Protection Paper 10 Rev, 1978.
29. Pesticide residues in food: 1977 evaluations. FAO Plant Production and Protection Paper 10 Suppl., 1978.
30. Pesticide residues in food – 1978. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues and Environment and the WHO Expert Group on Pesticide Residues. FAO Plant Production and Protection Paper 15, 1979.
31. Pesticide residues in food: 1978 evaluations. FAO Plant Production and Protection Paper 15 Suppl., 1979.
32. Pesticide residues in food – 1979. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Expert Group on Pesticide Residues. FAO Plant Production and Protection Paper 20, 1980.
33. Pesticide residues in food: 1979 evaluations. FAO Plant Production and Protection Paper 20 Suppl., 1980.
34. Pesticide residues in food – 1980. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Expert Group on Pesticide Residues. FAO Plant Production and Protection Paper 26, 1981.
35. Pesticide residues in food: 1980 evaluations. FAO Plant Production and Protection Paper 26 Suppl., 1981.

36. Pesticide residues in food – 1981. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Expert Group on Pesticide Residues. FAO Plant Production and Protection Paper 37, 1982.
37. Pesticide residues in food: 1981 evaluations. FAO Plant Production and Protection Paper 42, 1982.
38. Pesticide residues in food – 1982. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Expert Group on Pesticide Residues. FAO Plant Production and Protection Paper 46, 1982.
39. Pesticide residues in food: 1982 evaluations. FAO Plant Production and Protection Paper 49, 1983.
40. Pesticide residues in food – 1983. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Expert Group on Pesticide Residues. FAO Plant Production and Protection Paper 56, 1985.
41. Pesticide residues in food: 1983 evaluations. FAO Plant Production and Protection Paper 61, 1985.
42. Pesticide residues in food – 1984. Report of the Joint Meeting on Pesticide Residues. FAO Plant Production and Protection Paper 62, 1985.
43. Pesticide residues in food – 1984 evaluations. FAO Plant Production and Protection Paper 67, 1985.
44. Pesticide residues in food – 1985. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and a WHO Expert Group on Pesticide Residues. FAO Plant Production and Protection Paper 68, 1986.
45. Pesticide residues in food – 1985 evaluations. Part I. Residues. FAO Plant Production and Protection Paper 72/1, 1986.
46. Pesticide residues in food – 1985 evaluations. Part II. Toxicology. FAO Plant Production and Protection Paper 72/2, 1986.
47. Pesticide residues in food – 1986. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and a WHO Expert Group on Pesticide Residues. FAO Plant Production and Protection Paper 77, 1986.
48. Pesticide residues in food – 1986 evaluations. Part I. Residues. FAO Plant Production and Protection Paper 78, 1986.
49. Pesticide residues in food – 1986 evaluations. Part II. Toxicology. FAO Plant Production and Protection Paper 78/2, 1987.
50. Pesticide residues in food – 1987. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and a WHO Expert Group on Pesticide Residues. FAO Plant Production and Protection Paper 84, 1987.
51. Pesticide residues in food – 1987 evaluations. Part I. Residues. FAO Plant Production and Protection Paper 86/1, 1988.
52. Pesticide residues in food – 1987 evaluations. Part II. Toxicology. FAO Plant Production and Protection Paper 86/2, 1988.
53. Pesticide residues in food – 1988. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and a WHO Expert Group on Pesticide Residues. FAO Plant Production and Protection Paper 92, 1988.
54. Pesticide residues in food – 1988 evaluations. Part I. Residues. FAO Plant Production and Protection Paper 93/1, 1988.

55. Pesticide residues in food – 1988 evaluations. Part II. Toxicology. FAO Plant Production and Protection Paper 93/2, 1989.
56. Pesticide residues in food – 1989. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and a WHO Expert Group on Pesticide Residues. FAO Plant Production and Protection Paper 99, 1989.
57. Pesticide residues in food – 1989 evaluations. Part I. Residues. FAO Plant Production and Protection Paper 100, 1990.
58. Pesticide residues in food – 1989 evaluations. Part II. Toxicology. FAO Plant Production and Protection Paper 100/2, 1990.
59. Pesticide residues in food – 1990. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and a WHO Expert Group on Pesticide Residues. FAO Plant Production and Protection Paper 102, Rome, 1990.
60. Pesticide residues in food – 1990 evaluations. Part I. Residues. FAO Plant Production and Protection Paper 103/1, Rome, 1990.
61. Pesticide residues in food – 1990 evaluations. Part II. Toxicology. World Health Organization, WHO/PCS/91.47, Geneva, 1991.
62. Pesticide residues in food – 1991. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and a WHO Expert Group on Pesticide Residues. FAO Plant Production and Protection Paper 111, Rome, 1991.
63. Pesticide residues in food – 1991 evaluations. Part I. Residues. FAO Plant Production and Protection Paper 113/1, Rome, 1991.
64. Pesticide residues in food – 1991 evaluations. Part II. Toxicology. World Health Organization, WHO/PCS/92.52, Geneva, 1992.
65. Pesticide residues in food – 1992. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and a WHO Expert Group on Pesticide Residues. FAO Plant Production and Protection Paper 116, Rome, 1993.
66. Pesticide residues in food – 1992 evaluations. Part I. Residues. FAO Plant Production and Protection Paper 118, Rome, 1993.
67. Pesticide residues in food – 1992 evaluations. Part II. Toxicology. World Health Organization, WHO/PCS/93.34, Geneva, 1993.
68. Pesticide residues in food – 1993. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and a WHO Expert Group on Pesticide Residues. FAO Plant Production and Protection Paper 122, Rome, 1994.
69. Pesticide residues in food – 1993 evaluations. Part I. Residues. FAO Plant Production and Protection Paper 124, Rome, 1994.
70. Pesticide residues in food – 1993 evaluations. Part II. Toxicology. World Health Organization, WHO/PCS/94.4, Geneva, 1994.
71. Pesticide residues in food – 1994. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and a WHO Expert Group on Pesticide Residues. FAO Plant Production and Protection Paper 127, Rome, 1995.
72. Pesticide residues in food – 1994 evaluations. Part I. Residues. FAO Plant Production and Protection Paper 131/1 and 131/2 (2 volumes), Rome, 1995.
73. Pesticide residues in food – 1994 evaluations. Part II. Toxicology. World Health Organization, WHO/PCS/95.2, Geneva, 1995.

74. Pesticide residues in food – 1995. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Core Assessment Group. FAO Plant Production and Protection Paper 133, Rome, 1996.
75. Pesticide residues in food – 1995 evaluations. Part I. Residues. FAO Plant Production and Protection Paper 137, 1996.
76. Pesticide residues in food – 1995 evaluations. Part II. Toxicological and Environmental. World Health Organization, WHO/PCS/96.48, Geneva, 1996.
77. Pesticide residues in food – 1996. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Core Assessment Group. FAO Plant Production and Protection Paper, 140, 1997.
78. Pesticide residues in food – 1996 evaluations. Part I. Residues. FAO Plant Production and Protection Paper, 142, 1997.
79. Pesticide residues in food – 1996 evaluations. Part II. Toxicological. World Health Organization, WHO/PCS/97.1, Geneva, 1997.
80. Pesticide residues in food – 1997. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Core Assessment Group. FAO Plant Production and Protection Paper, 145, 1998.
81. Pesticide residues in food – 1997 evaluations. Part I. Residues. FAO Plant Production and Protection Paper, 146, 1998.
82. Pesticide residues in food – 1997 evaluations. Part II. Toxicological and Environmental. World Health Organization, WHO/PCS/98.6, Geneva, 1998.
83. Pesticide residues in food – 1998. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Core Assessment Group. FAO Plant Production and Protection Paper, 148, 1999.
84. Pesticide residues in food – 1998 evaluations. Part I. Residues. FAO Plant Production and Protection Paper, 152/1 and 152/2 (two volumes).
85. Pesticide residues in food – 1998 evaluations. Part II. Toxicological and Environmental. World Health Organization, WHO/PCS/99.18, Geneva, 1999.
86. Pesticide residues in food – 1999. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Core Assessment Group. FAO Plant Production and Protection Paper, 153, 1999.
87. Pesticide residues in food – 1999 evaluations. Part I. Residues. FAO Plant Production and Protection Paper, 157, 2000.
88. Pesticide residues in food – 1999 evaluations. Part II. Toxicological. World Health Organization, WHO/PCS/00.4, Geneva, 2000.
89. Pesticide residues in food – 2000. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Core Assessment Group. FAO Plant Production and Protection Paper, 163, 2001.
90. Pesticide residues in food – 2000 evaluations. Part I. Residues. FAO Plant Production and Protection Paper, 165, 2001.
91. Pesticide residues in food – 2000 evaluations. Part II. Toxicological. World Health Organization, WHO/PCS/01.3, 2001.
92. Pesticide residues in food – 2001. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Core Assessment Group. FAO Plant Production and Protection Paper, 167, 2001.

93. Pesticide residues in food – 2001 evaluations. Part I. Residues. FAO Plant Production and Protection Paper, 171, 2002.
94. Pesticide residues in food – 2001 evaluations. Part II. Toxicological. World Health Organization, WHO/PCS/02.1, 2002.
95. Pesticide residues in food – 2002. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Core Assessment Group. FAO Plant Production and Protection Paper, 172, 2002.
96. Pesticide residues in food – 2002 evaluations. Part I. Residues. FAO Plant Production and Protection Paper, 175/1 and 175/2 (two volumes).
97. Pesticide residues in food – 2002 evaluations. Part II. Toxicological. World Health Organization, WHO/PCS, 2003.
98. Pesticide residues in food – 2003. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Core Assessment Group. FAO Plant Production and Protection Paper, 176, 2004.
99. Pesticide residues in food – 2003 evaluations. Part I. Residues. FAO Plant Production and Protection Paper, 177, 2004.
100. Pesticide residues in food – 2003 evaluations. Part II. Toxicological. World Health Organization, WHO/PCS, 2004.
101. Pesticide residues in food – 2004. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Core Assessment Group. FAO Plant Production and Protection Paper, 178, 2004.
102. Pesticide residues in food – 2004 evaluations. Part I. Residues. FAO Plant Production and Protection Paper, 182, 2005.
103. Pesticide residues in food – 2004 evaluations. Part II. Toxicological. World Health Organization, WHO/PCS, 2005.
104. Pesticide residues in food – 2005. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Core Assessment Group. FAO Plant Production and Protection Paper, 183, 2005.
105. Pesticide residues in food – 2005 evaluations. Part I. Residues. FAO Plant Production and Protection Paper, 184, 2006.
106. Pesticide residues in food – 2005 evaluations. Part II. Toxicological. World Health Organization, WHO/PCS/07.1, 2006.
107. Pesticide residues in food – 2006. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Core Assessment Group. FAO Plant Production and Protection Paper, 187, 2007.
108. Pesticide residues in food – 2006 evaluations. Part I. Residues. FAO Plant Production and Protection Paper, 189/1 and 189/2 (two volumes), 2007.
109. Pesticide residues in food – 2006 evaluations. Part II. Toxicological. World Health Organization, 2008.
110. Pesticide residues in food – 2007. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Core Assessment Group. FAO Plant Production and Protection Paper, 191, 2008.
111. Pesticide residues in food – 2007 evaluations. Part I. Residues. FAO Plant Production and Protection Paper, 192, 2008.
112. Pesticide residues in food – 2007 evaluations. Part II. Toxicological. World Health Organization, 2009.

113. Pesticide residues in food – 2008. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Core Assessment Group. FAO Plant Production and Protection Paper, 193, 2009.
114. Pesticide residues in food – 2008 evaluations. Part I. Residues. FAO Plant Production and Protection Paper, 194, 2009.
115. Pesticide residues in food – 2008 evaluations. Part II. Toxicological. World Health Organization, 2010.
116. Pesticide residues in food – 2009. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Core Assessment Group. FAO Plant Production and Protection Paper, 196, 2010.
117. Pesticide residues in food – 2009 evaluations. Part I. Residues. FAO Plant Production and Protection Paper, 198, 2010.
118. Pesticide residues in food – 2009 evaluations. Part II. Toxicological. World Health Organization, 2011.
119. Pesticide residues in food – 2010. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Core Assessment Group. FAO Plant Production and Protection Paper, 200, 2011.
120. Pesticide residues in food – 2010 evaluations. Part I. Residues. FAO Plant Production and Protection Paper, 206, 2011.
121. Pesticide residues in food – 2010 evaluations. Part II. Toxicological. World Health Organization, 2011.
122. Pesticide residues in food – 2011. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Core Assessment Group on Pesticide Residues. FAO Plant Production and Protection Paper, 211, 2012.
123. Pesticide residues in food – 2011 evaluations. Part I. Residues. FAO Plant Production and Protection Paper, 212, 2012.
124. Pesticide residues in food – 2011 evaluations. Part II. Toxicological. World Health Organization, 2012.
125. Pesticide residues in food – 2012. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Core Assessment Group on Pesticide Residues. FAO Plant Production and Protection Paper, 215, 2013.
126. Pesticide residues in food – 2012 evaluations. Part I. Residues. FAO Plant Production and Protection Paper, 216, 2013.
127. Pesticide residues in food – 2012 evaluations. Part II. Toxicological. World Health Organization, 2013.
128. Pesticide residues in food – 2013. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Core Assessment Group on Pesticide Residues. FAO Plant Production and Protection Paper, 219, 2014.
129. Pesticide residues in food – 2013 evaluations. Part I. Residues. FAO Plant Production and Protection Paper, 220, 2014.
130. Pesticide residues in food – 2013 evaluations. Part II. Toxicological. World Health Organization, 2014.
131. Pesticide residues in food – 2014. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Core Assessment Group on Pesticide Residues. FAO Plant Production and Protection Paper, 221, 2014.

132. Pesticide residues in food – 2014 evaluations. Part I. Residues. FAO Plant Production and Protection Paper, 222, 2015.
133. Pesticide residues in food – 2014 evaluations. Part II. Toxicological. World Health Organization, 2015.
134. Pesticide residues in food – 2015. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Core Assessment Group on Pesticide Residues. FAO Plant Production and Protection Paper, 223, 2015.
135. Pesticide residues in food – 2015 evaluations. Part I. Residues. FAO Plant Production and Protection Paper, 226, 2016.
136. Pesticide residues in food – 2015 evaluations. Part II. Toxicological. World Health Organization, 2016.
137. Pesticide residues in food – 2016. Report of a Special Session of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Core Assessment Group on Pesticide Residues. FAO Plant Production and Protection Paper, 227, 2016.
138. Pesticide residues in food – 2016 evaluations (Special Session). Toxicological. World Health Organization, in preparation.
139. Pesticide residues in food – 2016. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Core Assessment Group on Pesticide Residues. FAO Plant Production and Protection Paper, 229, 2015.
140. Pesticide residues in food – 2016 evaluations. Part I. Residues. FAO Plant Production and Protection Paper, 231, 2017.
141. Pesticide residues in food – 2016 evaluations. Part II. Toxicological. World Health Organization, 2017.
142. Pesticides residues in food – 2017. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Core Assessment Group on Pesticide Residues. FAO Plant Production and Protection Paper, 232, 2017.
143. Pesticide residues in food – 2017 evaluations. Part I. Residues. FAO Plant Production and Protection Paper, 233, 2018.
144. Pesticide residues in food – 2017 evaluations. Part II. Toxicological. World Health Organization, 2018.
145. Pesticides residues in food – 2018. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Core Assessment Group on Pesticide Residues. FAO Plant Production and Protection Paper, 234, 2018.
146. Pesticide residues in food – 2018 evaluations. Part I. Residues. FAO Plant Production and Protection Paper, 235, 2019.
147. Pesticide residues in food – 2018 evaluations. Part II. Toxicological. World Health Organization, 2019.
148. Pesticides residues in food – 2019. Report of the extra Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Core Assessment Group on Pesticide Residues.
149. Pesticide residues in food – 2019 Extra Joint FAO/WHO Meeting on Pesticide Residues Evaluation Part I: Residues. FAO Plant Production and Protection Paper, 2019.
150. Pesticides residues in food – 2019. Report of the Joint Meeting of the FAO Panel of Experts on Pesticide Residues in Food and the Environment and the WHO Core Assessment Group on Pesticide Residues. FAO Plant Production and Protection Paper, 2020.

151. Pesticide residues in food – 2019 evaluations. Part I. Residues. FAO Plant Production and Protection Paper, 235, 2019.

Annex 6: Livestock dietary burdens

BIXAFEN (262) – ESTIMATED MAXIMUM DIETARY BURDEN													
BEEF CATTLE													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Wheat forage	AF/AS	7.3	HR	25	29.20		20	100			5.84	29.2	
Barley forage	AF/AS	7.3	HR	30	24.33		10				2.433		
Wheat asp gr fn	CM/CF	17.1	STM	85	20.12	5				1.006			
Soybean asp gr fn	SM	15.98	STM	85	18.80	5				0.940			
Barley hay	AF/AS	11	HR	88	12.50	15				1.875			
Swede roots	VR	0.068	HR	10	0.68		40				0.272		
Potato culls	VR	0.068	HR	20	0.34	30				0.102			
Sorghum, grain grain	GC	0.196	STM	86	0.23	40	30		35	0.091	0.068		0.08
Soybean hulls	SM	0.0952	STM	90	0.11	5			5	0.005			0.005
Barley grain	GC	0.08	STM	88	0.09				60				0.055
Total						100	100	100	100	4.019	8.614	29.2	0.14

BIXAFEN (262) – ESTIMATED MAXIMUM DIETARY BURDEN													
DAIRY CATTLE													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Wheat forage	AF/AS	7.3	HR	25	29.20	20	20	60		5.840	5.84	17.52	
Barley forage	AF/AS	7.3	HR	30	24.33		10				2.433		
Oat forage	AF/AS	7.3	HR	30	24.33	10		40	5	2.433		9.733	1.217
Swede roots	VR	0.068	HR	10	0.68		20				0.136		
Carrot culls	VR	0.068	HR	12	0.57	10				0.057			
Potato culls	VR	0.068	HR	20	0.34		10				0.034		
Sorghum, grain grain	GC	0.196	STM	86	0.23	45	40		30	0.103	0.091		0.068
Soybean hulls	SM	0.0952	STM	90	0.11	10				0.011			
Barley grain	GC	0.08	STM	88	0.09				10				0.009
Brewer's grain dried	SM	0.074	STM	92	0.08				40				0.032
Soybean seed	VD	0.034	STM	89	0.04	5			10	0.002			0.004
Corn, field grain	GC	0.02	STM	88	0.02				5				0.001
Total						100	100	100	100	8.445	8.534	27.25	1.33

BIXAFEN (262) – ESTIMATED MAXIMUM DIETARY BURDEN													
POULTRY BROILER													
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Swede roots	VR	0.068	HR	10	0.68		10				0.068		
Sorghum, grain grain	GC	0.196	STMR	86	0.23	75	70	70	65	0.171	0.16	0.16	0.148
Cassava/tapioca roots	VR	0.068	HR	37	0.18		10				0.018		
Soybean hulls	SM	0.0952	STMR	90	0.11		5				0.005		
Brewer's grain dried	SM	0.074	STMR	92	0.08		5				0.004		
Soybean seed	VD	0.034	STMR	89	0.04	20		15		0.008		0.006	
Corn, field grain	GC	0.02	STMR	88	0.02				35				0.008
Canola meal	SM	0.02	STMR	88	0.02	5		5		0.001		0.001	
Peanut meal	SM	0.0114	STMR	85	0.01			5				7E-04	
Soybean meal	SM	0.00476	STMR	92	0.01			5				3E-04	
Total						100	100	100	100	0.180	0.255	0.167	0.156

POULTRY LAYER													
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Wheat forage	AF/AS	7.3	HR	25	29.20		10				2.92		
Swede roots	VR	0.068	HR	10	0.68		10				0.068		
Sorghum, grain grain	GC	0.196	STMR	86	0.23	75	70	70	55	0.171	0.16	0.16	0.125
Cassava/tapioca roots	VR	0.068	HR	37	0.18		10				0.018		
Soybean seed	VD	0.034	STMR	89	0.04	20		15		0.008		0.006	
Corn, field grain	GC	0.02	STMR	88	0.02				45				0.01
Canola meal	SM	0.02	STMR	88	0.02	5		5		0.001		0.001	
Peanut meal	SM	0.0114	STMR	85	0.01			5				7E-04	
Soybean meal	SM	0.00476	STMR	92	0.01			5				3E-04	
Total						100	100	100	100	0.180	3.166	0.167	0.136

BIXAFEN (262) – ESTIMATED MEAN DIETARY BURDEN													
BEEF CATTLE													
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Wheat asp gr fn	CM/CF	17.1	STMR/STMR-P	85	20.12	5				1.0059			
Soybean asp gr fn	SM	15.98	STMR/STMR-P	85	18.80	5				0.94			
Wheat forage	AF/AS	3.5	STMR/STMR-P	25	14.00		20	100			2.8	14	
Barley forage	AF/AS	3.5	STMR/STMR-P	30	11.67		10				1.167		
Barley hay	AF/AS	2.2	STMR/STMR-P	88	2.50	15				0.375			
Swede roots	VR	0.034	STMR/STMR-P	10	0.34		40				0.136		
Sorghum, grain grain	GC	0.196	STMR/STMR-P	86	0.23	40	30		35	0.0912	0.068		0.08

BIXAFEN (262) – ESTIMATED MEAN DIETARY BURDEN													
BEEF CATTLE													
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Potato culls	VR	0.034	STMR/STMR-P	20	0.17	30				0.051			
Soybean hulls	SM	0.0952	STMR/STMR-P	90	0.11	5			5	0.005			0.005
Barley grain	GC	0.08	STMR/STMR-P	88	0.09				60				0.055
Total						100	100	100	100	2.468	4.171	14	0.14

DAIRY CATTLE													
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Wheat forage	AF/AS	3.5	STMR/STMR-P	25	14.00	20	20	60		2.8	2.8	8.4	
Barley forage	AF/AS	3.5	STMR/STMR-P	30	11.67	0	10			0	1.167		
Oat forage	AF/AS	3.5	STMR/STMR-P	30	11.67	10		40	5	1.1667		4.667	0.583
Swede roots	VR	0.034	STMR/STMR-P	10	0.34	0	20			0	0.068		
Carrot culls	VR	0.034	STMR/STMR-P	12	0.28	10				0.0283			
Sorghum, grain	GC	0.196	STMR/STMR-P	86	0.23	45	40		30	0.1026	0.091		0.068
Potato culls	VR	0.034	STMR/STMR-P	20	0.17	0	10			0	0.017		
Soybean hulls	SM	0.0952	STMR/STMR-P	90	0.11	10				0.0106			
Barley grain	GC	0.08	STMR/STMR-P	88	0.09	0			10	0			0.009
Brewer's grain dried	SM	0.074	STMR/STMR-P	92	0.08	0			40	0			0.032
Soybean seed	VD	0.034	STMR/STMR-P	89	0.04	5			10	0.0019			0.004
Corn, field grain	GC	0.02	STMR/STMR-P	88	0.02	0			5	0			0.001
Total						100	100	100	100	4.1101	4.143	13.07	0.698

BIXAFEN (262) – ESTIMATED MEAN DIETARY BURDEN													
POULTRY BROILER													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Swede roots	VR	0.034	STMR/STMR-P	10	0.34		10				0.034		
Sorghum, grain grain	GC	0.196	STMR/STMR-P	86	0.23	75	70	70	65	0.17	0.16	0.16	0.148
Soybean hulls	SM	0.0952	STMR/STMR-P	90	0.11		5				0.005		
Cassava/tapioca roots	VR	0.034	STMR/STMR-P	37	0.09		15				0.014		
Soybean seed	VD	0.034	STMR/STMR-P	89	0.04	20		15		0.01		0.006	
Corn, field grain	GC	0.02	STMR/STMR-P	88	0.02				35				0.008
Canola meal	SM	0.02	STMR/STMR-P	88	0.02	5		5		0.00		0.001	
Peanut meal	SM	0.0114	STMR/STMR-P	85	0.01			5				7E-04	
Soybean meal	SM	0.00476	STMR/STMR-P	92	0.01			5				3E-04	
Total						100	100	100	100	0.1797	0.213	0.167	0.156

POULTRY LAYER													MEAN
Commodity	CC	Residue (mg/kg)	Basis	D M (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Wheat forage	AF/AS	3.5	STMR/STMR-P	25	14.00		10				1.4		
Swede roots	VR	0.034	STMR/STMR-P	10	0.34		10				0.034		
Sorghum, grain grain	GC	0.196	STMR/STMR-P	86	0.23	75	70	70	55	0.1709	0.16	0.16	0.125
Soybean hulls	SM	0.0952	STMR/STMR-P	90	0.11		5				0.005		
Cassava/tapioca roots	VR	0.034	STMR/STMR-P	37	0.09		5				0.005		
Soybean seed	VD	0.034	STMR/STMR-P	89	0.04	20		15		0.0076		0.006	
Corn, field grain	GC	0.02	STMR/STMR-P	88	0.02				45				0.01
Canola meal	SM	0.02	STMR/STMR-P	88	0.02	5		5		0.001136		0.001	
Peanut meal	SM	0.0114	STMR/STMR-P	85	0.01			5				0.001	
Soybean meal	SM	0.00476	STMR/STMR-P	92	0.01			5				3E-04	
Total						100	100	100	100	0.1797	1.603	0.167	0.136

CLOTHIANIDIN (238) – ESTIMATED MAXIMUM DIETARY BURDEN													
BEEF CATTLE													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Grape pomace, wet	AB	0.23	STMR	15	1.53			20				0.307	
Swede roots	VR	0.15	HR	10	1.50		40	10			0.6	0.15	
Rice hulls	CM/CF	1.1	STMR	90	1.22			5				0.061	
Sugarcane tops	AM/AV	0.27	HR	25	1.08			50				0.54	
Wheat forage	AF/AS	0.21	HR	25	0.84		20	15			0.168	0.126	
Sorghum, grain forage	AF/AS	0.29	HR	35	0.83	15				0.124			
Barley straw	AF/AS	0.72	HR	89	0.81		10				0.081		
Potato culls	VR	0.15	HR	20	0.75	30				0.225			
Cabbage heads, leaves	AM/AV	0.08	HR	15	0.53		20				0.107		
Rice grain	GC	0.3	STMR	88	0.34	20				0.068			
Rice bran/pollard	CM/CF	0.28	STMR	90	0.31	15			20	0.047			0.062
Pea vines	AL	0.05	HR	25	0.20		10				0.02		
Rice straw	AF/AS	0.13	HR	90	0.14				55				0.079
Beet, sugar molasses	DM	0.064	STMR	75	0.09	10				0.009			
Corn, field forage/silage	AF/AS	0.021	HR	40	0.05	10				0.005			
Beet, sugar dried pulp	AB	0.034	STMR	88	0.04				5				0.002
Corn, field grain	GC	0.02	STMR	88	0.02				20				0.005
Total						100	100	100	100	0.478	0.976	1.184	0.148

DAIRY CATTLE													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Grape pomace, wet	AB	0.23	STMR	15	1.53			20				0.307	
Swede roots	VR	0.15	HR	10	1.50		20	10			0.3	0.15	
Carrot culls	VR	0.15	HR	12	1.25	10				0.125			
Rice hulls	CM/CF	1.1	STMR	90	1.22			10				0.122	
Sugarcane tops	AM/AV	0.27	HR	25	1.08			25				0.27	
Wheat forage	AF/AS	0.21	HR	25	0.84	20	20	35		0.168	0.168	0.294	
Sorghum, grain forage	AF/AS	0.29	HR	35	0.83	20			40	0.166			0.331
Barley straw	AF/AS	0.72	HR	89	0.81		10				0.081		
Potato culls	VR	0.15	HR	20	0.75		10				0.075		
Cabbage heads, leaves	AM/AV	0.08	HR	15	0.53		20				0.107		
Corn, sweet forage	AF/AS	0.22	HR	48	0.46	5				0.023			
Rice grain	GC	0.3	STMR	88	0.34	20				0.068			
Bean vines	AL	0.11	HR	35	0.31		20				0.063		
Rice bran/pollard	CM/CF	0.28	STMR	90	0.31	15			10	0.047			0.031
Pea vines	AL	0.05	HR	25	0.20	10				0.020			
Rice whole crop silage	AF/AS	0.025	HR	40	0.06				50				0.031
Total						100	100	100	100	0.616	0.793	1.143	0.394

CLOTHIANIDIN (238) – ESTIMATED MAXIMUM DIETARY BURDEN													
POULTRY BROILER													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Swede roots	VR	0.15	HR	10	1.50		10				0.15		
Rice grain	GC	0.3	STMR	88	0.34	20		50		0.068		0.17	
Rice bran/pollard	CM/CF	0.28	STMR	90	0.31	10	10	20	5	0.031	0.031	0.062	0.016
Bean seed	VD	0.02	STMR	88	0.02		20	30			0.005	0.007	
Corn, field grain	GC	0.02	STMR	88	0.02	70	60		70	0.016	0.014		0.016
Sorghum, grain grain	GC	0.015	STMR	86	0.02				25				0.004
Total						100	100	100	100	0.12	0.199	0.239	0.036

CLOTHIANIDIN (238) – ESTIMATED MAXIMUM DIETARY BURDEN													
POULTRY LAYER													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Swede roots	VR	0.15	HR	10	1.50		10				0.15		
Wheat forage	AF/AS	0.21	HR	25	0.84		10				0.084		
Cabbage heads, leaves	AM/AV	0.08	HR	15	0.53		5				0.027		
Rice grain	GC	0.3	STMR	88	0.34	20		50		0.068		0.17	
Rice bran/pollard	CM/CF	0.28	STMR	90	0.31	10	5	20	20	0.031	0.016	0.062	0.062
Pea vines	AL	0.05	HR	25	0.20		10				0.02		
Rape forage	AM/AV	0.027	HR	30	0.09		5				0.005		
Bean seed	VD	0.02	STMR	88	0.02		20	30			0.005	0.007	
Corn, field grain	GC	0.02	STMR	88	0.02	70	35		80	0.02	0.008		0.02
Total						100	100	100	100	0.115	0.313	0.239	0.08

CLOTHIANIDIN (238) – ESTIMATED MEAN DIETARY BURDEN													
BEEF CATTLE													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Grape pomace, wet	AB	0.23	STMR/STMR-P	15	1.53			20				0.307	
Rice hulls	CM/CF	1.1	STMR/STMR-P	90	1.22			5				0.061	
Sugarcane tops	AM/AV	0.19	STMR/STMR-P	25	0.76			50				0.38	
Sorghum, grain forage	AF/AS	0.19	STMR/STMR-P	35	0.54	15	20	25		0.0814	0.109	0.136	
Rice grain	GC	0.3	STMR/STMR-P	88	0.34	20				0.0682			
Rice bran/pollard	CM/CF	0.28	STMR/STMR-P	90	0.31	15			20	0.0467			0.062
Cabbage heads, leaves	AM/AV	0.03	STMR/STMR-P	15	0.20		20				0.04		
Pea vines	AL	0.05	STMR/STMR-P	25	0.20		20				0.04		

CLOTHIANIDIN (238) – ESTIMATED MEAN DIETARY BURDEN													
BEEF CATTLE													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Barley forage	AF/AS	0.057	STMR/STMR-P	30	0.19		10				0.019		
Cowpea forage	AL	0.05	STMR/STMR-P	30	0.17		30				0.05		
Potato culls	VR	0.02	STMR/STMR-P	20	0.10	30				0.030			
Beet, sugar molasses	DM	0.064	STMR/STMR-P	75	0.09	10				0.009			
Grass forage (fresh)	AF/AS	0.01	STMR/STMR-P	25	0.04				5				0.002
Beet, sugar dried pulp	AB	0.034	STMR/STMR-P	88	0.04	10			5	0.004			0.002
Rice straw	AF/AS	0.03	STMR/STMR-P	90	0.03				50				0.0167
Corn, field grain	GC	0.02	STMR/STMR-P	88	0.02				20				0.005
Total						100	100	100	100	0.2387	0.258	0.883	0.087

DAIRY CATTLE													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Grape pomace, wet	AB	0.23	STMR/STMR-P	15	1.53		0	20			0	0.307	
Rice hulls	CM/CF	1.1	STMR/STMR-P	90	1.22	0		10		0		0.122	
Sugarcane tops	AM/AV	0.19	STMR/STMR-P	25	0.76	0		25		0		0.19	
Sorghum, grain forage	AF/AS	0.19	STMR/STMR-P	35	0.54	40	20	45	40	0.2171	0.109	0.244	0.217
Rice grain	GC	0.3	STMR/STMR-P	88	0.34	20				0.0682			
Rice bran/pollard	CM/CF	0.28	STMR/STMR-P	90	0.31	15	20		10	0.0467	0.062		0.031
Bean vines	AL	0.075	STMR/STMR-P	35	0.21	0	20			0	0.043		
Cabbage heads, leaves	AM/AV	0.03	STMR/STMR-P	15	0.20	0	20			0	0.04		
Pea vines	AL	0.05	STMR/STMR-P	25	0.20	10				0.02			
Barley forage	AF/AS	0.057	STMR/STMR-P	30	0.19	0	20			0	0.038		
Corn, sweet forage	AF/AS	0.084	STMR/STMR-P	48	0.18	15				0.0263			
Beet, sugar dried pulp	AB	0.034	STMR/STMR-P	88	0.04	0			40	0			0.015
Corn, field forage/silage	AF/AS	0.01	STMR/STMR-P	40	0.03	0			10	0			0.003
Total						100	100	100	100	0.3782	0.292	0.863	0.266

CLOTHIANIDIN (238) – ESTIMATED MEAN DIETARY BURDEN													
POULTRY BROILER													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Rice grain	GC	0.3	STMR/STMR-P	88	0.34	20		50		0.0682		0.17	
Rice bran/pollard	CM/CF	0.28	STMR/STMR-P	90	0.31	10	10	20	5	0.0311	0.031	0.062	0.016
Carrot culls	VR	0.02	STMR/STMR-P	12	0.17		10				0.017		
Cassava/tapioca roots	VR	0.02	STMR/STMR-P	37	0.05		10				0.005		
Bean seed	VD	0.02	STMR/STMR-P	88	0.02		20	30			0.005	0.007	
Corn, field grain	GC	0.02	STMR/STMR-P	88	0.02	70	50		70	0.0159	0.011		0.016
Sorghum, grain grain	GC	0.015	STMR/STMR-P	86	0.02				25				0.004
Total						100	100	100	100	0.1152	0.069	0.239	0.036

POULTRY LAYER													
POULTRY LAYER													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Sorghum, grain forage	AF/AS	0.19	STMR/STMR-P	35	0.54		10				0.054		
Rice grain	GC	0.3	STMR/STMR-P	88	0.34	20		50		0.0682		0.17	
Rice bran/pollard	CM/CF	0.28	STMR/STMR-P	90	0.31	10	5	20	20	0.0311	0.016	0.062	0.062
Cabbage heads, leaves	AM/AV	0.03	STMR/STMR-P	15	0.20		5				0.01		
Pea vines	AL	0.05	STMR/STMR-P	25	0.20		10				0.02		
Carrot culls	VR	0.02	STMR/STMR-P	12	0.17		10				0.017		
Rape forage	AM/AV	0.02	STMR/STMR-P	30	0.07		5				0.003		
Cassava/tapioca roots	VR	0.02	STMR/STMR-P	37	0.05		5				0.003		
Bean seed	VD	0.02	STMR/STMR-P	88	0.02		20	30			0.005	0.007	
Corn, field grain	GC	0.02	STMR/STMR-P	88	0.02	70	30		80	0.0159	0.007		0.018
Total						100	100	100	100	0.1152	0.134	0.239	0.08

CYPRODINIL (207) ESTIMATED MAXIMUM DIETARY BURDEN													
BEEF CATTLE													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Kale leaves	AM/AV	8	HR	15	53.33		20				10.67		
Carrot culls	VR	1	HR	12	8.33		15	5			1.25	0.417	
Wheat straw	AF/AS	5.8	HR	88	6.59	10	20	80		0.659	1.318	5.273	
Barley straw	AF/AS	5.8	HR	89	6.52		10	15			0.652	0.978	
Apple pomace, wet	AB	1.8	STMR	40	4.50		20				0.9		
Pea vines	AL	0.56	HR	25	2.24		15				0.336		
Barley bran fractions	CM/CF	1.74	STMR	90	1.93				10				0.193
Barley grain	GC	0.58	STMR	88	0.66	50			70	0.330			0.461
Wheat milled bypds	CM/CF	0.21	STMR	88	0.24	40			20	0.095			0.048
Total						100	100	100	100	1.084	15.12	6.667	0.702

DAIRY CATTLE													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Kale leaves	AM/AV	8	HR	15	53.33		20	40			10.67	21.33	
Carrot culls	VR	1	HR	12	8.33	10	15	5		0.833	1.25	0.417	
Wheat straw	AF/AS	5.8	HR	88	6.59	10	20	20		0.659	1.318	1.318	
Barley straw	AF/AS	5.8	HR	89	6.52		10				0.652		
Triticale straw	AF/AS	5.8	HR	90	6.44			35				2.256	
Apple pomace, wet	AB	1.8	STMR	40	4.50	10	10			0.450	0.45		
Pea vines	AL	0.56	HR	25	2.24	10	20			0.224	0.448		
Barley grain	GC	0.58	STMR	88	0.66	45	5		40	0.297	0.033		0.264
Wheat milled bypds	CM/CF	0.21	STMR	88	0.24	15			45	0.036			0.107
Soybean seed	VD	0.085	STMR	89	0.10				10				0.01
Rape meal	SM	0.02	STMR	88	0.02				5				0.001
Total						100	100	100	100	2.499	14.82	25.32	0.382

POULTRY BROILER													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Carrot culls	VR	1	HR	12	8.33		10				0.833		
Barley bran fractions	CM/CF	1.74	STMR	90	1.93	8				0.155			
Barley grain	GC	0.58	STMR	88	0.66	75	70	15	10	0.494	0.461	0.099	0.066
Wheat milled bypds	CM/CF	0.21	STMR	88	0.24	17	20	20	5	0.041	0.048	0.048	0.012
Soybean seed	VD	0.085	STMR	89	0.10			15				0.014	
Wheat grain	GC	0.07	STMR	89	0.08			50				0.039	

POULTRY BROILER													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Rape meal	SM	0.02	STMR	88	0.02				5				0.001
Total						100	100	100	20	0.690	1.342	0.2	0.079

CYPRODINIL (207) ESTIMATED MAXIMUM DIETARY BURDEN													
POULTRY LAYER													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Carrot culls	VR	1	HR	12	8.33		10				0.833		
Wheat straw	AF/AS	5.8	HR	88	6.59		10				0.659		
Cabbage heads, leaves	AM/AV	0.36	HR	15	2.40		5				0.12		
Pea vines	AL	0.56	HR	25	2.24		10				0.224		
Barley bran fractions	CM/CF	1.74	STMR	90	1.93				5				0.097
Barley grain	GC	0.58	STMR	88	0.66	75	65	15		0.494	0.428	0.099	
Wheat milled bypds	CM/CF	0.21	STMR	88	0.24	25		20	25	0.060		0.048	0.06
Soybean seed	VD	0.085	STMR	89	0.10			15				0.014	
Wheat grain	GC	0.07	STMR	89	0.08			50				0.039	
Rape meal	SM	0.02	STMR	88	0.02				15				0.003
Total						100	100	100	45	0.554	2.265	0.2	0.16

CYPRODINIL (207) ESTIMATED MEAN DIETARY BURDEN													
BEEF CATTLE													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Apple pomace, wet	AB	1.8	STMR/STMR-P	40	4.50		20	20			0.9	0.9	
Kale leaves	AM/AV	0.37	STMR/STMR-P	15	2.47		20				0.493		
Barley bran fractions	CM/CF	1.74	STMR/STMR-P	90	1.93				10				0.193
Carrot culls	VR	0.195	STMR/STMR-P	12	1.63		15	5			0.244	0.081	
Barley grain	GC	0.58	STMR/STMR-P	88	0.66	50	45	75	70	0.3295	0.297	0.494	0.461
Wheat straw	AF/AS	0.395	STMR/STMR-P	88	0.45	10				0.0449			
Wheat milled bypds	CM/CF	0.21	STMR/STMR-P	88	0.24	40			20	0.0955			0.048
Total						100	100	100	100	0.4699	1.934	1.476	0.702

CYPRODINIL (207) ESTIMATED MEAN DIETARY BURDEN													
DAIRY CATTLE													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Apple pomace, wet	AB	1.8	STMR/STMR-P	40	4.50	10	10	10		0.4500	0.45	0.45	
Kale leaves	AM/AV	0.37	STMR/STMR-P	15	2.47	0	20	40		0	0.493	0.987	
Carrot culls	VR	0.195	STMR/STMR-P	12	1.63	10	15	5		0.1625	0.244	0.081	
Barley grain	GC	0.58	STMR/STMR-P	88	0.66	45	40	40	40	0.2966	0.264	0.264	0.264
Wheat straw	AF/AS	0.395	STMR/STMR-P	88	0.45	10	15	5		0.0449	0.067	0.022	
Pea vines	AL	0.1	STMR/STMR-P	25	0.40	10				0.04			
Wheat milled bypds	CM/CF	0.21	STMR/STMR-P	88	0.24	15			45	0.0358			0.107
Soybean seed	VD	0.085	STMR/STMR-P	89	0.10	0			10	0			0.01
Rape meal	SM	0.02	STMR/STMR-P	88	0.02	0			5	0			0.001
Total						100	100	100	100	1.0298	1.518	1.804	0.382

POULTRY BROILER													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Barley bran fractions	CM/CF	1.74	STMR/STMR-P	90	1.93	8				0.1547			
Carrot culls	VR	0.195	STMR/STMR-P	12	1.63		10				0.163		
Barley grain	GC	0.58	STMR/STMR-P	88	0.66	75	70	15	10	0.4943	0.461	0.099	0.066
Wheat milled bypds	CM/CF	0.21	STMR/STMR-P	88	0.24	17	20	20	5	0.0406	0.048	0.048	0.012
Soybean seed	VD	0.085	STMR/STMR-P	89	0.10			15				0.014	
Wheat grain	GC	0.07	STMR/STMR-P	89	0.08			50				0.039	
Rape meal	SM	0.02	STMR/STMR-P	88	0.02				5				0.001
Total						100	100	100	20	0.6896	0.672	0.2	0.079

CYPRODINIL (207) ESTIMATED MEAN DIETARY BURDEN													
POULTRY LAYER													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US- CAN	EU	AU	JP	US- CAN	EU	AU	JP
Barley bran fractions	CM/CF	1.74	STMR/STMR-P	90	1.93				5				0.097
Carrot culls	VR	0.195	STMR/STMR-P	12	1.63		10				0.163		
Barley grain	GC	0.58	STMR/STMR-P	88	0.66	75	90	15		0.4943	0.593	0.099	
Wheat milled bypds	CM/CF	0.21	STMR/STMR-P	88	0.24	25		20	25	0.0597		0.048	0.06
Soybean seed	VD	0.085	STMR/STMR-P	89	0.10			15				0.014	
Wheat grain	GC	0.07	STMR/STMR-P	89	0.08			50				0.039	
Rape meal	SM	0.02	STMR/STMR-P	88	0.02				15				0.003
Total						100	100	100	45	0.5540	0.756	0.2	0.16

DIFENOCONAZOLE (224) – ESTIMATED MAXIMUM DIETARY BURDEN													
BEEF CATTLE													
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Grape pomace, wet	AB	6.2	STMR	15	41.33			20				8.267	
Potato process waste	AB	3.8	STMR	12	31.67	30	40			9.500	12.67		
Rice straw	AF/AS	10	HR	90	11.11		10	60	55		1.111	6.667	6.111
Potato culls	VR	1.9	HR	20	9.50	30	30	10		2.850	2.85	0.95	
Soybean aspr gr fn	SM	6.22	STMR	85	7.32	5				0.366			
Beet, mangel fodder	AM/AV	0.95	HR	15	6.33		20				1.267		
Rice hulls	CM/CF	3.6	STMR	90	4.00			5				0.2	
Bean vines	AL	0.85	HR	35	2.43			5				0.121	
Wheat straw	AF/AS	1.2	HR	88	1.36	10				0.136			
Rice grain	GC	1.1	STMR	88	1.25	20				0.250			
Rice bran/pollard	CM/CF	0.76	STMR	90	0.84	5			20	0.042			0.169
Soybean hulls	SM	0.02	STMR	90	0.02				5				0.001
Soybean seed	VD	0.01	STMR	89	0.01				15				0.002
Soybean meal	SM	0.004	STMR	92	0.00				5				0.00
Total						100	100	100	100	13.144	17.89	16.2	6.283

DAIRY CATTLE													
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Grape pomace, wet	AB	6.2	STMR	15	41.33			20				8.267	
Potato process waste	AB	3.8	STMR	12	31.67	10	30			3.167	9.5		
Rice straw	AF/AS	10	HR	90	11.11		5	20	25		0.556	2.222	2.778
Potato culls	VR	1.9	HR	20	9.50	10	30	10		0.950	2.85	0.95	
Beet, mangel fodder	AM/AV	0.95	HR	15	6.33		25				1.583		
Rice hulls	CM/CF	3.6	STMR	90	4.00			10				0.4	
Bean vines	AL	0.85	HR	35	2.43		10	40			0.243	0.971	
Almond hulls	AM/AV	1.24	STMR	90	1.38	10				0.138			
Wheat straw	AF/AS	1.2	HR	88	1.36	10				0.136			
Rice grain	GC	1.1	STMR	88	1.25	20				0.250			
Rice bran/pollard	CM/CF	0.76	STMR	90	0.84	15			10	0.127			0.084
Canola meal	SM	0.03	STMR	88	0.03	10				0.003			
Cotton undelinted seed	SO	0.021	STMR	88	0.02	10				0.002			
Soybean hulls	SM	0.02	STMR	90	0.02	5				0.001			
Soybean seed	VD	0.01	STMR	89	0.01				10				0.001

DAIRY CATTLE													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Soybean meal	SM	0.004	STM	92	0.00				55				0.002
Total						100	100	100	100	4.774	14.73	12.81	2.866

DIFENOCONAZOLE (224) – ESTIMATED MAXIMUM DIETARY BURDEN													
POULTRY BROILER													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Potato culls	VR	1.9	HR	20	9.50		10				0.95		
Rice grain	GC	1.1	STM	88	1.25	20		50		0.250		0.625	
Rice bran/pollard	CM/CF	0.76	STM	90	0.84	10	10	20	5	0.084	0.084	0.169	0.042
Canola meal	SM	0.03	STM	88	0.03	15	18	5		0.005	0.006	0.002	
Soybean seed	VD	0.01	STM	89	0.01	20	20	15		0.002	0.002	0.002	
Sunflower meal	SM	0.01	STM	92	0.01	10		10		0.001		0.001	
Soybean meal	SM	0.004	STM	92	0.00		22		35		1E-03		0.002
Total						75	80	100	40	0.343	1.044	0.798	0.044

POULTRY LAYER													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Potato culls	VR	1.9	HR	20	9.50		10				0.95		
Wheat straw	AF/AS	1.2	HR	88	1.36		10				0.136		
Cabbage heads, leaves	AM/AV	0.19	HR	15	1.27		5				0.063		
Rice grain	GC	1.1	STM	88	1.25	20		50		0.250		0.625	
Rice bran/pollard	CM/CF	0.76	STM	90	0.84	10	5	20	20	0.084	0.042	0.169	0.169
Canola meal	SM	0.03	STM	88	0.03	15	10	5		0.005	0.003	0.002	
Soybean seed	VD	0.01	STM	89	0.01	20	15	15		0.002	0.002	0.002	
Sunflower meal	SM	0.01	STM	92	0.01	10		10		0.001		0.001	
Soybean meal	SM	0.004	STM	92	0.00		15		30		7E-04		0.00
Total						75	70	100	50	0.343	1.198	0.798	0.17

DIFENOCONAZOLE (224) – ESTIMATED MEAN DIETARY BURDEN													
BEEF CATTLE													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Grape pomace, wet	AB	6.2	STMR/STMR-P	15	41.33			20				8.267	
Potato process waste	AB	3.8	STMR/STMR-P	12	31.67	30	40			9.5	12.67		
Soybean asp gr fn	SM	6.22	STMR/STMR-P	85	7.32	5				0.3659			
Potato culls	VR	1.2	STMR/STMR-P	20	6.00	30	30	10		1.8	1.8	0.6	
Rice hulls	CM/CF	3.6	STMR/STMR-P	90	4.00			5				0.2	
Rice straw	AF/AS	2.2	STMR/STMR-P	90	2.44		10	60	55		0.244	1.467	1.344
Bean vines	AL	0.75	STMR/STMR-P	35	2.14			5				0.107	
Beet, mangel fodder	AM/AV	0.25	STMR/STMR-P	15	1.67		20				0.333		
Rice grain	GC	1.1	STMR/STMR-P	88	1.25	20				0.250			
Rice bran/pollard	CM/CF	0.76	STMR/STMR-P	90	0.84	15			20	0.1267			0.169
Soybean hulls	SM	0.02	STMR/STMR-P	90	0.02				5				0.001
Soybean seed	VD	0.01	STMR/STMR-P	89	0.01				15				0.002
Soybean meal	SM	0.004	STMR/STMR-P	92	0.00				5				2E-04
Total						100	100	100	100	12.0425	15.04	10.64	1.516

DAIRY CATTLE													
													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Grape pomace, wet	AB	6.2	STMR/STMR-P	15	41.33		0	20			0	8.267	
Potato process waste	AB	3.8	STMR/STMR-P	12	31.67	10	30			3.1667	9.5		
Potato culls	VR	1.2	STMR/STMR-P	20	6.00	10	30	10		0.6	1.8	0.6	
Rice hulls	CM/CF	3.6	STMR/STMR-P	90	4.00	0		10		0		0.4	
Rice straw	AF/AS	2.2	STMR/STMR-P	90	2.44	0	5	20	25	0	0.122	0.489	0.611
Bean vines	AL	0.75	STMR/STMR-P	35	2.14	0	20	40		0	0.429	0.857	
Beet, mangel fodder	AM/AV	0.25	STMR/STMR-P	15	1.67	0	15			0	0.25		
Almond hulls	AM/AV	1.24	STMR/STMR-P	90	1.38	10				0.1378			
Rice grain	GC	1.1	STMR/STMR-P	88	1.25	20				0.25			
Rice bran/pollard	CM/CF	0.76	STMR/STMR-P	90	0.84	15			10	0.1267			0.084
Wheat straw	AF/AS	0.685	STMR/STMR-P	88	0.78	10				0.0778			
Canola meal	SM	0.03	STMR/STMR-P	88	0.03	10				0.0034			
Cotton undelinted seed	SO	0.021	STMR/STMR-P	88	0.02	10				0.0024			
Soybean hulls	SM	0.02	STMR/STMR-P	90	0.02	5				0.0011			
Soybean seed	VD	0.01	STMR/STMR-P	89	0.01	0			10	0			0.00
Soybean meal	SM	0.004	STMR/STMR-P	92	0.00	0			55	0			0.002
Total						100	100	100	100	4.3659	12.1	10.61	0.699

DIFENOCONAZOLE (224) – ESTIMATED MEAN DIETARY BURDEN													
POULTRY BROILER													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Potato culls	VR	1.2	STMR/STMR-P	20	6.00		10				0.6		
Rice grain	GC	1.1	STMR/STMR-P	88	1.25	20		50		0.25		0.625	
Rice bran/pollard	CM/CF	0.76	STMR/STMR-P	90	0.84	10	10	20	5	0.08	0.084	0.169	0.042
Canola meal	SM	0.03	STMR/STMR-P	88	0.03	15	18	5		0.01	0.006	0.002	
Soybean seed	VD	0.01	STMR/STMR-P	89	0.01	20	20	15		0.00	0.002	0.002	
Sunflower meal	SM	0.01	STMR/STMR-P	92	0.01	10		10		0.00		0.001	
Soybean meal	SM	0.004	STMR/STMR-P	92	0.00		22		35		1E-03		0.002
Total						75	80	100	40	0.3429	0.694	0.798	0.044

POULTRY LAYER													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Potato culls	VR	1.2	STMR/STMR-P	20	6.00		10				0.6		
Rice grain	GC	1.1	STMR/STMR-P	88	1.25	20		50		0.25		0.625	
Rice bran/pollard	CM/CF	0.76	STMR/STMR-P	90	0.84	10	5	20	20	0.0844	0.042	0.169	0.169
Wheat straw	AF/AS	0.685	STMR/STMR-P	88	0.78		10				0.078		
Cabbage heads, leaves	AM/AV	0.035	STMR/STMR-P	15	0.23		5				0.012		
Canola meal	SM	0.03	STMR/STMR-P	88	0.03	15	10	5		0.0051	0.003	0.002	
Soybean seed	VD	0.01	STMR/STMR-P	89	0.01	20	15	15		0.0022	0.002	0.002	
Sunflower meal	SM	0.01	STMR/STMR-P	92	0.01	10		10		0.0011		0.001	
Soybean meal	SM	0.004	STMR/STMR-P	92	0.00		15		30		0.001		0.00
Total						75	70	100	50	0.3429	0.737	0.798	0.17

ETHIPROLE (304) – ESTIMATED MAXIMUM DIETARY BURDEN													
BEEF CATTLE													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Soybean asp gr fn	SM	4.97	STMR	85	5.85	5				0.292			
Rice hulls	CM/CF	0.62	STMR	90	0.69			5				0.034	
Rice bran/pollard	CM/CF	0.53	STMR	90	0.59	15		35	20	0.088		0.206	0.118
Rice grain	GC	0.44	STMR	88	0.50	20		40		0.100		0.2	
Soybean hulls	SM	0.045	STMR	90	0.05	10	10		5	0.005	0.005		0.003
Soybean seed	VD	0.0065	STMR	89	0.01	5	10	20	15	0.000	7E-04	0.001	0.001
Soybean meal	SM	0.002	STMR	92	0.00		10		60		2E-04		0.001
Total						55	30	100	100	0.486	0.006	0.442	0.123

DAIRY CATTLE													
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Rice hulls	CM/CF	0.62	STMR	90	0.69			10				0.069	
Rice bran/pollard	CM/CF	0.53	STMR	90	0.59	15	20	30	10	0.088	0.118	0.177	0.059
Rice grain	GC	0.44	STMR	88	0.50	20		20		0.100		0.1	
Soybean hulls	SM	0.045	STMR	90	0.05	10				0.005			
Soybean seed	VD	0.0065	STMR	89	0.01	10	10	20	10	0.001	7E-04	0.001	7E-04
Soybean meal	SM	0.002	STMR	92	0.00		25	15	60		5E-04	3E-04	0.001
Total						55	55	95	80	0.194	0.119	0.347	0.061

POULTRY BROILER													
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Rice bran/pollard	CM/CF	0.53	STMR	90	0.59	10	10	20	5	0.059	0.059	0.118	0.029
Rice grain	GC	0.44	STMR	88	0.50	20		50		0.100		0.25	
Soybean hulls	SM	0.045	STMR	90	0.05		5				0.003		
Soybean seed	VD	0.0065	STMR	89	0.01	20	20	15		0.001	0.001	0.001	
Soybean meal	SM	0.002	STMR	92	0.00	25	35	15	35	0.001	8E-04	3E-04	8E-04
Total						75	70	100	40	0.161	0.064	0.369	0.03

ETHIPROLE (304) – ESTIMATED MAXIMUM DIETARY BURDEN													
POULTRY LAYER													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Rice bran/pollard	CM/CF	0.53	STMR	90	0.59	10	5	20	20	0.059	0.029	0.118	0.118
Rice grain	GC	0.44	STMR	88	0.50	20		50		0.100		0.25	
Soybean hulls	SM	0.045	STMR	90	0.05		5				0.003		
Soybean seed	VD	0.0065	STMR	89	0.01	20	15	15		0.001	0.001	0.001	
Soybean meal	SM	0.002	STMR	92	0.00	25	20	15	30	0.001	4E-04	3E-04	7E-04
Total						75	45	100	50	0.161	0.033	0.369	0.118

ETHIPROLE (304) – ESTIMATED MEAN DIETARY BURDEN													
BEEF CATTLE													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Soybean asp gr fn	SM	4.97	STMR/STMR-P	85	5.85	5				0.2924			
Rice hulls	CM/CF	0.62	STMR/STMR-P	90	0.69			5				0.034	
Rice bran/pollard	CM/CF	0.53	STMR/STMR-P	90	0.59	15		35	20	0.0883		0.206	0.118
Rice grain	GC	0.44	STMR/STMR-P	88	0.50	20		40		0.1		0.2	
Soybean hulls	SM	0.045	STMR/STMR-P	90	0.05	10	10		5	0.005	0.005		0.003
Soybean seed	VD	0.0065	STMR/STMR-P	89	0.01	5	10	20	15	0.0004	7E-04	0.001	0.001
Soybean meal	SM	0.002	STMR/STMR-P	92	0.00		10		60		2E-04		0.001
Total						55	30	100	100	0.4861	0.006	0.442	0.123

DAIRY CATTLE													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Rice hulls	CM/CF	0.62	STMR/STMR-P	90	0.69		0	10			0	0.069	
Rice bran/pollard	CM/CF	0.53	STMR/STMR-P	90	0.59	15	20	30	10	0.0883	0.118	0.177	0.059
Rice grain	GC	0.44	STMR/STMR-P	88	0.50	20		20		0.1		0.1	
Soybean hulls	SM	0.045	STMR/STMR-P	90	0.05	10				0.005			
Soybean seed	VD	0.0065	STMR/STMR-P	89	0.01	10	10	20	10	0.0007	7E-04	0.001	7E-04
Soybean meal	SM	0.002	STMR/STMR-P	92	0.00	0	25	15	60	0	5E-04	3E-04	0.001
Total						55	55	95	80	0.1941	0.119	0.347	0.061

ETHIPROLE (304) – ESTIMATED MEAN DIETARY BURDEN													
POULTRY BROILER													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Rice bran/pollard	CM/CF	0.53	STMR/STMR-P	90	0.59	10	10	20	5	0.0589	0.059	0.118	0.029
Rice grain	GC	0.44	STMR/STMR-P	88	0.50	20		50		0.10		0.25	
Soybean hulls	SM	0.045	STMR/STMR-P	90	0.05		5				0.003		
Soybean seed	VD	0.0065	STMR/STMR-P	89	0.01	20	20	15		0.0015	0.001	0.001	
Soybean meal	SM	0.002	STMR/STMR-P	92	0.00	25	35	15	35	0.005	0.008	0.008	0.008
Total						75	70	100	40	0.1609	0.0636	0.3692	0.0302

POULTRY LAYER													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Rice bran/pollard	CM/CF	0.53	STMR/STMR-P	90	0.59	10	5	20	20	0.0589	0.0294	0.1178	0.1178
Rice grain	GC	0.44	STMR/STMR-P	88	0.50	20		50		0.1		0.25	
Soybean hulls	SM	0.045	STMR/STMR-P	90	0.05		5				0.0025		
Soybean seed	VD	0.0065	STMR/STMR-P	89	0.01	20	15	15		0.0015	0.001	0.001	
Soybean meal	SM	0.002	STMR/STMR-P	92	0.00	25	20	15	30	0.0005	0.004	0.003	0.007
Total						75	45	100	50	0.1609	0.0335	0.3692	0.1184

FENPICOXAMID (305) – ESTIMATED MAXIMUM DIETARY BURDEN													
BEEF CATTLE													
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Wheat straw	AF/AS	12	HR	88	13.64	10	20	80		1.364	2.727	10.91	
Wheat forage	AF/AS	2.2	HR	25	8.80			20				1.76	
Rye grain	GC	0.0215	STMR	88	0.02	20	40		35	0.005	0.01		0.009
Total						30	60	100	35	1.369	2.737	12.67	0.009

DAIRY CATTLE													
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Wheat straw	AF/AS	12	HR	88	13.64	10	20	20		1.364	2.727	2.727	
Wheat forage	AF/AS	2.2	HR	25	8.80	10		40		0.880		3.52	
Rye grain	GC	0.0215	STMR	88	0.02	20	40		15	0.005	0.01		0.004
Triticale grain	GC	0.0215	STMR	89	0.02			30				0.007	
Total						40	60	90	15	2.2	2.7	6.3	0.00

POULTRY BROILER													
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Rye grain	GC	0.0215	STMR	88	0.02	35	70	50		0.009	0.017	0.012	
Triticale grain	GC	0.0215	STMR	89	0.02	65				0.016			
Total						100	70	50		0.024	0.017	0.012	

POULTRY LAYER													
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Wheat straw	AF/AS	12	HR	88	13.64		10				1.364		
Rye grain	GC	0.0215	STMR	88	0.02	35	35	35		0.009	0.009	0.009	
Triticale grain	GC	0.0215	STMR	89	0.02	65				0.016			
Total						100	45	35		0.024	1.37	0.009	

FENPICOXAMID (305) – ESTIMATED MEAN DIETARY BURDEN													
BEEF CATTLE													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Wheat straw	AF/AS	4.7	STMR/STMR-P	88	5.34	10	20	80		0.5341	1.068	4.273	
Wheat forage	AF/AS	0.8	STMR/STMR-P	25	3.20			20				0.64	
Rye grain	GC	0.0215	STMR/STMR-P	88	0.02	20	40		35	0.0049	0.01		0.009
Total						30	60	100	35	0.539	1.078	4.913	0.009

FENPICOXAMID (305) – ESTIMATED MEAN DIETARY BURDEN													
DAIRY CATTLE													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Wheat straw	AF/AS	4.7	STMR/STMR-P	88	5.34	10	20	20		0.5341	1.068	1.068	
Wheat forage	AF/AS	0.8	STMR/STMR-P	25	3.20	10		40		0.32		1.28	
Rye grain	GC	0.0215	STMR/STMR-P	88	0.02	20	40		15	0.0049	0.01		0.004
Triticale grain	GC	0.0215	STMR/STMR-P	89	0.02	0		30		0		0.007	
Total						40	60	90	15	0.859	1.08	2.36	0.00

FENPICOXAMID (305) – ESTIMATED MEAN DIETARY BURDEN													
POULTRY BROILER													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Rye grain	GC	0.0215	STMR/STMR-P	88	0.02	35	70	50		0.01	0.017	0.012	
Triticale grain	GC	0.0215	STMR/STMR-P	89	0.02	65				0.02			
Total						100	70	50		0.0243	0.017	0.012	

FENPICOXAMID (305) – ESTIMATED MEAN DIETARY BURDEN													
POULTRY LAYER													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Wheat straw	AF/AS	4.7	STMR/STMR-P	88	5.34		10				0.534		
Rye grain	GC	0.0215	STMR/STMR-P	88	0.02	35	35	35		0.0086	0.009	0.009	
Triticale grain	GC	0.0215	STMR/STMR-P	89	0.02	65				0.0157			
Total						100	45	35		0.0243	0.54	0.009	

IMAZALIL (110) – ESTIMATED MAXIMUM DIETARY BURDEN													
BEEF CATTLE													
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Potato process waste	AB	6.6	STMR	12	55.00	30	40	5		16.500	22	2.75	
Potato culls	VR	4.6	HR	20	23.00	30	30	10		6.900	6.9	2.3	
Citrus dried pulp	AB	15	STMR	91	16.48			25				4.121	
Wheat forage	AF/AS	0.01	HR	25	0.04		20	60			0.008	0.024	
Barley forage	AF/AS	0.01	HR	30	0.03		10				0.003		
Barley straw	AF/AS	0.01	HR	89	0.01	10				0.001			
Barley grain	GC	0.01	STMR	88	0.01	30			70	0.003			0.008
Total						100	100	100	70	23.405	28.91	9.195	0.008

DAIRY CATTLE													
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Potato process waste	AB	6.6	STMR	12	55.00	10	30			5.500	16.5		
Potato culls	VR	4.6	HR	20	23.00	10	30	10		2.300	6.9	2.3	
Citrus dried pulp	AB	15	STMR	91	16.48			30				4.945	
Wheat forage	AF/AS	0.01	HR	25	0.04	20	20	60		0.008	0.008	0.024	
Barley forage	AF/AS	0.01	HR	30	0.03		20				0.007		
Barley grain	GC	0.01	STMR	88	0.01	45			40	0.005			0.005
Total						85	100	100	40	7.813	23.41	7.269	0.005

POULTRY BROILER													
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Potato culls	VR	4.6	HR	20	23.00		10				2.3		
Barley grain	GC	0.01	STMR	88	0.01	75	70	15	10	0.009	0.008	0.002	0.001
Wheat grain	GC	0.01	STMR	89	0.01			55				0.006	
Total						75	80	70	10	0.009	2.308	0.008	0.001

IMAZALIL (110) – ESTIMATED MAXIMUM DIETARY BURDEN													
POULTRY LAYER													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Potato culls	VR	4.6	HR	20	23.00		10				2.3		
Wheat forage	AF/AS	0.01	HR	25	0.04		10				0.004		
Barley grain	GC	0.01	STMR	88	0.01	75	80	15		0.009	0.009	0.002	
Wheat grain	GC	0.01	STMR	89	0.01			40				0.004	
Total						75	100	55		0.009	2.313	0.006	

IMAZALIL (110) – ESTIMATED MEAN DIETARY BURDEN													
BEEF CATTLE													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Potato process waste	AB	6.6	STMR/STMR-P	12	55.00	30	40	5		16.5	22	2.75	
Citrus dried pulp	AB	15	STMR/STMR-P	91	16.48			25				4.121	
Potato culls	VR	2.2	STMR/STMR-P	20	11.00	30	30	10		3.3	3.3	1.1	
Wheat forage	AF/AS	0.01	STMR/STMR-P	25	0.04		20	60			0.008	0.024	
Barley forage	AF/AS	0.01	STMR/STMR-P	30	0.03		10				0.003		
Barley straw	AF/AS	0.01	STMR/STMR-P	89	0.01	10				0.0011			
Barley grain	GC	0.01	STMR/STMR-P	88	0.01	30			70	0.0034			0.008
Total						100	100	100	70	19.805	25.31	7.995	0.008

DAIRY CATTLE													
DAIRY CATTLE													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Potato process waste	AB	6.6	STMR/STMR-P	12	55.00	10	30	0		5.5	16.5	0	
Citrus dried pulp	AB	15	STMR/STMR-P	91	16.48	0		30		0		4.945	
Potato culls	VR	2.2	STMR/STMR-P	20	11.00	10	30	10		1.1	3.3	1.1	
Wheat forage	AF/AS	0.01	STMR/STMR-P	25	0.04	20	20	60		0.008	0.008	0.024	
Barley forage	AF/AS	0.01	STMR/STMR-P	30	0.03	0	20			0	0.007		
Barley grain	GC	0.01	STMR/STMR-P	88	0.01	45			40	0.0051			0.005
Total						85	100	100	40	6.6131	19.81	6.069	0.005

IMAZALIL (110) – ESTIMATED MEAN DIETARY BURDEN													
POULTRY BROILER													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Potato culls	VR	2.2	STMR/STMR-P	20	11.00		10				1.1		
Barley grain	GC	0.01	STMR/STMR-P	88	0.01	75	70	15	10	0.0085	0.008	0.002	0.001
Wheat grain	GC	0.01	STMR/STMR-P	89	0.01			55				0.006	
Total						75	80	70	10	0.0085	1.108	0.008	0.001

POULTRY LAYER													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Potato culls	VR	2.2	STMR/STMR-P	20	11.00		10				1.1		
Wheat forage	AF/AS	0.01	STMR/STMR-P	25	0.04		10				0.004		
Barley grain	GC	0.01	STMR/STMR-P	88	0.01	75	80	15		0.0085	0.009	0.002	
Wheat grain	GC	0.01	STMR/STMR-P	89	0.01			40				0.004	
Total						75	100	55		0.0085	1.113	0.006	

ISOXAFLUTOLE (268) – ESTIMATED MAXIMUM DIETARY BURDEN													
BEEF CATTLE													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Corn, field forage/silage	AF/AS	0.34	HR	40	0.85	15	80	80		0.128	0.68	0.68	
Soybean asp gr fn	SM	0.12	STM	85	0.14	5				0.007			
Sugarcane tops	AM/AV	0.01	HR	25	0.04			20				0.008	
Soybean meal	SM	0.024	STM	92	0.03		20		65		0.005		0.017
Corn, field grain	GC	0.02	STM	88	0.02	80			35	0.018			0.008
Total						100	100	100	100	0.153	0.685	0.688	0.025

DAIRY CATTLE													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Corn, field forage/silage	AF/AS	0.34	HR	40	0.85	45	60	80	50	0.383	0.51	0.68	0.425
Sugarcane tops	AM/AV	0.01	HR	25	0.04			20				0.008	
Soybean meal	SM	0.024	STM	92	0.03	10	25		50	0.003	0.007		0.013
Corn, field grain	GC	0.02	STM	88	0.02	45	15			0.010	0.003		
Total						100	100	100	100	0.395	0.52	0.688	0.438

POULTRY BROILER													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Soybean meal	SM	0.024	STM	92	0.03	25	40	25	35	0.007	0.01	0.007	0.009
Corn, field grain	GC	0.02	STM	88	0.02	75	60		65	0.017	0.014		0.015
Soybean seed	VD	0.02	STM	89	0.02			15				0.003	
Total						100	100	40	100	0.024	0.024	0.01	0.024

POULTRY LAYER													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Soybean meal	SM	0.024	STM	92	0.03	25	25	25	30	0.007	0.007	0.007	0.008
Corn, field grain	GC	0.02	STM	88	0.02	75	70		70	0.017	0.016		0.016
Soybean seed	VD	0.02	STM	89	0.02		5	15			0.001	0.003	
Total						100	100	40	100	0.024	0.024	0.01	0.024

ISOXAFLUTOLE (268) – ESTIMATED MEAN DIETARY BURDEN													
BEEF CATTLE													
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Soybean asp gr fn	SM	0.12	STMR/STMR-P	85	0.14	5				0.0071			
Corn, field forage/silage	AF/AS	0.02	STMR/STMR-P	40	0.05	15	80	80		0.0075	0.04	0.04	
Soybean meal	SM	0.024	STMR/STMR-P	92	0.03		20	10	65		0.005	0.003	0.017
Corn, field grain	GC	0.02	STMR/STMR-P	88	0.02	80		10	35	0.0182		0.002	0.008
Total						100	100	100	100	0.0327	0.045	0.045	0.025

DAIRY CATTLE													
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Corn, field forage/silage	AF/AS	0.02	STMR/STMR-P	40	0.05	45	60	80	50	0.0225	0.03	0.04	0.025
Soybean meal	SM	0.024	STMR/STMR-P	92	0.03	10	25	15	50	0.0026	0.007	0.004	0.013
Corn, field grain	GC	0.02	STMR/STMR-P	88	0.02	45	15	5		0.0102	0.003	0.001	
Total						100	100	100	100	0.0353	0.04	0.045	0.038

POULTRY BROILER													
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Soybean meal	SM	0.024	STMR/STMR-P	92	0.03	25	40	25	35	0.0065	0.01	0.007	0.009
Corn, field grain	GC	0.02	STMR/STMR-P	88	0.02	75	60		65	0.0170	0.014		0.015
Soybean seed	VD	0.02	STMR/STMR-P	89	0.02			15				0.003	
Total						100	100	40	100	0.0236	0.024	0.01	0.024

POULTRY LAYER													
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Soybean meal	SM	0.024	STMR/STMR-P	92	0.03	25	25	25	30	0.0065	0.007	0.007	0.008
Corn, field grain	GC	0.02	STMR/STMR-P	88	0.02	75	70		70	0.0170	0.016		0.016
Soybean seed	VD	0.02	STMR/STMR-P	89	0.02		5	15			0.001	0.003	
Total						100	100	40	100	0.0236	0.024	0.01	0.024

MANDIPROPAMID (231) – ESTIMATED MAXIMUM DIETARY BURDEN													
BEEF CATTLE													
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Kale leaves	AM/AV	11.5	HR	15	76.67		20				15.33		
Bean vines	AL	9.27	HR	35	26.49			60				15.89	
Grape pomace, wet	AB	1.16	STMR	15	7.73			20				1.547	
Potato culls	VR	0.073	HR	20	0.37	30	30	10		0.110	0.11	0.037	
Potato process waste	AB	0.04	STMR	12	0.33	30	40			0.100	0.133		
Citrus dried pulp	AB	0.29	STMR	91	0.32			10				0.032	
Total						60	90	100		0.210	15.58	17.51	

DAIRY CATTLE													
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Kale leaves	AM/AV	11.5	HR	15	76.67		20	40			15.33	30.67	
Bean vines	AL	9.27	HR	35	26.49		20	60			5.297	18.54	
Potato culls	VR	0.073	HR	20	0.37	10	30			0.037	0.11		
Potato process waste	AB	0.04	STMR	12	0.33	10	30			0.033	0.1		
Total						20	100	100		0.070	20.84	49.21	

POULTRY BROILER													
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Potato culls	VR	0.073	HR	20	0.37		10				0.037		
Potato dried pulp	AB	0.019	STMR	88	0.02		20				0.004		
Total							30				0.041		

POULTRY LAYER													
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Cabbage heads, leaves	AM/AV	5.8	HR	15	38.67		5				1.933		
Potato culls	VR	0.073	HR	20	0.37		10				0.037		
Potato dried pulp	AB	0.019	STMR	88	0.02		15				0.003		
Total							30				1.973		

MANDIPROPAMID (231) – ESTIMATED MEAN DIETARY BURDEN													
BEEF CATTLE													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Kale leaves	AM/AV	5.65	STMR/STMR-P	15	37.67		20				7.533		
Bean vines	AL	4.35	STMR/STMR-P	35	12.43			60				7.457	
Grape pomace, wet	AB	1.16	STMR/STMR-P	15	7.73			20				1.547	
Potato process waste	AB	0.04	STMR/STMR-P	12	0.33	30	40			0.1	0.133		
Citrus dried pulp	AB	0.29	STMR/STMR-P	91	0.32			20				0.064	
Potato culls	VR	0.019	STMR/STMR-P	20	0.10	30	30			0.0285	0.029		
Total						60	90	100		0.1285	7.695	9.067	

DAIRY CATTLE													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Kale leaves	AM/AV	5.65	STMR/STMR-P	15	37.67		20	40			7.533	15.07	
Bean vines	AL	4.35	STMR/STMR-P	35	12.43	0	20	60		0	2.486	8.7	
Potato process waste	AB	0.04	STMR/STMR-P	12	0.33	10	30			0.0333	0.1		
Potato culls	VR	0.019	STMR/STMR-P	20	0.10	10	30			0.0095	0.029		
Total						20	100	100		0.0428	10.15	23.77	

POULTRY BROILER													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Potato culls	VR	0.019	STMR/STMR-P	20	0.10		10				0.01		
Potato dried pulp	AB	0.019	STMR/STMR-P	88	0.02		20				0.004		
Total							30				0.014		

MANDIPROPAMID (231) – ESTIMATED MEAN DIETARY BURDEN													
POULTRY LAYER													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US- CAN	EU	AU	JP	US- CAN	EU	AU	JP
Cabbage heads, leaves	AM/AV	3.55	STMR/STMR- P	15	23.67		5				1.183		
Potato culls	VR	0.019	STMR/STMR- P	20	0.10		10				0.01		
Potato dried pulp	AB	0.019	STMR/STMR- P	88	0.02		15				0.003		
Total							30				1.196		

PROTHIOCONAZOLE (232) – ESTIMATED MAXIMUM DIETARY BURDEN (relevant to the estimation of livestock residues according to the risk assessment definition)													
BEEF CATTLE													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US- CAN	EU	AU	JP	US- CAN	EU	AU	JP
Peanut hay	AL	27.84	HR	85	32.75			60				19.65	
Barley forage	AF/AS	8.64	HR	30	28.80		30	40			8.64	11.52	
Beet, sugar tops	AM/AV	6.24	HR	23	27.13		20				5.426		
Corn, field stover	AF/AS	14.07	HR	83	16.95	15				2.543			
Corn, field forage/silage	AF/AS	6.528	HR	40	16.32		50				8.16		
Wheat milled bypds	CM/CF	8.5	STM	88	9.66	40			55	3.864			5.313
Soybean asp gr fn	SM	6.375	STM	85	7.50	5				0.375			
Rice straw	AF/AS	4.94	HR	90	5.49				45				2.47
Cotton gin byproducts	AM/AV	4.68	HR	90	5.20	5				0.260			
Soybean seed	VD	0.085	STM	89	0.10	5				0.005			
Barley grain	GC	0.0595	STM	88	0.07	30				0.020			
Total						100	100	100	100	7.066	22.23	31.17	7.783

DAIRY CATTLE													
													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US- CAN	EU	AU	JP	US- CAN	EU	AU	JP
Peanut hay	AL	27.84	HR	85	32.75	15		60		4.913		19.65	
Barley forage	AF/AS	8.64	HR	30	28.80		30	40			8.64	11.52	
Beet, sugar tops	AM/AV	6.24	HR	23	27.13		30				8.139		
Corn, field stover	AF/AS	14.07	HR	83	16.95	15				2.543			
Corn, field forage/silage	AF/AS	6.528	HR	40	16.32	30	40		50	4.896	6.528		8.16
Corn, sweet forage	AF/AS	6.528	HR	48	13.60	40				5.440			
Wheat milled bypds	CM/CF	8.5	STM	88	9.66				45				4.347
Rice straw	AF/AS	4.94	HR	90	5.49				5				0.274
Total						100	100	100	100	17.792	23.31	31.17	12.78

POULTRY BROILER													
													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US- CAN	EU	AU	JP	US- CAN	EU	AU	JP
Wheat milled bypds	CM/CF	8.5	STM	88	9.66	50	20	20	5	4.830	1.932	1.932	0.483
Bean seed	VD	0.085	STM	88	0.10		20	70			0.019	0.068	
Cowpea seed	VD	0.085	STM	88	0.10	10				0.010			
Soybean seed	VD	0.085	STM	89	0.10	10				0.010			
Barley grain	GC	0.0595	STM	88	0.07	30	60	10	10	0.020	0.041	0.007	0.007
Rape meal	SM	0.04641	STM	88	0.05				5				0.003
Corn, field grain	GC	0.0306	STM	88	0.03				60				0.021
Corn, pop grain	GC	0.0306	STM	88	0.03				20				0.007
Total						100	100	100	100	4.869	1.992	2.006	0.52

PROTHIOCONAZOLE (232) – ESTIMATED MAXIMUM DIETARY BURDEN (relevant to the estimation of livestock residues according to the risk assessment definition)													
POULTRY LAYER													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US- CAN	EU	AU	JP	US- CAN	EU	AU	JP
Beet, sugar tops	AM/AV	6.24	HR	23	27.13		5				1.357		
Wheat milled bypdts	CM/CF	8.5	STMR	88	9.66	50	20	20	30	4.830	1.932	1.932	2.898
Wheat straw	AF/AS	4.94	HR	88	5.61		10				0.561		
Bean seed	VD	0.085	STMR	88	0.10		20	70			0.019	0.068	
Cowpea seed	VD	0.085	STMR	88	0.10	10				0.010			
Soybean seed	VD	0.085	STMR	89	0.10	10				0.010			
Barley grain	GC	0.0595	STMR	88	0.07	30	45	10		0.020	0.03	0.007	
Rape meal	SM	0.04641	STMR	88	0.05				15				0.008
Corn, field grain	GC	0.0306	STMR	88	0.03				55				0.02
Total						100	100	100	100	4.869	3.899	2.006	2.925

PROTHIOCONAZOLE (232) – ESTIMATED MEAN DIETARY BURDEN (relevant to the estimation of livestock residues according to the risk assessment definition)													
BEEF CATTLE													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US- CAN	EU	AU	JP	US-CAN	EU	AU	JP
Peanut hay	AL	9.792	STMR/STMR-P	85	11.52			60				6.912	
Beet, sugar tops	AM/AV	2.4	STMR/STMR-P	23	10.43		20				2.087		
Wheat milled bypdts	CM/CF	8.5	STMR/STMR-P	88	9.66	40	30	40	55	3.8636	2.898	3.864	5.313
Corn, field stover	AF/AS	7.308	STMR/STMR-P	83	8.80	15	25			1.3207	2.201		
Corn, field forage/silage	AF/AS	3.44	STMR/STMR-P	40	8.60		25				2.15		
Soybean asp gr fn	SM	6.375	STMR/STMR-P	85	7.50	5				0.375			
Cotton gin byproducts	AM/AV	2.86	STMR/STMR-P	90	3.18	5				0.1589			
Rice straw	AF/AS	1.69	STMR/STMR-P	90	1.88				45				0.845
Soybean seed	VD	0.085	STMR/STMR-P	89	0.10	5				0.005			
Barley grain	GC	0.0595	STMR/STMR-P	88	0.07	30				0.020			
Total						100	100	100	100	5.743	9.336	10.78	6.158

PROTHIOCONAZOLE (232) – ESTIMATED MEAN DIETARY BURDEN (relevant to the estimation of livestock residues according to the risk assessment definition)													
DAIRY CATTLE													
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US- CAN	EU	AU	JP	US-CAN	EU	AU	JP
Peanut hay	AL	9.792	STMR/STMR-P	85	11.52	15	0	60		1.728	0	6.912	
Beet, sugar tops	AM/AV	2.4	STMR/STMR-P	23	10.43	0	30			0	3.13		
Wheat milled bypds	CM/CF	8.5	STMR/STMR-P	88	9.66	30	30	40	45	2.8977	2.898	3.864	4.347
Corn, field stover	AF/AS	7.308	STMR/STMR-P	83	8.80	15	20			1.3207	1.761		
Corn, field forage/silage	AF/AS	3.44	STMR/STMR-P	40	8.60	40	20		50	3.44	1.72		4.3
Rice straw	AF/AS	1.69	STMR/STMR-P	90	1.88	0			5	0			0.094
Total						100	100	100	100	9.3864	9.509	10.78	8.74

POULTRY ROILER													
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US- CAN	EU	AU	JP	US-CAN	EU	AU	JP
Wheat milled bypds	CM/CF	8.5	STMR/STMR-P	88	9.66	50	20	20	5	4.830	1.932	1.932	0.483
Bean seed	VD	0.085	STMR/STMR-P	88	0.10		20	70			0.019	0.068	
Cowpea seed	VD	0.085	STMR/STMR-P	88	0.10	10				0.010			
Soybean seed	VD	0.085	STMR/STMR-P	89	0.10	10				0.010			
Barley grain	GC	0.0595	STMR/STMR-P	88	0.07	30	60	10	10	0.02	0.041	0.007	0.007
Rape meal	SM	0.04641	STMR/STMR-P	88	0.05				5				0.003
Corn, field grain	GC	0.0306	STMR/STMR-P	88	0.03				60				0.021
Corn, pop grain	GC	0.0306	STMR/STMR-P	88	0.03				20				0.007
Total						100	100	100	100	4.870	1.992	2.006	0.52

POULTRY LAYER													
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US- CAN	EU	AU	JP	US-CAN	EU	AU	JP
Beet, sugar tops	AM/AV	2.4	STMR/STMR-P	23	10.43		5				0.522		
Wheat milled bypds	CM/CF	8.5	STMR/STMR-P	88	9.66	50	20	20	30	4.8296	1.932	1.932	2.898
Wheat straw	AF/AS	1.69	STMR/STMR-P	88	1.92		10				0.192		
Bean seed	VD	0.085	STMR/STMR-P	88	0.10		20	70			0.019	0.068	
Cowpea seed	VD	0.085	STMR/STMR-P	88	0.10	10				0.0097			
Soybean seed	VD	0.085	STMR/STMR-P	89	0.10	10				0.0096			
Barley grain	GC	0.0595	STMR/STMR-P	88	0.07	30	45	10		0.0203	0.03	0.007	

POULTRY LAYER													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Rape meal	SM	0.04641	STMR/STMR-P	88	0.05				15				0.008
Corn, field grain	GC	0.0306	STMR/STMR-P	88	0.03				55				0.02
Total						100	100	100	100	4.8692	2.695	2.006	2.925

PROTHIOCONAZOLE (232) – ESTIMATED MAXIMUM DIETARY BURDEN (relevant to the estimation of maximum residue levels)													
BEEF CATTLE													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Barley forage	AF/AS	5.4	HR	30	18.00		30	50			5.4	9	
Beet, sugar tops	AM/AV	3.9	HR	23	16.96		20				3.391		
Peanut hay	AL	11.6	HR	85	13.65			50				6.824	
Corn, field forage/silage	AF/AS	4.08	HR	40	10.20	15	50			1.530	5.1		
Wheat milled bypds	CM/CF	5	STMR	88	5.68	40			55	2.273			3.125
Soybean asp gr fn	SM	3.75	STMR	85	4.41	5				0.221			
Cotton gin byproducts	AM/AV	1.8	HR	90	2.00	5				0.100			
Soybean seed	VD	0.05	STMR	89	0.06	5			15	0.003			0.008
Potato culls	VR	0.01	HR	20	0.05	30				0.015			
Barley grain	GC	0.035	STMR	88	0.04				30				0.012
Total						100	100	100	100	4.141	13.89	15.82	3.145

DAIRY CATTLE													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Barley forage	AF/AS	5.4	HR	30	18.00		30	50			5.4	9	
Beet, sugar tops	AM/AV	3.9	HR	23	16.96		30				5.087		
Peanut hay	AL	11.6	HR	85	13.65	15		50		2.047		6.824	
Corn, field forage/silage	AF/AS	4.08	HR	40	10.20	45	40		50	4.590	4.08		5.1
Corn, sweet forage	AF/AS	4.08	HR	48	8.50	40				3.400			
Wheat milled bypds	CM/CF	5	STMR	88	5.68				45				2.557
Soybean seed	VD	0.05	STMR	89	0.06				5				0.003
Total						100	100	100	100	10.037	14.57	15.82	7.66

PROTHIOCONAZOLE (232) – ESTIMATED MAXIMUM DIETARY BURDEN (relevant to the estimation of maximum residue levels)													
POULTRY BROILER													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US- CAN	EU	AU	JP	US- CAN	EU	AU	JP
Wheat milled bypdts	CM/CF	5	STMR	88	5.68	50	20	20	5	2.841	1.136	1.136	0.284
Bean seed	VD	0.05	STMR	88	0.06		20	70			0.011	0.04	
Cowpea seed	VD	0.05	STMR	88	0.06	10				0.006			
Pea seed	VD	0.05	STMR	90	0.06	10				0.006			
Potato culls	VR	0.01	HR	20	0.05		10				0.005		
Barley grain	GC	0.035	STMR	88	0.04	30	50	10	10	0.012	0.02	0.004	0.004
Rape meal	SM	0.0273	STMR	88	0.03				5				0.002
Corn, field grain	GC	0.018	STMR	88	0.02				60				0.012
Corn, pop grain	GC	0.018	STMR	88	0.02				20				0.004
Total						100	100	100	100	2.864	1.173	1.18	0.306

POULTRY LAYER													
POULTRY LAYER													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US- CAN	EU	AU	JP	US- CAN	EU	AU	JP
Beet, sugar tops	AM/AV	3.9	HR	23	16.96		5				0.848		
Wheat milled bypdts	CM/CF	5	STMR	88	5.68	50	20	20	30	2.841	1.136	1.136	1.705
Wheat straw	AF/AS	1.9	HR	88	2.16		10				0.216		
Bean seed	VD	0.05	STMR	88	0.06		20	70			0.011	0.04	
Cowpea seed	VD	0.05	STMR	88	0.06	10				0.006			
Pea seed	VD	0.05	STMR	90	0.06	10				0.006			
Potato culls	VR	0.01	HR	20	0.05		10				0.005		
Barley grain	GC	0.035	STMR	88	0.04	30	35	10		0.012	0.014	0.004	
Rape meal	SM	0.0273	STMR	88	0.03				15				0.00
Corn, field grain	GC	0.018	STMR	88	0.02				55				0.011
Total						100	100	100	100	2.864	2.23	1.18	1.72

PROTHIOCONAZOLE (232) – ESTIMATED MEAN DIETARY BURDEN (relevant to the estimation of maximum residue levels)													
BEEF CATTLE													
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US- CAN	EU	AU	JP	US-CAN	EU	AU	JP
Beet, sugar tops	AM/AV	1.5	STMR/STMR-P	23	6.52		20				1.304		
Wheat milled bypds	CM/CF	5	STMR/STMR-P	88	5.68	40	30	40	55	2.2727	1.705	2.273	3.125
Corn, field forage/silage	AF/AS	2.15	STMR/STMR-P	40	5.38	15	50	60		0.8063	2.688	3.225	
Soybean asp gr fn	SM	3.75	STMR/STMR-P	85	4.41	5				0.2206			
Cotton gin byproducts	AM/AV	1.1	STMR/STMR-P	90	1.22	5				0.0611			
Soybean seed	VD	0.05	STMR/STMR-P	89	0.06	5			15	0.0028			0.008
Potato culls	VR	0.01	STMR/STMR-P	20	0.05	30				0.015			
Barley grain	GC	0.035	STMR/STMR-P	88	0.04				30				0.012
Total						100	100	100	100	3.3785	5.696	5.498	3.145

DAIRY CATTLE													
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US- CAN	EU	AU	JP	US-CAN	EU	AU	JP
Beet, sugar tops	AM/AV	1.5	STMR/STMR-P	23	6.52		30	0			1.957	0	
Wheat milled bypds	CM/CF	5	STMR/STMR-P	88	5.68	30	30	40	45	1.7046	1.705	2.273	2.557
Corn, field forage/silage	AF/AS	2.15	STMR/STMR-P	40	5.38	45	40	60	50	2.4188	2.15	3.225	2.688
Peanut hay	AL	4.08	STMR/STMR-P	85	4.80	15				0.72			
Soybean seed	VD	0.05	STMR/STMR-P	89	0.06	10			5	0.0056			0.003
Total						100	100	100	100	4.8489	5.811	5.498	5.247

POULTRY BROILER													
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US- CAN	EU	AU	JP	US-CAN	EU	AU	JP
Wheat milled bypds	CM/CF	5	STMR/STMR-P	88	5.68	50	20	20	5	2.8409	1.136	1.136	0.284
Bean seed	VD	0.05	STMR/STMR-P	88	0.06		20	70			0.011	0.04	
Cowpea seed	VD	0.05	STMR/STMR-P	88	0.06	10				0.0057			
Pea seed	VD	0.05	STMR/STMR-P	90	0.06	10				0.0056			
Potato culls	VR	0.01	STMR/STMR-P	20	0.05		10				0.005		
Barley grain	GC	0.035	STMR/STMR-P	88	0.04	30	50	10	10	0.0119	0.02	0.004	0.004
Rape meal	SM	0.0273	STMR/STMR-P	88	0.03				5				0.002
Corn, field grain	GC	0.018	STMR/STMR-P	88	0.02				60				0.012

POULTRY BROILER													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Corn, pop grain	GC	0.018	STMR/STMR-P	88	0.02				20				0.004
Total						100	100	100	100	2.8641	1.173	1.18	0.306

PROTHIOCONAZOLE (232) – ESTIMATED MEAN DIETARY BURDEN
(relevant to the estimation of maximum residue levels)

POULTRY LAYER													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Beet, sugar tops	AM/AV	1.5	STMR/STMR-P	23	6.52		5				0.326		
Wheat milled bypds	CM/CF	5	STMR/STMR-P	88	5.68	50	20	20	30	2.8409	1.136	1.136	1.705
Wheat straw	AF/AS	0.65	STMR/STMR-P	88	0.74		10				0.074		
Bean seed	VD	0.05	STMR/STMR-P	88	0.06		20	70			0.011	0.04	
Cowpea seed	VD	0.05	STMR/STMR-P	88	0.06	10				0.0057			
Pea seed	VD	0.05	STMR/STMR-P	90	0.06	10				0.0056			
Potato culls	VR	0.01	STMR/STMR-P	20	0.05		10				0.005		
Barley grain	GC	0.035	STMR/STMR-P	88	0.04	30	35	10		0.0119	0.014	0.004	
Rape meal	SM	0.0273	STMR/STMR-P	88	0.03				15				0.00
Corn, field grain	GC	0.018	STMR/STMR-P	88	0.02				55				0.011
Total						100	100	100	100	2.8641	1.567	1.18	1.72

PYDIFLUMETOFEN (309) – ESTIMATED MAXIMUM DIETARY BURDEN													
BEEF CATTLE													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Rye straw	AF/AS	40	HR	88	45.45	10	20	20		4.545	9.091	9.091	
Triticale hay	AF/AS	40	HR	88	45.45	5		80		2.273		36.36	
Barley straw	AF/AS	40	HR	100	40.00		10				4		
Sorghum, grain asp gr fn	CM/CF	31.9	STMR	85	37.53	5				1.876			
Beet, sugar tops	AM/AV	8	HR	23	34.78		20				6.957		
Turnip tops (leaves)	AM/AV	8	HR	30	26.67		20				5.333		
Trefoil hay	AL	15	HR	85	17.65	15	20			2.647	3.529		
Vetch hay	AL	15	HR	85	17.65		10		5		1.765		0.882
Soybean asp gr fn	SM	3.9	STMR	85	4.59	5				0.229			
Sorghum, grain grain	GC	0.515	STMR	86	0.60	40			35	0.240			0.21
Potato process waste	AB	0.063	STMR	12	0.53	20				0.105			
Wheat milled bypds	CM/CF	0.38	STMR	88	0.43				55				0.238
Rice straw	AF/AS	0.28	HR	90	0.31				5				0.016
Total						100	100	100	100	11.916	30.67	45.45	1.35

DAIRY CATTLE													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Rye straw	AF/AS	40	HR	88	45.45	10	20	20	5	4.545	9.091	9.091	2.273
Triticale hay	AF/AS	40	HR	88	45.45	10		50		4.545		22.73	
Oat straw	AF/AS	40	HR	90	44.44			30				13.33	
Barley straw	AF/AS	40	HR	100	40.00		10				4		
Oat hay	AF/AS	40	HR	100	40.00	10				4.000			
Beet, sugar tops	AM/AV	8	HR	23	34.78		30				10.43		
Turnip tops (leaves)	AM/AV	8	HR	30	26.67	30				8.000			
Soybean hay	AL	15	HR	85	17.65	20				3.529			
Trefoil hay	AL	15	HR	85	17.65	20	40			3.529	7.059		
Vetch hay	AL	15	HR	85	17.65				25				4.412
Corn, field forage/silage	AF/AS	4.9	HR	40	12.25				45				5.513
Sorghum, grain forage	AF/AS	0.73	HR	35	2.09				25				0.521
Total						100	100	100	100	28.150	30.58	45.15	12.72

PYDIFLUMETOFEN (309) – ESTIMATED MAXIMUM DIETARY BURDEN													
POULTRY BROILER													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Swede roots	VR	0.25	HR	10	2.50		10				0.25		
Sorghum, grain	GC	0.515	STMR	86	0.60	75	70	70	65	0.449	0.419	0.419	0.389
Wheat milled bypds	CM/CF	0.38	STMR	88	0.43	25	20	20	5	0.108	0.086	0.086	0.022
Corn, field grain	GC	0.03	STMR	88	0.03				30				0.01
Soybean seed	VD	0.028	STMR	89	0.03			10				0.003	
Total						100	100	100	100	0.557	0.756	0.509	0.421

POULTRY LAYER													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Wheat hay	AF/AS	40	HR	88	45.45		10				4.545		
Beet, sugar tops	AM/AV	8	HR	23	34.78		5				1.739		
Soybean hay	AL	15	HR	85	17.65		10				1.765		
Swede roots	VR	0.25	HR	10	2.50		10				0.25		
Sorghum, grain	GC	0.515	STMR	86	0.60	75	65	70	55	0.449	0.389	0.419	0.329
Wheat milled bypds	CM/CF	0.38	STMR	88	0.43	25		20	30	0.108		0.086	0.13
Corn, field grain	GC	0.03	STMR	88	0.03				15				0.005
Soybean seed	VD	0.028	STMR	89	0.03			10				0.003	
Total						100	100	100	100	0.56	8.689	0.509	0.46

PYDIFLUMETOFEN (309) – ESTIMATED MEAN DIETARY BURDEN													
BEEF CATTLE													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Sorghum, grain asp gr fn	CM/CF	31.9	STMR/STMR-P	85	37.53	5		20		1.8765		7.506	
Peanut hay	AL	15	STMR/STMR-P	100	15.00			60				9	
Soybean hay	AL	9.2	STMR/STMR-P	85	10.82			20				2.165	
Trefoil hay	AL	9.2	STMR/STMR-P	85	10.82	15	20			1.6235	2.165		
Vetch hay	AL	9.2	STMR/STMR-P	85	10.82		5		5		0.541		0.541
Wheat forage	AF/AS	2.7	STMR/STMR-P	25	10.80		20				2.16		
Rye straw	AF/AS	9.2	STMR/STMR-P	88	10.45	10				1.0455			
Triticale hay	AF/AS	9.2	STMR/STMR-P	88	10.45	5				0.5227			
Barley straw	AF/AS	9.2	STMR/STMR-P	100	9.20		10				0.92		

PYDIFLUMETOFEN (309) – ESTIMATED MEAN DIETARY BURDEN													
BEEF CATTLE													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
			P										
Cowpea hay	AL	9.2	STMR/STMR-P	100	9.20		10				0.92		
Beet, sugar tops	AM/AV	1.5	STMR/STMR-P	23	6.52		20				1.304		
Barley forage	AF/AS	1.95	STMR/STMR-P	30	6.50		15				0.975		
Soybean aspr fin	SM	3.9	STMR/STMR-P	85	4.59	5				0.229			
Sorghum, grain grain	GC	0.515	STMR/STMR-P	86	0.60	40			35	0.240			0.21
Potato process waste	AB	0.063	STMR/STMR-P	12	0.53	20				0.1050			
Wheat milled bypds	CM/CF	0.38	STMR/STMR-P	88	0.43				55				0.238
Beet, sugar dried pulp	AB	0.24	STMR/STMR-P	88	0.27				5				0.014
Total						100	100	100	100	5.6421	8.985	18.67	1.002

DAIRY CATTLE													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Peanut hay	AL	15	STMR/STMR-P	100	15.00	15	0	60		2.25	0	9	
Soybean hay	AL	9.2	STMR/STMR-P	85	10.82	5				0.5412			
Trefoil hay	AL	9.2	STMR/STMR-P	85	10.82	20	40			2.1647	4.329		
Vetch hay	AL	9.2	STMR/STMR-P	85	10.82	0			25	0			2.706
Wheat forage	AF/AS	2.7	STMR/STMR-P	25	10.80	20	20	40		2.16	2.16	4.32	
Rye straw	AF/AS	9.2	STMR/STMR-P	88	10.45	0			5	0			0.523
Barley straw	AF/AS	9.2	STMR/STMR-P	100	9.20	0	10			0	0.92		
Cowpea hay	AL	9.2	STMR/STMR-P	100	9.20	0	30			0	2.76		
Oat hay	AF/AS	9.2	STMR/STMR-P	100	9.20	10				0.92			
Turnip tops (leaves)	AM/AV	1.5	STMR/STMR-P	30	5.00	30				1.5			
Corn, field forage/silage	AF/AS	1.15	STMR/STMR-P	40	2.88	0			45	0			1.294
Sorghum, grain forage	AF/AS	0.23	STMR/STMR-P	35	0.66	0			25	0			0.164
Total						100	100	100	100	9.5359	10.17	13.32	4.687

PYDIFLUMETOFEN (309) – ESTIMATED MEAN DIETARY BURDEN													
POULTRY BROILER													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Swede roots	VR	0.08	STMR/STMR-P	10	0.80		10				0.08		
Sorghum, grain grain	GC	0.515	STMR/STMR-P	86	0.60	75	70	70	65	0.45	0.419	0.419	0.389
Wheat milled bypds	CM/CF	0.38	STMR/STMR-P	88	0.43	25	20	20	5	0.11	0.086	0.086	0.022
Corn, field grain	GC	0.03	STMR/STMR-P	88	0.03				30				0.01
Soybean seed	VD	0.028	STMR/STMR-P	89	0.03			10				0.003	
Total						100	100	100	100	0.56	0.586	0.509	0.421

POULTRY LAYER													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Soybean hay	AL	9.2	STMR/STMR-P	85	10.82		10				1.082		
Wheat forage	AF/AS	2.7	STMR/STMR-P	25	10.80		10				1.08		
Beet, sugar tops	AM/AV	1.5	STMR/STMR-P	23	6.52		5				0.326		
Swede roots	VR	0.08	STMR/STMR-P	10	0.80		10				0.08		
Sorghum, grain grain	GC	0.515	STMR/STMR-P	86	0.60	75	65	70	55	0.4491	0.389	0.419	0.329
Wheat milled bypds	CM/CF	0.38	STMR/STMR-P	88	0.43	25		20	30	0.1080		0.086	0.13
Corn, field grain	GC	0.03	STMR/STMR-P	88	0.03				15				0.005
Soybean seed	VD	0.028	STMR/STMR-P	89	0.03			10				0.003	
Total						100	100	100	100	0.5571	2.958	0.509	0.46

S-METHOPRENE (147) – ESTIMATED MAXIMUM DIETARY BURDEN													
BEEF CATTLE													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Rice hulls	CM/CF	22.3	STMR	90	24.78			5				1.239	
Wheat milled bypds	CM/CF	13.6	STMR	88	15.45	40	30	35	55	6.182	4.636	5.409	8.5
Corn gluten feed	CM/CF	4.85	STMR	40	12.13	60				7.275			
Peanut meal	SM	5	STMR	85	5.88		20	10			1.176	0.588	
Sorghum, grain grain	GC	4.85	STMR	86	5.64		40	50	35		2.256	2.82	1.974
Barley grain	GC	4.85	STMR	88	5.51		10		10		0.551		0.551
Total						100	100	100	100	13.457	8.62	10.06	11.02

DAIRY CATTLE													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Rice hulls	CM/CF	22.3	STMR	90	24.78			10				2.478	
Wheat milled bypds	CM/CF	13.6	STMR	88	15.45	30	30	30	45	4.636	4.636	4.636	6.955
Peanut meal	SM	5	STMR	85	5.88	10	10	15		0.588	0.588	0.882	
Sorghum, grain grain	GC	4.85	STMR	86	5.64	45	40	45	30	2.538	2.256	2.538	1.692
Barley grain	GC	4.85	STMR	88	5.51				25				1.378
Corn, field hominy meal	CM/CF	4.85	STMR	88	5.51	15				0.827			
Brewer's grain dried	SM	4.85	STMR	92	5.27		5				0.264		
Soybean seed	VD	2.4	STMR	89	2.70		10				0.27		
Soybean meal	SM	2.4	STMR	92	2.61		5				0.13		
Total						100	100	100	100	8.589	8.144	10.53	10.02

POULTRY BROILER													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Wheat milled bypds	CM/CF	13.6	STMR	88	15.45	50	20	20	5	7.727	3.091	3.091	0.773
Peanut meal	SM	5	STMR	85	5.88	25	10	10		1.471	0.588	0.588	
Corn, field milled bypds	CM/CF	4.85	STMR	85	5.71	25	40			1.426	2.282		
Sorghum, grain grain	GC	4.85	STMR	86	5.64		30	70	65		1.692	3.948	3.666
Corn, field grain	GC	4.85	STMR	88	5.51				30				1.653
Total						100	100	100	100	10.624	7.653	7.627	6.092

S-METHOPRENE (147) – ESTIMATED MAXIMUM DIETARY BURDEN													
POULTRY LAYER													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Wheat milled bypds	CM/CF	13.6	STMR	88	15.45	50	20	20	30	7.727	3.091	3.091	4.636
Peanut meal	SM	5	STMR	85	5.88	25	10	10		1.471	0.588	0.588	
Corn, field milled bypds	CM/CF	4.85	STMR	85	5.71	25	30			1.426	1.712		
Sorghum, grain grain	GC	4.85	STMR	86	5.64		40	70	55		2.256	3.948	3.102
Corn, field grain	GC	4.85	STMR	88	5.51				15				0.827
Total						100	100	100	100	10.624	7.647	7.627	8.565

S-METHOPRENE (147) – ESTIMATED MEAN DIETARY BURDEN													
BEEF CATTLE													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Rice hulls	CM/CF	22.3	STMR/STMR-P	90	24.78			5				1.239	
Wheat milled bypds	CM/CF	13.6	STMR/STMR-P	88	15.45	40	30	35	55	6.1818	4.636	5.409	8.5
Corn gluten feed	CM/CF	4.85	STMR/STMR-P	40	12.13	60				7.275			
Peanut meal	SM	5	STMR/STMR-P	85	5.88		20	10			1.176	0.588	
Sorghum, grain grain	GC	4.85	STMR/STMR-P	86	5.64		40	50	35		2.256	2.82	1.974
Barley grain	GC	4.85	STMR/STMR-P	88	5.51		10		10		0.551		0.551
Total						100	100	100	100	13.4568	8.62	10.06	11.02

DAIRY CATTLE													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Rice hulls	CM/CF	22.3	STMR/STMR-P	90	24.78		0	10			0	2.478	
Wheat milled bypds	CM/CF	13.6	STMR/STMR-P	88	15.45	30	30	30	45	4.6364	4.636	4.636	6.955
Peanut meal	SM	5	STMR/STMR-P	85	5.88	10	10	15		0.5882	0.588	0.882	
Sorghum, grain grain	GC	4.85	STMR/STMR-P	86	5.64	45	40	45	30	2.5378	2.256	2.538	1.692
Barley grain	GC	4.85	STMR/STMR-P	88	5.51	0			25	0			1.378
Corn, field hominy meal	CM/CF	4.85	STMR/STMR-P	88	5.51	15				0.8267			
Brewer's grain dried	SM	4.85	STMR/STMR-P	92	5.27	0	5			0	0.264		
Soybean seed	VD	2.4	STMR/STMR-P	89	2.70	0	10			0	0.27		
Soybean	SM	2.4	STMR/STMR-P	92	2.61	0	5			0	0.13		

DAIRY CATTLE													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
meal			P										
Total						100	100	100	100	8.5891	8.144	10.53	10.02

S-METHOPRENE (147) – ESTIMATED MEAN DIETARY BURDEN

POULTRY BROILER													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Wheat milled bypds	CM/CF	13.6	STMR/STMR-P	88	15.45	50	20	20	5	7.73	3.091	3.091	0.773
Peanut meal	SM	5	STMR/STMR-P	85	5.88	25	10	10		1.47	0.588	0.588	
Corn, field milled bypds	CM/CF	4.85	STMR/STMR-P	85	5.71	25	40			1.43	2.282		
Sorghum, grain grain	GC	4.85	STMR/STMR-P	86	5.64		30	70	65		1.692	3.948	3.666
Corn, field grain	GC	4.85	STMR/STMR-P	88	5.51				30				1.653
Total						100	100	100	100	10.62	7.653	7.627	6.092

POULTRY LAYER													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Wheat milled bypds	CM/CF	13.6	STMR/STMR-P	88	15.45	50	20	20	30	7.7273	3.091	3.091	4.636
Peanut meal	SM	5	STMR/STMR-P	85	5.88	25	10	10		1.4706	0.588	0.588	
Corn, field milled bypds	CM/CF	4.85	STMR/STMR-P	85	5.71	25	30			1.4265	1.712		
Sorghum, grain grain	GC	4.85	STMR/STMR-P	86	5.64		40	70	55		2.256	3.948	3.102
Corn, field grain	GC	4.85	STMR/STMR-P	88	5.51				15				0.827
Total						100	100	100	100	10.6243	7.647	7.627	8.565

THIAMETHOXAM (245) – ESTIMATED MAXIMUM DIETARY BURDEN													
BEEF CATTLE													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Rice hulls	CM/CF	95.2	STMR	90	105.78			5				5.289	
Cabbage heads, leaves	AM/AV	3	HR	15	20.00		20				4		
Rice grain	GC	17	STMR	88	19.32	20		40		3.864		7.727	
Rice bran/pollard	CM/CF	9.35	STMR	90	10.39	15		35	20	1.558		3.636	2.078
Barley hay	AF/AS	6.7	HR	88	7.61	15		20		1.142		1.523	
Triticale hay	AF/AS	6.7	HR	88	7.61		20				1.523		
Barley straw	AF/AS	6.7	HR	89	7.53		10				0.753		
Swede roots	VR	0.2	HR	10	2.00		40				0.8		
Rice straw	AF/AS	1	HR	90	1.11				55				0.611
Potato culls	VR	0.2	HR	20	1.00	30				0.300			
Sorghum, grain forage	AF/AS	0.3	HR	35	0.86		10				0.086		
Barley grain	GC	0.12	STMR	88	0.14	20			25	0.027			0.034
Total						100	100	100	100	6.891	7.161	18.18	2.723

DAIRY CATTLE													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Rice hulls	CM/CF	95.2	STMR	90	105.78			10				10.58	
Cabbage heads, leaves	AM/AV	3	HR	15	20.00		20				4		
Rice grain	GC	17	STMR	88	19.32	20		20		3.864		3.864	
Rice bran/pollard	CM/CF	9.35	STMR	90	10.39	15	20	30	10	1.558	2.078	3.117	1.039
Barley hay	AF/AS	6.7	HR	88	7.61	20		40		1.523		3.045	
Triticale hay	AF/AS	6.7	HR	88	7.61		20				1.523		
Barley straw	AF/AS	6.7	HR	89	7.53		10				0.753		
Oat hay	AF/AS	6.7	HR	90	7.44	10			5	0.744			0.372
Corn, sweet forage	AF/AS	2.8	HR	48	5.83	35				2.042			
Bean vines	AL	1.4	HR	35	4.00		20				0.8		
Wheat forage	AF/AS	0.73	HR	25	2.92		10				0.292		
Rice straw	AF/AS	1	HR	90	1.11				20				0.222
Sorghum, grain forage	AF/AS	0.3	HR	35	0.86				15				0.13
Barley grain	GC	0.12	STMR	88	0.14				40				0.055
Corn, field forage/silage	AF/AS	0.05	HR	40	0.13				10				0.013
Total						100	100	100	100	9.731	9.445	20.6	1.829

THIAMETHOXAM (245) – ESTIMATED MAXIMUM DIETARY BURDEN													
POULTRY BROILER													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Rice grain	GC	17	STMR	88	19.32	20		50		3.864		9.659	
Rice bran/pollard	CM/CF	9.35	STMR	90	10.39	10	10	20	5	1.039	1.039	2.078	0.519
Swede roots	VR	0.2	HR	10	2.00		10				0.2		
Cassava/tapioca roots	VR	0.2	HR	37	0.54		10				0.054		
Barley grain	GC	0.12	STMR	88	0.14	70	70		10	0.095	0.095		0.014
Sorghum, grain grain	GC	0.079	STMR	86	0.09			30	55			0.028	0.051
Corn, field grain	GC	0.02	STMR	88	0.02				30				0.007
Total						100	100	100	100	4.998	1.388	11.76	0.59

THIAMETHOXAM (245) – ESTIMATED MAXIMUM DIETARY BURDEN													
POULTRY LAYER													MAX
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Cabbage heads, leaves	AM/AV	3	HR	15	20.00		5				1		
Rice grain	GC	17	STMR	88	19.32	20		50		3.864		9.659	
Rice bran/pollard	CM/CF	9.35	STMR	90	10.39	10	5	20	20	1.039	0.519	2.078	2.078
Wheat hay	AF/AS	6.7	HR	88	7.61		10				0.761		
Swede roots	VR	0.2	HR	10	2.00		10				0.2		
Cassava/tapioca roots	VR	0.2	HR	37	0.54		5				0.027		
Pea vines	AL	0.1	HR	25	0.40		10				0.04		
Barley grain	GC	0.12	STMR	88	0.14	70	55			0.095	0.075		
Sorghum, grain grain	GC	0.079	STMR	86	0.09			30	55			0.028	0.05
Corn, field grain	GC	0.02	STMR	88	0.02				25				0.006
Total						100	100	100	100	4.998	2.623	11.76	2.134

THIAMETHOXAM (245) – ESTIMATED MEAN DIETARY BURDEN													
BEEF CATTLE													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Rice hulls	CM/CF	95.2	STMR/STMR-P	90	105.78			5				5.289	
Rice grain	GC	17	STMR/STMR-P	88	19.32	20		40		3.8636		7.727	
Rice bran/pollard	CM/CF	9.35	STMR/STMR-P	90	10.39	15		35	20	1.5583		3.636	2.078
Cabbage heads, leaves	AM/AV	0.78	STMR/STMR-P	15	5.20		20				1.04		
Corn, sweet forage	AF/AS	1.5	STMR/STMR-P	48	3.13			20				0.625	
Wheat forage	AF/AS	0.53	STMR/STMR-P	25	2.12		20				0.424		

THIAMETHOXAM (245) – ESTIMATED MEAN DIETARY BURDEN													
BEEF CATTLE													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
Wheat straw	AF/AS	0.53	STMR/STMR-P	88	0.60	10				0.0602			
Barley straw	AF/AS	0.53	STMR/STMR-P	89	0.60		10				0.06		
Rice straw	AF/AS	0.47	STMR/STMR-P	90	0.52				55				0.287
Apple pomace, wet	AB	0.11	STMR/STMR-P	40	0.28		20				0.055		
Barley hay	AF/AS	0.2	STMR/STMR-P	88	0.23	5				0.011			
Pea vines	AL	0.04	STMR/STMR-P	25	0.16		20				0.032		
Sorghum, grain stover	AF/AS	0.14	STMR/STMR-P	88	0.16		10				0.016		
Barley grain	GC	0.12	STMR/STMR-P	88	0.14	30			25	0.041			0.034
Sorghum, grain	GC	0.079	STMR/STMR-P	86	0.09	20				0.0184			
Total						100	100	100	100	5.5528	1.626	17.28	2.399

DAIRY CATTLE													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Rice hulls	CM/CF	95.2	STMR/STMR-P	90	105.78		0	10			0	10.58	
Rice grain	GC	17	STMR/STMR-P	88	19.32	20		20		3.8636		3.864	
Rice bran/pollard	CM/CF	9.35	STMR/STMR-P	90	10.39	15	20	30	10	1.5583	2.078	3.117	1.039
Cabbage heads, leaves	AM/AV	0.78	STMR/STMR-P	15	5.20	0	20			0	1.04		
Corn, sweet forage	AF/AS	1.5	STMR/STMR-P	48	3.13	45		40		1.4063		1.25	
Bean vines	AL	0.88	STMR/STMR-P	35	2.51	0	20			0	0.503		
Wheat forage	AF/AS	0.53	STMR/STMR-P	25	2.12	0	20			0	0.424		
Barley straw	AF/AS	0.53	STMR/STMR-P	89	0.60	0	20			0	0.119		
Oat straw	AF/AS	0.53	STMR/STMR-P	90	0.59	0			5	0			0.029
Rice straw	AF/AS	0.47	STMR/STMR-P	90	0.52	0			20	0			0.104
Apple pomace, wet	AB	0.11	STMR/STMR-P	40	0.28	10				0.0275			
Sorghum, grain forage	AF/AS	0.06	STMR/STMR-P	35	0.17	0			15	0			0.026
Rape forage	AM/AV	0.05	STMR/STMR-P	30	0.17	10				0.0167			
Barley grain	GC	0.12	STMR/STMR-P	88	0.14	0			40	0			0.055
Sorghum, grain grain	GC	0.079	STMR/STMR-P	86	0.09	0			10	0.00			0.01
Total						100	100	100	100	6.8724	4.164	18.81	1.262

THIAMETHOXAM (245) – ESTIMATED MEAN DIETARY BURDEN													
POULTRY BROILER													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Rice grain	GC	17	STMR/STMR-P	88	19.32	20		50		3.86		9.659	
Rice bran/pollard	CM/CF	9.35	STMR/STMR-P	90	10.39	10	10	20	5	1.04	1.039	2.078	0.519
Barley grain	GC	0.12	STMR/STMR-P	88	0.14	70	70		10	0.10	0.095		0.014
Oat grain	GC	0.112	STMR/STMR-P	89	0.13		20				0.025		
Sorghum, grain grain	GC	0.079	STMR/STMR-P	86	0.09			30	55			0.028	0.051
Corn, field grain	GC	0.02	STMR/STMR-P	88	0.02				30				0.007
Total						100	100	100	100	5.00	1.16	11.76	0.59

POULTRY LAYER													MEAN
Commodity	CC	Residue (mg/kg)	Basis	DM (%)	Residue dw (mg/kg)	Diet content (%)				Residue Contribution (ppm)			
						US-CAN	EU	AU	JP	US-CAN	EU	AU	JP
Rice grain	GC	17	STMR/STMR-P	88	19.32	20		50		3.8636		9.659	
Rice bran/pollard	CM/CF	9.35	STMR/STMR-P	90	10.39	10	5	20	20	1.0389	0.519	2.078	2.078
Cabbage heads, leaves	AM/AV	0.78	STMR/STMR-P	15	5.20		5				0.26		
Wheat forage	AF/AS	0.53	STMR/STMR-P	25	2.12		10				0.212		
Rape forage	AM/AV	0.05	STMR/STMR-P	30	0.17		5				0.008		
Pea vines	AL	0.04	STMR/STMR-P	25	0.16		10				0.016		
Barley grain	GC	0.12	STMR/STMR-P	88	0.14	70	65			0.0955	0.089		
Sorghum, grain	GC	0.079	STMR/STMR-P	86	0.09			30	55			0.028	0.051
Corn, field grain	GC	0.02	STMR/STMR-P	88	0.02				25				0.01
Total						100	100	100	100	4.9980	1.104	11.76	2.134



An Extra Joint Meeting of the Food and Agriculture Organization of the United Nations (FAO) Panel of Experts on Pesticide Residues in Food and the Environment and the World Health Organization (WHO) Core Assessment Group on Pesticide Residues (JMPR) was held virtually over two sessions from 17 to 21 May and 7 to 11 June 2021. The Meeting was held in pursuance of recommendations made by previous Meetings and accepted by the governing bodies of FAO and WHO that studies should be undertaken jointly by experts to evaluate possible hazards to humans arising from the occurrence of pesticide residues in foods. During the meeting the FAO Panel of Experts was responsible for reviewing pesticide use patterns (use of good agricultural practices), data on the chemistry and composition of the pesticides and methods of analysis for pesticide residues and for estimating the maximum residue levels that might occur as a result of the use of the pesticides according to good agricultural use practices. The WHO Core Assessment Group was responsible for reviewing toxicological and related data and for estimating, where possible and appropriate, acceptable daily intakes (ADIs) and acute reference doses (ARfDs) of the pesticides for humans. This report contains information on ADIs, ARfDs, maximum residue levels, and general principles for the evaluation of pesticides. The recommendations of the Joint Meeting, including further research and information, are proposed for use by Member governments of the respective agencies and other interested parties.

ISBN 978-92-5-135028-7



9 789251 350287

CB6975EN/1/10.21