

Bernoulli Distribution

Simplest non-constant distribution


Foundation of many others

μ

V

σ

Repeated experiments



Jacob Bernoulli, 1655-1705

Theology → mathematics

Calculus Integrals

“Euler” number

$$e = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n$$

$e \rightarrow b$

Ars Conjectandi

First law of large numbers

Mentored brother Johann Medicine → Math Dynasty

The simplest Distribution

Simplest

One value

5

Constant


Always same

Trivial

Simplest non-trivial

Two values

Simplest values



0 and 1



Bernoulli Coin!


Bernoulli Distribution

B_p $0 \leq p \leq 1$

Two values	0	1
Probability	$1-p$	p

failure success

\bar{p} q



Σ Will it add?

$$p(0) + p(1) = (1-p) + p = 1$$

YES IT ADDS!

$X \sim B_p$ Bernoulli random variable, coin, experiment, trial

Who Cares About Two Values?

Everyone!

Binary version of complex events

Products: 80 good, 20 defective

Select one, good or not

$\sim B_{.8}$

Next child will be a boy $\sim B_{.5}$

Generalizes to more complex variables

Patient has one of three diseases

Repeated trials yield # successes

Many important distributions

Binomial, Geometric, Poisson, Normal

Mean

$$X \sim B_p$$

$$p(0) = 1-p$$

$$p(1) = p$$

$$EX = \sum p(x) \cdot x = (1-p) \cdot 0 + p \cdot 1 = p$$

$$X \sim B_{0.8}$$

$$EX = 0.8$$

$$EX = P(X=1)$$

Fraction of times expect to see 1

Variance

$$X \sim B_p$$

$$EX = p$$

Variance

Easy route

$$0^2 = 0$$

$$1^2 = 1$$

$$X^2 = X$$

$$E(X^2) = EX = p$$

$$V(X) = E(X^2) - (EX)^2 = p - p^2 = p(1-p) = pq$$


Standard Deviation

$$\sigma = \sqrt{pq}$$

B_p varies most
when $p = 1/2$

E, V, σ for
various p

p	EX	V(X)	σ
0	0	0	0
1	1	0	0
1/2	1/2	1/4	1/2



Independent Trials

Much of B_p importance stems from multiple trials

Most common Independent \perp

$$0 \leq p \leq 1 \quad X_1, X_2, X_3 \sim B_p \quad \perp$$

$$q \stackrel{\text{def}}{=} 1-p \quad P(110) = p^2 q = P(101) = P(011)$$

Generally $X_1, X_2, \dots, X_n \sim B_p \quad \perp$

$$x^n = x_1, x_2, \dots, x_n \in \{0, 1\}^n \quad n_0 \text{ 0's and } n_1 \text{ 1's}$$

$$P(x_1, \dots, x_n) = p^{n_1} q^{n_0} \quad P(10101) = p^{n_1} q^{n_0} = p^3 q^2$$



Typical Samples

Distribution	Typical seq.	Description	Probability
B_0	0000000000	constant 0	$1^{10} = 1$
B_1	1111111111	constant 1	$1^{10} = 1$
$B_{0.8}$	1110111011	80% 1's	$0.8^8 \cdot 0.2^2$
$B_{0.5}$	1011010010	50% 1's	0.5^{10}

Fair coin flip

Not most probable
Most probable: 1...1
Unlikely to be seen

Bernoulli Distribution

Simplest non-constant distribution

$$B_p \quad 0 \leq p \leq 1$$

$$0 \text{ and } 1 \quad p(1) = p \quad p(0) = 1 - p = q$$

$$\mu = p \quad V = pq \quad \sigma = \sqrt{pq}$$

Foundation of many other distributions



Binomial

