

Cartogram Mapping and its Application to Cancer Data Visualisation

Stephanie Kobakian, Jessie Roberts and Dianne Cook

Contents

1	Introduction	1
2	Disease mapping methods	2
2.1	Current best practice map displays for cancer data	2
2.2	Public face	4
2.3	Cartograms	8
2.4	Centroid displays	12
3	A critique of mapping methods	12
4	Animation and Interactivity	12
5	Acknowledgements	13
6	References	13

Abstract

1 Introduction

Maps have been adopted to present geospatial statistics for centuries. Maps connect data and statistics to the geographical representation of areas that are generally familiar to the audience. However, it is not enough for areas on maps to be recognisable, if information regarding the distribution of the statistic cannot be gleaned. In situations where the people are of interest, not the land they live on, it is reasonable to explore views that enhance the communication of the cancer statistics. Identifying and explaining spatial structures, patterns, and processes involves considering the individuals in communities and organising communities into representable units (Moore and Carpenter 1999). This has spurred innovations over the previous centuries to enhance the shapes and maps presented to effectively communicate cancer outcomes, and health outcomes more broadly. Cancer statistics directly represent and relate to the people living within individual geographical areas.

The visualisation methods used to present cancer statistics will depend on the intended message and users of the map. Transforming statistics to visualisations considers individual observations or observation points aggregated into communities and geographical units. Visualising diseases on maps is often the first step in exploratory spatial data analysis and effectively helps in the formulation of hypotheses (Jahan et al. 2018). Disease maps help to present geographic patterns that may have been overlooked in a table, obscuring the geospatially related statistics (Moore and Carpenter 1999). By providing a visual representation of cancer outcomes, geographic patterns of disease are able to be identified and effectively addressed.




Figure 1: A choropleth map of the United States of America.

2 Disease mapping methods

2.1 Current best practice map displays for cancer data

A choropleth map is used to display the spatial characteristics of a relationship, or variability, of measurements. As an alternative storage device to a table, it preserves locations for geographically ordered data, with usage dating back to the 1800s (Berry, Morrill, and Tobler, n.d.). A choropleth is constructed by drawing the geographic or political boundaries, and filling the shapes with colours to represent values of a measured variable (Tufte 1990). Early versions of choropleth maps used symbols or patterns instead of colour. Bell et al. (2006) discusses their use in visualising cancer data, and Walter (2001) gives an overview of the development of these maps for displaying disease data. Just as geographers are no longer the only creators of maps, Bell et al. (2006) suggests the audiences of spatial health data analysis have extended beyond researchers to the public, policymakers and the media. A choropleth map is a true map of the topology, constructed for visual inspection of spatial patterns across a familiar geographic form. Figure 1 shows a choropleth map where each state of the United States of America is coloured by the average annual rate of new cases of lung and bronchus cancer from years 2012 to 2016.

Utilising the familiar state boundaries can make a map intuitive to read (Brewster and Subramanian 2010), and allow viewers to visually infer the spatial relationships in the data, i.e. how cancer rate differs across the states. The familiarity of the geography is a worthy consideration when presenting results of spatial analysis. Just as geographers are no longer the only creators of maps, Bell et al. (2006) suggests the audiences of spatial health data analysis have extended beyond researchers to the public, policymakers and the media. While the areas are recognisable shapes, they are often politically driven boundaries with individual areas being of non-uniform size, containing different population densities and subject to change over time. The different population and geographical sizes of administrative areas can attract attention to the shades of the unpopulated but large areas (Tufte 1990). Choropleths can inhibit visual inference when presenting human related statistics as the display may draw attention from the ‘potentially more important results in the more populous communities’ (Exeter 2017).


Choropleth maps can be useful devices for communicating information to public on a familiar map

base. In epidemiology, choropleths are often used as a tool to study the spatial distribution of cancer incidence and mortality. A cancer atlas is a choropleth map, or collection of maps, representing cancer incidence and mortality for a country, or group of countries. d’Onofrio et al. (2016) provide the definition and reports that cancer atlases began with Haviland’s maps in 1875. The data collection methods of cancer mortality rates across regions, and the administrative control within regions lends itself to choropleth visualisation. The increasing development and use of disease maps can be attributed to the availability of geographic information system software (Exeter 2017). Early work in US cancer atlases can be attributed to Burbank ???, and UK cancer atlases to Howe 1963 ??. The choropleth maps presented levels via hatchings or dots on a black and white scale. These atlases were key to developing hypotheses regarding areas with unusually high rates, geographic correlations, work related exposures, and high risk diets d’Onofrio et al. (2016).

Almost 100 years of cancer mapping in the United States and the United Kingdom has seen increased effectiveness in the presentation cancer statistics. Mortality rates are now often presented as relative rates of risk across the population, and age adjusted to correct for the the higher prevalence of cancers in older populations. Howe (1989) describes Stock’s development of the standardised mortality ratios through the 1930s. Table 1 presents summarises the measures presented in published cancer atlases, and provides a definition of each measure.

Table 1: Measures used to report cancer statistics

Measure	Details
1. IR (Incidence Ratio)	$(IR)_i = \frac{(Incidence\ Rate)_i}{Average\ Incidence\ Rate}$, Cancer incidence rate in region i over the average cancer incidence rate for the total region
2. SIR (Standardised Incidence Ratio)	IR standardised by age structure in each region i
3. RER (Relative Excess Risk)	$RER = \frac{(Cancer\ related\ mortality)_i}{Average\ cancer\ related\ mortality}$ Represents the estimate of cancer related mortality within five years of diagnosis Also referred to as ‘excess hazard ratio’
4. Age Adjusted Relative Risk	RR standardised by age structure in each region i
5. Rate per 100,000	Cancer incidence per 100,000 population
6. Age Adjusted Rate per 100,000	#5 standardised by age structure or region
7. New cancer cases per 100,000	Specific methods could not be found
8. Count	Crude cancer counts
9. Below or above Expected	Alternative expression of the SIR



2.1.1 Supporting material in cancer atlases

A map communicates quickly and draws attention to prominent geographies, but an atlas is often supplemented with supporting statistics. Cruickshank's (1947) as cited by Walter (2001), discusses using visuals as 'formal statistical assessment of the spatial pattern'. However, when presenting cancer atlases, d'Onofrio et al. (2016) believes the intuition derived must be 'validated by rigorous statistical analyses'.

These additional statistics often include a measure of the statistical uncertainty of the values of the statistics presented in a choropleth. In the review of atlases in the public domain, atlases were considered to report uncertainty to the non-expert user if they included a measure of statistical uncertainty either within or alongside the map. Maps that only reported this information within the supplementary material were not considered to have directly attempted to report uncertainty. The maps considered used standard and well known measures including credible intervals and standard deviation, statistical significance, box plots and distributions. These maps ranged from static pdfs or infographics, to interactive online resources.

The interactivity of the more modern maps enabled supplementary information to be incorporated without cluttering the screen, such as in a tool tip feature.

2.2 Public face

Cancer maps are effective communication tools for a general or non-expert audience. They are commonly used in the public domain to communicate results of sophisticated statistical analyses. The




Figure 2: Example of CI visualisation for uncertainty representation in cancer mapping (1/3).
Source: Alberta Health IHDA Geographic. (2012)




Figure 3: Example of CI visualisation for uncertainty representation in cancer mapping (2/3).
Source: Pennsylvania Cancer Atlas

heavy use of chloropleth maps within the research literature is reflected in the types of maps that are found in the public domain. A grey literature review conducted by (ref) identified 33 cancer atlases published on the internet between January 2010 and November 2015. These chloropleth maps were mostly published by non-commercial organisations, including not-for-profits (NFPs), government, research organisations, advocacy groups or a partnership between an NFP & government. Only one map was published by a commercial entity.

The cancer atlases covered geographies from all around the world, four were global atlases. Most focussed on single nations, the United States was considered by eleven atlases, the United Kingdom by seven, followed by three of Australia, two of Canada, and one of each from Switzerland, Germany, Norway. One atlas covered the European Union. Not all maps had a national focus and ten covered a region or state rather than an entire nation. The states or counties/regions covered were South Australia (AUS), Queensland (AUS), Ontario (CAN), Valencia (Spain), Pennsylvania county Massachusetts (US), New Hampshire (US), Cape Cod (US), Missouri (US), Florida (US), New York State (US) and Arizona (US).

2.2.1 Examples of supporting statistics in public facing atlases

Close to half of the atlases identified (42%, n=14) included some measure of uncertainty. The most common measure used to represent uncertainty were credible or confidence intervals (CIs).



Figure 4: Example of CI visualisation for uncertainty representation in cancer mapping (3/3).
Source: Centres for Disease Control and Prevention (CDC). United States Cancer Statistics: An Interactive Cancer Atlas (InCA)

Data Table		Glossary		Further Info	
County	Rate	Lower CI	Upper CI	Count	Population
Adams	60.4	55.3	65.9	518	801,717
Allegheny	61.4	60.1	62.7	9,390	11,676,746
Armstrong	58.9	53.8	64.4	506	656,638
Beaver	56.4	53.3	59.7	1,247	1,654,797
Bedford	52.3	46.3	58.8	287	446,092
Berks	60.1	57.6	62.6	2,287	3,306,811
Blair	67.4	63.2	71.7	1,012	1,171,282
Bradford	54.3	48.8	60.3	363	563,934
Bucks	61.3	59.1	63.4	3,174	5,298,961
Butler	61.5	57.8	65.4	1,042	1,537,647
Cambria	65.9	62.3	69.7	1,291	1,400,022
Cameron	76.7	57.7	102	57	54,504
Carbon	65.0	59.2	71.4	472	530,218
Centre	52.1	47.6	56.9	502	1,214,956
Chester	56.2	53.8	58.7	2,015	3,797,562
Clarion	64.5	57	72.8	273	376,923

Figure 5: Example of an interactive data table with CI upper and lower bounds used in cancer mapping. Source: Pennsylvania Cancer Atlas




Figure 6: Example of standard deviation visualised in cancer mapping




Figure 7: Example of boxplot used in cancer mapping. Source: Pennsylvania Cancer Atlas

Methods for representing sources of uncertainty information can be visualised or communicated in different ways, examples identified through this grey literature review are listed below.

Table 2: (#tab:method-exp) Implicit and explicit measures of uncertainty.

Measure	Example
CI Interval	Figures 3.1, 3.2, 3.3
Statistical Significance	Textured overlay on top of coloured regions used to indicate statistical significance
Distribution	Figure 3.4
Boxplots	Figures 3.4, 3.5
Sample Size	Textured overlay or lack of colour on a region, was used to show regions with small sample size

Measure	Example
Standard deviation	Figure 3.6 - the second map in the bottom right corner shows standard deviation

2.3 Cartograms

A cartogram alters the map base with the intention of improving the presentation of the statistic of interest. For a single variable of interest, each map area is changed to emphasise the distribution by representing the corresponding value, in comparison to the value of the other areas (Dougenik, Chrisman, and Niemeyer 1985).

Choropleths may be considered true topological maps, however, if the land mass displayed covers enough of the globe, there must be a transformation or distortion to display the land in 2D. The amount of distortion is related to the distance covered by the landmass displayed Tobler (1963). World map projections reflect the frequent perspectives used to view the earth. Choropleth maps will always be distorted if they cover enough of the globe, just as photographs of the globe from space. Choropleth creation requires choosing a map projection that shows a favourable distortion of the geography for presenting the set of spatial information. Diagrams that do not specify a projection can be considered to have some unknown projection.

If the statistic presented on the map base relies on physical distance and is influenced by the topology there is no transformation needed beyond choosing a projection. The purposeful distortion of the map space is beneficial when a uniform density of the map base is desired. When a map base is transformed according to population density, population becomes a uniformly distributed background for the statistic presented (Berry, Morrill, and Tobler, n.d.).

If the statistic presented on the map base relies on physical distance and is influenced by the topology there is no transformation needed beyond choosing a projection. The purposeful distortion of the map space is beneficial when a uniform density of the map base is desired, Dorling (2011) suggests ‘population distribution is often extremely uneven in former British colonies’, this makes the distortion necessary. When a map base is transformed according to population density, population becomes a uniformly distributed background for the statistic presented (Berry, Morrill, and Tobler, n.d.). Using choropleth maps for population characteristics requires graphic distortions when the population concerned varies greatly in density (Griffin 1980). When implementing a distortion of the geographical shape according to population, an area cartogram (Olson 1976), population-by-area cartograms (Levison and Haddon Jr 1965), or iso-demographic map is the result.

Cartograms provide an alternative visualisation method for statistical and geographical information. The key difference between a choropleth and a cartogram is the desirable augmentation of the size, shape or distance of geographical areas (Dorling 2011). Cartograms may be seen as an extension of map transformations and projections. The favourable distortion is proportional to a value other than actual earth size area (Olson 1976).

There have been many algorithms presented, Nusrat and Kobourov (2016) provided a framework to investigate implementations and the “statistical accuracy, geographical accuracy, and topological accuracy”.

The creation of cartograms was largely in the hands of cartographers. Dorling (2011) discusses early approaches including John Hunter and Jonathan Young (1968) and Durham’s wooden tile method, Skoda and Robertson’s (1972) steel ball bearing approach and Tobler’s (1973) computer programs.




Figure 8: “Choropleth maps of the United States of America using four coordinate reference systems.”

There are many alternatives to consider, the intended audience of the map, and its purpose are key points in cartogram use and creation. Dorling (2011) reiterates: ‘There is no “best” cartogram or method of creating cartograms just as there is no “best” map’ (Monmonier and Schnell, 1988).

2.3.1 Contiguous

A contiguous cartogram maintains connectivity of the map regions while areas are resized according to a statistic. This transformation often occurs at the expense of the shape (Kocmoud and House 1998, @NAC, @TAAM). From a computer graphics perspective, Min Ouyang and Revesz (2000) believe it is a problem of ‘map deformation’ to account for the value assigned to each area, they provide three methods for creating value-by-area cartograms. Examples include Tobler’s Pseudo-Cartogram Method, Dorling’s Cellular Automaton Method (2011), Radial Expansion Method of Selvin et al., Rubber Sheet Method of Dougenik et al., Gusein-Zade and Tikunov’s Line Integral Method, Constraint-Based Method (Kocmoud and House) (1998).

describes the process of creating a population cartogram for Canada, using the 1966 Census. An intentional goal of this project was to maintain contiguity, while attempting to keep the actual shape of places. The end result was a ‘very accurate isodemo-graphic map of Canada’. This intentional design goal coincided with the rising interest in urban geography.

To be able to recognise the significant changes, a reader will usually have to know the initial geography to find the differences in the new cartogram layout (Olson 1976). Tobler’s Conformal mapping means to preserve angles locally so that the shapes of very small areas on a traditional map and a cartogram would be similar. Kocmoud and House (1998) presents this issue as conflicting tasks or aims, to adjust region sizes and retain region shapes. Distortion of region shapes on the contiguous cartogram presents an additional hurdle to visual recognition and this hurdle is not only eliminated on the noncontiguous cartogram but is replaced by the meaningful empty-space property (Olson 1976, @ECGC).




Figure 9: “A contiguous cartogram of the United States of America”

2.3.2 Non-Contiguous

Dorling (2011) puts forward a simple question:

If, for instance, it is desirable that areas on a map have boundaries which are as simple as possible, why not draw the areas as simple shapes in the first place?

He answers this with his implementation of maps created with ‘the simplest of all shapes’. While contiguous cartograms may be a ‘more sophisticated’ method, they produce ‘very complex shapes’. Circular cartograms use the same shape for every region represented, and size them according to the statistic represented or the population for a base map. To produce a compelling map, a gravity model is applied to avoid overlaps, and keep spatial relationships with neighbouring areas over many iterations. This implementation can work for up to ‘one hundred thousand’ areas.

‘In Australia the urban federal constituencies occupy only a tenth of the land, but contain nine tenths of the people. It would be almost unthinkable to show election results for that country on a conventional equal land area map.’ This 1966 cartogram uses mostly straight lines, and the result looks very little like the geographical shape of Australia.

‘Given the increasingly uneven population distribution of the United States and the growing social divides between the populations of neighbourhoods living at different densities, the need for cartograms like this is greater now than ever.’

Used in displays of the UK by Howe in 1986 cited by Howe (1989)

Tobler’s method and the many implementations that ‘elaborated’ on it are derived from ‘numerical approximations to a pair of equations’(Dorling 2011). They all operate through incremental adjustments, and can produce wildly different outcomes from small changes in the inputs.

Tobler (2004) Value-Area Cartograms. In these cartograms a region,country, or continent is subdivided into small regions, each of which is represented by a rectangle. This rectangle is




Figure 10: “A Non - contiguous cartogram of the Unites States of America”




Figure 11: “A dorling cartogram of the Unites States of America”

proportionate in area to the value which it represents in certain statistical distributions. The regions are grouped in approximately the same positions as they are on the map.

Computer generated map examples: Howe (1989) (Hopps et al. 1968; Armstrong 1972). There has followed a flood of disease atlases, mainly concentrating on the modern problems of cancer and degenerative diseases from countries as scattered as the United States (Burbank 1971; Mason et al. 1975, 1976; Pickle et al. 1987), the Soviet Union (Levin 1980), Japan (Shigematsu 1977), the Federal Republic of Germany

Cano et al. (2015) define the term ‘mosaic cartograms’. Compare amount of tiles to contrast population of regions. ‘Cartograms show a data value per input region by scaling each region such that its area is proportional to its data value. Mosaic cartograms show data in multiples of tiles, hence the input data must consist of, or be cast into, small integer units.’

2.4 Centroid displays

Dot plot: one dot for each region, coloured, and placed at centroid.

Olson (1976) gives an example of

Plotting centroids on top of geographies. (Size is kept constant)

Replies on the idea that every area is important, no matter the size. This gives equal emphasis to every area, allows distributions and relationships between neighbours to become more clear.

3 A critique of mapping methods

designing a map tailored to precise goals [is] easier than forcing a single map to accommodate diverse objectives - Bell et al. (2006)

Waldo Tobler (2012) explores many graphical techniques, and suggests there are particular methods for particular purposes. To choose an appropriate map display, the map creator must consider the intended user, and message the map will communicate. It is the objectives of the investigator that will drive the choice of representation (Bell et al. 2006).

There are two keys to drive the choice of display: the properties of the visualisation, and the ease or accuracy of information extraction for map users Moore and Carpenter (1999).

(only worth including if it is possible for us to implement) Tabular form comparing and contrasting
- Relationship to geography

- Show using cancer examples

4 Animation and Interactivity

Recent developments of technology allowed interactive

‘Where control of the message is important, static maps will continue to be the most effective, although good tables, graphs, and explanatory text are still needed in order to ensure that different people will see the same thing in the maps’ Bell et al. (2006)

lends to temporal pattern exploration

Keim et al. (2002) ?? Highlight the value of animating contiguous to see changes over time, US can be recognisable but animation aides interpretation

5 Acknowledgements

6 References

Wickham (2017) for data analysis, Bivand, Nowosad, and Lovelace (2019) and Pebesma (2018) to implement plotting of spatial data. Kobakian and Cook (2019) to create hexagon tesselation. Arnold (2019) to enhance plot displays. Jeworutzki (2018) for contiguous and non-contiguous cartogram displays.

Arnold, Jeffrey B. 2019. *Ggthemes: Extra Themes, Scales and Geoms for 'Ggplot2'*. <https://CRAN.R-project.org/package=ggthemes>.

Bell, B Sue, Richard E Hoskins, Linda Williams Pickle, and Daniel Wartenberg. 2006. “Current Practices in Spatial Analysis of Cancer Data: Mapping Health Statistics to Inform Policymakers and the Public.” *International Journal of Health Geographics* 5: 49. <https://doi.org/10.1186/1476-072X-5-49>.

Berry, Brian J. L., Richard L. Morrill, and Waldo R. Tobler. n.d. “GEOGRAPHIC Ordering of Information: NEW Opportunities” 16 (4). Association of American Geographers: 39–44. https://doi.org/10.1111/j.0033-0124.1964.039_q.x.

Bivand, Roger, Jakub Nowosad, and Robin Lovelace. 2019. *SpData: Datasets for Spatial Analysis*. <https://CRAN.R-project.org/package=spData>.

Brewster, Mark B., and S. V. Subramanian. 2010. “Cartographic Insights into the Burden of Mortality in the United Kingdom: A Review of ‘The Grim Reaper’s Road Map’” Journal Article. *International Journal of Epidemiology* 39 (4): 1120–2. <https://doi.org/10.1093/ije/dyp395>.

Cano, R. G., K. Buchin, T. Castermans, A. Pieterse, W. Sonke, and B. Speckmann. 2015. “Mosaic Drawings and Cartograms.” *Computer Graphics Forum* 34 (3): 361–70.

d’Onofrio, Alberto, Chiara Mazzetta, Chris Robertson, Michel Smans, Peter Boyle, and Mathieu Boniol. 2016. “Maps and Atlases of Cancer Mortality: A Review of a Useful Tool to Trigger New Questions.” Journal Article. *Ecancermedicalscience* 10: 670–70. <https://doi.org/10.3332/ecancer.2016.670>.

Dorling, Daniel. 2011. “Area Cartograms: Their Use and Creation.” In *Concepts and Techniques in Modern Geography (CATMOG)*, 59:252–60. <https://doi.org/10.1002/9780470979587.ch33>.

———. 2012. *The Visualisation of Spatial Social Structure*. John Wiley & Sons Ltd.

Dougenik, James A., Nicholas R. Chrisman, and Duane R. Niemeyer. 1985. “AN Algorithm to Construct Continuous Area Cartograms.” *The Professional Geographer* 37 (1). Routledge: 75–81. <https://doi.org/10.1111/j.0033-0124.1985.00075.x>.

Exeter, Daniel J. 2017. “Spatial Epidemiology.” Journal Article, 1–4. <https://doi.org/10.1002/9781118786352.wbieg0283>.

Griffin, T.L.C. 1980. “Cartographic Transformation of the Thematic Map Base.” *Cartography* 11 (3). Taylor & Francis: 163–74. <https://doi.org/10.1080/00690805.1980.10438102>.

- Howe, G. M. 1989. "Historical Evolution of Disease Mapping in General and Specifically of Cancer Mapping." In *Cancer Mapping*, edited by Peter Boyle, Calum S. Muir, and Ekkehard Grundmann, 1–21. Berlin, Heidelberg: Springer Berlin Heidelberg.
- Jahan, Farzana, Earl Duncan, Susanna Cramb, Peter Baade, and Kerrie Mengersen. 2018. "Making More of Spatial Maps: A Bayesian Meta-Analysis Approach." In.
- Jeworutzki, Sebastian. 2018. *Cartogram: Create Cartograms with R*. <https://CRAN.R-project.org/package=cartogram>.
- Keim, D.A, S.C North, C Panse, and J Schneidewind. 2002. "Efficient Cartogram Generation: A Comparison." In *IEEE Symposium on Information Visualization, 2002. INFOVIS 2002*, 2002-:33–36. IEEE.
- Kobakian, Stephanie, and Dianne Cook. 2019. *Sugarbag: Create Tessellated Hexagon Maps*. <https://CRAN.R-project.org/package=sugarbag>.
- Kocmoud, Christopher J., and Donald H. House. 1998. "A Constraint-Based Approach to Constructing Continuous Cartograms." In.
- Levison, M. E., and W. Haddon Jr. 1965. "THE Area Adjusted Map. AN Epidemiologic Device." Journal Article. *Public Health Reports* 80: 55–59.
- Min Ouyang, and P. Revesz. 2000. "Algorithms for Cartogram Animation." In *Proceedings 2000 International Database Engineering and Applications Symposium (Cat. No.PR00789)*, 231–35. <https://doi.org/10.1109/IDEAS.2000.880581>.
- Moore, Dale A., and Tim E. Carpenter. 1999. "Spatial Analytical Methods and Geographic Information Systems: Use in Health Research and Epidemiology." *Epidemiologic Reviews* 21 (2): 143–61. <https://doi.org/10.1093/oxfordjournals.epirev.a017993>.
- Nusrat, Sabrina, and Stephen G. Kobourov. 2016. "The State of the Art in Cartograms." *CoRR* abs/1605.08485. <http://arxiv.org/abs/1605.08485>.
- Olson, Judy M. 1976. "NONCONTIGUOUS Area Cartograms." *The Professional Geographer* 28 (4). Routledge: 371–80. <https://doi.org/10.1111/j.0033-0124.1976.00371.x>.
- Pebesma, Edzer. 2018. "Simple Features for R: Standardized Support for Spatial Vector Data." *The R Journal* 10 (1): 439–46. <https://doi.org/10.32614/RJ-2018-009>.
- Tobler, Waldo. 1963. "Geographic Area and Map Projections," January. American Geographical Society, 59–78. <https://www.jstor.org/stable/212809>.
- . 2004. "Thirty Five Years of Computer Cartograms." *Annals of the Association of American Geographers* 94 (1). Taylor & Francis Group: 58–73.
- Tufte, Edward R. 1990. *Envisioning Information*. Graphics Press.
- Walter, S. D. 2001. *Disease Mapping: A Historical Perspective*. Book. Oxford University Press: Oxford.
- Wickham, Hadley. 2017. *Tidyverse: Easily Install and Load the 'Tidyverse'*. <https://CRAN.R-project.org/package=tidyverse>.