ILLUMINATING A DARK LENS: A TYPE IA SUPERNOVA MAGNIFIED BY THE FRONTIER FIELDS GALAXY CLUSTER ABELL 2744

Steven A. Rodney^{1,2}, Brandon Patel³, Daniel Scolnic⁴, Ryan J. Foley^{5,6}, Alberto Molino^{7,8}, Gabriel Brammer⁹, Mathilde Jauzac^{10,11}, Maruša Bradač¹², Dan Coe⁹, Tom Broadhurst^{13,14}, Jose M. Diego¹⁵, Or Graur^{16,17}, Jens Hjorth¹⁸, Austin Hoag¹², Saurabh W. Jha³, Traci L. Johnson¹⁹, Patrick Kelly²⁰, Daniel Lam²¹, Curtis McCully^{22,23}, Elinor Medezinski^{1,24}, Massimo Meneghetti^{25,26,27}, Julian Merten²⁸, Johan Richard²⁹, Adam G. Riess^{1,9}, Keren Sharon¹⁹, Louis-Gregory Strolger^{9,30}, Tommaso Treu^{31,32}, Xin Wang²³, Liliya L. R. Williams³³, and Adi Zitrin^{34,2}

Draft version July 17, 2015

ABSTRACT

SN HFF14Tom is a Type Ia Supernova (SN) discovered at $z = 1.3457 \pm 0.0001$ behind the galaxy cluster Abell 2744 (z = 0.308). In a cosmology-independent analysis, we find that HFF14Tom is 0.77 ± 0.15 magnitudes brighter than unlensed Type Ia SNe at similar redshift, implying a lensing magnification of $\mu_{\rm obs} = 2.03 \pm 0.29$. This observed magnification provides a rare opportunity for a direct empirical test of galaxy cluster lens models. Here we test 17 lens models, 13 of which were generated before the SN magnification was known, qualifying as pure "blind tests". The models are collectively fairly accurate: 8 of the models deliver median magnifications that are consistent with the measured μ to within 1σ . However, there is a subtle systematic bias: the significant disagreements all involve models overpredicting the magnification. We evaluate possible causes for this mild bias, and find no single physical or methodological explanation to account for it. We do find that model accuracy can be improved to some extent with stringent quality cuts on multiply-imaged systems, such as requiring that a large fraction have spectroscopic redshifts. In addition to testing model accuracies as we have done here, Type Ia SN magnifications could also be used as inputs for future lens models of Abell 2744 and other clusters, providing valuable constraints in regions where traditional strong- and weak-lensing information is unavailable.

Subject headings: supernovae: general, supernovae: individual: HFF14Tom, galaxies: clusters: general, galaxies: clusters: individual: Abell 2744, gravitational lensing: strong, gravitational lensing: weak

- ¹ Department of Physics and Astronomy, The Johns Hopkins University, 3400 N. Charles St., Baltimore, MD 21218, USA
- Hubble Fellow
- ³ Department of Physics and Astronomy, Rutgers, The State University of New Jersey, Piscataway, NJ 08854, USA
- Department of Physics, The University of Chicago, Chicago, IL 60637, USA
- Department of Physics, University of Illinois at Urbana-Champaign, 1110 W. Green Street, Urbana, IL 61801, USA
- Astronomy Department, University of Illinois at Urbana-Champaign, 1002 W. Green Street, Urbana, IL 61801, USA
- ⁷ Instituto de Astrofísica de Andalucía (CSIC), E-18080 Granada, Spain
- Instituto de Astronomia, Geofísica e Ciências Atmosféricas, Universidade de São Paulo, Cidade Universitária, 05508-090, São Paulo, Brazil
- ⁹ Space Telescope Science Institute, 3700 San Martin Dr., Baltimore, MD 21218, USA
- ¹⁰ Institute for Computational Cosmology, Durham Univer-
- sity, South Road, Durham DH1 3LE, UK

 11 Astrophysics and Cosmology Research Unit, School of Mathematical Sciences, University of KwaZulu-Natal, Durban
- 4041, South Africa

 12 University of California Davis, 1 Shields Avenue, Davis, CA
- ¹³ Fisika Teorikoa, Zientzia eta Teknologia Fakultatea, Euskal Herriko Unibertsitatea UPV/EHU
- ¹⁴ IKERBASQUE, Basque Foundation for Science, Alameda Urquijo, 36-5 48008 Bilbao, Spain
- IFCA, Instituto de Física de Cantabria (UC-CSIC), Av. de Los Castros s/n, 39005 Santander, Spain
- ¹⁶ Center for Cosmology and Particle Physics, New York University, New York, NY 10003, USA
- ¹⁷ Department of Astrophysics, American Museum of Natural History, Central Park West and 79th Street, New York, NY 10024, USA

 18 Dark Cosmology Centre, Niels Bohr Institute, University

- of Copenhagen, Juliane Maries Vej 30, DK-2100 Copenhagen,
- ¹⁹ Department of Astronomy, University of Michigan, 1085 S. University Avenue, Ann Arbor, MI 48109, USA
- ²⁰ Department of Astronomy, University of California, Berke-
- ley, CA 94720-3411, USA 21 Department of Physics, The University of Hong Kong, Pokfulam Road, Hong Kong
- ²² Las Cumbres Observatory Global Telescope Network, 6740 Cortona Dr., Suite 102, Goleta, California 93117, USA

 ²³ Department of Physics, University of California, Santa Bar-
- bara, CA 93106-9530, USA $^{24}\,\mathrm{The~Hebrew~University,~The~Edmond~J.~Safra~Campus~-}$
- Giyat Ram, Jerusalem 9190401, Israel ⁵ INAF, Osservatorio Astronomico di Bologna, via Ranzani
- 1, I-40127 Bologna, Italy ²⁶ Jet Propulsion Laboratory, California Institute of Technol-
- ogy, 4800 Oak Grove Drive, Pasadena, CA 91109, USA ²⁷ INFN, Sezione di Bologna, Viale Berti Pichat 6/2, I-40127
- Bologna, Italy
 ²⁸ Department of Physics, University of Oxford, Keble Road, Oxford OX1 3RH, UK
- ²⁹ CRAL, Observatoire de Lyon, Université Lyon 1, 9 Avenue Ch. André, F-69561 Saint Genis Laval Cedex, France

 30 Department of Physics, Western Kentucky University,
- Bowling Green, KY 42101, USA

 31 Department of Physics and Astronomy, University of Cali-
- fornia, Los Angeles, CA 90095
 - ³² Packard Fellow
- ³³ School of Physics and Astronomy, University of Minnesota, 116 Church Street SE, Minneapolis, MN 55455, USA ³⁴ California Institute of Technology, 1200 East California
- Boulevard, Pasadena, CA 91125

1. INTRODUCTION

Galaxy clusters can be used as cosmic telescopes to magnify distant background objects through gravitational lensing, which can substantially increase the reach of deep imaging surveys. The lensing magnification enables the study of objects that would otherwise be unobservable because they are either intrinsically faint (e.g. ??) or extremely distant (e.g. ??????????). Background galaxies are also spatially magnified, allowing for studies of the internal structure of galaxies in the early universe with resolutions of ~ 100 pc (e.g. ?????).

Gravitational lensing can also provide a powerful window onto the *transient* sky through an appropriately cadenced imaging survey. The flux magnification from strong-lensing clusters is especially valuable for the study of z > 1.5 supernovae (SNe) (e.g. ???????), which are still extremely difficult to characterize in unlensed fields (e.g. ???????).

In the case of lensed SNe, we can also reverse the experimental setup: instead of using strong-lensing clusters to study distant SNe, we can use the SNe as tools for examining the lenses (?). The most valuable transients for testing and improving cluster lens models would be strongly lensed SNe that are resolved into multiple images (??). Transients that are lensed into multiple images can also become cosmological tools, as the measurement of time delays between the images can provide cosmographic information to constrain the Hubble parameter (?) and other cosmological parameters (?). We have recently observed the first example of a multiply-imaged SN (?). We expect to detect the reappearance of this object (called "SN Refsdal") within the next year (???), delivering a precise test of lens model predictions. Although detections of such objects are currently very unlikely (?), they will become much more common in the next decade (??), and may be developed into an important new cosmological tool (??).

In addition to time delays from multiply-imaged SNe, we can also put cluster mass models to the test with the much more common category of Type Ia SNe that are magnified but not multiply-imaged. ?, hereafter P14 and ? presented independent analyses of three lensed SNe, of which at least 2 are securely classified as Type Ia SNe all found in the Cluster Lensing and Supernova survey with Hubble (CLASH, PI:Postman, HST Program ID 12068, ?). Both groups demonstrated that these standard candles can be used to provide accurate and precise measurements of the true absolute magnification along a random sight line through the cluster. Although in these cases the SNe were used to test the cluster mass models, one could in principle incorporate the measured magnifications of Type Ia SNe into the cluster as additional model constraints. In that role, Type Ia SNe have the particular value that they can be found anywhere in the cluster field. Thus, they can deliver model constraints in regions of "middle distance" from the cluster core, where both strong- and weak-lensing constraints are unavailable. Moreover, given enough time, multiple background SNe Ia could be measured behind the same cluster, each providing a new magnification constraint.

One of the key values in observing standard candles behind gravitational lenses is in addressing the problem of the mass-sheet degeneracy (??). This degeneracy arises

because one can introduce into a lens model an unassociated sheet of uniform mass in front of or behind the lens, without disturbing the primary observable quantities. For example, take a lens model with a given surface mass density κ , and then transform the surface mass density to $\kappa' = (1 - \lambda)\kappa + \lambda$ for any arbitrary value λ . Both the κ and κ' models will produce exactly the same values for all positional and shear constraints from strong and weak lensing (?). When lensed background sources are available across a wide range of redshifts (as is the case for the Abell 2744 cluster discussed here), it should in principle be possible to break this degeneracy (??). However, there are more complex versions of positional constraint degeneracies (??). Such degeneracies do not extend to the absolute magnification of a background source's flux and size. In the case of the simple mass sheet degeneracy described above, the magnification scales as $\mu \propto (1 - \lambda)^{-2}$ (see e.g. ?). Therefore, one can break these fundamental degeneracies with an absolute measurement of magnification from a standard candle (?) or a standard ruler (?).

In Section 2 we present the discovery and follow-up observations of SN HFF14Tom at z=1.3457, discovered behind the galaxy cluster Abell 2744. Section 3 examines the SN host galaxy. Sections 4 and 5 describe the spectroscopy and photometry of this SN, leading to a classification of the object as a normal Type Ia SN. In Section 6 we make a direct measurement of the magnification of this source due to gravitational lensing. Section 7 discusses the tension between our magnification measurement and the lens models. Our conclusions are summarized in Section 8, along with a discussion of future prospects.

2. DISCOVERY, FOLLOW-UP, AND DATA PROCESSING

SN HFF14Tom was discovered in Hubble Space Telescope (HST) observations with the Advanced Camera for Surveys (ACS) in the F606W and F814W bands (V and I), collected on UT 2014 May 15 as part of the Hubble Frontier Fields (HFF) survey (PI:J.Lotz, HST-PID:13495).³⁵ The HFF program is a 3-year Director's discretionary initiative that is collecting 140 orbits of HST imaging (roughly 340 ksec) on six massive galaxy clusters, plus 6 accompanying parallel fields. Each field is observed in 3 optical bands (ACS F435W, F606W and F814W) and 4 infrared (IR) bands (WFC3-IR F105W, F125W, F140W, and F160W), although the optical and IR imaging campaigns are separated by ~ 6 months. Abell 2744 was the first cluster observed, with IR imaging spanning 2013 October-November, and optical imaging from 2014 May-July. A composite image of the HFF data showing the SN is presented in Figure 1, and the locations of the cluster center, the SN, and the presumed host galaxy are given in Table 1. The SN detection was made in difference images constructed using template imaging of Abell 2744 from HST+ACS observations taken in 2009 (PI:Dupke, HST-PID:11689).

Upon discovery, HST target-of-opportunity observations were triggered from the FrontierSN program (PI:Rodney, HST-PID:13386), which aims to discover and follow transient sources in the HFF cluster and parallel fields. The FrontierSN observations provided WFC3-

³⁵ http://www.stsci.edu/hst/campaigns/frontier-fields




Figure 1. SN HFF14Tom in the Abell 2744 field. The left panel shows a UV/Optical/IR color composite image constructed from all available HST imaging of the Abell 2744 cluster field. The inset panels on the right show F814W imaging of the immediate vicinity of SN HFF14Tom, approximately 40" from the center of the cluster. The top panel shows the template image, combining all data prior to the SN appearance. Labeled ellipses mark the nearest galaxies and their spectroscopic redshift constraints, with the most likely host galaxy marked in blue, and a background galaxy in yellow. The bottom panel is constructed from all HFF F814W imaging taken while the SN was detectable, and marks the SN location with an arrow. (Left panel image credit: NASA, ESA, and J. Lotz, M. Mountain, A. Koekemoer, and the HFF Team (STScI))

| Object | R.A. | Decl. | R.A. | Decl. |
|-------------------------|-------------|-------------|----------|------------|
| | (h:m:s) | (d:m:s) | (deg) | (deg) |
| HFF14Tom | 00:14:17.87 | -30:23:59.7 | 3.574458 | -30.399917 |
| Host galaxy | 00:14:17.88 | -30:24:00.6 | 3.574483 | -30.400175 |
| Abell 2744 ^a | 00:14:21.20 | -30:23:50.1 | 3.588333 | -30.397250 |

^a Coordinates of the HFF field center, approximately at the center of the cluster.

IR imaging as well as spectroscopy of the SN itself using the ACS G800L grism, supplementing the rapid-cadence optical imaging from HST+ACS already being provided by the HFF program. The last detections in the IR F105W and F140W bands came from the direct-imaging component of the GLASS program. Difference images for the IR follow-up data were generated using templates constructed from the HFF WFC3-IR imaging campaign, which concluded in November, 2013.

All of the imaging data were processed using the sndrizpipe pipeline,³⁶ a custom data reduction package in Python that employs the DrizzlePac tools from the Space Telescope Science Institute (STScI) (?). Photometry was collected using the PyPhot software package,³⁷ a pure-Python implementation of the photometry algorithms from the IDL AstroLib package (?), which in turn are based on the DAOPHOT program (?). For the IR

bands we used point spread function (PSF) fitting on the difference images, and in the ACS optical bands we collected photometry with a 0".3 aperture. Table 2 presents the list of observations, along with measured photometry from all available imaging data.

3. HOST GALAXY

The most probable host galaxy for SN HFF14Tom is a faint and diffuse galaxy immediately to the south-east of the SN location. With photometry of the host galaxy collected from the template images, we fit the spectral energy distribution (SED) using the BPZ code – a Bayesian photometric redshift estimator (?). From the BPZ analysis, we found the host to be most likely an actively star-forming galaxy at a redshift of $z=1.5\pm0.2$. This photoz was subsequently refined to a spectroscopic redshift of $z=1.3457\pm0.0001$, based on optical spectroscopy of the host (Mahler et al. in prep.) that shows two significant (> 10σ) emission lines at 8740.2 and 8746.7 Angstroms, consistent with the [OII] $\lambda\lambda$ 3726-3729 Å doublet.

The next nearest galaxy detected in HST imaging is 2.2''northeast of the SN position. It has a redshift of z=1.742 determined from a spectrum taken with the G141 grism of the HST WFC3-IR camera, collected as part of the Grism Lens-Amplified Survey from Space (GLASS, PI:Treu, PID:13459, ??). As we will see in Sections 4 and 5, both the spectroscopic and photometric data from the SN itself are consistent with the redshift of the fainter galaxy at z=1.3457, and incompatible with z=1.742 from this brighter galaxy. This means that the latter galaxy is a background object and therefore has no impact on the SN magnification.

³⁶ https://github.com/srodney/sndrizpipe DOI:10.5281/zenodo.10731

³⁷ https://github.com/djones1040/PyPhot

Table 2
HFF14Tom Observations and Photometry

| Obs. Date | Camera | Filter | Exp. Time | Flux | Flux Err | $AB Mag^a$ | Mag Err | AB Zero Point | $\Delta \mathrm{ZP^b}$ |
|-----------|-------------|----------------|-----------|-----------------|---------------|---------------|---------------------|---------------|------------------------|
| (MJD) | | or grism | (sec) | (counts/sec) | (counts/sec) | | | | (Vega-AB) |
| 56820.06 | ACS | F435W | 5083 | -0.027 | 0.053 | 27.66 | | 25.665 | -0.102 |
| 56821.85 | ACS | F435W | 5083 | 0.105 | 0.053 | 28.11 | 0.55 | 25.665 | -0.102 |
| 56823.77 | ACS | F435W | 5083 | 0.103 0.022 | 0.053 | 29.80 | 2.59 | 25.665 | -0.102 |
| 56824.97 | ACS | F435W | 5083 | 0.022 | 0.053 | 29.85 | $\frac{2.39}{2.72}$ | 25.665 | -0.102 |
| | ACS | | | -0.148 | | | 2.12 | | |
| 56828.68 | | F435W | 5083 | | 0.053 | 27.65 | | 25.665 | -0.102 |
| 56830.87 | ACS | F435W | 5083 | 0.100 -0.080 | 0.054 | 28.16 | 0.58 | 25.665 | -0.102 |
| 56832.86 | ACS | F435W | 5083 | | 0.053 | 27.66 | | 25.665 | -0.102 |
| 56833.86 | ACS | F435W | 5083 | -0.002 | 0.053 | 27.67 | • • • | 25.665 | -0.102 |
| 56839.50 | ACS | F435W | 5083 | -0.022 | 0.052 | 27.68 | • • • | 25.665 | -0.102 |
| 56792.06 | ACS | F606W | 5046 | 0.363 | 0.083 | 27.59 | 0.25 | 26.493 | -0.086 |
| 56792.98 | ACS | F606W | 3586 | 0.692 | 0.095 | 26.89 | 0.15 | 26.493 | -0.086 |
| 56797.10 | ACS | F606W | 4977 | 0.968 | 0.087 | 26.53 | 0.10 | 26.493 | -0.086 |
| 56800.08 | ACS | F606W | 4977 | 0.844 | 0.085 | 26.68 | 0.11 | 26.493 | -0.086 |
| 56804.99 | ACS | F606W | 5046 | 0.977 | 0.086 | 26.52 | 0.10 | 26.493 | -0.086 |
| 56792.99 | ACS | F814W | 3652 | 1.639 | 0.104 | 25.41 | 0.07 | 25.947 | -0.424 |
| 56797.11 | ACS | F814W | 4904 | 3.376 | 0.141 | 24.63 | 0.05 | 25.947 | -0.424 |
| 56798.95 | ACS | F814W | 5046 | 3.951 | 0.156 | 24.46 | 0.03 | 25.947 | -0.424 |
| 56800.10 | ACS | F814W | 4904 | 3.854 | 0.155 | 24.48 | 0.04 | 25.947 | -0.424 |
| 56801.89 | ACS | F814W | 10092 | 4.102 | 0.153 | 24.40 24.41 | 0.04 0.04 | 25.947 | -0.424 |
| 56802.95 | ACS | F814W | 10092 | 4.102 4.325 | 0.160 | 24.41 24.36 | 0.04 0.04 | 25.947 | -0.424 |
| | ACS | | | 4.402 | | 24.30 24.34 | $0.04 \\ 0.04$ | | |
| 56803.93 | | F814W | 15138 | | 0.160 | | | 25.947 | -0.424 |
| 56804.08 | ACS | F814W | 5046 | 4.658 | 0.178 | 24.28 | 0.04 | 25.947 | -0.424 |
| 56812.08 | ACS | F814W | 637 | 4.705 | 0.258 | 24.27 | 0.06 | 25.947 | -0.424 |
| 56815.93 | ACS | F814W | 446 | 4.026 | 0.285 | 24.43 | 0.08 | 25.947 | -0.424 |
| 56820.07 | ACS | F814W | 5044 | 3.508 | 0.142 | 24.58 | 0.04 | 25.947 | -0.424 |
| 56821.87 | ACS | F814W | 5044 | 3.541 | 0.144 | 24.57 | 0.04 | 25.947 | -0.424 |
| 56823.79 | ACS | F814W | 5044 | 2.876 | 0.124 | 24.80 | 0.05 | 25.947 | -0.424 |
| 56824.99 | ACS | F814W | 5044 | 3.060 | 0.129 | 24.73 | 0.05 | 25.947 | -0.424 |
| 56828.70 | ACS | F814W | 5044 | 2.777 | 0.121 | 24.84 | 0.05 | 25.947 | -0.424 |
| 56830.89 | ACS | F814W | 5044 | 2.395 | 0.111 | 25.00 | 0.05 | 25.947 | -0.424 |
| 56832.88 | ACS | F814W | 5044 | 2.331 | 0.108 | 25.03 | 0.05 | 25.947 | -0.424 |
| 56833.88 | ACS | F814W | 5044 | 2.389 | 0.111 | 25.00 | 0.05 | 25.947 | -0.424 |
| 56839.52 | ACS | F814W | 5044 | 1.673 | 0.093 | 25.39 | 0.06 | 25.947 | -0.424 |
| 56833.14 | WFC3-IR | F105W | 756 | 7.504 | 0.239 | 24.08 | 0.03 | 26.269 | -0.645 |
| 56841.82 | WFC3-IR | F105W | 756 | 5.822 | 0.208 | 24.36 | 0.04 | 26.269 | -0.645 |
| 56850.06 | WFC3-IR | F105W | 756 | 3.952 | 0.207 | 24.78 | 0.06 | 26.269 | -0.645 |
| 56860.62 | WFC3-IR | F105W | 1159 | 2.899 | 0.167 | 25.11 | 0.06 | 26.269 | -0.645 |
| 56886.63 | WFC3-IR | F105W | 1159 | 1.216 | 0.147 | 26.06 | 0.13 | 26.269 | -0.645 |
| 56891.67 | WFC3-IR | F105W | 356 | 0.971 | 0.324 | 26.30 | 0.36 | 26.269 | -0.645 |
| 56893.20 | WFC3-IR | F105W | 712 | 0.954 | 0.242 | 26.32 | 0.28 | 26.269 | -0.645 |
| 56954.64 | WFC3-IR | F105W | 356 | 0.521 | 0.388 | 26.98 | 0.81 | 26.269 | -0.645 |
| | WFC3-IR | F125W | 1206 | | 0.191 | 23.91 | 0.02 | 26.230 | |
| 56817.08 | | F125W F125W | | 8.459 | | 23.91 24.01 | | | -0.901 |
| 56833.15 | WFC3-IR | | 756 | 7.753 | 0.255 | | 0.04 | 26.230 | -0.901 |
| 56841.83 | WFC3-IR | F125W | 806 | 6.015 | 0.227 | 24.28 | 0.04 | 26.230 | -0.901 |
| 56850.07 | WFC3-IR | F125W | 806 | 4.343 | 0.224 | 24.64 | 0.06 | 26.230 | -0.901 |
| 56891.86 | WFC3-IR | F140W | 712 | 2.578 | 0.344 | 25.42 | 0.14 | 26.452 | -1.076 |
| 56893.06 | WFC3-IR | F140W | 712 | 3.026 | 0.363 | 25.25 | 0.13 | 26.452 | -1.076 |
| 56955.58 | WFC3-IR | F140W | 1424 | 1.218 | 0.269 | 26.24 | 0.24 | 26.452 | -1.076 |
| 56817.09 | WFC3-IR | F160W | 1206 | 4.831 | 0.263 | 24.24 | 0.06 | 25.946 | -1.251 |
| 56833.21 | WFC3-IR | F160W | 756 | 3.965 | 0.241 | 24.45 | 0.00 | 25.946 | -1.251 |
| 56841.84 | WFC3-IR | F160W | 756 | 3.011 | 0.234 | 24.45 24.75 | 0.07 | 25.946 | -1.251 |
| 56850.08 | WFC3-IR | F160W | 756 | 2.744 | 0.223 | 24.75 | 0.08 | 25.946 | -1.251 |
| 56860.67 | WFC3-IR | F160W | 1159 | 1.895 | 0.223 0.177 | 25.25 | 0.09 | 25.946 | -1.251 |
| 56886.64 | WFC3-IR | F160W | 1159 | 1.677 | 0.177 | 25.25 25.38 | $0.10 \\ 0.12$ | 25.946 | -1.251 |
| 50000.04 | M I O 9-111 | 1. 100 44 | 1109 | 1.077 | 0.131 | 20.00 | 0.12 | 20.340 | -1.201 |
| F0010 = | . ~~ | Cocci | 2.40- | | | | | | |
| 56812.0 | ACS | G800L | 3490 | • • • | • • • | • • • | • • • | • • • | • • • |
| 56815.7 | ACS | G800L | 6086 | • • • | • • • | • • • | • • • | • • • | • • • |
| | | | | | | | | | |

 $[\]overline{}^a$ For non-positive flux values we report the magnitude as a 3- σ upper limit b Zero point difference: the magnitude shift for conversion from AB to Vega magnitude units.

The galaxy identified as the host is not close enough to the cluster core to necessarily be multiply-imaged, but it is still possible that the host galaxy is one of the outer images of a multiple image system. In such a case, and given the position of the SN host, one would expect that another image of the galaxy would be present at a similar brightness, and would therefore be detectable in HST imaging. To date, no plausible candidate for a counterimage has been identified.

To measure the stellar mass of the host galaxy we use Eq. 8 of ?, which relates the rest-frame (g-i) color and iband luminosity to the total stellar mass. To derive these values, we fixed the redshift at z=1.3457 and repeated the SED fitting using BPZ. From the best-fit SED we extracted rest-frame optical magnitudes, and corrected them for lensing using a magnification factor of $\mu=2.0$ – a value that we will derive from the SN itself in Section 6. From this we determine the host galaxy mass to be $10^{9.8}$ ${\rm M}_{\odot}$.

4. Spectroscopy

A spectrum of SN HFF14Tom was collected with the ACS G800L grism on 2014 June 4 and 7, when the SN was within 3 observer-frame days of the observed peak brightness in the F814W band. The observations – listed at the bottom of Table 2 – used 5 HST orbits from the FrontierSN program for a total spectroscopic exposure time of $\sim \! \! 10$ ksec. The grism data were processed and the target spectrum was extracted using a custom pipeline (?), which was developed for the 3D-HST program (PI:Van Dokkum; PID:12177, 12328) and also used by the GLASS team.

Figure 2 shows the composite 1-D ACS grism spectrum, combining all available G800L exposures, overlaid with SN model fits that will be described below. The spectrum is largely free of contamination, because the orientation was chosen to avoid nearby bright sources and the host galaxy is diffuse and optically faint. Thus, the SN spectral features can be unambiguously identified, most notably the red slope of the continuum and a prominent absorption feature at $\sim 8700 \,\text{Å}$.

Spectra of Type I SNe (including all sub-classes Ia, Ib and Ic) are dominated by broad absorption features, which cannot in general be used to directly extract a spectroscopic redshift (see e.g. ?). As described below, we fit template spectra to the SN HFF14Tom data in two steps. First we determine a spectral classification – and get a preliminary estimate of the redshift and age – using the SuperNova IDentification (SNID) software (?). Second, we refine the redshift and age measurement using a custom Type Ia spectral template matching program.

4.1. Classification with SNID

The SNID program is designed to estimate the type, redshift, and age of a SN spectrum through cross-correlation matching with a library of template spectra, using the algorithm of ?. To account for possible distortions in the broad shape of the SN pseudo-continuum due to dust or instrumental calibration effects, SNID divides each SED by a smooth cubic spline fit. This effectively removes the shape of the SN pseudo-continuum to leave behind a flat SED superimposed with spectral absorption and emission features. It is these features which drive the cross-correlation fit, so the SNID approach is




Figure 2. Redshift and age determination from spectral template matching to the the SN HFF14Tom maximum light spectrum. The y axis plots flux in arbitrary units, and the x axis marks wavelength in Å with the observer-frame on the bottom and rest-frame on the top. The HFF14Tom spectrum observed with the HST ACS G800L grism is shown in black, overlaid with model fits derived from a library of Type Ia templates that have extended rest-frame UV coverage. The top two panels show the best matching templates when using a smooth 3rd-order polynomial to warp the shape of the template pseudo-continuum, with the redshift fixed at z=1.3457 and then allowed to float as a free parameter. The lower three panels show matches found when the templates are not warped, both with and without fixing the redshift. The bottom panel shows the best match in this set, although at z=0.98 it is inconsistent with the host galaxy redshift prior and the light curve.

insensitive to the overall color of the SED. We used v2.0 of the SNID template library, which includes template SEDs covering all Type Ia and core collapse sub-classes, and has recently been updated with corrections and improvements to the Type Ib/c templates (?).

In SNID the goodness of fit is evaluated primarily through the rlap parameter, which measures the degree of wavelength overlap and the strength of the cross-correlation peak. Typically, an rlap value > 5 is required to be considered an acceptable match.

To match the SN HFF14Tom spectrum we use conservative constraints on age and redshift: limiting the age to ± 5 rest-frame days from peak brightness and 0.8 < z < 1.8, consistent with the SN light curve and the two plausible host galaxies. With these constraints we find that the only acceptable match is a normal Type Ia SN near z=1.3. The best match has rlap=8.7, using the normal Type Ia SN 2005cf at z=1.35 and age=-2.2 rest-frame days before peak. In contrast, the best non-Ia matches all have rlap<2.5.

Using SNID we can find an acceptable CCSN match only when we remove all age and redshift constraints. In this case the best non-Ia match is the Type IcSN 1997ef, which delivers rlap=6.8 at z=0.51 and age=47.3 restframe days past peak (71 observer-frame days). This is not as good a fit as the best Type Ia models, is at odds with the host galaxy redshift prior, and is strongly disfavored by the shape and colors of the SN light curve (see Section 5).

From the preceding analysis, we conclude that HFF14Tom is a Type Ia SN at $z\approx 1.35$. At this redshift, the absorption at $\sim\!8700\text{Å}$ corresponds to the blended Ca II H&K and Si II $\lambda 3858$ features. Generally referred to as the Ca H&K feature, this absorption is commonly seen in Type Ia SN spectra near maximum light, although it is also prominent in the spectra of Type Ib and Ic core collapse SNe (CC SNe). The red color of the HFF14Tom SED is qualitatively consistent with a redshift of z>1 – although this information was not used by SNID for the template matching. As we will see in Section 5, this spectral classification of SN HFF14Tom is reinforced by the photometric information, which also supports classification as a Type Ia SN at $z\approx 1.35$.

4.2. Spectral Fitting with UV Type Ia Templates

To refine the redshift and phase constraints on HFF14Tom, we next fit the spectrum with a custom spectral matching program that employs a library of Type Ia SN SEDs. This library is similar to the Type Ia spectral set used by SNID, but also includes more recent SNe with well-observed spectral time series that extend to rest-frame UV wavelengths (e.g. SN 2011fe and 2014J). We first use an approach similar to the SNID algorithm: warping the pseudo-continuum of each template spectrum by dividing out a 3rd-order polynomial to match the observed SED of SN HFF14Tom. This approach will find templates that have similar abundances and photospheric velocities. With the redshift fixed at z = 1.3457we find the best fit is a spectrum from the normal Type Ia SN 2014J (?), with a χ^2 per degree of freedom ν of $\chi^2/\nu = 121.8/89$ shown in the top panel of Figure 2. The excess variance in this fit may be attributed to the intrinsic variation of Type Ia SN spectra, which is more

prominent at UV wavelengths (e.g. ??) and is not fully represented in the available template library. Allowing the redshift as an additional free parameter, we still find results that are consistent with the SNID fits and $< 2\sigma$ from the host galaxy redshift: $z = 1.31 \pm 0.02$ and a phase of 0 ± 3 rest-frame days. The best-fitting spectral template in this case is the normal Type Ia SN 2012cg (?) at z = 1.31 with $\chi^2/\nu = 98.5/88$ (second panel of Figure 2).

Next, we repeat the fitting, but without any warping of the templates to account for differences in the continuum shape. In this iteration we only allow each template SED to be scaled in flux coherently at all wavelengths, so the fits are more sensitive to the overall color of the SED.³⁸ Fixing the redshift to z=1.3457, we find the best match is from the normal Type Ia SN 2011fe (?), though the fit is quite poor, with $\chi^2/\nu=313.2/91$ (third panel of Figure 2). When the redshift is allowed as a free parameter the HFF14Tom SED is still matched best by a SN 2011fe template, now at redshift z=1.31 with $\chi^2/\nu=240.8/90$ (fourth panel of Figure 2). SN 2011fe had effectively no dust reddening (e.g. ??), so the fact that SN 2011fe provides the best un-warped template match is further evidence that SN HFF14Tom suffers from very little dust extinction.

Without the continuum warping, an alternative fit also arises: the fast-declining (91bg-like) Type Ia SN 2011iv (?) at $z=0.98\pm0.01$. Formally, this match provides a slightly better fit to the unwarped HFF14Tom spectrum ($\chi^2/\nu=220.6/90$, bottom panel of Figure 2), although the fit is notably poorer at $\sim 8700\text{Å}$ where the most significant absorption feature is found. Furthermore, a redshift $z\sim1$ is at odds with the spectroscopic redshift of the nearest galaxy (z=1.3457), and we will see in the following section that the photometric data is also incompatible with a Type Ia SN at $z\sim1.0$.

Setting aside the $z\sim 1$ solution, all other template matches provide a consistent redshift constraint of $z=1.31\pm0.02$, regardless of whether the templates are warped to match the SN HFF14Tom continuum shape. The inferred age from these fits is 0 ± 3 rest-frame days from peak brightness, which is also consistent with the observed light curve. Taken together with the host galaxy spectroscopic redshift, these fits suggest that SN HFF14Tom is a normal Type Ia SN at z=1.3457 with an SED color close to SN 2011fe, but spectral absorption features similar to SN 2014J.

5. PHOTOMETRIC CLASSIFICATION

Relative to other SNe at z>1, the SN HFF14Tom light curve was unusually well sampled at rest-frame ultraviolet wavelengths, due to the rapid cadence of the HFF imaging campaign. These ACS observations therefore provide a tight constraint on the time of peak brightness and the evolution of the SN color. Supplemental observations with the WFC3-IR camera provided critical rest-frame optical photometry, enabling a measurement of the apparent luminosity distance through light curve fitting.

As a check on the spectral classification of SN

³⁸ Note that gravitational lensing does not affect the color of background sources at all, so the expected lensing magnification of HFF14Tom does not affect this analysis.




Figure 3. Maximum likelihood model for each SN sub-class, derived from Bayesian model selection using the photometric data alone. Grey points show the observed SN HFF14Tom photometry with error bars, though these are typically smaller than the size of the marker. The Type Ia model (orange solid line) is drawn from the SALT2 template at z=1.35. The best match from all Type Ib/Ic models is based on the Type Ic SN SDSS-14475 at z=0.695 (green dashed line). For the Type II class, the best match is from the Type II-L SN 2007pg at z=1.8 (blue dash-dot line). The Type Ia model is by far the best match, and the only one that is consistent with both the photo-z of the probable host galaxy and the spectroscopic redshift from the SN spectrum. The date of the HST spectral observations is marked with a thin grey vertical line.

HFF14Tom (Section 4.1), we independently classified the SN using a Bayesian photometric classifier. We use the sncosmo software package³⁹ to simulate SN light curves from z = 0.3 to 2.3 and evaluate the classification probability using traditional Bayesian model selection (as in ????). In this analysis we represent normal Type Ia

The CCSN models have free parameters for date of peak brightness $(t_{\rm pk})$, amplitude, and redshift (z). Due to the expected impact of gravitational lensing magnification, we do not include any prior on the intrinsic luminosity for any SN sub-class. We also do not assign a prior for the SN redshift. This allows our photometric analysis to provide an independent check on the host galaxy photo-z and spectroscopic redshift (Sections 2 and 4.2).

For Type Ia SNe, the SALT2 model has two additional parameters that control the shape (x_1) and color (c) of the light curve. We use conservative priors here, defined to encompass a range of Type Ia SN shapes and colors that is broader than typically allowed in cosmological analyses (see e.g., ???). For x_1 the prior is a bifurcated Gaussian distribution with mean $\bar{x}_1 = 0$, dispersion $\sigma_{x_1}^+ = 0.9$ and $\sigma_{x_1}^- = -1.5$. The bifurcated Gaussian prior for the color parameter c has $\bar{c} = 0.0$, $\sigma_c^- = 0.08$, and $\sigma_c^+ = 0.54$. The c parameter in SALT2 combines intrinsic SN color and extinction due to dust, so the large red tail of this distribution allows for the possibility of several magnitudes of dust extinction along the HFF14Tom line of sight.

We also assign a class prior for each of the three primary SN sub-classes (Type Ia, Ib/c, and II), using a fixed relative fraction for each sub-class as determined at z=0 by ? and ?. A more rigorous classification would extrapolate these local SN class fractions to higher redshift using models or measurements of the volumetric SN rate. For simplicity, we do not vary the class priors with redshift, and in practice these priors do not have any significant impact on the resulting classification.

The final photometric classification probability for SN HFF14Tom is $p(\text{Ia}|\mathbf{D})=1.0$, with the classification probability from all CCSN sub-classes totaling less than 10^{-32} . Although this Bayesian classification utilizes the full posterior probability distribution, for illustration we highlight in Figure 3 a single best-fit model for each subclass. This demonstrates how the CCSN models fail to adequately match the observed photometry. In particular, only the Type Ia model can simultaneously provide an acceptable fit to the well-sampled rising light curve in F814W and the F814W-F160W color near peak.

The marginal posterior distribution in redshift for the Type Ia model is sharply peaked at $z=1.35\pm0.02$, which is fully consistent with the redshift of the presumed host galaxy (z=1.3457) as well as the redshift of $z=1.31\pm0.02$ derived from the SN spectrum in Section 4.2. The time of peak brightness is also tightly constrained at $t_{\rm pk}=56816.3\pm0.3$, which means the HST grism observations were collected within 2 rest-frame days of the epoch of peak brightness – also consistent with our spectroscopic analysis.

SNe with the SALT2 model (?), and CCSNe with 42 discrete templates (26 Type II and 16 Type Ib/c) drawn from the template library of the SuperNova Analysis software package (SNANA, ?).⁴⁰ Likelihoods are defined by comparing the observed fluxes to model predictions in all passbands where the model is defined. In practice, this means we exclude the SN detections in the F435W and F606W bands, which are too blue for our models at z > 0.85.

³⁹ http://sncosmo.github.io/

⁴⁰ Throughout this work we use SNANA v10₋35g.

8

| Fitter | Distance | Modulus | Measured |
|------------------|--|---------------------------------|---------------------------------|
| | HFF14Tom | Control | Magnification |
| MLCS2k2 SALT2 | 44.205 ± 0.12 44.177 ± 0.18 | $44.97 \pm 0.06 44.92 \pm 0.06$ | 2.03 ± 0.29 1.99 ± 0.38 |

6. DISTANCE MODULUS AND MAGNIFICATION

With the type and redshift securely defined as a normal Type Ia SN at z = 1.3457, we now turn to fitting the light curve with Type Ia templates to measure the distance modulus (§6.1) and then derive the gravitational lensing magnification ($\S6.2$). In this process, we would like to avoid introducing systematic uncertainties inherent to any assumed cosmological model. To that end, in section 6.2 we follow? and define the magnification by comparing the measured distance modulus of SN HFF14Tom against an average distance modulus derived from a "control sample" of unlessed Type Ia SNe at similar redshift. This allows us to make only the minimal assumption that the redshift-distance relationship for Type Ia SN is smooth and approximately linear over a small redshift span, which should be true for any plausible cosmological model.

6.1. Light Curve Fitting

As in ?, we derive a distance modulus for SN HFF14Tom and all SNe in our control sample using two independent light curve fitters: the SALT2 model described above and the MLCS2k2 model (?). With both fitters we find light curve shape and color parameters for HFF14Tom that are fully consistent with a normal Type Ia SN. For SALT2, with the redshift fixed at z=1.3457, we find a light curve shape parameter of $x_1=0.135\pm0.199$ and a color parameter of $c=-0.127\pm0.025$, yielding a χ^2 value of 45.0 for 36 degrees of freedom, ν . With the MLCS2k2 fitter the best-fit shape parameter is $\Delta=-0.082\pm0.070$ and the color term is $A_{\rm V}=0.011\pm0.025$, giving $\chi^2/\nu=22.8/36$.

The MLCS2k2 fitter returns a distance modulus⁴¹ dm_{MLCS2k2} directly, as it is defined to be one of the free parameters in the model. To derive dm from the SALT2 fit, we use

$$dm_{SALT2} = m_B^* - M + \alpha(s-1) - \beta C. \tag{1}$$

Here the parameters for light curve shape s and color C correspond to the SiFTO light curve fitter (?), so we first use the formulae from ? to convert from SALT2 (x_1 and c) into the equivalent SiFTO parameters. We also add an offset of 0.27 mag to the value of m_B^* returned by SNANA, in order to match the arbitrary normalization of the SALT2 fitter used by ? and ?. This conversion from SALT2 to SiFTO is necessary, as it allows us to adopt values for the constants M, α , and β from ?, which have been calibrated using 472 SNe from the SNLS3 sample (?): $M = -19.12 \pm 0.03$, $\alpha = 1.367 \pm 0.086$, and $\beta = 3.179 \pm 0.101$.

The SNANA version of the MLCS2k2 fitter returns a value for the distance modulus ($dm_{MLCS2k2}$) that has an arbitrary zero point offset relative to the SALT2 distances (dm_{SALT2}). To put the two distances onto the same reference frame we add a zeropoint correction of 0.20 mag to the MLCS2k2 distances as in ?. This correction was derived by applying both fitters to a sample of Type Ia SNe from the SDSS survey (??), with the extinction law R_V fixed at 1.9.

The total uncertainty in the distance modulus is

$$\sigma_{\text{tot}} = \sqrt{\sigma_{\text{stat}}^2 + \sigma_{int}^2}.$$
 (2)

The $\sigma_{\rm stat}$ term is the statistical uncertainty, which encapsulates uncertainties from the data and the model, and $\sigma_{\rm int}$ accounts for the remaining unmodeled scatter. This latter term is derived by finding the amount of additional distance modulus scatter that needs to be added to a Type Ia SN population to get χ^2 per degree of freedom equal to 1 for a fiducial cosmological model fit to the SN Ia Hubble diagram (distance modulus vs redshift). Thus, $\sigma_{\rm int}$ is designed to account for any unknown sources of scatter in the Type Ia SN population, including unidentified errors in the data analysis as well as the natural scatter in intrinsic Type Ia SN luminosities. The value of $\sigma_{\rm int}$ may be expected to vary as a function of redshift and also from survey to survey.

Recent work has highlighted the inadequacy of this simplistic approach for handling intrinsic scatter in the Type Ia SN population (?????), but a full consideration of those alternative approaches is beyond the scope of this work. We adopt the simple approach of using a single empirically defined value for $\sigma_{\rm int}$, reflecting principally an intrinsic scatter in Type Ia SN intrinsic luminosity that is fixed across time and phase. Measurements of $\sigma_{\rm int}$ range from 0.08 mag (??) to 0.15 mag (??). We adopt a value of $\sigma_{\rm int} = 0.08$ mag, as derived by ? and ?. Although this is on the low end of the range reported in the literature, this value is the most appropriate to apply to our analysis for two reasons. First and foremost, for the SALT2 fitter we are using light curve fit parameters (α,β) with associated uncertainties that have been derived from the joint analysis of? and?. Similarly, the implementation of MLCS2k2 that we have used is based on the uncertainty model derived from model training in ?. Inflating σ_{int} beyond 0.08 mag would therefore be equivalent to driving the reduced χ^2 of the Type Ia SN Hubble diagram to < 1. Second, the value of 0.08 mag determined in? is specific to the HST SN sample, from ? and ?, which is the SN subset that is most similar to HFF14Tom in terms of redshift and data analysis. Larger values for $\sigma_{\rm int}$ are typically associated with SN samples at significantly lower redshifts that have less homogeneous data collection and analysis.

Final values for the SN distance modulus are shown in Table 3, adopting the spectroscopic redshift (z=1.3457). In addition to modifying the distance modulus uncertainty for SN HFF14Tom, we also add $\sigma_{\rm int}=0.08$ mag in quadrature to the uncertainty for every SN in the control sample. Note however that the uncertainty on the control sample value at z=1.3457 is only 0.06 mag, smaller than the intrinsic dispersion of any single SN because it reflects our measurement error on the mean

 $^{^{41}}$ We use 'dm' to indicate the distance modulus to avoid confusion, reserving the symbol μ to refer to the lensing magnification. This 'dm' is a standard distance modulus, defined as $\mathrm{dm} = 5\log_{10}d_L + 25,$ where d_L is the luminosity distance in Mpc.




Figure 4. Type Ia light curve fits to SN HFF14Tom using the MLCS2k2 (top row) and SALT2 (bottom row) fitters. The model redshifts are set to z=1.3457 as determined from combined spectroscopic and photometric constraints. Solid lines denote the best-fit model and shaded lines show the range allowed by 1- σ uncertainties on the model parameters. Observed fluxes are shown as diamonds, scaled to an AB magnitude zero point of 25. Error bars are plotted, but most are commensurate with the size of the points. The left-most panel includes observations in the F435W and F606W filters, although these were not used for the fit, as they are bluer than the minimum wavelength for the model. The lower axis marks time in observer frame days, while the top axis shows the time in the rest frame relative to the epoch of peak brightness.

distance modulus of the population. The distance moduli derived from the SALT2 and MLCS2k2 light curve fitters are fully consistent within the uncertainties.

6.1.1. Host Galaxy Mass Correction

It is now an accepted practice in cosmological analyses using Type Ia SN to apply a correction to the luminosity of each SN based on the stellar mass of its host galaxy. Typically this is described as a simple bifurcation of the SN population: SNe that appear in more massive hosts are observed to be $\sim\!0.08$ mag brighter (after corrections for light curve shape and color) than SNe in low-mass hosts (??). The dividing line for this purely empirical "mass step" correction is generally set around $10^{10} \rm M_{\odot}$. Although this threshold value is somewhat arbitrary (see e.g. ?), it happens to be very close to the -SN HFF14Tom host galaxy mass of $10^{9.8} \rm M_{\odot}(Section~3)$.

The physical mechanism that drives the mass step is not yet understood, but may be related to the metallicity or age of the SN progenitor systems. In either case, the significance of this effect should decrease with redshift, as metal-rich passive galaxies become much less common at z>1 (see e.g. ??). Indeed, when the size of the mass step correction is allowed to vary with redshift, there is no significant evidence that a non-zero mass step is required at z>1 (???). Given the absence of a clear physical model and the lack of empirical support for a high-z mass step, we do not apply any correction to SN HFF14Tom or the control sample.

If we were to apply the correction to this sample using the standard approach, then the SN HFF14Tom distance modulus would not be adjusted, because its host galaxy mass (Section 3) is just below $10^{10}~\rm M_{\odot}$. Note that there is some circularity here, as we used the SN magnification in deriving the host galaxy mass, and now use the host mass to inform the magnification. However, regardless of whether the HFF14Tom host mass falls above the mass threshold or below, this correction is not substantial enough to account for the observed magnification tension. The appropriate correction to be applied to a SN above the mass threshold in this redshift range would be <0.04 mag (??), which results in a change to the inferred magnification of HFF14Tom that is much less than the observed discrepancy of \sim 0.24 mag.

6.2. Control Sample Comparison

The unlensed sample comprises 22 spectroscopically confirmed Type Ia SNe in the range 1.1 < z < 1.6 from three HST surveys: 11 from the GOODS Higher-z SN search⁴² (??), 7 from SCP⁴³ (?), and 4 from CAN-DELS⁴⁴ (??). Using the SALT2 and MLCS2k2 fitters as described above, we get distance modulus measures for every object in this control sample. We then fit a linear relationship for distance modulus vs. redshift, and derive a prediction for the distance modulus of a normal Type Ia SN at the redshift of SN HFF14Tom (Figure 5). This predicted value is given in Table 3 under the "Control"

 $^{^{42}}$ GOODS: the Great Observatories Origins Deep Survey, PI:Giavalisco, HST-PID:9425,9583

⁴³ SCP: the Supernova Cosmology Project, PI:Perlmutter

 $^{^{44}}$ CANDELS: the Cosmic Assembly Near-infrared Deep Extragalactic Legacy Survey, PI:Faber & Ferguson




Figure 5. Measurement of the lensing magnification from comparison of the HFF14Tom distance modulus to a sample of unlensed field SN. SNe from the GOODS program are plotted as squares, those from the SCP survey are shown as circles, and the CANDELS objects are triangles. The distance modulus for each SN is derived from light curve fits using the MLCS2k2 fitter (top panel) and the SALT2 fitter (bottom panel). Grey lines with shading show a linear fit to the unlensed control sample is shown, anchored at the redshift of SN HFF14Tom, with fit parameters given at the top. The derived magnification is reported at the bottom of each panel.

column. The difference between the observed distance of SN HFF14Tom and this control sample value is attributed to the magnification from gravitational lensing:

$$dm_{control} - dm_{HFF14Tom} = 2.5 \log_{10} \mu. \tag{3}$$

The inferred magnifications for the two different fitters are reported in the final column of Table 3. Although we have chosen to use a cosmology-independent approach to determine the magnifications, we note that these results are fully consistent with the values that would be determined by comparing the observed SN HFF14Tom distance modulus against the predicted value from a flat Λ CDM cosmology. Adopting the cosmological parameters used in ? and derived in ? (H_0 =71.6, $\Omega_{\rm m}$ =0.27, Ω_{Λ} =0.73) would give a predicted distance modulus $dm_{\Lambda CDM}=44.90$ and a measured magnification $\mu=1.9$. Substituting alternative cosmological parameters (e.g., ??) would not significantly change the inferred magnification or affect our conclusions.

7. COMPARISON TO MODEL PREDICTIONS

Before the Frontier Fields observations began, the Space Telescope Science Institute (STScI) issued a call for lens modeling teams to generate mass models of all 6 Frontier Field clusters, using a shared collection of all imaging and spectroscopic data available at the time. In response to this opportunity, five teams generated seven models for Abell 2744. These models necessarily relied on pre-HFF data, and were required to be complete be-

fore the HFF program began, in order to enable the estimation of magnifications for any new lensed background sources revealed by the HFF imaging. An interactive web tool was created by D. Coe and hosted at STScI, to extract magnification estimates and uncertainties from each model for any given redshift and position. In this work we also consider 8 additional models that were created later. Some of these are updates of the original models produced in response to the lens modeling call, and some of them include new multiply-imaged galaxies discovered in the HFF imaging as well as new redshifts for lensed background galaxies. Magnifications derived from several of these later models are also available on the interactive web tool hosted by STScI. The full list of models and details on their construction are given in Table 4.

Table 5 gives each model's predicted magnification and uncertainty for a source at z=1.3457 and at the position of HFF14Tom. For thirteen of these models the SN magnification qualifies as a true "blind test", as they were completed before the SN HFF14Tom magnification measurement was known. The final four models (Bradac-v2, Lam-v2, CATSv2.1 and 2.2) are technically not blind, although none of these modelers used the SN magnification as a constraint, and the modelers did not have access to the final SN magnification value when constructing their model.

These models represent a broad sampling of the techniques and assumptions that can be applied to the modeling of mass distributions in galaxy clusters. We discuss some of the modeling choices here, but a more rigorous comparison of these diverse lens modeling techniques is beyond the scope of this work. For a complete discussion of each model's methodology the reader is directed to the listed references. Eight of these are so-called parametric⁴⁵ models, and the remining seven are free-form models. Broadly speaking, the parametric models use parameterized density distributions to describe the arrangement of mass within the cluster. Therefore, parametric models rely (to varying degrees) on the assumption that the cluster's dark matter can be described by analytic forms such as NFW halos (?), or pseudo isothermal elliptical mass distributions (PIEMD?).

The seven free-form models divide the cluster field into a grid, generally using a multi-scale grid to get better sampling in regions with a higher density of information (e.g. density of multiple images). Each grid cell is assigned a mass or a potential, and then the mass values are iteratively refined to match the observed lensing constraints. In some cases an adaptive grid is used so that the grid spacing itself can also be modified as the model is iterated (e.g. ???).

As usual, there is a tradeoff between a model's flexibility, the strength of the model assumptions, and the resulting uncertainties. In a probabilistic framework, the posterior distribution function of the desired quantities depends on all the priors, including model assumptions like parametrization. In general, free-form methods tend to be more flexibile than simply parameterized models,

⁴⁵ Although this nomenclature is becoming standard in the literature, it is somewhat misleading, as the pixels or grid cells in free-form models are effectively parameters as well. Perhaps "simply-parameterized" would be more accurate, though we adopt the more common usage here.

Table 4 Tested lens models for Abell 2744.

| Model | N _{sys} ^a | $N_{\mathrm{im}}{}^{\mathrm{b}}$ | $N_{\rm spec}{}^{\rm c}$ | $N_{\mathrm{phot}}^{\mathrm{d}}$ | References | Description | |
|--------------------|---|----------------------------------|--------------------------|----------------------------------|------------------------|---|--|
| Bradac(v1) | 16 | 56 | 2 | 14 | HFF ^e ; ? | SWUnited ^f : Free-form, strong+weak-lensing based model. Errors from bootstrap resampling only weak-lensing constraints. | |
| CATS(v1) | 17 | 60 | 2 | 14 | HFF; ? | CATSg team implementation of LENSTOOL ^h parametric strong-lensing based model. | |
| CATS(v1.1) | 17 | 60 | 2 | 14 | ? | CATS team LENSTOOL parametric model with both strong and weak lensing constraints. | |
| Merten | 16 | 56 | 2 | 14 | HFF; ? | Sawlens, Grid-based free-form strong+weak lensing based model using adaptive mesh refinement. | |
| Sharon(v1) | 17 | 60 | 2 | 14 | HFF; | LENSTOOL parametric, strong-lensing based model | |
| Sharon(v2) | 15 | 47 | 3 | 11 | ? | LENSTOOL parametric, strong-lensing based model. Includes cosmological parameter varia- tions in uncertainty estimates. | |
| Zitrin-LTM | 10 | 44 | 2 | 0 | HFF; ? | Parametric strong-lensing model, adopts the Light-Traces-Mass assumption for both the lumi- | |
| Zitrin-NFW | 10 | 44 | 2 | 0 | HFF; ? | nous and dark matter. Parametric strong-lensing model using PIEMD profiles for galaxies and NFW profiles for dark matter halos. | |
| Williams | 10 | 40 | 2 | 8 | HFF | GRALE ^j : Free-form strong-lensing model using a genetic algorithm. | |
| | Post-HFF models : include data from the HFF program | | | | | | |
| CATS(v2) | 50 | 151 | 4 | 1 | ? | Updated version of the CATSv1 model, adds 33 new multiply-imaged galaxies, for a total of 159 individual lensed images. | |
| Diego ^k | 15 | 48 | 4 | 11 | ? | WSLAP+ ¹ : Free-form strong-lensing model using a grid-based method, supplemented by deflections fixed to cluster member galaxies. | |
| GLAFIC | 24 | 67 | 3 | 12 | ? | Parametric strong-lensing model using v1.0 of the GLAFIC code. ^m | |
| Lam(v1) | 21 | 65 | 4 | 17 | ? | Alternative implementation of the WSLAP+ model, using a different set of strong-lensing constraints and redshifts. | |
| | | | Unblind | models: | generated after the SN | magnification was known | |
| Bradac(v2) | 25 | 72 | 7 | 18 | ? | Updated version of the SWUnited model with new strong-lensing constraints from HFF imaging and GLASS spectra. Errors from bootstrap resam- | |
| Lam(v2) | 10 | 32 | 5 | 5 | ? | pling only strong-lensing constraints. Updated version of the WSLAP+ model, using more selective strong-lensing constraints and | |
| CATS(v2.1) | 55 | 154 | 8 | 1 | ? | GALFIT ⁿ models for galaxy mass. Updated version of the CATSv2 model, adopting the spec-z constraints used for the Bradac(v2) | |
| CATS(v2.2) | 25 | 72 | 8 | 1 | ? | model. Updated version of the CATSv2 model, adopting the spec-z constraints and multiple-image definitions used for the Bradac(v2) model. | |

^a Number of multiply imaged systems used as strong-lensing constraints.

^b Total number of multiple images used.

^c Number of multiply imaged systems with spectroscopic redshifts.

d Number of multiply imaged systems with photometric redshifts.

e Lens models with the reference code "HFF" were produced as part of the Hubble Frontier Fields lens modeling program, using arcs identified in HST archival imaging from ?, spectroscopic redshifts from ?, and ground-based imaging from ?????. Details on the model construction and an interactive model magnification web interface are available at http://archive.stsci.edu/prepds/ frontier/lensmodels/

f Swunited: Strong and Weak lensing United; ?
g CATS: Clusters As TelescopeS lens modeling team. PI's: J.-P. Kneib & P. Natarajan

h LENSTOOL: ?; http://projects.lam.fr/repos/lenstool/wiki

ⁱ SaWLENS: ?; Strong and Weak LENSing analysis code. http://www.julianmerten.net/codes.html

j GRALE: GRAvitational LEnsing; ???

k Abell 2744 model available at http://www.ifca.unican.es/users/jdiego/LensExplorer. No uncertainty estimates were available for the Diego implementation of the WSLAP+ model, so we adopt the uncertainties from the closely related Lam model.

¹ WSLAP+: ?; Weak and Strong Lensing Analysis Package plus member galaxies (Note: no weak-lensing constraints used for Abell

 $^{^{\}rm 2744})$ $^{\rm m}$ GLAFIC: ?; http://www.slac.stanford.edu/~oguri/glafic/

ⁿ GALFIT: Two-dimensional galaxy fitting algorithm (?)

| Table 5 |
|--|
| Lens model predictions for SN HFF14Tom |
| magnification |

| Modela | $\mathrm{Best^b}$ | Median ^c | 68% Conf. Range ^d |
|--------------------|-------------------|---------------------|------------------------------|
| Bradac(v1) | 3.19 | 2.48 | 2.31 - 2.66 |
| CATS(v1) | 2.28 | 2.29 | 2.25 - 2.34 |
| CATS(v1sw) | | 2.62 | 2.44 - 2.80 |
| Merten | 2.33 | 2.24 | 2.04 - 2.92 |
| Sharon(v1) | 2.56 | 2.60 | 2.44 - 2.78 |
| Sharon(v2) | 2.74 | 2.59 | 2.42 - 2.85 |
| Zitrin-LTM | 2.67 | 2.99 | 2.61 - 3.77 |
| Zitrin-NFW | 2.09 | 2.29 | 2.07 - 2.52 |
| Williams | 2.70 | 2.81 | 1.65 - 5.54 |
| CATS(v2) | | 3.42 | 3.27 - 3.58 |
| Diego | | 1.80 | 1.44 - 2.16 |
| $GL \breve{A} FIC$ | 2.34 | 2.29 | 2.19 - 2.37 |
| Lam(v1) | | 2.79 | 2.42 - 3.16 |
| Bradac(v2)* | 2.23 | 2.26 | 2.30 - 2.23 |
| Lam(v2)* | 1.86 | 1.91 | 1.54 - 2.28 |
| $CATS(v2.1)^*$ | | 3.06 | 2.92 - 3.19 |
| $CATS(v2.2)^*$ | | 3.07 | 2.94 - 3.20 |
| | | | |


^a Models above the line are from the pre-HFF set, and those below incorporate HFF data. The final two, marked by an asterisk, were not part of the blind test as they include modifications made after the measured magnification of the SN was known.

and thus tend to result in larger error bars. In brief, if the free form methods are too flexible, then they will result in overestimated error bars. Conversely, if the simply parameterized models are too inflexible, they will result in underestimated error bars.

In Figure 6 the model predictions are plotted along-side the observed magnification of SN HFF14Tom, derived in Section 6. To first order, this comparison shows that these 17 models are largely consistent with each other and with the observed magnification of SN HFF14Tom. The "naive mean" of the full set of models is $\mu_{\rm pre}=2.6\pm0.4$. This is an unweighted mean (i.e. we ignore all quoted uncertainties) derived by naively treating each as an independent prediction for the magnification (this is clearly incorrect, as several models are represented multiple times as different versions). The naive mean is separated from the observed SN magnification by $\delta\mu/\mu=28\%$, which is approximately a 1.5σ difference.

This is approximately consistent with the results of ? and ?, where model predictions were found to be in reasonably good agreement with a set of 3 lensed SNe from the CLASH program. The general agreement between the model predictions and the SN measurement is especially encouraging for these Abell 2744 models. This is a merging cluster with a complex mass distribution, and the SN is located outside of the strong-lensing region where the models are most tightly constrained.

However, beyond this first-order agreement, there is a small systematic bias apparent. All but two of the lens models return median magnifications that are *higher* than the observed value, and six of the models are dis-



Comparison of the observed lensing magnification to predictions from lens models. The vertical blue line shows the constraints from SN HFF14Tom derived in Section 6 using the MLCS2k2 fitter, with a shaded region marking the total uncertainty. Markers with horizontal error bars show the median magnification and 68% confidence region from each of the 17 lensing models. Circles indicate models that use only strong-lensing constraints, while diamonds denote those that also incorporate weaklensing measurements. Models using a "free-form" approach are shown as open markers, while those in the "parametric" family are given filled markers. The top half, with points in black, shows the nine models that were constructed using only data available before the start of the Frontier Fields observations. A black dashed line marks the unweighted mean for these models, at $\mu = 2.6$. The lower six models in green used additional input constraints, including new multiply imaged systems and redshifts. The final two points, with square orange outlines, are the "unblind" models that were generated after the magnification of the SN was known. The green dashed line marks the unweighted mean for these six models, at $\mu = 2.3$.

crepant by more than 1.5σ . These six discrepant models are all biased to higher magnifications. They are found in both the pre-HFF and post-HFF models, in the parametric and free-form families, and among the strong-lensing-only and the strong+weak subsets. It is important to emphasize that SN HFF14Tom only samples a single line of sight through the cluster, and this bias to higher magnifications is minor. Nevertheless, a systematic shift of this nature is surprising, given the wide range of modeling strategies, input data, and physical assumptions represented by this set of models. In the following sub-

^b The magnification returned for the optimal version of each model, as independently defined by each lens modeling team.

 $^{^{\}rm c}$ Median magnification from 100-600 Monte Carlo realizations of the model.

^d Confidence ranges about the median, enclosing 68% of the realized values.

sections we examine possible explanations for this small but nearly universal bias. We first consider whether a misinterpretation of the data on the SN itself can account for the observed systematic bias, and then examine the lens models.

7.1. Possible Errors in Supernova Analysis 7.1.1. Redshift Error

If the redshift of the SN derived in Section 4 were incorrect, then one would derive a different value for the magnification, both from the SN measurement and the lens model predictions. Conceivably, this could resolve the tension between the measurement and the models. It is often the case in SN surveys that redshifts are assigned based on a host galaxy association, typically inferred from the projected separation between the SN and nearby galaxies. In this case the redshift is strongly supported by evidence from the SN itself: we find a consistent redshift from both the SN spectrum (Section 4.2) and the light curve Section 5, which are both within 1σ of the spectroscopic redshift for the nearest detected galaxy: z = 1.3457. This appears to be a solid and self-consistent picture, so the evidence strongly disfavors any redshift that is significantly different from z = 1.35.

We have adopted the most precise redshift of z=1.3457 from the host galaxy as our baseline for the magnification comparison. If instead we adopt the spectroscopic redshift from the SN itself (z=1.31; Section 4.2) then we find no significant change in the inferred magnifications or in the suggestion of a small systematic bias.

7.1.2. Foreground Dust and SN Color

All SN sight-lines must intersect some amount of foreground dust from the immediate circumstellar environment, the host galaxy, and the intergalactic medium (IGM). In the case of SN HFF14Tom one might posit some dust extinction from the intra-cluster medium (ICM) of Abell 2744, although measurements of rich clusters suggest that the ICM has only a negligible dust content (???). When fitting the HFF14Tom light curve we account for dust by including corrections that modify the inferred luminosity distance based on the SN color. If after applying these dust corrections we were still underestimating the effect of dust along this sight-line, then the SN would appear dimmer than it really is, the inferred distance modulus would be higher, and the measured magnification would be biased to an artifically low value. Thus, an underestimation of dust would be in the right direction to match discrepancy we observe.

In Section 6.1 we found that SN HFF14Tom is on the blue end of the normal range of Type Ia SN colors. With the SALT2 fitter we measured a color parameter $c=-0.127\pm0.025$, and with MLCS2k2 we found the host galaxy dust extinction to be $A_{\rm V}=0.011\pm0.025$ magnitudes. These colors are tightly constrained, as we are fitting to photometry that covers a rest-frame wavelength range from $\sim 3500-7000 \rm \mathring{A}$ and extends to ~ 30 days past maximum brightness. This leaves little room for the luminosity measurement to be biased by dust, as the dimming of HFF14Tom would also necessarily be accompanied by some degree of reddening.

Nevertheless, one might suppose that a bias could be introduced if we have adopted incorrect values for the color correction parameter β or extinction law R_V in the SALT2 and MLCS2k2 fits, respectively. The appropriate value to use for this color correction and how it affects inferences about the intrinsic scatter in Type Ia SN luminosities is a complex question that is beyond the scope of this work (see e.g. ????). However, we can already rule this out as a solution for the magnification discrepancy. Our error on the HFF14Tom distance modulus already includes an uncertainty in the extinction law and a related error to account for the intrinsic luminosity scatter. These are well vetted parameters, based on observations of ~ 500 SNe extending to $z \sim 1.5$ (?). Furthermore, there is no reason to propose that SN HFF14Tom is uniquely affected by a peculiar type of dust. Thus, any change in the color correction applied to SN HFF14Tom would require the same adjustment to be applied to the unlensed SNe at similar redshift that make up our comparison sample, largely negating the effect on the inferred magnification.

The traditional color corrections as formalized in SN light curve fitters are designed to account for a dust component that lies in the rest frame of the SN. The inferred luminosity of a SN can also be affected by the presence of foreground dust with a different redshift and possibly a different reddening law (?). However, the magnitude of such a bias is insufficient to account for the observed discrepancy, ? estimate the opacity of the universe as $\langle A_V \rangle \sim 0.03$ mag up to z=0.5. While this can have a measurable impact on precise cosmological constraints, it is far less than the 0.23 mag difference between the observed magnification of HFF14Tom and the mean of the model predictions.

Although the very blue color of SN HFF14Tom is helpful to rule out dust as a possible explanation for the observed small systematic bias, it is possible that this very blue color is itself leading to a bias in the distance measurement. ? measured a small bias in the SALT2 fitter for Type Ia SNe that have a color parameter derived from the light curve fit c < -0.1 (see, e.g., their Figure 10 and Section 5.1). For HFF14Tom we have measured c = -0.13, which would correspond to a bias of roughly +0.05 mag in the SALT2 distance measurement. If we were to apply a -0.05 mag correction to the distance modulus from the SALT2 fit, then this would increase the inferred magnification to $\mu_{\text{SALT2}} = 2.08 \pm 0.36$. This would slightly reduce the tension between the SALT2 measurement and the model predictions, though it would not be enough to completely alleviate it.

$7.1.3.\ Misclassification$

Is it possible that HFF14Tom is an example of a peculiar stellar explosion that does not follow the relationship between light curve shape and luminosity observed in normal Type Ia SNe? In Sections 4.1 and 5 we classified SN HFF14Tom as a normal Type Ia SN based on both the spectroscopic and photometric evidence. This rules out the possibility that HFF14Tom belongs to a different class of normal SN explosions (Type Ib, Ic, or II). In Section 6.1 we fit the HFF14Tom light curve to determine a luminosity distance and found that the light curve shape and color are consistent with a Type Ia SN of average light curve width, with minimal dust extinction. This excludes the possibility that HFF14Tom is one of the sub-class of faint and fast-declining Type Ia SNe like




Figure 7. Comparing HFF14Tom observations to the color curves of SN 2006bt and a normal Type Ia SN model. Grey points show the observed colors of SN HFF14Tom, k-corrected to restframe UBVr bands. Solid orange lines show the observed color curves for the peculiar Type Ia SN 2006bt. Dashed blue lines show the color curves for a normal Type Ia SN, derived from the SALT2 model with a color parameter c=-0.13, set to match the best-fit color for SN HFF14Tom.

SN 1991bg.

One remaining possibility is that HFF14Tom could be part of a rare sub-category of peculiar Type Ia SNe that masquerade as their normal cousins, epitomized by the prototype SN 2006bt (?). These objects appear to have a normal Type Ia light curve shape, except for the absence of a secondary maximum or "shoulder" in near-IR bands. The reddest filter available for the HFF14Tom light curve is F160W, which has an effective rest-frame wavelength of 6623Å at z=1.31, making it close to the rest-frame r band. Although the near-IR shoulder is more prominent in the rest-frame i band, we can see in Figure 4 that the F160W light curve may suggest a weak or delayed near-IR shoulder. The second-to-last observation in F160W is $\sim 1\sigma$ lower than predicted by the best-fit SALT2 and MLCS2k2 models. This is tenuous evidence, but it would be consistent with a 2006bt-like light curve.

In this case, however, the color of SN HFF14Tom can rule out a classification as a 2006bt-like object. All members of this peculiar sub-class exhibit very red colors across all optical and near-IR bands, consistent with a very cool photosphere (?). Figure 7 shows that HFF14Tom is bluer than the SN 2006bt prototype by at least 0.25 magnitudes in every color. The observed HFF14Tom colors are fully consistent with a normal, blue Type Ia SN (the SALT2 model is shown).

7.2. Possible Errors in Lens Modeling

Having found no evidence for misinterpetations of the SN data, we are left to seek an explanation for the mild tension by scrutinizing the lens models. Here we evaluate four possible ways in which the input data or assump-

tions of these lens models could lead to a systematic bias in the magnification.

7.2.1. Cosmic Weak Lensing

In addition to the dominant gravitational lensing from the foreground cluster, a distant source like SN HFF14Tom will also be subject to the cumulative weak lensing effects due to uncorrelated large scale structure along the line of sight, sometimes called "Cosmic Weak Lensing" (CWL; ???????). This CWL effect is most important for very high redshift sources (z > 5), which have a much longer path length over which to encounter large scale structure and for highly magnified sources very near to the cluster lensing critical curve (because CWL can perturb the position of the critical curve). For SN HFF14Tom at the modest redshift of z = 1.35 and far from the critical curve, CWL should increase the scatter in the lensing magnification at a level much less than 10% (?). The CWL effect is therefore unable to account for the observed discrepancy of $\sim 23\%$ by itself.

In addition, some sources of uncertainty are taken into account only by a subset of models. For example, the Willams model (using GRALE; Liesenborgs et al. 2006, 2007; Mohammed et al. 2014) accounts for mass sheet degeneracy by introducing a free parameter in the reconstruction that represents an arbitrarily scaled mass sheet. The flexibility of the Williams model and the inclusion of this source of uncertainty contribute to its relatively large uncertainties compared to the other models.

7.2.2. Nearby Cluster Member

A component of the Abell 2744 lens that is particularly relevant to SN HFF14Tom is the cluster member galaxy that lies just 5".8 north of the SN position (see Figure 1). If the mass-to-light ratio or the profile of the dark matter halo for this galaxy were significantly different from other cluster member galaxies, then its proximity to the SN sight-line might drive a bias in the magnification. We tested this hypothesis using the CATS(v2) model by allowing the mass of the nearby cluster member galaxy to vary as a free parameter in the model. We found that the change in the SN HFF14Tom magnification prediction was less than $\Delta \mu = 0.1$. This additional dispersion is already included in the uncertainties quoted for that model in Table 5 and Figure 6. Furthermore, an erroneous M/L value for the nearby cluster member galaxy could not explain the systematic shift of all lens models, because some do not incorporate cluster member galaxies into their constraints at all (the free-form Bradac, Williams and Merten models).

7.2.3. Cosmological Parameter Uncertainty

Most lens models make a fixed assumption for the values of cosmological parameters in a standard Λ CDM cosmology (typically: $\Omega_{\rm m}=0.3,\Omega_{\Lambda}=0.7,H_0=70$ km s⁻¹ Mpc⁻¹). This can introduce a systematic magnification error that is comparable in magnitude to that of the statistical uncertainties (??). Incorporating this cosmological parameter uncertainty would increase the model magnification errors, and therefore reduce the tension between models and observation, but would not resolve the overall systematic shift. ? find that for Abell 2744

the additional cosmological uncertainty for magnifications $\mu \approx 2$ is < 10% which would still leave 6 models discrepant by at least 1.5 σ .

7.2.4. Misidentification of a multiple image

? proposed a correction for the location of image 3.3, the third component of a triply-imaged galaxy (?) at $z=3.98\pm0.02$ (?). This possibly specious multiple image is only ~10" from the SN HFF14Tom position, meaning a change in its location could have a large impact on the predicted magnification for SN HFF14Tom (see ?, for a quantitative discussion of this effect). However, ? also evaluated a version of the CATS(v2) model in which this single source is left in the original position and found that the location of image 3.3 does not significantly affect the predicted magnifications. Furthermore, dividing our 17 models into those using the original location vs those that adopt the new position, we find that both groups include models that are within 1σ of the observed μ as well as significantly discrepant points.

7.2.5. Source Plane Minimization

A general tendency towards high model magnification can result when lens models are minimized in the source plane (rather than the image plane). For a given model, modifying the surface mass density to have a shallower profile (declining less quickly with radius) will lead to an increase in the predicted magnifications across the field. As the magnification applies to both flux and area, the shallower profile also leads to a smaller source plane area, which necessarily brings delensed images of the same source closer together in the source plane. Thus, models that are optimized in the source plane may have a tendency toward shallower mass profiles and larger magnifications, which force all sources toward the same location in the source plane.

There are several strategies for mitigating this well known bias. Optimizing a model in the image plane can avoid this bias entirely (??). The Sharon(v2), Zitrin-LTM, Zitrin-NFW, and CATS(v2-2.2) models all use image plane minimization. Alternatively, one may guard against this bias by using a modified source plane minimization that is constrained not to generate overly small de-lensed sources (??). In our test set, this second strategy is employed by the Diego model. Finally, the effect of this bias should be reduced when one uses many strong lensing constraints over a wide range of redshifts. The CATS(v2) and CATS(v2.1) models have the largest number of strong-lensing constraints, using 50 and 55 multiply-imaged systems, respectively.

Although many of the tested models utilize these strategies to avoid a source plane minimization bias, Table 5 and Figure 6 show that these models do not uniformly deliver more accurate magnification predictions. The most direct test of this is in the Sharon and CATS model series. In both cases the v1 model used the faster source plane minimization approach, but the v2 model was constructed with the more costly image plane minimization. Neither of those models show a significant improvement in model accuracy from v1 to v2. This provides a strong indication that the observed tension between models and the SN observation cannot be entirely attributed to a bias arising from source plane minimization.

7.3. Quantity and Quality of Strong Lensing Constraints

None of the possible errors in lens model inputs described above can completely account for the mild tension between the observed and predicted magnifications. However, we do have several models that agree with the measured SN magnification. In this section we take a simple pragmatic approach and seek to identify any shared characteristics of the models that are most accurate for this particular sight-line. In this way SN HFF14Tom may serve as a guide toward optimizing future lens models for magnification predictions, particularly in regions where strong-lensing constratins may be scarce.

The addition of deeper imaging and new spectroscopic constraints from the HFF program and affiliated efforts might be expected to improve the accuracy of magnification predictions. However, in Figure 6 we saw no evidence for a universal improvement in model accuracy when moving from the pre- to the post-HFF models. In Figure 8 we attempt to refine this hypothesis by isolating the quantity and quality of strong lensing constraints.

The left panel plots the tension between each model prediction and the measured SN magnification against the number of multiply-imaged galaxies used as stronglensing constraints. The Bradac, Lam and CATSv2 model series are highlighted here with connecting lines, as three examples where the number of systems used has changed substantially from one version to the next, but the model construction has remained fundamentally unchanged. There is no clear correlation here, and in fact for both the Lam and CATSv2 model series, we see the tension actually *increases* for a larger number of stronglensing constraints.

In the right panel of Figure 8 we examine an alternative explanation by plotting the model tension against the fraction of strong-lensing constraints that have a spectroscopic redshift. For the SN HFF14Tom magnification prediction, we see that all three of the testable model series (Bradac,Lam,CATSv2) become *more* accurate as the spec-z fraction increases. ⁴⁶ The spec-z fraction here serves as a crude stand-in for measuring how stringently the strong-lensing constraints are selected in these models.

The SN HFF14Tom comparison therefore supports the notion that the quality of strong-lensing constraints is just as important as the quantity – at least for computing magnifications of sources outside the strong-lensing region. Once a sufficient number of robust multiply-imaged systems have been included (perhaps ~ 10) most models will be dominated by systematic uncertainties such as those described in Section 7.2. To improve the accuracy of magnification predictions with a large number of strong-lensing constraints, one—should employ a strict vetting of the multiply-imaged systems. Each system used should ideally have a spectroscopic redshift, or at least a very well constrained photometric redshift. The primary value in this strict vetting is to avoid erroneous redshifts being assigned to multiply-imaged systems that

⁴⁶ Of course this apparent correlation does not imply any direct causation. Consider the limit of just a single multiply-imaged galaxy constraint: the spec-z fraction could be 1.0 but the model would be wildly unconstrained.




Figure 8. The model tension as a function of strong-lensing constraints. The y axis in both panels marks the number of standard deviations in discrepancy between the measured SN magnification and the prediction from each lens model. The left panel plots this tension against the total number of multiply imaged systems used, and the right panel plots the fraction of multiply-imaged systems that have a spectroscopic redshift constraint. Marker shapes and colors are the same as in Figure 6. In both panels, three lens model sequences are highlighted with connecting lines: the Bradac, Lam, and CATS(v2) models, which are the three examples where the number of strong lensing constraints and the fraction with spectroscopic redshifts changed substantially from one model version to the next.

are unconstrained in redshift space. Such redshift errors are especially problematic for relatively low-z background systems: at lower redshifts the model-derived deflection angle and magnification show a more rapid variation with the redshift of the background galaxy than for similarly positioned high redshift galaxies. This conclusion should be unsurprising, as for example, ? have shown that cosmological uncertainties are smaller for models that use a large fraction of strong-lensing constraints with spectroscopic redshifts, and ? have shown a case study where adding spectroscopic redshift information changes the inferred magnification by $\sim 10\%$.

It is relevant to note that the three model series highlighted here (Bradac, CATSv2 and Lam) also happen to be the three models that are not truly "blind," in that the modelers were aware of preliminary analysis of the SN magnification when constructing the latest version. However, none of these models used the SN magnification as an input constraint, and none were consciously tuned to match the SN magnification. In fact, for the Bradac(v2) and Lam(v2) cases, the modelers did not have access to a final measurement of the SN magnification at the time their models were generated.

8. SUMMARY AND CONCLUSIONS

The appearance of a Type Ia SN behind a massive galaxy cluster provides a rare opportunity to use a standard candle for a direct measurement of the absolute magnification due to gravitational lensing. The discovery of SN HFF14Tom in the HFF imaging of Abell 2744 offers the first chance to apply this test on a cluster with multiple publicly-available lens models. We have found that the spectrum and light curve of SN HFF14Tom are well matched by templates of a normal Type Ia SN at z=1.3457. Using the two most prevalent SN Ia light

curve fitters, SALT2 and MLCS2k2, we get a consistent measurement of the distance modulus (Table 3). Using a cosmology-independent comparison against a sample of unlensed SNe Ia at similar redshifts, we find that SN HFF14Tom is ~ 0.7 magnitudes brighter than the field sample would predict. Attributing this difference to the gravitational lensing magnification (and accounting for the intrinsic scatter in luminosity of the SN Ia population), we have derived a consistent measured magnification of $\mu_{\rm SALT2}=1.99\pm0.38.~\mu_{\rm MLCS2k2}=2.03\pm0.29,$ from the two light curve fitters.

Taking advantage of the availability of 17 well-constrained lens models for the Abell 2744 cluster, we have used SN HFF14Tom to ask how accurately these lens models can predict the magnification along this line of sight. We find that these models are consistent, and fairly accurate, collectively predicting $\mu=2.5\pm0.4$, within 1σ of the measured value. This is encouraging, and reinforces the quality and value of these public lens models for studying magnified background objects. However, we note that all but two of the tested models provide a median magnification prediction that is larger than the measurement from the SN, and all the models with significant discrepancy are too high. This may be an indication that there is a small systematic bias at work, at least in some of the models.

We have speculated on what could be the origin for such a systematic bias, first considering and rejecting three possible explanations that presume an error in the interpretation of the available SN data.

1. Redshift: We reject the possibility that a redshift error is the primary cause of the discrepancy, as the redshift evidence is well supported by multiple lines of evidence from both the SN and the presumed host galaxy. Changing the redshift within the constraints of these complementary observations does not resolve the tension.

- Dust: We find it implausible that there is sufficient dust in the cluster or elsewhere along the line of sight to account for the magnification discrepancy. A dust-induced bias is also disfavored by the very blue color of SN HFF14Tom.
- 3. Misclassification: The combination of spectroscopic and photometric evidence strongly supports our classification of SN HFF14Tom as a normal Type Ia SN. The most plausible mis-classification would be that the object is a peculiar Type Ia of the SN 2006bt-like sub-class. Although the light curve shape could allow this possibility, the blue color of HFF14Tom can once again reject this alternative.

Turning to the lens models, we have considered 5 ways in which a bias might be introduced into the lens models:

- 1. Cosmic weak lensing
- 2. Peculiar mass for a nearby cluster member
- 3. Cosmological parameter uncertainty
- 4. Misidentification of a multiple image
- 5. Source plane minimization

Although each of these could plausibly introduce a small systematic bias, we find that none are likely to be significant enough (or universal enough) to completely resolve the tension between models and observations.

Finally, we have examined whether there is a simple prescription for the kind of strong lensing constraints that are most likely to deliver an accurate magnification prediction for this sight-line. We find that the number of multiply imaged systems used is not in itself predictive of model accuracy. Rather, it is the fraction of multiply imaged systems that have spectroscopic redshifts that is most correlated with model accuracy. A reasonable interpretation – at least for this cluster and this particular set of models – is that the spectroscopic redshift fraction serves as an effective proxy for the "quality" of the strong lensing constraints. This quality of the input data appears to be a key ingredient for deriving accurate magnifications outside the strong-lensing regime.

We have evaluated here only a single object behind a single cluster, with a minor tension between the observed magnification and the model predictions. This is not in and of itself a cause for alarm. Previous analyses of lensed SN Ia found no significant discrepancy between the observed SN Ia magnifications and the predictions from lens models (?; ?), albeit with a much smaller set of lens models being tested. The observed systematic bias for HFF14Tom is small, and many of the lens models that deviate from the measured magnification are preliminary models that have not been updated to include all of the HFF data. Future revisions of the lens models for Abell 2744 could either incorporate the observed magnification of HFF14Tom as a new model constraint, or can revisit this test to evaluate whether the bias persists.

The general problem of combining predictions from many independent models has a rich history in the astronomical and statistical literature (e.g., ???). In a recent example, ? examined the photometric redshifts from multiple galaxy SED fitting codes using the same input data. They found that the photo-z accuracy and precision can be substantially improved by combining the outputs using a sum of probability distributions, a hierarchical Bayesian method, or even a technique as simple as taking the straight median from all available models. It may be that lensing magnifications are similar, in that they can in general be improved through a simple combination of lens model predictions. Such model averaging will not, however, resolve a systematic bias that is shared by many models.

A promising avenue for exploring the origins of such systematic biases in cluster lens models is through the use of simulated lensing data. One can start with very deep high-resolution multi-band imaging on an unlensed field that has a fairly complete spectroscopic redshift catalog, such as the Hubble Ultra Deep Field. Then a simulated galaxy cluster is placed in the field, and the background galaxies are distorted into arcs and multiple images using a well-defined lensing prescription. The artificially lensed images can then be distributed to lens modeling teams who attempt to reconstruct the (known) mass profile of the simulated cluster. This exercise has recently been pursued with a set of synthetic clusters similar to those observed in the HFF program (Meneghetti et al. in prep), using techniques similar to ??. Preliminary analysis of this simulation comparison suggests that magnifications can be systematically overestimated for sources that lie outside the strong-lensing region along the minor axis direction of a simulated cluster that is very elongated, similar to Abell 2744. This indicates that some lens models significantly underestimate the ellipticity of the mass distribution on large scales for that simulated cluster, which is consistent with a recent lens model comparison using 25 real clusters from the CLASH program (?). Our analysis of SN HFF14Tom here indicates that such simulation efforts and head-to-head lens model comparisons will be an important step for moving toward precision science with cluster-lensed sources.

Increasing the sample of SNe behind clusters like those in the HFF program – with rich lensing constraints and deep imaging – would allow the test described here to be repeated and refined. With 10 or 100 such objects, it would be possible to see whether the SN HFF14Tom μ discrepancy is simply an outlier, or an indication of a more pernicious systematic error. Any cluster that has been vetted by pencil-beam magnification tests using lensed SNIa will be able to provide a more reliable measure of the magnifications for very high redshift objects. Similarly, a broad sample of SNe like HFF14Tom would help to define the preferred lens modeling methodology by highlighting any models that consistently perform well in SN Ia lensing tests. The ongoing FrontierSN program will discover and follow any more highly magnified SNe that appear behind the Frontier Field clusters. Unfortunately, the HFF survey is not designed with highz transient discovery as a primary science goal, so the FrontierSN effort will likely add no more than 1-3 new lensed SNIa. Further imaging of strong-lensing clusters with HST or the James Webb Space Telescope (JWST)

could enable a larger sample to be collected, especially if the filters and cadence are optimized for detection of SN Ia at z>1. Massive clusters such as Abell 2744 will continue to be attractive as cosmic telescopes, allowing the next generation of telescopes to reach the faintest objects in the very early universe. The puzzling bias revealed by SN HFF14Tom supports a concerted effort to improve these lenses with further examination of lensing systematics through simulations and collection of a larger sample of magnified SNe.

Acknowledgments:

This work is dedicated to our colleague and friend Tomas Dahlen, for whom this supernova has been named in memorium. He is dearly missed.

We thank the Hubble Frontier Fields team at STScI for their substantial efforts to make the HFF program successful. In particular, thanks are due to Matt Mountain for the allocation of discretionary orbits for the HFF program; to Jennifer Lotz, Norman Grogin and Patricia Royle for accommodations in strategy and implementation to make the FrontierSN program possible; to Anton Koekemoer for HFF data processing support. We also thank the CLASH team, led by Marc Postman, for observations, catalogs, and high level science products that were of significant value for this analysis. Thanks to Jonatan Selsing for helpful comments on the manuscript.

This work utilizes gravitational lensing models produced by modeling teams that were funded as part of the HST Frontier Fields program conducted by STScI. STScI is operated by the Association of Universities for Research in Astronomy, Inc. under NASA contract NAS 5-26555. The lens models were obtained from the Mikulski Archive for Space Telescopes (MAST).

Financial support for this work was provided to S.A.R. by NASA through grants HST-HF-51312 and HST-GO-13386 from STScI, which is operated by Associated Universities for Research in Astronomy, Inc. (AURA), under NASA contract NAS 5-26555. A.M. acknowledge the financial support of the Brazilian funding agency FAPESP (Post-doc fellowship - process number 2014/11806-9). Support for this research at Rutgers University was provided in part by NSF CAREER award AST-0847157 to SWJ. The Dark Cosmology Centre is supported by the Danish National Research Foundation. J.M.D acknowledges support of the consolider project CSD2010-00064 and AYA2012-39475-C02-01 funded by the Ministerio de Economia y Competitividad. J.M. contributed to this research from the Jet Propulsion Laboratory, California Institute of Technology, under a contract with NASA and acknowledges support from NASA Grants HST-GO-13343.05-A and HST-GO-13386.13-A. The research leading to these results has received funding from the People Programme (Marie Curie Actions) of the European Union's Seventh Framework Programme (FP7/2007-2013) under REA grant agreement number 627288. A.Z. acknowledges financial support from NASA through grant HST-HF-51334.01-A awarded by STScI and operated by AURA. TT acknowledges support by the Packard Foundation in the form of Packard Research Fellowship. GLASS is funded by NASA through HST grant GO-13459. LLRW acknowledges the support of the Minnesota Supercomputing Institute.

REFERENCES

Facilities: HST (WFC3)