


**DATE :- 29-08-2021**

**TIME : 10.30 AM TO 11.50 AM**


1. For the reaction



The equilibrium constant cannot be disturbed by

- Addition of A
- Addition of D
- Increasing of pressure
- Increasing of temperature

**Ans. c**




Pressure has no effect on equilibrium state if  $\Delta n = 0$


2. An organic compound 'X' on treatment with PCC in dichloromethane gives the compound Y. Compound 'Y' reacts with  $I_2$  and alkali to form yellow precipitate of triiodomethane. The compound X is

- $CH_3CHO$
- $CH_3COCH_3$
- $CH_3CH_2OH$
- $CH_3COOH$

**Ans. c**




3. A compound 'A' ( $C_7H_8O$ ) is insoluble in  $NaHCO_3$  solution but dissolve in  $NaOH$  and give a characteristic colour with neutral  $FeCl_3$  solution. When treated with Bromine water compound 'A' forms the compound B with the formula  $C_7H_5OBr_3$ . 'A' is




**Ans. b**


**Sol.**



4. In set of reactions, identify D


$$\text{CH}_3\text{COOH} \xrightarrow{\text{SOCl}_2} \text{A} \xrightarrow[\text{Anh AlCl}_3]{\text{Benzene}} \text{B} \xrightarrow{\text{HCN}} \text{C} \xrightarrow{\text{H}_2\text{O}} \text{D}$$


**Sol.**  $\text{CH}_3 - \text{COOH} \xrightarrow{\text{SOCl}_2} \text{CH}_3 \overset{\text{A}}{\text{C}}\text{OCl} \xrightarrow[\text{anhydrous AlCl}_3]{\text{C}_6\text{H}_6} \text{C}_6\text{H}_5 \overset{\text{B}}{\text{C}}\text{OCH}_3$




a)  $\text{H}_2\text{SO}_3$                       b)  $\text{HNO}_2$   
c)  $\text{CH}_3\text{COOH}$                     d)  $\text{HCN}$

**Sol.** Acidic strength  $\propto K_a$   
The conjugate base of a weakest acid is strongest  $\therefore \text{HCN} (K_a = 4 \times 10^{-10})$  (least value)




**Sol.**  $A \rightarrow CH \equiv CH \xrightarrow[H_2SO_4]{HgSO_4} CH_2 = CH - OH \xrightleftharpoons{\text{Tautomerises}} CH_3CHO$




a)  $\text{LiAlH}_4$  / ether      b)  $\text{H}_2$ , Pt  
c)  $\text{NaBH}_4$                 d) Na and  $\text{C}_2\text{H}_5\text{OH}$

**Sol.**  $\text{CH}_3\text{COOH} \xrightarrow{\text{LiAlH}_4} \text{CH}_3\text{CH}_2\text{OH}$



**Sol.** 
$$\text{CH}_3\text{CHO} \xrightarrow[\text{(ii) H}_3\text{O}^+]{\text{(i) CH}_3\text{MgBr}} \text{CH}_3 - \underset{\text{A}}{\overset{\text{CH}_3}{\text{CH}}} - \text{OH} \xrightarrow[\Delta]{\text{conc. H}_2\text{SO}_4} \text{CH}_3 - \text{CH} = \text{CH}_2$$




$\therefore$  A and C are position isomers.

- addition of oxygen
- addition of electronegative element
- removal of hydrogen
- removal of electronegative element

**Sol.** Conceptual

10. Which is the most suitable reagent for the following conversion ?



- Tollen's reagent
- Benzoyl peroxide
- $\text{I}_2$  and NaOH solution with subsequent acidification
- Sn and NaOH solution


**Ans. c**

**Sol.** Conceptual

11.  $\text{C}_6\text{H}_5\text{CH}_2\text{Cl} \xrightarrow{\text{alc. NH}_3} \text{A} \xrightarrow{2\text{CH}_3\text{Cl}} \text{B}$ . The product B is

- N, N-Dimethyl phenyl methanamine
- N, N-Dimethyl benzenamine
- N-Benzyl-N-methyl methanamine
- phenyl-N-N-dimethyl methanamine

**Ans. a**



12. The method by which aniline cannot be prepared is

- Nitration of benzene followed by reduction with Sn and con. HCl
- Degradation of benzamide with bromine in alkaline solution
- Reduction of nitrobenzene with  $\text{H}_2/\text{Pd}$  in ethanol
- Potassium salt of phthalimide treated with chlorobenzene followed by the hydrolysis with aqueous NaOH solution

**Ans. d**

**Sol.** Conceptual


13. Permanent hardness cannot be removed by

- Using washing soda
- Calgon's method
- Clark's method
- Ion exchange method

**Ans. c**

**Sol.** Conceptual

14. A hydrocarbon A ( $\text{C}_4\text{H}_8$ ) on reaction with HCl gives a compound B ( $\text{C}_4\text{H}_9\text{Cl}$ ) which on reaction with 1 mol of  $\text{NH}_3$  gives compound C ( $\text{C}_4\text{H}_{10}\text{N}$ ). On reacting with  $\text{NaNO}_2$  and HCl followed by treatment with water, compound C yields an optically active compound D. The is



**Ans. b**

**Sol.** Optically active compound is option B.

15. RNA and DNA are chiral molecules, their chirality is due to the presence of

- D-sugar component
- L-sugar component
- Chiral bases
- Chiral phosphate ester unit

**Ans. a**

**Sol.** Conceptual

16. The property of the alkaline earth metals that increases with their atomic number is

- Ionisation enthalpy
- Electronegativity
- Solubility of their hydroxide in water
- Solubility of their sulphate in water

**Ans. c**

**Sol.** Conceptual

17. Primary structure in a nucleic acid contains bases as GATGC ... The chain which is complementary to this chain is

- G G T G A ...
- T G A A G ...
- C T A C G ...
- T T T A G ...

**Ans. c**

**Sol.** Conceptual

18. In the detection of II group acid radical, the salt containing chloride is treated with concentrated sulphuric acid, the colourless gas is liberated. The name of the gas is  
 a) Hydrogen chloride gas  
 b) Chlorine gas  
 c) Sulphur dioxide gas  
 d) Hydrogen gas

**Ans. a**

**Sol.** Conceptual

19. The number of six membered and five membered rings in Buckminster Fullerene respectively is  
 a) 20, 12    b) 12, 20    c) 14, 18    d) 14, 11

**Ans. a**

**Sol.** Conceptual

20. In chrysoberyl, a compound containing Beryllium, Aluminium and oxygen, oxide ions form cubic close packed structure. Aluminium ions occupy  $\frac{1}{4}$ th of octahedral voids. The formula of the compound is  
 a)  $\text{BeAlO}_4$     b)  $\text{BeAl}_2\text{O}_4$     c)  $\text{Be}_2\text{AlO}_2$     d)  $\text{BeAlO}_2$

**Ans. b**

Be   Al   O   N = No. of oxide ions involved in CCP

**Sol.**  $\frac{N}{4} : \frac{2N}{4} : N \therefore \text{octahedral voids} = N$

1 : 2 : 4 Tetrahedral voids = 2N

21. The correct statement regarding defects in solid is  
 a) Frenkel defect is a vacancy defect  
 b) Schottky defect is a dislocation defect  
 c) Trapping of an electron in the lattice leads to the formation of F-centre  
 d) Schottky defect has no effect on density

**Ans. c**

**Sol.** Frenkel defect – dislocation defect  
 Schottky defect – decreases density  
 F-centre – Trapping of on electrons in the lattices

22. A metal crystallises in BCC lattice with unit cell edge length of 300 pm and density  $6.15 \text{ g cm}^{-3}$ . The molar mass of the metal is  
 a)  $50 \text{ g mol}^{-1}$     b)  $60 \text{ g mol}^{-1}$   
 c)  $40 \text{ g mol}^{-1}$     d)  $70 \text{ g mol}^{-1}$

**Ans. a**

**Sol.**  $d = \frac{ZM}{a^3 N_A}$

$$M = \frac{d a^3 N_A}{Z} = \frac{6.15 \times (300 \times 10^{-10})^3 \times 6 \times 10^{23}}{2} \cong 50 \text{ g mol}^{-1}$$

23. Henry's law constant for the solubility of  $\text{N}_2$  gas in water at 298K is  $1.0 \times 10^5 \text{ atm}$ . The mole fraction of  $\text{N}_2$  in air is 0.8 The number of moles of  $\text{N}_2$  from air dissolved in 10 moles of water at 298K and 5 atm pressure is  
 a)  $4.0 \times 10^{-4}$     b)  $4.0 \times 10^{-5}$   
 c)  $5.0 \times 10^{-4}$     d)  $4.0 \times 10^{-6}$

**Ans. a**

**Sol.**  $P_{\text{N}_2} = X_{\text{N}_2} \cdot P_{\text{total}}$   
 $= 0.8 \times 5 = 4 \text{ atm}$

$$P_{\text{N}_2} = K_H \cdot X_{\text{N}_2}$$

$$4 = 10^5 \cdot X_{\text{N}_2}$$

$$X_{\text{N}_2} = 4 \times 10^{-5}$$

$$X_{\text{N}_2} = \frac{n_{\text{N}_2}}{n_{\text{N}_2} + n_{\text{H}_2\text{O}}} \quad (n_{\text{N}_2} \lll n_{\text{H}_2\text{O}})$$

$$4 \times 10^{-5} = \frac{n_{\text{N}_2}}{10}$$

$$n_{\text{N}_2} = 4 \times 10^{-4}$$

24. A pure compound contains 2.4g of C,  $1.2 \times 10^{23}$  atoms of H, 0.2 moles of oxygen atoms. Its empirical formula is  
 a)  $\text{C}_2\text{HO}$     b)  $\text{C}_2\text{H}_2\text{O}_2$     c)  $\text{CH}_2\text{O}$     d)  $\text{CHO}$

**Ans. d**

**Sol.**  $2.4 \text{ g C} = \frac{2.4}{12} = 0.2 \text{ mol}$

$$1.2 \times 10^{23} \text{ atoms of H} = \frac{1.2 \times 10^{23}}{6 \times 10^{23}} = 0.2 \text{ mol}$$

0.2 mole of 'O' atoms

$\therefore$  simplest ratio = C : H : O

$$0.2 : 0.2 : 0.2$$

$$= \text{CHO}$$

25. Choose the correct statement
- $K_H$  value is same for a gas in any solution
  - Higher the  $K_H$  value more the solubility of gas
  - $K_H$  value increases on increasing the temperature of the solution
  - Easily liquefiable gases usually has lesser  $K_H$  values

**Ans. c**

**Sol.**  $K_H$  value changes with solvent nature  
Higher the  $K_H$  less is solubility  
 $K_H$  value increase with increase of 'T'  
Easily liquefied gases have high  $K_H$  value

26. The  $K_H$  value (K bar) of Argon (I), Carbondioxide (II) formoldehyde (III) and methane (IV) are respectively 40.3, 167,  $1.83 \times 10^{-5}$  and 0.413 at 298 K. The increasing order of solubility of gas in liquid is
- $I < II < IV < III$
  - $III < IV < II < I$
  - $I < III < II < IV$
  - $I < IV < II < III$

**Ans. a**


**Sol.**  $P_H = K_H \times$   
$$K_H \propto \frac{1}{X(\text{solubility})}$$
  
 $\therefore$  more is the  $K_H$  less is the solubility

27. The vapour pressure of pure liquids A and B are 450 and 700 mm of Hg at 350 K respectively. If the total vapour pressure of the mixture is 600 mm of Hg, the composition of the mixture in the solution is
- $x_A = 0.4, x_B = 0.6$
  - $x_A = 0.6, x_B = 0.4$
  - $x_A = 0.3, x_B = 0.7$
  - $x_A = 0.7, x_B = 0.3$

**Ans. a**

**Sol.**  $P_{\text{total}} = P_A^0 X_A + P_B^0 X_B$   
$$= P_A^0 X_A + P_B^0 (1 - X_A)$$
  
$$600 = 450 X_A + 700 (1 - X_A)$$
  
$$X_A = 0.4$$
  
$$X_B = 1 - 0.4 = 0.6$$

28. Consider the following electrodes



$E^\circ \text{Zn} / \text{Zn}^{2+} = -0.76\text{V}$  electrode potentials of the above electrodes in volts are in the order

- $P > S > R > Q$
- $S > R > Q > P$
- $Q > R > S > P$
- $P > Q > R > S$

**Ans. c**

**Sol.**  $\text{Zn}^{2+}_{(\text{aq})} + 2e^- \rightarrow \text{Zn}_{(\text{s})}$

$$E_{\text{red}} = E_{\text{red}}^\ominus - \frac{0.059}{n} \log \frac{1}{[\text{Zn}^{2+}]}$$

$$E_{\text{red}} = -0.76 + \frac{0.059}{2} \log [\text{Zn}^{2+}]$$

$$\text{as } [\text{Zn}^{2+}] \uparrow E_{\text{red}} \uparrow$$

29. The number of angular and radial nodes in 3p orbital respectively are

- 3,1
- 1,1
- 2,1
- 2,3

**Ans. b**

**Sol.** No. of angular nodes =  $l = 1$  (3p)  
No. of radial nodes =  $n - l - 1 = 3 - 1 - 1 = 1$

30. The resistance of 0.01 m KCl solution at 298 K is  $1500 \Omega$ . If the conductivity of 0.01 m KCl solution at 298 K is  $0.1466 \times 10^{-3} \text{S cm}^{-1}$ . The cell constant of the conductivity cell in  $\text{cm}^{-1}$  is

- 0.219
- 0.291
- 0.301
- 0.194

**Ans. a**

**Sol.**  $G^* = KR$   
$$= 0.146 \times 10^{-3} \times 1500$$
  
$$= 0.219$$

31.  $\text{H}_{2(\text{g})} + 2\text{AgCl}_{(\text{s})} \rightleftharpoons 2\text{Ag}_{(\text{s})} + 2\text{HCl}_{(\text{aq})}$

$E_{\text{cell}}^\circ$  at  $25^\circ\text{C}$  for the cell is 0.22 V. The equilibrium constant at  $25^\circ\text{C}$  is

- $2.8 \times 10^7$
- $5.2 \times 10^8$
- $2.8 \times 10^5$
- $5.2 \times 10^4$

**Ans. a**

**Sol.**  $\log K_c = \frac{E_{\text{cell}}^\ominus \times n}{0.059} = \frac{0.22 \times 2}{0.059} = 7.45$

$$K_c = \text{Antilog}(7.45) = 2.8 \times 10^7$$

32. For a reaction  $A + 2B \rightarrow \text{Products}$ , when concentration of B alone is increased half life remains the same. If concentration of A alone is doubled, rate remains the same. The unit of rate constant for the reaction is

- a)  $\text{S}^{-1}$                                       b)  $\text{L mol}^{-1} \text{S}^{-1}$   
c)  $\text{mol L}^{-1} \text{S}^{-1}$                               d)  $\text{atm}^{-1}$

**Ans. a**

**Sol.** As  $[B]$  increase,  $t_{\frac{1}{2}}$  remains same

i.e. 1<sup>st</sup> order with respect to 'B'

$$\text{rate} = k[A]^0[B]^1$$

overall order = 1

$\therefore$  units of  $k = \text{S}^{-1}$

33. The third ionisation enthalpy is highest in

- a) Alkali metals  
b) Alkaline earth metals  
c) Chalcogens  
d) Pnictogens

**Ans. b**

**Sol.** Conceptual

34. If the rate constant for a first order reaction is  $k$ , the time( $t$ ) required for the completion of 99% of the reaction is given by

- a)  $t = \frac{4.606}{k}$                                       b)  $t = \frac{2.303}{k}$   
c)  $t = \frac{0.693}{k}$                                       d)  $t = \frac{6.909}{k}$

**Ans. a**

**Sol.** 
$$t = \frac{2.303}{k} \log \frac{[R]_0}{[R]}$$
  
$$= \frac{2.303}{k} \log \left( \frac{100}{1} \right)$$
  
$$= \frac{4.606}{k}$$

35. The rate of a gaseous reaction is given by the expression  $k[A][B]^2$ . If the volume of vessel is reduced to one half of the initial volume, the reaction rate as compared to original rate is


- a)  $\frac{1}{16}$                       b)  $\frac{1}{8}$                       c) 8                      d) 16

**Ans. c**

**Sol.**  $\text{rate} = K[A]^1[B]^2$

$$= K \left[ \frac{n}{v} \right]_A^1 \left[ \frac{n}{v} \right]_B^2$$
  
$$= K \left[ \frac{n}{\frac{v}{2}} \right]_A^1 \left[ \frac{n}{\frac{v}{2}} \right]_B^2$$
  
$$= '8' K \left[ \frac{n}{v} \right]_A^1 \left[ \frac{n}{v} \right]_B^2$$
  
$$'8' \text{ times increases}$$

36. The correct IUPAC name of



- a) 4-Ethyl-1-Fluoro-2-nitrobenzene  
b) 1-Ethyl-4-Fluoro-3-nitrobenzene  
c) 3-Ethyl-6-Fluoronitrobenzene  
d) 5-Ethyl-2-Fluoronitrobenzene

**Ans. a**

**Sol.** Conceptual

37. Higher order (>3) reactions are rare due to
- Shifting of equilibrium towards reactants due to elastic collisions
  - Loss of active species on collision
  - Low probability of simultaneous collision of all reacting species
  - Increase in entropy as more molecules are involved

**Ans. c**

**Sol.** Conceptual

38. Arrange benzene, n-hexane and ethyne in decreasing order of their acidic behaviour

- a) Benzene > n-hexane > ethyne  
b) n-hexane > Benzene > ethyne  
c) ethyne > n-hexane > Benzene  
d) ethyne > Benzene > n-hexane

**Ans. d**

**Sol.** Conceptual

39. A colloidal solution is subjected to an electric field than colloidal particles move towards anode. The amount of electrolytes of  $\text{BaCl}_2$ ,  $\text{AlCl}_3$  and  $\text{NaCl}$  required to coagulate the given colloid is in the order
- $\text{NaCl} > \text{BaCl}_2 > \text{AlCl}_3$
  - $\text{BaCl}_2 < \text{AlCl}_3 > \text{NaCl}$
  - $\text{AlCl}_3 = \text{NaCl} = \text{BaCl}_2$
  - $\text{AlCl}_3 > \text{BaCl}_2 > \text{NaCl}$

**Ans. a**

**Sol.** As ions are moving toward anode i.e. negatively charged colloid

$$\text{Coagulation value} \propto \frac{1}{\text{coagulating power}}$$

$$\therefore \text{Na}^{+1} > \text{Ba}^{+2} > \text{Al}^{+3}$$

40. Which of the following is an incorrect statement?
- Hydrogen bonding is stronger than dispersion forces
  - Sigma bonds are stronger than  $\pi$ -bonds
  - Ionic bonding is non-directional
  - $\sigma$ -electrons are referred to as mobile electrons

**Ans. d**

**Sol.** Conceptual

41. Zeta potential is
- Potential required to bring about coagulation of a colloidal sol.
  - Potential required to give the particle a speed of  $1 \text{ cm s}^{-1}$
  - Potential difference between fixed charged layer and the diffused layer having opposite charges
  - Potential energy of the colloidal particles.

**Ans. c**

**Sol.** Conceptual

42. Which of the following compound on heating gives  $\text{N}_2\text{O}$ ?
- $\text{Pb}(\text{NO}_3)_2$
  - $\text{NH}_4\text{NO}_3$
  - $\text{NH}_4\text{NO}_2$
  - $\text{NaNO}_3$

**Ans. b**

**Sol.** Conceptual

43. Which of the following property is true for the given sequence  
 $\text{NH}_3 > \text{PH}_3 > \text{AsH}_3 > \text{SbH}_3 > \text{BiH}_3$ ?
- Reducing property
  - Thermal stability
  - Bond angle
  - Acidic character

**Ans. b**

**Sol.** Conceptual

44. The correct order of boiling point in the following compounds is
- $\text{HF} > \text{H}_2\text{O} > \text{NH}_3$
  - $\text{H}_2\text{O} > \text{HF} > \text{NH}_3$
  - $\text{NH}_3 > \text{H}_2\text{O} > \text{HF}$
  - $\text{NH}_3 > \text{HF} > \text{H}_2\text{O}$


**Ans. b**

**Sol.** Conceptual

45.  $\text{XeF}_6$  on partial hydrolysis gives a compound X, which has square pyramidal geometry 'X' is
- $\text{XeO}_3$
  - $\text{XeO}_4$
  - $\text{XeOF}_4$
  - $\text{XeO}_2\text{F}_2$


**Ans. c**

**Sol.**  $\text{XeF}_6 + \text{H}_2\text{O} \rightarrow \text{XeOF}_4 + 2\text{HF}$  ( $\text{XeOF}_4$  – square pyramidal)



46. A colourless, neutral, paramagnetic oxide of Nitrogen 'P' on oxidation gives reddish brown gas Q. Q on cooling gives colourless gas R. R on reaction with P gives blue solid S. Identify P, Q, R, S respectively
- $\text{N}_2\text{O}$   $\text{NO}$   $\text{NO}_2$   $\text{N}_2\text{O}_5$
  - $\text{N}_2\text{O}$   $\text{NO}_2$   $\text{N}_2\text{O}_4$   $\text{N}_2\text{O}_3$
  - $\text{NO}$   $\text{NO}_2$   $\text{N}_2\text{O}_4$   $\text{N}_2\text{O}_3$
  - $\text{NO}$   $\text{NONO}_2$   $\text{N}_2\text{O}_4$   $\text{N}_2\text{O}_5$

**Ans. c**



**Sol.**

47. Which of the following does not represent property stated against it?
- $\text{CO}^{+2} < \text{Fe}^{+2} < \text{Mn}^{+2}$  - Ionic size
  - $\text{Ti} < \text{V} < \text{Mn}$  - Number of oxidation states
  - $\text{Cr}^{+2} < \text{Mn}^{+2} < \text{Fe}^{+2}$  - Paramagnetic behaviour
  - $\text{Sc} > \text{Cr} > \text{Fe}$  - Density

**Ans. c**

**Sol.** Conceptual

48. Which one of the following is correct for all elements from Sc to Cu?
- The lowest oxidation state shown by them is +2
  - 4s orbital is completely filled in the ground state
  - 3d orbital is not completely filled in the ground state
  - The ions in +2 oxidation states are paramagnetic

**Ans. d**

**Sol.** Conceptual

49. When the absolute temperature of ideal gas is doubled and pressure is halved, the volume of gas
- will be half of original volume
  - will be 4 times the original volume
  - will be 2 times the original volume
  - will be  $1/4^{\text{th}}$  times the original volume

**Ans. b**

**Sol.**  $PV = nRT$

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2} \quad \frac{P \times V_1}{T} = \frac{P}{2} \times \frac{V_2}{2T} \quad V_2 = 4V_1$$

50. Which of the following pairs has both the ions coloured in aqueous solution? [Atomic numbers of  
[Sc = 21, Ti = 22, Ni = 28, Cu = 29, Mn = 25 ]
- Sc<sup>3+</sup>, Mn<sup>2+</sup>
  - Ni<sup>2+</sup>, Ti<sup>4+</sup>
  - Ti<sup>3+</sup>, Cu<sup>+</sup>
  - Mn<sup>2+</sup>, Ti<sup>3+</sup>

**Ans. d**

**Sol.** Conceptual

51. For the crystal field splitting in octahedral complexes,
- the energy of the  $e_g$  orbitals will decrease by  $(3/5)\Delta_0$  and that of the  $t_{2g}$  will increase by  $(2/5)\Delta_0$
  - the energy of the  $e_g$  orbitals will increase by  $(3/5)\Delta_0$  and that of the  $t_{2g}$  will decrease by  $(2/5)\Delta_0$
  - the energy of the  $e_g$  orbitals will increase by  $(3/5)\Delta_0$  and that of the  $t_{2g}$  will increase by  $(2/5)\Delta_0$
  - the energy of the  $e_g$  orbitals will decrease by  $(3/5)\Delta_0$  and that of the  $t_{2g}$  will decrease by  $(2/5)\Delta_0$

**Ans. b**

**Sol.** Conceptual

52. Peroxide effect is observed with the addition of HBr but not with the addition of HI to unsymmetrical alkene because
- H-I bond is stronger than H-Br and is not cleaved by the free radical
  - H-I bond is weaker than H-Br bond so that iodine free radicals combine to form iodine molecules
  - Bond strength of HI and HBr are same but free radicals are formed in HBr
  - All of these

**Ans. b**

**Sol.** Conceptual

53. The IUPAC name of  $[\text{Co}(\text{NH}_3)_5(\text{CO}_3)]\text{Cl}$  is
- Pentaamminecarbonatocobalt (III) Chloride
  - Carbonatopentamminecobalt (III) Chloride
  - Pentaamminecarbonatocobaltate (III) Chloride
  - Pentaammine cobalt (III) Carbonate Chloride

**Ans. a**

**Sol.** Conceptual

54. Homoleptic complexes among the following are
- A)  $\text{K}_3[\text{Al}(\text{C}_2\text{O}_4)_3]$ , B)  $[\text{CoCl}_2(\text{en})_2]^+$
- C)  $\text{K}_2[\text{Zn}(\text{OH})_4]$
- A only
  - A and B only
  - A and C only
  - C only

**Ans. c**


**Sol.** Conceptual

55. The correct order for wavelengths of light absorbed in the complex ions  $[\text{CoCl}(\text{NH}_3)_5]^{2+}$ ,  $[\text{Co}(\text{NH}_3)_6]^{3+}$  and  $[\text{Co}(\text{CN})_6]^{3-}$  is
- $[\text{CoCl}(\text{NH}_3)_5]^{2+} > [\text{Co}(\text{NH}_3)_6]^{3+} > [\text{Co}(\text{CN})_6]^{3-}$
  - $[\text{Co}(\text{NH}_3)_6]^{3+} > [\text{Co}(\text{CN})_6]^{3-} > [\text{CoCl}(\text{NH}_3)_5]^{2+}$
  - $[\text{Co}(\text{CN})_6]^{3-} > [\text{CoCl}(\text{NH}_3)_5]^{2+} > [\text{Co}(\text{NH}_3)_6]^{3+}$
  - $[\text{Co}(\text{NH}_3)_6]^{3+} > [\text{CoCl}(\text{NH}_3)_5]^{2+} > [\text{Co}(\text{CN})_6]^{3-}$





**Ans. a**

**Sol.** Wave length of light absorbed is inversely proportional to strength of the ligand.





The compound A (major product) is

- a) 
- b) 
- c) 
- d) 

**Ans. b**

**Sol.** Free radical substitution of alkane part. (Benzyl free radical) takes place.

57. Bond enthalpies of  $A_2$ ,  $B_2$  and  $AB$  are in the ratio 2 : 1 : 2. If bond enthalpy of formation of  $AB$  is  $-100 \text{ kJ mol}^{-1}$ . The bond enthalpy of  $B_2$  is


- a)  $100 \text{ kJ mol}^{-1}$       b)  $50 \text{ kJ mol}^{-1}$   
c)  $200 \text{ kJ mol}^{-1}$       d)  $150 \text{ kJ mol}^{-1}$

**Ans. c**

**Sol.** Assume bond strength of  $A_2 = 2X$ , then,

$$B_2 = X, AB = 2X$$





$$\Delta_f H^\ominus = \sum \Delta_{\text{diss}} H^\ominus_{(R)} - \sum \Delta_{\text{diss}} H^\ominus_{(P)}$$



$$-100 = \frac{1}{2} 2X + \frac{1}{2} X - 2$$

$$X = 200$$

58. The order of reactivity of the compounds  $C_6H_5CH_2Br$ ,  $C_6H_5CH(C_6H_5)Br$ ,  $C_6H_5CH(CH_3)Br$  and  $C_6H_5C(CH_3)(C_6H_5)Br$  in  $S_N^2$  reaction is

- a) 
- b) 
- c) 
- d) 

**Ans. a**


**Sol.** Conceptual

59. The major product of the following reaction is  $CH_2 = CH - CH_2 - OH \xrightarrow[\text{Excess}]{HBr}$  Product

- a)  $CH_3 - CHBr - CH_2Br$   
b)  $CH_2 = CH - CH_2Br$   
c)  $CH_3 - CHBr - CH_2 - OH$   
d)  $CH_3 - CHOH - CH_2OH$

**Ans. a**

**Sol.**  $CH_2 = CH - CH_2OH \xrightarrow[\text{Excess}]{HBr} CH_3CH(Br) - CH_2OH \xrightarrow{HBr} CH_3CH(Br) - CH_2Br$



The product 'A' gives white precipitate when treated with bromine water. The product 'B' is treated with Barium hydroxide to give the product C. The compound C is heated strongly to form product D. The product D is

a) 4-Methylpent-3-en-2-one  
b) But-2 enal  
c) 3-Methylpent-3-en-2-one  
d) 2-Methylbut-2-enal

**Ans. a**

**Sol.** A is phenol

B is acetone

Acetone when treated with  $Ba(OH)_2$  Undergoes aldol condensation and followed by heating gives 4-methylpent -3-en-2-one