

# Naive Bayes Classification

Naive Bayes method is a supervised learning algorithm based on Bayes' theorem

It is taking the “naive” assumption of independence between every pair of features.

# Baye's Rule

$$P(b \mid a) = \frac{P(a \mid b)P(b)}{P(a)}$$

# Naïve Bayes Classifier

$$P(y|X) = \frac{P(X|y)P(y)}{P(X)}$$

where,  $y$  is class variable and  $X$  is a dependent feature vector (of size  $n$ ) where:

$$X = (x_1, x_2, x_3, \dots, x_n)$$

Now, if any two events  $A$  and  $B$  are independent, then,

$$P(A, B) = P(A)P(B)$$

Hence, we reach to the result:

$$P(y|x_1, \dots, x_n) = \frac{P(x_1|y)P(x_2|y)\dots P(x_n|y)P(y)}{P(x_1)P(x_2)\dots P(x_n)}$$

which can be expressed as:

$$P(y|x_1, \dots, x_n) = \frac{P(y) \prod_{i=1}^n P(x_i|y)}{P(x_1)P(x_2)\dots P(x_n)}$$

Now, as the denominator remains constant for a given input, we can remove that term:

$$P(y|x_1, \dots, x_n) \propto P(y) \prod_{i=1}^n P(x_i|y)$$

we need to classify whether players will play or not based on weather condition

Weather	Play
Sunny	No
Overcast	Yes
Rainy	Yes
Sunny	Yes
Sunny	Yes
Overcast	Yes
Rainy	No
Rainy	No
Sunny	Yes
Rainy	Yes
Sunny	No
Overcast	Yes
Overcast	Yes
Rainy	No

**Problem:** Players will play if weather is sunny. Is this statement is correct?

Weather	Play
Sunny	No
Overcast	Yes
Rainy	Yes
Sunny	Yes
Sunny	Yes
Overcast	Yes
Rainy	No
Rainy	No
Sunny	Yes
Rainy	Yes
Sunny	No
Overcast	Yes
Overcast	Yes
Rainy	No

Frequency Table		
Weather	No	Yes
Overcast		4
Rainy	3	2
Sunny	2	3
Grand Total	5	9

Likelihood table				
Weather	No	Yes		
Overcast		4	=4/14	0.29
Rainy	3	2	=5/14	0.36
Sunny	2	3	=5/14	0.36
All	5	9		
	=5/14	=9/14		
	0.36	0.64		

$$P(\text{Yes} \mid \text{Sunny}) = P(\text{Sunny} \mid \text{Yes}) * P(\text{Yes}) / P(\text{Sunny})$$

Here we have  $P(\text{Sunny} \mid \text{Yes}) = 3/9 = 0.33$ ,  $P(\text{Sunny}) = 5/14 = 0.36$ ,  $P(\text{Yes}) = 9/14 = 0.64$

Now,  $P(\text{Yes} \mid \text{Sunny}) = 0.33 * 0.64 / 0.36 = 0.60$ , which has higher probability.

# Baye's Rule

The diagram shows the formula for Bayes' Rule:  $P(c|x) = \frac{P(x|c)P(c)}{P(x)}$ . Four blue arrows point from descriptive labels to the components of the formula: 'Likelihood' points to  $P(x|c)$ , 'Class Prior Probability' points to  $P(c)$ , 'Posterior Probability' points to  $P(c|x)$ , and 'Predictor Prior Probability' points to  $P(x)$ .

$$P(c|x) = \frac{P(x|c)P(c)}{P(x)}$$

Labels and their corresponding parts in the formula:

- Likelihood:  $P(x|c)$
- Class Prior Probability:  $P(c)$
- Posterior Probability:  $P(c|x)$
- Predictor Prior Probability:  $P(x)$

# Naive Bayes Classifier

- When to use:
  - Moderate or large training set available
  - Attributes that describe instances are conditionally independent given classification
- Successful applications:
  - Diagnosis
  - Classifying text documents