

CS 446: Machine Learning

Homework 5

Due on Tuesday, February 20, 2018, 11:59 a.m. Central Time

1. [6 points] Multiclass Classification Basics

- (a) Which of the following is the most suitable application for multiclass classification? Which is the most suitable application for binary classification?
- Predicting tomorrow's stock price;
 - Recognizing flower species from photos;
 - Deciding credit card approval for a bank;
 - Assigning captions to pictures.

Your answer:

- (b) Suppose in an n -dimensional Euclidean space where $n \geq 3$, we have n samples $x^{(i)} = e_i$ for $i = 1 \dots n$ (which means $x^{(1)} = (1, 0, \dots, 0)_n, x^{(2)} = (0, 1, \dots, 0)_n, \dots, x^{(n)} = (0, 0, \dots, 1)_n$), with $x^{(i)}$ having class i . What are the numbers of binary SVM classifiers we need to train, to get 1-vs-all and 1-vs-1 multiclass classifiers?

Your answer:

- (c) Suppose we have trained a 1-vs-1 multiclass classifier from non-regularized binary SVM classifiers on the samples of the previous question. What are the regions in the Euclidean space that will receive the same number of majority votes from more than one classes? **Note:** You can assume for the binary SVM always outputs the *perpendicular bisector* of AB as decision boundary, when trained on two samples A and B . You can ignore samples on the decision boundary of any binary SVM.

Your answer:

2. [8 points] Multiclass SVM

Consider the objective function of multiclass SVM as

$$\min_{w, \xi^{(i)} \geq 0} \frac{C}{2} \|w\|^2 + \sum_{i=1}^n \xi^{(i)}$$

$$\text{s.t. } w_{y^{(i)}} \phi(x^{(i)}) - w_{\hat{y}} \phi(x^{(i)}) \geq 1 - \xi^{(i)} \quad \forall i = 1 \dots n, \hat{y} = 0 \dots K-1, \hat{y} \neq y_i$$

Let $n = K = 3$, $d = 2$, $x^{(1)} = (0, -1)$, $x^{(2)} = (1, 0)$, $x^{(3)} = (0, 1)$, $y^{(1)} = 0$, $y^{(2)} = 1$, $y^{(3)} = 2$, and $\phi(x) = x$.

- (a) Rewrite the objective function with w being a Kd -dimensional vector $(w_1, w_2, w_3, w_4, w_5, w_6)^\top$ and with the specific choices of x , y and ϕ , following the definition of slide No.23 of Lecture 7.

Your answer:

- (b) Rewrite the objective function you get in (a) such that there are no slack variables $\xi^{(i)}$.

Your answer:

- (c) Let $w_t = (1, 1, 1, 2, 1, -1)^\top$. Compute the derivative of the objective function you get in (b) w.r.t. w_2 , at w_t , where w_2 is the weight of second dimension on Class 0 (in case you used non-conventional definition of w in (a)).

Your answer:

- (d) Prove that

$$\max_{\hat{y}} \left(1 + w_{\hat{y}}^\top \phi(x) \right) = \lim_{\epsilon \rightarrow 0} \epsilon \ln \sum_{\hat{y}} \exp \left(\frac{1 + w_{\hat{y}}^\top \phi(x)}{\epsilon} \right).$$

Your answer: