

Supplemental Information

Whole-Genome CRISPR Screening Identifies Genetic Modifiers of Stem Cell-Derived Islet Transplantation

Running Head: CRISPR Screening on Transplanted SC-Islets

Marlie M. Maestas^{1,2}, Kameron Bradley^{2,3}, Mira Shunkarova², Noyonika Mukherjee², Matthew Ishahak², James Lu², and Jeffrey R. Millman^{1,2,3*}

¹Roy and Diana Vagelos Division of Biology and Biomedical Sciences

Washington University School of Medicine, MSC 8127-057-08

660 South Euclid Avenue, St. Louis, MO 63110, USA

²Division of Endocrinology, Metabolism, and Lipid Research

Washington University School of Medicine, MSC 8127-057-08

660 South Euclid Avenue, St. Louis, MO 63110, USA

³Department of Biomedical Engineering

Washington University in St. Louis


Washington University School of Medicine, MSC 8127-057-08

660 South Euclid Avenue, St. Louis, MO 63110, USA

*To whom correspondence should be addressed: Jeffrey R. Millman, jmillman@wustl.edu,

Division of Endocrinology, Metabolism, and Lipid Research, Washington University School of

Medicine, MSC 8127-057-08, 660 South Euclid Avenue, St. Louis, MO 63110, USA



Supplementary Figure 1. Single-cell sequencing of *FCAMR* OE in vitro.

a. Pearson correlation between CTRL and *FCAMR* OE cells across cell types. **b-e.** Volcano plots of differentially expressed genes between CTRL and *FCAMR* OE in (b) β -cells, (c) EC-cells, (d) Stressed cells, (e) proliferating cells. **f.** Violin plot of stress-associated genes between CTRL and *FCAMR* OE.