MDPI

Article

Tweedie Evolutionary Credibility with Autoregressive Latent Risk

Nahyun Kang[1] and Himchan Jeong[1,2]*

[1] Department of Statistics and Actuarial Science, Simon Fraser University, Burnaby, BC V5A 1S6, Canada
[2] Department of Next Generation Smart Energy System Convergence, Gachon University, Republic of Korea
* Correspondence: himchan_jeong@sfu.ca

Abstract

Claim costs combine a mass at zero, a heavy right tail and serial dependence within policies. We develop evolutionary credibility for Tweedie claim costs in a GLM rating structure, where a multiplicative latent risk factor follows a second-order stationary process. For any latent autocovariance $\gamma(k)$, the best linear premium is the GLM premium times a closed-form credibility factor that depends only on $\gamma(\cdot)$ and a single nugget constant; the naive premium and the exact Tweedie credibility of Ohlsson and Johansson (2006) are special cases. For autoregressive latent risk, the premium has an updating-type Kalman recursion with positive weights; it is always positive, and the maximal bonus is bounded by the persistence parameter. Because every input is identified by second moments of standardized residuals, we propose a semi-parametric two-step GMM estimator that requires no distributional assumption on the latent process. The estimator comes with stage-1-corrected standard errors, a Hansen specification test and Wald tests of static versus evolutionary credibility. In simulations with estimated parameters, the feasible premium captures 80–97% of the oracle gain whenever latent heterogeneity is non-negligible, and a fitted static premium is never materially worse than the naive premium. Interval estimates of the latent variance are less reliable when the latent signal is weak or the residuals are very heavy-tailed.

Keywords: evolutionary credibility; Tweedie distribution; latent autoregressive process; Kalman filter; generalized method of moments; bonus–malus; experience rating

MSC: 62P05; 91G05; 62M10; 62F12

1. Introduction

Insurance claim costs have a large mass at zero, a heavy right tail and a variance that grows with the mean. The Tweedie compound Poisson–gamma family with power parameter $1 < p < 2$ captures all three features in a single exponential dispersion model and, combined with a log-link generalized linear model (GLM), is a standard tool for pricing with observable rating factors [1–4]. Rating factors leave persistent unexplained heterogeneity, and experience rating, or credibility, uses each policyholder's claim history to correct the GLM premium [5].

Classical credibility treats the latent risk of a policyholder as constant over time, so past years are exchangeable and receive equal weight. In practice, driving habits, occupation, health and exposure change, so the latent risk evolves. Recent experience is then more informative than old experience. Evolutionary credibility models make this

Received:
Revised:
Accepted:
Published:

Copyright: © 2026 by the authors. Submitted to *Mathematics* for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license.

explicit by letting the latent risk follow a stationary process [5–8]. In frequency models with autoregressive random effects, the resulting premiums discount the age of claims [9,10].

This paper develops evolutionary credibility for Tweedie claim costs inside a GLM rating structure. Conditional on a multiplicative latent factor θ_{it} with mean one, the claim cost Z_{it} of policyholder i in year t is Tweedie with mean $\mu_{it}\theta_{it}$, where $\mu_{it} = \exp(\boldsymbol{x}_{it}^\top \boldsymbol{\beta})$. The latent factor is only assumed to be second-order stationary with autocovariance $\gamma(k)$. The contributions are as follows.

1. **A closed-form premium for any stationary latent process (Theorem 1).** The best linear predictor of next year's claim cost is the GLM premium times a credibility factor computed from $\gamma(\cdot)$ and a single nugget constant $a = \phi\,\mathbb{E}[\theta^p]$. The within-policy noise enters through a heteroscedastic, covariate-dependent diagonal term $a\mu_{it}^{p-2}$. The independent, exchangeable (static), θ-scale AR(1) and log-normal AR(1) cases are corollaries. The exchangeable case reproduces the Tweedie credibility estimator of Ohlsson and Johansson [11].

2. **Structural results for AR(1) dependence (Proposition 1).** The premium has an updating-type (Kalman) recursion that costs $O(T)$ operations per policy. All weights are positive and add up to less than the autoregressive parameter ρ. The premium is therefore always positive, and the maximal bonus is bounded by ρ. We also give conditions under which recent years receive more weight than older ones, and show that this ordering can fail when the GLM mean changes over time.

3. **Semi-parametric GMM estimation and tests (Section 5).** Every input of the premium is identified by second moments of the standardized residuals $e_{it} = Z_{it}/\mu_{it} - 1$. Our two-step generalized method of moments (GMM) estimator needs no distributional assumption on the latent process. It has closed-form, precision-weighted moments, stays inside the parameter space, and comes with a sandwich variance that corrects for first-stage estimation of $\boldsymbol{\beta}$. It also provides a Hansen [12] specification test and Wald tests of static versus evolutionary credibility and of no dependence.

4. **Simulation evidence with estimated parameters (Section 6).** A simulation study with time-varying covariates shows that the estimator is accurate, the tests have close to nominal size, and the feasible premium recovers most of the gain of the infeasible oracle-parameter premium. Exact population mean squared errors serve as a unit test of the simulation engine.

The paper is organized as follows. Section 2 reviews related work. Section 3 presents the model. Section 4 derives the premium and its properties. Section 5 develops estimation and inference, and Section 6 reports the simulation study. Section 7 concludes. Proofs and computational details are in the appendices.

2. Related Literature

Tweedie models and credibility. Tweedie GLMs are widely used for pure premiums because the compound Poisson–gamma representation produces exact zeros and continuous positive amounts [2,3]. Jewell [13] showed that linear credibility is exact Bayesian for conjugate exponential families. Ohlsson and Johansson [11] extended exactness to Tweedie models with a multiplicative random effect. The resulting credibility weights are proportional to μ^{2-p}, the natural Tweedie weights. Related work includes bonus–malus systems with separate frequency and severity components [14], Tweedie models with distribution-free random effects [15], multivariate Tweedie credibility [16] and orderings of credibility factors [17]. These works treat the latent effect as constant over time.

Evolutionary credibility. Gerber and Jones [6] introduced credibility formulas of the updating type, and Sundt [7] developed recursive credibility estimation. In these, the premium is revised each year from the previous premium and the latest observation, which is a Kalman filter. Albrecht [8] derived credibility for Poisson claim counts driven by a weakly stationary latent sequence, with recursive formulas for the coefficients. This is the closest antecedent of Theorem 1. Chapter 9 of Bühlmann and Gisler [5] presents evolutionary models and their recursive (Kalman) computation in general. Pinquet et al. [9] modelled the age of claims in bonus–malus systems. Bolancé et al. [10] estimated and tested autoregressive specifications of random effects in frequency models by moment methods, which is the spirit of our Section 5. Pinquet [18] studied positivity of credibility weights under long-memory ARFIMA$(0, d, 0)$ specifications, which is related to our Proposition 1. Taylor [19] extended evolutionary credibility to hierarchical structures.

Dependence in insurance losses. Copula and multilevel models capture dependence across coverages and over time [20,21]. Multivariate Tweedie constructions have been used for capital and reserving [22–24]. These models describe dependence in the joint distribution of losses. Our focus is the design of linear credibility premiums that exploit the serial dependence of the latent risk.

Positioning. Relative to Albrecht [8] and its successors, this paper adds four elements. First, the conditional response is Tweedie with power variance, so the nugget $a\mu_{it}^{p-2}$ is heteroscedastic and depends on covariates inside a GLM rating structure. Second, the exchangeable special case coincides exactly with Ohlsson and Johansson [11]. Third, we prove positivity and bonus bounds. Fourth, the GMM estimator needs no distributional assumption on θ and comes with tests of static versus evolutionary credibility. The paper builds on the first author's MSc project [25].

3. Model

3.1. Notation and Assumptions

Policyholders $i = 1, \dots, I$ are observed in years $t = 1, \dots, T_i$, and the target is year $T_i + 1$. For exposition we take $T_i = T$; unbalanced panels are discussed in Section 5.5. Table 1 collects the notation. Each symbol has one meaning throughout the paper. In particular, (v, ρ) always refer to the θ scale, and (σ_W^2, ρ_W) to the log scale.

Table 1. Notation.

Symbol	Meaning
$\mu_{it} = \exp(\boldsymbol{x}_{it}^\top \boldsymbol{\beta})$, $\boldsymbol{\mu}_i$	GLM mean and its vector $(\mu_{i1}, \dots, \mu_{iT})^\top$
θ_{it}	latent multiplicative risk factor, $\mathbb{E}[\theta_{it}] = 1$
$v = \mathrm{Var}(\theta_{it})$	latent variance (θ scale)
$\gamma(k)$, $r(k) = \gamma(k)/v$	latent autocovariance $\mathrm{Cov}(\theta_{it}, \theta_{i,t+k})$ and autocorrelation; $\gamma(0) = v$
$a = \phi\, \mathbb{E}[\theta_{it}^p]$	nugget (within-variance) constant
$q_{it} = \mu_{it}^{p-2}$	nugget scale: $\mathrm{Var}(e_{it} \mid \boldsymbol{\theta}_i)$ has mean aq_{it}
$e_{it} = Z_{it}/\mu_{it} - 1$, $\boldsymbol{e}_i$	standardized residual and its vector
$\Gamma_T = [\gamma(\lvert t-s \rvert)]_{t,s=1}^T$	latent covariance matrix of years $1, \dots, T$
$\boldsymbol{c}_T = (\gamma(T), \dots, \gamma(1))^\top$	latent cross-covariance with year $T+1$
$D_i = \mathrm{diag}(aq_{i1}, \dots, aq_{iT})$	nugget matrix
ρ	θ-scale AR(1) parameter: $\gamma(k) = v\rho^k$
$W_{it} = \log \theta_{it}$, σ_W^2, ρ_W	log-scale variance and AR(1) parameter (log-normal AR(1) example)

Assumption 1. *For each policyholder i, conditional on the covariates:*

(A1) $Z_{it} \mid \boldsymbol{\theta}_i \sim \mathcal{T}w(\mu_{it}\theta_{it}, \phi, p)$ *with* $1 < p < 2$, *so that* $\mathbb{E}[Z_{it} \mid \boldsymbol{\theta}_i] = \mu_{it}\theta_{it}$ *and* $\mathrm{Var}(Z_{it} \mid \boldsymbol{\theta}_i) = \phi(\mu_{it}\theta_{it})^p$. *The* Z_{it} *are conditionally independent over* t *given the latent path* $\boldsymbol{\theta}_i = (\theta_{i1}, \theta_{i2}, \dots)$.

(A2) $\theta_{it} > 0$, $\mathbb{E}[\theta_{it}] = 1$ *and* $\mathbb{E}[\theta_{it}^2] < \infty$.

(A3) $(\theta_{it})_t$ *is second-order stationary:* $\mathrm{Cov}(\theta_{it}, \theta_{is}) = \gamma(|t-s|)$ *with* $v = \gamma(0) > 0$.

(A4) $\mathbb{E}[\theta_{it}^p]$ *does not depend on* t *(this holds, e.g., under strict stationarity).*

(A5) Policyholders are mutually independent.

The normalization $\mathbb{E}[\theta_{it}] = 1$ makes μ_{it} the marginal mean, $\mathbb{E}[Z_{it}] = \mu_{it}$, and separates the latent factor from the GLM intercept. By Jensen's and Lyapunov's inequalities, $1 \le \mathbb{E}[\theta^p] \le (1+v)^{p-1}$ for $1 < p < 2$, so a is finite under (A2). Assumptions (A1)–(A5) restrict only the first two moments of the latent process and the mean–variance relation of the response. No distributional assumption on θ is needed. As shown below, the fractional moment $\mathbb{E}[\theta^p]$ enters only through a, and a is identified from the data.

Lemma 1 (Second-moment structure)**.** *Under Assumption 1,* $\mathbb{E}[e_{it}] = 0$ *and*

$$\mathrm{Cov}(e_{it}, e_{is}) = \gamma(|t-s|) + \mathbb{1}\{t = s\}\, a\, q_{it}, \qquad t, s \ge 1. \tag{1}$$

Proof. Write $e_{it} = \xi_{it} + \varepsilon_{it}$ with $\xi_{it} = \theta_{it} - 1$ and $\varepsilon_{it} = (Z_{it} - \mu_{it}\theta_{it})/\mu_{it}$. By (A1), $\mathbb{E}[\varepsilon_{it} \mid \boldsymbol{\theta}_i] = 0$, $\mathrm{Var}(\varepsilon_{it} \mid \boldsymbol{\theta}_i) = \phi\theta_{it}^p \mu_{it}^{p-2}$ and $\mathrm{Cov}(\varepsilon_{it}, \varepsilon_{is} \mid \boldsymbol{\theta}_i) = 0$ for $t \ne s$. Hence the ε_{it} are uncorrelated with each other and with the latent path, and $\mathrm{Var}(\varepsilon_{it}) = \phi\, \mathbb{E}[\theta_{it}^p]\, \mu_{it}^{p-2} = a q_{it}$ by (A4). The result follows from (A3). □

Intuitively, the decomposition $e_{it} = \xi_{it} + \varepsilon_{it}$ separates the observable relative surprise e_{it}, i.e., how far the claim cost deviates from its GLM premium, into a persistent signal $\xi_{it} = \theta_{it} - 1$, the unobserved deviation of the policyholder's true risk level, and a transient noise ε_{it}, the luck of the claim draw around that level, which is uncorrelated over time and whose variance $a q_{it}$ shrinks as μ_{it} grows; credibility weighting amounts to extracting the signal from the noisy residuals.

3.2. Examples of Latent Processes

Any positive-semidefinite autocovariance $\gamma(\cdot)$ is allowed. The following cases are used below.

- *Independence:* $\gamma(k) = 0$ for $k \ge 1$.
- *Exchangeable (static):* $\theta_{it} = \theta_i$, so $\gamma(k) = v$ for all k.
- *θ-scale AR(1):* $\gamma(k) = v\rho^k$, $0 \le \rho < 1$. Any positive process with $\mathbb{E}[\theta_{t+1} \mid \theta_t, \theta_{t-1}, \dots] = 1 - \rho + \rho\theta_t$ has this autocovariance, for example the gamma autoregressive sequences of Gaver and Lewis [26].
- *Log-normal AR(1):* $W_{it} = \log\theta_{it}$ is a stationary Gaussian AR(1) with variance σ_W^2, autocorrelation ρ_W and mean $-\sigma_W^2/2$. Then

$$v = e^{\sigma_W^2} - 1, \qquad \gamma(k) = \exp(\sigma_W^2 \rho_W^k) - 1, \qquad \mathbb{E}[\theta^p] = \exp\{\sigma_W^2 p(p-1)/2\}. \tag{2}$$

  The θ-scale autocorrelation $r(k) = \{\exp(\sigma_W^2\rho_W^k) - 1\}/\{\exp(\sigma_W^2) - 1\}$ is *not* geometric. By convexity, $r(k) \le \rho_W^k$. For given $(v, r(1))$, the log-scale parameters are $\sigma_W^2 = \log(1+v)$ and $\rho_W = \log\{1 + v\, r(1)\}/\sigma_W^2$.
- *Other structures:* ARMA, long-memory (ARFIMA) and regime-switching processes for $\log\theta_{it}$ or for θ_{it}. Each implies its own $\gamma(\cdot)$, and Theorem 1 applies unchanged.

4. The Evolutionary Credibility Premium

4.1. General Stationary Latent Processes

We suppress the index i. Let $\mathcal{F}_T = \{Z_1, \dots, Z_T\}$, with the covariates $\boldsymbol{x}_1, \dots, \boldsymbol{x}_{T+1}$ treated as known. The (inhomogeneous) linear credibility premium is the best linear predictor

$$P_{T+1} = \widehat{\alpha}_0 + \sum_{t=1}^{T} \widehat{\alpha}_t Z_t, \qquad (\widehat{\alpha}_0, \dots, \widehat{\alpha}_T) = \operatorname*{arg\,min}_{\alpha \in \mathbb{R}^{T+1}} \mathbb{E}\Big[\Big(Z_{T+1} - \alpha_0 - \sum_{t=1}^{T} \alpha_t Z_t\Big)^2\Big].$$

Theorem 1. *Under Assumption 1, the linear credibility premium for year $T+1$ is*

$$P_{T+1} = \mu_{T+1}\Big\{1 + \boldsymbol{c}_T^\top (\Gamma_T + D_T)^{-1} \boldsymbol{e}_T\Big\} = \mu_{T+1}\Big(1 + \sum_{t=1}^{T} w_t e_t\Big), \qquad \boldsymbol{w} = (\Gamma_T + D_T)^{-1} \boldsymbol{c}_T, \tag{3}$$

where $\Gamma_T = [\gamma(|t-s|)]_{t,s \le T}$, $D_T = \operatorname{diag}(a\mu_1^{p-2}, \dots, a\mu_T^{p-2})$, $\boldsymbol{c}_T = (\gamma(T), \dots, \gamma(1))^\top$ and $\boldsymbol{e}_T = (e_1, \dots, e_T)^\top$. The mean squared error is

$$\mathbb{E}[(Z_{T+1} - P_{T+1})^2] = \mu_{T+1}^2 \{v + a\mu_{T+1}^{p-2} - \boldsymbol{c}_T^\top (\Gamma_T + D_T)^{-1} \boldsymbol{c}_T\}.$$

Proof. Since $Z_t = \mu_t(1 + e_t)$ with known μ_t, linear predictors in $(1, Z_1, \dots, Z_T)$ and in $(1, e_1, \dots, e_T)$ coincide. Because $\mathbb{E}[\boldsymbol{e}_T] = \boldsymbol{0}$, the best linear predictor of e_{T+1} given $\boldsymbol{e}_T$ is

$$\operatorname{Cov}(e_{T+1}, \boldsymbol{e}_T)\ \operatorname{Var}(\boldsymbol{e}_T)^{-1} \boldsymbol{e}_T.$$

By Lemma 1, $\operatorname{Var}(\boldsymbol{e}_T) = \Gamma_T + D_T$, which is positive definite because Γ_T is positive semi-definite and D_T has a positive diagonal. Also $\operatorname{Cov}(e_{T+1}, e_t) = \gamma(T+1-t)$. Multiplying by μ_{T+1} gives (3), and the MSE is the usual prediction error variance. □

The premium is a multiplicative experience correction to the GLM premium. The weight of each past year depends on the covariance of the latent risk with year $T+1$ and on the precision of that year's experience. A year with a large expected claim cost μ_t has a smaller nugget $a\mu_t^{p-2}$, because $p < 2$, and therefore carries more information about θ_t.

Corollary 1 (Special cases). *(i)* Independence. *If $\gamma(k) = 0$ for $k \ge 1$, then $P_{T+1} = \mu_{T+1}$, the naive (GLM) premium.*

(ii) Exchangeable. *If $\gamma(k) = v$ for all $k \ge 0$, then, with $\kappa = a/v$,*

$$P_{T+1} = \mu_{T+1}\, \frac{\kappa + \sum_{t=1}^{T} Z_t\, \mu_t^{1-p}}{\kappa + \sum_{t=1}^{T} \mu_t^{2-p}}. \tag{4}$$

This is the Tweedie credibility estimator of Ohlsson and Johansson [11]: a Bühlmann–Straub-type estimator with natural weights μ_t^{2-p} and credibility constant κ.

(iii) θ-scale AR(1). *If $\gamma(k) = v\rho^k$, then $\boldsymbol{w} = (R_T + v^{-1} D_T)^{-1} (\rho^T, \dots, \rho)^\top$ with $[R_T]_{ts} = \rho^{|t-s|}$.*

(iv) Log-normal AR(1). *If $\log \theta_t$ is a Gaussian AR(1), (3) holds with $\gamma(k)$ and a given by (2).*

Proof. (i) $\boldsymbol{c}_T = \boldsymbol{0}$. (ii) $\Gamma_T = v\boldsymbol{1}\boldsymbol{1}^\top$ and $\boldsymbol{c}_T = v\boldsymbol{1}$. By the Sherman–Morrison formula, $\boldsymbol{w} = vD_T^{-1}\boldsymbol{1} / (1 + v\boldsymbol{1}^\top D_T^{-1}\boldsymbol{1})$, so $w_t = \mu_t^{2-p} / (\kappa + \sum_s \mu_s^{2-p})$. Using $\mu_t^{2-p}(1 + e_t) = Z_t \mu_t^{1-p}$ gives (4). (iii) and (iv) are direct substitutions. □

Remark 1 (Which moments of θ matter)**.** *The premium depends on the latent law only through $\gamma(\cdot)$ and the fractional moment $\mathbb{E}[\theta^p]$. The latter enters only through the product $a = \phi\,\mathbb{E}[\theta^p]$, which is identified from the diagonal second moments of e_{it} (Section 5). Neither ϕ nor $\mathbb{E}[\theta^p]$ needs to be known separately, and no distributional assumption on θ is required. Formulas such as $\mathbb{E}[\theta^p] = (1+v)^{p(p-1)/2}$ hold only for specific constructions (here, a log-normal marginal with variance v) and are not used for estimation.*

4.2. AR(1) Dependence: Recursion, Positivity and Bounds

For the θ-scale AR(1) structure $\gamma(k) = v\rho^k$, write $\xi_t = \theta_t - 1$ and $d_t = a\mu_t^{p-2}$. Then $\zeta_t = \xi_{t+1} - \rho\xi_t$ is uncorrelated with ξ_s for $s \le t$ and with every ε_s, and $\mathrm{Var}(\zeta_t) = v(1-\rho^2)$. Hence

$$\xi_{t+1} = \rho\xi_t + \zeta_t, \qquad e_t = \xi_t + \varepsilon_t, \qquad \mathrm{Var}(\varepsilon_t) = d_t,$$

is a linear state-space model in the second-order sense. The Kalman recursions therefore compute linear projections exactly, without any Gaussian assumption [27, Chapter 12]; see also Harvey [28].

Proposition 1. *Let $\gamma(k) = v\rho^k$ with $0 < \rho < 1$, and let $d_t = a\mu_t^{p-2} > 0$.*

(a) Updating-type recursion. *Set $\hat{\xi}_1 = 0$, $V_1 = v$ and, for $t = 1, \dots, T$,*

$$K_t = \frac{V_t}{V_t + d_t}, \qquad \hat{\xi}_{t+1} = \rho\{(1-K_t)\hat{\xi}_t + K_t e_t\}, \qquad V_{t+1} = \rho^2(1-K_t)V_t + (1-\rho^2)v.$$

Then $P_{T+1} = \mu_{T+1}(1+\hat{\xi}_{T+1})$ coincides with (3), and V_{T+1} is the prediction MSE of θ_{T+1}.

(b) Weights. *$w_t = \rho^{T+1-t}K_t\prod_{s=t+1}^{T}(1-K_s) > 0$ for $t = 1, \dots, T$.*

(c) Total credibility. *$S_T = \sum_{t=1}^{T} w_t$ satisfies $S_t = \rho\{(1-K_t)S_{t-1} + K_t\}$ with $S_0 = 0$, and $0 < S_T < \rho$.*

(d) Positivity and bonus bound. *$P_{T+1} \ge \mu_{T+1}(1-S_T) > (1-\rho)\mu_{T+1} > 0$. The maximal bonus, attained by a claim-free history, is $S_T < \rho$, so the premium discount never exceeds $100\rho\%$. The malus is unbounded.*

(e) Comparative statics. *Every gain K_t, the most recent weight $w_T = \rho K_T$ and the total credibility S_T are nondecreasing in v. Because the weights depend on $(v, a, \boldsymbol{\mu})$ only through d_t/v, the same holds under a proportional increase of all μ_t.*

(f) Recency. *If $d_t - d_{t-1} \le v(1-\rho)/\rho$ for $t = 2, \dots, T$, then $w_1 < w_2 < \cdots < w_T$. This holds, in particular, when μ_t is constant or nondecreasing in t. Without such a condition the ordering can fail.*

(g) Limits and steady state. *As $\rho \to 0$, $P_{T+1} \to \mu_{T+1}$. As $\rho \to 1$, P_{T+1} converges to the exchangeable premium (4), and the recursion becomes the Bühlmann–Straub recursion $V_{t+1} = (1-K_t)V_t$. If $d_t \equiv d$, then $K_t \to K^* = u^*/(u^*+\delta)$ with $\delta = d/v$ and*

$$u^* = \tfrac{1}{2}\Big[-(1-\rho^2)(\delta-1) + \sqrt{(1-\rho^2)^2(\delta-1)^2 + 4(1-\rho^2)\delta}\Big].$$

For long histories the premium is exponential smoothing: the weight of year $T+1-j$ tends to $\rho K^*\{\rho(1-K^*)\}^{j-1}$, and $S_T \to \rho K^*/\{1-\rho(1-K^*)\} < \rho$.

The proof is in Appendix A. Part (d) has two practical consequences. The fitted premium is always strictly positive, so the Tweedie deviance is always defined. The persistence parameter ρ also directly caps the bonus a claim-free history can earn. Part (f) is

the "age of claims" property of Pinquet et al. [9]. A counterexample to unrestricted recency is $T = 2$, $v = 0.5$, $\rho = 0.8$, $d = (0.01, 25)$: an old year with a large expected cost (small nugget) receives weight $w_1 = 0.623$, while a recent year with a small expected cost receives $w_2 = 0.006$. Part (e) is in the spirit of the orderings of credibility factors in Ahn et al. [17]. Individual older weights need not be monotone in v, because a larger v also shifts weight toward recent years.

For a general $\gamma(\cdot)$, the Durbin–Levinson or innovations algorithm provides an analogous $O(T^2)$ recursion [27, Chapter 5]. Positivity of the weights is then not guaranteed for non-monotone autocovariances [cf. 18]. Under the log-normal AR(1) structure, positivity held in all our simulated portfolios, but this is an empirical observation rather than a theorem.

5. Semi-Parametric Estimation and Inference

5.1. Moment Identities and Identification

By Lemma 1, the standardized residuals satisfy

$$\mathbb{E}[e_{it}e_{i,t+k} \mid \boldsymbol{x}_i] = \gamma(k), \qquad k = 1, \dots, T-1, \tag{5}$$

$$\mathbb{E}[e_{it}^2 \mid \boldsymbol{x}_i] = \gamma(0) + a\, q_{it}, \qquad q_{it} = \mu_{it}^{p-2}. \tag{6}$$

The lag moments (5) do not depend on ϕ, p or the law of θ. The premium (3) depends on the data-generating process only through $(\boldsymbol{\beta}, p)$, $\gamma(0), \dots, \gamma(T)$ and a. Hence:

- $\gamma(1), \dots, \gamma(T-1)$ are identified nonparametrically by (5);
- $\gamma(0)$ and a are identified separately by (6) whenever μ_{it}^{p-2} varies across observations, which is the case with any nontrivial rating structure;
- $\gamma(T)$, which enters $\boldsymbol{c}_T$, cannot be identified nonparametrically from a T-year panel and requires a parametric structure or an extrapolation rule.

5.2. Two-Step Estimator

Stage 1: mean structure. Fit the Tweedie quasi-likelihood GLM under working independence, i.e., solve $\sum_{i,t} \boldsymbol{x}_{it}(Z_{it} - \mu_{it})\mu_{it}^{1-p} = \boldsymbol{0}$. The estimator $\widehat{\boldsymbol{\beta}}$ is consistent because $\mathbb{E}[Z_{it}] = \mu_{it}$ [29], whatever the serial dependence. Set $\hat{\mu}_{it} = \exp(\boldsymbol{x}_{it}^\top \widehat{\boldsymbol{\beta}})$, $\hat{e}_{it} = Z_{it}/\hat{\mu}_{it} - 1$ and $\hat{q}_{it} = \hat{\mu}_{it}^{p-2}$.

Stage 2: weighted moment contributions. Collect the second-moment parameters in

$$\boldsymbol{\eta} = (\gamma(0), a, \gamma(1), \dots, \gamma(T-1))^\top \in \mathbb{R}^{T+1}.$$

The per-policy moment contributions are

$$g_{ik}(\boldsymbol{\eta}) = \sum_{t=1}^{T-k} w_{itk}\{\hat{e}_{it}\hat{e}_{i,t+k} - \gamma(k)\}, \qquad k = 1, \dots, T-1, \tag{7}$$

$$h_{ij}(\boldsymbol{\eta}) = \sum_{t=1}^{T} \omega_{it}\, z_{itj}\{\hat{e}_{it}^2 - \gamma(0) - a\hat{q}_{it}\}, \qquad \boldsymbol{z}_{it} = (1, \hat{q}_{it})^\top, \tag{8}$$

with precision weights

$$w_{itk} = (\hat{s}_{it}^2\, \hat{s}_{i,t+k}^2)^{-1}, \qquad \omega_{it} = \hat{s}_{it}^{-4}, \qquad \hat{s}_{it}^2 = \hat{a}\,\hat{q}_{it} + \hat{v}.$$

Here $\hat{s}_{it}^2$ estimates $\mathrm{Var}(e_{it})$. The weights are computed from preliminary estimates and updated two or three times, starting from equal weights. They down-weight policy-years with small μ_{it}, whose residuals consist mostly of Tweedie noise. This matters when μ is very

heterogeneous. Since the weights depend only on covariates and preliminary estimates, they do not affect consistency.

Let $\boldsymbol{m}_i(\boldsymbol{\eta}) = (h_{i1}, h_{i2}, g_{i1}, \ldots, g_{i,T-1})^\top$ and $\bar{\boldsymbol{m}}(\boldsymbol{\eta}) = I^{-1}\sum_i \boldsymbol{m}_i(\boldsymbol{\eta})$. The moments are *linear* in $\boldsymbol{\eta}$, so $\bar{\boldsymbol{m}}(\boldsymbol{\eta}) = \bar{\boldsymbol{b}} - \bar{A}\boldsymbol{\eta}$ for a block-diagonal matrix $\bar{A}$ and a vector $\bar{\boldsymbol{b}}$ of weighted sums. The *nonparametric* (NP) estimator solves $\bar{\boldsymbol{m}}(\widehat{\boldsymbol{\eta}}) = \mathbf{0}$ in closed form: each $\hat{\gamma}(k)$ is a weighted average of lag-k cross-products, and $(\hat{\gamma}(0), \hat{a})$ is a weighted regression of $\hat{e}_{it}^2$ on $(1, \hat{q}_{it})$.

Parametric latent structures. A structure $\boldsymbol{\eta} = \boldsymbol{\eta}(\boldsymbol{\psi})$ specifies $\gamma(k; \boldsymbol{\psi})$ and a. We use

- S0 (independence, naive): $\gamma(k) = 0$ for $k \geq 1$;
- S1 (θ-scale AR(1)): $\boldsymbol{\psi} = (v, \rho, a)$, $\gamma(k) = v\rho^k$; Proposition 1 applies;
- S2 (log-normal AR(1)): $\boldsymbol{\psi} = (\sigma_W^2, \rho_W, a)$, $\gamma(k) = \exp(\sigma_W^2 \rho_W^k) - 1$;
- S3 (exchangeable): $\boldsymbol{\psi} = (v, a)$, $\gamma(k) = v$, which gives a *fitted* static benchmark through Corollary 1(ii);
- NP: $\widehat{\boldsymbol{\eta}}$ itself, with the tail $\hat{\gamma}(T) = \hat{\gamma}(T-1)\min\{\hat{\gamma}(T-1)/\hat{\gamma}(T-2), 1\}$ when both are positive, and 0 otherwise.

The efficient two-step GMM estimator is

$$\widehat{\boldsymbol{\psi}} = \arg\min_{\boldsymbol{\psi}} \; \bar{\boldsymbol{m}}(\boldsymbol{\eta}(\boldsymbol{\psi}))^\top \widehat{\Omega}^{-1} \bar{\boldsymbol{m}}(\boldsymbol{\eta}(\boldsymbol{\psi})), \tag{9}$$

where $\widehat{\Omega}$ is the covariance matrix of the per-policy contributions (clustered by policyholder) after the stage-1 correction below. Because $\bar{\boldsymbol{m}}$ is linear in $\boldsymbol{\eta}$, (9) is equivalent to the minimum-distance problem

$$\widehat{\boldsymbol{\psi}} = \arg\min_{\boldsymbol{\psi}} \{\widehat{\boldsymbol{\eta}} - \boldsymbol{\eta}(\boldsymbol{\psi})\}^\top \widehat{V}^{-1} \{\widehat{\boldsymbol{\eta}} - \boldsymbol{\eta}(\boldsymbol{\psi})\}, \qquad \widehat{V} = I^{-1}\bar{A}^{-1}\widehat{\Omega}\bar{A}^{-\top}. \tag{10}$$

All structures are therefore fitted to the same $T+1$ summary statistics with their estimated covariance, which is fast and numerically stable. We reparametrize $v = e^\nu$, $\sigma_W^2 = e^\nu$, $\rho = \text{logit}^{-1}(\kappa)$, $\rho_W = \text{logit}^{-1}(\kappa)$ and $a = e^\alpha$, so that estimates always lie in the parameter space. We also report two robustness variants of S1, each of which removes one ingredient of the estimator. The *lag-only* variant fits (v, ρ) to the lag moments (5) alone and takes $\hat{a}$ from the NP diagonal block. The lag moments are free of the claim noise and do not depend on p, ϕ or the variance function, whereas the diagonal moment (6) mixes signal and noise, relies on the nugget being proportional to μ_{it}^{p-2} and is driven by the heavy-tailed $\hat{e}_{it}^2$. Comparing the two fits therefore shows how much the diagonal moment adds, and a large discrepancy flags a misspecified variance function or p. The *equal-weight* variant sets $w_{itk} = \omega_{it} = 1$. Because the moment conditions hold for any weights that depend only on covariates, the weights affect efficiency rather than consistency, and this variant measures how much precision they buy, which is expected to matter most when μ_{it} is very heterogeneous.

Plug-in premium. The feasible premium inserts $\hat{\mu}_{i,T+1}$, $\hat{e}_{it}$, $\hat{q}_{it}$, $\gamma(k; \widehat{\boldsymbol{\psi}})$ and $\hat{a}$ into (3), or into the recursion of Proposition 1 for S1.

5.3. Asymptotic Theory

Let $\boldsymbol{s}_i(\boldsymbol{\beta}) = \sum_t \boldsymbol{x}_{it}(Z_{it} - \mu_{it})\mu_{it}^{1-p}$ be the stage-1 quasi-score and $S = \mathbb{E}[\partial \boldsymbol{s}_i / \partial \boldsymbol{\beta}^\top]$. Stage 1 matters for inference: since $\partial e_{it}/\partial \boldsymbol{\beta} = -\boldsymbol{x}_{it}(1 + e_{it})$, we have $\mathbb{E}[\partial(e_{it}e_{is})/\partial\boldsymbol{\beta}] = -(\boldsymbol{x}_{it} + \boldsymbol{x}_{is})\gamma(|t-s|) \neq \mathbf{0}$ for $t \neq s$. Let $G_\beta = \mathbb{E}[\partial \boldsymbol{m}_i / \partial \boldsymbol{\beta}^\top]$ and define the corrected contributions $\tilde{\boldsymbol{m}}_i = \boldsymbol{m}_i - G_\beta S^{-1}\boldsymbol{s}_i$. This is the influence function of the stacked estimator [30, Section 6].

Proposition 2. *Suppose Assumption 1 holds with $(\boldsymbol{x}_i, \boldsymbol{\theta}_i, \boldsymbol{Z}_i)$ i.i.d. across i, $\mathbb{E}[\theta_{it}^4] < \infty$, bounded covariates and a nonsingular $\mathbb{E}[\sum_t \omega_{it} \boldsymbol{z}_{it} \boldsymbol{z}_{it}^\top]$. Let T be fixed and $I \to \infty$. Then*

(i) $\sqrt{I}(\widehat{\boldsymbol{\eta}} - \boldsymbol{\eta}_0) \to N(\boldsymbol{0}, V_\infty)$ *with* $V_\infty = \bar{A}^{-1} \Omega \bar{A}^{-\top}$ *and* $\Omega = \mathrm{Var}(\tilde{\boldsymbol{m}}_i)$;

(ii) if $\boldsymbol{\eta}_0 = \boldsymbol{\eta}(\boldsymbol{\psi}_0)$ *with* $\boldsymbol{\psi}_0$ *interior,* $\boldsymbol{\eta}(\cdot)$ *is injective and continuously differentiable, and* $H = \partial \boldsymbol{\eta} / \partial \boldsymbol{\psi}^\top$ *has full column rank at* $\boldsymbol{\psi}_0$*, then* $\sqrt{I}(\widehat{\boldsymbol{\psi}} - \boldsymbol{\psi}_0) \to N(\boldsymbol{0}, (H^\top V_\infty^{-1} H)^{-1})$;

(iii) the Hansen statistic $J = \{\widehat{\boldsymbol{\eta}} - \boldsymbol{\eta}(\widehat{\boldsymbol{\psi}})\}^\top \widehat{V}^{-1} \{\widehat{\boldsymbol{\eta}} - \boldsymbol{\eta}(\widehat{\boldsymbol{\psi}})\}$ *converges in distribution to* $\chi^2_{T+1-\dim \boldsymbol{\psi}}$.

Sketch. (i) follows from a first-order expansion of $\bar{\boldsymbol{m}}(\widehat{\boldsymbol{\eta}}; \widehat{\boldsymbol{\beta}}) = \boldsymbol{0}$ around $(\boldsymbol{\eta}_0, \boldsymbol{\beta}_0)$ combined with $\widehat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0 = -S^{-1} I^{-1} \sum_i \boldsymbol{s}_i + o_p(I^{-1/2})$. The moment conditions hold for any weights that are functions of covariates, so estimating the weights does not affect the limit. (ii) and (iii) are standard results for optimal minimum distance and GMM [12,30]. □

In practice, G_β is computed by numerical differentiation of $\bar{\boldsymbol{m}}$ with the weights held fixed, and $\widehat{\Omega}$ is the sample covariance of the $\tilde{\boldsymbol{m}}_i$. A policyholder-level (cluster) bootstrap is a simple alternative. Delta-method intervals for v are formed on the log scale.

5.4. Tests

- *Specification.* The J statistic of Proposition 2(iii) tests a parametric structure, with $T - 2$ degrees of freedom for S1 and S2 and $T - 1$ for S3.
- *Static versus evolutionary credibility.* The Wald statistic for $H_0 : \gamma(1) = \cdots = \gamma(T-1)$, computed from the NP estimates, is asymptotically χ^2_{T-2}. It does not require testing $\rho = 1$ on the boundary of S1, and it does not involve the diagonal moments, so it is robust to misspecification of the variance function. In spirit it parallels the tests of Bolancé et al. [10] for frequency models.
- *No dependence.* The Wald statistic for $H_0 : \gamma(1) = \cdots = \gamma(T-1) = 0$ from the NP estimates is asymptotically χ^2_{T-1}. The NP estimates are unrestricted, so there is no boundary problem. Testing $v = 0$ within S1 requires the boundary theory of Andrews [31].

5.5. Practical Extensions

Choice of p. The marginal variance is $\mathrm{Var}\, Z_{it} = a\mu_{it}^p + v\mu_{it}^2$, so a marginal Tweedie profile likelihood estimates a working value of p that is biased toward 2. Instead, p can be chosen to minimize the J statistic of the diagonal moments (6) with the extra instrument $\hat{q}_{it} \log \hat{\mu}_{it}$ (one overidentifying restriction), or estimated jointly with $\boldsymbol{\eta}$. Sensitivity to p should be reported; see Section 6.4.

Exposure. For pure premiums $Y_{it} = Z_{it}/\omega_{it}$ with exposure ω_{it} and $Y_{it} \mid \theta_{it} \sim \mathcal{T}w(\mu_{it}\theta_{it}, \phi/\omega_{it}, p)$, set $e_{it} = Y_{it}/\mu_{it} - 1$. The diagonal moment becomes $\gamma(0) + a\mu_{it}^{p-2}/\omega_{it}$, and q_{it} and the weights change accordingly.

Unbalanced panels. Entry, exit and gaps need no special treatment. The sums in (7)–(8) run over observed years and observed pairs, and in (3) unobserved years are dropped. Equivalently, their nugget is set to infinity, which in Proposition 1 means $K_t = 0$.

Computation. The estimator requires one GLM fit, $O(IT^2)$ sums for the moment contributions and a low-dimensional minimum-distance problem. No numerical integration or MCMC is needed. For a portfolio of $I = 10{,}000$ policies and $T = 5$, the whole pipeline (GLM, moments, S1 fit and premiums) takes well under one second on a single CPU core.

6. Simulation Study

6.1. Design

Data-generating process. Each replication simulates $I = 2000$ policyholders over $T = 5$ training years and one hold-out year $T+1$. Covariates are $\boldsymbol{x}_{it} = (1, x_{1,it}, x_{2,i})^\top$, where $x_{1,it} = b_i + u_{it}$ has a time-invariant part $b_i \sim N(0, 0.6^2)$ and a time-varying part $u_{it} \sim N(0, 0.8^2)$, and $x_{2,i} \sim \text{Bernoulli}(0.5)$ is fixed over time. The GLM mean is $\mu_{it} = \exp(2.5 + 0.5x_{1,it} - 0.5x_{2,i})$. Time-varying covariates exercise the time-varying nugget $a\mu_{it}^{p-2}$ that the premium is built for, and they help separate v from a in (6). The latent factor follows the log-normal AR(1) model of Section 3.2, and $Z_{it} \mid \theta_{it} \sim \mathcal{T}w(\mu_{it}\theta_{it}, \phi = 10, p = 1.5)$. The share of zero claims is 54–60% across cells (Table 2).

Grid. The grid is indexed by interpretable θ-scale quantities: $v \in \{0.1, 0.25, 0.5, 1\}$, with $v = 2$ as a stress case, and $r(1) \in \{0.3, 0.5, 0.7, 0.9\}$. Each pair maps to the log-normal AR(1) by $\sigma_W^2 = \log(1+v)$ and $\rho_W = \log\{1 + v\,r(1)\}/\sigma_W^2$. Under this data-generating process, S2 is correctly specified, whereas S1 is a (mildly) misspecified working model, because the θ-scale autocorrelation is not exactly geometric. Each of the 20 cells uses 500 replications.

Competitors. All feasible methods use parameters estimated from the training years only.

- *Naive*: the GLM premium $\hat{\mu}_{i,T+1}$.
- *Static (fitted)*: structure S3 and Corollary 1(ii). This is a fitted exchangeable model that estimates the covariance best matching the observed moments, not the full Var θ.
- *S1*: structure S1 and the recursion of Proposition 1 (the main proposal).
- *S2* and *NP*: structure S2 and the nonparametric autocovariances, plugged into Theorem 1.
- *S1 lag-only* and *S1 equal weights*: the robustness variants of Section 5.
- *Oracle parameters*: Theorem 1 with the true $(\boldsymbol{\beta}, \gamma, a)$, i.e., the best linear premium.
- *Best exchangeable*: the static premium of Corollary 1(ii) with the true $\boldsymbol{\beta}$ and a and the credibility constant chosen to minimize the population MSE under the true $\gamma(\cdot)$ (Appendix B).
- *Oracle* θ: $\mu_{i,T+1}\theta_{i,T+1}$, the true conditional mean of next year's claim cost, computed from the true $\boldsymbol{\beta}$ and the realized, unobservable latent factor rather than from past claims. Its premium error is zero by construction, so its out-of-sample MSE equals the irreducible Tweedie noise $\mathbb{E}[a\mu_{i,T+1}^p]$, a lower bound for every premium. It is infeasible in practice and serves only to show how much of the MSE no premium can remove.

Metrics. For prediction, we report the paired relative reduction of out-of-sample MSE versus the naive premium, averaged over replications with Monte Carlo standard errors. We also report the share of the oracle gain,

$$\text{share}(M) = \frac{\text{PE}_{\text{naive}} - \text{PE}_M}{\text{PE}_{\text{naive}} - \text{PE}_{\text{oracle param}}}, \qquad \text{PE}_M = \mathbb{E}[(\mu_{T+1}\theta_{T+1} - P_{T+1}^M)^2],$$

which uses the premium error to remove the irreducible Tweedie noise (Appendix B); its standard error is computed by bootstrapping replications. We also report the fraction of replications in which a method beats the naive premium, and the mean Tweedie deviance. For estimation, we report medians and interquartile ranges (the sampling distributions are skewed), coverage of 95% Wald intervals, rejection rates of the J and Wald tests at the 5% level, and the frequency of estimates near the boundary ($\hat{\rho} > 0.99$ or $\hat{v} < 10^{-3}$).

Unit test. Under correct moments the oracle-parameter premium is the best linear predictor. Its exact population MSE (A1) is therefore a lower bound for every linear premium in expectation, and it must match the simulated oracle-parameter MSE within

Monte Carlo error. Table 2 confirms this for all cells. In expectation, the ordering is oracle parameters $\leq$ best exchangeable $\leq$ naive.

Table 2. Exact population quantities and the unit test. Zero: share of zero claims (%). Exact MSE of the naive premium, and exact relative MSE reduction (%) of the oracle-parameter premium (OP) and of the best exchangeable premium (BE) relative to naive. The last two columns compare the exact OP MSE with its simulated value (Monte Carlo SE in parentheses; 500 replications of $I = 2000$).

v	$r(1)$	Zero (%)	Naive MSE	OP gain (%)	BE gain (%)	OP MSE (exact)	OP MSE (simulated)
0.1	0.3	54	446.6	0.01	0.00	446.6	446.8 (2.1)
0.1	0.5	54	446.6	0.04	0.02	446.4	446.4 (2.1)
0.1	0.7	54	446.6	0.12	0.09	446.1	447.7 (2.2)
0.1	0.9	54	446.6	0.32	0.32	445.2	444.5 (2.0)
0.25	0.3	55	492.9	0.07	0.02	492.5	495.6 (2.8)
0.25	0.5	55	492.9	0.21	0.12	491.8	493.3 (2.8)
0.25	0.7	55	492.9	0.56	0.44	490.1	486.4 (2.6)
0.25	0.9	55	492.9	1.48	1.44	485.6	483.3 (2.7)
0.5	0.3	56	566.6	0.20	0.07	565.5	566.1 (3.9)
0.5	0.5	56	566.6	0.64	0.35	563.0	565.1 (3.8)
0.5	0.7	56	566.6	1.59	1.25	557.6	552.5 (3.5)
0.5	0.9	56	566.6	3.93	3.80	544.3	540.8 (3.5)
1	0.3	58	705.3	0.51	0.19	701.7	701.5 (7.3)
1	0.5	58	705.3	1.60	0.89	694.0	690.4 (5.9)
1	0.7	58	705.3	3.81	2.93	678.5	682.3 (6.3)
1	0.9	58	705.3	8.68	8.33	644.1	652.9 (6.3)
2	0.3	60	961.1	1.10	0.43	950.5	930.8 (12.3)
2	0.5	60	961.1	3.36	1.86	928.8	908.3 (10.8)
2	0.7	60	961.1	7.57	5.67	888.4	900.3 (11.5)
2	0.9	60	961.1	15.93	15.14	808.0	813.1 (14.2)

6.2. Study A: Estimation and Tests

Tables 4 and 5 and Figure 3 summarize the estimator. Five findings stand out.

Point estimates. Both parametric fits recover the latent variance and the lag-1 correlation well whenever there is a non-negligible latent signal. In the central cell ($v = 0.5$, $r(1) = 0.7$), the medians are $\hat{v} = 0.50$ (S1) and 0.51 (S2), and $\hat{r}(1) = 0.68$ and 0.67. The S1 working model is misspecified for $k \geq 2$, yet its $\hat{v}$ and $\hat{\rho}$ are almost indistinguishable from the S2 estimates of v and $r(1)$. At $v = 2$, the median S1 estimate of v is 3–13% below the truth (S2: between -14% and $+9\%$), a consequence of the very heavy tails of e_{it}^2 ($\mathbb{E}[\theta^4] \approx 735$). At $v = 0.1$, the signal is weak relative to the nugget ($v/(aq) \approx 0.03$ for a typical policy). Here $\hat{v}$ is biased upward, and $\hat{r}(1)$ is shrunk toward zero when $r(1)$ is small.

Coverage. With $0.25 \leq v \leq 1$ and $r(1) \geq 0.5$, Wald intervals for v and $r(1)$ under S2 cover the truth in 80–97% of replications. Coverage drops to 72–88% in the weak-signal cells ($v = 0.1$) and, for v, in the heavy-tailed cells ($v = 2$). Intervals for the nonparametric $\gamma(1)$, a linear moment, are closer to nominal: 89–95% for $v \leq 1$ and 78–90% at $v = 2$. Table 3 reports a cluster bootstrap of the whole two-stage procedure. In the weak-signal cell it restores near-nominal coverage (94.5% for v and 92% for $r(1)$, against 72% and 78.5% for Wald intervals), and bias correction removes the upward bias of $\hat{v}$ (median 0.10 against 0.15). In the heavy-tailed cell ($v = 2$, $r(1) = 0.9$) it does not help for v: coverage stays near 75%, and the median $\hat{v}$ remains 18–20% below the truth, with or without bias correction, in these 200 replications. There, the sampling distribution of $\hat{v}$ is driven by rare extreme residuals that a resample of 2000 policies does not reproduce. Intervals for $r(1)$ are reliable in all three cells.

Specification test. The J test of the correctly specified S2 has an empirical size of 4–8% for $v \leq 0.5$, rising to 9–10% for $v = 1$ and 12–15% for $v = 2$. The over-rejection at large v again reflects the heavy tails of the squared residuals. The J test of S1 rejects almost exactly

as often as that of S2. With $T = 5$ the test therefore has essentially no power against the gap between θ-scale and log-scale AR(1), which is also inconsequential for prediction (Study B). The J test of the exchangeable S3 has power that increases with v and is highest at moderate $r(1)$.

Static and no-dependence tests. The Wald test of a static latent risk rejects in 8–94% of replications across the grid. Its power is highest for large v and moderate $r(1)$ and lowest at $r(1) = 0.9$, where $\gamma(1), \ldots, \gamma(4)$ are nearly flat over a five-year window. The no-dependence test rejects in 7% of replications in the weakest cell and in 96–100% whenever $v \geq 0.5$ and $r(1) \geq 0.5$.

Boundary. Fits with $\hat{\rho} > 0.99$ or $\hat{v} < 10^{-3}$ occur in up to 28% of replications at $v = 0.1$, but in at most 2% when $v \geq 0.5$ and $r(1) \leq 0.7$. Thanks to the reparametrization, such fits remain usable; they correspond to static-like premiums with a small credibility weight.

Table 3. Cluster bootstrap (policyholder resampling of the full two-stage procedure, $B = 199$, fixed minimum-distance weight matrix) versus delta-method Wald intervals for S2: coverage (%) of nominal 95% intervals for v and $r(1)$, and median $\hat{v}$ without and with bootstrap bias correction. 200 replications per cell.

		Coverage of v		Coverage of $r(1)$		Median $\hat{v}$	
v	$r(1)$	Wald	Bootstrap	Wald	Bootstrap	Plain	Bias-corrected
0.1	0.3	72.0	94.5	78.5	92.0	0.15	0.10
0.5	0.7	93.0	92.5	89.0	92.5	0.51	0.48
2	0.9	71.5	75.5	95.5	96.0	1.65	1.60

Table 4. Study A: sampling distribution of the latent-process estimates, as median [interquartile range] over 500 replications. S1 is the θ-scale AR(1) working model and S2 is the correctly specified log-normal AR(1) model. The truth is $(v, r(1))$ in the first two columns.

v	$r(1)$	S1: $\hat{v}$	S1: $\hat{r}(1) = \hat{\rho}$	S2: $\hat{v}$	S2: $\hat{r}(1)$
0.1	0.3	0.13 [0.04, 0.29]	0.18 [0.03, 0.53]	0.13 [0.04, 0.29]	0.19 [0.03, 0.54]
0.1	0.5	0.15 [0.06, 0.30]	0.33 [0.11, 0.63]	0.15 [0.06, 0.30]	0.34 [0.12, 0.64]
0.1	0.7	0.13 [0.08, 0.25]	0.55 [0.31, 0.86]	0.14 [0.08, 0.25]	0.55 [0.31, 0.86]
0.1	0.9	0.12 [0.08, 0.20]	0.79 [0.58, 1.00]	0.12 [0.09, 0.20]	0.79 [0.58, 1.00]
0.25	0.3	0.27 [0.13, 0.49]	0.27 [0.12, 0.49]	0.27 [0.13, 0.49]	0.26 [0.12, 0.48]
0.25	0.5	0.27 [0.17, 0.45]	0.46 [0.29, 0.63]	0.28 [0.17, 0.46]	0.44 [0.28, 0.62]
0.25	0.7	0.27 [0.18, 0.38]	0.67 [0.50, 0.80]	0.28 [0.19, 0.40]	0.65 [0.48, 0.80]
0.25	0.9	0.26 [0.21, 0.32]	0.88 [0.79, 0.97]	0.26 [0.21, 0.32]	0.87 [0.78, 0.97]
0.5	0.3	0.52 [0.33, 0.80]	0.29 [0.17, 0.42]	0.53 [0.33, 0.80]	0.27 [0.17, 0.41]
0.5	0.5	0.52 [0.37, 0.74]	0.49 [0.35, 0.61]	0.53 [0.38, 0.79]	0.47 [0.32, 0.59]
0.5	0.7	0.50 [0.40, 0.62]	0.68 [0.60, 0.76]	0.51 [0.41, 0.64]	0.67 [0.58, 0.75]
0.5	0.9	0.49 [0.42, 0.58]	0.89 [0.82, 0.94]	0.50 [0.42, 0.58]	0.88 [0.81, 0.95]
1	0.3	1.02 [0.73, 1.47]	0.29 [0.20, 0.39]	1.07 [0.76, 1.54]	0.27 [0.19, 0.36]
1	0.5	0.99 [0.77, 1.24]	0.51 [0.41, 0.59]	1.07 [0.79, 1.33]	0.47 [0.37, 0.56]
1	0.7	0.94 [0.80, 1.14]	0.69 [0.63, 0.76]	0.98 [0.81, 1.20]	0.67 [0.59, 0.75]
1	0.9	0.95 [0.82, 1.10]	0.88 [0.83, 0.93]	0.95 [0.82, 1.11]	0.88 [0.82, 0.93]
2	0.3	1.95 [1.40, 2.68]	0.30 [0.22, 0.40]	2.17 [1.56, 3.01]	0.26 [0.19, 0.34]
2	0.5	1.77 [1.43, 2.26]	0.52 [0.43, 0.59]	1.97 [1.51, 2.63]	0.46 [0.37, 0.55]
2	0.7	1.80 [1.53, 2.15]	0.70 [0.65, 0.75]	1.88 [1.53, 2.30]	0.67 [0.60, 0.74]
2	0.9	1.74 [1.46, 2.03]	0.89 [0.84, 0.94]	1.72 [1.42, 2.02]	0.89 [0.84, 0.95]

Table 5. Study A: inference. Coverage (%) of nominal 95% Wald intervals for v (log scale) and $r(1)$ under S2, and for $\gamma(1)$ under NP. Rejection rates (%) at the 5% level of the J tests of S1, S2 and S3 and of the Wald test of H_0: $\gamma(1) = \cdots = \gamma(T-1)$ (static). Bnd.: share (%) of S1 fits with $\hat{\rho} > 0.99$ or $\hat{v} < 10^{-3}$.

		Coverage (%)			Rejection rate (%)				
v	$r(1)$	v (S2)	$r(1)$ (S2)	$\gamma(1)$ (NP)	J S1	J S2	J S3	Static	Bnd.
0.1	0.3	71.6	81.6	93.2	7.4	7.2	8.8	8.2	20.0
0.1	0.5	79.4	74.6	94.6	5.6	5.6	8.8	8.4	14.8
0.1	0.7	86.0	80.4	94.4	6.4	6.6	10.4	11.2	20.2
0.1	0.9	87.8	84.8	92.8	5.2	4.8	7.2	7.6	27.8
0.25	0.3	86.2	80.0	95.2	6.6	6.8	20.4	18.0	6.0
0.25	0.5	90.8	81.4	93.8	4.4	4.4	25.0	24.2	5.4
0.25	0.7	91.2	89.2	93.8	5.0	5.0	25.0	24.8	7.8
0.25	0.9	94.8	95.2	93.0	6.0	5.8	12.2	10.4	19.8
0.5	0.3	87.6	82.4	93.2	8.0	7.8	38.2	38.0	1.6
0.5	0.5	87.8	80.4	92.2	5.8	6.4	55.0	54.4	0.4
0.5	0.7	93.0	90.0	92.4	5.2	5.8	48.2	50.6	1.0
0.5	0.9	97.2	95.6	94.0	6.0	7.6	13.6	14.2	14.4
1	0.3	85.4	81.4	93.4	9.8	10.4	74.6	71.0	0.4
1	0.5	88.8	85.4	92.8	8.0	9.6	83.2	83.0	0.0
1	0.7	89.0	87.4	89.8	7.2	8.8	71.6	72.4	0.0
1	0.9	91.0	92.4	89.4	7.2	9.2	25.8	24.8	9.4
2	0.3	87.8	79.2	90.2	9.0	12.2	92.6	90.8	0.0
2	0.5	85.4	80.8	85.0	9.2	11.6	93.6	94.2	0.0
2	0.7	82.2	83.0	85.4	8.6	14.0	81.0	83.2	0.6
2	0.9	75.2	92.6	78.4	9.8	14.8	30.8	27.8	7.8

6.3. Study B: Predictive Performance

Table 6 and Figures 1–2 report predictive performance in year $T+1$.

Size of the attainable gain. Even the infeasible oracle-parameter premium reduces the out-of-sample MSE of claim costs only by 0.01–16% (Table 2). Most of the MSE is irreducible Tweedie noise around $\mu_{T+1}\theta_{T+1}$. On the premium-error scale, which is what a pricing actuary controls, the oracle gains are much larger: 1.3–45% (Figure 2). As theory predicts, the gain increases with both v and $r(1)$. With correct moments, the ordering oracle parameters $\leq$ best exchangeable $\leq$ naive holds exactly in every cell.

Feasible premium. The S1 premium with estimated parameters captures 80–97% of the oracle gain whenever $v \geq 0.25$ and $r(1) \geq 0.7$, or $v \geq 0.5$ and $r(1) \geq 0.5$ (Figure 1). In the central cell, for example, S1 reduces the premium error by 10.2% against an oracle value of 11.1%. In those cells it beats the naive premium in 97–100% of replications. When the oracle gain itself is tiny ($v = 0.1$ with $r(1) \leq 0.5$, an oracle MSE gain of at most 0.04%), estimation noise consumes the gain. The S1 premium is then about as good as the naive premium (relative MSE differences within $\pm 0.02\%$), but the share of the gain is not meaningful. S1 and S2 are practically indistinguishable, so the convenient θ-scale AR(1) structure, with the positivity and recursion of Proposition 1, loses nothing against the correctly specified log-normal structure.

Static benchmark. The fitted static premium is never materially worse than the naive premium; its largest loss is 0.03% of MSE. This contradicts the view that pooling past claims can be harmful under serial dependence: that conclusion arises only when the full $\mathrm{Var}\,\theta$ is plugged into an exchangeable formula, whereas a fitted exchangeable model estimates the average covariance. The static premium captures substantially less of the gain than S1 at low and moderate persistence (e.g., 37% versus 84% at $v = 0.5$, $r(1) = 0.5$). At $r(1) = 0.9$, the two are close, because a five-year window of a highly persistent process is nearly exchangeable. Even then, S1 has the smaller premium error in 58–76% of the replications with $v \geq 0.25$.

Variants. Table 7 compares estimator variants. Joint GMM (S1) gains most over the lag-only variant where the lag moments are weak: by 16–28 percentage points of the oracle gain at $r(1) = 0.3$ (for $v \geq 0.25$) and by up to 14 points at $r(1) = 0.5$. There, the diagonal moments carry real information about v. At $r(1) \geq 0.7$ the differences are small (-1 to 8 points). In the baseline design, equal weights lose 0–8 points, because the residuals are already standardized and μ is moderately heterogeneous. In the more heterogeneous design, the precision weights matter more (Section 6.4). The nonparametric plug-in premium is unreliable. Its noisy $\hat{\gamma}(k)$ and extrapolated $\hat{\gamma}(T)$ occasionally produce extreme or negative premiums (up to 567 nonpositive premiums out of 10^6 in a cell), which destroys its mean deviance. The NP estimates are therefore best used for testing, not pricing. Neither S1 (by Proposition 1) nor S2 (empirically) produced a single nonpositive premium. On the deviance scale, S1 improves on the naive premium in every cell with $v \geq 0.25$.

Table 6. Study B: relative reduction (%) of out-of-sample MSE versus the naive premium (paired, mean over 500 replications, with Monte Carlo SE in parentheses), share (%) of the oracle gain captured on the premium-error scale, and the share (%) of replications in which S1 has a smaller premium error than naive. OP: oracle parameters.

		MSE reduction vs naive (%)				Share of oracle gain (%)			S1 beats
v	$r(1)$	Static	S1	S2	OP	Static	S1	S2	naive (%)
0.1	0.3	−0.03 (0.01)	−0.01 (0.01)	−0.01 (0.01)	0.06 (0.01)	−35	−10	−10	67
0.1	0.5	−0.01 (0.01)	0.01 (0.01)	0.01 (0.01)	0.10 (0.01)	−9	15	16	76
0.1	0.7	0.06 (0.01)	0.07 (0.01)	0.08 (0.01)	0.16 (0.01)	32	43	43	88
0.1	0.9	0.29 (0.02)	0.26 (0.02)	0.26 (0.02)	0.36 (0.02)	76	67	68	98
0.25	0.3	−0.03 (0.01)	0.04 (0.01)	0.04 (0.01)	0.10 (0.01)	−13	35	36	77
0.25	0.5	0.04 (0.02)	0.14 (0.02)	0.14 (0.02)	0.21 (0.02)	29	68	69	89
0.25	0.7	0.32 (0.03)	0.41 (0.03)	0.41 (0.03)	0.52 (0.03)	63	80	80	99
0.25	0.9	1.41 (0.05)	1.40 (0.05)	1.40 (0.05)	1.55 (0.05)	92	91	91	100
0.5	0.3	0.00 (0.02)	0.14 (0.02)	0.14 (0.02)	0.22 (0.02)	9	65	66	87
0.5	0.5	0.25 (0.03)	0.57 (0.04)	0.57 (0.04)	0.68 (0.04)	37	84	85	97
0.5	0.7	1.16 (0.06)	1.50 (0.06)	1.50 (0.06)	1.65 (0.06)	71	92	92	100
0.5	0.9	3.39 (0.10)	3.48 (0.09)	3.48 (0.09)	3.65 (0.10)	94	95	95	100
1	0.3	0.09 (0.02)	0.41 (0.03)	0.42 (0.03)	0.50 (0.03)	19	81	83	92
1	0.5	0.69 (0.06)	1.46 (0.06)	1.47 (0.06)	1.57 (0.06)	45	93	93	98
1	0.7	2.80 (0.10)	3.69 (0.11)	3.70 (0.11)	3.89 (0.11)	73	95	95	100
1	0.9	8.20 (0.19)	8.39 (0.19)	8.40 (0.19)	8.65 (0.19)	95	97	97	100
2	0.3	0.32 (0.04)	1.01 (0.06)	1.03 (0.05)	1.15 (0.06)	30	91	92	94
2	0.5	1.62 (0.09)	3.06 (0.12)	3.03 (0.12)	3.26 (0.13)	49	95	94	98
2	0.7	5.29 (0.21)	7.12 (0.23)	7.05 (0.23)	7.47 (0.23)	72	96	95	99
2	0.9	13.60 (0.35)	13.96 (0.34)	14.07 (0.34)	14.53 (0.33)	94	97	98	99




Figure 1. Share of the oracle-parameter gain captured by the feasible S1 premium, on the premium-error scale, over the $(v, r(1))$ grid; 500 replications per cell.




Figure 2. Relative reduction in premium error versus the naive premium, against the lag-1 latent correlation, for each latent variance v. Solid lines with markers: feasible S1 premium; dashed lines: oracle-parameter premium.




Figure 3. Sampling distributions of $\hat{v}/v$ and $\hat{r}(1)$ for S1 and S2 at $r(1) = 0.7$. Boxes show quartiles and whiskers the 5th and 95th percentiles; the dotted line marks the truth.

https://doi.org/10.3390/math1010000

Table 7. Estimator variants: share (%) of the oracle gain captured, and relative reduction (%) of mean Tweedie deviance versus naive. Hetero.: central cell with $\beta_1 = 1.5$ and $\beta_0 = 2$ (more heterogeneous μ).

		Share of oracle gain (%)						Deviance reduction (%)		
v	$r(1)$	Static	S1	S1 lag	S1 equal	S2	NP	Static	S1	NP
0.1	0.7	32	43	39	37	43	−44	0.04	0.04	−0.05
0.25	0.7	63	80	78	77	80	56	0.31	0.36	0.24
0.5	0.7	71	92	90	90	92	81	1.01	1.21	1.09
1	0.7	73	95	92	94	95	88	2.80	3.24	1.96
2	0.7	72	96	88	95	95	< -100	6.04	6.86	−21.84
0.1	0.9	76	67	68	59	68	28	0.22	0.19	0.10
0.25	0.9	92	91	90	88	91	78	1.07	1.05	0.91
0.5	0.9	94	95	94	94	95	87	2.81	2.82	2.66
1	0.9	95	97	94	95	97	80	6.81	6.79	−17.63
2	0.9	94	97	92	96	98	5	12.55	12.35	−79.89
hetero.		58	98	102	75	98	95	1.18	1.47	−1.52

6.4. Sensitivity Analysis

Table 8 varies one design factor at a time around the central cell, and Table 9 reports size and power of the tests.

Sample size and panel length. The estimator behaves as Proposition 2 predicts. With $I = 10{,}000$, S1 captures 98% of the oracle gain, the J test of S2 has a size of 5.6%, and the S2 Wald intervals cover v and $r(1)$ in 94% of replications. With $I = 500$, the share falls to 75%, and the J test over-rejects (12%). A short panel ($T = 3$) reduces the share to 84% and produces more boundary fits (18%), whereas $T = 10$ raises it to 97%. With $T = 10$, the static Wald test rejects in all but 0.4% of replications, and the fitted static premium captures only 43% of the gain, because exchangeability becomes less plausible over a longer window.

Dispersion. Less Tweedie noise ($\phi = 5$) raises the share to 98%, and more noise ($\phi = 20$) lowers it to 75%. The MSE reduction achieved by S1 falls accordingly, from 4.1% to 0.4%.

Misspecified power parameter. The moments depend on p only through the nugget scale $q_{it} = \mu_{it}^{p-2}$. With a true $p = 1.3$ and a working $p = 1.5$, the estimates of (v, ρ) are biased (medians 0.40 and 0.78), and the J test of S2 detects the misspecification in 85% of replications. Prediction is nevertheless robust: S1 still captures 95% of the gain. The reverse error is costlier. With a true $p = 1.7$ and a working $p = 1.5$, part of the Tweedie variance is attributed to v (median $\hat{v} = 2.31$, $\hat{\rho} = 0.16$), the share falls to 58%, and the J test rejects in only 17% of replications. Profiling p by the diagonal J statistic (Section 5.5) locates p correctly on average (median $\hat{p} = 1.35$ and 1.65), but it is imprecise with $I = 2000$ (interquartile range 1.10–1.55 and 1.10–1.90) and recovers only part of the loss. In this design, a working p that is too large is much less harmful than one that is too small.

Unbalanced panels and heterogeneity. With 20% of policy-years missing at random, the share is 87%. In the more heterogeneous design ($\beta_1 = 1.5$), precision weighting matters: the joint GMM captures 98% of the gain, the equal-weight variant only 75%.

Null designs. Under a static latent risk, the static Wald test has a size of 6.4%, and the J test of S3 a size of 6.2%. The fitted static and S1 premiums capture 99% and 97% of the oracle gain, so little is lost by using the evolutionary premium when the truth is static. Under independence, the no-dependence Wald test has a size of 8.8%, and the static Wald test a size of 7.0%. Estimating a dependence structure that is absent costs S1 less than 0.01% of MSE relative to the naive premium (SE 0.003%). The J tests of S1 and S2 over-reject in this design (11.8%) because their true parameters lie on the boundary ($\rho = 0$), where the χ^2 approximation does not apply [31].

Table 8. One-at-a-time sensitivity around the central cell $v = 0.5$, $r(1) = 0.7$ (500 replications each): median S1 estimates, J rejection rate (%) of S2, share (%) of the oracle gain captured, and relative MSE reduction (%) of S1 versus naive (SE in parentheses).

Design	S1 $\hat{v}$	S1 $\hat{\rho}$	J S2	Static share	S1 share	S2 share	S1 MSE red.
Central ($I = 2000$, $T = 5$, $\phi = 10$, $p = 1.5$)	0.50	0.68	5.8	71	92	92	1.50 (0.06)
$I = 500$	0.58	0.59	12.0	63	75	75	1.25 (0.11)
$I = 10{,}000$	0.50	0.70	5.6	72	98	98	1.56 (0.03)
$T = 3$	0.53	0.66	11.2	79	84	85	1.31 (0.06)
$T = 10$	0.48	0.70	7.8	43	97	97	1.59 (0.06)
$\phi = 5$	0.49	0.69	7.0	69	98	98	4.13 (0.09)
$\phi = 20$	0.61	0.60	6.8	63	75	76	0.40 (0.03)
True $p = 1.3$, working $p = 1.5$	0.40	0.78	85.4	69	95	94	3.53 (0.08)
True $p = 1.3$, p profiled (median $\hat{p}$=1.35)		—		66	94	94	3.51 (0.08)
True $p = 1.7$, working $p = 1.5$	2.31	0.16	17.4	62	58	62	0.29 (0.03)
True $p = 1.7$, p profiled (median $\hat{p}$=1.65)		—		50	63	66	0.34 (0.03)
20% of years missing	0.50	0.68	8.2	70	87	88	1.28 (0.05)
Heterogeneous μ ($\beta_1 = 1.5$)	0.48	0.69	6.8	58	98	98	3.52 (0.60)

Table 9. Size and power of the tests (rejection rates in %, 5% level, 500 replications, $v = 0.5$, $I = 2000$, $T = 5$), and share (%) of the oracle gain captured by the fitted static and S1 premiums. Under independence there is no gain to capture.

Truth	Static Wald	No-dep. Wald	J S1	J S2	J S3	Static share	S1 share
Static truth ($\gamma(k) = v$)	6.4	100.0	8.0	10.8	6.2	99	97
Independence ($\gamma(k) = 0, k \geq 1$)	7.0	8.8	11.8	11.8	19.6	—	—
Central design	50.6	99.8	5.2	5.8	48.2	71	92

7. Discussion and Conclusions

This paper develops evolutionary credibility for Tweedie claim costs within a GLM rating structure. Theorem 1 gives the best linear premium for any second-order stationary latent process. It requires only the autocovariance $\gamma(\cdot)$ and a single nugget constant a, and it contains the naive premium and the Tweedie credibility estimator of Ohlsson and Johansson [11] as special cases. For AR(1) dependence, Proposition 1 provides an updating-type recursion with positive weights, a guaranteed positive premium and a bonus cap equal to the persistence parameter. Recency of the weights holds under a mild condition on the evolution of μ_t, but not in general. Because every input of the premium is identified by second moments of standardized residuals, a semi-parametric GMM estimator with closed-form moments, constraints built into the parametrization and stage-1-corrected inference makes the premium operational without distributional assumptions on the latent process.

Practical implications. Four points emerge from the simulation study. First, with estimated parameters the AR(1) premium captures most of the gain that is available to any linear premium, and it never produced a nonpositive premium. Second, a fitted static premium is a safe but conservative choice: it is never materially worse than the naive premium, but it leaves much of the gain unused unless persistence is high. Third, gains should be judged on the premium error, the deviance or rank-based ratemaking metrics rather than on the raw MSE of claim costs, which is dominated by irreducible Tweedie noise. Fourth, the Wald test of a static latent risk has limited power over a five-year window when persistence is very high, but in exactly those cases static and evolutionary premiums nearly coincide, so the cost of a wrong decision is small.

Limitations. The premium is linear and therefore not the exact Bayesian predictor. Its optimality is restricted to linear rules, and higher-order features of the latent law

(e.g., tail dependence) do not enter it. Second-moment inference is sensitive to the heavy tails of squared Tweedie residuals: when the latent variance is large, Wald intervals for v under-cover and the J test over-rejects. The policyholder-level bootstrap repairs the under-coverage caused by a weak signal, but not the under-coverage caused by heavy tails. Robust (e.g., trimmed or median-based) moment conditions are a natural remedy. Prediction is much less affected than inference. With $T = 5$, the J test cannot discriminate between θ-scale and log-scale AR(1) dependence, although this distinction turned out to be immaterial for prediction. The power parameter p must be chosen or profiled, and we have not yet applied the method to real data.

Extensions. Natural next steps are as follows:

- an application to a longitudinal policy-level portfolio with five or more years, such as the Wisconsin Local Government Property Insurance Fund data used by Frees et al. [21];
- robustness studies with latent processes that share $\gamma(\cdot)$ but differ in higher-order dependence, for example gamma autoregressions or copula-based Markov chains with tail dependence;
- an exact-Bayes benchmark by particle filtering, to quantify the cost of linearity;
- comparisons with machine-learning predictors that use lagged claims, such as gradient boosting with Tweedie loss, evaluated with ordered Lorenz curves and Gini indices;
- hierarchical [19], higher-order and long-memory [18] latent structures, all of which fit Theorem 1 and the GMM framework of Section 5 without modification.

Author Contributions: [To be completed by the authors.]

Funding: [To be completed by the authors.]

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: No new data were created. The simulation code (Python) that reproduces all tables and figures is available in the authors' repository [link to be added].

Acknowledgments: This article is based on the first author's MSc project at Simon Fraser University [25], supervised by the second author. [GenAI statement to be confirmed by the authors: During the preparation of this manuscript, the authors used Claude (Anthropic) to assist with drafting text, implementing the simulation code and preparing tables and figures. The authors have reviewed and edited the output and take full responsibility for the content of this publication.]

Conflicts of Interest: The authors declare no conflicts of interest.

Abbreviations

The following abbreviations are used in this manuscript:

AR(1)	first-order autoregressive
BLP	best linear predictor
GLM	generalized linear model
GMM	generalized method of moments
MD	minimum distance
MSE	mean squared error
NP	nonparametric (autocovariance)

Appendix A. Proofs

Proof of Proposition 1. *(a)* Let $\hat{\xi}_t$ be the best linear predictor of ξ_t given $(1, e_1, \ldots, e_{t-1})$, and let V_t be its MSE. Then $\hat{\xi}_1 = \mathbb{E}[\xi_1] = 0$ and $V_1 = v$. The innovation $e_t - \hat{\xi}_t = (\xi_t - \hat{\xi}_t) +$

ε_t has variance $V_t + d_t$, because ε_t is uncorrelated with ξ_t and with the past. Projecting ξ_t on the innovation gives $\hat{\xi}_{t|t} = \hat{\xi}_t + K_t(e_t - \hat{\xi}_t)$, with MSE $(1 - K_t)V_t$. Since $\zeta_t = \xi_{t+1} - \rho\xi_t$ is uncorrelated with $(e_1, \ldots, e_t)$, we get $\hat{\xi}_{t+1} = \rho\hat{\xi}_{t|t}$ and $V_{t+1} = \rho^2(1 - K_t)V_t + \mathrm{Var}(\zeta_t)$. Finally, ε_{T+1} is uncorrelated with $(e_1, \ldots, e_T)$, so the best linear predictor of e_{T+1} is $\hat{\xi}_{T+1}$. The best linear predictor is unique, so it coincides with (3). See Brockwell and Davis [27, Proposition 12.2.2] for the general wide-sense Kalman recursions.

(b) Unrolling $\hat{\xi}_{t+1} = \rho(1 - K_t)\hat{\xi}_t + \rho K_t e_t$ from $\hat{\xi}_1 = 0$ gives the stated coefficients. Since $V_1 = v > 0$, $V_{t+1} \geq (1 - \rho^2)v > 0$ and $0 < d_t < \infty$, we have $K_t \in (0, 1)$, so $w_t > 0$ when $\rho > 0$.

(c) Summing the coefficients of $\hat{\xi}_{t+1}$ gives $S_t = \rho(1 - K_t)S_{t-1} + \rho K_t$. If $0 \leq S_{t-1} < 1$, then $(1 - K_t)S_{t-1} + K_t = 1 - (1 - K_t)(1 - S_{t-1}) \in (0, 1)$, hence $0 < S_t < \rho$. Induction from $S_0 = 0$ completes the proof.

(d) Since $Z_t \geq 0$, we have $e_t \geq -1$. By (b), $\sum_t w_t e_t \geq -S_T$, with equality if and only if $Z_1 = \cdots = Z_T = 0$. The bound follows from (c).

(e) Let $\lambda = 1/v$, $u_t = V_t/v$ and $\delta_t = \lambda d_t$. Then $u_1 = 1$, $K_t = u_t/(u_t + \delta_t)$ and $u_{t+1} = \rho^2 h(u_t, \delta_t) + 1 - \rho^2$ with $h(x, y) = xy/(x + y)$. We show by induction that u_t/λ is nonincreasing in λ. This is clear for $t = 1$. Since h is homogeneous of degree one, $u_{t+1}/\lambda = \rho^2 h(u_t/\lambda, d_t) + (1 - \rho^2)/\lambda$. The function h is nondecreasing in its first argument, so the claim follows. Hence $K_t = \{1 + d_t/(u_t/\lambda)\}^{-1}$ is nonincreasing in λ, i.e., nondecreasing in v, and so is $w_T = \rho K_T$. Finally, $S_t = \rho\{S_{t-1} + K_t(1 - S_{t-1})\}$ is nondecreasing in both S_{t-1} and K_t because $S_{t-1} < 1$, so S_T is nondecreasing in v. Replacing every μ_t by $c\mu_t$ multiplies every d_t by c^{p-2}, which is equivalent to replacing v by $c^{2-p}v$.

(f) From (b), $w_{t-1}/w_t = \rho K_{t-1}(1 - K_t)/K_t = \rho K_{t-1}\delta_t/u_t$. Moreover, $(1 - K_{t-1})u_{t-1} = K_{t-1}\delta_{t-1}$, so $u_t = \rho^2 K_{t-1}\delta_{t-1} + 1 - \rho^2$. Hence $w_{t-1} < w_t$ if and only if

$$\rho K_{t-1}\{(1 - \rho)\delta_{t-1} + (\delta_t - \delta_{t-1})\} < 1 - \rho^2.$$

By induction, $u_t \leq 1$ for all t, since $u_{t+1} = \rho^2 h(u_t, \delta_t) + 1 - \rho^2$ and $h(u_t, \delta_t) < u_t$. Therefore $K_{t-1}\delta_{t-1} = h(u_{t-1}, \delta_{t-1}) < 1$ and $K_{t-1} < 1$, and the left-hand side is smaller than $\rho(1 - \rho) + \rho(\delta_t - \delta_{t-1})^+$. This is at most $1 - \rho^2$ whenever $(\delta_t - \delta_{t-1})^+ \leq (1 - \rho)/\rho$, that is, $d_t - d_{t-1} \leq v(1 - \rho)/\rho$. The counterexample in the main text violates this condition.

(g) As $\rho \to 0$, $\boldsymbol{c}_T \to \mathbf{0}$ while $\Gamma_T + D_T \to vI + D_T$ stays nonsingular. As $\rho \to 1$, $\Gamma_T \to v\mathbf{1}\mathbf{1}^\top$ and $\boldsymbol{c}_T \to v\mathbf{1}$, and $(\Gamma_T + D_T)^{-1}$ is continuous because the limit is positive definite. With $d_t \equiv d$, the map $f(u) = \rho^2 h(u, \delta) + 1 - \rho^2$ satisfies $0 < f'(u) = \rho^2\delta^2/(u + \delta)^2 < \rho^2$, so u_t converges to the unique fixed point u^*. The fixed point solves $u^2 + (1 - \rho^2)(\delta - 1)u - (1 - \rho^2)\delta = 0$, and u^* is its positive root. The limiting weights follow from (b) with $K_s \to K^*$, and the sum converges by dominated convergence, since $w_{T+1-j} \leq \rho^j$.

Appendix B. Exact Mean Squared Error of Linear Premiums

Let $P = \mu_{T+1}(1 + \boldsymbol{w}^\top \boldsymbol{e}_T)$, where $\boldsymbol{w}$ is any weight vector that depends on covariates only. Then $Z_{T+1} - P = \mu_{T+1}(e_{T+1} - \boldsymbol{w}^\top \boldsymbol{e}_T)$, and Lemma 1 gives

$$\mathbb{E}[(Z_{T+1} - P)^2 \mid \boldsymbol{x}] = \mu_{T+1}^2\{v + a\mu_{T+1}^{p-2} - 2\boldsymbol{w}^\top \boldsymbol{c}_T + \boldsymbol{w}^\top(\Gamma_T + D_T)\boldsymbol{w}\}. \tag{A1}$$

Averaging (A1) over the covariate distribution (here over 2×10^6 simulated policies) gives the exact population MSE of the naive premium ($\boldsymbol{w} = \mathbf{0}$), of the oracle-parameter premium ($\boldsymbol{w} = (\Gamma_T + D_T)^{-1}\boldsymbol{c}_T$) and of every exchangeable premium. The *best exchangeable* premium uses the true a and $\boldsymbol{\mu}$ and chooses the static constant v_s in Corollary 1(ii) to minimize this population MSE under the true $\gamma(\cdot)$; v_s generally differs from v and acts as an average

covariance across lags. Moreover, if P is any function of the training data, then $\mathbb{E}[Z_{T+1} - \mu_{T+1}\theta_{T+1} \mid \boldsymbol{\theta}, \mathcal{F}_T] = 0$ implies

$$\mathbb{E}[(Z_{T+1} - P)^2] = \mathbb{E}[(\mu_{T+1}\theta_{T+1} - P)^2] + \mathbb{E}\big[a\mu_{T+1}^{p}\big].$$

Differences in out-of-sample MSE between premiums therefore equal differences in the *premium error* $\mathbb{E}[(\mu_{T+1}\theta_{T+1} - P)^2]$. The premium error excludes the irreducible Tweedie noise and has a much smaller Monte Carlo variance.

Appendix C. Computational Details

Tweedie variates are generated from the compound Poisson–gamma representation: $N \sim \text{Poisson}(\mu^{2-p}/\{\phi(2-p)\})$, and given N, $Z \sim \text{Gamma}(N(2-p)/(p-1),\ \phi(p-1)\mu^{p-1})$, with $Z = 0$ if $N = 0$. Stage 1 uses Fisher scoring for the Tweedie quasi-score. Stage 2 starts from equal weights and recomputes the precision weights three times from $(\hat{\gamma}(0), \hat{a})$, floored at 0.02. G_β is obtained by central differences with step 10^{-5} and the weights held fixed. The minimum-distance problems (10) are solved by BFGS from three starting values derived from the NP estimates. The Proposition 1 recursion agrees with the direct solve of (3) to 10^{-11}, and the simulated MSE of the oracle-parameter premium agrees with (A1) within Monte Carlo error. The code is organized as `tweedie_gmm.py` (estimator, tests and premiums), `sim.py` (simulation design), `test_core.py` (unit tests) and `make_tables.py` (tables and figures).

References

1. Jørgensen, B. Exponential Dispersion Models. *Journal of the Royal Statistical Society: Series B* **1987**, *49*, 127–162.
2. Jørgensen, B.; de Souza, M.C.P. Fitting Tweedie's Compound Poisson Model to Insurance Claims Data. *Scandinavian Actuarial Journal* **1994**, *1994*, 69–93.
3. Smyth, G.K.; Jørgensen, B. Fitting Tweedie's Compound Poisson Model to Insurance Claims Data: Dispersion Modelling. *ASTIN Bulletin* **2002**, *32*, 143–157.
4. Klugman, S.A.; Panjer, H.H.; Willmot, G.E. *Loss Models: From Data to Decisions*, 5th ed.; Wiley: Hoboken, NJ, 2019.
5. Bühlmann, H.; Gisler, A. *A Course in Credibility Theory and Its Applications*; Springer: Berlin, 2005.
6. Gerber, H.U.; Jones, D.A. Credibility Formulas of the Updating Type. *Transactions of the Society of Actuaries* **1975**, *27*, 31–52.
7. Sundt, B. Recursive Credibility Estimation. *Scandinavian Actuarial Journal* **1981**, *1981*, 3–21.
8. Albrecht, P. An Evolutionary Credibility Model for Claim Numbers. *ASTIN Bulletin* **1985**, *15*, 1–17.
9. Pinquet, J.; Guillén, M.; Bolancé, C. Allowance for the Age of Claims in Bonus-Malus Systems. *ASTIN Bulletin* **2001**, *31*, 337–348.
10. Bolancé, C.; Guillén, M.; Pinquet, J. Time-Varying Credibility for Frequency Risk Models: Estimation and Tests for Autoregressive Specifications on the Random Effects. *Insurance: Mathematics and Economics* **2003**, *33*, 273–282.
11. Ohlsson, E.; Johansson, B. Exact Credibility and Tweedie Models. *ASTIN Bulletin* **2006**, *36*, 121–133.
12. Hansen, L.P. Large Sample Properties of Generalized Method of Moments Estimators. *Econometrica* **1982**, *50*, 1029–1054.
13. Jewell, W.S. Credible Means Are Exact Bayesian for Exponential Families. *ASTIN Bulletin* **1974**, *8*, 77–90.
14. Frangos, N.E.; Vrontos, S.D. Design of Optimal Bonus-Malus Systems with a Frequency and a Severity Component on an Individual Basis in Automobile Insurance. *ASTIN Bulletin* **2001**, *31*, 1–22.
15. Ma, R.; Yan, G.; Hasan, M.T. Tweedie Family of Generalized Linear Models with Distribution-Free Random Effects for Skewed Longitudinal Data. *Statistics in Medicine* **2018**, *37*, 3519–3532.
16. Jeong, H. Tweedie Multivariate Semi-Parametric Credibility with the Exchangeable Correlation. *Insurance: Mathematics and Economics* **2024**, *115*, 13–21.
17. Ahn, J.Y.; Jeong, H.; Lu, Y. On the Ordering of Credibility Factors. *Insurance: Mathematics and Economics* **2021**, *101*, 626–638.
18. Pinquet, J. Positivity Properties of the ARFIMA(0,d,0) Specifications and Credibility Analysis of Frequency Risks. *Insurance: Mathematics and Economics* **2020**, *95*, 159–165.
19. Taylor, G. Evolutionary Hierarchical Credibility. *ASTIN Bulletin* **2018**, *48*, 339–374.
20. Shi, P.; Feng, X.; Boucher, J.P. Multilevel Modeling of Insurance Claims Using Copulas. *The Annals of Applied Statistics* **2016**, *10*, 834–863.
21. Frees, E.W.; Lee, G.; Yang, L. Multivariate Frequency-Severity Regression Models in Insurance. *Risks* **2016**, *4*, 4.

22. Furman, E.; Landsman, Z. Multivariate Tweedie Distributions and Some Related Capital-at-Risk Analyses. *Insurance: Mathematics and Economics* **2010**, *46*, 351–361.
23. Avanzi, B.; Taylor, G.; Vu, P.A.; Wong, B. Stochastic Loss Reserving with Dependence: A Flexible Multivariate Tweedie Approach. *Insurance: Mathematics and Economics* **2016**, *71*, 63–78.
24. Alai, D.H.; Landsman, Z.; Sherris, M. Multivariate Tweedie Lifetimes: The Impact of Dependence. *Scandinavian Actuarial Journal* **2016**, *2016*, 692–712.
25. Kang, N. Tweedie Evolutionary Credibility for Insurance Pricing with AR(1) Latent Dependence. Msc project, Simon Fraser University, Burnaby, BC, Canada, 2026.
26. Gaver, D.P.; Lewis, P.A.W. First-Order Autoregressive Gamma Sequences and Point Processes. *Advances in Applied Probability* **1980**, *12*, 727–745.
27. Brockwell, P.J.; Davis, R.A. *Time Series: Theory and Methods*, 2nd ed.; Springer: New York, 1991.
28. Harvey, A.C. *Forecasting, Structural Time Series Models and the Kalman Filter*; Cambridge University Press: Cambridge, 1989.
29. Wedderburn, R.W.M. Quasi-Likelihood Functions, Generalized Linear Models, and the Gauss–Newton Method. *Biometrika* **1974**, *61*, 439–447.
30. Newey, W.K.; McFadden, D. Large Sample Estimation and Hypothesis Testing. In *Handbook of Econometrics*; Engle, R.F.; McFadden, D.L., Eds.; Elsevier: Amsterdam, 1994; Vol. 4, pp. 2111–2245.
31. Andrews, D.W.K. Estimation When a Parameter Is on a Boundary. *Econometrica* **1999**, *67*, 1341–1383.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.