


A


Contact map · 3 models · sum of binarized contacts
78 residues · 45 contacts in ≥2 models
Chain B
Chain A
High
Medium
Low
Fraction of models in contact


B


Complex stability
abin
model
Fraction buried
dASA (Å²)


C


Burial vs occupancy · 3 models
78 residues · 45 contacts in ≥2 models
H-bond
Salt bridge
Hydrophobic
A861 LYS
B454 ASN
B585 ASP
A758 PRO
B422 PRO
A919 THR
B601 LEU
Models
2/3
3/3
Mean relative dSASA
Contact occupancy (≤8 Å)

D


Residue occupancy · 3 models
78 residues · 45 contacts in ≥2 models
H-bond
Salt bridge
Hydrophobic
Contact
A883 GLY
A886 ILE
A885 HIS
A887 VAL
A884 THR
A926 ARG
A861 LYS
A757 ASP
B455 ASP
B529 SER
B513 PHE
B530 GLY
B458 TYR
B532 PRO
B454 ASN
B585 ASP

E


Residue-pair occupancy · 3 models · 1/2
78 residues · 45 contacts in ≥2 models
A861 LYS – B584 ASP
A862 GLN – B585 ASP
A861 LYS – B585 ASP
A861 LYS – B586 GLU
A861 LYS – B424 ILE
A887 VAL – B454 ASN
Contact occupancy (≤8 Å)
Residue-pair occupancy · 3 models · 2/2
78 residues · 45 contacts in ≥2 models
A861 LYS – B420 PRO
A861 LYS – B422 PRO
A861 LYS – B425 PRO
A920 ALA – B454 ASN
A858 ARG – B585 ASP
A859 THR – B586 GLU
Contact occupancy (≤8 Å)
H-bond
Salt bridge
Hydrophobic
Contact ≤8 Å