


A


Contact map · 11 models · sum of binarized contacts
309 residues · 631 contacts in ≥2 models
Chain B
Chain A
High
Medium
Low
Fraction of models in contact


B


Complex stability
abin
model
pdbeval
Fraction buried
dASA (Å²)


C


Burial vs occupancy · 11 models
309 residues · 631 contacts in ≥2 models
H-bond
Salt bridge
Hydrophobic
B446 VAL
B484 ARG
B543 LYS
B445 HIS
B434 PHE
B438 GLU
A200 ASP
B431 ARG
B443 ARG
Models
2/11
6/11
9/11
Mean relative dSASA
Contact occupancy (≤8 Å)

D


Residue occupancy · 11 models
309 residues · 631 contacts in ≥2 models
H-bond
Salt bridge
Hydrophobic
Contact
A356 ARG
A148 ARG
A200 ASP
A138 TYR
A142 GLN
A194 HIS
A195 LEU
A197 GLY
B438 GLU
B437 SER
B443 ARG
B435 GLN
B490 LYS
B491 ARG
B452 ARG
B439 SER

E


Residue-pair occupancy · 11 models · 1/2
309 residues · 631 contacts in ≥2 models
A363 ASP – B484 ARG
A150 GLU – B443 ARG
A45 GLU – B431 ARG
A157 GLU – B445 HIS
A362 LYS – B484 ARG
A371 GLN – B448 GLY
Contact occupancy (≤8 Å)
H-bond
Salt bridge
Hydrophobic
Contact ≤8 Å

Residue-pair occupancy · 11 models · 2/2
309 residues · 631 contacts in ≥2 models
A425 ASP – B543 LYS
A428 GLU – B543 LYS
A150 GLU – B450 HIS
A151 GLU – B443 ARG
A96 CYS – B431 ARG
A367 ARG – B482 GLY
Contact occupancy (≤8 Å)