


A


Contact map · 14 models · sum of binarized contacts
139 residues · 560 contacts in ≥2 models
Chain B
Chain A
High
Medium
Low
Fraction of models in contact


B


Complex stability
abin
model
pdbeval
Fraction buried
dASA (Å²)


C


Burial vs occupancy · 14 models
139 residues · 560 contacts in ≥2 models
H-bond
Salt bridge
Hydrophobic
A200 ASP
B50 LYS
B21 ARG
B65 LEU
A194 HIS
Models
2/14
8/14
14/14
Mean relative dSASA
Contact occupancy (≤8 Å)

D


Residue occupancy · 14 models
139 residues · 560 contacts in ≥2 models
H-bond
Salt bridge
Hydrophobic
Contact
A200 ASP 79% 79% · 100%
A348 GLU 79% 79% · 100%
A344 ASP 43% 7% · 100%
A352 ASN 50% · · 100%
A203 LEU 50% · 7% 100%
A201 LEU 14% · 14% 100%
A345 PHE 14% · · 100%
A194 HIS 86% · · 93%
B67 ASP · · · 93%
B48 ARG 43% 7% · 79%
B62 GLU 29% 7% · 71%
B64 ARG 29% 29% · 64%
B47 TYR · · · 64%
B65 LEU 29% · 29% 57%
B66 ASN 36% · · 57%
B2 TRP 29% · 50% 57%

E


Residue-pair occupancy · 14 models · 1/2
139 residues · 560 contacts in ≥2 models
A352 ASN – B67 ASP 57%
A348 GLU – B48 ARG 43%
A200 ASP – B62 GLU 43%
A352 ASN – B65 LEU 43%
A348 GLU – B67 ASP 43%
A200 ASP – B21 ARG 36%
Contact occupancy (≤8 Å)
H-bond
Salt bridge
Hydrophobic
Contact ≤8 Å

Residue-pair occupancy · 14 models · 2/2
139 residues · 560 contacts in ≥2 models
A200 ASP – B50 LYS 36%
A348 GLU – B64 ARG 36%
A352 ASN – B48 ARG 36%
A352 ASN – B66 ASN 36%
A351 GLU – B48 ARG 36%
A351 GLU – B66 ASN 36%
Contact occupancy (≤8 Å)