


A


Contact map · 5 models · sum of binarized contacts
84 residues · 173 contacts in ≥2 models
Chain B
Chain A
High
Medium
Low
Fraction of models in contact


B


Complex stability
model
Fraction buried
dASA (Å²)


C


Burial vs occupancy · 5 models
84 residues · 173 contacts in ≥2 models
H-bond
Salt bridge
Hydrophobic
B366 ASP
B335 GLU
A43 ARG
B355 GLU
B365 ASN
A150 GLU
A84 ARG
Models
2/5
4/5
5/5
Mean relative dSASA
Contact occupancy (≤8 Å)

D


Residue occupancy · 5 models
84 residues · 173 contacts in ≥2 models
H-bond
Salt bridge
Hydrophobic
Contact
A43 ARG
A150 GLU
A84 ARG
A154 LEU
A36 LYS
A151 GLU
A40 ARG
A158 ARG
B366 ASP
B335 GLU
B365 ASN
B364 ARG
B359 TYR
B357 GLU
B336 CYS
B334 ASP

E


Residue-pair occupancy · 5 models · 1/2
84 residues · 173 contacts in ≥2 models
A84 ARG – B335 GLU
A43 ARG – B335 GLU
A84 ARG – B359 TYR
A84 ARG – B366 ASP
A84 ARG – B363 CYS
A39 ILE – B335 GLU
Contact occupancy (≤8 Å)
H-bond
Salt bridge
Hydrophobic
Contact ≤8 Å

Residue-pair occupancy · 5 models · 2/2
84 residues · 173 contacts in ≥2 models
A40 ARG – B335 GLU
A43 ARG – B336 CYS
A43 ARG – B359 TYR
A43 ARG – B363 CYS
A35 VAL – B335 GLU
A36 LYS – B335 GLU
Contact occupancy (≤8 Å)