


A


Contact map · 11 models · sum of binarized contacts
152 residues · 318 contacts in ≥2 models
Chain B
Chain A
High
Medium
Low
Fraction of models in contact


B


Complex stability
abin
model
pdbeval
Fraction buried
dASA (Å²)


C


Burial vs occupancy · 11 models
152 residues · 318 contacts in ≥2 models
H-bond
Salt bridge
Hydrophobic
B787 PRO
B791 ASN
A200 ASP
A138 TYR
Models
2/11
7/11
11/11
Mean relative dSASA
Contact occupancy (≤8 Å)

D


Residue occupancy · 11 models
152 residues · 318 contacts in ≥2 models
H-bond
Salt bridge
Hydrophobic
Contact
A194 HIS 73% · · 100%
A344 ASP 64% · · 100%
A348 GLU 36% 9% · 100%
A191 TRP 9% · 18% 100%
A200 ASP 82% 18% · 91%
A138 TYR 64% · 55% 91%
A195 LEU 55% · · 91%
A142 GLN 55% · · 91%
B787 PRO 36% · 36% 64%
B783 ASN 36% · · 64%
B780 THR 36% · · 64%
B784 GLN 18% · · 64%
B679 GLN 9% · · 64%
B683 LEU 9% · · 64%
B785 LEU · · · 64%
B791 ASN 27% · · 55%

E


Residue-pair occupancy · 11 models · 1/2
152 residues · 318 contacts in ≥2 models
A200 ASP – B783 ASN 45%
A194 HIS – B776 GLN 36%
A194 HIS – B711 PHE 36%
A127 GLU – B679 GLN 36%
A195 LEU – B776 GLN 36%
A200 ASP – B711 PHE 36%
Contact occupancy (≤8 Å)
H-bond
Salt bridge
Hydrophobic
Contact ≤8 Å

Residue-pair occupancy · 11 models · 2/2
152 residues · 318 contacts in ≥2 models
A197 GLY – B711 PHE 36%
A199 ASP – B711 PHE 36%
A200 ASP – B680 GLN 36%
A194 HIS – B779 GLN 36%
A194 HIS – B780 THR 36%
A194 HIS – B784 GLN 36%
Contact occupancy (≤8 Å)