


A


Contact map · 12 models · sum of binarized contacts
142 residues · 395 contacts in ≥2 models
Chain B
Chain A
High
Medium
Low
Fraction of models in contact


B


Complex stability
abin
model
pdbeval
Fraction buried
dASA (Å²)


C


Burial vs occupancy · 12 models
142 residues · 395 contacts in ≥2 models
H-bond
Salt bridge
Hydrophobic
B445 HIS
A1078 THR
A1052 GLU
A1049 GLU
A1056 GLU
A1053 GLU
B446 VAL
Models
2/12
7/12
12/12
Mean relative dSASA
Contact occupancy (≤8 Å)

D


Residue occupancy · 12 models
142 residues · 395 contacts in ≥2 models
H-bond
Salt bridge
Hydrophobic
Contact
A1049 GLU 58% 33% · 100%
A1056 GLU 75% 33% · 100%
A1053 GLU 92% 42% · 100%
A1050 GLU 25% · · 100%
A1057 ASN 33% · · 100%
A1055 MET · · · 100%
A1052 GLU 83% 17% · 92%
A1051 ASN · · · 92%
B445 HIS 83% · · 83%
B446 VAL 75% · 8% 83%
B491 ARG 58% 33% · 83%
B434 VAL 50% · 42% 83%
B447 ALA 50% · · 83%
B438 GLU 25% · · 83%
B443 ARG 17% 17% · 83%
B444 PRO 42% · · 83%

E


Residue-pair occupancy · 12 models · 1/2
142 residues · 395 contacts in ≥2 models
A1053 GLU – B446 VAL 75%
A1052 GLU – B446 VAL 67%
A1049 GLU – B446 VAL 67%
A1049 GLU – B491 ARG 67%
A1050 GLU – B446 VAL 67%
A1056 GLU – B445 HIS 58%
Residue-pair occupancy · 12 models · 2/2
142 residues · 395 contacts in ≥2 models
A1049 GLU – B445 HIS 58%
A1052 GLU – B445 HIS 58%
A1079 PHE – B434 VAL 58%
A1048 THR – B445 HIS 58%
A1052 GLU – B433 ARG 58%
A1046 PRO – B445 HIS 58%
Contact occupancy (≤8 Å)
H-bond
Salt bridge
Hydrophobic
Contact ≤8 Å