
a
Primary analysis and drug-specific comparisons with SGLT2 inhibitors, type 2 diabetes
b
Tirzepatide vs semaglutide
Substance use
Alcohol use disorder
Opioid use disorder
Cannabis use disorder
Cocaine use disorder
Other stimulant use disorder
Nicotine dependence
Psychosis and bipolar
Schizophrenia
Bipolar disorder
Suicidality
Suicidal ideation
Suicide attempt
Depression, anxiety, stress, sleep
Major depressive disorder, single episode
Major depressive disorder, recurrent
Anxiety disorders
Stress-related and adjustment disorders
Insomnia
Eating, impulse, compulsive
Bulimia nervosa
Anorexia nervosa
Pathological gambling
Obsessive-compulsive disorder
1.00
0.88
0.88
0.85
0.78
0.90
0.75
0.95
0.87
0.86
0.92
0.91
0.99
0.93
0.97
0.58
0.58
1.03
0.25
0.5
1
2
3
Calibrated hazard ratio (95% CI)
Calibrated HR (95% CI)
Semaglutide or tirzepatide (primary)
Semaglutide
Tirzepatide
q ≥ 0.05
c
Semaglutide
Tirzepatide
SGLT2 inhibitor
New diagnoses per 1,000 (crude, 3 years)
10
1
Cocaine
Stimulant
Cannabis
Opioid
Nicotine
Alcohol
Schizophrenia
Bipolar
Suicidal ideation
Suicide attempt
d
r = 0.92
HR, semaglutide vs DPP-4i
HR, semaglutide vs SGLT2i
2
1
0.5
0.5
1
2