
Semaglutide vs no drug (obesity+OSA): uncalibrated and calibrated estimates
Calibrated HR (95% CI)
Substance use
Alcohol use disorder
1.03 (0.67–1.60)
Opioid use disorder
0.71 (0.46–1.09)
Cannabis use disorder
0.69 (0.44–1.06)
Cocaine use disorder
0.58 (0.36–0.92)
Other stimulant use disorder
0.64 (0.41–1.01)
Nicotine dependence
0.95 (0.62–1.46)
Psychosis and bipolar
Schizophrenia
0.46 (0.28–0.76)*
Bipolar disorder
0.71 (0.46–1.10)
Suicidality
Suicidal ideation
0.46 (0.29–0.71)*
Suicide attempt
0.38 (0.23–0.62)*
Depression, anxiety, stress, sleep
Major depressive disorder, single episode
0.82 (0.54–1.27)
Major depressive disorder, recurrent
0.91 (0.59–1.40)
Anxiety disorders
1.08 (0.70–1.66)
Stress-related and adjustment disorders
0.97 (0.63–1.48)
Insomnia
0.98 (0.64–1.51)
Eating, impulse and compulsive
Bulimia nervosa
0.70 (0.41–1.19)
Anorexia nervosa
0.64 (0.34–1.20)
Pathological gambling
1.00 (0.47–2.13)
Obsessive-compulsive disorder
0.91 (0.59–1.42)
Negative controls
Ingrowing nail
0.90 (0.84–0.97)
Otitis externa
1.04 (0.99–1.10)
Allergic rhinitis
0.96 (0.93–0.99)
Melanocytic naevi
1.20 (1.16–1.25)
Forearm fracture
0.75 (0.69–0.82)
Refraction disorders
0.67 (0.63–0.70)
Uncalibrated HR
Calibrated HR
0.25
0.5
1
2
4
Hazard ratio (95% CI), log scale