
E-values for significant inverse associations
Semaglutide vs SGLT2i
Cocaine use disorder
Other stimulant use disorder
Suicide attempt
Schizophrenia
Suicidal ideation
Cannabis use disorder
Alcohol use disorder
Bipolar disorder
Nicotine dependence
Opioid use disorder
Tirzepatide vs SGLT2i
Cocaine use disorder
Other stimulant use disorder
Schizophrenia
Suicide attempt
Suicidal ideation
Cannabis use disorder
Opioid use disorder
Bipolar disorder
Nicotine dependence
Semaglutide vs DPP-4i
Suicide attempt
Schizophrenia
Other stimulant use disorder
Cocaine use disorder
Suicidal ideation
Bipolar disorder
Cannabis use disorder
Opioid use disorder
Nicotine dependence
Major depressive disorder, single episode
Semaglutide vs no drug (T2D)
Cocaine use disorder
Schizophrenia
Other stimulant use disorder
Suicide attempt
Suicidal ideation
Cannabis use disorder
Tirzepatide vs no drug (T2D)
Schizophrenia
Cocaine use disorder
Suicide attempt
Other stimulant use disorder
Suicidal ideation
Opioid use disorder
Cannabis use disorder
Semaglutide vs no drug (obesity+OSA)
Suicide attempt
Suicidal ideation
Schizophrenia
Tirzepatide vs no drug (obesity+OSA)
Suicide attempt
Other stimulant use disorder
Schizophrenia
Suicidal ideation
Cocaine use disorder
Opioid use disorder
1.0
1.5
2.0
2.5
3.0
3.5
4.0
4.5
5.0
5.5
E-value (dark: CI limit; light: point estimate)