

 [Topics](#) [More](#) [CATEGORIES](#)[General](#)[Site Feedback](#) [All categories](#)

2025 USA-NA-AIO Round 1, Problem 1, Part 5

USAAIO 

Mar 2025

Part 5 (10 points, non-coding task)

Let us go back to our matrix A .

In the following eigenvalue equation

$$Ax = \lambda x,$$

compute two eigenvalues λ_0 and λ_1 whose values are in a descending order.

- Reasoning is required.

USAAIO 

Mar 2025

Misplaced '#'

We solve the following characteristic equation:

$$\det(A - \lambda I) = 0.$$

Thus, we compute λ that satisfies

$$(1 - \lambda)(-\lambda) - 1 \cdot 1 = 0.$$

Hence, all eigenvalues are

[Skip to main content](#)

$$\lambda_0 = \frac{1 + \sqrt{5}}{2}, \quad \lambda_1 = \frac{1 - \sqrt{5}}{2}.$$

"" END OF THIS PART ""

✦ Related topics

Topic	Replies	Activity
2025 USA-NA-AIO Round 1, Problem 1, Part 6	1	Mar 2025
2025 USA-NA-AIO Round 1, Problem 1, Part 2	1	Mar 2025
2025 USA-NA-AIO Round 1, Problem 1, Part 7	1	Mar 2025
2025 USA-NA-AIO Round 1, Problem 1, Part 8	3	22d
2025 USA-NA-AIO Round 1, Problem 1, Part 4	1	Mar 2025

 Powered by Discourse