

# 3D transformations

## The primitive building block transformations

$$S(S_x, S_y, S_z) = \begin{bmatrix} S_x & 0 & 0 & 0 \\ 0 & S_y & 0 & 0 \\ 0 & 0 & S_z & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$T(T_x, T_y, T_z) = \begin{bmatrix} 1 & 0 & 0 & T_x \\ 0 & 1 & 0 & T_y \\ 0 & 0 & 1 & T_z \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$R_z(\theta) = \begin{bmatrix} \cos\theta & -\sin\theta & 0 & 0 \\ \sin\theta & \cos\theta & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$R_y(\theta) = \begin{bmatrix} \cos\theta & 0 & \sin\theta & 0 \\ 0 & 1 & 0 & 0 \\ -\sin\theta & 0 & \cos\theta & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix},$$

$$R_x(\theta) = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos\theta & -\sin\theta & 0 \\ 0 & \sin\theta & \cos\theta & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Algorithm for  $R_i(\theta)$ :

where

i=0,1,2 for the X,Y,Z axes (respectively)

j = (i+0) mod 3

k = (i+2) mod 3

$R_{jj} = \cos\theta$

$R_{kk} = \cos\theta$

$R_{ii} = R_{33} = 1$

$R_{jk} = -\sin\theta$

$R_{kj} = \sin\theta$

*others*=0

# 3D Transformation Interactions


## Scale vs Rotate

left[ matrix{1# 0# 0# 0## 0# 1# 0# 0## 0# 0# 1# 0## 0# 0# 0# 1} right]

left[ matrix{2# 0# 0# 0## 0# 1# 0# 0## 0# 0# 1# 0## 0# 0# 0# 1} right]

left[ matrix{cos 45# -sin 45# 0# 0## sin 45# cos 45# 0# 0## 0# 0# 1# 0## 0# 0# 0# 1} right]

left[ matrix{1# 0# 0# 0## 0# 1# 0# 0## 0# 0# 1# 0## 0# 0# 0# 1} right]




In matrix form, these two series of transformations are:


$$\begin{bmatrix} \cos 45 & -\sin 45 & 0 & 0 \\ \sin 45 & \cos 45 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 2 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 2 \cos 45 & -\sin 45 & 0 & 0 \\ 2 \sin 45 & \cos 45 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos 45 & -\sin 45 & 0 & 0 \\ \sin 45 & \cos 45 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 2 \cos 45 & -2 \sin 45 & 0 & 0 \\ \sin 45 & \cos 45 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

## Scale vs Translate



## Translate vs Rotate



## Complex transformation example

Let's build a rotation by  $\theta$  around a vector  $v = (a, b)$ .

Since we know how to rotate around the axes, we must

1. rotate  $v$  to the X axis:  $R_z(-\phi)$
2. rotate by the desired angle around X:  $R_x(\theta)$
3. rotate  $v$  back to its starting direction:  $R_z(\phi)$

Final transformation is

$$R_z(\phi) R_x(\theta) R_z(-\phi)$$


What about  $\phi$ ?

We don't really need  $\phi$ , but only  $\cos(\phi)$  and  $\sin(\phi)$  to build  $R_z(\phi)$

A little trigonometry gets those two values:

$$\cos(\phi) = a / \sqrt{a^2 + b^2}$$

$$\sin(\phi) = b / \sqrt{a^2 + b^2}$$



## Eye transformation

Specified by

E: Eye position

Z: Direction of view

U: Approximate UP vector

Compute:

$$Y = U - \frac{U \cdot Z}{Z \cdot Z} Z \text{ as the "real" up vector}$$

$$X = Z \times Y$$

Normalize:

X, Y, and Z

Note that X, Y, and Z are **orthonormal**

Step 1: translate to origin:

Translate(-E)

Step 2: rotate:

$$R = \begin{bmatrix} X_x & X_y & X_z \\ Y_x & Y_y & Y_z \\ Z_x & Z_y & Z_z \end{bmatrix} \quad \text{Note that: } \begin{aligned} R X^T &= [1, 0, 0]^T \\ R Y^T &= [0, 1, 0]^T \\ R Z^T &= [0, 0, 1]^T \end{aligned}$$

Why is that called a "rotate"?

## Orthonormal bases and rigid transformations:

Rows of R, taken as vectors are:

normal:  $X \cdot X = Y \cdot Y = Z \cdot Z = 1$

mutually orthogonal:  $X \cdot Y = Y \cdot Z = Z \cdot X = 0$

Such transformations are called *rigid* because:

canonical orthonormal vectors transform to orthonormal vectors

Has a matrix interpretation:

Like this:  $R R^T = I$ ,

but:  $R R^{-1} = I$ ,

so also:  $R^T = R^{-1}$

This is a feature of all rotates:

$$R(\theta) = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$$

$$R(-\theta) = (R(\theta))^{-1} = (R(\theta))^T = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$$

And of all products of rotates:

If A, B are rigid transformations, then AB is also:

$$A^{-1} = A^T, \text{ and } B^{-1} = B^T$$

so

$$(AB)^{-1} = B^{-1} A^{-1} = B^T A^T = (AB)^T$$

## ***A more Interactive approach***

Model sits on a turntable, with controls:

$C$  : center of turntable

$\alpha$  : angle of turntable spin

$\beta$  : angle of turntable forward/backward tilt

$\gamma$  : perhaps angle of turntable spindle (up) projection

$d$  : distance of viewing

Transformations:

$$T(0,0,d) \cdot R_z(\gamma) \cdot R_x(\beta) \cdot R_z(\alpha) \cdot T(-C)$$