

# Bayes' Theorem: Basics

ให้หาความน่าจะเป็นของเบย์

- Total probability Theorem:

$$p(B) = \sum_i p(B|A_i)p(A_i)$$

- Bayes' Theorem:

$$p(H|X) = \frac{p(X|H)P(H)}{p(X)} \propto p(X|H)P(H)$$

posteriori probability      likelihood      prior probability

What we should choose      What we just see      What we knew previously

- **X**: a data sample (“evidence”)

Prediction can be done based on Bayes' Theorem:

- **H**: X belongs to class C

Classification is to derive the maximum posteriori

ความน่าจะเป็นของเบย์

# Naïve Bayes Classifier: Making a Naïve Assumption

- ❑ Practical difficulty of Naïve Bayes inference: It requires initial knowledge of many probabilities, which may not be available or involving significant computational cost
- ❑ A Naïve Special Case
  - ❑ Make an additional **assumption** to simplify the model, but achieve comparable performance.

*Attribute independence assumption*  
attributes are conditionally independent  
(i.e., no dependence relation between attributes)

$$p(X|C_i) = \prod_k p(x_k|C_i) = p(x_1|C_i) \cdot p(x_2|C_i) \cdot \dots \cdot p(x_n|C_i)$$

- ❑ Only need to count the class distribution w.r.t. features

# Naïve Bayes Classifier: Categorical vs. Continuous Valued Features

---

- If feature  $x_k$  is categorical,  $p(x_k = v_k | C_i)$  is the # of tuples in  $C_i$  with  $x_k = v_k$ , divided by  $|C_{i,D}|$  (# of tuples of  $C_i$  in  $D$ )

$$p(X|C_i) = \prod_k p(x_k|C_i) = p(x_1|C_i) \cdot p(x_2|C_i) \cdots p(x_n|C_i)$$

- If feature  $x_k$  is continuous-valued,  $p(x_k = v_k | C_i)$  is usually computed based on Gaussian distribution with a mean  $\mu$  and standard deviation  $\sigma$

$$p(x_k = v_k | C_i) = N(x_k | \mu_{C_i}, \sigma_{C_i}) = \frac{1}{\sqrt{2\pi}\sigma_{C_i}} e^{-\frac{(x - \mu_{C_i})^2}{2\sigma^2}}$$

# Naïve Bayes Classifier: Training Dataset

Class:

C1:buys\_computer = 'yes'

C2:buys\_computer = 'no'

Data to be classified:

X = (age <=30, Income = medium,

Student = yes, Credit\_rating = Fair)

Handwritten red notes: "ok" and "716408" with an arrow pointing to the text above.

| age     | income | student | credit_rating | buys_computer |
|---------|--------|---------|---------------|---------------|
| <=30    | high   | no      | fair          | no            |
| <=30    | high   | no      | excellent     | no            |
| 31...40 | high   | no      | fair          | yes           |
| >40     | medium | no      | fair          | yes           |
| >40     | low    | yes     | fair          | yes           |
| >40     | low    | yes     | excellent     | no            |
| 31...40 | low    | yes     | excellent     | yes           |
| <=30    | medium | no      | fair          | no            |
| <=30    | low    | yes     | fair          | yes           |
| >40     | medium | yes     | fair          | yes           |
| <=30    | medium | yes     | excellent     | yes           |
| 31...40 | medium | no      | excellent     | yes           |
| 31...40 | high   | yes     | fair          | yes           |
| >40     | medium | no      | excellent     | no            |

5

Training Data

$$P(H^y | X') = ?$$

$$P(H'^N | X') = ?$$

$$\rightarrow = P(X|H) P(H)^y \xrightarrow{\text{Training Data } \frac{9}{14}}$$

# Naïve Bayes Classifier: An Example

□  $P(C_i): P(\text{buys\_computer} = \text{"yes"}) = 9/14 = 0.643$

$P(\text{buys\_computer} = \text{"no"}) = 5/14 = 0.357$

□ Compute  $P(X|C_i)$  for each class

$P(\text{age} = \text{"<=30"} | \text{buys\_computer} = \text{"yes"}) = 2/9 = 0.222$

$P(\text{age} = \text{"<=30"} | \text{buys\_computer} = \text{"no"}) = 3/5 = 0.6$

$P(\text{income} = \text{"medium"} | \text{buys\_computer} = \text{"yes"}) = 4/9 = 0.444$

$P(\text{income} = \text{"medium"} | \text{buys\_computer} = \text{"no"}) = 2/5 = 0.4$

$P(\text{student} = \text{"yes"} | \text{buys\_computer} = \text{"yes"}) = 6/9 = 0.667$

$P(\text{student} = \text{"yes"} | \text{buys\_computer} = \text{"no"}) = 1/5 = 0.2$

$P(\text{credit\_rating} = \text{"fair"} | \text{buys\_computer} = \text{"yes"}) = 6/9 = 0.667$

$P(\text{credit\_rating} = \text{"fair"} | \text{buys\_computer} = \text{"no"}) = 2/5 = 0.4$

□  **$X = (\text{age} \leq 30, \text{income} = \text{medium}, \text{student} = \text{yes}, \text{credit\_rating} = \text{fair})$**

**$P(X|C_i): P(X | \text{buys\_computer} = \text{"yes"}) = 0.222 \times 0.444 \times 0.667 \times 0.667 = 0.044$**

$P(X | \text{buys\_computer} = \text{"no"}) = 0.6 \times 0.4 \times 0.2 \times 0.4 = 0.019$

**$P(X|C_i) \cdot P(C_i): P(X | \text{buys\_computer} = \text{"yes"}) \cdot P(\text{buys\_computer} = \text{"yes"}) = 0.028$**

$P(X | \text{buys\_computer} = \text{"no"}) \cdot P(\text{buys\_computer} = \text{"no"}) = 0.007$

**Therefore, X belongs to class ("buys\_computer = yes")**

| age     | income | student | credit_rating | buys_computer |
|---------|--------|---------|---------------|---------------|
| <=30    | high   | no      | fair          | no            |
| <=30    | high   | no      | excellent     | no            |
| 31...40 | high   | no      | fair          | yes           |
| >40     | medium | no      | fair          | yes           |
| >40     | low    | yes     | fair          | yes           |
| >40     | low    | yes     | excellent     | no            |
| 31...40 | low    | yes     | excellent     | yes           |
| <=30    | medium | no      | fair          | no            |
| <=30    | low    | yes     | fair          | yes           |
| >40     | medium | yes     | fair          | yes           |
| <=30    | medium | yes     | excellent     | yes           |
| 31...40 | medium | no      | excellent     | yes           |
| 31...40 | high   | yes     | fair          | yes           |
| >40     | medium | no      | excellent     | no            |

$$\hat{x} = \text{age} = 42, \text{student} = \text{yes}$$

$$P(x^y | \hat{x}) = ?$$


$$P(H=y | \text{age}=42, \text{student}=\text{yes}) = P(\text{age}=42 | y) P(\text{stu}=\text{yes} | y) P(y)$$

$$= \frac{2}{9} \times \frac{6}{9} \times \frac{9}{14}$$


$$P(H_{\text{boy}}=N | \text{age}=42, \text{stu}=\text{yes}) = P(\text{age}=42 | N) P(\text{stu}=\text{yes} | N) P(N)$$



$K'-N'N'$



test data



① 601 จุด จาก 1000 จุด  
 ฝึก space → test space  
 plot 96 จุด

② ระบายสีจุด  
 k = 3

สีของจุด  $k = 3$