Residual Lifetime Risk for Developing Hypertension in Middle-aged Women and Men

The Framingham Heart Study

Ramachandran S. Vasan, MD

Alexa Beiser, PhD

Sudha Seshadri, MD

Martin G. Larson, ScD

William B. Kannel, MD

Ralph B. D'Agostino, PhD

Daniel Levy, MD

IGH BLOOD PRESSURE IS A KEY risk factor for cardiovascular disease events.¹⁻³ The long-term risk for developing hypertension in an individual is best described by the lifetime risk statistic, ie, the probability that an individual will develop hypertension over the course of his or her remaining lifetime. Lifetime risk estimates for disease conditions are more easily understood by the general public compared with other measures of disease frequency, such as agespecific prevalence.4-6 For example, the American Cancer Society used the "1 in 9" statistic for the lifetime risk for breast cancer in the early 1990s to promote mammographic screening.5 Furthermore, lifetime risk estimates also may be used to assess temporal trends in longterm disease risk.4 Although lifetime risk estimates are available for several chronic disease conditions, including breast cancer, dementia, fractures, and coronary heart disease,5,7-9 the lifetime risk for developing hypertension has not been reported.

See also Patient Page.

Context The long-term risk for developing hypertension is best described by the lifetime risk statistic. The lifetime risk for hypertension and trends in this risk over time are unknown.

Objectives To estimate the residual lifetime risk for hypertension in older US adults and to evaluate temporal trends in this risk.

Design, Setting, and Participants Community-based prospective cohort study of 1298 participants from the Framingham Heart Study who were aged 55 to 65 years and free of hypertension at baseline (1976-1998).

Main Outcome Measures Residual lifetime risk (lifetime cumulative incidence not adjusted for competing causes of mortality) for hypertension, defined as blood pressure of 140/90 mm Hg or greater or use of antihypertensive medications.

Results The residual lifetime risks for developing hypertension and stage 1 high blood pressure or higher (≥140/90 mm Hg regardless of treatment) were 90% in both 55-and 65-year-old participants. The lifetime probability of receiving antihypertensive medication was 60%. The risk for hypertension remained unchanged for women, but it was approximately 60% higher for men in the contemporary 1976-1998 period compared with an earlier 1952-1975 period. In contrast, the residual lifetime risk for stage 2 high blood pressure or higher (≥160/100 mm Hg regardless of treatment) was considerably lower in both sexes in the recent period (35%-57% in 1952-1975 vs 35%-44% in 1976-1998), likely due to a marked increase in treatment of individuals with substantially elevated blood pressure.

Conclusion The residual lifetime risk for hypertension for middle-aged and elderly individuals is 90%, indicating a huge public health burden. Although the decline in lifetime risk for stage 2 high blood pressure or higher represents a major achievement, efforts should be directed at the primary prevention of hypertension.

JAMA. 2002;287:1003-1010

www.jama.com

Although the prevalence of hypertension is an indicator of burden of disease in the community, it does not tell us about the risk for developing hyper-

tension in individuals. Data from the serial National Health and Nutritional Examination Surveys (NHANES) indicate a decrease in the age-adjusted preva-

Author Affiliations: the National Heart, Lung, and Blood Institute's Framingham Heart Study (Drs Vasan, Seshadri, Larson, Kannel, and Levy), Framingham, Mass; Cardiology Section (Dr Vasan), Preventive Medicine and Epidemiology (Drs Vasan, Larson, Kannel, and Levy), Department of Neurology (Dr Seshadri), School of Medicine, Epidemiology and Biostatistics (Dr Beiser), School of Public Health, and the Department of Mathematics (Dr D'Agostino), Boston University, Boston, Mass; the Divisions of Cardiology and Clinical

Epidemiology, Beth Israel Deaconess Medical Center, Boston, Mass (Dr Levy); and the National Heart, Lung, and Blood Institute, Bethesda, Md (Dr Levy).

Financial Disclosure: Dr Kannel is a member of the speakers bureau for Bristol-Myers Squibb and Pfizer and consults for Sanofi Pharmaceuticals.

Corresponding Author and Reprints: Ramachandran S. Vasan, MD, Framingham Heart Study, 5 Thurber St, Framingham, MA 01702 (e-mail: vasan@fram.nhlbi.nih.gov).

©2002 American Medical Association. All rights reserved.

(Reprinted) JAMA, February 27, 2002—Vol 287, No. 8 1003

lence of hypertension in the community in the 1976-1991 period compared with surveys during the 1960-1974 period. 10 A trend of decreasing blood pressure levels has been noted also in regional cross-sectional surveys of cardiovascular disease risk factors11 and in longitudinal cohort studies. 12,13 Although these prevalence trends are encouraging, these data should not be interpreted to indicate that the risk for developing hypertension has decreased in recent times. It is possible that the lifetime risk for hypertension may have increased in recent years because of an increase in life expectancy14 and an increase in obesity in the United States.15 On the other hand, the lifetime risk for hypertension may have declined because of preventive efforts directed at patients with high-normal blood pressure.16 Accordingly, we estimated the residual lifetime risk for developing hypertension among Framingham Heart Study participants.

METHODS Study Participants

The Framingham Heart Study is a prospective longitudinal cohort study that began in 1948 with the enrollment of 5209 men and women who were free of cardiovascular disease and between the ages of 28 and 62 years. 17 These participants since have been examined every 2 years. Participants who were free of hypertension in 1975 and who attended subsequent biennial examinations were eligible for our investigation. We chose the period from 1976 onward up to 1998 so our estimates would reflect contemporary experience. We evaluated the residual lifetime risk for hypertension beginning at the baseline ages of 55 and 65 years because the risk for developing hypertension increases markedly during and after the sixth decade of life. 18 Additionally, sufficient numbers of younger participants were not available in our cohort to estimate risks for those younger than 55 years.

Participants could contribute blood pressure information at more than 1 baseline age provided they reached the next baseline age free of hypertension.

For example, a 55-year-old nonhypertensive subject attending a biennial examination in 1976 could contribute information over the next 10 years for this baseline age. If he or she was free of hypertension in 1986, the subject could provide additional information as a 65-year-old individual for the remainder of the time between 1986 and 1998.

Measurement of Blood Pressure

At each biennial examination, participants underwent a physical examination (with a medical history), laboratory assessment of cardiovascular disease risk factors, and routine electrocardiography. A physician measured the systolic and diastolic blood pressures of seated study participants using a mercury column sphygmomanometer and a standard protocol. The first and the fifth Korotkoff sounds were taken as indicative of systolic and diastolic blood pressure, respectively. The average of 2 such systolic and diastolic blood pressure readings was taken as the examination blood pressure. At each examination, participants were asked if they used antihypertensive medications for the specific purpose of lowering elevated blood pressure.

Blood Pressure on Follow Up

All eligible participants were followed up from the time of entry into the study until the end of the observation period, the development of hypertension, death (in the absence of hypertension), or the last follow-up examination. We examined separately the development of each of the following 5 blood pressure outcomes:

1. Hypertension defined according to the recommendations of the Joint National Committee on Detection, Evaluation, and Treatment of High Blood Pressure (JNC VI)¹⁹ and the World Health Organization-International Society of Hypertension¹ as a systolic blood pressure of 140 mm Hg or higher and/or a diastolic blood pressure of 90 mm Hg or higher or the use of medications for the purpose of treating high blood pressure at a qualifying examination;

- 2. Use of antihypertensive medication on follow-up for the treatment of hypertension;
- 3. Stage 1 high blood pressure or higher, defined as blood pressure of 140/90 mm Hg or higher regardless of the use of antihypertensive medications;
- 4. Stage 2 high blood pressure or higher, defined as blood pressure of 160/100 mm Hg or higher regardless of the use of antihypertensive medications;
- 5. Stage 2 high blood pressure or higher or use of antihypertensive medication on follow-up.

Statistical Methods

Residual Lifetime Risk: Definitions. The term *residual lifetime risk* has been defined in 2 ways in the literature: (1) unadjusted lifetime cumulative incidence that indicates the cumulative risk over the remaining lifetime but does not consider the impact of mortality due to competing causes; and (2) mortality-adjusted lifetime cumulative incidence.²⁰ We estimated both statistics in our study.

Residual Lifetime Risk for Developing Hypertension and Other Blood Pressure Outcomes. The residual lifetime risk for developing the blood pressure-related outcomes was estimated for study participants who attained the ages of 55 and 65 years free of hypertension during the 1976-1998 period. Sexspecific analyses were performed separately for these 2 baseline ages. For 55vear-old participants, we estimated the risk for developing hypertension through age 80 years. For 65-year-old participants, we estimated the risk for developing hypertension through age 85 years. These follow-up time intervals (25 years for 55-year-olds and 20 years for 65-year-olds) correspond to the current mean residual life expectancies for white individuals at these 2 ages in the United States.²¹

The residual lifetime risk for each blood pressure-related outcome (including hypertension) was calculated using the Practical Incidence Estimator macro detailed elsewhere.²⁰ This method uses survival age as the time


1004 JAMA, February 27, 2002—Vol 287, No. 8 (Reprinted)


scale, combines information on participants entering the observation periods at different ages, and accounts for varying durations of follow-up of individuals. Briefly, a modified Kaplan-Meier method was used with survival age as the time scale (FIGURE 1). For example, the remaining lifetime risk for hypertension for 55-year-old participants is simply the cumulative incidence of hypertension over 25 years (estimated residual life expectancy) thus:

Cumulative Incidence₅₅= $\sum_{i}(h_{i}S_{i-1})$

where h_i is the conditional probability of developing hypertension at age i years given survival beyond age i-1 years, S_{i-1} is the probability of survival beyond age i-1 years free of hypertension, and $h_i S_{i-1}$ is the unconditional probability of developing hypertension at age i years, where the summation is from i=55 to i=80. Thus, a person who attends an examination at age 61 years and is free of hypertension but who did not attend any examination at age 55 years (subject B in Figure 1) still can contribute to the es-

Figure 1. Estimation of Residual Lifetime Risk of Hypertension





Lifetime risk estimation uses age as the time scale (B) instead of calendar period in the standard Kaplan-Meier estimation (A). As an example, 5 participants enter the study in 1976 at different ages and contribute to lifetime risk estimates for baseline age 55 years. Subjects A, B, C, and E contribute to the estimation of lifetime risk for baseline age 65 years. The lifetime risk estimation method uses risk sets of individuals free of hypertension at different ages. The rates of hypertension are calculated for different age intervals with the number of people who develop hypertension in a given age interval as the numerator and the total person-years of observations of at-risk participants in that interval as the denominator. The person-years of observation obtained on these participants (in this case 61 person-years) is the same regardless of whether calendar period or age is used as the time scale. For instance, for the age interval 55 to 59 years, participants C and D each contribute 5 years to the observation for a total of 10 person-years of observation. For the age interval of 60 to 65 years, subjects B and C contribute 4 person-years of observation each while participant C provides 5 person-years of observation, a total of 13 person-years of observation for this age interval.

Table 1. Residual Lifetime Risk of Hypertension According to Baseline Age*

Time, y	Risk for Hypertension, % (95% Confidence Interval)					
	Women	ı, Age, y	Men, Age, y			
	55 (n = 709)	65 (n = 549)	55 (n = 589)	65 (n = 438)		
10	52 (46-58)	64 (60-69)	56 (49-63)	72 (67-78)		
15	72 (68-76)	81 (77-84)	78 (74-82)	85 (81-89)		
20	83 (80-86)	89 (86-92)	88 (85-91)	90 (87-93)		
25	91 (89-93)		93 (91-95)			

^{*}For 55-year-old subjects, the risk for developing hypertension over 25 years represents their lifetime risk. For 65-yearold subjects, the risk for developing hypertension over 20 years indicates their lifetime risk. Ellipses indicate not applicable.

timation of lifetime risk for baseline age 55 years. If the subject then attends an examination 4 years hence (at age 65 years) and is still free of hypertension, he or she would have contributed 4 person-years of observation as a nonhypertensive individual to this baseline age group. In similar fashion, person-years of observation on all individuals older than baseline ages of 55 and 65 years are summated. The 95% confidence intervals (CIs) of the lifetime risk were calculated as described previously.20 In the calculation of unadjusted cumulative incidence, participants who die free of hypertension were censored (treated as withdrawals from the risk sets). Additional analyses were performed in which the lifetime risk for hypertension was estimated adjusting for competing causes of mortality using double decrement life table analyses.20 In these analyses, nonhypertensive participants who died due to other competing causes were treated as "escapees" (ie, they cannot develop hypertension and do not contribute to the estimation of hypertension incidence).

Residual lifetime risks for developing hypertension for men and women at a baseline age were compared using a z test and by fitting Cox proportional hazards regression models²² with left truncation (SAS program PROC PHREG).²³ The assumption of proportionality of hazards was met, and there were no time-dependent covariates in the models. These Cox models intrinsically adjust for age and compare the risk for developing hypertension among groups over the entire follow-up period (up to time *t* assuming a constant

ratio of hazards) while the *z* tests compared lifetime risks at a specified point in time

Short-term Risks for Developing Hypertension

We also estimated the sex- and agespecific 10- and 15-year cumulative residual risks for developing hypertension (and other blood pressure outcomes) for both age groups. These short-term estimates are of immediate relevance to individuals and complement estimates of lifetime risk.

Temporal Trends in Residual Lifetime Risk for Developing Hypertension

We compared the residual lifetime risk for developing hypertension (and other blood pressure outcomes) in the 1976-1998 period with that in the 1952-1975 period. For this comparison, we focused on unadjusted lifetime cumulative incidence because this statistic is unaffected by changing rates of mortality due to other causes over time.²⁴

Subjects in the Framingham Heart Study who were not hypertensive at their third biennial examination (earliest eligible baseline for that time) contributed to the estimation of lifetime risk for developing hypertension during the 1952-1975 period. We compared the sex- and age-specific risks of developing hypertension and other blood pressure outcomes for the 2 periods with *z* tests (for lifetime risk estimates) and with the Cox models with left truncation. For 65-year-old men, only 15-year risk estimates were compared because few individuals in this age group

contributed information beyond this time interval in the 1952-1975 period. Two sets of analyses were performed: (1) models that included information from all eligible participants in each of the periods; participants who contributed to the 1952-1975 period could provide additional information in the later period (at a different age) provided they reached the later period free of hypertension and (2) models that compared incidence of blood pressure outcomes in participants who contributed to only the 1952-1975 period with that in participants who were eligible only in the 1976-1998 period.

Additionally, we compared sexspecific trends in body mass index (BMI), calculated as weight in kilograms divided by the square of height in meters, across the 2 periods for our study participants using repeated measures analysis of variance adjusting for age as a covariate. All *P* values reported are 2-sided, and a *P* value <.05 was considered statistically significant.

RESULTS

Residual Lifetime and Short-term Risks for Developing Hypertension

During the 1976-1998 period, 1298 participants (589 men and 709 women) provided 8469 person-years of observation. The residual lifetime risk for developing hypertension for study participants was 90% (TABLE 1). Lifetime risk for hypertension was similar for men and women and did not differ between participants aged 55 and 65 years (hazards ratio [HR] for women vs men aged 55 years, 0.91 [95% CI, 0.80-1.04]; for those aged 65 years, 0.88 [95% CI, 0.76-1.04]). More than half of the 55-year-old participants and about two thirds of the 65-year-old participants developed hypertension within 10 years.

On adjustment for competing causes of mortality, the mortality-adjusted lifetime risk remained unchanged in women (90% and 86%, respectively, for baseline ages 55 and 65 years) but decreased marginally in men (83% and 81%, respectively, for baseline ages 55 and 65 years).

1006 JAMA, February 27, 2002—Vol 287, No. 8 (Reprinted)

Lifetime Probability of Using Antihypertensive Medications

Nearly 60% of men and women received blood pressure–lowering drugs on follow-up (TABLE 2). Estimates were similar for both sexes (HR for women vs men aged 55 years, 0.98 [95% CI, 0.83-1.16]; for those aged 65 years, 0.92 [95% CI, 0.76-1.11]), and for the 2 baseline ages. A third of 55-year-old and nearly half of 65-year-old participants were likely to receive antihypertensive medication within 15 years.

Residual Lifetime Risk for High Blood Pressure

The lifetime risk for high blood pressure in our study participants is shown in TABLE 3. Nearly 85% of our study participants developed stage 1 high blood pressure or higher (≥140 mm Hg systolic or ≥90 mm Hg diastolic) during a follow-up period of 20 to 25 years. Lifetime risk estimates did not vary for the 2 sexes (HR for women vs men aged 55 years, 0.91 [95% CI, 0.80-1.04]; for those aged 65 years 0.88 [95% CI, 0.76-1.04]) or between the 2 baseline ages.

The lifetime risk for experiencing stage 2 high blood pressure or higher (≥160 mm Hg systolic or ≥100 mm Hg diastolic) in the 1976-1998 period varied from 35% to 44% for different agesex groups. The lifetime risk for developing stage 2 high blood pressure or higher or receiving antihypertensive medications was about 70% regardless of baseline age or sex.

Trends in Blood Pressure Outcomes Over Time

During the 1952-1975 period, 1740 participants (785 men, 955 women) contributed 12 338 person-years of observation for the estimation of the residual lifetime risk for hypertension. In women, the residual lifetime risk for hypertension did not differ between the 2 periods; risk estimates for the 1952-1975 period were 92% vs 91% for the 1976-1998 periods for baseline age 55 years (*P*=.68); and 93% vs 89% for the baseline age 65 years, respectively (*P*=.51; FIGURE 2A). However, for 55-year-old men, the residual

Table 2. Probability of Receiving Antihypertensive Treatment According to Baseline Age*

Probability of Treatment for Hypertension, % (95% Confidence Interval)

	Women, Age, y		Men, Age, y		
Time, y	55 (n = 709)	65 (n = 549)	55 (n = 589)	65 (n = 438)	
10	20 (15-25)	35 (30-39)	13 (9-17)	39 (33-44)	
15	36 (31-41)	48 (44-53)	33 (28-39)	52 (46-57)	
20	47 (43-52)	57 (53-62)	47 (41-52) 60 (5		
25	58 (54-63)		58 (53-63)		

^{*}For 55-year-old subjects, the probability of treatment over 25 years represents their lifetime risk. For 65-year-old subjects, the probability of treatment over 20 years indicates their lifetime risk. Ellipses indicate not applicable.

Table 3. Residual Lifetime Risk of High Blood Pressure According to Baseline Age*

	Risk for	Risk for High Blood Pressure, % (95% Confidence Interval)					
	Women	, Age, y	Men, Age, y				
Time, y	55 65 (n = 709) (n = 549)		55 (n = 589)	65 (n = 438)			
	Stage 1 High Blo	od Pressure or Highe	r (≥140/90 mm Hg)				
10	45 (39-51)	58 (53-63)	50 (43-57)	62 (56-67)			
15	64 (59-69)	75 (71-79)	70 (65-76)	76 (72-81)			
20	77 (73-80)	83 (80-87)	81 (77-85)	84 (80-88)			
25	86 (83-89)		88 (85-91)				
	Stage 2 High Bloo	od Pressure or Higher	(≥160/100 mm Hg)				
10	6 (3-9)	26 (22-30)	9 (5-13)	20 (16-24)			
15	18 (14-22)	35 (31-40)	20 (15-25)	29 (24-34)			
20	30 (26-35)	44 (39-48)	27 (22-32)	40 (34-46)			
25	39 (35-44)		35 (30-41)				
		od Pressure or Higher hypertensive Medicat					
10	24 (19-29)	46 (41-50)	20 (15-25)	46 (40-51)			
15	45 (40-50)	61 (56-65)	44 (38-49)	59 (53-64)			
20	59 (54-63)	70 (66-74)	56 (51-62)	68 (63-73)			

^{*}For 55-year-old subjects, the risk for developing high blood pressure over 25 years represents their lifetime risk. For 65-year old subjects, the risk for developing high blood pressure over 20 years indicates their lifetime risk. Ellipses indicates not applicable. High blood pressure stages are defined as systolic or diastolic measurements exceeding the specified threshold.

lifetime risk for hypertension was 93% in the 1976-1998 period vs 82% in the 1952-1975 period (P<.001). Similarly, the 15-year risk for hypertension in 65-year-old men increased from 65% in the 1952-1975 period to 85% in the 1976-1998 period (P<.001; Figure 2B). Cox regression analyses confirmed these observations (TABLE 4).

70 (66-74)

Women in the 1952-1975 period had a slightly higher risk for developing hypertension than men in that period (HR for women vs men aged 55 years, 1.15 [95% CI, 1.01-1.30]; for those aged 65 years, 1.23 [95% CI, 0.97-1.56]). As noted previously, point estimates of the

HRs suggested a lower risk in women compared with men in the 1976-1998 period although this was not statistically significant.

67 (62-72)

The residual lifetime probability of being treated with antihypertensive medications ranged between 16% and 36% in different age-sex groups in the 1952-1975 period (data not shown), increasing 2- to 3-fold in both sexes in the 1976-1998 period (P<.001 for all comparisons). These findings were supported by the results of Cox models (Table 4). The risk for stage 1 high blood pressure or higher was 92% for women (for both baseline ages) but var-

©2002 American Medical Association. All rights reserved.

(Reprinted) JAMA, February 27, 2002—Vol 287, No. 8 1007


ied from 65% (age 65 years) to 82% (age 55 years) for men in the 1952-1975 period. The risk for stage 1 high blood pressure or higher did not change across the 2 periods for women but increased by about 25% in men (Table 4). The risk for stage 2 high blood pressure or higher was 57% for women (for both baseline ages) and varied from 35% (age 65 years) to 48% (age 55 years) for men in the 1952-1975 period. The risk for stage 2 high blood pressure or higher declined in the 1976-1998 period both in women and in men (Table 4).

There were 786 participants (344 men, 442 women) who contributed information to both periods although at

different ages. Analyses were repeated after excluding these participants. Estimates of residual lifetime risk for hypertension in the 1976-1998 period were higher after these exclusions, in part because people who contributed to the 1952-1975 period and remained eligible in the 1976-1998 period (having reached this period free of hypertension) were likely participants with a lower risk for developing hypertension. A comparison of participants contributing to the 1952-1975 period only (n=954; 513 women, 441 men) with those providing information in the 1976-1998 period only (n=512; 267 women, 245 men) confirmed the increased risk for hypertension in men and the increased use of antihypertensive medications in both men and women in the 1976-1998 period. However, the increased risk for stage 1 high blood pressure or higher among men in the 1976-1998 period noted in earlier analyses did not achieve statistical significance, in part due to a decrease in sample size and statistical power.

The temporal trends we observed may be due, in part, to concomitant trends in obesity. In women, ageadjusted BMI decreased from 25.8 kg/m² in the 1952-1975 period to 25.3 kg/m² in the 1976-1998 period (*P*=.05). In contrast, age-adjusted BMI in-

Figure 2. Residual Lifetime Risk of Hypertension in Women and Men Aged 65 Years



Cumulative Incidence of hypertension in 65-year-old women and men. Data for 65-year-old men in the 1952-1975 period is truncated at 15 years since there were few participants in this age category who were followed up beyond this time interval.

Table 4. Changes in Risk of Hypertension Over Time*

	1976-1998 vs 1952-1975							
	Women, Age, y				Men, Age, y			
	55		65	55		65		
	HR (95% CI)	P Value	HR (95% CI)	P Value	HR (95% CI)	P Value	HR (95% CI)	P Value
Hypertension	1.13 (0.97-1.31)	.11	1.05 (0.85-1.29)	.67	1.56 (1.34-1.83)	<.001	1.62 (1.31-2.00)	<.001
Antihypertension medication	1.98 (1.62-2.42)	<.001	2.14 (1.67-2.75)	<.001	3.33 (2.59-4.27)	<.001	3.95 (2.87-5.41)	<.001
High blood pressure stage	0.95 (0.82-1.11)	.53	0.91 (0.74-1.12)	.38	1.26 (1.07-1.48)	.006	1.24 (1.00-1.54)	.05
2	0.63 (0.51-0.77)	<.001	0.69 (0.54-0.86)	.001	0.73 (0.57-0.94)	.01	0.85 (0.64-1.14)	.28
2 or antihypertension medication	1.28 (1.08-1.51)	.004	1.28 (1.04-1.56)	.02	1.61 (1.33-1.95)	<.001	1.90 (1.49-2.42)	<.001

^{*}These comparisons are between 1298 subjects in the 1976-1998 period and 1740 subjects in the 1952-1975 period. After exclusion of 786 subjects who contributed to both periods, the increased risk of hypertension in men and the increased use of antihypertensive medications in both men and women in the 1976-1998 period persisted. However, the increased risk of stage 1 high blood pressure or higher among men in the 1976-1998 period did not achieve statistical significance, in part due to a decrease in sample size and in statistical power. HR indicates hazards ratio; CI confidence interval. For definition of each stage of high blood pressure see "Methods" section.

creased in men from 25.8 kg/m² in the 1952-1975 period to 26.3 kg/m² in the 1976-1998 period (P=.02).

COMMENT

Principal Findings

Our investigation provides estimates of the residual lifetime and the shortterm risks for developing hypertension and examines trends in these risks over time. Whereas the short-term risk estimates are of more immediate value to individuals and, therefore, more likely to motivate behavioral changes, residual lifetime risk estimates permit a comparison of the likelihood of developing 1 or more disease conditions over a lifetime.

The residual lifetime risk for developing hypertension in our sample was about 90%. These risk estimates were similar for men and women and for participants aged 55 and 65 years. It is often assumed that the risk for hypertension increases with age because the incidence of hypertension increases with age. Although this is true of shortterm age-conditional risk estimates, this may not be true for lifetime risk. The lack of increase in lifetime risk with increasing age is consistent with similar observations made for other chronic diseases such as dementia⁷ and may be because the rising age-specific incidence of hypertension is offset by a decreasing residual life expectancy. Using a higher threshold to define high blood pressure (stage 2 high blood pressure or higher or use of medications) vielded lifetime risk estimates of about 70% in both women and men.

A comparison of lifetime risk estimates over the 2 periods showed that the risk for developing hypertension was unchanged for women but increased for men. The increase in lifetime risk for hypertension in men may be related in part to the trend for an increase in BMI over this period among men (but not women) in our sample. Other investigators have also reported favorable secular trends in blood pressure of women but less encouraging trends in men. ^{25,26} In contrast, in the 1976-1998 period, the lifetime prob-

ability of experiencing stage 2 high blood pressure or higher was considerably lower in both sexes, in tandem with and probably as a result of a 2- to 3-fold increased use of antihypertensive medications. These data are consistent with prior reports of a decrease in the prevalence of long-term sustained hypertension in our cohort 13,27 and are likely the result of an increasing emphasis on pharmacological treatment of elevated blood pressure and the progressive lowering of blood pressure treatment thresholds in national guidelines for hypertension treatment (along with changing definitions of hypertension).1 However, the residual lifetime risk for experiencing stage 1 high blood pressure or higher is markedly high at about 85% in the 1976-1998 period.

Trends in Risk for Hypertension: Comparison With Prior Reports

Our findings differ significantly from observations made in serial national cross-sectional surveys between 1960 and 1991, which suggest that the ageadjusted prevalence of hypertension in the United States declined from 36% in 1971 to 20% in 1988 through 1991, along with a downward shift in the average blood pressure of the population. 10 A principal reason our results differ from data from national surveys is that we evaluated trends in the incidence of hypertension while the NHANES report assessed trends in the cross-sectional prevalence of hypertension. It is important to note that our study and the NHANES report evaluated slightly different periods; our investigation began earlier and extended into 1998. It has been reported that some of the "gains" noted in the initial phase of the NHANES III (1988-1991) had leveled off by the time of the second phase of the survey (1991-1994). 19 Increases in the mean blood pressure of cohorts in Iowa and Minnesota in the late 1980s and 1990s, respectively, also have been reported.^{28,29} An additional explanation to be considered is that our sample was white and restricted in its age distribution while

that of NHANES was ethnically more diverse and encompassed a wider age spectrum.

Strengths and Limitations

Our study is based on the longitudinal surveillance of the same community-based cohort over a period of 50 years. Over this time, blood pressure measurements were obtained every 2 years using a standardized protocol, and information regarding the use of antihypertensive medications was gathered consistently and routinely. Additionally, we examined trends in the lifetime risk for several blood pressure outcomes that yielded complementary information.

Nevertheless, we acknowledge certain limitations of our investigation. Our data indicate the residual lifetime risk of hypertension for 55- and 65-yearold participants who were free of hypertension at these baseline ages. It is important to note that a considerable proportion of individuals with hypertension have onset of the condition before this age. Therefore, the actual lifetime risk for hypertension for younger individuals may be different. Moreover, these lifetime risk data are average estimates for our study participants. The risk for hypertension for a given individual will vary depending on the presence or absence of risk factors for hypertension (such as obesity, family history of high blood pressure, dietary sodium and potassium intake, and alcohol consumption). We did not examine the impact of these covariates. Categorizing people as hypertensive based on a reading obtained on a single occasion may overestimate risk for hypertension. For this reason, we assessed risk for several blood pressure outcomes. The 90% lifetime risk estimate likely represents the upper estimate of true risk because it is based on a single occasion measurement and not on the average of multiple visits (required by the JNC VI). The lower bound of the residual lifetime risk for hypertension is 70%, a figure obtained using a composite definition of stage 2 high blood pressure or higher or the use of antihypertensive medications. We

©2002 American Medical Association. All rights reserved.

(Reprinted) JAMA, February 27, 2002—Vol 287, No. 8 **1009**

may have underestimated the lifetime probability of experiencing stage 2 high blood pressure or higher because some participants experiencing this level of blood pressure on seeing a private physician between biennial Heart Study examinations may have had their blood pressure lowered with medications by the time of the next examination. Last, our lifetime risk estimates and trends may not be generalizable to other ethnic groups. Similar studies in other multiethnic cohorts are warranted to obtain information regarding risk for hypertension in different racial groups.

Public Health Implications

The finding that 9 out of 10 middle-aged and older adults are likely to develop elevated blood pressure over their remaining lifetime reemphasizes that hypertension poses a major public health burden. Furthermore, the lack of change in the lifetime risk over the 2 periods in women and the modest increase in risk in men is of concern. With the aging of the US population, ¹⁴ the societal burden of hypertension will increase further if the lifetime risk for the condition remains unattenuated.

The marked decline in the lifetime risk for stage 2 high blood pressure or higher in men and women in our study sample represents a major public health achievement. In contrast, the risk for stage 1 high blood pressure or higher remained unaltered and represents the next frontier. Although hypertension is easily diagnosed and treated, ¹⁹ prior data from Framingham and elsewhere suggest that stage 1 hypertension is inadequately controlled. ^{10,30,31} This is of public health importance because a substantial proportion of cardiovascular diseases are attributable to stage 1 hypertension. ^{2,32}

It is widely accepted that hypertension is preventable through lifestyle modification. ^{19,33} Our estimates of the short-term risk for hypertension indicate that more than half of the residual lifetime risk for hypertension for 55- and 65-year-old participants is experienced over the initial 10 years of follow-up. The immediacy and the magnitude of this risk should encourage

middle-aged nonhypertensive individuals to adopt lifestyle-related measures for maintaining optimal blood pressure and for preventing the development of hypertension. The approach of waiting for hypertension to develop and only then treating the elevated blood pressure is injudicious. As articulated by Stamler, ³⁴ such a strategy is "late, defensive, reactive, time consuming, associated with side effects, costly, only partially successful, and endless."

Author Contributions: Study concept and design: Vasan, Beiser, Seshadri, Kannel, D'Agostino, Levy. Analysis and interpretation of data: Vasan, Beiser, Seshadri, Larson, Kannel, D'Agostino, Levy. Drafting of the manuscript: Vasan, Beiser, Seshadri, Levy.

Critical revision of the manuscript for important intellectual content: Vasan, Beiser, Seshadri, Larson, Kannel, D'Agostino, Levy.

Statistical expertise: Beiser, Larson, D'Agostino. Study supervision: Vasan, Seshadri, Kannel, Levy. Funding/Support: This work was in part supported by grants N01-HC-38038 from the National Heart, Lung, and Blood Institute (NHLBI) and NS17950 from the National Institute of Neurological Disorders and Stroke. Dr Vasan was supported in part by grant K24 HL 04334-01A1 from the NHLBI.

REFERENCES

- **1.** 1999 World Health Organization-International Society of Hypertension Guidelines for the Management of Hypertension. Guidelines Subcommittee. *J Hypertens*. 1999; 17:151-183.
- **2.** Stamler J, Stamler R, Neaton JD. Blood pressure, systolic and diastolic, and cardiovascular risks: US population data. *Arch Intern Med.* 1993;153:598-615.
- 3. van den Hoogen PC, Feskens EJ, Nagelkerke NJ, Menotti A, Nissinen A, Kromhout D. The relation between blood pressure and mortality due to coronary heart disease among men in different parts of the world. *N Engl J Med.* 2000;342:1-8.
- **4.** Sasieni PD, Adams J. Standardized lifetime risk. *Am J Epidemiol*. 1999;149:869-875.
- **5.** Phillips KA, Glendon G, Knight JA. Putting the risk of breast cancer in perspective. *N Engl J Med*. 1999; 340:141-144.
- **6.** Fortin JM, Hirota LK, Bond BE, O'Connor AM, Col NF. Identifying patient preferences for communicating risk estimates: a descriptive pilot study. *BMC Med Inform Decis Mak*. 2001:1:2.
- 7. Seshadri S, Wolf PA, Beiser A, et al. Lifetime risk of dementia and Alzheimer's disease. *Neurology*. 1997; 49:1498-1504.
- **8.** Cummings SR, Black DM, Rubin SM. Lifetime risks of hip, Colles', or vertebral fracture and coronary heart disease among white postmenopausal women. *Arch Intern Med.* 1989:149:2445-2448.
- **9.** Lloyd-Jones DM, Larson MG, Beiser A, Levy D. Lifetime risk of developing coronary heart disease. *Lancet.* 1999;353:89-92.
- **10.** Burt VL, Cutler JA, Higgins M, et al. Trends in the prevalence, awareness, treatment, and control of hypertension in the adult US population. *Hypertension*. 1995; 26:60-69
- 11. McGovern PG, Burke GL, Sprafka JM, Xue S, Folsom AR, Blackburn H. Trends in mortality, morbidity, and risk factor levels for stroke from 1960 through 1990: the Minnesota Heart Survey. *JAMA*. 1992;268: 753-759.

- **12.** Glynn RJ, Field TS, Satterfield S, et al. Modification of increasing systolic blood pressure in the elderly during the 1980s. *Am J Epidemiol*. 1993;138: 365-379
- **13.** Mosterd A, D'Agostino RB, Silbershatz H, et al. Trends in the prevalence of hypertension, antihypertensive therapy, and left ventricular hypertrophy from 1950 to 1989. *N Engl J Med.* 1999;340:1221-1227.
- **14.** Kinsella K, Velkoff VA. *An Aging World: 2001*. Washington, DC: US Bureau of Census, US Government Printing Office; 2001;1-183. Series P95/01-1.
- **15.** Mokdad AH, Serdula MK, Dietz WH, Bowman BA, Marks JS, Koplan JP. The spread of the obesity epidemic in the United States, 1991-1998. *JAMA*. 1999; 282:1519-1522.
- **16.** National High Blood Pressure Education Program Working Group report on primary prevention of hypertension. *Arch Intern Med.* 1993;153:186-208.
- **17.** Dawber TR, Meadors GF, Moore FE. Epidemiologic approaches to heart disease. *Am J Public Health*. 1951;41:279-286.
- **18.** Dannenberg AL, Garrison RJ, Kannel WB. Incidence of hypertension in the Framingham Study. *Am J Public Health*. 1988;78:676-679.
- **19.** The sixth report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. *Arch Intern Med.* 1997;157: 2413-2446.
- **20.** Beiser A, D'Agostino RB Sr, Seshadri S, Sullivan LM, Wolf PA. Computing estimates of incidence, including lifetime risk. *Stat Med*. 2000;19:1495-1522.
- 21. Anderson RN. *United States Life Tables: National Vital Statistics Report*. Hyattsville, Md: National Center for Health Statistics; 1999:12-13. Series 47(28).
- **22.** Cox DR, Oakes D. *Analysis of Survival Data*. London, England: Chapman & Hall; 1984.
- 23. Sas/Stat Software: Changes and Enhancements, Through Release 6.11. Cary, NC: SAS Institute; 1996.
- **24.** Wun LM, Merrill RM, Feuer EJ. Estimating lifetime and age-conditional probabilities of developing cancer. *Lifetime Data Anal*. 1998;4:169-186.
- **25.** Garraway WM, Whisnant JP. The changing pattern of hypertension and the declining incidence of stroke. *JAMA*. 1987;258:214-217.
- **26.** Barker WH, Mullooly JP, Linton KL. Trends in hypertension prevalence, treatment, and control. *Hypertension*. 1998;31(1pt 2):552-559.
- 27. Sytkowski PA, D'Agostino RB, Belanger AJ, Kannel WB. Secular trends in long-term sustained hypertension, long-term treatment, and cardiovascular mortality. *Circulation*. 1996:93:697-703.
- **28.** Glynn RJ, Brock DB, Harris T, et al. Use of anti-hypertensive drugs and trends in blood pressure in the elderly. *Arch Intern Med.* 1995;155:1855-1860.
- **29.** Meissner I, Whisnant JP, Sheps SG, et al. Detection and control of high blood pressure in the community. *Hypertension*. 1999;34:466-471.
- **30.** Lloyd-Jones DM, Evans JC, Larson MG, O'Donnell CJ, Roccella EJ, Levy D. Differential control of systolic and diastolic blood pressure. *Hypertension*. 2000;36: 594-599
- **31.** Hyman DJ, Pavlik VN. Self-reported hypertension treatment practices among primary care physicians: blood pressure thresholds, drug choices, and the role of guidelines and evidence-based medicine. *Arch Intern Med.* 2000:160:2281-2286.
- **32.** Miura K, Daviglus ML, Dyer AR, et al. Relationship of blood pressure to 25-year mortality due to coronary heart disease, cardiovascular diseases, and all causes in young adult men. *Arch Intern Med.* 2001; 161:1501-1508
- **33.** Sacks FM, Svetkey LP, Vollmer WM, et al. Effects on blood pressure of reduced dietary sodium and the Dietary Approaches to Stop Hypertension (DASH) diet. *N Engl J Med*. 2001;344:3-10.
- **34.** Stamler J. Setting the TONE for ending the hypertension epidemic. *JAMA*. 1998;279:878-879.

1010 JAMA, February 27, 2002—Vol 287, No. 8 (Reprinted)