

Scalar subquery -returns a single value.

Correlated subquery - one that depends on a value or table name from the outer query to execute.

Uncorrelated subquery - has no reference to objects in the main query.

If the subquery below is run standalone , it returns 213707.3

The screenshot shows a PostgreSQL query editor window titled "analysis/postgres@PostgreSQL 15+ X". The query editor displays the following SQL code:

```
1
2
3  -- Example of an uncorrelated subquery
4
5
6
7  SELECT  county_name,
8          state_name,
9          pop_est_2019
10 FROM us_counties_2019 >=(
11     SELECT percentile_cont(.9) WITHIN GROUP (ORDER BY pop_est_2019)
12     FROM us_counties_pop_est_2019
13 )
14 ORDER BY pop_est_2019 DESC;
15
16
17 -- This query can be run as-is, and is not dependent on any other values;
18 -- It is embedded into the above query as an uncorrelated subquery:
19
20 SELECT percentile_cont(.9) WITHIN GROUP (ORDER BY pop_est_2019)
21 FROM us_counties_pop_est_2019;
22
```

The "Data Output" tab is active, showing the result of the query. The result is a single row with the following values:

	percentile_cont double precision
1	213707.30000000008

The interface also includes a toolbar with various icons for query execution and a status bar at the bottom right indicating "Showing rows: 1 to 1".