

METHODS OF MULTIVARIATE ANALYSIS

WILEY SERIES IN PROBABILITY AND STATISTICS

Established by WALTER A. SHEWHART and SAMUEL S. WILKS

Editors: *David J. Balding, Noel A. C. Cressie, Garrett M. Fitzmaurice, Harvey Goldstein, Iain M. Johnstone, Geert Molenberghs, David W. Scott, Adrian F. M. Smith, Ruey S. Tsay, Sanford Weisberg*

Editors Emeriti: *Vic Barnett, J. Stuart Hunter, Joseph B. Kadane, Jozef L. Teugels*

A complete list of the titles in this series appears at the end of this volume.

METHODS OF MULTIVARIATE ANALYSIS

Third Edition

Alvin C. Rencher

William F. Christensen

Department of Statistics
Brigham Young University
Provo, Utah



A JOHN WILEY & SONS, INC., PUBLICATION

Copyright © 2012 by John Wiley & Sons, Inc. All rights reserved

Published by John Wiley & Sons, Inc., Hoboken, New Jersey
Published simultaneously in Canada

No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, scanning, or otherwise, except as permitted under Section 107 or 108 of the 1976 United States Copyright Act, without either the prior written permission of the Publisher, or authorization through payment of the appropriate per-copy fee to the Copyright Clearance Center, Inc., 222 Rosewood Drive, Danvers, MA 01923, (978) 750-8400, fax (978) 750-4470, or on the web at www.copyright.com. Requests to the Publisher for permission should be addressed to the Permissions Department, John Wiley & Sons, Inc., 111 River Street, Hoboken, NJ 07030, (201) 748-6011, fax (201) 748-6008, or online at <http://www.wiley.com/go/permission>.

Limit of Liability/Disclaimer of Warranty: While the publisher and author have used their best efforts in preparing this book, they make no representations or warranties with respect to the accuracy or completeness of the contents of this book and specifically disclaim any implied warranties of merchantability or fitness for a particular purpose. No warranty may be created or extended by sales representatives or written sales materials. The advice and strategies contained herein may not be suitable for your situation. You should consult with a professional where appropriate. Neither the publisher nor author shall be liable for any loss of profit or any other commercial damages, including but not limited to special, incidental, consequential, or other damages.

For general information on our other products and services or for technical support, please contact our Customer Care Department within the United States at (800) 762-2974, outside the United States at (317) 572-3993 or fax (317) 572-4002.

Wiley also publishes its books in a variety of electronic formats. Some content that appears in print may not be available in electronic formats. For more information about Wiley products, visit our web site at www.wiley.com.

Library of Congress Cataloging-in-Publication Data:

Rencher, Alvin C., 1934—

Methods of multivariate analysis / Alvin C. Rencher, William F. Christensen, Department of Statistics, Brigham Young University, Provo, UT. — Third Edition.

pages cm. — (Wiley series in probability and statistics)

Includes index.

ISBN 978-0-470-17896-6 (hardback)

1. Multivariate analysis. I. Christensen, William F., 1970– II. Title.

QA278.R45 2012

519.5'35—dc23

2012009793

Printed in the United States of America.

10 9 8 7 6 5 4 3 2 1

CONTENTS

Preface	xvii
Acknowledgments	xxi
1 Introduction	1
1.1 WHY MULTIVARIATE ANALYSIS?	1
1.2 PREREQUISITES	3
1.3 OBJECTIVES	3
1.4 BASIC TYPES OF DATA AND ANALYSIS	4
2 Matrix Algebra	7
2.1 INTRODUCTION	7
2.2 NOTATION AND BASIC DEFINITIONS	8
2.2.1 Matrices, Vectors, and Scalars	8
2.2.2 Equality of Vectors and Matrices	9
2.2.3 Transpose and Symmetric Matrices	9
2.2.4 Special Matrices	10
2.3 OPERATIONS	11

2.3.1	Summation and Product Notation	11
2.3.2	Addition of Matrices and Vectors	12
2.3.3	Multiplication of Matrices and Vectors	13
2.4	PARTITIONED MATRICES	22
2.5	RANK	23
2.6	INVERSE	25
2.7	POSITIVE DEFINITE MATRICES	26
2.8	DETERMINANTS	28
2.9	TRACE	31
2.10	ORTHOGONAL VECTORS AND MATRICES	31
2.11	EIGENVALUES AND EIGENVECTORS	32
2.11.1	Definition	32
2.11.2	$\mathbf{I} + \mathbf{A}$ and $\mathbf{I} - \mathbf{A}$	34
2.11.3	$\text{tr}(\mathbf{A})$ and $ \mathbf{A} $	34
2.11.4	Positive Definite and Semidefinite Matrices	35
2.11.5	The Product \mathbf{AB}	35
2.11.6	Symmetric Matrix	35
2.11.7	Spectral Decomposition	35
2.11.8	Square Root Matrix	36
2.11.9	Square and Inverse Matrices	36
2.11.10	Singular Value Decomposition	37
2.12	KRONECKER AND VEC NOTATION	37
	Problems	39
3	Characterizing and Displaying Multivariate Data	47
3.1	MEAN AND VARIANCE OF A UNIVARIATE RANDOM VARIABLE	47
3.2	COVARIANCE AND CORRELATION OF BIVARIATE RANDOM VARIABLES	49
3.2.1	Covariance	49
3.2.2	Correlation	53
3.3	SCATTERPLOTS OF BIVARIATE SAMPLES	55
3.4	GRAPHICAL DISPLAYS FOR MULTIVARIATE SAMPLES	56
3.5	DYNAMIC GRAPHICS	58
3.6	MEAN VECTORS	63
3.7	COVARIANCE MATRICES	66
3.8	CORRELATION MATRICES	69

3.9	MEAN VECTORS AND COVARIANCE MATRICES FOR SUBSETS OF VARIABLES	71
3.9.1	Two Subsets	71
3.9.2	Three or More Subsets	73
3.10	LINEAR COMBINATIONS OF VARIABLES	75
3.10.1	Sample Properties	75
3.10.2	Population Properties	81
3.11	MEASURES OF OVERALL VARIABILITY	81
3.12	ESTIMATION OF MISSING VALUES	82
3.13	DISTANCE BETWEEN VECTORS	84
	Problems	85
4	The Multivariate Normal Distribution	91
4.1	MULTIVARIATE NORMAL DENSITY FUNCTION	91
4.1.1	Univariate Normal Density	92
4.1.2	Multivariate Normal Density	92
4.1.3	Generalized Population Variance	93
4.1.4	Diversity of Applications of the Multivariate Normal	93
4.2	PROPERTIES OF MULTIVARIATE NORMAL RANDOM VARIABLES	94
4.3	ESTIMATION IN THE MULTIVARIATE NORMAL	99
4.3.1	Maximum Likelihood Estimation	99
4.3.2	Distribution of \bar{y} and S	100
4.4	ASSESSING MULTIVARIATE NORMALITY	101
4.4.1	Investigating Univariate Normality	101
4.4.2	Investigating Multivariate Normality	106
4.5	TRANSFORMATIONS TO NORMALITY	108
4.5.1	Univariate Transformations to Normality	109
4.5.2	Multivariate Transformations to Normality	110
4.6	OUTLIERS	111
4.6.1	Outliers in Univariate Samples	112
4.6.2	Outliers in Multivariate Samples	113
	Problems	117
5	Tests on One or Two Mean Vectors	125
5.1	MULTIVARIATE VERSUS UNIVARIATE TESTS	125
5.2	TESTS ON μ WITH Σ KNOWN	126

5.2.1	Review of Univariate Test for $H_0: \mu = \mu_0$ with σ Known	126
5.2.2	Multivariate Test for $H_0: \mu = \mu_0$ with Σ Known	127
5.3	TESTS ON μ WHEN Σ IS UNKNOWN	130
5.3.1	Review of Univariate t -Test for $H_0: \mu = \mu_0$ with σ Unknown	130
5.3.2	Hotelling's T^2 -Test for $H_0: \mu = \mu_0$ with Σ Unknown	131
5.4	COMPARING TWO MEAN VECTORS	134
5.4.1	Review of Univariate Two-Sample t -Test	134
5.4.2	Multivariate Two-Sample T^2 -Test	135
5.4.3	Likelihood Ratio Tests	139
5.5	TESTS ON INDIVIDUAL VARIABLES CONDITIONAL ON REJECTION OF H_0 BY THE T^2-TEST	139
5.6	COMPUTATION OF T^2	143
5.6.1	Obtaining T^2 from a MANOVA Program	143
5.6.2	Obtaining T^2 from Multiple Regression	144
5.7	PAIRED OBSERVATIONS TEST	145
5.7.1	Univariate Case	145
5.7.2	Multivariate Case	147
5.8	TEST FOR ADDITIONAL INFORMATION	149
5.9	PROFILE ANALYSIS	152
5.9.1	One-Sample Profile Analysis	152
5.9.2	Two-Sample Profile Analysis	154
	Problems	161
6	Multivariate Analysis of Variance	169
6.1	ONE-WAY MODELS	169
6.1.1	Univariate One-Way Analysis of Variance (ANOVA)	169
6.1.2	Multivariate One-Way Analysis of Variance Model (MANOVA)	171
6.1.3	Wilks' Test Statistic	174
6.1.4	Roy's Test	178
6.1.5	Pillai and Lawley–Hotelling Tests	179
6.1.6	Unbalanced One-Way MANOVA	181
6.1.7	Summary of the Four Tests and Relationship to T^2	182
6.1.8	Measures of Multivariate Association	186
6.2	COMPARISON OF THE FOUR MANOVA TEST STATISTICS	189
6.3	CONTRASTS	191

6.3.1	Univariate Contrasts	191
6.3.2	Multivariate Contrasts	192
6.4	TESTS ON INDIVIDUAL VARIABLES FOLLOWING REJECTION OF H_0 BY THE OVERALL MANOVA TEST	195
6.5	TWO-WAY CLASSIFICATION	198
6.5.1	Review of Univariate Two-Way ANOVA	198
6.5.2	Multivariate Two-Way MANOVA	201
6.6	OTHER MODELS	207
6.6.1	Higher-Order Fixed Effects	207
6.6.2	Mixed Models	208
6.7	CHECKING ON THE ASSUMPTIONS	210
6.8	PROFILE ANALYSIS	211
6.9	REPEATED MEASURES DESIGNS	215
6.9.1	Multivariate Versus Univariate Approach	215
6.9.2	One-Sample Repeated Measures Model	219
6.9.3	k -Sample Repeated Measures Model	222
6.9.4	Computation of Repeated Measures Tests	224
6.9.5	Repeated Measures with Two Within-Subjects Factors and One Between-Subjects Factor	224
6.9.6	Repeated Measures with Two Within-Subjects Factors and Two Between-Subjects Factors	230
6.9.7	Additional Topics	232
6.10	GROWTH CURVES	232
6.10.1	Growth Curve for One Sample	232
6.10.2	Growth Curves for Several Samples	239
6.10.3	Additional Topics	241
6.11	TESTS ON A SUBVECTOR	241
6.11.1	Test for Additional Information	241
6.11.2	Stepwise Selection of Variables	243
	Problems	244
7	Tests on Covariance Matrices	259
7.1	INTRODUCTION	259
7.2	TESTING A SPECIFIED PATTERN FOR Σ	259
7.2.1	Testing $H_0: \Sigma = \Sigma_0$	260
7.2.2	Testing Sphericity	261
7.2.3	Testing $H_0: \Sigma = \sigma^2[(1 - \rho)\mathbf{I} + \rho\mathbf{J}]$	263
7.3	TESTS COMPARING COVARIANCE MATRICES	265

X CONTENTS

7.3.1	Univariate Tests of Equality of Variances	265
7.3.2	Multivariate Tests of Equality of Covariance Matrices	266
7.4	TESTS OF INDEPENDENCE	269
7.4.1	Independence of Two Subvectors	269
7.4.2	Independence of Several Subvectors	271
7.4.3	Test for Independence of All Variables	275
	Problems	276
8	Discriminant Analysis: Description of Group Separation	281
8.1	INTRODUCTION	281
8.2	THE DISCRIMINANT FUNCTION FOR TWO GROUPS	282
8.3	RELATIONSHIP BETWEEN TWO-GROUP DISCRIMINANT ANALYSIS AND MULTIPLE REGRESSION	286
8.4	DISCRIMINANT ANALYSIS FOR SEVERAL GROUPS	288
8.4.1	Discriminant Functions	288
8.4.2	A Measure of Association for Discriminant Functions	292
8.5	STANDARDIZED DISCRIMINANT FUNCTIONS	292
8.6	TESTS OF SIGNIFICANCE	294
8.6.1	Tests for the Two-Group Case	294
8.6.2	Tests for the Several-Group Case	295
8.7	INTERPRETATION OF DISCRIMINANT FUNCTIONS	298
8.7.1	Standardized Coefficients	298
8.7.2	Partial <i>F</i> -Values	299
8.7.3	Correlations Between Variables and Discriminant Functions	300
8.7.4	Rotation	301
8.8	SCATTERPLOTS	301
8.9	STEPWISE SELECTION OF VARIABLES	303
	Problems	306
9	Classification Analysis: Allocation of Observations to Groups	309
9.1	INTRODUCTION	309
9.2	CLASSIFICATION INTO TWO GROUPS	310
9.3	CLASSIFICATION INTO SEVERAL GROUPS	314
9.3.1	Equal Population Covariance Matrices: Linear Classification Functions	315
9.3.2	Unequal Population Covariance Matrices: Quadratic Classification Functions	317

9.4	ESTIMATING MISCLASSIFICATION RATES	318
9.5	IMPROVED ESTIMATES OF ERROR RATES	320
9.5.1	Partitioning the Sample	321
9.5.2	Holdout Method	322
9.6	SUBSET SELECTION	322
9.7	NONPARAMETRIC PROCEDURES	326
9.7.1	Multinomial Data	326
9.7.2	Classification Based on Density Estimators	327
9.7.3	Nearest Neighbor Classification Rule	330
9.7.4	Classification Trees	331
	Problems	336
10	Multivariate Regression	339
10.1	INTRODUCTION	339
10.2	MULTIPLE REGRESSION: FIXED x 's	340
10.2.1	Model for Fixed x 's	340
10.2.2	Least Squares Estimation in the Fixed- x Model	342
10.2.3	An Estimator for σ^2	343
10.2.4	The Model Corrected for Means	344
10.2.5	Hypothesis Tests	346
10.2.6	R^2 in Fixed- x Regression	349
10.2.7	Subset Selection	350
10.3	MULTIPLE REGRESSION: RANDOM x 's	354
10.4	MULTIVARIATE MULTIPLE REGRESSION: ESTIMATION	354
10.4.1	The Multivariate Linear Model	354
10.4.2	Least Squares Estimation in the Multivariate Model	356
10.4.3	Properties of Least Squares Estimator $\hat{\mathbf{B}}$	358
10.4.4	An Estimator for Σ	360
10.4.5	Model Corrected for Means	361
10.4.6	Estimation in the Seemingly Unrelated Regressions (SUR) Model	362
10.5	MULTIVARIATE MULTIPLE REGRESSION: HYPOTHESIS TESTS	364
10.5.1	Test of Overall Regression	364
10.5.2	Test on a Subset of the x 's	367
10.6	MULTIVARIATE MULTIPLE REGRESSION: PREDICTION	370
10.6.1	Confidence Interval for $E(y_0)$	370
10.6.2	Prediction Interval for a Future Observation y_0	371

10.7	MEASURES OF ASSOCIATION BETWEEN THE y 's AND THE x 's	372
10.8	SUBSET SELECTION	374
10.8.1	Stepwise Procedures	374
10.8.2	All Possible Subsets	377
10.9	MULTIVARIATE REGRESSION: RANDOM x 's Problems	380 381
11	Canonical Correlation	385
11.1	INTRODUCTION	385
11.2	CANONICAL CORRELATIONS AND CANONICAL VARIATES	385
11.3	PROPERTIES OF CANONICAL CORRELATIONS	390
11.4	TESTS OF SIGNIFICANCE	391
11.4.1	Tests of No Relationship Between the y 's and the x 's	391
11.4.2	Test of Significance of Succeeding Canonical Correlations After the First	393
11.5	INTERPRETATION	395
11.5.1	Standardized Coefficients	396
11.5.2	Correlations between Variables and Canonical Variates	397
11.5.3	Rotation	397
11.5.4	Redundancy Analysis	398
11.6	RELATIONSHIPS OF CANONICAL CORRELATION ANALYSIS TO OTHER MULTIVARIATE TECHNIQUES	398
11.6.1	Regression	398
11.6.2	MANOVA and Discriminant Analysis	400
	Problems	402
12	Principal Component Analysis	405
12.1	INTRODUCTION	405
12.2	GEOMETRIC AND ALGEBRAIC BASES OF PRINCIPAL COMPONENTS	406
12.2.1	Geometric Approach	406
12.2.2	Algebraic Approach	410
12.3	PRINCIPAL COMPONENTS AND PERPENDICULAR REGRESSION	412
12.4	PLOTTING OF PRINCIPAL COMPONENTS	414

12.5	PRINCIPAL COMPONENTS FROM THE CORRELATION MATRIX	419
12.6	DECIDING HOW MANY COMPONENTS TO RETAIN	423
12.7	INFORMATION IN THE LAST FEW PRINCIPAL COMPONENTS	427
12.8	INTERPRETATION OF PRINCIPAL COMPONENTS	427
12.8.1	Special Patterns in S or R	427
12.8.2	Rotation	429
12.8.3	Correlations Between Variables and Principal Components	429
12.9	SELECTION OF VARIABLES	430
	Problems	432
13	Exploratory Factor Analysis	435
13.1	INTRODUCTION	435
13.2	ORTHOGONAL FACTOR MODEL	437
13.2.1	Model Definition and Assumptions	437
13.2.2	Nonuniqueness of Factor Loadings	441
13.3	ESTIMATION OF LOADINGS AND COMMUNALITIES	442
13.3.1	Principal Component Method	443
13.3.2	Principal Factor Method	448
13.3.3	Iterated Principal Factor Method	450
13.3.4	Maximum Likelihood Method	452
13.4	CHOOSING THE NUMBER OF FACTORS, m	453
13.5	ROTATION	457
13.5.1	Introduction	457
13.5.2	Orthogonal Rotation	458
13.5.3	Oblique Rotation	462
13.5.4	Interpretation	465
13.6	FACTOR SCORES	466
13.7	VALIDITY OF THE FACTOR ANALYSIS MODEL	470
13.8	RELATIONSHIP OF FACTOR ANALYSIS TO PRINCIPAL COMPONENT ANALYSIS	475
	Problems	476
14	Confirmatory Factor Analysis	479
14.1	INTRODUCTION	479
14.2	MODEL SPECIFICATION AND IDENTIFICATION	480

14.2.1	Confirmatory Factor Analysis Model	480
14.2.2	Identified Models	482
14.3	PARAMETER ESTIMATION AND MODEL ASSESSMENT	487
14.3.1	Maximum Likelihood Estimation	487
14.3.2	Least Squares Estimation	488
14.3.3	Model Assessment	489
14.4	INFERENCE FOR MODEL PARAMETERS	492
14.5	FACTOR SCORES	495
	Problems	496
15	Cluster Analysis	501
15.1	INTRODUCTION	501
15.2	MEASURES OF SIMILARITY OR DISSIMILARITY	502
15.3	HIERARCHICAL CLUSTERING	505
15.3.1	Introduction	505
15.3.2	Single Linkage (Nearest Neighbor)	506
15.3.3	Complete Linkage (Farthest Neighbor)	508
15.3.4	Average Linkage	511
15.3.5	Centroid	514
15.3.6	Median	514
15.3.7	Ward's Method	517
15.3.8	Flexible Beta Method	520
15.3.9	Properties of Hierarchical Methods	521
15.3.10	Divisive Methods	529
15.4	NONHIERARCHICAL METHODS	531
15.4.1	Partitioning	532
15.4.2	Other Methods	540
15.5	CHOOSING THE NUMBER OF CLUSTERS	544
15.6	CLUSTER VALIDITY	546
15.7	CLUSTERING VARIABLES	547
	Problems	548
16	Graphical Procedures	555
16.1	MULTIDIMENSIONAL SCALING	555
16.1.1	Introduction	555
16.1.2	Metric Multidimensional Scaling	556
16.1.3	Nonmetric Multidimensional Scaling	560
16.2	CORRESPONDENCE ANALYSIS	565

16.2.1	Introduction	565
16.2.2	Row and Column Profiles	566
16.2.3	Testing Independence	570
16.2.4	Coordinates for Plotting Row and Column Profiles	572
16.2.5	Multiple Correspondence Analysis	576
16.3	BIPLOTS	580
16.3.1	Introduction	580
16.3.2	Principal Component Plots	581
16.3.3	Singular Value Decomposition Plots	583
16.3.4	Coordinates	583
16.3.5	Other Methods	585
	Problems	588
	Appendix A: Tables	597
	Appendix B: Answers and Hints to Problems	637
	Appendix C: Data Sets and SAS Files	727
	References	728
	Index	745

Preface

We have long been fascinated by the interplay of variables in multivariate data and by the challenge of unraveling the effect of each variable. Our continuing objective in the third edition has been to present the power and utility of multivariate analysis in a highly readable format.

Practitioners and researchers in all applied disciplines often measure several variables on each subject or experimental unit. In some cases, it may be productive to isolate each variable in a system and study it separately. Typically, however, the variables are not only correlated with each other, but each variable is influenced by the other variables as it affects a test statistic or descriptive statistics. Thus, in many instances, the variables are intertwined in such a way that when analyzed individually they yield little information about the system. Using multivariate analysis, the variables can be examined simultaneously in order to access the key features of the process that produced them. The multivariate approach enables us to (1) explore the joint performance of the variables and (2) determine the effect of each variable in the presence of the others.

Multivariate analysis provides both descriptive and inferential procedures—we can search for patterns in the data or test hypotheses about patterns of a priori interest. With multivariate descriptive techniques, we can peer beneath the tangled web of variables on the surface and extract the essence of the system. Multivariate inferential procedures include hypothesis tests that (1) process any number of variables

without inflating the Type I error rate and (2) allow for whatever intercorrelations the variables possess. A wide variety of multivariate descriptive and inferential procedures is readily accessible in statistical software packages.

Our selection of topics for this volume reflects years of consulting with researchers in many fields of inquiry. A brief overview of multivariate analysis is given in Chapter 1. Chapter 2 reviews the fundamentals of matrix algebra. Chapters 3 and 4 give an introduction to sampling from multivariate populations. Chapters 5, 6, 7, 10, and 11 extend univariate procedures with one dependent variable (including *t*-tests, analysis of variance, tests on variances, multiple regression, and multiple correlation) to analogous multivariate techniques involving several dependent variables. A review of each univariate procedure is presented before covering the multivariate counterpart. These reviews may provide key insights that the student has missed in previous courses.

Chapters 8, 9, 12, 13, 14, 15, and 16 describe multivariate techniques that are not extensions of univariate procedures. In Chapters 8 and 9, we find functions of the variables that discriminate among groups in the data. In Chapters 12, 13, and 14 (new in the third edition), we find functions of the variables that reveal the basic dimensionality and characteristic patterns of the data, and we discuss procedures for finding the underlying latent variables of a system. In Chapters 15 and 16, we give methods for searching for groups in the data, and we provide plotting techniques that show relationships in a reduced dimensionality for various kinds of data.

In Appendix A, tables are provided for many multivariate distributions and tests. These enable the reader to conduct an exact test in many cases for which software packages provide only approximate tests. Appendix B gives answers and hints for most of the problems in the book.

Appendix C describes an ftp site that contains (1) all data sets and (2) SAS command files for all examples in the text. These command files can be adapted for use in working problems or in analyzing data sets encountered in applications.

To illustrate multivariate applications, we have provided many examples and exercises based on 60 real data sets from a wide variety of disciplines. A practitioner or consultant in multivariate analysis gains insights and acumen from long experience in working with data. It is not expected that a student can achieve this kind of seasoning in a one-semester class. However, the examples provide a good start, and further development is gained by working problems with the data sets. For example, in Chapters 12–14, the exercises cover several typical patterns in the covariance or correlation matrix. The student's intuition is expanded by associating these covariance patterns with the resulting configuration of the principal components or factors.

Although this is a methods book, we have included a few derivations. For some readers, an occasional proof provides insights obtainable in no other way. We hope that instructors who do not wish to use proofs will not be deterred by their presence. The proofs can easily be disregarded when reading the book.

Our objective has been to make the book accessible to readers who have taken as few as two statistical methods courses. The students in our classes in multivariate analysis include majors in statistics and majors from other departments. With the applied researcher in mind, we have provided careful intuitive explanations of the

concepts and have included many insights typically available only in journal articles or in the minds of practitioners.

Our overriding goal in preparation of this book has been clarity of exposition. We hope that students and instructors alike will find this multivariate text more comfortable than most. In the final stages of development of each edition, we asked our students for written reports on their initial reaction as they read each day's assignment. They made many comments that led to improvements in the manuscript. We will be very grateful if readers will take the time to notify us of errors or of other suggestions they might have for improvements.

We have tried to use standard mathematical and statistical notation as far as possible and to maintain consistency of notation throughout the book. We have refrained from the use of abbreviations and mnemonic devices. These save space when one is reading a book page by page, but they are annoying to those using a book as a reference.

Equations are numbered sequentially throughout a chapter; for example, (3.75) indicates the 75th numbered equation in Chapter 3. Tables and figures are also numbered sequentially throughout a chapter in the form "Table 3.9" or "Figure 3.1." Examples are not numbered sequentially; each example is identified by the same number as the section in which it appears and is placed at the end of the section.

When citing references in the text, we have used the standard format involving the year of publication. For a journal article, the year alone suffices, for example, Fisher (1936). But for books, we have included a page number, as in Seber (1984, p. 216).

This is the first volume of a two-volume set on multivariate analysis. The second volume is entitled *Multivariate Statistical Inference and Applications* by Alvin Rencher (Wiley, 1998). The two volumes are not necessarily sequential; they can be read independently. Al adopted the two-volume format in order to (1) provide broader coverage than would be possible in a single volume and (2) offer the reader a choice of approach.

The second volume includes proofs of many techniques covered in the first 13 chapters of the present volume and also introduces additional topics. The present volume includes many examples and problems using actual data sets, and there are fewer algebraic problems. The second volume emphasizes derivations of the results and contains fewer examples and problems with real data. The present volume has fewer references to the literature than the other volume, which includes a careful review of the latest developments and a more comprehensive bibliography. In this third edition, we have occasionally referred the reader to "Rencher (1998)" to note that added coverage of a certain subject is available in the second volume.

We are indebted to many individuals in the preparation of the first two editions of this book. Al's initial exposure to multivariate analysis came in courses taught by Rolf Bargmann at the University of Georgia and D.R. Jensen at Virginia Tech. Additional impetus to probe the subtleties of this field came from research conducted with Bruce Brown at BYU. William's interest and training in multivariate statistics were primarily influenced by Alvin Rencher and Yasuo Amemiya. We thank these mentors and colleagues and also thank Bruce Brown, Deane Branstetter, Del Scott,

Robert Smidt, and Ingram Olkin for reading various versions of the manuscript and making valuable suggestions.

We are grateful to the following colleagues and students at BYU who helped with computations and typing: Mitchell Tolland, Tawnia Newton, Marianne Matis Mohr, Gregg Littlefield, Suzanne Kimball, Wendy Nielsen, Tiffany Nordgren, David Whiting, Karla Wasden, Rachel Jones, Lonette Stoddard, Candace B. McNaughton, J. D. Williams, and Jonathan Christensen. We are grateful to the many readers who have pointed out errors or made suggestions for improvements. The book is better for their caring and their efforts.

THIRD EDITION

For the third edition, we have added a new chapter covering confirmatory factor analysis (Chapter 14). We have added new sections discussing Kronecker products and vec notation (Section 2.12), dynamic graphics (Section 3.5), transformations to normality (Section 4.5), classification trees (Section 9.7.4), seemingly unrelated regressions (Section 10.4.6), and prediction for multivariate multiple regression (Section 10.6). Additionally, we have updated and revised the graphics throughout the book and have substantially expanded the section discussing estimation for multivariate multiple regression (Section 10.4).

Many other additions and changes have been made in an effort to broaden the book's scope and improve its exposition. Additional problems have been added to accompany the new material. The new ftp site for the third edition can be found at: ftp://ftp.wiley.com/public/sci_tech_med/multivariate_analysis_3e.

We thank Jonathan Christensen for the countless ways he contributed to the revision, from updated graphics to technical and formatting assistance. We are grateful to Byron Gajewski, Scott Grimshaw, and Jeremy Yorgason, who have provided useful feedback on sections of the book that are new to this edition. We are also grateful to Scott Simpson and Storm Atwood for invaluable editing assistance and to the students of William's multivariate statistics course at Brigham Young University for pointing out errors and making suggestions for improvements in this edition. We are deeply appreciative of the support provided by the BYU Department of Statistics for the writing of the book.

Al dedicates this volume to his wife, LaRue, who has supplied much needed support and encouragement. William dedicates this volume to his wife Mary Elizabeth, who has provided both encouragement and careful feedback throughout the writing process.

ALVIN C. RENCHER & WILLIAM F. CHRISTENSEN

*Department of Statistics
Brigham Young University
Provo, Utah*

Acknowledgments

We wish to thank the authors, editors, and owners of copyrights for permission to reproduce the following materials:

- Figure 3.8 and Table 3.2, Kleiner and Hartigan (1981), Reprinted by permission of *Journal of the American Statistical Association*
- Table 3.3, Colvin et al. (1997), Reprinted by permission of T. S. Colvin and D. L. Karlen
- Table 3.4, Kramer and Jensen (1969a), Reprinted by permission of *Journal of Quality Technology*
- Table 3.5, Reaven and Miller (1979), Reprinted by permission of *Diabetologia*
- Table 3.6, Timm (1975), Reprinted by permission of Elsevier North-Holland Publishing Company
- Table 3.7, Elston and Grizzle (1962), Reprinted by permission of *Biometrics*
- Table 3.8, Frets (1921), Reprinted by permission of *Genetica*
- Table 3.9, O'Sullivan and Mahan (1966), Reprinted by permission of *American Journal of Clinical Nutrition*

- Table 4.2, Royston (1983), Reprinted by permission of *Applied Statistics*
- Table 5.1, Beall (1945), Reprinted by permission of *Psychometrika*
- Table 5.2, Hummel and Sligo (1971), Reprinted by permission of *Psychological Bulletin*
- Table 5.3, Kramer and Jensen (1969b), Reprinted by permission of *Journal of Quality Technology*
- Table 5.5, Lubischew (1962), Reprinted by permission of *Biometrics*
- Table 5.6, Travers (1939), Reprinted by permission of *Psychometrika*
- Table 5.7, Andrews and Herzberg (1985), Reprinted by permission of Springer-Verlag
- Table 5.8, Tintner (1946), Reprinted by permission of *Journal of the American Statistical Association*
- Table 5.9, Kramer (1972), Reprinted by permission of the author
- Table 5.10, Cameron and Pauling (1978), Reprinted by permission of National Academy of Sciences
- Table 6.2, Andrews and Herzberg (1985), Reprinted by permission of Springer-Verlag
- Table 6.3, Rencher and Scott (1990), Reprinted by permission of *Communications in Statistics: Simulation and Computation*
- Table 6.6, Posten (1962), Reprinted by permission of the author
- Table 6.8, Crowder and Hand (1990, pp. 21-29), Reprinted by permission of Routledge Chapman and Hall
- Table 6.12, Cochran and Cox (1957), Timm (1980), Reprinted by permission of John Wiley and Sons and Elsevier North-Holland Publishing Company
- Table 6.14, Timm (1980), Reprinted by permission of Elsevier North-Holland Publishing Company
- Table 6.16, Potthoff and Roy (1964), Reprinted by permission of Biometrika Trustees
- Table 6.17, Baten, Tack, and Baeder (1958), Reprinted by permission of *Quality Progress*
- Table 6.18, Keuls et al. (1984), Reprinted by permission of *Scientia Horticulturae*
- Table 6.19, Burdick (1979), Reprinted by permission of the author

- Table 6.20, Box (1950), Reprinted by permission of *Biometrics*
- Table 6.21, Rao (1948), Reprinted by permission of Biometrika Trustees
- Table 6.22, Cameron and Pauling (1978), Reprinted by permission of National Academy of Sciences
- Table 6.23, Williams and Izenman (1981), Reprinted by permission of Colorado State University
- Table 6.24, Beauchamp and Hoel (1973), Reprinted by permission of *Journal of Statistical Computation and Simulation*
- Table 6.25, Box (1950), Reprinted by permission of *Biometrics*
- Table 6.26, Grizzle and Allen (1969), Reprinted by permission of *Biometrics*
- Table 6.27, Crepeau et. al (1985), Reprinted by permission of *Biometrics*
- Table 6.28, Zerbe (1979a), Reprinted by permission of *Journal of the American Statistical Association*
- Table 6.29, Timm (1980), Reprinted by permission of Elsevier North-Holland Publishing Company
- Table 7.1, Siotani et al. (1963), Reprinted by permission of Institute of Statistical Mathematics
- Table 7.2, Reprinted by permission of R. J. Freund.
- Table 8.1, Kramer and Jensen (1969a), Reprinted by permission of *Journal of Quality Technology*
- Table 8.3, Reprinted by permission of G. R. Bryce and R. M. Barker
- Table 10.1, Box and Youle (1955), Reprinted by permission of *Biometrics*
- Tables 12.2, 12.3, and 12.4, Jeffers (1967), Reprinted by permission of *Applied Statistics*
- Table 13.1, Brown et al. (1984), Reprinted by permission of *Journal of Pascal, Ada, and Modula*
- Correlation matrix in Example 13.6, Brown, Strong, and Rencher (1973), Reprinted by permission of *The Journal of the Acoustical Society of America*
- Table 15.1, Hartigan (1975a), Reprinted by permission of John Wiley and Sons
- Table 15.3, Dawkins (1989), Reprinted by permission of *The American Statistician*
- Table 15.7, Hand et al. (1994), Reprinted by permission of D. J. Hand

- Table 15.12, Sokol and Rohlf (1981), Reprinted by permission of W. H. Freeman and Co.
- Table 15.13, Hand et al. (1994), Reprinted by permission of D. J. Hand
- Table 16.1, Kruskal and Wish (1978), Reprinted by permission of Sage Publications
- Tables 16.2 and 16.5, Hand et al. (1994), Reprinted by permission of D. J. Hand
- Table 16.13, Edwards and Kreiner (1983), Reprinted by permission of *Biometrika*
- Table 16.15, Hand et al. (1994), Reprinted by permission of D. J. Hand
- Table 16.16, Everitt (1987), Reprinted by permission of the author
- Table 16.17, Andrews and Herzberg (1985), Reprinted by permission of Springer Verlag
- Table 16.18, Clausen (1998), Reprinted by permission of Sage Publications
- Table 16.19, Andrews and Herzberg (1985), Reprinted by permission of Springer Verlag
- Table A.1, Mulholland (1977), Reprinted by permission of Biometrika Trustees
- Table A.2, D'Agostino and Pearson (1973), Reprinted by permission of Biometrika Trustees
- Table A.3, D'Agostino and Tietjen (1971), Reprinted by permission of Biometrika Trustees
- Table A.4, D'Agostino (1972), Reprinted by permission of Biometrika Trustees
- Table A.5, Mardia (1970, 1974), Reprinted by permission of Biometrika Trustees
- Table A.6, Barnett and Lewis (1978), Reprinted by permission of John Wiley and Sons
- Table A.7, Kramer and Jensen (1969a), Reprinted by permission of *Journal of Quality Technology*
- Table A.8, Bailey (1977), Reprinted by permission of *Journal of the American Statistical Association*
- Table A.9, Wall (1967), Reprinted by permission of the author, Albuquerque, NM
- Table A.10, Pearson and Hartley (1972) and Pillai (1964, 1965), Reprinted by permission of Biometrika Trustees

- Table A.11, Schuurmann et al. (1975), Reprinted by permission of *Journal of Statistical Computation and Simulation*
- Table A.12, Davis (1970a,b, 1980), Reprinted by permission of Biometrika Trustees
- Table A.13, Kleinbaum, Kupper, and Muller (1988), Reprinted by permission of PWS-KENT Publishing Company
- Table A.14, Lee et al. (1977), Reprinted by permission of Elsevier North-Holland Publishing Company
- Table A.15, Mathai and Katiyar (1979), Reprinted by permission of Biometrika Trustees

CHAPTER 1

INTRODUCTION

1.1 WHY MULTIVARIATE ANALYSIS?

Multivariate analysis consists of a collection of methods that can be used when several measurements are made on each individual or object in one or more samples. We will refer to the measurements as *variables* and to the individuals or objects as *units* (research units, sampling units, or experimental units) or *observations*. In practice, multivariate data sets are common, although they are not always analyzed as such. But the exclusive use of univariate procedures with such data is no longer excusable, given the availability of multivariate techniques and inexpensive computing power to carry them out.

Historically, the bulk of applications of multivariate techniques have been in the behavioral and biological sciences. However, interest in multivariate methods has now spread to numerous other fields of investigation. For example, we have collaborated on multivariate problems with researchers in education, chemistry, environmental science, physics, geology, medicine, engineering, law, business, literature, religion, public broadcasting, nursing, mining, linguistics, biology, psychology, and many other fields. Table 1.1 shows some examples of multivariate observations.

Table 1.1 Examples of Multivariate Data

Units	Variables
1. Students	Several exam scores in a single course
2. Students	Grades in mathematics, history, music, art, physics
3. People	Height, weight, percentage of body fat, resting heart rate
4. Skulls	Length, width, cranial capacity
5. Companies	Expenditures for advertising, labor, raw materials
6. Manufactured items	Various measurements to check on compliance with specifications
7. Applicants for bank loans	Income, education level, length of residence, savings account, current debt load
8. Segments of literature	Sentence length, frequency of usage of certain words and style characteristics
9. Human hairs	Composition of various elements
10. Birds	Lengths of various bones

The reader will notice that in some cases all the variables are measured in the same scale (see 1 and 2 in Table 1.1). In other cases, measurements are in different scales (see 3 in Table 1.1). In a few techniques such as profile analysis (Sections 5.9 and 6.8), the variables must be commensurate, that is, similar in scale of measurement; however, most multivariate methods do not require this.

Ordinarily the variables are measured simultaneously on each sampling unit. Typically, these variables are correlated. If this were not so, there would be little use for many of the techniques of multivariate analysis. We need to untangle the overlapping information provided by correlated variables and peer beneath the surface to see the underlying structure. Thus the goal of many multivariate approaches is *simplification*. We seek to express “what is going on” in terms of a reduced set of dimensions. Such multivariate techniques are *exploratory*; they essentially generate hypotheses rather than test them.

On the other hand, if our goal is a formal hypothesis test, we need a technique that will (1) allow several variables to be tested and still preserve the significance level and (2) do this for any intercorrelation structure of the variables. Many such tests are available.

As the two preceding paragraphs imply, multivariate analysis is concerned generally with two areas, *descriptive* and *inferential* statistics. In the descriptive realm, we often obtain optimal linear combinations of variables. The optimality criterion varies from one technique to another, depending on the goal in each case. Although linear combinations may seem too simple to reveal the underlying structure, we use them for two obvious reasons: (1) mathematical tractability (linear approximations are used throughout all science for the same reason) and (2) they often perform well in practice. These linear functions may also be useful as a follow-up to inferential procedures. When we have a statistically significant test result that compares sev-

eral groups, for example, we can find the linear combination (or combinations) of variables that led to rejection. Then the contribution of each variable to these linear combinations is of interest.

In the inferential area, many multivariate techniques are extensions of univariate procedures. In such cases we review the univariate procedure before presenting the analogous multivariate approach.

Multivariate inference is especially useful in curbing the researcher's natural tendency to read too much into the data. Total control is provided for experimentwise error rate; that is, no matter how many variables are tested simultaneously, the value of α (the significance level) remains at the level set by the researcher.

Some authors warn against applying the common multivariate techniques to data for which the measurement scale is not interval or ratio. It has been found, however, that many multivariate techniques give reliable results when applied to ordinal data.

For many years the applications lagged behind the theory because the computations were beyond the power of the available desk-top calculators. However, with modern computers, virtually any analysis one desires, no matter how many variables or observations are involved, can be quickly and easily carried out. Perhaps it is not premature to say that multivariate analysis has come of age.

1.2 PREREQUISITES

The mathematical prerequisite for reading this book is matrix algebra. Calculus is not used [with a brief exception in equation (4.29)]. But the basic tools of matrix algebra are essential, and the presentation in Chapter 2 is intended to be sufficiently complete so that the reader with no previous experience can master matrix manipulation up to the level required in this book.

The statistical prerequisites are basic familiarity with the normal distribution, t -tests, confidence intervals, multiple regression, and analysis of variance. These techniques are reviewed as each is extended to the analogous multivariate procedure.

This is a multivariate methods text. Most of the results are given without proof. In a few cases proofs are provided, but the major emphasis is on heuristic explanations. Our goal is an intuitive grasp of multivariate analysis, in the same mode as other statistical methods courses. Some problems are algebraic in nature, but the majority involve data sets to be analyzed.

1.3 OBJECTIVES

We have formulated three objectives that we hope this book will achieve for the reader. These objectives are based on long experience teaching a course in multivariate methods, consulting on multivariate problems with researchers in many fields, and guiding statistics graduate students as they consulted with similar clients.

The first objective is to gain a thorough understanding of the details of various multivariate techniques, their purposes, their assumptions, their limitations, and so

on. Many of these techniques are related, yet they differ in some essential ways. These similarities and differences are emphasized.

The second objective is to be able to select one or more appropriate techniques for a given multivariate data set. Recognizing the essential nature of a multivariate data set is the first step in a meaningful analysis. Basic types of multivariate data are introduced in Section 1.4.

The third objective is to be able to interpret the results of a computer analysis of a multivariate data set. Reading the manual for a particular program package is not enough to make an intelligent appraisal of the output. Achievement of the first objective and practice on data sets in the text should help achieve the third objective.

1.4 BASIC TYPES OF DATA AND ANALYSIS

We will list four basic types of (continuous) multivariate data and then briefly describe some possible analyses. Some writers would consider this an oversimplification and might prefer elaborate tree diagrams of data structure. However, many data sets can fit into one of these categories, and the simplicity of this structure makes it easier to remember. The four basic data types are as follows:

1. A single sample with several variables measured on each sampling unit (subject or object).
2. A single sample with two sets of variables measured on each unit.
3. Two samples with several variables measured on each unit.
4. Three or more samples with several variables measured on each unit.

Each data type has extensions, and various combinations of the four are possible. A few examples of analyses for each case will now be given:

1. A single sample with several variables measured on each sampling unit:
 - a. Test the hypothesis that the means of the variables have specified values.
 - b. Test the hypothesis that the variables are uncorrelated and have a common variance.
 - c. Find a small set of linear combinations of the original variables that summarizes most of the variation in the data (principal components).
 - d. Express the original variables as linear functions of a smaller set of underlying variables that account for the original variables and their inter-correlations (factor analysis).
2. A single sample with two sets of variables measured on each unit:
 - a. Determine the number, the size, and the nature of relationships between the two sets of variables (canonical correlation). For example, we may

wish to relate a set of interest variables to a set of achievement variables. How much overall correlation is there between these two sets?

- b.** Find a model to predict one set of variables from the other set (multivariate multiple regression).
- 3.** Two samples with several variables measured on each unit:
- a.** Compare the means of the variables across the two samples (Hotelling's T^2 -test).
 - b.** Find a linear combination of the variables that best separates the two samples (discriminant analysis).
 - c.** Find a function of the variables that will accurately allocate the units into the two groups (classification analysis).
- 4.** Three or more samples with several variables measured on each unit:
- a.** Compare the means of the variables across the groups (multivariate analysis of variance).
 - b.** Extension of 3b to more than two groups.
 - c.** Extension of 3c to more than two groups.

CHAPTER 2

MATRIX ALGEBRA

2.1 INTRODUCTION

This chapter introduces the basic elements of matrix algebra used in the remainder of this book. It is essentially a review of the requisite matrix tools and is not intended to be a complete development. However, it is sufficiently self-contained so that those with no previous exposure to the subject should need no other reference. Anyone unfamiliar with matrix algebra should plan to work most of the problems entailing numerical illustrations. It would also be helpful to explore some of the problems involving general matrix manipulation.

With the exception of a few derivations that seemed instructive, most of the results are given without proof. Some additional proofs are requested in the problems. For the remaining proofs, see any general text on matrix theory or one of the specialized matrix texts oriented to statistics, such as Graybill (1969), Searle (1982), or Harville (1997).

2.2 NOTATION AND BASIC DEFINITIONS

2.2.1 Matrices, Vectors, and Scalars

A *matrix* is a rectangular or square array of numbers or variables arranged in rows and columns. We use uppercase boldface letters to represent matrices. All entries in matrices will be real numbers or variables representing real numbers. The elements of a matrix are displayed in brackets. For example, the ACT score and GPA for three students can be conveniently listed in the following matrix:

$$\mathbf{A} = \begin{pmatrix} 23 & 3.54 \\ 29 & 3.81 \\ 18 & 2.75 \end{pmatrix}. \quad (2.1)$$

The elements of \mathbf{A} can also be variables, representing possible values of ACT and GPA for three students:

$$\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \\ a_{31} & a_{32} \end{pmatrix}. \quad (2.2)$$

In this double-subscript notation for the elements of a matrix, the first subscript indicates the row; the second identifies the column. The matrix \mathbf{A} in (2.2) could also be expressed as

$$\mathbf{A} = (a_{ij}), \quad (2.3)$$

where a_{ij} is a general element.

With three rows and two columns, the matrix \mathbf{A} in (2.1) or (2.2) is said to be 3×2 . In general, if a matrix \mathbf{A} has n rows and p columns, it is said to be $n \times p$. Alternatively, we say the *size* of \mathbf{A} is $n \times p$.

A *vector* is a matrix with a single column or row. The following could be the test scores of a student in a course in multivariate analysis:

$$\mathbf{x} = \begin{pmatrix} 98 \\ 86 \\ 93 \\ 97 \end{pmatrix}. \quad (2.4)$$

Variable elements in a vector can be identified by a single subscript:

$$\mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{pmatrix}. \quad (2.5)$$

We use lowercase boldface letters for column vectors. Row vectors are expressed as

$$\mathbf{x}' = (x_1, x_2, x_3, x_4) \text{ or as } \mathbf{x}' = (x_1 \ x_2 \ x_3 \ x_4),$$

where \mathbf{x}' indicates the *transpose* of \mathbf{x} . The transpose operation is defined in Section 2.2.3.

Geometrically, a vector with p elements identifies a point in a p -dimensional space. The elements in the vector are the coordinates of the point. In (2.35) in Section 2.3.3, we define the distance from the origin to the point. In Section 3.13, we define the distance between two vectors. In some cases, we will be interested in a directed line segment or arrow from the origin to the point.

A single real number is called a *scalar*, to distinguish it from a vector or matrix. Thus 2, -4, and 125 are scalars. A variable representing a scalar will usually be denoted by a lowercase nonbolded letter, such as $a = 5$. A product involving vectors and matrices may reduce to a matrix of size 1×1 , which then becomes a scalar.

2.2.2 Equality of Vectors and Matrices

Two matrices are equal if they are the same size and the elements in corresponding positions are equal. Thus if $\mathbf{A} = (a_{ij})$ and $\mathbf{B} = (b_{ij})$, then $\mathbf{A} = \mathbf{B}$ if $a_{ij} = b_{ij}$ for all i and j . For example, let

$$\mathbf{A} = \begin{pmatrix} 3 & -2 & 4 \\ 1 & 3 & 7 \end{pmatrix}, \quad \mathbf{B} = \begin{pmatrix} 3 & 1 \\ -2 & 3 \\ 4 & 7 \end{pmatrix},$$

$$\mathbf{C} = \begin{pmatrix} 3 & -2 & 4 \\ 1 & 3 & 7 \end{pmatrix}, \quad \mathbf{D} = \begin{pmatrix} 3 & -2 & 4 \\ 1 & 3 & 6 \end{pmatrix}.$$

Then $\mathbf{A} = \mathbf{C}$. But even though \mathbf{A} and \mathbf{B} have the same elements, $\mathbf{A} \neq \mathbf{B}$ because the two matrices are not the same size. Likewise, $\mathbf{A} \neq \mathbf{D}$ because $a_{23} \neq d_{23}$. Thus two matrices of the same size are unequal if they differ in a single position.

2.2.3 Transpose and Symmetric Matrices

The *transpose* of a matrix \mathbf{A} , denoted by \mathbf{A}' , is obtained from \mathbf{A} by interchanging rows and columns. Thus the columns of \mathbf{A}' are the rows of \mathbf{A} , and the rows of \mathbf{A}' are the columns of \mathbf{A} . The following examples illustrate the transpose of a matrix or vector:

$$\mathbf{A} = \begin{pmatrix} -5 & 2 & 4 \\ 3 & 6 & -2 \end{pmatrix}, \quad \mathbf{A}' = \begin{pmatrix} -5 & 3 \\ 2 & 6 \\ 4 & -2 \end{pmatrix},$$

$$\mathbf{B} = \begin{pmatrix} 2 & -3 \\ 4 & 1 \end{pmatrix}, \quad \mathbf{B}' = \begin{pmatrix} 2 & 4 \\ -3 & 1 \end{pmatrix},$$

$$\mathbf{a} = \begin{pmatrix} 2 \\ -3 \\ 1 \end{pmatrix}, \quad \mathbf{a}' = (2, -3, 1).$$

The transpose operation does not change a scalar, since it has only one row and one column.

If the transpose operator is applied twice to any matrix, the result is the original matrix:

$$(\mathbf{A}')' = \mathbf{A}. \quad (2.6)$$

If the transpose of a matrix is the same as the original matrix, the matrix is said to be *symmetric*; that is, \mathbf{A} is symmetric if $\mathbf{A} = \mathbf{A}'$. For example,

$$\mathbf{A} = \begin{pmatrix} 3 & -2 & 4 \\ -2 & 10 & -7 \\ 4 & -7 & 9 \end{pmatrix}, \quad \mathbf{A}' = \begin{pmatrix} 3 & -2 & 4 \\ -2 & 10 & -7 \\ 4 & -7 & 9 \end{pmatrix}.$$

Clearly, all symmetric matrices are square.

2.2.4 Special Matrices

The *diagonal* of a $p \times p$ square matrix \mathbf{A} consists of the elements $a_{11}, a_{22}, \dots, a_{pp}$. For example, in the matrix

$$\mathbf{A} = \begin{pmatrix} 5 & -2 & 4 \\ 7 & 9 & 3 \\ -6 & 8 & 1 \end{pmatrix},$$

the elements 5, 9, and 1 lie on the diagonal. If a matrix contains zeros in all off-diagonal positions, it is said to be a *diagonal matrix*. An example of a diagonal matrix is

$$\mathbf{D} = \begin{pmatrix} 10 & 0 & 0 & 0 \\ 0 & -3 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 7 \end{pmatrix}.$$

This matrix could also be denoted as

$$\mathbf{D} = \text{diag}(10, -3, 0, 7). \quad (2.7)$$

A diagonal matrix can be formed from any square matrix by replacing off-diagonal elements by 0's. This is denoted by $\text{diag}(\mathbf{A})$. Thus for the above matrix \mathbf{A} , we have

$$\text{diag}(\mathbf{A}) = \text{diag}\left(\begin{pmatrix} 5 & -2 & 4 \\ 7 & 9 & 3 \\ -6 & 8 & 1 \end{pmatrix} \right) = \begin{pmatrix} 5 & 0 & 0 \\ 0 & 9 & 0 \\ 0 & 0 & 1 \end{pmatrix}. \quad (2.8)$$

A diagonal matrix with a 1 in each diagonal position is called an *identity matrix* and is denoted by \mathbf{I} . For example, a 3×3 identity matrix is given by

$$\mathbf{I} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}. \quad (2.9)$$

An *upper triangular matrix* is a square matrix with zeros below the diagonal, for example,

$$\mathbf{T} = \begin{pmatrix} 8 & 3 & 4 & 7 \\ 0 & 0 & -2 & 3 \\ 0 & 0 & 5 & 1 \\ 0 & 0 & 0 & 6 \end{pmatrix}. \quad (2.10)$$

A *lower triangular matrix* is defined similarly.

A vector of 1's will be denoted by \mathbf{j} :

$$\mathbf{j} = \begin{pmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{pmatrix}. \quad (2.11)$$

A square matrix of 1's is denoted by \mathbf{J} . For example, a 3×3 matrix \mathbf{J} is given by

$$\mathbf{J} = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}. \quad (2.12)$$

Finally, we denote a vector of zeros by $\mathbf{0}$ and a matrix of zeros by \mathbf{O} . For example,

$$\mathbf{0} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}, \quad \mathbf{O} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}. \quad (2.13)$$

2.3 OPERATIONS

2.3.1 Summation and Product Notation

For completeness, we review the standard mathematical notation for sums and products. The sum of a sequence of numbers a_1, a_2, \dots, a_n is indicated by

$$\sum_{i=1}^n a_i = a_1 + a_2 + \cdots + a_n.$$

If the n numbers are all the same, then $\sum_{i=1}^n a = a + a + \cdots + a = na$. The sum of all the numbers in an array with double subscripts, such as

$$\begin{array}{ccc} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23}, \end{array}$$

is indicated by

$$\sum_{i=1}^2 \sum_{j=1}^3 a_{ij} = a_{11} + a_{12} + a_{13} + a_{21} + a_{22} + a_{23}.$$

This is sometimes abbreviated to

$$\sum_{i=1}^2 \sum_{j=1}^3 a_{ij} = \sum_{ij} a_{ij}.$$

The product of a sequence of numbers a_1, a_2, \dots, a_n is indicated by

$$\prod_{i=1}^n a_i = (a_1)(a_2) \cdots (a_n).$$

If the n numbers are all equal, the product becomes $\prod_{i=1}^n a = (a)(a) \cdots (a) = a^n$.

2.3.2 Addition of Matrices and Vectors

If two matrices (or two vectors) are the same size, their *sum* is found by adding corresponding elements, that is, if \mathbf{A} is $n \times p$ and \mathbf{B} is $n \times p$, then $\mathbf{C} = \mathbf{A} + \mathbf{B}$ is also $n \times p$ and is found as $(c_{ij}) = (a_{ij} + b_{ij})$. For example,

$$\begin{pmatrix} -2 & 5 \\ 3 & 1 \\ 7 & -6 \end{pmatrix} + \begin{pmatrix} 3 & -2 \\ 4 & 5 \\ 10 & -3 \end{pmatrix} = \begin{pmatrix} 1 & 3 \\ 7 & 6 \\ 17 & -9 \end{pmatrix},$$

$$\begin{pmatrix} 1 \\ 3 \\ 7 \end{pmatrix} + \begin{pmatrix} 5 \\ -1 \\ 3 \end{pmatrix} = \begin{pmatrix} 6 \\ 2 \\ 10 \end{pmatrix}.$$

Similarly, the *difference* between two matrices or two vectors of the same size is found by subtracting corresponding elements. Thus $\mathbf{C} = \mathbf{A} - \mathbf{B}$ is found as $(c_{ij}) = (a_{ij} - b_{ij})$. For example,

$$(3 \quad 9 \quad -4) - (5 \quad -4 \quad 2) = (-2 \quad 13 \quad -6).$$

If two matrices are identical, their difference is a zero matrix; that is, $\mathbf{A} = \mathbf{B}$ implies $\mathbf{A} - \mathbf{B} = \mathbf{O}$. For example,

$$\begin{pmatrix} 3 & -2 & 4 \\ 6 & 7 & 5 \end{pmatrix} - \begin{pmatrix} 3 & -2 & 4 \\ 6 & 7 & 5 \end{pmatrix} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

Matrix addition is commutative:

$$\mathbf{A} + \mathbf{B} = \mathbf{B} + \mathbf{A}. \quad (2.14)$$

The transpose of the sum (difference) of two matrices is the sum (difference) of the transposes:

$$(\mathbf{A} + \mathbf{B})' = \mathbf{A}' + \mathbf{B}', \quad (2.15)$$

$$(\mathbf{A} - \mathbf{B})' = \mathbf{A}' - \mathbf{B}', \quad (2.16)$$

$$(\mathbf{x} + \mathbf{y})' = \mathbf{x}' + \mathbf{y}', \quad (2.17)$$

$$(\mathbf{x} - \mathbf{y})' = \mathbf{x}' - \mathbf{y}'. \quad (2.18)$$

2.3.3 Multiplication of Matrices and Vectors

In order for the product \mathbf{AB} to be defined, the number of columns in \mathbf{A} must be the same as the number of rows in \mathbf{B} , in which case \mathbf{A} and \mathbf{B} are said to be *conformable*. Then the (ij) th element of $\mathbf{C} = \mathbf{AB}$ is

$$c_{ij} = \sum_k a_{ik} b_{kj}. \quad (2.19)$$

Thus c_{ij} is the sum of products of the i th row of \mathbf{A} and the j th column of \mathbf{B} . We therefore multiply each row of \mathbf{A} by each column of \mathbf{B} , and the size of \mathbf{AB} consists of the number of rows of \mathbf{A} and the number of columns of \mathbf{B} . Thus, if \mathbf{A} is $n \times m$ and \mathbf{B} is $m \times p$, then $\mathbf{C} = \mathbf{AB}$ is $n \times p$. For example, if

$$\mathbf{A} = \begin{pmatrix} 2 & 1 & 3 \\ 4 & 6 & 5 \\ 7 & 2 & 3 \\ 1 & 3 & 2 \end{pmatrix} \quad \text{and} \quad \mathbf{B} = \begin{pmatrix} 1 & 4 \\ 2 & 6 \\ 3 & 8 \end{pmatrix},$$

then

$$\begin{aligned} \mathbf{C} = \mathbf{AB} &= \begin{pmatrix} 2 \cdot 1 + 1 \cdot 2 + 3 \cdot 3 & 2 \cdot 4 + 1 \cdot 6 + 3 \cdot 8 \\ 4 \cdot 1 + 6 \cdot 2 + 5 \cdot 3 & 4 \cdot 4 + 6 \cdot 6 + 5 \cdot 8 \\ 7 \cdot 1 + 2 \cdot 2 + 3 \cdot 3 & 7 \cdot 4 + 2 \cdot 6 + 3 \cdot 8 \\ 1 \cdot 1 + 3 \cdot 2 + 2 \cdot 3 & 1 \cdot 4 + 3 \cdot 6 + 2 \cdot 8 \end{pmatrix} \\ &= \begin{pmatrix} 13 & 38 \\ 31 & 92 \\ 20 & 64 \\ 13 & 38 \end{pmatrix}. \end{aligned}$$

Note that \mathbf{A} is 4×3 , \mathbf{B} is 3×2 , and \mathbf{AB} is 4×2 . In this case, \mathbf{AB} is of a different size than either \mathbf{A} or \mathbf{B} .

If \mathbf{A} and \mathbf{B} are both $n \times n$, then \mathbf{AB} is also $n \times n$. Clearly, \mathbf{A}^2 is defined only if \mathbf{A} is square.

In some cases \mathbf{AB} is defined, but \mathbf{BA} is not defined. In the above example, \mathbf{BA} cannot be found because \mathbf{B} is 3×2 and \mathbf{A} is 4×3 and a row of \mathbf{B} cannot be multiplied by a column of \mathbf{A} . Sometimes \mathbf{AB} and \mathbf{BA} are both defined but are different in size. For example, if \mathbf{A} is 2×4 and \mathbf{B} is 4×2 , then \mathbf{AB} is 2×2 and \mathbf{BA} is 4×4 . If \mathbf{A} and \mathbf{B} are square and the same size, then \mathbf{AB} and \mathbf{BA} are both defined. However,

$$\mathbf{AB} \neq \mathbf{BA}, \quad (2.20)$$

except for a few special cases. For example, let

$$\mathbf{A} = \begin{pmatrix} 1 & 3 \\ 2 & 4 \end{pmatrix}, \quad \mathbf{B} = \begin{pmatrix} 1 & -2 \\ 3 & 5 \end{pmatrix}.$$

Then

$$\mathbf{AB} = \begin{pmatrix} 10 & 13 \\ 14 & 16 \end{pmatrix}, \quad \mathbf{BA} = \begin{pmatrix} -3 & -5 \\ 13 & 29 \end{pmatrix}.$$

Thus we must be careful to specify the order of multiplication. If we wish to multiply both sides of a matrix equation by a matrix, we must multiply “on the left” or “on the right” and be consistent on both sides of the equation.

Multiplication is distributive over addition or subtraction:

$$\mathbf{A}(\mathbf{B} + \mathbf{C}) = \mathbf{AB} + \mathbf{AC}, \quad (2.21)$$

$$\mathbf{A}(\mathbf{B} - \mathbf{C}) = \mathbf{AB} - \mathbf{AC}, \quad (2.22)$$

$$(\mathbf{A} + \mathbf{B})\mathbf{C} = \mathbf{AC} + \mathbf{BC}, \quad (2.23)$$

$$(\mathbf{A} - \mathbf{B})\mathbf{C} = \mathbf{AC} - \mathbf{BC}. \quad (2.24)$$

Note that, in general, because of (2.20),

$$\mathbf{A}(\mathbf{B} + \mathbf{C}) \neq \mathbf{BA} + \mathbf{CA}. \quad (2.25)$$

Using the distributive law, we can expand products such as $(\mathbf{A} - \mathbf{B})(\mathbf{C} - \mathbf{D})$ to obtain

$$\begin{aligned} (\mathbf{A} - \mathbf{B})(\mathbf{C} - \mathbf{D}) &= (\mathbf{A} - \mathbf{B})\mathbf{C} - (\mathbf{A} - \mathbf{B})\mathbf{D} && [\text{by (2.22)}] \\ &= \mathbf{AC} - \mathbf{BC} - \mathbf{AD} + \mathbf{BD} && [\text{by (2.24)}]. \end{aligned} \quad (2.26)$$

The transpose of a product is the product of the transposes in reverse order:

$$(\mathbf{AB})' = \mathbf{B}'\mathbf{A}'. \quad (2.27)$$

Note that (2.27) holds as long as \mathbf{A} and \mathbf{B} are conformable. They need not be square.

Multiplication involving vectors follows the same rules as for matrices. Suppose \mathbf{A} is $n \times p$, \mathbf{a} is $p \times 1$, \mathbf{b} is $p \times 1$, and \mathbf{c} is $n \times 1$. Then some possible products are \mathbf{Ab} , $\mathbf{c}'\mathbf{A}$, $\mathbf{a}'\mathbf{b}$, $\mathbf{b}'\mathbf{a}$, and \mathbf{ab}' . For example, let

$$\mathbf{A} = \begin{pmatrix} 3 & -2 & 4 \\ 1 & 3 & 5 \end{pmatrix}, \quad \mathbf{a} = \begin{pmatrix} 1 \\ -2 \\ 3 \end{pmatrix}, \quad \mathbf{b} = \begin{pmatrix} 2 \\ 3 \\ 4 \end{pmatrix}, \quad \mathbf{c} = \begin{pmatrix} 2 \\ -5 \end{pmatrix}.$$

Then

$$\begin{aligned}
 \mathbf{Ab} &= \begin{pmatrix} 3 & -2 & 4 \\ 1 & 3 & 5 \end{pmatrix} \begin{pmatrix} 2 \\ 3 \\ 4 \end{pmatrix} = \begin{pmatrix} 16 \\ 31 \end{pmatrix} \\
 \mathbf{c}'\mathbf{A} &= (2 \quad -5) \begin{pmatrix} 3 & -2 & 4 \\ 1 & 3 & 5 \end{pmatrix} = (1 \quad -19 \quad -17) \\
 \mathbf{c}'\mathbf{Ab} &= (2 \quad -5) \begin{pmatrix} 3 & -2 & 4 \\ 1 & 3 & 5 \end{pmatrix} \begin{pmatrix} 2 \\ 3 \\ 4 \end{pmatrix} \\
 &= (2 \quad -5) \begin{pmatrix} 16 \\ 31 \end{pmatrix} = -123 \\
 \mathbf{a}'\mathbf{b} &= (1 \quad -2 \quad 3) \begin{pmatrix} 2 \\ 3 \\ 4 \end{pmatrix} = 8 \\
 \mathbf{b}'\mathbf{a} &= (2 \quad 3 \quad 4) \begin{pmatrix} 1 \\ -2 \\ 3 \end{pmatrix} = 8 \\
 \mathbf{ab}' &= \begin{pmatrix} 1 \\ -2 \\ 3 \end{pmatrix} (2 \quad 3 \quad 4) = \begin{pmatrix} 2 & 3 & 4 \\ -4 & -6 & -8 \\ 6 & 9 & 12 \end{pmatrix} \\
 \mathbf{ac}' &= \begin{pmatrix} 1 \\ -2 \\ 3 \end{pmatrix} (2 \quad -5) = \begin{pmatrix} 2 & -5 \\ -4 & 10 \\ 6 & -15 \end{pmatrix}.
 \end{aligned}$$

Note that \mathbf{Ab} is a column vector, $\mathbf{c}'\mathbf{A}$ is a row vector, $\mathbf{c}'\mathbf{Ab}$ is a scalar, and $\mathbf{a}'\mathbf{b} = \mathbf{b}'\mathbf{a}$. The triple product $\mathbf{c}'\mathbf{Ab}$ was obtained as $\mathbf{c}'(\mathbf{Ab})$. The same result would be obtained if we multiplied in the order $(\mathbf{c}'\mathbf{A})\mathbf{b}$:

$$(\mathbf{c}'\mathbf{A})\mathbf{b} = (1 \quad -19 \quad -17) \begin{pmatrix} 2 \\ 3 \\ 4 \end{pmatrix} = -123.$$

This is true in general for a triple product:

$$\mathbf{ABC} = \mathbf{A}(\mathbf{BC}) = (\mathbf{AB})\mathbf{C}. \quad (2.28)$$

Thus multiplication of three matrices can be defined in terms of the product of two matrices, since (fortunately) it does not matter which two are multiplied first. Note that \mathbf{A} and \mathbf{B} must be conformable for multiplication, and \mathbf{B} and \mathbf{C} must be conformable. For example, if \mathbf{A} is $n \times p$, \mathbf{B} is $p \times q$, and \mathbf{C} is $q \times m$, then both multiplications are possible and the product \mathbf{ABC} is $n \times m$.

We can sometimes factor a sum of triple products on both the right and left sides. For example,

$$\mathbf{ABC} + \mathbf{ADC} = \mathbf{A}(\mathbf{B} + \mathbf{D})\mathbf{C}. \quad (2.29)$$

As another illustration, let \mathbf{X} be $n \times p$ and \mathbf{A} be $n \times n$. Then

$$\mathbf{X}'\mathbf{X} - \mathbf{X}'\mathbf{AX} = \mathbf{X}'(\mathbf{X} - \mathbf{AX}) = \mathbf{X}'(\mathbf{I} - \mathbf{A})\mathbf{X}. \quad (2.30)$$

If \mathbf{a} and \mathbf{b} are both $n \times 1$, then

$$\mathbf{a}'\mathbf{b} = a_1b_1 + a_2b_2 + \cdots + a_nb_n \quad (2.31)$$

is a sum of products and is a scalar. On the other hand, \mathbf{ab}' is defined for any size \mathbf{a} and \mathbf{b} and is a matrix, either rectangular or square:

$$\begin{aligned} \mathbf{ab}' &= \begin{pmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{pmatrix} \begin{pmatrix} b_1 & b_2 & \cdots & b_p \end{pmatrix} \\ &= \begin{pmatrix} a_1b_1 & a_1b_2 & \cdots & a_1b_p \\ a_2b_1 & a_2b_2 & \cdots & a_2b_p \\ \vdots & \vdots & & \vdots \\ a_nb_1 & a_nb_2 & \cdots & a_nb_p \end{pmatrix}. \end{aligned} \quad (2.32)$$

Similarly,

$$\mathbf{a}'\mathbf{a} = a_1^2 + a_2^2 + \cdots + a_n^2, \quad (2.33)$$

$$\mathbf{aa}' = \begin{pmatrix} a_1^2 & a_1a_2 & \cdots & a_1a_n \\ a_2a_1 & a_2^2 & \cdots & a_2a_n \\ \vdots & \vdots & & \vdots \\ a_na_1 & a_na_2 & \cdots & a_n^2 \end{pmatrix}. \quad (2.34)$$

Thus $\mathbf{a}'\mathbf{a}$ is a sum of squares and \mathbf{aa}' is a square (symmetric) matrix. The products $\mathbf{a}'\mathbf{a}$ and \mathbf{aa}' are sometimes referred to as the *dot product* and *matrix product*, respectively. The square root of the sum of squares of the elements of \mathbf{a} is the *distance* from the origin to the point \mathbf{a} and is also referred to as the *length* of \mathbf{a} :

$$\text{Length of } \mathbf{a} = \sqrt{\mathbf{a}'\mathbf{a}} = \sqrt{\sum_{i=1}^n a_i^2}. \quad (2.35)$$

As special cases of (2.33) and (2.34), note that if \mathbf{j} is $n \times 1$, then

$$\mathbf{j}'\mathbf{j} = n, \quad \mathbf{jj}' = \begin{pmatrix} 1 & 1 & \cdots & 1 \\ 1 & 1 & \cdots & 1 \\ \vdots & \vdots & & \vdots \\ 1 & 1 & \cdots & 1 \end{pmatrix} = \mathbf{J}, \quad (2.36)$$

where \mathbf{j} and \mathbf{J} were defined in (2.11) and (2.12). If \mathbf{a} is $n \times 1$ and \mathbf{A} is $n \times p$, then

$$\mathbf{a}'\mathbf{j} = \mathbf{j}'\mathbf{a} = \sum_{i=1}^n a_i, \quad (2.37)$$

$$\mathbf{j}'\mathbf{A} = (\sum_i a_{i1}, \sum_i a_{i2}, \dots, \sum_i a_{ip}), \quad \mathbf{A}\mathbf{j} = \begin{pmatrix} \sum_j a_{1j} \\ \sum_j a_{2j} \\ \vdots \\ \sum_j a_{nj} \end{pmatrix}. \quad (2.38)$$

Thus $\mathbf{a}'\mathbf{j}$ is the sum of the elements in \mathbf{a} , $\mathbf{j}'\mathbf{A}$ contains the column sums of \mathbf{A} , and $\mathbf{A}\mathbf{j}$ contains the row sums of \mathbf{A} . In $\mathbf{a}'\mathbf{j}$, the vector \mathbf{j} is $n \times 1$; in $\mathbf{j}'\mathbf{A}$, the vector \mathbf{j} is $n \times 1$; and in $\mathbf{A}\mathbf{j}$, the vector \mathbf{j} is $p \times 1$.

Since $\mathbf{a}'\mathbf{b}$ is a scalar, it is equal to its transpose:

$$\mathbf{a}'\mathbf{b} = (\mathbf{a}'\mathbf{b})' = \mathbf{b}'(\mathbf{a}')' = \mathbf{b}'\mathbf{a}. \quad (2.39)$$

This allows us to write $(\mathbf{a}'\mathbf{b})^2$ in the form

$$(\mathbf{a}'\mathbf{b})^2 = (\mathbf{a}'\mathbf{b})(\mathbf{a}'\mathbf{b}) = (\mathbf{a}'\mathbf{b})(\mathbf{b}'\mathbf{a}) = \mathbf{a}'(\mathbf{b}\mathbf{b}')\mathbf{a}. \quad (2.40)$$

From (2.18), (2.26), and (2.39) we obtain

$$(\mathbf{x} - \mathbf{y})'(\mathbf{x} - \mathbf{y}) = \mathbf{x}'\mathbf{x} - 2\mathbf{x}'\mathbf{y} + \mathbf{y}'\mathbf{y}. \quad (2.41)$$

Note that in analogous expressions with matrices, however, the two middle terms cannot be combined:

$$\begin{aligned} (\mathbf{A} - \mathbf{B})'(\mathbf{A} - \mathbf{B}) &= \mathbf{A}'\mathbf{A} - \mathbf{A}'\mathbf{B} - \mathbf{B}'\mathbf{A} + \mathbf{B}'\mathbf{B}, \\ (\mathbf{A} - \mathbf{B})^2 &= (\mathbf{A} - \mathbf{B})(\mathbf{A} - \mathbf{B}) = \mathbf{A}^2 - \mathbf{AB} - \mathbf{BA} + \mathbf{B}^2. \end{aligned}$$

If \mathbf{a} and $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$ are all $p \times 1$ and \mathbf{A} is $p \times p$, we obtain the following factoring results as extensions of (2.21) and (2.29):

$$\sum_{i=1}^n \mathbf{a}'\mathbf{x}_i = \mathbf{a}' \sum_{i=1}^n \mathbf{x}_i, \quad (2.42)$$

$$\sum_{i=1}^n \mathbf{A}\mathbf{x}_i = \mathbf{A} \sum_{i=1}^n \mathbf{x}_i, \quad (2.43)$$

$$\sum_{i=1}^n (\mathbf{a}'\mathbf{x}_i)^2 = \mathbf{a}' \left(\sum_{i=1}^n \mathbf{x}_i \mathbf{x}_i' \right) \mathbf{a} \quad [\text{by (2.40)}], \quad (2.44)$$

$$\sum_{i=1}^n \mathbf{A}\mathbf{x}_i (\mathbf{A}\mathbf{x}_i)' = \mathbf{A} \left(\sum_{i=1}^n \mathbf{x}_i \mathbf{x}_i' \right) \mathbf{A}'. \quad (2.45)$$

We can express matrix multiplication in terms of row vectors and column vectors. If \mathbf{a}'_i is the i th row of \mathbf{A} and \mathbf{b}_j is the j th column of \mathbf{B} , then the (ij) th element of

\mathbf{AB} is $\mathbf{a}_i' \mathbf{b}_j$. For example, if \mathbf{A} has three rows and \mathbf{B} has two columns,

$$\mathbf{A} = \begin{pmatrix} \mathbf{a}_1' \\ \mathbf{a}_2' \\ \mathbf{a}_3' \end{pmatrix}, \quad \mathbf{B} = (\mathbf{b}_1, \mathbf{b}_2),$$

then the product \mathbf{AB} can be written as

$$\mathbf{AB} = \begin{pmatrix} \mathbf{a}_1' \mathbf{b}_1 & \mathbf{a}_1' \mathbf{b}_2 \\ \mathbf{a}_2' \mathbf{b}_1 & \mathbf{a}_2' \mathbf{b}_2 \\ \mathbf{a}_3' \mathbf{b}_1 & \mathbf{a}_3' \mathbf{b}_2 \end{pmatrix}. \quad (2.46)$$

This can be expressed in terms of the rows of \mathbf{A} :

$$\mathbf{AB} = \begin{pmatrix} \mathbf{a}_1'(\mathbf{b}_1, \mathbf{b}_2) \\ \mathbf{a}_2'(\mathbf{b}_1, \mathbf{b}_2) \\ \mathbf{a}_3'(\mathbf{b}_1, \mathbf{b}_2) \end{pmatrix} = \begin{pmatrix} \mathbf{a}_1' \mathbf{B} \\ \mathbf{a}_2' \mathbf{B} \\ \mathbf{a}_3' \mathbf{B} \end{pmatrix} = \begin{pmatrix} \mathbf{a}_1' \\ \mathbf{a}_2' \\ \mathbf{a}_3' \end{pmatrix} \mathbf{B}. \quad (2.47)$$

Note that the first column of \mathbf{AB} in (2.46) is

$$\begin{pmatrix} \mathbf{a}_1' \mathbf{b}_1 \\ \mathbf{a}_2' \mathbf{b}_1 \\ \mathbf{a}_3' \mathbf{b}_1 \end{pmatrix} = \begin{pmatrix} \mathbf{a}_1' \\ \mathbf{a}_2' \\ \mathbf{a}_3' \end{pmatrix} \mathbf{b}_1 = \mathbf{Ab}_1$$

and likewise the second column is \mathbf{Ab}_2 . Thus \mathbf{AB} can be written in the form

$$\mathbf{AB} = \mathbf{A}(\mathbf{b}_1, \mathbf{b}_2) = (\mathbf{Ab}_1, \mathbf{Ab}_2).$$

This result holds in general:

$$\mathbf{AB} = \mathbf{A}(\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_p) = (\mathbf{Ab}_1, \mathbf{Ab}_2, \dots, \mathbf{Ab}_p). \quad (2.48)$$

To further illustrate matrix multiplication in terms of rows and columns, let $\mathbf{A} = \begin{pmatrix} \mathbf{a}_1' \\ \mathbf{a}_2' \end{pmatrix}$ be a $2 \times p$ matrix, \mathbf{x} be a $p \times 1$ vector, and \mathbf{S} be a $p \times p$ matrix. Then

$$\mathbf{Ax} = \begin{pmatrix} \mathbf{a}_1' \\ \mathbf{a}_2' \end{pmatrix} \mathbf{x} = \begin{pmatrix} \mathbf{a}_1' \mathbf{x} \\ \mathbf{a}_2' \mathbf{x} \end{pmatrix}, \quad (2.49)$$

$$\mathbf{ASA}' = \begin{pmatrix} \mathbf{a}_1' \mathbf{Sa}_1 & \mathbf{a}_1' \mathbf{Sa}_2 \\ \mathbf{a}_2' \mathbf{Sa}_1 & \mathbf{a}_2' \mathbf{Sa}_2 \end{pmatrix}. \quad (2.50)$$

Any matrix can be multiplied by its transpose. If \mathbf{A} is $n \times p$, then

\mathbf{AA}' is $n \times n$ and is obtained as products of rows of \mathbf{A} [see (2.52)].

Similarly,

$\mathbf{A}'\mathbf{A}$ is $p \times p$ and is obtained as products of columns of \mathbf{A} [see (2.54)].

From (2.6) and (2.27), it is clear that both $\mathbf{A}\mathbf{A}'$ and $\mathbf{A}'\mathbf{A}$ are symmetric.

In the above illustration for \mathbf{AB} in terms of row and column vectors, the rows of \mathbf{A} were denoted by \mathbf{a}'_i and the columns of \mathbf{B} by \mathbf{b}_j . If both rows and columns of a matrix \mathbf{A} are under discussion, as in \mathbf{AA}' and $\mathbf{A}'\mathbf{A}$, we will use the notation \mathbf{a}'_i for rows and $\mathbf{a}_{(j)}$ for columns. To illustrate, if \mathbf{A} is 3×4 , we have

$$\begin{aligned}\mathbf{A} &= \begin{pmatrix} a_{11} & a_{12} & a_{13} & a_{14} \\ a_{21} & a_{22} & a_{23} & a_{24} \\ a_{31} & a_{32} & a_{33} & a_{34} \end{pmatrix} \\ &= \begin{pmatrix} \mathbf{a}'_1 \\ \mathbf{a}'_2 \\ \mathbf{a}'_3 \end{pmatrix} = (\mathbf{a}_{(1)}, \mathbf{a}_{(2)}, \mathbf{a}_{(3)}, \mathbf{a}_{(4)}),\end{aligned}$$

where, for example,

$$\mathbf{a}'_2 = (a_{21} \ a_{22} \ a_{23} \ a_{24}),$$

$$\mathbf{a}_{(3)} = \begin{pmatrix} a_{13} \\ a_{23} \\ a_{33} \end{pmatrix}.$$

With this notation for rows and columns of \mathbf{A} , we can express the elements of $\mathbf{A}'\mathbf{A}$ or of \mathbf{AA}' as products of the rows of \mathbf{A} or of the columns of \mathbf{A} . Thus if we write \mathbf{A} in terms of its rows as

$$\mathbf{A} = \begin{pmatrix} \mathbf{a}'_1 \\ \mathbf{a}'_2 \\ \vdots \\ \mathbf{a}'_n \end{pmatrix},$$

then we have

$$\mathbf{A}'\mathbf{A} = (\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n) \begin{pmatrix} \mathbf{a}'_1 \\ \mathbf{a}'_2 \\ \vdots \\ \mathbf{a}'_n \end{pmatrix} = \sum_{i=1}^n \mathbf{a}_i \mathbf{a}'_i, \quad (2.51)$$

$$\begin{aligned}\mathbf{AA}' &= \begin{pmatrix} \mathbf{a}'_1 \\ \mathbf{a}'_2 \\ \vdots \\ \mathbf{a}'_n \end{pmatrix} (\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n) \\ &= \begin{pmatrix} \mathbf{a}'_1 \mathbf{a}_1 & \mathbf{a}'_1 \mathbf{a}_2 & \cdots & \mathbf{a}'_1 \mathbf{a}_n \\ \mathbf{a}'_2 \mathbf{a}_1 & \mathbf{a}'_2 \mathbf{a}_2 & \cdots & \mathbf{a}'_2 \mathbf{a}_n \\ \vdots & \vdots & & \vdots \\ \mathbf{a}'_n \mathbf{a}_1 & \mathbf{a}'_n \mathbf{a}_2 & \cdots & \mathbf{a}'_n \mathbf{a}_n \end{pmatrix}.\end{aligned} \quad (2.52)$$

Similarly, if we express \mathbf{A} in terms of columns as

$$\mathbf{A} = (\mathbf{a}_{(1)}, \mathbf{a}_{(2)}, \dots, \mathbf{a}_{(p)}),$$

then

$$\mathbf{AA}' = (\mathbf{a}_{(1)}, \mathbf{a}_{(2)}, \dots, \mathbf{a}_{(p)}) \begin{pmatrix} \mathbf{a}'_{(1)} \\ \mathbf{a}'_{(2)} \\ \vdots \\ \mathbf{a}'_{(p)} \end{pmatrix} = \sum_{j=1}^p \mathbf{a}_{(j)} \mathbf{a}'_{(j)}, \quad (2.53)$$

$$\begin{aligned} \mathbf{A}'\mathbf{A} &= \begin{pmatrix} \mathbf{a}'_{(1)} \\ \mathbf{a}'_{(2)} \\ \vdots \\ \mathbf{a}'_{(p)} \end{pmatrix} (\mathbf{a}_{(1)}, \mathbf{a}_{(2)}, \dots, \mathbf{a}_{(p)}) \\ &= \begin{pmatrix} \mathbf{a}'_{(1)}\mathbf{a}_{(1)} & \mathbf{a}'_{(1)}\mathbf{a}_{(2)} & \cdots & \mathbf{a}'_{(1)}\mathbf{a}_{(p)} \\ \mathbf{a}'_{(2)}\mathbf{a}_{(1)} & \mathbf{a}'_{(2)}\mathbf{a}_{(2)} & \cdots & \mathbf{a}'_{(2)}\mathbf{a}_{(p)} \\ \vdots & \vdots & & \vdots \\ \mathbf{a}'_{(p)}\mathbf{a}_{(1)} & \mathbf{a}'_{(p)}\mathbf{a}_{(2)} & \cdots & \mathbf{a}'_{(p)}\mathbf{a}_{(p)} \end{pmatrix}. \end{aligned} \quad (2.54)$$

Let $\mathbf{A} = (a_{ij})$ be an $n \times n$ matrix and \mathbf{D} be a diagonal matrix, $\mathbf{D} = \text{diag}(d_1, d_2, \dots, d_n)$. Then, in the product \mathbf{DA} , the i th row of \mathbf{A} is multiplied by d_i , and in \mathbf{AD} , the j th column of \mathbf{A} is multiplied by d_j . For example, if $n = 3$, we have

$$\begin{aligned} \mathbf{DA} &= \begin{pmatrix} d_1 & 0 & 0 \\ 0 & d_2 & 0 \\ 0 & 0 & d_3 \end{pmatrix} \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \\ &= \begin{pmatrix} d_1 a_{11} & d_1 a_{12} & d_1 a_{13} \\ d_2 a_{21} & d_2 a_{22} & d_2 a_{23} \\ d_3 a_{31} & d_3 a_{32} & d_3 a_{33} \end{pmatrix}, \end{aligned} \quad (2.55)$$

$$\begin{aligned} \mathbf{AD} &= \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix} \begin{pmatrix} d_1 & 0 & 0 \\ 0 & d_2 & 0 \\ 0 & 0 & d_3 \end{pmatrix} \\ &= \begin{pmatrix} d_1 a_{11} & d_2 a_{12} & d_3 a_{13} \\ d_1 a_{21} & d_2 a_{22} & d_3 a_{23} \\ d_1 a_{31} & d_2 a_{32} & d_3 a_{33} \end{pmatrix}, \end{aligned} \quad (2.56)$$

$$\mathbf{DAD} = \begin{pmatrix} d_1^2 a_{11} & d_1 d_2 a_{12} & d_1 d_3 a_{13} \\ d_2 d_1 a_{21} & d_2^2 a_{22} & d_2 d_3 a_{23} \\ d_3 d_1 a_{31} & d_3 d_2 a_{32} & d_3^2 a_{33} \end{pmatrix}. \quad (2.57)$$

In the special case where the diagonal matrix is the identity, we have

$$\mathbf{IA} = \mathbf{AI} = \mathbf{A}. \quad (2.58)$$

If \mathbf{A} is rectangular, (2.58) still holds, but the two identities are of different sizes.

The product of a scalar and a matrix is obtained by multiplying each element of the matrix by the scalar:

$$c\mathbf{A} = (ca_{ij}) = \begin{pmatrix} ca_{11} & ca_{12} & \cdots & ca_{1m} \\ ca_{21} & ca_{22} & \cdots & ca_{2m} \\ \vdots & \vdots & & \vdots \\ ca_{n1} & ca_{n2} & \cdots & ca_{nm} \end{pmatrix}. \quad (2.59)$$

For example,

$$c\mathbf{I} = \begin{pmatrix} c & 0 & \cdots & 0 \\ 0 & c & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & c \end{pmatrix}, \quad (2.60)$$

$$c\mathbf{x} = \begin{pmatrix} cx_1 \\ cx_2 \\ \vdots \\ cx_n \end{pmatrix}. \quad (2.61)$$

Since $ca_{ij} = a_{ij}c$, the product of a scalar and a matrix is commutative:

$$c\mathbf{A} = \mathbf{A}c. \quad (2.62)$$

Multiplication of vectors or matrices by scalars permits the use of linear combinations, such as

$$\sum_{i=1}^k a_i \mathbf{x}_i = a_1 \mathbf{x}_1 + a_2 \mathbf{x}_2 + \cdots + a_k \mathbf{x}_k,$$

$$\sum_{i=1}^k a_i \mathbf{B}_i = a_1 \mathbf{B}_1 + a_2 \mathbf{B}_2 + \cdots + a_k \mathbf{B}_k.$$

If \mathbf{A} is a symmetric matrix and \mathbf{x} and \mathbf{y} are vectors, the product

$$\mathbf{y}' \mathbf{A} \mathbf{y} = \sum_i a_{ii} y_i^2 + \sum_{i \neq j} a_{ij} y_i y_j \quad (2.63)$$

is called a *quadratic form*, while

$$\mathbf{x}' \mathbf{A} \mathbf{y} = \sum_{ij} a_{ij} x_i y_j \quad (2.64)$$

is called a *bilinear form*. Either of these is, of course, a scalar and can be treated as such. Expressions such as $\mathbf{x}' \mathbf{A} \mathbf{y} / \sqrt{\mathbf{x}' \mathbf{A} \mathbf{x}}$ are permissible (assuming \mathbf{A} is positive definite; see Section 2.7).

2.4 PARTITIONED MATRICES

It is sometimes convenient to partition a matrix into submatrices. For example, a partitioning of a matrix \mathbf{A} into four submatrices could be indicated symbolically as follows:

$$\mathbf{A} = \begin{pmatrix} \mathbf{A}_{11} & \mathbf{A}_{12} \\ \mathbf{A}_{21} & \mathbf{A}_{22} \end{pmatrix}.$$

For example, a 4×5 matrix \mathbf{A} could be partitioned as

$$\mathbf{A} = \left(\begin{array}{ccc|cc} 2 & 1 & 3 & 8 & 4 \\ -3 & 4 & 0 & 2 & 7 \\ 9 & 3 & 6 & 5 & -2 \\ \hline 4 & 8 & 3 & 1 & 6 \end{array} \right) = \begin{pmatrix} \mathbf{A}_{11} & \mathbf{A}_{12} \\ \mathbf{A}_{21} & \mathbf{A}_{22} \end{pmatrix},$$

where

$$\mathbf{A}_{11} = \begin{pmatrix} 2 & 1 & 3 \\ -3 & 4 & 0 \\ 9 & 3 & 6 \end{pmatrix}, \quad \mathbf{A}_{12} = \begin{pmatrix} 8 & 4 \\ 2 & 7 \\ 5 & -2 \end{pmatrix},$$

$$\mathbf{A}_{21} = (4 \ 8 \ 3), \quad \mathbf{A}_{22} = (1 \ 6).$$

If two matrices \mathbf{A} and \mathbf{B} are conformable and \mathbf{A} and \mathbf{B} are partitioned so that the submatrices are appropriately conformable, then the product \mathbf{AB} can be found by following the usual row-by-column pattern of multiplication on the submatrices as if they were single elements; for example,

$$\begin{aligned} \mathbf{AB} &= \begin{pmatrix} \mathbf{A}_{11} & \mathbf{A}_{12} \\ \mathbf{A}_{21} & \mathbf{A}_{22} \end{pmatrix} \begin{pmatrix} \mathbf{B}_{11} & \mathbf{B}_{12} \\ \mathbf{B}_{21} & \mathbf{B}_{22} \end{pmatrix} \\ &= \begin{pmatrix} \mathbf{A}_{11}\mathbf{B}_{11} + \mathbf{A}_{12}\mathbf{B}_{21} & \mathbf{A}_{11}\mathbf{B}_{12} + \mathbf{A}_{12}\mathbf{B}_{22} \\ \mathbf{A}_{21}\mathbf{B}_{11} + \mathbf{A}_{22}\mathbf{B}_{21} & \mathbf{A}_{21}\mathbf{B}_{12} + \mathbf{A}_{22}\mathbf{B}_{22} \end{pmatrix}. \end{aligned} \quad (2.65)$$

It can be seen that this formulation is equivalent to the usual row-by-column definition of matrix multiplication. For example, the $(1, 1)$ element of \mathbf{AB} is the product of the first row of \mathbf{A} and the first column of \mathbf{B} . In the $(1, 1)$ element of $\mathbf{A}_{11}\mathbf{B}_{11}$ we have the sum of products of part of the first row of \mathbf{A} and part of the first column of \mathbf{B} . In the $(1, 1)$ element of $\mathbf{A}_{12}\mathbf{B}_{21}$ we have the sum of products of the rest of the first row of \mathbf{A} and the remainder of the first column of \mathbf{B} .

Multiplication of a matrix and a vector can also be carried out in partitioned form. For example,

$$\mathbf{Ab} = (\mathbf{A}_1, \mathbf{A}_2) \begin{pmatrix} \mathbf{b}_1 \\ \mathbf{b}_2 \end{pmatrix} = \mathbf{A}_1\mathbf{b}_1 + \mathbf{A}_2\mathbf{b}_2, \quad (2.66)$$

where the partitioning of the columns of \mathbf{A} corresponds to the partitioning of the elements of \mathbf{b} . Note that the partitioning of \mathbf{A} into two sets of columns is indicated by a comma, $\mathbf{A} = (\mathbf{A}_1, \mathbf{A}_2)$.

The partitioned multiplication in (2.66) can be extended to individual columns of \mathbf{A} and individual elements of \mathbf{b} :

$$\begin{aligned}\mathbf{Ab} &= (\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_p) \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_p \end{pmatrix} \\ &= b_1 \mathbf{a}_1 + b_2 \mathbf{a}_2 + \cdots + b_p \mathbf{a}_p.\end{aligned}\quad (2.67)$$

Thus \mathbf{Ab} is expressible as a linear combination of the columns of \mathbf{A} , the coefficients being elements of \mathbf{b} . For example, let

$$\mathbf{A} = \begin{pmatrix} 3 & -2 & 1 \\ 2 & 1 & 0 \\ 4 & 3 & 2 \end{pmatrix} \quad \text{and} \quad \mathbf{b} = \begin{pmatrix} 4 \\ 2 \\ 3 \end{pmatrix}.$$

Then

$$\mathbf{Ab} = \begin{pmatrix} 11 \\ 10 \\ 28 \end{pmatrix}.$$

Using a linear combination of columns of \mathbf{A} as in (2.67), we obtain

$$\begin{aligned}\mathbf{Ab} &= b_1 \mathbf{a}_1 + b_2 \mathbf{a}_2 + b_3 \mathbf{a}_3 \\ &= 4 \begin{pmatrix} 3 \\ 2 \\ 4 \end{pmatrix} + 2 \begin{pmatrix} -2 \\ 1 \\ 3 \end{pmatrix} + 3 \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix} \\ &= \begin{pmatrix} 12 \\ 8 \\ 16 \end{pmatrix} + \begin{pmatrix} -4 \\ 2 \\ 6 \end{pmatrix} + \begin{pmatrix} 3 \\ 0 \\ 6 \end{pmatrix} = \begin{pmatrix} 11 \\ 10 \\ 28 \end{pmatrix}.\end{aligned}$$

We note that if \mathbf{A} is partitioned as in (2.66), $\mathbf{A} = (\mathbf{A}_1, \mathbf{A}_2)$, the transpose is not equal to $(\mathbf{A}'_1, \mathbf{A}'_2)$, but rather

$$\mathbf{A}' = (\mathbf{A}_1, \mathbf{A}_2)' = \begin{pmatrix} \mathbf{A}'_1 \\ \mathbf{A}'_2 \end{pmatrix}. \quad (2.68)$$

2.5 RANK

Before defining the rank of a matrix, we first introduce the notion of linear independence and dependence. A set of vectors $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n$ is said to be *linearly dependent* if constants c_1, c_2, \dots, c_n (not all zero) can be found such that

$$c_1 \mathbf{a}_1 + c_2 \mathbf{a}_2 + \cdots + c_n \mathbf{a}_n = \mathbf{0}. \quad (2.69)$$

If no constants c_1, c_2, \dots, c_n can be found satisfying (2.69), the set of vectors is said to be *linearly independent*.

If (2.69) holds, then at least one of the vectors \mathbf{a}_i can be expressed as a linear combination of the other vectors in the set. Thus linear dependence of a set of vectors implies redundancy in the set. Among linearly independent vectors there is no redundancy of this type.

The *rank* of any square or rectangular matrix \mathbf{A} is defined as

$$\begin{aligned}\text{rank}(\mathbf{A}) &= \text{number of linearly independent rows of } \mathbf{A} \\ &= \text{number of linearly independent columns of } \mathbf{A}.\end{aligned}$$

It can be shown that the number of linearly independent rows of a matrix is always equal to the number of linearly independent columns.

If \mathbf{A} is $n \times p$, the maximum possible rank of \mathbf{A} is the smaller of n and p , in which case \mathbf{A} is said to be of *full rank* (sometimes said *full row rank* or *full column rank*). For example,

$$\mathbf{A} = \begin{pmatrix} 1 & -2 & 3 \\ 5 & 2 & 4 \end{pmatrix}$$

has rank 2 because the two rows are linearly independent (neither row is a multiple of the other). However, even though \mathbf{A} is full rank, the columns are linearly dependent because rank 2 implies there are only two linearly independent columns. Thus, by (2.69), there exist constants c_1, c_2 , and c_3 such that

$$c_1 \begin{pmatrix} 1 \\ 5 \end{pmatrix} + c_2 \begin{pmatrix} -2 \\ 2 \end{pmatrix} + c_3 \begin{pmatrix} 3 \\ 4 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}. \quad (2.70)$$

By (2.67), we can write (2.70) in the form

$$\begin{pmatrix} 1 & -2 & 3 \\ 5 & 2 & 4 \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \\ c_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

or

$$\mathbf{Ac} = \mathbf{0}. \quad (2.71)$$

A solution vector to (2.70) or (2.71) is given by any multiple of $\mathbf{c} = (14, -11, -12)'$. Hence we have the interesting result that a product of a matrix \mathbf{A} and a vector \mathbf{c} is equal to $\mathbf{0}$, even though $\mathbf{A} \neq \mathbf{0}$ and $\mathbf{c} \neq \mathbf{0}$. This is a direct consequence of the linear dependence of the column vectors of \mathbf{A} .

Another consequence of the linear dependence of rows or columns of a matrix is the possibility of expressions such as $\mathbf{AB} = \mathbf{CB}$, where $\mathbf{A} \neq \mathbf{C}$. For example, let

$$\mathbf{A} = \begin{pmatrix} 1 & 3 & 2 \\ 2 & 0 & -1 \end{pmatrix}, \quad \mathbf{B} = \begin{pmatrix} 1 & 2 \\ 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \mathbf{C} = \begin{pmatrix} 2 & 1 & 1 \\ 5 & -6 & -4 \end{pmatrix}.$$

Then

$$\mathbf{AB} = \mathbf{CB} = \begin{pmatrix} 3 & 5 \\ 1 & 4 \end{pmatrix}.$$

All three of the matrices \mathbf{A} , \mathbf{B} , and \mathbf{C} are full rank; but being rectangular, they have a rank deficiency in either rows or columns, which permits us to construct $\mathbf{AB} = \mathbf{CB}$ with $\mathbf{A} \neq \mathbf{C}$. Thus in a matrix equation, we cannot, in general, cancel matrices from both sides of the equation.

There are two exceptions to this rule. One involves a nonsingular matrix to be defined in Section 2.6. The other special case occurs when the expression holds for all possible values of the matrix common to both sides of the equation. For example,

$$\text{If } \mathbf{Ax} = \mathbf{Bx} \text{ for all possible values of } \mathbf{x}, \text{ then } \mathbf{A} = \mathbf{B}. \quad (2.72)$$

To see this, let $\mathbf{x} = (1, 0, \dots, 0)'$. Then the first column of \mathbf{A} equals the first column of \mathbf{B} . Now let $\mathbf{x} = (0, 1, 0, \dots, 0)'$, and the second column of \mathbf{A} equals the second column of \mathbf{B} . Continuing in this fashion, we obtain $\mathbf{A} = \mathbf{B}$.

Suppose a rectangular matrix \mathbf{A} is $n \times p$ of rank p , where $p < n$. We typically shorten this statement to “ \mathbf{A} is $n \times p$ of rank $p < n$.”

2.6 INVERSE

If a matrix \mathbf{A} is square and of full rank, then \mathbf{A} is said to be *nonsingular*, and \mathbf{A} has a unique *inverse*, denoted by \mathbf{A}^{-1} , with the property that

$$\mathbf{AA}^{-1} = \mathbf{A}^{-1}\mathbf{A} = \mathbf{I}. \quad (2.73)$$

For example, let

$$\mathbf{A} = \begin{pmatrix} 3 & 4 \\ 2 & 6 \end{pmatrix}.$$

Then

$$\mathbf{A}^{-1} = \begin{pmatrix} .6 & -.4 \\ -.2 & .3 \end{pmatrix},$$

$$\mathbf{AA}^{-1} = \begin{pmatrix} 3 & 4 \\ 2 & 6 \end{pmatrix} \begin{pmatrix} .6 & -.4 \\ -.2 & .3 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

If \mathbf{A} is square and of less than full rank, then an inverse does not exist, and \mathbf{A} is said to be *singular*. Note that rectangular matrices do not have inverses as in (2.73), even if they are full rank.

If \mathbf{A} and \mathbf{B} are the same size and nonsingular, then the inverse of their product is the product of their inverses in reverse order,

$$(\mathbf{AB})^{-1} = \mathbf{B}^{-1}\mathbf{A}^{-1}. \quad (2.74)$$

Note that (2.74) holds only for nonsingular matrices. Thus, for example, if \mathbf{A} is $n \times p$ of rank $p < n$, then $\mathbf{A}'\mathbf{A}$ has an inverse, but $(\mathbf{A}'\mathbf{A})^{-1}$ is not equal to $\mathbf{A}^{-1}(\mathbf{A}')^{-1}$ because \mathbf{A} is rectangular and does not have an inverse.

If a matrix is nonsingular, it can be canceled from both sides of an equation, provided it appears on the left (right) on both sides. For example, if \mathbf{B} is nonsingular, then

$$\mathbf{AB} = \mathbf{CB} \text{ implies } \mathbf{A} = \mathbf{C},$$

since we can multiply on the right by \mathbf{B}^{-1} to obtain

$$\mathbf{A}\mathbf{B}\mathbf{B}^{-1} = \mathbf{C}\mathbf{B}\mathbf{B}^{-1},$$

$$\mathbf{AI} = \mathbf{CI},$$

$$\mathbf{A} = \mathbf{C}.$$

Otherwise, if \mathbf{A} , \mathbf{B} , and \mathbf{C} are rectangular or square and singular, it is easy to construct $\mathbf{AB} = \mathbf{CB}$, with $\mathbf{A} \neq \mathbf{C}$, as illustrated near the end of Section 2.5.

The inverse of the transpose of a nonsingular matrix is given by the transpose of the inverse:

$$(\mathbf{A}')^{-1} = (\mathbf{A}^{-1})'. \quad (2.75)$$

If the symmetric nonsingular matrix \mathbf{A} is partitioned in the form

$$\mathbf{A} = \begin{pmatrix} \mathbf{A}_{11} & \mathbf{a}_{12} \\ \mathbf{a}'_{12} & a_{22} \end{pmatrix},$$

then the inverse is given by

$$\mathbf{A}^{-1} = \frac{1}{b} \begin{pmatrix} b\mathbf{A}_{11}^{-1} + \mathbf{A}_{11}^{-1}\mathbf{a}_{12}\mathbf{a}'_{12}\mathbf{A}_{11}^{-1} & -\mathbf{A}_{11}^{-1}\mathbf{a}_{12} \\ -\mathbf{a}'_{12}\mathbf{A}_{11}^{-1} & 1 \end{pmatrix}, \quad (2.76)$$

where $b = a_{22} - \mathbf{a}'_{12}\mathbf{A}_{11}^{-1}\mathbf{a}_{12}$. A nonsingular matrix of the form $\mathbf{B} + \mathbf{cc}'$, where \mathbf{B} is nonsingular, has as its inverse

$$(\mathbf{B} + \mathbf{cc}')^{-1} = \mathbf{B}^{-1} - \frac{\mathbf{B}^{-1}\mathbf{cc}'\mathbf{B}^{-1}}{1 + \mathbf{c}'\mathbf{B}^{-1}\mathbf{c}}. \quad (2.77)$$

2.7 POSITIVE DEFINITE MATRICES

The symmetric matrix \mathbf{A} is said to be *positive definite* if $\mathbf{x}'\mathbf{Ax} > 0$ for all possible vectors \mathbf{x} (except $\mathbf{x} = \mathbf{0}$). Similarly, \mathbf{A} is *positive semidefinite* if $\mathbf{x}'\mathbf{Ax} \geq 0$ for all $\mathbf{x} \neq \mathbf{0}$. [A quadratic form $\mathbf{x}'\mathbf{Ax}$ was defined in (2.63).]

The diagonal elements a_{ii} of a positive definite matrix are positive. To see this, let $\mathbf{x}' = (0, \dots, 0, 1, 0, \dots, 0)$ with a 1 in the i th position. Then $\mathbf{x}'\mathbf{Ax} = a_{ii} > 0$. Similarly, for a positive semidefinite matrix \mathbf{A} , $a_{ii} \geq 0$ for all i .

One way to obtain a positive definite matrix is as follows:

If $\mathbf{A} = \mathbf{B}'\mathbf{B}$, where \mathbf{B} is $n \times p$ of rank $p < n$, then $\mathbf{B}'\mathbf{B}$ is positive definite.

$$(2.78)$$

This is easily shown:

$$\mathbf{x}'\mathbf{Ax} = \mathbf{x}'\mathbf{B}'\mathbf{Bx} = (\mathbf{Bx})'(\mathbf{Bx}) = \mathbf{z}'\mathbf{z},$$

where $\mathbf{z} = \mathbf{B}\mathbf{x}$. Thus, $\mathbf{x}'\mathbf{A}\mathbf{x} = \sum_{i=1}^n z_i^2$, which is positive ($\mathbf{B}\mathbf{x}$ cannot be $\mathbf{0}$ unless $\mathbf{x} = \mathbf{0}$, because \mathbf{B} is full rank). If \mathbf{B} is less than full rank, then by a similar argument, $\mathbf{B}'\mathbf{B}$ is positive semidefinite.

Note that $\mathbf{A} = \mathbf{B}'\mathbf{B}$ is analogous to $a = b^2$ in real numbers, where the square of any number (including negative numbers) is positive.

In another analogy to positive real numbers, a positive definite matrix can be factored into a “square root” in two ways. We give one method below in (2.79) and the other in Section 2.11.8.

A positive definite matrix \mathbf{A} can be factored into

$$\mathbf{A} = \mathbf{T}'\mathbf{T}, \quad (2.79)$$

where \mathbf{T} is a nonsingular upper triangular matrix. One way to obtain \mathbf{T} is the *Cholesky decomposition*, which can be carried out in the following steps.

Let $\mathbf{A} = (a_{ij})$ and $\mathbf{T} = (t_{ij})$ be $n \times n$. Then the elements of \mathbf{T} are found as follows:

$$\begin{aligned} t_{11} &= \sqrt{a_{11}}, \quad t_{1j} = \frac{a_{1j}}{t_{11}} \quad && 2 \leq j \leq n \\ t_{ii} &= \sqrt{a_{ii} - \sum_{k=1}^{i-1} t_{ki}^2} \quad && 2 \leq i \leq n \\ t_{ij} &= \frac{a_{ij} - \sum_{k=1}^{i-1} t_{ki} t_{kj}}{t_{ii}} \quad && 2 \leq i < j \leq n \\ t_{ij} &= 0 \quad && 1 \leq j < i \leq n \end{aligned}$$

For example, let

$$\mathbf{A} = \begin{pmatrix} 3 & 0 & -3 \\ 0 & 6 & 3 \\ -3 & 3 & 6 \end{pmatrix}.$$

Then by the Cholesky method, we obtain

$$\mathbf{T} = \begin{pmatrix} \sqrt{3} & 0 & -\sqrt{3} \\ 0 & \sqrt{6} & \sqrt{1.5} \\ 0 & 0 & \sqrt{1.5} \end{pmatrix},$$

$$\begin{aligned} \mathbf{T}'\mathbf{T} &= \begin{pmatrix} \sqrt{3} & 0 & 0 \\ 0 & \sqrt{6} & 0 \\ -\sqrt{3} & \sqrt{1.5} & \sqrt{1.5} \end{pmatrix} \begin{pmatrix} \sqrt{3} & 0 & -\sqrt{3} \\ 0 & \sqrt{6} & \sqrt{1.5} \\ 0 & 0 & \sqrt{1.5} \end{pmatrix} \\ &= \begin{pmatrix} 3 & 0 & -3 \\ 0 & 6 & 3 \\ -3 & 3 & 6 \end{pmatrix} = \mathbf{A}. \end{aligned}$$

2.8 DETERMINANTS

The *determinant* of an $n \times n$ matrix \mathbf{A} is defined as the sum of all $n!$ possible products of n elements such that

1. Each product contains one element from every row and every column, and
2. The factors in each product are written so that the column subscripts appear in order of magnitude and each product is then preceded by a plus or minus sign according to whether the number of inversions in the row subscripts is even or odd.

An *inversion* occurs whenever a larger number precedes a smaller one. The symbol $n!$ is defined as

$$n! = n(n - 1)(n - 2) \cdots 2 \cdot 1. \quad (2.80)$$


The determinant of \mathbf{A} is a scalar denoted by $|\mathbf{A}|$ or by $\det(\mathbf{A})$. The preceding definition is not useful in evaluating determinants, except in the case of 2×2 or 3×3 matrices. For larger matrices, other methods are available for manual computation, but determinants are typically evaluated by computer. For a 2×2 matrix, the determinant is found by

$$|\mathbf{A}| = \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} = a_{11}a_{22} - a_{21}a_{12}. \quad (2.81)$$


For a 3×3 matrix, the determinant is given by

$$\begin{aligned} |\mathbf{A}| = & a_{11}a_{22}a_{33} + a_{12}a_{23}a_{31} + a_{13}a_{21}a_{32} \\ & - a_{31}a_{22}a_{13} - a_{32}a_{23}a_{11} - a_{33}a_{12}a_{21}. \end{aligned} \quad (2.82)$$

This can be found by the following scheme. The three positive terms are obtained by



and the three negative terms by



The determinant of a diagonal matrix is the product of the diagonal elements; that is, if $\mathbf{D} = \text{diag}(d_1, d_2, \dots, d_n)$, then

$$|\mathbf{D}| = \prod_{i=1}^n d_i. \quad (2.83)$$

As a special case of (2.83), suppose all diagonal elements are equal, say,

$$\mathbf{D} = \text{diag}(c, c, \dots, c) = c\mathbf{I}.$$

Then

$$|\mathbf{D}| = |c\mathbf{I}| = \prod_{i=1}^n c = c^n. \quad (2.84)$$

The extension of (2.84) to any square matrix \mathbf{A} is

$$|c\mathbf{A}| = c^n |\mathbf{A}|. \quad (2.85)$$

Because the determinant is a scalar, we can carry out operations such as

$$|\mathbf{A}|^2, \quad |\mathbf{A}|^{1/2}, \quad \frac{1}{|\mathbf{A}|},$$

provided that $|\mathbf{A}| > 0$ for $|\mathbf{A}|^{1/2}$ and that $|\mathbf{A}| \neq 0$ for $1/|\mathbf{A}|$.

If the square matrix \mathbf{A} is singular, its determinant is 0:

$$|\mathbf{A}| = 0 \text{ if } \mathbf{A} \text{ is singular.} \quad (2.86)$$

If \mathbf{A} is *near singular*, then there exists a linear combination of the columns that is close to $\mathbf{0}$, and $|\mathbf{A}|$ is also close to 0. If \mathbf{A} is nonsingular, its determinant is nonzero:

$$|\mathbf{A}| \neq 0 \text{ if } \mathbf{A} \text{ is nonsingular.} \quad (2.87)$$

If \mathbf{A} is positive definite, its determinant is positive:

$$|\mathbf{A}| > 0 \text{ if } \mathbf{A} \text{ is positive definite.} \quad (2.88)$$

If \mathbf{A} and \mathbf{B} are square and the same size, the determinant of the product is the product of the determinants:

$$|\mathbf{AB}| = |\mathbf{A}||\mathbf{B}|. \quad (2.89)$$

For example, let

$$\mathbf{A} = \begin{pmatrix} 1 & 2 \\ -3 & 5 \end{pmatrix} \quad \text{and} \quad \mathbf{B} = \begin{pmatrix} 4 & 2 \\ 1 & 3 \end{pmatrix}.$$

Then

$$\mathbf{AB} = \begin{pmatrix} 6 & 8 \\ -7 & 9 \end{pmatrix}, \quad |\mathbf{AB}| = 110,$$

$$|\mathbf{A}| = 11, \quad |\mathbf{B}| = 10, \quad |\mathbf{A}||\mathbf{B}| = 110.$$

The determinant of the transpose of a matrix is the same as the determinant of the matrix, and the determinant of the inverse of a matrix is the reciprocal of the determinant:

$$|\mathbf{A}'| = |\mathbf{A}|, \quad (2.90)$$

$$|\mathbf{A}^{-1}| = \frac{1}{|\mathbf{A}|} = |\mathbf{A}|^{-1}. \quad (2.91)$$

If a partitioned matrix has the form

$$\mathbf{A} = \begin{pmatrix} \mathbf{A}_{11} & \mathbf{O} \\ \mathbf{O} & \mathbf{A}_{22} \end{pmatrix},$$

where \mathbf{A}_{11} and \mathbf{A}_{22} are square, but not necessarily the same size, then

$$|\mathbf{A}| = \begin{vmatrix} \mathbf{A}_{11} & \mathbf{O} \\ \mathbf{O} & \mathbf{A}_{22} \end{vmatrix} = |\mathbf{A}_{11}| |\mathbf{A}_{22}|. \quad (2.92)$$

For a general partitioned matrix,

$$\mathbf{A} = \begin{pmatrix} \mathbf{A}_{11} & \mathbf{A}_{12} \\ \mathbf{A}_{21} & \mathbf{A}_{22} \end{pmatrix},$$

where \mathbf{A}_{11} and \mathbf{A}_{22} are square and nonsingular (not necessarily the same size), the determinant is given by either of the following two expressions:

$$\begin{vmatrix} \mathbf{A}_{11} & \mathbf{A}_{12} \\ \mathbf{A}_{21} & \mathbf{A}_{22} \end{vmatrix} = |\mathbf{A}_{11}| |\mathbf{A}_{22} - \mathbf{A}_{21}\mathbf{A}_{11}^{-1}\mathbf{A}_{12}| \quad (2.93)$$

$$= |\mathbf{A}_{22}| |\mathbf{A}_{11} - \mathbf{A}_{12}\mathbf{A}_{22}^{-1}\mathbf{A}_{21}|. \quad (2.94)$$

Note the analogy of (2.93) and (2.94) to the case of the determinant of a 2×2 matrix as given by (2.81):

$$\begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} = a_{11}a_{22} - a_{21}a_{12} \\ = a_{11} \left(a_{22} - \frac{a_{21}a_{12}}{a_{11}} \right) \\ = a_{22} \left(a_{11} - \frac{a_{12}a_{21}}{a_{22}} \right).$$

If \mathbf{B} is nonsingular and \mathbf{c} is a vector, then

$$|\mathbf{B} + \mathbf{cc}'| = |\mathbf{B}|(1 + \mathbf{c}'\mathbf{B}^{-1}\mathbf{c}). \quad (2.95)$$

2.9 TRACE

A simple function of an $n \times n$ matrix \mathbf{A} is the *trace*, denoted by $\text{tr}(\mathbf{A})$ and defined as the sum of the diagonal elements of \mathbf{A} ; that is, $\text{tr}(\mathbf{A}) = \sum_{i=1}^n a_{ii}$. The trace is, of course, a scalar. For example, suppose

$$\mathbf{A} = \begin{pmatrix} 5 & 4 & 4 \\ 2 & -3 & 1 \\ 3 & 7 & 9 \end{pmatrix}.$$

Then

$$\text{tr}(\mathbf{A}) = 5 + (-3) + 9 = 11.$$

The trace of the sum of two square matrices is the sum of the traces of the two matrices:

$$\text{tr}(\mathbf{A} + \mathbf{B}) = \text{tr}(\mathbf{A}) + \text{tr}(\mathbf{B}). \quad (2.96)$$

An important result for the product of two matrices is

$$\text{tr}(\mathbf{AB}) = \text{tr}(\mathbf{BA}). \quad (2.97)$$

This result holds for any matrices \mathbf{A} and \mathbf{B} where \mathbf{AB} and \mathbf{BA} are both defined. It is not necessary that \mathbf{A} and \mathbf{B} be square or that \mathbf{AB} equal \mathbf{BA} . For example, let

$$\mathbf{A} = \begin{pmatrix} 1 & 3 \\ 2 & -1 \\ 4 & 6 \end{pmatrix}, \quad \mathbf{B} = \begin{pmatrix} 3 & -2 & 1 \\ 2 & 4 & 5 \end{pmatrix}.$$

Then

$$\mathbf{AB} = \begin{pmatrix} 9 & 10 & 16 \\ 4 & -8 & -3 \\ 24 & 16 & 34 \end{pmatrix}, \quad \mathbf{BA} = \begin{pmatrix} 3 & 17 \\ 30 & 32 \end{pmatrix},$$

$$\text{tr}(\mathbf{AB}) = 9 - 8 + 34 = 35, \quad \text{tr}(\mathbf{BA}) = 3 + 32 = 35.$$

From (2.52) and (2.54), we obtain

$$\text{tr}(\mathbf{A}'\mathbf{A}) = \text{tr}(\mathbf{AA}') = \sum_{i=1}^n \sum_{j=1}^p a_{ij}^2, \quad (2.98)$$

where the a_{ij} 's are elements of the $n \times p$ matrix \mathbf{A} .

2.10 ORTHOGONAL VECTORS AND MATRICES

Two vectors \mathbf{a} and \mathbf{b} of the same size are said to be *orthogonal* if

$$\mathbf{a}'\mathbf{b} = a_1 b_1 + a_2 b_2 + \cdots + a_n b_n = 0. \quad (2.99)$$

Geometrically, orthogonal vectors are perpendicular [see (3.14) and the comments following (3.14)]. If $\mathbf{a}'\mathbf{a} = 1$, the vector \mathbf{a} is said to be *normalized*. The vector \mathbf{a} can always be normalized by dividing by its length, $\sqrt{\mathbf{a}'\mathbf{a}}$. Thus

$$\mathbf{c} = \frac{\mathbf{a}}{\sqrt{\mathbf{a}'\mathbf{a}}} \quad (2.100)$$

is normalized so that $\mathbf{c}'\mathbf{c} = 1$.

A matrix $\mathbf{C} = (\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_p)$ whose columns are normalized and mutually orthogonal is called an *orthogonal* matrix. Since the elements of $\mathbf{C}'\mathbf{C}$ are products of columns of \mathbf{C} [see (2.54)], which have the properties $\mathbf{c}_i'\mathbf{c}_i = 1$ for all i and $\mathbf{c}_i'\mathbf{c}_j = 0$ for all $i \neq j$, we have

$$\mathbf{C}'\mathbf{C} = \mathbf{I}. \quad (2.101)$$

If \mathbf{C} satisfies (2.101), it necessarily follows that

$$\mathbf{C}\mathbf{C}' = \mathbf{I}, \quad (2.102)$$

from which we see that the rows of \mathbf{C} are also normalized and mutually orthogonal. It is clear from (2.101) and (2.102) that $\mathbf{C}^{-1} = \mathbf{C}'$ for an orthogonal matrix \mathbf{C} .

We illustrate the creation of an orthogonal matrix by starting with

$$\mathbf{A} = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & -1 \\ 1 & -2 & 0 \end{pmatrix},$$

whose columns are mutually orthogonal. To normalize the three columns, we divide by the respective lengths, $\sqrt{3}$, $\sqrt{6}$, and $\sqrt{2}$, to obtain

$$\mathbf{C} = \begin{pmatrix} 1/\sqrt{3} & 1/\sqrt{6} & 1/\sqrt{2} \\ 1/\sqrt{3} & 1/\sqrt{6} & -1/\sqrt{2} \\ 1/\sqrt{3} & -2/\sqrt{6} & 0 \end{pmatrix}.$$

Note that the rows also became normalized and mutually orthogonal so that \mathbf{C} satisfies both (2.101) and (2.102).

Multiplication by an orthogonal matrix has the effect of rotating axes; that is, if a point \mathbf{x} is transformed to $\mathbf{z} = \mathbf{Cx}$, where \mathbf{C} is orthogonal, then

$$\mathbf{z}'\mathbf{z} = (\mathbf{Cx})'(\mathbf{Cx}) = \mathbf{x}'\mathbf{C}'\mathbf{C}\mathbf{x} = \mathbf{x}'\mathbf{I}\mathbf{x} = \mathbf{x}'\mathbf{x}, \quad (2.103)$$

and the distance from the origin to \mathbf{z} is the same as the distance to \mathbf{x} .

2.11 EIGENVALUES AND EIGENVECTORS

2.11.1 Definition

For every square matrix \mathbf{A} , a scalar λ and a nonzero vector \mathbf{x} can be found such that

$$\mathbf{Ax} = \lambda\mathbf{x}. \quad (2.104)$$

In (2.104), λ is called an *eigenvalue* of \mathbf{A} and \mathbf{x} is an *eigenvector*. To find λ and \mathbf{x} , we write (2.104) as

$$(\mathbf{A} - \lambda\mathbf{I})\mathbf{x} = \mathbf{0}. \quad (2.105)$$

If $|\mathbf{A} - \lambda\mathbf{I}| \neq 0$, then $(\mathbf{A} - \lambda\mathbf{I})$ has an inverse and $\mathbf{x} = \mathbf{0}$ is the only solution. Hence, in order to obtain nontrivial solutions, we set $|\mathbf{A} - \lambda\mathbf{I}| = 0$ to find values of λ that can be substituted into (2.105) to find corresponding values of \mathbf{x} . Alternatively, (2.69) and (2.71) require that the columns of $\mathbf{A} - \lambda\mathbf{I}$ be linearly dependent. Thus in $(\mathbf{A} - \lambda\mathbf{I})\mathbf{x} = \mathbf{0}$, the matrix $\mathbf{A} - \lambda\mathbf{I}$ must be singular in order to find a solution vector \mathbf{x} that is not $\mathbf{0}$.

The equation $|\mathbf{A} - \lambda\mathbf{I}| = 0$ is called the *characteristic equation*. If \mathbf{A} is $n \times n$, the characteristic equation will have n roots; that is, \mathbf{A} will have n eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$. The λ 's will not necessarily all be distinct or all nonzero. However, if \mathbf{A} arises from computations on real (continuous) data and is nonsingular, the λ 's will all be distinct (with probability 1). After finding $\lambda_1, \lambda_2, \dots, \lambda_n$, the accompanying eigenvectors $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$ can be found using (2.105).

If we multiply both sides of (2.105) by a scalar k and note by (2.62) that k and $\mathbf{A} - \lambda\mathbf{I}$ commute, we obtain

$$(\mathbf{A} - \lambda\mathbf{I})k\mathbf{x} = k\mathbf{0} = \mathbf{0}. \quad (2.106)$$

Thus if \mathbf{x} is an eigenvector of \mathbf{A} , $k\mathbf{x}$ is also an eigenvector, and eigenvectors are unique only up to multiplication by a scalar. Hence we can adjust the length of \mathbf{x} , but the direction from the origin is unique; that is, the relative values of (ratios of) the components of $\mathbf{x} = (x_1, x_2, \dots, x_n)'$ are unique. Typically, the eigenvector \mathbf{x} is scaled so that $\mathbf{x}'\mathbf{x} = 1$.

To illustrate, we will find the eigenvalues and eigenvectors for the matrix

$$\mathbf{A} = \begin{pmatrix} 1 & 2 \\ -1 & 4 \end{pmatrix}.$$

The characteristic equation is

$$|\mathbf{A} - \lambda\mathbf{I}| = \begin{vmatrix} 1 - \lambda & 2 \\ -1 & 4 - \lambda \end{vmatrix} = (1 - \lambda)(4 - \lambda) + 2 = 0,$$

$$\lambda^2 - 5\lambda + 6 = (\lambda - 3)(\lambda - 2) = 0,$$

from which $\lambda_1 = 3$ and $\lambda_2 = 2$. To find the eigenvector corresponding to $\lambda_1 = 3$, we use (2.105),

$$(\mathbf{A} - \lambda\mathbf{I})\mathbf{x} = \mathbf{0},$$

$$\begin{pmatrix} 1 - 3 & 2 \\ -1 & 4 - 3 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix},$$

$$-2x_1 + 2x_2 = 0$$

$$-x_1 + x_2 = 0.$$

As expected, either equation is redundant in the presence of the other, and there remains a single equation with two unknowns, $x_1 = x_2$. The solution vector can be written with an arbitrary constant,

$$\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = x_1 \begin{pmatrix} 1 \\ 1 \end{pmatrix} = c \begin{pmatrix} 1 \\ 1 \end{pmatrix}.$$

If c is set equal to $1/\sqrt{2}$ to normalize the eigenvector, we obtain

$$\mathbf{x}_1 = \begin{pmatrix} 1/\sqrt{2} \\ 1/\sqrt{2} \end{pmatrix}.$$

Similarly, corresponding to $\lambda_2 = 2$, we have

$$\mathbf{x}_2 = \begin{pmatrix} 2/\sqrt{5} \\ 1/\sqrt{5} \end{pmatrix}.$$

2.11.2 $\mathbf{I} + \mathbf{A}$ and $\mathbf{I} - \mathbf{A}$

If λ is an eigenvalue of \mathbf{A} and \mathbf{x} is the corresponding eigenvector, then $1 + \lambda$ is an eigenvalue of $\mathbf{I} + \mathbf{A}$ and $1 - \lambda$ is an eigenvalue of $\mathbf{I} - \mathbf{A}$. In either case, \mathbf{x} is the corresponding eigenvector.

We demonstrate this for $\mathbf{I} + \mathbf{A}$:

$$\begin{aligned} \mathbf{Ax} &= \lambda\mathbf{x}, \\ \mathbf{x} + \mathbf{Ax} &= \mathbf{x} + \lambda\mathbf{x}, \\ (\mathbf{I} + \mathbf{A})\mathbf{x} &= (1 + \lambda)\mathbf{x}. \end{aligned}$$

2.11.3 $\text{tr}(\mathbf{A})$ and $|\mathbf{A}|$

For any square matrix \mathbf{A} with eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$, we have

$$\text{tr}(\mathbf{A}) = \sum_{i=1}^n \lambda_i, \tag{2.107}$$

$$|\mathbf{A}| = \prod_{i=1}^n \lambda_i. \tag{2.108}$$

Note that by the definition in Section 2.9, $\text{tr}(\mathbf{A})$ is also equal to $\sum_{i=1}^n a_{ii}$, but $a_{ii} \neq \lambda_i$.

We illustrate (2.107) and (2.108) using the matrix

$$\mathbf{A} = \begin{pmatrix} 1 & 2 \\ -1 & 4 \end{pmatrix}$$

from the illustration in Section 2.11.1, for which $\lambda_1 = 3$ and $\lambda_2 = 2$. Using (2.107), we obtain

$$\text{tr}(\mathbf{A}) = \lambda_1 + \lambda_2 = 3 + 2 = 5,$$

and from (2.108), we have

$$|\mathbf{A}| = \lambda_1 \lambda_2 = 3(2) = 6.$$

By definition, we obtain

$$\text{tr}(\mathbf{A}) = 1 + 4 = 5 \quad \text{and} \quad |\mathbf{A}| = (1)(4) - (-1)(2) = 6.$$

2.11.4 Positive Definite and Semidefinite Matrices

The eigenvalues and eigenvectors of positive definite and positive semidefinite matrices have the following properties:

1. The eigenvalues of a positive definite matrix are all positive.
2. The eigenvalues of a positive semidefinite matrix are positive or zero, with the number of positive eigenvalues equal to the rank of the matrix.

It is customary to list the eigenvalues of a positive definite matrix in descending order: $\lambda_1 > \lambda_2 > \dots > \lambda_p$. The eigenvectors $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$ are listed in the same order; \mathbf{x}_1 corresponds to λ_1 , \mathbf{x}_2 corresponds to λ_2 , and so on.

The following result, known as the Perron–Frobenius theorem, is of interest in Chapter 12: If all elements of the positive definite matrix \mathbf{A} are positive, then all elements of the first eigenvector are positive. (The first eigenvector is the one associated with the first eigenvalue, λ_1 .)

2.11.5 The Product \mathbf{AB}

If \mathbf{A} and \mathbf{B} are square and the same size, the eigenvalues of \mathbf{AB} are the same as those of \mathbf{BA} , although the eigenvectors are usually different. This result also holds if \mathbf{AB} and \mathbf{BA} are both square but of different sizes, as when \mathbf{A} is $n \times p$ and \mathbf{B} is $p \times n$. (In this case, the nonzero eigenvalues of \mathbf{AB} and \mathbf{BA} will be the same.)

2.11.6 Symmetric Matrix

The eigenvectors of an $n \times n$ symmetric matrix \mathbf{A} are mutually orthogonal. It follows that if the n eigenvectors of \mathbf{A} are normalized and inserted as columns of a matrix $\mathbf{C} = (\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n)$, then \mathbf{C} is orthogonal.

2.11.7 Spectral Decomposition

It was noted in Section 2.11.6 that if the matrix $\mathbf{C} = (\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n)$ contains the normalized eigenvectors of an $n \times n$ symmetric matrix \mathbf{A} , then \mathbf{C} is orthogonal. Therefore, by (2.102), $\mathbf{I} = \mathbf{CC}'$, which we can multiply by \mathbf{A} to obtain

$$\mathbf{A} = \mathbf{ACC}'.$$

We now substitute $\mathbf{C} = (\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n)$:

$$\begin{aligned}\mathbf{A} &= \mathbf{A}(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n)\mathbf{C}' \\ &= (\mathbf{Ax}_1, \mathbf{Ax}_2, \dots, \mathbf{Ax}_n)\mathbf{C}' && [\text{by (2.48)}] \\ &= (\lambda_1\mathbf{x}_1, \lambda_2\mathbf{x}_2, \dots, \lambda_n\mathbf{x}_n)\mathbf{C}' && [\text{by (2.104)}] \\ &= \mathbf{CDC}' && [\text{by (2.56)}],\end{aligned}\quad (2.109)$$

where

$$\mathbf{D} = \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix}. \quad (2.110)$$

The expression $\mathbf{A} = \mathbf{CDC}'$ in (2.109) for a symmetric matrix \mathbf{A} in terms of its eigenvalues and eigenvectors is known as the *spectral decomposition* of \mathbf{A} .

Since \mathbf{C} is orthogonal and $\mathbf{C}'\mathbf{C} = \mathbf{CC}' = \mathbf{I}$, we can multiply (2.109) on the left by \mathbf{C}' and on the right by \mathbf{C} to obtain

$$\mathbf{C}'\mathbf{AC} = \mathbf{D}. \quad (2.111)$$

Thus a symmetric matrix \mathbf{A} can be *diagonalized* by an orthogonal matrix containing normalized eigenvectors of \mathbf{A} , and by (2.110) the resulting diagonal matrix contains eigenvalues of \mathbf{A} .

2.11.8 Square Root Matrix

If \mathbf{A} is positive definite, the spectral decomposition of \mathbf{A} in (2.109) can be modified by taking the square roots of the eigenvalues to produce a *square root matrix*,

$$\mathbf{A}^{1/2} = \mathbf{CD}^{1/2}\mathbf{C}', \quad (2.112)$$

where

$$\mathbf{D}^{1/2} = \begin{pmatrix} \sqrt{\lambda_1} & 0 & \cdots & 0 \\ 0 & \sqrt{\lambda_2} & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & \sqrt{\lambda_n} \end{pmatrix}. \quad (2.113)$$

The square root matrix $\mathbf{A}^{1/2}$ is symmetric and serves as the square root of \mathbf{A} :

$$\mathbf{A}^{1/2}\mathbf{A}^{1/2} = (\mathbf{A}^{1/2})^2 = \mathbf{A}. \quad (2.114)$$

2.11.9 Square and Inverse Matrices

Other functions of \mathbf{A} have spectral decompositions analogous to (2.112). Two of these are the square and inverse of \mathbf{A} . If the square matrix \mathbf{A} has eigenvalues

$\lambda_1, \lambda_2, \dots, \lambda_n$ and accompanying eigenvectors $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$, then \mathbf{A}^2 has eigenvalues $\lambda_1^2, \lambda_2^2, \dots, \lambda_n^2$ and eigenvectors $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$. If \mathbf{A} is nonsingular, then \mathbf{A}^{-1} has eigenvalues $1/\lambda_1, 1/\lambda_2, \dots, 1/\lambda_n$ and eigenvectors $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$. If \mathbf{A} is also symmetric, then

$$\mathbf{A}^2 = \mathbf{CD}^2\mathbf{C}', \quad (2.115)$$

$$\mathbf{A}^{-1} = \mathbf{CD}^{-1}\mathbf{C}', \quad (2.116)$$

where $\mathbf{C} = (\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n)$ has as columns the normalized eigenvectors of \mathbf{A} (and of \mathbf{A}^2 and \mathbf{A}^{-1}), $\mathbf{D}^2 = \text{diag}(\lambda_1^2, \lambda_2^2, \dots, \lambda_n^2)$, and $\mathbf{D}^{-1} = \text{diag}(1/\lambda_1, 1/\lambda_2, \dots, 1/\lambda_n)$.

2.11.10 Singular Value Decomposition

In Section 2.11.7 we expressed a symmetric matrix in terms of its eigenvalues and eigenvectors in the spectral decomposition. In a similar manner, we can express any (real) matrix \mathbf{A} in terms of eigenvalues and eigenvectors of $\mathbf{A}'\mathbf{A}$ and \mathbf{AA}' . Let \mathbf{A} be an $n \times p$ matrix of rank k . Then the *singular value decomposition* of \mathbf{A} can be expressed as

$$\mathbf{A} = \mathbf{UDV}', \quad (2.117)$$

where \mathbf{U} is $n \times k$, \mathbf{D} is $k \times k$, and \mathbf{V} is $p \times k$. The diagonal elements of the non-singular diagonal matrix $\mathbf{D} = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_k)$ are the positive square roots of $\lambda_1^2, \lambda_2^2, \dots, \lambda_k^2$, which are the nonzero eigenvalues of $\mathbf{A}'\mathbf{A}$ or of \mathbf{AA}' . The values $\lambda_1, \lambda_2, \dots, \lambda_k$ are called the *singular values* of \mathbf{A} . The k columns of \mathbf{U} are the normalized eigenvectors of \mathbf{AA}' corresponding to the eigenvalues $\lambda_1^2, \lambda_2^2, \dots, \lambda_k^2$. The k columns of \mathbf{V} are the normalized eigenvectors of $\mathbf{A}'\mathbf{A}$ corresponding to the eigenvalues $\lambda_1^2, \lambda_2^2, \dots, \lambda_k^2$. Since the columns of \mathbf{U} and \mathbf{V} are (normalized) eigenvectors of symmetric matrices, they are mutually orthogonal (see Section 2.11.6), and we have $\mathbf{U}'\mathbf{U} = \mathbf{V}'\mathbf{V} = \mathbf{I}$.

2.12 KRONECKER AND VEC NOTATION

When manipulating matrices with block structure, it is often convenient to use Kronecker and vec notation. The Kronecker product of an $m \times n$ matrix \mathbf{A} with a $p \times q$ matrix \mathbf{B} is an $mp \times nq$ matrix that is denoted $\mathbf{A} \otimes \mathbf{B}$ and is defined as

$$\mathbf{A} \otimes \mathbf{B} = \begin{pmatrix} a_{11}\mathbf{B} & a_{12}\mathbf{B} & \cdots & a_{1n}\mathbf{B} \\ a_{21}\mathbf{B} & a_{22}\mathbf{B} & \cdots & a_{2n}\mathbf{B} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1}\mathbf{B} & a_{m2}\mathbf{B} & \cdots & a_{mn}\mathbf{B} \end{pmatrix}. \quad (2.118)$$

For example, because of the block structure in the matrix

$$\mathbf{C} = \begin{pmatrix} 1 & 2 & 3 & 0 & 0 & 0 \\ 4 & 5 & 6 & 0 & 0 & 0 \\ 2 & 4 & 6 & 10 & 20 & 30 \\ 8 & 10 & 12 & 40 & 50 & 60 \end{pmatrix},$$

we can write

$$\mathbf{C} = \begin{pmatrix} 1 \cdot \mathbf{B} & 0 \cdot \mathbf{B} \\ 2 \cdot \mathbf{B} & 10 \cdot \mathbf{B} \end{pmatrix} = \mathbf{A} \otimes \mathbf{B},$$

where

$$\mathbf{A} = \begin{pmatrix} 1 & 0 \\ 2 & 10 \end{pmatrix}, \quad \mathbf{B} = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix}.$$

If \mathbf{A} is an $m \times n$ matrix with columns $\mathbf{a}_1, \dots, \mathbf{a}_n$, then we can refer to the elements of \mathbf{A} in vector (or “vec”) form using

$$\begin{aligned} \text{vec } \mathbf{A} &= \begin{pmatrix} \mathbf{a}_1 \\ \mathbf{a}_2 \\ \vdots \\ \mathbf{a}_n \end{pmatrix} \\ &= (a_{11}, a_{21}, \dots, a_{m1}, a_{12}, a_{22}, \dots, a_{m2}, \dots, a_{1n}, a_{2n}, \dots, a_{mn})', \end{aligned} \tag{2.119}$$

so that $\text{vec } \mathbf{A}$ is an $mn \times 1$ vector.

If \mathbf{A} is an $m \times m$ symmetric matrix, then the m^2 elements in $\text{vec } \mathbf{A}$ will include $m(m - 1)/2$ pairs of identical elements (since $a_{ij} = a_{ji}$). In such settings it is often useful to denote the vector half (or “vech”) of a symmetric matrix in order to include only the unique elements of the matrix. If we separate elements from different columns using semicolons, we can define the “vech” operator with

$$\text{vech } \mathbf{A} = (a_{11}, a_{21}, \dots, a_{m1}; a_{22}, a_{32}, \dots, a_{m2}; \dots; a_{mm})', \tag{2.120}$$

so that $\text{vech } \mathbf{A}$ is an $m(m + 1)/2 \times 1$ vector. Note that $\text{vech } \mathbf{A}$ can be obtained by finding $\text{vec } \mathbf{A}$ and then eliminating the $m(m - 1)/2$ elements above the diagonal of \mathbf{A} .

Assuming that all matrix dimensions for matrices \mathbf{A} , \mathbf{B} , \mathbf{C} , and \mathbf{D} are appropriate for matrix multiplication, the following important properties related to Kronecker products will hold true:

$$(i) \quad (\mathbf{A} \otimes \mathbf{B})(\mathbf{C} \otimes \mathbf{D}) = (\mathbf{AC}) \otimes (\mathbf{BD}) \tag{2.121}$$

$$(ii) \quad \text{vec } \mathbf{ABC} = (\mathbf{C}' \otimes \mathbf{A}) \text{ vec } \mathbf{B} \tag{2.122}$$

$$(iii) \quad (\mathbf{A} \otimes \mathbf{B})' = \mathbf{A}' \otimes \mathbf{B}' \tag{2.123}$$

$$(iv) \quad (\mathbf{A} \otimes \mathbf{B})^{-1} = \mathbf{A}^{-1} \otimes \mathbf{B}^{-1} \tag{2.124}$$

Finally, following the discussion in Fuller (1987, Section 4.3), we can also define an $m^2 \times m(m + 1)/2$ matrix \mathbf{H}_m such that

$$\mathbf{H}_m \text{vech } \mathbf{A} = \text{vec } \mathbf{A}. \tag{2.125}$$

Further, we can define a generalized inverse $\mathbf{H}_m^+ = (\mathbf{H}'_m \mathbf{H}_m)^{-1} \mathbf{H}'_m$ so that

$$\begin{aligned}\mathbf{H}_m^+ \text{vec } \mathbf{A} &= \mathbf{H}_m^+ \mathbf{H}_m \text{vech } \mathbf{A} \\ &= \text{vech } \mathbf{A}.\end{aligned}\quad (2.126)$$

Consider, for example, the 2×2 symmetric matrix

$$\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}.$$

Then

$$\underbrace{\begin{pmatrix} a_{11} \\ a_{21} \\ a_{12} \\ a_{22} \end{pmatrix}}_{\text{vec } \mathbf{A}} = \underbrace{\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}}_{\mathbf{H}_2} \underbrace{\begin{pmatrix} a_{11} \\ a_{21} \\ a_{22} \end{pmatrix}}_{\text{vech } \mathbf{A}},$$

and

$$\underbrace{\begin{pmatrix} a_{11} \\ a_{21} \\ a_{22} \end{pmatrix}}_{\text{vech } \mathbf{A}} = \underbrace{\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1/2 & 1/2 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}}_{\mathbf{H}_2^+} \underbrace{\begin{pmatrix} a_{11} \\ a_{21} \\ a_{12} \\ a_{22} \end{pmatrix}}_{\text{vec } \mathbf{A}}.$$

PROBLEMS

2.1 Let

$$\mathbf{A} = \begin{pmatrix} 4 & 2 & 3 \\ 7 & 5 & 8 \end{pmatrix} \quad \text{and} \quad \mathbf{B} = \begin{pmatrix} 3 & -2 & 4 \\ 6 & 9 & -5 \end{pmatrix}.$$

- (a) Find $\mathbf{A} + \mathbf{B}$ and $\mathbf{A} - \mathbf{B}$.
- (b) Find $\mathbf{A}'\mathbf{A}$ and $\mathbf{A}\mathbf{A}'$.

2.2 Use the matrices \mathbf{A} and \mathbf{B} in Problem 2.1:

- (a) Find $(\mathbf{A} + \mathbf{B})'$ and $\mathbf{A}' + \mathbf{B}'$ and compare them, thus illustrating (2.15).
- (b) Show that $(\mathbf{A}')' = \mathbf{A}$, thus illustrating (2.6).

2.3 Let

$$\mathbf{A} = \begin{pmatrix} 1 & 3 \\ 2 & -1 \end{pmatrix}, \quad \mathbf{B} = \begin{pmatrix} 2 & 0 \\ 1 & 5 \end{pmatrix}.$$

- (a) Find \mathbf{AB} and \mathbf{BA} .
- (b) Find $|\mathbf{AB}|$, $|\mathbf{A}|$, and $|\mathbf{B}|$ and verify that (2.89) holds in this case.

2.4 Use the matrices \mathbf{A} and \mathbf{B} in Problem 2.3:

- (a) Find $\mathbf{A} + \mathbf{B}$ and $\text{tr}(\mathbf{A} + \mathbf{B})$.
 (b) Find $\text{tr}(\mathbf{A})$ and $\text{tr}(\mathbf{B})$ and show that (2.96) holds for these matrices.

2.5 Let

$$\mathbf{A} = \begin{pmatrix} 1 & 2 & 3 \\ 2 & -1 & 1 \end{pmatrix}, \quad \mathbf{B} = \begin{pmatrix} 3 & -2 \\ 2 & 0 \\ -1 & 1 \end{pmatrix}.$$

- (a) Find \mathbf{AB} and \mathbf{BA} .
 (b) Compare $\text{tr}(\mathbf{AB})$ and $\text{tr}(\mathbf{BA})$ and confirm that (2.97) holds here.

2.6 Let

$$\mathbf{A} = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \\ 5 & 10 & 15 \end{pmatrix}, \quad \mathbf{B} = \begin{pmatrix} -1 & 1 & -2 \\ -1 & 1 & -2 \\ 1 & -1 & 2 \end{pmatrix}.$$

- (a) Show that $\mathbf{AB} = \mathbf{O}$.
 (b) Find a vector \mathbf{x} such that $\mathbf{Ax} = \mathbf{0}$.
 (c) Show that $|\mathbf{A}| = 0$.

2.7 Let

$$\mathbf{A} = \begin{pmatrix} 1 & -1 & 4 \\ -1 & 1 & 3 \\ 4 & 3 & 2 \end{pmatrix}, \quad \mathbf{B} = \begin{pmatrix} 3 & -2 & 4 \\ 7 & 1 & 0 \\ 2 & 3 & 5 \end{pmatrix},$$

$$\mathbf{x} = \begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix}, \quad \mathbf{y} = \begin{pmatrix} 3 \\ 2 \\ 1 \end{pmatrix}.$$

Find the following:

- | | | |
|------------------------------|------------------------------|-----------------------------|
| (a) \mathbf{Bx} | (d) $\mathbf{x}'\mathbf{Ay}$ | (g) \mathbf{xx}' |
| (b) $\mathbf{y}'\mathbf{B}$ | (e) $\mathbf{x}'\mathbf{x}$ | (h) \mathbf{xy}' |
| (c) $\mathbf{x}'\mathbf{Ax}$ | (f) $\mathbf{x}'\mathbf{y}$ | (i) $\mathbf{B}'\mathbf{B}$ |

2.8 Use \mathbf{x} , \mathbf{y} , and \mathbf{A} as defined in Problem 2.7:

- (a) Find $\mathbf{x} + \mathbf{y}$ and $\mathbf{x} - \mathbf{y}$.
 (b) Find $(\mathbf{x} - \mathbf{y})'\mathbf{A}(\mathbf{x} - \mathbf{y})$.

2.9 Using \mathbf{B} and \mathbf{x} in Problem 2.7, find \mathbf{Bx} as a linear combination of columns of \mathbf{B} as in (2.67) and compare with \mathbf{Bx} found in Problem 2.7(a).

2.10 Let

$$\mathbf{A} = \begin{pmatrix} 2 & 1 \\ 1 & 3 \end{pmatrix}, \quad \mathbf{B} = \begin{pmatrix} 1 & 4 & 2 \\ 5 & 0 & 3 \end{pmatrix}, \quad \mathbf{I} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

- (a) Show that $(\mathbf{AB})' = \mathbf{B}'\mathbf{A}'$ as in (2.27).
 (b) Show that $\mathbf{AI} = \mathbf{A}$ and that $\mathbf{IB} = \mathbf{B}$.
 (c) Find $|\mathbf{A}|$.

2.11 Let

$$\mathbf{a} = \begin{pmatrix} 1 \\ -3 \\ 2 \end{pmatrix} \quad \text{and} \quad \mathbf{b} = \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix}.$$

- (a) Find $\mathbf{a}'\mathbf{b}$ and $(\mathbf{a}'\mathbf{b})^2$.
 (b) Find \mathbf{bb}' and $\mathbf{a}'(\mathbf{bb}')\mathbf{a}$.
 (c) Compare $(\mathbf{a}'\mathbf{b})^2$ with $\mathbf{a}'(\mathbf{bb}')\mathbf{a}$ and thus illustrate (2.40).

2.12 Let

$$\mathbf{A} = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix}, \quad \mathbf{D} = \begin{pmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{pmatrix}.$$

Find \mathbf{DA} , \mathbf{AD} , and \mathbf{DAD} .

2.13 Let the matrices \mathbf{A} and \mathbf{B} be partitioned as follows:

$$\mathbf{A} = \left(\begin{array}{cc|c} 2 & 1 & 2 \\ 3 & 2 & 0 \\ \hline 1 & 0 & 1 \end{array} \right), \quad \mathbf{B} = \left(\begin{array}{ccc|c} 1 & 1 & 1 & 0 \\ 2 & 1 & 1 & 2 \\ \hline 2 & 3 & 1 & 2 \end{array} \right).$$

- (a) Find \mathbf{AB} as in (2.65) using the indicated partitioning.
 (b) Check by finding \mathbf{AB} in the usual way, ignoring the partitioning.

2.14 Let

$$\mathbf{A} = \begin{pmatrix} 1 & 3 & 2 \\ 2 & 0 & -1 \end{pmatrix}, \quad \mathbf{B} = \begin{pmatrix} 1 & 2 \\ 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \mathbf{C} = \begin{pmatrix} 2 & 1 & 1 \\ 5 & -6 & -4 \end{pmatrix}.$$

Find \mathbf{AB} and \mathbf{CB} . Are they equal? What is the rank of \mathbf{A} , \mathbf{B} , and \mathbf{C} ?

2.15 Let

$$\mathbf{A} = \begin{pmatrix} 5 & 4 & 4 \\ 2 & -3 & 1 \\ 3 & 7 & 2 \end{pmatrix}, \quad \mathbf{B} = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 2 & 3 \end{pmatrix}.$$

- (a) Find $\text{tr}(\mathbf{A})$ and $\text{tr}(\mathbf{B})$.
 (b) Find $\mathbf{A} + \mathbf{B}$ and $\text{tr}(\mathbf{A} + \mathbf{B})$. Is $\text{tr}(\mathbf{A} + \mathbf{B}) = \text{tr}(\mathbf{A}) + \text{tr}(\mathbf{B})$?
 (c) Find $|\mathbf{A}|$ and $|\mathbf{B}|$.
 (d) Find \mathbf{AB} and $|\mathbf{AB}|$. Is $|\mathbf{AB}| = |\mathbf{A}||\mathbf{B}|$?

2.16 Let

$$\mathbf{A} = \begin{pmatrix} 3 & 4 & 3 \\ 4 & 8 & 6 \\ 3 & 6 & 9 \end{pmatrix}.$$

- (a) Show that $|\mathbf{A}| > 0$.
- (b) Using the Cholesky decomposition in Section 2.7, find an upper triangular matrix \mathbf{T} such that $\mathbf{A} = \mathbf{T}'\mathbf{T}$.

2.17 Let

$$\mathbf{A} = \begin{pmatrix} 3 & -5 & -1 \\ -5 & 13 & 0 \\ -1 & 0 & 1 \end{pmatrix}.$$

- (a) Show that $|\mathbf{A}| > 0$.
- (b) Using the Cholesky decomposition in Section 2.7, find an upper triangular matrix \mathbf{T} such that $\mathbf{A} = \mathbf{T}'\mathbf{T}$.

2.18 The columns of the following matrix are mutually orthogonal:

$$\mathbf{A} = \begin{pmatrix} 1 & -1 & 1 \\ 2 & 1 & 0 \\ 1 & -1 & -1 \end{pmatrix}.$$

- (a) Normalize the columns of \mathbf{A} by dividing each column by its length; denote the resulting matrix by \mathbf{C} .
- (b) Show that \mathbf{C} is an orthogonal matrix, that is, $\mathbf{C}'\mathbf{C} = \mathbf{CC}' = \mathbf{I}$.

2.19 Let

$$\mathbf{A} = \begin{pmatrix} 1 & 1 & -2 \\ -1 & 2 & 1 \\ 0 & 1 & -1 \end{pmatrix}.$$

- (a) Find the eigenvalues and associated normalized eigenvectors.
- (b) Find $\text{tr}(\mathbf{A})$ and $|\mathbf{A}|$ and show that $\text{tr}(\mathbf{A}) = \sum_{i=1}^3 \lambda_i$ and $|\mathbf{A}| = \prod_{i=1}^3 \lambda_i$.

2.20 Let

$$\mathbf{A} = \begin{pmatrix} 3 & 1 & 1 \\ 1 & 0 & 2 \\ 1 & 2 & 0 \end{pmatrix}.$$

- (a) The eigenvalues of \mathbf{A} are $1, 4, -2$. Find the normalized eigenvectors and use them as columns in an orthogonal matrix \mathbf{C} .
- (b) Show that $\mathbf{C}'\mathbf{AC} = \mathbf{D}$ as in (2.111), where \mathbf{D} is diagonal with the eigenvalues of \mathbf{A} on the diagonal.
- (c) Show that $\mathbf{A} = \mathbf{CDC}'$ as in (2.109).

2.21 For the positive definite matrix

$$\mathbf{A} = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix},$$

calculate the eigenvalues and eigenvectors and find the square root matrix $\mathbf{A}^{1/2}$ as in (2.112). Check by showing that $(\mathbf{A}^{1/2})^2 = \mathbf{A}$.

2.22 Let

$$\mathbf{A} = \begin{pmatrix} 3 & 6 & -1 \\ 6 & 9 & 4 \\ -1 & 4 & 3 \end{pmatrix}.$$

- (a) Find the spectral decomposition of \mathbf{A} as in (2.109).
- (b) Find the spectral decomposition of \mathbf{A}^2 and show that the diagonal matrix of eigenvalues is equal to the square of the matrix \mathbf{D} found in part (a), thus illustrating (2.115).
- (c) Find the spectral decomposition of \mathbf{A}^{-1} and show that the diagonal matrix of eigenvalues is equal to the inverse of the matrix \mathbf{D} found in part (a), thus illustrating (2.116).

2.23 Find the singular value decomposition of \mathbf{A} as in (2.117), where

$$\mathbf{A} = \begin{pmatrix} 4 & -5 & -1 \\ 7 & -2 & 3 \\ -1 & 4 & -3 \\ 8 & 2 & 6 \end{pmatrix}.$$

2.24 If \mathbf{j} is a vector of 1's, as defined in (2.11), show that the following hold:

- (a) $\mathbf{j}'\mathbf{a} = \mathbf{a}'\mathbf{j} = \sum_i a_i$ as in (2.37)
- (b) $\mathbf{j}'\mathbf{A}$ is a row vector whose elements are the column sums of \mathbf{A} as in (2.38)
- (c) $\mathbf{A}\mathbf{j}$ is a column vector whose elements are the row sums of \mathbf{A} as in (2.38)

2.25 Verify (2.41); that is, show that $(\mathbf{x} - \mathbf{y})'(\mathbf{x} - \mathbf{y}) = \mathbf{x}'\mathbf{x} - 2\mathbf{x}'\mathbf{y} + \mathbf{y}'\mathbf{y}$.

2.26 Show that $\mathbf{A}'\mathbf{A}$ is symmetric, where \mathbf{A} is $n \times p$.

2.27 If \mathbf{a} and $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$ are all $p \times 1$ and \mathbf{A} is $p \times p$, show that (2.42)–(2.45) hold:

- (a) $\sum_{i=1}^n \mathbf{a}'\mathbf{x}_i = \mathbf{a}' \sum_{i=1}^n \mathbf{x}_i$
- (b) $\sum_{i=1}^n \mathbf{A}\mathbf{x}_i = \mathbf{A} \sum_{i=1}^n \mathbf{x}_i$
- (c) $\sum_{i=1}^n (\mathbf{a}'\mathbf{x}_i)^2 = \mathbf{a}'(\sum_{i=1}^n \mathbf{x}_i\mathbf{x}_i')\mathbf{a}$
- (d) $\sum_{i=1}^n \mathbf{A}\mathbf{x}_i(\mathbf{A}\mathbf{x}_i)' = \mathbf{A}(\sum_{i=1}^n \mathbf{x}_i\mathbf{x}_i')\mathbf{A}'$

2.28 Assume that $\mathbf{A} = \begin{pmatrix} \mathbf{a}'_1 \\ \mathbf{a}'_2 \end{pmatrix}$ is $2 \times p$, \mathbf{x} is $p \times 1$, and \mathbf{S} is $p \times p$.

(a) Show that

$$\mathbf{A}\mathbf{x} = \begin{pmatrix} \mathbf{a}'_1\mathbf{x} \\ \mathbf{a}'_2\mathbf{x} \end{pmatrix}$$

as in (2.49).

(b) Show that

$$\mathbf{A}\mathbf{S}\mathbf{A}' = \begin{pmatrix} \mathbf{a}'_1\mathbf{S}\mathbf{a}_1 & \mathbf{a}'_1\mathbf{S}\mathbf{a}_2 \\ \mathbf{a}'_2\mathbf{S}\mathbf{a}_1 & \mathbf{a}'_2\mathbf{S}\mathbf{a}_2 \end{pmatrix}$$

as in (2.50).

2.29 (a) If the rows of \mathbf{A} are denoted by \mathbf{a}'_i , show that $\mathbf{A}'\mathbf{A} = \sum_{i=1}^n \mathbf{a}_i\mathbf{a}'_i$ as in (2.51).

(b) If the columns of \mathbf{A} are denoted by $\mathbf{a}_{(j)}$, show that $\mathbf{A}\mathbf{A}' = \sum_{j=1}^p \mathbf{a}_{(j)}\mathbf{a}'_{(j)}$ as in (2.53).

2.30 Show that $(\mathbf{A}')^{-1} = (\mathbf{A}^{-1})'$ as in (2.75).

2.31 Show that the inverse of the partitioned matrix given in (2.76) is correct by multiplying by

$$\begin{pmatrix} \mathbf{A}_{11} & \mathbf{a}_{12} \\ \mathbf{a}'_{12} & a_{22} \end{pmatrix}$$

to obtain an identity.

2.32 Show that the inverse of $\mathbf{B} + \mathbf{c}\mathbf{c}'$ given in (2.77) is correct by multiplying by $\mathbf{B} + \mathbf{c}\mathbf{c}'$ to obtain an identity.

2.33 Show that $|c\mathbf{A}| = c^n |\mathbf{A}|$ as in (2.85).

2.34 Show that $|\mathbf{A}^{-1}| = 1/|\mathbf{A}|$ as in (2.91).

2.35 If \mathbf{B} is nonsingular and \mathbf{c} is a vector, show that $|\mathbf{B} + \mathbf{c}\mathbf{c}'| = |\mathbf{B}|(1 + \mathbf{c}'\mathbf{B}^{-1}\mathbf{c})$ as in (2.95).

2.36 Show that $\text{tr}(\mathbf{A}'\mathbf{A}) = \text{tr}(\mathbf{A}\mathbf{A}') = \sum_{ij} a_{ij}^2$ as in (2.98).

2.37 Show that $\mathbf{C}\mathbf{C}' = \mathbf{I}$ in (2.102) follows from $\mathbf{C}'\mathbf{C} = \mathbf{I}$ in (2.101).

2.38 Show that the eigenvalues of \mathbf{AB} are the same as those of \mathbf{BA} , as noted in Section 2.11.5.

2.39 If $\mathbf{A}^{1/2}$ is the square root matrix defined in (2.112), show that

(a) $(\mathbf{A}^{1/2})^2 = \mathbf{A}$ as in (2.114),

(b) $|\mathbf{A}^{1/2}|^2 = |\mathbf{A}|$,

(c) $|\mathbf{A}^{1/2}| = |\mathbf{A}|^{1/2}$.

- 2.40** Let \mathbf{A} and \mathbf{B} be $n \times k$ and $p \times p$ matrices, respectively. Use the properties of Kronecker products in Section 2.12 to show that

$$\left[(\mathbf{I}_p \otimes \mathbf{A})' (\mathbf{B} \otimes \mathbf{I}_n)^{-1} (\mathbf{I}_p \otimes \mathbf{A}) \right]^{-1} = \mathbf{B} \otimes (\mathbf{A}' \mathbf{A})^{-1}.$$

CHAPTER 3

CHARACTERIZING AND DISPLAYING MULTIVARIATE DATA

We review some univariate and bivariate procedures in Sections 3.1, 3.2, and 3.3 and then extend them to vectors of higher dimension in the remainder of the chapter.

3.1 MEAN AND VARIANCE OF A UNIVARIATE RANDOM VARIABLE

Informally, a *random variable* may be defined as a variable whose value depends on the outcome of a chance experiment. Generally, we will consider only *continuous* random variables. Some types of multivariate data are only approximations to this ideal, such as test scores or a seven-point semantic differential (Likert) scale consisting of ordered responses ranging from “strongly disagree” to “strongly agree.” Special techniques have been developed for such data, but in many cases, the usual methods designed for continuous data work almost as well.

The *density function* $f(y)$ indicates the relative frequency of occurrence of the random variable y . (We do not use Y to denote the random variable for reasons given at the beginning of Section 3.6.) Thus if $f(y_1) > f(y_2)$, then points in the neighborhood of y_1 are more likely to occur than points in the neighborhood of y_2 .

The *population mean* of a random variable y is defined (informally) as the mean of all possible values of y and is denoted by μ . The mean is also referred to as the *expected value* of y or $E(y)$. If the density $f(y)$ is known, the mean can sometimes be found using methods of calculus, but we will not use these techniques in this text.

If $f(y)$ is unknown, the population mean μ will ordinarily remain unknown unless it has been established from extensive past experience with a stable population. If a large random sample from the population represented by $f(y)$ is available, it is highly probable that the mean of the sample is close to μ .

The *sample mean* of a random sample of n observations y_1, y_2, \dots, y_n is given by the ordinary arithmetic average

$$\bar{y} = \frac{1}{n} \sum_{i=1}^n y_i. \quad (3.1)$$

Generally, \bar{y} will never be equal to μ ; by this we mean that the probability is zero that a sample will ever arise in which \bar{y} is exactly equal to μ . However, \bar{y} is considered a good estimator for μ because $E(\bar{y}) = \mu$ and $\text{var}(\bar{y}) = \sigma^2/n$, where σ^2 is the variance of y . In other words, \bar{y} is an unbiased estimator of μ and has a smaller variance than a single observation y . The variance σ^2 is defined below. The notation $E(\bar{y})$ indicates the mean of all possible values of \bar{y} ; that is, conceptually, every possible sample is obtained from the population, the mean of each is found, and the average of all these sample means is calculated.

If every y in the population is multiplied by a constant a , the expected value is also multiplied by a :

$$E(ay) = aE(y) = a\mu. \quad (3.2)$$

The sample mean has a similar property. If $z_i = ay_i$ for $i = 1, 2, \dots, n$, then

$$\bar{z} = a\bar{y}. \quad (3.3)$$

The *variance* of the population is defined as $\text{var}(y) = \sigma^2 = E(y - \mu)^2$. This is the average squared deviation from the mean and is thus an indication of the extent to which the values of y are spread or scattered. It can be shown that $\sigma^2 = E(y^2) - \mu^2$.

The *sample variance* is defined as

$$s^2 = \frac{\sum_{i=1}^n (y_i - \bar{y})^2}{n-1}, \quad (3.4)$$

which can be shown to be equal to

$$s^2 = \frac{\sum_{i=1}^n y_i^2 - n\bar{y}^2}{n-1}. \quad (3.5)$$

The sample variance s^2 is generally never equal to the population variance σ^2 (the probability of such an occurrence is zero), but it is an unbiased estimator for σ^2 ; that is, $E(s^2) = \sigma^2$. Again the notation $E(s^2)$ indicates the mean of all possible sample variances. The square root of either the population variance or the sample variance is called the *standard deviation*.

Table 3.1 Height and Weight for a Sample of 20 College-Age Males

Person	Height, x	Weight, y	Person	Height, x	Weight, y
1	69	153	11	72	140
2	74	175	12	79	265
3	68	155	13	74	185
4	70	135	14	67	112
5	72	172	15	66	140
6	67	150	16	71	150
7	66	115	17	74	165
8	70	137	18	75	185
9	76	200	19	75	210
10	68	130	20	76	220

If each y is multiplied by a constant a , the population variance is multiplied by a^2 , or $\text{var}(ay) = a^2\sigma^2$. Similarly, if $z_i = ay_i, i = 1, 2, \dots, n$, then the sample variance of z is given by

$$s_z^2 = a^2 s^2. \quad (3.6)$$

3.2 COVARIANCE AND CORRELATION OF BIVARIATE RANDOM VARIABLES

3.2.1 Covariance

If two variables x and y are measured on each research unit (object or subject), we have a *bivariate random variable* (x, y) . Often x and y will tend to covary; if one is above its mean, the other is more likely to be above its mean, and vice versa. For example, height and weight were observed for a sample of 20 college-age males. The data are given in Table 3.1.

The values of height x and weight y from Table 3.1 are both plotted in the vertical direction in Figure 3.1. The tendency for x and y to stay on the same side of the mean is clear in Figure 3.1. This illustrates positive covariance. With negative covariance the points would tend to deviate simultaneously to opposite sides of the mean.

The *population covariance* is defined as $\text{cov}(x, y) = \sigma_{xy} = E[(x - \mu_x)(y - \mu_y)]$, where μ_x and μ_y are the means of x and y , respectively. Thus if x and y are usually both above their means or both below their means, the product $(x - \mu_x)(y - \mu_y)$ will typically be positive and the average value of the product will be positive. Conversely, if x and y tend to fall on opposite sides of their respective means, the product will usually be negative and the average product will be negative. It can be shown that $\sigma_{xy} = E(xy) - \mu_x\mu_y$.




Figure 3.1 Two variables with a tendency to covary.

If the two random variables x and y in a bivariate random variable are added or multiplied, a new random variable is obtained. The mean of $x+y$ or xy is as follows:

$$E(x+y) = E(x) + E(y) \quad (3.7)$$

$$E(xy) = E(x)E(y) \quad \text{if } x, y \text{ independent.} \quad (3.8)$$

Formally, x and y are independent if their joint density factors into the product of their individual densities: $f(x, y) = g(x)h(y)$. Informally, x and y are independent if the random behavior of either of the variables is not affected by the behavior of the other. Note that (3.7) is true whether or not x and y are independent, but (3.8) holds only for x and y independently distributed.

The notion of independence of x and y is more general than that of zero covariance. The covariance σ_{xy} measures linear relationship only, whereas if two random variables are independent, they are not related either linearly or nonlinearly. Independence implies $\sigma_{xy} = 0$, but $\sigma_{xy} = 0$ does not imply independence. It is easy to show that if x and y are independent, then $\sigma_{xy} = 0$:

$$\begin{aligned} \sigma_{xy} &= E(xy) - \mu_x\mu_y \\ &= E(x)E(y) - \mu_x\mu_y \quad [\text{by (3.8)}] \\ &= \mu_x\mu_y - \mu_x\mu_y = 0. \end{aligned}$$

One way to demonstrate that the converse is not true is to construct examples of bivariate x and y that have zero covariance and yet are related in a nonlinear way (the relationship will have zero slope). This is illustrated in Figure 3.2.




Figure 3.2 A sample from a population where x and y have zero covariance and yet are dependent.

If x and y have a bivariate normal distribution (see Chapter 4), then zero covariance implies independence. This is because (1) the covariance measures only linear relationships and (2) in the bivariate normal case, the mean of y given x (or x given y) is a straight line.

The *sample covariance* is defined as

$$s_{xy} = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{n - 1}. \quad (3.9)$$

It can be shown that

$$s_{xy} = \frac{\sum_{i=1}^n x_i y_i - n \bar{x} \bar{y}}{n - 1}. \quad (3.10)$$

Note that s_{xy} is essentially never equal to σ_{xy} (for continuous data), that is, the probability is zero that s_{xy} will equal σ_{xy} . It is true, however, that s_{xy} is an unbiased estimator for σ_{xy} , that is, $E(s_{xy}) = \sigma_{xy}$.

Since $s_{xy} \neq \sigma_{xy}$ in any given sample, this is also true when $\sigma_{xy} = 0$. Thus when the population covariance is zero, no random sample from the population will have zero covariance. The only way a sample from a continuous bivariate distribution will have zero covariance is for the experimenter to choose the values of x and y so that $s_{xy} = 0$. (Such a sample would not be a random sample.) One way to achieve this is to place the values in the form of a grid. This is illustrated in Figure 3.3.

The sample covariance measures only linear relationships. If the points in a bivariate sample follow a curved trend, as, for example, in Figure 3.2, the sample




Figure 3.3 A sample of (x, y) values with zero covariance.

covariance will not measure the strength of the relationship. To see that s_{xy} measures only linear relationships, note that the slope of a simple linear regression line is

$$\hat{\beta}_1 = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sum_{i=1}^n (x_i - \bar{x})^2} = \frac{s_{xy}}{s_x^2}. \quad (3.11)$$

Thus s_{xy} is proportional to the slope, which shows only the linear relationship between y and x .

Variables with zero sample covariance can be said to be *orthogonal*. By (2.99), two sets of numbers a_1, a_2, \dots, a_n and b_1, b_2, \dots, b_n are orthogonal if $\sum_{i=1}^n a_i b_i = 0$. This is true for the centered variables $x_i - \bar{x}$ and $y_i - \bar{y}$ when the sample covariance is zero, that is, $\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y}) = 0$.

■ EXAMPLE 3.2.1

To obtain the sample covariance for the height and weight data in Table 3.1, we first calculate \bar{x} , \bar{y} , and $\sum_i x_i y_i$, where x is height and y is weight:

$$\bar{x} = \frac{69 + 74 + \cdots + 76}{20} = 71.45,$$

$$\bar{y} = \frac{153 + 175 + \cdots + 220}{20} = 164.7,$$

$$\sum_{i=1}^{20} x_i y_i = (69)(153) + (74)(175) + \cdots + (76)(220) = 237,805.$$

Now, by (3.10), we have

$$\begin{aligned}s_{xy} &= \frac{\sum_{i=1}^n x_i y_i - n\bar{x}\bar{y}}{n-1} \\&= \frac{237,805 - (20)(71.45)(164.7)}{19} = 128.88.\end{aligned}$$

By itself, the sample covariance 128.88 is not very meaningful. We are not sure if this represents a small, moderate, or large amount of relationship between y and x . A method of standardizing the covariance is given in the next section.

□

3.2.2 Correlation

Since the covariance depends on the scale of measurement of x and y , it is difficult to compare covariances between different pairs of variables. For example, if we change a measurement from inches to centimeters, the covariance will change. To find a measure of linear relationship that is invariant to changes of scale, we can standardize the covariance by dividing by the standard deviations of the two variables. This standardized covariance is called a *correlation*. The *population correlation* of two random variables x and y is

$$\rho_{xy} = \text{corr}(x, y) = \frac{\sigma_{xy}}{\sigma_x \sigma_y} = \frac{E[(x - \mu_x)(y - \mu_y)]}{\sqrt{E(x - \mu_x)^2} \sqrt{E(y - \mu_y)^2}}, \quad (3.12)$$

and the *sample correlation* is

$$r_{xy} = \frac{s_{xy}}{s_x s_y} = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum_{i=1}^n (x_i - \bar{x})^2} \sqrt{\sum_{i=1}^n (y_i - \bar{y})^2}}. \quad (3.13)$$

Either of these correlations will range between -1 and 1 .

The sample correlation r_{xy} is related to the cosine of the angle between two vectors. Let θ be the angle between vectors \mathbf{a} and \mathbf{b} in Figure 3.4. The vector from the terminal point of \mathbf{a} to the terminal point of \mathbf{b} can be represented as $\mathbf{c} = \mathbf{b} - \mathbf{a}$. Then the law of cosines can be stated in vector form as

$$\begin{aligned}\cos \theta &= \frac{\mathbf{a}'\mathbf{a} + \mathbf{b}'\mathbf{b} - (\mathbf{b} - \mathbf{a})'(\mathbf{b} - \mathbf{a})}{2\sqrt{(\mathbf{a}'\mathbf{a})(\mathbf{b}'\mathbf{b})}} \\&= \frac{\mathbf{a}'\mathbf{a} + \mathbf{b}'\mathbf{b} - (\mathbf{b}'\mathbf{b} + \mathbf{a}'\mathbf{a} - 2\mathbf{a}'\mathbf{b})}{2\sqrt{(\mathbf{a}'\mathbf{a})(\mathbf{b}'\mathbf{b})}} \\&= \frac{\mathbf{a}'\mathbf{b}}{\sqrt{(\mathbf{a}'\mathbf{a})(\mathbf{b}'\mathbf{b})}}.\end{aligned} \quad (3.14)$$

Since $\cos(90^\circ) = 0$, we see from (3.14) that $\mathbf{a}'\mathbf{b} = 0$ when $\theta = 90^\circ$. Thus \mathbf{a} and \mathbf{b} are *perpendicular* when $\mathbf{a}'\mathbf{b} = 0$. By (2.99), two vectors \mathbf{a} and \mathbf{b} , such that




Figure 3.4 Vectors \mathbf{a} and \mathbf{b} in 3-space.

$\mathbf{a}'\mathbf{b} = 0$, are also said to be *orthogonal*. Hence orthogonal vectors are perpendicular in a geometric sense.

To express the correlation in the form given in (3.14), let the n observation vectors $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$ in two dimensions be represented as two vectors $\mathbf{x}' = (x_1, x_2, \dots, x_n)$ and $\mathbf{y}' = (y_1, y_2, \dots, y_n)$ in n dimensions, and let \mathbf{x} and \mathbf{y} be centered as $\mathbf{x} - \bar{x}\mathbf{j}$ and $\mathbf{y} - \bar{y}\mathbf{j}$. Then the cosine of the angle θ between them [see (3.14)] is equal to the sample correlation between x and y :

$$\begin{aligned}\cos \theta &= \frac{(\mathbf{x} - \bar{x}\mathbf{j})'(\mathbf{y} - \bar{y}\mathbf{j})}{\sqrt{[(\mathbf{x} - \bar{x}\mathbf{j})'(\mathbf{x} - \bar{x}\mathbf{j})][(y - \bar{y}\mathbf{j})'(\mathbf{y} - \bar{y}\mathbf{j})]}} \\ &= \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum_{i=1}^n (x_i - \bar{x})^2 \sum_{i=1}^n (y_i - \bar{y})^2}} \\ &= r_{xy}.\end{aligned}\tag{3.15}$$

Thus if the angle θ between the two centered vectors $\mathbf{x} - \bar{x}\mathbf{j}$ and $\mathbf{y} - \bar{y}\mathbf{j}$ is small so that $\cos \theta$ is near 1, r_{xy} will be close to 1. If the two vectors are perpendicular, $\cos \theta$ and r_{xy} will be zero. If the two vectors have nearly opposite directions, r_{xy} will be close to -1.

■ EXAMPLE 3.2.2

To obtain the correlation for the height and weight data of Table 3.1, we first calculate the sample variance of x :




Figure 3.5 Bivariate scatterplot of the data in Figure 3.1.

$$\begin{aligned}s_x^2 &= \frac{\sum_{i=1}^n x_i^2 - n\bar{x}^2}{n-1} \\&= \frac{102,379 - (20)(71.45)^2}{19} = 14.576.\end{aligned}$$

Then $s_x = \sqrt{14.576} = 3.8179$ and, similarly, $s_y = 37.964$. By (3.13), we have

$$r_{xy} = \frac{s_{xy}}{s_x s_y} = \frac{128.88}{(3.8179)(37.964)} = .889.$$

□

3.3 SCATTERPLOTS OF BIVARIATE SAMPLES

Figures 3.2 and 3.3 are examples of *scatterplots* of bivariate samples. In Figure 3.1, the two variables x and y were plotted separately for the data in Table 3.1. Figure 3.5 shows a bivariate scatterplot of the same data.

If the origin is shifted to (\bar{x}, \bar{y}) , as indicated by the dashed lines, then the first and third quadrants contain most of the points. Scatterplots for correlated data typically show a substantial positive or negative slope.

A hypothetical sample of the uncorrelated variables height and IQ is shown in Figure 3.6. We could change the shape of the swarm of points by altering the scale on either axis. But because of the independence assumed for these variables, each




Figure 3.6 A sample of data from a population where x and y are uncorrelated.

quadrant is likely to have as many points as any other quadrant. A tall person is as likely to have a high IQ as a low IQ. A person of low IQ is as likely to be short as to be tall.

3.4 GRAPHICAL DISPLAYS FOR MULTIVARIATE SAMPLES

It is a relatively simple procedure to plot bivariate samples as in Section 3.3. The position of a point shows at once the value of both variables. However, for three or more variables it is a challenge to show graphically the values of all the variables in an observation vector y . On a two-dimensional plot, the value of a third variable could be indicated by color or intensity or size of the plotted point. Four dimensions might be represented by starting with a two-dimensional scatterplot and adding two additional dimensions as line segments at right angles, as in Figure 3.7. The “corner point” represents y_1 and y_2 , whereas y_3 and y_4 are given by the lengths of the two line segments.

We will now describe various methods proposed for representing p dimensions in a plot of an observation vector, where $p > 2$.

Profiles represent each point by p vertical bars, with the heights of the bars depicting the values of the variables. Sometimes the profile is outlined by a polygonal line rather than bars.




Figure 3.7 Four-dimensional plot.

Stars portray the value of each (normalized) variable as a point along a ray from the center to the outside of a circle. The points on the rays are usually joined to form a polygon.

Glyphs (Anderson 1960) are circles of fixed size with rays whose lengths represent the values of the variables. Anderson suggested using only three lengths of rays, thus rounding the variable values to three levels.

Faces (Chernoff 1973) depict each variable as a feature on a face, such as length of nose, size of eyes, shape of eyes, and so on. Flury and Riedwyl (1981) suggested using asymmetric faces, thus increasing the number of representable variables.

Boxes (Hartigan 1975b) show each variable as the length of a dimension of a box. For more than three variables, the dimensions are partitioned into segments.

Among these five methods, Chambers and Kleiner (1982) prefer the star plots because they “combine a reasonably distinctive appearance with computational simplicity and ease of interpretation.” Commenting on the other methods, they state, “Profiles are not so easy to compare as a general shape. Faces are memorable but they are more complex to draw and one must be careful in assigning variables to parameters and in choosing parameter ranges. Faces to some extent disguise the data in the sense that individual data values may not be directly comparable from the plot.”

Table 3.2 Percentage of Republican Votes in Presidential Elections in Six Southern States for Selected Years

State	1932	1936	1940	1960	1964	1968
Missouri	35	38	48	50	36	45
Maryland	36	37	41	46	35	42
Kentucky	40	40	42	54	36	44
Louisiana	7	11	14	29	57	23
Mississippi	4	3	4	25	87	14
South Carolina	2	1	4	49	59	39

■ EXAMPLE 3.4

The data in Table 3.2 are from Kleiner and Hartigan (1981). For these data, the preceding five graphical devices are illustrated in Figure 3.8. The relative magnitudes of the variables can be compared more readily using stars or profiles than faces. \square

3.5 DYNAMIC GRAPHICS

In the previous section, we discussed several methods for static graphical displays. Over the past 25 years, there have been several new methods for visualization of multivariate data that employ computer animation and/or interactive approaches. Because these approaches have profoundly affected the world of statistical graphics, we discuss some dynamic graphics tools here.

For bivariate data, visualization of a data set is fairly straightforward with the standard scatterplot that shows the location of each observation in a two-dimensional space. For p -dimensional data, the problem is much more challenging because there is no simple way to show the locations of objects (observation vectors) in an arbitrarily large p -dimensional space. Finding ways to easily digest the “shape” of a high-dimensional data set is one of the primary goals of dynamic graphical methods. Of great importance is the use of low-dimensional projections of high-dimensional data. For example, we can create a two-dimensional projection (or “shadow”) of an $n \times p$ data matrix \mathbf{X} by post-multiplying it by a $p \times 2$ projection matrix \mathbf{A} .

The *grand tour* of a data set is a set of random projections of the data with a smooth interpolation between projections via intermediate projection matrices (Cook and Swayne, 2007; Asimov, 1985; Buja, Cook, Asimov, and Hurley, 2005). Most commonly, the data set is projected onto a two-dimensional space. When the projections are viewed in sequence as a movie, we see the data points so that the series of projections resembles a swarm of insects. The nature of the movements of the data swarm can provide profound insights about the essential dimensionality of the data as well as the existence of outliers or clusters of observations. Algorithms for the grand tour have been implemented in software packages such as ggobi (Swayne,




Figure 3.8 Profiles, stars, glyphs, faces, and boxes of percentage of Republican votes in six presidential elections in six southern states. The radius of the circles in the stars is 50%. Assignments of variables to facial features are 1932, shape of face; 1936, length of nose; 1940, curvature of mouth; 1960, width of mouth; 1964, slant of eyes; and 1968, length of eyebrows. (From the *Journal of the American Statistical Association*, 1981, p. 262.)

Temple Lang, Buja, and Cook, 2003; Cook and Swayne, 2007) and R (Wickham, Cook, Hofmann, and Buja, 2011; R Development Core Team, 2010).

Interactive methods such as linked brushing of scatterplots and/or parallel coordinates plots have also proven to yield unique insights about the nature of a data set. For each of the n observations in the data set, the *parallel coordinates plot* (Inselberg, 1985; Wegman, 1990) places the p variables at equal intervals along one axis and on the other axis plots the scaled value of each variable as a dot. The p dots for a given observation are then connected to form a multi-segment line. If the variables are time points, this plot is identical to the profile plots used in Section 6.8. (For example, see Figure 6.3.) These plots are useful in illustrating the general correlation structure and highlighting those outliers that do not conform to the typical structure.

In linked brushing, multiple plots of a data set are connected (e.g., two scatterplots are linked together or a scatterplot is linked to a parallel coordinates plot). Observations of interest in one plot are manually “brushed” to change the observations’ appearance. In a scatterplot, the brush will change the points’ color or glyph (symbol). In a parallel coordinates plot, the brush will change the lines’ color or line type. The power of linked brushing is that the same observations in the linked plots will be

appropriately altered, allowing the user to thoroughly explore the nature of outliers and the multidimensional dependencies among variables in the data set. While the static graphical methods discussed in Section 3.4 can illuminate many of the fundamental properties of a data set, dynamic graphics and linked brushing often provide additional insights about complex and often nonlinear relationships among variables.

■ EXAMPLE 3.5

The data set in Table 3.3 contains yield and soil quality measurements at each of 215 locations in a 16-hectare field. The Baker field (Colvin et al., 1997) consists of 8 transects (y) with 28 locations (x) on each transect. The first transect ($y = 1$) lies along the southern boundary of the field, and the last transect ($y = 8$) lies along the northern boundary, with rows roughly 48 meters apart. The locations within a transect run from west ($x = 1$) to east ($x = 28$) and are approximately 12.2 meters apart. Data at nine of the locations in the 8 by 28 layout are missing. At each location, the data set consists of corn yield in bushels per acre for 1997 (Corn97BU) and richness of 10 soil nutrients in parts per million (B, Ca, Cu, Fe, K, Mg, Mn, Na, P, and Zn).

We begin by considering a grand tour of the 10 soil richness variables. In the tour, we observe that many (but not all) of the projections indicate collinearity in the data [compare panels (a) and (b) of Figure 3.9 with panel (c) of the same figure]. Also apparent in a large number of the projections observed during the grand tour is the presence of several outliers. By brushing the five outliers, we change the points from dots to X's. Linking the grand tour to a scatterplot of x (location within transect) versus y (transect) in Figure 3.9(d), we note that the 5 outliers are clustered near the western end of the fourth and fifth transect, which corresponds to a low elevation point in the field (Colvin et al., 1997). Linking these locations to a parallel coordinates plot of the soil richness measures (Figure 3.10), we note the ways in which the profiles for the five outliers deviate from the typical profile. Specifically, the outliers are characterized by unusually high amounts of Ca, Mg, and Na coupled with comparatively low amounts of Fe.

To further illustrate the complicated dependence structures that can be decoded using linked brushing, consider the linked plots in Figure 3.11. Panel (a) of this figure gives a dotplot of Ca with random jitter added in the horizontal direction for ease of visualization. Panel (b) plots log(Fe) against log(yield) (i.e., log of Corn97BU) and indicates a pronounced nonlinear relationship between these variables. By brushing the highest values of Ca, we uncover a complicated dependency between Fe and yield. When Ca is high, the linear relationship between log(Fe) and log(yield) is very weak. However, among low-Ca measurements, the linear relationship between log(Fe) and log(yield) is a relatively strong positive one. □

Table 3.3 Baker Corn Field Measurements of Yield and Soil Richness

x	y	Corn97BU	B	Ca	Cu	Fe	K	Mg	Mn	Na	P	Zn
1	1	174.73	2.74	2608	1.44	192	88	442	62	14	16	0.74
2	1	147.43	4.23	3167	1.73	180	81	522	66	16	20	1.48
3	1	155.39	2.78	2359	1.27	249	110	413	85	15	24	1.33
4	1	141.46	1.92	1766	0.76	189	117	314	63	15	21	0.95
5	1	148.54	3.95	1647	0.48	188	99	304	73	15	17	0.64
6	1	135.30	1.99	1505	0.44	147	72	263	58	13	14	0.60
7	1	145.80	1.87	1755	0.52	125	111	302	47	13	16	0.62
9	1	150.83	2.00	1902	0.86	188	112	325	76	15	23	1.00
10	1	146.21	1.89	1970	0.96	236	105	327	93	12	20	1.16
11	1	167.43	1.73	2069	0.92	209	108	343	85	13	20	0.92
12	1	147.65	4.64	2306	1.15	208	101	381	73	15	25	1.37
13	1	155.80	3.21	2827	1.73	230	85	454	87	13	18	1.37
14	1	152.11	3.07	3187	2.02	201	94	491	95	15	23	2.34
15	1	161.02	2.54	3952	2.44	202	112	628	117	15	36	2.85
16	1	171.58	2.81	2973	1.78	191	82	444	94	14	18	1.75
17	1	160.72	4.35	1963	1.50	260	95	307	88	15	19	1.04
18	1	151.71	4.79	1612	1.15	197	85	259	69	16	21	0.69
19	1	137.44	4.12	1650	0.89	142	86	290	41	16	13	0.35
20	1	136.81	4.10	1759	0.97	139	77	322	63	16	11	0.72
21	1	126.73	4.34	1765	0.98	116	65	342	75	14	9	0.35
22	1	130.15	4.07	1858	0.87	142	79	356	60	17	13	0.36
23	1	139.24	2.35	2152	1.32	220	99	363	110	16	19	1.04
24	1	163.20	6.35	2172	1.21	271	82	342	114	17	16	1.26
25	1	159.59	3.96	2191	1.52	247	46	341	88	12	12	1.02
26	1	147.29	4.65	3179	2.10	298	117	528	91	17	26	1.33
27	1	167.77	5.52	3096	2.14	332	78	519	63	17	20	1.31
28	1	181.79	5.74	3931	2.59	248	86	651	73	19	21	2.16
1	2	148.83	3.31	1997	0.92	212	199	330	75	16	41	0.89
2	2	152.27	4.79	2589	1.32	161	79	360	67	17	19	1.17
3	2	148.18	6.76	2874	1.58	177	129	431	65	19	45	1.99
4	2	171.04	2.66	3697	2.23	255	141	521	92	15	46	3.20
:	:	:	:	:	:	:	:	:	:	:	:	:
25	8	144.88	3.67	2541	1.81	191	88	412	77	14	40	1.65
26	8	159.65	5.99	1952	1.46	228	118	310	84	18	54	1.52
27	8	135.49	3.05	1702	1.34	230	167	267	91	14	61	1.36
28	8	151.52	8.01	1525	1.14	270	158	228	97	18	63	1.17




Figure 3.9 Grand tour of the soil richness measurements in Table 3.3. Panels (a), (b), and (c) show three different projections observed during the grand tour, and panel (d) indicates the geographic locations of the five outliers discovered by the grand tour and brushed to become X's instead of dots.




Figure 3.10 Parallel coordinates plot of the soil richness measurements in Table 3.3. Highlighted lines are linked to the points denoted as outliers in Figure 3.9.




Figure 3.11 Linked and brushed plots of the Baker field data in Table 3.3. The dotplot of Ca in panel (a) is linked to the scatterplot of $\log(\text{Fe})$ versus $\log(\text{yield})$ in panel (b), with the points associated with high Ca brushed to appear as X's.

3.6 MEAN VECTORS

It is a common practice in many texts to use an uppercase letter for a variable name and the corresponding lowercase letter for a particular value or observed value of the random variable, for example, $P(Y > y)$. This notation is convenient in some univariate contexts, but it is often confusing in multivariate analysis, where we use uppercase letters for matrices. In the belief that it is easier to distinguish between a

random vector and an observed value than between a vector and a matrix, throughout this text we follow the notation established in Chapter 2. Uppercase boldface letters are used for matrices of random variables or constants, lowercase boldface letters represent vectors of random variables or constants, and univariate random variables or constants are usually represented by lowercase nonbolded letters.

Let \mathbf{y} represent a random vector of p variables measured on a sampling unit (subject or object). If there are n individuals in the sample, the n *observation vectors* are denoted by $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n$, where

$$\mathbf{y}_i = \begin{pmatrix} y_{i1} \\ y_{i2} \\ \vdots \\ y_{ip} \end{pmatrix}.$$

The *sample mean* vector $\bar{\mathbf{y}}$ can be found either as the average of the n observation vectors or by calculating the average of each of the p variables separately:

$$\bar{\mathbf{y}} = \frac{1}{n} \sum_{i=1}^n \mathbf{y}_i = \begin{pmatrix} \bar{y}_1 \\ \bar{y}_2 \\ \vdots \\ \bar{y}_p \end{pmatrix}, \quad (3.16)$$

where, for example, $\bar{y}_2 = \sum_{i=1}^n y_{i2}/n$. Thus \bar{y}_1 is the mean of the n observations on the first variable, \bar{y}_2 is the mean of the second variable, and so on.

All n observation vectors $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n$ can be transposed to row vectors and listed in the *data matrix* \mathbf{Y} as follows:

$$\mathbf{Y} = \begin{pmatrix} \mathbf{y}'_1 \\ \mathbf{y}'_2 \\ \vdots \\ \mathbf{y}'_i \\ \vdots \\ \mathbf{y}'_n \end{pmatrix} = (\text{units}) \begin{matrix} & & & & & & & (\text{variables}) \\ & 1 & 2 & \cdots & j & \cdots & p \\ 1 & \left(\begin{array}{cccccc} y_{11} & y_{12} & \cdots & y_{1j} & \cdots & y_{1p} \\ y_{21} & y_{22} & \cdots & y_{2j} & \cdots & y_{2p} \\ \vdots & \vdots & & \vdots & & \vdots \\ y_{i1} & y_{i2} & \cdots & y_{ij} & \cdots & y_{ip} \\ \vdots & \vdots & & \vdots & & \vdots \\ y_{n1} & y_{n2} & \cdots & y_{nj} & \cdots & y_{np} \end{array} \right) \\ 2 \\ i \\ n \end{matrix}. \quad (3.17)$$

Since n is usually greater than p , the data can be more conveniently tabulated by entering the observation vectors as rows rather than columns. Note that the first subscript i corresponds to units (subjects or objects) and the second subscript j refers to variables. This convention will be followed whenever possible.

In addition to the two ways of calculating $\bar{\mathbf{y}}$ given in (3.16), we can obtain $\bar{\mathbf{y}}$ from \mathbf{Y} . We sum the n entries in each column of \mathbf{Y} and divide by n , which gives $\bar{\mathbf{y}}'$. This

can be indicated in matrix notation using (2.38),

$$\bar{\mathbf{y}}' = \frac{1}{n} \mathbf{j}' \mathbf{Y}, \quad (3.18)$$

where \mathbf{j}' is a vector of 1's, as defined in (2.11). For example, the second element of $\mathbf{j}' \mathbf{Y}$ is

$$(1, 1, \dots, 1) \begin{pmatrix} y_{12} \\ y_{22} \\ \vdots \\ y_{n2} \end{pmatrix} = \sum_{i=1}^n y_{i2}.$$

We can transpose (3.18) to obtain

$$\bar{\mathbf{y}} = \frac{1}{n} \mathbf{Y}' \mathbf{j}. \quad (3.19)$$

We now turn to populations. The mean of \mathbf{y} over all possible values in the population is called the *population mean vector* or *expected value* of \mathbf{y} . It is defined as a vector of expected values of each variable,

$$E(\mathbf{y}) = E \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_p \end{pmatrix} = \begin{pmatrix} E(y_1) \\ E(y_2) \\ \vdots \\ E(y_p) \end{pmatrix} = \begin{pmatrix} \mu_1 \\ \mu_2 \\ \vdots \\ \mu_p \end{pmatrix} = \boldsymbol{\mu}, \quad (3.20)$$

where μ_j is the population mean of the j th variable.

It can be shown that the expected value of each \bar{y}_j in $\bar{\mathbf{y}}$ is μ_j , that is, $E(\bar{y}_j) = \mu_j$. Thus the expected value of $\bar{\mathbf{y}}$ (over all possible samples) is

$$E(\bar{\mathbf{y}}) = E \begin{pmatrix} \bar{y}_1 \\ \bar{y}_2 \\ \vdots \\ \bar{y}_p \end{pmatrix} = \begin{pmatrix} E(\bar{y}_1) \\ E(\bar{y}_2) \\ \vdots \\ E(\bar{y}_p) \end{pmatrix} = \begin{pmatrix} \mu_1 \\ \mu_2 \\ \vdots \\ \mu_p \end{pmatrix} = \boldsymbol{\mu}. \quad (3.21)$$

Therefore, $\bar{\mathbf{y}}$ is an unbiased estimator of $\boldsymbol{\mu}$. We emphasize again that $\bar{\mathbf{y}}$ is never equal to $\boldsymbol{\mu}$.

■ EXAMPLE 3.6

Table 3.4 gives partial data from Kramer and Jensen (1969a). Three variables were measured (in milliequivalents per 100 g) at 10 different locations in the South. The variables are

y_1 = available soil calcium

y_2 = exchangeable soil calcium

y_3 = turnip green calcium

Table 3.4 Calcium in Soil and Turnip Greens

Location Number	y_1	y_2	y_3
1	35	3.5	2.80
2	35	4.9	2.70
3	40	30.0	4.38
4	10	2.8	3.21
5	6	2.7	2.73
6	20	2.8	2.81
7	35	4.6	2.88
8	35	10.9	2.90
9	35	8.0	3.28
10	30	1.6	3.20

To find the mean vector $\bar{\mathbf{y}}$, we simply calculate the average of each column and obtain

$$\bar{\mathbf{y}}' = (28.1, 7.18, 3.089).$$

□

3.7 COVARIANCE MATRICES

The *sample covariance matrix* $\mathbf{S} = (s_{jk})$ is the matrix of sample variances and covariances of the p variables:

$$\mathbf{S} = (s_{jk}) = \begin{pmatrix} s_{11} & s_{12} & \dots & s_{1p} \\ s_{21} & s_{22} & \dots & s_{2p} \\ \vdots & \vdots & & \vdots \\ s_{p1} & s_{p2} & \dots & s_{pp} \end{pmatrix}. \quad (3.22)$$

In \mathbf{S} the sample variances of the p variables are on the diagonal, and all possible pairwise sample covariances appear off the diagonal. The j th row (column) contains the covariances of y_j with the other $p - 1$ variables.

We give three approaches to obtaining \mathbf{S} . The first of these is to simply calculate the individual elements s_{jk} . The sample variance of the j th variable, $s_{jj} = s_j^2$, is calculated as in (3.4) and (3.5), using the j th column of \mathbf{Y} :

$$s_{jj} = s_j^2 = \frac{1}{n-1} \sum_{i=1}^n (y_{ij} - \bar{y}_j)^2 \quad (3.23)$$

$$= \frac{1}{n-1} \left(\sum_i y_{ij}^2 - n\bar{y}_j^2 \right), \quad (3.24)$$

where \bar{y}_j is the mean of the j th variable, as in (3.16). The sample covariance of the j th and k th variables, s_{jk} , is calculated as in (3.9) or (3.10), using the j th and k th columns of \mathbf{Y} :

$$s_{jk} = \frac{1}{n-1} \sum_{i=1}^n (y_{ij} - \bar{y}_j)(y_{ik} - \bar{y}_k) \quad (3.25)$$

$$= \frac{1}{n-1} \left(\sum_i y_{ij}y_{ik} - n\bar{y}_j\bar{y}_k \right). \quad (3.26)$$

Note that in (3.23) the variance s_{jj} is expressed as s_j^2 , the square of the standard deviation s_j , and that \mathbf{S} is symmetric because $s_{jk} = s_{kj}$ in (3.25). Other names used for the covariance matrix are *variance matrix*, *variance-covariance matrix*, and *dispersion matrix*.

By way of notational clarification, we note that in the univariate case, the sample variance is denoted by s^2 . But in the multivariate case, we denote the sample covariance matrix as \mathbf{S} , not as \mathbf{S}^2 .

The sample covariance matrix \mathbf{S} can also be expressed in terms of the observation vectors:

$$\mathbf{S} = \frac{1}{n-1} \sum_{i=1}^n (\mathbf{y}_i - \bar{\mathbf{y}})(\mathbf{y}_i - \bar{\mathbf{y}})' \quad (3.27)$$

$$= \frac{1}{n-1} \left(\sum_{i=1}^n \mathbf{y}_i \mathbf{y}_i' - n\bar{\mathbf{y}} \bar{\mathbf{y}}' \right). \quad (3.28)$$

Since $(\mathbf{y}_i - \bar{\mathbf{y}})' = (y_{i1} - \bar{y}_1, y_{i2} - \bar{y}_2, \dots, y_{ip} - \bar{y}_p)$, the element in the (1,1) position of $(\mathbf{y}_i - \bar{\mathbf{y}})(\mathbf{y}_i - \bar{\mathbf{y}})'$ is $(y_{i1} - \bar{y}_1)^2$, and when this is summed over i as in (3.27), the result is the numerator of s_{11} in (3.23). Similarly, the (1, 2) element of $(\mathbf{y}_i - \bar{\mathbf{y}})(\mathbf{y}_i - \bar{\mathbf{y}})'$ is $(y_{i1} - \bar{y}_1)(y_{i2} - \bar{y}_2)$, which sums to the numerator of s_{12} in (3.25). Thus (3.27) is equivalent to (3.23) and (3.25), and likewise (3.28) produces (3.24) and (3.26).

We can also obtain \mathbf{S} directly from the data matrix \mathbf{Y} in (3.17), which provides a third approach. The first term in the right side of (3.26), $\sum_i y_{ij}y_{ik}$, is the product of the j th and k th columns of \mathbf{Y} , whereas the second term, $n\bar{y}_j\bar{y}_k$, is the (jk) th element of $n\bar{\mathbf{y}} \bar{\mathbf{y}}'$. It was noted in (2.54) that $\mathbf{Y}'\mathbf{Y}$ is obtained as products of columns of \mathbf{Y} . By (3.18) and (3.19), $\bar{\mathbf{y}} = \mathbf{Y}'\mathbf{j}/n$ and $\bar{\mathbf{y}}' = \mathbf{j}'\mathbf{Y}/n$; and using (2.36), we have $n\bar{\mathbf{y}} \bar{\mathbf{y}}' = \mathbf{Y}'(\mathbf{J}/n)\mathbf{Y}$. Thus \mathbf{S} can be written as

$$\begin{aligned} \mathbf{S} &= \frac{1}{n-1} \left[\mathbf{Y}'\mathbf{Y} - \mathbf{Y}' \left(\frac{1}{n}\mathbf{J} \right) \mathbf{Y} \right] \\ &= \frac{1}{n-1} \mathbf{Y}' \left(\mathbf{I} - \frac{1}{n}\mathbf{J} \right) \mathbf{Y} \quad [\text{by (2.30)}]. \end{aligned} \quad (3.29)$$

Expression (3.29) is a convenient representation of \mathbf{S} , since it makes direct use of the data matrix \mathbf{Y} . However, the matrix $\mathbf{I} - \mathbf{J}/n$ is $n \times n$ and may be unwieldy in computation if n is large.

If \mathbf{y} is a random vector taking on any possible value in a multivariate population, the *population covariance matrix* is defined as

$$\Sigma = \text{cov}(\mathbf{y}) = \begin{pmatrix} \sigma_{11} & \sigma_{12} & \cdots & \sigma_{1p} \\ \sigma_{21} & \sigma_{22} & \cdots & \sigma_{2p} \\ \vdots & \vdots & & \vdots \\ \sigma_{p1} & \sigma_{p2} & \cdots & \sigma_{pp} \end{pmatrix}. \quad (3.30)$$

The diagonal elements $\sigma_{jj} = \sigma_j^2$ are the population variances of the y 's, and the off-diagonal elements σ_{jk} are the population covariances of all possible pairs of y 's.

The notation Σ for the covariance matrix is widely used and seems natural because Σ is the uppercase version of σ . It should not be confused with the same symbol used for summation of a series. The difference should always be apparent from the context. To help further distinguish the two uses, the covariance matrix Σ will differ in typeface and in size from the summation symbol \sum . Also, whenever they appear together, the summation symbol will have an index of summation, such as $\sum_{i=1}^n$.

The population covariance matrix in (3.30) can also be found as

$$\Sigma = E[(\mathbf{y} - \boldsymbol{\mu})(\mathbf{y} - \boldsymbol{\mu})'], \quad (3.31)$$

which is analogous to (3.27) for the sample covariance matrix. The $p \times p$ matrix $(\mathbf{y} - \boldsymbol{\mu})(\mathbf{y} - \boldsymbol{\mu})'$ is a random matrix. The expected value of a random matrix is defined as the matrix of expected values of the corresponding elements. To see that (3.31) produces population variances and covariances of the p variables as in (3.30), note that

$$\begin{aligned} \Sigma &= E[(\mathbf{y} - \boldsymbol{\mu})(\mathbf{y} - \boldsymbol{\mu})'] = E \begin{pmatrix} y_1 - \mu_1 \\ y_2 - \mu_2 \\ \vdots \\ y_p - \mu_p \end{pmatrix} (y_1 - \mu_1, y_2 - \mu_2, \dots, y_p - \mu_p)' \\ &= E \begin{pmatrix} (y_1 - \mu_1)^2 & (y_1 - \mu_1)(y_2 - \mu_2) & \cdots & (y_1 - \mu_1)(y_p - \mu_p) \\ (y_2 - \mu_2)(y_1 - \mu_1) & (y_2 - \mu_2)^2 & \cdots & (y_2 - \mu_2)(y_p - \mu_p) \\ \vdots & \vdots & & \vdots \\ (y_p - \mu_p)(y_1 - \mu_1) & (y_p - \mu_p)(y_2 - \mu_2) & \cdots & (y_p - \mu_p)^2 \end{pmatrix} \\ &= \begin{pmatrix} E(y_1 - \mu_1)^2 & E(y_1 - \mu_1)(y_2 - \mu_2) & \cdots & E(y_1 - \mu_1)(y_p - \mu_p) \\ E(y_2 - \mu_2)(y_1 - \mu_1) & E(y_2 - \mu_2)^2 & \cdots & E(y_2 - \mu_2)(y_p - \mu_p) \\ \vdots & \vdots & & \vdots \\ E(y_p - \mu_p)(y_1 - \mu_1) & E(y_p - \mu_p)(y_2 - \mu_2) & \cdots & E(y_p - \mu_p)^2 \end{pmatrix} \\ &= \begin{pmatrix} \sigma_{11} & \sigma_{12} & \cdots & \sigma_{1p} \\ \sigma_{21} & \sigma_{22} & \cdots & \sigma_{2p} \\ \vdots & \vdots & & \vdots \\ \sigma_{p1} & \sigma_{p2} & \cdots & \sigma_{pp} \end{pmatrix}. \end{aligned}$$

It can be easily shown that Σ can be expressed in a form analogous to (3.28):

$$\Sigma = E(\mathbf{y}\mathbf{y}') - \boldsymbol{\mu}\boldsymbol{\mu}' . \quad (3.32)$$

Since $E(s_{jk}) = \sigma_{jk}$ for all j, k , the sample covariance matrix \mathbf{S} is an unbiased estimator for Σ :

$$E(\mathbf{S}) = \Sigma . \quad (3.33)$$

As in the univariate case, we note that it is the average of all possible values of \mathbf{S} that is equal to Σ . Generally, \mathbf{S} will never be equal to Σ .

■ EXAMPLE 3.7

To calculate the sample covariance matrix for the calcium data of Table 3.4 using the computational forms (3.24) and (3.26), we need the sum of squares of each column and the sum of products of each pair of columns. We illustrate the computation of s_{13} .

$$\sum_{i=1}^{10} y_{i1}y_{i3} = (35)(2.80) + (35)(2.70) + \cdots + (30)(3.20) = 885.48.$$

From Example 3.6 we have $\bar{y}_1 = 28.1$ and $\bar{y}_3 = 3.089$. By (3.26), we obtain

$$s_{13} = \frac{1}{10-1}[885.48 - 10(28.1)(3.089)] = \frac{17.471}{9} = 1.9412.$$

Continuing in this fashion, we obtain

$$\mathbf{S} = \begin{pmatrix} 140.54 & 49.68 & 1.94 \\ 49.68 & 72.25 & 3.68 \\ 1.94 & 3.68 & .25 \end{pmatrix} .$$

□

3.8 CORRELATION MATRICES

The sample correlation between the j th and k th variables is defined in (3.13) as

$$r_{jk} = \frac{s_{jk}}{\sqrt{s_{jj}s_{kk}}} = \frac{s_{jk}}{s_js_k} . \quad (3.34)$$

The *sample correlation matrix* is analogous to the covariance matrix with correlations in place of covariances:

$$\mathbf{R} = (r_{jk}) = \begin{pmatrix} 1 & r_{12} & \cdots & r_{1p} \\ r_{21} & 1 & \cdots & r_{2p} \\ \vdots & \vdots & & \vdots \\ r_{p1} & r_{p2} & \cdots & 1 \end{pmatrix} . \quad (3.35)$$

The second row, for example, contains the correlation of y_2 with each of the y 's (including the correlation of y_2 with itself, which is 1). Of course, the matrix \mathbf{R} is symmetric, since $r_{jk} = r_{kj}$.

The correlation matrix can be obtained from the covariance matrix and vice versa. Define

$$\begin{aligned}\mathbf{D}_s &= \text{diag}(\sqrt{s_{11}}, \sqrt{s_{22}}, \dots, \sqrt{s_{pp}}) \\ &= \text{diag}(s_1, s_2, \dots, s_p) \\ &= \begin{pmatrix} s_1 & 0 & \cdots & 0 \\ 0 & s_2 & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & s_p \end{pmatrix}. \end{aligned} \quad (3.36)$$

Then by (2.57)

$$\mathbf{R} = \mathbf{D}_s^{-1} \mathbf{S} \mathbf{D}_s^{-1}, \quad (3.37)$$

$$\mathbf{S} = \mathbf{D}_s \mathbf{R} \mathbf{D}_s. \quad (3.38)$$

The *population correlation matrix* analogous to (3.35) is defined as

$$\mathbf{P}_\rho = (\rho_{jk}) = \begin{pmatrix} 1 & \rho_{12} & \cdots & \rho_{1p} \\ \rho_{21} & 1 & \cdots & \rho_{2p} \\ \vdots & \vdots & & \vdots \\ \rho_{p1} & \rho_{p2} & \cdots & 1 \end{pmatrix}, \quad (3.39)$$

where

$$\rho_{jk} = \frac{\sigma_{jk}}{\sigma_j \sigma_k},$$

as in (3.12).

■ EXAMPLE 3.8

In Example 3.7, we obtained the sample covariance matrix \mathbf{S} for the calcium data in Table 3.4. To obtain the sample correlation matrix for the same data, we can calculate the individual elements using (3.34) or use the direct matrix operation in (3.37). The diagonal matrix \mathbf{D}_s can be found by taking the square roots of the diagonal elements of \mathbf{S} ,

$$\mathbf{D}_s = \begin{pmatrix} 11.8551 & 0 & 0 \\ 0 & 8.4999 & 0 \\ 0 & 0 & .5001 \end{pmatrix}$$

(note that we have used the unrounded version of \mathbf{S} for computation). Then by (3.37),

$$\mathbf{R} = \mathbf{D}_s^{-1} \mathbf{S} \mathbf{D}_s^{-1} = \begin{pmatrix} 1.000 & .493 & .327 \\ .493 & 1.000 & .865 \\ .327 & .865 & 1.000 \end{pmatrix}.$$

Note that $.865 > .493 > .327$, which is a different order than that of the covariances in \mathbf{S} in Example 3.7. Thus we cannot compare covariances, even within the same matrix \mathbf{S} . \square

3.9 MEAN VECTORS AND COVARIANCE MATRICES FOR SUBSETS OF VARIABLES

3.9.1 Two Subsets

Sometimes a researcher is interested in two different kinds of variables, both measured on the same sampling unit. This corresponds to type 2 data in Section 1.4. For example, several classroom behaviors are observed for students, and during the same time period (the basic experimental unit) several teacher behaviors are also observed. The researcher wishes to study the relationships between the pupil variables and the teacher variables.

We will denote the two subvectors by \mathbf{y} and \mathbf{x} , with p variables in \mathbf{y} and q variables in \mathbf{x} . Thus each observation vector in a sample is partitioned as

$$\begin{pmatrix} \mathbf{y}_i \\ \mathbf{x}_i \end{pmatrix} = \begin{pmatrix} y_{i1} \\ \vdots \\ y_{ip} \\ x_{i1} \\ \vdots \\ x_{iq} \end{pmatrix}, \quad i = 1, 2, \dots, n. \quad (3.40)$$

Hence there are $p + q$ variables in each of n observation vectors. In Chapter 10 we will discuss regression of the y 's on the x 's, and in Chapter 11 we will define a measure of correlation between the y 's and the x 's.

For the sample of n observation vectors, the mean vector and covariance matrix have the form

$$\begin{pmatrix} \bar{\mathbf{y}} \\ \bar{\mathbf{x}} \end{pmatrix} = \begin{pmatrix} \bar{y}_1 \\ \vdots \\ \bar{y}_p \\ \bar{x}_1 \\ \vdots \\ \bar{x}_q \end{pmatrix}, \quad (3.41)$$

$$\mathbf{S} = \begin{pmatrix} \mathbf{S}_{yy} & \mathbf{S}_{yx} \\ \mathbf{S}_{xy} & \mathbf{S}_{xx} \end{pmatrix}, \quad (3.42)$$

where \mathbf{S}_{yy} is $p \times p$, \mathbf{S}_{yx} is $p \times q$, \mathbf{S}_{xy} is $q \times p$, and \mathbf{S}_{xx} is $q \times q$. Note that because of the symmetry of \mathbf{S} ,

$$\mathbf{S}_{xy} = \mathbf{S}'_{yx}. \quad (3.43)$$

Thus (3.42) could be written

$$\mathbf{S} = \begin{pmatrix} \mathbf{S}_{yy} & \mathbf{S}_{yx} \\ \mathbf{S}'_{yx} & \mathbf{S}_{xx} \end{pmatrix}. \quad (3.44)$$

To illustrate (3.41) and (3.42), let $p = 2$ and $q = 3$. Then

$$\begin{pmatrix} \bar{\mathbf{y}} \\ \bar{\mathbf{x}} \end{pmatrix} = \begin{pmatrix} \bar{y}_1 \\ \bar{y}_2 \\ \hline \bar{x}_1 \\ \bar{x}_2 \\ \bar{x}_3 \end{pmatrix},$$

$$\mathbf{S} = \begin{pmatrix} \mathbf{S}_{yy} & \mathbf{S}_{yx} \\ \mathbf{S}_{xy} & \mathbf{S}_{xx} \end{pmatrix} = \left(\begin{array}{cc|ccc} s_{y_1}^2 & s_{y_1 y_2} & s_{y_1 x_1} & s_{y_1 x_2} & s_{y_1 x_3} \\ s_{y_2 y_1} & s_{y_2}^2 & s_{y_2 x_1} & s_{y_2 x_2} & s_{y_2 x_3} \\ \hline s_{x_1 y_1} & s_{x_1 y_2} & s_{x_1}^2 & s_{x_1 x_2} & s_{x_1 x_3} \\ s_{x_2 y_1} & s_{x_2 y_2} & s_{x_2 x_1} & s_{x_2}^2 & s_{x_2 x_3} \\ s_{x_3 y_1} & s_{x_3 y_2} & s_{x_3 x_1} & s_{x_3 x_2} & s_{x_3}^2 \end{array} \right).$$

The pattern in each of \mathbf{S}_{yy} , \mathbf{S}_{yx} , \mathbf{S}_{xy} , and \mathbf{S}_{xx} is clearly seen in this illustration. For example, the first row of \mathbf{S}_{yx} has the covariance of y_1 with each of x_1, x_2, x_3 ; the second row exhibits covariances of y_2 with the three x 's. On the other hand, \mathbf{S}_{xy} has as its first row the covariances of x_1 with y_1 and y_2 , and so on. Thus $\mathbf{S}_{xy} = \mathbf{S}'_{yx}$ as noted in (3.43).

The analogous population results for a partitioned random vector are

$$E\begin{pmatrix} \mathbf{y} \\ \mathbf{x} \end{pmatrix} = \begin{pmatrix} E(\mathbf{y}) \\ E(\mathbf{x}) \end{pmatrix} = \begin{pmatrix} \boldsymbol{\mu}_y \\ \boldsymbol{\mu}_x \end{pmatrix}, \quad (3.45)$$

$$\text{cov}\begin{pmatrix} \mathbf{y} \\ \mathbf{x} \end{pmatrix} = \boldsymbol{\Sigma} = \begin{pmatrix} \boldsymbol{\Sigma}_{yy} & \boldsymbol{\Sigma}_{yx} \\ \boldsymbol{\Sigma}_{xy} & \boldsymbol{\Sigma}_{xx} \end{pmatrix}, \quad (3.46)$$

where $\boldsymbol{\Sigma}_{xy} = \boldsymbol{\Sigma}'_{yx}$. The submatrix $\boldsymbol{\Sigma}_{yy}$ is a $p \times p$ covariance matrix containing the variances of y_1, y_2, \dots, y_p on the diagonal and the covariance of each y_j with each y_k off the diagonal. Similarly, $\boldsymbol{\Sigma}_{xx}$ is the $q \times q$ covariance matrix of x_1, x_2, \dots, x_q . The matrix $\boldsymbol{\Sigma}_{yx}$ is $p \times q$ and contains the covariance of each y_j with each x_k . The covariance matrix $\boldsymbol{\Sigma}_{yx}$ is also denoted by $\text{cov}(\mathbf{y}, \mathbf{x})$, that is,

$$\text{cov}(\mathbf{y}, \mathbf{x}) = \boldsymbol{\Sigma}_{yx}. \quad (3.47)$$

Note the difference in meaning between $\text{cov}(\mathbf{x}) = \boldsymbol{\Sigma}$ in (3.46) and $\text{cov}(\mathbf{y}, \mathbf{x}) = \boldsymbol{\Sigma}_{yx}$ in (3.47); $\text{cov}(\mathbf{x})$ involves a single vector containing $p + q$ variables, and $\text{cov}(\mathbf{y}, \mathbf{x})$ involves two vectors.

If \mathbf{x} and \mathbf{y} are independent, then $\boldsymbol{\Sigma}_{yx} = \mathbf{O}$. This means that each y_j is uncorrelated with each x_k so that $\sigma_{y_j x_k} = 0$ for $j = 1, 2, \dots, p$; $k = 1, 2, \dots, q$.

■ EXAMPLE 3.9.1

Reaven and Miller (1979; see also Andrews and Herzberg 1985, pp. 215–219) measured five variables in a comparison of normal patients and diabetics. In Table 3.5 we give partial data for normal patients only. The three variables of major interest were

$$\begin{aligned}x_1 &= \text{glucose intolerance} \\x_2 &= \text{insulin response to oral glucose} \\x_3 &= \text{insulin resistance}\end{aligned}$$

The two additional variables of minor interest were

$$\begin{aligned}y_1 &= \text{relative weight} \\y_2 &= \text{fasting plasma glucose}\end{aligned}$$

The mean vector, partitioned as in (3.41), is

$$\left(\begin{array}{c} \bar{\mathbf{y}} \\ \bar{\mathbf{x}} \end{array} \right) = \left(\begin{array}{c} \bar{y}_1 \\ \bar{y}_2 \\ \hline \bar{x}_1 \\ \bar{x}_2 \\ \bar{x}_3 \end{array} \right) = \left(\begin{array}{c} .918 \\ 90.41 \\ \hline 340.83 \\ 171.37 \\ 97.78 \end{array} \right).$$

The covariance matrix, partitioned as in the illustration following (3.44), is

$$\begin{aligned}\mathbf{S} &= \left(\begin{array}{cc} \mathbf{S}_{yy} & \mathbf{S}_{yx} \\ \mathbf{S}_{xy} & \mathbf{S}_{xx} \end{array} \right) \\&= \left(\begin{array}{cc|ccc} .0162 & .2160 & .7872 & -.2138 & 2.189 \\ .2160 & 70.56 & 26.23 & -23.96 & -20.84 \\ \hline .7872 & 26.23 & 1106 & 396.7 & 108.4 \\ -.2138 & -23.96 & 396.7 & 2382 & 1143 \\ 2.189 & -20.84 & 108.4 & 1143 & 2136 \end{array} \right).\end{aligned}$$

Note that \mathbf{S}_{yy} and \mathbf{S}_{xx} are symmetric and that \mathbf{S}_{xy} is the transpose of \mathbf{S}_{yx} . \square

3.9.2 Three or More Subsets

In some cases, three or more subsets of variables are of interest. If the observation vector \mathbf{y} is partitioned as

$$\mathbf{y} = \left(\begin{array}{c} \mathbf{y}_1 \\ \mathbf{y}_2 \\ \vdots \\ \mathbf{y}_k \end{array} \right),$$

Table 3.5 Relative Weight, Blood Glucose, and Insulin Levels

Patient Number	y_1	y_2	x_1	x_2	x_3
1	.81	80	356	124	55
2	.95	97	289	117	76
3	.94	105	319	143	105
4	1.04	90	356	199	108
5	1.00	90	323	240	143
6	.76	86	381	157	165
7	.91	100	350	221	119
8	1.10	85	301	186	105
9	.99	97	379	142	98
10	.78	97	296	131	94
11	.90	91	353	221	53
12	.73	87	306	178	66
13	.96	78	290	136	142
14	.84	90	371	200	93
15	.74	86	312	208	68
16	.98	80	393	202	102
17	1.10	90	364	152	76
18	.85	99	359	185	37
19	.83	85	296	116	60
20	.93	90	345	123	50
21	.95	90	378	136	47
22	.74	88	304	134	50
23	.95	95	347	184	91
24	.97	90	327	192	124
25	.72	92	386	279	74
26	1.11	74	365	228	235
27	1.20	98	365	145	158
28	1.13	100	352	172	140
29	1.00	86	325	179	145
30	.78	98	321	222	99
31	1.00	70	360	134	90
32	1.00	99	336	143	105
33	.71	75	352	169	32
34	.76	90	353	263	165
35	.89	85	373	174	78
36	.88	99	376	134	80
37	1.17	100	367	182	54
38	.85	78	335	241	175
39	.97	106	396	128	80
⋮	⋮	⋮	⋮	⋮	⋮
46	.91	86	328	106	56

where \mathbf{y}_1 has p_1 variables, \mathbf{y}_2 has p_2, \dots, \mathbf{y}_k has p_k , with $p = p_1 + p_2 + \dots + p_k$, then the sample mean vector and covariance matrix are given by

$$\bar{\mathbf{y}} = \begin{pmatrix} \bar{\mathbf{y}}_1 \\ \bar{\mathbf{y}}_2 \\ \vdots \\ \bar{\mathbf{y}}_k \end{pmatrix}, \quad (3.48)$$

$$\mathbf{S} = \begin{pmatrix} \mathbf{S}_{11} & \mathbf{S}_{12} & \cdots & \mathbf{S}_{1k} \\ \mathbf{S}_{21} & \mathbf{S}_{22} & \cdots & \mathbf{S}_{2k} \\ \vdots & \vdots & & \vdots \\ \mathbf{S}_{k1} & \mathbf{S}_{k2} & \cdots & \mathbf{S}_{kk} \end{pmatrix}. \quad (3.49)$$

The $p_2 \times p_k$ submatrix \mathbf{S}_{2k} , for example, contains the covariances of the variables in \mathbf{y}_2 with the variables in \mathbf{y}_k .

The corresponding population results are

$$\boldsymbol{\mu} = \begin{pmatrix} \mu_1 \\ \mu_2 \\ \vdots \\ \mu_k \end{pmatrix}, \quad (3.50)$$

$$\boldsymbol{\Sigma} = \begin{pmatrix} \boldsymbol{\Sigma}_{11} & \boldsymbol{\Sigma}_{12} & \cdots & \boldsymbol{\Sigma}_{1k} \\ \boldsymbol{\Sigma}_{21} & \boldsymbol{\Sigma}_{22} & \cdots & \boldsymbol{\Sigma}_{2k} \\ \vdots & \vdots & & \vdots \\ \boldsymbol{\Sigma}_{k1} & \boldsymbol{\Sigma}_{k2} & \cdots & \boldsymbol{\Sigma}_{kk} \end{pmatrix}. \quad (3.51)$$

3.10 LINEAR COMBINATIONS OF VARIABLES

3.10.1 Sample Properties

We are frequently interested in linear combinations of the variables y_1, y_2, \dots, y_p . For example, two of the types of linear functions we use in later chapters are (1) linear combinations that maximize some function and (2) linear combinations that compare variables, for example, $y_1 - y_3$. In this section, we investigate the means, variances, and covariances of linear combinations.

Let a_1, a_2, \dots, a_p be constants and consider the linear combination of the elements of the vector \mathbf{y} ,

$$z = a_1 y_1 + a_2 y_2 + \cdots + a_p y_p = \mathbf{a}' \mathbf{y}, \quad (3.52)$$

where $\mathbf{a}' = (a_1, a_2, \dots, a_p)$. If the same coefficient vector \mathbf{a} is applied to each \mathbf{y}_i in a sample, we have

$$\begin{aligned} z_i &= a_1 y_{i1} + a_2 y_{i2} + \cdots + a_p y_{ip} \\ &= \mathbf{a}' \mathbf{y}_i, \quad i = 1, 2, \dots, n. \end{aligned} \quad (3.53)$$

The sample mean of z can be found either by averaging the n values $z_1 = \mathbf{a}'\mathbf{y}_1, z_2 = \mathbf{a}'\mathbf{y}_2, \dots, z_n = \mathbf{a}'\mathbf{y}_n$ or as a linear combination of $\bar{\mathbf{y}}$, the sample mean vector of $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n$:

$$\bar{z} = \frac{1}{n} \sum_{i=1}^n z_i = \mathbf{a}'\bar{\mathbf{y}}. \quad (3.54)$$

The result in (3.54) is analogous to the univariate result (3.3), $\bar{z} = a\bar{y}$, where $z_i = ay_i, i = 1, 2, \dots, n$.

Similarly, the sample variance of $z_i = \mathbf{a}'\mathbf{y}_i, i = 1, 2, \dots, n$, can be found as the sample variance of z_1, z_2, \dots, z_n or directly from \mathbf{a} and \mathbf{S} , where \mathbf{S} is the sample covariance matrix of $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n$:

$$s_z^2 = \frac{\sum_{i=1}^n (z_i - \bar{z})^2}{n - 1} = \mathbf{a}'\mathbf{S}\mathbf{a}. \quad (3.55)$$

Note that $s_z^2 = \mathbf{a}'\mathbf{S}\mathbf{a}$ is the multivariate analogue of the univariate result in (3.6), $s_z^2 = a^2 s^2$, where $z_i = ay_i, i = 1, 2, \dots, n$, and s^2 is the variance of y_1, y_2, \dots, y_n .

Since a variance is always nonnegative, we have $s_z^2 \geq 0$, and therefore $\mathbf{a}'\mathbf{S}\mathbf{a} \geq 0$, for every \mathbf{a} . Hence \mathbf{S} is at least positive semidefinite (see Section 2.7). If the variables are continuous and are not linearly related, and if $n - 1 > p$ (so that \mathbf{S} is full rank), then \mathbf{S} is positive definite (with probability 1).

If we define another linear combination $w = \mathbf{b}'\mathbf{y} = b_1y_1 + b_2y_2 + \dots + b_py_p$, where $\mathbf{b}' = (b_1, b_2, \dots, b_p)$ is a vector of constants different from \mathbf{a}' , then the sample covariance of z and w is given by

$$s_{zw} = \frac{\sum_{i=1}^n (z_i - \bar{z})(w_i - \bar{w})}{n - 1} = \mathbf{a}'\mathbf{S}\mathbf{b}. \quad (3.56)$$

The sample correlation between z and w is readily obtained as

$$r_{zw} = \frac{s_{zw}}{\sqrt{s_z^2 s_w^2}} = \frac{\mathbf{a}'\mathbf{S}\mathbf{b}}{\sqrt{(\mathbf{a}'\mathbf{S}\mathbf{a})(\mathbf{b}'\mathbf{S}\mathbf{b})}}. \quad (3.57)$$

We now denote the two constant vectors \mathbf{a} and \mathbf{b} as \mathbf{a}_1 and \mathbf{a}_2 to facilitate later expansion to more than two such vectors. Let

$$\mathbf{A} = \begin{pmatrix} \mathbf{a}'_1 \\ \mathbf{a}'_2 \end{pmatrix}$$

and define

$$\mathbf{z} = \begin{pmatrix} \mathbf{a}'_1\mathbf{y} \\ \mathbf{a}'_2\mathbf{y} \end{pmatrix} = \begin{pmatrix} z_1 \\ z_2 \end{pmatrix}.$$

Then we can factor \mathbf{y} from this expression by (2.49):

$$\mathbf{z} = \begin{pmatrix} \mathbf{a}'_1 \\ \mathbf{a}'_2 \end{pmatrix} \mathbf{y} = \mathbf{Ay}.$$

If we evaluate the bivariate \mathbf{z}_i for each p -variate \mathbf{y}_i in the sample, we obtain $\mathbf{z}_i = \mathbf{A}\mathbf{y}_i, i = 1, 2, \dots, n$, and the average of \mathbf{z} over the sample can be found from $\bar{\mathbf{y}}$:

$$\bar{\mathbf{z}} = \begin{pmatrix} \bar{z}_1 \\ \bar{z}_2 \end{pmatrix} = \begin{pmatrix} \mathbf{a}'_1 \bar{\mathbf{y}} \\ \mathbf{a}'_2 \bar{\mathbf{y}} \end{pmatrix} \quad [\text{by (3.54)}] \quad (3.58)$$

$$= \begin{pmatrix} \mathbf{a}'_1 \\ \mathbf{a}'_2 \end{pmatrix} \bar{\mathbf{y}} = \mathbf{A}\bar{\mathbf{y}} \quad [\text{by (2.49)}]. \quad (3.59)$$

We can use (3.55) and (3.56) to construct the sample covariance matrix for \mathbf{z} :

$$\begin{aligned} \mathbf{S}_z &= \begin{pmatrix} s_{z_1}^2 & s_{z_1 z_2} \\ s_{z_2 z_1} & s_{z_2}^2 \end{pmatrix} \\ &= \begin{pmatrix} \mathbf{a}'_1 \mathbf{S} \mathbf{a}_1 & \mathbf{a}'_1 \mathbf{S} \mathbf{a}_2 \\ \mathbf{a}'_2 \mathbf{S} \mathbf{a}_1 & \mathbf{a}'_2 \mathbf{S} \mathbf{a}_2 \end{pmatrix}. \end{aligned} \quad (3.60)$$

By (2.50), this factors into

$$\mathbf{S}_z = \begin{pmatrix} \mathbf{a}'_1 \\ \mathbf{a}'_2 \end{pmatrix} \mathbf{S}(\mathbf{a}_1, \mathbf{a}_2) = \mathbf{A} \mathbf{S} \mathbf{A}' \quad (3.61)$$

The bivariate results in (3.59) and (3.61) can be readily extended to more than two linear combinations. (See principal components in Chapter 12, for instance, where we attempt to transform the y 's to a few dimensions that capture most of the information in the y 's.) If we have k linear transformations, they can be expressed as

$$\begin{aligned} z_1 &= a_{11}y_1 + a_{12}y_2 + \cdots + a_{1p}y_p = \mathbf{a}'_1 \mathbf{y} \\ z_2 &= a_{21}y_1 + a_{22}y_2 + \cdots + a_{2p}y_p = \mathbf{a}'_2 \mathbf{y} \\ &\vdots && \vdots \\ z_k &= a_{k1}y_1 + a_{k2}y_2 + \cdots + a_{kp}y_p = \mathbf{a}'_k \mathbf{y} \end{aligned}$$

or in matrix notation

$$\mathbf{z} = \begin{pmatrix} z_1 \\ z_2 \\ \vdots \\ z_k \end{pmatrix} = \begin{pmatrix} \mathbf{a}'_1 \mathbf{y} \\ \mathbf{a}'_2 \mathbf{y} \\ \vdots \\ \mathbf{a}'_k \mathbf{y} \end{pmatrix} = \begin{pmatrix} \mathbf{a}'_1 \\ \mathbf{a}'_2 \\ \vdots \\ \mathbf{a}'_k \end{pmatrix} \mathbf{y} = \mathbf{A}\mathbf{y} \quad [\text{by (2.47)}],$$

where \mathbf{z} is $k \times 1$, \mathbf{A} is $k \times p$, and \mathbf{y} is $p \times 1$ (we typically have $k \leq p$). If $\mathbf{z}_i = \mathbf{A}\mathbf{y}_i$ is evaluated for all $\mathbf{y}_i, i = 1, 2, \dots, n$, then by (3.54) and (2.49), the sample mean vector of the \mathbf{z} 's is

$$\bar{\mathbf{z}} = \begin{pmatrix} \mathbf{a}'_1 \bar{\mathbf{y}} \\ \mathbf{a}'_2 \bar{\mathbf{y}} \\ \vdots \\ \mathbf{a}'_k \bar{\mathbf{y}} \end{pmatrix} = \begin{pmatrix} \mathbf{a}'_1 \\ \mathbf{a}'_2 \\ \vdots \\ \mathbf{a}'_k \end{pmatrix} \bar{\mathbf{y}} = \mathbf{A}\bar{\mathbf{y}}. \quad (3.62)$$

By an extension of (3.60), the sample covariance matrix of the \mathbf{z} 's becomes

$$\begin{aligned} \mathbf{S}_z &= \begin{pmatrix} \mathbf{a}'_1 \mathbf{S} \mathbf{a}_1 & \mathbf{a}'_1 \mathbf{S} \mathbf{a}_2 & \cdots & \mathbf{a}'_1 \mathbf{S} \mathbf{a}_k \\ \mathbf{a}'_2 \mathbf{S} \mathbf{a}_1 & \mathbf{a}'_2 \mathbf{S} \mathbf{a}_2 & \cdots & \mathbf{a}'_2 \mathbf{S} \mathbf{a}_k \\ \vdots & \vdots & & \vdots \\ \mathbf{a}'_k \mathbf{S} \mathbf{a}_1 & \mathbf{a}'_k \mathbf{S} \mathbf{a}_2 & \cdots & \mathbf{a}'_k \mathbf{S} \mathbf{a}_k \end{pmatrix} & (3.63) \\ &= \begin{pmatrix} \mathbf{a}'_1 (\mathbf{S} \mathbf{a}_1, \mathbf{S} \mathbf{a}_2, \dots, \mathbf{S} \mathbf{a}_k) \\ \mathbf{a}'_2 (\mathbf{S} \mathbf{a}_1, \mathbf{S} \mathbf{a}_2, \dots, \mathbf{S} \mathbf{a}_k) \\ \vdots \\ \mathbf{a}'_k (\mathbf{S} \mathbf{a}_1, \mathbf{S} \mathbf{a}_2, \dots, \mathbf{S} \mathbf{a}_k) \end{pmatrix} \\ &= \begin{pmatrix} \mathbf{a}'_1 \\ \mathbf{a}'_2 \\ \vdots \\ \mathbf{a}'_k \end{pmatrix} (\mathbf{S} \mathbf{a}_1, \mathbf{S} \mathbf{a}_2, \dots, \mathbf{S} \mathbf{a}_k) \quad [\text{by (2.47)}] \\ &= \begin{pmatrix} \mathbf{a}'_1 \\ \mathbf{a}'_2 \\ \vdots \\ \mathbf{a}'_k \end{pmatrix} \mathbf{S} (\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_k) \quad [\text{by (2.48)}] \\ &= \mathbf{A} \mathbf{S} \mathbf{A}' . \end{aligned} \quad (3.64)$$

Note that by (3.63) and (3.64), we have

$$\text{tr}(\mathbf{A} \mathbf{S} \mathbf{A}') = \sum_{i=1}^k \mathbf{a}'_i \mathbf{S} \mathbf{a}_i . \quad (3.65)$$

A slightly more general linear transformation is

$$\mathbf{z}_i = \mathbf{A} \mathbf{y}_i + \mathbf{b}, \quad i = 1, 2, \dots, n. \quad (3.66)$$

The sample mean vector and covariance matrix of \mathbf{z} are given by

$$\bar{\mathbf{z}} = \mathbf{A} \bar{\mathbf{y}} + \mathbf{b}, \quad (3.67)$$

$$\mathbf{S}_z = \mathbf{A} \mathbf{S} \mathbf{A}' . \quad (3.68)$$

■ EXAMPLE 3.10.1

Timm (1975, p. 233; 1980, p. 47) reported the results of an experiment in which subjects responded to “probe words” at five positions in a sentence. The variables are response times for the j th probe word, $y_j, j = 1, 2, \dots, 5$. The data are given in Table 3.6.

These variables are commensurate (same measurement units and similar means and variances), and the researcher may wish to examine some simple

Table 3.6 Response Times for Five Probe Word Positions

Subject Number	y_1	y_2	y_3	y_4	y_5
1	51	36	50	35	42
2	27	20	26	17	27
3	37	22	41	37	30
4	42	36	32	34	27
5	27	18	33	14	29
6	43	32	43	35	40
7	41	22	36	25	38
8	38	21	31	20	16
9	36	23	27	25	28
10	26	31	31	32	36
11	29	20	25	26	25

linear combinations. Consider the following linear combination for illustrative purposes:

$$\begin{aligned} z &= 3y_1 - 2y_2 + 4y_3 - y_4 + y_5 \\ &= (3, -2, 4, -1, 1)\mathbf{y} = \mathbf{a}'\mathbf{y}. \end{aligned}$$

If z is calculated for each of the 11 observations, we obtain $z_1 = 288, z_2 = 155, \dots, z_{11} = 146$ with mean $\bar{z} = 197.0$ and variance $s_z^2 = 2084.0$. These same results can be obtained using (3.54) and (3.55). The sample mean vector and covariance matrix for the data are

$$\bar{\mathbf{y}} = \begin{pmatrix} 36.09 \\ 25.55 \\ 34.09 \\ 27.27 \\ 30.73 \end{pmatrix}, \quad \mathbf{S} = \begin{pmatrix} 65.09 & 33.65 & 47.59 & 36.77 & 25.43 \\ 33.65 & 46.07 & 28.95 & 40.34 & 28.36 \\ 47.59 & 28.95 & 60.69 & 37.37 & 41.13 \\ 36.77 & 40.34 & 37.37 & 62.82 & 31.68 \\ 25.43 & 28.36 & 41.13 & 31.68 & 58.22 \end{pmatrix}.$$

Then, by (3.54),

$$\bar{z} = \mathbf{a}'\bar{\mathbf{y}} = (3, -2, 4, -1, 1) \begin{pmatrix} 36.09 \\ 25.55 \\ 34.09 \\ 27.27 \\ 30.73 \end{pmatrix} = 197.0,$$

and by (3.55), $s_z^2 = \mathbf{a}'\mathbf{S}\mathbf{a} = 2084.0$.

We now define a second linear combination:

$$\begin{aligned} w &= y_1 + 3y_2 - y_3 + y_4 - 2y_5 \\ &= (1, 3, -1, 1, -2)\mathbf{y} = \mathbf{b}'\mathbf{y}. \end{aligned}$$

The sample mean and variance of w are $\bar{w} = \mathbf{b}'\bar{\mathbf{y}} = 44.45$ and $s_w^2 = \mathbf{b}'\mathbf{S}\mathbf{b} = 605.67$. The sample covariance of z and w is, by (3.56), $s_{zw} = \mathbf{a}'\mathbf{S}\mathbf{b} = 40.2$.

Using (3.57), we find the sample correlation between z and w to be

$$r_{zw} = \frac{s_{zw}}{\sqrt{s_z^2 s_w^2}} = \frac{40.2}{\sqrt{(2084)(605.67)}} = .0358.$$

We now define the three linear functions

$$\begin{aligned} z_1 &= y_1 + y_2 + y_3 + y_4 + y_5 \\ z_2 &= 2y_1 - 3y_2 + y_3 - 2y_4 - y_5 \\ z_3 &= -y_1 - 2y_2 + y_3 - 2y_4 + 3y_5, \end{aligned}$$

which can be written in matrix form as

$$\begin{pmatrix} z_1 \\ z_2 \\ z_3 \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 \\ 2 & -3 & 1 & -2 & -1 \\ -1 & -2 & 1 & -2 & 3 \end{pmatrix} \begin{pmatrix} y_1 \\ y_2 \\ y_3 \\ y_4 \\ y_5 \end{pmatrix}$$

or

$$\mathbf{z} = \mathbf{Ay}.$$

The sample mean vector for \mathbf{z} is given by (3.62) as

$$\bar{\mathbf{z}} = \mathbf{A}\bar{\mathbf{y}} = \begin{pmatrix} 153.73 \\ -55.64 \\ -15.45 \end{pmatrix},$$

and the sample covariance matrix of \mathbf{z} is given by (3.64) as

$$\mathbf{S}_z = \mathbf{ASA}' = \begin{pmatrix} 995.42 & -502.09 & -211.04 \\ -502.09 & 811.45 & 268.08 \\ -211.04 & 268.08 & 702.87 \end{pmatrix}.$$

The covariance matrix \mathbf{S}_z can be converted to a correlation matrix by use of (3.37):

$$\mathbf{R}_z = \mathbf{D}_z^{-1} \mathbf{S}_z \mathbf{D}_z^{-1} = \begin{pmatrix} 1.00 & -.56 & -.25 \\ -.56 & 1.00 & .35 \\ -.25 & .35 & 1.00 \end{pmatrix},$$

where

$$\mathbf{D}_z = \begin{pmatrix} 31.55 & 0 & 0 \\ 0 & 28.49 & 0 \\ 0 & 0 & 26.51 \end{pmatrix}$$

is obtained from the square roots of the diagonal elements of \mathbf{S}_z . □

3.10.2 Population Properties

The sample results in Section 3.10.1 for linear combinations have population counterparts. Let $z = \mathbf{a}'\mathbf{y}$, where \mathbf{a} is a vector of constants. Then the *population mean* of z is

$$E(z) = E(\mathbf{a}'\mathbf{y}) = \mathbf{a}'E(\mathbf{y}) = \mathbf{a}'\boldsymbol{\mu}, \quad (3.69)$$

and the *population variance* is

$$\sigma_z^2 = \text{var}(\mathbf{a}'\mathbf{y}) = \mathbf{a}'\Sigma\mathbf{a}. \quad (3.70)$$

Let $w = \mathbf{b}'\mathbf{y}$, where \mathbf{b} is a vector of constants different from \mathbf{a} . The *population covariance* of $z = \mathbf{a}'\mathbf{y}$ and $w = \mathbf{b}'\mathbf{y}$ is

$$\text{cov}(z, w) = \sigma_{zw} = \mathbf{a}'\Sigma\mathbf{b}. \quad (3.71)$$

By (3.12) the *population correlation* of z and w is

$$\begin{aligned} \rho_{zw} &= \text{corr}(\mathbf{a}'\mathbf{y}, \mathbf{b}'\mathbf{y}) = \frac{\sigma_{zw}}{\sigma_z \sigma_w} \\ &= \frac{\mathbf{a}'\Sigma\mathbf{b}}{\sqrt{(\mathbf{a}'\Sigma\mathbf{a})(\mathbf{b}'\Sigma\mathbf{b})}}. \end{aligned} \quad (3.72)$$

If $\mathbf{A}\mathbf{y}$ represents several linear combinations, the *population mean vector* and *covariance matrix* are given by

$$E(\mathbf{A}\mathbf{y}) = \mathbf{A}E(\mathbf{y}) = \mathbf{A}\boldsymbol{\mu}, \quad (3.73)$$

$$\text{cov}(\mathbf{A}\mathbf{y}) = \mathbf{A}\Sigma\mathbf{A}'. \quad (3.74)$$

The more general linear transformation $\mathbf{z} = \mathbf{A}\mathbf{y} + \mathbf{b}$ has population mean vector and covariance matrix

$$E(\mathbf{A}\mathbf{y} + \mathbf{b}) = \mathbf{A}E(\mathbf{y}) + \mathbf{b} = \mathbf{A}\boldsymbol{\mu} + \mathbf{b}, \quad (3.75)$$

$$\text{cov}(\mathbf{A}\mathbf{y} + \mathbf{b}) = \mathbf{A}\Sigma\mathbf{A}'. \quad (3.76)$$

3.11 MEASURES OF OVERALL VARIABILITY

The covariance matrix contains the variances of the p variables and the covariances between all pairs of variables and is thus a multifaceted picture of the overall variation in the data. Sometimes it is desirable to have a single numerical value for the overall multivariate scatter. One such measure is the *generalized sample variance*, defined as the determinant of the covariance matrix:

$$\text{Generalized sample variance} = |\mathbf{S}|. \quad (3.77)$$

The generalized sample variance has a geometric interpretation. The extension of an ellipse to more than two dimensions is called a *hyperellipsoid*. A p -dimensional hyperellipsoid $(\mathbf{y} - \bar{\mathbf{y}})' \mathbf{S}^{-1} (\mathbf{y} - \bar{\mathbf{y}}) = a^2$, centered at $\bar{\mathbf{y}}$ and based on \mathbf{S}^{-1} to standardize the distance to the center, contains a proportion of the observations $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n$ in the sample (if \mathbf{S} were replaced by Σ , the value of a^2 could be determined by tables of the chi-square distribution; see property 3 in Section 4.2). The ellipsoid $(\mathbf{y} - \bar{\mathbf{y}})' \mathbf{S}^{-1} (\mathbf{y} - \bar{\mathbf{y}})$ has axes proportional to the square roots of the eigenvalues of \mathbf{S} . It can be shown that the volume of the ellipsoid is proportional to $|\mathbf{S}|^{1/2}$. If the smallest eigenvalue λ_p is zero, there is no axis in that direction, and the ellipsoid lies wholly in a $(p - 1)$ -dimensional subspace of p -space. Consequently, the volume in p -space is zero. This can also be seen by (2.108), $|\mathbf{S}| = \lambda_1 \lambda_2 \cdots \lambda_p$. Hence, if $\lambda_p = 0$, $|\mathbf{S}| = 0$. A zero eigenvalue indicates a redundancy in the form of a linear relationship among the variables. (As will be seen in Section 12.7, the eigenvector corresponding to the zero eigenvalue reveals the form of the linear dependency.) One solution to the dilemma when $\lambda_p = 0$ is to remove one or more variables.

Another measure of overall variability, the *total sample variance*, is simply the trace of \mathbf{S} :

$$\text{Total sample variance} = s_{11} + s_{22} + \cdots + s_{pp} = \text{tr}(\mathbf{S}). \quad (3.78)$$

This measure of overall variation ignores covariance structure altogether but is found useful for comparison purposes in techniques such as principal components (Chapter 12).

In general, for both $|\mathbf{S}|$ and $\text{tr}(\mathbf{S})$, relatively large values reflect a broad scatter of $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_p$ about $\bar{\mathbf{y}}$, whereas lower values indicate closer concentration about $\bar{\mathbf{y}}$. In the case of $|\mathbf{S}|$, however, as noted above, an extremely small value of $|\mathbf{S}|$ or $|\mathbf{R}|$ may indicate either small scatter or *multicollinearity*, a term indicating near-linear relationships in a set of variables. Multicollinearity may be due to high pairwise correlations or to a high multiple correlation between one variable and several of the other variables. For other measures of intercorrelation, see Rencher (1998, Section 1.7).

3.12 ESTIMATION OF MISSING VALUES

It is not uncommon to find missing measurements in an observation vector, that is, missing values for one or more variables. A small number of rows with missing entries in the data matrix \mathbf{Y} [see (3.17)] does not constitute a serious problem; we can simply discard each row that has a missing value. However, with this procedure, a small portion of missing data, if widely distributed, would lead to a substantial loss of data. For example, in a large data set with $n = 550$ and $p = 85$, only about 1.5% of the $550 \times 85 = 46,750$ measurements were missing. However, nearly half of the observation vectors (rows of \mathbf{Y}) turned out to be incomplete.

The distribution of missing values in a data set is an important consideration. Randomly missing variable values scattered throughout a data matrix are less serious

than a pattern of missing values that depends to some extent on the values of the missing variables.

We discuss two methods of estimating the missing values or “filling the holes” in the data matrix, also called *imputation*. Both of the procedures presume that the missing values occur at random. If the occurrence or non-occurrence of missing values is related to the values of some of the variables, then the techniques may not estimate the missing responses very well.

The first method is very simple: Substitute a mean for each missing value, specifically the average of the available data in the column of the data matrix in which the unknown value lies. Replacing an observation by its mean reduces the variance and the absolute value of the covariance. Therefore, the sample covariance matrix S computed from the data matrix \mathbf{Y} in (3.17) with means imputed for missing values is biased. However, it is positive definite.

The second technique is a regression approach. The data matrix \mathbf{Y} is partitioned into two parts, one containing all rows with missing entries and the other comprising all the complete rows. Suppose y_{ij} is the only missing entry in the i th row of \mathbf{Y} . Then using the data in the submatrix with complete rows, y_j is regressed on the other variables to obtain a prediction equation $\hat{y}_j = b_0 + b_1 y_1 + \cdots + b_{j-1} y_{j-1} + b_{j+1} y_{j+1} + \cdots + b_p y_p$. Then the nonmissing entries in the i th row are entered as independent variables in the regression equation to obtain the predicted value, \hat{y}_{ij} . The regression method was first proposed by Buck (1960) and is a special case of the EM algorithm (Dempster, Laird, and Rubin 1977).

The regression method can be improved by iteration, carried out, for example, in the following way. Estimate all missing entries in the data matrix using regression. After filling in the missing entries, use the full data matrix to obtain new prediction equations. Use these prediction equations to calculate new predicted values \hat{y}_{ij} for missing entries. Use the new data matrix to obtain revised prediction equations and new predicted values \hat{y}_{ij} . Continue this process until the predicted values stabilize.

A modification may be needed if the missing entries are so pervasive that it is difficult to find data to estimate the initial regression equations. In this case, the process could be started by using means as in the first method and then beginning the iteration.

The regression approach will ordinarily yield better results than the method of inserting means. However, if the other variables are not very highly correlated with the one to be predicted, the regression technique is essentially equivalent to imputing means. The regression method underestimates the variances and covariances, though to a lesser extent than the method based on means.

■ EXAMPLE 3.12

We illustrate the iterated regression method of estimating missing values. Consider the calcium data of Table 3.4 as reproduced here and suppose the entries in parentheses are missing:

Location Number	y_1	y_2	y_3
1	35	(3.5)	2.80
2	35	4.9	(2.70)
3	40	30.0	4.38
4	10	2.8	3.21
5	6	2.7	2.73
6	20	2.8	2.81
7	35	4.6	2.88
8	35	10.9	2.90
9	35	8.0	3.28
10	30	1.6	3.20

We first regress y_2 on y_1 and y_3 for observations 3–10 and obtain $\hat{y}_2 = b_0 + b_1 y_1 + b_3 y_3$. When this is evaluated for the two nonmissing entries in the first row ($y_1 = 35$ and $y_3 = 2.80$), we obtain $\hat{y}_2 = 4.097$. Similarly, we regress y_3 on y_1 and y_2 for observations 3–10 to obtain $\hat{y}_3 = c_0 + c_1 y_1 + c_2 y_2$. Evaluating this for the two nonmissing entries in the second row yields $\hat{y}_3 = 3.011$. We now insert these estimates for the missing values and calculate the regression equations based on all 10 observations. Using the revised equation $\hat{y}_2 = b_0 + b_1 y_1 + b_3 y_3$, we obtain a new predicted value, $\hat{y}_2 = 3.698$. Similarly, we obtain a revised regression equation for y_3 that gives a new predicted value, $\hat{y}_3 = 2.981$. With these values inserted, we calculate new equations and obtain new predicted values, $\hat{y}_2 = 3.672$ and $\hat{y}_3 = 2.976$. At the third iteration we obtain $\hat{y}_2 = 3.679$ and $\hat{y}_3 = 2.975$. There is very little change in subsequent iterations. These values are closer to the actual values, $y_2 = 3.5$ and $y_3 = 2.70$, than the initial regression estimates, $\hat{y}_2 = 4.097$ and $\hat{y}_3 = 3.011$. They are also much better estimates than the means of the second and third columns, $\bar{y}_2 = 7.589$ and $\bar{y}_3 = 3.132$. \square

3.13 DISTANCE BETWEEN VECTORS

In a univariate setting, the distance between two points is simply the difference (or absolute difference) between their values. For statistical purposes, this difference may not be very informative. For example, we do not want to know how many centimeters apart two means are, but rather how many standard deviations apart they are. Thus we examine the standardized or statistical distances, such as

$$\frac{|\mu_1 - \mu_2|}{\sigma} \quad \text{or} \quad \frac{|\bar{y} - \mu|}{\sigma_{\bar{y}}}.$$

To obtain a useful distance measure in a multivariate setting, we must consider not only the variances of the variables but also their covariances or correlations. The simple (squared) Euclidean distance between two vectors, $(y_1 - y_2)'(y_1 - y_2)$, is not useful in some situations because there is no adjustment for the variances or the

covariances. For a statistical distance, we standardize by inserting the inverse of the covariance matrix:

$$d^2 = (\mathbf{y}_1 - \mathbf{y}_2)' \mathbf{S}^{-1} (\mathbf{y}_1 - \mathbf{y}_2). \quad (3.79)$$

Other examples are

$$D^2 = (\bar{\mathbf{y}} - \boldsymbol{\mu})' \mathbf{S}^{-1} (\bar{\mathbf{y}} - \boldsymbol{\mu}) \quad (3.80)$$

$$\Delta^2 = (\bar{\mathbf{y}} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1} (\bar{\mathbf{y}} - \boldsymbol{\mu}) \quad (3.81)$$

$$\Delta^2 = (\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2)' \boldsymbol{\Sigma}^{-1} (\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2). \quad (3.82)$$

These (squared) distances between two vectors were first proposed by Mahalanobis (1936) and are often referred to as *Mahalanobis distances*. If a random variable has a larger variance than another, it receives relatively less weight in a Mahalanobis distance. Similarly, two highly correlated variables do not contribute as much as two variables that are less correlated. In essence, then, the use of the inverse of the covariance matrix in a Mahalanobis distance has the effect of (1) standardizing all variables to the same variance and (2) eliminating correlations. To illustrate this, we use the square root matrix defined in (2.112) to rewrite (3.81) as

$$\begin{aligned} \Delta^2 &= (\bar{\mathbf{y}} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1} (\bar{\mathbf{y}} - \boldsymbol{\mu}) = (\bar{\mathbf{y}} - \boldsymbol{\mu})' (\boldsymbol{\Sigma}^{1/2} \boldsymbol{\Sigma}^{1/2})^{-1} (\bar{\mathbf{y}} - \boldsymbol{\mu}) \\ &= [(\boldsymbol{\Sigma}^{1/2})^{-1} (\bar{\mathbf{y}} - \boldsymbol{\mu})]' [(\boldsymbol{\Sigma}^{1/2})^{-1} (\bar{\mathbf{y}} - \boldsymbol{\mu})] = \mathbf{z}' \mathbf{z}, \end{aligned}$$

where $\mathbf{z} = (\boldsymbol{\Sigma}^{1/2})^{-1} (\bar{\mathbf{y}} - \boldsymbol{\mu}) = (\boldsymbol{\Sigma}^{1/2})^{-1} \bar{\mathbf{y}} - (\boldsymbol{\Sigma}^{1/2})^{-1} \boldsymbol{\mu}$. Now, by (3.76) it can be shown that

$$\text{cov}(\mathbf{z}) = \frac{1}{n} \mathbf{I}. \quad (3.83)$$

Hence the transformed variables z_1, z_2, \dots, z_p are uncorrelated, and each has variance equal to $1/n$. If the appropriate covariance matrix for the random vector were used in a Mahalanobis distance, the variances would reduce to 1. For example, if $\text{cov}(\bar{\mathbf{y}}) = \boldsymbol{\Sigma}/n$ were used above in place of $\boldsymbol{\Sigma}$, we would obtain $\text{cov}(\mathbf{z}) = \mathbf{I}$.

PROBLEMS

- 3.1** If $z_i = ay_i$ for $i = 1, 2, \dots, n$, show that $\bar{z} = a\bar{y}$ as in (3.3).
- 3.2** If $z_i = ay_i$ for $i = 1, 2, \dots, n$, show that $s_z^2 = a^2 s^2$ as in (3.6).
- 3.3** For the data in Figure 3.3, show that $\sum_i (x_i - \bar{x})(y_i - \bar{y}) = 0$.
- 3.4** Show that $(\mathbf{x} - \bar{x}\mathbf{j})' (\mathbf{y} - \bar{y}\mathbf{j}) = \sum_i (x_i - \bar{x})(y_i - \bar{y})$, thus verifying (3.15).
- 3.5** For $p = 3$ show that

$$\frac{1}{n-1} \sum_{i=1}^n (\mathbf{y}_i - \bar{\mathbf{y}})(\mathbf{y}_i - \bar{\mathbf{y}})' = \begin{pmatrix} s_{11} & s_{12} & s_{13} \\ s_{21} & s_{22} & s_{23} \\ s_{31} & s_{32} & s_{33} \end{pmatrix},$$

which illustrates (3.27).

- 3.6** Show that $\bar{z} = \mathbf{a}'\bar{\mathbf{y}}$ as in (3.54), where $z_i = \mathbf{a}'\mathbf{y}_i, i = 1, 2, \dots, n$.
- 3.7** Show that $s_z^2 = \mathbf{a}'\mathbf{S}\mathbf{a}$ as in (3.55), where $z_i = \mathbf{a}'\mathbf{y}_i, i = 1, 2, \dots, n$.
- 3.8** Show that $\text{tr}(\mathbf{A}\mathbf{S}\mathbf{A}') = \sum_{i=1}^k \mathbf{a}'_i \mathbf{S} \mathbf{a}_i$ as in (3.65).
- 3.9** Use (3.76) to verify (3.83), $\text{cov}(\mathbf{z}) = \mathbf{I}/n$, where $\mathbf{z} = (\boldsymbol{\Sigma}^{1/2})^{-1}(\bar{\mathbf{y}} - \boldsymbol{\mu})$.
- 3.10** Use the calcium data in Table 3.4:
- Calculate \mathbf{S} using the data matrix \mathbf{Y} as in (3.29).
 - Obtain \mathbf{R} by calculating r_{12}, r_{13} , and r_{23} , as in (3.34) and (3.35).
 - Find \mathbf{R} using (3.37).
- 3.11** Use the calcium data in Table 3.4:
- Find the generalized sample variance $|\mathbf{S}|$ as in (3.77).
 - Find the total sample variance $\text{tr}(\mathbf{S})$ as in (3.78).
- 3.12** Use the probe word data of Table 3.6:
- Find the generalized sample variance $|\mathbf{S}|$ as in (3.77).
 - Find the total sample variance $\text{tr}(\mathbf{S})$ as in (3.78).
- 3.13** For the probe word data in Table 3.6, find \mathbf{R} using (3.37).
- 3.14** For the variables in Table 3.4, define $z = 3y_1 - y_2 + 2y_3 = (3, -1, 2)\mathbf{y}$. Find \bar{z} and s_z^2 in two ways:
- Evaluate z for each row of Table 3.4 and find \bar{z} and s_z^2 directly from z_1, z_2, \dots, z_{10} using (3.1) and (3.5).
 - Use $\bar{z} = \mathbf{a}'\bar{\mathbf{y}}$ and $s_z^2 = \mathbf{a}'\mathbf{S}\mathbf{a}$, as in (3.54) and (3.55).
- 3.15** For the variables in Table 3.4, define $w = -2y_1 + 3y_2 + y_3$ and define z as in Problem 3.14. Find r_{zw} in two ways:
- Evaluate z and w for each row of Table 3.4 and find r_{zw} from the 10 pairs $(z_i, w_i), i = 1, 2, \dots, 10$, using (3.10) and (3.13).
 - Find r_{zw} using (3.57).
- 3.16** For the variables in Table 3.4, find the correlation between y_1 and $\frac{1}{2}(y_2 + y_3)$ using (3.57).

Table 3.7 Ramus Bone Length at Four Ages for 20 Boys

Individual	Age			
	8 yr (y_1)	$8\frac{1}{2}$ yr (y_2)	9 yr (y_3)	$9\frac{1}{2}$ yr (y_4)
1	47.8	48.8	49.0	49.7
2	46.4	47.3	47.7	48.4
3	46.3	46.8	47.8	48.5
4	45.1	45.3	46.1	47.2
5	47.6	48.5	48.9	49.3
6	52.5	53.2	53.3	53.7
7	51.2	53.0	54.3	54.5
8	49.8	50.0	50.3	52.7
9	48.1	50.8	52.3	54.4
10	45.0	47.0	47.3	48.3
11	51.2	51.4	51.6	51.9
12	48.5	49.2	53.0	55.5
13	52.1	52.8	53.7	55.0
14	48.2	48.9	49.3	49.8
15	49.6	50.4	51.2	51.8
16	50.7	51.7	52.7	53.3
17	47.2	47.7	48.4	49.5
18	53.3	54.6	55.1	55.3
19	46.2	47.5	48.1	48.4
20	46.3	47.6	51.3	51.8

3.17 Define the following linear combinations for the variables in Table 3.4:

$$\begin{aligned}z_1 &= y_1 + y_2 + y_3 \\z_2 &= 2y_1 - 3y_2 + 2y_3 \\z_3 &= -y_1 - 2y_2 - 3y_3\end{aligned}$$

- (a) Find \bar{z} and \mathbf{S}_z using (3.62) and (3.64).
- (b) Find \mathbf{R}_z from \mathbf{S}_z using (3.37).

3.18 The data in Table 3.7 (Elston and Grizzle 1962) consist of measurements y_1, y_2, y_3 , and y_4 of the ramus bone at four different ages on each of 20 boys.

- (a) Find $\bar{\mathbf{y}}$, \mathbf{S} , and \mathbf{R} .
- (b) Find $|\mathbf{S}|$ and $\text{tr}(\mathbf{S})$.

3.19 For the data in Table 3.7, define $z = y_1 + 2y_2 + y_3 - 3y_4$ and $w = -2y_1 + 3y_2 - y_3 + 2y_4$.

- (a) Find \bar{z}, \bar{w}, s_z^2 , and s_w^2 using (3.54) and (3.55).

Table 3.8 Measurements on the First and Second Adult Sons in a Sample of 25 Families

First Son		Second Son	
Head Length, y_1	Head Breadth, y_2	Head Length, x_1	Head Breadth, x_2
191	155	179	145
195	149	201	152
181	148	185	149
183	153	188	149
176	144	171	142
208	157	192	152
189	150	190	149
197	159	189	152
188	152	197	159
192	150	187	151
179	158	186	148
183	147	174	147
174	150	185	152
190	159	195	157
188	151	187	158
163	137	161	130
195	155	183	158
186	153	173	148
181	145	182	146
175	140	165	137
192	154	185	152
174	143	178	147
176	139	176	143
197	167	200	158
190	163	187	150

(b) Find s_{zw} and r_{zw} using (3.56) and (3.57).

3.20 For the data in Table 3.7 define

$$\begin{aligned}z_1 &= 2y_1 + 3y_2 - y_3 + 4y_4 \\z_2 &= -2y_1 - y_2 + 4y_3 - 2y_4 \\z_3 &= 3y_1 - 2y_2 - y_3 + 3y_4\end{aligned}$$

Find $\bar{\mathbf{z}}$, \mathbf{S}_z , and \mathbf{R}_z , using (3.62), (3.64), and (3.37), respectively.

3.21 The data in Table 3.8 consist of head measurements on first and second sons (Frets 1921). Define y_1 and y_2 as the measurements on the first son and x_1 and x_2 for the second son.

- (a) Find the mean vector for all four variables and partition it into $\begin{pmatrix} \bar{\mathbf{y}} \\ \bar{\mathbf{x}} \end{pmatrix}$ as in (3.41).
- (b) Find the covariance matrix for all four variables and partition it into

$$\mathbf{S} = \begin{pmatrix} \mathbf{S}_{yy} & \mathbf{S}_{yx} \\ \mathbf{S}_{xy} & \mathbf{S}_{xx} \end{pmatrix}$$

as in (3.42).

- 3.22** Table 3.9 contains data from O'Sullivan and Mahan (1966; see also Andrews and Herzberg 1985, p. 214) with measurements of blood glucose levels on three occasions for 50 women. The y 's represent fasting glucose measurements on the three occasions; the x 's are glucose measurements 1 hour after sugar intake. Find the mean vector and covariance matrix for all six variables and partition them into $\begin{pmatrix} \bar{\mathbf{y}} \\ \bar{\mathbf{x}} \end{pmatrix}$, as in (3.41), and

$$\mathbf{S} = \begin{pmatrix} \mathbf{S}_{yy} & \mathbf{S}_{yx} \\ \mathbf{S}_{xy} & \mathbf{S}_{xx} \end{pmatrix}$$

as in (3.42).

Table 3.9 Blood Glucose Measurements on Three Occasions

Fasting			One Hour After Sugar Intake		
y_1	y_2	y_3	x_1	x_2	x_3
60	69	62	97	69	98
56	53	84	103	78	107
80	69	76	66	99	130
55	80	90	80	85	114
62	75	68	116	130	91
74	64	70	109	101	103
64	71	66	77	102	130
73	70	64	115	110	109
68	67	75	76	85	119
69	82	74	72	133	127
60	67	61	130	134	121
70	74	78	150	158	100
66	74	78	150	131	142
83	70	74	99	98	105
68	66	90	119	85	109
78	63	75	164	98	138
103	77	77	160	117	121
77	68	74	144	71	153
66	77	68	77	82	89
70	70	72	114	93	122
75	65	71	77	70	109
91	74	93	118	115	150
66	75	73	170	147	121
75	82	76	153	132	115
74	71	66	143	105	100
76	70	64	114	113	129
74	90	86	73	106	116
74	77	80	116	81	77
67	71	69	63	87	70
67	71	69	63	87	70
78	75	80	105	132	80
64	66	71	83	94	133
71	80	76	81	87	86
63	75	73	120	89	59
:	:	:	:	:	:
52	70	76	92	94	100

Note: Measurements are in mg/100 ml.

CHAPTER 4

THE MULTIVARIATE NORMAL DISTRIBUTION

4.1 MULTIVARIATE NORMAL DENSITY FUNCTION

Many univariate tests and confidence intervals are based on the univariate normal distribution. Similarly, the majority of multivariate procedures have the multivariate normal distribution as their underpinning.

The following are some of the useful features of the multivariate normal distribution (see Section 4.2): (1) the distribution can be completely described using only means, variances, and covariances; (2) bivariate plots of multivariate data show linear trends; (3) if the variables are uncorrelated, they are independent; (4) linear functions of multivariate normal variables are also normal; (5) as in the univariate case, the convenient form of the density function lends itself to derivation of many properties and test statistics; and (6) even when the data are not multivariate normal, the multivariate normal may serve as a useful approximation, especially in inferences involving sample mean vectors, which are approximately multivariate normal by the central limit theorem (see Section 4.3.2).

Since the multivariate normal density is an extension of the univariate normal density and shares many of its features, we review the univariate normal density function




Figure 4.1 The normal density curve.

in Section 4.1.1. We then describe the multivariate normal density in Sections 4.1.2–4.1.4.

4.1.1 Univariate Normal Density

If a random variable y , with mean μ and variance σ^2 , is normally distributed, its density is given by

$$f(y) = \frac{1}{\sqrt{2\pi}\sqrt{\sigma^2}} e^{-(y-\mu)^2/2\sigma^2}, \quad -\infty < y < \infty. \quad (4.1)$$

When y has the density (4.1), we say that y is distributed as $N(\mu, \sigma^2)$, or simply y is $N(\mu, \sigma^2)$. This function is represented by the familiar bell-shaped curve illustrated in Figure 4.1 for $\mu = 10$ and $\sigma = 2.5$.

4.1.2 Multivariate Normal Density

If \mathbf{y} has a multivariate normal distribution with mean vector $\boldsymbol{\mu}$ and covariance matrix $\boldsymbol{\Sigma}$, the density is given by

$$g(\mathbf{y}) = \frac{1}{(\sqrt{2\pi})^p |\boldsymbol{\Sigma}|^{1/2}} e^{-(\mathbf{y}-\boldsymbol{\mu})'\boldsymbol{\Sigma}^{-1}(\mathbf{y}-\boldsymbol{\mu})/2}, \quad (4.2)$$

where p is the number of variables. When \mathbf{y} has the density (4.2), we say that \mathbf{y} is distributed as $N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, or simply \mathbf{y} is $N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$.

The term $(y - \mu)^2/\sigma^2 = (y - \mu)(\sigma^2)^{-1}(y - \mu)$ in the exponent of the univariate normal density (4.1) measures the squared distance from y to μ in standard deviation units. Similarly, the term $(y - \mu)' \Sigma^{-1} (y - \mu)$ in the exponent of the multivariate normal density (4.2) is the squared generalized distance from y to μ , or the Mahalanobis distance,

$$\Delta^2 = (y - \mu)' \Sigma^{-1} (y - \mu). \quad (4.3)$$

The characteristics of this distance between y and μ were discussed in Section 3.13. Note that Δ , the square root of (4.3), is not in standard deviation units as is $(y - \mu)/\sigma$. The distance Δ increases with p , the number of variables (see Problem 4.4).

In the coefficient of the exponential function in (4.2), $|\Sigma|^{1/2}$ appears as the analogue of $\sqrt{\sigma^2}$ in (4.1). In the next section, we discuss the effect of $|\Sigma|$ on the density.

4.1.3 Generalized Population Variance

In Section 3.11, we referred to $|S|$ as a generalized sample variance. Analogously, $|\Sigma|$ is a *generalized population variance*. If σ^2 is small in the univariate normal, the y values are concentrated near the mean. Similarly, a small value of $|\Sigma|$ in the multivariate case indicates that the y 's are concentrated close to μ in p -space or that there is multicollinearity among the variables. The term *multicollinearity* indicates that the variables are highly intercorrelated, in which case the effective dimensionality is less than p . (See Chapter 12 for a method of finding a reduced number of new dimensions that represent the data.) In the presence of multicollinearity, one or more eigenvalues of Σ will be near zero and $|\Sigma|$ will be small, since $|\Sigma|$ is the product of the eigenvalues [see (2.108)].

Figure 4.2 shows, for the bivariate case, a comparison of a distribution with small $|\Sigma|$ and a distribution with larger $|\Sigma|$. An alternative way to portray the concentration of points in the bivariate normal distribution is with contour plots. Figure 4.3 shows contour plots for the two distributions in Figure 4.2. Each ellipse contains a different proportion of observation vectors y . The contours in Figure 4.3 can be found by setting the density function equal to a constant and solving for y , as illustrated in Figure 4.4. The bivariate normal density surface sliced at a constant height traces an ellipse, which contains a given proportion of the observations [see Rencher (1998, Section 2.1.3)].

In both Figures 4.2 and 4.3, small $|\Sigma|$ appears on the left and large $|\Sigma|$ appears on the right. In Figure 4.3(a), there is a larger correlation between y_1 and y_2 . In Figure 4.3(b), the variances are larger (in the natural directions). In general, for any number of variables p , a decrease in intercorrelations among the variables or an increase in the variances will lead to a larger $|\Sigma|$.

4.1.4 Diversity of Applications of the Multivariate Normal

Nearly all the inferential procedures we discuss in this book are based on the multivariate normal distribution. We acknowledge that a major motivation for the widespread use of the multivariate normal is its mathematical tractability. From the mul-




Figure 4.2 Bivariate normal densities.




Figure 4.3 Contour plots for the distributions in Figure 4.2.

tivariate normal assumption, a host of useful procedures can be derived, and many of these are available in software packages. Practical alternatives to the multivariate normal are fewer than in the univariate case. Because it is not as simple to order (or rank) multivariate observation vectors as it is univariate observations, there are not as many nonparametric procedures available for multivariate data.

Although real data may not often be exactly multivariate normal, the multivariate normal will frequently serve as a useful approximation to the true distribution. Tests and graphical procedures are available for assessing normality (see Sections 4.4 and 4.6). Fortunately, many of the procedures based on multivariate normality are robust to departures from normality.

4.2 PROPERTIES OF MULTIVARIATE NORMAL RANDOM VARIABLES

We list some of the properties of a random $p \times 1$ vector \mathbf{y} from a multivariate normal distribution $N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$:

1. Normality of linear combinations of the variables in \mathbf{y}




Figure 4.4 Constant density contour for bivariate normal.

- a. If \mathbf{a} is a vector of constants, the linear function $\mathbf{a}'\mathbf{y} = a_1y_1 + a_2y_2 + \dots + a_py_p$ is univariate normal:

If \mathbf{y} is $N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, then $\mathbf{a}'\mathbf{y}$ is $N(\mathbf{a}'\boldsymbol{\mu}, \mathbf{a}'\boldsymbol{\Sigma}\mathbf{a})$.

The mean and variance of $\mathbf{a}'\mathbf{y}$ were given previously in (3.69) and (3.70) as $E(\mathbf{a}'\mathbf{y}) = \mathbf{a}'\boldsymbol{\mu}$ and $\text{var}(\mathbf{a}'\mathbf{y}) = \mathbf{a}'\boldsymbol{\Sigma}\mathbf{a}$ for any random vector \mathbf{y} . We now have the additional attribute that $\mathbf{a}'\mathbf{y}$ has a (univariate) normal distribution if \mathbf{y} is $N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$.

- b. If \mathbf{A} is a constant $q \times p$ matrix of rank q , where $q \leq p$, the q linear combinations in \mathbf{Ay} have a multivariate normal distribution:

If \mathbf{y} is $N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, then \mathbf{Ay} is $N_q(\mathbf{A}\boldsymbol{\mu}, \mathbf{A}\boldsymbol{\Sigma}\mathbf{A}')$.

Here, again, $E(\mathbf{Ay}) = \mathbf{A}\boldsymbol{\mu}$ and $\text{cov}(\mathbf{Ay}) = \mathbf{A}\boldsymbol{\Sigma}\mathbf{A}'$, in general, as given in (3.73) and (3.74). But we now have the additional feature that the q variables in \mathbf{Ay} have a multivariate normal distribution.

2. Standardized variables

A *standardized vector* \mathbf{z} can be obtained in two ways:

$$\mathbf{z} = (\mathbf{T}')^{-1}(\mathbf{y} - \boldsymbol{\mu}), \quad (4.4)$$

where $\Sigma = \mathbf{T}'\mathbf{T}$ is factored using the Cholesky procedure in Section 2.7, or

$$\mathbf{z} = (\Sigma^{1/2})^{-1}(\mathbf{y} - \boldsymbol{\mu}), \quad (4.5)$$

where $\Sigma^{1/2}$ is the symmetric square root matrix of Σ defined in (2.112) such that $\Sigma = \Sigma^{1/2}\Sigma^{1/2}$. In either (4.4) or (4.5), the standardized vector of random variables has all means equal to 0, all variances equal to 1, and all correlations equal to 0. In either case, it follows from property 1b that \mathbf{z} is multivariate normal:

$$\text{If } \mathbf{y} \text{ is } N_p(\boldsymbol{\mu}, \Sigma), \text{ then } \mathbf{z} \text{ is } N_p(\mathbf{0}, \mathbf{I}),$$

3. Chi-square distribution

A *chi-square random variable* with p degrees of freedom is defined as the sum of squares of p independent standard normal random variables. Thus if \mathbf{z} is the standardized vector defined in (4.4) or (4.5), then $\sum_{j=1}^p z_j^2 = \mathbf{z}'\mathbf{z}$ has the χ^2 -distribution with p degrees of freedom, denoted as χ_p^2 or $\chi^2(p)$. From either (4.4) or (4.5) we obtain $\mathbf{z}'\mathbf{z} = (\mathbf{y} - \boldsymbol{\mu})'\Sigma^{-1}(\mathbf{y} - \boldsymbol{\mu})$. Hence,

$$\text{If } \mathbf{y} \text{ is } N_p(\boldsymbol{\mu}, \Sigma), \text{ then } (\mathbf{y} - \boldsymbol{\mu})'\Sigma^{-1}(\mathbf{y} - \boldsymbol{\mu}) \text{ is } \chi_p^2. \quad (4.6)$$

4. Normality of marginal distributions

- a. Any subset of the y 's in \mathbf{y} has a multivariate normal distribution, with mean vector consisting of the corresponding subvector of $\boldsymbol{\mu}$ and covariance matrix composed of the corresponding submatrix of Σ . To illustrate, let $\mathbf{y}_1 = (y_1, y_2, \dots, y_r)'$ denote the subvector containing the first r elements of \mathbf{y} and $\mathbf{y}_2 = (y_{r+1}, \dots, y_p)'$ consist of the remaining $p - r$ elements. Thus \mathbf{y} , $\boldsymbol{\mu}$, and Σ are partitioned as

$$\mathbf{y} = \begin{pmatrix} \mathbf{y}_1 \\ \mathbf{y}_2 \end{pmatrix}, \quad \boldsymbol{\mu} = \begin{pmatrix} \boldsymbol{\mu}_1 \\ \boldsymbol{\mu}_2 \end{pmatrix}, \quad \Sigma = \begin{pmatrix} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{21} & \Sigma_{22} \end{pmatrix},$$

where \mathbf{y}_1 and $\boldsymbol{\mu}_1$ are $r \times 1$ and Σ_{11} is $r \times r$. Then \mathbf{y}_1 is multivariate normal:

$$\text{If } \mathbf{y} \text{ is } N_p(\boldsymbol{\mu}, \Sigma), \text{ then } \mathbf{y}_1 \text{ is } N_r(\boldsymbol{\mu}_1, \Sigma_{11}).$$

Here, again, $E(\mathbf{y}_1) = \boldsymbol{\mu}_1$ and $\text{cov}(\mathbf{y}_1) = \Sigma_{11}$ hold for any random vector partitioned in this way. But if \mathbf{y} is p -variate normal, then \mathbf{y}_1 is r -variate normal.

- b. As a special case of the preceding result, each y_j in \mathbf{y} has the univariate normal distribution:

$$\text{If } \mathbf{y} \text{ is } N_p(\boldsymbol{\mu}, \Sigma), \text{ then } y_j \text{ is } N(\mu_j, \sigma_{jj}), j = 1, 2, \dots, p.$$

The converse of this is not true. If the density of each y_j in \mathbf{y} is normal, it does not necessarily follow that \mathbf{y} is multivariate normal.

In the next three properties, let the observation vector be partitioned into two subvectors denoted by \mathbf{y} and \mathbf{x} , where \mathbf{y} is $p \times 1$ and \mathbf{x} is $q \times 1$. Or, alternatively, let \mathbf{x} represent some additional variables to be considered along with those in \mathbf{y} . Then, as in (3.45) and (3.46),

$$E\begin{pmatrix} \mathbf{y} \\ \mathbf{x} \end{pmatrix} = \begin{pmatrix} \boldsymbol{\mu}_y \\ \boldsymbol{\mu}_x \end{pmatrix}, \quad \text{cov}\begin{pmatrix} \mathbf{y} \\ \mathbf{x} \end{pmatrix} = \begin{pmatrix} \boldsymbol{\Sigma}_{yy} & \boldsymbol{\Sigma}_{yx} \\ \boldsymbol{\Sigma}_{xy} & \boldsymbol{\Sigma}_{xx} \end{pmatrix}.$$

In properties 5, 6, and 7, we assume that

$$\begin{pmatrix} \mathbf{y} \\ \mathbf{x} \end{pmatrix} \text{ is } N_{p+q} \left[\begin{pmatrix} \boldsymbol{\mu}_x \\ \boldsymbol{\mu}_y \end{pmatrix}, \begin{pmatrix} \boldsymbol{\Sigma}_{yy} & \boldsymbol{\Sigma}_{yx} \\ \boldsymbol{\Sigma}_{xy} & \boldsymbol{\Sigma}_{xx} \end{pmatrix} \right].$$

5. Independence

- a. The subvectors \mathbf{y} and \mathbf{x} are independent if $\boldsymbol{\Sigma}_{yx} = \mathbf{O}$.
- b. Two individual variables y_j and y_k are independent if $\sigma_{jk} = 0$. Note that this is not true for many nonnormal random variables, as illustrated in Section 3.2.1.

6. Conditional distribution

If \mathbf{y} and \mathbf{x} are not independent, then $\boldsymbol{\Sigma}_{yx} \neq \mathbf{O}$, and the conditional distribution of \mathbf{y} given \mathbf{x} , $f(\mathbf{y}|\mathbf{x})$, is multivariate normal with

$$E(\mathbf{y}|\mathbf{x}) = \boldsymbol{\mu}_y + \boldsymbol{\Sigma}_{yx} \boldsymbol{\Sigma}_{xx}^{-1} (\mathbf{x} - \boldsymbol{\mu}_x), \quad (4.7)$$

$$\text{cov}(\mathbf{y}|\mathbf{x}) = \boldsymbol{\Sigma}_{yy} - \boldsymbol{\Sigma}_{yx} \boldsymbol{\Sigma}_{xx}^{-1} \boldsymbol{\Sigma}_{xy}. \quad (4.8)$$

Note that $E(\mathbf{y}|\mathbf{x})$ is a vector of linear functions of \mathbf{x} , while $\text{cov}(\mathbf{y}|\mathbf{x})$ does not depend on \mathbf{x} . The linear trend in (4.7) holds for any pair of variables. Thus to use (4.7) as a check on normality, one can examine bivariate scatter-plots of all pairs of variables and look for any nonlinear trends. In (4.7), we have the justification for using the covariance or correlation to measure the relationship between two bivariate normal random variables. As noted in Section 3.2.1, the covariance and correlation are good measures of relationship only for variables with linear trends and are generally unsuitable for nonnormal random variables with a curvilinear relationship. The matrix $\boldsymbol{\Sigma}_{yx} \boldsymbol{\Sigma}_{xx}^{-1}$ in (4.7) is called the *matrix of regression coefficients* because it relates $E(\mathbf{y}|\mathbf{x})$ to \mathbf{x} . The sample counterpart of this matrix appears in (10.57).

7. Distribution of the sum of two subvectors

If \mathbf{y} and \mathbf{x} are the same size (both $p \times 1$) and independent, then

$$\mathbf{y} + \mathbf{x} \text{ is } N_p(\boldsymbol{\mu}_y + \boldsymbol{\mu}_x, \boldsymbol{\Sigma}_{yy} + \boldsymbol{\Sigma}_{xx}), \quad (4.9)$$

$$\mathbf{y} - \mathbf{x} \text{ is } N_p(\boldsymbol{\mu}_y - \boldsymbol{\mu}_x, \boldsymbol{\Sigma}_{yy} + \boldsymbol{\Sigma}_{xx}). \quad (4.10)$$

In the remainder of this section, we illustrate property 6 for the special case of the bivariate normal. Let

$$\mathbf{u} = \begin{pmatrix} y \\ x \end{pmatrix}$$

have a bivariate normal distribution with

$$E(\mathbf{u}) = \begin{pmatrix} \mu_y \\ \mu_x \end{pmatrix}, \quad \text{cov}(\mathbf{u}) = \boldsymbol{\Sigma} = \begin{pmatrix} \sigma_y^2 & \sigma_{yx} \\ \sigma_{yx} & \sigma_x^2 \end{pmatrix}.$$

By definition $f(y|x) = g(y,x)/h(x)$, where $g(y,x)$ is the joint density of y and x and $h(x)$ is the density of x . Hence

$$g(y,x) = f(y|x)h(x),$$

and because the right side is a product, we seek a function of y and x that is independent of x and whose density can serve as $f(y|x)$. Since linear functions of y and x are normal by property 1a, we consider $y - \beta x$ and seek the value of β so that $y - \beta x$ and x are independent.

Since $z = y - \beta x$ and x are normal and independent, $\text{cov}(x,z) = 0$. To find $\text{cov}(x,z)$, we express x and z as functions of \mathbf{u} ,

$$\begin{aligned} x &= (0, 1) \begin{pmatrix} y \\ x \end{pmatrix} = (0, 1)\mathbf{u} = \mathbf{a}'\mathbf{u}, \\ z &= y - \beta x = (1, -\beta)\mathbf{u} = \mathbf{b}'\mathbf{u}. \end{aligned}$$

Now

$$\begin{aligned} \text{cov}(x, z) &= \text{cov}(\mathbf{a}'\mathbf{u}, \mathbf{b}'\mathbf{u}) \\ &= \mathbf{a}'\boldsymbol{\Sigma}\mathbf{b} \quad [\text{by (3.71)}] \\ &= (0, 1) \begin{pmatrix} \sigma_y^2 & \sigma_{yx} \\ \sigma_{yx} & \sigma_x^2 \end{pmatrix} \begin{pmatrix} 1 \\ -\beta \end{pmatrix} = (\sigma_{yx}, \sigma_x^2) \begin{pmatrix} 1 \\ -\beta \end{pmatrix} \\ &= \sigma_{yx} - \beta\sigma_x^2. \end{aligned}$$

Since $\text{cov}(x, z) = 0$, we obtain $\beta = \sigma_{yx}/\sigma_x^2$ and $z = y - \beta x$ becomes

$$z = y - \frac{\sigma_{yx}}{\sigma_x^2}x.$$

By property 1a above, the density of $y - (\sigma_{yx}/\sigma_x^2)x$ is normal with

$$\begin{aligned} E\left(y - \frac{\sigma_{yx}}{\sigma_x^2}x\right) &= \mu_y - \frac{\sigma_{yx}}{\sigma_x^2}\mu_x, \\ \text{var}\left(y - \frac{\sigma_{yx}}{\sigma_x^2}x\right) &= \text{var}(\mathbf{b}'\mathbf{u}) = \mathbf{b}'\boldsymbol{\Sigma}\mathbf{b} \\ &= \left(1, -\frac{\sigma_{yx}}{\sigma_x^2}\right) \begin{pmatrix} \sigma_y^2 & \sigma_{yx} \\ \sigma_{yx} & \sigma_x^2 \end{pmatrix} \begin{pmatrix} 1 \\ -\frac{\sigma_{yx}}{\sigma_x^2} \end{pmatrix} \\ &= \sigma_y^2 - \frac{\sigma_{yx}^2}{\sigma_x^2}. \end{aligned}$$

For a given value of x , we can express y as $y = \beta x + (y - \beta x)$, where βx is a fixed quantity corresponding to the given value of x and $y - \beta x$ is a random deviation. Then $f(y|x)$ is normal, with

$$\begin{aligned} E(y|x) &= \beta x + E(y - \beta x) = \beta x + \mu_y - \beta \mu_x \\ &= \mu_y + \beta(x - \mu_x) = \mu_y + \frac{\sigma_{yx}}{\sigma_x^2}(x - \mu_x), \\ \text{var}(y|x) &= \sigma_y^2 - \frac{\sigma_{yx}^2}{\sigma_x^2}. \end{aligned}$$

4.3 ESTIMATION IN THE MULTIVARIATE NORMAL

4.3.1 Maximum Likelihood Estimation

When a distribution such as the multivariate normal is assumed to hold for a population, estimates of the parameters are often found by the method of *maximum likelihood*. This technique is conceptually simple: The observation vectors $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n$ are considered to be known, and values of $\boldsymbol{\mu}$ and $\boldsymbol{\Sigma}$ are sought that maximize the joint density of the \mathbf{y} 's, called the *likelihood function*. For the multivariate normal, the maximum likelihood estimates of $\boldsymbol{\mu}$ and $\boldsymbol{\Sigma}$ are

$$\hat{\boldsymbol{\mu}} = \bar{\mathbf{y}}, \quad (4.11)$$

$$\begin{aligned} \hat{\boldsymbol{\Sigma}} &= \frac{1}{n} \sum_{i=1}^n (\mathbf{y}_i - \bar{\mathbf{y}})(\mathbf{y}_i - \bar{\mathbf{y}})' \\ &= \frac{1}{n} \mathbf{W} \\ &= \frac{n-1}{n} \mathbf{S}, \end{aligned} \quad (4.12)$$

where $\mathbf{W} = \sum_{i=1}^n (\mathbf{y}_i - \bar{\mathbf{y}})(\mathbf{y}_i - \bar{\mathbf{y}})'$ and \mathbf{S} is the sample covariance matrix defined in (3.22) and (3.27). Since $\hat{\boldsymbol{\Sigma}}$ has divisor n instead of $n-1$, it is biased [see (3.33)], and we usually use \mathbf{S} in place of $\hat{\boldsymbol{\Sigma}}$.

We now give a justification of $\bar{\mathbf{y}}$ as the maximum likelihood estimator of $\boldsymbol{\mu}$. Because the \mathbf{y}_i 's constitute a random sample, they are independent, and the joint density is the product of the densities of the \mathbf{y} 's. The likelihood function is, therefore,

$$\begin{aligned} L(\boldsymbol{\mu}, \boldsymbol{\Sigma}; \mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n) &= \prod_{i=1}^n f(\mathbf{y}_i, \boldsymbol{\mu}, \boldsymbol{\Sigma}) \\ &= \prod_{i=1}^n \frac{1}{(\sqrt{2\pi})^p |\boldsymbol{\Sigma}|^{1/2}} e^{-(\mathbf{y}_i - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1} (\mathbf{y}_i - \boldsymbol{\mu})/2} \\ &= \frac{1}{(\sqrt{2\pi})^{np} |\boldsymbol{\Sigma}|^{n/2}} e^{-\sum_{i=1}^n (\mathbf{y}_i - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1} (\mathbf{y}_i - \boldsymbol{\mu})/2}. \end{aligned} \quad (4.13)$$

To see that $\hat{\mu} = \bar{y}$ maximizes the likelihood function, we write the exponent of (4.13) in terms of \bar{y} . By adding and subtracting \bar{y} , the exponent in (4.13) becomes

$$-\frac{1}{2} \sum_{i=1}^n (\mathbf{y}_i - \bar{\mathbf{y}} + \bar{\mathbf{y}} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1} (\mathbf{y}_i - \bar{\mathbf{y}} + \bar{\mathbf{y}} - \boldsymbol{\mu}).$$

When this is expanded in terms of $\mathbf{y}_i - \bar{\mathbf{y}}$ and $\bar{\mathbf{y}} - \boldsymbol{\mu}$, two of the four resulting terms vanish because $\sum_i (\mathbf{y}_i - \bar{\mathbf{y}}) = \mathbf{0}$, and (4.13) becomes

$$L = \frac{1}{(\sqrt{2\pi})^{np} |\boldsymbol{\Sigma}|^{n/2}} e^{-\sum_{i=1}^n (\mathbf{y}_i - \bar{\mathbf{y}})' \boldsymbol{\Sigma}^{-1} (\mathbf{y}_i - \bar{\mathbf{y}})/2 - n(\bar{\mathbf{y}} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1} (\bar{\mathbf{y}} - \boldsymbol{\mu})/2}. \quad (4.14)$$

Since $\boldsymbol{\Sigma}^{-1}$ is positive definite, we have $-n(\bar{\mathbf{y}} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1} (\bar{\mathbf{y}} - \boldsymbol{\mu})/2 \leq 0$ and $0 < e^{-n(\bar{\mathbf{y}} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1} (\bar{\mathbf{y}} - \boldsymbol{\mu})/2} \leq 1$, with the maximum occurring when the exponent is 0. Therefore, L is maximized when $\hat{\mu} = \bar{y}$.

The maximum likelihood estimator of the population correlation matrix \mathbf{P}_p [see (3.39)] is the sample correlation matrix, that is,

$$\hat{\mathbf{P}}_p = \mathbf{R}.$$

Relationships among multinormal variables are linear, as can be seen in (4.7). Thus the estimators \mathbf{S} and \mathbf{R} serve well for the multivariate normal because they measure only linear relationships (see Sections 3.2.1 and 4.2). These estimators are not as useful for some nonnormal distributions.

4.3.2 Distribution of $\bar{\mathbf{y}}$ and \mathbf{S}

For the distribution of $\bar{\mathbf{y}} = \sum_{i=1}^n \mathbf{y}_i/n$, we can distinguish two cases:

- (a) When $\bar{\mathbf{y}}$ is based on a random sample $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n$ from a multivariate normal distribution $N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, then $\bar{\mathbf{y}}$ is $N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma}/n)$.
- (b) When $\bar{\mathbf{y}}$ is based on a random sample $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n$ from a nonnormal multivariate population with mean vector $\boldsymbol{\mu}$ and covariance matrix $\boldsymbol{\Sigma}$, then for large n , $\bar{\mathbf{y}}$ is approximately $N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma}/n)$. More formally, this result is known as the *multivariate central limit theorem*: If $\bar{\mathbf{y}}$ is the mean vector of a random sample $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n$ from a population with mean vector $\boldsymbol{\mu}$ and covariance matrix $\boldsymbol{\Sigma}$, then as $n \rightarrow \infty$, the distribution of $\sqrt{n}(\bar{\mathbf{y}} - \boldsymbol{\mu})$ approaches $N_p(\mathbf{0}, \boldsymbol{\Sigma})$.

There are p variances in \mathbf{S} and $\binom{p}{2}$ covariances, for a total of

$$p + \binom{p}{2} = p + p(p-1)/2 = p(p+1)/2$$

distinct entries. The joint distribution of these $p(p+1)/2$ distinct variables in $\mathbf{W} = (n-1)\mathbf{S} = \sum_i (\mathbf{y}_i - \bar{\mathbf{y}})(\mathbf{y}_i - \bar{\mathbf{y}})'$ is the Wishart distribution, denoted by $W_p(n-1, \boldsymbol{\Sigma})$, where $n-1$ is the degrees of freedom.

The Wishart distribution is the multivariate analogue of the χ^2 -distribution, and it has similar uses. As noted in property 3 of Section 4.2, a χ^2 random variable is defined formally as the sum of squares of independent standard normal (univariate) random variables:

$$\sum_{i=1}^n z_i^2 = \sum_{i=1}^n \frac{(y_i - \mu)^2}{\sigma^2} \quad \text{is } \chi^2(n).$$

If \bar{y} is substituted for μ , then $\sum_i (y_i - \bar{y})^2 / \sigma^2 = (n-1)s^2 / \sigma^2$ is $\chi^2(n-1)$. Similarly, the formal definition of a Wishart random variable is

$$\sum_{i=1}^n (\mathbf{y}_i - \boldsymbol{\mu})(\mathbf{y}_i - \boldsymbol{\mu})' \quad \text{is } W_p(n, \boldsymbol{\Sigma}), \quad (4.15)$$

where $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n$ are independently distributed as $N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$. When $\bar{\mathbf{y}}$ is substituted for $\boldsymbol{\mu}$, the distribution remains Wishart with one less degree of freedom:

$$(n-1)\mathbf{S} = \sum_{i=1}^n (\mathbf{y}_i - \bar{\mathbf{y}})(\mathbf{y}_i - \bar{\mathbf{y}})' \quad \text{is } W_p(n-1, \boldsymbol{\Sigma}). \quad (4.16)$$

Finally, we note that when sampling from a multivariate normal distribution, $\bar{\mathbf{y}}$ and \mathbf{S} are independent.

4.4 ASSESSING MULTIVARIATE NORMALITY

Many tests and graphical procedures have been suggested for evaluating whether a data set likely originated from a multivariate normal population. One possibility is to check each variable separately for univariate normality. Excellent reviews for both the univariate and multivariate cases have been given by Gnanadesikan (1997, pp. 178–220) and Seber (1984, pp. 141–155). We give a representative sample of univariate and multivariate methods in Sections 4.4.1 and 4.4.2, respectively.

4.4.1 Investigating Univariate Normality

When we have several variables, checking each for univariate normality should not be the sole approach, because (1) the variables are correlated and (2) normality of the individual variables does not guarantee joint normality. On the other hand, multivariate normality implies individual normality. Hence, if even one of the separate variables is not normal, the vector is not multivariate normal. An initial check on the individual variables may therefore be useful.

A basic graphical approach for checking normality is the $Q-Q$ plot comparing quantiles of a sample against the population quantiles of the univariate normal. If the points are close to a straight line, there is no indication of departure from normality. Deviation from a straight line indicates nonnormality (at least for a large sample). In




Figure 4.5 Typical $Q-Q$ plots for nonnormal data.

fact, the type of nonlinear pattern may reveal the type of departure from normality. Some possibilities are illustrated in Figure 4.5.

Quantiles are similar to the more familiar percentiles, which are expressed in terms of percent; a test score at the 90th percentile, for example, is above 90% of the test scores and below 10% of them. Quantiles are expressed in terms of fractions or proportions. Thus the 90th percentile score becomes the .9 quantile score.

The sample quantiles for the $Q-Q$ plot are obtained as follows. First we rank the observations y_1, y_2, \dots, y_n and denote the ordered values by $y_{(1)}, y_{(2)}, \dots, y_{(n)}$; thus $y_{(1)} \leq y_{(2)} \leq \dots \leq y_{(n)}$. Then the point $y_{(i)}$ is the i/n sample quantile. For example, if $n = 20$, $y_{(7)}$ is the $\frac{7}{20} = .35$ quantile, because .35 of the sample is less than or equal to $y_{(7)}$. The fraction i/n is often changed to $(i - \frac{1}{2})/n$ as a continuity correction. If $n = 20$, $(i - \frac{1}{2})/n$ ranges from .025 to .975 and more evenly covers the interval from 0 to 1. With this convention, $y_{(i)}$ is designated as the $(i - \frac{1}{2})/n$ sample quantile.

The population quantiles for the $Q-Q$ plot are similarly defined corresponding to $(i - \frac{1}{2})/n$. If we denote these by q_1, q_2, \dots, q_n , then q_i is the value below which a proportion $(i - \frac{1}{2})/n$ of the observations in the population lie; that is, $(i - \frac{1}{2})/n$ is the probability of getting an observation less than or equal to q_i . Formally, q_i can be found for the standard normal random variable y with distribution $N(0, 1)$ by solving

$$\Phi(q_i) = P(y < q_i) = \frac{i - \frac{1}{2}}{n}, \quad (4.17)$$

which would require numerical integration or tables of the cumulative standard normal distribution, $\Phi(x)$. Another benefit of using $(i - \frac{1}{2})/n$ instead of i/n is that $n/n = 1$ would make $q_n = \infty$.

The population need not have the same mean and variance as the sample, since changes in mean and variance merely change the slope and intercept of the plotted line in the $Q-Q$ plot. Therefore, we use the standard normal distribution, and the q_i values can easily be found from a table of cumulative standard normal probabilities. We then plot the pairs $(q_i, y_{(i)})$ and examine the resulting $Q-Q$ plot for linearity.

Special graph paper, called normal probability paper, is available that eliminates the need to look up the q_i values. We need only plot $(i - \frac{1}{2})/n$ in place of q_i , that is, plot the pairs $[(i - \frac{1}{2})/n, y_{(i)}]$ and look for linearity as before. As an even easier alternative, most general-purpose statistical software programs provide normal probability plots of the pairs $(q_i, y_{(i)})$.

The $Q-Q$ plots provide a good visual check on normality and are considered to be adequate for this purpose by many researchers. For those who desire a more objective procedure, several hypothesis tests are available. We give three of these that have good properties and are computationally tractable.

We discuss first a classical approach based on the following measures of skewness and kurtosis:

$$\sqrt{b_1} = \frac{\sqrt{n} \sum_{i=1}^n (y_i - \bar{y})^3}{[\sum_{i=1}^n (y_i - \bar{y})^2]^{3/2}}, \quad (4.18)$$

$$b_2 = \frac{n \sum_{i=1}^n (y_i - \bar{y})^4}{[\sum_{i=1}^n (y_i - \bar{y})^2]^2}. \quad (4.19)$$

These are sample estimates of the population skewness and kurtosis parameters $\sqrt{\beta_1}$ and β_2 , respectively. When the population is normal, $\sqrt{\beta_1} = 0$ and $\beta_2 = 3$. If




Figure 4.6 A distribution with positive skewness.

$\sqrt{\beta_1} < 0$, we have negative skewness; if $\sqrt{\beta_1} > 0$, the skewness is positive. Positive skewness is illustrated in Figure 4.6. If $\beta_2 < 3$, we have negative kurtosis, and if $\beta_2 > 3$, there is positive kurtosis. A distribution with negative kurtosis is characterized by being flatter than the normal distribution, that is, less peaked, with heavier flanks and thinner tails. A distribution with positive kurtosis has a higher peak than the normal, with an excess of values near the mean and in the tails but with thinner flanks. Positive and negative kurtosis are illustrated in Figure 4.7.

The test of normality can be carried out using the exact percentage points for $\sqrt{b_1}$ in Table A.1 for $4 \leq n \leq 25$, as given by Mulholland (1977). Alternatively, for $n \geq 8$ the function g as defined by

$$g(\sqrt{b_1}) = \delta \sinh^{-1}\left(\frac{\sqrt{b_1}}{\lambda}\right) \quad (4.20)$$

is approximately $N(0, 1)$, where

$$\sinh^{-1}(x) = \ln(x + \sqrt{x^2 + 1}). \quad (4.21)$$

Table A.2, from D'Agostino and Pearson (1973), gives values for δ and $1/\lambda$. To use b_2 as a test of normality, we can use Table A.3, from D'Agostino and Tietjen (1971), which gives simulated percentiles of b_2 for selected values of n in the range $7 \leq n \leq 50$. Charts of percentiles of b_2 for $20 \leq n \leq 200$ can be found in D'Agostino and Pearson (1973).




Figure 4.7 Distributions with positive and negative kurtosis compared to the normal.

Our second test for normality was given by D'Agostino (1971). The observations y_1, y_2, \dots, y_n are ordered as $y_{(1)} \leq y_{(2)} \leq \dots \leq y_{(n)}$, and we calculate

$$D = \frac{\sum_{i=1}^n [i - \frac{1}{2}(n+1)] y_{(i)}}{\sqrt{n^3 \sum_{i=1}^n (y_i - \bar{y})^2}}, \quad (4.22)$$

$$Y = \frac{\sqrt{n}[D - (2\sqrt{\pi})^{-1}]}{.02998598}. \quad (4.23)$$

A table of percentiles for Y , given by D'Agostino (1972) for $10 \leq n \leq 250$, is provided in Table A.4.

The final test we report is by Lin and Mudholkar (1980). The test statistic is

$$z = \tanh^{-1}(r) = \frac{1}{2} \ln \left(\frac{1+r}{1-r} \right), \quad (4.24)$$

where r is the sample correlation of the n pairs (y_i, x_i) , $i = 1, 2, \dots, n$, with x_i defined as

$$x_i = \frac{1}{n} \left[\sum_{j \neq i} y_j^2 - \frac{\left(\sum_{j \neq i} y_j \right)^2}{n-1} \right]^{1/3}. \quad (4.25)$$

If the \mathbf{y} 's are normal, z is approximately $N(0, 3/n)$. A more accurate upper 100α percentile is given by

$$z_\alpha = \sigma_n \left[u_\alpha + \frac{1}{24} (u_\alpha^3 - 3u_\alpha) \gamma_{2n} \right], \quad (4.26)$$

with

$$\begin{aligned}\sigma_n^2 &= \frac{3}{n} - \frac{7.324}{n^2} + \frac{53.005}{n^3}, \quad u_\alpha = \Phi^{-1}(\alpha), \\ \gamma_{2n} &= -\frac{11.70}{n} + \frac{55.06}{n^2},\end{aligned}$$

where Φ is the distribution function of the $N(0, 1)$ distribution; that is, $\Phi(x)$ is the probability of an observation less than or equal to x , as in (4.17). The inverse function Φ^{-1} is essentially a quantile. For example, $u_{.05} = -1.645$ and $u_{.95} = 1.645$.

4.4.2 Investigating Multivariate Normality

Checking for multivariate normality is conceptually not as straightforward as assessing univariate normality, and consequently the state of the art is not as well developed. The complexity of this issue can be illustrated in the context of a goodness-of-fit test for normality. For a goodness-of-fit test in the univariate case, the range covered by a sample y_1, y_2, \dots, y_n is divided into several intervals, and we count how many y 's fall into each interval. These observed frequencies (counts) are compared to the expected frequencies under the assumption that the sample came from a normal distribution with the same mean and variance as the sample. If the n observations $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n$ are multivariate, however, the procedure is not so simple. We now have a p -dimensional region that would have to be divided into many more subregions than in the univariate case, and the expected frequencies for these subregions would be less easily obtained. With so many subregions, relatively few would contain observations.

Thus because of the inherent “sparseness” of multivariate data, a goodness-of-fit test would be impractical. The points $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n$ are more distant from each other in p -space than in any one of the p individual dimensions. Unless n is very large, a multivariate sample may not provide a very complete picture of the distribution from which it was taken.

As a consequence of the sparseness of the data in p -space, the tests for multivariate normality may not be very powerful. However, some check on the distribution is often desirable. Numerous procedures have been proposed for assessing multivariate normality. We now discuss three of these.

The first procedure is based on the standardized distance from each \mathbf{y}_i to $\bar{\mathbf{y}}$,

$$D_i^2 = (\mathbf{y}_i - \bar{\mathbf{y}})' \mathbf{S}^{-1} (\mathbf{y}_i - \bar{\mathbf{y}}), \quad i = 1, 2, \dots, n. \quad (4.27)$$

Gnanadesikan and Kettenring (1972) showed that if the \mathbf{y}_i 's are multivariate normal, then

$$u_i = \frac{n D_i^2}{(n-1)^2} \quad (4.28)$$

has a beta distribution, which is related to the F . To obtain a $Q-Q$ plot, the values u_1, u_2, \dots, u_n are ranked to give $u_{(1)} \leq u_{(2)} \leq \dots \leq u_{(n)}$, and we plot $(u_{(i)}, v_i)$, where the quantiles v_i of the beta are given by the solution to

$$\int_0^{v_i} \frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} x^{a-1} (1-x)^{b-1} dx = \frac{i-\alpha}{n-\alpha-\beta+1}, \quad (4.29)$$

where $a = \frac{1}{2} p$, $b = \frac{1}{2} (n-p-1)$,

$$\alpha = \frac{p-2}{2p}, \quad (4.30)$$

$$\beta = \frac{n-p-3}{2(n-p-1)}. \quad (4.31)$$

A nonlinear pattern in the plot would indicate a departure from normality. The quantities of the beta are easily obtained in many software packages. A formal significance test is also available for $D_{(n)}^2 = \max_i D_i^2$. Table A.6 gives the upper 5% and 1% critical values for $p = 2, 3, 4, 5$ from Barnett and Lewis (1978).

Some writers have suggested that the distribution of D_i^2 in (4.27) can be adequately approximated by a χ_p^2 since $(\mathbf{y} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1} (\mathbf{y} - \boldsymbol{\mu})$ is χ_p^2 [see (4.6)]. However, in Section 5.3.2, it is shown that this approximation is very poor for even moderate values of p . Small (1978) showed that plots of D_i^2 vs. χ^2 quantiles are misleading.

The second procedure involves scatterplots in two dimensions. If p is not too high, the bivariate plots of each pair of variables are often reduced in size and shown on one page, arranged to correspond to the entries in a correlation matrix. In this visual matrix, the eye readily picks out those pairs of variables that show a curved trend, outliers, or other nonnormal appearance. This plot is illustrated in Example 4.6.2 in Section 4.6.2. The procedure is based on properties 4 and 6 of Section 4.2, from which we infer that (1) each pair of variables has a bivariate normal distribution and (2) bivariate normal variables follow a straight-line trend.

An extension of this method for assessing multivariate normality is the use of the grand tour discussed in Section 3.5. In the grand tour, we consider a series of random two-dimensional projections of the p -variate data. For the purposes of normality assessment, properties 1 and 6 from Section 4.2 imply that each projection viewed during the grand tour should exhibit bivariate normality, with no indications of nonlinearity in the projected data. Any deviation from elliptical clouds of points is an indication of potential nonnormality.

The third procedure for assessing multivariate normality is a generalization of the univariate test based on the skewness and kurtosis measures $\sqrt{b_1}$ and b_2 as given by (4.18) and (4.19). The test is due to Mardia (1970). Let \mathbf{y} and \mathbf{x} be independent and identically distributed with mean vector $\boldsymbol{\mu}$ and covariance matrix $\boldsymbol{\Sigma}$. Then skewness and kurtosis for multivariate populations are defined by Mardia as

$$\beta_{1,p} = E[(\mathbf{y} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1} (\mathbf{x} - \boldsymbol{\mu})]^3, \quad (4.32)$$

$$\beta_{2,p} = E[(\mathbf{y} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1} (\mathbf{y} - \boldsymbol{\mu})]^2. \quad (4.33)$$

Since third-order central moments for the multivariate normal distribution are zero, $\beta_{1,p} = 0$ when \mathbf{y} is $N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$. It can also be shown that if \mathbf{y} is $N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, then

$$\beta_{2,p} = p(p+2). \quad (4.34)$$

To estimate $\beta_{1,p}$ and $\beta_{2,p}$ using a sample $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_p$, we first define

$$g_{ij} = (\mathbf{y}_i - \bar{\mathbf{y}})' \hat{\boldsymbol{\Sigma}}^{-1} (\mathbf{y}_j - \bar{\mathbf{y}}), \quad (4.35)$$

where $\hat{\boldsymbol{\Sigma}} = \sum_{i=1}^n (\mathbf{y}_i - \bar{\mathbf{y}})(\mathbf{y}_i - \bar{\mathbf{y}})' / n$ is the maximum likelihood estimator (4.12). Then estimates of $\beta_{1,p}$ and $\beta_{2,p}$ are given by

$$b_{1,p} = \frac{1}{n^2} \sum_{i=1}^n \sum_{j=1}^n g_{ij}^3, \quad (4.36)$$

$$b_{2,p} = \frac{1}{n} \sum_{i=1}^n g_{ii}^2. \quad (4.37)$$

Table A.5 (Mardia 1970, 1974) gives percentage points of $b_{1,p}$ and $b_{2,p}$ for $p = 2, 3, 4$, which can be used in testing for multivariate normality. For other values of p or when $n \geq 50$, the following approximate tests are available. For $b_{1,p}$, the statistic

$$z_1 = \frac{(p+1)(n+1)(n+3)}{6[(n+1)(p+1)-6]} b_{1,p} \quad (4.38)$$

is approximately χ^2 with $\frac{1}{6}p(p+1)(p+2)$ degrees of freedom. Reject the hypothesis of multivariate normality if $z_1 \geq \chi^2_{0.05}$. With $b_{2,p}$, on the other hand, we wish to reject for large values (distribution too peaked) or small values (distribution too flat). For the upper 2.5% points of $b_{2,p}$ use

$$z_2 = \frac{b_{2,p} - p(p+2)}{\sqrt{8p(p+2)/n}}, \quad (4.39)$$

which is approximately $N(0, 1)$. For the lower 2.5% points we have two cases: (1) when $50 \leq n \leq 400$, use

$$z_3 = \frac{b_{2,p} - p(p+2)(n+p+1)/n}{\sqrt{8p(p+2)/(n-1)}}, \quad (4.40)$$

which is approximately $N(0, 1)$; (2) when $n \geq 400$, use z_2 as given by (4.39).

4.5 TRANSFORMATIONS TO NORMALITY

As noted in Section 4.3.2, the asymptotic distribution of $\bar{\mathbf{y}}$ is normal, regardless of the distribution of \mathbf{y} . However, there are many scenarios in which we wish to

draw inferences about statistics other than the mean, and/or we do not have a large enough sample size to appeal to the central limit theorem. For such situations, it is often useful to consider transformations of the data that exhibit greater normality than the original data. In many cases, transformations will be very natural from a phenomenological standpoint. For example, log abundance of bacteria in a replicated experiment may not only appear more normal, but will often be a more natural way of quantifying the growth process of interest.

Although our aim here is to find a multivariate transformation $g(\cdot)$ that makes $g(y)$ look as close to p -variate normality as possible, it is often the case that transforming each of the p variables to a marginal normal distribution will yield a multivariate transformation that is close to optimal. Because univariate transformations are much less complicated, we primarily focus on the univariate approach developed by Box and Cox (1964). We then briefly consider an extension for multivariate transformations based on Andrews, Gnanadesikan, and Warner (1971).

4.5.1 Univariate Transformations to Normality

When the histogram or normal probability plot for a variable y exhibits skewness, a transformation will usually yield a modified distribution that is more normal in appearance. For increasingly right-skewed observations, it is useful to use transformations such as $\log(y)$, y^{-1} , y^{-2} , etc. For increasingly left-skewed observations, it is useful to use transformations such as y^2 , y^3 , y^4 , etc. One can choose an appropriate transformation “by eye” using the trial-and-error approach of choosing a transformation $g(\cdot)$, evaluating the histogram of $g(y)$, and repeating the process until the “best” transformation has been obtained. Because this process is neither systematic nor objective, Box and Cox (1964) present the family of transformations defined by

$$y^{(\lambda)} = \begin{cases} \frac{y^\lambda - 1}{\lambda} & \text{for } \lambda \neq 0, \\ \ln(y) & \text{for } \lambda = 0. \end{cases} \quad (4.41)$$

The value of λ is chosen by maximizing

$$\ell(\lambda) = -\frac{n}{2} \ln s_\lambda^2 + (\lambda - 1) \sum_{i=1}^n \ln(y_i), \quad (4.42)$$

where

$$s_\lambda^2 = 1/n \sum_{i=1}^n (y_i^{(\lambda)} - \bar{y}^{(\lambda)})^2$$

is the maximum likelihood estimate of the variance of $y^{(\lambda)}$ and $\bar{y}^{(\lambda)}$ is the sample mean of the n transformed observations. Note that although the transformations (y^1, y^2, y^3, \dots) are not substantively different from the Box–Cox transformations $(y^{(1)}, y^{(2)}, y^{(3)}, \dots)$ defined by (4.41), we use the Box–Cox transformations because the criterion to be maximized [$\ell(\lambda)$] is continuous in λ , even when λ approaches 0. Thus, the Box–Cox transformations allow us to objectively evaluate the comparative utility

of any possible transform $y^{(\lambda)}$. The optimal λ can be found by explicitly maximizing the function in (4.42), but it is more common that $\ell(\lambda)$ is calculated for a wide range of values for λ , and then λ is plotted against $\ell(\lambda)$ to find the optimal value.

4.5.2 Multivariate Transformations to Normality

An extension of the Box–Cox method for transforming to p -variate normality was proposed by Andrews, Gnanadesikan, and Warner (1971). The multivariate approach is motivated by the fact that the optimal transformation to p -variate normality is not necessarily based on the collection of p transformations obtained when targeting marginal normality for each of the p variables. In practice, the collection of univariately transformed variables will often be very close to the optimal p -variate transformation, but a more precise criterion for transforming to multivariate normality is to find the collection of transformations $\boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_p)$ such that the function

$$\ell(\boldsymbol{\lambda}) = -\frac{n}{2} \ln |\mathbf{S}_{\boldsymbol{\lambda}}| + \sum_{j=1}^p \left[(\lambda_j - 1) \sum_{i=1}^n \ln(y_{ij}) \right] \quad (4.43)$$

is maximized. In (4.43), y_{ij} is the i th measurement on the j th variable, $\mathbf{S}_{\boldsymbol{\lambda}}$ is the maximum likelihood estimate of the covariance matrix for the transformed data with its (j, k) element equal to

$$\frac{1}{n} \sum_{i=1}^n (y_{ij}^{(\lambda_j)} - \overline{y_j^{(\lambda_j)}})(y_{ik}^{(\lambda_k)} - \overline{y_k^{(\lambda_k)}}),$$

and $\overline{y_k^{(\lambda_k)}}$ is the sample mean of the n transformed observations on the k th variable. As in the univariate case in Section 4.5.1, the optimal value for $\boldsymbol{\lambda}$ can be found by explicitly maximizing the function in (4.43), or the value of $\ell(\boldsymbol{\lambda})$ can be found for every choice of $\boldsymbol{\lambda}$ within a p -dimensional array of candidate values.

■ EXAMPLE 4.5.2

We use the soil richness data in Table 3.3 to illustrate the process of transforming data to appear more normal. The top row of Figure 4.8 illustrates distributions for the concentrations of Fe, K, and Mg. Note that Fe appears relatively normal, with Mg exhibiting moderate right-skewness and K exhibiting extreme right-skewness. Applying the Box–Cox approach discussed in Section 4.5.1, we obtain λ values of 1.0, -0.3 , and 0.1, respectively. The middle row of Figure 4.8 shows the plot of $\ell(\lambda)$ versus λ for each variable. The distribution for iron (Fe) concentrations is at the near-optimal transformation for appearing normal. Potassium (K) requires a transformation that is more extreme than the log transformation, while magnesium (Mg) needs a transformation that is slightly less extreme than the log transformation. The bottom row of Figure 4.8 shows the distributions of the three elements after applying the optimal univariate Box–Cox transformation.




Figure 4.8 Transforming data from Table 3.3 to near-normality. Top row: histograms for concentrations of Fe, K, and Mg. Middle row: plot of λ against $\ell(\lambda)$ for Fe, K, and Mg. Bottom row: histograms for the transformed concentrations of Fe, K, and Mg when using λ values of 1.0 (no transformation), -0.3, and 0.1, respectively.

We apply the multivariate Box–Cox approach discussed in Section 4.5.2 by considering an array of values for $\lambda = (\lambda_{\text{Fe}}, \lambda_K, \lambda_{\text{Mg}})$ such that $\lambda_{\text{Fe}} \in (0.5, 0.6, \dots, 1.5)$, $\lambda_K \in (-0.8, -0.7, \dots, 0.2)$, and $\lambda_{\text{Mg}} \in (-0.4, -0.3, \dots, 0.6)$. The value of λ that maximizes $\ell(\lambda)$ in (4.43) is $\lambda = (1.0, -0.4, 0.2)$, which is predictably similar to the transformation obtained by applying the optimal univariate transformation to each of the three variables. \square

4.6 OUTLIERS

The detection of outliers has been of concern to statisticians and other scientists for over a century. Some authors have claimed that the researcher can typically expect

up to 10% of the observations to have errors in measurement or recording. Occasional stray observations from a different population than the target population are also fairly common. We review some major concepts and suggested procedures for univariate outliers in Section 4.6.1 before moving to the multivariate case in Section 4.6.2. An alternative to detection of outliers is to use robust estimators of μ and Σ (see Rencher 1998, Section 1.10) that are less sensitive to extreme observations than are the standard estimators \bar{y} and S .

4.6.1 Outliers in Univariate Samples

Excellent treatments of outliers have been given by Beckman and Cook (1983), Hawkins (1980), and Barnett and Lewis (1978). We abstract a few highlights from Beckman and Cook. Many techniques have been proposed for detecting outliers in the residuals from regression or designed experiments, but we will be concerned only with simple random samples from the normal distribution.

There are two principal approaches for dealing with outliers. The first is *identification*, which usually involves deletion of the outlier(s) but may also provide important information about the model or the data. The second method is *accommodation*, in which the method of analysis or the model is modified. Robust methods, in which the influence of outliers is reduced, provide an example of modification of the analysis. An example of a correction to the model is a mixture model that combines two normals with different variances. For example, Marks and Rao (1978) accommodated a particular type of outlier by a mixture of two normal distributions.

In small or moderate-sized univariate samples, visual methods of identifying outliers are the most frequently used. Tests are also available if a less subjective approach is desired.

Two types of *slippage* models have been proposed to account for outliers. Under the *mean slippage* model, all observations have the same variance, but one or more of the observations arise from a distribution with a different (population) mean. In the *variance slippage* model, one or more of the observations arise from a model with larger (population) variance but the same mean. Thus in the mean slippage model, the bulk of the observations arise from $N(\mu, \sigma^2)$, whereas the outliers originate from $N(\mu + \theta, \sigma^2)$. For the variance slippage model, the main distribution would again be $N(\mu, \sigma^2)$, with the outliers coming from $N(\mu, a\sigma^2)$ where $a > 1$. These models have led to the development of tests for rejection of outliers. We now briefly discuss some of these tests.

For a single outlier in a sample y_1, y_2, \dots, y_n , most tests are based on the maximum studentized residual,

$$\max_i \tau_i = \max_i \left| \frac{y_i - \bar{y}}{s} \right|. \quad (4.44)$$

If the largest or smallest observation is rejected, one could then examine the $n - 1$ remaining observations for another possible outlier, and so on. This procedure is called a *consecutive test*. However, if there are two or more outliers, the less extreme

ones will often make it difficult to detect the most extreme one, due to inflation of both mean and variance. This effect is called *masking*.

Ferguson (1961) showed that the maximum studentized residual (4.44) is more powerful than most other techniques for detecting intermediate or large shifts in the mean and gave the following guidelines for small shifts:

1. For outliers with small positive shifts in the mean, tests based on sample skewness are best.
2. For outliers with small shifts in the mean in either direction, tests based on the sample kurtosis are best.
3. For outliers with small positive shifts in the variance, tests based on the sample kurtosis are best.

Because of the masking problem in consecutive tests, *block tests* have been proposed for simultaneous rejection of $k > 1$ outliers. These tests work well if k is known, but in practice, k is usually not known. If the value we conjecture for k is too small, we incur the risk of failing to detect any outliers because of masking. If we set k too large, there is a high risk of rejecting more outliers than there really are, an effect known as *swamping*.

4.6.2 Outliers in Multivariate Samples

In the case of multivariate data, the problems in detecting outliers are intensified for several reasons:

1. For $p > 2$ the data cannot be readily plotted to pinpoint the outliers.
2. Multivariate data cannot be ordered as can a univariate sample, where extremes show up readily on either end.
3. An observation vector may have a large recording error in one of its components or smaller errors in several components.
4. A multivariate outlier may reflect slippage in mean, variance, or correlation. This is illustrated in Figure 4.9. Observation 1 causes a small shift in means and variances of both y_1 and y_2 but has little effect on the correlation. Observation 2 has little effect on means and variances, but it reduces the correlation somewhat. Observation 3 has a major effect on means, variances, and correlation.

One approach to multivariate outlier identification or accommodation is to use robust methods of estimation. Such methods minimize the influence of outliers in estimation or model fitting. However, an outlier sometimes furnishes valuable information, and the specific pursuit of outliers can be very worthwhile.

We present two methods of multivariate outlier identification, both of which are related to methods of assessing multivariate normality. (A third approach based




Figure 4.9 Bivariate sample showing three types of outliers.

on principal components is given in Section 12.4.) The first method, due to Wilks (1963), is designed for detection of a single outlier. Wilks' statistic is

$$w = \min_i \frac{|(n-2)\mathbf{S}_{-i}|}{|(n-1)\mathbf{S}|}, \quad (4.45)$$

where \mathbf{S} is the usual sample covariance matrix and \mathbf{S}_{-i} is obtained from the same sample with the i th observation deleted. The statistic w can also be expressed in terms of $D_{(n)}^2 = \max_i (\mathbf{y}_i - \bar{\mathbf{y}})' \mathbf{S}^{-1} (\mathbf{y}_i - \bar{\mathbf{y}})$ as

$$w = 1 - \frac{n D_{(n)}^2}{(n-1)^2}, \quad (4.46)$$

thus basing a test for an outlier on the distances D_i^2 used in Section 4.4.2 in a graphical procedure for checking multivariate normality. Table A.6 gives the upper 5% and 1% critical values for $D_{(n)}^2$ from Barnett and Lewis (1978).

Yang and Lee (1987) provide an F -test of w as given by (4.46). Define

$$F_i = \frac{n-p-1}{p} \left[\frac{1}{1 - n D_i^2 / (n-1)^2} - 1 \right], \quad i = 1, 2, \dots, n. \quad (4.47)$$

Then the F_i 's are independently and identically distributed as $F_{p, n-p-1}$, and a test can be constructed in terms of $\max_i F_i$:

$$P\left(\max_i F_i > f\right) = 1 - P(\text{all } F_i \leq f) = 1 - [P(F \leq f)]^n.$$

Therefore, the test can be carried out using an F -table. Note that

$$\max_i F_i = F_{(n)} = \frac{n-p-1}{p} \left(\frac{1}{w} - 1 \right), \quad (4.48)$$

where w is given in (4.46).

The second test we discuss is designed for detection of several outliers. Schwager and Margolin (1982) showed that the locally best invariant test for mean slippage is based on Mardia's (1970) sample kurtosis $b_{2,p}$ as defined by (4.35) and (4.37). To be more specific, among all tests invariant to a class of transformations of the type $\mathbf{z} = \mathbf{Ay} + \mathbf{b}$, where \mathbf{A} is nonsingular (see Problem 4.8), the test using $b_{2,p}$ is most powerful for small shifts in the mean vector. This result holds if the proportion of outliers is no more than 21.13%. With some restrictions on the pattern of the outliers, the permissible fraction of outliers can go as high as $33\frac{1}{3}\%$. The hypothesis is H_0 : no outliers are present. This hypothesis is rejected for large values of $b_{2,p}$.

A table of critical values of $b_{2,p}$ and some approximate tests were described in Section 4.4.2 following (4.37). Thus the test doubles as a check for multivariate normality and for the presence of outliers. One advantage of this test for outliers is that we do not have to specify the number of outliers and run the attendant risk of masking or swamping. Schwager and Margolin (1982) pointed out that this feature "increases the importance of performing an overall test that is sensitive to a broad range of outlier configurations. There is also empirical evidence that the kurtosis test performs well in situations of practical interest when compared with other inferential outlier procedures."

Sinha (1984) extended the result of Schwager and Margolin to cover the general case of elliptically symmetric distributions. An *elliptically symmetric distribution* is one in which $f(\mathbf{y}) = |\Sigma|^{-1/2} g[(\mathbf{y} - \boldsymbol{\mu})' \Sigma^{-1} (\mathbf{y} - \boldsymbol{\mu})]$. By varying the function g , distributions with shorter or longer tails than the normal can be obtained. The critical value of $b_{2,p}$ must be adjusted to correspond to the distribution, but rejection for large values would be a locally best invariant test.


■ EXAMPLE 4.6.2

We use the ramus bone data set of Table 3.7 to illustrate a search for multivariate outliers, while at the same time checking for multivariate normality. An examination of each column of Table 3.7 does not reveal any apparent univariate outliers. To check for multivariate outliers, we first calculate D_i^2 in (4.27) for each observation vector. The results are given in Table 4.1. We see that D_9^2 , D_{12}^2 , and D_{20}^2 seem to stand out as possible outliers. In Table A.6, the upper 5% critical value for the maximum value, $D_{(20)}^2$, is given as 11.63. In our case, the largest D_i^2 is $D_9^2 = 11.03$, which does not exceed the critical value. This does not surprise us, since the test was designed to detect a single outlier, and we may have as many as three.

We compute u_i and v_i in (4.28) and (4.29) and plot them in Figure 4.10. The figure shows a departure from linearity due to three values and possibly a fourth.

Table 4.1 Values of D_i^2 for the Ramus Bone Data in Table 3.7

Observation Number	D_i^2	Observation Number	D_i^2
1	0.7588	11	2.8301
2	1.2980	12	10.5718
3	1.7591	13	2.5941
4	3.8539	14	0.6594
5	0.8706	15	0.3246
6	2.8106	16	0.8321
7	4.2915	17	1.1083
8	7.9897	18	4.3633
9	11.0301	19	2.1088
10	5.3519	20	10.0931

**Figure 4.10** $Q-Q$ plot of u_i and v_i for the ramus bone data of Table 3.7.

We next calculate $b_{1,p}$ and $b_{2,p}$ as given by (4.36) and (4.37):

$$b_{1,p} = 11.338, \quad b_{2,p} = 28.884.$$

In Table A.5, the upper .01 critical value for $b_{1,p}$ is 9.9; the upper .005 critical value for $b_{2,p}$ is 27.1. Thus both $b_{1,p}$ and $b_{2,p}$ exceed their critical values, and we have significant skewness and kurtosis, apparently caused by the three observations with large values of D_i^2 .




Figure 4.11 Scatterplots for the ramus bone data in Table 3.7, with density curves for each variable.

The bivariate scatterplots are given in Figure 4.11. Three values are clearly separate from the other observations in the plot of y_1 versus y_4 . In Table 3.7, the 9th, 12th, and 20th values of y_4 are not unusual, nor are the 9th, 12th, and 20th values of y_1 . However, the increase from y_1 to y_4 is exceptional in each case. If these values are not due to errors in recording the data and if this sample is representative, then we appear to have a mixture of two populations. This should be taken into account in making inferences. \square

PROBLEMS

- 4.1** Consider the two covariance matrices

$$\Sigma_1 = \begin{pmatrix} 14 & 8 & 3 \\ 8 & 5 & 2 \\ 3 & 2 & 1 \end{pmatrix}, \quad \Sigma_2 = \begin{pmatrix} 6 & 6 & 1 \\ 6 & 8 & 2 \\ 1 & 2 & 1 \end{pmatrix}.$$

Show that $|\Sigma_2| > |\Sigma_1|$ and that $\text{tr}(\Sigma_2) < \text{tr}(\Sigma_1)$. Thus the generalized variance of population 2 is greater than the generalized variance of population 1, even though the total variance is less. Comment on why this is true in terms of the variances and correlations.

- 4.2** For $\mathbf{z} = (\mathbf{T}')^{-1}(\mathbf{y} - \boldsymbol{\mu})$ in (4.4), show that $E(\mathbf{z}) = \mathbf{0}$ and $\text{cov}(\mathbf{z}) = \mathbf{I}$.
- 4.3** Show that the form of the likelihood function in (4.13) follows from the previous expression.
- 4.4** For $(\mathbf{y} - \boldsymbol{\mu})'\boldsymbol{\Sigma}^{-1}(\mathbf{y} - \boldsymbol{\mu})$ in (4.3) and (4.6), show that $E[(\mathbf{y} - \boldsymbol{\mu})'\boldsymbol{\Sigma}^{-1}(\mathbf{y} - \boldsymbol{\mu})] = p$. Assume $E(\mathbf{y}) = \boldsymbol{\mu}$ and $\text{cov}(\mathbf{y}) = \boldsymbol{\Sigma}$. Normality is not required.
- 4.5** Show that by adding and subtracting $\bar{\mathbf{y}}$, the exponent of (4.13) has the form given in (4.14), that is,

$$\begin{aligned} & \frac{1}{2} \sum_{i=1}^n (\mathbf{y}_i - \bar{\mathbf{y}} + \bar{\mathbf{y}} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1} (\mathbf{y}_i - \bar{\mathbf{y}} + \bar{\mathbf{y}} - \boldsymbol{\mu}) \\ &= \frac{1}{2} \sum_{i=1}^n (\mathbf{y}_i - \bar{\mathbf{y}})' \boldsymbol{\Sigma}^{-1} (\mathbf{y}_i - \bar{\mathbf{y}}) + \frac{n}{2} (\bar{\mathbf{y}} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1} (\bar{\mathbf{y}} - \boldsymbol{\mu}). \end{aligned}$$

- 4.6** Show that $\sqrt{b_1}$ and b_2 , as given in (4.18) and (4.19), are invariant to the transformation $z_i = ay_i + b$.
- 4.7** Show that if \mathbf{y} is $N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, then $\beta_{2,p} = p(p+2)$ as in (4.34).
- 4.8** Show that $b_{1,p}$ and $b_{2,p}$, as given by (4.36) and (4.37), are invariant under the transformation $\mathbf{z}_i = \mathbf{A}\mathbf{y}_i + \mathbf{b}$, where \mathbf{A} is nonsingular. Thus $b_{1,p}$ and $b_{2,p}$ do not depend on the units of measurement.
- 4.9** Show that $F_{(n)} = [(n-p-1)/p](1/w-1)$ as in (4.48).
- 4.10** Suppose \mathbf{y} is $N_3(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, where
- $$\boldsymbol{\mu} = \begin{pmatrix} 3 \\ 1 \\ 4 \end{pmatrix}, \quad \boldsymbol{\Sigma} = \begin{pmatrix} 6 & 1 & -2 \\ 1 & 13 & 4 \\ -2 & 4 & 4 \end{pmatrix}.$$
- (a) Find the distribution of $z = 2y_1 - y_2 + 3y_3$.
- (b) Find the joint distribution of $z_1 = y_1 + y_2 + y_3$ and $z_2 = y_1 - y_2 + 2y_3$.
- (c) Find the distribution of y_2 .
- (d) Find the joint distribution of y_1 and y_3 .
- (e) Find the joint distribution of y_1 , y_3 , and $\frac{1}{2}(y_1 + y_2)$.
- 4.11** Suppose \mathbf{y} is $N_3(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, with $\boldsymbol{\mu}$ and $\boldsymbol{\Sigma}$ given in Problem 4.10.

- (a) Find a vector \mathbf{z} such that $\mathbf{z} = (\mathbf{T}')^{-1}(\mathbf{y} - \boldsymbol{\mu})$ is $N_3(\mathbf{0}, \mathbf{I})$ as in (4.4).
 (b) Find a vector \mathbf{z} such that $\mathbf{z} = (\boldsymbol{\Sigma}^{1/2})^{-1}(\mathbf{y} - \boldsymbol{\mu})$ is $N_3(\mathbf{0}, \mathbf{I})$ as in (4.5).
 (c) What is the distribution of $(\mathbf{y} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1}(\mathbf{y} - \boldsymbol{\mu})$?

4.12 Suppose \mathbf{y} is $N_4(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, where

$$\boldsymbol{\mu} = \begin{pmatrix} -2 \\ 3 \\ -1 \\ 5 \end{pmatrix}, \quad \boldsymbol{\Sigma} = \begin{pmatrix} 11 & -8 & 3 & 9 \\ -8 & 9 & -3 & -6 \\ 3 & -3 & 2 & 3 \\ 9 & -6 & 3 & 9 \end{pmatrix}.$$

- (a) Find the distribution of $z = 4y_1 - 2y_2 + y_3 - 3y_4$.
 (b) Find the joint distribution of $z_1 = y_1 + y_2 + y_3 + y_4$ and $z_2 = -2y_1 + 3y_2 + y_3 - 2y_4$.
 (c) Find the joint distribution of $z_1 = 3y_1 + y_2 - 4y_3 - y_4$, $z_2 = -y_1 - 3y_2 + y_3 - 2y_4$, and $z_3 = 2y_1 + 2y_2 + 4y_3 - 5y_4$.
 (d) What is the distribution of y_3 ?
 (e) What is the joint distribution of y_2 and y_4 ?
 (f) Find the joint distribution of y_1 , $\frac{1}{2}(y_1 + y_2)$, $\frac{1}{3}(y_1 + y_2 + y_3)$, and $\frac{1}{4}(y_1 + y_2 + y_3 + y_4)$.

4.13 Suppose \mathbf{y} is $N_4(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ with $\boldsymbol{\mu}$ and $\boldsymbol{\Sigma}$ given in Problem 4.12.

- (a) Find a vector \mathbf{z} such that $\mathbf{z} = (\mathbf{T}')^{-1}(\mathbf{y} - \boldsymbol{\mu})$ is $N_4(\mathbf{0}, \mathbf{I})$ as in (4.4).
 (b) Find a vector \mathbf{z} such that $\mathbf{z} = (\boldsymbol{\Sigma}^{1/2})^{-1}(\mathbf{y} - \boldsymbol{\mu})$ is $N_4(\mathbf{0}, \mathbf{I})$ as in (4.5).
 (c) What is the distribution of $(\mathbf{y} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1}(\mathbf{y} - \boldsymbol{\mu})$?

4.14 Suppose \mathbf{y} is $N_3(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, with

$$\boldsymbol{\mu} = \begin{pmatrix} 2 \\ -3 \\ 4 \end{pmatrix}, \quad \boldsymbol{\Sigma} = \begin{pmatrix} 4 & -3 & 0 \\ -3 & 6 & 0 \\ 0 & 0 & 5 \end{pmatrix}.$$

Which of the following random variables are independent?

- (a) y_1 and y_2 (d) (y_1, y_2) and y_3
 (b) y_1 and y_3 (e) (y_1, y_3) and y_2
 (c) y_2 and y_3

4.15 Suppose \mathbf{y} is $N_4(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, with

$$\boldsymbol{\mu} = \begin{pmatrix} -4 \\ 2 \\ 5 \\ -1 \end{pmatrix}, \quad \boldsymbol{\Sigma} = \begin{pmatrix} 8 & 0 & -1 & 0 \\ 0 & 3 & 0 & 2 \\ -1 & 0 & 5 & 0 \\ 0 & 2 & 0 & 7 \end{pmatrix}.$$

Which of the following random variables are independent?

- | | | |
|---------------------|----------------------------|-----------------------------------|
| (a) y_1 and y_2 | (f) y_3 and y_4 | (k) y_1 and y_2 and y_3 |
| (b) y_1 and y_3 | (g) (y_1, y_2) and y_3 | (l) y_1 and y_2 and y_4 |
| (c) y_1 and y_4 | (h) (y_1, y_2) and y_4 | (m) (y_2, y_2) and (y_3, y_4) |
| (d) y_2 and y_3 | (i) (y_1, y_3) and y_4 | (n) (y_1, y_3) and (y_2, y_4) |
| (e) y_2 and y_4 | (j) y_1 and (y_2, y_4) | |

4.16 Assume \mathbf{y} and \mathbf{x} are subvectors, each 2×1 , where $\begin{pmatrix} \mathbf{y} \\ \mathbf{x} \end{pmatrix}$ is $N_4(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ with

$$\boldsymbol{\mu} = \begin{pmatrix} 2 \\ -1 \\ 3 \\ 1 \end{pmatrix}, \quad \boldsymbol{\Sigma} = \left(\begin{array}{cc|cc} 7 & 3 & -3 & 2 \\ 3 & 6 & 0 & 4 \\ \hline -3 & 0 & 5 & -2 \\ 2 & 4 & -2 & 4 \end{array} \right).$$

- (a) Find $E(\mathbf{y}|\mathbf{x})$ by (4.7).
(b) Find $\text{cov}(\mathbf{y}|\mathbf{x})$ by (4.8).

4.17 Suppose \mathbf{y} and \mathbf{x} are subvectors, such that \mathbf{y} is 2×1 and \mathbf{x} is 3×1 , with $\boldsymbol{\mu}$ and $\boldsymbol{\Sigma}$ partitioned accordingly:

$$\boldsymbol{\mu} = \begin{pmatrix} 3 \\ -2 \\ 4 \\ -3 \\ 5 \end{pmatrix}, \quad \boldsymbol{\Sigma} = \left(\begin{array}{cc|ccc} 14 & -8 & 15 & 0 & 3 \\ -8 & 18 & 8 & 6 & -2 \\ \hline 15 & 8 & 50 & 8 & 5 \\ 0 & 6 & 8 & 4 & 0 \\ 3 & -2 & 5 & 0 & 1 \end{array} \right).$$

Assume that $\begin{pmatrix} \mathbf{y} \\ \mathbf{x} \end{pmatrix}$ is distributed as $N_5(\boldsymbol{\mu}, \boldsymbol{\Sigma})$.

- (a) Find $E(\mathbf{y}|\mathbf{x})$ by (4.7).
(b) Find $\text{cov}(\mathbf{y}|\mathbf{x})$ by (4.8).

4.18 Suppose that $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n$ is a random sample from a nonnormal multivariate population with mean $\boldsymbol{\mu}$ and covariance matrix $\boldsymbol{\Sigma}$. If n is large, what is the approximate distribution of each of the following?

- (a) $\sqrt{n}(\bar{\mathbf{y}} - \boldsymbol{\mu})$
(b) $\bar{\mathbf{y}}$

- 4.19** For the ramus bone data treated in Example 4.6.2, check each of the four variables for univariate normality using the following techniques:
- (a) $Q-Q$ plots
 - (b) $\sqrt{b_1}$ and b_2 as given by (4.18) and (4.19)
 - (c) D'Agostino's test using D and Y given in (4.22) and (4.23)
 - (d) The test by Lin and Mudholkar using z defined in (4.24)
- 4.20** For the calcium data in Table 3.4, check for multivariate normality and outliers using the following tests:
- (a) Calculate D_i^2 as in (4.27) for each observation.
 - (b) Compare the largest value of D_i^2 with the critical value in Table A.6.
 - (c) Compute u_i and v_i in (4.28) and (4.29) and plot them. Is there an indication of nonlinearity or outliers?
 - (d) Calculate $b_{1,p}$ and $b_{2,p}$ in (4.36) and (4.37) and compare them with critical values in Table A.5.
- 4.21** For the probe word data in Table 3.6, check each of the five variables for univariate normality and outliers using the following tests:
- (a) $Q-Q$ plots
 - (b) $\sqrt{b_1}$ and b_2 as given by (4.18) and (4.19)
 - (c) D'Agostino's test using D and Y given in (4.22) and (4.23)
 - (d) The test by Lin and Mudholkar using z defined in (4.24)
- 4.22** For the probe word data in Table 3.6, check for multivariate normality and outliers using the following tests:
- (a) Calculate D_i^2 as in (4.27) for each observation.
 - (b) Compare the largest value of D_i^2 with the critical value in Table A.6.
 - (c) Compute u_i and v_i in (4.28) and (4.29) and plot them. Is there an indication of nonlinearity or outliers?
 - (d) Calculate $b_{1,p}$ and $b_{2,p}$ in (4.36) and (4.37) and compare them with critical values in Table A.5.
- 4.23** Six hematology variables were measured on 51 workers (Royston 1983):
- | | |
|-----------------------------------------|-----------------------------------------|
| $y_1 = \text{hemoglobin concentration}$ | $y_4 = \text{lymphocyte count}$ |
| $y_2 = \text{packed cell volume}$ | $y_5 = \text{neutrophil count}$ |
| $y_3 = \text{white blood cell count}$ | $y_6 = \text{serum lead concentration}$ |

Table 4.2 Hematology Data

Observation Number	y_1	y_2	y_3	y_4	y_5	y_6
1	13.4	39	4100	14	25	17
2	14.6	46	5000	15	30	20
3	13.5	42	4500	19	21	18
4	15.0	46	4600	23	16	18
5	14.6	44	5100	17	31	19
6	14.0	44	4900	20	24	19
7	16.4	49	4300	21	17	18
8	14.8	44	4400	16	26	29
9	15.2	46	4100	27	13	27
10	15.5	48	8400	34	42	36
11	15.2	47	5600	26	27	22
12	16.9	50	5100	28	17	23
13	14.8	44	4700	24	20	23
14	16.2	45	5600	26	25	19
15	14.7	43	4000	23	13	17
16	14.7	42	3400	9	22	13
17	16.5	45	5400	18	32	17
18	15.4	45	6900	28	36	24
19	15.1	45	4600	17	29	17
20	14.2	46	4200	14	25	28
21	15.9	46	5200	8	34	16
22	16.0	47	4700	25	14	18
23	17.4	50	8600	37	39	17
24	14.3	43	5500	20	31	19
25	14.8	44	4200	15	24	29
26	14.9	43	4300	9	32	17
27	15.5	45	5200	16	30	20
28	14.5	43	3900	18	18	25
29	14.4	45	6000	17	37	23
30	14.6	44	4700	23	21	27
31	15.3	45	7900	43	23	23
:	:	:	:	:	:	:
51	14.7	44	7800	38	34	16

The data are given in Table 4.2. Check each of the six variables for univariate normality using the following tests:

- (a) $Q-Q$ plots
- (b) $\sqrt{b_1}$ and b_2 as given by (4.18) and (4.19)
- (c) D'Agostino's test using D and Y given in (4.22) and (4.23)
- (d) The test by Lin and Mudholkar using z defined in (4.24)

4.24 For the hematology data in Table 4.2, check for multivariate normality using the following techniques:

- Calculate D_i^2 as in (4.27) for each observation.
- Compare the largest value of D_i^2 with the critical value in Table A.6 (extrapolate).
- Compute u_i and v_i in (4.28) and (4.29) and plot them. Is there an indication of nonlinearity or outliers?
- Calculate $b_{1,p}$ and $b_{2,p}$ in (4.36) and (4.37) and compare them with critical values in Table A.5.

4.25 Use the glucose data in Table 3.9.

- Use the methods in Section 4.5.1 to find the optimal univariate transformation to normality for each of the glucose measurements obtained one hour after sugar intake (x_1 , x_2 , and x_3).
- Use the methods in Section 4.5.2 to find the optimal multivariate transformation to 3-variate normality for the glucose measurements obtained one hour after sugar intake (x_1 , x_2 , and x_3).
- How do the transformations obtained from the two approaches compare?

4.26 Use the Baker corn field data in Table 3.3.

- Use the methods in Section 4.5.1 to find the optimal univariate transformation to normality for the B, Ca, and Cu measurements.
- Use the methods in Section 4.5.2 to find the optimal multivariate transformation to 3-variate normality for the B, Ca, and Cu measurements.
- How do the transformations obtained from the two approaches compare?

CHAPTER 5

TESTS ON ONE OR TWO MEAN VECTORS

5.1 MULTIVARIATE VERSUS UNIVARIATE TESTS

Hypothesis testing in a multivariate context is more complex than in a univariate setting. The number of parameters may be staggering. The p -variate normal distribution, for example, has p means, p variances, and $\binom{p}{2}$ covariances, where $\binom{p}{2}$ represents the number of pairs among the p variables. The total number of parameters is

$$p + p + \binom{p}{2} = \frac{1}{2}p(p + 3).$$

For $p = 10$, for example, the number of parameters is 65, for each of which a hypothesis could be formulated. Additionally, we might be interested in testing hypotheses about subsets of these parameters or about functions of them. In some cases, we have the added dilemma of choosing among competing test statistics (see Chapter 6).

We first discuss the motivation for testing p variables multivariately rather than (or in addition to) univariately, as for example, in hypotheses about $\mu_1, \mu_2, \dots, \mu_p$ in μ . There are at least four arguments for a multivariate approach to hypothesis testing:

1. The use of p univariate tests inflates the Type I error rate, α , whereas the multivariate test preserves the exact α level. For example, if we do $p = 10$ separate univariate tests at the .05 level, the probability of at least one false rejection is greater than .05. If the variables were independent (they rarely are), we would have (under H_0)

$$\begin{aligned} P(\text{at least one rejection}) &= 1 - P(\text{all 10 tests accept } H_0) \\ &= 1 - (.95)^{10} = .40. \end{aligned}$$

The resulting overall α of .40 is not an acceptable error rate. Typically, the 10 variables are correlated, and the overall α would lie somewhere between .05 and .40.

2. The univariate tests completely ignore the correlations among the variables, whereas the multivariate tests make direct use of the correlations.
3. The multivariate test is more powerful in many cases. The *power* of a test is the probability of rejecting H_0 when it is false. In some cases, all p of the univariate tests fail to reach significance, but the multivariate test is significant because small effects on some of the variables combine to jointly indicate significance. However, for a given sample size, there is a limit to the number of variables a multivariate test can handle without losing power. This is discussed further in Section 5.3.2.
4. Many multivariate tests involving means have as a by-product the construction of a linear combination of variables that reveals more about how the variables unite to reject the hypothesis.

5.2 TESTS ON μ WITH Σ KNOWN

The test on a mean vector assuming a known Σ is introduced to illustrate the issues involved in multivariate testing and to serve as a foundation for the unknown Σ case. We first review the univariate case, in which we work with a single variable y that is distributed as $N(\mu, \sigma^2)$.

5.2.1 Review of Univariate Test for $H_0: \mu = \mu_0$ with σ Known

The hypothesis of interest is that the mean of y is equal to a given value, μ_0 , versus the alternative that it is not equal to μ_0 :

$$H_0: \mu = \mu_0 \quad \text{vs.} \quad H_1: \mu \neq \mu_0.$$

We do not consider one-sided alternative hypotheses because they do not readily generalize to multivariate tests. We assume a random sample of n observations

y_1, y_2, \dots, y_n from $N(\mu, \sigma^2)$ with σ^2 known. We calculate $\bar{y} = \sum_{i=1}^n y_i/n$ and compare it to μ_0 using the test statistic

$$z = \frac{\bar{y} - \mu_0}{\sigma_{\bar{y}}} = \frac{\bar{y} - \mu_0}{\sigma/\sqrt{n}}, \quad (5.1)$$

which is distributed as $N(0, 1)$ if H_0 is true. For $\alpha = .05$, we reject H_0 if $|z| \geq 1.96$. Equivalently, we can use z^2 , which is distributed as χ^2 with one degree of freedom, and reject H_0 if $z^2 \geq (1.96)^2 = 3.84$. If n is large, we are assured by the central limit theorem that z is approximately normal, even if the observations are not from a normal distribution.

5.2.2 Multivariate Test for $H_0: \mu = \mu_0$ with Σ Known

In the multivariate case we have several variables measured on each sampling unit, and we wish to hypothesize a value for the mean of each variable, $H_0: \mu = \mu_0$ vs. $H_1: \mu \neq \mu_0$. More explicitly, we have

$$H_0: \begin{pmatrix} \mu_1 \\ \mu_2 \\ \vdots \\ \mu_p \end{pmatrix} = \begin{pmatrix} \mu_{01} \\ \mu_{02} \\ \vdots \\ \mu_{0p} \end{pmatrix}, \quad H_1: \begin{pmatrix} \mu_1 \\ \mu_2 \\ \vdots \\ \mu_p \end{pmatrix} \neq \begin{pmatrix} \mu_{01} \\ \mu_{02} \\ \vdots \\ \mu_{0p} \end{pmatrix},$$

where each μ_{0j} is specified from previous experience or is a target value. The vector equality in H_0 implies $\mu_j = \mu_{0j}$ for all $j = 1, 2, \dots, p$. The vector inequality in H_1 implies at least one $\mu_j \neq \mu_{0j}$. Thus, for example, if $\mu_j = \mu_{0j}$ for all j except 2, for which $\mu_2 \neq \mu_{02}$, then we wish to reject H_0 .

To test H_0 , we use a random sample of n observation vectors $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n$ from $N_p(\mu, \Sigma)$, with Σ known, and calculate $\bar{\mathbf{y}} = \sum_{i=1}^n \mathbf{y}_i/n$. The test statistic is

$$Z^2 = n(\bar{\mathbf{y}} - \mu_0)' \Sigma^{-1} (\bar{\mathbf{y}} - \mu_0). \quad (5.2)$$

If H_0 is true, Z^2 is distributed as χ_p^2 by (4.6), and we therefore reject H_0 if $Z^2 > \chi_{\alpha,p}^2$. Note that for one variable, z^2 [the square of (5.1)] has a chi-square distribution with one degree of freedom, whereas for p variables, Z^2 in (5.2) is distributed as a chi-square with p degrees of freedom.

If Σ is unknown, we could use \mathbf{S} in its place in (5.2), and Z^2 would have an approximate χ^2 -distribution. But n would have to be larger than in the analogous univariate situation, in which $t = (\bar{y} - \mu_0)/(s/\sqrt{n})$ is approximately $N(0, 1)$ for $n > 30$. The value of n needed for $n(\bar{\mathbf{y}} - \mu_0)' \mathbf{S}^{-1} (\bar{\mathbf{y}} - \mu_0)$ to have an approximate χ^2 -distribution depends on p . This is clarified further in Section 5.3.2.

■ EXAMPLE 5.2.2

In Table 3.1, height and weight were given for a sample of 20 college-age males. Let us assume that this sample originated from the bivariate normal




Figure 5.1 Elliptical acceptance region.

$N_2(\boldsymbol{\mu}, \Sigma)$, where

$$\Sigma = \begin{pmatrix} 20 & 100 \\ 100 & 1000 \end{pmatrix}.$$

Suppose we wish to test $H_0: \boldsymbol{\mu} = (70, 170)'$. From Example 3.2.1, $\bar{y}_1 = 71.45$ and $\bar{y}_2 = 164.7$. We thus have

$$\begin{aligned} Z^2 &= n(\bar{y} - \boldsymbol{\mu}_0)' \Sigma^{-1} (\bar{y} - \boldsymbol{\mu}_0) \\ &= (20) \begin{pmatrix} 71.45 - 70 \\ 164.7 - 170 \end{pmatrix}' \begin{pmatrix} 20 & 100 \\ 100 & 1000 \end{pmatrix}^{-1} \begin{pmatrix} 71.45 - 70 \\ 164.7 - 170 \end{pmatrix} \\ &= (20)(1.45, -5.3) \begin{pmatrix} .1 & -.01 \\ -.01 & .002 \end{pmatrix} \begin{pmatrix} 1.45 \\ -5.3 \end{pmatrix} = 8.4026. \end{aligned}$$

Using $\alpha = .05$, $\chi^2_{.05, 2} = 5.99$, and we therefore reject $H_0: \boldsymbol{\mu} = (70, 170)'$ because $Z^2 = 8.4026 > 5.99$.

The rejection region for $\bar{y} = (\bar{y}_1, \bar{y}_2)'$ is on or outside the ellipse in Figure 5.1; that is, the test statistic Z^2 is greater than 5.99 if and only if \bar{y} is outside the ellipse. If \bar{y} falls inside the ellipse, H_0 is accepted. Thus distance from $\boldsymbol{\mu}_0$ as well as direction must be taken into account. When the distance is standardized by Σ^{-1} , all points on the curve are “statistically equidistant” from the center.

Note that the test is sensitive to the covariance structure. If $\text{cov}(y_1, y_2)$ were negative, y_2 would tend to decrease as y_1 increases, and the ellipse would be tilted in the other direction. In this case, \bar{y} would be in the acceptance region.




Figure 5.2 Acceptance and rejection regions for univariate and multivariate tests.

Let us now investigate the consequence of testing each variable separately. Using $z_{\alpha/2} = 1.96$ for $\alpha = .05$, we have

$$z_1 = \frac{\bar{y}_1 - \mu_{01}}{\sigma_1 / \sqrt{n}} = 1.450 < 1.96,$$

$$z_2 = \frac{\bar{y}_2 - \mu_{02}}{\sigma_2 / \sqrt{n}} = -.7495 > -1.96.$$

Thus both tests accept the hypothesis. In this case neither of the \bar{y} 's is far enough from the hypothesized value to cause rejection. But when the positive correlation between y_1 and y_2 is taken into account in the multivariate test, the two evidences against μ_0 combine to cause rejection. This illustrates the third advantage of multivariate tests given in Section 5.1.

Figure 5.2 shows the rectangular acceptance region for the univariate tests superimposed on the elliptical multivariate acceptance region. The rectangle was obtained by calculating the two acceptance regions

$$\mu_{01} - 1.96 \frac{\sigma_1}{\sqrt{n}} < \bar{y}_1 < \mu_{01} + 1.96 \frac{\sigma_1}{\sqrt{n}},$$

$$\mu_{02} - 1.96 \frac{\sigma_2}{\sqrt{n}} < \bar{y}_2 < \mu_{02} + 1.96 \frac{\sigma_2}{\sqrt{n}}.$$

Points inside the ellipse but outside the rectangle will be rejected in at least one univariate dimension but will be accepted multivariately. This illustrates the inflation of α resulting from univariate tests, as discussed in the first motive

for multivariate testing in Section 5.1. This phenomenon has been referred to as Rao's paradox. For further discussion see Rao (1966), Healy (1969), and Morrison (1990, p. 174). Points outside the ellipse but inside the rectangle will be rejected multivariately but accepted univariately in both dimensions. This illustrates the third reason for multivariate testing given in Section 5.1, namely, that the multivariate test is more powerful in some situations.

Thus in either case represented by the shaded areas, we should use the multivariate test result, not the univariate results. In the one case, the multivariate test is more powerful than the univariate tests; in the other case, the multivariate test preserves α whereas the univariate tests inflate α . Consequently, when the multivariate and univariate results disagree, our tendency is to trust the multivariate result. In Section 5.5, we discuss various procedures for ascertaining the contribution of the individual variables after the multivariate test has rejected the hypothesis. \square

5.3 TESTS ON μ WHEN Σ IS UNKNOWN

In Section 5.2, we said little about properties of the tests, because the tests discussed were of slight practical consequence because of the assumption that Σ is known. We will be more concerned with test properties in Sections 5.3 and 5.4, first in the one-sample case and then in the two-sample case. The reader may wonder why we include one-sample tests, since we seldom, if ever, have need of a test for $H_0: \mu = \mu_0$. However, we will cover this case for two reasons:

1. Many general principles are more easily illustrated in the one-sample framework than in the two-sample case.
2. Some very useful tests can be cast in the one-sample framework. Two examples are (1) $H_0: \mu_d = 0$ used in the paired comparison test covered in Section 5.7 and (2) $H_0: C\mu = 0$ used in profile analysis in Section 5.9, in analysis of repeated measures in Section 6.9, and in growth curves in Section 6.10.

5.3.1 Review of Univariate t -Test for $H_0: \mu = \mu_0$ with σ Unknown

We first review the familiar one-sample t -test in the univariate case, with only one variable measured on each sampling unit. We assume that a random sample y_1, y_2, \dots, y_n is available from $N(\mu, \sigma^2)$. We estimate μ by \bar{y} and σ^2 by s^2 , where \bar{y} and s^2 are given by (3.1) and (3.4). To test $H_0: \mu = \mu_0$ vs. $H_1: \mu \neq \mu_0$, we use

$$t = \frac{\bar{y} - \mu_0}{s/\sqrt{n}} = \frac{\sqrt{n}(\bar{y} - \mu_0)}{s}. \quad (5.3)$$

If H_0 is true, t is distributed as t_{n-1} , where $n - 1$ is the degrees of freedom. We reject H_0 if $|\sqrt{n}(\bar{y} - \mu_0)/s| \geq t_{\alpha/2, n-1}$, where $t_{\alpha/2, n-1}$ is a critical value from the t -table.

The first expression in (5.3), $t = (\bar{y} - \mu_0)/(s/\sqrt{n})$, is the *characteristic form* of the t -statistic, which represents a sample standardized distance between \bar{y} and μ_0 . In this form, the hypothesized mean is subtracted from \bar{y} and the difference is divided by $s_{\bar{y}} = s/\sqrt{n}$. Since y_1, y_2, \dots, y_n is a random sample from $N(\mu, \sigma^2)$, the random variables \bar{y} and s are independent. We will see an analogous characteristic form for the T^2 -statistic in the multivariate case in Section 5.3.2.

5.3.2 Hotelling's T^2 -Test for $H_0: \mu = \mu_0$ with Σ Unknown

We now move to the multivariate case in which p variables are measured on each sampling unit. We assume that a random sample $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n$ is available from $N_p(\mu, \Sigma)$, where \mathbf{y}_i contains the p measurements on the i th sampling unit (subject or object). We estimate μ by $\bar{\mathbf{y}}$ and Σ by \mathbf{S} , where $\bar{\mathbf{y}}$ and \mathbf{S} are given by (3.16), (3.19), (3.22), (3.27), and (3.29). In order to test $H_0: \mu = \mu_0$ versus $H_1: \mu \neq \mu_0$, we use an extension of the univariate t -statistic in (5.3). In squared form, the univariate t can be rewritten as

$$t^2 = \frac{n(\bar{y} - \mu_0)^2}{s^2} = n(\bar{y} - \mu_0)(s^2)^{-1}(\bar{y} - \mu_0). \quad (5.4)$$

When $\bar{y} - \mu_0$ and s^2 are replaced by $\bar{\mathbf{y}} - \mu_0$ and \mathbf{S} , we obtain the test statistic

$$T^2 = n(\bar{\mathbf{y}} - \mu_0)' \mathbf{S}^{-1} (\bar{\mathbf{y}} - \mu_0). \quad (5.5)$$

Alternatively, T^2 can be obtained from Z^2 in (5.2) by replacing Σ with \mathbf{S} .

The distribution of T^2 was obtained by Hotelling (1931), assuming H_0 is true and sampling is from $N_p(\mu, \Sigma)$. The distribution is indexed by two parameters, the dimension p and degrees of freedom $\nu = n - 1$. We reject H_0 if $T^2 > T_{\alpha, p, n-1}^2$ and accept H_0 otherwise. Critical values of the T^2 -distribution are found in Table A.7, taken from Kramer and Jensen (1969a).

Note that the terminology “accept H_0 ” is used for expositional convenience to describe our decision when we do not reject the hypothesis. Strictly speaking, we do not accept H_0 in the sense of actually believing it is true. If the sample size were extremely large and we accepted H_0 , we could be reasonably certain that the true μ is close to the hypothesized value μ_0 . Otherwise, “accepting H_0 ” means only that we have failed to reject H_0 .

The T^2 -statistic can be viewed as the sample standardized distance between the observed sample mean vector and the hypothetical mean vector. If the sample mean vector is notably distant from the hypothetical mean vector, we become suspicious of the hypothetical mean vector and wish to reject H_0 .

The test statistic is a scalar quantity, since $T^2 = n(\bar{\mathbf{y}} - \mu_0)' \mathbf{S}^{-1} (\bar{\mathbf{y}} - \mu_0)$ is a quadratic form. As with the χ^2 -distribution of Z^2 , the density of T^2 is skewed because the lower limit is zero and there is no upper limit.

The *characteristic form* of the T^2 -statistic (5.5) is

$$T^2 = (\bar{\mathbf{y}} - \mu_0)' \left(\frac{\mathbf{S}}{n} \right)^{-1} (\bar{\mathbf{y}} - \mu_0). \quad (5.6)$$

The characteristic form has two features:

1. S/n is the sample covariance matrix of \bar{y} and serves as a standardizing matrix in the distance function.
2. Since y_1, y_2, \dots, y_n are distributed as $N_p(\mu, \Sigma)$, it follows that \bar{y} is $N_p(\mu, \frac{1}{n} \Sigma)$, $(n - 1)S$ is $W(n - 1, \Sigma)$, and \bar{y} and S are independent (see Section 4.3.2).

In (5.3), the univariate t -statistic represents the number of standard deviations \bar{y} is separated from μ_0 . In appearance, the T^2 -statistic (5.6) is similar, but no such simple interpretation is possible. If we add a variable, the distance in (5.6) increases. (By analogy, the hypotenuse of a right triangle is longer than either of the legs.) Thus we need a test statistic that indicates the significance of the distance from \bar{y} to μ_0 , while allowing for the number of dimensions (see comment 3 below about the T^2 -table). Since the resulting T^2 -statistic cannot be readily interpreted in terms of the number of standard deviations \bar{y} is from μ_0 , we do not have an intuitive feel for its significance as we do with the univariate t . We must compare the calculated value of T^2 with the table value. In addition, the T^2 -table provides some insights into the behavior of the T^2 -distribution. Four of these insights are noted at the end of this section.

If a test leads to rejection of $H_0: \mu = \mu_0$, the question arises as to which variable or variables contributed most to the rejection. This issue is discussed in Section 5.5 for the two-sample T^2 -test of $H_0: \mu_1 = \mu_2$, and the results there can be easily adapted to the one-sample test of $H_0: \mu = \mu_0$. For confidence intervals on the individual μ_j 's in μ , see Rencher (1998, Section 3.4).

The following are some key properties of the T^2 -test:

1. We must have $n > p$. Otherwise, S is singular and T^2 cannot be computed.
2. In both the one-sample and two-sample cases, the degrees of freedom for the T^2 -statistic will be the same as for the analogous univariate t -test, that is, $\nu = n - 1$ for one sample and $\nu = n_1 + n_2 - 2$ for two samples (see Section 5.4.2).
3. The alternative hypothesis is two-sided. Because the space is multidimensional, we do not consider one-sided alternative hypotheses, such as $\mu > \mu_0$. However, even though the alternative hypothesis $H_1: \mu \neq \mu_0$ is essentially two-sided, the critical region is one-tailed (we reject for large values of T^2). This is typical of many multivariate tests.
4. In the univariate case, $t_{n-1}^2 = F_{1,n-1}$. The statistic T^2 can also be converted to an F -statistic as follows:

$$\frac{\nu - p + 1}{\nu p} T_{p,\nu}^2 = F_{p,\nu-p+1}. \quad (5.7)$$

Note that the dimension p (number of variables) of the T^2 -statistic becomes the first of the two degrees-of-freedom parameters of the F . The degrees of

freedom for T^2 is denoted by ν , and the F -transformation is given in terms of a general ν , since other applications of T^2 will have ν different from $n - 1$ (see, for example, Sections 5.4.2 and 6.3.2).

Equation (5.7) gives an easy way to find critical values for the T^2 -test. However, we have provided critical values of T^2 in Table A.7 because of the insights they provide into the behavior of the T^2 -distribution in particular and multivariate tests in general. The following are some insights that can readily be gleaned from the T^2 -tables:

1. The first column of Table A.7 contains squared values from the t table; that is, $T_{\alpha,1,\nu}^2 = t_{\alpha/2,\nu}^2$. (We use $t_{\alpha/2,\nu}^2$ because the univariate test of $H_0: \mu = \mu_0$ versus $H_1: \mu \neq \mu_0$ is two-tailed.) Thus for $p = 1$, T^2 reduces to t^2 . This can easily be seen by comparing (5.5) with (5.4).
2. The last row of each page of Table A.7 contains χ^2 critical values, that is, $T_{p,\infty}^2 = \chi_p^2$. Thus as n increases, \mathbf{S} approaches Σ , and

$$T^2 = n(\bar{\mathbf{y}} - \boldsymbol{\mu}_0)' \mathbf{S}^{-1} (\bar{\mathbf{y}} - \boldsymbol{\mu}_0)$$

approaches $Z^2 = n(\bar{\mathbf{y}} - \boldsymbol{\mu}_0)' \boldsymbol{\Sigma}^{-1} (\bar{\mathbf{y}} - \boldsymbol{\mu}_0)$ in (5.2), which is distributed as χ_p^2 .

3. The values increase along each row of Table A.7; that is, for a fixed ν , the critical value $T_{\alpha,p,\nu}^2$ increases with p . It was noted above that in any given sample, the calculated value of T^2 increases if a variable is added. However, since the critical value also increases, a variable should not be added unless it adds a significant amount to T^2 .
4. As p increases, larger values of ν are required for the distribution of T^2 to approach χ^2 . In the univariate case, t in (5.3) is considered a good approximation to the standard normal z in (5.1) when $\nu = n - 1$ is at least 30. In the first column ($p = 1$) of Table A.7, we see $T_{.05,1,30}^2 = 4.171$ and $T_{.05,1,\infty}^2 = 3.841$, with a ratio of $4.171/3.841 = 1.086$. For $p = 5$, ν must be 100 to obtain the same ratio: $T_{.05,5,100}^2/T_{.05,5,\infty}^2 = 1.086$. For $p = 10$, we need $\nu = 200$ to obtain a similar value of the ratio: $T_{.05,10,200}^2/T_{.05,10,\infty}^2 = 1.076$. Thus one must be very cautious in stating that T^2 has an approximate χ^2 -distribution for large n . The α level (Type I error rate) could be substantially inflated. For example, suppose $p = 10$ and we assume that $n = 30$ is sufficiently large for a χ^2 -approximation to hold. Then we would reject H_0 for $T^2 \geq 18.307$ with a target α -level of .05. However, the correct critical value is 34.044, and the misuse of 18.307 would yield an actual α of $P(T_{10,29}^2 \geq 18.307) = .314$.

■ EXAMPLE 5.3.2

In Table 3.4 we have $n = 10$ observations on $p = 3$ variables. Desirable levels for y_1 and y_2 are 15.0 and 6.0, respectively, and the expected level of y_3 is

2.85. We can, therefore, test the hypothesis

$$H_0: \boldsymbol{\mu} = \begin{pmatrix} 15.0 \\ 6.0 \\ 2.85 \end{pmatrix}.$$

In Examples 3.6 and 3.7, $\bar{\mathbf{y}}$ and \mathbf{S} were obtained as

$$\bar{\mathbf{y}} = \begin{pmatrix} 28.1 \\ 7.18 \\ 3.09 \end{pmatrix}, \quad \mathbf{S} = \begin{pmatrix} 140.54 & 49.68 & 1.94 \\ 49.68 & 72.25 & 3.68 \\ 1.94 & 3.68 & .25 \end{pmatrix}.$$

To test H_0 , we use (5.5):

$$\begin{aligned} T^2 &= n(\bar{\mathbf{y}} - \boldsymbol{\mu}_0)' \mathbf{S}^{-1} (\bar{\mathbf{y}} - \boldsymbol{\mu}_0) \\ &= 10 \begin{pmatrix} 28.1 - 15.0 \\ 7.18 - 6.0 \\ 3.09 - 2.85 \end{pmatrix}' \begin{pmatrix} 140.54 & 49.68 & 1.94 \\ 49.68 & 72.25 & 3.68 \\ 1.94 & 3.68 & .25 \end{pmatrix}^{-1} \begin{pmatrix} 28.1 - 15.0 \\ 7.18 - 6.0 \\ 3.09 - 2.85 \end{pmatrix} \\ &= 24.559. \end{aligned}$$

From Table A.7, we obtain the critical value $T_{.05,3,9}^2 = 16.766$. Since the observed value of T^2 exceeds the critical value, we reject the hypothesis. \square

5.4 COMPARING TWO MEAN VECTORS

We first review the univariate two-sample t -test and then proceed with the analogous multivariate test.

5.4.1 Review of Univariate Two-Sample t -Test

In the one-variable case we obtain a random sample $y_{11}, y_{12}, \dots, y_{1n_1}$ from $N(\mu_1, \sigma_1^2)$ and a second random sample $y_{21}, y_{22}, \dots, y_{2n_2}$ from $N(\mu_2, \sigma_2^2)$. We assume that the two samples are independent and that $\sigma_1^2 = \sigma_2^2 = \sigma^2$, say, with σ^2 unknown. [The assumptions of independence and equal variances are necessary in order for the t -statistic below in (5.8) to have a t -distribution.] From the two samples we calculate $\bar{y}_1, \bar{y}_2, SS_1 = \sum_{i=1}^{n_1} (y_{1i} - \bar{y}_1)^2 = (n_1 - 1)s_1^2, SS_2 = \sum_{i=1}^{n_2} (y_{2i} - \bar{y}_2)^2 = (n_2 - 1)s_2^2$, and the pooled variance

$$s_{\text{pl}}^2 = \frac{SS_1 + SS_2}{n_1 + n_2 - 2} = \frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2},$$

where $n_1 + n_2 - 2$ is the sum of the weights $n_1 - 1$ and $n_2 - 1$ in the numerator. With this denominator, s_{pl}^2 is an unbiased estimator for the common variance, σ^2 , that is, $E(s_{\text{pl}}^2) = \sigma^2$.

To test

$$H_0: \mu_1 = \mu_2 \quad \text{vs.} \quad H_1: \mu_1 \neq \mu_2,$$

we use

$$t = \frac{\bar{y}_1 - \bar{y}_2}{s_{\text{pl}} \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}, \quad (5.8)$$

which has a t -distribution with $n_1 + n_2 - 2$ degrees of freedom when H_0 is true. We therefore reject H_0 if $|t| \geq t_{\alpha/2, n_1+n_2-2}$.

Note that (5.8) exhibits the *characteristic form* of a t -statistic. In this form, the denominator is the sample standard deviation of the numerator, that is,

$$s_{\text{pl}} \sqrt{1/n_1 + 1/n_2}$$

is an estimate of

$$\begin{aligned} \sigma_{\bar{y}_1 - \bar{y}_2} &= \sqrt{\text{var}(\bar{y}_1 - \bar{y}_2)} = \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}} \\ &= \sqrt{\frac{\sigma^2}{n_1} + \frac{\sigma^2}{n_2}} = \sigma \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}. \end{aligned}$$

5.4.2 Multivariate Two-Sample T^2 -Test

We now consider the case where p variables are measured on each sampling unit in two samples. We wish to test

$$H_0: \mu_1 = \mu_2 \quad \text{vs.} \quad H_1: \mu_1 \neq \mu_2.$$

We obtain a random sample $\mathbf{y}_{11}, \mathbf{y}_{12}, \dots, \mathbf{y}_{1n_1}$ from $N_p(\mu_1, \Sigma_1)$ and a second random sample $\mathbf{y}_{21}, \mathbf{y}_{22}, \dots, \mathbf{y}_{2n_2}$ from $N_p(\mu_2, \Sigma_2)$. We assume that the two samples are independent and that $\Sigma_1 = \Sigma_2 = \Sigma$, say, with Σ unknown. These assumptions are necessary in order for the T^2 -statistic in (5.9) to have a T^2 -distribution. A test of $H_0: \Sigma_1 = \Sigma_2$ is given in Section 7.3.2. For an approximate test of $H_0: \mu_1 = \mu_2$ that can be used when $\Sigma_1 \neq \Sigma_2$, see Rencher (1998, Section 3.10).

The sample mean vectors are $\bar{\mathbf{y}}_1 = \sum_{i=1}^{n_1} \mathbf{y}_{1i}/n_1$ and $\bar{\mathbf{y}}_2 = \sum_{i=1}^{n_2} \mathbf{y}_{2i}/n_2$. Define \mathbf{W}_1 and \mathbf{W}_2 to be the matrices of sums of squares and cross products for the two samples:

$$\begin{aligned} \mathbf{W}_1 &= \sum_{i=1}^{n_1} (\mathbf{y}_{1i} - \bar{\mathbf{y}}_1)(\mathbf{y}_{1i} - \bar{\mathbf{y}}_1)' = (n_1 - 1)\mathbf{S}_1, \\ \mathbf{W}_2 &= \sum_{i=1}^{n_2} (\mathbf{y}_{2i} - \bar{\mathbf{y}}_2)(\mathbf{y}_{2i} - \bar{\mathbf{y}}_2)' = (n_2 - 1)\mathbf{S}_2. \end{aligned}$$

Since $(n_1 - 1)\mathbf{S}_1$ is an unbiased estimator of $(n_1 - 1)\Sigma$ and $(n_2 - 1)\mathbf{S}_2$ is an unbiased estimator of $(n_2 - 1)\Sigma$, we can pool them to obtain an unbiased estimator of the

common population covariance matrix, Σ :

$$\begin{aligned}\mathbf{S}_{\text{pl}} &= \frac{1}{n_1 + n_2 - 2} (\mathbf{W}_1 + \mathbf{W}_2) \\ &= \frac{1}{n_1 + n_2 - 2} [(n_1 - 1)\mathbf{S}_1 + (n_2 - 1)\mathbf{S}_2].\end{aligned}$$

Thus $E(\mathbf{S}_{\text{pl}}) = \Sigma$.

The square of the univariate t -statistic (5.8) can be expressed as

$$t^2 = \frac{n_1 n_2}{n_1 + n_2} (\bar{y}_1 - \bar{y}_2)(s_{\text{pl}}^2)^{-1} (\bar{y}_1 - \bar{y}_2).$$

This can be generalized to p variables by substituting $\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2$ for $\bar{y}_1 - \bar{y}_2$ and \mathbf{S}_{pl} for s_{pl}^2 to obtain

$$T^2 = \frac{n_1 n_2}{n_1 + n_2} (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{\text{pl}}^{-1} (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2), \quad (5.9)$$

which is distributed as $T_{p, n_1 + n_2 - 2}^2$ when $H_0: \mu_1 = \mu_2$ is true. To carry out the test, we collect the two samples, calculate T^2 by (5.9), and reject H_0 if $T^2 \geq T_{\alpha, p, n_1 + n_2 - 2}^2$. Critical values of T^2 are found in Table A.7. For tables of the power of the T^2 -test (probability of rejecting H_0 when it is false) and illustrations of their use, see Rencher (1998, Section 3.11).

The T^2 -statistic (5.9) can be expressed in *characteristic form* as the standardized distance between $\bar{\mathbf{y}}_1$ and $\bar{\mathbf{y}}_2$:

$$T^2 = (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \left[\left(\frac{1}{n_1} + \frac{1}{n_2} \right) \mathbf{S}_{\text{pl}} \right]^{-1} (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2), \quad (5.10)$$

where $(1/n_1 + 1/n_2)\mathbf{S}_{\text{pl}}$ is the sample covariance matrix for $\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2$ and \mathbf{S}_{pl} is independent of $\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2$ because of sampling from the multivariate normal. For a discussion of robustness of T^2 to departures from the assumptions of multivariate normality and homogeneity of covariance matrices ($\Sigma_1 = \Sigma_2$), see Rencher (1998, Section 3.8).

Some key properties of the two-sample T^2 -test are given in the following list:

1. It is necessary that $n_1 + n_2 - 2 > p$ for \mathbf{S}_{pl} to be nonsingular.
2. The statistic T^2 is, of course, a scalar. The $3p + p(p - 1)/2$ quantities in $\bar{\mathbf{y}}_1, \bar{\mathbf{y}}_2$, and \mathbf{S}_{pl} have been reduced to a single scale on which T^2 is large if the sample evidence favors $H_1: \mu_1 \neq \mu_2$ and small if the evidence supports $H_0: \mu_1 = \mu_2$; we reject H_0 if the standardized distance between $\bar{\mathbf{y}}_1$ and $\bar{\mathbf{y}}_2$ is large.
3. Since the lower limit of T^2 is zero and there is no upper limit, the density is skewed. In fact, as noted in (5.11) below, T^2 is directly related to F , which is a well-known skewed distribution.
4. For degrees of freedom of T^2 we have $n_1 + n_2 - 2$, which is the same as for the corresponding univariate t -statistic (5.8).

5. The alternative hypothesis $H_1: \mu_1 \neq \mu_2$ is two sided. The critical region $T^2 > T_{\alpha}^2$ is one-tailed, however, as is typical of many multivariate tests.
6. The T^2 -statistic can be readily transformed to an F -statistic using (5.7):

$$\frac{n_1 + n_2 - p - 1}{(n_1 + n_2 - 2)} T^2 = F_{p, n_1 + n_2 - p - 1}, \quad (5.11)$$

where again the dimension p of the T^2 -statistic becomes the first degree-of-freedom parameter for the F -statistic.

■ EXAMPLE 5.4.2

Four psychological tests were given to 32 men and 32 women. The data are recorded in Table 5.1 (Beall 1945). The variables are

$$\begin{array}{ll} y_1 = \text{pictorial inconsistencies} & y_3 = \text{tool recognition} \\ y_2 = \text{paper form board} & y_4 = \text{vocabulary} \end{array}$$

The mean vectors and covariance matrices of the two samples are

$$\bar{\mathbf{y}}_1 = \begin{pmatrix} 15.97 \\ 15.91 \\ 27.19 \\ 22.75 \end{pmatrix}, \quad \bar{\mathbf{y}}_2 = \begin{pmatrix} 12.34 \\ 13.91 \\ 16.66 \\ 21.94 \end{pmatrix},$$

$$\mathbf{S}_1 = \begin{pmatrix} 5.192 & 4.545 & 6.522 & 5.250 \\ 4.545 & 13.18 & 6.760 & 6.266 \\ 6.522 & 6.760 & 28.67 & 14.47 \\ 5.250 & 6.266 & 14.47 & 16.65 \end{pmatrix},$$

$$\mathbf{S}_2 = \begin{pmatrix} 9.136 & 7.549 & 4.864 & 4.151 \\ 7.549 & 18.60 & 10.22 & 5.446 \\ 4.864 & 10.22 & 30.04 & 13.49 \\ 4.151 & 5.446 & 13.49 & 28.00 \end{pmatrix}.$$

The sample covariance matrices do not appear to indicate a disparity in the population covariance matrices. (A significance test to check this assumption is carried out in Example 7.3.2, and the hypothesis $H_0: \Sigma_1 = \Sigma_2$ is not rejected.) The pooled covariance matrix is

$$\begin{aligned} \mathbf{S}_{\text{pl}} &= \frac{1}{32 + 32 - 2} [(32 - 1)\mathbf{S}_1 + (32 - 1)\mathbf{S}_2] \\ &= \begin{pmatrix} 7.164 & 6.047 & 5.693 & 4.701 \\ 6.047 & 15.89 & 8.492 & 5.856 \\ 5.693 & 8.492 & 29.36 & 13.98 \\ 4.701 & 5.856 & 13.98 & 22.32 \end{pmatrix}. \end{aligned}$$

Table 5.1 Four Psychological Test Scores on 32 Males and 32 Females

Males				Females			
y_1	y_2	y_3	y_4	y_1	y_2	y_3	y_4
15	17	24	14	13	14	12	21
17	15	32	26	14	12	14	26
15	14	29	23	12	19	21	21
13	12	10	16	12	13	10	16
20	17	26	28	11	20	16	16
15	21	26	21	12	9	14	18
15	13	26	22	10	13	18	24
13	5	22	22	10	8	13	23
14	7	30	17	12	20	19	23
17	15	30	27	11	10	11	27
17	17	26	20	12	18	25	25
17	20	28	24	14	18	13	26
15	15	29	24	14	10	25	28
18	19	32	28	13	16	8	14
18	18	31	27	14	8	13	25
15	14	26	21	13	16	23	28
18	17	33	26	16	21	26	26
10	14	19	17	14	17	14	14
18	21	30	29	16	16	15	23
18	21	34	26	13	16	23	24
13	17	30	24	2	6	16	21
16	16	16	16	14	16	22	26
11	15	25	23	17	17	22	28
16	13	26	16	16	13	16	14
16	13	23	21	15	14	20	26
18	18	34	24	12	10	12	9
16	15	28	27	14	17	24	23
15	16	29	24	13	15	18	20
18	19	32	23	11	16	18	28
18	16	33	23	7	7	19	18
17	20	21	21	12	15	7	28
19	19	30	28	6	5	6	13

By (5.9), we obtain

$$T^2 = \frac{n_1 n_2}{n_1 + n_2} (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{\text{pl}}^{-1} (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2) = 97.6015.$$

From interpolation in Table A.7, we obtain $T_{0.01, 4, 62}^2 = 15.373$, and we therefore reject $H_0: \mu_1 = \mu_2$. See Example 5.5 for a discussion of which variables contribute most to separation of the two groups. \square

5.4.3 Likelihood Ratio Tests

The maximum likelihood approach to estimation was introduced in Section 4.3.1. As noted there, the likelihood function is the joint density of y_1, y_2, \dots, y_n . The values of the parameters that maximize the likelihood function are the maximum likelihood estimators.

The *likelihood ratio* method of test construction uses the ratio of the maximum value of the likelihood function assuming H_0 is true to the maximum under H_1 , which is essentially unrestricted. Likelihood ratio tests usually have good power and sometimes have optimum power over a wide class of alternatives.

When applied to multivariate normal samples and $H_0: \mu_1 = \mu_2$, the likelihood ratio approach leads directly to Hotelling's T^2 -test in (5.9). Similarly, in the one-sample case, the T^2 -statistic in (5.5) is the likelihood ratio test. Thus the T^2 -test, which we introduced rather informally, is the best test according to certain criteria.

5.5 TESTS ON INDIVIDUAL VARIABLES CONDITIONAL ON REJECTION OF H_0 BY THE T^2 -TEST

If the hypothesis $H_0: \mu_1 = \mu_2$ is rejected, the implication is that $\mu_{1j} \neq \mu_{2j}$ for at least one $j = 1, 2, \dots, p$. But there is no guarantee that $H_0: \mu_{1j} = \mu_{2j}$ will be rejected for some j by a univariate test. However, if we consider a linear combination of the variables, $z = \mathbf{a}'\mathbf{y}$, then there is at least one coefficient vector \mathbf{a} for which

$$t(\mathbf{a}) = \frac{\bar{z}_1 - \bar{z}_2}{\sqrt{(1/n_1 + 1/n_2)s_z^2}} \quad (5.12)$$

will reject the corresponding hypothesis $H_0: \mu_{z_1} = \mu_{z_2}$ or $H_0: \mathbf{a}'\mu_1 = \mathbf{a}'\mu_2$. By (3.54), $\bar{z}_1 = \mathbf{a}'\bar{\mathbf{y}}_1$ and $\bar{z}_2 = \mathbf{a}'\bar{\mathbf{y}}_2$, and from (3.55) the variance estimator s_z^2 is the pooled estimator $\mathbf{a}'\mathbf{S}_{\text{pl}}\mathbf{a}$. Thus (5.12) can be written as

$$t(\mathbf{a}) = \frac{\mathbf{a}'\bar{\mathbf{y}}_1 - \mathbf{a}'\bar{\mathbf{y}}_2}{\sqrt{[(n_1 + n_2)/n_1 n_2]\mathbf{a}'\mathbf{S}_{\text{pl}}\mathbf{a}}}. \quad (5.13)$$

Since $t(\mathbf{a})$ can be negative, we work with $t^2(\mathbf{a})$. The linear function $z = \mathbf{a}'\mathbf{y}$ is a projection of \mathbf{y} onto a line through the origin. We seek the line (direction) on which the difference $\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2$ is maximized when projected. The projected difference $\mathbf{a}'(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)$ [standardized by $\mathbf{a}'\mathbf{S}_{\text{pl}}\mathbf{a}$ as in (5.13)] will be less in any other direction than that parallel to the line joining $\bar{\mathbf{y}}_1$ and $\bar{\mathbf{y}}_2$. The value of \mathbf{a} that projects onto this line, or equivalently, maximizes $t^2(\mathbf{a})$ in (5.13), is (any multiple of)

$$\mathbf{a} = \mathbf{S}_{\text{pl}}^{-1}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2). \quad (5.14)$$

Since \mathbf{a} in (5.14) projects $\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2$ onto a line parallel to the line joining $\bar{\mathbf{y}}_1$ and $\bar{\mathbf{y}}_2$, we would expect that $t^2(\mathbf{a}) = T^2$, and this is indeed the case (see Problem 5.3).

When $\mathbf{a} = \mathbf{S}_{\text{pl}}^{-1}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)$, then $z = \mathbf{a}'\mathbf{y}$ is called the *discriminant function*. Sometimes the vector \mathbf{a} itself in (5.14) is loosely referred to as the discriminant function.

If $H_0: \mu_1 = \mu_2$ is rejected by T^2 in (5.9), the discriminant function $\mathbf{a}'\mathbf{y}$ will lead to rejection of $H_0: \mathbf{a}'\mu_1 = \mathbf{a}'\mu_2$ using (5.13), with $\mathbf{a} = \mathbf{S}_{\text{pl}}^{-1}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)$. We can then examine each a_j in \mathbf{a} for an indication of the contribution of the corresponding y_j to rejection of H_0 . This follow-up examination of each a_j should be done only if $H_0: \mu_1 = \mu_2$ is rejected by T^2 . The discriminant function will appear again in Section 5.6.2 and in Chapters 8 and 9.

We list these and other procedures that could be used to check each variable following rejection of H_0 by a two-sample T^2 -test:

1. Univariate t -tests, one for each variable,

$$t_j = \frac{\bar{y}_{1j} - \bar{y}_{2j}}{\sqrt{[(n_1 + n_2)/n_1 n_2]s_{jj}}}, \quad j = 1, 2, \dots, p, \quad (5.15)$$

where s_{jj} is the j th diagonal element of \mathbf{S}_{pl} . Reject $H_0: \mu_{1j} = \mu_{2j}$ if $|t_j| > t_{\alpha/2, n_1+n_2-2}$. For confidence intervals on $\mu_{1j} - \mu_{2j}$, see Rencher (1998, Section 3.7).

2. To adjust the α -level resulting from performing the p tests in (5.15), we could use a Bonferroni critical value $t_{\alpha/2p, n_1+n_2-2}$ for (5.15) (Bonferroni 1936). A critical value $t_{\alpha/2p}$ is much greater than the corresponding $t_{\alpha/2}$, and the resulting overall α -level is conservative. Bonferroni critical values $t_{\alpha/2p, \nu}$ are given in Table A.8, from Bailey (1977).
3. Another critical value that could be used with (5.15) is T_{α, p, n_1+n_2-2} , where T_α is the square root of T_α^2 from Table A.7; that is,

$$T_{\alpha, p, n_1+n_2-2} = \sqrt{T_{\alpha, p, n_1+n_2-2}^2}$$

This allows for all p variables to be tested as well as all possible linear combinations, as in (5.13), even linear combinations chosen after seeing the data. Consequently, the use of T_α is even more conservative than using $t_{\alpha/2p}$; that is, $T_{\alpha, p, n_1+n_2-2} > t_{\alpha/2p, n_1+n_2-2}$.

4. Partial F - or t -tests [test of each variable adjusted for the other variables; see (5.32) in Section 5.8]
5. Standardized discriminant function coefficients (see Section 8.5)
6. Correlations between the variables and the discriminant function (see Section 8.7.3)
7. Stepwise discriminant analysis (see Section 8.9)

The first three methods are univariate approaches that do not use covariances or correlations among the variables in the computation of the test statistic. The last four methods are multivariate in the sense that the correlation structure is explicitly taken into account in the computation.

Method 6, involving the correlation between each variable and the discriminant function, is recommended in many texts and software packages. However, Rencher (1988) has shown that these correlations are proportional to individual t - or F -tests (see Section 8.7.3). Thus this method is equivalent to method 1 and is a univariate rather than a multivariate approach. Method 7 is often used to identify a subset of important variables or even to rank the variables according to order of entry. But Rencher and Larson (1980) have shown that stepwise methods have a high risk of selecting spurious variables, unless the sample size is very large.

We now consider the univariate procedures 1, 2, and 3. The probability of rejecting one or more of the p univariate tests when H_0 is true is called the *overall α* or *experimentwise error rate*. If we do univariate tests only, with no T^2 -test, then the tests based on $t_{\alpha/2p}$ and T_α in procedures 2 and 3 are conservative (overall α too low), and tests based on $t_{\alpha/2}$ in procedure 1 are liberal (overall α too high). However, when these tests are carried out *only* after rejection by the T^2 -test (such tests are sometimes called protected tests), the experimentwise error rates change. Obviously the tests will reject less often (under H_0) if they are carried out only if T^2 rejects. Thus the tests using $t_{\alpha/2p}$ and T_α become even more conservative, and the test using $t_{\alpha/2}$ becomes more acceptable.

Hummel and Sligo (1971) studied the experimentwise error rate for univariate t -tests following rejection of H_0 by the T^2 -test (protected tests). Using $\alpha = .05$, they found that using $t_{\alpha/2}$ for a critical value yields an overall α acceptably close to the nominal .05. In fact, it is slightly conservative, making this the preferred univariate test (within the limits of their study). They also compared this procedure with that of performing univariate tests without a prior T^2 -test (unprotected tests). For this case, the overall α is too high, as expected. Table 5.2 gives an excerpt of Hummel and Sligo's results. The sample size is for each of the two samples; the r^2 in common is for every pair of variables.

Hummel and Sligo therefore recommended performing the multivariate T^2 -test followed by univariate t -tests. This procedure appears to have the desired overall α level and will clearly have better power than tests using T_α or $t_{\alpha/2p}$ as a critical value. Table 5.2 also highlights the importance of using univariate t -tests *only* if the multivariate T^2 -test is significant. The inflated α 's resulting if t -tests are used without regard to the outcome of the T^2 -test are clearly evident. Thus among the three univariate procedures (procedures 1, 2, and 3 above), the first appears to be preferred.

Among the multivariate approaches (procedures 4, 5, and 7 above), we prefer the fifth procedure, which compares the (absolute value of) coefficients in the discriminant function to find the effect of each variable in separating the two groups of observations. These coefficients will often tell a different story from the univariate tests, because the univariate tests do not take into account the correlations among the variables or the effect of each variable on T^2 in the presence of the other variables. A variable will typically have a different effect in the presence of other variables than it has by itself. In the discriminant function $z = \mathbf{a}'\mathbf{y} = a_1y_1 + a_2y_2 + \cdots + a_py_p$, where $\mathbf{a} = \mathbf{S}_{pl}^{-1}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)$, the coefficients a_1, a_2, \dots, a_p indicate the relative importance of the variables in a multivariate context, something the univariate t -tests

Table 5.2 Comparison of Experimentwise Error Rates
(Nominal $\alpha = 0.05$)

Sample Size	Number of Variables	Common r^2			
		.10	.30	.50	.70
<i>Univariate Tests Only^a</i>					
10	3	.145	.112	.114	.077
10	6	.267	.190	.178	.111
10	9	.348	.247	.209	.129
30	3	.115	.119	.117	.085
30	6	.225	.200	.176	.115
30	9	.296	.263	.223	.140
50	3	.138	.124	.102	.083
50	6	.230	.190	.160	.115
50	9	.324	.258	.208	.146
<i>Multivariate Test Followed by Univariate Tests^b</i>					
10	3	.044	.029	.035	.022
10	6	.046	.029	.030	.017
10	9	.050	.026	.025	.018
30	3	.037	.044	.029	.025
30	6	.037	.037	.032	.021
30	9	.042	.042	.030	.021
50	3	.038	.041	.033	.028
50	6	.037	.039	.028	.027
50	9	.036	.038	.026	.020

^a Ignoring multivariate tests.^b Carried out only if multivariate test rejects.

cannot do. If the variables are not commensurate (similar in scale and variance), the coefficients should be standardized, as in Section 8.5; this allows for more valid comparisons among the variables. Rencher and Scott (1990) provide a decomposition of the information in the standardized discriminant function coefficients. For a detailed analysis of the effect of each variable in the presence of the other variables, see Rencher (1993; 1998, Sections 3.3.5 and 3.5.3).

■ EXAMPLE 5.5

For the psychological data in Table 5.1, we obtained \bar{y}_1 , \bar{y}_2 , and S_{pl} in Example 5.4.2. The discriminant function coefficient vector is obtained from (5.14) as

$$\mathbf{a} = S_{pl}^{-1}(\bar{y}_1 - \bar{y}_2) = \begin{pmatrix} .5104 \\ -.2033 \\ .4660 \\ -.3097 \end{pmatrix}.$$

Thus the linear combination that best separates the two groups is

$$\mathbf{a}'\mathbf{y} = .5104y_1 - .2033y_2 + .4660y_3 - .3097y_4,$$

in which y_1 and y_3 appear to contribute most to separation of the two groups. (After standardization, the relative contribution of the variables changes somewhat; see the answer to Problem 8.7 in Appendix B.) \square

5.6 COMPUTATION OF T^2

If one has a program available with matrix manipulation capability, it is a simple matter to compute T^2 using (5.9). However, this approach is somewhat cumbersome for those not accustomed to the use of such a programming language, and many would prefer a more automated procedure. But very few general-purpose statistical programs provide for direct calculation of the two-sample T^2 -statistic, perhaps because it is so easy to obtain from other procedures. We will discuss two types of widely available procedures that can be used to compute T^2 .

5.6.1 Obtaining T^2 from a MANOVA Program

Multivariate analysis of variance (MANOVA) is discussed in Chapter 6, and the reader may wish to return to the present section after becoming familiar with that material. One-way MANOVA involves a comparison of mean vectors from several samples. Typically, the number of samples is three or more, but the procedure will also accommodate two samples. The two-sample T^2 test is thus a special case of MANOVA.

Four common tests of significance are defined in Section 6.1: Wilks' Λ , the Lawley-Hotelling $U^{(s)}$, Pillai's $V^{(s)}$, and Roy's largest root θ . Without concerning ourselves here with how these are defined or calculated, we show how to use each to obtain the two-sample T^2 :

$$T^2 = (n_1 + n_2 - 2) \frac{1 - \Lambda}{\Lambda}, \quad (5.16)$$

$$T^2 = (n_1 + n_2 - 2) U^{(s)}, \quad (5.17)$$

$$T^2 = (n_1 + n_2 - 2) \frac{V^{(s)}}{1 - V^{(s)}}, \quad (5.18)$$

$$T^2 = (n_1 + n_2 - 2) \frac{\theta}{1 - \theta}. \quad (5.19)$$

(For the special case of two groups, $V^{(s)} = \theta$.) These relationships are demonstrated in Section 6.1.7. If the MANOVA program gives eigenvectors of $\mathbf{E}^{-1}\mathbf{H}$ (\mathbf{E} and \mathbf{H} are defined in Section 6.1.2), the eigenvector corresponding to the largest eigenvalue will be equal to (a constant multiple of) the discriminant function $\mathbf{S}_{\text{pl}}^{-1}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)$.

5.6.2 Obtaining T^2 from Multiple Regression

In this section, the y 's become independent variables in a regression model. For each observation vector \mathbf{y}_{1i} and \mathbf{y}_{2i} in a two-sample T^2 , define a "dummy" group variable as

$$\begin{aligned} w_i &= \frac{n_2}{n_1 + n_2} \text{ for each of } \mathbf{y}_{11}, \mathbf{y}_{12}, \dots, \mathbf{y}_{1n_1} \text{ in sample 1} \\ &= -\frac{n_1}{n_1 + n_2} \text{ for each of } \mathbf{y}_{21}, \mathbf{y}_{22}, \dots, \mathbf{y}_{2n_2} \text{ in sample 2.} \end{aligned}$$

Then $\bar{w} = 0$ for all $n_1 + n_2$ observations. The prediction equation for the regression of w on the y 's can be written as

$$\hat{w}_i = b_0 + b_1 y_{i1} + b_2 y_{i2} + \cdots + b_p y_{ip},$$

where i ranges over all $n_1 + n_2$ observations and the least squares estimate b_0 is [see (10.15)]

$$b_0 = \bar{w} - b_1 \bar{y}_1 - b_2 \bar{y}_2 - \cdots - b_p \bar{y}_p.$$

Substituting this into the regression equation, we obtain

$$\begin{aligned} \hat{w}_i &= \bar{w} + b_1(y_{i1} - \bar{y}_1) + b_2(y_{i2} - \bar{y}_2) + \cdots + b_p(y_{ip} - \bar{y}_p) \\ &= b_1(y_{i1} - \bar{y}_1) + b_2(y_{i2} - \bar{y}_2) + \cdots + b_p(y_{ip} - \bar{y}_p) \quad (\text{since } \bar{w} = 0). \end{aligned}$$

Let $\mathbf{b}' = (b_1, b_2, \dots, b_p)$ be the vector of regression coefficients and R^2 the squared multiple correlation. Then we have the following relationships:

$$T^2 = (n_1 + n_2 - 2) \frac{R^2}{1 - R^2}, \tag{5.20}$$

$$\mathbf{a} = \mathbf{S}_{\text{pl}}^{-1}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2) = \left[\frac{n_1 + n_2}{n_1 n_2} (n_1 + n_2 - 2 + T^2) \right] \mathbf{b}. \tag{5.21}$$

Thus with ordinary multiple regression, one can easily obtain T^2 and the discriminant function $\mathbf{S}_{\text{pl}}^{-1}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)$. We simply define w_i as above for each of the $n_1 + n_2$ observations, regress the w 's on the y 's, and use the resulting R^2 in (5.20). For \mathbf{b} , delete the intercept from the regression coefficients for use in (5.21). Actually, since only the relative values of the elements of $\mathbf{a} = \mathbf{S}_{\text{pl}}^{-1}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)$ are of interest, it is not necessary to convert from \mathbf{b} to \mathbf{a} in (5.21). We can use \mathbf{b} directly or standardize the values b_1, b_2, \dots, b_p as in Section 8.5.

■ EXAMPLE 5.6.2

We illustrate the regression approach to computation of T^2 using the psychological data in Table 5.1. We set $w = n_2/(n_1 + n_2) = \frac{32}{64} = \frac{1}{2}$ for each observation in the first group (males) and equal to $-n_1/(n_1 + n_2) = -\frac{1}{2}$ in

the second group (females). When w is regressed on the 64 y 's, we obtain

$$\begin{pmatrix} b_0 \\ b_1 \\ b_2 \\ b_3 \\ b_4 \end{pmatrix} = \begin{pmatrix} -.751 \\ .051 \\ -.020 \\ .047 \\ -.031 \end{pmatrix}, \quad R^2 = .6115.$$

By (5.20),

$$T^2 = (n_1 + n_2 - 2) \frac{R^2}{1 - R^2} = \frac{62(.6115)}{1 - .6115} = 97.601,$$

as was obtained before in Example 5.4.2. Note that $\mathbf{b}' = (b_1, b_2, b_3, b_4) = (.051, -.020, .047, -.031)$, with the intercept deleted, is proportional to the discriminant function vector \mathbf{a} from Example 5.5, as we would expect from (5.21). \square

5.7 PAIRED OBSERVATIONS TEST

As usual, we begin with the univariate case to set the stage for the multivariate presentation.

5.7.1 Univariate Case

Suppose two samples are not independent because there exists a natural pairing between the i th observation y_i in the first sample and the i th observation x_i in the second sample for all i , as, for example, when a treatment is applied twice to the same individual or when subjects are matched according to some criterion, such as IQ or family background. With such pairing, the samples are often referred to as *paired observations* or *matched pairs*. The two samples thus obtained are correlated, and the two-sample test statistic in (5.8) is not appropriate because the samples must be independent in order for (5.8) to have a t -distribution. [The two-sample test in (5.8) is somewhat robust to heterogeneity of variances and to lack of normality but not to dependence.] We reduce the two samples to one by working with the differences between the paired observations, as in the following layout for two treatments applied to the same subject:

Pair Number	Treatment 1	Treatment 2	Difference $d_i = y_i - x_i$
1	y_1	x_1	d_1
2	y_2	x_2	d_2
\vdots	\vdots	\vdots	\vdots
n	y_n	x_n	d_n

To obtain a t -test, it is not sufficient to assume individual normality for each of y and x . To allow for the covariance between y and x , we need the additional assumption that y and x have a bivariate normal distribution with

$$\boldsymbol{\mu} = \begin{pmatrix} \mu_y \\ \mu_x \end{pmatrix}, \quad \boldsymbol{\Sigma} = \begin{pmatrix} \sigma_y^2 & \sigma_{yx} \\ \sigma_{yx} & \sigma_x^2 \end{pmatrix}.$$

It then follows by property 1a in Section 4.2 that $d_i = y_i - x_i$ is $N(\mu_y - \mu_x, \sigma_d^2)$, where $\sigma_d^2 = \sigma_y^2 - 2\sigma_{yx} + \sigma_x^2$. From d_1, d_2, \dots, d_n we calculate

$$\bar{d} = \frac{1}{n} \sum_{i=1}^n d_i \quad \text{and} \quad s_d^2 = \frac{1}{n-1} \sum_{i=1}^n (d_i - \bar{d})^2.$$

To test $H_0: \mu_y = \mu_x$, that is, $H_0: \mu_d = 0$, we use the one-sample statistic

$$t = \frac{\bar{d}}{s_d/\sqrt{n}}, \tag{5.22}$$

which is distributed as t_{n-1} if H_0 is true. We reject H_0 in favor of $H_1: \mu_d \neq 0$ if $|t| > t_{\alpha/2, n-1}$. It is not necessary to assume $\sigma_y^2 = \sigma_x^2$ because there are no restrictions on $\boldsymbol{\Sigma}$.

This test has only $n - 1$ degrees of freedom compared with $2(n - 1)$ for the two independent sample t -test (5.8). In general, the pairing reduces the within-sample variation s_d and thereby increases the power.

If we mistakenly treated the two samples as independent and used (5.8) with $n_1 = n_2 = n$, we would have

$$t = \frac{\bar{y} - \bar{x}}{s_{\text{pl}}\sqrt{2/n}} = \frac{\bar{y} - \bar{x}}{\sqrt{2s_{\text{pl}}^2/n}}.$$

However,

$$E\left(\frac{2s_{\text{pl}}^2}{n}\right) = 2E\left[\frac{(n-1)s_y^2 + (n-1)s_x^2}{(n+n-2)n}\right] = (\sigma_y^2 + \sigma_x^2)/n,$$

whereas $\text{var}(\bar{y} - \bar{x}) = (\sigma_y^2 + \sigma_x^2 - 2\sigma_{yx})/n$. Thus if the test for independent samples (5.8) is used for paired data, it does not have a t -distribution, and in fact underestimates the true average t -value (assuming H_0 is false), since $\sigma_y^2 + \sigma_x^2 > \sigma_y^2 + \sigma_x^2 - 2\sigma_{yx}$ if $\sigma_{yx} > 0$, which would be typical in this situation. One could therefore use

$$t = \frac{\bar{y} - \bar{x}}{\sqrt{(s_y^2 + s_x^2 - 2s_{yx})/n}}, \tag{5.23}$$

but $t = \sqrt{n}\bar{d}/s_d$ in (5.22) is equal to it and somewhat simpler to use.

5.7.2 Multivariate Case

Here we assume the same natural pairing of sampling units as in the univariate case, but we measure p variables on each sampling unit. Thus \mathbf{y}_i from the first sample is paired with \mathbf{x}_i from the second sample, $i = 1, 2, \dots, n$. In terms of two treatments applied to each sampling unit, this situation is as follows:

Pair Number	Treatment 1	Treatment 2	Difference $\mathbf{d}_i = \mathbf{y}_i - \mathbf{x}_i$
1	\mathbf{y}_1	\mathbf{x}_1	\mathbf{d}_1
2	\mathbf{y}_2	\mathbf{x}_2	\mathbf{d}_2
\vdots	\vdots	\vdots	\vdots
n	\mathbf{y}_n	\mathbf{x}_n	\mathbf{d}_n

In Section 5.7.1, we made the assumption that y and x have a bivariate normal distribution, in which y and x are correlated. Here we assume y and x are correlated and have a multivariate normal distribution:

$$\begin{pmatrix} \mathbf{y} \\ \mathbf{x} \end{pmatrix} \text{ is } N_{2p} \left[\begin{pmatrix} \boldsymbol{\mu}_y \\ \boldsymbol{\mu}_x \end{pmatrix}, \begin{pmatrix} \boldsymbol{\Sigma}_{yy} & \boldsymbol{\Sigma}_{yx} \\ \boldsymbol{\Sigma}_{xy} & \boldsymbol{\Sigma}_{xx} \end{pmatrix} \right].$$

To test $H_0: \boldsymbol{\mu}_d = \mathbf{0}$, which is equivalent to $H_0: \boldsymbol{\mu}_y = \boldsymbol{\mu}_x$ since $\boldsymbol{\mu}_d = E(\mathbf{y} - \mathbf{x}) = \boldsymbol{\mu}_y - \boldsymbol{\mu}_x$, we calculate

$$\bar{\mathbf{d}} = \frac{1}{n} \sum_{i=1}^n \mathbf{d}_i \quad \text{and} \quad \mathbf{S}_d = \frac{1}{n-1} \sum_{i=1}^n (\mathbf{d}_i - \bar{\mathbf{d}})(\mathbf{d}_i - \bar{\mathbf{d}})'.$$

We then have

$$T^2 = \bar{\mathbf{d}}' \left(\frac{\mathbf{S}_d}{n} \right)^{-1} \bar{\mathbf{d}} = n \bar{\mathbf{d}}' \mathbf{S}_d^{-1} \bar{\mathbf{d}}. \quad (5.24)$$

Under H_0 , this paired comparison T^2 -statistic is distributed as $T_{p,n-1}^2$. We reject H_0 if $T^2 > T_{\alpha,p,n-1}^2$. Note that \mathbf{S}_d estimates $\text{cov}(\mathbf{y} - \mathbf{x}) = \boldsymbol{\Sigma}_{yy} - \boldsymbol{\Sigma}_{yx} - \boldsymbol{\Sigma}_{xy} + \boldsymbol{\Sigma}_{xx}$, for which an equivalent estimator would be $\mathbf{S}_{yy} - \mathbf{S}_{yx} - \mathbf{S}_{xy} + \mathbf{S}_{xx}$ [see (3.42)].

The cautions expressed in Section 5.7.1 for univariate paired observation data also apply here. If the two samples of multivariate observations are correlated because of a natural pairing of sampling units, the test in (5.24) should be used rather than the two-sample T^2 -test in (5.9), which assumes two independent samples. Misuse of (5.9) in place of (5.24) will lead to loss of power.

Since the assumption $\boldsymbol{\Sigma}_{yy} = \boldsymbol{\Sigma}_{xx}$ is not needed for (5.24) to have a T^2 -distribution, this test can be used for independent samples when $\boldsymbol{\Sigma}_1 \neq \boldsymbol{\Sigma}_2$ (as long as $n_1 = n_2$). The observations in the two samples would be paired in the order they were obtained or in an arbitrary order. However, in the case of independent samples, the pairing achieves no gain in power to offset the loss of $n - 1$ degrees of freedom.

By analogy with (5.14), the discriminant function for paired observation data becomes

$$\mathbf{a} = \mathbf{S}_d^{-1} \bar{\mathbf{d}}. \quad (5.25)$$

Table 5.3 Maximum Depth of Pits and Number of Pits of Coated Pipes

Location	Coating 1		Coating 2		Difference	
	Depth y_1	Number y_2	Depth x_1	Number x_2	Depth d_1	Number d_2
1	73	31	51	35	22	-4
2	43	19	41	14	2	5
3	47	22	43	19	4	3
4	53	26	41	29	12	-3
5	58	36	47	34	11	2
6	47	30	32	26	15	4
7	52	29	24	19	28	10
8	38	36	43	37	-5	-1
9	61	34	53	24	8	10
10	56	33	52	27	4	6
11	56	19	57	14	-1	5
12	34	19	44	19	-10	0
13	55	26	57	30	-2	-4
14	65	15	40	7	25	8
15	75	18	68	13	7	5

For tests on individual variables, we have

$$t_j = \frac{\bar{d}_j}{\sqrt{s_{d,jj}/n}}, \quad j = 1, 2, \dots, p. \quad (5.26)$$

The critical value for t_j is $t_{\alpha/2p, n-1}$ or $t_{\alpha/2, n-1}$ depending on whether a T^2 -test is carried out first (see Section 5.5).

■ EXAMPLE 5.7.2

To compare two types of coating for resistance to corrosion, 15 pieces of pipe were coated with each type of coating (Kramer and Jensen 1969b). Two pipes, one with each type of coating, were buried together and left for the same length of time at 15 different locations, providing a natural pairing of the observations. Corrosion for the first type of coating was measured by two variables,

y_1 = maximum depth of pit in thousandths of an inch

y_2 = number of pits

with x_1 and x_2 defined analogously for the second coating. The data and differences are given in Table 5.3. Thus we have, for example, $\mathbf{y}'_1 = (73, 31)$, $\mathbf{x}'_1 = (51, 35)$, and $\mathbf{d}'_1 = \mathbf{y}'_1 - \mathbf{x}'_1 = (22, -4)$. For the 15 difference vectors, we obtain

$$\bar{\mathbf{d}} = \begin{pmatrix} 8.000 \\ 3.067 \end{pmatrix}, \quad \mathbf{S}_d = \begin{pmatrix} 121.571 & 17.071 \\ 17.071 & 21.781 \end{pmatrix}.$$

By (5.24),

$$T^2 = (15)(8.000, 3.067) \begin{pmatrix} 121.571 & 17.071 \\ 17.071 & 21.781 \end{pmatrix}^{-1} \begin{pmatrix} 8.000 \\ 3.067 \end{pmatrix} = 10.819.$$

Since $T^2 = 10.819 > T_{0.05, 2, 14}^2 = 8.197$, we reject $H_0: \mu_d = 0$ and conclude that the two coatings differ in their effect on corrosion. \square

5.8 TEST FOR ADDITIONAL INFORMATION

In this section, we are again considering two independent samples as in Section 5.4.2. We start with a basic $p \times 1$ vector \mathbf{y} of measurements on each sampling unit and ask whether a $q \times 1$ subvector \mathbf{x} measured in addition to \mathbf{y} (on the same unit) will significantly increase the separation of the two samples as shown by T^2 . It is not necessary that we add new variables. We may be interested in determining whether some of the variables we already have are redundant in the presence of other variables in terms of separating the groups. We have designated the subset of interest by \mathbf{x} for notational convenience.

It is assumed that the two samples are from multivariate normal populations with a common covariance matrix, that is,

$$\begin{pmatrix} \mathbf{y}_{11} \\ \mathbf{x}_{11} \end{pmatrix}, \begin{pmatrix} \mathbf{y}_{12} \\ \mathbf{x}_{12} \end{pmatrix}, \dots, \begin{pmatrix} \mathbf{y}_{1n_1} \\ \mathbf{x}_{1n_1} \end{pmatrix} \text{ are from } N_{p+q}(\boldsymbol{\mu}_1, \Sigma),$$

$$\begin{pmatrix} \mathbf{y}_{21} \\ \mathbf{x}_{21} \end{pmatrix}, \begin{pmatrix} \mathbf{y}_{22} \\ \mathbf{x}_{22} \end{pmatrix}, \dots, \begin{pmatrix} \mathbf{y}_{2n_2} \\ \mathbf{x}_{2n_2} \end{pmatrix} \text{ are from } N_{p+q}(\boldsymbol{\mu}_2, \Sigma),$$

where

$$\boldsymbol{\mu}_1 = E\begin{pmatrix} \mathbf{y}_{1i} \\ \mathbf{x}_{1i} \end{pmatrix} = \begin{pmatrix} \boldsymbol{\mu}_{1y} \\ \boldsymbol{\mu}_{1x} \end{pmatrix}, \quad \boldsymbol{\mu}_2 = E\begin{pmatrix} \mathbf{y}_{2i} \\ \mathbf{x}_{2i} \end{pmatrix} = \begin{pmatrix} \boldsymbol{\mu}_{2y} \\ \boldsymbol{\mu}_{2x} \end{pmatrix},$$

$$\Sigma = \text{cov}\begin{pmatrix} \mathbf{y}_{1i} \\ \mathbf{x}_{1i} \end{pmatrix} = \text{cov}\begin{pmatrix} \mathbf{y}_{2i} \\ \mathbf{x}_{2i} \end{pmatrix} = \begin{pmatrix} \Sigma_{yy} & \Sigma_{yx} \\ \Sigma_{xy} & \Sigma_{xx} \end{pmatrix}.$$

We partition the sample mean vectors and covariance matrix accordingly:

$$\begin{pmatrix} \bar{\mathbf{y}}_1 \\ \bar{\mathbf{x}}_1 \end{pmatrix}, \begin{pmatrix} \bar{\mathbf{y}}_2 \\ \bar{\mathbf{x}}_2 \end{pmatrix}, \quad \mathbf{S}_{pl} = \begin{pmatrix} \mathbf{S}_{yy} & \mathbf{S}_{yx} \\ \mathbf{S}_{xy} & \mathbf{S}_{xx} \end{pmatrix},$$

where \mathbf{S}_{pl} is the pooled sample covariance matrix from the two samples.

We wish to test the hypothesis that \mathbf{x}_1 and \mathbf{x}_2 are redundant for separating the two groups, that is, that the extra q variables do not contribute anything significant beyond the information already available in \mathbf{y}_1 and \mathbf{y}_2 for separating the groups. This is in the spirit of a *full and reduced model* test in regression [see (5.31) and Section 10.2.5b]. However, here we are working with a subset of dependent variables as contrasted to the subset of independent variables in the regression setting. Thus

both \mathbf{y} and \mathbf{x} are subvectors of dependent variables. In this setting, the independent variables would be grouping variables 1 and 2 corresponding to μ_1 and μ_2 .

We are not asking whether the x 's can significantly separate the two groups by themselves, but whether they provide additional separation beyond the separation already achieved by the y 's. If the x 's were independent of the y 's, we would have $T_{p+q}^2 = T_p^2 + T_q^2$, but this does not hold, because they are correlated. We must compare T_{p+q}^2 for the full set of variables $(y_1, \dots, y_p, x_1, \dots, x_q)$ with T_p^2 based on the reduced set (y_1, \dots, y_p) . We are inquiring whether the increase from T_p^2 to T_{p+q}^2 is significant.

By definition, the T^2 -statistic based on the full set of $p + q$ variables is given by

$$T_{p+q}^2 = \frac{n_1 n_2}{n_1 + n_2} \left[\left(\begin{array}{c} \bar{\mathbf{y}}_1 \\ \bar{\mathbf{x}}_1 \end{array} \right) - \left(\begin{array}{c} \bar{\mathbf{y}}_2 \\ \bar{\mathbf{x}}_2 \end{array} \right) \right]' \mathbf{S}_{\text{pl}}^{-1} \left[\left(\begin{array}{c} \bar{\mathbf{y}}_1 \\ \bar{\mathbf{x}}_1 \end{array} \right) - \left(\begin{array}{c} \bar{\mathbf{y}}_2 \\ \bar{\mathbf{x}}_2 \end{array} \right) \right], \quad (5.27)$$

whereas T^2 for the reduced set of p variables is

$$T_p^2 = \frac{n_1 n_2}{n_1 + n_2} (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{yy}^{-1} (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2). \quad (5.28)$$

Then the test statistic for the significance of the increase from T_p^2 to T_{p+q}^2 is given by

$$T^2(\mathbf{x}|\mathbf{y}) = (\nu - p) \frac{T_{p+q}^2 - T_p^2}{\nu + T_p^2}, \quad (5.29)$$

which is distributed as $T_{q,\nu-p}^2$. We reject the hypothesis of redundancy of \mathbf{x} if $T^2(\mathbf{x}|\mathbf{y}) \geq T_{\alpha,q,\nu-p}^2$.

By (5.7), $T^2(\mathbf{x}|\mathbf{y})$ can be converted to an F -statistic:

$$F = \frac{\nu - p - q + 1}{q} \frac{T_{p+q}^2 - T_p^2}{\nu + T_p^2}, \quad (5.30)$$

which is distributed as $F_{q,\nu-p-q+1}$, and we reject the hypothesis of redundancy if $F \geq F_{\alpha,q,\nu-p-q+1}$.

In both cases $\nu = n_1 + n_2 - 2$. Note that the first degrees-of-freedom parameter in both (5.29) and (5.30) is q , the number of x 's. The second parameter in (5.29) is $\nu - p$ because the statistic is adjusted for the p variables in \mathbf{y} .

To prove directly that the statistic defined in (5.30) has an F -distribution, we can use a basic relationship from multiple regression [see (10.33)]:

$$F_{q,\nu-p-q+1} = \frac{(R_{p+q}^2 - R_p^2)(\nu - p - q + 1)}{(1 - R_{p+q}^2)q}, \quad (5.31)$$

where R_{p+q}^2 is the squared multiple correlation from the full model with $p + q$ independent variables and R_p^2 is from the reduced model with p independent variables. If we solve for R^2 in terms of T^2 from (5.20) and substitute this into (5.31), we readily obtain the test statistic in (5.30).

If we are interested in the effect of adding a single x , then $q = 1$, and both (5.29) and (5.30) reduce to

$$t^2(x|\mathbf{y}) = (\nu - p) \frac{T_{p+1}^2 - T_p^2}{\nu + T_p^2}, \quad (5.32)$$

and we reject the hypothesis of redundancy of x if $t^2(x|\mathbf{y}) \geq t_{\alpha/2, \nu-p}^2 = F_{\alpha, 1, \nu-p}$.

■ EXAMPLE 5.8

We use the psychological data of Table 5.1 to illustrate tests on subvectors. We begin by testing the significance of y_3 and y_4 above and beyond y_1 and y_2 . (In the notation of the present section, y_3 and y_4 become x_1 and x_2 .) For these subvectors, $p = 2$ and $q = 2$. The value of T_{p+q}^2 for all four variables as given by (5.27) was obtained in Example 5.4.2 as 97.6015. For y_1 and y_2 , we obtain, by (5.28),

$$\begin{aligned} T_p^2 &= \frac{n_1 n_2}{n_1 + n_2} (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{yy}^{-1} (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2) \\ &= \frac{(32)^2}{32 + 32} \begin{pmatrix} 15.97 - 12.34 \\ 15.91 - 13.91 \end{pmatrix}' \begin{pmatrix} 7.16 & 6.05 \\ 6.05 & 15.89 \end{pmatrix}^{-1} \begin{pmatrix} 15.97 - 12.34 \\ 15.91 - 13.91 \end{pmatrix} \\ &= 31.0126. \end{aligned}$$

By (5.29), the test statistic is

$$\begin{aligned} T^2(y_3, y_4|y_1, y_2) &= (\nu - p) \frac{T_{p+q}^2 - T_p^2}{\nu + T_p^2} \\ &= (62 - 2) \frac{97.6015 - 31.0126}{62 + 31.0126} = 42.955. \end{aligned}$$

We reject the hypothesis that $\mathbf{x} = (y_3, y_4)'$ is redundant, since $42.955 > T_{0.01, 2, 60}^2 = 10.137$. We conclude that $\mathbf{x} = (y_3, y_4)'$ adds a significant amount of separation to $\mathbf{y} = (y_1, y_2)'$.

To test the effect of each variable adjusted for the other three, we use (5.32). In this case, $p = 3$, $\nu = 62$, and $\nu - p = 59$. The results are given below, where $T_{p+1}^2 = 97.6015$ and T_p^2 in each case is based on the three variables, excluding the variable in question. For example, $T_p^2 = 90.8348$ for y_2 is based on y_1, y_3 , and y_4 , and $t^2(y_2|y_1, y_2, y_3) = 2.612$:

Variable	T_p^2	$(\nu - p) \frac{T_{p+1}^2 - T_p^2}{\nu + T_p^2}$
y_1	78.8733	7.844
y_2	90.8348	2.612
y_3	32.6253	40.513
y_4	74.5926	9.938

When we compare these four test statistic values with the critical value $t_{0.025, 59}^2 = 4.002$, we see that each variable makes a significant contribution to T^2 except y_2 . Note that y_3 contributes most, followed by y_4 and then y_1 . This order

differs from that given by the raw discriminant function in Example 5.5 but agrees with the order for the standardized discriminant function given in the answer to Problem 8.7 in Appendix B. \square

5.9 PROFILE ANALYSIS

If \mathbf{y} is $N_p(\boldsymbol{\mu}, \Sigma)$ and the variables in \mathbf{y} are commensurate (measured in the same units and with approximately equal variances as, for example, in the probe word data in Table 3.6), we may wish to compare the means $\mu_1, \mu_2, \dots, \mu_p$ in $\boldsymbol{\mu}$. This might be of interest when a measurement is taken on the same research unit at p successive times. Such situations are often referred to as *repeated measures* designs or *growth curves*, which are discussed in some generality in Sections 6.9 and 6.10. In the present section, we discuss one- and two-sample *profile analysis*. Profile analysis for several samples is covered in Section 6.8.

The pattern obtained by plotting $\mu_1, \mu_2, \dots, \mu_p$ as ordinates and connecting the points is called a *profile*; we usually draw straight lines connecting the points $(1, \mu_1), (2, \mu_2), \dots, (p, \mu_p)$. Profile analysis is an analysis of the profile or a comparison of two or more profiles. Profile analysis is often discussed in the context of administering a battery of p psychological or other tests.

In growth curve analysis where the variables are measured at time intervals, the responses have a natural order. In profile analysis where the variables arise from test scores, there is ordinarily no natural order. A distinction is not always made between repeated measures of the same variable through time and profile analysis of several different commensurate variables on the same individual.

5.9.1 One-Sample Profile Analysis

We begin with a discussion of the profile of the mean vector $\boldsymbol{\mu}$ from a single sample. A plot of $\boldsymbol{\mu}$ might appear as in Figure 5.3, where we plot $(1, \mu_1), (2, \mu_2), \dots, (p, \mu_p)$ and connect the points.

In order to compare the means $\mu_1, \mu_2, \dots, \mu_p$ in $\boldsymbol{\mu}$, the basic hypothesis is that the profile is *level* or *flat*:

$$H_0: \mu_1 = \mu_2 = \dots = \mu_p \quad \text{vs.} \quad H_1: \mu_j \neq \mu_k \quad \text{for some } j \neq k.$$

The data matrix \mathbf{Y} is given in (3.17). We cannot use univariate analysis of variance to test H_0 because the columns in \mathbf{Y} are not independent. For a multivariate approach that allows for correlated variables, we first express H_0 as $p - 1$ comparisons,

$$H_0: \begin{pmatrix} \mu_1 - \mu_2 \\ \mu_2 - \mu_3 \\ \vdots \\ \mu_{p-1} - \mu_p \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix},$$




Figure 5.3 Profile of a mean vector.

or as

$$H_0: \begin{pmatrix} \mu_1 - \mu_2 \\ \mu_1 - \mu_3 \\ \vdots \\ \mu_1 - \mu_p \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix},$$

These two expressions can be written in the form $H_0 : \mathbf{C}_1\boldsymbol{\mu} = \mathbf{0}$ and $H_0 : \mathbf{C}_2\boldsymbol{\mu} = \mathbf{0}$, where \mathbf{C}_1 and \mathbf{C}_2 are the $(p-1) \times p$ matrices:

$$\mathbf{C}_1 = \begin{pmatrix} 1 & -1 & 0 & \cdots & 0 \\ 0 & 1 & -1 & \cdots & 0 \\ \vdots & \vdots & \vdots & & \vdots \\ 0 & 0 & 0 & \cdots & -1 \end{pmatrix}, \quad \mathbf{C}_2 = \begin{pmatrix} 1 & -1 & 0 & \cdots & 0 \\ 1 & 0 & -1 & \cdots & 0 \\ \vdots & \vdots & \vdots & & \vdots \\ 1 & 0 & 0 & \cdots & -1 \end{pmatrix}.$$

In fact, any $(p-1) \times p$ matrix \mathbf{C} of rank $p-1$ such that $\mathbf{C}\mathbf{j} = \mathbf{0}$ can be used in $H_0 : \mathbf{C}\boldsymbol{\mu} = \mathbf{0}$ to produce $H_0 : \mu_1 = \mu_2 = \cdots = \mu_p$. If $\mathbf{C}\mathbf{j} = \mathbf{0}$, each row \mathbf{c}'_i of \mathbf{C} sums to zero by (2.38), and $\mathbf{C}\boldsymbol{\mu}$ is a set of $p-1$ contrasts in the μ 's. A linear combination $\mathbf{c}'_i\boldsymbol{\mu} = c_{i1}\mu_1 + c_{i2}\mu_2 + \cdots + c_{ip}\mu_p$ is called a *contrast* in the μ 's if the coefficients sum to zero, that is, if $\sum_j c_{ij} = 0$. The contrasts in $\mathbf{C}\boldsymbol{\mu}$ must be linearly independent in order to express $H_0 : \mu_1 = \mu_2 = \cdots = \mu_p$ as $H_0 : \mathbf{C}\boldsymbol{\mu} = \mathbf{0}$. Thus $\text{rank}(\mathbf{C}) = p-1$.

From a sample $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n$, we obtain estimates $\bar{\mathbf{y}}$ and \mathbf{S} of population parameters $\boldsymbol{\mu}$ and Σ . To test $H_0 : \mathbf{C}\boldsymbol{\mu} = \mathbf{0}$, we transform each $\mathbf{y}_i, i = 1, 2, \dots, n$, to $\mathbf{z}_i = \mathbf{C}\mathbf{y}_i$, which is $(p-1) \times 1$. By (3.62) and (3.64), the sample mean vector




Figure 5.4 Comparison of two profiles under the hypothesis of parallelism.

and covariance matrix of $\mathbf{z}_i = \mathbf{C}\mathbf{y}_i, i = 1, 2, \dots, n$, are $\bar{\mathbf{z}} = \mathbf{C}\bar{\mathbf{y}}$ and $\mathbf{S}_z = \mathbf{C}\mathbf{S}\mathbf{C}'$, respectively. If \mathbf{y} is $N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, then by property 1b in Section 4.2, $\mathbf{z} = \mathbf{C}\mathbf{y}$ is $N_{p-1}(\mathbf{C}\boldsymbol{\mu}, \mathbf{C}\boldsymbol{\Sigma}\mathbf{C}')$. Thus when $H_0: \mathbf{C}\boldsymbol{\mu} = \mathbf{0}$ is true, $\mathbf{C}\bar{\mathbf{y}}$ is $N_{p-1}(\mathbf{0}, \mathbf{C}\boldsymbol{\Sigma}\mathbf{C}'/n)$, and

$$T^2 = (\mathbf{C}\bar{\mathbf{y}})'(\mathbf{C}\mathbf{S}\mathbf{C}'/n)^{-1}(\mathbf{C}\bar{\mathbf{y}}) = n(\mathbf{C}\bar{\mathbf{y}})'(\mathbf{C}\mathbf{S}\mathbf{C}')^{-1}(\mathbf{C}\bar{\mathbf{y}}) \quad (5.33)$$

is distributed as $T_{p-1,n-1}^2$. We reject $H_0: \mathbf{C}\boldsymbol{\mu} = \mathbf{0}$ if $T^2 \geq T_{\alpha,p-1,n-1}^2$. The dimension $p-1$ corresponds to the number of rows of \mathbf{C} . Thus $\bar{\mathbf{z}} = \mathbf{C}\bar{\mathbf{y}}$ is $(p-1) \times 1$ and $\mathbf{S}_z = \mathbf{C}\mathbf{S}\mathbf{C}'$ is $(p-1) \times (p-1)$. Note that the \mathbf{C} 's in (5.33) don't "cancel" because \mathbf{C} is $(p-1) \times p$ and does not have an inverse. In fact, T^2 in (5.33) is less than $T^2 = n\bar{\mathbf{y}}'\mathbf{S}^{-1}\bar{\mathbf{y}}$ [see Rencher (1998, p. 84)].

If the variables have a natural ordering, as, for example, in the ramus bone data in Table 3.7, we could test for a linear trend or polynomial curve in the means by suitably choosing the rows of \mathbf{C} . This is discussed in connection with growth curves in Section 6.10. Otherwise, any comparisons of interest can be made as long as they are linearly independent.

5.9.2 Two-Sample Profile Analysis

Suppose two independent groups or samples receive the same set of p tests or measurements. If these tests are comparable, for example, all on a scale of 0 to 100, the variables will often be commensurate.

Rather than testing the hypothesis that $\mu_1 = \mu_2$, we wish to be more specific in comparing the profiles obtained by connecting the points $(j, \mu_{1j}), j = 1, 2, \dots, p$, and $(j, \mu_{2j}), j = 1, 2, \dots, p$. There are three hypotheses of interest in comparing the profiles of two samples. The first of these hypotheses addresses the question "Are the two profiles similar in appearance, or more precisely, are they parallel?" We illustrate this hypothesis in Figure 5.4. If the two profiles are parallel, then one group scored uniformly better than the other group on all p tests.

The parallelism hypothesis can be defined in terms of the slopes. The two profiles are parallel if the two slopes for each segment are the same. If the two profiles are parallel, the two increments for each segment are the same, and it is not necessary to use the actual slopes to express the hypothesis. We can simply compare the increase from one point to the next. The hypothesis can thus be expressed as $H_{01}: \mu_{1j} - \mu_{1,j-1} = \mu_{2j} - \mu_{2,j-1}$ for $j = 2, 3, \dots, p$, or

$$H_{01}: \begin{pmatrix} \mu_{12} - \mu_{11} \\ \mu_{13} - \mu_{12} \\ \vdots \\ \mu_{1p} - \mu_{1,p-1} \end{pmatrix} = \begin{pmatrix} \mu_{22} - \mu_{21} \\ \mu_{23} - \mu_{22} \\ \vdots \\ \mu_{2p} - \mu_{2,p-1} \end{pmatrix},$$

which can be written as $H_{01}: \mathbf{C}\boldsymbol{\mu}_1 = \mathbf{C}\boldsymbol{\mu}_2$, using the contrast matrix

$$\mathbf{C} = \begin{pmatrix} -1 & 1 & 0 & \cdots & 0 \\ 0 & -1 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & & \vdots \\ 0 & 0 & 0 & \cdots & 1 \end{pmatrix}.$$

From two samples, $\mathbf{y}_{11}, \mathbf{y}_{12}, \dots, \mathbf{y}_{1n_1}$ and $\mathbf{y}_{21}, \mathbf{y}_{22}, \dots, \mathbf{y}_{2n_2}$, we obtain $\bar{\mathbf{y}}_1, \bar{\mathbf{y}}_2$, and \mathbf{S}_{pl} as estimates of $\boldsymbol{\mu}_1, \boldsymbol{\mu}_2$, and $\boldsymbol{\Sigma}$. As in the two-sample T^2 -test, we assume that each \mathbf{y}_{1i} in the first sample is $N_p(\boldsymbol{\mu}_1, \boldsymbol{\Sigma})$, and each \mathbf{y}_{2i} in the second sample is $N_p(\boldsymbol{\mu}_2, \boldsymbol{\Sigma})$. If \mathbf{C} is a $(p-1) \times p$ contrast matrix as before, then $\mathbf{C}\bar{\mathbf{y}}_{1i}$ and $\mathbf{C}\bar{\mathbf{y}}_{2i}$ are distributed as $N_{p-1}(\mathbf{C}\boldsymbol{\mu}_1, \mathbf{C}\boldsymbol{\Sigma}\mathbf{C}')$ and $N_{p-1}(\mathbf{C}\boldsymbol{\mu}_2, \mathbf{C}\boldsymbol{\Sigma}\mathbf{C}')$, respectively. Under $H_{01}: \mathbf{C}\boldsymbol{\mu}_1 - \mathbf{C}\boldsymbol{\mu}_2 = \mathbf{0}$, the random vector $\mathbf{C}\bar{\mathbf{y}}_1 - \mathbf{C}\bar{\mathbf{y}}_2$ is $N_{p-1}[\mathbf{0}, \mathbf{C}\boldsymbol{\Sigma}\mathbf{C}'(1/n_1 + 1/n_2)]$ and

$$\begin{aligned} T^2 &= (\mathbf{C}\bar{\mathbf{y}}_1 - \mathbf{C}\bar{\mathbf{y}}_2)' \left[\left(\frac{1}{n_1} + \frac{1}{n_2} \right) \mathbf{C}\mathbf{S}_{\text{pl}}\mathbf{C}' \right]^{-1} (\mathbf{C}\bar{\mathbf{y}}_1 - \mathbf{C}\bar{\mathbf{y}}_2) \\ &= \frac{n_1 n_2}{n_1 + n_2} (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{C}' [\mathbf{C}\mathbf{S}_{\text{pl}}\mathbf{C}']^{-1} \mathbf{C} (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2) \end{aligned} \quad (5.34)$$

is distributed as T^2_{p-1, n_1+n_2-2} . Note that the dimension $p-1$ is the number of rows of \mathbf{C} .

By analogy with the discussion in Section 5.5, if H_{01} is rejected, we can follow up with univariate tests on the individual components of $\mathbf{C}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)$. Or alternatively we can calculate the discriminant function

$$\mathbf{a} = (\mathbf{C}\mathbf{S}_{\text{pl}}\mathbf{C}')^{-1} \mathbf{C} (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2) \quad (5.35)$$

as an indication of which slope differences contributed most to rejection of H_{01} in the presence of the other components of $\mathbf{C}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)$. There should be less need in this case to standardize the components of \mathbf{a} as suggested in Section 5.5, because the variables are assumed to be commensurate. The vector \mathbf{a} is $(p-1) \times 1$, corresponding to the $p-1$ segments of the profile. Thus if the second component of \mathbf{a} , for example,

Table 5.4 Data Layout for Two-Sample Profile Analysis

		Tests (variables)			
		1	2	...	p
Group 1					
\mathbf{y}'_{11}	=	$(y_{111} \quad y_{112} \quad \cdots \quad y_{11p})'$			
\mathbf{y}'_{12}	=	$(y_{121} \quad y_{122} \quad \cdots \quad y_{12p})'$			
\vdots		\vdots	\vdots		\vdots
\mathbf{y}'_{1n_1}	=	$(y_{1n_11} \quad y_{1n_12} \quad \cdots \quad y_{1n_1p})'$			
Group 2					
\mathbf{y}'_{21}	=	$(y_{211} \quad y_{212} \quad \cdots \quad y_{21p})'$			
\mathbf{y}'_{22}	=	$(y_{221} \quad y_{222} \quad \cdots \quad y_{22p})'$			
\vdots		\vdots	\vdots		\vdots
\mathbf{y}'_{2n_2}	=	$(y_{2n_21} \quad y_{2n_22} \quad \cdots \quad y_{2n_2p})'$			

is largest in absolute value, the divergence in slopes between the two profiles on the second segment contributes most to rejection of H_{01} .

If the data are arranged as in Table 5.4, we see an analogy to a two-way ANOVA model. A plot of the means is often made in a two-way ANOVA; a lack of parallelism corresponds to interaction between the two factors. Thus the hypothesis H_{01} is analogous to the group by test (variable) interaction hypothesis.

However, the usual ANOVA assumption of independence of observations does not hold here because the variables (tests) are correlated. The ANOVA assumption of independence and homogeneity of variances would require $\text{cov}(\mathbf{y}) = \Sigma = \sigma^2 \mathbf{I}$. Hence the test of H_{01} cannot be carried out using a univariate ANOVA approach, since $\Sigma \neq \sigma^2 \mathbf{I}$. We therefore proceed with the multivariate approach using T^2 .

The second hypothesis of interest in comparing two profiles is “Are the two populations or groups at the same *level*?” This hypothesis corresponds to a group (population) main effect in the ANOVA analogy. We can express this hypothesis in terms of the average level of group 1 compared to the average level of group 2:

$$H_{02}: \frac{\mu_{11} + \mu_{12} + \cdots + \mu_{1p}}{p} = \frac{\mu_{21} + \mu_{22} + \cdots + \mu_{2p}}{p}.$$

By (2.37), this can be expressed as

$$H_{02}: \mathbf{j}'\boldsymbol{\mu}_1 = \mathbf{j}'\boldsymbol{\mu}_2.$$

If H_{01} is true, H_{02} can be pictured as in Figure 5.5(a). If H_{02} is false, then the two profiles differ by a constant (given that H_{01} is true), as in Figure 5.5(b).

The hypothesis H_{02} can be true when H_{01} does not hold. Thus the average level of population 1 can equal the average level of population 2 without the two profiles being parallel, as illustrated in Figure 5.6. In this case, the “group main effect” is




Figure 5.5 Hypothesis H_{02} of equal group effect, assuming parallelism.




Figure 5.6 Hypothesis H_{02} of equal group effect without parallelism.

somewhat harder to interpret, as is the case in the analogous two-way ANOVA, where main effects are more difficult to describe in the presence of significant interaction. However, the test may still furnish useful information if a careful description of the results is provided.

To test $H_{02} : \mathbf{j}'(\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2) = 0$, we estimate $\mathbf{j}'(\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2)$ by $\mathbf{j}'(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)$, which is $N[0, \mathbf{j}'\Sigma\mathbf{j}(1/n_1 + 1/n_2)]$ when H_{02} is true. We can therefore use

$$t = \frac{\mathbf{j}'(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)}{\sqrt{\mathbf{j}'\mathbf{S}_p\mathbf{j}(1/n_1 + 1/n_2)}} \quad (5.36)$$




Figure 5.7 Hypothesis H_{03} of equal tests (variables) assuming parallelism.

and reject H_{02} if $|t| \geq t_{\alpha/2, n_1 + n_2 - 2}$.

The third hypothesis of interest, corresponding to the test (or variable) main effect, is “Are the profiles flat?” Assuming parallelism (assuming H_{01} is true), the “flatness” hypothesis can be pictured as in Figure 5.7. If H_{01} is not true, the test could be carried out separately for each group using the test in Section 5.9.1. If H_{01} and H_{02} are true, the two profiles in Figure 5.7(a) and Figure 5.7(b) will be coincident.

To express the third hypothesis in a form suitable for testing, we note from Figure 5.7(a) that the average of the two group means is the same for each test:

$$H_{03}: \frac{1}{2}(\mu_{11} + \mu_{21}) = \frac{1}{2}(\mu_{12} + \mu_{22}) = \cdots = \frac{1}{2}(\mu_{1p} + \mu_{2p}) \quad (5.37)$$

or

$$H_{03}: \frac{1}{2} \mathbf{C}(\boldsymbol{\mu}_1 + \boldsymbol{\mu}_2) = \mathbf{0}, \quad (5.38)$$

where \mathbf{C} is a $(p - 1) \times p$ matrix such that $\mathbf{C}\mathbf{j} = \mathbf{0}$. From Figure 5.7(a), we see that if H_{01} is true, H_{03} could also be expressed as $\mu_{11} = \mu_{12} = \cdots = \mu_{1p}$ and $\mu_{21} = \mu_{22} = \cdots = \mu_{2p}$ or

$$H_{03}: \mathbf{C}\boldsymbol{\mu}_1 = \mathbf{0} \quad \text{and} \quad \mathbf{C}\boldsymbol{\mu}_2 = \mathbf{0}.$$

To estimate $\frac{1}{2}(\boldsymbol{\mu}_1 + \boldsymbol{\mu}_2)$, we use the sample grand mean vector based on a weighted average:

$$\bar{\mathbf{y}} = \frac{n_1 \bar{\mathbf{y}}_1 + n_2 \bar{\mathbf{y}}_2}{n_1 + n_2}.$$

It can easily be shown that under H_{03} , $E(\mathbf{C}\bar{\mathbf{y}}) = \mathbf{0}$ and $\text{cov}(\bar{\mathbf{y}}) = \boldsymbol{\Sigma}/(n_1 + n_2)$. Therefore, $\mathbf{C}\bar{\mathbf{y}}$ is $N_{p-1}[0, \mathbf{C}\boldsymbol{\Sigma}\mathbf{C}'/(n_1 + n_2)]$, and

$$\begin{aligned} T^2 &= (\mathbf{C}\bar{\mathbf{y}})' \left(\frac{\mathbf{C}\mathbf{S}_{\text{pl}}\mathbf{C}'}{n_1 + n_2} \right)^{-1} (\mathbf{C}\bar{\mathbf{y}}) \\ &= (n_1 + n_2)(\mathbf{C}\bar{\mathbf{y}})' (\mathbf{C}\mathbf{S}_{\text{pl}}\mathbf{C}')^{-1} \mathbf{C}\bar{\mathbf{y}} \end{aligned} \quad (5.39)$$

is distributed as $T_{p-1, n_1 + n_2 - 2}^2$ when H_{03} is true. It can be readily shown that H_{03} is unaffected by a difference in the profile levels (unaffected by the status of H_{02}).




Figure 5.8 Profiles for the psychological data in Table 5.1.

■ EXAMPLE 5.9.2

We use the psychological data in Table 5.1 to illustrate two-sample profile analysis. The values of \bar{y}_1 , \bar{y}_2 , and S_{pl} are given in Example 5.4.2. The profiles of the two mean vectors \bar{y}_1 and \bar{y}_2 are plotted in Figure 5.8. There appears to be a lack of parallelism.

To test for parallelism, $H_{01}: \mathbf{C}\mu_1 = \mathbf{C}\mu_2$, we use the matrix

$$\mathbf{C} = \begin{pmatrix} -1 & 1 & 0 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 0 & -1 & 1 \end{pmatrix}$$

and obtain

$$\mathbf{C}(\bar{y}_1 - \bar{y}_2) = \begin{pmatrix} -1.62 \\ 8.53 \\ -9.72 \end{pmatrix}, \quad \mathbf{C}S_{pl}\mathbf{C}' = \begin{pmatrix} 10.96 & -7.05 & -1.64 \\ -7.05 & 27.26 & -12.74 \\ -1.64 & -12.74 & 23.72 \end{pmatrix}.$$

Then, by (5.34),

$$T^2 = \frac{(32)(32)}{32+32} (\bar{y}_1 - \bar{y}_2)' \mathbf{C}' (\mathbf{C}S_{pl}\mathbf{C}')^{-1} \mathbf{C} (\bar{y}_1 - \bar{y}_2) = 74.240.$$

Upon comparison of this value with $T^2_{01,3,62} = 12.796$ (obtained by interpolation in Table A.7), we reject the hypothesis of parallelism.

In Figure 5.8 the lack of parallelism is most notable in the second and third segments. This can also be seen in the relatively large values of the second and

third components of

$$\mathbf{C}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2) = \begin{pmatrix} -1.62 \\ 8.53 \\ -9.72 \end{pmatrix}.$$

To see which of these made the greatest statistical contribution, we can examine the discriminant function coefficient vector given in (5.35) as

$$\mathbf{a} = (\mathbf{CS}_{\text{pl}}\mathbf{C}')^{-1}\mathbf{C}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2) = \begin{pmatrix} -.136 \\ .104 \\ -.363 \end{pmatrix}.$$

Thus the third segment contributed most to rejection in the presence of the other two segments.

To test for equal levels, $H_{02}: \mathbf{j}'\boldsymbol{\mu}_1 = \mathbf{j}'\boldsymbol{\mu}_2$, we use (5.36),

$$\begin{aligned} t &= \frac{\mathbf{j}'(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)}{\sqrt{\mathbf{j}'\mathbf{S}_{\text{pl}}\mathbf{j}(1/n_1 + 1/n_2)}} \\ &= \frac{16.969}{\sqrt{(164.276)(1/32 + 1/32)}} = 5.2957. \end{aligned}$$

Comparing this with $t_{.005,62} = 2.658$, we reject the hypothesis of equal levels.

To test the flatness hypothesis, $H_{03}: \frac{1}{2}\mathbf{C}(\boldsymbol{\mu}_1 + \boldsymbol{\mu}_2) = \mathbf{0}$, we first calculate

$$\bar{\mathbf{y}} = \frac{32\bar{\mathbf{y}}_1 + 32\bar{\mathbf{y}}_2}{32 + 32} = \frac{\bar{\mathbf{y}}_1 + \bar{\mathbf{y}}_2}{2} = \begin{pmatrix} 14.16 \\ 14.91 \\ 21.92 \\ 22.34 \end{pmatrix}.$$

Using

$$\mathbf{C} = \begin{pmatrix} 1 & -1 & 0 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 1 & -1 \end{pmatrix},$$

we obtain, by (5.39),

$$T^2 = (32 + 32)(\mathbf{C}\bar{\mathbf{y}})'(\mathbf{CS}_{\text{pl}}\mathbf{C})^{-1}\mathbf{C}\bar{\mathbf{y}} = 254.004,$$

which exceeds $T_{.01,3,62}^2 = 12.796$, so we reject the hypothesis of flatness. However, since the parallelism hypothesis was rejected, a more appropriate approach would be to test each of the two groups separately for flatness using the test of Section 5.9.1. By (5.33), we obtain

$$\begin{aligned} T^2 &= n_1(\mathbf{C}\bar{\mathbf{y}}_1)'(\mathbf{CS}_1\mathbf{C})^{-1}(\mathbf{C}\bar{\mathbf{y}}_1) = 221.126, \\ T^2 &= n_2(\mathbf{C}\bar{\mathbf{y}}_2)'(\mathbf{CS}_2\mathbf{C})^{-1}(\mathbf{C}\bar{\mathbf{y}}_2) = 103.483. \end{aligned}$$

Both of these exceed $T_{.01,3,31}^2 = 14.626$, and we have significant lack of flatness. \square

PROBLEMS

- 5.1** Show that the characteristic form of T^2 in (5.6) is the same as the original form in (5.5).
- 5.2** Show that the T^2 -statistic in (5.9) can be expressed in the characteristic form given in (5.10).
- 5.3** Show that $t^2(\mathbf{a}) = T^2$, where $t(\mathbf{a})$ is given by (5.13), T^2 is given by (5.9), and $\mathbf{a} = \mathbf{S}_{\text{pl}}^{-1}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)$ as in (5.14).
- 5.4** Show that the paired observation t -test in (5.22), $t = \bar{d}/(s_d/\sqrt{n})$, has the t_{n-1} distribution.
- 5.5** Show that $s_d^2 = \sum_{i=1}^n (d_i - \bar{d})^2/(n-1) = s_y^2 + s_x^2 - 2s_{yx}$, as in a comparison of (5.22) and (5.23).
- 5.6** Show that $T^2 = n\bar{\mathbf{d}}' \mathbf{S}_d^{-1} \bar{\mathbf{d}}$ in (5.24) has the characteristic form given by $T^2 = \bar{\mathbf{d}}' (\mathbf{S}_d/n)^{-1} \bar{\mathbf{d}}$.
- 5.7** Use (5.7) to show that $T^2(\mathbf{x}|\mathbf{y})$ in (5.29) can be converted to F as in (5.30).
- 5.8** Show that the test statistic in (5.30) for additional information in \mathbf{x} above and beyond \mathbf{y} has an F -distribution by solving for R^2 in terms of T^2 from (5.20) and substituting this into (5.31).
- 5.9** In Section 5.9.2, show that under H_{03} and H_{01} , $E(\mathbf{C}\bar{\mathbf{y}}) = \mathbf{0}$ and $\text{cov}(\bar{\mathbf{y}}) = \Sigma/(n_1 + n_2)$, where $\bar{\mathbf{y}} = (n_1\bar{\mathbf{y}}_1 + n_2\bar{\mathbf{y}}_2)/(n_1 + n_2)$ and Σ is the common covariance matrix of the two populations from which $\bar{\mathbf{y}}_1$ and $\bar{\mathbf{y}}_2$ are sampled.
- 5.10** Verify that $T^2 = (n_1 + n_2)(\mathbf{C}\bar{\mathbf{y}})'(\mathbf{C}\mathbf{S}_{\text{pl}}\mathbf{C}')^{-1}\mathbf{C}\bar{\mathbf{y}}$ in (5.39) is distributed as $T_{p-1, n_1+n_2-2}^2$.
- 5.11** Test $H_0: \boldsymbol{\mu}' = (6, 11)$ using the data
- $$\mathbf{Y} = \begin{pmatrix} 3 & 10 \\ 6 & 12 \\ 5 & 14 \\ 10 & 9 \end{pmatrix}.$$
- 5.12** Use the probe word data in Table 3.6:
- Test $H_0: \boldsymbol{\mu} = (30, 25, 40, 25, 30)'$.
 - If H_0 is rejected, test each variable separately, using (5.3).
- 5.13** For the probe word data in Table 3.6, test $H_0: \mu_1 = \mu_2 = \dots = \mu_5$, using T^2 in (5.33).
- 5.14** Use the ramus bone data in Table 3.7:

Table 5.5 Four Measurements on Two Species of Flea Beetles

<i>Haltica oleracea</i>					<i>Haltica carduorum</i>				
Experiment Number	y_1	y_2	y_3	y_4	Experiment Number	y_1	y_2	y_3	y_4
1	189	245	137	163	1	181	305	184	209
2	192	260	132	217	2	158	237	133	188
3	217	276	141	192	3	184	300	166	231
4	221	299	142	213	4	171	273	162	213
5	171	239	128	158	5	181	297	163	224
6	192	262	147	173	6	181	308	160	223
7	213	278	136	201	7	177	301	166	221
8	192	255	128	185	8	198	308	141	197
9	170	244	128	192	9	180	286	146	214
10	201	276	146	186	10	177	299	171	192
11	195	242	128	192	11	176	317	166	213
12	205	263	147	192	12	192	312	166	209
13	180	252	121	167	13	176	285	141	200
14	192	283	138	183	14	169	287	162	214
15	200	294	138	188	15	164	265	147	192
16	192	277	150	177	16	181	308	157	204
17	200	287	136	173	17	192	276	154	209
18	181	255	146	183	18	181	278	149	235
19	192	287	141	198	19	175	271	140	192
					20	197	303	170	205

(a) Test $H_0: \mu = (48, 49, 50, 51)'$.

(b) If H_0 is rejected, test each variable separately, using (5.3).

5.15 For the ramus bone data in Table 3.7, test $H_0: \mu_1 = \mu_2 = \mu_3 = \mu_4$, using T^2 in (5.33).

5.16 Four measurements were made on two species of flea beetles (Lubischew 1962). The variables were

y_1 = distance of transverse groove from posterior border of prothorax (μm)

y_2 = length of elytra (in 0.01 mm)

y_3 = length of second antennal joint (μm)

y_4 = length of third antennal joint (μm)

The data are given in Table 5.5.

(a) Test $H_0: \mu_1 = \mu_2$ using T^2 .

Table 5.6 Comparison of Six Tests on Engineer Apprentices and Pilots

Engineer Apprentices						Pilots					
y_1	y_2	y_3	y_4	y_5	y_6	y_1	y_2	y_3	y_4	y_5	y_6
121	22	74	223	54	254	132	17	77	232	50	249
108	30	80	175	40	300	123	32	79	192	64	315
122	49	87	266	41	223	129	31	96	250	55	319
77	37	66	178	80	209	131	23	67	291	48	310
140	35	71	175	38	261	110	24	96	239	42	268
108	37	57	241	59	245	47	22	87	231	40	217
124	39	52	194	72	242	125	32	87	227	30	324
130	34	89	200	85	242	129	29	102	234	58	300
149	55	91	198	50	277	130	26	104	256	58	270
129	38	72	162	47	268	147	47	82	240	30	322
154	37	87	170	60	244	159	37	80	227	58	317
145	33	88	208	51	228	135	41	83	216	39	306
112	40	60	232	29	279	100	35	83	183	57	242
120	39	73	159	39	233	149	37	94	227	30	240
118	21	83	152	88	233	149	38	78	258	42	271
141	42	80	195	36	241	153	27	89	283	66	291
135	49	73	152	42	249	136	31	83	257	31	311
151	37	76	223	74	268	97	36	100	252	30	225
97	46	83	164	31	243	141	37	105	250	27	243
109	42	82	188	57	267	164	32	76	187	30	264

- (b) If the T^2 -test in part (a) rejects H_0 , carry out a t -test on each variable, as in (5.15).
- (c) Calculate the discriminant function coefficient vector $\mathbf{a} = \mathbf{S}_{\text{pl}}^{-1}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)$.
- (d) Show that if the vector \mathbf{a} found in part (c) is substituted into $t^2(\mathbf{a})$ from (5.13), the result is the same as the value of T^2 found in part (a).
- (e) Obtain T^2 using the regression approach in Section 5.6.2.
- (f) Test the significance of each variable adjusted for the other three.
- (g) Test the significance of y_3 and y_4 adjusted for y_1 and y_2 .

5.17 Carry out a profile analysis on the beetle data in Table 5.5.

5.18 Twenty engineer apprentices and 20 pilots were given six tests (Travers 1939). The variables were

$$\begin{array}{ll} y_1 = \text{intelligence} & y_4 = \text{dotting} \\ y_2 = \text{form relations} & y_5 = \text{sensory motor coordination} \\ y_3 = \text{dynamometer} & y_6 = \text{perseveration} \end{array}$$

The data are given in Table 5.6.

- (a) Test $H_0: \mu_1 = \mu_2$.
- (b) If the T^2 -test in part (a) rejects H_0 , carry out a t -test for each variable, as in (5.15).
- (c) Test each variable adjusted for the other five.
- (d) Test the significance of y_4, y_5, y_6 adjusted for y_1, y_2, y_3 .
- 5.19** Data were collected in an attempt to find a screening procedure to detect carriers of Duchenne muscular dystrophy, a disease transmitted from female carriers to some of their male offspring (Andrews and Herzberg 1985, pp. 223–228). The following variables were measured on a sample of noncarriers and a sample of carriers:

y_1 = age

y_2 = month in which measurements are taken

y_3 = creatine kinase

y_4 = hemopexin

y_5 = lactate dehydrogenase

y_6 = pyruvate kinase

The data are given in Table 5.7.

- (a) Test $H_0: \mu_1 = \mu_2$ using y_3, y_4, y_5 , and y_6 .
- (b) The variables y_3 and y_4 are relatively inexpensive to measure compared to y_5 and y_6 . Do y_5 and y_6 contribute an important amount to T^2 above and beyond y_3 and y_4 ?
- (c) The levels of y_3, y_4, y_5 , and y_6 may depend on age and season, y_1 and y_2 . Do y_1 and y_2 contribute a significant amount to T^2 when adjusted for y_3, y_4, y_5 , and y_6 ?
- 5.20** Various aspects of economic cycles were measured for consumer goods and producer goods by Tintner (1946). The variables are

y_1 = length of cycle

y_2 = percentage of rising prices

y_3 = cyclical amplitude

y_4 = rate of change

The data for several items are given in Table 5.8.

- (a) Test $H_0: \mu_1 = \mu_2$ using T^2 .
- (b) Calculate the discriminant function coefficient vector.
- (c) Test for significance of each variable adjusted for the other three.

Table 5.7 Comparison of Carriers and Noncarriers of Muscular Dystrophy

Noncarriers						Carriers					
y_1	y_2	y_3	y_4	y_5	y_6	y_1	y_2	y_3	y_4	y_5	y_6
22	6	52	83.5	10.9	176	30	10	167	89.0	25.6	364
32	8	20	77.0	11.0	200	41	10	104	81.0	26.8	245
36	7	28	86.5	13.2	171	22	8	30	108.0	8.8	284
22	11	30	104.0	22.6	230	22	8	44	104.0	17.4	172
23	1	40	83.0	15.2	205	20	10	65	87.0	23.8	198
30	5	24	78.8	9.6	151	42	9	440	107.0	20.2	239
27	8	15	87.0	13.5	232	59	8	58	88.2	11.0	259
30	11	22	91.0	17.5	198	35	9	129	93.1	18.3	188
25	10	42	65.5	13.3	216	36	6	104	87.5	16.7	256
26	2	130	80.3	17.1	211	35	2	122	88.5	21.6	263
26	3	48	85.2	22.7	160	29	4	265	83.5	16.1	136
27	7	31	86.5	6.9	162	27	4	285	79.5	36.4	245
26	10	47	53.0	14.6	131	27	9	25	91.0	49.1	209
27	3	36	56.0	18.2	105	28	4	124	92.0	32.2	298
27	7	24	57.5	5.6	130	29	8	53	76.0	14.0	174
31	4	34	92.7	7.9	140	30	2	46	71.0	16.9	197
31	9	38	96.0	12.6	158	30	7	40	85.5	12.7	201
35	10	40	104.6	16.1	209	30	8	41	90.0	9.7	342
28	4	59	88.0	9.9	128	31	6	657	104.0	110.0	358
28	8	75	81.0	10.1	177	32	2	465	86.5	63.7	412
28	9	72	66.3	16.4	156	32	5	485	83.5	73.0	382
27	7	42	77.0	15.3	163	37	2	168	82.5	23.3	261
27	3	30	80.2	8.1	100	38	6	286	109.5	31.9	260
28	6	24	87.0	3.5	132	39	1	388	91.0	41.6	204
24	9	26	84.5	20.7	145	39	9	148	105.2	18.8	221
23	8	65	75.0	19.9	187	34	6	73	105.5	17.0	285
27	3	34	86.3	11.8	120	35	4	36	92.8	22.0	308
25	2	37	73.3	13.0	254	58	8	19	100.5	10.9	196
34	3	73	57.4	7.4	107	58	2	34	98.5	19.9	299
34	7	87	76.3	6.0	87	38	1	113	97.0	18.8	216
25	7	35	71.0	8.8	186	30	8	57	105.0	12.9	155
20	7	31	61.5	9.9	172	42	8	78	118.0	15.5	212
20	5	62	81.0	10.2	181	43	11	73	104.0	20.6	201
31	6	48	79.0	16.8	182	29	3	69	111.0	16.0	175
31	7	40	82.5	6.4	151						
26	7	55	85.5	10.9	216						
26	7	32	73.8	8.6	147						
21	11	26	79.3	16.4	123						
27	6	25	91.0	10.3	135						

Table 5.8 Cyclical Measurements of Consumer Goods and Producer Goods

Consumer Goods					Producer Goods				
Item	y_1	y_2	y_3	y_4	Item	y_1	y_2	y_3	y_4
1	72	50	8	0.5	1	57	57	12.5	0.9
2	66.5	48	15	1.0	2	100	54	17	0.5
3	54	57	14	1.0	3	100	32	16.5	0.7
4	67	60	15	0.9	4	96.5	65	20.5	0.9
5	44	57	14	0.3	5	79	51	18	0.9
6	41	52	18	1.9	6	78.5	53	18	1.2
7	34.5	50	4	0.5	7	48	50	21	1.6
8	34.5	46	8.5	1.0	8	155	44	20.5	1.4
9	24	54	3	1.2	9	84	64	13	0.8
					10	105	35	17	1.8

- 5.21** Each of 15 students wrote an informal and a formal essay Kramer (1972, p. 100). The variables recorded were the number of words and the number of verbs:

y_1 = number of words in the informal essay

y_2 = number of verbs in the informal essay

x_1 = number of words in the formal essay

x_2 = number of verbs in the formal essay

The data are given in Table 5.9. Since each student wrote both types of essays, the observation vectors are paired, and we use the paired comparison test.

- (a) Test $H_0: \mu_d = 0$.
- (b) Find the discriminant function coefficient vector.
- (c) Do a univariate t -test on each d_j .

- 5.22** A number of patients with bronchus cancer were treated with ascorbate and compared with matched patients who received no ascorbate (Cameron and Pauling 1978). The data are given in Table 5.10. The variables measured were

y_1, x_1 = survival time (days) from date of first hospital admission

y_2, x_2 = survival time from date of untreatability

Compare y_1 and y_2 with x_1 and x_2 using a paired comparison T^2 -test.

- 5.23** Use the glucose data in Table 3.9:

- (a) Test $H_0: \mu_y = \mu_x$ using a paired comparison test.
- (b) Test the significance of each variable adjusted for the other two.

Table 5.9 Number of Words and Number of Verbs

Student	Informal		Formal		$d_1 = y_1 - x_1$	$d_2 = y_2 - x_2$
	Words y_1	Verbs y_2	Words x_1	Verbs x_2		
1	148	20	137	15	+11	+5
2	159	24	164	25	-5	-1
3	144	19	224	27	-80	-8
4	103	18	208	33	-105	-15
5	121	17	178	24	-57	-7
6	89	11	128	20	-39	-9
7	119	17	154	18	-35	-1
8	123	13	158	16	-35	-3
9	76	16	102	21	-26	-5
10	217	29	214	25	+3	+4
11	148	22	209	24	-61	-2
12	151	21	151	16	0	+5
13	83	7	123	13	-40	-6
14	135	20	161	22	-26	-2
15	178	15	175	23	+3	-8

Table 5.10 Survival Times for Bronchus Cancer Patients and Matched Controls

Ascorbate Patients		Matched Controls	
y_1	y_2	x_1	x_2
81	74	72	33
461	423	134	18
20	16	84	20
450	450	98	58
246	87	48	13
166	115	142	49
63	50	113	38
64	50	90	24
155	113	30	18
151	38	260	34
166	156	116	20
37	27	87	27
223	218	69	32
138	138	100	27
72	39	315	39
245	231	188	65

CHAPTER 6

MULTIVARIATE ANALYSIS OF VARIANCE

In this chapter we extend univariate analysis of variance to multivariate analysis of variance, in which we measure more than one variable on each experimental unit. For multivariate analysis of covariance, see Rencher (1998, Section 4.10).

6.1 ONE-WAY MODELS

We begin with a review of univariate analysis of variance (ANOVA) before covering multivariate analysis of variance (MANOVA) with several dependent variables.

6.1.1 Univariate One-Way Analysis of Variance (ANOVA)

In the balanced one-way ANOVA, we have a random sample of n observations from each of k normal populations with equal variances, as in the following layout:

	Sample 1 from $N(\mu_1, \sigma^2)$	Sample 2 from $N(\mu_2, \sigma^2)$...	Sample k from $N(\mu_k, \sigma^2)$
	y_{11}	y_{21}	...	y_{k1}
	y_{12}	y_{22}	...	y_{k2}
	\vdots	\vdots		\vdots
	y_{1n}	y_{2n}	...	y_{kn}
Total	$y_{1\cdot}$	$y_{2\cdot}$...	$y_{k\cdot}$
Mean	$\bar{y}_{1\cdot}$	$\bar{y}_{2\cdot}$...	$\bar{y}_{k\cdot}$
Variance	s_1^2	s_2^2	...	s_k^2

The k samples or the populations from which they arise are sometimes referred to as *groups*. The groups may correspond to *treatments* applied by the researcher in an experiment. We have used the “dot” notation for totals and means for each group:

$$y_{i\cdot} = \sum_{j=1}^n y_{ij}, \quad \bar{y}_{i\cdot} = \sum_{j=1}^n \frac{y_{ij}}{n}. \quad (6.1)$$

The k samples are assumed to be independent. The assumptions of independence and common variance are necessary to obtain an F -test.

The model for each observation is

$$\begin{aligned} y_{ij} &= \mu + \alpha_i + \varepsilon_{ij} \\ &= \mu_i + \varepsilon_{ij}, \quad i = 1, 2, \dots, k; j = 1, 2, \dots, n; \end{aligned} \quad (6.2)$$

where $\mu_i = \mu + \alpha_i$ is the mean of the i th population. We wish to compare the sample means $\bar{y}_{i\cdot}$, $i = 1, 2, \dots, k$, to see if they are sufficiently different to lead us to believe the population means differ. The hypothesis can be expressed as $H_0: \mu_1 = \mu_2 = \dots = \mu_k$. Note that the notation for subscripts differs from that of previous chapters, in which the subscript i represented the observation. In this chapter, we use the last subscript in a model such as (6.2) to represent the observation.

If the hypothesis is true, all y_{ij} are from the same population, $N(\mu, \sigma^2)$, and we can obtain two estimates of σ^2 , one based on the sample variances $s_1^2, s_2^2, \dots, s_k^2$ [see (3.4) and (3.5)] and the other based on the sample means $\bar{y}_1, \bar{y}_2, \dots, \bar{y}_k$. The pooled “within-sample” estimator of σ^2 is

$$s_e^2 = \frac{1}{k} \sum_{i=1}^k s_i^2 = \frac{\sum_{i=1}^k \sum_{j=1}^n (y_{ij} - \bar{y}_{i\cdot})^2}{k(n-1)} = \frac{\sum_{ij} y_{ij}^2 - \sum_i y_{i\cdot}^2/n}{k(n-1)}. \quad (6.3)$$

Our second estimate of σ^2 (under H_0) is based on the variance of the sample means,

$$s_{\bar{y}}^2 = \frac{\sum_{i=1}^k (\bar{y}_{i\cdot} - \bar{y}_{..})^2}{k-1}, \quad (6.4)$$

where $\bar{y}_{..} = \sum_{i=1}^k \bar{y}_{i\cdot}/k$ is the overall mean. If H_0 is true, $s_{\bar{y}}^2$ estimates $\sigma_{\bar{y}}^2 = \sigma^2/n$ [see remarks following (3.1) in Section 3.1], and therefore $E(ns_{\bar{y}}^2) = n(\sigma^2/n) =$

σ^2 , from which the estimate of σ^2 is

$$ns_y^2 = \frac{n \sum_{i=1}^k (\bar{y}_{i\cdot} - \bar{y}_{..})^2}{k-1} = \frac{\sum_i y_{i\cdot}^2/n - y_{..}^2/kn}{k-1}, \quad (6.5)$$

where $y_{..} = \sum_i y_{i\cdot} = \sum_{ij} y_{ij}$ is the overall total. If H_0 is false, $E(ns_y^2) = \sigma^2 + n \sum_i \alpha_i^2/(k-1)$, and ns_y^2 will tend to reflect a larger spread in $\bar{y}_{1\cdot}, \bar{y}_{2\cdot}, \dots, \bar{y}_{k\cdot}$. Since s_e^2 is based on variability within each sample, it estimates σ^2 whether or not H_0 is true; thus $E(s_e^2) = \sigma^2$ in either case.

When sampling from normal distributions, s_e^2 , a pooled estimator based on the k values of s_i^2 , is independent of s_y^2 , which is based on the $\bar{y}_{i\cdot}$'s. We can justify this assertion by noting that $\bar{y}_{i\cdot}$ and s_i^2 are independent in each sample (when sampling from the normal distribution) and that the k samples are independent of each other. Since ns_y^2 and s_e^2 are independent and both estimate σ^2 , their ratio forms an F -statistic (see Section 7.3.1):

$$\begin{aligned} F &= \frac{ns_y^2}{s_e^2} = \frac{(\sum_i y_{i\cdot}^2/n - y_{..}^2/kn)/(k-1)}{(\sum_{ij} y_{ij}^2 - \sum_i y_{i\cdot}^2/n)/[k(n-1)]} \\ &= \frac{\text{SSH}/(k-1)}{\text{SSE}/[k(n-1)]} \end{aligned} \quad (6.6)$$

$$= \frac{\text{MSH}}{\text{MSE}}, \quad (6.7)$$

where $\text{SSH} = \sum_i y_{i\cdot}^2/n - y_{..}^2/kn$ and $\text{SSE} = \sum_{ij} y_{ij}^2 - \sum_i y_{i\cdot}^2/n$ are the “between”-sample sum of squares (due to the means) and “within”-sample sum of squares, respectively, and MSH and MSE are the corresponding sample mean squares. The F -statistic (6.6) is distributed as $F_{k-1, k(n-1)}$ when H_0 is true. We reject H_0 if $F > F_\alpha$. The F -statistic (6.6) can be shown to be a simple function of the likelihood ratio.

6.1.2 Multivariate One-Way Analysis of Variance Model (MANOVA)

We often measure several dependent variables on each experimental unit instead of just one variable. In the multivariate case, we assume that k independent random samples of size n are obtained from p -variate normal populations with equal covariance matrices, as in the following layout for balanced one-way multivariate analysis of variance (MANOVA). (In practice, the observation vectors \mathbf{y}_{ij} would ordinarily be listed in row form, and sample 2 would appear below sample 1, and so on. See, for example, Table 6.2.)

	Sample 1 from $N_p(\mu_1, \Sigma)$	Sample 2 from $N_p(\mu_2, \Sigma)$...	Sample k from $N_p(\mu_k, \Sigma)$
	\mathbf{y}_{11}	\mathbf{y}_{21}	...	\mathbf{y}_{k1}
	\mathbf{y}_{12}	\mathbf{y}_{22}	...	\mathbf{y}_{k2}
	\vdots	\vdots		\vdots
	\mathbf{y}_{1n}	\mathbf{y}_{2n}	...	\mathbf{y}_{kn}
Total	$\mathbf{y}_{1\cdot}$	$\mathbf{y}_{2\cdot}$...	$\mathbf{y}_{k\cdot}$
Mean	$\bar{\mathbf{y}}_{1\cdot}$	$\bar{\mathbf{y}}_{2\cdot}$...	$\bar{\mathbf{y}}_{k\cdot}$

Totals and means are defined as follows:

$$\text{Total of the } i\text{th sample: } \mathbf{y}_{i\cdot} = \sum_{j=1}^n \mathbf{y}_{ij}.$$

$$\text{Overall total: } \mathbf{y}_{\cdot\cdot} = \sum_{i=1}^k \sum_{j=1}^n \mathbf{y}_{ij}.$$

$$\text{Mean of the } i\text{th sample: } \bar{\mathbf{y}}_{i\cdot} = \mathbf{y}_{i\cdot}/n.$$

$$\text{Overall mean: } \bar{\mathbf{y}}_{\cdot\cdot} = \mathbf{y}_{\cdot\cdot}/kn.$$

The model for each observation vector is

$$\begin{aligned} \mathbf{y}_{ij} &= \boldsymbol{\mu} + \boldsymbol{\alpha}_i + \boldsymbol{\varepsilon}_{ij} \\ &= \boldsymbol{\mu}_i + \boldsymbol{\varepsilon}_{ij}, \quad i = 1, 2, \dots, k; \quad j = 1, 2, \dots, n. \end{aligned} \tag{6.8}$$

In terms of the p variables in \mathbf{y}_{ij} , (6.8) becomes

$$\begin{pmatrix} y_{ij1} \\ y_{ij2} \\ \vdots \\ y_{ijp} \end{pmatrix} = \begin{pmatrix} \mu_1 \\ \mu_2 \\ \vdots \\ \mu_p \end{pmatrix} + \begin{pmatrix} \alpha_{i1} \\ \alpha_{i2} \\ \vdots \\ \alpha_{ip} \end{pmatrix} + \begin{pmatrix} \varepsilon_{ij1} \\ \varepsilon_{ij2} \\ \vdots \\ \varepsilon_{ijp} \end{pmatrix} = \begin{pmatrix} \mu_{i1} \\ \mu_{i2} \\ \vdots \\ \mu_{ip} \end{pmatrix} + \begin{pmatrix} \varepsilon_{ij1} \\ \varepsilon_{ij2} \\ \vdots \\ \varepsilon_{ijp} \end{pmatrix},$$

so that the model for the r th variable ($r = 1, 2, \dots, p$) in each vector \mathbf{y}_{ij} is

$$y_{ijr} = \mu_r + \alpha_{ir} + \varepsilon_{ijr} = \mu_{ir} + \varepsilon_{ijr}.$$

We wish to compare the mean vectors of the k samples for significant differences. The hypothesis is therefore

$$H_0: \boldsymbol{\mu}_1 = \boldsymbol{\mu}_2 = \cdots = \boldsymbol{\mu}_k \quad \text{vs.} \quad H_1: \text{at least two } \boldsymbol{\mu}\text{'s are unequal.}$$

Equality of the mean vectors implies that the k means are equal for each variable; that is, $\mu_{1r} = \mu_{2r} = \cdots = \mu_{kr}$ for $r = 1, 2, \dots, p$. If two means differ for just one variable, for example, $\mu_{23} \neq \mu_{43}$, then H_0 is false and we wish to reject it. We can see this by examining the elements of the population mean vectors:

$$H_0: \begin{pmatrix} \mu_{11} \\ \mu_{12} \\ \vdots \\ \mu_{1p} \end{pmatrix} = \begin{pmatrix} \mu_{21} \\ \mu_{22} \\ \vdots \\ \mu_{2p} \end{pmatrix} = \cdots = \begin{pmatrix} \mu_{k1} \\ \mu_{k2} \\ \vdots \\ \mu_{kp} \end{pmatrix}.$$

Thus H_0 implies p sets of equalities:

$$\begin{aligned}\mu_{11} &= \mu_{21} = \cdots = \mu_{k1} \\ \mu_{12} &= \mu_{22} = \cdots = \mu_{k2} \\ &\vdots & \vdots & \vdots \\ \mu_{1p} &= \mu_{2p} = \cdots = \mu_{kp}.\end{aligned}$$

All $p(k - 1)$ equalities must hold for H_0 to be true; failure of only one equality will falsify the hypothesis.

In the univariate case, we have “between” and “within” sums of squares SSH and SSE. By (6.3), (6.5), and (6.6), these are given by

$$\begin{aligned}\text{SSH} &= n \sum_{i=1}^k (\bar{y}_{i\cdot} - \bar{y}_{..})^2 = \sum_{i=1}^k \frac{y_{i\cdot}^2}{n} - \frac{y_{..}^2}{kn}, \\ \text{SSE} &= \sum_{i=1}^k \sum_{j=1}^n (y_{ij} - \bar{y}_{i\cdot})^2 = \sum_{ij} y_{ij}^2 - \sum_i \frac{y_{i\cdot}^2}{n}.\end{aligned}$$

By analogy, in the multivariate case, we have “between” and “within” matrices \mathbf{H} and \mathbf{E} , defined as

$$\mathbf{H} = n \sum_{i=1}^k (\bar{\mathbf{y}}_{i\cdot} - \bar{\mathbf{y}}_{..})(\bar{\mathbf{y}}_{i\cdot} - \bar{\mathbf{y}}_{..})' \quad (6.9)$$

$$= \sum_{i=1}^k \frac{1}{n} \mathbf{y}_{i\cdot} \mathbf{y}'_{i\cdot} - \frac{1}{kn} \mathbf{y}_{..} \mathbf{y}'_{..},$$

$$\mathbf{E} = \sum_{i=1}^k \sum_{j=1}^n (\mathbf{y}_{ij} - \bar{\mathbf{y}}_{i\cdot})(\mathbf{y}_{ij} - \bar{\mathbf{y}}_{i\cdot})' \quad (6.10)$$

$$= \sum_{ij} \mathbf{y}_{ij} \mathbf{y}'_{ij} - \sum_i \frac{1}{n} \mathbf{y}_{i\cdot} \mathbf{y}'_{i\cdot}$$

The $p \times p$ “hypothesis” matrix \mathbf{H} has a “between” sum of squares on the diagonal for each of the p variables. Off-diagonal elements are analogous sums of products for each pair of variables. Assuming there are no linear dependencies in the variables, the rank of \mathbf{H} is the smaller of p and ν_H , $\min(p, \nu_H)$, where ν_H represents the degrees of freedom for hypothesis; in the one-way case $\nu_H = k - 1$. Thus \mathbf{H} can be singular. The $p \times p$ “error” matrix \mathbf{E} has a “within” sum of squares for each variable on the diagonal with analogous sums of products off-diagonal. The rank of \mathbf{E} is p , unless ν_E is less than p .

Thus \mathbf{H} has the form

$$\mathbf{H} = \begin{pmatrix} \text{SSH}_{11} & \text{SPH}_{12} & \cdots & \text{SPH}_{1p} \\ \text{SPH}_{12} & \text{SSH}_{22} & \cdots & \text{SPH}_{2p} \\ \vdots & \vdots & & \vdots \\ \text{SPH}_{1p} & \text{SPH}_{2p} & \cdots & \text{SSH}_{pp} \end{pmatrix}, \quad (6.11)$$

where, for example,

$$\begin{aligned} \text{SSH}_{22} &= n \sum_{i=1}^k (\bar{y}_{i.2} - \bar{y}_{..2})^2 = \sum_i \frac{y_{i.2}^2}{n} - \frac{\bar{y}_{..2}^2}{kn}, \\ \text{SPH}_{12} &= n \sum_{i=1}^k (\bar{y}_{i.1} - \bar{y}_{..1})(\bar{y}_{i.2} - \bar{y}_{..2}) = \sum_i \frac{y_{i.1}y_{i.2}}{n} - \frac{\bar{y}_{..1}\bar{y}_{..2}}{kn}. \end{aligned}$$

In these expressions, the subscript 1 or 2 indicates the first or second variable. Thus, for example, $\bar{y}_{i.2}$ is the second element in $\bar{\mathbf{y}}_i$:

$$\bar{\mathbf{y}}_i = \begin{pmatrix} \bar{y}_{i.1} \\ \bar{y}_{i.2} \\ \vdots \\ \bar{y}_{i.p} \end{pmatrix}.$$

The matrix \mathbf{E} can be expressed in a form similar to (6.11):

$$\mathbf{E} = \begin{pmatrix} \text{SSE}_{11} & \text{SPE}_{12} & \cdots & \text{SPE}_{1p} \\ \text{SPE}_{12} & \text{SSE}_{22} & \cdots & \text{SPE}_{2p} \\ \vdots & \vdots & & \vdots \\ \text{SPE}_{1p} & \text{SPE}_{2p} & \cdots & \text{SSE}_{pp} \end{pmatrix}, \quad (6.12)$$

where, for example,

$$\begin{aligned} \text{SSE}_{22} &= \sum_{i=1}^k \sum_{j=1}^n (y_{ij2} - \bar{y}_{i.2})^2 = \sum_{ij} y_{ij2}^2 - \sum_i \frac{y_{i.2}^2}{n}, \\ \text{SPE}_{12} &= \sum_{i=1}^k \sum_{j=1}^n (y_{ij1} - \bar{y}_{i.1})(y_{ij2} - \bar{y}_{i.2}) = \sum_{ij} y_{ij1}y_{ij2} - \sum_i \frac{y_{i.1}y_{i.2}}{n}. \end{aligned}$$

Note that the elements of \mathbf{E} are sums of squares and products, not variances and covariances. To estimate Σ , we use $\mathbf{S}_{p1} = \mathbf{E}/(nk - k)$, so that

$$E\left(\frac{\mathbf{E}}{nk - k}\right) = \Sigma.$$

6.1.3 Wilks' Test Statistic

The likelihood ratio test of $H_0: \mu_1 = \mu_2 = \cdots = \mu_k$ is given by

$$\Lambda = \frac{|\mathbf{E}|}{|\mathbf{E} + \mathbf{H}|}, \quad (6.13)$$

which is known as Wilks' Λ . (It has also been called Wilks' U .) We reject H_0 if $\Lambda \leq \Lambda_{\alpha, p, \nu_H, \nu_E}$. Note that rejection is for small values of Λ . Exact critical values $\Lambda_{\alpha, p, \nu_H, \nu_E}$ for Wilks' Λ are found in Table A.9, taken from Wall (1967). The parameters in Wilks' Λ distribution are

$$\begin{aligned} p &= \text{number of variables (dimension),} \\ \nu_H &= \text{degrees of freedom for hypothesis,} \\ \nu_E &= \text{degrees of freedom for error.} \end{aligned}$$

Wilks' Λ compares the "within" sum of squares and products matrix \mathbf{E} to the "total" sum of squares and products matrix $\mathbf{E} + \mathbf{H}$. This is similar to the univariate F -statistic in (6.6) that compares the between sum of squares to the within sum of squares. By using determinants, the test statistic Λ is reduced to a scalar. Thus the multivariate information in \mathbf{E} and \mathbf{H} about separation of mean vectors $\bar{\mathbf{y}}_1, \bar{\mathbf{y}}_2, \dots, \bar{\mathbf{y}}_k$ is channeled into a single scale, on which we can determine whether the separation of mean vectors is significant. This is typical of multivariate tests in general.

The mean vectors occupy a space of dimension $s = \min(p, \nu_H)$, and within this space various configurations of these mean vectors are possible. This suggests the possibility that another test statistic may be more powerful than Wilks' Λ . Competing test statistics are discussed in Sections 6.1.4 and 6.1.5.

Some of the properties and characteristics of Wilks' Λ are as follows:

1. In order for the determinants in (6.13) to be positive, it is necessary that $\nu_E \geq p$.
2. For any MANOVA model, the degrees of freedom ν_H and ν_E are always the same as in the analogous univariate case. In the balanced one-way model, for example, $\nu_H = k - 1$ and $\nu_E = k(n - 1)$.
3. The parameters p and ν_H can be interchanged; the distribution of $\Lambda_{p, \nu_H, \nu_E}$ is the same as that of $\Lambda_{\nu_H, p, \nu_E + \nu_H - p}$.
4. Wilks' Λ in (6.13) can be expressed in terms of the eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_s$ of $\mathbf{E}^{-1}\mathbf{H}$, as follows:

$$\Lambda = \prod_{i=1}^s \frac{1}{1 + \lambda_i}. \quad (6.14)$$

The number of nonzero eigenvalues of $\mathbf{E}^{-1}\mathbf{H}$ is $s = \min(p, \nu_H)$, which is the rank of \mathbf{H} . The matrix $\mathbf{H}\mathbf{E}^{-1}$ has the same eigenvalues as $\mathbf{E}^{-1}\mathbf{H}$ (see Section 2.11.5) and could be used in its place to obtain Λ . However, we prefer $\mathbf{E}^{-1}\mathbf{H}$ because we will use its eigenvectors later.

5. The range of Λ is $0 \leq \Lambda \leq 1$, and the test based on Wilks' Λ is an inverse test in the sense that we reject H_0 for small values of Λ . If the sample mean

Table 6.1 Transformations of Wilks' Λ to Exact Upper Tail F -Tests

Parameters p, ν_H	Statistic Having F -Distribution	Degrees of Freedom
Any $p, \nu_H = 1$	$\frac{1 - \Lambda}{\Lambda} \frac{\nu_E - p + 1}{p}$	$p, \nu_E - p + 1$
Any $p, \nu_H = 2$	$\frac{1 - \sqrt{\Lambda}}{\sqrt{\Lambda}} \frac{\nu_E - p + 1}{p}$	$2p, 2(\nu_E - p + 1)$
$p = 1, \text{any } \nu_H$	$\frac{1 - \Lambda}{\Lambda} \frac{\nu_E}{\nu_H}$	ν_H, ν_E
$p = 2, \text{any } \nu_H$	$\frac{1 - \sqrt{\Lambda}}{\sqrt{\Lambda}} \frac{\nu_E - 1}{\nu_H}$	$2\nu_H, 2(\nu_E - 1)$

vectors were equal, we would have $\mathbf{H} = \mathbf{O}$ and $\Lambda = |\mathbf{E}|/|\mathbf{E} + \mathbf{O}| = 1$. On the other hand, as the sample mean vectors become more widely spread apart compared to the within-sample variation, \mathbf{H} becomes much “larger” than \mathbf{E} , and Λ approaches zero.

6. In Table A.9, the critical values decrease for increasing p . Thus the addition of variables will reduce the power unless the variables contribute to rejection of the hypothesis by causing a significant reduction in Λ .
7. When $\nu_H = 1$ or 2 or when $p = 1$ or 2, Wilks' Λ transforms to an exact F -statistic. The transformations from Λ to F for these special cases are given in Table 6.1. The hypothesis is rejected when the transformed value of Λ exceeds the upper α -level percentage point of the F -distribution, with degrees of freedom as shown.
8. For values of p and ν_H other than those in Table 6.1, an approximate F -statistic is given by

$$F = \frac{1 - \Lambda^{1/t}}{\Lambda^{1/t}} \frac{df_2}{df_1}, \quad (6.15)$$

with df_1 and df_2 degrees of freedom, where

$$df_1 = p\nu_H, \quad df_2 = wt - \frac{1}{2}(p\nu_H - 2),$$

$$w = \nu_E + \nu_H - \frac{1}{2}(p + \nu_H + 1), \quad t = \sqrt{\frac{p^2\nu_H^2 - 4}{p^2 + \nu_H^2 - 5}}.$$

When $p\nu_H = 2$, t is set equal to 1. The approximate F in (6.15) reduces to the exact F -values given in Table 6.1, when either ν_H or p is 1 or 2.

A (less accurate) approximate test is given by

$$\chi^2 = -[\nu_E - \frac{1}{2}(p - \nu_H + 1)] \ln \Lambda, \quad (6.16)$$




Figure 6.1 Three samples with significant Wilks' Λ but nonsignificant F 's.

which has an approximate χ^2 -distribution with $p\nu_H$ degrees of freedom. We reject H_0 if $\chi^2 > \chi_{\alpha}^2$. This approximation is accurate to three decimal places when $p^2 + \nu_H^2 \leq \frac{1}{3}f$, where $f = \nu_E - \frac{1}{2}(p - \nu_H + 1)$.

Although it is more expensive computationally, there is also a method for obtaining exact p -values for Λ . This exact method is due to Lee (1972) and Davis (1979) and is available in modern statistical computing packages. For example, in SAS Version 9 (SAS Institute Inc., 2009) exact tests can be requested using the "MSTAT = EXACT" option of the MANOVA statement in PROC GLM.

9. If the multivariate test based on Λ rejects H_0 , it could be followed by an F -test as in (6.6) on each of the p individual y 's. We can formulate a hypothesis comparing the means across the k groups for each variable, namely, $H_{0r}: \mu_{1r} = \mu_{2r} = \dots = \mu_{kr}, r = 1, 2, \dots, p$. It does not necessarily follow that any of the F -tests on the p individual variables will reject the corresponding H_{0r} . Conversely, it is possible that one or more of the F 's will reject H_{0r} when the Λ -test accepts H_0 . In either case, where the multivariate test and the univariate tests disagree, we use the multivariate test result rather than the univariate results. This is similar to the relationship between Z^2 -tests and z -tests shown in Figure 5.2.

In the three bivariate samples plotted in Figure 6.1, we illustrate the case where Λ rejects $H_0: \mu_1 = \mu_2 = \mu_3$ but the F 's accept both of $H_{0r}: \mu_{1r} = \mu_{2r} = \mu_{3r}, r = 1, 2$, that is, for y_1 and y_2 . There is no significant separation of the

three samples in either the y_1 or y_2 direction alone. Other follow-up procedures are given in Sections 6.1.4 and 6.4.

- 10.** The Wilks' Λ -test is the likelihood ratio test. Other approaches to test construction lead to different tests. Three such tests are given in Sections 6.1.4 and 6.1.5.

6.1.4 Roy's Test

In the *union-intersection* approach, we seek the linear combination $z_{ij} = \mathbf{a}'\mathbf{y}_{ij}$ that maximizes the spread of the transformed means $\bar{z}_{i\cdot} = \mathbf{a}'\bar{\mathbf{y}}_{i\cdot}$ relative to the within-sample spread of points. Thus we seek the vector \mathbf{a} that maximizes

$$F = \frac{n \sum_{i=1}^k (\bar{z}_{i\cdot} - \bar{z}_{..})^2 / (k-1)}{\sum_{i=1}^k \sum_{j=1}^n (z_{ij} - \bar{z}_{i\cdot})^2 / (kn-k)}, \quad (6.17)$$

which, by analogy to $s_z^2 = \mathbf{a}'\mathbf{S}\mathbf{a}$ in (3.55), can be written as

$$F = \frac{\mathbf{a}'\mathbf{H}\mathbf{a}/(k-1)}{\mathbf{a}'\mathbf{E}\mathbf{a}/(kn-k)}. \quad (6.18)$$

This is maximized by \mathbf{a}_1 , the eigenvector corresponding to λ_1 , the largest eigenvalue of $\mathbf{E}^{-1}\mathbf{H}$ (see Section 8.4.1), and we have

$$\max_{\mathbf{a}} F = \frac{\mathbf{a}_1'\mathbf{H}\mathbf{a}_1/(k-1)}{\mathbf{a}_1'\mathbf{E}\mathbf{a}_1/(kn-k)} = \frac{k(n-1)}{k-1} \lambda_1. \quad (6.19)$$

Since $\max_{\mathbf{a}} F$ in (6.19) is maximized over all possible linear functions, it no longer has an F -distribution. To test $H_0: \mu_1 = \mu_2 = \dots = \mu_k$ based on λ_1 , we use Roy's *union-intersection test*, also called *Roy's largest root test*. The test statistic is given by

$$\theta = \frac{\lambda_1}{1 + \lambda_1}. \quad (6.20)$$

Critical values for θ are given in Table A.10 (Pearson and Hartley 1972, Pillai 1964, 1965). We reject $H_0: \mu_1 = \mu_2 = \dots = \mu_k$ if $\theta \geq \theta_{\alpha,s,m,N}$. The parameters s , m , and N are defined as

$$s = \min(\nu_H, p), \quad m = \frac{1}{2}(|\nu_H - p| - 1), \quad N = \frac{1}{2}(\nu_E - p - 1).$$

For $s = 1$, use (6.33) and (6.36) in Section 6.1.7 to obtain an F -test.

The eigenvector \mathbf{a}_1 corresponding to λ_1 is used in the *discriminant function*, $z = \mathbf{a}_1'\mathbf{y}$. Since this is the function that best separates the transformed means $\bar{z}_{i\cdot} = \mathbf{a}'\bar{\mathbf{y}}_{i\cdot}$, $i = 1, 2, \dots, k$ [relative to the within-sample spread, see (6.17)], the coefficients $a_{11}, a_{12}, \dots, a_{1p}$ in the linear combination $z = \mathbf{a}_1'\mathbf{y}$ can be examined for an indication of which variables contribute most to separating the means. The discriminant function is discussed further in Sections 6.1.8 and 6.4 and in Chapter 8.

We do not have a satisfactory F -approximation for θ or λ_1 , but an “upper bound” on F that is provided in some software programs is given by

$$F = \frac{(\nu_E - d - 1)\lambda_1}{d}, \quad (6.21)$$

with degrees of freedom d and $\nu_E - d - 1$, where $d = \max(p, \nu_H)$. The term “upper bound” indicates that the F in (6.21) is greater than the “true F "; that is, $F > F_{d, \nu_E - d - 1}$. Therefore, we feel safe if H_0 is accepted by (6.21), but if rejection of H_0 is indicated, the information is virtually worthless.

Fortunately, there exists a method due to Davis (1972) and Pillai and Flury (1984) that allows us to calculate an exact p -value associated with Roy's largest root. This exact method has been implemented in SAS Version 9 (SAS Institute Inc., 2009) by using the “MSTAT = EXACT” option of the MANOVA statement in PROC GLM.

Some computer programs do not provide eigenvalues of nonsymmetric matrices, such as $\mathbf{E}^{-1}\mathbf{H}$. However, the eigenvalues of $\mathbf{E}^{-1}\mathbf{H}$ are the same as the eigenvalues of the symmetric matrices $(\mathbf{E}^{1/2})^{-1}\mathbf{H}(\mathbf{E}^{1/2})^{-1}$ and $(\mathbf{U}')^{-1}\mathbf{H}\mathbf{U}'^{-1}$, where $\mathbf{E}^{1/2}$ is the square root matrix of \mathbf{E} given in (2.112) and $\mathbf{U}'\mathbf{U} = \mathbf{E}$ is the Cholesky factorization of \mathbf{E} (Section 2.7). We demonstrate this for the Cholesky approach. We first multiply the defining relationship $(\mathbf{E}^{-1}\mathbf{H} - \lambda\mathbf{I})\mathbf{a} = \mathbf{0}$ by \mathbf{E} to obtain

$$(\mathbf{H} - \lambda\mathbf{E})\mathbf{a} = \mathbf{0}. \quad (6.22)$$

Then substituting $\mathbf{E} = \mathbf{U}'\mathbf{U}$ into (6.22), multiplying by $(\mathbf{U}')^{-1}$, and inserting $\mathbf{U}'^{-1}\mathbf{U} = \mathbf{I}$, we have

$$\begin{aligned} & (\mathbf{H} - \lambda\mathbf{U}'\mathbf{U})\mathbf{a} = \mathbf{0}, \\ & (\mathbf{U}')^{-1}(\mathbf{H} - \lambda\mathbf{U}'\mathbf{U})\mathbf{a} = (\mathbf{U}')^{-1}\mathbf{0} = \mathbf{0}, \\ & [(\mathbf{U}')^{-1}\mathbf{H} - \lambda\mathbf{U}]\mathbf{U}'^{-1}\mathbf{U}\mathbf{a} = \mathbf{0}, \\ & [(\mathbf{U}')^{-1}\mathbf{H}\mathbf{U}'^{-1} - \lambda\mathbf{I}]\mathbf{U}\mathbf{a} = \mathbf{0}. \end{aligned} \quad (6.23)$$

Thus $(\mathbf{U}')^{-1}\mathbf{H}\mathbf{U}'^{-1}$ has the same eigenvalues as $\mathbf{E}^{-1}\mathbf{H}$ and has eigenvectors of the form $\mathbf{U}\mathbf{a}$, where \mathbf{a} is an eigenvector of $\mathbf{E}^{-1}\mathbf{H}$. Note that $(\mathbf{U}')^{-1}\mathbf{H}\mathbf{U}'^{-1}$ is positive semidefinite, and thus $\lambda_i \geq 0$ for all eigenvalues of $\mathbf{E}^{-1}\mathbf{H}$.

6.1.5 Pillai and Lawley–Hotelling Tests

There are two additional test statistics for $H_0: \mu_1 = \mu_2 = \dots = \mu_k$ based on the eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_s$ of $\mathbf{E}^{-1}\mathbf{H}$. The *Pillai statistic* is given by

$$V^{(s)} = \text{tr}[(\mathbf{E} + \mathbf{H})^{-1}\mathbf{H}] = \sum_{i=1}^s \frac{\lambda_i}{1 + \lambda_i}. \quad (6.24)$$

We reject H_0 for $V^{(s)} \geq V_\alpha^{(s)}$. The upper percentage points, $V_\alpha^{(s)}$, are given in Table A.11 (Schuurmann et al. 1975), indexed by s , m , and N , which are defined as in

Section 6.1.4 for Roy's test. For $s = 1$, use (6.33) and (6.36) in Section 6.1.7 to obtain an F -test.

Pillai's test statistic in (6.24) is an extension of Roy's statistic $\theta = \lambda_1/(1 + \lambda_1)$. If the mean vectors do not lie in one dimension, the information in the additional terms $\lambda_i/(1 + \lambda_i)$, $i = 2, 3, \dots, s$, may be helpful in rejecting H_0 .

For parameter values not included in Table A.11, we can use an approximate F -statistic:

$$F_1 = \frac{(2N + s + 1)V^{(s)}}{(2m + s + 1)(s - V^{(s)})}, \quad (6.25)$$

which is approximately distributed as $F_{s(2m+s+1), s(2N+s+1)}$. The approximation in (6.25) can also be given as

$$F_2 = \frac{(\nu_E - p + s)V^{(s)}}{d(s - V^{(s)})}, \quad (6.26)$$

with sd and $s(\nu_E - p + s)$ degrees of freedom, where $d = \max(p, \nu_H)$. A more recent F -approximation due to Muller (1998) has been shown to yield better properties than F_1 and F_2 , but is more computationally expensive. Although it is too complicated to present in detail here, it has been implemented in software packages such as SAS Version 9 (SAS Institute Inc., 2009).

The *Lawley–Hotelling statistic* (Lawley 1938, Hotelling 1951) is defined as

$$U^{(s)} = \text{tr}(\mathbf{E}^{-1}\mathbf{H}) = \sum_{i=1}^s \lambda_i \quad (6.27)$$

and is also known as *Hotelling's generalized T^2 -statistic* (see a comment at the end of Section 6.1.7). Table A.12 (Davis 1970a,b, 1980) gives upper percentage points of the test statistic

$$\frac{\nu_E}{\nu_H} U^{(s)}.$$

We reject H_0 for large values of the test statistic. Note that in Table A.12, $p \leq \nu_H$ and $p \leq \nu_E$. If $p > \nu_H$, use $(\nu_H, p, \nu_E + \nu_H - p)$ in place of (p, ν_H, ν_E) . (This same pattern in the parameters is found in Wilks' Λ ; see property 3 in Section 6.1.3.) If $\nu_H = 1$ and $p > 1$, use the relationship $U^{(1)} = T^2/\nu_E$ [see (6.38) in Section 6.1.7]. For other values of the parameters not included in Table A.12, we can use an approximate F -statistic:

$$F_1 = \frac{U^{(s)}}{c}, \quad (6.28)$$

which is approximately distributed as $F_{a,b}$, where

$$a = p\nu_H, \quad b = 4 + \frac{a + 2}{B - 1}, \quad c = \frac{a(b - 2)}{b(\nu_E - p - 1)},$$

$$B = \frac{(\nu_E + \nu_H - p - 1)(\nu_E - 1)}{(\nu_E - p - 3)(\nu_E - p)}.$$

An alternative F -approximation is given by

$$F_2 = \frac{2(sN + 1)U^{(s)}}{s^2(2m + s + 1)}, \quad (6.29)$$

with $s(2m + s + 1)$ and $2(sN + 1)$ degrees of freedom. The approximation in (6.29) can also be given by

$$F_3 = \frac{[s(\nu_E - p - 1) + 2]U^{(s)}}{sp\nu_H}, \quad (6.30)$$

with $p\nu_H$ and $s(\nu_E - p - 1) + 2$ degrees of freedom.

An exact p -value based on $U^{(s)}$ has been given by Davis (1970c, 1980) and has been implemented with the other multivariate exact tests in SAS Version 9 (SAS Institute Inc., 2009).

6.1.6 Unbalanced One-Way MANOVA

The balanced one-way model can be easily extended to the unbalanced case, in which there are n_i observation vectors in the i th group. The model in (6.8) becomes

$$\mathbf{y}_{ij} = \boldsymbol{\mu} + \boldsymbol{\alpha}_i + \boldsymbol{\varepsilon}_{ij} = \boldsymbol{\mu}_i + \boldsymbol{\varepsilon}_{ij}, \quad i = 1, 2, \dots, k; j = 1, 2, \dots, n_i.$$

The mean vectors become $\bar{\mathbf{y}}_{i\cdot} = \sum_{j=1}^{n_i} \mathbf{y}_{ij}/n_i$ and $\bar{\mathbf{y}}_{\cdot\cdot} = \sum_{i=1}^k \sum_{j=1}^{n_i} \mathbf{y}_{ij}/N$, where $N = \sum_{i=1}^k n_i$. Similarly, the total vectors are defined as $\mathbf{y}_{i\cdot} = \sum_{j=1}^{n_i} \mathbf{y}_{ij}$ and $\mathbf{y}_{\cdot\cdot} = \sum_{ij} \mathbf{y}_{ij}$. The \mathbf{H} and \mathbf{E} matrices are calculated as

$$\mathbf{H} = \sum_{i=1}^k n_i (\bar{\mathbf{y}}_{i\cdot} - \bar{\mathbf{y}}_{\cdot\cdot})(\bar{\mathbf{y}}_{i\cdot} - \bar{\mathbf{y}}_{\cdot\cdot})' = \sum_{i=1}^k \frac{1}{n_i} \mathbf{y}_{i\cdot} \mathbf{y}_{i\cdot}' - \frac{1}{N} \mathbf{y}_{\cdot\cdot} \mathbf{y}_{\cdot\cdot}', \quad (6.31)$$

$$\mathbf{E} = \sum_{i=1}^k \sum_{j=1}^{n_i} (\mathbf{y}_{ij} - \bar{\mathbf{y}}_{i\cdot})(\mathbf{y}_{ij} - \bar{\mathbf{y}}_{i\cdot})' = \sum_{i=1}^k \sum_{j=1}^{n_i} \mathbf{y}_{ij} \mathbf{y}_{ij}' - \sum_{i=1}^k \frac{1}{n_i} \mathbf{y}_{i\cdot} \mathbf{y}_{i\cdot}'. \quad (6.32)$$

Wilks' Λ and the other tests have the same form as in Sections 6.1.3–6.1.5 using \mathbf{H} and \mathbf{E} from (6.31) and (6.32). In each test we have

$$\nu_H = k - 1, \quad \nu_E = N - k = \sum_{i=1}^k n_i - k.$$

Note that $N = \sum_i n_i$ differs from N used as a parameter in Roy's and Pillai's tests in Sections 6.1.4 and 6.1.5.

6.1.7 Summary of the Four Tests and Relationship to T^2

We compare the four test statistics in terms of the eigenvalues $\lambda_1 > \lambda_2 > \dots > \lambda_s$ of $\mathbf{E}^{-1}\mathbf{H}$, where $s = \min(\nu_H, p)$:

$$\begin{aligned}\text{Pillai } V^{(s)} &= \sum_{i=1}^s \frac{\lambda_i}{1 + \lambda_i} \\ \text{Lawley-Hotelling } U^{(s)} &= \sum_{i=1}^s \lambda_i \\ \text{Wilks' lambda } \Lambda &= \prod_{i=1}^s \frac{1}{1 + \lambda_i} \\ \text{Roy's largest root } \theta &= \frac{\lambda_1}{1 + \lambda_1}\end{aligned}$$

Note that for all four tests we must have $\nu_E \geq p$. As noted in Section 6.1.3 and elsewhere, p is the number of variables, ν_H is the degrees of freedom for the hypothesis, and ν_E is the degrees of freedom for error.

Why do we use four different tests? All four are exact tests; that is, when H_0 is true, each test has probability α of rejecting H_0 . However, the tests are not equivalent, and in a given sample they may lead to different conclusions even when H_0 is true; some may reject H_0 while others accept H_0 . This is due to the multidimensional nature of the space in which the mean vectors $\mu_1, \mu_2, \dots, \mu_k$ lie. A comparison of power and other properties of the tests is given in Section 6.2.

When $\nu_H = 1$, then s is also equal to 1, and there is only one nonzero eigenvalue λ_1 . In this case, all four test statistics are functions of each other and give equivalent results. In terms of θ , for example, the other three become

$$U^{(1)} = \lambda_1 = \frac{\theta}{1 - \theta}, \quad (6.33)$$

$$V^{(1)} = \theta, \quad (6.34)$$

$$\Lambda = 1 - \theta. \quad (6.35)$$

In the case of $\nu_H = 1$, all four statistics can be transformed to an exact F using

$$F = \frac{\nu_E - p + 1}{p} U^{(1)}, \quad (6.36)$$

which is distributed as $F_{p, \nu_E - p + 1}$.

The equivalence of all four test statistics to Hotelling's T^2 when $\nu_H = 1$ was noted in Section 5.6.1. We now demonstrate the relationship $T^2 = (n_1 + n_2 - 2)U^{(1)}$ in (5.17). For \mathbf{H} and \mathbf{E} , we use (6.31) and (6.32), which allow unequal n_i , since we do not require $n_1 = n_2$ in T^2 . In this case, with only two groups, $\mathbf{H} = \sum_{i=1}^2 n_i (\bar{\mathbf{y}}_{i\cdot} - \bar{\mathbf{y}}_{..})(\bar{\mathbf{y}}_{i\cdot} - \bar{\mathbf{y}}_{..})'$ can be expressed as

$$\mathbf{H} = \frac{n_1 n_2}{n_1 + n_2} (\bar{\mathbf{y}}_{1\cdot} - \bar{\mathbf{y}}_{2\cdot})(\bar{\mathbf{y}}_{1\cdot} - \bar{\mathbf{y}}_{2\cdot})' = c(\bar{\mathbf{y}}_{1\cdot} - \bar{\mathbf{y}}_{2\cdot})(\bar{\mathbf{y}}_{1\cdot} - \bar{\mathbf{y}}_{2\cdot})', \quad (6.37)$$

where $c = n_1 n_2 / (n_1 + n_2)$. Then by (6.33) and (6.27), $U^{(1)}$ becomes

$$\begin{aligned} U^{(1)} &= \lambda_1 = \text{tr}(\mathbf{E}^{-1} \mathbf{H}) \\ &= \text{tr}[c \mathbf{E}^{-1} (\bar{\mathbf{y}}_{1\cdot} - \bar{\mathbf{y}}_{2\cdot})(\bar{\mathbf{y}}_{1\cdot} - \bar{\mathbf{y}}_{2\cdot})'] \quad [\text{by (6.37)}] \\ &= c \text{tr}[(\bar{\mathbf{y}}_{1\cdot} - \bar{\mathbf{y}}_{2\cdot})' \mathbf{E}^{-1} (\bar{\mathbf{y}}_{1\cdot} - \bar{\mathbf{y}}_{2\cdot})] \quad [\text{by (2.97)}] \\ &= \frac{c}{n_1 + n_2 - 2} (\bar{\mathbf{y}}_{1\cdot} - \bar{\mathbf{y}}_{2\cdot})' \left(\frac{\mathbf{E}}{n_1 + n_2 - 2} \right)^{-1} (\bar{\mathbf{y}}_{1\cdot} - \bar{\mathbf{y}}_{2\cdot}) \\ &= \frac{T^2}{n_1 + n_2 - 2}, \end{aligned} \quad (6.38)$$

since $\mathbf{E}/(n_1 + n_2 - 2) = \mathbf{S}_{\text{pl}}$ (see Section 5.4.2). From this result, (5.16), (5.18), and (5.19) follow immediately using (6.33)–(6.35).

Because of the direct relationship in (6.38) between $U^{(1)}$ and T^2 for the case of two groups, the Lawley–Hotelling statistic $U^{(s)}$ is often called the *generalized T^2 -statistic*.

■ EXAMPLE 6.1.7

In a classical experiment carried out from 1918 to 1934, apple trees of different rootstocks were compared (Andrews and Herzberg 1985, pp. 357–360). The data for eight trees from each of six rootstocks are given in Table 6.2. The variables are

- y_1 = trunk girth at 4 years ($\text{mm} \times 100$)
- y_2 = extension growth at 4 years (m)
- y_3 = trunk girth at 15 years ($\text{mm} \times 100$)
- y_4 = weight of tree above ground at 15 years ($\text{lb} \times 1000$)

The matrices \mathbf{H} , \mathbf{E} , and $\mathbf{E} + \mathbf{H}$ are given by

$$\begin{aligned} \mathbf{H} &= \begin{pmatrix} .074 & .537 & .332 & .208 \\ .537 & 4.200 & 2.355 & 1.637 \\ .332 & 2.355 & 6.114 & 3.781 \\ .208 & 1.637 & 3.781 & 2.493 \end{pmatrix}, \\ \mathbf{E} &= \begin{pmatrix} .320 & 1.697 & .554 & .217 \\ 1.697 & 12.143 & 4.364 & 2.110 \\ .554 & 4.364 & 4.291 & 2.482 \\ .217 & 2.110 & 2.482 & 1.723 \end{pmatrix}, \\ \mathbf{E} + \mathbf{H} &= \begin{pmatrix} .394 & 2.234 & .886 & .426 \\ 2.234 & 16.342 & 6.719 & 3.747 \\ .886 & 6.719 & 10.405 & 6.263 \\ .426 & 3.747 & 6.263 & 4.216 \end{pmatrix}. \end{aligned}$$

In this case, the mean vectors represent six points in four-dimensional space. We can compare the mean vectors for significant differences using Wilks' Λ as

Table 6.2 Rootstock Data

Rootstock	y_1	y_2	y_3	y_4
1	1.11	2.569	3.58	.760
1	1.19	2.928	3.75	.821
1	1.09	2.865	3.93	.928
1	1.25	3.844	3.94	1.009
1	1.11	3.027	3.60	.766
1	1.08	2.336	3.51	.726
1	1.11	3.211	3.98	1.209
1	1.16	3.037	3.62	.750
2	1.05	2.074	4.09	1.036
2	1.17	2.885	4.06	1.094
2	1.11	3.378	4.87	1.635
2	1.25	3.906	4.98	1.517
2	1.17	2.782	4.38	1.197
2	1.15	3.018	4.65	1.244
2	1.17	3.383	4.69	1.495
2	1.19	3.447	4.40	1.026
3	1.07	2.505	3.76	.912
3	.99	2.315	4.44	1.398
3	1.06	2.667	4.38	1.197
3	1.02	2.390	4.67	1.613
3	1.15	3.021	4.48	1.476
3	1.20	3.085	4.78	1.571
3	1.20	3.308	4.57	1.506
3	1.17	3.231	4.56	1.458
4	1.22	2.838	3.89	.944
4	1.03	2.351	4.05	1.241
4	1.14	3.001	4.05	1.023
4	1.01	2.439	3.92	1.067
4	.99	2.199	3.27	.693
4	1.11	3.318	3.95	1.085
4	1.20	3.601	4.27	1.242
4	1.08	3.291	3.85	1.017
5	.91	1.532	4.04	1.084
5	1.15	2.552	4.16	1.151
5	1.14	3.083	4.79	1.381
5	1.05	2.330	4.42	1.242
5	.99	2.079	3.47	.673
5	1.22	3.366	4.41	1.137
5	1.05	2.416	4.64	1.455
5	1.13	3.100	4.57	1.325
6	1.11	2.813	3.76	.800
6	.75	.840	3.14	.606
6	1.05	2.199	3.75	.790
6	1.02	2.132	3.99	.853
6	1.05	1.949	3.34	.610
6	1.07	2.251	3.21	.562
6	1.13	3.064	3.63	.707
6	1.11	2.469	3.95	.952

given by (6.13):

$$\Lambda = \frac{|\mathbf{E}|}{|\mathbf{E} + \mathbf{H}|} = \frac{.6571}{4.2667} = .154.$$

In this case, the parameters of the Wilks' Λ distribution are $p = 4$, $\nu_H = 6 - 1 = 5$, and $\nu_E = 6(8 - 1) = 42$. We reject $H_0: \boldsymbol{\mu}_1 = \boldsymbol{\mu}_2 = \dots = \boldsymbol{\mu}_6$ because

$$\Lambda = .154 < \Lambda_{.05,4,5,40} = .455.$$

Note the use of $\nu_E = 40$ in place of $\nu_E = 42$. This is a conservative approach that allows a table value to be used without interpolation.

To obtain an approximate F , we first calculate

$$t = \sqrt{\frac{p^2\nu_H^2 - 4}{p^2 + \nu_H^2 - 5}} = \sqrt{\frac{4^25^2 - 4}{4^2 + 5^2 - 5}} = 3.3166,$$

$$w = \nu_E + \nu_H - \frac{1}{2}(p + \nu_H + 1) = 42 + 5 - \frac{1}{2}(4 + 5 + 1) = 42,$$

$$df_1 = p\nu_H = 4(5) = 20, df_2 = wt - \frac{1}{2}(p\nu_H - 2) = 130.3.$$

Then the approximate F is given by (6.15),

$$F = \frac{1 - \Lambda^{1/t}}{\Lambda^{1/t}} \frac{df_2}{df_1} = \frac{1 - (.154)^{1/3.3166}}{(.154)^{1/3.3166}} \frac{130.3}{20} = 4.937,$$

which exceeds $F_{.001,20,120} = 2.53$, and we reject H_0 .

The four eigenvalues of $\mathbf{E}^{-1}\mathbf{H}$ are 1.876, .791, .229, and .026. With these we can calculate the other three test statistics. For Pillai's statistic we have, by (6.24),

$$V^{(s)} = \sum_{i=1}^4 \frac{\lambda_i}{1 + \lambda_i} = 1.305.$$

To find a critical value for $V^{(s)}$ in Table A.11, we need

$$s = \min(\nu_H, p) = 4, \quad m = \frac{1}{2}(|\nu_H - p| - 1) = 0,$$

$$N = \frac{1}{2}(\nu_E - p - 1) = 18.5.$$

Then $V_{.05}^{(s)} = .645$ (by interpolation). Since $1.305 > .645$, we reject H_0 .

For the Lawley–Hotelling statistic we obtain, by (6.27),

$$U^{(s)} = \sum_{i=1}^s \lambda_i = 2.921.$$

To make the test, we calculate the test statistic

$$\frac{\nu_E}{\nu_H} U^{(s)} = \frac{42}{5} (2.921) = 24.539.$$

The .05 critical value for $\nu_E U^{(s)} / \nu_H$ is given in Table A.12 as 7.6188 (using $\nu_E = 40$), and we therefore reject H_0 .

Roy's test statistic is given by (6.20) as

$$\theta = \frac{\lambda_1}{1 + \lambda_1} = \frac{1.876}{1 + 1.876} = .652,$$

which exceeds the .05 critical value .377 obtained (by interpolation) from Table A.10, and we reject H_0 . \square

6.1.8 Measures of Multivariate Association

In multiple regression, a measure of association between the dependent variable y and the independent variables x_1, x_2, \dots, x_q is given by the *squared multiple correlation*

$$R^2 = \frac{\text{regression sum of squares}}{\text{total sum of squares}}. \quad (6.39)$$

Similarly, in one-way univariate ANOVA, Fisher's *correlation ratio* η^2 is defined as

$$\eta^2 = \frac{\text{between sum of squares}}{\text{total sum of squares}}.$$

This is a measure of model fit similar to R^2 and gives the proportion of variation in the dependent variable y attributable to differences among the means of the groups. It answers the question "How well can we predict y by knowing what group it is from?" Thus η^2 can be considered to be a measure of association between the dependent variable y and the grouping variable i associated with μ_i or α_i in the model (6.2). In fact, if the grouping variable is represented by $k - 1$ *dummy* variables (also called *indicator* or *categorical* variables), then we have a dependent variable related to several independent variables as in multiple regression.

A dummy variable takes on the value 1 for sampling units in a group (sample) and 0 for all other sampling units. (Values other than 0 and 1 could be used.) Thus for k samples (groups), the $k - 1$ dummy variables are

$$x_i = \begin{cases} 1 & \text{if sampling unit is in } i\text{th group} \\ 0 & \text{otherwise} \end{cases} \quad i = 1, 2, \dots, k - 1.$$

Only $k - 1$ dummy variables are needed because if $x_1 = x_2 = \dots = x_{k-1} = 0$, the sampling unit must be from the k th group (see Section 11.6.2 for an illustration). The dependent variable y can be regressed on the $k - 1$ dummy variables x_1, x_2, \dots, x_{k-1} to produce results equivalent to the usual ANOVA calculations.

In (one-way) MANOVA, we need to measure the strength of the association between several dependent variables and several independent (grouping) variables. Various measurements of multivariate association have been proposed. Wilks (1932) suggested a "generalized η^2 :

$$\text{MANOVA } \eta^2 = \eta_\Lambda^2 = 1 - \Lambda, \quad (6.40)$$

based on the use of $|\mathbf{E}|$ and $|\mathbf{E} + \mathbf{H}|$ as generalizations of sums of squares. We use $1 - \Lambda$ because Λ is small if the spread in the means is large.

We now consider an η^2 based on Roy's statistic, θ . We noted in Section 6.1.4 that the discriminant function is the linear function $z = \mathbf{a}_1' \mathbf{y}$ that maximizes the spread of the means $\bar{z}_i = \mathbf{a}_1' \bar{\mathbf{y}}_i, i = 1, 2, \dots, k$, where \mathbf{a}_1 is the eigenvector of $\mathbf{E}^{-1} \mathbf{H}$ corresponding to the largest eigenvalue λ_1 . We measure the spread among the means by $\text{SSH} = n \sum_{i=1}^k (\bar{z}_i - \bar{z}_{..})^2$, divided by the within-sample spread $\text{SSE} = \sum_{ij} (z_{ij} - \bar{z}_i)^2$. The maximum value of this ratio is given by λ_1 [see (6.19)]. Thus

$$\lambda_1 = \frac{\text{SSH}(z)}{\text{SSE}(z)},$$

and by (6.20),

$$\theta = \frac{\lambda_1}{1 + \lambda_1} = \frac{\text{SSH}(z)}{\text{SSE}(z) + \text{SSH}(z)}. \quad (6.41)$$

Hence θ serves directly as a measure of multivariate association:

$$\eta_\theta^2 = \theta = \frac{\lambda_1}{1 + \lambda_1}. \quad (6.42)$$

It can be shown that the square root of this quantity,

$$\eta_\theta = \sqrt{\frac{\lambda_1}{1 + \lambda_1}}, \quad (6.43)$$

is the maximum correlation between a linear combination of the p dependent variables and a linear combination of the $k - 1$ dummy group variables (see Section 11.6.2). This type of correlation is often called a *canonical correlation* (see Chapter 11) and is defined for each eigenvalue $\lambda_1, \lambda_2, \dots, \lambda_s$ as $r_i = \sqrt{\lambda_i / (1 + \lambda_i)}$.

We now consider some measures of multivariate association suggested by Cramer and Nicewander (1979) and Muller and Peterson (1984). It is easily shown (Section 11.6.2) that Λ can be expressed as

$$\Lambda = \prod_{i=1}^s \frac{1}{1 + \lambda_i} = \prod_{i=1}^s (1 - r_i^2), \quad (6.44)$$

where $r_i^2 = \lambda_i / (1 + \lambda_i)$ is the i th squared canonical correlation described above. The *geometric mean* of a set of positive numbers a_1, a_2, \dots, a_n is defined as $(a_1 a_2 \dots a_n)^{1/n}$. Thus $\Lambda^{1/s}$ is the geometric mean of the $(1 - r_i^2)$'s, and another measure of multivariate association based on Λ , in addition to that in (6.40), is

$$A_\Lambda = 1 - \Lambda^{1/s}. \quad (6.45)$$

In fact, as noted by Muller and Peterson, the F -approximation given in (6.15),

$$F = \frac{1 - \Lambda^{1/t}}{\Lambda^{1/t}} \frac{df_2}{df_1},$$

is very similar to the univariate F -statistic (10.31) for testing significance in multiple regression,

$$F = \frac{R^2/(\text{df model})}{(1 - R^2)/(\text{df error})}, \quad (6.46)$$

based on R^2 in (6.39).

Pillai's statistic is easily expressible as the sum of the squared canonical correlations:

$$V^{(s)} = \sum_{i=1}^s \frac{\lambda_i}{1 + \lambda_i} = \sum_{i=1}^s r_i^2, \quad (6.47)$$

and the average of the r_i^2 can be used as a measure of multivariate association:

$$A_P = \frac{\sum_{i=1}^s r_i^2}{s} = \frac{V^{(s)}}{s}. \quad (6.48)$$

In terms of A_P the F -approximation given in (6.26) becomes

$$F_2 = \frac{A_p/p\nu_H}{(1 - A_p)/s(\nu_E - \nu_H + s)}, \quad (6.49)$$

which has an obvious parallel to (6.46).

For the Lawley–Hotelling statistic $U^{(s)}$, a multivariate measure of association can be defined as

$$A_{\text{LH}} = \frac{U^{(s)}/s}{1 + U^{(s)}/s}. \quad (6.50)$$

If $s = 1$, (6.50) reduces to (6.42). In fact, (6.42) is a special case of (6.50) because $U^{(s)}/s = \sum_{i=1}^s \lambda_i/s$ is the arithmetic average of the λ_i 's. It is easily seen that the F -approximation F_3 for the Lawley–Hotelling statistic given in (6.30) can be expressed in terms of A_{LH} as

$$F_3 = \frac{A_{\text{LH}}/p\nu_H}{(1 - A_{\text{LH}})/[s(\nu_E - p - 1) + 2]}, \quad (6.51)$$

which resembles (6.46).

■ EXAMPLE 6.1.8

We illustrate some measures of association for the rootstock data in Table 6.2:

$$\eta_\Lambda^2 = 1 - \Lambda = .846,$$

$$\eta_\theta^2 = \theta = .652,$$

$$A_\Lambda = 1 - \Lambda^{1/4} = 1 - (.154)^{1/4} = .374,$$

$$A_P = V^{(s)}/s = 1.305/4 = .326,$$

$$A_{\text{LH}} = \frac{U^{(s)}/s}{1 + U^{(s)}/s} = \frac{2.921/4}{1 + 2.921/4} = .422.$$

There is a wide range of values among these measures of association. □




Figure 6.2 Two possible configurations for three mean vectors in 3-space.

6.2 COMPARISON OF THE FOUR MANOVA TEST STATISTICS

When $H_0: \mu_1 = \mu_2 = \cdots = \mu_k$ is true, all the mean vectors are at the same point. Therefore, all four MANOVA test statistics have the same Type I error rate, α , as noted in Section 6.1.7; that is, all have the same probability of rejection when H_0 is true. However, when H_0 is false, the four tests have different probabilities of rejection. We noted in Section 6.1.7 that in a given sample the four tests need not agree, even if H_0 is true. One test could reject H_0 and the others accept H_0 , for example.

Historically, Wilks' Λ has played the dominant role in MANOVA significance tests because it was the first to be derived and has well-known χ^2 - and F -approximations. It can also be partitioned in certain ways we will find useful later. However, it is not always the most powerful among the four tests. The probability of rejecting H_0 when it is false is known as the *power* of the test.

In univariate ANOVA with $p = 1$, the means $\mu_1, \mu_2, \dots, \mu_k$ can be uniquely ordered along a line in one dimension, and the usual F -test is uniformly most powerful. In the multivariate case, on the other hand, with $p > 1$, the mean vectors are points in $s = \min(p, \nu_H)$ dimensions. We have four tests, not one of which is uniformly most powerful. The relative powers of the four test statistics depend on the configuration of the mean vectors $\mu_1, \mu_2, \dots, \mu_k$ in the s -dimensional space. A given test will be more powerful for one configuration of mean vectors than another.

If $\nu_H < p$, then $s = \nu_H$ and the mean vectors lie in an s -dimensional subspace of the p -dimensional space of the observations. The points may, in fact, occupy a subspace of the s dimensions. For example, they may be confined to a line (one dimension) or a plane (two dimensions). This is illustrated in Figure 6.2.

An indication of the pattern of the mean vectors is given by the eigenvalues of $\mathbf{E}^{-1}\mathbf{H}$. If there is one large eigenvalue and the others are small, the mean vectors lie close to a line. If there are two large eigenvalues, the mean vectors lie mostly in two dimensions, and so on.

Because Roy's test uses only the largest eigenvalue of $\mathbf{E}^{-1}\mathbf{H}$, it is more powerful than the others if the mean vectors are collinear. The other three tests have greater power than Roy's when the mean vectors are diffuse (spread out in several dimensions).

In terms of power, the tests are ordered $\theta \geq U^{(s)} \geq \Lambda \geq V^{(s)}$ for the collinear case. In the diffuse case and for intermediate structure between collinear and diffuse, the ordering of power is reversed, $V^{(s)} \geq \Lambda \geq U^{(s)} \geq \theta$. The latter ordering also holds for accuracy of the Type I error rate when the population covariance matrices $\Sigma_1, \Sigma_2, \dots, \Sigma_k$ are not equal. These orderings are comparisons of power. For actual computation of power in a given experimental setting or to find the sample size needed to yield a desired level of power, see Rencher (1998, Section 4.4).

Generally, if group sizes are equal, the tests are sufficiently robust with respect to heterogeneity of covariance matrices so that we need not worry. If the n_i 's are unequal and we have heterogeneity, then the α -level of the MANOVA test may be affected as follows. If the larger variances and covariances are associated with the larger samples, the true α -level is reduced and the tests become conservative. On the other hand, if the larger variances and covariances come from the smaller samples, α is inflated, and the tests become liberal. Box's M -test in Section 7.3.2 can be used to test for homogeneity of covariance matrices.

In conclusion, the use of Roy's θ is not recommended in any situation except the collinear case under standard assumptions. In the diffuse case its performance is inferior to that of the other three tests, both when the assumptions hold and when they do not. If the data come from nonnormal populations exhibiting skewness or positive kurtosis, any of the other three tests perform acceptably well. Among these three, $V^{(s)}$ is superior to the other two when there is heterogeneity of covariance matrices. Indeed $V^{(s)}$ is first in all rankings except those for the collinear case. However, Λ is not far behind, except when there is severe heterogeneity of covariance matrices. It seems likely that Wilks' Λ will continue its dominant role because of its flexibility and historical precedence.

In practice, most MANOVA software programs routinely calculate all four test statistics, and they usually reach the same conclusion. In those cases when they differ as to acceptance or rejection of the hypothesis, one can examine the eigenvalues and covariance matrices and evaluate the conflicting conclusions in light of the test properties discussed previously.

■ EXAMPLE 6.2

We inspect the eigenvalues of $\mathbf{E}^{-1}\mathbf{H}$ for the rootstock data of Table 6.2 for an indication of the configuration of the six mean vectors in a four-dimensional space. The eigenvalues are 1.876, .791, .229, .026. The first eigenvalue 1.876 constitutes a proportion

$$\frac{1.876}{1.876 + .791 + .229 + .026} = .642$$

of the sum of the eigenvalues. Therefore, the first eigenvalue does not dominate the others and the mean vectors are not collinear. The first two eigenvalues account for a proportion

$$\frac{1.876 + .791}{1.876 + \dots + .026} = .913$$

of the sum of the eigenvalues, and thus the six mean vectors lie largely in two dimensions. Since the mean vectors are not collinear, the test statistics Λ , $V^{(s)}$, and $U^{(s)}$ will be more appropriate than θ in this case. \square

6.3 CONTRASTS

As in Sections 6.1.1–6.1.5, we consider only the balanced model where $n_1 = n_2 = \dots = n_k = n$. We begin with a review of contrasts in the univariate setting before moving to the multivariate case.

6.3.1 Univariate Contrasts

A *contrast* in the population means is defined as a linear combination

$$\delta = c_1\mu_1 + c_2\mu_2 + \dots + c_k\mu_k, \quad (6.52)$$

where the coefficients satisfy

$$\sum_{i=1}^k c_i = 0. \quad (6.53)$$

An unbiased estimator of δ is given by

$$\hat{\delta} = c_1\bar{y}_{1.} + c_2\bar{y}_{2.} + \dots + c_k\bar{y}_{k.}. \quad (6.54)$$

The sample means $\bar{y}_{i.}$ were defined in (6.1). Since the $\bar{y}_{i.}$'s are independent with variance σ^2/n , the variance of $\hat{\delta}$ is

$$\text{var}(\hat{\delta}) = \frac{\sigma^2}{n} \sum_{i=1}^k c_i^2,$$

which can be estimated by

$$s_{\hat{\delta}}^2 = \frac{\text{MSE}}{n} \sum_{i=1}^k c_i^2, \quad (6.55)$$

where MSE was defined in (6.6) and (6.7) as $\text{SSE}/k(n - 1)$.

The usual hypothesis to be tested by a contrast is

$$H_0: \delta = c_1\mu_1 + c_2\mu_2 + \dots + c_k\mu_k = 0.$$

For example, suppose $k = 4$ and that a contrast of interest to the researcher is $3\mu_1 - \mu_2 - \mu_3 - \mu_4$. If this contrast is set equal to zero, we have

$$3\mu_1 = \mu_2 + \mu_3 + \mu_4 \quad \text{or} \quad \mu_1 = \frac{1}{3}(\mu_2 + \mu_3 + \mu_4),$$

and the experimenter is comparing the first mean with the average of the other three. A contrast is often called a *comparison* among the treatment means.

Assuming normality, $H_0 : \delta = 0$ can be tested by

$$t = \frac{\hat{\delta} - 0}{s_{\hat{\delta}}}, \quad (6.56)$$

which is distributed as t_{ν_E} . Alternatively, since $t^2 = F$, we can use

$$\begin{aligned} F &= \frac{\hat{\delta}^2}{s_{\hat{\delta}}^2} = \frac{\left(\sum_{i=1}^k c_i \bar{y}_{i\cdot} \right)^2}{\text{MSE} \sum_{i=1}^k c_i^2 / n} \\ &= \frac{n (\sum_i c_i \bar{y}_{i\cdot})^2 / \sum_i c_i^2}{\text{MSE}}, \end{aligned} \quad (6.57)$$

which is distributed as F_{1, ν_E} . The numerator of (6.57) is often referred to as the sum of squares for the contrast.

If two contrasts $\delta = \sum_i a_i \mu_i$ and $\gamma = \sum_i b_i \mu_i$ are such that $\sum_i a_i b_i = 0$, the contrasts are said to be *orthogonal*. The two estimated contrasts can be written in the form $\sum_i a_i \bar{y}_{i\cdot} = \mathbf{a}' \bar{\mathbf{y}}$ and $\sum_i b_i \bar{y}_{i\cdot} = \mathbf{b}' \bar{\mathbf{y}}$, where $\bar{\mathbf{y}} = (\bar{y}_1, \bar{y}_2, \dots, \bar{y}_k)'$. Then $\sum_i a_i b_i = \mathbf{a}' \mathbf{b} = 0$, and by the discussion following (3.14), the coefficient vectors \mathbf{a} and \mathbf{b} are perpendicular.

When two contrasts are orthogonal, the two corresponding sums of squares are independent. In fact, for k treatments, we can find $k - 1$ orthogonal contrasts that partition the treatment sum of squares SSH into $k - 1$ independent sums of squares, each with one degree of freedom. In the unbalanced case (Section 6.1.6), orthogonal contrasts such that $\sum_i a_i b_i = 0$ no longer partition SSH into $k - 1$ independent sums of squares. For a discussion of contrasts in the unbalanced case, see Rencher (1998, Sections 4.8.2 and 4.8.3) or Rencher and Schaalje (2008, Section 15.2.2).

6.3.2 Multivariate Contrasts

There are two usages of contrasts in a multivariate setting. We have previously encountered one use in Section 5.9.1, where we considered the hypothesis $H_0: \mathbf{C}\boldsymbol{\mu} = \mathbf{0}$ with $\mathbf{C}\mathbf{j} = \mathbf{0}$. Each row of \mathbf{C} sums to zero, and $\mathbf{C}\boldsymbol{\mu}$ is therefore a set of contrasts comparing the elements $\mu_1, \mu_2, \dots, \mu_p$ of $\boldsymbol{\mu}$ with each other. In this section, on the other hand, we consider contrasts comparing several mean vectors, not the elements within a vector.

A contrast among the population mean vectors is defined as

$$\boldsymbol{\delta} = c_1 \boldsymbol{\mu}_1 + c_2 \boldsymbol{\mu}_2 + \cdots + c_k \boldsymbol{\mu}_k, \quad (6.58)$$

where $\sum_{i=1}^k c_i = 0$. An unbiased estimator of δ is given by the corresponding contrast in the sample mean vectors:

$$\hat{\delta} = c_1 \bar{y}_{1\cdot} + c_2 \bar{y}_{2\cdot} + \cdots + c_k \bar{y}_{k\cdot}. \quad (6.59)$$

The sample mean vectors $\bar{y}_{1\cdot}, \bar{y}_{2\cdot}, \dots, \bar{y}_{k\cdot}$ as defined in Section 6.1.2 were assumed to be independent and to have common covariance matrix, $\text{cov}(\bar{y}_{i\cdot}) = \Sigma/n$. Thus the covariance matrix for $\hat{\delta}$ is given by

$$\text{cov}(\hat{\delta}) = c_1^2 \frac{\Sigma}{n} + c_2^2 \frac{\Sigma}{n} + \cdots + c_k^2 \frac{\Sigma}{n} = \frac{\Sigma}{n} \sum_{i=1}^k c_i^2, \quad (6.60)$$

which can be estimated by

$$\frac{S_{pl}}{n} \sum_{i=1}^k c_i^2 = \left(\frac{E}{\nu_E} \right) \left(\frac{\sum_{i=1}^k c_i^2}{n} \right),$$

where $S_{pl} = E/\nu_E$ is an unbiased estimator of Σ .

The hypothesis $H_0 : \delta = \mathbf{0}$ or $H_0 : c_1 \mu_1 + c_2 \mu_2 + \cdots + c_k \mu_k = \mathbf{0}$ makes comparisons among the population mean vectors. For example, $\mu_1 - 2\mu_2 + \mu_3 = \mathbf{0}$ is equivalent to

$$\mu_2 = \frac{1}{2}(\mu_1 + \mu_3),$$

and we are comparing μ_2 to the average of μ_1 and μ_3 . Of course, this implies that every element of μ_2 must equal the corresponding element of $\frac{1}{2}(\mu_1 + \mu_3)$:

$$\begin{pmatrix} \mu_{21} \\ \mu_{22} \\ \vdots \\ \mu_{2p} \end{pmatrix} = \begin{pmatrix} \frac{1}{2}(\mu_{11} + \mu_{31}) \\ \frac{1}{2}(\mu_{12} + \mu_{32}) \\ \vdots \\ \frac{1}{2}(\mu_{1p} + \mu_{3p}) \end{pmatrix}.$$

Under appropriate multivariate normality assumptions, $H_0 : c_1 \mu_1 + c_2 \mu_2 + \cdots + c_k \mu_k = \mathbf{0}$ or $H_0 : \delta = \mathbf{0}$ can be tested with the one-sample T^2 -statistic

$$\begin{aligned} T^2 &= \hat{\delta}' \left(\frac{S_{pl}}{n} \sum_{i=1}^k c_i^2 \right)^{-1} \hat{\delta} \\ &= \frac{n}{\sum_{i=1}^k c_i^2} \left(\sum_{i=1}^k c_i \bar{y}_{i\cdot} \right)' \left(\frac{E}{\nu_E} \right)^{-1} \left(\sum_{i=1}^k c_i \bar{y}_{i\cdot} \right), \end{aligned} \quad (6.61)$$

which is distributed as T_{p, ν_E}^2 . In the one-way model under discussion here, $\nu_E = k(n-1)$.

An equivalent test of H_0 can be made with Wilks' Λ . By analogy with the numerator of (6.57), the hypothesis matrix due to the contrast is given by

$$\mathbf{H}_1 = \frac{n}{\sum_{i=1}^k c_i^2} \left(\sum_{i=1}^k c_i \bar{y}_{i\cdot} \right) \left(\sum_{i=1}^k c_i \bar{y}_{i\cdot} \right)'. \quad (6.62)$$

The rank of \mathbf{H}_1 is 1, and the test statistic is

$$\Lambda = \frac{|\mathbf{E}|}{|\mathbf{E} + \mathbf{H}_1|}, \quad (6.63)$$

which is distributed as $\Lambda_{p,1,\nu_E}$. The other three MANOVA test statistics can also be applied here using the single nonzero eigenvalue of $\mathbf{E}^{-1}\mathbf{H}_1$. Because $\nu_H = 1$ in this case, all four MANOVA statistics and T^2 give the same results; that is, all five transform to the same F -value using the formulations in Section 6.1.7. If $k - 1$ orthogonal contrasts are used, they partition the \mathbf{H} matrix into $k - 1$ independent matrices $\mathbf{H}_1, \mathbf{H}_2, \dots, \mathbf{H}_{k-1}$. Each \mathbf{H}_i matrix has one degree of freedom because $\text{rank}(\mathbf{H}_i) = 1$.

■ EXAMPLE 6.3.2

We consider the following two orthogonal contrasts for the rootstock data in Table 6.2:

$$\begin{matrix} 2 & -1 & -1 & -1 & -1 & 2 \\ 1 & 0 & 0 & 0 & 0 & -1 \end{matrix}$$

The first compares μ_1 and μ_6 with the other four mean vectors. The second compares μ_1 vs. μ_6 . Thus $H_{01}: 2\mu_1 - \mu_2 - \mu_3 - \mu_4 - \mu_5 + 2\mu_6 = \mathbf{0}$ can be written as

$$H_{01}: 2\mu_1 + 2\mu_6 = \mu_2 + \mu_3 + \mu_4 + \mu_5.$$

Dividing both sides by 4 so as to express this in terms of averages, we obtain

$$H_{01}: \frac{1}{2}(\mu_1 + \mu_6) = \frac{1}{4}(\mu_2 + \mu_3 + \mu_4 + \mu_5).$$

Similarly, the hypothesis for the second contrast can be expressed as

$$H_{02}: \mu_1 = \mu_6.$$

The mean vectors are given by

$\bar{\mathbf{y}}_1.$	$\bar{\mathbf{y}}_2.$	$\bar{\mathbf{y}}_3.$	$\bar{\mathbf{y}}_4.$	$\bar{\mathbf{y}}_5.$	$\bar{\mathbf{y}}_6.$
1.14	1.16	1.11	1.10	1.08	1.04
2.98	3.11	2.82	2.88	2.56	2.21
3.74	4.52	4.46	3.91	4.31	3.60
.87	1.28	1.39	1.04	1.18	.74

For the first contrast, we obtain \mathbf{H}_1 from (6.62) as

$$\begin{aligned} \mathbf{H}_1 &= \frac{n}{\sum_i c_i^2} \left(\sum_i c_i \bar{\mathbf{y}}_i. \right) \left(\sum_i c_i \bar{\mathbf{y}}_i. \right)' \\ &= \frac{8}{12} (2\bar{\mathbf{y}}_1. - \bar{\mathbf{y}}_2. - \cdots + 2\bar{\mathbf{y}}_6.) (2\bar{\mathbf{y}}_1. - \bar{\mathbf{y}}_2. - \cdots + 2\bar{\mathbf{y}}_6.)' \\ &= \frac{8}{12} \begin{pmatrix} -.095 \\ -.978 \\ -2.519 \\ -1.680 \end{pmatrix} (-.095, -.978, -2.519, -1.680) \end{aligned}$$

$$= \begin{pmatrix} .006 & .062 & .160 & .106 \\ .062 & .638 & 1.642 & 1.095 \\ .160 & 1.642 & 4.229 & 2.820 \\ .106 & 1.095 & 2.820 & 1.881 \end{pmatrix}.$$

Then

$$\Lambda = \frac{|\mathbf{E}|}{|\mathbf{E} + \mathbf{H}_1|} = \frac{.6571}{1.4824} = .443,$$

which is less than $\Lambda_{.05, 4, 1, 40} = .779$ from Table A.9. We therefore reject H_{01} .

To test the significance of the second contrast, we have

$$\begin{aligned} \mathbf{H}_2 &= \frac{8}{2} (\bar{\mathbf{y}}_{1\cdot} - \bar{\mathbf{y}}_{6\cdot})(\bar{\mathbf{y}}_{1\cdot} - \bar{\mathbf{y}}_{6\cdot})' \\ &= \frac{8}{2} \begin{pmatrix} .101 \\ .762 \\ .142 \\ .136 \end{pmatrix} (.101, .762, .142, .136) \\ &= \begin{pmatrix} .041 & .309 & .058 & .055 \\ .309 & 2.326 & .435 & .415 \\ .058 & .435 & .081 & .078 \\ .055 & .415 & .078 & .074 \end{pmatrix}. \end{aligned}$$

Then

$$\Lambda = \frac{|\mathbf{E}|}{|\mathbf{E} + \mathbf{H}_2|} = \frac{.6571}{.8757} = .750,$$

which is less than $\Lambda_{.05, 4, 1, 40} = .779$, and we reject H_{02} . \square

6.4 TESTS ON INDIVIDUAL VARIABLES FOLLOWING REJECTION OF H_0 BY THE OVERALL MANOVA TEST

In Section 6.1, we considered tests of equality of mean vectors, $H_0: \boldsymbol{\mu}_1 = \boldsymbol{\mu}_2 = \dots = \boldsymbol{\mu}_k$, which implies equality of means for each of the p variables:

$$H_{0r}: \mu_{1r} = \mu_{2r} = \dots = \mu_{kr}, \quad r = 1, 2, \dots, p.$$

This hypothesis could be tested for each variable by itself with an ordinary univariate ANOVA F -test, as noted in property 9 in Section 6.1.3. For example, if there are three mean vectors,

$$\boldsymbol{\mu}_1 = \begin{pmatrix} \mu_{11} \\ \mu_{12} \\ \vdots \\ \mu_{1p} \end{pmatrix}, \quad \boldsymbol{\mu}_2 = \begin{pmatrix} \mu_{21} \\ \mu_{22} \\ \vdots \\ \mu_{2p} \end{pmatrix}, \quad \boldsymbol{\mu}_3 = \begin{pmatrix} \mu_{31} \\ \mu_{32} \\ \vdots \\ \mu_{3p} \end{pmatrix},$$

we have $H_{01}: \mu_{11} = \mu_{21} = \mu_{31}$, $H_{02}: \mu_{12} = \mu_{22} = \mu_{32}, \dots$, $H_{0p}: \mu_{1p} = \mu_{2p} = \mu_{3p}$. Each of these p hypotheses can be tested with a simple ANOVA F -test.

If an F -test is made on each of the p variables regardless of whether the overall MANOVA test of $H_0: \mu_1 = \mu_2 = \mu_3$ rejects H_0 , then the overall α -level will increase beyond the nominal value because we are making p tests. As in Section 5.5, we define the *overall α* or *experimentwise error rate* as the probability of rejecting one or more of the p univariate tests when $H_0: \mu_1 = \mu_2 = \mu_3$ is true. We could “protect” against inflation of the experimentwise error rate by performing tests on individual variables *only* if the overall MANOVA test of $H_0: \mu_1 = \mu_2 = \mu_3$ is rejected. In this procedure, the probability of rejection for the tests on individual variables is reduced, and these tests become more conservative.

Rencher and Scott (1990) compared the above two procedures for testing the individual variables in a one-way MANOVA model. Since the focus was on α -levels, only the case where H_0 is true was considered. Specifically, the two procedures were as follows:

1. A univariate F -test is made on each variable, testing $H_{0r}: \mu_{1r} = \mu_{2r} = \dots = \mu_{kr}, r = 1, 2, \dots, p$. In this context, the p univariate tests constitute an experiment and one or more rejections are counted as one experimentwise error. No multivariate test is made.
2. The overall MANOVA hypothesis $H_0: \mu_1 = \mu_2 = \dots = \mu_k$ is tested with Wilks' Λ , and if H_0 is rejected, p univariate F -tests on $H_{01}, H_{02}, \dots, H_{0p}$ are carried out. Again, one or more rejections among the F -tests are counted as one experimentwise error.

The amount of intercorrelation among the multivariate normal variables was indicated by $\sum_{i=1}^p (1/\lambda_i)/p$, where $\lambda_1, \lambda_2, \dots, \lambda_p$ are the eigenvalues of the population correlation matrix \mathbf{P}_ρ . Note that $\sum_i (1/\lambda_i)/p = 1$ for the uncorrelated case ($\mathbf{P}_\rho = \mathbf{I}$) and $\sum_i (1/\lambda_i)/p > 1$ for the correlated case ($\mathbf{P}_\rho \neq \mathbf{I}$). When the variables are highly intercorrelated, one or more of the eigenvalues will be near zero (see Section 4.1.3), and $\sum_i (1/\lambda_i)/p$ will be large.

The error rates of these two procedures were investigated for several values of p , n , k , and $\sum_i (1/\lambda_i)/p$, where p is the number of variables, n is the number of observation vectors in each group, k is the number of groups, and $\sum_i (1/\lambda_i)/p$ is the measure of intercorrelation defined above. In procedure 1, the probability of rejecting one or more univariate tests when H_0 is true varied from .09 to .31 (α was .05 in each test). Such experimentwise error rates are clearly unacceptable when the nominal value of α is .05. However, this approach is commonly used when the researcher is not familiar with the MANOVA approach or does not have access to appropriate software.

Table 6.3 contains the error rates for procedure 2, univariate F -tests following a rejection by Wilks' Λ . The values range from .022 to .057, comfortably close to the target value of .05. No apparent trends or patterns are seen; the values do not seem to depend on p , k , n , or amount of intercorrelation as measured by $\sum_i (1/\lambda_i)/p$. Thus when univariate tests are made *only* following a rejection of the overall test, the experimentwise error rate is about right.

Table 6.3 Experimentwise Error Rates for Procedure 2: Univariate F -Tests Following Rejection by Wilks' Λ

		$\sum_i(1/\lambda_i)/p$							
n	p	1		10		100		1000	
		$k = 3$	$k = 5$	$k = 3$	$k = 5$	$k = 3$	$k = 5$	$k = 3$	$k = 5$
5	3	.043	.037	.022	.035	.046	.039	.022	.029
5	5	.041	.037	.039	.057	.038	.035	.027	.039
5	7	.030	.042	.035	.045	.039	.037	.026	.048
10	3	.047	.041	.030	.033	.043	.045	.026	.032
10	5	.047	.037	.026	.049	.041	.026	.027	.029
10	7	.034	.054	.037	.047	.047	.040	.040	.044
20	3	.050	.043	.032	.054	.048	.039	.040	.032
20	5	.045	.055	.042	.051	.037	.044	.050	.043
20	7	.055	.051	.029	.040	.033	.051	.039	.033

Based on these results, we recommend making an overall MANOVA test followed by F -tests on the individual variables (at the same α -level as the MANOVA test) only if the MANOVA test leads to rejection of H_0 .

Another procedure that can be used following rejection of the MANOVA test is an examination of the discriminant function coefficients. The discriminant function was defined in Section 6.1.4 as $z = \mathbf{a}_1' \mathbf{y}$, where \mathbf{a}_1 is the eigenvector associated with the largest eigenvalue λ_1 of $\mathbf{E}^{-1} \mathbf{H}$. Additionally, there are other discriminant functions using eigenvectors corresponding to the other eigenvalues. Since the first discriminant function maximally separates the groups, we can examine its coefficients for the contribution of each variable to group separation. Thus in $z = a_{11}y_1 + a_{12}y_2 + \dots + a_{1p}y_p$, if a_{12} is larger than the other a_{1r} 's, we believe y_2 contributes more than any of the other variables to separation of groups. A method of standardization of the a_{1r} 's to adjust for differences in the scale among the variables is given in Section 8.5.

The information in a_{1r} (from $z = \mathbf{a}_1' \mathbf{y}$) about the contribution of y_r to separation of the groups is fundamentally different from the information provided in a univariate F -test that considers y_r alone (see property 9 in Section 6.1.3). The relative size of a_{1r} shows the contribution of y_r in the presence of the other variables and takes into account (1) the correlation of y_r with the others y 's and (2) the contribution of y_r to Wilks' Λ above and beyond the contribution of the other y 's. In contrast, the individual F -test on y_r ignores the presence of the other variables. Because we are primarily interested in the collective behavior of the variables, the discriminant function coefficients provide more pertinent information than the tests on individual variables. For a detailed analysis of the effect of each variable in the presence of other variables, see Rencher (1993; 1998, Section 4.1.6).

Huberty (1975) compared the standardized coefficients to some correlations that can be shown to be related to individual variable tests (see Section 8.7.3). In a limited simulation, the discriminant coefficients were found to be more valid than the univariate tests in identifying those variables that contribute least to separation of groups. Considerable variation was found from sample to sample in ranking the relative potency of the variables.

■ EXAMPLE 6.4

In Example 6.1.7, the hypothesis $H_0: \mu_1 = \mu_2 = \dots = \mu_6$ was rejected for the rootstock data of Table 6.2. We can therefore test the four individual variables using the .05 level of significance. For the first variable, $y_1 = 4\text{-year trunk girth}$, we obtain the following ANOVA table:

Source	Sum of Squares	df	Mean Square	F
Rootstocks	.073560	5	.014712	1.93
Error	.319988	42	.007619	
Total	.393548	47		

For $F = 1.93$ the p -value is .1094, and we do not reject H_0 . For the other three variables we have

Variable	F	p-Value
$y_2 = 4\text{-year extension growth}$	2.91	.024
$y_3 = 15\text{-year trunk girth}$	11.97	< .0001
$y_4 = 15\text{-year weight}$	12.16	< .0001

Thus for three of the four variables, the six means differ significantly. We examine the standardized discriminant function coefficients for this set of data in Chapter 8 (Problem 8.12). \square

6.5 TWO-WAY CLASSIFICATION

We consider only balanced models, where each cell in the model has the same number of observations, n . For the unbalanced case with unequal cell sizes, see Rencher (1998, Section 4.8).

6.5.1 Review of Univariate Two-Way ANOVA

In the univariate two-way model, we measure one dependent variable y on each experimental unit. The balanced two-way fixed-effects model with factors A and B is

$$y_{ijk} = \mu + \alpha_i + \beta_j + \gamma_{ij} + \varepsilon_{ijk} \quad (6.64)$$

$$= \mu_{ij} + \varepsilon_{ijk}, \quad (6.65)$$

$$i = 1, 2, \dots, a, \quad j = 1, 2, \dots, b, \quad k = 1, 2, \dots, n,$$

where α_i is the effect (on y_{ijk}) of the i th level of A , β_j is the effect of the j th level of B , γ_{ij} is the corresponding interaction effect, and μ_{ij} is the population mean for the i th level of A and the j th level of B . In order to obtain F -tests, we further assume that the ε_{ijk} are independently distributed as $N(0, \sigma^2)$.

Let $\bar{\mu}_{i..} = \sum_j \mu_{ij}/b$ be the mean at the i th level of A and define $\bar{\mu}_{..j}$ and $\bar{\mu}_{...}$ similarly. Then if we use side conditions $\sum_i \alpha_i = \sum_j \beta_j = \sum_i \gamma_{ij} = \sum_j \gamma_{ij} = 0$, the effect of the i th level of A can be defined as $\alpha_i = \bar{\mu}_{i..} - \bar{\mu}_{..}$, with similar definitions of β_j and γ_{ij} . We can show that $\sum_i \alpha_i = 0$ if $\alpha_i = \bar{\mu}_{i..} - \bar{\mu}_{..}$ as follows:

$$\begin{aligned}\sum_{i=1}^a \alpha_i &= \sum_{i=1}^a (\bar{\mu}_{i..} - \bar{\mu}_{..}) = \sum_i \bar{\mu}_{i..} - a\bar{\mu}_{..} \\ &= a\bar{\mu}_{..} - a\bar{\mu}_{..} = 0.\end{aligned}$$

Many texts recommend that the interaction AB be tested first, and if it is found to be significant, then the main effects should not be tested. However, with the side conditions imposed above (side conditions are not necessary in order to obtain tests), the effect of A is defined as the average effect over the levels of B and similarly for the effect of B . With this definition of main effects, the tests for A and B make sense even if AB is significant. Admittedly, interpretation requires more care, and the effect of a factor may vary if the number of levels of the other factor is altered. But in many cases, useful information can be gained about the main effects in the presence of interaction.

We illustrate the preceding statement that $\alpha_i = \bar{\mu}_{i..} - \bar{\mu}_{..}$ represents the effect of the i th level of A averaged over the levels of B . Suppose A has two levels and B has three. We represent the means of the six cells as follows:

		B			Mean
		1	2	3	
A	1	μ_{11}	μ_{12}	μ_{13}	$\bar{\mu}_{1..}$
	2	μ_{21}	μ_{22}	μ_{23}	$\bar{\mu}_{2..}$
Mean		$\bar{\mu}_{.1}$	$\bar{\mu}_{.2}$	$\bar{\mu}_{.3}$	$\bar{\mu}_{..}$

The means of the rows (corresponding to levels of A) and columns (levels of B) are also given. Then $\alpha_i = \bar{\mu}_{i..} - \bar{\mu}_{..}$ can be expressed as the average of the effect of the i th level of A at the three levels of B . For example,

$$\begin{aligned}\alpha_1 &= \frac{1}{3}[(\mu_{11} - \bar{\mu}_{.1}) + (\mu_{12} - \bar{\mu}_{.2}) + (\mu_{13} - \bar{\mu}_{.3})] \\ &= \frac{1}{3}(\mu_{11} + \mu_{12} + \mu_{13}) - \frac{1}{3}(\bar{\mu}_{.1} + \bar{\mu}_{.2} + \bar{\mu}_{.3}) = \bar{\mu}_{1..} - \bar{\mu}_{..}.\end{aligned}$$

To estimate α_i , we can use $\hat{\alpha}_i = \bar{y}_{i..} - \bar{y}_{..}$, with similar estimates for β_j and γ_{ij} . The notation $\bar{y}_{i..}$ indicates that y_{ijk} is averaged over the levels of j and k to obtain the mean of all nb observations at the i th level of A , namely, $\bar{y}_{i..} = \sum_{jk} y_{ijk}/nb$. The means $\bar{y}_{..j}$, $\bar{y}_{ij..}$, and $\bar{y}_{...}$ have analogous definitions.

To construct tests for the significance of factors A and B and the interaction AB , we use the usual sums of squares and degrees of freedom in Table 6.4. Computational forms of the sums of squares can be found in many standard (univariate) methods texts.

Table 6.4 Univariate Two-Way Analysis of Variance

Source	Sum of Squares	df
A	$SSA = nb \sum_i (\bar{y}_{i..} - \bar{y}_{...})^2$	$a - 1$
B	$SSB = na \sum_j (\bar{y}_{.j} - \bar{y}_{...})^2$	$b - 1$
AB	$SSAB = n \sum_{ij} (\bar{y}_{ij} - \bar{y}_{i..} - \bar{y}_{.j} + \bar{y}_{...})^2$	$(a - 1)(b - 1)$
Error	$SSE = \sum_{ijk} (y_{ijk} - \bar{y}_{ijk})^2$	$ab(n - 1)$
Total	$SST = \sum_{ijk} (y_{ijk} - \bar{y}_{...})^2$	$abn - 1$

The sums of squares in Table 6.4 (for the balanced model) have the relationship

$$SST = SSA + SSB + SSAB + SSE,$$

and the four sums of squares on the right are independent. The sums of squares are divided by their corresponding degrees of freedom to obtain mean squares MSA, MSB, MSAB, and MSE. For the fixed-effects model, each of MSA, MSB, and MSAB is divided by MSE to obtain an F -test. In the case of factor A, for example, the hypothesis can be expressed as

$$H_{0A}: \alpha_1 = \alpha_2 = \cdots = \alpha_a = 0,$$

and the test statistic is $F = MSA/MSE$, which is distributed as $F_{a-1, ab(n-1)}$.

In order to define contrasts among the levels of each main effect, we can conveniently use the model in the form given in (6.65),

$$y_{ijk} = \mu_{ij} + \varepsilon_{ijk}.$$

A contrast among the levels of A is defined as $\sum_{i=1}^a c_i \bar{\mu}_i$, where $\sum_i c_i = 0$. An estimate of the contrast is given by $\sum_i c_i \bar{y}_{i..}$, with variance $\sigma^2 \sum_i c_i^2 / nb$, since each $\bar{y}_{i..}$ is based on nb observations and the $\bar{y}_{i..}$'s are independent. To test $H_0: \sum_i c_i \bar{\mu}_i = 0$, we can use an F -statistic corresponding to (6.57),

$$F = \frac{nb (\sum_{i=1}^a c_i \bar{y}_{i..})^2 / \sum_{i=1}^a c_i^2}{MSE}, \quad (6.66)$$

with 1 and ν_E degrees of freedom. To preserve the experimentwise error rate, significance tests for more than one contrast could be carried out in the spirit of Section 6.4; that is, contrasts should be chosen prior to seeing the data and tests should be made only if the overall F -test for factor A rejects H_{0A} .

Contrasts $\sum_j c_j \bar{\mu}_{.j}$ among the levels of B are tested in an entirely analogous manner.

Table 6.5 Multivariate Two-Way Analysis of Variance

Source	Sum of Squares and Products Matrix	df
<i>A</i>	$\mathbf{H}_A = nb \sum_i (\bar{\mathbf{y}}_{i..} - \bar{\mathbf{y}}_{...})(\bar{\mathbf{y}}_{i..} - \bar{\mathbf{y}}_{...})'$	$a - 1$
<i>B</i>	$\mathbf{H}_B = na \sum_j (\bar{\mathbf{y}}_{.j.} - \bar{\mathbf{y}}_{...})(\bar{\mathbf{y}}_{.j.} - \bar{\mathbf{y}}_{...})'$	$b - 1$
<i>AB</i>	$\mathbf{H}_{AB} = n \sum_{ij} (\bar{\mathbf{y}}_{ij.} - \bar{\mathbf{y}}_{i..} - \bar{\mathbf{y}}_{.j.} + \bar{\mathbf{y}}_{...}) \times (\bar{\mathbf{y}}_{ij.} - \bar{\mathbf{y}}_{i..} - \bar{\mathbf{y}}_{.j.} + \bar{\mathbf{y}}_{...})'$	$(a - 1)(b - 1)$
Error	$\mathbf{E} = \sum_{ijk} (\mathbf{y}_{ijk} - \bar{\mathbf{y}}_{ij.})(\mathbf{y}_{ijk} - \bar{\mathbf{y}}_{ij.})'$	$ab(n - 1)$
Total	$\mathbf{T} = \sum_{ijk} (\mathbf{y}_{ijk} - \bar{\mathbf{y}}_{...})(\mathbf{y}_{ijk} - \bar{\mathbf{y}}_{...})'$	$abn - 1$

6.5.2 Multivariate Two-Way MANOVA

A balanced two-way fixed-effects MANOVA model for p dependent variables can be expressed in vector form analogous to (6.64) and (6.65):

$$\mathbf{y}_{ijk} = \boldsymbol{\mu} + \boldsymbol{\alpha}_i + \boldsymbol{\beta}_j + \boldsymbol{\gamma}_{ij} + \boldsymbol{\varepsilon}_{ijk} = \boldsymbol{\mu}_{ij} + \boldsymbol{\varepsilon}_{ijk}, \quad (6.67)$$

$$i = 1, 2, \dots, a, \quad j = 1, 2, \dots, b, \quad k = 1, 2, \dots, n,$$

where $\boldsymbol{\alpha}_i$ is the effect of the i th level of A on each of the p variables in \mathbf{y}_{ijk} , $\boldsymbol{\beta}_j$ is the effect of the j th level of B , and $\boldsymbol{\gamma}_{ij}$ is the AB interaction effect. We use side conditions $\sum_i \boldsymbol{\alpha}_i = \sum_j \boldsymbol{\beta}_j = \sum_i \boldsymbol{\gamma}_{ij} = \sum_j \boldsymbol{\gamma}_{ij} = \mathbf{0}$ and assume the $\boldsymbol{\varepsilon}_{ijk}$'s are independently distributed as $N_p(\mathbf{0}, \boldsymbol{\Sigma})$. Under the side condition $\sum_i \boldsymbol{\alpha}_i = \mathbf{0}$, the effect of A is averaged over the levels of B ; that is, $\boldsymbol{\alpha}_i = \bar{\boldsymbol{\mu}}_i - \bar{\boldsymbol{\mu}}_{..}$, where $\bar{\boldsymbol{\mu}}_i = \sum_j \boldsymbol{\mu}_{ij}/b$ and $\bar{\boldsymbol{\mu}}_{..} = \sum_{ij} \boldsymbol{\mu}_{ij}/ab$. There are similar definitions for $\boldsymbol{\beta}_j$ and $\boldsymbol{\gamma}_{ij}$.

As in the univariate usage, the mean vector $\bar{\mathbf{y}}_{i..}$ indicates an average over the subscripts replaced by dots, that is, $\bar{\mathbf{y}}_{i..} = \sum_{jk} \mathbf{y}_{ijk}/nb$. The means $\bar{\mathbf{y}}_{.j.}$, $\bar{\mathbf{y}}_{ij.}$, and $\bar{\mathbf{y}}_{...}$ have analogous definitions: $\bar{\mathbf{y}}_{.j.} = \sum_{ik} \mathbf{y}_{ijk}/na$, $\bar{\mathbf{y}}_{ij.} = \sum_k \mathbf{y}_{ijk}/n$, $\bar{\mathbf{y}}_{...} = \sum_{ijk} \mathbf{y}_{ijk}/nab$. The sum of squares and products matrices are given in Table 6.5. Note that the degrees of freedom in Table 6.5 are the same as in the univariate case in Table 6.4. For the two-way model with balanced data, the total sum of squares and products matrix is partitioned as

$$\mathbf{T} = \mathbf{H}_A + \mathbf{H}_B + \mathbf{H}_{AB} + \mathbf{E}. \quad (6.68)$$

The structure of any of the hypothesis matrices is similar to that of \mathbf{H} in (6.11). For example, \mathbf{H}_A has on the diagonal the sum of squares for factor A for each of the p variables. The off-diagonal elements of \mathbf{H}_A are corresponding sums of products for all pairs of variables. Thus the r th diagonal element of \mathbf{H}_A corresponding to the r th variable, $r = 1, 2, \dots, p$, is given by

$$h_{Arr} = nb \sum_{i=1}^a (\bar{y}_{i..r} - \bar{y}_{...r})^2 = \sum_{i=1}^a \frac{y_{i..r}^2}{nb} - \frac{y_{...r}^2}{nab}, \quad (6.69)$$

where $\bar{y}_{i..r}$ and $\bar{y}_{...r}$ represent the r th components of $\bar{\mathbf{y}}_{i..}$ and $\bar{\mathbf{y}}_{...}$, respectively, and $y_{i..r}$ and $y_{...r}$ are totals corresponding to $\bar{y}_{i..r}$ and $\bar{y}_{...r}$. The (rs) th off-diagonal element of \mathbf{H}_A is

$$h_{Ars} = nb \sum_{i=1}^a (\bar{y}_{i..r} - \bar{y}_{...r})(\bar{y}_{i..s} - \bar{y}_{...s}) = \sum_{i=1}^a \frac{y_{i..r}y_{i..s}}{nb} - \frac{y_{...r}y_{...s}}{nab}. \quad (6.70)$$

From (6.68) and Table 6.5, we obtain

$$\begin{aligned} h_{ABrr} &= \sum_{ij} \frac{y_{ij.r}^2}{n} - \frac{y_{...r}^2}{nab} - h_{Arr} - h_{Brr} \\ h_{ABrs} &= \sum_{ij} \frac{y_{ij.r}y_{ij.s}}{n} - \frac{y_{...r}y_{...s}}{nab} - h_{Ars} - h_{Brs}. \end{aligned} \quad (6.71)$$

For the \mathbf{E} matrix, computational formulas are based on (6.68):

$$\mathbf{E} = \mathbf{T} - \mathbf{H}_A - \mathbf{H}_B - \mathbf{H}_{AB}.$$

Thus the elements of \mathbf{E} have the form

$$\begin{aligned} e_{rr} &= \sum_{ijk} y_{ijk.r}^2 - \frac{y_{...r}^2}{nab} - h_{Arr} - h_{Brr} - h_{ABrr}, \\ e_{rs} &= \sum_{ijk} y_{ijk.r}y_{ijk.s} - \frac{y_{...r}y_{...s}}{nab} - h_{Ars} - h_{Brs} - h_{ABrs}. \end{aligned} \quad (6.72)$$

The hypotheses matrices for interaction and main effects in this fixed-effects model can be compared to \mathbf{E} to make a test. Thus for Wilks' Λ , we use \mathbf{E} to test each of A , B , and AB :

$$\begin{aligned} \Lambda_A &= \frac{|\mathbf{E}|}{|\mathbf{E} + \mathbf{H}_A|} \text{ is } \Lambda_{p,a-1,ab(n-1)}, \\ \Lambda_B &= \frac{|\mathbf{E}|}{|\mathbf{E} + \mathbf{H}_B|} \text{ is } \Lambda_{p,b-1,ab(n-1)}, \\ \Lambda_{AB} &= \frac{|\mathbf{E}|}{|\mathbf{E} + \mathbf{H}_{AB}|} \text{ is } \Lambda_{p,(a-1)(b-1),ab(n-1)}. \end{aligned}$$

In each case, the indicated distribution holds when H_0 is true. To calculate the other three MANOVA test statistics for A , B , and AB , we use the eigenvalues of $\mathbf{E}^{-1}\mathbf{H}_A$, $\mathbf{E}^{-1}\mathbf{H}_B$, and $\mathbf{E}^{-1}\mathbf{H}_{AB}$.

If the interaction is not significant, interpretation of the main effects is simpler. However, the comments in Section 6.5.1 about testing main effects in the presence of interaction apply to the multivariate model as well. If we define each main effect as the average effect over the levels of the other factor, then main effects can be tested even if the interaction is significant. One must be more careful with the interpretation in case of a significant interaction, but there is information to be gained.

By analogy with the univariate two-way ANOVA in Section 6.5.1, a contrast among the levels of factor A can be defined in terms of the mean vectors as follows: $\sum_{i=1}^a c_i \bar{\mu}_{i..}$, where $\sum_i c_i = 0$ and $\bar{\mu}_{i..} = \sum_j \mu_{ij}/b$. Similarly, $\sum_{j=1}^b c_j \bar{\mu}_{.j}$ represents a contrast among the levels of B . The hypothesis that these contrasts are 0 can be tested by T^2 or any of the four MANOVA test statistics, as in (6.61), (6.62), and (6.63). To test $H_0: \sum_i c_i \bar{\mu}_{i..} = 0$, for example, we can use

$$T^2 = \frac{nb}{\sum_{i=1}^a c_i^2} \left(\sum_{i=1}^a c_i \bar{\mathbf{y}}_{i..} \right)' \left(\frac{\mathbf{E}}{\nu_E} \right)^{-1} \left(\sum_{i=1}^a c_i \bar{\mathbf{y}}_{i..} \right), \quad (6.73)$$

which is distributed as T_{p,ν_E}^2 when H_0 is true. Alternatively, the hypothesis matrix

$$\mathbf{H}_1 = \frac{nb}{\sum_{i=1}^a c_i^2} \left(\sum_{i=1}^a c_i \bar{\mathbf{y}}_{i..} \right) \left(\sum_{i=1}^a c_i \bar{\mathbf{y}}_{i..} \right)' \quad (6.74)$$

can be used in

$$\Lambda = \frac{|\mathbf{E}|}{|\mathbf{E} + \mathbf{H}_1|},$$

which, under H_0 , is $\Lambda_{p,1,\nu_E}$, with $\nu_E = ab(n - 1)$ in the two-way model. The other three MANOVA test statistics can also be constructed from $\mathbf{E}^{-1}\mathbf{H}_1$. All five test statistics will give equivalent results because $\nu_H = 1$.

If follow-up tests on individual variables are desired, we can infer from Rencher and Scott (1990), as reported in Section 6.4, that if the MANOVA test on factor A or B leads to rejection of H_0 , then we can proceed with the univariate F -tests on the individual variables with assurance that the experimentwise error rate will be close to α .

To determine the contribution of each variable in the presence of the others, we can examine the first discriminant function obtained from eigenvectors of $\mathbf{E}^{-1}\mathbf{H}_A$ or $\mathbf{E}^{-1}\mathbf{H}_B$, as in Section 6.4 for one-way MANOVA. The first discriminant function for $\mathbf{E}^{-1}\mathbf{H}_A$, for example, is $z = \mathbf{a}'\mathbf{y}$, where \mathbf{a} is the eigenvector associated with the largest eigenvalue of $\mathbf{E}^{-1}\mathbf{H}_A$. In $z = a_1 y_1 + a_2 y_2 + \dots + a_p y_p$, if a_r is larger than the other a 's, then y_r contributes more than the other variables to the significance of Λ_A . (In many cases, the a_r 's should be standardized as in Section 8.5.) Note that the first discriminant function obtained from $\mathbf{E}^{-1}\mathbf{H}_A$ will not have the same pattern as the first discriminant function from $\mathbf{E}^{-1}\mathbf{H}_B$. This is not surprising since we expect that the relative contribution of the variables to separating the levels of factor A will be different from the relative contribution to separating the levels of B .

A randomized block design or a two-way MANOVA without replication can easily be analyzed in a manner similar to that for the two-way model with replication given here; therefore, no specific details will be given.

■ EXAMPLE 6.5.2

Table 6.6 contains data reported by Posten (1962) and analyzed by Kramer and Jensen (1970). The experiment involved a 2×4 design with 4 replications, for

Table 6.6 Two-Way Classification of Measurements on Bar Steel

Lubricant	A_1		A_2	
	y_1	y_2	y_1	y_2
B_1	7.80	90.4	7.12	85.1
	7.10	88.9	7.06	89.0
	7.89	85.9	7.45	75.9
	7.82	88.8	7.45	77.9
B_2	9.00	82.5	8.19	66.0
	8.43	92.4	8.25	74.5
	7.65	82.4	7.45	83.1
	7.70	87.4	7.45	86.4
B_3	7.28	79.6	7.15	81.2
	8.96	95.1	7.15	72.0
	7.75	90.2	7.70	79.9
	7.80	88.0	7.45	71.9
B_4	7.60	94.1	7.06	81.2
	7.00	86.6	7.04	79.9
	7.82	85.9	7.52	86.4
	7.80	88.8	7.70	76.4

a total of 32 observation vectors. The factors were rotational velocity [A_1 (fast) and A_2 (slow)] and lubricants (four types). The experimental units were 32 homogeneous pieces of bar steel. Two variables were measured on each piece of bar steel:

y_1 = ultimate torque,

y_2 = ultimate strain

We display the totals for each variable for use in computations. The numbers inside the box are cell totals (over the four replications), and the marginal totals are for each level of A and B :

Totals for y_1			Totals for y_2		
	A_1	A_2		A_1	A_2
B_1	30.61	29.08	59.69	354.0	327.9
B_2	32.61	31.34	64.12	344.7	310.0
B_3	31.79	29.45	61.24	352.9	305.0
B_4	30.22	29.32	59.54	355.4	323.9
	125.40	119.19	244.59	1407.0	1266.8
					2673.8

Using computational forms for h_{Arr} in (6.69), the (1, 1) element of \mathbf{H}_A (corresponding to y_1) is given by

$$h_{A11} = \frac{(125.40)^2 + (119.19)^2}{(4)(4)} - \frac{(244.59)^2}{(4)(4)(2)} = 1.205.$$

For the (2, 2) element of \mathbf{H}_A (corresponding to y_2), we have

$$h_{A22} = \frac{(1407.0)^2 + (1266.8)^2}{16} - \frac{(2673.8)^2}{32} = 614.25.$$

For the (1, 2) element of \mathbf{H}_A (corresponding to $y_1 y_2$), we use (6.70) for h_{Ars} to obtain

$$\begin{aligned} h_{A12} &= \frac{(125.40)(1407.0) + (119.19)(1266.8)}{16} - \frac{(244.59)(2673.8)}{32} \\ &= 27.208. \end{aligned}$$

Thus

$$\mathbf{H}_A = \begin{pmatrix} 1.205 & 27.208 \\ 27.208 & 614.251 \end{pmatrix}.$$

We obtain \mathbf{H}_B similarly:

$$\begin{aligned} h_{B11} &= \frac{(59.69)^2 + \dots + (59.54)^2}{(4)(2)} - \frac{(244.59)^2}{32} = 1.694, \\ h_{B22} &= \frac{(681.9)^2 + \dots + (679.3)^2}{8} - \frac{(2673.8)^2}{32} = 74.874, \\ h_{B12} &= \frac{(59.69)(681.9) + \dots + (59.54)(679.3)}{8} - \frac{(244.59)(2673.8)}{32} \\ &= -9.862, \\ \mathbf{H}_B &= \begin{pmatrix} 1.694 & -9.862 \\ -9.862 & 74.874 \end{pmatrix}. \end{aligned}$$

For \mathbf{H}_{AB} we have by (6.71)

$$\begin{aligned} h_{AB11} &= \frac{(30.61)^2 + \dots + (29.32)^2}{4} - \frac{(244.59)^2}{32} - 1.205 - 1.694 = .132, \\ h_{AB22} &= \frac{(354.0)^2 + \dots + (323.9)^2}{4} - \frac{(2673.8)^2}{32} - 614.25 - 74.874 \\ &= 32.244, \\ h_{AB12} &= \frac{(30.61)(354.0) + \dots + (29.32)(323.9)}{4} - \frac{(244.59)(2673.8)}{32} \\ &\quad - 27.208 - (-9.862) = 1.585, \\ \mathbf{H}_{AB} &= \begin{pmatrix} .132 & 1.585 \\ 1.585 & 32.244 \end{pmatrix}. \end{aligned}$$

The error matrix \mathbf{E} is obtained using the computational forms given for e_{rr} and e_{rs} in (6.72). For example, e_{11} and e_{12} are computed as

$$\begin{aligned} e_{11} &= (7.80)^2 + (7.10)^2 + \cdots + (7.70)^2 - \frac{(244.59)^2}{32} - 1.205 \\ &\quad - 1.694 - .132 = 4.897, \\ e_{12} &= (7.80)(90.4) + \cdots + (7.70)(76.4) - \frac{(244.59)(2673.8)}{32} - 27.208 \\ &\quad - (-9.862) - 1.585 = -1.890. \end{aligned}$$

Proceeding in this fashion, we obtain

$$\mathbf{E} = \begin{pmatrix} 4.897 & -1.890 \\ -1.890 & 736.390 \end{pmatrix},$$

with $\nu_E = ab(n - 1) = (2)(4)(4 - 1) = 24$.

To test the main effect of A with Wilks' Λ , we compute

$$\Lambda_A = \frac{|\mathbf{E}|}{|\mathbf{E} + \mathbf{H}_A|} = \frac{3602.2}{7600.2} = .474 < \Lambda_{.05,2,1,24} = .771,$$

and we conclude that velocity has a significant effect on y_1 or y_2 or both.

For the B main effect, we have

$$\Lambda_B = \frac{|\mathbf{E}|}{|\mathbf{E} + \mathbf{H}_B|} = \frac{3602.2}{5208.6} = .6916 > \Lambda_{.05,2,3,24} = .591.$$

We conclude that the effect of lubricants is not significant.

For the AB interaction, we obtain

$$\Lambda_{AB} = \frac{|\mathbf{E}|}{|\mathbf{E} + \mathbf{H}_{AB}|} = \frac{3602.2}{3865.3} = .932 > \Lambda_{.05,2,3,24} = .591.$$

Hence we conclude that the interaction effect is not significant.

We now obtain the other three MANOVA test statistics for each test. For A , the only nonzero eigenvalue of $\mathbf{E}^{-1}\mathbf{H}_A$ is 1.110. Thus

$$\begin{aligned} V^{(s)} &= \frac{\lambda_1}{1 + \lambda_1} = .526, \quad U^{(s)} = \lambda_1 = 1.110, \\ \theta &= \frac{\lambda_1}{1 + \lambda_1} = .526. \end{aligned}$$

In this case, all three tests give results equivalent to that of Λ_A because $\nu_H = s = 1$.

For B , $\nu_H = 3$ and $p = s = 2$. The eigenvalues of $\mathbf{E}^{-1}\mathbf{H}_B$ are .418 and .020, and we obtain

$$V^{(s)} = \sum_{i=1}^s \frac{\lambda_i}{1 + \lambda_i} = .314,$$

$$U^{(s)} = \sum_{i=1}^s \lambda_i = .438, \quad \frac{\nu_E}{\nu_H} U^{(s)} = 3.502,$$

$$\theta = \frac{\lambda_1}{1 + \lambda_1} = .295.$$

With $s = 2$, $m = 0$, and $N = 10.5$, we have $V^{(s)} = .439$ and $\theta_{.05} = .364$. The .05 critical value of $\nu_E U^{(s)} / \nu_H$ is 5.1799. Thus $V^{(s)}$, $U^{(s)}$, and θ lead to acceptance of H_0 , as does Λ . Of the four tests, θ appears to be closer to rejection. This is because $\lambda_1 / (\lambda_1 + \lambda_2) = .418 / (.418 + .020) = .954$, indicating that the mean vectors for factor B are essentially collinear, in which case Roy's test is more powerful. If the mean vectors $\bar{y}_{.1.}$, $\bar{y}_{.2.}$, $\bar{y}_{.3.}$, and $\bar{y}_{.4.}$ for the four levels of B were a little further apart, we would have a situation in which the four MANOVA tests do not lead to the same conclusion.

For AB , the eigenvalues of $\mathbf{E}^{-1}\mathbf{H}_{AB}$ are .0651 and .0075, from which

$$V^{(s)} = \sum_{i=1}^s \frac{\lambda_i}{1 + \lambda_i} = .0685, \quad U^{(s)} = .0726, \quad \frac{\nu_E}{\nu_H} U^{(s)} = .580,$$

$$\theta = \frac{\lambda_1}{1 + \lambda_1} = .0611.$$

The critical values remain the same as for factor B , and all three tests accept H_0 , as does Wilks' Λ . With a nonsignificant interaction, interpretation of the main effects is simplified. \square

6.6 OTHER MODELS

6.6.1 Higher-Order Fixed Effects

A higher-order (balanced) fixed-effects model or factorial experiment presents no new difficulties. As an illustration, consider a three-way classification with three factors A , B , and C and all interactions AB , AC , BC , and ABC . The observation vector \mathbf{y} has p variables as usual. The MANOVA model allowing for main effects and interactions can be written as

$$\mathbf{y}_{ijkl} = \mu + \alpha_i + \beta_j + \gamma_k + \delta_{ij} + \eta_{ik} + \tau_{jk} + \phi_{ijk} + \varepsilon_{ijkl}, \quad (6.75)$$

where, for example, α_i is the effect of the i th level of factor A on each of the p variables in \mathbf{y}_{ijkl} and δ_{ij} is the AB interaction effect on each of the p variables. Similarly, η_{ik} , τ_{jk} , and ϕ_{ijk} represent the AC , BC , and ABC interactions on each of the p variables.

The matrices of sums of squares and products for main effects, interactions, and error are defined in a manner similar to that for the matrices detailed for the two-way model in Section 6.5.2. The sum of squares (on the diagonal) for each variable is calculated exactly the same as in a univariate ANOVA for a three-way model. The sums of products (off-diagonal) are obtained analogously. Test construction parallels that for the two-way model, using the matrix for error to test all factors and interactions.

Degrees of freedom for each factor are the same as in the corresponding three-way univariate model. All four MANOVA test statistics can be computed for each test. Contrasts can be defined and tested in a manner similar to that in Section 6.5.2. Follow-up procedures on the individual variables (F -tests and discriminant functions) can be used as discussed for the one-way or two-way models in Sections 6.4 and 6.5.2.

6.6.2 Mixed Models

There is a MANOVA counterpart for every univariate ANOVA design. This applies to fixed, random, and mixed models and to experimental structures that are crossed, nested, or a combination. Roebrick (1982) has provided a formal proof that univariate mixed models can be generalized to multivariate mixed models. Schott and Saw (1984) have shown that for the one-way multivariate random-effects model, the approach leads to the same test statistics involving the eigenvalues of $\mathbf{E}^{-1}\mathbf{H}$ as in the fixed-effects model.

In the (balanced) MANOVA mixed model, the expected mean square matrices have exactly the same pattern as expected mean squares for the corresponding univariate ANOVA model. Thus a table of expected mean squares for the terms in the corresponding univariate model provides direction for choosing the appropriate error matrix to test each term in the MANOVA model. However, if the matrix indicated for "error" has fewer degrees of freedom than p , it will not have an inverse and the test cannot be made.

To illustrate, suppose we have a (balanced) two-way MANOVA model with A fixed and B random. Then the (univariate) expected mean squares (EMS) and Wilks' Λ -tests are as follows:

Source	EMS	Λ
A	$\sigma^2 + n\sigma_{AB}^2 + nb\sigma_A^{*2}$	$ \mathbf{H}_{AB} / \mathbf{H}_{AB} + \mathbf{H}_A $
B	$\sigma^2 + na\sigma_B^2$	$ \mathbf{E} / \mathbf{E} + \mathbf{H}_B $
AB	$\sigma^2 + n\sigma_{AB}^2$	$ \mathbf{E} / \mathbf{E} + \mathbf{H}_{AB} $
Error	σ^2	

In the expected mean square for factor A , we have used the notation σ_A^{*2} in place of $\sum_{i=1}^a \alpha_i^2/(a-1)$. The test for A using \mathbf{H}_{AB} for error matrix will be indeterminate (of the form 0/0) if $\nu_{AB} \leq p$, where $\nu_{AB} = (a-1)(b-1)$. In this case, ν_{AB} will often fail to exceed p . For example, suppose A has two levels and B has three. Then $\nu_{AB} = 2$, which will ordinarily be less than p . In such a case, we would have little recourse except to compute univariate tests on the p individual variables.

Table 6.7 Wilks' Λ Tests for a Typical Split-Plot Design

Source	df	Expected Mean Squares	Wilks' Λ
A	$a - 1$	$\sigma^2 + ce\sigma_B^2 + bce\sigma_A^{*2}$	$ \mathbf{H}_B / \mathbf{H}_A + \mathbf{H}_B $
B	$a(b - 1)$	$\sigma^2 + ce\sigma_B^2$	$ \mathbf{E} / \mathbf{H}_B + \mathbf{E} $
C	$c - 1$	$\sigma^2 + eo\sigma_{BC}^2 + abeo\sigma_C^{*2}$	$ \mathbf{H}_{BC} / \mathbf{H}_C + \mathbf{H}_{BC} $
AC	$(a - 1)(c - 1)$	$\sigma^2 + eo\sigma_{BC}^2 + beo\sigma_{AC}^{*2}$	$ \mathbf{H}_{BC} / \mathbf{H}_{AC} + \mathbf{H}_{BC} $
BC	$a(b - 1)(c - 1)$	$\sigma^2 + eo\sigma_{BC}^2$	$ \mathbf{E} / \mathbf{H}_{BC} + \mathbf{E} $
Error	$abc(e - 1)$	σ^2	

However, we would not have the multivariate test to protect against carrying out too many univariate tests and thereby inflating the experimentwise α (see Section 6.4). To protect against inflation of α when making p tests, we could use a Bonferroni correction, as in procedure 2 in Section 5.5. In the case of F -tests, we do not have a table of Bonferroni critical values, as we do for t -tests (Table A.8), but we can achieve an equivalent result by comparing the p -values for the F -tests against α/p instead of against α .

As another illustration, consider the analysis for a (balanced) multivariate split-plot design. For simplicity, we show the associated univariate model in place of the multivariate model. We use the factor names A, B, AC, \dots to indicate parameters in the model:

$$y_{ijkl} = \mu + A_i + B_{(i)j} + C_k + AC_{ik} + BC_{(i)jk} + \varepsilon_{(ijk)l},$$

where A and C are fixed and B is random. Nesting is indicated by bracketed subscripts; for example, B and BC are nested in A . Table 6.7 shows the expected mean squares and corresponding Wilks tests.

Since we use \mathbf{H}_B and \mathbf{H}_{BC} , as well as \mathbf{E} , to make tests, the following must hold:

$$a(b - 1) \geq p, \quad a(b - 1)(c - 1) \geq p, \quad abc(e - 1) \geq p.$$

To construct the other three MANOVA tests, we use eigenvalues of the following matrices:

Source	Matrix
A	$\mathbf{H}_B^{-1}\mathbf{H}_A$
B	$\mathbf{E}^{-1}\mathbf{H}_B$
C	$\mathbf{H}_{BC}^{-1}\mathbf{H}_C$
AC	$\mathbf{H}_{BC}^{-1}\mathbf{H}_{AC}$
BC	$\mathbf{E}^{-1}\mathbf{H}_{BC}$

With a table of expected mean squares, such as those in Table 6.7, it is a simple matter to determine the error matrix in each case. For a given factor or interaction, such as A, B , or AC , the appropriate error matrix is ordinarily the one whose expected mean square matches that of the given factor except for the last term. For

example, factor C , with expected mean square $\sigma^2 + e\sigma_{BC}^2 + abe\sigma_C^{*2}$, is tested by BC , whose expected mean square is $\sigma^2 + e\sigma_{BC}^2$. If $H_0: \sigma_C^{*2} = 0$ is true, then C and BC have the same expected mean square.

In some mixed and random models, certain terms have no available error term. When this happens in the univariate case, we can construct an approximate test using Satterthwaite's (1941) or other synthetic mean square approach. For a similar approach in the multivariate case, see Khuri, Mathew, and Nel (1994).

6.7 CHECKING ON THE ASSUMPTIONS

In Section 6.2 we discussed the robustness of the four MANOVA test statistics to nonnormality and heterogeneity of covariance matrices. The MANOVA tests (except for Roy's) are rather robust to these departures from the assumptions, although, in general, as dimensionality increases, robustness decreases.

Even though MANOVA procedures are fairly robust to departures from multivariate normality, we may want to check for gross violations of this assumption. Any of the tests or plots from Section 4.4 could be used. For a two-way design, for example, the tests could be applied separately to the n observations in each individual cell (if n is sufficiently large) or to all the residuals. The residual vectors after fitting the model $y_{ijk} = \mu_{ij} + \varepsilon_{ijk}$ would be

$$\hat{\varepsilon}_{ijk} = y_{ijk} - \bar{y}_{ij.}, \quad i = 1, 2, \dots, a, \quad j = 1, 2, \dots, b \quad k = 1, 2, \dots, n.$$

It is also advisable to check for outliers, which can lead to either a Type I or a Type II error. The tests of Section 4.6 can be run separately for each cell (for sufficiently large n) or for all of the abn residuals, $\hat{\varepsilon}_{ijk} = y_{ijk} - \bar{y}_{ij..}$.

A test of the equality of covariance matrices can be made using Box's M -test given in Section 7.3.2. Note the cautions expressed there about the sensitivity of this test to nonnormality and unequal sample sizes.

The assumption of independence of the observation vectors y_{ijk} is even more important than the assumptions of normality and equality of covariance matrices. We are referring, of course, to independence from one observation vector to another. The variables within a vector are assumed to be correlated, as usual. In the univariate case, Barcikowski (1981) showed that a moderate amount of dependence among the observations produces an actual α much greater than the nominal α . This effect is to be expected, since the dependence leads to an underestimate of the variance, so that MSE is reduced and the F -statistic is inflated. We can assume that this effect on error rates carries over to MANOVA.

In univariate ANOVA, a simple measure of dependence among the kn observations in a one-way model is the so-called *intraclass correlation*:

$$r_c = \frac{\text{MSB} - \text{MSE}}{\text{MSB} + (n-1)\text{MSE}}, \quad (6.76)$$

where MSB and MSE are the between and within mean squares for the variable and n is the number of observations per group. This could be calculated for each variable in a MANOVA to check for independence.

In many experimental settings, we do not anticipate a lack of independence. But in certain cases the observations are dependent. For example, if the sampling units are people, they may influence each other as they interact together. In some educational studies, researchers must use entire classrooms as sampling units rather than individual students. Another example of dependence is furnished by observations that are *serially correlated*, as in a *time series*, for example. Each observation depends to a certain extent on the preceding one, and its random movement is somewhat dampened as a result.

6.8 PROFILE ANALYSIS

The two-sample profile analysis of Section 5.9.2 can be extended to k groups. Again we assume that the variables are commensurate, as, for example, when each subject is given a battery of tests. Other assumptions, cautions, and comments expressed in Section 5.9.2 apply here as well.

The basic model is the balanced one-way MANOVA:

$$\mathbf{y}_{ij} = \boldsymbol{\mu}_i + \boldsymbol{\varepsilon}_{ij}, \quad i = 1, 2, \dots, k, \quad j = 1, 2, \dots, n.$$

To test $H_0: \boldsymbol{\mu}_1 = \boldsymbol{\mu}_2 = \dots = \boldsymbol{\mu}_k$, we use the usual \mathbf{H} and \mathbf{E} matrices given in (6.9) and (6.10). If the variables are commensurate, we can be more specific and extend H_0 to an examination of the k profiles obtained by plotting the p values $\mu_{i1}, \mu_{i2}, \dots, \mu_{ip}$ in each $\boldsymbol{\mu}_i$, as was done with two $\boldsymbol{\mu}_i$'s in Section 5.9.2. We are interested in the same three hypotheses as before:

- H_{01} : The k profiles are parallel.
- H_{02} : The k profiles are all at the same level.
- H_{03} : The k profiles are flat.

The hypothesis of parallelism for two groups was expressed in Section 5.9.2 as $H_{01}: \mathbf{C}\boldsymbol{\mu}_1 = \mathbf{C}\boldsymbol{\mu}_2$, where \mathbf{C} is any $(p - 1) \times p$ matrix of rank $p - 1$ such that $\mathbf{C}\mathbf{j} = \mathbf{0}$, for example,

$$\mathbf{C} = \begin{pmatrix} 1 & -1 & 0 & \cdots & 0 \\ 0 & 1 & -1 & \cdots & 0 \\ \vdots & \vdots & \vdots & & \vdots \\ 0 & 0 & 0 & \cdots & -1 \end{pmatrix}.$$

For k groups, the analogous hypothesis of parallelism is

$$H_{01}: \mathbf{C}\boldsymbol{\mu}_1 = \mathbf{C}\boldsymbol{\mu}_2 = \dots = \mathbf{C}\boldsymbol{\mu}_k. \quad (6.77)$$

The hypothesis (6.77) is equivalent to the hypothesis $H_0: \boldsymbol{\mu}_{z1} = \boldsymbol{\mu}_{z2} = \dots = \boldsymbol{\mu}_{zk}$ in a one-way MANOVA on the transformed variables $\mathbf{z}_{ij} = \mathbf{C}\mathbf{y}_{ij}$. Since \mathbf{C} has $p - 1$ rows, $\mathbf{C}\mathbf{y}_{ij}$ is $(p - 1) \times 1$, $\mathbf{C}\boldsymbol{\mu}_i$ is $(p - 1) \times 1$, and $\mathbf{C}\Sigma\mathbf{C}'$ is $(p - 1) \times (p - 1)$. By property 1b in Section 4.2, \mathbf{z}_{ij} is distributed as $N_{p-1}(\mathbf{C}\boldsymbol{\mu}_i, \mathbf{C}\Sigma\mathbf{C}')$.

By analogy with (3.64), the hypothesis and error matrices for testing H_{01} in (6.77) are

$$\mathbf{H}_z = \mathbf{CHC}', \quad \mathbf{E}_z = \mathbf{CEC}'.$$

We thus have

$$\Lambda = \frac{|\mathbf{CEC}'|}{|\mathbf{CEC}' + \mathbf{CHC}'|} = \frac{|\mathbf{CEC}'|}{|\mathbf{C}(\mathbf{E} + \mathbf{H})\mathbf{C}'|}, \quad (6.78)$$

which is distributed as $\Lambda_{p-1, \nu_H, \nu_E}$, where $\nu_H = k - 1$ and $\nu_E = k(n - 1)$. The other three MANOVA test statistics can be obtained from the eigenvalues of $(\mathbf{CEC}')^{-1}(\mathbf{CHC}')$. The test for H_{01} can easily be adjusted for unbalanced data, as in Section 6.1.6. We would calculate \mathbf{H} and \mathbf{E} by (6.31) and (6.32) and use $\nu_E = \sum_i n_i - k$.

The hypothesis that two profiles are at the same level is $H_{02}: \mathbf{j}'\boldsymbol{\mu}_1 = \mathbf{j}'\boldsymbol{\mu}_2$ (see Section 5.9.2), which generalizes immediately to k profiles at the same level,

$$H_{02}: \mathbf{j}'\boldsymbol{\mu}_1 = \mathbf{j}'\boldsymbol{\mu}_2 = \cdots = \mathbf{j}'\boldsymbol{\mu}_k. \quad (6.79)$$

For two groups we used a univariate t , as defined in (5.36), to test H_{02} . Similarly, for k groups we can employ an F -test for one-way ANOVA comparing k groups with observations $\mathbf{j}'\mathbf{y}_{ij}$. Alternatively, we can utilize (6.78) with $\mathbf{C} = \mathbf{j}'$,

$$\Lambda = \frac{\mathbf{j}'\mathbf{E}\mathbf{j}}{\mathbf{j}'\mathbf{E}\mathbf{j} + \mathbf{j}'\mathbf{H}\mathbf{j}}, \quad (6.80)$$

which is distributed as $\Lambda_{1, \nu_H, \nu_E}$ ($p = 1$ because $\mathbf{j}'\mathbf{y}_{ij}$ is a scalar). This is, of course, equivalent to the F -test on $\mathbf{j}'\mathbf{y}_{ij}$, since by Table 6.1 in Section 6.1.3,

$$F = \frac{1 - \Lambda}{\Lambda} \frac{\nu_E}{\nu_H} \quad (6.81)$$

is distributed as F_{ν_H, ν_E} .

The third hypothesis, that of “flatness,” essentially states that the average of the k group means is the same for each variable [see (5.37)]:

$$\begin{aligned} H_{03}: \frac{\mu_{11} + \mu_{21} + \cdots + \mu_{k1}}{k} &= \frac{\mu_{12} + \mu_{22} + \cdots + \mu_{k2}}{k} \\ &= \cdots = \frac{\mu_{1p} + \mu_{2p} + \cdots + \mu_{kp}}{k}, \end{aligned}$$

or by analogy with (5.38),

$$H_{03}: \frac{\mathbf{C}(\boldsymbol{\mu}_1 + \boldsymbol{\mu}_2 + \cdots + \boldsymbol{\mu}_k)}{k} = \mathbf{0}, \quad (6.82)$$

where \mathbf{C} is a $(p - 1) \times p$ matrix of rank $p - 1$ such that $\mathbf{C}\mathbf{j} = \mathbf{0}$ [see (6.77)]. If H_{01} is true, the flatness hypothesis can also be stated as, the means of all p variables in each group are the same, or $\mu_{i1} = \mu_{i2} = \cdots = \mu_{ip}, i = 1, 2, \dots, k$. This can be expressed as $H_{03}: \mathbf{C}\boldsymbol{\mu}_1 = \mathbf{C}\boldsymbol{\mu}_2 = \cdots = \mathbf{C}\boldsymbol{\mu}_k = \mathbf{0}$.

Table 6.8 Weight of Guinea Pigs Under 3 Levels of Vitamin E Supplements

Group	Animal	Week 1	Week 3	Week 4	Week 5	Week 6	Week 7
1	1	455	460	510	504	436	466
1	2	467	565	610	596	542	587
1	3	445	530	580	597	582	619
1	4	485	542	594	583	611	612
1	5	480	500	550	528	562	576
2	6	514	560	565	524	552	597
2	7	440	480	536	484	567	569
2	8	495	570	569	585	576	677
2	9	520	590	610	637	671	702
2	10	503	555	591	605	649	675
3	11	496	560	622	622	632	670
3	12	498	540	589	557	568	609
3	13	478	510	568	555	576	605
3	14	545	565	580	601	633	649
3	15	472	498	540	524	532	583

To test H_{03} as given by (6.82), we can extend the T^2 -test in (5.39). The grand mean vector $(\mu_1 + \mu_2 + \dots + \mu_k)/k$ in (6.82) can be estimated as in Section 6.1.2 by

$$\bar{\mathbf{y}}_{..} = \sum_{ij} \frac{\mathbf{y}_{ij}}{kn}.$$

Under H_{03} , $\mathbf{C}\bar{\mathbf{y}}_{..}$ is $N_{p-1}(\mathbf{0}, \mathbf{C}\Sigma\mathbf{C}'/kn)$, and H_{03} can be tested by

$$T^2 = kn(\mathbf{C}\bar{\mathbf{y}}_{..})'(\mathbf{C}\mathbf{E}\mathbf{C}'/\nu_E)^{-1}\mathbf{C}\bar{\mathbf{y}}_{..}, \quad (6.83)$$

where \mathbf{E}/ν_E is an estimate of Σ . As in the two-sample case, H_{03} is unaffected by the status of H_{02} . When H_{03} is true, T^2 in (6.83) is distributed as T_{p-1, ν_E}^2 .

■ EXAMPLE 6.8

Three vitamin E diet supplements with levels zero, low, and high were compared for their effect on growth of guinea pigs (Crowder and Hand 1990, pp. 21–29). Five guinea pigs received each supplement level, and their weights were recorded at the end of weeks 1, 3, 4, 5, 6, and 7. These weights are given in Table 6.8.

The three mean vectors are

$$\bar{\mathbf{y}}'_{1.} = (466.4, 519.4, 568.8, 561.6, 546.6, 572.0)$$

$$\bar{\mathbf{y}}'_{2.} = (494.4, 551.0, 574.2, 567.0, 603.0, 644.0)$$

$$\bar{\mathbf{y}}'_{3.} = (497.8, 534.6, 579.8, 571.8, 588.2, 623.2)$$

and the overall mean vector is

$$\bar{\mathbf{y}}'_{..} = (486.2, 535.0, 574.3, 566.8, 579.3, 613.1).$$




Figure 6.3 Profile of the three groups for the guinea pig data of Table 6.8.

A profile plot of the means \bar{y}_1 , \bar{y}_2 , and \bar{y}_3 is given in Figure 6.3. There is a high degree of parallelism in the three profiles, with the possible exception of week 6 for group 1.

The **E** and **H** matrices are as follows:

$$\mathbf{E} = \begin{pmatrix} 8481.2 & 8538.8 & 4819.8 & 8513.6 & 8710.0 & 8468.2 \\ 8538.8 & 17170.4 & 13293.0 & 19476.4 & 17034.2 & 20035.4 \\ 4819.8 & 13293.0 & 12992.4 & 17077.4 & 17287.8 & 17697.2 \\ 8513.6 & 19476.4 & 17077.4 & 28906.0 & 26226.4 & 28625.2 \\ 8710.0 & 17034.2 & 17287.8 & 26226.4 & 36898.0 & 31505.8 \\ 8468.2 & 20035.4 & 17697.2 & 28625.2 & 31505.8 & 33538.8 \end{pmatrix},$$

$$\mathbf{H} = \begin{pmatrix} 2969.2 & 2177.2 & 859.4 & 813.0 & 4725.2 & 5921.6 \\ 2177.2 & 2497.6 & 410.0 & 411.6 & 4428.8 & 5657.6 \\ 859.4 & 410.0 & 302.5 & 280.4 & 1132.1 & 1392.5 \\ 813.0 & 411.6 & 280.4 & 260.4 & 1096.4 & 1352.0 \\ 4725.2 & 4428.8 & 1132.1 & 1096.4 & 8550.9 & 10830.9 \\ 5921.6 & 5657.6 & 1392.5 & 1352.0 & 10830.9 & 13730.1 \end{pmatrix}.$$

Using

$$\mathbf{C} = \begin{pmatrix} 1 & -1 & 0 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 & 0 & 0 \\ 0 & 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 0 & 1 & -1 \end{pmatrix}$$

in the test statistic (6.78), we have, as a test for parallelism,

$$\begin{aligned}\Lambda &= \frac{|\mathbf{CEC}'|}{|\mathbf{C}(\mathbf{E} + \mathbf{H})\mathbf{C}'|} = \frac{3.8238 \times 10^{18}}{2.1355 \times 10^{19}} \\ &= .1791 > \Lambda_{.05,5,2,12} = .153.\end{aligned}$$

Thus we do not reject the parallelism hypothesis.

To test the hypothesis that the three profiles are at the same level, we use (6.80),

$$\begin{aligned}\Lambda &= \frac{\mathbf{j}'\mathbf{Ej}}{\mathbf{j}'\mathbf{Ej} + \mathbf{j}'\mathbf{Hj}} = \frac{632,605.2}{632,605.2 + 111,288.1} \\ &= .8504 > \Lambda_{.05,1,2,12} = .607.\end{aligned}$$

Hence we do not reject the levels hypothesis. This can also be seen by using (6.81) to transform Λ to F ,

$$F = \frac{(1 - \Lambda)\nu_E}{\Lambda\nu_H} = \frac{(1 - .8504)12}{(.8504)2} = 1.0555,$$

Which is clearly nonsignificant ($p = .378$).

To test the flatness hypothesis, we use (6.83):

$$\begin{aligned}T^2 &= kn(\mathbf{C}\bar{\mathbf{y}}_{..})'(\mathbf{CEC}'/\nu_E)^{-1}\mathbf{C}\bar{\mathbf{y}}_{..} \\ &= 15 \begin{pmatrix} -48.80 \\ -39.27 \\ 7.47 \\ -12.47 \\ -33.80 \end{pmatrix}' \begin{pmatrix} 714.5 & -13.2 & 207.5 & -219.9 & 270.2 \\ -13.2 & 298.1 & -174.9 & 221.0 & -216.0 \\ 207.5 & -174.9 & 645.3 & -240.8 & 165.8 \\ -219.9 & 221.0 & -240.8 & 1112.6 & -649.2 \\ 270.2 & -216.0 & 165.8 & -649.2 & 618.8 \end{pmatrix}^{-1} \\ &\quad \times \begin{pmatrix} -48.80 \\ -39.27 \\ 7.47 \\ -12.47 \\ -33.80 \end{pmatrix} \\ &= 297.13 > T^2_{.01,5,12} = 49.739.\end{aligned}$$

Thus only the flatness hypothesis is rejected in this case. \square

6.9 REPEATED MEASURES DESIGNS

6.9.1 Multivariate Versus Univariate Approach

In *repeated measures* designs, the research unit is typically a human or animal subject. Each subject is measured under several treatments or at different points of time.

Table 6.9 Data Layout for k -Groups Repeated Measures Experiment

Factor B (Group)	Subjects	Factor A (Repeated Measures)				
		A_1	A_2	...	A_p	
B_1	S_{11}	$(y_{111} \quad y_{112} \quad \cdots \quad y_{11p})$				$= \mathbf{y}'_{11}$
	S_{12}	$(y_{121} \quad y_{122} \quad \cdots \quad y_{12p})$				$= \mathbf{y}'_{12}$
	\vdots	\vdots	\vdots			\vdots
	S_{1n}	$(y_{1n1} \quad y_{1n2} \quad \cdots \quad y_{1np})$				$= \mathbf{y}'_{1n}$
B_2	S_{21}	$(y_{211} \quad y_{212} \quad \cdots \quad y_{21p})$				$= \mathbf{y}'_{21}$
	S_{22}	$(y_{221} \quad y_{222} \quad \cdots \quad y_{22p})$				$= \mathbf{y}'_{22}$
	\vdots	\vdots	\vdots			\vdots
	S_{2n}	$(y_{2n1} \quad y_{2n2} \quad \cdots \quad y_{2np})$				$= \mathbf{y}'_{2n}$
\vdots	\vdots	\vdots	\vdots			\vdots
B_k	S_{k1}	$(y_{k11} \quad y_{k12} \quad \cdots \quad y_{k1p})$				$= \mathbf{y}'_{k1}$
	S_{k2}	$(y_{k21} \quad y_{k22} \quad \cdots \quad y_{k2p})$				$= \mathbf{y}'_{k2}$
	\vdots	\vdots	\vdots			\vdots
	S_{kn}	$(y_{kn1} \quad y_{kn2} \quad \cdots \quad y_{knp})$				$= \mathbf{y}'_{kn}$

The treatments might be tests, drug levels, various kinds of stimuli, and so on. If the treatments are such that the order of presentation to the various subjects can be varied, then the order should be randomized to avoid an ordering bias. If subjects are measured at successive time points, it may be of interest to determine the degree of polynomial required to fit the curve. This is treated in Section 6.10 as part of an analysis of growth curves.

When comparing means of the treatments applied to each subject, we are examining the *within-subjects* factor. There will also be a *between-subjects* factor if there are several groups of subjects that we wish to compare. In Sections 6.9.2–6.9.6, we consider designs up to a complexity level of two within-subjects factors and two between-subjects factors.

We now discuss univariate and multivariate approaches to hypothesis testing in repeated measures designs. As a framework for this discussion, consider the layout in Table 6.9 for a repeated measures design with one repeated measures (within-subjects) factor, A , and one grouping (between-subjects) factor, B .

This design has often been analyzed as a univariate mixed-model nested design, also called a split-plot design, with subjects nested in factor B (whole-plot), which is crossed with factor A (repeated measures or split-plot). The univariate model for each y_{ijr} would be

$$y_{ijr} = \mu + B_i + S_{(i)j} + A_r + BA_{ir} + \varepsilon_{ijr}, \quad (6.84)$$

where the factor level designations (B , S , A , and BA) from (6.84) and Table 6.9 are used as parameter values, and the subscript $(i)j$ on S indicates that subjects are nested in factor B . In Table 6.9, the observations y_{ijr} for $r = 1, 2, \dots, p$ are enclosed in parentheses and denoted by \mathbf{y}'_{ij} to emphasize that these p variables are measured on one subject and thus constitute a vector of correlated variables. The ranges of the subscripts can be seen in Table 6.9: $i = 1, 2, \dots, k$; $j = 1, 2, \dots, n$; and $r = 1, 2, \dots, p$. With factors A and B fixed and subjects random, the univariate ANOVA is given in Table 6.10.

However, our initial reaction would be to rule out the univariate ANOVA because the y 's in each row are correlated and the assumption of independence is critical, as noted in Section 6.7. We will discuss below some assumptions under which the univariate analysis would be appropriate.

In the multivariate approach, the p responses $y_{ij1}, y_{ij2}, \dots, y_{ijp}$ (repeated measures) for subject S_{ij} constitute a vector \mathbf{y}_{ij} , as shown in Table 6.9. The multivariate model for \mathbf{y}_{ij} is a simple one-way MANOVA model,

$$\mathbf{y}_{ij} = \boldsymbol{\mu} + \boldsymbol{\beta}_i + \boldsymbol{\varepsilon}_{ij}, \quad (6.85)$$

where $\boldsymbol{\beta}_i$ is a vector of p main effects (corresponding to the p variables in \mathbf{y}_{ij}) for factor B , and $\boldsymbol{\varepsilon}_{ij}$ is an error vector for subject S_{ij} . This model seems to include only factor B , but we show in Section 6.9.3 how to use an approach similar to profile analysis in Section 6.8 to obtain tests on factor A and the BA interaction. The MANOVA assumption that $\text{cov}(\mathbf{y}_{ij}) = \boldsymbol{\Sigma}$ for all i and j allows the p repeated measures to be correlated in any pattern, since $\boldsymbol{\Sigma}$ is completely general. On the other hand, the ANOVA assumptions of independence and homogeneity of variances can be expressed as $\text{cov}(\mathbf{y}_{ij}) = \sigma^2 \mathbf{I}$. We would be very surprised if repeated measurements on the same subject were independent.

The univariate ANOVA approach has been found to be appropriate under less stringent conditions than $\boldsymbol{\Sigma} = \sigma^2 \mathbf{I}$. Wilks (1946) showed that the ordinary F -tests of ANOVA remain valid for a covariance structure of the form

$$\begin{aligned} \text{cov}(\mathbf{y}_{ij}) &= \boldsymbol{\Sigma} = \sigma^2 \begin{pmatrix} 1 & \rho & \rho & \cdots & \rho \\ \rho & 1 & \rho & \cdots & \rho \\ \vdots & \vdots & \vdots & & \vdots \\ \rho & \rho & \rho & \cdots & 1 \end{pmatrix} \\ &= \sigma^2[(1 - \rho)\mathbf{I} + \rho\mathbf{J}], \end{aligned} \quad (6.86)$$

where \mathbf{J} is a square matrix of 1's, as defined in (2.12) [see Rencher and Schaalje (2008, pp. 166–167)]. The covariance pattern (6.86) is variously known as *uniformity*, *compound symmetry*, or the *intraclass correlation* model. It allows for the variables to be correlated but restricts every variable to have the same variance and every pair of variables to have the same covariance. In a carefully designed experiment with appropriate randomization, this assumption may hold under the hypothesis of no A effect. Alternatively, we could use a test of the hypothesis that $\boldsymbol{\Sigma}$ has the pattern (6.86) (see Section 7.2.3). If this hypothesis is accepted, one could proceed with the usual ANOVA F -tests.

Table 6.10 Univariate ANOVA for Data Layout in Table 6.9

Source	df	MS	F
B (between)	$k - 1$	MSB	MSB/MSS
S (subjects)	$k(n - 1)$	MSS	
A (within or repeated)	$p - 1$	MSA	MSA/MSE
BA	$(k - 1)(p - 1)$	MSBA	MSBA/MSE
Error (SA interaction)	$k(n - 1)(p - 1)$	MSE	

Bock (1963) and Huynh and Feldt (1970) showed that the most general condition under which univariate F -tests remain valid is that

$$\mathbf{C}\Sigma\mathbf{C}' = \sigma^2\mathbf{I}, \quad (6.87)$$

where \mathbf{C} is a $(p - 1) \times p$ matrix whose rows are *orthonormal* contrasts (orthogonal contrasts that have been normalized to unit length). We can construct \mathbf{C} by choosing any $p - 1$ orthogonal contrasts among the means $\mu_1, \mu_2, \dots, \mu_p$ of the repeated measures factor and dividing each contrast by $\sqrt{\sum_{r=1}^p c_r^2}$. (This matrix \mathbf{C} is different from \mathbf{C} used in Section 6.8 and in the remainder of Section 6.9, whose rows are contrasts that are not normalized to unit length.) It can be shown that (6.86) is a special case of (6.87). The condition (6.87) is sometimes referred to as *sphericity*, although this term can also refer to the covariance pattern $\Sigma = \sigma^2\mathbf{I}$ on the untransformed \mathbf{y}_{ij} (see Section 7.2.2).

A simple way to test the hypothesis that (6.87) holds is to transform the data by $\mathbf{z}_{ij} = \mathbf{C}\mathbf{y}_{ij}$ and test $H_0: \Sigma_z = \sigma^2\mathbf{I}$, as in Section 7.2.2, using $\mathbf{CS}_{p1}\mathbf{C}'$ in place of $\mathbf{S}_{p1} = \mathbf{E}/\nu_E$.

Thus one procedure for repeated measures designs is to make a preliminary test for (6.86) or (6.87) and, if the hypothesis is accepted, use univariate F -tests, as in Table 6.10. Fehlberg (1980) investigated the use of larger α -values with a preliminary test of structure of the covariance matrix, as in (6.87). He concludes that using $\alpha = .40$ sufficiently controls the problem of falsely accepting sphericity so as to justify the use of a preliminary test.

If the univariate test for the repeated measures factor A is appropriate, it is more powerful because it has more degrees of freedom for error than the corresponding multivariate test. However, even mild departures from (6.87) seriously inflate the Type 1 error rate of the univariate test for factor A (Box 1954, Davidson 1972, Boik 1981). Because such departures can be easily missed in a preliminary test, Boik (1981) concludes that “on the whole, the ordinary F tests have nothing to recommend them” (p. 248) and emphasized that “there is no justification for employing ordinary univariate F tests for repeated measures treatment contrasts” (p. 254).

Another approach to analysis of repeated measures designs is to adjust the univariate F -test for the amount of departure from sphericity. Box (1954) and Greenhouse and Geisser (1959) showed that when $\Sigma \neq \sigma^2\mathbf{I}$, an approximate F -test for effects involving the repeated measures is obtained by reducing the degrees of freedom for

Table 6.11 Data Layout for a Single-Sample Repeated Measures Design

Subjects	Factor A (Repeated Measures)				
	A ₁	A ₂	A ₃	A ₄	
S ₁	(y ₁₁	y ₁₂	y ₁₃	y ₁₄)	= \mathbf{y}'_1
S ₂	(y ₂₁	y ₂₂	y ₂₃	y ₂₄)	= \mathbf{y}'_2
:	:	:	:	:	:
S _n	(y _{n1}	y _{n2}	y _{n3}	y _{n4})	= \mathbf{y}'_n

both numerator and denominator by a factor of ε

$$\varepsilon = \frac{[\text{tr}(\Sigma - \mathbf{J}\Sigma/p)]^2}{(p-1)\text{tr}(\Sigma - \mathbf{J}\Sigma/p)^2}, \quad (6.88)$$

where \mathbf{J} is a $p \times p$ matrix of 1's defined in (2.12). For example, in Table 6.10 the F -value for the BA interaction would be compared to F_α with $\varepsilon(k-1)(p-1)$ and $\varepsilon k(n-1)(p-1)$ degrees of freedom. An estimate $\hat{\varepsilon}$ can be obtained by substituting $\hat{\Sigma} = \mathbf{E}/\nu_E$ in (6.88). Greenhouse and Geisser (1959) showed that ε and $\hat{\varepsilon}$ vary between $1/(p-1)$ and 1, with $\varepsilon = 1$ when sphericity holds and $\varepsilon \geq 1/(p-1)$ for other values of Σ . Thus ε is a measure of nonsphericity. For a conservative test, Greenhouse and Geisser recommend dividing numerator and denominator degrees of freedom by $p-1$. Huynh and Feldt (1976) provided an improved estimator of ε .

The behavior of the approximate univariate F -test with degrees of freedom adjusted by $\hat{\varepsilon}$ has been investigated by Collier et al. (1967), Huynh (1978), Davidson (1972), Rogan et al. (1979), and Maxwell and Avery (1982). In these studies, the true α level turned out to be close to the nominal α , and the power was close to that of the multivariate test. However, since the ε -adjusted F -test is only approximate and has no power advantage over the exact multivariate test, there appears to be no compelling reason to use it. The only case in which we need to fall back on a univariate test is when there are insufficient degrees of freedom to perform a multivariate test, that is, when $p > \nu_E$.

In Sections 6.9.2–6.9.6, we discuss the multivariate approach to repeated measures. We will cover several models, beginning with the simple one-sample design.

6.9.2 One-Sample Repeated Measures Model

We illustrate some of the procedures in this section with $p = 4$. A one-sample design with four repeated measures on n subjects would appear as in Table 6.11. There is a superficial resemblance to a univariate randomized block design. However, in the repeated measures design, the observations y_{i1}, y_{i2}, y_{i3} , and y_{i4} are correlated because they are measured on the same subject (experimental unit), whereas in a randomized block design y_{i1}, y_{i2}, y_{i3} , and y_{i4} would be measured on four different experimental units. Thus we have a single sample of n observation vectors $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n$.

To test for significance of factor A , we compare the means of the four variables in \mathbf{y}_i ,

$$E(\mathbf{y}_i) = \boldsymbol{\mu} = \begin{pmatrix} \mu_1 \\ \mu_2 \\ \mu_3 \\ \mu_4 \end{pmatrix}.$$

The hypothesis is $H_0: \mu_1 = \mu_2 = \mu_3 = \mu_4$, which can be reexpressed as $H_0: \mu_1 - \mu_2 = \mu_2 - \mu_3 = \mu_3 - \mu_4 = 0$ or $\mathbf{C}_1\boldsymbol{\mu} = \mathbf{0}$, where

$$\mathbf{C}_1 = \begin{pmatrix} 1 & -1 & 0 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 1 & -1 \end{pmatrix}.$$

To test $H_0: \mathbf{C}_1\boldsymbol{\mu} = \mathbf{0}$ for a general p (p repeated measures on n subjects), we calculate $\bar{\mathbf{y}}$ and \mathbf{S} from $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n$ and extend \mathbf{C}_1 to $p - 1$ rows. Then when H_0 is true, $\mathbf{C}_1\bar{\mathbf{y}}$ is $N_{p-1}(\mathbf{0}, \mathbf{C}_1\boldsymbol{\Sigma}\mathbf{C}'_1/n)$, and

$$T^2 = n(\mathbf{C}_1\bar{\mathbf{y}})'(\mathbf{C}_1\mathbf{S}\mathbf{C}'_1)^{-1}(\mathbf{C}_1\bar{\mathbf{y}}) \quad (6.89)$$

is distributed as $T^2_{p-1,n-1}$. We reject $H_0: \mathbf{C}_1\boldsymbol{\mu} = \mathbf{0}$ if $T^2 \geq T^2_{\alpha,p-1,n-1}$. Note that the dimension is $p - 1$ because $\mathbf{C}_1\bar{\mathbf{y}}$ is $(p - 1) \times 1$ [see (5.33)].

The multivariate approach involving transformed observations $\mathbf{z}_i = \mathbf{C}_1\mathbf{y}_i$ was first suggested by Hsu (1938) and has been discussed further by Williams (1970) and Morrison (1972). Note that in $\mathbf{C}_1\bar{\mathbf{y}}$ (for $p = 4$), we work with contrasts on the elements $\bar{y}_1, \bar{y}_2, \bar{y}_3$, and \bar{y}_4 within the vector

$$\bar{\mathbf{y}} = \begin{pmatrix} \bar{y}_1 \\ \bar{y}_2 \\ \bar{y}_3 \\ \bar{y}_4 \end{pmatrix},$$

as opposed to the contrasts involving comparisons of several mean vectors themselves, as, for example, in Section 6.3.2.

The hypothesis $H_0: \mu_1 = \mu_2 = \mu_3 = \mu_4$ can also be expressed as $H_0: \mu_1 - \mu_4 = \mu_2 - \mu_4 = \mu_3 - \mu_4 = 0$, or $\mathbf{C}_2\boldsymbol{\mu} = \mathbf{0}$, where

$$\mathbf{C}_2 = \begin{pmatrix} 1 & 0 & 0 & -1 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & -1 \end{pmatrix}.$$

The matrix \mathbf{C}_1 can be obtained from \mathbf{C}_2 by simple row operations, for example, subtracting the second row from the first and the third row from the second. Hence, $\mathbf{C}_1 = \mathbf{AC}_2$, where

$$\mathbf{A} = \begin{pmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{pmatrix}.$$

In fact, $H_0 : \mu_1 = \mu_2 = \cdots = \mu_p$ can be expressed as $\mathbf{C}\boldsymbol{\mu} = \mathbf{0}$ for any full-rank $(p - 1) \times p$ matrix \mathbf{C} such that $\mathbf{C}\mathbf{j} = \mathbf{0}$, and the same value of T^2 in (6.89) will result. The contrasts in \mathbf{C} can be either linearly independent or orthogonal.

The hypothesis $H_0 : \mu_1 = \mu_2 = \cdots = \mu_p = \mu$, say, can also be expressed as

$$H_0 : \boldsymbol{\mu} = \mu\mathbf{j},$$

where $\mathbf{j} = (1, 1, \dots, 1)'$. The maximum likelihood estimate of μ is

$$\hat{\mu} = \frac{\bar{\mathbf{y}}' \mathbf{S}^{-1} \mathbf{j}}{\mathbf{j}' \mathbf{S}^{-1} \mathbf{j}}. \quad (6.90)$$

The likelihood ratio test of H_0 is a function of

$$\bar{\mathbf{y}}' \mathbf{S}^{-1} \bar{\mathbf{y}} - \frac{(\bar{\mathbf{y}}' \mathbf{S}^{-1} \mathbf{j})^2}{\mathbf{j}' \mathbf{S}^{-1} \mathbf{j}}.$$

Williams (1970) showed that for any $(p - 1) \times p$ matrix \mathbf{C} of rank $p - 1$ such that $\mathbf{C}\mathbf{j} = \mathbf{0}$,

$$\bar{\mathbf{y}}' \mathbf{S}^{-1} \bar{\mathbf{y}} - \frac{(\bar{\mathbf{y}}' \mathbf{S}^{-1} \mathbf{j})^2}{\mathbf{j}' \mathbf{S}^{-1} \mathbf{j}} = (\mathbf{C}\bar{\mathbf{y}})' (\mathbf{C}\mathbf{S}\mathbf{C}')^{-1} (\mathbf{C}\bar{\mathbf{y}}),$$

and thus the T^2 -test in (6.89) is equivalent to the likelihood ratio test.

■ EXAMPLE 6.9.2

The data in Table 6.12 were given by Cochran and Cox (1957, p. 130). As rearranged by Timm (1980), the observations constitute a one-sample repeated measures design with two within-subjects factors. Factor A is a comparison of two tasks; factor B is a comparison of two types of calculators. The measurements are speed of calculation.

To test the hypothesis $H_0 : \mu_1 = \mu_2 = \mu_3 = \mu_4$, we use the contrast matrix

$$\mathbf{C} = \begin{pmatrix} 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & -1 & 1 \end{pmatrix},$$

where the first row compares the two levels of A , the second row compares the two levels of B , and the third row corresponds to the AB interaction. From the five observation vectors in Table 6.12, we obtain

$$\bar{\mathbf{y}} = \begin{pmatrix} 23.2 \\ 15.6 \\ 20.0 \\ 11.6 \end{pmatrix}, \quad \mathbf{S} = \begin{pmatrix} 51.7 & 29.8 & 9.2 & 7.4 \\ 29.8 & 46.8 & 16.2 & -8.7 \\ 9.2 & 16.2 & 8.5 & -10.5 \\ 7.4 & -8.7 & -10.5 & 24.3 \end{pmatrix}.$$

For the overall test of equality of means, we have, by (6.89),

$$T^2 = n(\mathbf{C}\bar{\mathbf{y}})' (\mathbf{C}\mathbf{S}\mathbf{C}')^{-1} (\mathbf{C}\bar{\mathbf{y}}) = 29.736 < T^2_{.05, 3, 4} = 114.986.$$

Table 6.12 Calculator Speed Data

Subjects	A ₁		A ₂	
	B ₁	B ₂	B ₁	B ₂
S ₁	30	21	21	14
S ₂	22	13	22	5
S ₃	29	13	18	17
S ₄	12	7	16	14
S ₅	23	24	23	8

Since the T^2 -test is not significant, we would ordinarily not proceed with tests based on the individual rows of \mathbf{C} . We will do so, however, for illustrative purposes. (Note that the T^2 -test has very low power in this case, because $n - 1 = 4$ is very small.)

To test A , B , and AB , we test each row of \mathbf{C} , in which case

$$T^2 = n(\mathbf{c}'_i \bar{\mathbf{y}})' (\mathbf{c}'_i \mathbf{S} \mathbf{c}_i)^{-1} \mathbf{c}'_i \bar{\mathbf{y}}$$

is the square of the t -statistic

$$t_i = \frac{\sqrt{n} \mathbf{c}'_i \bar{\mathbf{y}}}{\sqrt{\mathbf{c}'_i \mathbf{S} \mathbf{c}_i}}, \quad i = 1, 2, 3,$$

where \mathbf{c}'_i is the i th row of \mathbf{C} .

The three results are as follows:

$$\text{Factor } A \quad t_1 = 1.459 < t_{.025,4} = 2.776,$$

$$\text{Factor } B \quad t_2 = 5.247 > t_{.005,4} = 4.604,$$

$$\text{Interaction } AB \quad t_3 = -.152$$

Thus only the main effect for B is significant. Note that in all but one case in Table 6.12, the value for B_1 is greater than that for B_2 . \square

6.9.3 *k*-Sample Repeated Measures Model

We turn now to the k -sample repeated measures design depicted in Table 6.9. As noted in Section 6.9.1, the multivariate approach to this repeated measures design uses the one-way MANOVA model $\mathbf{y}_{ij} = \boldsymbol{\mu} + \boldsymbol{\beta}_i + \boldsymbol{\varepsilon}_{ij} = \boldsymbol{\mu}_i + \boldsymbol{\varepsilon}_{ij}$. From the k groups of n observation vectors each, we calculate $\bar{\mathbf{y}}_1, \bar{\mathbf{y}}_2, \dots, \bar{\mathbf{y}}_k$ and the error matrix \mathbf{E} .

The layout in Table 6.9 is similar to that of a k -sample profile analysis in Section 6.8. To test (the within-subjects) factor A , we need to compare the means of the variables y_1, y_2, \dots, y_p within \mathbf{y} averaged across the levels of factor B . The p variables correspond to the levels of factor A . In the model $\mathbf{y}_{ij} = \boldsymbol{\mu}_i + \boldsymbol{\varepsilon}_{ij}$, the

mean vectors $\mu_1, \mu_2, \dots, \mu_k$ correspond to the levels of factor B and are estimated by $\bar{y}_{1..}, \bar{y}_{2..}, \dots, \bar{y}_{k..}$. To compare the means of y_1, y_2, \dots, y_p averaged across the levels of B , we use $\bar{\mu}_.. = \sum_{i=1}^k \mu_i/k$, which is estimated by $\bar{y}_.. = \sum_{i=1}^k \bar{y}_{i..}/k$. The hypothesis $H_0: \bar{\mu}_{..1} = \bar{\mu}_{..2} = \dots = \bar{\mu}_{..p}$ comparing the means of y_1, y_2, \dots, y_p (for factor A) can be expressed using contrasts:

$$H_0: \mathbf{C}\bar{\mu}_.. = \mathbf{0}, \quad (6.91)$$

where \mathbf{C} is any $(p - 1) \times p$ full-rank contrast matrix with $\mathbf{C}\mathbf{j} = \mathbf{0}$. This is equivalent to the “flatness” test of profile analysis, the third test in Section 6.8. Under H_0 , the vector $\mathbf{C}\bar{y}_..$ is distributed as $N_{p-1}(0, \mathbf{C}\Sigma\mathbf{C}'/N)$, where $N = \sum_i n_i$ for an unbalanced design and $N = kn$ in the balanced case. We can therefore test H_0 with

$$T^2 = N(\mathbf{C}\bar{y}_..)'(\mathbf{C}\mathbf{S}_{pl}\mathbf{C}')^{-1}(\mathbf{C}\bar{y}_..), \quad (6.92)$$

where $\mathbf{S}_{pl} = \mathbf{E}/\nu_E$. The T^2 -statistic in (6.92) is distributed as T^2_{p-1, ν_E} when H_0 is true, where $\nu_E = N - k$ [see (6.83) and the comments following]. Note that the dimension of T^2 is $p - 1$ because $\mathbf{C}\bar{y}_..$ is $(p - 1) \times 1$.

For the grouping or between-subjects factor B , we wish to compare the means for the k levels of B . The mean response for the i th level of B (averaged over the levels of A) is $\sum_{r=1}^p \mu_{ir}/p = \mathbf{j}'\mu_i/p$. The hypothesis can be expressed as

$$H_0: \mathbf{j}'\mu_1 = \mathbf{j}'\mu_2 = \dots = \mathbf{j}'\mu_k, \quad (6.93)$$

which is analogous to (6.79), the “levels” hypothesis in profile analysis. This is easily tested by calculating a univariate F -statistic for a one-way ANOVA on $z_{ij} = \mathbf{j}'\mathbf{y}_{ij}$, $i = 1, 2, \dots, k$; $j = 1, 2, \dots, n_i$. There is a z_{ij} corresponding to each subject, S_{ij} . The observation vector for each subject is thus reduced to a single scalar observation, and we have a one-way ANOVA comparing the means $\mathbf{j}'\bar{y}_1, \mathbf{j}'\bar{y}_2, \dots, \mathbf{j}'\bar{y}_k$. (Note that $\mathbf{j}'\bar{y}_i/p$ is an average over the p levels of A .)

The AB interaction hypothesis is equivalent to the parallelism hypothesis in profile analysis [see (6.77)],

$$H_0: \mathbf{C}\mu_1 = \mathbf{C}\mu_2 = \dots = \mathbf{C}\mu_k. \quad (6.94)$$

In other words, differences or contrasts among the levels of factor A are the same across all levels of factor B . This is easily tested by performing a one-way MANOVA on $\mathbf{z}_{ij} = \mathbf{C}\mathbf{y}_{ij}$ or directly by

$$\Lambda = \frac{|\mathbf{C}\mathbf{E}\mathbf{C}'|}{|\mathbf{C}(\mathbf{E} + \mathbf{H})\mathbf{C}'|} \quad (6.95)$$

[see (6.77)], which is distributed as $\Lambda_{p-1, \nu_H, \nu_E}$, with $\nu_H = k - 1$ and $\nu_E = N - k$; that is, $\nu_E = \sum_i (n_i - 1)$ for the unbalanced model or $\nu_E = k(n - 1)$ in the balanced model.

6.9.4 Computation of Repeated Measures Tests

Some statistical software packages have automated repeated measures procedures that are easily implemented. However, if one is unsure as to how the resulting tests correspond to the tests in Section 6.9.3, there are two ways to obtain the tests directly. One approach is to calculate (6.92) and (6.95) outright using a matrix manipulation routine. We would need to have available the \mathbf{E} and \mathbf{H} matrices of a one-way MANOVA using a data layout as in Table 6.9.

The second approach uses simple data transformations available in virtually all programs. To test (6.91) for factor A , we would transform each \mathbf{y}_{ij} to $\mathbf{z}_{ij} = \mathbf{C}\mathbf{y}_{ij}$ by using the rows of \mathbf{C} . For example, if

$$\mathbf{C} = \begin{pmatrix} 1 & -1 & 0 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 0 & 1 & -1 \end{pmatrix},$$

then each $\mathbf{y}' = (y_1, y_2, y_3, y_4)$ becomes $\mathbf{z}' = (y_1 - y_2, y_2 - y_3, y_3 - y_4)$. We then test $H_0: \boldsymbol{\mu}_z = \mathbf{0}$ using a one-sample T^2 on all N of the \mathbf{z}_{ij} 's,

$$T^2 = N\bar{\mathbf{z}}'\mathbf{S}_z\bar{\mathbf{z}},$$

where $N = \sum_i n_i$, $\bar{\mathbf{z}} = \sum_{ij} \mathbf{z}_{ij}/N$, and $\mathbf{S}_z = \mathbf{E}_z/\nu_E$ is the pooled covariance matrix. Reject H_0 if $T^2 \geq T_{\alpha, p-1, \nu_E}^2$.

To test (6.93) for factor B , we sum the components of each observation vector to obtain $z_{ij} = \mathbf{j}'\mathbf{y}_{ij} = y_{ij1} + y_{ij2} + \dots + y_{ijp}$ and compare the means $\bar{z}_1, \bar{z}_2, \dots, \bar{z}_k$ by an F -test as in one-way ANOVA.

To test the interaction hypothesis (6.94), we transform each \mathbf{y}_{ij} to $\mathbf{z}_{ij} = \mathbf{C}\mathbf{y}_{ij}$ using the rows of \mathbf{C} as above. Note that \mathbf{z}_{ij} is $(p-1) \times 1$. We then do a one-way MANOVA on \mathbf{z}_{ij} to obtain

$$\Lambda = \frac{|\mathbf{E}_z|}{|\mathbf{E}_z + \mathbf{H}_z|}. \quad (6.96)$$

6.9.5 Repeated Measures with Two Within-Subjects Factors and One Between-Subjects Factor

The repeated measures model with two within-subjects factors A and B and one between-subjects factor C corresponds to a one-way MANOVA design in which each observation vector includes measurements on a two-way factorial arrangement of treatments. Thus each subject receives all treatment combinations of the two factors A and B . As usual, the sequence of administration of treatment combinations should be randomized for each subject. A design of this type is illustrated in Table 6.13.

Each \mathbf{y}_{ij} in Table 6.13 has nine elements, consisting of responses to the nine treatment combinations $A_1B_1, A_1B_2, \dots, A_3B_3$. We are interested in the same hypotheses as in a univariate split-plot design, but we use a multivariate approach to allow

Table 6.13 Data Layout for Repeated Measures with Two Within-Subjects Factors and One Between-Subjects Factor

Between-Subjects		Within-Subjects Factors								
Subjects		A ₁			A ₂			A ₃		
Factor	Subjects	B ₁	B ₂	B ₃	B ₁	B ₂	B ₃	B ₁	B ₂	B ₃
C ₁	S ₁₁	(y ₁₁₁	y ₁₁₂	y ₁₁₃	y ₁₁₄	y ₁₁₅	y ₁₁₆	y ₁₁₇	y ₁₁₈	y ₁₁₉) = \mathbf{y}'_{11}
	S ₁₂					\mathbf{y}'_{12}				
	⋮					⋮				
	S _{1n₁}					\mathbf{y}'_{1n_1}				
C ₂	S ₂₁					\mathbf{y}'_{21}				
	S ₂₂					\mathbf{y}'_{22}				
	⋮					⋮				
	S _{2n₂}					\mathbf{y}'_{2n_2}				
C ₃	S ₃₁					\mathbf{y}'_{31}				
	S ₃₂					\mathbf{y}'_{32}				
	⋮					⋮				
	S _{3n₃}					\mathbf{y}'_{3n_3}				

for correlated y 's. The model for the observation vectors is the one-way MANOVA model

$$\mathbf{y}_{ij} = \boldsymbol{\mu} + \boldsymbol{\gamma}_i + \boldsymbol{\varepsilon}_{ij} = \boldsymbol{\mu}_i + \boldsymbol{\varepsilon}_{ij},$$

where $\boldsymbol{\gamma}_i$ is the C effect.

To test factors A , B , and AB in Table 6.13, we use contrasts in the y 's. An example of contrast matrices would be

$$\mathbf{A} = \begin{pmatrix} 2 & 2 & 2 & -1 & -1 & -1 & -1 & -1 & -1 \\ 0 & 0 & 0 & 1 & 1 & 1 & -1 & -1 & -1 \end{pmatrix}, \quad (6.97)$$

$$\mathbf{B} = \begin{pmatrix} 2 & -1 & -1 & 2 & -1 & -1 & 2 & -1 & -1 \\ 0 & 1 & -1 & 0 & 1 & -1 & 0 & 1 & -1 \end{pmatrix}, \quad (6.98)$$

$$\mathbf{G} = \begin{pmatrix} 4 & -2 & -2 & -2 & 1 & 1 & -2 & 1 & 1 \\ 0 & 2 & -2 & 0 & -1 & 1 & 0 & -1 & 1 \\ 0 & 0 & 0 & 2 & -1 & -1 & -2 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 & -1 & 0 & -1 & 1 \end{pmatrix}. \quad (6.99)$$

Note that the matrices \mathbf{A} , \mathbf{B} , and \mathbf{G} can be formed by turning the desired comparisons for each k -level factor into a $(k-1) \times k$ contrast matrix and then using Kronecker products (see Section 2.12). For example, the rows of \mathbf{A} are orthogonal

contrasts with two comparisons:

$$\begin{aligned} A_1 &\text{ vs. } A_2 \text{ and } A_3, \\ A_2 &\text{ vs. } A_3. \end{aligned}$$

Similarly, \mathbf{B} compares

$$\begin{aligned} B_1 &\text{ vs. } B_2 \text{ and } B_3, \\ B_2 &\text{ vs. } B_3. \end{aligned}$$

So if factor A has a levels and factor B has b levels, we obtain \mathbf{A} , \mathbf{B} , and \mathbf{G} by constructing matrices \mathbf{A}^* and \mathbf{B}^*

$$\mathbf{A}^* = \begin{pmatrix} -2 & 1 & 1 \\ 0 & -1 & 1 \end{pmatrix} \quad \text{and} \quad \mathbf{B}^* = \begin{pmatrix} -2 & 1 & 1 \\ 0 & -1 & 1 \end{pmatrix},$$

and defining $\mathbf{A} = \mathbf{A}^* \otimes \mathbf{1}_b'$, $\mathbf{B} = \mathbf{1}_a' \otimes \mathbf{B}^*$, and $\mathbf{G} = \mathbf{A}^* \otimes \mathbf{B}^*$. Other orthogonal (or linearly independent) contrasts could be used for A and B . Note that the matrix \mathbf{G} is for the AB interaction and can also be obtained from products of the corresponding elements of the rows of \mathbf{A} and the rows of \mathbf{B} .

As before, we define $\bar{\mathbf{y}}_{..} = \sum_{ij} \mathbf{y}_{ij}/N$, $\mathbf{S}_{\text{pl}} = \mathbf{E}/\nu_E$, and $N = \sum_i n_i$. If there were k levels of C in Table 6.13 with mean vectors $\boldsymbol{\mu}_1, \boldsymbol{\mu}_2, \dots, \boldsymbol{\mu}_k$, then $\bar{\boldsymbol{\mu}}_{..} = \sum_{i=1}^k \boldsymbol{\mu}_i/k$, and the A main effect corresponding to $H_0: \mathbf{A}\bar{\boldsymbol{\mu}}_{..} = \mathbf{0}$ could be tested with

$$T^2 = N(\mathbf{A}\bar{\mathbf{y}}_{..})'(\mathbf{A}\mathbf{S}_{\text{pl}}\mathbf{A}')^{-1}(\mathbf{A}\bar{\mathbf{y}}_{..}), \quad (6.100)$$

which is distributed as T_{2,ν_E}^2 under H_0 , where $\nu_E = \sum_{i=1}^k (n_i - 1)$. The dimension is 2, corresponding to the two rows of \mathbf{A} .

Similarly, to test $H_0: \mathbf{B}\bar{\boldsymbol{\mu}}_{..} = \mathbf{0}$ and $H_0: \mathbf{G}\bar{\boldsymbol{\mu}}_{..} = \mathbf{0}$ for the B main effect and the AB interaction, respectively, we have

$$T^2 = N(\mathbf{B}\bar{\mathbf{y}}_{..})'(\mathbf{B}\mathbf{S}_{\text{pl}}\mathbf{B}')^{-1}(\mathbf{B}\bar{\mathbf{y}}_{..}), \quad (6.101)$$

$$T^2 = N(\mathbf{G}\bar{\mathbf{y}}_{..})'(\mathbf{G}\mathbf{S}_{\text{pl}}\mathbf{G}')^{-1}(\mathbf{G}\bar{\mathbf{y}}_{..}), \quad (6.102)$$

which are distributed as T_{2,ν_E}^2 and T_{4,ν_E}^2 , respectively. In general, if factor A has a levels and factor B has b levels, then \mathbf{A} has $a - 1$ rows, \mathbf{B} has $b - 1$ rows, and \mathbf{G} has $(a - 1)(b - 1)$ rows. The T^2 -statistics are then distributed as T_{a-1,ν_E}^2 , T_{b-1,ν_E}^2 , and $T_{(a-1)(b-1),\nu_E}^2$, respectively.

Factors A , B , and AB can be tested with Wilks' Λ as well as T^2 . Define $\mathbf{H}^* = N\bar{\mathbf{y}}_{..}\bar{\mathbf{y}}'_{..}$ from the partitioning $\sum_{ij} \mathbf{y}_{ij}\mathbf{y}'_{ij} = \mathbf{E} + \mathbf{H} + N\bar{\mathbf{y}}_{..}\bar{\mathbf{y}}'_{..}$. This can be used to test $H_0: \bar{\boldsymbol{\mu}}_{..} = \mathbf{0}$ (not usually a hypothesis of interest) by means of

$$\Lambda = \frac{|\mathbf{E}|}{|\mathbf{E} + \mathbf{H}^*|}, \quad (6.103)$$

which is $\Lambda_{p,1,\nu_E}$ if H_0 is true. Then the hypothesis of interest, $H_0: \mathbf{A}\bar{\boldsymbol{\mu}}_{..} = \mathbf{0}$ for factor A , can be tested with

$$\Lambda = \frac{|\mathbf{A}\mathbf{E}\mathbf{A}'|}{|\mathbf{A}(\mathbf{E} + \mathbf{H}^*)\mathbf{A}'|}, \quad (6.104)$$

which is distributed as $\Lambda_{a-1,1,\nu_E}$ when H_0 is true, where a is the number of levels of factor A . There are similar expressions for testing factors B and AB . Note that the dimension of Λ in (6.104) is $a - 1$, because $\mathbf{A}\mathbf{E}\mathbf{A}'$ is $(a - 1) \times (a - 1)$.

The T^2 and Wilks' Λ expressions in (6.100) and (6.104) are related by

$$\Lambda = \frac{\nu_E}{\nu_E + T^2}, \quad (6.105)$$

$$T^2 = \nu_E \frac{1 - \Lambda}{\Lambda}. \quad (6.106)$$

We can establish (6.105) as follows. By (6.27),

$$\begin{aligned} U^{(s)} &= \sum_{i=1}^s \lambda_i = \text{tr}[(\mathbf{A}\mathbf{E}\mathbf{A}')^{-1}(\mathbf{A}\mathbf{H}^*\mathbf{A}')] \\ &= \text{tr}[(\mathbf{A}\mathbf{E}\mathbf{A}')^{-1}\mathbf{A}N\bar{\mathbf{y}}_{..}\bar{\mathbf{y}}'_{..}\mathbf{A}'] \\ &= N \text{tr}[(\mathbf{A}\mathbf{E}\mathbf{A}')^{-1}\mathbf{A}\bar{\mathbf{y}}_{..}(\mathbf{A}\bar{\mathbf{y}}_{..})'] \\ &= N \text{tr}[(\mathbf{A}\bar{\mathbf{y}}_{..})'(\mathbf{A}\mathbf{E}\mathbf{A}')^{-1}\mathbf{A}\bar{\mathbf{y}}_{..}] \\ &= \frac{N}{\nu_E} (\mathbf{A}\bar{\mathbf{y}}_{..})'(\mathbf{A}\mathbf{S}_{\text{pl}}\mathbf{A}')^{-1}\mathbf{A}\bar{\mathbf{y}}_{..} \\ &= \frac{T^2}{\nu_E}. \end{aligned}$$

Since $\text{rank}(\mathbf{H}^*) = 1$, only λ_1 is nonzero, and

$$U^{(1)} = \sum_{i=1}^s \lambda_i = \lambda_1.$$

By (6.14),

$$\Lambda = \frac{1}{1 + \lambda_1} = \frac{1}{1 + U^{(1)}} = \frac{1}{1 + T^2/\nu_E},$$

which is the same as (6.105).

Factor C is tested exactly as factor B in Section 6.9.3. The hypothesis is

$$H_0: \mathbf{j}'\boldsymbol{\mu}_1 = \mathbf{j}'\boldsymbol{\mu}_2 = \cdots = \mathbf{j}'\boldsymbol{\mu}_k$$

as in (6.93), and we perform a univariate F -test on $z_{ij} = \mathbf{j}'\mathbf{y}_{ij}$ in a one-way ANOVA layout.

The AC , BC , and ABC interactions are tested as follows:

AC Interaction. The AC interaction hypothesis is

$$H_0: \mathbf{A}\boldsymbol{\mu}_1 = \mathbf{A}\boldsymbol{\mu}_2 = \cdots = \mathbf{A}\boldsymbol{\mu}_k,$$

which states that contrasts in factor A are the same across all levels of factor C . This can be tested by

$$\Lambda = \frac{|\mathbf{A}\mathbf{E}\mathbf{A}'|}{|\mathbf{A}(\mathbf{E} + \mathbf{H})\mathbf{A}'|}, \quad (6.107)$$

which is distributed as Λ_{2,ν_H,ν_E} , where $a - 1 = 2$ is the number of rows of \mathbf{A} and ν_H and ν_E are from the multivariate one-way model. Alternatively, the test can be carried out by transforming \mathbf{y}_{ij} to $\mathbf{z}_{ij} = \mathbf{Ay}_{ij}$ and doing a one-way MANOVA on \mathbf{z}_{ij} .

BC Interaction. The BC interaction hypothesis

$$H_0: \mathbf{B}\boldsymbol{\mu}_1 = \mathbf{B}\boldsymbol{\mu}_2 = \cdots = \mathbf{B}\boldsymbol{\mu}_k$$

is tested by

$$\Lambda = \frac{|\mathbf{BEB}'|}{|\mathbf{B}(\mathbf{E} + \mathbf{H})\mathbf{B}'|},$$

which is Λ_{2,ν_H,ν_E} , where $b - 1 = 2$; H_0 can also be tested by doing MANOVA on $\mathbf{z}_{ij} = \mathbf{By}_{ij}$.

ABC Interaction. The ABC interaction hypothesis

$$H_0: \mathbf{G}\boldsymbol{\mu}_1 = \mathbf{G}\boldsymbol{\mu}_2 = \cdots = \mathbf{G}\boldsymbol{\mu}_k$$

is tested by

$$\Lambda = \frac{|\mathbf{GEG}'|}{|\mathbf{G}(\mathbf{E} + \mathbf{H})\mathbf{G}'|},$$

which is Λ_{4,ν_H,ν_E} , or by doing MANOVA on $\mathbf{z}_{ij} = \mathbf{Gy}_{ij}$. In this case the dimension is $(a - 1)(b - 1) = 4$.

The above tests for AC , BC , or ABC can be also carried out with the other three MANOVA test statistics using eigenvalues of the appropriate matrices. For example, for AC we would use $(\mathbf{AEA}')^{-1}(\mathbf{AHA}')$.

■ EXAMPLE 6.9.5

The data in Table 6.14 represent a repeated measures design with two within-subjects factors and one between-subjects factor (Timm 1980). Since A and B have three levels each, as in the illustration in this section, we will use the \mathbf{A} , \mathbf{B} , and \mathbf{G} matrices in (6.97), (6.98), and (6.99). The \mathbf{E} and \mathbf{H} matrices are 9×9 and will not be shown. The overall mean vector is given by

$$\bar{\mathbf{y}}_{..} = (46.45, 39.25, 31.70, 38.85, 45.40, 40.15, 34.55, 36.90, 39.15).$$

By (6.100), the test for factor A is

$$\begin{aligned} T^2 &= N(\mathbf{A}\bar{\mathbf{y}}_{..})'(\mathbf{AS}_{\text{pl}}\mathbf{A}')^{-1}(\mathbf{A}\bar{\mathbf{y}}_{..}) \\ &= 20(-.20, 13.80) \begin{pmatrix} 2138.4 & 138.6 \\ 138.6 & 450.4 \end{pmatrix}^{-1} \begin{pmatrix} -.20 \\ 13.80 \end{pmatrix} \\ &= 8.645 > T^2_{.05, 2, 18} = 7.606. \end{aligned}$$

Table 6.14 Data from a Repeated Measures Experiment with Two Within-Subjects Factors and One Between-Subjects Factor

Between-Subjects		Within-Subjects Factors								
		A ₁			A ₂			A ₃		
Factor	Subjects	B ₁	B ₂	B ₃	B ₁	B ₂	B ₃	B ₁	B ₂	B ₃
<i>C</i> ₁	S ₁₁	20	21	21	32	42	37	32	32	32
	S ₁₂	67	48	29	43	56	48	39	40	41
	S ₁₃	37	31	25	27	28	30	31	33	34
	S ₁₄	42	40	38	37	36	28	19	27	35
	S ₁₅	57	45	32	27	21	25	30	29	29
	S ₁₆	39	39	38	46	54	43	31	29	28
	S ₁₇	43	32	20	33	46	44	42	37	31
	S ₁₈	35	34	34	39	43	39	35	39	42
	S ₁₉	41	32	23	37	51	39	27	28	30
	S _{1,10}	39	32	24	30	35	31	26	29	32
<i>C</i> ₂	S ₂₁	47	36	25	31	36	29	21	24	27
	S ₂₂	53	43	32	40	48	47	46	50	54
	S ₂₃	38	35	33	38	42	45	48	48	49
	S ₂₄	60	51	41	54	67	60	53	52	50
	S ₂₅	37	36	35	40	45	40	34	40	46
	S ₂₆	59	48	37	45	52	44	36	44	52
	S ₂₇	67	50	33	47	61	46	31	41	50
	S ₂₈	43	35	27	32	36	35	33	33	32
	S ₂₉	64	59	53	58	62	51	40	42	43
	S _{2,10}	41	38	34	41	47	42	37	41	46

For factor *B*, we use (6.101) to obtain

$$\begin{aligned}
 T^2 &= N(\mathbf{B}\bar{\mathbf{y}}_{..})'(\mathbf{B}\mathbf{S}_{\text{pl}}\mathbf{B}')^{-1}(\mathbf{B}\bar{\mathbf{y}}_{..}) \\
 &= 20(7.15, 10.55) \begin{pmatrix} 305.7 & 94.0 \\ 94.0 & 69.8 \end{pmatrix}^{-1} \begin{pmatrix} 7.15 \\ 10.55 \end{pmatrix} \\
 &= 37.438 > T^2_{.01, 2, 18} = 12.943.
 \end{aligned}$$

By (6.102), the test for the *AB* interaction is given by

$$\begin{aligned}
 T^2 &= N(\mathbf{G}\bar{\mathbf{y}}_{..})'(\mathbf{G}\mathbf{S}_{\text{pl}}\mathbf{G}')^{-1}(\mathbf{G}\bar{\mathbf{y}}_{..}) \\
 &= 61.825 > T^2_{.01, 4, 18} = 23.487.
 \end{aligned}$$

To test factor *C*, we carry out a one-way ANOVA on $z_{ij} = \mathbf{j}'\mathbf{y}_{ij}/9$:

Source	Sum of Squares	df	Mean Square	F
Between	3042.22	1	3042.22	8.54
Error	6408.98	18	356.05	

The observed F , 8.54, has a p -value of .0091 and is therefore significant.

The AC interaction is tested by (6.107) as

$$\begin{aligned}\Lambda &= \frac{|\mathbf{A}\mathbf{E}\mathbf{A}'|}{|\mathbf{A}(\mathbf{E} + \mathbf{H})\mathbf{A}'|} = \frac{3.058 \times 10^8}{3.092 \times 10^8} \\ &= .9889 > \Lambda_{.05,2,1,18} = .703.\end{aligned}$$

For the BC interaction, we have

$$\begin{aligned}\Lambda &= \frac{|\mathbf{B}\mathbf{E}\mathbf{B}'|}{|\mathbf{B}(\mathbf{E} + \mathbf{H})\mathbf{B}'|} = \frac{4.053 \times 10^6}{4.170 \times 10^6} \\ &= .9718 > \Lambda_{.05,2,1,18} = .703.\end{aligned}$$

For ABC , we obtain

$$\begin{aligned}\Lambda &= \frac{|\mathbf{G}\mathbf{E}\mathbf{G}'|}{|\mathbf{G}(\mathbf{E} + \mathbf{H})\mathbf{G}'|} = \frac{2.643 \times 10^{12}}{2.927 \times 10^{12}} \\ &= .9029 > \Lambda_{.05,4,1,18} = .551.\end{aligned}$$

In summary, factors A , B , and C and the AB interaction are significant. \square

6.9.6 Repeated Measures with Two Within-Subjects Factors and Two Between-Subjects Factors

In this section we consider a balanced two-way MANOVA design in which each observation vector arises from a two-way factorial arrangement of treatments. This is illustrated in Table 6.15 for a balanced design with three levels of all factors. Each \mathbf{y}_{ijk} has nine elements, consisting of responses to the nine treatment combinations $A_1B_1, A_2B_2, \dots, A_3B_3$ (see Table 6.13).

To test A , B , and AB , we can use the same contrast matrices \mathbf{A} , \mathbf{B} , and \mathbf{G} as in (6.97)–(6.99). We define a grand mean vector $\bar{\mathbf{y}}_{...} = \sum_{ijk} \mathbf{y}_{ijk}/N$, where N is the total number of observation vectors; in this illustration, $N = 27$. In general, $N = cdn$, where c and d are the number of levels of factors C and D and n is the number of replications in each cell (in the illustration, $n = 3$). The test statistics for A , B , and AB are as follows, where $\mathbf{S}_{pl} = \mathbf{E}/\nu_E$ and the \mathbf{E} matrix is obtained from the two-way MANOVA with $\nu_E = cd(n - 1)$ degrees of freedom.

Factor A

$$T^2 = N(\mathbf{A}\bar{\mathbf{y}}_{...})'(\mathbf{A}\mathbf{S}_{pl}\mathbf{A}')^{-1}(\mathbf{A}\bar{\mathbf{y}}_{...})$$

is distributed as T^2_{a-1, ν_E} .

Factor B

$$T^2 = N(\mathbf{B}\bar{\mathbf{y}}_{...})'(\mathbf{B}\mathbf{S}_{pl}\mathbf{B}')^{-1}(\mathbf{B}\bar{\mathbf{y}}_{...})$$

is distributed as T^2_{b-1, ν_E} .

Table 6.15 Data Layout for Repeated Measures with Two Within-Subjects Factors and Two Between-Subjects Factors

Between-Subjects			Within-Subjects Factors								
Factors		Subject	A ₁			A ₂			A ₃		
C	D		B ₁	B ₂	B ₃	B ₁	B ₂	B ₃	B ₁	B ₂	B ₃
C ₁	D ₁	S ₁₁₁							y' ₁₁₁		
		S ₁₁₂							y' ₁₁₂		
		S ₁₁₃							y' ₁₁₃		
	D ₂	S ₁₂₁							y' ₁₂₁		
		S ₁₂₂							y' ₁₂₂		
		S ₁₂₃							y' ₁₂₃		
	D ₃	S ₁₃₁							y' ₁₃₁		
		S ₁₃₂							y' ₁₃₂		
		S ₁₃₃							y' ₁₃₃		
C ₂	D ₁	S ₂₁₁							y' ₂₁₁		
		S ₂₁₂							y' ₂₁₂		
		S ₂₁₃							y' ₂₁₃		
	D ₂	S ₂₂₁							y' ₂₂₁		
		D ₃	:						:		
C ₃	D ₁		:						:		
	D ₂		:						:		
	D ₃	S ₃₃₃							y' ₃₃₃		

AB Interaction

$$T^2 = N(\mathbf{G}\bar{\mathbf{y}}_{...})'(\mathbf{G}\mathbf{S}_{\text{p}1}\mathbf{G}')^{-1}(\mathbf{G}\bar{\mathbf{y}}_{...})$$

is distributed as $T^2_{(a-1)(b-1), \nu_E}$.

To test factors C , D , and CD , we transform to $z_{ijk} = \mathbf{j}'\mathbf{y}_{ijk}$ and carry out univariate F -tests on a two-way ANOVA design.

To test factors AC , AD , and ACD , we perform a two-way MANOVA on \mathbf{Ay}_{ijk} . Then, the C main effect on \mathbf{Ay}_{ijk} compares the levels of C on \mathbf{Ay}_{ijk} , which is an effective description of the AC interaction. Similarly, the D main effect on \mathbf{Ay}_{ijk} yields the AD interaction, and the CD interaction on \mathbf{Ay}_{ijk} gives the ACD interaction.

To test factors BC , BD , and BCD , we carry out a two-way MANOVA on \mathbf{By}_{ijk} . The C main effect on \mathbf{By}_{ijk} gives the BC interaction, the D main effect on \mathbf{By}_{ijk} yields the BD interaction, and the CD interaction on \mathbf{By}_{ijk} corresponds to the BCD interaction.

Finally, to test factors ABC , ABD , and $ABCD$, we perform a two-way MANOVA on \mathbf{Gy}_{ijk} . Then the C main effect on \mathbf{Gy}_{ijk} gives the ABC interaction, the D main

effect on \mathbf{Gy}_{ijk} yields the ABD interaction, and the CD interaction on \mathbf{Gy}_{ijk} corresponds to the $ABCD$ interaction.

6.9.7 Additional Topics

Wang (1983) and Timm (1980) give a method for obtaining univariate mixed-model sums of squares from the multivariate \mathbf{E} and \mathbf{H} matrices. Crepeau et al. (1985) consider repeated measures experiments with missing data. Federer (1986) discusses the planning of repeated measures designs, emphasizing such aspects as determining the length of treatment period, eliminating carry-over effects, the nature of pre- and posttreatment, the nature of a response to a treatment, treatment sequences, and the choice of a model. Vonesh (1986) discusses sample size requirements to achieve a given power level in repeated measures designs. Patel (1986) presents a model that accommodates both within- and between-subjects covariates in repeated measures designs. Jensen (1982) compares the efficiency and robustness of various procedures.

A *multivariate* or *multiresponse* repeated measurement design will result if more than one variable is measured on each subject at each treatment combination. Such designs are discussed by Timm (1980), Reinsel (1982), Wang (1983), and Thomas (1983). Bock (1975) refers to observations of this type as *doubly multivariate* data.

6.10 GROWTH CURVES

When the subject responds to a treatment or stimulus at successive time periods, the pattern of responses is often referred to as a *growth curve*. As in repeated measures experiments, subjects are usually human or animal. We consider estimation and testing hypotheses about the form of the response curve for a single sample in Section 6.10.1 and extend to growth curves for several samples in Section 6.10.2.

6.10.1 Growth Curve for One Sample

The data layout for a single sample growth curve experiment is analogous to Table 6.11, with the levels of factor A representing time periods. Thus we have a sample of n observation vectors $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n$, for which we compute $\bar{\mathbf{y}}$ and \mathbf{S} . The usual approach is to approximate the shape of the growth curve by a polynomial function of time. If the time points are equally spaced, we can use orthogonal polynomials. This approach will be described first, followed by a method suitable for unequal time intervals.

Orthogonal polynomials are special contrasts that are often used in testing for linear, quadratic, cubic, and higher-order trends in quantitative factors. For a more complete description and derivation see Guttman (1982, pp. 194–207), Morrison (1983, pp. 182–188), or Rencher and Schaalje (2008, pp. 363–371). Here we give only a heuristic introduction to the use of these contrasts.

Suppose we administer a drug to some subjects and measure a certain reaction at 3-minute intervals. Let $\mu_1, \mu_2, \mu_3, \mu_4$, and μ_5 designate the average responses at 0, 3, 6, 9, and 12 minutes, respectively. To test the hypothesis that there are no trends in the μ_j 's, we could test $H_0: \mu_1 = \mu_2 = \cdots = \mu_5$ or $H_0: \mathbf{C}\boldsymbol{\mu} = \mathbf{0}$ using the contrast matrix

$$\mathbf{C} = \begin{pmatrix} -2 & -1 & 0 & 1 & 2 \\ 2 & -1 & -2 & -1 & 2 \\ -1 & 2 & 0 & -2 & 1 \\ 1 & -4 & 6 & -4 & 1 \end{pmatrix} \quad (6.108)$$

in $T^2 = n(\mathbf{C}\bar{\mathbf{y}})'(\mathbf{C}\mathbf{S}\mathbf{C}')^{-1}(\mathbf{C}\bar{\mathbf{y}})$ as in (6.89). The four rows of \mathbf{C} are orthogonal polynomials that test for linear, quadratic, cubic, and quartic trends in the means. As noted in Section 6.9.2, any set of orthogonal contrasts in \mathbf{C} will give the same value of T^2 to test $H_0: \mu_1 = \mu_2 = \cdots = \mu_5$. However, in this case we will be interested in using a subset of the rows of \mathbf{C} to determine the shape of the response curve.

Table A.13 (Kleinbaum, Kupper, and Muller 1988) gives orthogonal polynomials for $p = 3, 4, \dots, 10$. The $p - 1$ entries for each value of p constitute the matrix \mathbf{C} . Some software programs will generate these automatically.

As with all orthogonal contrasts, the rows of \mathbf{C} in (6.108) sum to zero and are mutually orthogonal. It is also apparent that the coefficients in each row increase and decrease in conformity with the desired pattern. Thus the entries in the first row, $(-2, -1, 0, 1, 2)$, increase steadily in a straight-line trend. The values in the second row dip down and back up in a quadratic-type bend. The third row entries increase, decrease, then increase in a cubic pattern with two bends. The fourth row bends three times in a quartic curve.

To further illustrate how the orthogonal polynomials pinpoint trends in the means when testing $H_0: \mathbf{C}\boldsymbol{\mu} = \mathbf{0}$, consider the three different patterns for $\boldsymbol{\mu}$ depicted in Figure 6.4, where $\boldsymbol{\mu}'_a = (8, 8, 8, 8, 8)$, $\boldsymbol{\mu}'_b = (20, 16, 12, 8, 4)$, and $\boldsymbol{\mu}'_c = (5, 12, 15, 12, 5)$. Let us denote the rows of \mathbf{C} in (6.108) as $\mathbf{c}'_1, \mathbf{c}'_2, \mathbf{c}'_3$, and \mathbf{c}'_4 . It is clear that $\mathbf{c}'_i \boldsymbol{\mu}_a = 0$ for $i = 1, 2, 3, 4$; that is, when $H_0: \mu_1 = \cdots = \mu_5$ is true, all four comparisons confirm it. If $\boldsymbol{\mu}$ has the pattern $\boldsymbol{\mu}_b$, only $\mathbf{c}'_1 \boldsymbol{\mu}_b$ is nonzero. The other rows are not sensitive to a linear pattern. We illustrate this for \mathbf{c}'_1 and \mathbf{c}'_2 :

$$\begin{aligned} \mathbf{c}'_1 \boldsymbol{\mu}_b &= (-2)(20) + (-1)(16) + (0)(12) + (1)(8) + (2)(4) = -44, \\ \mathbf{c}'_2 \boldsymbol{\mu}_b &= 2(20) - 16 - 2(12) - 8 + 2(4) = 0. \end{aligned}$$

For $\boldsymbol{\mu}_c$, only $\mathbf{c}'_2 \boldsymbol{\mu}_c$ is nonzero. For example,

$$\begin{aligned} \mathbf{c}'_1 \boldsymbol{\mu}_c &= -2(5) - 12 + 12 + 2(5) = 0, \\ \mathbf{c}'_2 \boldsymbol{\mu}_c &= 2(5) - 12 - 2(15) - 12 + 2(5) = -19. \end{aligned}$$

Thus each orthogonal polynomial independently detects the type of curvature it is designed for and ignores other types. Of course, real curves generally exhibit a mixture of more than one type of curvature, and in practice more than one orthogonal polynomial contrast may be significant.




Figure 6.4 Three different patterns for μ .

To test hypotheses about the shape of the curve, we therefore use the appropriate rows of \mathbf{C} in (6.108). Suppose we suspected a priori that there would be a combined linear and quadratic trend. Then we would partition \mathbf{C} as follows:

$$\begin{aligned}\mathbf{C}_1 &= \begin{pmatrix} -2 & -1 & 0 & 1 & 2 \\ 2 & -1 & -2 & -1 & 2 \end{pmatrix}, \\ \mathbf{C}_2 &= \begin{pmatrix} -1 & 2 & 0 & -2 & 1 \\ 1 & -4 & 6 & -4 & 1 \end{pmatrix}.\end{aligned}$$

We would test $H_0: \mathbf{C}_1\boldsymbol{\mu} = \mathbf{0}$ by

$$T^2 = n(\mathbf{C}_1\bar{\mathbf{y}})'(\mathbf{C}_1\mathbf{S}\mathbf{C}_1')^{-1}(\mathbf{C}_1\bar{\mathbf{y}}),$$

which is distributed as $T_{2,n-1}^2$, where 2 is the number of rows of \mathbf{C}_1 , n is the number of subjects in the sample, and $\bar{\mathbf{y}}$ and \mathbf{S} are the mean vector and covariance matrix for the sample. Similarly, $H_0: \mathbf{C}_2\boldsymbol{\mu} = \mathbf{0}$ is tested by

$$T^2 = n(\mathbf{C}_2\bar{\mathbf{y}})'(\mathbf{C}_2\mathbf{S}\mathbf{C}_2')^{-1}(\mathbf{C}_2\bar{\mathbf{y}}),$$

which is $T_{2,n-1}^2$. In this case we might expect the first to reject H_0 and the second to accept H_0 .

If we have no a priori expectations as to the shape of the curve, we could proceed as follows. Test the overall hypothesis $H_0: \mathbf{C}\boldsymbol{\mu} = \mathbf{0}$, and if H_0 rejected, use each of the rows of \mathbf{C} separately to test $H_0: \mathbf{c}_i'\boldsymbol{\mu} = 0$, $i = 1, 2, 3, 4$. The respective test statistics are

$$T^2 = n(\mathbf{C}\bar{\mathbf{y}})'(\mathbf{C}\mathbf{S}\mathbf{C}')^{-1}(\mathbf{C}\bar{\mathbf{y}}),$$

which is $T_{4,n-1}^2$, and

$$t_i = \frac{\mathbf{c}_i'\bar{\mathbf{y}}}{\sqrt{\mathbf{c}_i'\mathbf{S}\mathbf{c}_i/n}}, \quad i = 1, 2, 3, 4,$$

each of which is distributed as t_{n-1} (see Example 6.9.2).

In a case where p is large so that $\boldsymbol{\mu}$ has a large number of levels, say 10 or more, we would likely want to stop testing after the first four or five rows of \mathbf{C} and test the remaining rows in one group. However, for larger values of p , most tables of orthogonal polynomials give only the first few rows and omit those corresponding to higher degrees of curvature. We can find a matrix whose rows are orthogonal to the rows of a given matrix as follows. Suppose $p = 11$ so that \mathbf{C} is 10×11 and \mathbf{C}_1 contains the first five orthogonal polynomials. Then a matrix \mathbf{C}_2 , with rows orthogonal to those of \mathbf{C}_1 , can be obtained by selecting five linearly independent rows of

$$\mathbf{B} = \mathbf{I} - \mathbf{C}_1'(\mathbf{C}_1\mathbf{C}_1')^{-1}\mathbf{C}_1, \quad (6.109)$$

whose rows can easily be shown to be orthogonal to those of \mathbf{C}_1 . The matrix \mathbf{B} is not full rank, and some care must be exercised in choosing linearly independent rows. However, if an incorrect choice of \mathbf{C}_2 is made, the computer algorithm should indicate this as it attempts to invert $\mathbf{C}_2\mathbf{S}\mathbf{C}_2'$ in $T^2 = n(\mathbf{C}_2\bar{\mathbf{y}})'(\mathbf{C}_2\mathbf{S}\mathbf{C}_2')^{-1}(\mathbf{C}_2\bar{\mathbf{y}})$.

Alternatively, to check for significant curvature beyond the rows of \mathbf{C}_1 without finding \mathbf{C}_2 , we can use the test for additional information in a subset of variables in Section 5.8. We need not find \mathbf{C}_2 in order to find the overall T^2 , since, as noted in Section 6.9.2, any full rank $(p-1) \times p$ matrix \mathbf{C} such that $\mathbf{C}\mathbf{j} = \mathbf{0}$ will give the same value in the overall T^2 -test of $H_0: \mathbf{C}\boldsymbol{\mu} = \mathbf{0}$. We can conveniently use a simple contrast matrix such as

$$\mathbf{C} = \begin{pmatrix} 1 & -1 & 0 & \cdots & 0 \\ 0 & 1 & -1 & \cdots & 0 \\ \vdots & \vdots & \vdots & & \vdots \\ 0 & 0 & 0 & \cdots & -1 \end{pmatrix}$$

in

$$T^2 = n(\mathbf{C}\bar{\mathbf{y}})'(\mathbf{C}\mathbf{S}\mathbf{C}')^{-1}(\mathbf{C}\bar{\mathbf{y}}), \quad (6.110)$$

which is $T_{p-1,n-1}^2$. Let p_1 be the number of orthogonal polynomials in \mathbf{C}_1 and p_2 be the number of rows of \mathbf{C}_2 if it were available; that is $p_1 + p_2 = p - 1$. Then the test statistic for the p_1 orthogonal polynomials in \mathbf{C}_1 is

$$T_1^2 = n(\mathbf{C}_1\bar{\mathbf{y}})'(\mathbf{C}_1\mathbf{S}\mathbf{C}_1')^{-1}(\mathbf{C}_1\bar{\mathbf{y}}), \quad (6.111)$$

which is $T_{p_1, n-1}^2$. We wish to compare T_1^2 in (6.111) to T^2 in (6.110) to check for significant curvature beyond the rows of \mathbf{C}_1 . However, the test for additional information in a subset of variables in Section 5.8 was for the two-sample case. We can adapt (5.29) for use with the one-sample case as follows. The test for significance of any curvature remaining after that accounted for in \mathbf{C}_1 is made by comparing

$$(n - p_1 - 1) \frac{T^2 - T_1^2}{n - 1 + T_1^2}$$

with the critical value $T_{\alpha, p_2, n-p_1-1}^2$.

We now describe an approach that can be used when the time points are not equally spaced. It may also be of interest in the equal-time-increment case because it provides an estimate of the response function.

Suppose we observe the response of the subject at p time points t_1, t_2, \dots, t_p and that the average response μ at any time point t is a polynomial in t of degree $k < p$:

$$\mu = \beta_0 + \beta_1 t + \beta_2 t^2 + \cdots + \beta_k t^k.$$

This holds for each point t_r and the corresponding average response μ_r . Thus our hypothesis becomes

$$H_0: \begin{bmatrix} \mu_1 = \beta_0 + \beta_1 t_1 + \beta_2 t_1^2 + \cdots + \beta_k t_1^k \\ \mu_2 = \beta_0 + \beta_1 t_2 + \beta_2 t_2^2 + \cdots + \beta_k t_2^k \\ \vdots \\ \mu_p = \beta_0 + \beta_1 t_p + \beta_2 t_p^2 + \cdots + \beta_k t_p^k \end{bmatrix}, \quad (6.112)$$

which can be expressed in matrix notation as

$$H_0: \boldsymbol{\mu} = \mathbf{A}\boldsymbol{\beta}, \quad (6.113)$$

where

$$\mathbf{A} = \begin{pmatrix} 1 & t_1 & t_1^2 & \cdots & t_1^k \\ 1 & t_2 & t_2^2 & \cdots & t_2^k \\ \vdots & \vdots & \vdots & & \vdots \\ 1 & t_p & t_p^2 & \cdots & t_p^k \end{pmatrix} \quad \text{and} \quad \boldsymbol{\beta} = \begin{pmatrix} \beta_0 \\ \beta_1 \\ \vdots \\ \beta_k \end{pmatrix}.$$

In practice, it may be useful to transform the t_r 's by subtracting the mean or the smallest value in order to reduce their size for computational purposes.

The following method of testing H_0 is due to Rao (1959, 1973). The model $\boldsymbol{\mu} = \mathbf{A}\boldsymbol{\beta}$ is similar to a regression model $E(\mathbf{y}) = \mathbf{X}\boldsymbol{\beta}$ (see Section 10.2.1). However, in this case, we have $\text{cov}(\mathbf{y}) = \boldsymbol{\Sigma}$ rather than $\sigma^2 \mathbf{I}$, as in the standard regression assumption. In place of the usual regression approach of seeking $\hat{\boldsymbol{\beta}}$ to minimize $\text{SSE} = (\mathbf{y} - \mathbf{X}\hat{\boldsymbol{\beta}})'(\mathbf{y} - \mathbf{X}\hat{\boldsymbol{\beta}})$ [see (10.4) and (10.6)], we use a standardized distance as in (3.80), $(\bar{\mathbf{y}} - \mathbf{A}\hat{\boldsymbol{\beta}})' \mathbf{S}^{-1} (\bar{\mathbf{y}} - \mathbf{A}\hat{\boldsymbol{\beta}})$. The value of $\hat{\boldsymbol{\beta}}$ that minimizes $(\bar{\mathbf{y}} - \mathbf{A}\hat{\boldsymbol{\beta}})' \mathbf{S}^{-1} (\bar{\mathbf{y}} - \mathbf{A}\hat{\boldsymbol{\beta}})$ is

$$\hat{\beta} = (\mathbf{A}'\mathbf{S}^{-1}\mathbf{A})^{-1}\mathbf{A}'\mathbf{S}^{-1}\bar{\mathbf{y}} \quad (6.114)$$

[see Rencher and Schaalje (2008, Section 7.8.1)], and $H_0: \boldsymbol{\mu} = \mathbf{A}\boldsymbol{\beta}$ can be tested by

$$T^2 = n(\bar{\mathbf{y}} - \mathbf{A}\hat{\beta})'\mathbf{S}^{-1}(\bar{\mathbf{y}} - \mathbf{A}\hat{\beta}), \quad (6.115)$$

which is distributed as $T_{p-k-1, n-1}^2$. The dimension of T^2 is reduced from p to $p - k - 1$ because $k + 1$ parameters have been estimated in $\hat{\beta}$. The T^2 -statistic in (6.115) is usually given in the equivalent form

$$T^2 = n(\bar{\mathbf{y}}'\mathbf{S}^{-1}\bar{\mathbf{y}} - \bar{\mathbf{y}}'\mathbf{S}^{-1}\mathbf{A}\hat{\beta}). \quad (6.116)$$

The mean response at the r th time point,

$$\begin{aligned} \mu_r &= \beta_0 + \beta_1 t_r + \beta_2 t_r^2 + \cdots + \beta_k t_r^k \\ &= (1, t_r, t_r^2, \dots, t_r^k)\boldsymbol{\beta} = \mathbf{a}_r'\boldsymbol{\beta}, \end{aligned}$$

can be estimated by

$$\hat{\mu}_r = \mathbf{a}_r'\hat{\boldsymbol{\beta}}. \quad (6.117)$$

Simultaneous confidence intervals for all possible $\mathbf{a}'\boldsymbol{\beta}$ are given by

$$\mathbf{a}'\hat{\boldsymbol{\beta}} \pm \frac{T_\alpha}{\sqrt{n}} \sqrt{\mathbf{a}'(\mathbf{A}'\mathbf{S}^{-1}\mathbf{A})^{-1}\mathbf{a}} \left(1 + \frac{T^2}{n-1}\right), \quad (6.118)$$

where $T_\alpha = \sqrt{T_{\alpha, k+1, n-1}^2}$ is from Table A.7 and T^2 is given by (6.115) or (6.116). The intervals in (6.118) for $\mathbf{a}'\boldsymbol{\beta}$ include, of course, $\mathbf{a}_r'\boldsymbol{\beta}$ for the p rows of \mathbf{A} , that is, confidence intervals for the p time points. If $\mathbf{a}_r'\boldsymbol{\beta}$, $r = 1, 2, \dots, p$, are the only values of interest, we can shorten the intervals in (6.118) by using a Bonferroni coefficient $t_{\alpha/2p}$ in place of T_α :

$$\mathbf{a}_r'\hat{\boldsymbol{\beta}} \pm \frac{t_{\alpha/2p}}{\sqrt{n}} \sqrt{\mathbf{a}_r'(\mathbf{A}'\mathbf{S}^{-1}\mathbf{A})^{-1}\mathbf{a}_r} \left(1 + \frac{T^2}{n-1}\right), \quad (6.119)$$

where $t_{\alpha/2p} = t_{\alpha/2p, n-1}$. Bonferroni critical values $t_{\alpha/2p, \nu}$ are given in Table A.8. See procedures 2 and 3 in Section 5.5 for additional comments on the use of $t_{\alpha/2p}$ and T_α .

■ EXAMPLE 6.10.1

Potthoff and Roy (1964) reported measurements in a dental study on boys and girls from ages 8 to 14. The data are given in Table 6.16.

To illustrate the methods of this section, we use the data for the boys alone. In Example 6.10.2 we will compare the growth curves of the boys with those of

Table 6.16 Dental Measurements

Subject	Girls' Age in Years				Subject	Boys' Age in Years			
	8	10	12	14		8	10	12	14
1	21.0	20.0	21.5	23.0	1	26.0	25.0	29.0	31.0
2	21.0	21.5	24.0	25.5	2	21.5	22.5	23.0	26.5
3	20.5	24.0	24.5	26.0	3	23.0	22.5	24.0	27.5
4	23.5	24.5	25.0	26.5	4	25.5	27.5	26.5	27.0
5	21.5	23.0	22.5	23.5	5	20.0	23.5	22.5	26.0
6	20.0	21.0	21.0	22.5	6	24.5	25.5	27.0	28.5
7	21.5	22.5	23.0	25.0	7	22.0	22.0	24.5	26.5
8	23.0	23.0	23.5	24.0	8	24.0	21.5	24.5	25.5
9	20.0	21.0	22.0	21.5	9	23.0	20.5	31.0	26.0
10	16.5	19.0	19.0	19.5	10	27.5	28.0	31.0	31.5
11	24.5	25.0	28.0	28.0	11	23.0	23.0	23.5	25.0
					12	21.5	23.5	24.0	28.0
					13	17.0	24.5	26.0	29.5
					14	22.5	25.5	25.5	26.0
					15	23.0	24.5	26.0	30.0
					16	22.0	21.5	23.5	25.0

the girls. We first test the overall hypothesis $H_0: \mathbf{C}\boldsymbol{\mu} = \mathbf{0}$, where \mathbf{C} contains orthogonal polynomials for linear, quadratic, and cubic effects:

$$\mathbf{C} = \begin{pmatrix} -3 & -1 & 1 & 3 \\ 1 & -1 & -1 & 1 \\ -1 & 3 & -3 & 1 \end{pmatrix}. \quad (6.120)$$

From the 16 observation vectors we obtain

$$\bar{\mathbf{y}} = \begin{pmatrix} 22.88 \\ 23.81 \\ 25.72 \\ 27.47 \end{pmatrix}, \quad \mathbf{S} = \begin{pmatrix} 6.02 & 2.29 & 3.63 & 1.61 \\ 2.29 & 4.56 & 2.19 & 2.81 \\ 3.63 & 2.19 & 7.03 & 3.24 \\ 1.61 & 2.81 & 3.24 & 4.35 \end{pmatrix}.$$

To test $H_0: \mathbf{C}\boldsymbol{\mu} = \mathbf{0}$, we calculate

$$T^2 = n(\mathbf{C}\bar{\mathbf{y}})'(\mathbf{C}\mathbf{S}\mathbf{C}')^{-1}(\mathbf{C}\bar{\mathbf{y}}) = 77.957,$$

which exceeds $T^2_{.01,3,15} = 19.867$. We now test $H_0: \mathbf{c}'_i\boldsymbol{\mu} = 0$ for each row of \mathbf{C} to determine the shape of the growth curve. For the linear effect, using the first row, \mathbf{c}'_1 , we obtain

$$t_1 = \frac{\mathbf{c}'_1\bar{\mathbf{y}}}{\sqrt{\mathbf{c}'_1\mathbf{S}\mathbf{c}_1/n}} = 7.722 > t_{.005,15} = 2.947.$$

The test of significance of the quadratic component using the second row yields

$$t_2 = \frac{\mathbf{c}'_2 \bar{\mathbf{y}}}{\sqrt{\mathbf{c}'_2 \mathbf{S} \mathbf{c}_2/n}} = 1.370 < t_{.025, 15} = 2.131.$$

To test for a cubic trend, we use the third row of \mathbf{C} :

$$t_3 = \frac{\mathbf{c}'_3 \bar{\mathbf{y}}}{\sqrt{\mathbf{c}'_3 \mathbf{S} \mathbf{c}_3/n}} = -.511 > -t_{.025, 15} = -2.131.$$

Thus only the linear trend is needed to describe the growth curve.

To model the curve for each variable, we use (6.112),

$$\begin{aligned}\mu_r &= \beta_0 + \beta_1 t_r, \quad r = 1, 2, 3, 4, \text{ or} \\ \boldsymbol{\mu} &= \mathbf{A} \boldsymbol{\beta},\end{aligned}$$

where

$$\mathbf{A} = \begin{pmatrix} 1 & -3 \\ 1 & -1 \\ 1 & 1 \\ 1 & 3 \end{pmatrix}, \quad \boldsymbol{\beta} = \begin{pmatrix} \beta_0 \\ \beta_1 \end{pmatrix}.$$

The values in the second column of \mathbf{A} are obtained as $t = \text{age} - 11$. By (6.114), we obtain

$$\hat{\boldsymbol{\beta}} = (\mathbf{A}' \mathbf{S}^{-1} \mathbf{A})^{-1} \mathbf{A}' \mathbf{S}^{-1} \bar{\mathbf{y}} = \begin{pmatrix} 25.002 \\ .834 \end{pmatrix},$$

and our prediction equation is

$$\begin{aligned}\hat{\mu} &= 25.002 + .834t = 25.002 + .834(\text{age} - 11) \\ &= 15.828 + .834(\text{age}).\end{aligned}$$

□

6.10.2 Growth Curves for Several Samples

For the case of several samples or groups, the data layout would be similar to that in Table 6.9, where the p levels of factor A represent time points. Assuming the time points are equally spaced, we can use orthogonal polynomials in the $(p-1) \times p$ contrast matrix \mathbf{C} and express the basic hypothesis in the form $H_0: \mathbf{C} \bar{\boldsymbol{\mu}} = \mathbf{0}$, where $\bar{\boldsymbol{\mu}} = \sum_{i=1}^k \boldsymbol{\mu}_i/k$. This is equivalent to $H_0: \bar{\mu}_{.1} = \bar{\mu}_{.2} = \dots = \bar{\mu}_{.p}$, which compares the means of the p time points averaged across groups. As in Section 6.9.3, let us denote the sample mean vectors for the k groups as $\bar{\mathbf{y}}_1, \bar{\mathbf{y}}_2, \dots, \bar{\mathbf{y}}_k$, with grand mean $\bar{\mathbf{y}}..$ and pooled covariance matrix $\mathbf{S}_{\text{pl}} = \mathbf{E}/\nu_E$. For the overall test of $H_0: \mathbf{C} \bar{\boldsymbol{\mu}} = \mathbf{0}$ we use the test statistic

$$T^2 = N(\mathbf{C} \bar{\mathbf{y}}..)' (\mathbf{C} \mathbf{S}_{\text{pl}} \mathbf{C}')^{-1} (\mathbf{C} \bar{\mathbf{y}}..), \quad (6.121)$$

which is T_{p-1,ν_E}^2 as in (6.92), where $N = \sum_{i=1}^k n_i$ for unbalanced data or $N = kn$ for balanced data. The corresponding degrees of freedom for error is $\nu_E = N - k$ or $\nu_E = k(n - 1)$. A test that the average growth curve (averaged over groups) has a particular form can be tested with \mathbf{C}_1 , containing a subset of the rows of \mathbf{C} :

$$T^2 = N(\mathbf{C}_1 \bar{\mathbf{y}}_{..})'(\mathbf{C}_1 \mathbf{S}_{\text{pl}} \mathbf{C}_1')^{-1}(\mathbf{C}_1 \bar{\mathbf{y}}_{..}), \quad (6.122)$$

which is distributed as T_{p_1,ν_E}^2 , where p_1 is the number of rows in \mathbf{C}_1 .

The growth curves for the k groups can be compared by the interaction or parallelism test of Section 6.9.3 using either \mathbf{C} or \mathbf{C}_1 . We do a one-way MANOVA on $\mathbf{C}\mathbf{y}_{ij}$ or $\mathbf{C}_1\mathbf{y}_{ij}$, or equivalently calculate by (6.95),

$$\Lambda = \frac{|\mathbf{C}\mathbf{E}\mathbf{C}'|}{|\mathbf{C}(\mathbf{E} + \mathbf{H})\mathbf{C}'|} \quad \text{or} \quad \Lambda = \frac{|\mathbf{C}_1\mathbf{E}\mathbf{C}_1'|}{|\mathbf{C}_1(\mathbf{E} + \mathbf{H})\mathbf{C}_1'|}, \quad (6.123)$$

which are distributed as $\Lambda_{p-1,k-1,\nu_E}$ and $\Lambda_{p_1,k-1,\nu_E}$, respectively.

■ EXAMPLE 6.10.2

In Example 6.10.1, we found a linear trend for the growth curve for dental measurements of boys in Table 6.16. We now consider the growth curve for the combined group and also compare the girls' group with the boys' group.

The two sample sizes are unequal and we use (6.32) to calculate the \mathbf{E} matrix for the two groups,

$$\mathbf{E} = \begin{pmatrix} 135.39 & 67.88 & 97.76 & 67.76 \\ 67.88 & 103.76 & 72.86 & 82.71 \\ 97.76 & 72.86 & 161.39 & 103.27 \\ 67.76 & 82.71 & 103.27 & 124.64 \end{pmatrix},$$

from which we obtain $\mathbf{S}_{\text{pl}} = \mathbf{E}/\nu_E$. Using the \mathbf{C} matrix in (6.120), we can test the basic hypothesis of equal means for the combined samples, $H_0 : \mathbf{C}\bar{\mu}_{..} = \mathbf{0}$, using (6.121):

$$\begin{aligned} T^2 &= N(\mathbf{C}\bar{\mathbf{y}}_{..})'(\mathbf{C}\mathbf{S}_{\text{pl}}\mathbf{C}')^{-1}(\mathbf{C}\bar{\mathbf{y}}_{..}) \\ &= 118.322 > T_{.01,3,25}^2 = 15.538. \end{aligned}$$

To test for a linear trend, we use the first row of \mathbf{C} in (6.122):

$$\begin{aligned} T^2 &= N(\mathbf{c}_1'\bar{\mathbf{y}}_{..})'(\mathbf{c}_1'\mathbf{S}_{\text{pl}}\mathbf{c}_1)^{-1}(\mathbf{c}_1'\bar{\mathbf{y}}_{..}) \\ &= 99.445 > T_{.01,1,25}^2 = 7.770. \end{aligned}$$

This is, of course, the square of a t -statistic, but in the T^2 form it can readily be compared with the T^2 above using all three rows of \mathbf{C} . The linear trend is seen to dominate the relationship among the means.

We now compare the growth curves of the two groups using (6.123). For \mathbf{C} , we obtain

$$\begin{aligned}\Lambda &= \frac{|\mathbf{C}\mathbf{E}\mathbf{C}'|}{|\mathbf{C}(\mathbf{E} + \mathbf{H})\mathbf{C}'|} = \frac{1.3996 \times 10^8}{1.9025 \times 10^8} \\ &= .736 > \Lambda_{.05,3,1,25} = .717.\end{aligned}$$

For the linear trend, we have

$$\begin{aligned}\Lambda &= \frac{|\mathbf{c}'_1 \mathbf{E} \mathbf{c}_1|}{|\mathbf{c}'_1 (\mathbf{E} + \mathbf{H}) \mathbf{c}_1|} = \frac{1184.2}{1427.9} \\ &= .829 < \Lambda_{.05,1,1,25} = .855.\end{aligned}$$

Thus the overall comparison does not reach significance, but the more specific comparison of linear trends does give a significant result. \square

6.10.3 Additional Topics

Jackson and Bryce (1981) presented methods of analyzing growth curves based on univariate linear models. Snee (1972) and Snee et al. (1979) proposed the use of eigenvalues and eigenvectors of a matrix derived from residuals after fitting the model. If one of the eigenvalues is dominant, certain simplifications result. Bryce (1980) discussed a similar simplification for the two-group case. Geisser (1980) and Fearn (1975, 1977) gave the Bayesian approach to growth curves, including estimation and prediction. Zerbe (1979a,b) provided a randomization test requiring fewer assumptions than normal-based tests.

6.11 TESTS ON A SUBVECTOR

6.11.1 Test for Additional Information

In Section 5.8, we considered tests of significance of the additional information in a subvector when comparing two groups. We now extend these concepts to several groups and use similar notation.

Let \mathbf{y} be a $p \times 1$ vector of measurements and \mathbf{x} be a $q \times 1$ vector measured in addition to \mathbf{y} . We are interested in determining whether \mathbf{x} makes a significant contribution to the test of $H_0: \mu_1 = \mu_2 = \dots = \mu_k$ above and beyond \mathbf{y} . Another way to phrase the question is: Can the separation of groups achieved by \mathbf{x} be predicted from the separation achieved by \mathbf{y} ? It is not necessary, of course, that \mathbf{x} represent new variables. It may be that (\mathbf{y}, \mathbf{x}) is a partitioning of the present variables, and we wish to know if the variables in \mathbf{x} can be deleted because they do not contribute to rejecting H_0 .

We consider here only the one-way MANOVA, but the results could be extended to higher-order designs, where various possibilities arise. In a two-way context, for

example, it may happen that \mathbf{x} contributes nothing to the A main effect but does contribute significantly to the B main effect.

It is assumed that we have k samples,

$$\begin{pmatrix} \mathbf{y}_{ij} \\ \mathbf{x}_{ij} \end{pmatrix}, \quad i = 1, 2, \dots, k; j = 1, 2, \dots, n,$$

from which we calculate

$$\mathbf{E} = \begin{pmatrix} \mathbf{E}_{yy} & \mathbf{E}_{yx} \\ \mathbf{E}_{xy} & \mathbf{E}_{xx} \end{pmatrix}, \quad \mathbf{H} = \begin{pmatrix} \mathbf{H}_{yy} & \mathbf{H}_{yx} \\ \mathbf{H}_{xy} & \mathbf{H}_{xx} \end{pmatrix},$$

where \mathbf{E} and \mathbf{H} are $(p+q) \times (p+q)$ and \mathbf{E}_{yy} and \mathbf{H}_{yy} are $p \times p$.

Then

$$\Lambda(\mathbf{y}, \mathbf{x}) = \frac{|\mathbf{E}|}{|\mathbf{E} + \mathbf{H}|} \quad (6.124)$$

is distributed as $\Lambda_{p+q, \nu_H, \nu_E}$ and tests the significance of group separation using the full vector (\mathbf{y}, \mathbf{x}) . In the balanced one-way model, the degrees of freedom are $\nu_H = k - 1$ and $\nu_E = k(n - 1)$. To test group separation using the reduced vector \mathbf{y} , we can compute

$$\Lambda(\mathbf{y}) = \frac{|\mathbf{E}_{yy}|}{|\mathbf{E}_{yy} + \mathbf{H}_{yy}|}, \quad (6.125)$$

which is distributed as $\Lambda_{p, \nu_H, \nu_E}$.

To test the hypothesis that the extra variables in \mathbf{x} do not contribute anything significant to separating the groups beyond the information already available in \mathbf{y} , we calculate

$$\Lambda(\mathbf{x}|\mathbf{y}) = \frac{\Lambda(\mathbf{y}, \mathbf{x})}{\Lambda(\mathbf{y})}, \quad (6.126)$$

which is distributed as $\Lambda_{q, \nu_H, \nu_E - p}$. Note that the dimension of $\Lambda(\mathbf{x}|\mathbf{y})$ is q , the number of x 's. The error degrees of freedom $\nu_E - p$ has been adjusted for the p y 's. Thus to test for the contribution of additional variables to separation of groups, we take the ratio of Wilks' Λ for the full set of variables in (6.124) to Wilks' Λ for the reduced set in (6.125). If the addition of \mathbf{x} makes $\Lambda(\mathbf{y}, \mathbf{x})$ sufficiently smaller than $\Lambda(\mathbf{y})$, then $\Lambda(\mathbf{x}|\mathbf{y})$ in (6.126) will be small enough to reject the hypothesis.

If we are interested in the effect of adding a single x , then $q = 1$, and (6.126) becomes

$$\Lambda(x|\mathbf{y}_1, \dots, \mathbf{y}_p) = \frac{\Lambda(y_1, \dots, y_p, x)}{\Lambda(y_1, \dots, y_p)}, \quad (6.127)$$

which is distributed as $\Lambda_{1, \nu_H, \nu_E - p}$. In this test we are inquiring whether x reduces the overall Λ by a significant amount. With a dimension of 1, the Λ -statistic in (6.127) has an exact F -transformation from Table 6.1,

$$F = \frac{1 - \Lambda}{\Lambda} \frac{\nu_E - p}{\nu_H}, \quad (6.128)$$

which is distributed as $F_{\nu_H, \nu_E - p}$. The statistic (6.127) is often referred to as a *partial Λ -statistic*, and correspondingly, (6.128) is called a *partial F -statistic*.

In (6.127) and (6.128), we have a test of the significance of a variable in the presence of the other variables. For a breakdown of precisely how the contribution of a variable depends on the other variables, see Rencher (1993; 1998, Section 4.1.6).

We can rewrite (6.127) as

$$\Lambda(y_1, \dots, y_p, x) = \Lambda(x|y_1, \dots, y_p)\Lambda(y_1, \dots, y_p) \leq \Lambda(y_1, \dots, y_p), \quad (6.129)$$

which shows that Wilks' Λ can only decrease with an additional variable.

■ EXAMPLE 6.11.1

We use the rootstock data of Table 6.2 to illustrate tests on subvectors. From Example 6.1.7, we have, for all four variables, $\Lambda(y_1, y_2, y_3, y_4) = .1540$. For the first two variables, we obtain $\Lambda(y_1, y_2) = .6990$. Then to test the significance of y_3 and y_4 adjusted for y_1 and y_2 , we have by (6.126),

$$\Lambda(y_3, y_4|y_1, y_2) = \frac{\Lambda(y_1, y_2, y_3, y_4)}{\Lambda(y_1, y_2)} = \frac{.1540}{.6990} = .2203,$$

which is less than the critical value $\Lambda_{.05, 2, 5, 40} = .639$.

Similarly, the test for y_4 adjusted for y_1 , y_2 , and y_3 is given by (6.127) as

$$\begin{aligned} \Lambda(y_4|y_1, y_2, y_3) &= \frac{\Lambda(y_1, y_2, y_3, y_4)}{\Lambda(y_1, y_2, y_3)} = \frac{.1540}{.2460} \\ &= .6261 < \Lambda_{.05, 1, 5, 39} = .759. \end{aligned}$$

For each of the other variables, we have a similar test:

$$\begin{aligned} y_3 : \quad \Lambda(y_3|y_1, y_2, y_4) &= \frac{.1540}{.2741} = .5618 < \Lambda_{.05, 1, 5, 39} = .759 \\ y_2 : \quad \Lambda(y_2|y_1, y_3, y_4) &= \frac{.1540}{.1922} = .8014 > \Lambda_{.05, 1, 5, 39} = .759 \\ y_1 : \quad \Lambda(y_1|y_2, y_3, y_4) &= \frac{.1540}{.1599} = .9630 > \Lambda_{.05, 1, 5, 39} = .759. \end{aligned}$$

Thus the two variables y_3 and y_4 , either individually or together, contribute a significant amount to separation of the six groups. \square

6.11.2 Stepwise Selection of Variables

If there are no variables for which we have a priori interest in testing for significance, we can do a data-directed search for the variables that best separate the groups. Such a strategy is often called *stepwise discriminant analysis*, although it could more aptly be called stepwise MANOVA. The procedure appears in many software packages.

We first describe an approach that is usually called *forward selection*. At the first step calculate $\Lambda(y_i)$ for each individual variable and choose the one with minimum $\Lambda(y_i)$ (or maximum associated F). At the second step calculate $\Lambda(y_i|y_1)$ for each of

the $p - 1$ variables not entered at the first step, where y_1 indicates the first variable entered. For the second variable we choose the one with minimum $\Lambda(y_i|y_1)$ (or maximum associated partial F), that is, the variable that adds the maximum separation to the one entered at step 1. Denote the variable entered at step 2 by y_2 . At the third step calculate $\Lambda(y_i|y_1, y_2)$ for each of the $p - 2$ remaining variables and choose the one that minimizes $\Lambda(y_i|y_1, y_2)$ (or maximizes the associated partial F). Continue this process until the F falls below some predetermined threshold value, say F_{in} .

A *stepwise* procedure follows a similar sequence, except that after a variable has entered, the variables previously selected are reexamined to see if each still contributes a significant amount. The variable with smallest partial F will be removed if the partial F is less than a second threshold value, F_{out} . If F_{out} is the same as F_{in} , there is a very small possibility that the procedure will cycle continuously without stopping. This possibility can be eliminated by using a value of F_{out} slightly less than F_{in} . For an illustration of the stepwise procedure, see Example 8.9.

PROBLEMS

6.1 Verify the computational forms given in (6.3) and (6.5); that is, show that

$$(a) \sum_{ij} (y_{ij} - \bar{y}_{i.})^2 = \sum_{ij} y_{ij}^2 - \sum_i y_{i.}^2/n$$

$$(b) n \sum_i (\bar{y}_{i.} - \bar{y}_{..})^2 = \sum_i y_{i.}^2/n - y_{..}^2/kn$$

6.2 Show that Wilks' Λ can be expressed in terms of the eigenvalues of $\mathbf{E}^{-1}\mathbf{H}$ as in (6.14).

6.3 Show that the eigenvalues of the matrices $\mathbf{E}^{-1}\mathbf{H}$ and $(\mathbf{E}^{1/2})^{-1}\mathbf{H}(\mathbf{E}^{1/2})^{-1}$ are the same, as noted in Section 6.1.4, where $\mathbf{E}^{1/2}$ is the square root matrix defined in (2.112).

6.4 Show that F_2 in (6.26) is the same as F_1 in (6.25).

6.5 Show that F_3 in (6.30) is the same as F_2 in (6.29).

6.6 Show that if there is only one nonzero eigenvalue λ_1 , then $U^{(1)}$, $V^{(1)}$, and Λ can be expressed in terms of θ , as in (6.33)–(6.35).

6.7 Show that (5.16), (5.18), and (5.19), which relate T^2 to Λ , $V^{(s)}$, and θ , follow from (6.33)–(6.35) and (6.38), $U^{(1)} = T^2/(n_1 + n_2 - 2)$.

6.8 Verify the computational forms of \mathbf{H} and \mathbf{E} in (6.31) and (6.32); that is, show that

$$(a) \sum_{i=1}^k n_i (\bar{y}_{i.} - \bar{y}_{..}) (\bar{y}_{i.} - \bar{y}_{..})' = \sum_{i=1}^k \mathbf{y}_{i.} \mathbf{y}_{i.}' / n_i - \mathbf{y}_{..} \mathbf{y}_{..}' / N$$

$$(b) \sum_{i=1}^k \sum_{j=1}^{n_i} (\mathbf{y}_{ij} - \bar{y}_{i.}) (\mathbf{y}_{ij} - \bar{y}_{i.})' = \sum_{i=1}^k \sum_{j=1}^{n_i} \mathbf{y}_{ij} \mathbf{y}_{ij}' - \sum_{i=1}^k \mathbf{y}_{i.} \mathbf{y}_{i.}' / n_i$$

- 6.9** Show that for two groups, $\mathbf{H} = \sum_{i=1}^2 n_i (\bar{\mathbf{y}}_{i\cdot} - \bar{\mathbf{y}}_{..})(\bar{\mathbf{y}}_{i\cdot} - \bar{\mathbf{y}}_{..})'$ can be expressed as $\mathbf{H} = [n_1 n_2 / (n_1 + n_2)] (\bar{\mathbf{y}}_{1\cdot} - \bar{\mathbf{y}}_{2\cdot})(\bar{\mathbf{y}}_{1\cdot} - \bar{\mathbf{y}}_{2\cdot})'$, thus verifying (6.37). Note that

$$\bar{\mathbf{y}}_{..} = \frac{n_1 \bar{\mathbf{y}}_{1\cdot} + n_2 \bar{\mathbf{y}}_{2\cdot}}{n_1 + n_2}.$$

- 6.10** Show that θ can be expressed as $\theta = \text{SSH}(z)/[\text{SSE}(z) + \text{SSH}(z)]$ as in (6.41).

- 6.11** Show that

$$\prod_{i=1}^s \frac{1}{1 + \lambda_i} = \prod_{i=1}^s (1 - r_i^2),$$

as in (6.44), where $r_i^2 = \lambda_i / (1 + \lambda_i)$.

- 6.12** Show that the F -approximation based on A_P in (6.49) reduces to (6.26) if $A_P = V^{(s)}/s$, as in (6.48).

- 6.13** Show that if $s = 1$, A_{LH} in (6.50) reduces to (6.42).

- 6.14** Show that the F -approximation denoted by F_3 in (6.30) is equivalent to (6.51).

- 6.15** Show that $\text{cov}(\hat{\boldsymbol{\delta}}) = \frac{\Sigma}{n} \sum_{i=1}^k c_i^2$ as in (6.60).

- 6.16** If $\mathbf{z}_{ij} = \mathbf{C}\mathbf{y}_{ij}$, where \mathbf{C} is $(p-1) \times p$, show that $\mathbf{H}_z = \mathbf{C}\mathbf{H}\mathbf{C}'$ and $\mathbf{E}_z = \mathbf{C}\mathbf{E}\mathbf{C}'$, as used in (6.78).

- 6.17** Why do \mathbf{C} and \mathbf{C}' not “cancel out” of Wilks’ Λ in (6.78)?

- 6.18** Show that under H_{03} and H_{01} , $\mathbf{C}\bar{\mathbf{y}}_{..}$ is $N_{p-1}(\mathbf{0}, \mathbf{C}\Sigma\mathbf{C}'/kn)$, as noted preceding (6.83).

- 6.19** Show that $T^2 = kn(\mathbf{C}\bar{\mathbf{y}}_{..})'(\mathbf{C}\mathbf{E}\mathbf{C}'/\nu_E)^{-1}\mathbf{C}\bar{\mathbf{y}}_{..}$ in (6.83) is distributed as T_{p-1, ν_E}^2 .

- 6.20** For ε defined by (6.88), show that $\varepsilon = 1$ when $\Sigma = \sigma^2 \mathbf{I}$.

- 6.21** Give a justification of the Wilks’ Λ test of $H_0: \bar{\boldsymbol{\mu}}_i = \mathbf{0}$ in (6.103).

- 6.22** Provide an alternative derivation of (6.105), $\Lambda = \nu_E / (\nu_E + T^2)$, starting with (6.104).

- 6.23** Obtain T^2 in terms of Λ in (6.106) from (6.105).

- 6.24** Show that the rows of \mathbf{C}_1 are orthogonal to those of $\mathbf{B} = \mathbf{I} - \mathbf{C}_1'(\mathbf{C}_1\mathbf{C}_1')^{-1}\mathbf{C}_1$ in (6.109).

- 6.25** Show that $\hat{\boldsymbol{\beta}}$ in (6.114) minimizes $(\bar{\mathbf{y}} - \mathbf{A}\hat{\boldsymbol{\beta}})' \mathbf{S}^{-1} (\bar{\mathbf{y}} - \mathbf{A}\hat{\boldsymbol{\beta}})$.

- 6.26** Show that T^2 in (6.116) is equivalent to T^2 in (6.115).

- 6.27** Baten, Tack, and Baeder (1958) compared judges’ scores on fish prepared by three methods. Twelve fish were cooked by each method, and several judges

Table 6.17 Judges' Scores on Fish Prepared by Three Methods

Method 1				Method 2				Method 3			
y_1	y_2	y_3	y_4	y_1	y_2	y_3	y_4	y_1	y_2	y_3	y_4
5.4	6.0	6.3	6.7	5.0	5.3	5.3	6.5	4.8	5.0	6.5	7.0
5.2	6.2	6.0	5.8	4.8	4.9	4.2	5.6	5.4	5.0	6.0	6.4
6.1	5.9	6.0	7.0	3.9	4.0	4.4	5.0	4.9	5.1	5.9	6.5
4.8	5.0	4.9	5.0	4.0	5.1	4.8	5.8	5.7	5.2	6.4	6.4
5.0	5.7	5.0	6.5	5.6	5.4	5.1	6.2	4.2	4.6	5.3	6.3
5.7	6.1	6.0	6.6	6.0	5.5	5.7	6.0	6.0	5.3	5.8	6.4
6.0	6.0	5.8	6.0	5.2	4.8	5.4	6.0	5.1	5.2	6.2	6.5
4.0	5.0	4.0	5.0	5.3	5.1	5.8	6.4	4.8	4.6	5.7	5.7
5.7	5.4	4.9	5.0	5.9	6.1	5.7	6.0	5.3	5.4	6.8	6.6
5.6	5.2	5.4	5.8	6.1	6.0	6.1	6.2	4.6	4.4	5.7	5.6
5.8	6.1	5.2	6.4	6.2	5.7	5.9	6.0	4.5	4.0	5.0	5.9
5.3	5.9	5.8	6.0	5.1	4.9	5.3	4.8	4.4	4.2	5.6	5.5

Source: Baten, Tack, and Baeder (1958, p. 8).

tasted fish samples and rated each on four variables: y_1 = aroma, y_2 = flavor, y_3 = texture, and y_4 = moisture. The data are in Table 6.17. Each entry is an average score for the judges on that fish.

- (a) Compare the three methods using all four MANOVA tests.
- (b) Compute the following measures of multivariate association from Section 6.1.8 : η_A^2 , η_θ^2 , A_A , A_{LH} , A_P .
- (c) Based on the eigenvalues, is the essential dimensionality of the space containing the mean vectors equal to 1 or 2?
- (d) Using contrasts, test the following two comparisons: 1 and 2 vs. 3, and 1 vs. 2.
- (e) If any of the four tests in (a) is significant, run an ANOVA F -test on each y_i and examine the discriminant function $z = \mathbf{a}'\mathbf{y}$ (Section 6.4).
- (f) Test the significance of y_3 and y_4 adjusted for y_1 and y_2 .
- (g) Test the significance of each variable adjusted for the other three.

6.28 Table 6.18 from Keuls et al. (1984) gives data from a two-way (fixed-effects) MANOVA on snap beans showing the results of four variables: y_1 = yield earliness, y_2 = specific leaf area (SLA) earliness, y_3 = total yield, and y_4 = average SLA. The factors are sowing date (S) and variety (V).

- (a) Test for main effects and interaction using all four MANOVA statistics.
- (b) In previous experiments, the second variety gave higher yields. Compare variety 2 with varieties 1 and 3 by means of a test on a contrast.

Table 6.18 Snap Bean Data

<i>S</i>	<i>V</i>		<i>y</i> ₁	<i>y</i> ₂	<i>y</i> ₃	<i>y</i> ₄	<i>S</i>	<i>V</i>		<i>y</i> ₁	<i>y</i> ₂	<i>y</i> ₃	<i>y</i> ₄
1	1	1	59.3	4.5	38.4	295	3	1	1	68.1	3.4	42.2	280
		2	60.3	4.5	38.6	302			2	68.0	2.9	42.4	284
		3	60.9	5.3	37.2	318			3	68.5	3.3	41.5	286
		4	60.6	5.8	38.1	345			4	68.6	3.1	41.9	284
		5	60.4	6.0	38.8	325			5	68.6	3.3	42.1	268
1	2	1	59.3	6.7	37.9	275	3	2	1	64.0	3.6	40.9	233
		2	59.4	4.8	36.6	290			2	63.4	3.9	41.4	248
		3	60.0	5.1	38.7	295			3	63.5	3.7	41.6	244
		4	58.9	5.8	37.5	296			4	63.4	3.7	41.4	266
		5	59.5	4.8	37.0	330			5	63.5	4.1	41.1	244
1	3	1	59.4	5.1	38.7	299	3	3	1	68.0	3.7	42.3	293
		2	60.2	5.3	37.0	315			2	68.7	3.5	41.6	284
		3	60.7	6.4	37.4	304			3	68.7	3.8	40.7	277
		4	60.5	7.1	37.0	302			4	68.4	3.5	42.0	299
		5	60.1	7.8	36.9	308			5	68.6	3.4	42.4	285
2	1	1	63.7	5.4	39.5	271	4	1	1	69.8	1.4	48.4	265
		2	64.1	5.4	39.2	284			2	69.5	1.3	47.8	247
		3	63.4	5.4	39.0	281			3	69.5	1.3	46.9	231
		4	63.2	5.3	39.0	291			4	69.9	1.3	47.5	268
		5	63.2	5.0	39.0	270			5	70.3	1.1	47.1	247
2	2	1	60.6	6.8	38.1	248	4	2	1	66.6	1.8	45.7	205
		2	61.0	6.5	38.6	264			2	66.5	1.7	46.8	239
		3	60.7	6.8	38.8	257			3	67.1	1.7	46.3	230
		4	60.6	7.1	38.6	260			4	65.8	1.8	46.3	235
		5	60.3	6.0	38.5	261			5	65.6	1.9	46.1	220
2	3	1	63.8	5.7	40.5	282	4	3	1	70.1	1.7	48.1	253
		2	63.2	6.1	40.2	284			2	72.3	0.7	47.8	249
		3	63.3	6.0	40.0	291			3	69.7	1.5	46.7	226
		4	63.2	5.9	40.0	299			4	69.9	1.3	47.1	248
		5	63.1	5.4	39.7	295			5	69.8	1.4	46.7	236

- (c) Test linear, quadratic, and cubic contrasts for sowing date. (Interpretation of these for mean vectors is not as straightforward as for univariate means.)
- (d) If any of the tests in part (a) rejects H_0 , carry out ANOVA F -tests on the four variables.
- (e) Test the significance of y_3 and y_4 adjusted for y_1 and y_2 in main effects and interaction.
- (f) Test the significance of each variable adjusted for the other three in main effects and interaction.

Table 6.19 Blood Data

Subject	Reagent 1			Reagent 2			Reagent 3			Reagent 4		
	y_1	y_2	y_3									
1	8.0	3.96	12.5	8.0	3.93	12.7	7.9	3.86	13.0	7.9	3.87	13.2
2	4.0	5.37	16.9	4.2	5.35	17.2	4.1	5.39	17.2	4.0	5.35	17.3
3	6.3	5.47	17.1	6.3	5.39	17.5	6.0	5.39	17.2	6.1	5.41	17.4
4	9.4	5.16	16.2	9.4	5.16	16.7	9.4	5.17	16.7	9.1	5.16	16.7
5	8.2	5.16	17.0	8.0	5.13	17.5	8.1	5.10	17.4	7.8	5.12	17.5
6	11.0	4.67	14.3	10.7	4.60	14.7	10.6	4.52	14.6	10.5	4.58	14.7
7	6.8	5.20	16.2	6.8	5.16	16.7	6.9	5.13	16.8	6.7	5.19	16.8
8	9.0	4.65	14.7	9.0	4.57	15.0	8.9	4.58	15.0	8.6	4.55	15.1
9	6.1	5.22	16.3	6.0	5.16	16.9	6.1	5.14	16.9	6.0	5.21	16.9
10	6.4	5.13	15.9	6.4	5.11	16.4	6.4	5.11	16.4	6.3	5.07	16.3
11	5.6	4.47	13.3	5.5	4.45	13.6	5.3	4.46	13.6	5.3	4.44	13.7
12	8.2	5.22	16.0	8.2	5.14	16.5	8.0	5.14	16.5	7.8	5.16	16.5
13	5.7	5.10	14.9	5.6	5.05	15.3	5.5	5.02	15.4	5.4	5.05	15.5
14	9.8	5.25	16.1	9.8	5.15	16.6	8.1	5.10	13.8	9.4	5.16	16.6
15	5.9	5.28	15.8	5.8	5.25	16.4	5.7	5.26	16.4	5.6	5.29	16.2
16	6.6	4.65	12.8	6.4	4.59	13.2	6.3	4.58	13.1	6.4	4.57	13.2
17	5.7	4.42	14.5	5.5	4.31	14.9	5.5	4.30	14.9	5.4	4.32	14.8
18	6.7	4.38	13.1	6.5	4.32	13.4	6.5	4.32	13.6	6.5	4.31	13.5
19	6.8	4.67	15.6	6.6	4.57	15.8	6.5	4.55	16.0	6.5	4.56	15.9
20	9.6	5.64	17.0	9.5	5.58	17.5	9.3	5.50	17.4	9.2	5.46	17.5

- 6.29** The bar steel data in Table 6.6 was analyzed in Example 6.5.2 as a two-way fixed-effects design. Consider lubricants to be random so that we have a mixed model. Test for main effects and interaction.
- 6.30** In Table 6.19, we have a comparison of four reagents (Burdick 1979). The first reagent is the one presently in use and the other three are less expensive reagents that we wish to compare with the first. All four reagents are used with a blood sample from each patient. The three variables measured for each reagent are y_1 = white blood count, y_2 = red blood count, and y_3 = hemoglobin count.
- (a) Analyze as a randomized block design with subjects as blocks.
 - (b) Compare the first reagent with the other three using a contrast.
- 6.31** The data in Table 6.20 from Box (1950) show the amount of fabric wear y_1 , y_2 , and y_3 in three successive periods: (1) the first 1000 revolutions, (2) the second 1000 revolutions, and (3) the third 1000 revolutions of the abrasive wheel. There were three factors: type of abrasive surface, type of filler, and proportion of filler. There were two replications. Carry out a three-way MANOVA, testing for main effects and interactions. (Ignore the repeated measures aspects of the data.)

Table 6.20 Wear of Coated Fabrics in Three Periods (mg)

Surface Treatment	Filler	Proportion of Filler								
		P_1 (25%)			P_2 (50%)			P_3 (75%)		
		y_1	y_2	y_3	y_1	y_2	y_3	y_1	y_2	y_3
T_0	F_1	194	192	141	233	217	171	265	252	207
		208	188	165	241	222	201	269	283	191
	F_2	239	127	90	224	123	79	243	117	100
		187	105	85	243	123	110	226	125	75
T_1	F_1	155	169	151	198	187	176	235	225	166
		173	152	141	177	196	167	229	270	183
	F_2	137	82	77	129	94	78	155	76	92
		160	82	83	98	89	48	132	105	67

Table 6.21 Weights of Cork Borings (cg) in Four Directions for 28 Trees

Tree	N	E	S	W	Tree	N	E	S	W
1	72	66	76	77	15	91	79	100	75
2	60	53	66	63	16	56	68	47	50
3	56	57	64	58	17	79	65	70	61
4	41	29	36	38	18	81	80	68	58
5	32	32	35	36	19	78	55	67	60
6	30	35	34	26	20	46	38	37	38
7	39	39	31	27	21	39	35	34	37
8	42	43	31	25	22	32	30	30	32
9	37	40	31	25	23	60	50	67	54
10	33	29	27	36	24	35	37	48	39
11	32	30	34	28	25	39	36	39	31
12	63	45	74	63	26	50	34	37	40
13	54	46	60	52	27	43	37	39	50
14	47	51	52	43	28	48	54	57	43

6.32 The fabric wear data in Table 6.20 can be considered to be a growth curve model, with the three periods (y_1, y_2, y_3) representing repeated measurements on the same specimen. We thus have one within-subjects factor, to which we should assign polynomial contrasts $(-1, 0, 1)$ and $(-1, 2, -1)$, and a three-way between-subjects classification. Test for period and the interaction of period with the between-subjects factors and interactions.

6.33 Carry out a profile analysis on the fish data in Table 6.17, testing for parallelism, equal levels, and flatness.

- 6.34** Rao (1948) measured the weight of cork borings taken from the north (N), east (E), south (S), and west (W) directions of 28 trees. The data are given in Table 6.21. It is of interest to compare the bark thickness (and hence weight) in the four directions. This can be done by analyzing the data as a one-sample repeated measures design. Since the primary comparison of interest is north and south vs. east and west, use the contrast matrix

$$\mathbf{C} = \begin{pmatrix} 1 & -1 & 1 & -1 \\ 1 & 0 & -1 & 0 \\ 0 & 1 & 0 & -1 \end{pmatrix}.$$

- (a) Test $H_0: \mu_N = \mu_E = \mu_S = \mu_W$ using the entire matrix \mathbf{C} .
 - (b) If the test in (a) rejects H_0 , test each row of \mathbf{C} .
- 6.35** Analyze the glucose data in Table 3.9 as a one-sample repeated measures design with two within-subjects factors. Factor A is a comparison of fasting test vs. 1 hour posttest. The three levels of factor B are y_1 (and x_1), y_2 (and x_2), and y_3 (and x_3).
- 6.36** Table 6.22 gives survival times for cancer patients (Cameron and Pauling 1978; see also Andrews and Herzberg 1985, pp. 203–206). The factors in this two-way design are gender (1 = male, 2 = female) and type of cancer (1 = stomach, 2 = bronchus, 3 = colon, 4 = rectum, 5 = bladder, 6 = kidney). The variables (repeated measures) are y_1 = survival time (days) of patient treated with ascorbate measured from date of first hospital attendance, y_2 = mean survival time for the patient's 10 matched controls (untreated with ascorbate), y_3 = survival time after ascorbate treatment ceased, and y_4 = mean survival time after all treatment ceased for the patient's 10 matched controls. Analyze as a repeated measures design with one within-subjects factor (y_1, y_2, y_3, y_4) and a two-way (unbalanced) design between subjects. Since the two-way classification of subjects is unbalanced, you will need to use a program that allows for this or delete some observations to achieve a balanced design.
- 6.37** Analyze the ramus bone data of Table 3.9 as a one-sample growth curve design.
- (a) Using a matrix \mathbf{C} of orthogonal polynomial contrasts, test the hypothesis of overall equality of means, $H_0: \mathbf{C}\boldsymbol{\mu} = \mathbf{0}$.
 - (b) If the overall hypothesis in (a) is rejected, find the degree of growth curve by testing each row of \mathbf{C} .
- 6.38** Table 6.23 contains the weights of 13 male mice measured every 3 days from birth to weaning. The data set was reported and analyzed by Williams and Izenman (1981) and by Izenman and Williams (1989) and has been further analyzed by Rao (1984, 1987) and by Lee (1988). Analyze as a one-sample growth curve design.

Table 6.22 Survival Times for Cancer Patients

Type of Cancer	Gender	Age	y_1	y_2	y_3	y_4
1	2	61	124	264	124	38
1	1	69	42	62	12	18
1	2	62	25	149	19	36
1	2	66	45	18	45	12
1	1	63	412	180	257	64
1	1	79	51	142	23	20
1	1	76	1112	35	128	13
1	1	54	46	299	46	51
1	1	62	103	85	90	10
1	1	46	146	361	123	52
1	1	57	340	269	310	28
1	2	59	396	130	359	55
2	1	74	81	72	74	33
2	1	74	461	134	423	18
2	1	66	20	84	16	20
2	1	52	450	98	450	58
2	2	48	246	48	87	13
2	2	64	166	142	115	49
2	1	70	63	113	50	38
2	1	77	64	90	50	24
2	1	71	155	30	113	18
2	1	39	151	260	38	34
2	1	70	166	116	156	20
2	1	70	37	87	27	27
2	1	55	223	69	218	32
2	1	74	138	100	138	27
2	1	69	72	315	39	39
2	1	73	245	188	231	65
3	2	76	248	292	135	18
3	2	58	377	492	50	30
3	1	49	189	462	189	65
3	1	69	1843	235	1267	17
⋮	⋮	⋮	⋮	⋮	⋮	⋮
6	1	69	1685	1056	46	15

- (a) Using a matrix \mathbf{C} of orthogonal polynomial contrasts, test the hypothesis of overall equality of means, $H_0: \mathbf{C}\boldsymbol{\mu} = \mathbf{0}$.
- (b) If the overall hypothesis in (a) is rejected, find the degree of growth curve by testing each row of \mathbf{C} .
- 6.39** In Table 6.24, we have measurements of proportions of albumin at four time points on three groups of trout (Beauchamp and Hoel 1973).

Table 6.23 Weights of 13 Male Mice Measured at Successive Intervals of 3 Days over 21 Days from Birth to Weaning

Mouse	Day 3	Day 6	Day 9	Day 12	Day 15	Day 18	Day 21
1	.190	.388	.621	.823	1.078	1.132	1.191
2	.218	.393	.568	.729	.839	.852	1.004
3	.211	.394	.549	.700	.783	.870	.925
4	.209	.419	.645	.850	1.001	1.026	1.069
5	.193	.362	.520	.530	.641	.640	.751
6	.201	.361	.502	.530	.657	.762	.888
7	.202	.370	.498	.650	.795	.858	.910
8	.190	.350	.510	.666	.819	.879	.929
9	.219	.399	.578	.699	.709	.822	.953
10	.225	.400	.545	.690	.796	.825	.836
11	.224	.381	.577	.756	.869	.929	.999
12	.187	.329	.441	.525	.589	.621	.796
13	.278	.471	.606	.770	.888	1.001	1.105

- (a) Using a matrix \mathbf{C} of orthogonal contrasts, test the hypothesis of overall equality of means, $H_0: \mathbf{C}\bar{\mu} = \mathbf{0}$, for the combined samples, as in Section 6.10.2.
- (b) If the overall hypothesis is rejected, find the degree of growth curve for the combined samples by testing each row of \mathbf{C} .
- (c) Compare the three groups using the entire matrix \mathbf{C} .
- (d) Compare the three groups using each row of \mathbf{C} .

Table 6.24 Measurements of Trout

Group	Time Point			
	1	2	3	4
1	.257	.288	.328	.358
1	.266	.282	.315	.464
1	.256	.303	.293	.261
1	.272	.456	.288	.261
2	.312	.300	.273	.253
2	.253	.220	.314	.261
2	.239	.261	.279	.224
2	.254	.243	.304	.254
3	.272	.279	.259	.295
3	.246	.292	.279	.302
3	.262	.311	.263	.264
3	.292	.261	.314	.244

Table 6.25 Weekly Gains in Weight for 27 Rats

Rat	Group 1				Group 2				Group 3					
	y_1	y_2	y_3	y_4	Rat	y_1	y_2	y_3	y_4	Rat	y_1	y_2	y_3	y_4
1	29	28	25	33	11	26	36	35	35	18	25	23	11	9
2	33	30	23	31	12	17	19	20	28	19	21	21	10	11
3	25	34	33	41	13	19	33	43	38	20	26	21	6	27
4	18	33	29	35	14	26	31	32	29	21	29	12	11	11
5	25	23	17	30	15	15	25	23	24	22	24	26	22	17
6	24	32	29	22	16	21	24	19	24	23	24	17	8	19
7	20	23	16	31	17	18	35	33	33	24	22	17	8	5
8	28	21	18	24						25	11	24	21	24
9	18	23	22	28						26	15	17	12	17
10	25	28	29	30						27	19	17	15	18

6.40 Table 6.25 contains weight gains for three groups of rats (Box 1950). The variables are y_i = gain in i th week, $i = 1, 2, 3, 4$. The groups are 1 = controls, 2 = thyroxin added to drinking water, and 3 = thiouracil added to drinking water.

- (a) Using a matrix \mathbf{C} of orthogonal contrasts, test the hypothesis of overall equality of means, $H_0: \mathbf{C}\bar{\mu}_e = \mathbf{0}$, for the combined samples, as in Section 6.10.2.
- (b) If the overall hypothesis is rejected, find the degree of growth curve for the combined samples by testing each row of \mathbf{C} .
- (c) Compare the three groups using the entire matrix \mathbf{C} .
- (d) Compare the three groups using each row of \mathbf{C} .

6.41 Table 6.26 contains measurements of coronary sinus potassium at 2-minute intervals after coronary occlusion on four groups of dogs (Grizzle and Allen 1969). The groups are 1 = control dogs, 2 = dogs with extrinsic cardiac denervation 3 weeks prior to coronary occlusion, 3 = dogs with extrinsic cardiac denervation immediately prior to coronary occlusion, and 4 = dogs with bilateral thoracic sympathectomy and stellectomy 3 weeks prior to coronary occlusion.

- (a) Using a matrix \mathbf{C} of orthogonal contrasts, test the hypothesis of overall equality of means, $H_0: \mathbf{C}\bar{\mu}_e = \mathbf{0}$, for the combined samples, as in Section 6.10.2.
- (b) If the overall hypothesis is rejected, find the degree of growth curve for the combined samples by testing each row of \mathbf{C} .
- (c) Compare the four groups using the entire matrix \mathbf{C} .
- (d) Compare the four groups using each row of \mathbf{C} .

Table 6.26 Coronary Sinus Potassium Measured at 2-Minute Intervals on Dogs

Group	Time						
	1	3	5	7	9	11	13
1	4.0	4.0	4.1	3.6	3.6	3.8	3.1
1	4.2	4.3	3.7	3.7	4.8	5.0	5.2
1	4.3	4.2	4.3	4.3	4.5	5.8	5.4
1	4.2	4.4	4.6	4.9	5.3	5.6	4.9
1	4.6	4.4	5.3	5.6	5.9	5.9	5.3
1	3.1	3.6	4.9	5.2	5.3	4.2	4.1
1	3.7	3.9	3.9	4.8	5.2	5.4	4.2
1	4.3	4.2	4.4	5.2	5.6	5.4	4.7
1	4.6	4.6	4.4	4.6	5.4	5.9	5.6
2	3.4	3.4	3.5	3.1	3.1	3.7	3.3
2	3.0	3.2	3.0	3.0	3.1	3.2	3.1
2	3.0	3.1	3.2	3.0	3.3	3.0	3.0
2	3.1	3.2	3.2	3.2	3.3	3.1	3.1
2	3.8	3.9	4.0	2.9	3.5	3.5	3.4
2	3.0	3.6	3.2	3.1	3.0	3.0	3.0
2	3.3	3.3	3.3	3.4	3.6	3.1	3.1
2	4.2	4.0	4.2	4.1	4.2	4.0	4.0
2	4.1	4.2	4.3	4.3	4.2	4.0	4.2
2	4.5	4.4	4.3	4.5	5.3	4.4	4.4
3	3.2	3.3	3.8	3.8	4.4	4.2	3.7
3	3.3	3.4	3.4	3.7	3.7	3.6	3.7
3	3.1	3.3	3.2	3.1	3.2	3.1	3.1
3	3.6	3.4	3.5	4.6	4.9	5.2	4.4
3	4.5	4.5	5.4	5.7	4.9	4.0	4.0
3	3.7	4.0	4.4	4.2	4.6	4.8	5.4
3	3.5	3.9	5.8	5.4	4.9	5.3	5.6
3	3.9	4.0	4.1	5.0	5.4	4.4	3.9
4	3.1	3.5	3.5	3.2	3.0	3.0	3.2
4	3.3	3.2	3.6	3.7	3.7	4.2	4.4
4	3.5	3.9	4.7	4.3	3.9	3.4	3.5
4	3.4	3.4	3.5	3.3	3.4	3.2	3.4
4	3.7	3.8	4.2	4.3	3.6	3.8	3.7
4	4.0	4.6	4.8	4.9	5.4	5.6	4.8
4	4.2	3.9	4.5	4.7	3.9	3.8	3.7
4	4.1	4.1	3.7	4.0	4.1	4.6	4.7
4	3.5	3.6	3.6	4.2	4.8	4.9	5.0

Table 6.27 Blood Pressure Data

Group	Number of Minutes after Ligation					
	1	5	10	15	30	60
1	112.5	100.5	102.5	102.5	107.5	107.5
1	92.5	102.5	105.0	100.0	110.0	117.5
1	132.5	125.0	115.0	112.5	110.0	110.0
1	102.5	107.5	107.5	102.5	90.0	112.5
1	110.0	130.0	115.0	105.0	112.5	110.0
1	97.5	97.5	80.0	82.5	82.5	102.5
1	90.0	70.0	85.0	85.0	92.5	97.5
2	115.0	115.0	107.5	107.5	112.5	107.5
2	125.0	125.0	120.0	120.0	117.5	125.0
2	95.0	90.0	95.0	90.0	100.0	107.5
2	87.5	65.5	85.0	90.0	105.0	90.0
2	90.0	87.5	97.5	95.0	100.0	95.0
2	97.5	92.5	57.5	55.0	90.0	97.5
2	107.5	107.5	145.0	110.0	105.0	112.5
2	102.5	130.0	85.0	80.0	127.5	97.5
3	107.5	107.5	102.5	102.5	102.5	97.5
3	97.5	108.5	94.5	102.5	102.5	107.5
3	100.0	105.0	105.0	105.0	110.0	110.0
3	95.0	95.0	90.0	100.0	100.0	100.0
3	85.0	92.5	92.5	92.5	90.0	110.0
3	82.5	77.5	75.0	65.5	65.0	72.5
3	62.5	75.0	115.0	110.0	100.0	100.0
4	70.0	67.5	67.5	77.5	77.5	77.5
4	45.0	37.5	45.0	45.0	47.5	45.0
4	52.5	22.5	90.0	65.0	60.0	65.5
4	100.0	100.0	100.0	100.0	97.5	92.5
4	115.0	110.0	100.0	110.0	105.0	105.0
4	97.5	97.5	97.5	105.0	95.0	92.5
4	95.0	125.0	130.0	125.0	115.0	117.5
4	72.5	87.5	65.0	57.5	92.5	82.5
4	105.0	105.0	105.0	105.0	102.5	100.0

6.42 Table 6.27 contains blood pressure measurements at intervals after inducing a heart attack for four groups of rats: group 1 is the controls and groups 2–4 have been exposed to halothane concentrations of .25%, .50%, 1.0%, respectively (Crepeau et al. 1985).

- (a) Find the degree of growth curve for the combined sample using the methods in (6.112)–(6.117).
- (b) Repeat (a) for group 1.
- (c) Repeat (a) for groups 2–4 combined.

Table 6.28 Plasma Inorganic Phosphate (mg/dl)

Patient	Hours after Glucose Challenge							
	0	$\frac{1}{2}$	1	$1\frac{1}{2}$	2	3	4	5
Control								
1	4.3	3.3	3.0	2.6	2.2	2.5	3.4	4.4 ^a
2	3.7	2.6	2.6	1.9	2.9	3.2	3.1	3.9
3	4.0	4.1	3.1	2.3	2.9	3.1	3.9	4.0
4	3.6	3.0	2.2	2.8	2.9	3.9	3.8	4.0
5	4.1	3.8	2.1	3.0	3.6	3.4	3.6	3.7
6	3.8	2.2	2.0	2.6	3.8	3.6	3.0	3.5
7	3.8	3.0	2.4	2.5	3.1	3.4	3.5	3.7
8	4.4	3.9	2.8	2.1	3.6	3.8	4.0	3.9
9	5.0	4.0	3.4	3.4	3.3	3.6	4.0	4.3
:	:	:	:	:	:	:	:	:
13	4.7	3.1	3.2	3.3	3.2	4.2	3.7	4.3
Obese								
1	4.3	3.3	3.0	2.6	2.2	2.5	2.4	3.4 ^a
2	5.0	4.9	4.1	3.7	3.7	4.1	4.7	4.9
3	4.6	4.4	3.9	3.9	3.7	4.2	4.8	5.0
4	4.3	3.9	3.1	3.1	3.1	3.1	3.6	4.0
5	3.1	3.1	3.3	2.6	2.6	1.9	2.3	2.7
6	4.8	5.0	2.9	2.8	2.2	3.1	3.5	3.6
7	3.7	3.1	3.3	2.8	2.9	3.6	4.3	4.4
8	5.4	4.7	3.9	4.1	2.8	3.7	3.5	3.7
9	3.0	2.5	2.3	2.2	2.1	2.6	3.2	3.5
:	:	:	:	:	:	:	:	:
20	4.6	4.4	3.8	3.8	3.8	3.6	3.8	3.8

^aThe similarity in the data for patient 1 in the control group and patient 1 in the obese group is coincidental.

- 6.43** Table 6.28 from Zerbe (1979a) compares 13 control and 20 obese patients on a glucose tolerance test using plasma inorganic phosphate. Delete the observations corresponding to $\frac{1}{2}$ and $1\frac{1}{2}$ hours so that the time points are equally spaced.

- (a) For the control group, use orthogonal polynomials to find the degree of growth curve.
- (b) Repeat (a) for the obese group.
- (c) Find the degree of growth curve for the combined groups, and compare the growth curves of the two groups.

Table 6.29 Mandible Measurements

Group	Subject	Activator Treatment								
		1			2			3		
		y_1	y_2	y_3	y_1	y_2	y_3	y_1	y_2	y_3
1	1	117.0	117.5	118.5	59.0	59.0	60.0	10.5	16.5	16.5
	2	109.0	110.5	111.0	60.0	61.5	61.5	30.5	30.5	30.5
	3	117.0	120.0	120.5	60.0	61.5	62.0	23.5	23.5	23.5
	4	122.0	126.0	127.0	67.5	70.5	71.5	33.0	32.0	32.5
	5	116.0	118.5	119.5	61.5	62.5	63.5	24.5	24.5	24.5
	6	123.0	126.0	127.0	65.5	61.5	67.5	22.0	22.0	22.0
	7	130.5	132.0	134.5	68.5	69.5	71.0	33.0	32.5	32.0
	8	126.5	128.5	130.5	69.0	71.0	73.0	20.0	20.0	20.0
	9	113.0	116.5	118.0	58.0	59.0	60.5	25.0	25.0	24.5
2	1	128.0	129.0	131.5	67.0	67.5	69.0	24.0	24.0	24.0
	2	116.5	120.0	121.5	63.5	65.0	66.0	28.5	29.5	29.5
	3	121.5	125.5	127.0	64.5	67.5	69.0	26.5	27.0	27.0
	4	109.5	112.0	114.0	54.0	55.5	57.0	18.0	18.5	19.0
	5	133.0	136.0	137.5	72.0	73.5	75.5	34.5	34.5	34.5
	6	120.0	124.5	126.0	62.5	65.0	66.0	26.0	26.0	26.0
	7	129.5	133.5	134.5	65.0	68.0	69.0	18.5	18.5	18.5
	8	122.0	124.0	125.5	64.5	65.5	66.0	18.5	18.5	18.5
	9	125.0	127.0	128.0	65.5	66.5	67.0	21.5	21.5	21.6

6.44 Consider the complete data from Table 6.28 including the observations corresponding to $\frac{1}{2}$ and $1\frac{1}{2}$ hours. Use the methods in (6.112)–(6.117) for unequally spaced time points to analyze each group separately and the combined groups.

6.45 Table 6.29 contains mandible measurements (Timm 1980). There were two groups of subjects. Each subject was measured at three time points y_1 , y_2 , and y_3 for each of three types of activator treatment. Analyze as a repeated measures design with two within-subjects factors and one between-subjects factor. Use linear and quadratic contrasts for time (growth curve).

CHAPTER 7

TESTS ON COVARIANCE MATRICES

7.1 INTRODUCTION

We now consider tests of hypotheses involving the variance–covariance structure. These tests are often carried out to check assumptions pertaining to other tests. In Sections 7.2–7.4, we cover three basic types of hypotheses: (1) the covariance matrix has a particular structure, (2) two or more covariance matrices are equal, and (3) certain elements of the covariance matrix are zero, thus implying independence of the corresponding (multivariate normal) random variables. In most cases we use the likelihood ratio approach (Section 5.4.3). The resulting test statistics often involve the ratio of the determinants of the sample covariance matrix under the null hypothesis and under the alternative hypothesis.

7.2 TESTING A SPECIFIED PATTERN FOR Σ

In this section, the discussion is in terms of a sample covariance matrix S from a single sample. However, the tests can be applied to a sample covariance matrix $S_{pl} = E/\nu_E$ obtained by pooling across several samples. To allow for either possibility,

the degrees-of-freedom parameter has been indicated by ν . For a single sample, $\nu = n - 1$; for a pooled covariance matrix, $\nu = \sum_{i=1}^k (n_i - 1) = \sum_{i=1}^k n_i - k = N - k$.

7.2.1 Testing $H_0: \Sigma = \Sigma_0$

We begin with the basic hypothesis $H_0: \Sigma = \Sigma_0$ versus $H_1: \Sigma \neq \Sigma_0$. The hypothesized covariance matrix Σ_0 is a target value for Σ or a nominal value from previous experience. Note that Σ_0 is completely specified in H_0 , whereas μ is not specified.

To test H_0 , we obtain a random sample of n observation vectors y_1, y_2, \dots, y_n from $N_p(\mu, \Sigma)$ and calculate S . To see if S is significantly different from Σ_0 , we use the following test statistic, which is a modification of the likelihood ratio (Section 5.4.3):

$$u = \nu[\ln |\Sigma_0| - \ln |S| + \text{tr}(S\Sigma_0^{-1}) - p], \quad (7.1)$$

where ν represents the degrees of freedom of S (see comments at the beginning of Section 7.2), \ln is the natural logarithm (base e), and tr is the trace of a matrix (Section 2.9). Note that if $S = \Sigma_0$, then $u = 0$; otherwise u increases with the "distance" between S and Σ_0 [see (7.4) and the comment following].

When ν is large, the statistic u in (7.1) is approximately distributed as $\chi^2[\frac{1}{2}p(p+1)]$ if H_0 is true. For moderate size ν ,

$$u' = \left[1 - \frac{1}{6\nu - 1} \left(2p + 1 - \frac{2}{p+1} \right) \right] u \quad (7.2)$$

is a better approximation to the $\chi^2[\frac{1}{2}p(p+1)]$ distribution. We reject H_0 if u or u' is greater than $\chi^2[\alpha, \frac{1}{2}p(p+1)]$. Note that the degrees of freedom for the χ^2 -statistic, $\frac{1}{2}p(p+1)$, is the number of distinct parameters in Σ .

We can express u in terms of the eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_p$ of $S\Sigma_0^{-1}$ by noting that $\text{tr}(S\Sigma_0^{-1})$ and $\ln |\Sigma_0| - \ln |S|$ become

$$\begin{aligned} \text{tr}(S\Sigma_0^{-1}) &= \sum_{i=1}^p \lambda_i && [\text{by (2.107)}], \\ \ln |\Sigma_0| - \ln |S| &= -\ln |\Sigma_0|^{-1} - \ln |S| \\ &= -\ln |S\Sigma_0^{-1}| && [\text{by (2.89) and (2.91)}] \\ &= -\ln \left(\prod_{i=1}^p \lambda_i \right) && [\text{by (2.108)}], \end{aligned} \quad (7.3)$$

from which (7.1) can be written as

$$\begin{aligned} u &= \nu \left[-\ln \left(\prod_{i=1}^p \lambda_i \right) + \sum_{i=1}^p \lambda_i - p \right] \\ &= \nu \left[\sum_{i=1}^p (\lambda_i - \ln \lambda_i) - p \right]. \end{aligned} \quad (7.4)$$

A plot of $y = x - \ln x$ will show that $x - \ln x \geq 1$ for all $x > 0$, with equality holding only for $x = 1$. Thus $\sum_{i=1}^p (\lambda_i - \ln \lambda_i) > p$ and $u > 0$.

The hypothesis that the variables are independent and have unit variance,

$$H_0: \Sigma = \mathbf{I},$$

can be tested by simply setting $\Sigma_0 = \mathbf{I}$ in (7.1).

7.2.2 Testing Sphericity

The hypothesis that the variables y_1, y_2, \dots, y_p in \mathbf{y} are independent and have the same variance can be expressed as $H_0: \Sigma = \sigma^2 \mathbf{I}$ versus $H_1: \Sigma \neq \sigma^2 \mathbf{I}$, where σ^2 is the unknown common variance. This hypothesis is of interest in repeated measures (see Section 6.9.1). Under H_0 , the ellipsoid $(\mathbf{y} - \boldsymbol{\mu})' \Sigma^{-1} (\mathbf{y} - \boldsymbol{\mu}) = c^2$ reduces to $(\mathbf{y} - \boldsymbol{\mu})' (\mathbf{y} - \boldsymbol{\mu}) = \sigma^2 c^2$, the equation of a sphere; hence the term *sphericity* is applied to the covariance structure $\Sigma = \sigma^2 \mathbf{I}$. Another sphericity hypothesis of interest in repeated measures is $H_0: \mathbf{C} \Sigma \mathbf{C}' = \sigma^2 \mathbf{I}$, where \mathbf{C} is any full-rank $(p-1) \times p$ matrix of orthonormal contrasts (see Section 6.9.1).

For a random sample $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n$ from $N_p(\boldsymbol{\mu}, \Sigma)$, the likelihood ratio for testing $H_0: \Sigma = \sigma^2 \mathbf{I}$ is

$$\text{LR} = \left[\frac{|\mathbf{S}|}{(\text{tr}\mathbf{S}/p)^p} \right]^{n/2}. \quad (7.5)$$

In some cases that we have considered previously, the likelihood ratio is a simple function of a test statistic such as F , T^2 , Wilks' Λ , and so on. However, LR in (7.5) does not reduce to a standard statistic, and we resort to an approximation for its distribution. It has been shown that for a general likelihood ratio statistic LR,

$$-2 \ln(\text{LR}) \text{ is approximately } \chi_\nu^2 \quad (7.6)$$

for large n , where ν is the total number of parameters minus the number estimated under the restrictions imposed by H_0 .

For the likelihood ratio statistic in (7.5), we obtain

$$-2 \ln(\text{LR}) = -n \ln \left[\frac{|\mathbf{S}|}{(\text{tr}\mathbf{S}/p)^p} \right] = -n \ln u,$$

where

$$u = (\text{LR})^{2/n} = \frac{p^p |\mathbf{S}|}{(\text{tr}\mathbf{S})^p}. \quad (7.7)$$

By (2.107) and (2.108), u becomes

$$u = \frac{p^p \prod_{i=1}^p \lambda_i}{(\sum_{i=1}^p \lambda_i)^p}, \quad (7.8)$$

where $\lambda_1, \lambda_2, \dots, \lambda_p$ are the eigenvalues of \mathbf{S} . An improvement over $-n \ln u$ is given by

$$u' = - \left(\nu - \frac{2p^2 + p + 2}{6p} \right) \ln u, \quad (7.9)$$

where ν is the degrees of freedom for \mathbf{S} (see comments at the beginning of Section 7.2). The statistic u' has an approximate χ^2 -distribution with $\frac{1}{2}p(p+1)-1$ degrees of freedom. We reject H_0 if $u' \geq \chi^2[\alpha, \frac{1}{2}p(p+1)-1]$. As noted above, the degrees of freedom in the χ^2 approximation is equal to the total number of parameters minus the number of parameters estimated under H_0 . The number of parameters in Σ is $p + \binom{p}{2} = \frac{1}{2}p(p+1)$ and the loss of one degree of freedom is due to estimation of σ^2 .

We see from (7.8) and (7.9) that if the sample λ_i 's are all equal, $u = 1$ and $u' = 0$. Hence, this statistic also tests the hypothesis of equality of the population eigenvalues.

To test $H_0: \mathbf{C}\Sigma\mathbf{C}' = \sigma^2\mathbf{I}$, use \mathbf{CSC}' in place of \mathbf{S} in (7.7) and use $p-1$ in place of p in (7.7)–(7.9) and in the degrees of freedom for χ^2 .

The likelihood ratio (7.5) was first given by Mauchly (1940), and his name is often associated with this test. Nagarsenker and Pillai (1973) gave the exact distribution of u and provided a table for $p = 4, 5, \dots, 10$. Venables (1976) showed that u can be obtained by a union–intersection approach (Section 6.1.4).

■ EXAMPLE 7.2.2

We use the probe word data in Table 3.6 to illustrate tests of sphericity. The five variables appear to be commensurate, and the hypothesis $H_0: \mu_1 = \mu_2 = \dots = \mu_5$ may be of interest. We would expect the variables to be correlated, and H_0 would ordinarily be tested using a multivariate approach, as in Sections 5.9.1 and 6.9.2. However, if $\Sigma = \sigma^2\mathbf{I}$ or $\mathbf{C}\Sigma\mathbf{C}' = \sigma^2\mathbf{I}$, then the hypothesis $H_0: \mu_1 = \mu_2 = \dots = \mu_5$ can be tested with a univariate ANOVA F -test (see Section 6.9.1).

We first test $H_0: \Sigma = \sigma^2\mathbf{I}$. The sample covariance matrix \mathbf{S} was obtained in Example 3.10.1. By (7.7),

$$u = \frac{p^p |\mathbf{S}|}{(\text{tr}\mathbf{S})^p} = \frac{5^5(27,236,586)}{(292.891)^5} = .0395.$$

Then by (7.9), with $n = 11$ and $p = 5$, we have

$$u' = -\left(n-1 - \frac{2p^2 + p + 2}{6p}\right) \ln u = 26.177.$$

The approximate χ^2 -test has $\frac{1}{2}p(p+1)-1 = 14$ degrees of freedom. We therefore compare $u' = 26.177$ with $\chi^2_{.05,14} = 23.68$ and reject $H_0: \Sigma = \sigma^2\mathbf{I}$.

To test $H_0: \mathbf{C}\Sigma\mathbf{C}' = \sigma^2\mathbf{I}$, we use the following matrix of orthonormalized contrasts:

$$\mathbf{C} = \begin{pmatrix} 4/\sqrt{20} & -1/\sqrt{20} & -1/\sqrt{20} & -1/\sqrt{20} & -1/\sqrt{20} \\ 0 & 3/\sqrt{12} & -1/\sqrt{12} & -1/\sqrt{12} & -1/\sqrt{12} \\ 0 & 0 & 2/\sqrt{6} & -1/\sqrt{6} & -1/\sqrt{6} \\ 0 & 0 & 0 & 1/\sqrt{2} & -1/\sqrt{2} \end{pmatrix}.$$

Then using $\mathbf{C}\mathbf{S}\mathbf{C}'$ in place of \mathbf{S} and with $p - 1 = 4$ for the four rows of \mathbf{C} , we obtain

$$u = \frac{(p-1)^{p-1}|\mathbf{C}\mathbf{S}\mathbf{C}'|}{[\text{tr}(\mathbf{C}\mathbf{S}\mathbf{C}')]^{p-1}} = \frac{4^4(144039.8)}{(93.6)^4} = .480,$$

$$u' = 6.170.$$

For degrees of freedom, we now have $\frac{1}{2}(4)(5) - 1 = 9$, and the critical value is $\chi^2_{0.05,9} = 16.92$. Hence, we do not reject $H_0: \mathbf{C}\Sigma\mathbf{C}' = \sigma^2\mathbf{I}$, and a univariate F -test of $H_0: \mu_1 = \mu_2 = \dots = \mu_5$ may be justified. \square

7.2.3 Testing $H_0: \Sigma = \sigma^2[(1 - \rho)\mathbf{I} + \rho\mathbf{J}]$

In Section 6.9.1, it was noted that univariate ANOVA remains valid if

$$\Sigma = \sigma^2 \begin{pmatrix} 1 & \rho & \rho & \dots & \rho \\ \rho & 1 & \rho & \dots & \rho \\ \vdots & \vdots & \vdots & & \vdots \\ \rho & \rho & \rho & \dots & 1 \end{pmatrix} \quad (7.10)$$

$$= \sigma^2[(1 - \rho)\mathbf{I} + \rho\mathbf{J}], \quad (7.11)$$

where \mathbf{J} is a square matrix of 1's, as defined in (2.12), and ρ is the population correlation between any two variables. This pattern of equal variances and equal covariances in Σ is variously referred to as *uniformity*, *compound symmetry*, or the *intraclass correlation model*.

We now consider the hypothesis that (7.10) holds:

$$H_0: \Sigma = \begin{pmatrix} \sigma^2 & \sigma^2\rho & \dots & \sigma^2\rho \\ \sigma^2\rho & \sigma^2 & \dots & \sigma^2\rho \\ \vdots & \vdots & & \vdots \\ \sigma^2\rho & \sigma^2\rho & \dots & \sigma^2 \end{pmatrix}.$$

From a sample we obtain the sample covariance matrix \mathbf{S} . Estimates of σ^2 and $\sigma^2\rho$ under H_0 are given by

$$s^2 = \frac{1}{p} \sum_{j=1}^p s_{jj} \quad \text{and} \quad s^2r = \frac{1}{p(p-1)} \sum_{j \neq k} s_{jk}, \quad (7.12)$$

respectively, where s_{jj} and s_{jk} are from \mathbf{S} . Thus s^2 is an average of the variances on the diagonal of \mathbf{S} and s^2r is an average of the off-diagonal covariances in \mathbf{S} . An estimate of ρ can be obtained as $r = s^2r/s^2$. Using s^2 and s^2r in (7.12), the estimate of Σ under H_0 is then

$$\mathbf{S}_0 = \begin{pmatrix} s^2 & s^2r & \dots & s^2r \\ s^2r & s^2 & \dots & s^2r \\ \vdots & \vdots & & \vdots \\ s^2r & s^2r & \dots & s^2 \end{pmatrix} = s^2[(1 - r)\mathbf{I} + r\mathbf{J}]. \quad (7.13)$$

To compare \mathbf{S} and \mathbf{S}_0 , we use the following function of the likelihood ratio:

$$u = \frac{|\mathbf{S}|}{|\mathbf{S}_0|}, \quad (7.14)$$

which can be expressed in the alternative form

$$u = \frac{|\mathbf{S}|}{(s^2)^p (1-r)^{p-1} [1 + (p-1)r]}. \quad (7.15)$$

By analogy with (7.9), the test statistic is given by

$$u' = - \left[\nu - \frac{p(p+1)^2(2p-3)}{6(p-1)(p^2+p-4)} \right] \ln u, \quad (7.16)$$

where ν is the degrees of freedom of \mathbf{S} (see comments at the beginning of Section 7.2). The statistic u' is approximately $\chi^2[\frac{1}{2}p(p+1)-2]$, and we reject H_0 if $u' > \chi^2[\alpha, \frac{1}{2}p(p+1)-2]$. Note that two degrees of freedom are lost due to estimation of σ^2 and ρ .

An alternative approximate test that is more precise when p is large and ν is relatively small is given by

$$F = \frac{-(\gamma_2 - \gamma_2 c_1 - \gamma_1)\nu}{\gamma_1 \gamma_2} \ln u,$$

where

$$\begin{aligned} c_1 &= \frac{p(p+1)^2(2p-3)}{6\nu(p-1)(p^2+p-4)}, & c_2 &= \frac{p(p^2-1)(p+2)}{6\nu^2(p^2+p-4)}, \\ \gamma_1 &= \frac{1}{2}p(p+1)-2, & \gamma_2 &= \frac{\gamma_1+2}{c_2 - c_1^2}. \end{aligned}$$

We reject $H_0 : \Sigma = \sigma^2[(1-\rho)\mathbf{I} + \rho\mathbf{J}]$ if $F > F_{\alpha, \gamma_1, \gamma_2}$.

■ EXAMPLE 7.2.3

To illustrate this test, we use the cork data of Table 6.21. In Problem 6.34, a comparison is made of average thickness, and hence weight, in the four directions. A standard ANOVA approach to this repeated measures design would be valid if (7.10) holds. To check this assumption, we test $H_0 : \Sigma = \sigma^2[(1-\rho)\mathbf{I} + \rho\mathbf{J}]$. The sample covariance matrix is given by

$$\mathbf{S} = \begin{pmatrix} 290.41 & 223.75 & 288.44 & 226.27 \\ 223.75 & 219.93 & 229.06 & 171.37 \\ 288.44 & 229.06 & 350.00 & 259.54 \\ 226.27 & 171.37 & 259.54 & 226.00 \end{pmatrix},$$

from which we obtain

$$|\mathbf{S}| = 25,617,563.28, \quad s^2 = \frac{1}{p} \sum_{j=1}^p s_{jj} = 271.586,$$

$$s^2 r = \frac{1}{p(p-1)} \sum_{j \neq k} s_{jk} = 233.072, \quad r = \frac{s^2 r}{s^2} = \frac{233.072}{271.586} = .858.$$

From (7.15) and (7.16), we now have

$$u = \frac{25,617,563.28}{(271.586)^4 (1 - .858)^3 [1 + (3)(.858)]} = .461,$$

$$u' = - \left[\nu - \frac{p(p+1)^2(2p-3)}{6(p-1)(p^2+p-4)} \right] \ln u$$

$$= \left[27 - \frac{(4)(25)(5)}{(6)(3)(16)} \right] 0.774 = 19.511.$$

Since $19.511 > \chi^2_{0.05,8} = 15.5$, we reject H_0 and conclude that Σ does not have the pattern in (7.10). \square

7.3 TESTS COMPARING COVARIANCE MATRICES

An assumption for T^2 or MANOVA tests comparing two or more mean vectors is that the corresponding population covariance matrices are equal: $\Sigma_1 = \Sigma_2 = \dots = \Sigma_k$. Under this assumption, the sample covariance matrices $\mathbf{S}_1, \mathbf{S}_2, \dots, \mathbf{S}_k$ reflect a common population Σ and are therefore pooled to obtain an estimate of Σ . If $\Sigma_1 = \Sigma_2 = \dots = \Sigma_k$ is not true, large differences in $\mathbf{S}_1, \mathbf{S}_2, \dots, \mathbf{S}_k$ may possibly lead to rejection of $H_0: \mu_1 = \mu_2 = \dots = \mu_k$. However, the T^2 and MANOVA tests are fairly robust to heterogeneity of covariance matrices as long as the sample sizes are large and equal. For other cases it is useful to have available a test of equality of covariance matrices. We begin with a review of the univariate case.

7.3.1 Univariate Tests of Equality of Variances

The two-sample univariate hypothesis $H_0: \sigma_1^2 = \sigma_2^2$ versus $H_1: \sigma_1^2 \neq \sigma_2^2$ is tested with

$$F = \frac{s_1^2}{s_2^2}, \tag{7.17}$$

where s_1^2 and s_2^2 are the variances of the two samples. If H_0 is true, F is distributed as F_{ν_1, ν_2} , where ν_1 and ν_2 are the degrees of freedom of s_1^2 and s_2^2 (typically, $n_1 - 1$ and $n_2 - 1$). Note that s_1^2 and s_2^2 must be independent, which will hold if the two samples are independent.

For the several-sample case, various procedures have been proposed. We present Bartlett's (1937) test of homogeneity of variances because it has been extended to

the multivariate case. To test

$$H_0: \sigma_1^2 = \sigma_2^2 = \cdots = \sigma_k^2,$$

we calculate

$$c = 1 + \frac{1}{3(k-1)} \left[\sum_{i=1}^k \frac{1}{\nu_i} - \frac{1}{\sum_{i=1}^k \nu_i} \right], \quad s^2 = \frac{\sum_{i=1}^k \nu_i s_i^2}{\sum_{i=1}^k \nu_i},$$

$$m = \left(\sum_{i=1}^k \nu_i \right) \ln s^2 - \sum_{i=1}^k \nu_i \ln s_i^2,$$

where $s_1^2, s_2^2, \dots, s_k^2$ are independent sample variances with $\nu_1, \nu_2, \dots, \nu_k$ degrees of freedom, respectively. Then

$$\frac{m}{c} \text{ is approximately } \chi_{k-1}^2.$$

We reject H_0 if $m/c > \chi_{\alpha, k-1}^2$.

For an F -approximation, we use c and m above and calculate in addition

$$a_1 = k-1, \quad a_2 = \frac{k+1}{(c-1)^2}, \quad b = \frac{a_2}{2-c+2/a_2}.$$

Then

$$F = \frac{a_2 m}{a_1(b-m)} \text{ is approximately } F_{a_1, a_2}.$$

We reject H_0 if $F > F_\alpha$.

Note that an assumption for either form of the above test is independence of $s_1^2, s_2^2, \dots, s_k^2$, which will hold for random samples from k distinct populations. This test would therefore be inappropriate for comparing $s_{11}, s_{22}, \dots, s_{pp}$ from the diagonal of \mathbf{S} , since the s_{jj} 's are correlated.

7.3.2 Multivariate Tests of Equality of Covariance Matrices

For k multivariate populations, the hypothesis of equality of covariance matrices is

$$H_0: \Sigma_1 = \Sigma_2 = \cdots = \Sigma_k. \tag{7.18}$$

The test of $H_0: \Sigma_1 = \Sigma_2$ for two groups is easily obtained as a special case by setting $k = 2$. Thus there is no exact test as there is in the analogous univariate case [see (7.17)]. We assume independent samples of size n_1, n_2, \dots, n_k from multivariate normal distributions, as in an unbalanced one-way MANOVA, for example. To make the test, we calculate

$$M = \frac{|\mathbf{S}_1|^{\nu_1/2} |\mathbf{S}_2|^{\nu_2/2} \cdots |\mathbf{S}_k|^{\nu_k/2}}{|\mathbf{S}_{\text{pl}}|^{\sum_i \nu_i/2}}, \tag{7.19}$$

in which $\nu_i = n_i - 1$, \mathbf{S}_i is the covariance matrix of the i th sample, and \mathbf{S}_{pl} is the pooled sample covariance matrix

$$\mathbf{S}_{\text{pl}} = \frac{\sum_{i=1}^k \nu_i \mathbf{S}_i}{\sum_{i=1}^k \nu_i} = \frac{\mathbf{E}}{\nu_E}, \quad (7.20)$$

where \mathbf{E} is given by (6.32) and $\nu_E = \sum_{i=1}^k \nu_i = \sum_i n_i - k$. It is clear that we must have every $\nu_i > p$; otherwise M would be zero because $|\mathbf{S}_i| = 0$ for some i . Exact upper percentage points of $-2 \ln M = \nu(k \ln |\mathbf{S}_{\text{pl}}| - \sum_i \ln |\mathbf{S}_i|)$ for the special case of $\nu_1 = \nu_2 = \dots = \nu_k = \nu$ are given in Table A.14 for $p = 2, 3, 4, 5$ and $k = 2, 3, \dots, 10$ (Lee et al. 1977). We can easily modify (7.19) and (7.20) to compare covariance matrices for the cells of a two-way model using $\nu_{ij} = n_{ij} - 1$.

The statistic M is a modification of the likelihood ratio and varies between 0 and 1, with values near 1 favoring H_0 in (7.18) and values near 0 leading to rejection of H_0 . It is not immediately obvious that M in (7.19) behaves in this way, and we offer the following heuristic argument. First we note that (7.19) can be expressed as

$$M = \left(\frac{|\mathbf{S}_1|}{|\mathbf{S}_{\text{pl}}|} \right)^{\nu_1/2} \left(\frac{|\mathbf{S}_2|}{|\mathbf{S}_{\text{pl}}|} \right)^{\nu_2/2} \cdots \left(\frac{|\mathbf{S}_k|}{|\mathbf{S}_{\text{pl}}|} \right)^{\nu_k/2}. \quad (7.21)$$

If $\mathbf{S}_1 = \mathbf{S}_2 = \dots = \mathbf{S}_k = \mathbf{S}_{\text{pl}}$, then $M = 1$. As the disparity among $\mathbf{S}_1, \mathbf{S}_2, \dots, \mathbf{S}_k$ increases, M approaches zero. To see this, note that the determinant of the pooled covariance matrix, $|\mathbf{S}_{\text{pl}}|$, lies somewhere near the “middle” of the $|\mathbf{S}_i|$ ’s and that as a set of variables z_1, z_2, \dots, z_n increases in spread, $z_{(1)}/\bar{z}$ reduces the product more than $z_{(n)}/\bar{z}$ increases it, where $z_{(1)}$ and $z_{(n)}$ are the minimum and maximum values, respectively. We illustrate this with the two sets of numbers 4, 5, 6 and 1, 5, 9, which have the same mean but different spread. If we assume $\nu_1 = \nu_2 = \nu_3 = \nu$, then for the first set,

$$M_1 = \left[\left(\frac{4}{5} \right) \left(\frac{5}{5} \right) \left(\frac{6}{5} \right) \right]^{\nu/2} = [(0.8)(1)(1.2)]^{\nu/2} = (.96)^{\nu/2}$$

and for the second set,

$$M_2 = \left[\left(\frac{1}{5} \right) \left(\frac{5}{5} \right) \left(\frac{9}{5} \right) \right]^{\nu/2} = [(0.2)(1)(1.8)]^{\nu/2} = (.36)^{\nu/2}.$$

In M_2 , the smallest value, .2, reduces the product proportionally more than the largest value, 1.8, increases it. Another illustration is found in Problem 7.9.

Box (1949, 1950) gave χ^2 - and F -approximations for the distribution of M . Either of these approximate tests is referred to as *Box’s M-test*. For the χ^2 -approximation, calculate

$$c_1 = \left[\sum_{i=1}^k \frac{1}{\nu_i} - \frac{1}{\sum_{i=1}^k \nu_i} \right] \left[\frac{2p^2 + 3p - 1}{6(p+1)(k-1)} \right]. \quad (7.22)$$

Then

$$u = -2(1 - c_1) \ln M \text{ is approximately } \chi^2 \left[\frac{1}{2}(k-1)p(p+1) \right], \quad (7.23)$$

where M is defined in (7.19), and

$$\ln M = \frac{1}{2} \sum_{i=1}^k \nu_i \ln |\mathbf{S}_i| - \frac{1}{2} \left(\sum_{i=1}^k \nu_i \right) \ln |\mathbf{S}_{\text{pl}}|. \quad (7.24)$$

We reject H_0 if $u > \chi^2_\alpha$. If $\nu_1 = \nu_2 = \dots = \nu_k = \nu$, then c_1 simplifies to

$$c_1 = \frac{(k+1)(2p^2 + 3p - 1)}{6k\nu(p+1)}. \quad (7.25)$$

To justify the degrees of freedom of the χ^2 -approximation, note that the total number of parameters estimated under H_1 is $k[\frac{1}{2}p(p+1)]$, whereas under H_0 we estimate only Σ , which has $p + \binom{p}{2} = \frac{1}{2}p(p+1)$ parameters. The difference is $(k-1)[\frac{1}{2}p(p+1)]$. The quantity $k[\frac{1}{2}p(p+1)]$ arises from the assumption that all Σ_i , $i = 1, 2, \dots, k$, are different. Technically, H_1 can be stated as $\Sigma_i \neq \Sigma_j$ for some $i \neq j$. However, the most general case is all Σ_i different, and the distribution of M is derived accordingly.

For the F -approximation, we use c_1 from (7.22) and calculate, additionally,

$$c_2 = \frac{(p-1)(p+2)}{6(k-1)} \left[\sum_{i=1}^k \frac{1}{\nu_i^2} - \frac{1}{\left(\sum_{i=1}^k \nu_i \right)^2} \right], \quad (7.26)$$

$$a_1 = \frac{1}{2}(k-1)p(p+1), \quad a_2 = \frac{a_1 + 2}{|c_2 - c_1^2|},$$

$$b_1 = \frac{1 - c_1 - a_1/a_2}{a_1}, \quad b_2 = \frac{1 - c_1 - 2/a_2}{a_2}.$$

If $c_2 > c_1^2$,

$$F = -2b_1 \ln M \quad \text{is approximately } F_{a_1, a_2}. \quad (7.27)$$

If $c_2 < c_1^2$,

$$F = -\frac{a_2 b_2 \ln M}{a_1(1 + 2b_2 \ln M)} \quad \text{is approximately } F_{a_1, a_2}. \quad (7.28)$$

In either case, we reject H_0 if $F > F_\alpha$. If $\nu_1 = \nu_2 = \dots = \nu_k = \nu$, then c_1 simplifies as in (7.25) and c_2 simplifies to

$$c_2 = \frac{(p-1)(p+2)(k^2 + k + 1)}{6k^2\nu^2}. \quad (7.29)$$

Box's M -test is calculated routinely in many computer programs for MANOVA. However, Olson (1974) showed that the M -test with equal ν_i may detect some forms of heterogeneity that have only minor effects on the MANOVA tests. The test is also sensitive to some forms of nonnormality. For example, it is sensitive to kurtosis, for which the MANOVA tests are rather robust. Thus the M -test may signal covariance heterogeneity in some cases where it is not damaging to the MANOVA tests. Hence we may not wish to automatically rule out standard MANOVA tests if the M -test leads to rejection of H_0 . Olson showed that the skewness and kurtosis statistics $b_{1,p}$ and $b_{2,p}$ (see Section 4.4.2) have similar shortcomings.

■ EXAMPLE 7.3.2

We test the hypothesis $H_0: \Sigma_1 = \Sigma_2$ for the psychological data of Table 5.1. The covariance matrices \mathbf{S}_1 , \mathbf{S}_2 , and \mathbf{S}_{pl} were given in Example 5.4.2. Using these, we obtain, by (7.24),

$$\begin{aligned}\ln M &= \frac{1}{2}[\nu_1 \ln |\mathbf{S}_1| + \nu_2 \ln |\mathbf{S}_2|] - \frac{1}{2}(\nu_1 + \nu_2) \ln |\mathbf{S}_{\text{pl}}| \\ &= \frac{1}{2}[(31) \ln(7917.7) + (31) \ln(58958.1)] \\ &\quad - \frac{1}{2}(31 + 31) \ln(27325.2) = -7.2803.\end{aligned}$$

For an exact test, we compare

$$-2 \ln M = 14.561$$

with 19.74, its critical value from Table A.14.

For the χ^2 -approximation, we compute, by (7.25) and (7.23),

$$\begin{aligned}c_1 &= \frac{(2+1)[2(4^2) + 3(4) - 1]}{6(2)(31)(4+1)} = .06935, \\ u &= -2(1 - c_1) \ln M = 13.551 < \chi^2_{.05,10} = 18.307.\end{aligned}$$

For an approximate F -test, we first calculate the following:

$$\begin{aligned}c_2 &= \frac{(4-1)(4+2)}{6(2-1)} \left[\frac{1}{31^2} + \frac{1}{31^2} - \frac{1}{(31+31)^2} \right] = .005463 \\ a_1 &= \frac{1}{2}(2-1)(4)(4+1) = 10 \\ a_2 &= \frac{10+2}{|.005463 - .06935^2|} = 18377.7 \\ b_1 &= \frac{1 - .06935 - 10/18377.7}{10} = .0930 \\ b_2 &= \frac{1 - .06935 - 2/18377.7}{18377.7} = 5.0634 \times 10^{-5}.\end{aligned}$$

Since $c_2 = .005463 > c_1^2 = .00481$, we use (7.27) to obtain

$$F = -2b_1 \ln M = 1.354 < F_{.05,10,\infty} = 1.83.$$

Thus all three tests accept H_0 . □

7.4 TESTS OF INDEPENDENCE

7.4.1 Independence of Two Subvectors

Suppose the observation vector is partitioned into two subvectors of interest, which we label \mathbf{y} and \mathbf{x} , as in Section 3.9.1, where \mathbf{y} is $p \times 1$ and \mathbf{x} is $q \times 1$. By (3.46), the

corresponding partitioning of the population covariance matrix is

$$\Sigma = \begin{pmatrix} \Sigma_{yy} & \Sigma_{yx} \\ \Sigma_{xy} & \Sigma_{xx} \end{pmatrix},$$

with analogous partitioning of \mathbf{S} and \mathbf{R} as in (3.42):

$$\mathbf{S} = \begin{pmatrix} \mathbf{S}_{yy} & \mathbf{S}_{yx} \\ \mathbf{S}_{xy} & \mathbf{S}_{xx} \end{pmatrix}, \quad \mathbf{R} = \begin{pmatrix} \mathbf{R}_{yy} & \mathbf{R}_{yx} \\ \mathbf{R}_{xy} & \mathbf{R}_{xx} \end{pmatrix}.$$

The hypothesis of independence of \mathbf{y} and \mathbf{x} can be expressed as

$$H_0: \Sigma = \begin{pmatrix} \Sigma_{yy} & \mathbf{O} \\ \mathbf{O} & \Sigma_{xx} \end{pmatrix} \quad \text{or} \quad H_0: \Sigma_{yx} = \mathbf{O}.$$

Thus independence of \mathbf{y} and \mathbf{x} means that every variable in \mathbf{y} is independent of every variable in \mathbf{x} . Note that there is no restriction on Σ_{yy} or Σ_{xx} .

The likelihood ratio test statistic for $H_0: \Sigma_{yx} = \mathbf{O}$ is given by

$$\Lambda = \frac{|\mathbf{S}|}{|\mathbf{S}_{yy}||\mathbf{S}_{xx}|} = \frac{|\mathbf{R}|}{|\mathbf{R}_{yy}||\mathbf{R}_{xx}|}, \quad (7.30)$$

which is distributed as $\Lambda_{p,q,n-1-q}$. We reject H_0 if $\Lambda \leq \Lambda_\alpha$. We thus have an exact test for $H_0: \Sigma_{yx} = \mathbf{O}$. Critical values for Wilks' Λ are given in Table A.9 using $\nu_H = q$ and $\nu_E = n - 1 - q$. The test statistic in (7.30) is equivalent (when H_0 is true) to the Λ -statistic (10.64) in Section 10.5.1 for testing the significance of the regression of \mathbf{y} on \mathbf{x} .

By the symmetry of

$$\frac{|\mathbf{S}|}{|\mathbf{S}_{yy}||\mathbf{S}_{xx}|} = \frac{|\mathbf{S}|}{|\mathbf{S}_{xx}||\mathbf{S}_{yy}|},$$

Λ in (7.30) is also distributed as $\Lambda_{q,p,n-1-p}$. This is equivalent to property 3 in Section 6.1.3.

Note that $|\mathbf{S}_{yy}||\mathbf{S}_{xx}|$ in (7.30) is an estimate of $|\Sigma_{yy}||\Sigma_{xx}|$, which by (2.92) is the determinant of Σ when $\Sigma_{yx} = \mathbf{O}$. Thus Wilks' Λ compares an estimate of Σ without restrictions to an estimate of Σ under $H_0: \Sigma_{yx} = \mathbf{O}$. We can see intuitively that $|\mathbf{S}| < |\mathbf{S}_{yy}||\mathbf{S}_{xx}|$ by noting from (2.94) that $|\mathbf{S}| = |\mathbf{S}_{xx}||\mathbf{S}_{yy} - \mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}|$, and since $\mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}$ is positive definite, $|\mathbf{S}_{yy} - \mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}| < |\mathbf{S}_{yy}|$. This can be illustrated for the case $p = q = 1$:

$$|\mathbf{S}| = \begin{vmatrix} s_y^2 & s_{yx} \\ s_{yx} & s_x^2 \end{vmatrix} = s_y^2 s_x^2 - (s_{yx})^2 < s_y^2 s_x^2.$$

As s_{yx}^2 increases, $|\mathbf{S}|$ decreases.

Wilks' Λ in (7.30) can be expressed in terms of eigenvalues:

$$\Lambda = \prod_{i=1}^s (1 - r_i^2), \quad (7.31)$$

where $s = \min(p, q)$ and the r_i^2 's are the nonzero eigenvalues of $\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}\mathbf{S}_{yy}^{-1}\mathbf{S}_{yx}$. We could also use $\mathbf{S}_{yy}^{-1}\mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}$, since the (nonzero) eigenvalues of $\mathbf{S}_{yy}^{-1}\mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}$ are the same as those of $\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}\mathbf{S}_{yy}^{-1}\mathbf{S}_{yx}$ (these two matrices are of the form AB and BA ; see Section 2.11.5). The number of nonzero eigenvalues is $s = \min(p, q)$, since s is the rank of both $\mathbf{S}_{yy}^{-1}\mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}$ and $\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}\mathbf{S}_{yy}^{-1}\mathbf{S}_{yx}$. The eigenvalues are designated r_i^2 because they are the squared *canonical correlations* between \mathbf{y} and \mathbf{x} (see Chapter 11). In the special case $p = 1$, (7.31) becomes

$$\Lambda = 1 - r_1^2 = 1 - R^2,$$

where R^2 is the square of the multiple correlation between y and (x_1, x_2, \dots, x_q) .

The other test statistics, $U^{(s)}$, $V^{(s)}$, and Roy's θ , can also be defined in terms of the r_i^2 's (see Section 11.4.1).

■ EXAMPLE 7.4.1

Consider the diabetes data in Table 3.5. There is a natural partitioning in the variables, with y_1 and y_2 of minor interest and x_1, x_2 , and x_3 of major interest. We test independence of the y 's and the x 's, that is, $H_0: \Sigma_{yx} = \mathbf{O}$. From Example 3.9.1, the partitioned covariance matrix is

$$\mathbf{S} = \left(\begin{array}{cc} \mathbf{S}_{yy} & \mathbf{S}_{yx} \\ \mathbf{S}_{xy} & \mathbf{S}_{xx} \end{array} \right) = \left(\begin{array}{cc|cc|c} .0162 & .2160 & .7872 & -.2138 & 2.189 \\ .2160 & 70.56 & 26.23 & -23.96 & -20.84 \\ \hline .7872 & 26.23 & 1106 & 396.7 & 108.4 \\ -.2138 & -23.96 & 396.7 & 2382 & 1143 \\ 2.189 & -20.84 & 108.4 & 1143 & 2136 \end{array} \right).$$

To make the test, we compute

$$\Lambda = \frac{|\mathbf{S}|}{|\mathbf{S}_{yy}||\mathbf{S}_{xx}|} = \frac{3.108 \times 10^9}{(1.095)(3.920 \times 10^9)} = .724 < \Lambda_{.05, 2, 3, 40} = .730.$$

Thus we reject the hypothesis of independence. Note the use of 40 in $\Lambda_{.05, 2, 3, 40}$ in place of $n - 1 - q = 46 - 1 - 3 = 42$. This is a conservative approach that allows the use of a table value without interpolation. \square

7.4.2 Independence of Several Subvectors

Let there be k sets of variates so that \mathbf{y} and Σ are partitioned as

$$\mathbf{y} = \begin{pmatrix} \mathbf{y}_1 \\ \mathbf{y}_2 \\ \vdots \\ \mathbf{y}_k \end{pmatrix} \quad \text{and} \quad \Sigma = \begin{pmatrix} \Sigma_{11} & \Sigma_{12} & \cdots & \Sigma_{1k} \\ \Sigma_{21} & \Sigma_{22} & \cdots & \Sigma_{2k} \\ \vdots & \vdots & & \vdots \\ \Sigma_{k1} & \Sigma_{k2} & \cdots & \Sigma_{kk} \end{pmatrix},$$

with p_i variables in \mathbf{y}_i , where $p_1 + p_2 + \cdots + p_k = p$. Note that $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_k$ represents a partitioning of \mathbf{y} , not a random sample of independent vectors. The

hypothesis that the subvectors $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_k$ are mutually independent can be expressed as $H_0: \Sigma_{ij} = \mathbf{O}$ for all $i \neq j$, or

$$H_0: \Sigma = \begin{pmatrix} \Sigma_{11} & \mathbf{O} & \cdots & \mathbf{O} \\ \mathbf{O} & \Sigma_{22} & \cdots & \mathbf{O} \\ \vdots & \vdots & & \vdots \\ \mathbf{O} & \mathbf{O} & \cdots & \Sigma_{kk} \end{pmatrix}. \quad (7.32)$$

The likelihood ratio statistic is

$$u = \frac{|\mathbf{S}|}{|\mathbf{S}_{11}||\mathbf{S}_{22}| \cdots |\mathbf{S}_{kk}|} \quad (7.33)$$

$$= \frac{|\mathbf{R}|}{|\mathbf{R}_{11}||\mathbf{R}_{22}| \cdots |\mathbf{R}_{kk}|}, \quad (7.34)$$

where \mathbf{S} and \mathbf{R} are obtained from a random sample of n observations and are partitioned as Σ above, conforming to $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_k$. Note that the denominator of (7.33) is the determinant of \mathbf{S} restricted by H_0 , that is, with $\mathbf{S}_{ij} = \mathbf{O}$ for all $i \neq j$ [see (2.92)]. The statistic u does not have Wilks' Λ -distribution as it does in (7.30) when $k = 2$, but a good χ^2 -approximation to its distribution is given by

$$u' = -\nu c \ln u, \quad (7.35)$$

where

$$c = 1 - \frac{1}{12f\nu}(2a_3 + 3a_2), \quad (7.36)$$

$$f = \frac{1}{2}a_2, \quad a_2 = p^2 - \sum_{i=1}^k p_i^2, \quad a_3 = p^3 - \sum_{i=1}^k p_i^3,$$

and ν is the degrees of freedom of \mathbf{S} or \mathbf{R} (see comments at the beginning of Section 7.2). We reject the independence hypothesis if $u' > \chi_{\alpha,f}^2$.

The degrees of freedom, $f = \frac{1}{2}a_2$, arises from the following consideration. The number of parameters in Σ unrestricted by the hypothesis is $\frac{1}{2}p(p+1)$. Under the hypothesis (7.32), the number of parameters in each Σ_{ii} is $\frac{1}{2}p_i(p_i+1)$, for a total of $\frac{1}{2}\sum_{i=1}^k p_i(p_i+1)$. The difference is

$$\begin{aligned} f &= \frac{1}{2}p(p+1) - \frac{1}{2}\sum_{i=1}^k p_i(p_i+1) = \frac{1}{2}\left(p^2 + p - \sum_i p_i^2 - \sum_i p_i\right) \\ &= \frac{1}{2}\left(p^2 + p - \sum_i p_i^2 - p\right) = \frac{1}{2}\left(p^2 - \sum_i p_i^2\right) = \frac{a_2}{2}. \end{aligned}$$

■ EXAMPLE 7.4.2

For 30 brands of Japanese Seishu wine, Siotani et al. (1963) studied the relationship between

$$y_1 = \text{taste}$$

$$y_2 = \text{odor}$$

and

$$x_1 = \text{pH}$$

$$x_5 = \text{direct reducing sugar}$$

$$x_2 = \text{acidity 1}$$

$$x_6 = \text{total sugar}$$

$$x_3 = \text{acidity 2}$$

$$x_7 = \text{alcohol}$$

$$x_4 = \text{sake meter}$$

$$x_8 = \text{formyl-nitrogen}.$$

The data are in Table 7.1.

We test independence of the following four subsets of variables:

$$(y_1, y_2), (x_1, x_2, x_3), (x_4, x_5, x_6), (x_7, x_8).$$

The sample covariance matrix is

$$\mathbf{S} = \begin{pmatrix} \mathbf{S}_{11} & \mathbf{S}_{12} & \mathbf{S}_{13} & \mathbf{S}_{14} \\ \mathbf{S}_{21} & \mathbf{S}_{22} & \mathbf{S}_{23} & \mathbf{S}_{24} \\ \mathbf{S}_{31} & \mathbf{S}_{32} & \mathbf{S}_{33} & \mathbf{S}_{34} \\ \mathbf{S}_{41} & \mathbf{S}_{42} & \mathbf{S}_{43} & \mathbf{S}_{44} \end{pmatrix} = \begin{pmatrix} .16 & .10 & .01 & .006 & .02 & -.04 & .02 & .01 & -.02 & 1.44 \\ .10 & .19 & -.01 & .009 & .02 & -.16 & .03 & .05 & .04 & 1.03 \\ .01 & -.01 & .03 & .004 & .03 & -.11 & -.03 & -.03 & -.01 & 4.45 \\ .006 & .009 & .004 & .024 & .020 & -.012 & -.009 & .0004 & .038 & 2.23 \\ .02 & .02 & .03 & .020 & .07 & -.18 & -.03 & -.03 & .05 & 9.03 \\ -.04 & -.16 & -.11 & -.012 & -.18 & 5.02 & -.35 & -.67 & -.12 & -23.11 \\ .02 & .03 & -.03 & -.009 & -.03 & -.35 & .13 & .15 & .05 & -4.26 \\ .01 & .05 & -.03 & .0004 & -.03 & -.67 & .15 & .26 & .13 & -3.47 \\ -.02 & .04 & -.01 & .038 & .05 & -.12 & .05 & .13 & .35 & 6.73 \\ 1.44 & 1.03 & 4.45 & 2.23 & 9.03 & -23.11 & -4.26 & -3.47 & 6.73 & 1541 \end{pmatrix}$$

where \mathbf{S}_{11} is 2×2 , \mathbf{S}_{22} is 3×3 , \mathbf{S}_{33} is 3×3 , and \mathbf{S}_{44} is 2×2 . We first obtain

$$u = \frac{|\mathbf{S}|}{|\mathbf{S}_{11}||\mathbf{S}_{22}||\mathbf{S}_{33}||\mathbf{S}_{44}|} = \frac{2.925 \times 10^{-7}}{(.0210)(.0000158)(.0361)(496.04)} = .01627.$$

Table 7.1 Seishu Measurements

y_1	y_2	x_1	x_2	x_3	x_4	x_5	x_6	x_7	x_8
1.0	.8	4.05	1.68	.85	3.0	3.97	5.00	16.90	122.0
.1	.2	3.81	1.39	.30	.6	3.62	4.52	15.80	62.0
.5	.0	4.20	1.63	.92	-2.3	3.48	4.46	15.80	139.0
.7	.7	4.35	1.43	.97	-1.6	3.45	3.98	15.40	150.0
-.1	-1.1	4.35	1.53	.87	-2.0	3.67	4.22	15.40	138.0
.4	.5	4.05	1.84	.95	-2.5	3.61	5.00	16.78	123.0
.2	-.3	4.20	1.61	1.09	-1.7	3.25	4.15	15.81	172.0
.3	-.1	4.32	1.43	.93	-5.0	4.16	5.45	16.78	144.0
.7	.4	4.21	1.74	.95	-1.5	3.40	4.25	16.62	153.0
.5	-.1	4.17	1.72	.92	-1.2	3.62	4.31	16.70	121.0
-.1	.1	4.45	1.78	1.19	-2.0	3.09	3.92	16.50	176.0
.5	-.5	4.45	1.48	.86	-2.0	3.32	4.09	15.40	128.0
.5	.8	4.25	1.53	.83	-3.0	3.48	4.54	15.55	126.0
.6	.2	4.25	1.49	.86	2.0	3.13	3.45	15.60	128.0
.0	-.5	4.05	1.48	.30	.0	3.67	4.52	15.38	99.0
-.2	-.2	4.22	1.64	.90	-2.2	3.59	4.49	16.37	122.8
.0	-.2	4.10	1.55	.85	1.8	3.02	3.62	15.31	114.0
.2	.2	4.28	1.52	.75	-4.8	3.64	4.93	15.77	125.0
-.1	-.2	4.32	1.54	.83	-2.0	3.17	4.62	16.60	119.0
.6	.1	4.12	1.68	.84	-2.1	3.72	4.83	16.93	111.0
.8	.5	4.30	1.50	.92	-1.5	2.98	3.92	15.10	68.0
.5	.2	4.55	1.50	1.14	.9	2.60	3.45	15.70	197.0
.4	.7	4.15	1.62	.78	-7.0	4.11	5.55	15.50	106.0
.6	-.3	4.15	1.32	.31	.8	3.56	4.42	15.40	49.5
-.7	-.3	4.25	1.77	1.12	.5	2.84	4.15	16.65	164.0
-.2	.0	3.95	1.36	.25	1.0	3.67	4.52	15.98	29.5
.3	-.1	4.35	1.42	.96	-2.5	3.40	4.12	15.30	131.0
.1	.4	4.15	1.17	1.06	-4.5	3.89	5.00	16.79	168.2
.4	.5	4.16	1.61	.91	-2.1	3.93	4.35	15.70	118.0
-.6	-.3	3.85	1.32	.30	.7	3.61	4.29	15.71	48.0

For the χ^2 -approximation, we calculate

$$a_2 = p^2 - \sum_{i=1}^4 p_i^2 = 10^2 - (2^2 + 3^2 + 3^2 + 2^2) = 74,$$

$$a_3 = p^3 - \sum_{i=1}^4 p_i^3 = 930, \quad f = \frac{1}{2}a_2 = 37,$$

$$c = 1 - \frac{1}{12f\nu}(2a_3 + 3a_2) = 1 - \frac{2(930) + 3(74)}{12(37)(29)} = .838.$$

Then,

$$u' = -\nu c \ln u = -(29)(.838) \ln(.01627) = 100.122,$$

which exceeds $\chi^2_{.001,37} = 69.35$, and we reject the hypothesis of independence of the four subsets. \square

7.4.3 Test for Independence of All Variables

If all $p_i = 1$ in the hypothesis (7.32) in Section 7.4.2, we have the special case in which all the variables are mutually independent, $H_0 : \sigma_{jk} = 0$ for $j \neq k$, or

$$H_0 : \Sigma = \begin{pmatrix} \sigma_{11} & 0 & \cdots & 0 \\ 0 & \sigma_{22} & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & \sigma_{pp} \end{pmatrix}.$$

There is no restriction on the σ_{jj} 's. With $\sigma_{jk} = 0$ for all $j \neq k$, the corresponding ρ_{jk} 's are also 0, and an equivalent form of the hypothesis is $H_0 : \mathbf{P}_\rho = \mathbf{I}$, where \mathbf{P}_ρ is the population correlation matrix defined in (3.37).

Since all $p_i = 1$, the statistics (7.33) and (7.34) reduce to

$$u = \frac{|\mathbf{S}|}{s_{11}s_{22} \cdots s_{pp}} = |\mathbf{R}|, \quad (7.37)$$

and (7.35) becomes

$$u' = -[\nu - \frac{1}{6}(2p + 5)] \ln u, \quad (7.38)$$

which has an approximate χ^2_f -distribution, where ν is the degrees of freedom of \mathbf{S} or \mathbf{R} (see comments at the beginning of Section 7.2) and $f = \frac{1}{2}p(p - 1)$ is the degrees of freedom of χ^2 . We reject H_0 if $u' > \chi^2_{\alpha,f}$. Exact percentage points of u' for selected values of n and p are given in Table A.15 (Mathai and Katiyar 1979). Percentage points for the limiting χ^2 -distribution are also given for comparison.

Note that $|\mathbf{R}|$ in (7.37) varies between 0 and 1. If the variables were uncorrelated (in the sample), we would have $\mathbf{R} = \mathbf{I}$ and $|\mathbf{R}| = 1$. On the other hand, if two or more variables were linearly related, \mathbf{R} would not be full rank and we would have $|\mathbf{R}| = 0$. If the variables were highly correlated, $|\mathbf{R}|$ would be close to 0; if the correlations were all small, $|\mathbf{R}|$ would be close to 1. This can be illustrated for $p = 2$:

$$|\mathbf{R}| = \begin{vmatrix} 1 & r \\ r & 1 \end{vmatrix} = 1 - r^2.$$

■ EXAMPLE 7.4.3

To test the hypothesis $H_0 : \sigma_{jk} = 0$, $j \neq k$, for the probe word data from Table 3.6, we calculate

$$\mathbf{R} = \begin{pmatrix} 1.000 & .614 & .757 & .575 & .413 \\ .614 & 1.000 & .547 & .750 & .548 \\ .757 & .547 & 1.000 & .605 & .692 \\ .575 & .750 & .605 & 1.000 & .524 \\ .413 & .548 & .692 & .524 & 1.000 \end{pmatrix}.$$

Then by (7.37) and (7.38),

$$u = |\mathbf{R}| = .0409,$$

$$u' = -[n - 1 - \frac{1}{6}(2p + 5)] \ln u = 23.97.$$

The exact .01 critical value for u' from Table A.15 is 23.75, and we therefore reject H_0 . The approximate χ^2 critical value for u' is $\chi^2_{.01,10} = 23.21$, with which we also reject H_0 . \square

PROBLEMS

- 7.1** Show that if $\mathbf{S} = \Sigma_0$ in (7.1), then $u = 0$.
- 7.2** Verify (7.3); that is, show that $\ln |\Sigma_0| - \ln |\mathbf{S}| = -\ln |\mathbf{S}\Sigma_0^{-1}|$.
- 7.3** Verify (7.4); that is, show that $-\ln(\prod_{i=1}^p \lambda_i) + \sum_{i=1}^p \lambda_i = \sum_{i=1}^p (\lambda_i - \ln \lambda_i)$.
- 7.4** Show that the likelihood ratio for $H_0: \Sigma = \sigma^2 \mathbf{I}$ is given by (7.5), $\text{LR} = [|\mathbf{S}|/(\text{tr } \mathbf{S}/p)^p]^{n/2}$.
- 7.5** Show that $u = 1$ and $u' = 0$ if all the λ_i 's are equal, as noted in Section 7.2.2, where u is given by (7.8) and u' by (7.9).
- 7.6** Show that the covariance matrix in (7.10) can be written in the form $\sigma^2[(1 - \rho)\mathbf{I} + \rho\mathbf{J}]$, as given in (7.11).
- 7.7** Obtain (7.15) from (7.14) as follows:
- Show that the $p \times p$ matrix \mathbf{J} has a single nonzero eigenvalue equal to p and corresponding eigenvector proportional to \mathbf{j} .
 - Show that $\mathbf{S}_0 = s^2[(1 - r)\mathbf{I} + r\mathbf{J}]$ in (7.13) can be written in the form $\mathbf{S}_0 = s^2(1 - r)(\mathbf{I} + \frac{r}{1-r} \mathbf{J})$.
 - Use Section 2.11.2 and (2.108) to obtain (7.15).
- 7.8** Show that M in (7.19) can be expressed in the form given in (7.21).
- 7.9** (a) Calculate M as given in (7.21) for

$$\mathbf{S}_1 = \begin{pmatrix} 2 & 1 \\ 1 & 4 \end{pmatrix}, \quad \mathbf{S}_2 = \begin{pmatrix} 4 & 3 \\ 3 & 6 \end{pmatrix}.$$

Assume $\nu_1 = \nu_2 = 5$.

(b) Calculate M for

$$\mathbf{S}_1 = \begin{pmatrix} 2 & 1 \\ 1 & 4 \end{pmatrix}, \quad \mathbf{S}_2 = \begin{pmatrix} 10 & 15 \\ 15 & 30 \end{pmatrix}.$$

Assume $\nu_1 = \nu_2 = 5$.

In (b), \mathbf{S}_1 and \mathbf{S}_2 differ more than in (a) and M is accordingly much smaller. This illustrates the comments following (7.21).

- 7.10** Obtain (7.31), $\Lambda = \prod_{i=1}^s (1 - r_i^2)$, by using (2.94) to write $|\mathbf{S}|$ in the form $|\mathbf{S}| = |\mathbf{S}_{xx}| |\mathbf{S}_{yy} - \mathbf{S}_{yx} \mathbf{S}_{xx}^{-1} \mathbf{S}_{xy}|$.
- 7.11** Show that the forms of u in (7.33) and (7.34) reduce to (7.37) when all $p_i = 1$.
- 7.12** Show that when all $p_i = 1$, c in (7.36) reduces to $1 - (2p + 5)/6\nu$, so that (7.35) becomes (7.38).
- 7.13** Give a justification for the degrees of freedom $f = \frac{1}{2} p(p - 1)$ for the approximate χ^2 test statistic u' in (7.38).
- 7.14** In Example 5.2.2, we assumed that for the height and weight data of Table 3.1, the population covariance matrix is

$$\Sigma = \begin{pmatrix} 20 & 100 \\ 100 & 1000 \end{pmatrix}.$$

Test this as a hypothesis using (7.2).

- 7.15** Test $H_0: \Sigma = \sigma^2 \mathbf{I}$ and $H_0: \mathbf{C} \Sigma \mathbf{C}' = \sigma^2 \mathbf{I}$ for the calculator speed data of Table 6.12.
- 7.16** Test $H_0: \Sigma = \sigma^2 \mathbf{I}$ and $H_0: \mathbf{C} \Sigma \mathbf{C}' = \sigma^2 \mathbf{I}$ for the ramus bone data of Table 3.7.
- 7.17** Test $H_0: \Sigma = \sigma^2 \mathbf{I}$ and $H_0: \mathbf{C} \Sigma \mathbf{C}' = \sigma^2 \mathbf{I}$ for the cork data of Table 6.21.
- 7.18** Test $H_0: \Sigma = \sigma^2[(1 - \rho)\mathbf{I} + \rho\mathbf{J}]$ for the probe word data in Table 3.6. Use both χ^2 - and F -approximations.
- 7.19** Test $H_0: \Sigma = \sigma^2[(1 - \rho)\mathbf{I} + \rho\mathbf{J}]$ for the calculator speed data in Table 6.12. Use both χ^2 - and F -approximations.
- 7.20** Test $H_0: \Sigma = \sigma^2[(1 - \rho)\mathbf{I} + \rho\mathbf{J}]$ for the ramus bone data in Table 3.7. Use both χ^2 - and F -approximations.
- 7.21** Test $H_0: \Sigma_1 = \Sigma_2$ for the beetles data of Table 5.5. Use an exact critical value from Table A.14 as well as χ^2 - and F -approximations.
- 7.22** Test $H_0: \Sigma_1 = \Sigma_2$ for the engineer data of Table 5.6. Use an exact critical value from Table A.14 as well as χ^2 - and F -approximations.

- 7.23** Test $H_0: \Sigma_1 = \Sigma_2$ for the dystrophy data of Table 5.7. Use an exact critical value from Table A.14 as well as χ^2 - and F -approximations.
- 7.24** Test $H_0: \Sigma_1 = \Sigma_2$ for the cyclical data of Table 5.8. Use an exact critical value from Table A.14 as well as χ^2 - and F -approximations.
- 7.25** Test $H_0: \Sigma_1 = \Sigma_2 = \Sigma_3$ for the fish data of Table 6.17. Use an exact critical value from Table A.14 as well as χ^2 - and F -approximations.
- 7.26** Test $H_0: \Sigma_1 = \Sigma_2 = \dots = \Sigma_6$ for the rootstock data in Table 6.2. Use an exact critical value from Table A.14 as well as χ^2 - and F -approximations.
- 7.27** Test $H_0: \Sigma_{11} = \Sigma_{12} = \dots = \Sigma_{43}$ for the snap bean data in Table 6.18. Use both χ^2 - and F -approximations.
- 7.28** Test independence of (y_1, y_2) and (x_1, x_2) for the sons data in Table 3.8.
- 7.29** Test independence of (y_1, y_2, y_3) and (x_1, x_2, x_3) for the glucose data in Table 3.9.
- 7.30** Test independence of (y_1, y_2) and (x_1, x_2, \dots, x_8) for the Seishu data of Table 7.1.
- 7.31** The data in Table 7.2 relate temperature, humidity, and evaporation (courtesy of R. J. Freund). The variables are

- y_1 = maximum daily air temperature
- y_2 = minimum daily air temperature
- y_3 = integrated area under daily air temperature curve, that is,
a measure of average air temperature
- y_4 = maximum daily soil temperature
- y_5 = minimum daily soil temperature
- y_6 = integrated area under soil temperature curve
- y_7 = maximum daily relative humidity
- y_8 = minimum daily relative humidity
- y_9 = integrated area under daily humidity curve
- y_{10} = total wind, measured in miles per day
- y_{11} = evaporation

Test independence of the following five groups of variables: (y_1, y_2, y_3) , (y_4, y_5, y_6) , (y_7, y_8, y_9) , y_{10} , and y_{11} .

- 7.32** Test the independence of all the variables for the calcium data of Table 3.4.
- 7.33** Test the independence of all the variables for the calculator speed data of Table 6.12.

Table 7.2 Temperature, Humidity, and Evaporation

y_1	y_2	y_3	y_4	y_5	y_6	y_7	y_8	y_9	y_{10}	y_{11}
84	65	147	85	59	151	95	40	398	273	30
84	65	149	86	61	159	94	28	345	140	34
79	66	142	83	64	152	94	41	368	318	33
81	67	147	83	65	158	94	50	406	282	26
84	68	167	88	69	180	93	46	379	311	41
74	66	131	77	67	147	96	73	478	446	4
73	66	131	78	69	159	96	72	462	294	5
75	67	134	84	68	159	95	70	464	313	20
84	68	161	89	71	195	95	63	430	455	31
86	72	169	91	76	206	93	56	406	604	36
88	73	176	91	76	206	94	55	393	610	43
90	74	187	94	76	211	94	51	385	520	47
88	72	171	94	75	211	96	54	405	663	45
58	72	171	92	70	201	95	51	392	467	45
81	69	154	87	68	167	95	61	448	184	11
79	68	149	83	68	162	95	59	436	177	10
84	69	160	87	66	173	95	42	392	173	30
84	70	160	87	68	177	94	44	392	76	29
84	70	168	88	70	169	95	48	396	72	23
77	67	147	83	66	170	97	60	431	183	16
87	67	166	92	67	196	96	44	379	76	37
89	69	171	92	72	199	94	48	393	230	50
89	72	180	94	72	204	95	48	394	193	36
93	72	186	92	73	201	94	47	386	400	54
93	74	188	93	72	206	95	47	389	339	44
94	75	199	94	72	208	96	45	370	172	41
93	74	193	95	73	214	95	50	396	238	45
93	74	196	95	70	210	96	45	380	118	42
96	75	198	95	71	207	93	40	365	93	50
95	76	202	95	69	202	93	39	357	269	48
84	73	173	96	69	173	94	58	418	128	17
91	71	170	91	69	168	94	44	420	423	20
⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮
96	76	202	96	71	208	94	40	368	139	50

- 7.34** Test the independence of all the variables for the ramus bone data of Table 3.7.
- 7.35** Test the independence of all the variables for the cork data of Table 6.21.

CHAPTER 8

DISCRIMINANT ANALYSIS: DESCRIPTION OF GROUP SEPARATION

8.1 INTRODUCTION

We use the term *group* to represent either a population or a sample from the population. There are two major objectives in separation of groups:

1. Description of group separation, in which linear functions of the variables (discriminant functions) are used to describe or elucidate the differences between two or more groups. The goals of descriptive discriminant analysis include identifying the relative contribution of the p variables to separation of the groups and finding the optimal plane on which the points can be projected to best illustrate the configuration of the groups.
2. Prediction or allocation of observations to groups, in which linear or quadratic functions of the variables (classification functions) are employed to assign an individual sampling unit to one of the groups. The measured values in the observation vector for an individual or object are evaluated by the classification functions to find the group to which the individual most likely belongs.

For consistency we will use the term *discriminant analysis* only in connection with objective 1. We will refer to all aspects of objective 2 as *classification analysis*, which is the subject of Chapter 9. Unfortunately, there is no general agreement with regard to usage of the terms “discriminant analysis” and “discriminant functions.” Many writers, perhaps the majority, use the term “discriminant analysis” in connection with the second objective, prediction or allocation. The linear functions contributing to the first objective, description of group separation, are often referred to as canonical variates or discriminant coordinates. To avoid confusion, we prefer to reserve the term *canonical* for canonical correlation analysis in Chapter 11.

Discriminant functions are linear combinations of variables that best separate groups. They were introduced in Section 5.5 for two groups and in Sections 6.1.4 and 6.4 for several groups. In those sections, interest was centered on follow-up to Hotelling’s T^2 -tests and MANOVA tests. In this chapter, we further develop these useful multivariate tools.

8.2 THE DISCRIMINANT FUNCTION FOR TWO GROUPS

We assume that the two populations to be compared have the same covariance matrix Σ but distinct mean vectors μ_1 and μ_2 . We work with samples $\mathbf{y}_{11}, \mathbf{y}_{12}, \dots, \mathbf{y}_{1n_1}$ and $\mathbf{y}_{21}, \mathbf{y}_{22}, \dots, \mathbf{y}_{2n_2}$ from the two populations. As usual, each vector \mathbf{y}_{ij} consists of measurements on p variables. The discriminant function is the linear combination of these p variables that maximizes the distance between the two (transformed) group mean vectors. A linear combination $z = \mathbf{a}'\mathbf{y}$ transforms each observation vector to a scalar:

$$\begin{aligned} z_{1i} &= \mathbf{a}'\mathbf{y}_{1i} = a_1y_{1i1} + a_2y_{1i2} + \dots + a_py_{1ip}, & i &= 1, 2, \dots, n_1 \\ z_{2i} &= \mathbf{a}'\mathbf{y}_{2i} = a_1y_{2i1} + a_2y_{2i2} + \dots + a_py_{2ip}, & i &= 1, 2, \dots, n_2. \end{aligned}$$

Hence the $n_1 + n_2$ observation vectors in the two samples,

$$\begin{array}{ll} \mathbf{y}_{11} & \mathbf{y}_{21} \\ \mathbf{y}_{12} & \mathbf{y}_{22} \\ \vdots & \vdots \\ \mathbf{y}_{1n_1} & \mathbf{y}_{2n_2}, \end{array}$$

are transformed to scalars,

$$\begin{array}{ll} z_{11} & z_{21} \\ z_{12} & z_{22} \\ \vdots & \vdots \\ z_{1n_1} & z_{2n_2}. \end{array}$$

We find the means $\bar{z}_1 = \sum_{i=1}^{n_1} z_{1i}/n_1 = \mathbf{a}'\bar{\mathbf{y}}_1$ and $\bar{z}_2 = \mathbf{a}'\bar{\mathbf{y}}_2$ by (3.54), where $\bar{\mathbf{y}}_1 = \sum_{i=1}^{n_1} \mathbf{y}_{1i}/n_1$ and $\bar{\mathbf{y}}_2 = \sum_{i=1}^{n_2} \mathbf{y}_{2i}/n_2$. We then wish to find the vector \mathbf{a} that maximizes the standardized difference $(\bar{z}_1 - \bar{z}_2)/s_z$. Since $(\bar{z}_1 - \bar{z}_2)/s_z$ can be

negative, we use the squared distance $(\bar{z}_1 - \bar{z}_2)^2/s_z^2$, which, by (3.54) and (3.55), can be expressed as

$$\frac{(\bar{z}_1 - \bar{z}_2)^2}{s_z^2} = \frac{[\mathbf{a}'(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)]^2}{\mathbf{a}'\mathbf{S}_{pl}\mathbf{a}}. \quad (8.1)$$

The maximum of (8.1) occurs when

$$\mathbf{a} = \mathbf{S}_{pl}^{-1}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2), \quad (8.2)$$

or when \mathbf{a} is any multiple of $\mathbf{S}_{pl}^{-1}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)$. Thus the maximizing vector \mathbf{a} is not unique. However, its "direction" is unique; that is, the relative values or ratios of a_1, a_2, \dots, a_p are unique, and $z = \mathbf{a}'\mathbf{y}$ projects points \mathbf{y} onto the line on which $(\bar{z}_1 - \bar{z}_2)^2/s_z^2$ is maximized. Note that in order for \mathbf{S}_{pl}^{-1} to exist, we must have $n_1 + n_2 - 2 > p$.

The optimum direction given by $\mathbf{a} = \mathbf{S}_{pl}^{-1}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)$ is effectively parallel to the line joining $\bar{\mathbf{y}}_1$ and $\bar{\mathbf{y}}_2$, because the squared distance $(\bar{z}_1 - \bar{z}_2)^2/s_z^2$ is equivalent to the standardized distance between $\bar{\mathbf{y}}_1$ and $\bar{\mathbf{y}}_2$. This can be seen by substituting (8.2) into (8.1) to obtain

$$\frac{(\bar{z}_1 - \bar{z}_2)^2}{s_z^2} = (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{pl}^{-1} (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2) \quad (8.3)$$

for $z = \mathbf{a}'\mathbf{y}$ with $\mathbf{a} = \mathbf{S}_{pl}^{-1}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)$. Since $\mathbf{a}' = (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{pl}^{-1}$, (8.3) can be written as $(\bar{z}_1 - \bar{z}_2)^2/s_z^2 = \mathbf{a}'(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)$, and any direction other than that represented by $\mathbf{a} = \mathbf{S}_{pl}^{-1}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)$ would yield a smaller difference between $\mathbf{a}'\bar{\mathbf{y}}_1$ and $\mathbf{a}'\bar{\mathbf{y}}_2$ (see Section 5.5).




Figure 8.1 Two-group discriminant analysis.




Figure 8.2 Separation achieved by the discriminant function.

Figure 8.1 illustrates the separation of two bivariate normal ($p = 2$) groups along the single dimension represented by the discriminant function $z = \mathbf{a}'\mathbf{y}$, where \mathbf{a} is given by (8.2). In this illustration the population covariance matrices are equal. The linear combinations $z_{1i} = \mathbf{a}'\mathbf{y}_{1i} = a_1y_{1i1} + a_2y_{1i2}$ and $z_{2i} = \mathbf{a}'\mathbf{y}_{2i} = a_1y_{2i1} + a_2y_{2i2}$ project the points \mathbf{y}_{1i} and \mathbf{y}_{2i} onto the line of optimum separation of the two groups. Since the two variables y_1 and y_2 are bivariate normal, a linear combination $z = a_1y_1 + a_2y_2 = \mathbf{a}'\mathbf{y}$ is univariate normal (see property 1a in Section 4.2). We have therefore indicated this by two univariate normal densities along the line representing z .

The point where the line joining the points of intersection of the two ellipses intersects the discriminant function line z is the point of maximum separation (minimum overlap) of points projected onto the line. If the two populations are multivariate normal with common covariance matrix Σ , as illustrated in Figure 8.1, it can be shown that all possible group separation is expressed in a single new dimension.

In Figure 8.2, we illustrate the optimum separation achieved by the discriminant function. Projection in another direction denoted by z' gives a smaller standardized distance between the transformed means \bar{z}'_1 and \bar{z}'_2 and also a larger overlap between the projected points.

■ EXAMPLE 8.2

Samples of steel produced at two different rolling temperatures are compared in Table 8.1 (Kramer and Jensen 1969a). The variables are y_1 = yield point

Table 8.1 Yield Point and Ultimate Strength of Steel Produced at Two Rolling Temperatures

Temperature 1		Temperature 2	
y_1	y_2	y_1	y_2
33	60	35	57
36	61	36	59
35	64	38	59
38	63	39	61
40	65	41	63
		43	65
		41	59

and y_2 = ultimate strength. From the data, we calculate

$$\bar{y}_1 = \begin{pmatrix} 36.4 \\ 62.6 \end{pmatrix}, \quad \bar{y}_2 = \begin{pmatrix} 39.0 \\ 60.4 \end{pmatrix}, \quad S_{Pl} = \begin{pmatrix} 7.92 & 5.68 \\ 5.68 & 6.29 \end{pmatrix}.$$

A plot of the data appears in Figure 8.3. We see that if the points were projected on either the y_1 or the y_2 axis, there would be considerable overlap. In fact, when the two groups are compared by means of a t -statistic for each




Figure 8.3 Ultimate strength and yield point for steel rolled at two temperatures.

Table 8.2 Discriminant Function $z = -1.633y_1 + 1.820y_2$ Evaluated for Data in Table 8.1

Temperature 1	Temperature 2
55.29	46.56
52.20	48.57
59.30	45.30
52.58	47.30
52.95	47.68
	48.05
	40.40

variable separately, both t 's are nonsignificant:

$$t_1 = \frac{\bar{y}_{11} - \bar{y}_{21}}{\sqrt{s_{11}(1/n_1 + 1/n_2)}} = -1.58,$$

$$t_2 = \frac{\bar{y}_{12} - \bar{y}_{22}}{\sqrt{s_{22}(1/n_1 + 1/n_2)}} = 1.48.$$

However, it is clear in Figure 8.3 that the two groups can be separated. If they are projected in an appropriate direction as in Figure 8.1, there will be no overlap. The single dimension onto which the points would be projected is the discriminant function

$$z = \mathbf{a}'\mathbf{y} = a_1y_1 + a_2y_2 = -1.633y_1 + 1.820y_2,$$

where \mathbf{a} is obtained as

$$\mathbf{a} = \mathbf{S}_{\text{pl}}^{-1}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2) = \begin{pmatrix} -1.633 \\ 1.820 \end{pmatrix}.$$

The values of the projected points are found by calculating z for each observation vector \mathbf{y} in the two groups. The results are given in Table 8.2, where the separation provided by the discriminant function is clearly evident. \square

8.3 RELATIONSHIP BETWEEN TWO-GROUP DISCRIMINANT ANALYSIS AND MULTIPLE REGRESSION

The mutual connection between multiple regression and two-group discriminant analysis was introduced as a computational device in Section 5.6.2. The roles of independent and dependent variables are reversed in the two models. The dependent variables (y 's) of discriminant analysis become the independent variables in regression.

Let w be a grouping variable (identifying groups 1 and 2) such that $\bar{w} = 0$ and define $\mathbf{b} = (b_1, b_2, \dots, b_p)'$ as the vector of regression coefficients when w is fit to the y 's. Then by (5.21), \mathbf{b} is proportional to the discriminant function coefficient vector $\mathbf{a} = \mathbf{S}_{\text{pl}}^{-1}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)$:

$$\mathbf{b} = \frac{n_1 n_2}{(n_1 + n_2)(n_1 + n_2 - 2 + T^2)} \mathbf{a}, \quad (8.4)$$

where $T^2 = [n_1 n_2 / (n_1 + n_2)](\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{\text{pl}}^{-1}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)$ as in (5.9). From (5.20) the squared multiple correlation R^2 is related to T^2 by

$$R^2 = (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{b} = \frac{T^2}{n_1 + n_2 - 2 + T^2}. \quad (8.5)$$

The test statistic (5.29) for the hypothesis that q of the $p + q$ variables are redundant for separating the groups can also be obtained in terms of regression by (5.31) as

$$F = \frac{n_1 + n_2 - p - q - 1}{q} \frac{R_{p+q}^2 - R_p^2}{1 - R_{p+q}^2}, \quad (8.6)$$

where R_{p+q}^2 and R_p^2 are from regressions with $p + q$ and p variables, respectively.

The link between two-group discriminant analysis and multiple regression was first noted by Fisher (1936). Flury and Riedwyl (1985) give further insights into the relationship.

■ EXAMPLE 8.3

In Example 5.6.2, the psychological data of Table 5.1 were used in an illustration of the regression approach to computation of \mathbf{a} and T^2 . We use the same data to obtain \mathbf{b} and R^2 from \mathbf{a} and T^2 .

From the results of Examples 5.4.2 and 5.5, we have

$$T^2 = 97.6015,$$

$$\mathbf{a} = \begin{pmatrix} .5104 \\ -.2033 \\ .4660 \\ -.3097 \end{pmatrix}.$$

To find \mathbf{b} from \mathbf{a} and T^2 , we use (8.4):

$$\mathbf{b} = \frac{(32)(32)}{(32 + 32)(32 + 32 - 2 + 97.6015)} \mathbf{a} = \begin{pmatrix} .051 \\ -.020 \\ .047 \\ -.031 \end{pmatrix}.$$

To find R^2 , we use (8.5):

$$R^2 = (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{b} = \begin{pmatrix} 3.625 \\ 2.000 \\ 10.531 \\ .812 \end{pmatrix}' \begin{pmatrix} .051 \\ -.020 \\ .047 \\ -.031 \end{pmatrix} = .611.$$

We can also use the relationship with T^2 in (8.5):

$$R^2 = \frac{T^2}{n_1 + n_2 - 2 + T^2} = \frac{97.6105}{32 + 32 - 2 + 97.6015} = .611.$$

□

8.4 DISCRIMINANT ANALYSIS FOR SEVERAL GROUPS

8.4.1 Discriminant Functions

In discriminant analysis for several groups, we are concerned with finding linear combinations of variables that best separate the k groups of multivariate observations. Discriminant analysis for several groups may serve any one of various purposes:

1. Examine group separation in a two-dimensional plot. When there are more than two groups, it requires more than one discriminant function to describe group separation. If the points in the p -dimensional space are projected onto a 2-dimensional space represented by the first two discriminant functions, we obtain the best possible view of how the groups are separated.
2. Find a subset of the original variables that separates the groups almost as well as the original set. This topic was introduced in Section 6.11.2.
3. Rank the variables in terms of their relative contribution to group separation. This use for discriminant functions has been mentioned in Sections 5.5, 6.1.4, 6.1.8, and 6.4. In Section 8.5, we discuss standardized discriminant function coefficients that provide a more valid comparison of the variables.
4. Interpret the new dimensions represented by the discriminant functions.
5. Follow up to fixed-effects MANOVA.

Purposes 3 and 4 are closely related. Any of the first four goals can be used to accomplish purpose 5. Methods of achieving these five goals of discriminant analysis are discussed in subsequent sections. In the present section we review discriminant functions for the several-group case and discuss attendant assumptions. For alternative estimators of discriminant functions that may be useful in the presence of multicollinearity or outliers, see Rencher (1998, Section 5.11).

For k groups (samples) with n_i observations in the i th group, we transform each observation vector \mathbf{y}_{ij} to obtain $z_{ij} = \mathbf{a}'\mathbf{y}_{ij}$, $i = 1, 2, \dots, k$; $j = 1, 2, \dots, n_i$, and find the means $\bar{z}_i = \mathbf{a}'\bar{\mathbf{y}}_i$, where $\bar{\mathbf{y}}_i = \sum_{j=1}^{n_i} \mathbf{y}_{ij}/n_i$. As in the two-group case, we seek the vector \mathbf{a} that maximally separates $\bar{z}_1, \bar{z}_2, \dots, \bar{z}_k$. To express separation among $\bar{z}_1, \bar{z}_2, \dots, \bar{z}_k$, we extend the separation criterion (8.1) to the k -group case. Since $\mathbf{a}'(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2) = (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)'\mathbf{a}$, we can express (8.1) in the form

$$\frac{(\bar{z}_1 - \bar{z}_2)^2}{s_z^2} = \frac{[\mathbf{a}'(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)]^2}{\mathbf{a}'\mathbf{S}_{\text{pl}}\mathbf{a}} = \frac{\mathbf{a}'(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)'\mathbf{a}}{\mathbf{a}'\mathbf{S}_{\text{pl}}\mathbf{a}}. \quad (8.7)$$

To extend (8.7) to k groups, we use the \mathbf{H} matrix from MANOVA in place of $(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)'$ [see (6.37)] and \mathbf{E} in place of \mathbf{S}_{pl} to obtain

$$\lambda = \frac{\mathbf{a}'\mathbf{H}\mathbf{a}}{\mathbf{a}'\mathbf{E}\mathbf{a}}, \quad (8.8)$$

which can also be expressed as

$$\lambda = \frac{\text{SSH}(z)}{\text{SSE}(z)}, \quad (8.9)$$

where $\text{SSH}(z)$ and $\text{SSE}(z)$ are the between and within sums of squares for z .

We can write (8.8) in the form

$$\begin{aligned} \mathbf{a}'\mathbf{H}\mathbf{a} &= \lambda\mathbf{a}'\mathbf{E}\mathbf{a}, \\ \mathbf{a}'(\mathbf{H}\mathbf{a} - \lambda\mathbf{E}\mathbf{a}) &= 0. \end{aligned} \quad (8.10)$$

We examine values of λ and \mathbf{a} that are solutions of (8.10) in a search for the value of \mathbf{a} that results in maximum λ . The solution $\mathbf{a}' = \mathbf{0}'$ is not permissible because it gives $\lambda = 0/0$ in (8.8). Other solutions are found from

$$\mathbf{H}\mathbf{a} - \lambda\mathbf{E}\mathbf{a} = \mathbf{0}, \quad (8.11)$$

which can be written in the form

$$(\mathbf{E}^{-1}\mathbf{H} - \lambda\mathbf{I})\mathbf{a} = \mathbf{0}. \quad (8.12)$$

The solutions of (8.12) are the eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_s$ and associated eigenvectors $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_s$ of $\mathbf{E}^{-1}\mathbf{H}$. As in previous discussions of eigenvalues, we consider them to be ranked $\lambda_1 > \lambda_2 > \dots > \lambda_s$. The number of (nonzero) eigenvalues s is the rank of \mathbf{H} , which can be found as the smaller of $k - 1$ or p . Thus the largest eigenvalue λ_1 is the maximum value of $\lambda = \mathbf{a}'\mathbf{H}\mathbf{a}/\mathbf{a}'\mathbf{E}\mathbf{a}$ in (8.8), and the coefficient vector that produces the maximum is the corresponding eigenvector \mathbf{a}_1 . (This can be verified using calculus.) Hence the discriminant function that maximally separates the means is $z_1 = \mathbf{a}'_1\mathbf{y}$; that is, z_1 represents the dimension or direction that maximally separates the means.

From the s eigenvectors $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_s$ of $\mathbf{E}^{-1}\mathbf{H}$ corresponding to $\lambda_1, \lambda_2, \dots, \lambda_s$, we obtain s discriminant functions $z_1 = \mathbf{a}'_1\mathbf{y}, z_2 = \mathbf{a}'_2\mathbf{y}, \dots, z_s = \mathbf{a}'_s\mathbf{y}$, which show the dimensions or directions of differences among $\bar{\mathbf{y}}_1, \bar{\mathbf{y}}_2, \dots, \bar{\mathbf{y}}_k$. These discriminant functions are uncorrelated but are not orthogonal [see Rencher (1998, pp. 203–204)]. These discriminant functions are uncorrelated, but they are not orthogonal ($\mathbf{a}'_i\mathbf{a}_j = 0$ for $i \neq j$) because $\mathbf{E}^{-1}\mathbf{H}$ is not symmetric. Note that the numbering z_1, z_2, \dots, z_s corresponds to the eigenvalues, not to the k groups as was done earlier in this section.

The relative importance of each discriminant function z_i can be assessed by considering its eigenvalue as a proportion of the total:

$$\frac{\lambda_i}{\sum_{j=1}^s \lambda_j}. \quad (8.13)$$

By this criterion, two or three discriminant functions will often suffice to describe the group differences. The discriminant functions associated with small eigenvalues can be neglected. A test of significance for each discriminant function is also available (see Section 8.6).

The matrix $\mathbf{E}^{-1}\mathbf{H}$ is not symmetric. Many algorithms for computation of eigenvalues and eigenvectors accept only symmetric matrices. In Section 6.1.4, it was shown that the eigenvalues of the symmetric matrix $(\mathbf{U}^{-1})'\mathbf{H}\mathbf{U}^{-1}$ are the same as those of $\mathbf{E}^{-1}\mathbf{H}$, where $\mathbf{E} = \mathbf{U}'\mathbf{U}$ is the Cholesky factorization of \mathbf{E} . However, an adjustment is needed for the eigenvectors. If \mathbf{b} is an eigenvector of $(\mathbf{U}^{-1})'\mathbf{H}\mathbf{U}^{-1}$, then $\mathbf{a} = \mathbf{U}^{-1}\mathbf{b}$ is an eigenvector of $\mathbf{E}^{-1}\mathbf{H}$.

The discussion above was presented in terms of unequal sample sizes n_1, n_2, \dots, n_k . In applications, this situation is common and can be handled with no difficulty. Ideally, the smallest n_i should exceed the number of variables, p . This is not required mathematically, but will lead to more stable discriminant functions.

■ EXAMPLE 8.4.1

The data in Table 8.3 were collected by G. R. Bryce and R. M. Barker (Brigham Young University) as part of a preliminary study of a possible link between football helmet design and neck injuries.

Six head measurements were made on each subject. There were 30 subjects in each of three groups: high school football players (group 1), college football players (group 2), and non-football players (group 3). The six variables are

WDIM = head width at widest dimension

CIRCUM = head circumference

FBEYE = front-to-back measurement at eye level

EYEHHD = eye-to-top-of-head measurement

EARHD = ear-to-top-of-head measurement

JAW = jaw width

The eigenvalues of $\mathbf{E}^{-1}\mathbf{H}$ are $\lambda_1 = 1.9178$ and $\lambda_2 = .1159$. The corresponding eigenvectors are

$$\mathbf{a}_1 = \begin{pmatrix} -.948 \\ .004 \\ .006 \\ .647 \\ .504 \\ .829 \end{pmatrix}, \quad \mathbf{a}_2 = \begin{pmatrix} -1.407 \\ .001 \\ .029 \\ -.540 \\ .384 \\ 1.529 \end{pmatrix}.$$

The first eigenvalue accounts for a substantial proportion of the total:

$$\frac{\lambda_1}{\lambda_1 + \lambda_2} = \frac{1.9178}{1.9178 + .1159} = .94.$$

Table 8.3 Head Measurements for Three Groups

Group	WDIM	CIRCUM	FBEYE	EYEHD	EARHD	JAW
1	13.5	57.2	19.5	12.5	14.0	11.0
1	15.5	58.4	21.0	12.0	16.0	12.0
1	14.5	55.9	19.0	10.0	13.0	12.0
1	15.5	58.4	20.0	13.5	15.0	12.0
1	14.5	58.4	20.0	13.0	15.5	12.0
1	14.0	61.0	21.0	12.0	14.0	13.0
1	15.0	58.4	19.5	13.5	15.5	13.0
1	15.0	58.4	21.0	13.0	14.0	13.0
1	15.5	59.7	20.5	13.5	14.5	12.5
1	15.5	59.7	20.5	13.0	15.0	13.0
1	15.0	57.2	19.0	14.0	14.5	11.5
1	15.5	59.7	21.0	13.0	16.0	12.5
1	16.0	57.2	19.0	14.0	14.5	12.0
1	15.5	62.2	21.5	14.0	16.0	12.0
1	15.5	57.2	19.5	13.5	15.0	12.0
1	14.0	61.0	20.0	15.0	15.0	12.0
1	14.5	58.4	20.0	12.0	14.5	12.0
1	15.0	56.9	19.0	13.0	14.0	12.5
1	15.5	59.7	20.0	12.5	14.0	12.5
1	15.0	57.2	19.5	12.0	14.0	11.0
1	15.0	56.9	19.0	12.0	13.0	12.0
1	15.5	56.9	19.5	14.5	14.5	13.0
1	17.5	63.5	21.5	14.0	15.5	13.5
1	15.5	57.2	19.0	13.0	15.5	12.5
1	15.5	61.0	20.5	12.0	13.0	12.5
1	15.5	61.0	21.0	14.5	15.5	12.5
1	15.5	63.5	21.8	14.5	16.5	13.5
1	14.5	58.4	20.5	13.0	16.0	10.5
1	15.5	56.9	20.0	13.5	14.0	12.0
1	16.0	61.0	20.0	12.5	14.5	12.5
2	15.5	60.0	21.1	10.3	13.4	12.4
2	15.4	59.7	20.0	12.8	14.5	11.3
2	15.1	59.7	20.2	11.4	14.1	12.1
2	14.3	56.9	18.9	11.0	13.4	11.0
2	14.8	58.0	20.1	9.6	11.1	11.7
⋮	⋮	⋮	⋮	⋮	⋮	⋮
3	17.3	61.7	20.7	11.9	13.3	13.3

Thus the mean vectors lie largely in one dimension, and one discriminant function suffices to describe most of the separation among the three groups. \square

8.4.2 A Measure of Association for Discriminant Functions

Measures of association between the dependent variables y_1, y_2, \dots, y_p and the independent grouping variable i associated with $\mu_i, i = 1, 2, \dots, k$, were presented in Section 6.1.8. These measures attempt to answer the question, How well do the variables separate the groups? It was noted that Roy's statistic θ serves as an R^2 -like measure of association, since it is the ratio of between sum of squares to total sum of squares for the first discriminant function, $z_1 = \mathbf{a}'_1 \mathbf{y}$:

$$\eta_\theta^2 = \theta = \frac{\lambda_1}{1 + \lambda_1} = \frac{\text{SSH}(z_1)}{\text{SSE}(z_1) + \text{SSH}(z_1)}$$

[see (6.41) and (8.9)]. Another interpretation of η_θ^2 is the maximum squared correlation between the first discriminant function and the best linear combination of the $k - 1$ (dummy) group membership variables [see comments following (6.39) in Section 6.1.8]. Dummy variables were defined in the first two paragraphs of Section 6.1.8. The maximum correlation is called the (first) canonical correlation (see Chapter 11). The squared canonical correlation can be calculated for each discriminant function:

$$r_i^2 = \frac{\lambda_i}{1 + \lambda_i}, i = 1, 2, \dots, s. \quad (8.14)$$

The average squared canonical correlation was used as a measure of association in (6.48).

■ EXAMPLE 8.4.2

For the football data of Table 8.3, we obtain the squared canonical correlation between each of the two discriminant functions and the grouping variables,

$$r_1^2 = \frac{\lambda_1}{1 + \lambda_1} = \frac{1.9178}{1 + 1.9178} = .657,$$

$$r_2^2 = \frac{\lambda_2}{1 + \lambda_2} = \frac{.1159}{1 + .1159} = .104.$$

\square

8.5 STANDARDIZED DISCRIMINANT FUNCTIONS

In Section 5.5, it was noted that in the two-group case the relative contribution of the y 's to separation of the two groups can best be assessed by comparing the coefficients $a_r, r = 1, 2, \dots, p$, in the discriminant function

$$z = \mathbf{a}' \mathbf{y} = a_1 y_1 + a_2 y_2 + \dots + a_p y_p.$$

Similar comments appeared in Section 6.1.4, 6.1.8, and 6.4 concerning the use of discriminant functions to assess contribution of the y 's to separation of several groups. However, such comparisons are informative only if the y 's are commensurate, that is, measured on the same scale and with comparable variances. If the y 's are not commensurate, we need coefficients a_r^* that are applicable to standardized variables.

Consider the case of two groups. For the i th observation vector \mathbf{y}_{1i} or \mathbf{y}_{2i} in group 1 or 2, we can express the discriminant function in terms of standardized variables as

$$\begin{aligned} z_{1i} &= a_1^* \frac{y_{1i1} - \bar{y}_{11}}{s_1} + a_2^* \frac{y_{1i2} - \bar{y}_{12}}{s_2} + \cdots + a_p^* \frac{y_{1ip} - \bar{y}_{1p}}{s_p}, \\ i &= 1, 2, \dots, n_1, \\ z_{2i} &= a_1^* \frac{y_{2i1} - \bar{y}_{21}}{s_1} + a_2^* \frac{y_{2i2} - \bar{y}_{22}}{s_2} + \cdots + a_p^* \frac{y_{2ip} - \bar{y}_{2p}}{s_p}, \\ i &= 1, 2, \dots, n_2, \end{aligned} \quad (8.15)$$

where $\bar{\mathbf{y}}'_1 = (\bar{y}_{11}, \bar{y}_{12}, \dots, \bar{y}_{1p})$ and $\bar{\mathbf{y}}'_2 = (\bar{y}_{21}, \bar{y}_{22}, \dots, \bar{y}_{2p})$ are the mean vectors for the two groups and s_r is the within-sample standard deviation of the r th variable, obtained as the square root of the r th diagonal element of \mathbf{S}_{pl} . Clearly, these standardized coefficients must be of the form

$$a_r^* = s_r a_r, \quad r = 1, 2, \dots, p. \quad (8.16)$$

In vector form, this becomes

$$\mathbf{a}^* = (\text{diag } \mathbf{S}_{\text{pl}})^{1/2} \mathbf{a}. \quad (8.17)$$

For the several-group case, we can standardize the discriminant functions in an analogous fashion. If we denote the r th coefficient in the m th discriminant function by a_{mr} , $m = 1, 2, \dots, s$; $r = 1, 2, \dots, p$, then the standardized form is

$$a_{mr}^* = s_r a_{mr},$$

where s_r is the within-group standard deviation obtained from the diagonal of $\mathbf{S}_{\text{pl}} = \mathbf{E}/\nu_E$. Note that a_{mr}^* has two subscripts because there are several discriminant functions, while a_r^* in (8.16) above has only the one subscript because there is one discriminant function for two groups.

Alternatively, since the m th eigenvector is unique only up to multiplication by a scalar, we can simplify the standardization by using

$$a_{mr}^* = \sqrt{e_{rr}} a_{mr}, \quad r = 1, 2, \dots, p,$$

where e_{rr} is the r th diagonal element of \mathbf{E} . For further discussion of the use of standardized discriminant function coefficients to gauge the relative contribution of the variables to group separation, see Section 8.7.1 [see also Rencher and Scott (1990) and Rencher (1998, Section 5.4)].

■ EXAMPLE 8.5

In Example 8.4.1, we obtained the discriminant function coefficient vectors \mathbf{a}_1 and \mathbf{a}_2 for the football data of Table 8.3. Since $\lambda_1/(\lambda_1 + \lambda_2) = .94$, we concentrate on \mathbf{a}_1 . To standardize \mathbf{a}_1 , we need the within-sample standard deviations of the variables. The pooled covariance matrix is given by

$$\mathbf{S}_{\text{pl}} = \frac{\mathbf{E}}{87} = \begin{pmatrix} .428 & .578 & .158 & .084 & .125 & .228 \\ .578 & 3.161 & 1.020 & .653 & .340 & .505 \\ .158 & 1.020 & .546 & .077 & .129 & .159 \\ .084 & .653 & .077 & 1.232 & .315 & .024 \\ .125 & .340 & .129 & .315 & .618 & .009 \\ .228 & .505 & .159 & .024 & .009 & .376 \end{pmatrix}.$$

Using the square roots of the diagonal elements of \mathbf{S}_{pl} , we obtain

$$\mathbf{a}_1^* = \begin{pmatrix} \sqrt{.428}(-.948) \\ \sqrt{3.161}(.004) \\ \vdots \\ \sqrt{.376}(.829) \end{pmatrix} = \begin{pmatrix} -.621 \\ .007 \\ .005 \\ .719 \\ .397 \\ .508 \end{pmatrix}.$$

Thus the fourth, first, sixth, and fifth variables contribute most to separating the groups, in that order. The second and third variables are not useful (in the presence of the others) in distinguishing groups. \square

8.6 TESTS OF SIGNIFICANCE

In order to test hypotheses, we need the assumption of multivariate normality. This was not explicitly required for the development of discriminant functions.

8.6.1 Tests for the Two-Group Case

By (8.3) we see that the separation of transformed means, $(\bar{z}_1 - \bar{z}_2)^2/s_z^2$, achieved by the discriminant function $z = \mathbf{a}'\mathbf{y}$ is equivalent to the standardized distance between the mean vectors $\bar{\mathbf{y}}_1$ and $\bar{\mathbf{y}}_2$. This standardized distance is proportional to the two-group T^2 in (5.9) in Section 5.4.2. Hence the discriminant function coefficient vector \mathbf{a} is significantly different from $\mathbf{0}$ if T^2 is significant. More formally, if the population discriminant function coefficient vector is expressed as $\boldsymbol{\alpha} = \boldsymbol{\Sigma}^{-1}(\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2)$, then $H_0 : \boldsymbol{\alpha} = \mathbf{0}$ is equivalent to $H_0 : \boldsymbol{\mu}_1 = \boldsymbol{\mu}_2$.

To test the significance of a subset of the discriminant function coefficients, we can use the test of the corresponding subset of \mathbf{y} 's given in Section 5.9. To test the hypothesis that the population discriminant function has a specified form $\mathbf{a}'_0\mathbf{y}$, see Rencher (1998, Section 5.5.1).

8.6.2 Tests for the Several-Group Case

In Section 8.4.1 we noted that the discriminant criterion $\lambda = \mathbf{a}'\mathbf{H}\mathbf{a}/\mathbf{a}'\mathbf{E}\mathbf{a}$ is maximized by λ_1 , the largest eigenvalue of $\mathbf{E}^{-1}\mathbf{H}$, and that the remaining eigenvalues $\lambda_2, \dots, \lambda_s$ correspond to other discriminant dimensions. These eigenvalues are the same as those in the Wilks' Λ -test in (6.14) for significant differences among mean vectors,

$$\Lambda_1 = \prod_{i=1}^s \frac{1}{1 + \lambda_i}, \quad (8.18)$$

which is distributed as $\Lambda_{p,k-1,N-k}$, where $N = \sum_i n_i$ for an unbalanced design or $N = kn$ in the balanced case. Since Λ_1 is small if one or more λ_i 's are large, Wilks' Λ will test for significance of the eigenvalues and thereby for the discriminant functions. The s eigenvalues represent s dimensions of separation of the mean vectors $\bar{\mathbf{y}}_1, \bar{\mathbf{y}}_2, \dots, \bar{\mathbf{y}}_k$. We are interested in which, if any, of these dimensions are significant. In the context of discriminant functions, Wilks' Λ is more useful than the other three MANOVA test statistics, because it can be used on a subset of eigenvalues, as we see below.

In addition to the exact test provided by the critical values for Λ found in Table A.9, we can use the χ^2 -approximation for Λ_1 given in (6.16), with $\nu_E = N - k = \sum_i n_i - k$ and $\nu_H = k - 1$:

$$\begin{aligned} V_1 &= -[\nu_E - \frac{1}{2}(p - \nu_H + 1)] \ln \Lambda_1 \\ &= -[N - 1 - \frac{1}{2}(p + k)] \ln \prod_{i=1}^s \frac{1}{1 + \lambda_i} \\ &= [N - 1 - \frac{1}{2}(p + k)] \sum_{i=1}^s \ln(1 + \lambda_i), \end{aligned} \quad (8.19)$$

which is approximately χ^2 with $p(k - 1)$ degrees of freedom. The test statistic Λ_1 and its approximation (8.19) test the significance of all of $\lambda_1, \lambda_2, \dots, \lambda_s$. If the test leads to rejection of H_0 , we conclude that at least one of the λ 's is significantly different from zero, and therefore there is at least one dimension of separation of mean vectors. Since λ_1 is the largest, we are only sure of its significance, along with that of $z_1 = \mathbf{a}_1' \mathbf{y}$.

To test the significance of $\lambda_2, \lambda_3, \dots, \lambda_s$, we delete λ_1 from Wilks' Λ and the associated χ^2 -approximation to obtain

$$\Lambda_2 = \prod_{i=2}^s \frac{1}{1 + \lambda_i}, \quad (8.20)$$

$$\begin{aligned} V_2 &= -[N - 1 - \frac{1}{2}(p + k)] \ln \Lambda_2 \\ &= [N - 1 - \frac{1}{2}(p + k)] \sum_{i=2}^s \ln(1 + \lambda_i), \end{aligned} \quad (8.21)$$

which is approximately χ^2 with $(p-1)(k-2)$ degrees of freedom. If this test leads to rejection of H_0 , we conclude that at least λ_2 is significant along with the associated discriminant function $z_2 = \mathbf{a}_2' \mathbf{y}$. We can continue in this fashion, testing each λ_i in turn until a test fails to reject H_0 . (To compensate for making several tests, an adjustment to the α -level of each test could be made as in procedure 2, Section 5.5.) The test statistic at the m th step is

$$\Lambda_m = \prod_{i=m}^s \frac{1}{1 + \lambda_i}, \quad (8.22)$$

which is distributed as $\Lambda_{p-m+1, k-m, N-k-m+1}$. The statistic

$$\begin{aligned} V_m &= -[N - 1 - \frac{1}{2}(p+k)] \ln \Lambda_m \\ &= [N - 1 - \frac{1}{2}(p+k)] \sum_{i=m}^s \ln(1 + \lambda_i) \end{aligned} \quad (8.23)$$

has an approximate χ^2 -distribution with $(p-m+1)(k-m)$ degrees of freedom. In some cases, more λ 's will be statistically significant than the researcher considers to be of practical importance. If $\lambda_i / \sum_j \lambda_j$ is small, the associated discriminant function may not be of interest, even if it is significant.

We can also use an F -approximation for each Λ_i . For

$$\Lambda_1 = \prod_{i=1}^s \frac{1}{1 + \lambda_i},$$

we use (6.15), with $\nu_E = N - k$ and $\nu_H = k - 1$:

$$F = \frac{1 - \Lambda_1^{1/t}}{\Lambda_1^{1/t}} \frac{df_2}{df_1}, \quad (8.24)$$

where

$$\begin{aligned} t &= \sqrt{\frac{p^2(k-1)^2 - 4}{p^2 + (k-1)^2 - 5}}, & w &= N - 1 - \frac{1}{2}(p+k), \\ df_1 &= p(k-1), & df_2 &= wt - \frac{1}{2}[p(k-1) - 2]. \end{aligned}$$

For

$$\Lambda_m = \prod_{i=m}^s \frac{1}{1 + \lambda_i}, \quad m = 2, 3, \dots, s,$$

we use

$$F = \frac{1 - \Lambda_m^{1/t}}{\Lambda_m^{1/t}} \frac{df_2}{df_1} \quad (8.25)$$

with $p - m + 1$ and $k - m$ in place of p and $k - 1$:

$$\begin{aligned} t &= \sqrt{\frac{(p-m+1)^2(k-m)^2-4}{(p-m+1)^2+(k-m)^2-5}}, \\ w &= N-1-\frac{1}{2}(p+k), \\ df_1 &= (p-m+1)(k-m), \\ df_2 &= wt-\frac{1}{2}[(p-m+1)(k-m)-2]. \end{aligned}$$

■ EXAMPLE 8.6.2

We test the significance of the two discriminant functions obtained in Example 8.4.1 for the football data. For the overall test we have, by (8.18),

$$\Lambda_1 = \prod_{i=1}^2 \frac{1}{1+\lambda_i} = \frac{1}{1+1.9178} \frac{1}{1+.1159} = .307.$$

With $p = 6$, $k = 3$, and $N - k = 87$, the critical value from Table A.9 is $\Lambda_{.05,6,2,80} = .762$. By (8.19), the χ^2 -approximation is

$$\begin{aligned} V_1 &= -[N-1-\frac{1}{2}(p+k)] \ln \Lambda_1 \\ &= -[90-1-\frac{1}{2}(6+3)] \ln (.307) = 99.75, \end{aligned}$$

which exceeds the critical value $\chi^2_{.01,12} = 26.217$. Thus at least the first discriminant function is significant.

To test the second discriminant function, we have, by (8.20),

$$\Lambda_2 = \frac{1}{1+.1159} = .896.$$

With $m = 2$, the (conservative) critical value is $\Lambda_{.05,5,1,80} = .867$. Since this is close to $\Lambda = .896$, we interpolate in Table A.9 to obtain $\Lambda_{.05,5,1,86} = .875$. By (8.21), the χ^2 -approximation is

$$\begin{aligned} V_2 &= -[N-1-\frac{1}{2}(p+k)] \ln \Lambda_2 \\ &= -[90-1-\frac{1}{2}(6+3)] \ln \frac{1}{1+.1159} = 9.27 < \chi^2_{.05,5} = 11.070. \end{aligned}$$

For the F -approximation for Λ_1 , we obtain by (8.24)

$$t = \sqrt{\frac{p^2(k-1)^2-4}{p^2+(k-1)^2-5}} = \sqrt{\frac{6^22^2-4}{6^2+2^2-5}} = 2,$$

$$w = N-1-\frac{1}{2}(p+k) = 90-1-\frac{1}{2}(6+3) = 84.5,$$

$$df_1 = p(k-1) = 6(2) = 12,$$

$$df_2 = wt - \frac{1}{2}[p(k-1)-2] = (84.5)(2) - \frac{1}{2}[6(2)-2] = 164,$$

$$F = \frac{1-\Lambda_1^{1/2}}{\Lambda_1^{1/2}} \frac{df_2}{df_1} = \frac{1-.307^{1/2}}{.307^{1/2}} \frac{164}{12} = 10.994.$$

The p -value for $F = 10.994$ is less than .0001. For the F -approximation for Λ_2 , we reduce p and k by 1 and obtain by (8.25)

$$\begin{aligned} t &= \sqrt{\frac{5^2 1^2 - 4}{5^2 + 1^2 - 5}} = 1, \quad w = 90 - 1 - \frac{1}{2}(6 + 3) = 84.5, \\ df_1 &= 5(1) = 5, \quad df_2 = 84.5(1) - \frac{1}{2}[5(1) - 2] = 83, \\ F &= \frac{1 - \Lambda_2}{\Lambda_2} \cdot \frac{df_2}{df_1} = \frac{1 - .896}{.896} \frac{83}{5} = 1.924. \end{aligned}$$

The p -value for $F = 1.924$ is .099. Thus only the first discriminant function significantly separates groups. The exact test above using Λ_2 appears to be somewhat closer to rejection than the approximate tests. \square

8.7 INTERPRETATION OF DISCRIMINANT FUNCTIONS

There is a close correspondence between interpreting discriminant functions and determining the contribution of each variable, and we shall not always make a distinction. In interpretation, the signs of the coefficients are taken into account; in ascertaining the contribution, the signs are ignored and the coefficients are ranked in absolute value. (We discuss this distinction further in Section 8.7.1.) We are more commonly interested in assessing the contribution of the variables than in interpreting the function.

In the next three sections, we cover three common approaches to assessing the contribution of each variable (in the presence of the other variables) to separating the groups. The three methods are (1) examine the standardized discriminant function coefficients, (2) calculate a partial F -test for each variable, and (3) calculate a correlation between each variable and the discriminant function. The third method is the most widely recommended, but we note in Section 8.7.3 that it is the least useful.

8.7.1 Standardized Coefficients

To offset differing scales among the variables, the discriminant function coefficients can be standardized using (8.16) or (8.17), in which the coefficients have been adjusted so that they apply to standardized variables. For the observations in the first of two groups, for example, we have by (8.15),

$$z_{1i} = a_1^* \frac{y_{1i1} - \bar{y}_{11}}{s_1} + a_2^* \frac{y_{1i2} - \bar{y}_{12}}{s_2} + \cdots + a_p^* \frac{y_{1ip} - \bar{y}_{1p}}{s_p},$$

$$i = 1, 2, \dots, n_1.$$

The standardized variables $(y_{1ir} - \bar{y}_{1r})/s_r$ are scale free, and the standardized coefficients $a_r^* = s_r a_r$, $r = 1, 2, \dots, p$, therefore correctly reflect the joint contribution of the variables to the discriminant function z as it maximally separates the groups. For the case of several groups, each discriminant function coefficient vector

$\mathbf{a} = (a_1, a_2, \dots, a_p)'$ is an eigenvector of $\mathbf{E}^{-1}\mathbf{H}$, and as such, it takes into account the sample correlations among the variables as well as the influence of each variable in the presence of the others.

As noted in Section 8.5, this standardization is carried out for each of the s discriminant functions. Typically, each will have a different interpretation; that is, the pattern of the coefficients a_r^* will vary from one function to another.

The absolute values of the coefficients can be used to rank the variables in order of their contribution to separating the groups. If we wish to go further and interpret or "name" a discriminant function, the signs can be taken into account. Thus, for example, $z_1 = .8y_1 - .9y_2 + .5y_3$ has a different meaning than $z_2 = .8y_1 + .9y_2 + .5y_3$, since z_1 depends on the difference between y_1 and y_2 , while z_2 is related to the sum of y_1 and y_2 .

The discriminant function is subject to the same limitations as other linear combinations such as a regression equation: for example, (1) the coefficient for a variable may change notably if variables are added or deleted and (2) the coefficients may not be stable from sample to sample unless the sample size is large relative to the number of variables. With regard to limitation 1, we note that the coefficients reflect the contribution of each variable in the presence of the particular variables at hand. This is, in fact, what we want the coefficients to do. As to limitation 2, the processing of a substantial number of variables is not "free." More stable estimates will be obtained from 50 observations on 2 variables than from 50 observations on 20 variables. In other words, if N/p is too small, the variables that rank high in one sample may emerge as less important in another sample.

8.7.2 Partial F -Values

For any variable y_r we can calculate a partial F -test showing the significance of y_r after adjusting for the other variables, that is, the separation provided by y_r in addition to that due to the other variables. After computing the partial F for each variable, the variables can then be ranked.

In the case of two groups, the partial F is given by (5.32) as

$$F = (\nu - p + 1) \frac{T_p^2 - T_{p-1}^2}{\nu + T_{p-1}^2}, \quad (8.26)$$

where T_p^2 is the two-sample Hotelling T^2 with all p variables, T_{p-1}^2 is the T^2 -statistic with all variables except y_r , and $\nu = n_1 + n_2 - 2$. The F -statistic in (8.26) is distributed as $F_{1,\nu-p+1}$.

For the several-group case, the partial Λ for y_r adjusted for the other $p - 1$ variables is given by (6.127) as

$$\Lambda(y_r | y_1, \dots, y_{r-1}, y_{r+1}, \dots, y_p) = \frac{\Lambda_p}{\Lambda_{p-1}}, \quad (8.27)$$

where Λ_p is Wilks' Λ for all p variables and Λ_{p-1} involves all variables except y_r . The corresponding partial F is given by (6.128) as

$$F = \frac{1 - \Lambda}{\Lambda} \frac{\nu_E - p + 1}{\nu_H}, \quad (8.28)$$

where Λ is defined in (8.27), $\nu_E = N - k$, and $\nu_H = k - 1$. The partial Λ -statistic in (8.27) is distributed as $\Lambda_{1,\nu_H,\nu_E-p+1}$, and the partial F in (8.28) is distributed as F_{ν_H,ν_E-p+1} .

The partial F -values in (8.26) and (8.28) are not associated with a single dimension of group separation as are the standardized discriminant function coefficients. For example, y_2 will have a different contribution in each of the s discriminant functions, but the partial F for y_2 constitutes an overall index of the contribution of y_2 to group separation taking into account all dimensions. However, the partial F -values will often rank the variables in the same order as the standardized coefficients for the first discriminant function, especially if $\lambda_1 / \sum_j \lambda_j$ is very large so that the first function accounts for most of the available separation.

A partial index of association for y_r similar to the overall measure for all y 's given in (6.40), $\eta_\Lambda^2 = 1 - \Lambda$, can be defined by

$$R_r^2 = 1 - \Lambda_r \quad r = 1, 2, \dots, p, \quad (8.29)$$

where Λ_r is the partial Λ in (8.27) for y_r . This partial R^2 is a measure of association between the grouping variables and y_i after adjusting for the other $p - 1$ y 's.

8.7.3 Correlations Between Variables and Discriminant Functions

Many textbooks and research papers assert that the best measure of variable importance is the correlation between each variable and a discriminant function, $r_{y_i z_j}$. It is claimed that these correlations are more informative than standardized coefficients with respect to the joint contribution of the variables to the discriminant functions. The correlations are often referred to as loadings or structure coefficients and are routinely provided in many major programs. However, Rencher (1988; 1992b; 1998, Section 5.7) has shown that the correlations in question show the contribution of each variable in a univariate context rather than in a multivariate one. The correlations actually reproduce the t or F for each variable, and consequently they only show how each variable by itself separates the groups, ignoring the presence of the other variables. Hence these correlations provide no information about how the variables contribute jointly to separation of the groups. They become misleading if used for interpretation of discriminant functions.

Upon reflection, we could have anticipated this failure of the correlations to provide multivariate information. The objection to standardized coefficients is based on the argument that they are "unstable" because they change if some variables are deleted and others added. However, we actually want them to behave this way, so as to reflect the mutual influence of the variables on each other. In a multivariate analysis, interest is centered on the joint performance of the set of variables at hand.

To ask for the contribution of each variable independent of all other variables is to request a univariate index that ignores the other variables. The correlations $r_{y_i z_j}$ are “stable” and do not change when variables are added or deleted; this should be a clear signal that they are univariate in nature. There is no middle ground between the univariate and multivariate realms.

8.7.4 Rotation

Rotation of the discriminant function coefficients is sometimes recommended. This is a procedure (see Section 13.5) that attempts to produce a pattern with (absolute values of) coefficients closer to 0 or 1. Discriminant functions with such coefficients might be easier to interpret, but they have two deficiencies: they no longer maximize group separation, and they are correlated.

Accordingly, for interpretation of discriminant functions we recommend standardized coefficients rather than correlations or rotated coefficients.

8.8 SCATTERPLOTS

One benefit of the dimension reduction effected by discriminant analysis is the potential for plotting. It was noted in Section 6.2 that the number of large eigenvalues of $\mathbf{E}^{-1}\mathbf{H}$ reflects the dimensionality of the space occupied by the mean vectors. In many data sets, the first two discriminant functions account for most of $\lambda_1 + \lambda_2 + \dots + \lambda_s$, and consequently the pattern of the mean vectors can be effectively portrayed in a two-dimensional plot. If the essential dimensionality is greater than 2, there may be some distortion in intergroup configuration in a two-dimensional plot; that is, some groups that overlap in two dimensions may be well separated in a third dimension.

To plot the first two discriminant functions for the individual observation vectors \mathbf{y}_{ij} , simply calculate $z_{1ij} = \mathbf{a}'_1 \mathbf{y}_{ij}$ and $z_{2ij} = \mathbf{a}'_2 \mathbf{y}_{ij}$ for $i = 1, 2, \dots, k$; $j = 1, 2, \dots, n_i$, and plot a scatterplot of the $N = \sum_i n_i$ values of

$$\mathbf{z}_{ij} = \begin{pmatrix} z_{1ij} \\ z_{2ij} \end{pmatrix} = \begin{pmatrix} \mathbf{a}'_1 \mathbf{y}_{ij} \\ \mathbf{a}'_2 \mathbf{y}_{ij} \end{pmatrix} = \begin{pmatrix} \mathbf{a}'_1 \\ \mathbf{a}'_2 \end{pmatrix} \mathbf{y}_{ij} = \mathbf{A} \mathbf{y}_{ij}. \quad (8.30)$$

The transformed mean vectors,

$$\bar{\mathbf{z}}_i = \begin{pmatrix} \bar{z}_{1i} \\ \bar{z}_{2i} \end{pmatrix} = \begin{pmatrix} \mathbf{a}'_1 \\ \mathbf{a}'_2 \end{pmatrix} \bar{\mathbf{y}}_i = \mathbf{A} \bar{\mathbf{y}}_i, \quad i = 1, 2, \dots, k \quad (8.31)$$

should be plotted along with the individual values, \mathbf{z}_{ij} . In some cases, a plot would show only the transformed mean vectors $\bar{\mathbf{z}}_1, \bar{\mathbf{z}}_2, \dots, \bar{\mathbf{z}}_k$. For confidence regions for $\boldsymbol{\mu}_{z_i} = \mathbf{A} \boldsymbol{\mu}_i$, see Rencher (1998, Section 5.8).

We note that the eigenvalues of $\mathbf{E}^{-1}\mathbf{H}$ reveal the dimensionality of the mean vectors, not of the individual points. The dimensionality of the individual observations is p , although the essential dimensionality may be less because the variables are correlated. (The dimensionality of the observation vectors is the concern of principal




Figure 8.4 Scatterplot of discriminant function values for the football data of Table 8.3.

components; see Chapter 12.) If $s = 2$, for example, so that the mean vectors occupy only two dimensions, the individual observation vectors ordinarily lie in more than two-dimensions, and their inclusion in a plot constitutes a projection onto the two-dimensional plane of the mean vectors.

It was noted in Section 8.4.1 that the discriminant functions are uncorrelated but not orthogonal. Thus the angle between \mathbf{a}_1 and \mathbf{a}_2 as given by (3.14) is not 90° (that is, $\mathbf{a}_1' \mathbf{a}_2 \neq 0$). In practice, however, the usual procedure is to plot discriminant functions on a rectangular coordinate system. The resulting distortion is generally not serious.

■ EXAMPLE 8.8

Figure 8.4 contains a scatterplot of (z_1, z_2) for the observations in the football data of Table 8.3. Each observation in group 1 is denoted by a circle, observations in group 2 are denoted by triangles, and observations in group 3 are indicated by + signs. We see that the first discriminant function z_1 (the horizontal direction) effectively separates group 1 from groups 2 and 3, while the second discriminant function z_2 (the vertical direction) is less successful in separating group 2 from group 3.

The group mean vectors are indicated by solid circles. They are almost collinear, as we would expect since $\lambda_1 = 1.92$ dominates $\lambda_2 = .12$. \square

8.9 STEPWISE SELECTION OF VARIABLES

In many applications, a large number of dependent variables are available and the experimenter would like to discard those that are redundant (in the presence of the other variables) for separating the groups. Our discussion is limited to procedures that delete or add variables one at a time. We emphasize that we are selecting *dependent* variables (y 's) and therefore the basic model (one-way MANOVA) does not change. In subset selection in regression, on the other hand, we select *independent* variables, with a consequent alteration of the model.

A *forward selection* method was discussed in Section 6.11.2. We begin with a single variable, the one that maximally separates the groups by itself. Then the variable entered at each step is the one that maximizes the partial F -statistic based on Wilks' Λ , thus obtaining the maximal additional separation of groups above and beyond the separation already attained by the other variables. Since we choose the variable with maximum partial F at each step, the proportion of these maximum F 's that exceed F_α is greater than α . This bias is discussed in Rencher and Larson (1980) and Rencher (1998, Section 5.10).

Backward elimination is a similar operation in which we begin with all the variables and then at each step the variable that contributes least is deleted, as indicated by the partial F .

Stepwise selection is a combination of the forward and backward approaches. Variables are added one at a time, and at each step the variables are reexamined to see if any variable that entered earlier has become redundant in the presence of recently added variables. The procedure stops when the largest partial F among the variables available for entry fails to exceed a preset threshold value. The stepwise procedure has long been popular with practitioners. Some detail about the steps in this procedure was given in Section 6.11.2.

All the preceding procedures are commonly referred to as *stepwise discriminant analysis*. However, as noted in Section 6.11.2, we are actually doing stepwise MANOVA. No discriminant functions are calculated in the selection process. After the subset selection is completed, we can calculate discriminant functions for the selected variables. We could also use the variables in a classification analysis as described in Chapter 9.

■ EXAMPLE 8.9

We use the football data of Table 8.3 to illustrate the stepwise procedure outlined in this section and in Section 6.11.2. At the first step, we carry out a univariate F (using ordinary ANOVA) for each variable to determine which variable best separates the three groups by itself:

Variable	<i>F</i>	<i>p</i> -Value
WDIM	2.550	.0839
CIRCUM	6.231	.0030
FB EYE	1.668	.1947
EYEHD	58.162	1.11×10^{-16}
EARHD	22.427	1.40×10^{-8}
JAW	4.511	.0137

Thus EYEHD is the first variable to “enter.” The Wilks’ Λ value equivalent to $F = 58.162$ is $\Lambda(y_1) = .4279$ (see Table 6.1 with $p = 1$). At the second step we calculate a partial Λ and accompanying partial F using (8.27) and (8.28):

$$\Lambda(y_r|y_1) = \frac{\Lambda(y_1, y_r)}{\Lambda(y_1)},$$

$$F = \frac{1 - \Lambda(y_r|y_1)}{\Lambda(y_r|y_1)} \frac{\nu_E - 1}{\nu_H},$$

where y_1 indicates the variable selected at step 1 (EYEHD) and y_r represents each of the five variables to be examined at step 2. The results are

Variable	Partial Λ	Partial F	<i>p</i> -Value
WDIM	.9355	2.964	.0569
CIRCUM	.9997	.012	.9881
FB EYE	.9946	.235	.7911
EARHD	.9525	2.143	.1235
JAW	.9540	2.072	.1322

The variable WDIM would enter at this step, since it has the largest partial F . With a *p*-value of .0569, entering this variable may be questionable, but we will continue the procedure for illustrative purposes. We next check to see if EYEHD is still significant now that WDIM has entered. The partial Λ and F for EYEHD adjusted for WDIM are $\Lambda = .424$ and $F = 58.47$. Thus EYEHD stays “in.” The overall Wilks’ Λ for EYEHD and WDIM is $\Lambda(y_1, y_2) = .4003$.

At step 3 we check each of the four remaining variables for possible entry using

$$\Lambda(y_r|y_1, y_2) = \frac{\Lambda(y_1, y_2, y_r)}{\Lambda(y_1, y_2)},$$

$$F = \frac{1 - \Lambda(y_r|y_1, y_2)}{\Lambda(y_r|y_1, y_2)} \frac{\nu_E - 2}{\nu_H},$$

where $y_1 = \text{EYEHD}$, $y_2 = \text{WDIM}$, and y_r represents each of the other four variables. The results are

Variable	Partial Λ	Partial F	<i>p</i> -Value
CIRCUM	.9774	.981	.3793
FB EYE	.9748	1.098	.3381
EARHD	.9292	3.239	.0441
JAW	.8451	7.791	.0008

The indicated variable for entry at this step is JAW. To determine whether one of the first two variables should be removed after JAW has entered, we calculate the partial Λ and F for each, adjusted for the other two:

Variable	Partial Λ	Partial F	p -Value
WDIM	.8287	8.787	.0003
EYEHD	.4634	49.211	6.33×10^{-15}

Thus both previously entered variables remain in the model. The overall Wilks' Λ for EYEHD, WDIM, and JAW is $\Lambda(y_1, y_2, y_3) = .3383$.

At step 4 there are three candidate variables for entry. The partial Λ - and F -statistics are

$$\Lambda(y_r|y_1, y_2, y_3) = \frac{\Lambda(y_1, y_2, y_3, y_r)}{\Lambda(y_1, y_2, y_3)},$$

$$F = \frac{1 - \Lambda(y_r|y_1, y_2, y_3)}{\Lambda(y_r|y_1, y_2, y_3)} \frac{\nu_E - 3}{\nu_H},$$

where y_1 , y_2 , and y_3 are the three variables already entered and y_r represents each of the other three remaining variables. The results are

Variable	Partial Λ	Partial F	p -Value
CIRCUM	.9987	.055	.9462
FBEYE	.9955	.189	.8282
EARHD	.9080	4.257	.0173

Hence EARHD enters at this step, and we check to see if any of the three previously entered variables has now become redundant. The partial Λ and partial F for each of these three are

Variable	Partial Λ	Partial F	p -Value
WDIM	.7889	11.237	4.74×10^{-15}
EYEHD	.6719	20.508	5.59×10^{-8}
JAW	.8258	8.861	.0003

Consequently, all three variables are retained. The overall Wilks' Λ for all four variables is now $\Lambda(y_1, y_2, y_3, y_4) = .3072$.

At step 5, the partial Λ - and F -values are

Variable	Partial Λ	Partial F	p -Value
CIRCUM	.9999	.003	.9971
FBEYE	.9999	.004	.9965

Thus no more variables will enter.

We summarize the selection process as follows:

Step	Variable Entered	Overall Λ	Partial Λ	Partial F	p -Value
1	EYEHD	.4279	.4279	58.162	1.11×10^{-16}
2	WDIM	.4003	.9355	2.964	.0569
3	JAW	.3383	.8451	7.791	.0008
4	EARHD	.3072	.9080	4.257	.0173

□

PROBLEMS

- 8.1** Show that if $\mathbf{a} = \mathbf{S}_{p1}^{-1}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)$ is substituted into $[\mathbf{a}'(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)]^2 / \mathbf{a}'\mathbf{S}_{p1}\mathbf{a}$, the result is (8.3).
- 8.2** Verify (8.4) for the relationship between \mathbf{b} and \mathbf{a} .
- 8.3** Verify the relationship between R^2 and T^2 shown in (8.5).
- 8.4** Show that $[\mathbf{a}'(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)]^2 = \mathbf{a}'(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)'\mathbf{a}$ as in (8.7).
- 8.5** Show that $\mathbf{H}\mathbf{a} - \lambda\mathbf{E}\mathbf{a} = \mathbf{0}$ can be written in the form $(\mathbf{E}^{-1}\mathbf{H} - \lambda\mathbf{I})\mathbf{a} = \mathbf{0}$, as in (8.12).
- 8.6** Verify (8.16) by substituting $a_r^* = s_r a_r$ into (8.15) to obtain $z_{1i} = a_1 y_{1i1} + a_2 y_{1i2} + \cdots + a_p y_{1ip} - \mathbf{a}'\bar{\mathbf{y}}_1$.
- 8.7** For the psychological data in Table 5.1, the discriminant function coefficient vector was given in Example 5.5.
- (a) Find the standardized coefficients.
 - (b) Calculate t -tests for the individual variables.
 - (c) Compare the results of (a) and (b) as to the contribution of the variables to separation of the two groups.
 - (d) Find the partial F for each variable, as in (8.26), and compare with the standardized coefficients.
- 8.8** Using the beetle data of Table 5.5, do the following:
- (a) Find the discriminant function coefficient vector.
 - (b) Find the standardized coefficients.
 - (c) Calculate t -tests for individual variables.
 - (d) Compare the results of (b) and (c) as to the contribution of each variable to separation of the groups.
 - (e) Find the partial F for each variable, as in (8.26). Do the partial F 's rank the variables in the same order of importance as the standardized coefficients?

8.9 Using the dystrophy data of Table 5.7, do the following:

- (a) Find the discriminant function coefficient vector.
- (b) Find the standardized coefficients.
- (c) Calculate *t*-tests for individual variables.
- (d) Compare the results of (b) and (c) as to the contribution of each variable to separation of the groups.
- (e) Find the partial *F* for each variable, as in (8.26). Do the partial *F*'s rank the variables in the same order of importance as the standardized coefficients?

8.10 For the cyclical data of Table 5.8, do the following:

- (a) Find the discriminant function coefficient vector.
- (b) Find the standardized coefficients.
- (c) Calculate *t*-tests for individual variables.
- (d) Compare the results of (b) and (c) as to the contribution of each variable to separation of the groups.
- (e) Find the partial *F* for each variable, as in (8.26). Do the partial *F*'s rank the variables in the same order of importance as the standardized coefficients?

8.11 Using the fish data in Table 6.17, do the following:

- (a) Find the eigenvectors of $\mathbf{E}^{-1}\mathbf{H}$.
- (b) Carry out tests of significance for the discriminant functions and find the relative importance of each as in (8.13), $\lambda_i / \sum_j \lambda_j$. Do these two procedures agree as to the number of important discriminant functions?
- (c) Find the standardized coefficients and comment on the contribution of the variables to separation of groups.
- (d) Find the partial *F* for each variable, as in (8.28). Do they rank the variables in the same order as the standardized coefficients for the first discriminant function?
- (e) Plot the first two discriminant functions for each observation and for the mean vectors.

8.12 For the rootstock data of Table 6.2, do the following:

- (a) Find the eigenvalues and eigenvectors of $\mathbf{E}^{-1}\mathbf{H}$.
- (b) Carry out tests of significance for the discriminant functions and find the relative importance of each as in (8.13), $\lambda_i / \sum_j \lambda_j$. Do these two procedures agree as to the number of important discriminant functions?

- (c) Find the standardized coefficients and comment on the contribution of the variables to separation of groups.
- (d) Find the partial F for each variable, as in (8.28). Do they rank the variables in the same order as the standardized coefficients for the first discriminant function?
- (e) Plot the first two discriminant functions for each observation and for the mean vectors.

8.13 Carry out a stepwise selection of variables on the rootstock data of Table 6.2.

8.14 Carry out a stepwise selection of variables on the engineer data of Table 5.6.

8.15 Carry out a stepwise selection of variables on the fish data of Table 6.17.

CHAPTER 9

CLASSIFICATION ANALYSIS: ALLOCATION OF OBSERVATIONS TO GROUPS

9.1 INTRODUCTION

The *descriptive* aspect of discriminant analysis, in which group separation is characterized by means of discriminant functions, was covered in Chapter 8. We turn now to *allocation* of observations to groups, which is the *predictive* aspect of discriminant analysis. We prefer to call this *classification analysis* to clearly distinguish it from the descriptive aspect. However, classification is often referred to simply as discriminant analysis. In engineering and computer science, classification is usually called *pattern recognition*. Some writers use the term “classification analysis” to describe *cluster analysis*, in which the observations are clustered according to variable values rather than into predefined groups (see Chapter 15).

In classification, a sampling unit (subject or object) whose group membership is unknown is assigned to a group on the basis of the vector of p measured values, \mathbf{y} , associated with the unit. To classify the unit, we must have available a previously obtained sample of observation vectors from each group. Then one approach is to compare \mathbf{y} with the mean vectors $\bar{\mathbf{y}}_1, \bar{\mathbf{y}}_2, \dots, \bar{\mathbf{y}}_k$ of the k samples and assign the unit to the group whose $\bar{\mathbf{y}}_i$ is closest to \mathbf{y} .

In this chapter, the term “groups” may refer to either the k samples or the k populations from which they were taken. It should be clear from the context which of the two uses is intended in every case.

We give some examples to illustrate the classification technique:

1. A university admissions committee wants to classify applicants as likely to succeed or likely to fail. The variables available are high school grades in various subject areas, standardized test scores, rating of high school, number of advanced placement courses, etc.
2. A psychiatrist gives a battery of diagnostic tests in order to assign a patient to the appropriate mental illness category.
3. A college student takes aptitude and interest tests in order to determine which vocational area his or her profile best matches.
4. African, or “killer,” bees cannot be distinguished visually from ordinary domestic honey bees. Ten variables based on chromatograph peaks can be used to readily identify them (Lavine and Carlson 1987).
5. The Air Force wishes to classify each applicant into the training program where he or she has the most potential.
6. Twelve of the *Federalist Papers* were claimed by both Madison and Hamilton. Can we identify authorship by measuring frequencies of word usage (Mosteller and Wallace 1984)?
7. Variables such as availability of fingerprints, availability of eyewitnesses, and time until police arrive can be used to classify burglaries into solvable and unsolvable.
8. One approach to speech recognition by computer consists of an attempt to identify phonemes based on the energy levels in speech waves.
9. A number of variables are measured at five weather stations. Based on these variables, we wish to predict the ceiling at a particular airport in 2 hours. The ceiling categories are closed, low instrument, high instrument, low open, and high open (Lachenbruch 1975, p. 2).

9.2 CLASSIFICATION INTO TWO GROUPS

In the case of two populations, we have a sampling unit (subject or object) to be classified into one of two populations. The information we have available consists of the p variables in the observation vector \mathbf{y} measured on the sampling unit. In the first illustration in Section 9.1, for example, we have an applicant with high school grades and various test scores recorded in \mathbf{y} . We do not know whether the applicant will succeed or fail at the university, but we have data on previous students

at the university for whom it is now known whether they succeeded or failed. By comparing \mathbf{y} with $\bar{\mathbf{y}}_1$ for those who succeeded and $\bar{\mathbf{y}}_2$ for those who failed, we attempt to predict the group the applicant will eventually belong to.

When there are two populations, we can use a classification procedure due to Fisher (1936). The principal assumption for Fisher's procedure is that the two populations have the same covariance matrix. Normality is not required. We obtain a sample from each of the two populations and compute $\bar{\mathbf{y}}_1$, $\bar{\mathbf{y}}_2$, and \mathbf{S}_{pl} . A simple procedure for classification can be based on the discriminant function,

$$z = \mathbf{a}'\mathbf{y} = (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{\text{pl}}^{-1} \mathbf{y} \quad (9.1)$$

(see Sections 5.5, 5.6, 8.2, and 8.5), where \mathbf{y} is the vector of measurements on a new sampling unit that we wish to classify into one of the two groups (populations). For convenience we speak of "classifying \mathbf{y} " rather than classifying the subject or object associated with \mathbf{y} .

To determine whether \mathbf{y} is closer to $\bar{\mathbf{y}}_1$ or $\bar{\mathbf{y}}_2$, we check to see if z in (9.1) is closer to the transformed mean \bar{z}_1 or to \bar{z}_2 . We evaluate (9.1) for each observation \mathbf{y}_{1i} from the first sample and obtain $z_{11}, z_{12}, \dots, z_{1n_1}$, from which, by (3.54), $\bar{z}_1 = \sum_{i=1}^{n_1} z_{1i}/n_1 = \mathbf{a}'\bar{\mathbf{y}}_1 = (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{\text{pl}}^{-1} \bar{\mathbf{y}}_1$. Similarly, $\bar{z}_2 = \mathbf{a}'\bar{\mathbf{y}}_2$. Denote the two groups by G_1 and G_2 . Fisher's (1936) *linear classification procedure* assigns \mathbf{y} to G_1 if $z = \mathbf{a}'\mathbf{y}$ is closer to \bar{z}_1 than to \bar{z}_2 and assigns \mathbf{y} to G_2 if z is closer to \bar{z}_2 . This is illustrated in Figure 9.1.

For the configuration in Figure 9.1, we see that z is closer to \bar{z}_1 if

$$z > \frac{1}{2}(\bar{z}_1 + \bar{z}_2). \quad (9.2)$$

This is true in general because \bar{z}_1 is always greater than \bar{z}_2 , which can easily be shown as follows:

$$\bar{z}_1 - \bar{z}_2 = \mathbf{a}'(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2) = (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{\text{pl}}^{-1} (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2) > 0, \quad (9.3)$$

because $\mathbf{S}_{\text{pl}}^{-1}$ is positive definite. Thus $\bar{z}_1 > \bar{z}_2$. [If \mathbf{a} were of the form $\mathbf{a}' = (\bar{\mathbf{y}}_2 - \bar{\mathbf{y}}_1)' \mathbf{S}_{\text{pl}}^{-1}$, then $\bar{z}_2 - \bar{z}_1$ would be positive.] Since $\frac{1}{2}(\bar{z}_1 + \bar{z}_2)$ is the midpoint, $z > \frac{1}{2}(\bar{z}_1 + \bar{z}_2)$ implies that z is closer to \bar{z}_1 . By (9.3) the distance from \bar{z}_1 to \bar{z}_2 is the same as that from $\bar{\mathbf{y}}_1$ to $\bar{\mathbf{y}}_2$.

To express the classification rule in terms of \mathbf{y} , we first write $\frac{1}{2}(\bar{z}_1 + \bar{z}_2)$ in the form

$$\frac{1}{2}(\bar{z}_1 + \bar{z}_2) = \frac{1}{2}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{\text{pl}}^{-1} (\bar{\mathbf{y}}_1 + \bar{\mathbf{y}}_2). \quad (9.4)$$

Then the classification rule becomes: Assign \mathbf{y} to G_1 if

$$\mathbf{a}'\mathbf{y} = (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{\text{pl}}^{-1} \mathbf{y} > \frac{1}{2}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{\text{pl}}^{-1} (\bar{\mathbf{y}}_1 + \bar{\mathbf{y}}_2), \quad (9.5)$$

and assign \mathbf{y} to G_2 if

$$\mathbf{a}'\mathbf{y} = (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{\text{pl}}^{-1} \mathbf{y} < \frac{1}{2}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{\text{pl}}^{-1} (\bar{\mathbf{y}}_1 + \bar{\mathbf{y}}_2). \quad (9.6)$$




Figure 9.1 Fisher's procedure for classification into two groups.

This *linear classification rule* employs the same discriminant function $z = \mathbf{a}'\mathbf{y}$ used in Section 8.2 in connection with descriptive separation of groups. Thus in the two-group case the discriminant function serves as a linear classification function as well. However, in the several-group case in Section 9.3, we use classification functions that are different from the descriptive discriminant functions in Section 8.4.

Fisher's (1936) approach using (9.5) and (9.6) is essentially nonparametric because no distributional assumptions were made. However, if the two populations are normal with equal covariance matrices, then this method is (asymptotically) optimal; that is, the probability of misclassification is minimized [see comments following (9.9)].

If *prior probabilities* p_1 and p_2 are known for the two populations, the classification rule can be modified to exploit this additional information. We define the prior probabilities as follows: p_1 is the proportion of observations in G_1 and p_2 is the proportion in G_2 , where $p_2 = 1 - p_1$. For example, suppose that at a certain university 70% of entering freshmen ultimately graduate. Then $p_1 = .7$ and $p_2 = .3$.

If relative *costs of misclassification* are known for the two populations, the classification rule can be further modified. Let c_{12} be the cost of misclassifying an observation from group G_1 into group G_2 , and let c_{21} be the cost of misclassifying an observation from group G_2 into group G_1 . In practice, all we need to know is the relative costs. For example, it would only be necessary for us to know that c_{21} is twice the value of c_{12} (i.e., $c_{21}/c_{12} = 2$).

In order to use the prior probabilities and misclassification costs, the density functions for the two populations, $f(\mathbf{y}|G_1)$ and $f(\mathbf{y}|G_2)$, must also be known. Then the

optimal classification rule (Welch 1939) that minimizes the cost of misclassification is: Assign \mathbf{y} to G_1 if

$$p_1 c_{12} f(\mathbf{y}|G_1) > c_{21} p_2 f(\mathbf{y}|G_2) \quad (9.7)$$

and to G_2 otherwise. If misclassification costs are unknown or equal, we can say that the rule assigning \mathbf{y} to G_1 when

$$p_1 f(\mathbf{y}|G_1) > p_2 f(\mathbf{y}|G_2) \quad (9.8)$$

will minimize the probability of misclassification. Note that $f(\mathbf{y}|G_1)$ is a convenient notation for the density when sampling from the population represented by G_1 . It does not represent a conditional distribution in the usual sense (Section 4.2).

Assuming that the two densities are multivariate normal with equal covariance matrices, namely, $f(\mathbf{y}|G_1) = N_p(\boldsymbol{\mu}_1, \Sigma)$ and $f(\mathbf{y}|G_2) = N_p(\boldsymbol{\mu}_2, \Sigma)$, then from (9.7) we obtain the following rule (with estimates in place of $\boldsymbol{\mu}_1$, $\boldsymbol{\mu}_2$, and Σ): Assign \mathbf{y} to G_1 if

$$\mathbf{a}'\mathbf{y} = (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{\text{pl}}^{-1} \mathbf{y} > \frac{1}{2} (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{\text{pl}}^{-1} (\bar{\mathbf{y}}_1 + \bar{\mathbf{y}}_2) + \ln\left(\frac{p_2 c_{21}}{p_1 c_{12}}\right) \quad (9.9)$$

and to G_2 otherwise [see Rencher (1998, p. 231)]. Because we have substituted estimates for the parameters, the rule in (9.9) is no longer optimal as is (9.7). However, it is *asymptotically optimal* (approaches optimality as the sample size increases).

If $p_1 = p_2$ and $c_{12} = c_{21}$, the normal-based classification rule in (9.9) becomes the same as Fisher's procedure given in (9.5) and (9.6). Thus Fisher's rule, which is not based on a normality assumption, has optimal properties when the data come from multivariate normal populations with $\Sigma_1 = \Sigma_2$, $p_1 = p_2$, and $c_{12} = c_{21}$. [For the case when $\Sigma_1 \neq \Sigma_2$, see Rencher (1998, Section 6.2.2).] Hence, even though Fisher's method is nonparametric, it works better for normally distributed populations or other populations with linear trends. For example, suppose two populations have 95% contours, as in Figure 9.2. If the points are projected in any direction onto a straight line, there will be almost total overlap. A linear discriminant procedure will not successfully separate the two populations.

■ EXAMPLE 9.2

For the psychological data of Table 5.1, $\bar{\mathbf{y}}_1$, $\bar{\mathbf{y}}_2$, and \mathbf{S}_{pl} were obtained in Example 5.4.2. The discriminant function coefficients were obtained in Example 5.5 as $\mathbf{a}' = (.5104, -.2032, .4660, -.3097)$. For G_1 (the male group), we find

$$\begin{aligned} \bar{z}_1 &= \mathbf{a}'\bar{\mathbf{y}}_1 = .5104(15.97) - .2032(15.91) + .4660(27.19) - .3097(22.75) \\ &= 10.5427. \end{aligned}$$

Similarly, for G_2 (the female group), $\bar{z}_2 = \mathbf{a}'\bar{\mathbf{y}}_2 = 4.4426$. Thus we assign an observation vector \mathbf{y} to G_1 if

$$z = \mathbf{a}'\mathbf{y} > \frac{1}{2}(\bar{z}_1 + \bar{z}_2) = 7.4927$$




Figure 9.2 Two populations with nonlinear separation.

and assign \mathbf{y} to G_2 if $z < 7.4927$.

There are no new observations available, so we will illustrate the procedure by classifying two of the observations in G_1 . For $\mathbf{y}'_{11} = (15, 17, 24, 14)$, the first observation in G_1 , we have $z_{11} = \mathbf{a}'\mathbf{y}_{11} = .5104(15) - .2032(17) + .4660(24) - .3097(14) = 11.0498$, which is greater than 7.4927, and \mathbf{y}_{11} would be correctly classified as belonging to G_1 . For $\mathbf{y}'_{14} = (13, 12, 10, 16)$, the fourth observation in G_1 , we find $z_{14} = 3.9016$, which would misclassify \mathbf{y}_{14} into G_2 . \square

9.3 CLASSIFICATION INTO SEVERAL GROUPS

In this section we discuss classification rules for several groups. As in the two-group case, we use a sample from each of the k groups to find the sample mean vectors $\bar{\mathbf{y}}_1, \bar{\mathbf{y}}_2, \dots, \bar{\mathbf{y}}_k$. For a vector \mathbf{y} whose group membership is unknown, one approach is to use a distance function to find the mean vector that \mathbf{y} is closest to and assign \mathbf{y} to the corresponding group.

Note that for the k -group case with $k > 2$, incorporating misclassification costs into classification rules becomes complicated because there are $k(k - 1)$ different costs to consider. Hence, we assume equal misclassification costs for the discussion in this section.

9.3.1 Equal Population Covariance Matrices: Linear Classification Functions

In this section we assume $\Sigma_1 = \Sigma_2 = \dots = \Sigma_k$. We can estimate the common population covariance matrix by a pooled sample covariance matrix

$$\mathbf{S}_{\text{pl}} = \frac{1}{N - k} \sum_{i=1}^k (n_i - 1) \mathbf{S}_i = \frac{\mathbf{E}}{N - k},$$

where n_i and \mathbf{S}_i are the sample size and covariance matrix of the i th group, \mathbf{E} is the error matrix from one-way MANOVA, and $N = \sum_i n_i$. We compare \mathbf{y} to each $\bar{\mathbf{y}}_i, i = 1, 2, \dots, k$, by the distance function

$$D_i^2(\mathbf{y}) = (\mathbf{y} - \bar{\mathbf{y}}_i)' \mathbf{S}_{\text{pl}}^{-1} (\mathbf{y} - \bar{\mathbf{y}}_i) \quad (9.10)$$

and assign \mathbf{y} to the group for which $D_i^2(\mathbf{y})$ is smallest.

We can obtain a linear classification rule by expanding (9.10):

$$\begin{aligned} D_i^2(\mathbf{y}) &= \mathbf{y}' \mathbf{S}_{\text{pl}}^{-1} \mathbf{y} - \mathbf{y}' \mathbf{S}_{\text{pl}}^{-1} \bar{\mathbf{y}}_i - \bar{\mathbf{y}}_i' \mathbf{S}_{\text{pl}}^{-1} \mathbf{y} + \bar{\mathbf{y}}_i' \mathbf{S}_{\text{pl}}^{-1} \bar{\mathbf{y}}_i \\ &= \mathbf{y}' \mathbf{S}_{\text{pl}}^{-1} \mathbf{y} - 2\bar{\mathbf{y}}_i' \mathbf{S}_{\text{pl}}^{-1} \mathbf{y} + \bar{\mathbf{y}}_i' \mathbf{S}_{\text{pl}}^{-1} \bar{\mathbf{y}}_i. \end{aligned}$$

The term $\mathbf{y}' \mathbf{S}_{\text{pl}}^{-1} \mathbf{y}$ on the right can be neglected since it is not a function of i and, consequently, does not change from group to group. The second term is a linear function of \mathbf{y} , and the third does not involve \mathbf{y} . We thus delete $\mathbf{y}' \mathbf{S}_{\text{pl}}^{-1} \mathbf{y}$ and obtain a *linear classification function*, which we denote by $L_i(\mathbf{y})$. If we multiply by $-\frac{1}{2}$ so as to agree with the rule based on the normal distribution and prior probabilities given in (9.13) below, our linear classification rule becomes: Assign \mathbf{y} to the group for which

$$L_i(\mathbf{y}) = \bar{\mathbf{y}}_i' \mathbf{S}_{\text{pl}}^{-1} \mathbf{y} - \frac{1}{2} \bar{\mathbf{y}}_i' \mathbf{S}_{\text{pl}}^{-1} \bar{\mathbf{y}}_i, \quad i = 1, 2, \dots, k \quad (9.11)$$

is a *maximum* (we reversed the sign when multiplying by $-\frac{1}{2}$). To highlight the linearity of (9.11) as a function of \mathbf{y} , we can express it as

$$L_i(\mathbf{y}) = \mathbf{c}'_i \mathbf{y} + c_{i0} = c_{i1} y_1 + c_{i2} y_2 + \dots + c_{ip} y_p + c_{i0},$$

where $\mathbf{c}'_i = \bar{\mathbf{y}}_i' \mathbf{S}_{\text{pl}}^{-1}$ and $c_{i0} = -\frac{1}{2} \bar{\mathbf{y}}_i' \mathbf{S}_{\text{pl}}^{-1} \bar{\mathbf{y}}_i$. To assign \mathbf{y} to a group using this procedure, we calculate \mathbf{c}_i and c_{i0} for each of the k groups, evaluate $L_i(\mathbf{y}), i = 1, 2, \dots, k$, and allocate \mathbf{y} to the group for which $L_i(\mathbf{y})$ is largest. This will be the same group for which $D_i^2(\mathbf{y})$ in (9.10) is smallest, that is, the group whose mean vector $\bar{\mathbf{y}}_i$ is closest to \mathbf{y} .

For the case of several groups, the optimal rule in (9.8) extends to:

$$\text{Assign } \mathbf{y} \text{ to the group for which } p_i f(\mathbf{y}|G_i) \text{ is maximum.} \quad (9.12)$$

With this rule, the probability of misclassification is minimized. If we assume normality with equal covariance matrices and with prior probabilities of group membership, p_1, p_2, \dots, p_k , then $f(\mathbf{y}|G_i) = N_p(\boldsymbol{\mu}_i, \boldsymbol{\Sigma})$, and the rule in (9.12) becomes

(with estimates in place of parameters): Calculate

$$L'_i(\mathbf{y}) = \ln p_i + \bar{\mathbf{y}}'_i \mathbf{S}_{\text{pl}}^{-1} \mathbf{y} - \frac{1}{2} \bar{\mathbf{y}}'_i \mathbf{S}_{\text{pl}}^{-1} \bar{\mathbf{y}}_i, \quad i = 1, 2, \dots, k \quad (9.13)$$

and assign \mathbf{y} to the group with maximum value of $L'_i(\mathbf{y})$. Note that if $p_1 = p_2 = \dots = p_k$, then (9.13), which optimizes the classification rate for the normal distribution, reduces to (9.11), which was based on the heuristic approach of minimizing the distance of \mathbf{y} to $\bar{\mathbf{y}}_i$.

The linear functions $L_i(\mathbf{y})$ defined in (9.11) are called *linear classification functions* (many writers refer to them as *linear discriminant functions*). They are different from the linear discriminant functions in Sections 6.1.4, 6.4, and 8.4.1, whose coefficients are eigenvectors of $\mathbf{E}^{-1} \mathbf{H}$. In fact, there will be k classification functions and $s = \min(p, k - 1)$ discriminant functions, where k is the number of groups and p is the number of variables. In many cases we do not need all s discriminant functions to effectively describe group differences, whereas all k classification functions must be used in assigning observations to groups.

■ EXAMPLE 9.3.1

For the football data of Table 8.3, the mean vectors for the three groups are as follows:

$$\begin{aligned}\bar{\mathbf{y}}'_1 &= (15.2, 58.9, 20.1, 13.1, 14.7, 12.3), \\ \bar{\mathbf{y}}'_2 &= (15.4, 57.4, 19.8, 10.1, 13.5, 11.9), \\ \bar{\mathbf{y}}'_3 &= (15.6, 57.8, 19.8, 10.9, 13.7, 11.8).\end{aligned}$$

Using these values of $\bar{\mathbf{y}}_i$ and the pooled covariance matrix \mathbf{S}_{pl} , given in Example 8.5, the linear classification functions (9.11) become

$$\begin{aligned}L_1(\mathbf{y}) &= 7.6y_1 + 13.3y_2 + 4.2y_3 - 1.2y_4 + 14.6y_5 + 8.2y_6 - 641.1, \\ L_2(\mathbf{y}) &= 10.2y_1 + 13.3y_2 + 4.2y_3 - 3.4y_4 + 13.2y_5 + 6.1y_6 - 608.0, \\ L_3(\mathbf{y}) &= 10.9y_1 + 13.3y_2 + 4.1y_3 - 2.7y_4 + 13.1y_5 + 5.2y_6 - 614.6.\end{aligned}$$

We note that y_2 and y_3 have essentially the same coefficients in all three functions and hence do not contribute to classification of \mathbf{y} . These same two variables were eliminated in the stepwise discriminant analysis in Example 8.9.

We illustrate the use of these linear functions for the first and third observations in group 1. For the first observation, \mathbf{y}_{11} , we obtain

$$\begin{aligned}L_1(\mathbf{y}_{11}) &= 7.6(13.5) + 13.3(57.2) + 4.2(19.5) - 1.2(12.5) + 14.6(14.0) \\ &\quad + 8.2(11.0) - 641.1 = 582.124, \\ L_2(\mathbf{y}_{11}) &= 10.2(13.5) + 13.3(57.2) + 4.2(19.5) - 3.4(12.5) + 13.2(14.0) \\ &\quad + 6.1(11.0) - 608.0 = 578.099, \\ L_3(\mathbf{y}_{11}) &= 10.9(13.5) + 13.3(57.2) + 4.1(19.5) - 2.7(12.5) + 13.1(14.0) \\ &\quad + 5.2(11.0) - 614.6 = 578.760.\end{aligned}$$

We classify \mathbf{y}_{11} into group 1 since $L_1(\mathbf{y}_{11}) = 582.1$ exceeds $L_2(\mathbf{y}_{11})$ and $L_3(\mathbf{y}_{11})$.

For the third observation in group 1, \mathbf{y}_{13} , we obtain

$$L_1(\mathbf{y}_{13}) = 567.054, \quad L_2(\mathbf{y}_{13}) = 570.290, \quad L_3(\mathbf{y}_{13}) = 569.137.$$

This observation is misclassified into group 2 since $L_2(\mathbf{y}_{13}) = 570.290$ exceeds $L_1(\mathbf{y}_{13})$ and $L_3(\mathbf{y}_{13})$. \square

9.3.2 Unequal Population Covariance Matrices: Quadratic Classification Functions

The linear classification functions in Section 9.3.1 are based on the assumption $\Sigma_1 = \Sigma_2 = \dots = \Sigma_k$. The resulting classification rules are sensitive to heterogeneity of covariance matrices. Observations tend to be classified too frequently into groups whose covariance matrices have smaller variances on the diagonal.

As an example, consider a simple case with one y and two groups, where group G_1 has a normal distribution with a mean of -1 with a large standard deviation ($\sigma_1 = 2$) and group G_2 has a normal distribution with a mean of 5 with a small standard deviation ($\sigma_2 = 0.8$). The proper decision rule would utilize both the means and the standard deviations for the groups, assigning a new observation y to group G_1 if $y < 3.05$ and to group G_2 otherwise, where the cutoff at $y = 3.05$ corresponds to the location where $f_1(y|G_1) = f_2(y|G_2)$.

Figure 9.3 shows the distributions for the groups and illustrates the pitfall of erroneously assuming equality of covariance matrices in classification analysis. As illustrated by the hatched areas in Figure 9.3, when using the proper cutoff of roughly $y = 3.05$, about 2.1% of the observations from group G_1 will be misclassified with group G_2 and 0.7% of the observations from group G_2 will be misclassified with group G_1 . However, if the classification rules are calculated when erroneously assuming that the variances for the two groups are equal, the decision rule would assign a new observation to G_1 if $y < 2$ and to group G_2 otherwise, where the cutoff at $y = 2$ corresponds to the midpoint between the two group means. Figure 9.3 illustrates that this erroneous rule would result in 6.7% of the observations from group G_1 being misclassified with group G_2 and <0.1% of the observations from group G_2 being misclassified with group G_1 . Thus, erroneously assuming equality for the population covariance matrices will often result in classification rules that perform relatively poorly.

If $\Sigma_1 = \Sigma_2 = \dots = \Sigma_k$ does not hold, the classification rules can easily be altered to preserve optimality of classification rates. In place of (9.10), we can use

$$D_i^2(\mathbf{y}) = (\mathbf{y} - \bar{\mathbf{y}}_i)' \mathbf{S}_i^{-1} (\mathbf{y} - \bar{\mathbf{y}}_i), \quad i = 1, 2, \dots, k, \quad (9.14)$$

where \mathbf{S}_i is the sample covariance matrix for the i th group. As before, we would assign \mathbf{y} to the group for which $D_i^2(\mathbf{y})$ is smallest. With \mathbf{S}_i in place of \mathbf{S}_{pl} , (9.14) cannot be reduced to a linear function of \mathbf{y} as in (9.11) but remains a quadratic function. Hence rules based on \mathbf{S}_i are called *quadratic classification rules*.




Figure 9.3 Misclassification rates when correctly assuming variances are unequal (hatched area) and when incorrectly assuming variances are equal (gray area).

If we assume normality with unequal covariance matrices and with prior probabilities p_1, p_2, \dots, p_k , then $f(\mathbf{y}|G_i) = N_p(\boldsymbol{\mu}_i, \boldsymbol{\Sigma}_i)$, and the optimal rule in (9.12) based on $p_i f(\mathbf{y}|G_i)$ becomes: Assign \mathbf{y} to the group for which

$$Q_i(\mathbf{y}) = \ln p_i - \frac{1}{2} \ln |\mathbf{S}_i| - \frac{1}{2} (\mathbf{y} - \bar{\mathbf{y}}_i)' \mathbf{S}_i^{-1} (\mathbf{y} - \bar{\mathbf{y}}_i) \quad (9.15)$$

is maximum. If $p_1 = p_2 = \dots = p_k$ or if the p_i 's are unknown, the term $\ln p_i$ is deleted.

In order to use a quadratic classification rule based on \mathbf{S}_i , each n_i must be greater than p so that \mathbf{S}_i^{-1} will exist. This restriction does not apply to linear classification rules based on \mathbf{S}_{pl} . Since more parameters are estimated with quadratic classification functions, larger values of the n_i 's are needed for stability of estimates. Note the distinction between p , the number of variables, and p_i , the prior probability for the i th group.

9.4 ESTIMATING MISCLASSIFICATION RATES

In Chapter 8, we assessed the effectiveness of the discriminant functions in group separation by the use of significance tests or by examining $\lambda_i / \sum_j \lambda_j$. To judge the ability of classification procedures to predict group membership, we usually use the probability of misclassification, which is known as the *error rate*. We could also use its complement, the *correct classification rate*.

A simple estimate of the error rate can be obtained by trying out the classification procedure on the same data set that has been used to compute the classification

functions. This method is commonly referred to as *resubstitution*. Each observation vector \mathbf{y}_{ij} is submitted to the classification functions and assigned to a group. We then count the number of correct classifications and the number of misclassifications. The proportion of misclassifications resulting from resubstitution is called the *apparent error rate*. The results can be conveniently displayed in a *classification table* or *confusion matrix*, such as Table 9.1 for two groups.

Among the n_1 observations in G_1 , n_{11} are correctly classified into G_1 and n_{12} are misclassified into G_2 , where $n_1 = n_{11} + n_{12}$. Similarly, of the n_2 observations in G_2 , n_{21} are misclassified into G_1 and n_{22} are correctly classified into G_2 , where $n_2 = n_{21} + n_{22}$. Thus

$$\begin{aligned}\text{Apparent error rate} &= \frac{n_{12} + n_{21}}{n_1 + n_2} \\ &= \frac{n_{12} + n_{21}}{n_{11} + n_{12} + n_{21} + n_{22}}.\end{aligned}\quad (9.16)$$

Similarly, we can define

$$\text{Apparent correct classification rate} = \frac{n_{11} + n_{22}}{n_1 + n_2}.\quad (9.17)$$

Clearly,

$$\text{Apparent error rate} = 1 - \text{apparent correct classification rate}.$$

The method of resubstitution can be readily extended to the case of several groups.

The apparent error rate is easily obtained and is routinely provided by most classification software programs. It is an estimate of the probability that our classification functions based on the present sample will misclassify a future observation. This probability is called the *actual error rate*. Unfortunately, the apparent error rate underestimates the actual error rate because the data set used to compute the classification functions is also used to evaluate them. The classification functions are optimized for the particular sample and may be capitalizing on chance to some degree, especially for small samples. For other estimates of error rates, see Rencher (1998, Section 6.4). In Section 9.5 we consider some approaches to reducing the bias in the apparent error rate.

Table 9.1 Classification Table for Two Groups

Actual Group	Number of Observations	Predicted Group	
		1	2
1	n_1	n_{11}	n_{12}
2	n_2	n_{21}	n_{22}

Table 9.2 Classification Table for the Psychological Data of Table 5.1

Actual Group	Number of Observations	Predicted Group	
		Male	Female
Male	32	28	4
Female	32	4	28

■ EXAMPLE 9.4(a)

We use the psychological data of Table 5.1 to illustrate the apparent error rate obtained by the resubstitution method for two groups. The hypothesis $H_0 : \Sigma_1 = \Sigma_2$ was not rejected in Example 7.3.2, and we therefore classify each of the 64 observations using the linear classification procedure obtained in Example 9.2: Classify as G_1 if $\mathbf{a}'\mathbf{y} > 7.4927$ and as G_2 otherwise. The resulting classification table is given in Table 9.2. By (9.16),

$$\text{Apparent error rate} = \frac{n_{12} + n_{21}}{n_1 + n_2} = \frac{4 + 4}{32 + 32} = .125.$$

□

■ EXAMPLE 9.4(b)

We use the football data of Table 8.3 to illustrate the use of the resubstitution method for estimating the error rate in the case of more than two groups. The sample covariance matrices for the three groups are almost significantly different, and we will use both linear and quadratic classification functions.

The linear classification functions $L_i(\mathbf{y})$ from (9.11) were given in Example 9.3.1 for the football data. Using these, we classify each of the 90 observations. The results are shown in Table 9.3.

An examination of this data set in Example 8.8 showed that groups 2 and 3 are harder to separate than 1 and 2 or 1 and 3. This pattern is reflected here in the misclassifications. Only 4 of the observation vectors in group 1 are misclassified, while 10 observations in each of groups 2 and 3 are misclassified.

Using the quadratic classification functions $Q_i(\mathbf{y})$, $i = 1, 2, 3$, in (9.15) and assuming $p_1 = p_2 = p_3$, we obtain the classification results in Table 9.4. There is some improvement in the apparent error rate using quadratic classification functions. □

9.5 IMPROVED ESTIMATES OF ERROR RATES

For large samples, the apparent error rate has only a small amount of bias for estimating the actual error rate and can be used with little concern. For small samples,

Table 9.3 Classification Table for the Football Data of Table 8.3 Using Linear Classification Functions

Actual Group	Number of Observations	Predicted Group		
		1	2	3
1	30	26	1	3
2	30	1	20	9
3	30	2	8	20

$$\text{Apparent correct classification rate} = \frac{26 + 20 + 20}{90} = .733$$

$$\text{Apparent error rate} = 1 - .733 = .267$$

Table 9.4 Classification Table for the Football Data of Table 8.3 Using Quadratic Classification Functions

Actual Group	Number of Observations	Predicted Group		
		1	2	3
1	30	27	1	2
2	30	2	21	7
3	30	1	4	25

$$\text{Apparent correct classification rate} = \frac{27 + 21 + 25}{90} = .811$$

$$\text{Apparent error rate} = 1 - .811 = .189$$

however, it is overly optimistic (biased downward), as noted above. We discuss two techniques for reducing the bias in the apparent error rate, that is, increasing the apparent error rate to a more realistic level.

9.5.1 Partitioning the Sample

One way to avoid bias is to split the sample into two parts, a *training* sample used to construct the classification rule and a *validation* sample used to evaluate it. With the training sample, we calculate linear or quadratic classification functions. We then submit each observation vector in the validation sample to the classification functions obtained from the training sample. Since these observations are not used in calculating the classification functions, the resulting error rate is unbiased. To increase the information gained, we could also reverse the roles of the two samples so that the classification functions are obtained from the validation sample and evaluated on the training sample. The two estimates of error could then be averaged.

Partitioning the sample has at least two disadvantages:

1. It requires large samples that may not be available.
2. It does not evaluate the classification function we will use in practice. The estimate of error based on half the sample may vary considerably from that based on the entire sample. We prefer to use all or almost all of the data to construct the classification functions so as to minimize the variance of our error rate estimate.

9.5.2 Holdout Method

The *holdout method* is an improved version of the sample splitting procedure in Section 9.5.1. In the holdout procedure, all but one observation is used to compute the classification rule, and this rule is then used to classify the omitted observation. We repeat this procedure for each observation, so that, in a sample of size $N = \sum_i n_i$, each observation is classified by a function based on the other $N - 1$ observations. The computation load is increased because N distinct classification procedures have to be constructed. The holdout procedure is also referred to as the *leaving-one-out method* or as *cross-validation*. Note that this procedure is used to estimate error rates. The actual classification rule for future observations would be based on all N observations.

■ EXAMPLE 9.5.2

We use the football data of Table 8.3 to illustrate the holdout method for estimating the error rate. Each of the 90 observations is classified by linear classification functions based on the other 89 observations. To begin the procedure, the first observation in group 1 (y_{11}) is held out and the linear classification functions $L_i(\mathbf{y}), i = 1, 2, 3$, in (9.11) are calculated using the remaining 29 observations in group 1 and the 60 observations in groups 2 and 3. The observation y_{11} is now classified using $L_1(\mathbf{y}), L_2(\mathbf{y})$, and $L_3(\mathbf{y})$. Then y_{11} is reinserted in group 1, and y_{12} is held out. The functions $L_1(\mathbf{y}), L_2(\mathbf{y})$, and $L_3(\mathbf{y})$ are recomputed and y_{12} is then classified. This procedure is followed for each of the 90 observations, and the results are in Table 9.5.

As expected, the holdout error rate has increased somewhat from the apparent error rate based on resubstitution in Tables 9.3 and 9.4 in Example 9.4(b). An error rate of .300 is a less optimistic (more realistic) estimate of what the classification functions can do with future samples. \square

9.6 SUBSET SELECTION

The experimenter often has available a large number of variables and wishes to keep any that might aid in predicting group membership, but at the same time to delete any superfluous variables that do not contribute to allocation. A reduction in the number of redundant variables may in fact lead to improved error rates. As an additional con-

Table 9.5 Classification Table for the Football Data of Table 8.3 Using the Holdout Method Based on Linear Classification Functions

Actual Group	Number of Observations	Predicted Group		
		1	2	3
1	30	26	1	3
2	30	1	18	11
3	30	2	9	19

$$\text{Correct classification rate} = \frac{26 + 18 + 19}{90} = .700$$

$$\text{Error rate} = 1 - .700 = .300$$

sideration, there is an increase in robustness to nonnormality of linear and quadratic classification functions as p (the number of variables) decreases.

The majority of selection schemes for classification analysis are based on stepwise discriminant analysis or a similar approach (Section 8.9). One finds the subset of variables that best separates groups using Wilks' Λ , for example, and then uses these variables to construct classification functions. Most of the major statistical software packages offer this method. When the "best" subset is selected in this way, an optimistic bias in error rates is introduced. For a discussion of this bias, see Rencher (1992a; 1998, Section 6.7).

Another link between separation and classification is the use of error rates in an informal stopping rule in a stepwise discriminant analysis. Thus, for example, if a subset of 5 variables out of 10 gives a misclassification rate of 33% compared to 30% for the full set of variables, we may decide that the 5 variables are adequate for separating the groups. We could try several subsets of decreasing sizes to see when the error rate begins to escalate noticeably.

■ EXAMPLE 9.6(a)

In Example 8.9, a stepwise discriminant analysis based on a partial Wilks' Λ (or partial F) was carried out for the football data of Table 8.3. Four variables were selected: EYEHD, WDIM, JAW, and EARHD. These same four variables are indicated by the coefficients in the linear classification functions in Example 9.3.1. We now use these four variables to classify the observations using the method of resubstitution to obtain the apparent error rate.

The linear classification functions (9.11) are

$$\begin{aligned} \text{Group 1 } L_1(\mathbf{y}) &= \bar{\mathbf{y}}_1' \mathbf{S}_{\text{pl}}^{-1} \mathbf{y} - \frac{1}{2} \bar{\mathbf{y}}_1' \mathbf{S}_{\text{pl}}^{-1} \bar{\mathbf{y}}_1 \\ &= 18.67y_1 + 4.13y_2 + 17.67y_3 + 20.44y_4 - 425.50, \end{aligned}$$

$$\text{Group 2 } L_2(\mathbf{y}) = 21.13y_1 + 1.96y_2 + 16.24y_3 + 18.36y_4 - 392.75,$$

$$\text{Group 3 } L_3(\mathbf{y}) = 21.87y_1 + 2.67y_2 + 16.13y_3 + 17.46y_4 - 399.63.$$

Table 9.6 Classification Table for the Football Data of Table 8.3 Using Linear Classification Functions Based on Four Variables Chosen by Stepwise Selection

Actual Group	Number of Observations	Predicted Group		
		1	2	3
1	30	26	1	3
2	30	1	20	9
3	30	2	8	20

$$\text{Correct classification rate} = \frac{26 + 20 + 20}{90} = .733$$

$$\text{Error rate} = 1 - .733 = .267$$

When each observation vector is classified using these linear functions, we obtain the classification results in Table 9.6.

Table 9.6 is identical to Table 9.3 in Example 9.4(b) where all six variables were used. Thus the four selected variables can classify the sample as well as all six variables classify it. \square

■ EXAMPLE 9.6(b)

We illustrate the use of error rates as an informal stopping rule in a stepwise discriminant analysis. Fifteen teacher and pupil behaviors were observed during 5-minute intervals of reading instruction in elementary school classrooms (Rencher, Wadham, and Young 1978). The observations were recorded in rate of occurrences per minute for each variable. The variables were the following:

Teacher Behaviors

1. *Explains*—Explains the task to learner.
2. *Models*—Models the task response for the learner.
3. *Questions*—Asks a question to elicit a task response.
4. *Directs*—Gives a direct signal to elicit a task response.
5. *Controls*—Controls management behavior with direction statements or gestures.
6. *Positive*—Gives a positive (affirmative) statement or gesture.
7. *Negative*—Gives a negative statement or gesture.

Pupil Behaviors

8. *Overt delayed*—An overt learner response to task signals that cannot be judged correct or incorrect until later.

Table 9.7 Stepwise Selection Statistics for the Teacher Data

Number of Variables	Overall Wilks' Λ	Percentage of Correct Classification
15	.132	77.4
10	.159	72.4
9	.170	73.3
8	.182	70.6
7	.195	72.9
6	.211	70.1
5	.231	70.6
4	.256	65.6

9. *Correct*—A correct learner response with relationship to task signals.
10. *Incorrect*—An incorrect learner response with relationship to task signals.
11. *No response*—Learner gives no response with relationship to task signals.
12. *Asks*—Learner asks a question about the task.
13. *Statement*—Learner gives a positive statement or gestures not related to the task.
14. *Inappropriate*—Learner gives inappropriate management behavior.
15. *Other*—Other learner than one being observed gives responses as teacher directs task signals.

The teachers were grouped into four categories:

Group 1: Outstanding teachers

Group 2: Poor teachers

Group 3: First-year teachers

Group 4: Teacher aides

The sample sizes in groups 1-4 were 62, 61, 57, and 41, respectively. Because of the large values of N and p ($N = 221, p = 15$), the data are not given here.

The stepwise discriminant analysis was run several times with different threshold F -to-enter values so as to select subsets with different sizes. A classification analysis based on resubstitution was carried out with each of the resulting subsets of variables. In Table 9.7, we compare the overall Wilks' Λ and the apparent correct classification rate.

According to the correct classification rate, we would choose to stop at five variables because of the abrupt change from 5 to 4. On the other hand, the changes in Wilks' Λ are more gradual, and no clear stopping point is indicated. \square

9.7 NONPARAMETRIC PROCEDURES

We have previously discussed both parametric and nonparametric classification procedures. Welch's optional rule in (9.8) and (9.12) is parametric, whereas Fisher's linear classification rule for two groups as given in (9.5) and (9.6) is essentially nonparametric, since no distributional assumptions were involved in its derivation. However, Fisher's procedure also turns out to be equivalent to the optimal normal-based approach in (9.9). Nonparametric procedures for estimating error rate include the resubstitution and holdout methods. In the sections below, we discuss additional nonparametric classification procedures.

9.7.1 Multinomial Data

We now consider data in which an observation vector consists of responses on each of several categorical variables. The various combinations of categories constitute the possible outcomes of a multinomial random variable. For example, consider the following four categorical variables: gender (male or female), political party (Republican, Democrat, other), size of city of residence (under 10,000, between 10,000 and 100,000, over 100,000), and education (less than high school graduation, high school graduate, college graduate, advanced degree). An observation vector might be $(2, 1, 3, 4)$, that is, a female Republican who lives in a city of over 100,000 and has an advanced degree. The total number of possible outcomes in this multinomial distribution is the product of the number of states of the individual variables: $2 \times 3 \times 3 \times 4 = 72$. We will use this example to illustrate classification procedures for multinomial data. Suppose we are attempting to predict whether or not a person will vote. Then there are two groups, G_1 and G_2 , and we assign a person to one of the groups after observing which of the 72 possible outcomes he or she gives.

Welch's (1939) optimum rule given in (9.8) can be written as: Assign \mathbf{y} to G_1 if

$$\frac{f(\mathbf{y}|G_1)}{f(\mathbf{y}|G_2)} > \frac{p_2}{p_1} \quad (9.18)$$

and to G_2 otherwise. In our categorical example, $f(\mathbf{y}|G_1)$ is represented by $q_{1i}, i = 1, 2, \dots, 72$, and $f(\mathbf{y}|G_2)$ becomes $q_{2i}, i = 1, 2, \dots, 72$, where q_{1i} is the probability that a person in group 1 will give the i th outcome, with an analogous definition for q_{2i} . In terms of these multinomial probabilities, the classification rule in (9.18) becomes: If a person gives the i th outcome, assign him or her to G_1 if

$$\frac{q_{1i}}{q_{2i}} > \frac{p_2}{p_1} \quad (9.19)$$

and to G_2 otherwise. If the probabilities q_{1i} and q_{2i} were known, it would be easy to check (9.19) for each i and partition the 72 possible outcomes into two subsets, those for which the person would be assigned to G_1 and those corresponding to G_2 .

The values of q_{1i} and q_{2i} are usually unknown and must be estimated from a sample. Let n_{1i} and n_{2i} be the numbers of persons in groups 1 and 2 who give the i th outcome, $i = 1, 2, \dots, 72$. Then we estimate q_{1i} and q_{2i} by

$$\hat{q}_{1i} = \frac{n_{1i}}{N_1} \quad \text{and} \quad \hat{q}_{2i} = \frac{n_{2i}}{N_2}, \quad i = 1, 2, \dots, 72, \quad (9.20)$$

where $N_1 = \sum_i n_{1i}$ and $N_2 = \sum_i n_{2i}$. However, a large sample size would be required for stable estimates; in any given example, some of the n 's may be zero.

Multinomial data can also be classified by ordinary linear classification functions. We must distinguish between ordered and unordered categories. If all of the variables have ordered categories, the data can be submitted directly to an ordinary classification program. In the above example, city size and education are variables of this type. It is customary to assign ordered categories ranked values such as 1, 2, 3, 4. It has been shown that linear classification functions perform reasonably well on (ordered) discrete data of this type [see Lachenbruch (1975, p.45), Titterington et al. (1981), and Gilbert (1968)].

Unordered categorical variables cannot be handled this same way. Thus the political party variable in the example above should not be coded 1, 2, 3 and entered into the computation of the classification functions. However, an unordered categorical variable with k categories can be replaced by $k - 1$ dummy variables (see Sections 6.1.8 and 11.6.2) for use with linear classification functions. For example, the political preference variable with three categories can be converted to two dummy variables as follows:

$$y_1 = \begin{cases} 1 & \text{if Republican} \\ 0 & \text{otherwise} \end{cases} \quad y_2 = \begin{cases} 1 & \text{if Democrat} \\ 0 & \text{otherwise.} \end{cases}$$

Thus the (y_1, y_2) pair takes the value (1,0) for a Republican, (0,1) for a Democrat, and (0,0) for other. Many software programs will create dummy variables automatically. Note that if a subset selection program is used, the dummy variables for a given categorical variable must be kept together; that is, they must all be included in the chosen subset or all excluded, because all are necessary to describe the categorical variable.

In some cases, such as in medical data collection, there is a mixture of continuous and categorical variables. Various approaches to classification with such data have been discussed by Krzanowski (1975, 1976, 1977, 1979, 1980), Lachenbruch and Goldstein (1979), Tu and Han (1982), and Bayne et al. (1983). See Rencher (1998, Section 6.8) for a discussion of logistic and probit classification, which are useful for certain types of continuous and discrete data that are not normally distributed.

9.7.2 Classification Based on Density Estimators

In (9.9), (9.13), and (9.15) we have linear and quadratic classification rules based on the multivariate normal density and prior probabilities. These normal-based rules

arose from Welch's optimal rule that assigns \mathbf{y} to the group for which $p_i f(\mathbf{y}|G_i)$ is maximum. If the form of $f(\mathbf{y}|G_i)$ is nonnormal and unknown, the density can be estimated directly from the data. The approach we describe is known as the *kernel* estimator.

We first describe the kernel method for a univariate continuous random variable y . Suppose y has density $f(y)$, which we wish to estimate using a sample y_1, y_2, \dots, y_n . A simple estimate of $f(y_0)$ for an arbitrary point y_0 can be based on the proportion of points in the interval $(y_0 - h, y_0 + h)$. If the number of points in the interval is denoted by $N(y_0)$, then the proportion $N(y_0)/n$ is an estimate of $P(y_0 - h < y < y_0 + h)$, which is approximately equal to $2hf(y_0)$. Thus we estimate $f(y_0)$ by

$$\hat{f}(y_0) = \frac{N(y_0)}{2hn}. \quad (9.21)$$

We can express $\hat{f}(y_0)$ as a function of all y_i in the sample by defining

$$K(u) = \begin{cases} \frac{1}{2} & \text{for } |u| \leq 1 \\ 0 & \text{for } |u| > 1 \end{cases}, \quad (9.22)$$

so that $N(y_0) = 2 \sum_{i=1}^n K[(y_0 - y_i)/h]$, and (9.21) becomes

$$\hat{f}(y_0) = \frac{1}{hn} \sum_{i=1}^n K\left(\frac{y_0 - y_i}{h}\right). \quad (9.23)$$

The function $K(u)$ is called the *kernel*. In (9.23), $K[(y_0 - y_i)/h]$ is $\frac{1}{2}$ for any point y_i in the interval $(y_0 - h, y_0 + h)$ and is zero for points outside the interval. Points in the interval add $1/2hn$ to the density, and points outside the interval contribute nothing.

Kernel estimators were first proposed by Rosenblatt (1956) and Parzen (1962). A good review of nonparametric density estimation including kernel estimators has been given by Silverman (1986), who noted that classification analysis provided the initial motivation for the development of density estimation.

The kernel defined by (9.22) is rectangular and the graph of $\hat{f}(y_0)$ plotted as a function of y_0 will be a step function, since there will be a jump (or drop) whenever y_0 is a distance h from one of the y_i 's. (A moving average has a similar property.)

To obtain a smooth estimator of $f(y)$, we must choose a smooth kernel. Two possibilities are

$$K(u) = \frac{1}{\pi} \frac{\sin^2 u}{u^2}, \quad (9.24)$$

$$K(u) = \frac{1}{\sqrt{2\pi}} e^{-u^2/2}, \quad (9.25)$$

which have the property that all n sample points y_1, y_2, \dots, y_n contribute to $\hat{f}(y_0)$ with the closest points weighted heavier than the more distant points. Even though

$K(u)$ in (9.25) has the form of the normal distribution, this does not imply any assumption about the density $f(y)$. We have used the normal density function because it is symmetric and unimodal. Other density functions could be used as kernels.

Cacoullos (1966) provided kernel estimates for multivariate density functions; see also Scott (1992). If $\mathbf{y}'_0 = (y_{01}, y_{02}, \dots, y_{0p})$ is an arbitrary point whose density we wish to estimate, then the extension of (9.23) is

$$\hat{f}(\mathbf{y}_0) = \frac{1}{nh_1 h_2 \cdots h_p} \sum_{i=1}^n K\left(\frac{y_{01} - y_{i1}}{h_1}, \dots, \frac{y_{0p} - y_{ip}}{h_p}\right). \quad (9.26)$$

An estimate $\hat{f}(\mathbf{y}_0)$ based on a multivariate normal kernel is given by

$$\hat{f}(\mathbf{y}_0) = \frac{1}{nh^p |\mathbf{S}_{pl}|^{1/2}} \sum_{i=1}^n e^{-(\mathbf{y}_0 - \mathbf{y}_i)' \mathbf{S}_{pl}^{-1} (\mathbf{y}_0 - \mathbf{y}_i)/2h^2}, \quad (9.27)$$

where $h_1 = h_2 = \cdots = h_p = h$ and \mathbf{S}_{pl} is the pooled covariance matrix from the k groups in the sample. The covariance matrix \mathbf{S}_{pl} could be replaced by other forms. Two examples are (1) \mathbf{S}_i for the i th group and (2) a diagonal matrix.

The choice of the smoothing parameter h is critical in a kernel density estimator. The size of h determines how much each \mathbf{y}_i contributes to $\hat{f}(\mathbf{y}_0)$. If h is too small, $\hat{f}(\mathbf{y}_0)$ has a peak at each \mathbf{y}_i , and if h is too large, $\hat{f}(\mathbf{y}_0)$ is almost uniform (overly smoothed). Therefore, the value chosen for h must depend on the sample size n to avoid too much or too little smoothing; the larger the sample size, the smaller h should be. In practice, we could try several values of h and check the resulting error rates from the classification analysis.

To use the kernel method of density estimation in classification, we can apply it to each group to obtain $\hat{f}(\mathbf{y}_0|G_1), \hat{f}(\mathbf{y}_0|G_2), \dots, \hat{f}(\mathbf{y}_0|G_k)$, where \mathbf{y}_0 is the vector of measurements for an individual of unknown group membership. The classification rule then becomes: Assign \mathbf{y}_0 to the group G_i for which

$$p_i \hat{f}(\mathbf{y}_0|G_i) \text{ is maximum.} \quad (9.28)$$

Habbema et al. (1974) proposed a forward selection method for classification based on density estimation. Wegman (1972) and Habbema et al. (1978) found that the size of the h_i 's is more important than the shape of the kernel. The choice of h was investigated by Pfeiffer (1985) in a stepwise mode. Remme et al. (1980) compared linear, quadratic, and kernel classification methods for two groups and reported that for multivariate normal data with equal covariance matrices, the linear classifications were clearly superior. For some cases with departures from these assumptions, the kernel methods gave better results.

■ EXAMPLE 9.7.2

We illustrate the density estimation method of classification for the football data of Table 8.3. We use the multivariate normal kernel estimator in (9.27) with $h = 2$ to obtain $\hat{f}(\mathbf{y}_0|G_i)$, $i = 1, 2, 3$, for the three groups. Using

Table 9.8 Classification Table for the Football Data of Table 8.3 Using the Density Estimation Method of Classification with Multivariate Normal Kernel

Actual Group	Number of Observations	Predicted Group		
		1	2	3
1	30	25	1	4
2	30	0	12	18
3	30	0	3	27

$$\text{Apparent correct classification rate} = \frac{25 + 12 + 27}{90} = .711$$

$$\text{Apparent error rate} = 1 - .711 = .289$$

Table 9.9 Classification Table for the Football Data of Table 8.3 Using the Holdout Method Based on Density Estimation

Actual Group	Number of Observations	Predicted Group		
		1	2	3
1	30	24	1	5
2	30	0	10	20
3	30	1	3	26

$$\text{Correct classification rate} = \frac{24 + 10 + 26}{90} = .667$$

$$\text{Error rate} = 1 - .667 = .333$$

$p_1 = p_2 = p_3$, the rule in (9.28) becomes: Assign \mathbf{y}_0 to the group for which $\hat{f}(\mathbf{y}_0|G_i)$ is greatest. To obtain an apparent error rate, we follow this procedure for each of the 90 observations and obtain the classification results in Table 9.8.

Applying a holdout method in which the observation \mathbf{y}_{ij} being classified is excluded from computation of $\hat{f}(\mathbf{y}_{ij}|G_1)$, $\hat{f}(\mathbf{y}_{ij}|G_2)$, and $\hat{f}(\mathbf{y}_{ij}|G_3)$, we obtain the classification results in Table 9.9. As expected, the holdout error rate has increased somewhat from the apparent error rate in Table 9.8. \square

9.7.3 Nearest Neighbor Classification Rule

The earliest nonparametric classification method was the *nearest neighbor rule* of Fix and Hodges (1951), also known as the k nearest neighbor rule. The procedure is conceptually simple. We compute the distance from an observation \mathbf{y}_i to all other points \mathbf{y}_j using the distance function

$$(\mathbf{y}_i - \mathbf{y}_j)' \mathbf{S}_{\text{pl}}^{-1} (\mathbf{y}_i - \mathbf{y}_j) \quad j \neq i.$$

To classify \mathbf{y}_i into one of two groups, the k points nearest to \mathbf{y}_i are examined, and if the majority of the k points belong to G_1 , assign \mathbf{y}_i to G_1 ; otherwise assign \mathbf{y}_i to G_2 . If we denote the number of points from G_1 as k_1 , with the remaining k_2 points from G_2 , where $k = k_1 + k_2$, then the rule can be expressed as: Assign \mathbf{y}_i to G_1 if

$$k_1 > k_2 \quad (9.29)$$

and to G_2 otherwise. If the sample sizes n_1 and n_2 differ, we may wish to use proportions in place of counts: Assign \mathbf{y}_i to G_1 if

$$\frac{k_1}{n_1} > \frac{k_2}{n_2}. \quad (9.30)$$

A further refinement can be made by taking into account prior probabilities: Assign \mathbf{y}_i to G_1 if

$$\frac{k_1/n_1}{k_2/n_2} > \frac{p_2}{p_1}. \quad (9.31)$$

These rules are easily extended to more than two groups. For example, (9.30) becomes: Assign the observation to the group that has the highest proportion k_i/n_i , where k_i is the number of observations from G_i among the k nearest neighbors of the observation in question.

A decision must be made as to the value of k . Loftsgaarden and Quesenberry (1965) suggest choosing k near $\sqrt{n_i}$ for a typical n_i . In practice, one could try several values of k and use the one with the best error rate.

Reviews and extensions of the nearest neighbor method have been given by Hart (1968), Gates (1972), Hand and Batchelor (1978), Chidananda Gowda and Krishna (1979), Rogers and Wagner (1978), and Brown and Koplowitz (1979).

■ EXAMPLE 9.7.3

We use the football data of Table 8.3 to illustrate the k nearest neighbor method of estimating error rate, with $k = 5$. Since $n_1 = n_2 = n_3 = 30$ and the p_i 's are also assumed to be equal, we simply examine the five points closest to a point \mathbf{y} and classify \mathbf{y} into the group that has the most points among the five points. If there is a tie, we do not classify the point. For example, if the numbers from G_1 , G_2 , and G_3 were 1, 2, and 2, respectively, then we do not assign \mathbf{y} to either G_2 or G_3 .

For each point \mathbf{y}_{ij} , $i = 1, 2, 3$; $j = 1, 2, \dots, 30$, we find the five nearest neighbors and classify the point accordingly. Table 9.10 gives the classification results. As can be seen, there were 3 observations in group 1 that were not classified because of ties, 1 in group 2, and 3 in group 3. This left a total of 83 observations classified. \square

9.7.4 Classification Trees

Although the notion of using decision trees as a means for classification has been discussed for nearly fifty years, the methodology known as *classification and regression*

Table 9.10 Classification Table for the Football Data of Table 8.3 Using the k Nearest Neighbor Method with $k = 5$

Actual Group	Number of Observations	Predicted Group		
		1	2	3
1	30	26	0	1
2	30	1	19	9
3	30	1	4	22

$$\text{Correct classification rate} = \frac{26 + 19 + 22}{83} = .807$$

$$\text{Error rate} = 1 - .807 = .193$$

sion trees received its first complete development by Breiman, Friedman, Olshen, and Stone (1984). Classification trees are used to predict categorial responses, while regression trees are used to predict continuous responses. Because this chapter's focus is on classification of observations into discrete groups, we limit our discussion to a very simple type of classification tree. For a more complete discussion of this rich class of predictive tools, we refer the reader to the aforementioned landmark publication of Breiman, Friedman, Olshen, and Stone (1984), overviews of this class of methods including extensions and more recent developments (e.g., Hastie, Tibshirani, and Friedman, 2009), or documentation associated with the various software implementations of classification trees. We adopt much of our notation and terminology from the primer on the RPART routines by Therneau and Atkinson (1997). RPART comprises a collection of implementations of classification and regression trees within the R statistical computing environment (R Development Core Team, 2010).

We begin forming a classification tree with all n observations in one group or *root node* and seek to find a predictor variable x and an associated cutoff criterion such that splitting the n subjects into the two groups will minimize the *impurity* (diversity) within each of the two *child nodes*. This involves a consideration of all possible x variables at all possible split points. The process is then repeated for each of the child nodes. Thus the classification tree is formed by recursive binary splits of the observations in order to form an optimal partitioning of the observations into predictive groups.

The tree grows with child nodes splitting to create new generations of nodes. Of course, the splitting could continue until each observation is its own terminal (final) node, but such overfitting will generally lead to decision trees with poor out-of-sample predictive properties. Various stopping and pruning rules exist for selecting the optimal tree, but most are based on cross-validation. When the final tree is constructed, each observation falls into exactly one terminal node that will be associated with a single predicted group.

There are many ways in which we might split a parent node into two child nodes, and there are several criteria that could influence the way we define an optimal split.

The first criterion is the collection of prior probabilities p_1, \dots, p_k associated with each group. Allowing prior probabilities to influence the decision rules will lead to rules that tend to classify more observations into the more commonly observed groups.

A second criterion is the misclassification cost for classifying an observation from group G_i into group G_j (for $i, j = 1, \dots, k$). As stated previously, this can become extremely complicated for more than two groups, so misclassification costs will not be considered here.

A third criterion affecting the definition of an optimal splitting rule is the measure of impurity chosen for the classification. Employing the oft-used Gini index, we can define the impurity associated with the node A as

$$I_A = \sum_{i=1}^k p_{i|A}(1 - p_{i|A}) \quad (9.32)$$

where $p_{i|A}$ is the probability that an observation is in group G_i given that it is classified into node A . Note that the impurity of node A is close to zero when each of $p_{1|A}, p_{2|A}, \dots, p_{k|A}$ is either near zero or near 1—that is, when the observations in a node are predominantly from only one group. The probability $p_{i|A}$ is calculated from the data using

$$p_{i|A} = \frac{p_i(n_{iA}/n_i)}{\sum_{i=1}^k p_i(n_{iA}/n_i)}, \quad (9.33)$$

where p_i is the prior probability associated with group G_i , n_i is the number of observations in G_i , and n_{iA} is the number of observations from G_i that are in node A . Note that if we assume prior probabilities are proportional to the size of the group G_i in the training data (i.e., if $p_1/n_1 = p_2/n_2 = \dots = p_k/n_k$), then (9.33) simplifies to

$$p_{i|A} = \frac{n_{iA}}{n_A}, \quad (9.34)$$

where n_A is the number of observations in node A .

We quantify the probability of an observation being in node A using

$$p_A = \sum_{i=1}^k p_i(n_{iA}/n_i). \quad (9.35)$$

For the case of prior probabilities proportional to the size of the group, (9.35) simplifies to $p_A = n_A/n$.

Finally, using the quantities defined above, we can define the quality for the split of a parent node A into two child nodes A_L and A_R (where L and R denote the left and right children from the split). The change in total impurity (ΔI) associated with the split is defined by

$$\Delta I = p_A I_A - (p_{A_L} I_{A_L} + p_{A_R} I_{A_R}), \quad (9.36)$$

with ΔI guaranteed to be greater than or equal to 0. After considering all possible choices for the split of node A into child nodes A_L and A_R , the split at node A that maximizes ΔI is deemed the optimal split.




Figure 9.4 Classification tree for classifying subjects into one of three groups using the football data of Table 8.3

■ EXAMPLE 9.7.4

To explain the basic components of a classification tree, we use the football data in Table 8.3 to illustrate the use of classification trees for classification analysis. For this classification exercise, we use equal prior probabilities for the three groups (college players, high school players, and non-football players). For these data, this is equivalent to assuming priors proportional to size since the training data groups each have 30 subjects. Costs of misclassification are assumed to be equal, and the Gini index is used as the basis for the measure of impurity when selecting optimal splits for nodes. Optimal tree size is chosen based on cross-validation.

In Figure 9.4, we begin with all 90 subjects in the root node and seek to find the head dimension variable and associated cutoff criterion that best separates the three types of football players (college players, high school players, and non-players) into two distinct groups. At the root node the optimal rule was to form groups based on whether or not the eye-to-top-of-head measurement (eyehd) is less than 11.95 cm. Those observations with $\text{eyehd} < 11.95$ cm are considered to be in the left child node (denoted A_L), and all others are in the right child node (A_R). We say this is the optimal split because A_R consists of mostly high school players (2 college, 29 high school, and 6 non-players) and the left node is predominantly college players and non-football players (28 college, 1 high school, and 24 non-players).

Table 9.11 Comparison of Classification Error Rates for the Football Data of Table 8.3 Using Several Methods Including the Classification Tree

Classification Method	Misclassification Rate (based on cross-validation)
Linear	0.300
Quadratic	0.344
Density Estimation	
normal density, $h = 2$	0.333
normal density, $h = 5$	0.367
Nearest Neighbor	
$k = 5$	0.278
$k = 9$	0.367
Classification Tree	0.256

For each child node, the same process is recursively applied. For example, those observations with $\text{eyehd} < 11.95$ cm are then further split into groups based again on eyehd , but this time with a new cutoff of 10.35 cm. This serves to further split this group of 53 subjects into two terminal child nodes. The first contains all subjects with $\text{eyehd} < 10.35$ and includes 21 college, 1 high school, and 7 non-players. The second terminal child node formed by this split contains all subjects with eyehd measurements in the range [10.35, 11.95) and includes 7 college, 0 high school, and 17 non-players. The set of subjects with $\text{eyehd} \geq 11.95$ cm are also further subdivided, with those having ear-to-top-of-head (earhd) measurements greater than or equal to 13.8 cm placed in a terminal child node with 1 college, 27 high school, and 2 non-players. The complementary terminal node for those with $\text{eyehd} > 11.95$ and $\text{earhd} < 13.8$ cm contains 1 college, 2 high school, and 4 non-players.

Thus there is one large terminal node associated with each of the college and high school player groups, and there are two smaller terminal nodes associated with the non-football-player group. For some future (unclassified) observation, the classification tree in Figure 9.4 illustrates how one would predict group membership based on the eyehd and earhd variables.

Table 9.11 compares the misclassification rates for several different classification methods using the football data of Table 8.3. For this data set, the misclassification rate for the classification tree is lowest among the competing approaches, but the relative accuracy of the approaches will be different for different data sets. \square

PROBLEMS

- 9.1** Show that if $z_{1i} = \mathbf{a}'\mathbf{y}_{1i}, i = 1, 2, \dots, n_1$, and $z_{2i} = \mathbf{a}'\mathbf{y}_{2i}, i = 1, 2, \dots, n_2$, where z is the discriminant function defined in (9.1), then $\bar{z}_1 - \bar{z}_2 = (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{\text{pl}}^{-1} (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)$ as in (9.3).
- 9.2** With $z = \mathbf{a}'\mathbf{y}$ as in (9.1), and $\bar{z}_1 = \mathbf{a}'\bar{\mathbf{y}}_1, \bar{z}_2 = \mathbf{a}'\bar{\mathbf{y}}_2$, show that $\frac{1}{2}(\bar{z}_1 + \bar{z}_2) = \frac{1}{2}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{\text{pl}}^{-1} (\bar{\mathbf{y}}_1 + \bar{\mathbf{y}}_2)$ as in (9.4).
- 9.3** Obtain the normal-based classification rule in (9.9).
- 9.4** Derive the linear classification rule in (9.13).
- 9.5** Derive the quadratic classification function in (9.15).
- 9.6** Show that if we assume prior probabilities are proportional to the size of the group G_i in the training data, then (9.33) simplifies to (9.34).
- 9.7** Do a classification analysis on the beetle data in Table 5.5 as follows:
- Find the classification function $z = (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{\text{pl}}^{-1} \mathbf{y}$ and the cutoff point $\frac{1}{2}(\bar{z}_1 + \bar{z}_2)$.
 - Find the classification table using the linear classification function in part (a).
 - Find the classification table using the nearest neighbor method.
- 9.8** Do a classification analysis on the dystrophy data of Table 5.7 as follows:
- Find the classification function $z = (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{\text{pl}}^{-1} \mathbf{y}$ and the cutoff point $\frac{1}{2}(\bar{z}_1 + \bar{z}_2)$.
 - Find the classification table using the linear classification function in part (a).
 - Repeat part (b) using p_1 and p_2 proportional to sample sizes.
- 9.9** Do a classification analysis on the cyclical data of Table 5.8 as follows:
- Find the classification function $z = (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{\text{pl}}^{-1} \mathbf{y}$ and the cutoff point $\frac{1}{2}(\bar{z}_1 + \bar{z}_2)$.
 - Find the classification table using the linear classification function in part (a).
 - Find the classification table using the holdout method.
 - Find the classification table using a kernel density estimator method.
- 9.10** Using the engineer data of Table 5.6, carry out a classification analysis as follows:

- (a) Find the classification table using the linear classification function.
 - (b) Carry out a stepwise discriminant selection of variables (see Problem 8.14).
 - (c) Find the classification table for the variables selected in part (b).
- 9.11** Do a classification analysis on the fish data in Table 6.17 as follows. Assume $p_1 = p_2 = p_3$.
- (a) Find the linear classification functions.
 - (b) Find the classification table using the linear classification functions in part (a) (assuming $\Sigma_1 = \Sigma_2 = \Sigma_3$).
 - (c) Find the classification table using quadratic classification functions (assuming population covariance matrices are not equal).
 - (d) Find the classification table using linear classification functions and the holdout method.
 - (e) Find the classification table using a nearest neighbor method.

9.12 Do a classification analysis on the rootstock data of Table 6.2 as follows:

- (a) Find the linear classification functions.
- (b) Find the classification table using the linear classification functions in part (a) (assuming $\Sigma_1 = \Sigma_2 = \Sigma_3$).
- (c) Find the classification table using quadratic classification functions (assuming population covariance matrices are not equal).
- (d) Find the classification table using the nearest neighbor method.
- (e) Find the classification table using a kernel density estimator method.

CHAPTER 10

MULTIVARIATE REGRESSION

10.1 INTRODUCTION

In this chapter, we consider the linear relationship between one or more y 's (the *dependent* or *response* variables) and one or more x 's (the *independent* or *predictor* variables). We will use a linear model to relate the y 's to the x 's and will be concerned with estimation and testing of the parameters in the model. One aspect of interest will be choosing which variables to include in the model if this is not already known.

We can distinguish three cases according to the number of variables:

1. Simple linear regression: one y and one x . For example, suppose we wish to predict college grade point average (GPA) based on an applicant's high school GPA.
2. Multiple linear regression: one y and several x 's. We could attempt to improve our prediction of college GPA by using more than one independent variable, for example, high school GPA, standardized test scores (such as ACT or SAT), or rating of high school.
3. Multivariate multiple linear regression: several y 's and several x 's. In the above illustration, we may wish to predict several y 's (such as number of

years of college the person will complete or GPA in the sciences, arts, and humanities). As another example, suppose the Air Force wishes to predict several measures of pilot efficiency. These response variables could be regressed against independent variables (such as math and science skills, reaction time, eyesight acuity, and manual dexterity).

To further distinguish case 2 from case 3, we could designate case 2 as *univariate* multiple regression because there is only one y . Thus in case 3, “multivariate” indicates that there are several y ’s and “multiple” implies several x ’s. The term *multivariate regression* usually refers to case 3.

There are two basic types of independent variables, *fixed* and *random*. In the above illustrations, all x ’s are random variables and are therefore not under the control of the researcher. A person is chosen at random, and all of the y ’s and x ’s are measured, or observed, for that person. In some experimental situations, the x ’s are fixed, that is, under the control of the experimenter. For example, a researcher may wish to relate yield per acre and nutritional value to level of application of various chemical fertilizers. The experimenter can choose the amount of chemicals to be applied and then observe the changes in the yield and nutritional responses.

In order to provide a solid base for multivariate multiple regression, we review several aspects of multiple regression with fixed x ’s in Section 10.2. The random- x case for multiple regression is discussed briefly in Section 10.3.

10.2 MULTIPLE REGRESSION: FIXED x ’s

10.2.1 Model for Fixed x ’s

In the fixed- x regression model, we express each y in a sample of n observations as a linear function of the x ’s plus a random error, ε :

$$\begin{aligned} y_1 &= \beta_0 + \beta_1 x_{11} + \beta_2 x_{12} + \cdots + \beta_q x_{1q} + \varepsilon_1 \\ y_2 &= \beta_0 + \beta_1 x_{21} + \beta_2 x_{22} + \cdots + \beta_q x_{2q} + \varepsilon_2 \\ &\vdots \\ y_n &= \beta_0 + \beta_1 x_{n1} + \beta_2 x_{n2} + \cdots + \beta_q x_{nq} + \varepsilon_n. \end{aligned} \tag{10.1}$$

The number of x ’s is denoted by q . The β ’s in (10.1) are called *regression coefficients*. Additional assumptions that accompany the equations of the model are as follows:

1. $E(\varepsilon_i) = 0$ for all $i = 1, 2, \dots, n$.
2. $\text{var}(\varepsilon_i) = \sigma^2$ for all $i = 1, 2, \dots, n$.
3. $\text{cov}(\varepsilon_i, \varepsilon_j) = 0$ for all $i \neq j$.

Assumption 1 states that the model is linear and that no additional terms are needed to predict y ; all remaining variation in y is purely random and unpredictable. Thus

if $E(\varepsilon_i) = 0$ and the x 's are fixed, then $E(y_i) = \beta_0 + \beta_1 x_{i1} + \beta_2 x_{i2} + \cdots + \beta_q x_{iq}$, and the mean of y is expressible in terms of these q x 's with no others needed. In assumption 2, the variance of each ε_i is the same, which also implies that $\text{var}(y_i) = \sigma^2$, since the x 's are fixed. Assumption 3 imposes the condition that the error terms be uncorrelated, from which it follows that the y 's are also uncorrelated, that is, $\text{cov}(y_i, y_j) = 0$.

Thus the three assumptions can be restated in terms of y as follows:

1. $E(y_i) = \beta_0 + \beta_1 x_{i1} + \beta_2 x_{i2} + \cdots + \beta_q x_{iq}, i = 1, 2, \dots, n.$
2. $\text{var}(y_i) = \sigma^2, i = 1, 2, \dots, n.$
3. $\text{cov}(y_i, y_j) = 0, \text{ for all } i \neq j.$

Using matrix notation, the models for the n observations in (10.1) can be written much more concisely in the form

$$\begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{pmatrix} = \begin{pmatrix} 1 & x_{11} & x_{12} & \cdots & x_{1q} \\ 1 & x_{21} & x_{22} & \cdots & x_{2q} \\ \vdots & \vdots & \vdots & & \vdots \\ 1 & x_{n1} & x_{n2} & \cdots & x_{nq} \end{pmatrix} \begin{pmatrix} \beta_0 \\ \beta_1 \\ \vdots \\ \beta_q \end{pmatrix} + \begin{pmatrix} \varepsilon_1 \\ \varepsilon_2 \\ \vdots \\ \varepsilon_n \end{pmatrix} \quad (10.2)$$

or

$$\mathbf{y} = \mathbf{X}\boldsymbol{\beta} + \boldsymbol{\varepsilon}. \quad (10.3)$$

With this notation, the above three assumptions become


1. $E(\boldsymbol{\varepsilon}) = \mathbf{0}$
2. $\text{cov}(\boldsymbol{\varepsilon}) = \sigma^2 \mathbf{I}$,

which can be rewritten in terms of \mathbf{y} as

1. $E(\mathbf{y}) = \mathbf{X}\boldsymbol{\beta}$
2. $\text{cov}(\mathbf{y}) = \sigma^2 \mathbf{I}.$

Note that the second assumption in matrix form incorporates both the second and third assumptions in univariate form; that is, $\text{cov}(\mathbf{y}) = \sigma^2 \mathbf{I}$ implies $\text{var}(y_i) = \sigma^2$ and $\text{cov}(y_i, y_j) = 0$.

For estimation and testing purposes, we need to have $n > q + 1$. Therefore, the matrix expression (10.3) has the following typical pattern:



10.2.2 Least Squares Estimation in the Fixed- x Model

If the first assumption holds, we have $E(y_i) = \beta_0 + \beta_1 x_{i1} + \beta_2 x_{i2} + \cdots + \beta_q x_{iq}$. We seek to estimate the β 's and thereby estimate $E(y_i)$. If the estimates are denoted by $\hat{\beta}_0, \hat{\beta}_1, \dots, \hat{\beta}_q$, then $\hat{E}(y_i) = \hat{\beta}_0 + \hat{\beta}_1 x_{i1} + \hat{\beta}_2 x_{i2} + \cdots + \hat{\beta}_q x_{iq}$. However, $\hat{E}(y_i)$ is usually designated \hat{y}_i . Thus \hat{y}_i estimates $E(y_i)$, not y_i . We now consider the least squares estimates of the β 's.

The *least squares* estimates of $\beta_0, \beta_1, \dots, \beta_q$ minimize the sum of squares of deviations of the n observed y 's from their "modeled" values, that is, from their values \hat{y}_i predicted by the model. Thus we seek $\hat{\beta}_0, \hat{\beta}_1, \dots, \hat{\beta}_q$ that minimize

$$\begin{aligned} SSE &= \sum_{i=1}^n \hat{\varepsilon}_i^2 = \sum_{i=1}^n (y_i - \hat{y}_i)^2 \\ &= \sum_{i=1}^n (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_{i1} - \hat{\beta}_2 x_{i2} - \cdots - \hat{\beta}_q x_{iq})^2. \end{aligned} \quad (10.4)$$

The value of $\hat{\beta} = (\hat{\beta}_0, \hat{\beta}_1, \dots, \hat{\beta}_q)'$ that minimizes SSE in (10.4) is given by

$$\hat{\beta} = (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y}. \quad (10.5)$$

In (10.5), we assume that $\mathbf{X}'\mathbf{X}$ is nonsingular. This will ordinarily hold if $n > q + 1$ and no x_j is a linear combination of other x 's.

In expression (10.5), we see a characteristic pattern similar to that for $\hat{\beta}_1$ in simple linear regression given in (3.11), $\hat{\beta}_1 = s_{xy}/s_x^2$. The product $\mathbf{X}'\mathbf{y}$ can be used to compute the covariances of the x 's with y . The product $\mathbf{X}'\mathbf{X}$ can be used to obtain the covariance matrix of the x 's, which includes the variances and covariances of the x 's [see the comment following (10.16) about variances and covariances involving the fixed x 's]. Since $\mathbf{X}'\mathbf{X}$ is typically not diagonal, each $\hat{\beta}_j$ depends on $s_{x_j y}$ and $s_{x_j}^2$, as well as the relationship of x_j to the other x 's.

We now demonstrate algebraically that $\hat{\beta} = (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y}$ in (10.5) minimizes SSE (this can also be done readily with calculus). If we designate the i th row of \mathbf{X} as $\mathbf{x}'_i = (1, x_{i1}, x_{i2}, \dots, x_{iq})$, we can write (10.4) as

$$SSE = \sum_{i=1}^n (y_i - \mathbf{x}'_i \hat{\beta})^2.$$

The quantity $y_i - \mathbf{x}'_i \hat{\beta}$ is the i th element of the vector $\mathbf{y} - \mathbf{X}\hat{\beta}$. Hence, by (2.33),

$$SSE = (\mathbf{y} - \mathbf{X}\hat{\beta})'(\mathbf{y} - \mathbf{X}\hat{\beta}). \quad (10.6)$$

Let \mathbf{b} be an alternative estimate that may lead to a smaller value of SSE than $\hat{\beta}$. We add $\mathbf{X}(\hat{\beta} - \mathbf{b})$ to see if this reduces SSE.

$$SSE = [(\mathbf{y} - \mathbf{X}\hat{\beta}) + \mathbf{X}(\hat{\beta} - \mathbf{b})]'[(\mathbf{y} - \mathbf{X}\hat{\beta}) + \mathbf{X}(\hat{\beta} - \mathbf{b})].$$

We now expand this using the two terms $\mathbf{y} - \mathbf{X}\hat{\boldsymbol{\beta}}$ and $\mathbf{X}(\hat{\boldsymbol{\beta}} - \mathbf{b})$ to obtain

$$\begin{aligned} \text{SSE} &= (\mathbf{y} - \mathbf{X}\hat{\boldsymbol{\beta}})'(\mathbf{y} - \mathbf{X}\hat{\boldsymbol{\beta}}) + [\mathbf{X}(\hat{\boldsymbol{\beta}} - \mathbf{b})]' \mathbf{X}(\hat{\boldsymbol{\beta}} - \mathbf{b}) + 2[\mathbf{X}(\hat{\boldsymbol{\beta}} - \mathbf{b})]'(\mathbf{y} - \mathbf{X}\hat{\boldsymbol{\beta}}) \\ &= (\mathbf{y} - \mathbf{X}\hat{\boldsymbol{\beta}})'(\mathbf{y} - \mathbf{X}\hat{\boldsymbol{\beta}}) + (\hat{\boldsymbol{\beta}} - \mathbf{b})' \mathbf{X}' \mathbf{X}(\hat{\boldsymbol{\beta}} - \mathbf{b}) + 2(\hat{\boldsymbol{\beta}} - \mathbf{b})' \mathbf{X}'(\mathbf{y} - \mathbf{X}\hat{\boldsymbol{\beta}}) \\ &= (\mathbf{y} - \mathbf{X}\hat{\boldsymbol{\beta}})'(\mathbf{y} - \mathbf{X}\hat{\boldsymbol{\beta}}) + (\hat{\boldsymbol{\beta}} - \mathbf{b})' \mathbf{X}' \mathbf{X}(\hat{\boldsymbol{\beta}} - \mathbf{b}) + 2(\hat{\boldsymbol{\beta}} - \mathbf{b})' (\mathbf{X}'\mathbf{y} - \mathbf{X}'\mathbf{X}\hat{\boldsymbol{\beta}}). \end{aligned}$$

The third term vanishes if we substitute $\hat{\boldsymbol{\beta}} = (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y}$ into $\mathbf{X}'\mathbf{X}\hat{\boldsymbol{\beta}}$. The second term is a positive definite quadratic form, and SSE is therefore minimized when $\mathbf{b} = \hat{\boldsymbol{\beta}}$. Thus no value of \mathbf{b} can reduce SSE from the value given by $\hat{\boldsymbol{\beta}}$.

When $E(\mathbf{y}) = \mathbf{X}\boldsymbol{\beta}$ and $\text{cov}(\mathbf{y}) = \sigma^2\mathbf{I}$, it follows that the least squares estimator $\hat{\boldsymbol{\beta}} = (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y}$ in (10.5) has three important properties. The first property is that $\hat{\boldsymbol{\beta}}$ is an unbiased estimator for $\boldsymbol{\beta}$. That is, $E(\hat{\boldsymbol{\beta}}) = \boldsymbol{\beta}$. The second property is that $\text{cov}(\hat{\boldsymbol{\beta}}) = \sigma^2(\mathbf{X}'\mathbf{X})^{-1}$. The third important property is characterized by the Gauss–Markov theorem.

Gauss–Markov Theorem. If $E(\mathbf{y}) = \mathbf{X}\boldsymbol{\beta}$ and $\text{cov}(\mathbf{y}) = \sigma^2\mathbf{I}$, the least squares estimators $\hat{\boldsymbol{\beta}}_0, \hat{\boldsymbol{\beta}}_1, \dots, \hat{\boldsymbol{\beta}}_q$ have minimum variance among all linear unbiased estimators.

The Gauss–Markov theorem is often simplified to the following statement: If $E(\mathbf{y}) = \mathbf{X}\boldsymbol{\beta}$ and $\text{cov}(\mathbf{y}) = \sigma^2\mathbf{I}$, the least squares estimators are “best linear unbiased estimators” (BLUE), where “best” implies “minimum variance” and “linear” indicates that the estimators are linear functions of \mathbf{y} . For a review of properties of $\hat{\boldsymbol{\beta}}$ and an alternative derivation of $\hat{\boldsymbol{\beta}}$ based on the assumption that \mathbf{y} is normally distributed, see Rencher (1998, Chapter 7) or Rencher and Schaalje (2008, Chapter 7).

10.2.3 An Estimator for σ^2

It can be shown that

$$E(\text{SSE}) = \sigma^2[n - (q + 1)] = \sigma^2(n - q - 1). \quad (10.7)$$

We can therefore obtain an unbiased estimator of σ^2 as

$$s^2 = \frac{\text{SSE}}{n - q - 1} = \frac{1}{n - q - 1}(\mathbf{y} - \mathbf{X}\hat{\boldsymbol{\beta}})'(\mathbf{y} - \mathbf{X}\hat{\boldsymbol{\beta}}). \quad (10.8)$$

We can also express SSE in the form

$$\text{SSE} = \mathbf{y}'\mathbf{y} - \hat{\boldsymbol{\beta}}'\mathbf{X}'\mathbf{y}, \quad (10.9)$$

and we note that there are n terms in $\mathbf{y}'\mathbf{y}$ and $q + 1$ terms in $\hat{\boldsymbol{\beta}}'\mathbf{X}'\mathbf{y}$. The difference is the denominator of s^2 in (10.8). Thus the degrees of freedom (denominator) for SSE is reduced by $q + 1$.

The need for an adjustment of $q + 1$ to the degrees of freedom of SSE can be illustrated with a simple random sample of a random variable y from a population

with mean μ and variance σ^2 . The sum of squares $\sum_i(y_i - \mu)^2$ has n degrees of freedom, while $\sum_i(y_i - \bar{y})^2$ has $n - 1$. It is intuitively clear that

$$E \left[\sum_{i=1}^n (y_i - \mu)^2 \right] > E \left[\sum_{i=1}^n (y_i - \bar{y})^2 \right]$$

because \bar{y} fits the sample better than μ , which is the mean of the population but not of the sample. Thus (squared) deviations from \bar{y} will tend to be smaller than deviations from μ . In fact, it is easily shown that

$$\begin{aligned} \sum_{i=1}^n (y_i - \mu)^2 &= \sum_{i=1}^n (y_i - \bar{y} + \bar{y} - \mu)^2 \\ &= \sum_i (y_i - \bar{y})^2 + n(\bar{y} - \mu)^2, \end{aligned} \quad (10.10)$$

whence

$$\sum_i (y_i - \bar{y})^2 = \sum_i (y_i - \mu)^2 - n(\bar{y} - \mu)^2.$$

Thus $\sum_i(y_i - \bar{y})^2$ is expressible as a sum of n squares minus one square, which corresponds to $n - 1$ degrees of freedom. More formally, we have

$$E \left[\sum_i (y_i - \bar{y})^2 \right] = n\sigma^2 - \frac{n\sigma^2}{n} = (n - 1)\sigma^2.$$

10.2.4 The Model Corrected for Means

It is sometimes convenient to “center” the x ’s by subtracting their means, $\bar{x}_1 = \sum_{i=1}^n x_{i1}/n, \bar{x}_2 = \sum_{i=1}^n x_{i2}/n$, and so on [$\bar{x}_1, \bar{x}_2, \dots, \bar{x}_q$ are the means of the columns of \mathbf{X} in (10.2)]. In terms of centered x ’s, the model for each y_i in (10.1) becomes

$$y_i = \alpha + \beta_1(x_{i1} - \bar{x}_1) + \beta_2(x_{i2} - \bar{x}_2) + \cdots + \beta_q(x_{iq} - \bar{x}_q) + \varepsilon_i, \quad (10.11)$$

where

$$\alpha = \beta_0 + \beta_1\bar{x}_1 + \beta_2\bar{x}_2 + \cdots + \beta_q\bar{x}_q. \quad (10.12)$$

To estimate

$$\boldsymbol{\beta}_1 = \begin{pmatrix} \beta_1 \\ \beta_2 \\ \vdots \\ \beta_q \end{pmatrix},$$

we use the centered x 's in the matrix

$$\mathbf{X}_c = \begin{pmatrix} x_{11} - \bar{x}_1 & x_{12} - \bar{x}_2 & \cdots & x_{1q} - \bar{x}_q \\ x_{21} - \bar{x}_1 & x_{22} - \bar{x}_2 & \cdots & x_{2q} - \bar{x}_q \\ \vdots & \vdots & & \vdots \\ x_{n1} - \bar{x}_1 & x_{n2} - \bar{x}_2 & \cdots & x_{nq} - \bar{x}_q \end{pmatrix} = \begin{pmatrix} (\mathbf{x}_1 - \bar{\mathbf{x}})' \\ (\mathbf{x}_2 - \bar{\mathbf{x}})' \\ \vdots \\ (\mathbf{x}_n - \bar{\mathbf{x}})' \end{pmatrix}, \quad (10.13)$$

where $\mathbf{x}'_i = (x_{i1}, x_{i2}, \dots, x_{iq})$ and $\bar{\mathbf{x}}' = (\bar{x}_1, \bar{x}_2, \dots, \bar{x}_q)$. Then by analogy to (10.5), the least squares estimate of β_1 is

$$\hat{\beta}_1 = (\mathbf{X}'_c \mathbf{X}_c)^{-1} \mathbf{X}'_c \mathbf{y}. \quad (10.14)$$

If $E(y) = \beta_0 + \beta_1 x_1 + \cdots + \beta_q x_q$ is evaluated at $x_1 = \bar{x}_1, x_2 = \bar{x}_2, \dots, x_q = \bar{x}_q$, the result is the same as α in (10.12). Thus, we estimate α by \bar{y} :

$$\hat{\alpha} = \bar{y}.$$

In other words, if the origin of the x 's is shifted to $\bar{\mathbf{x}} = (\bar{x}_1, \bar{x}_2, \dots, \bar{x}_q)'$, then the intercept of the fitted model is \bar{y} . With $\hat{\alpha} = \bar{y}$, we obtain

$$\hat{\beta}_0 = \hat{\alpha} - \hat{\beta}_1 \bar{x}_1 - \hat{\beta}_2 \bar{x}_2 - \cdots - \hat{\beta}_q \bar{x}_q = \bar{y} - \hat{\beta}_1' \bar{\mathbf{x}} \quad (10.15)$$

as an estimate of β_0 in (10.12). Together, the estimators $\hat{\beta}_0$ and $\hat{\beta}_1$ in (10.15) and (10.14) are the same as the usual least squares estimator $\hat{\beta}$ in (10.5).

We can express $\hat{\beta}_1$ in (10.14) in terms of sample variances and covariances. The overall sample covariance matrix of y and the x 's is

$$\mathbf{S} = \left(\begin{array}{c|cccc} s_{yy} & s_{y1} & s_{y2} & \cdots & s_{yq} \\ \hline s_{1y} & s_{11} & s_{12} & \cdots & s_{1q} \\ \vdots & \vdots & \vdots & & \vdots \\ s_{qy} & s_{q1} & s_{q2} & \cdots & s_{qq} \end{array} \right) = \begin{pmatrix} s_{yy} & \mathbf{s}'_{yx} \\ \mathbf{s}_{yx} & \mathbf{S}_{xx} \end{pmatrix}, \quad (10.16)$$

where s_{yy} is the variance of y , s_{yj} is the covariance of y and x_j , s_{jj} is the variance of x_j , s_{jk} is the covariance of x_j and x_k , and $\mathbf{s}'_{yx} = (s_{y1}, s_{y2}, \dots, s_{yq})$. These sample variances and covariances are mathematically equivalent to analogous formulas (3.23) and (3.25) for random variables, where the sample variances and covariances were estimates of population variances and covariances. However, here the x 's are considered to be constants that remain fixed from sample to sample, and a formula such as $s_{11} = \sum_{i=1}^n (x_{i1} - \bar{x}_1)^2 / (n - 1)$ summarizes the spread in the n values of x_1 but does not estimate a population variance.

To express $\hat{\beta}_1$ in terms of \mathbf{S}_{xx} and \mathbf{s}_{yx} in (10.16), we note first that the diagonal elements of $\mathbf{X}'_c \mathbf{X}_c$ are corrected sums of squares. For example, in the second diagonal position, we have

$$\sum_{i=1}^n (x_{i2} - \bar{x}_2)^2 = (n - 1)s_{22}.$$

The off-diagonal elements of $\mathbf{X}'_c \mathbf{X}_c$ are analogous corrected sums of products; for example, the element in the (1, 2) position is

$$\sum_{i=1}^n (x_{i1} - \bar{x}_1)(x_{i2} - \bar{x}_2) = (n-1)s_{12}.$$

Thus

$$\frac{1}{n-1} \mathbf{X}'_c \mathbf{X}_c = \mathbf{S}_{xx}. \quad (10.17)$$

Similarly,

$$\frac{1}{n-1} \mathbf{X}'_c \mathbf{y} = \mathbf{s}_{yx}, \quad (10.18)$$

even though \mathbf{y} has not been centered. The second element of $\mathbf{X}'_c \mathbf{y}$, for example, is $\sum_i (x_{i2} - \bar{x}_2)y_i$, which is equal to $(n-1)s_{2y}$:

$$\begin{aligned} (n-1)s_{2y} &= \sum_{i=1}^n (x_{i2} - \bar{x}_2)(y_i - \bar{y}) \\ &= \sum_i (x_{i2} - \bar{x}_2)y_i - \sum_i (x_{i2} - \bar{x}_2)\bar{y} \\ &= \sum_i (x_{i2} - \bar{x}_2)y_i, \end{aligned}$$

since

$$\sum_i (x_{i2} - \bar{x}_2)\bar{y} = 0. \quad (10.19)$$

Now, multiplying and dividing by $n-1$ in (10.14), we obtain

$$\begin{aligned} \hat{\beta}_1 &= (n-1)(\mathbf{X}'_c \mathbf{X}_c)^{-1} \frac{\mathbf{X}'_c \mathbf{y}}{n-1} = \left(\frac{\mathbf{X}'_c \mathbf{X}}{n-1} \right)^{-1} \frac{\mathbf{X}'_c \mathbf{y}}{n-1} \\ &= \mathbf{S}_{xx}^{-1} \mathbf{s}_{yx} \quad [\text{by (10.17) and (10.18)}], \end{aligned} \quad (10.20)$$

and substituting this in (10.15) gives

$$\hat{\beta}_0 = \hat{\alpha} - \hat{\beta}_1' \bar{\mathbf{x}} = \bar{y} - \bar{\mathbf{s}}'_{yx} \mathbf{S}_{xx}^{-1} \bar{\mathbf{x}}. \quad (10.21)$$

10.2.5 Hypothesis Tests

In this section, we review two basic tests on the β 's. For other tests and confidence intervals, see Rencher (1998, Section 7.2.4) and Rencher and Schaalje (2008, Sections 8.4–8.7). In order to obtain F -tests, we assume that \mathbf{y} is $N_n(\mathbf{X}\beta, \sigma^2 \mathbf{I})$.

10.2.5a Test of Overall Regression

The overall regression hypothesis that none of the x 's predicts y can be expressed as $H_0: \beta_1 = \mathbf{0}$, since $\beta'_1 = (\beta_1, \beta_2, \dots, \beta_q)$. We do not include $\beta_0 = 0$ in the hypothesis so as not to restrict y to have an intercept of zero.

We can write $SSE = \mathbf{y}'\mathbf{y} - \hat{\beta}'\mathbf{X}'\mathbf{y}$ in (10.9) in the form

$$\mathbf{y}'\mathbf{y} = (\mathbf{y}'\mathbf{y} - \hat{\beta}'\mathbf{X}'\mathbf{y}) + \hat{\beta}'\mathbf{X}'\mathbf{y}, \quad (10.22)$$

which partitions $\mathbf{y}'\mathbf{y}$ into a part due to β and a part due to deviations from the fitted model.

To correct y for its mean and thereby avoid inclusion of $\beta_0 = 0$, we subtract $n\bar{y}^2$ from both sides to obtain

$$\begin{aligned} \mathbf{y}'\mathbf{y} - n\bar{y}^2 &= (\mathbf{y}'\mathbf{y} - \hat{\beta}'\mathbf{X}'\mathbf{y}) + (\hat{\beta}'\mathbf{X}'\mathbf{y} - n\bar{y}^2) \\ &= SSE + SSR, \end{aligned} \quad (10.23)$$

where $\mathbf{y}'\mathbf{y} - n\bar{y}^2 = \sum_i(y_i - \bar{y})^2$ is the total sum of squares adjusted for the mean and $SSR = \hat{\beta}'\mathbf{X}'\mathbf{y} - n\bar{y}^2$ is the overall regression sum of squares adjusted for the intercept.

We can test $H_0: \beta_1 = \mathbf{0}$ by means of

$$F = \frac{SSR/q}{SSE/(n-q-1)}, \quad (10.24)$$

which is distributed as $F_{q,n-q-1}$ when $H_0: \beta_1 = \mathbf{0}$ is true. We reject H_0 if $F > F_{\alpha,h,n-q-1}$.

10.2.5b Test on a Subset of the β 's

In an attempt to simplify the model, we may wish to test the hypothesis that some of the β 's are zero. For example, in the model

$$y = \beta_0 + \beta_1x_1 + \beta_2x_2 + \beta_3x_1^2 + \beta_4x_2^2 + \beta_5x_1x_2 + \varepsilon,$$

we may be interested in the hypothesis $H_0: \beta_3 = \beta_4 = \beta_5 = 0$. If H_0 is true, the model is linear in x_1 and x_2 . In other cases, we may want to ascertain whether a single β_j can be deleted.

For convenience of exposition, let the β 's that are candidates for deletion be rearranged to appear last in β and denote this subset of β 's by β_d , where d reminds us that these β 's are to be *deleted* if $H_0: \beta_d = \mathbf{0}$ is accepted. Let the subset to be *retained* in the *reduced* model be denoted by β_r . Thus β is partitioned into

$$\beta = \begin{pmatrix} \beta_r \\ \beta_d \end{pmatrix}.$$

Let h designate the number of parameters in β_d . Then there are $q+1-h$ parameters in β_r .

To test the hypothesis $H_0: \beta_d = \mathbf{0}$, we fit the full model containing all the β 's in $\boldsymbol{\beta}$ and then fit the reduced model containing only the β 's in β_r . Let \mathbf{X}_r be the columns of \mathbf{X} corresponding to β_r . Then the reduced model can be written as

$$\mathbf{y} = \mathbf{X}_r \boldsymbol{\beta}_r + \boldsymbol{\varepsilon}, \quad (10.25)$$

and $\hat{\boldsymbol{\beta}}_r = (\mathbf{X}'_r \mathbf{X}_r)^{-1} \mathbf{X}'_r \mathbf{y}$. To compare the fit of the full model and the reduced model, we calculate

$$\hat{\boldsymbol{\beta}}' \mathbf{X}' \mathbf{y} - \hat{\boldsymbol{\beta}}'_r \mathbf{X}'_r \mathbf{y}, \quad (10.26)$$

where $\hat{\boldsymbol{\beta}}' \mathbf{X}' \mathbf{y}$ is the regression sum of squares from the full model and $\hat{\boldsymbol{\beta}}'_r \mathbf{X}'_r \mathbf{y}$ is the regression sum of squares for the reduced model. The difference in (10.26) shows what β_d contributes "above and beyond" β_r . We can test $H_0: \beta_d = \mathbf{0}$ with an F -statistic:

$$F = \frac{(\hat{\boldsymbol{\beta}}' \mathbf{X}' \mathbf{y} - \hat{\boldsymbol{\beta}}'_r \mathbf{X}'_r \mathbf{y})/h}{(y'y - \hat{\boldsymbol{\beta}}' \mathbf{X}' \mathbf{y})/(n - q - 1)} \quad (10.27)$$

$$= \frac{(SSR_f - SSR_r)/h}{SSE_f/(n - q - 1)} = \frac{MSR}{MSE}, \quad (10.28)$$

where $SSR_f = \hat{\boldsymbol{\beta}}' \mathbf{X}' \mathbf{y}$ and $SSR_r = \hat{\boldsymbol{\beta}}'_r \mathbf{X}'_r \mathbf{y}$. The F -statistic in (10.27) and (10.28) is distributed as $F_{h, n-q-1}$ if H_0 is true. We reject H_0 if $F > F_{\alpha, h, n-q-1}$.

The test in (10.27) is easy to carry out in practice. We fit the full model and obtain the regression and error sums of squares $\hat{\boldsymbol{\beta}}' \mathbf{X}' \mathbf{y}$ and $y'y - \hat{\boldsymbol{\beta}}' \mathbf{X}' \mathbf{y}$, respectively. We then fit the reduced model and obtain its regression sum of squares $\hat{\boldsymbol{\beta}}'_r \mathbf{X}'_r \mathbf{y}$ to be subtracted from $\hat{\boldsymbol{\beta}}' \mathbf{X}' \mathbf{y}$. If a software package gives the regression sum of squares in corrected form, this can readily be used to obtain $\hat{\boldsymbol{\beta}}' \mathbf{X}' \mathbf{y} - \hat{\boldsymbol{\beta}}'_r \mathbf{X}'_r \mathbf{y}$, since

$$\hat{\boldsymbol{\beta}}' \mathbf{X}' \mathbf{y} - n\bar{y}^2 - (\hat{\boldsymbol{\beta}}'_r \mathbf{X}'_r \mathbf{y} - n\bar{y}^2) = \hat{\boldsymbol{\beta}}' \mathbf{X}' \mathbf{y} - \hat{\boldsymbol{\beta}}'_r \mathbf{X}'_r \mathbf{y}.$$

Alternatively, we can obtain $\hat{\boldsymbol{\beta}}' \mathbf{X}' \mathbf{y} - \hat{\boldsymbol{\beta}}'_r \mathbf{X}'_r \mathbf{y}$ as the difference between error sums of squares for the two models:

$$\begin{aligned} SSE_r - SSE_f &= y'y - \hat{\boldsymbol{\beta}}'_r \mathbf{X}'_r \mathbf{y} - (y'y - \hat{\boldsymbol{\beta}}' \mathbf{X}' \mathbf{y}) \\ &= \hat{\boldsymbol{\beta}}' \mathbf{X}' \mathbf{y} - \hat{\boldsymbol{\beta}}'_r \mathbf{X}'_r \mathbf{y}. \end{aligned}$$

A test for an individual β_j above and beyond the other β 's is readily obtained using (10.27). To test $H_0: \beta_j = 0$, we arrange β_j last in $\boldsymbol{\beta}$,

$$\boldsymbol{\beta} = \begin{pmatrix} \boldsymbol{\beta}_r \\ \beta_j \end{pmatrix},$$

where $\boldsymbol{\beta}_r = (\beta_0, \beta_1, \dots, \beta_{q-1})'$ contains all the β 's except β_j . By (10.27), the test statistic is

$$F = \frac{\hat{\boldsymbol{\beta}}' \mathbf{X}' \mathbf{y} - \hat{\boldsymbol{\beta}}'_r \mathbf{X}'_r \mathbf{y}}{SSE_f/(n - q - 1)}, \quad (10.29)$$

which is $F_{1,n-q-1}$. Note that $h = 1$. The test of $H_0: \beta_j = 0$ made by the F -statistic in (10.29) is called a *partial F-test*. A detailed breakdown of the effect of each variable in the presence of the others is given by Rencher (1993) and Rencher and Schaalje (2008, Section 10.6).

Since the F -statistic in (10.29) has 1 and $n - q - 1$ degrees of freedom, it is the square of a t -statistic. The t -statistic equivalent to (10.29) is

$$t = \frac{\hat{\beta}_j}{s\sqrt{g_{jj}}},$$

where g_{jj} is the j th diagonal element of $(\mathbf{X}'\mathbf{X})^{-1}$ and $s = \sqrt{\text{SSE}_f/(n - q - 1)}$ (Rencher and Schaalje 2008, Section 8.5.1).

10.2.6 R^2 in Fixed- x Regression

The proportion of the (corrected) total variation in the y 's that can be attributed to regression on the x 's is denoted by R^2 :

$$\begin{aligned} R^2 &= \frac{\text{regression sum of squares}}{\text{total sum of squares}} \\ &= \frac{\hat{\beta}' \mathbf{X}' \mathbf{y} - n\bar{y}^2}{\mathbf{y}' \mathbf{y} - n\bar{y}^2}. \end{aligned} \quad (10.30)$$

The ratio R^2 is called the *coefficient of multiple determination*, or more commonly the *squared multiple correlation*. The *multiple correlation* R is defined as the positive square root of R^2 .

The F -test for overall regression in (10.24) can be expressed in terms of R^2 as

$$F = \frac{n - q - 1}{q} \frac{R^2}{1 - R^2}. \quad (10.31)$$

For the reduced model (10.25), R^2 can be written as

$$R_r^2 = \frac{\hat{\beta}_r' \mathbf{X}_r' \mathbf{y} - n\bar{y}^2}{\mathbf{y}' \mathbf{y} - n\bar{y}^2}. \quad (10.32)$$

Then in terms of R^2 and R_r^2 , the full and reduced model test in (10.27) for $H_0: \beta_d = \mathbf{0}$ becomes

$$F = \frac{(R^2 - R_r^2)/h}{(1 - R^2)/(n - q - 1)} \quad (10.33)$$

[see (11.36)].

We can express R^2 in terms of sample variances, covariances, and correlations:

$$R^2 = \frac{\mathbf{s}'_{yx} \mathbf{S}_{xx}^{-1} \mathbf{s}_{yx}}{\mathbf{s}_{yy}} = \mathbf{r}'_{yx} \mathbf{R}_{xx}^{-1} \mathbf{r}_{yx}, \quad (10.34)$$

where s_{yy} , s_{yx} , and \mathbf{S}_{xx} are defined in (10.16) and \mathbf{r}_{yx} and \mathbf{R}_{xx} are from an analogous partitioning of the sample correlation matrix of y and the x 's:

$$\mathbf{R} = \left(\begin{array}{c|ccccc} 1 & r_{y1} & r_{y2} & \cdots & r_{yq} \\ \hline r_{1y} & 1 & r_{12} & \cdots & r_{1q} \\ \vdots & \vdots & \vdots & & \vdots \\ r_{qy} & r_{q1} & r_{q2} & \cdots & 1 \end{array} \right) = \begin{pmatrix} 1 & \mathbf{r}'_{yx} \\ \mathbf{r}_{yx} & \mathbf{R}_{xx} \end{pmatrix}. \quad (10.35)$$

10.2.7 Subset Selection

In practice, one often has more x 's than are needed for predicting y . Some of them may be redundant and could be discarded. In addition to logistical motivations for deleting variables, there are statistical incentives; for example, if an x is deleted from the fitted model, the variances of the $\hat{\beta}_j$'s and of the \hat{y}_i 's are reduced. Various aspects of model validation are reviewed by Rencher and Schaalje (2008, Section 7.9 and Chapter 9).

The two most popular approaches to subset selection are to (1) examine all possible subsets and (2) use a stepwise technique. We discuss these in the next two sections.

10.2.7a All Possible Subsets

The optimal approach to subset selection is to examine all possible subsets of the x 's. This may not be computationally feasible if the sample size and number of variables are large. Some programs take advantage of algorithms that find the optimum subset of each size without examining all of the subsets [see, for example, Furnival and Wilson (1974)].

We discuss three criteria for comparing subsets when searching for the best subset. To conform with established notation in the literature, the number of variables in a subset is denoted by $p - 1$, so that with the inclusion of an intercept, there are p parameters in the model. The corresponding total number of available variables from which a subset is to be selected is denoted by $k - 1$, with k parameters in the model.

1. R_p^2 . By its definition in (10.30) as the proportion of total (corrected) sum of squares accounted for by regression, R^2 is clearly a measure of model fit. The subscript p is an index of the subset size, since it indicates the number of parameters in the model, including an intercept. However, R_p^2 does not reach a maximum for any value of p less than k because it cannot decrease when a variable is added to the model. The usual procedure is to find the subset with largest R_p^2 for each of $p = 2, 3, \dots, k$ and then choose a value of p beyond which the increases in R^2 appear to be unimportant. This judgment is, of course, subjective.

2. s_p^2 . Another useful criterion is the variance estimator for each subset as defined in (10.8):

$$s_p^2 = \frac{\text{SSE}_p}{n - p}. \quad (10.36)$$

For each of $p = 2, 3, \dots, k$, we find the subset with smallest s_p^2 . If k is fairly large, a typical pattern as p approaches k is for the minimal s_p^2 to decrease to an overall minimum less than s_k^2 and then increase. The minimum value of s_p^2 can be less than s_k^2 if the decrease in SSE_p with an additional variable does not offset the loss of a degree of freedom in the denominator. It is often suggested that the researcher choose the subset with absolute minimum s_p^2 . However, as Hocking (1976, p.19) notes, this procedure may fit some noise unique to the sample and thereby include one or more extraneous predictor variables. An alternative suggestion is to choose p such that $\min_p s_p^2 = s_k^2$, or more precisely, choose the smallest value of p such that $\min_p s_p^2 < s_k^2$, since there will not be a $p < k$ such that $\min_p s_p^2$ is exactly equal to s_k^2 .

3. C_p . The C_p criterion is due to Mallows (1964, 1973). In the following development, we follow Myers (1990, pp. 180–182). The *expected squared error*, $E[\hat{y}_i - E(\hat{y}_i)]^2$, is used in formulating the C_p criterion because it incorporates a variance component and a bias component. The goal is to find a model that achieves a good balance between the bias and variance of the fitted values, \hat{y}_i . Bias arises when the \hat{y}_i values are based on an incorrect model, in which $E(\hat{y}_i) \neq E(y_i)$. If \hat{y}_i were based on the correct model, so that $E(\hat{y}_i) = E(y_i)$, then $E[\hat{y}_i - E(y_i)]^2$ would be equal to $\text{var}(\hat{y}_i)$. In general, however, as we examine many competing models, for various values of p , \hat{y}_i is not based on the correct model, and we have (see Problem 10.4)

$$\begin{aligned} E[\hat{y}_i - E(y_i)]^2 &= E[\hat{y}_i - E(\hat{y}_i) + E(\hat{y}_i) - E(y_i)]^2 \\ &= E[\hat{y}_i - E(\hat{y}_i)]^2 + [E(\hat{y}_i) - E(y_i)]^2 \end{aligned} \quad (10.37)$$

$$= \text{var}(\hat{y}_i) + (\text{bias in } \hat{y}_i)^2. \quad (10.38)$$

For a given value of p , the total expected squared error for the n observations in the sample, standardized by dividing by σ^2 , becomes

$$\frac{1}{\sigma^2} \sum_{i=1}^n [\hat{y}_i - E(y_i)]^2 = \frac{1}{\sigma^2} \sum_{i=1}^n \text{var}(\hat{y}_i) + \frac{1}{\sigma^2} \sum_{i=1}^n (\text{bias in } \hat{y}_i)^2. \quad (10.39)$$

Before defining C_p as an estimate of (10.39), we can achieve some simplification. We first show that $\sum_i \text{var}(\hat{y}_i)/\sigma^2$ is equal to p . Let the model for all n observations be designated by

$$\mathbf{y} = \mathbf{X}_p \boldsymbol{\beta}_p + \boldsymbol{\varepsilon}.$$

We assume that, in general, this prospective model is underspecified and that the true model (which produces σ^2) contains additional β 's and additional columns of the \mathbf{X} matrix. If we designate the i th row of \mathbf{X}_p by \mathbf{x}'_{pi} , then the first term on the right side

of (10.39) becomes (see also Problem 10.5)

$$\begin{aligned}\frac{1}{\sigma^2} \sum_{i=1}^n \text{var}(\hat{y}_i) &= \frac{1}{\sigma^2} \sum_{i=1}^n \text{var}(\mathbf{x}'_{pi} \hat{\beta}_p) \\ &= \frac{1}{\sigma^2} \sum_i \mathbf{x}'_{pi} [\sigma^2 (\mathbf{X}'_p \mathbf{X}_p)^{-1}] \mathbf{x}_{pi} \quad [\text{by (3.70)}] \\ &= \text{tr}[\mathbf{X}_p (\mathbf{X}'_p \mathbf{X}_p)^{-1} \mathbf{X}'_p] \quad [\text{by (3.65)}] \quad (10.40)\end{aligned}$$

$$\begin{aligned}&= \text{tr}[(\mathbf{X}'_p \mathbf{X}_p)^{-1} \mathbf{X}'_p \mathbf{X}_p] \quad [\text{by (2.97)}] \\ &= \text{tr}(\mathbf{I}_p) = p. \quad (10.41)\end{aligned}$$

It can be shown (Myers 1990, pp. 178–179) that

$$\sum_{i=1}^n (\text{bias in } \hat{y}_i)^2 = (n - p) E(s_p^2 - \sigma^2). \quad (10.42)$$

Using (10.41) and (10.42), the final simplified form of the (standardized) total expected squared error in (10.39) is

$$\frac{1}{\sigma^2} \sum_{i=1}^n [\hat{y}_i - E(y_i)]^2 = p + \frac{n - p}{\sigma^2} E(s_p^2 - \sigma^2). \quad (10.43)$$

In practice, σ^2 is usually estimated by s_k^2 , the MSE from the full model. We thus estimate (10.43) by

$$C_p = p + (n - p) \frac{s_p^2 - s_k^2}{s_k^2}. \quad (10.44)$$

An alternative form is

$$C_p = \frac{\text{SSE}_p}{s_k^2} - (n - 2p). \quad (10.45)$$

In (10.44), we see that if the bias is small for a particular model, C_p will be close to p . For this reason, the line $C_p = p$ is commonly plotted along with the C_p values of several candidate models. We look for small values of C_p that are near this line.

In a Monte Carlo study, Hilton (1983) compared several subset selection criteria based on MSE_p and C_p . The three best procedures were to choose (1) the subset with the smallest p such that $C_p < p$, (2) the subset with the smallest p such that $s_p^2 < s_k^2$, and (3) the subset with minimum s_p^2 . The first of these was found to give best results overall, with the second method close behind. The third method performed best in some cases where k was small.

10.2.7b Stepwise Selection

For many data sets, it may be impractical to examine all possible subsets, even with an efficient algorithm such as that of Furnival and Wilson (1974). In such cases, we can use the familiar stepwise approach, which is widely available and has virtually no limit as to the number of variables or observations. A related stepwise technique

was discussed in Sections 6.11.2 and 8.9 in connection with selection of dependent variables to separate groups in a MANOVA or discriminant analysis setting. In this section, we are concerned with selecting the independent variables (x 's) that best predict the dependent variable (y) in regression.

We first review the *forward* selection procedure, which typically uses an F -test at each step. At the first step, y is regressed on each x_j alone and the x with the largest F -value is “entered” into the model. At the second step, we search for the variable with the largest *partial F*-value for testing the significance of each variable in the presence of the variable first entered. Thus, if we denote the first variable to enter as x_1 , then at the second step we calculate the partial F -statistic

$$F = \frac{\text{MSR}(x_j|x_1)}{\text{MSE}(x_j, x_1)}$$

for each $j \neq 1$ and choose the variable that maximizes F , where $\text{MSR} = (\text{SSR}_f - \text{SSR}_r)/h$ and $\text{MSE} = \text{SSE}_f/(n-q-1)$ are the mean squares for regression and error, respectively, as in (10.28). In this case, $\text{SSR}_f = \text{SSR}(x_1, x_j)$ and $\text{SSR}_r = \text{SSR}(x_1)$. Note also that $h = 1$ because only one variable is being added, and MSE is calculated using only the variable already entered plus the candidate variable. This procedure continues at each step until the largest partial F for an entering variable falls below a preselected threshold F -value or until the corresponding p -value exceeds some predetermined level.

The *stepwise* selection procedure similarly seeks the best variable to enter at each step. Then after a variable has entered, each of the variables previously entered is examined by a partial F -test to see if it is no longer significant and can be dropped from the model.

The *backward elimination* procedure begins with all x 's in the model and deletes one at a time. The partial F -statistic for each variable in the presence of the others is calculated, and the variable with smallest F is eliminated. This continues until the smallest F at some step exceeds a preselected threshold value.

Since these sequential methods do not examine all subsets, they will often fail to find the optimum subset, especially if k is large. However, R_p^2 , s_p^2 , or C_p may not differ substantially between the optimum subset and the one found by stepwise selection. These sequential methods have been popular for at least a generation, and it is very likely they will continue to be used, even though increased computing power has put the optimal methods within reach for larger data sets.

There are some possible risks in the use of stepwise methods. The stepwise procedure may fail to detect a true predictor (an x_j for which $\beta_j \neq 0$) because s_p^2 is biased upward in an underspecified model, thus artificially reducing the partial F -value. On the other hand, a variable that is not a true predictor of y (an x_j for which $\beta_j = 0$) may enter because of chance correlations in a particular sample. In the presence of such “noise” variables, the partial F -statistic for the entering variable does not have an F -distribution because it is maximized at each step. The calculated p -values become optimistic. This problem intensifies when the sample size is relatively small compared to the number of variables. Rencher and Pun (1980) found that in such cases some surprisingly large values of R^2 can occur, even when there is no relation-

ship between y and the x 's in the population. In a related study, Flack and Chang (1987) included x 's that were authentic contributors as well as noise variables. They found that "for most samples, a large percentage of the selected variables is noise, particularly when the number of candidate variables is large relative to the number of observations. The adjusted R^2 of the selected variables is highly inflated" (p. 84).

10.3 MULTIPLE REGRESSION: RANDOM x 'S

In Section 10.2, it was assumed that the x 's were fixed and would have the same values if another sample were taken; that is, the same \mathbf{X} matrix would be used each time a y vector was observed. However, many regression applications involve x 's that are random variables.

Thus in the random- x case, the values of x_1, x_2, \dots, x_q are not under the control of the experimenter. They occur randomly along with y . On each subject we observe y, x_1, x_2, \dots, x_q .

If we assume that $(y, x_1, x_2, \dots, x_q)$ has a multivariate normal distribution, then $\hat{\beta}$, R^2 , and the F -tests have the same formulation as in the fixed- x case [for details, see Rencher (1998, Section 7.3) or Rencher and Schaalje (2008, Section 10.5)]. Thus with the multivariate normal assumption, we can proceed with estimation and testing the same way in the random- x case as with fixed x 's.

10.4 MULTIVARIATE MULTIPLE REGRESSION: ESTIMATION

In this section we extend the estimation results of Sections 10.2.2–10.2.4 to the multivariate y case. We assume the x 's are fixed.

10.4.1 The Multivariate Linear Model

We turn now to the *multivariate multiple regression model*, where *multivariate* refers to the dependent variables and *multiple* pertains to the independent variables. In this case, several y 's are measured corresponding to each set of x 's. Each of y_1, y_2, \dots, y_p is to be predicted by all of x_1, x_2, \dots, x_q .

The n observed values of the vector of y 's can be listed as rows in the following matrix:

$$\mathbf{Y} = \begin{pmatrix} y_{11} & y_{12} & \cdots & y_{1p} \\ y_{21} & y_{22} & \cdots & y_{2p} \\ \vdots & \vdots & & \vdots \\ y_{n1} & y_{n2} & \cdots & y_{np} \end{pmatrix} = \begin{pmatrix} \mathbf{y}'_1 \\ \mathbf{y}'_2 \\ \vdots \\ \mathbf{y}'_n \end{pmatrix}.$$

Thus each row of \mathbf{Y} contains the values of the p dependent variables measured on a subject. Each column of \mathbf{Y} consists of the n observations on one of the p variables and therefore corresponds to the y vector in the (univariate) regression model (10.3).

The n values of x_1, x_2, \dots, x_q can be placed in a matrix that turns out to be the same as the \mathbf{X} matrix in the multiple regression formulation in Section 10.2.1:

$$\mathbf{X} = \begin{pmatrix} 1 & x_{11} & x_{12} & \cdots & x_{1q} \\ 1 & x_{21} & x_{22} & \cdots & x_{2q} \\ \vdots & \vdots & \vdots & & \vdots \\ 1 & x_{n1} & x_{n2} & \cdots & x_{nq} \end{pmatrix}.$$

We assume that \mathbf{X} is fixed from sample to sample.

Since each of the p y 's will depend on the x 's in its own way, each column of \mathbf{Y} will need different β 's. Thus we have a column of β 's for each column of \mathbf{Y} , and these columns form a matrix $\mathbf{B} = (\beta_1, \beta_2, \dots, \beta_p)$. Our multivariate model is therefore

$$\mathbf{Y} = \mathbf{XB} + \boldsymbol{\Xi}, \quad (10.46)$$

where \mathbf{Y} is $n \times p$, \mathbf{X} is $n \times (q+1)$, and \mathbf{B} is $(q+1) \times p$. The notation $\boldsymbol{\Xi}$ (the uppercase version of ξ) is adopted here because of its resemblance to ε .

We illustrate the multivariate model with $p = 2, q = 3$:

$$\begin{pmatrix} y_{11} & y_{12} \\ y_{21} & y_{22} \\ \vdots & \vdots \\ y_{n1} & y_{n2} \end{pmatrix} = \begin{pmatrix} 1 & x_{11} & x_{12} & x_{13} \\ 1 & x_{21} & x_{22} & x_{23} \\ \vdots & \vdots & \vdots & \vdots \\ 1 & x_{n1} & x_{n2} & x_{n3} \end{pmatrix} \begin{pmatrix} \beta_{01} & \beta_{02} \\ \beta_{11} & \beta_{12} \\ \beta_{21} & \beta_{22} \\ \beta_{31} & \beta_{32} \end{pmatrix} + \begin{pmatrix} \varepsilon_{11} & \varepsilon_{12} \\ \varepsilon_{21} & \varepsilon_{22} \\ \vdots & \vdots \\ \varepsilon_{n1} & \varepsilon_{n2} \end{pmatrix}.$$

The model for the first column of \mathbf{Y} is

$$\begin{pmatrix} y_{11} \\ y_{21} \\ \vdots \\ y_{n1} \end{pmatrix} = \begin{pmatrix} 1 & x_{11} & x_{12} & x_{13} \\ 1 & x_{21} & x_{22} & x_{23} \\ \vdots & \vdots & \vdots & \vdots \\ 1 & x_{n1} & x_{n2} & x_{n3} \end{pmatrix} \begin{pmatrix} \beta_{01} \\ \beta_{11} \\ \beta_{21} \\ \beta_{31} \end{pmatrix} + \begin{pmatrix} \varepsilon_{11} \\ \varepsilon_{21} \\ \vdots \\ \varepsilon_{n1} \end{pmatrix},$$

and for the second column, we have

$$\begin{pmatrix} y_{12} \\ y_{22} \\ \vdots \\ y_{n2} \end{pmatrix} = \begin{pmatrix} 1 & x_{11} & x_{12} & x_{13} \\ 1 & x_{21} & x_{22} & x_{23} \\ \vdots & \vdots & \vdots & \vdots \\ 1 & x_{n1} & x_{n2} & x_{n3} \end{pmatrix} \begin{pmatrix} \beta_{02} \\ \beta_{12} \\ \beta_{22} \\ \beta_{32} \end{pmatrix} + \begin{pmatrix} \varepsilon_{12} \\ \varepsilon_{22} \\ \vdots \\ \varepsilon_{n2} \end{pmatrix}.$$

By analogy with the univariate case in Section 10.2.1, additional assumptions that lead to good estimates are as follows:

1. $E(\mathbf{Y}) = \mathbf{XB}$ or $E(\boldsymbol{\Xi}) = \mathbf{O}$.
2. $\text{cov}(\mathbf{y}_i) = \boldsymbol{\Sigma}$ for all $i = 1, 2, \dots, n$, where \mathbf{y}'_i is the i th row of \mathbf{Y} .
3. $\text{cov}(\mathbf{y}_i, \mathbf{y}_j) = \mathbf{O}$ for all $i \neq j$.

Assumption 1 states that the linear model is correct and that no additional x 's are needed to predict the y 's. Assumption 2 asserts that each of the n observation vectors (rows) in \mathbf{Y} has the same covariance matrix. Assumption 3 declares that observation vectors (rows of \mathbf{Y}) are uncorrelated with each other. Thus we assume that the y 's within an observation vector (row of \mathbf{Y}) are correlated with each other but independent of the y 's in any other observation vector.

The covariance matrix Σ in assumption 2 contains the variances and covariances of $y_{i1}, y_{i2}, \dots, y_{ip}$ in any \mathbf{y}_i :

$$\text{cov}(\mathbf{y}_i) = \Sigma = \begin{pmatrix} \sigma_{11} & \sigma_{12} & \cdots & \sigma_{1p} \\ \sigma_{21} & \sigma_{22} & \cdots & \sigma_{2p} \\ \vdots & \vdots & & \vdots \\ \sigma_{p1} & \sigma_{p2} & \cdots & \sigma_{pp} \end{pmatrix}.$$

The covariance matrix \mathbf{O} in assumption 3 contains the covariances of each of $y_{i1}, y_{i2}, \dots, y_{ip}$ with each of $y_{j1}, y_{j2}, \dots, y_{jp}$:

$$\begin{pmatrix} \text{cov}(y_{i1}, y_{j1}) & \text{cov}(y_{i1}, y_{j2}) & \cdots & \text{cov}(y_{i1}, y_{jp}) \\ \text{cov}(y_{i2}, y_{j1}) & \text{cov}(y_{i2}, y_{j2}) & \cdots & \text{cov}(y_{i2}, y_{jp}) \\ \vdots & \vdots & & \vdots \\ \text{cov}(y_{ip}, y_{j1}) & \text{cov}(y_{ip}, y_{j2}) & \cdots & \text{cov}(y_{ip}, y_{jp}) \end{pmatrix} = \begin{pmatrix} 0 & 0 & \cdots & 0 \\ 0 & 0 & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & 0 \end{pmatrix}.$$

To aid in the discussion of properties of multivariate multiple regression, we will often choose to consider the multivariate model (10.46) re-expressed in a form that is more directly analogous to the univariate model in (10.3). Using Kronecker and vec notation and property (2.122) discussed in Section 2.12, we re-write the model as

$$\begin{aligned} \text{vec } \mathbf{Y} &= \text{vec } \mathbf{XB} + \text{vec } \boldsymbol{\Xi} \\ &= (\mathbf{I}_p \otimes \mathbf{X}) \text{vec } \mathbf{B} + \text{vec } \boldsymbol{\Xi} \\ &= \mathbf{X}^* \boldsymbol{\beta}^* + \boldsymbol{\varepsilon}^*, \end{aligned} \tag{10.47}$$

where $\mathbf{X}^* = \mathbf{I}_p \otimes \mathbf{X}$ is a $pn \times p(q+1)$ block-diagonal matrix, $\boldsymbol{\beta}^* = \text{vec } \mathbf{B}$ is a vector of length $p(q+1)$, and $\boldsymbol{\varepsilon}^* = \text{vec } \boldsymbol{\Xi}$ is a vector of length pn . Note that model (10.47) resembles the univariate model presented in (10.3), with $\text{vec } \mathbf{Y}$ as our vector of responses, \mathbf{X}^* as our full-column-rank matrix of predictors, $\boldsymbol{\beta}^*$ as our vector of regression coefficients, and $\boldsymbol{\varepsilon}^*$ as our vector of random errors.

10.4.2 Least Squares Estimation in the Multivariate Model

By analogy with the univariate case in (10.5), we estimate \mathbf{B} with

$$\hat{\mathbf{B}} = (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{Y}. \tag{10.48}$$

We call $\hat{\mathbf{B}}$ the *least squares estimator* for \mathbf{B} because it “minimizes” $\mathbf{E} = \hat{\boldsymbol{\Xi}}'\hat{\boldsymbol{\Xi}}$, a matrix analogous to SSE:

$$\mathbf{E} = \hat{\boldsymbol{\Xi}}'\hat{\boldsymbol{\Xi}} = (\mathbf{Y} - \mathbf{X}\hat{\mathbf{B}})'(\mathbf{Y} - \mathbf{X}\hat{\mathbf{B}}).$$

The matrix $\hat{\mathbf{B}}$ minimizes \mathbf{E} in the following sense. If we let \mathbf{B}_0 be an estimate that may possibly be better than $\hat{\mathbf{B}}$ and add $\mathbf{X}\hat{\mathbf{B}} - \mathbf{X}\mathbf{B}_0$ to $\mathbf{Y} - \mathbf{X}\hat{\mathbf{B}}$, we find that this adds a positive definite matrix to $\mathbf{E} = (\mathbf{Y} - \mathbf{X}\hat{\mathbf{B}})'(\mathbf{Y} - \mathbf{X}\hat{\mathbf{B}})$ (Rencher 1998, Section 7.4.2). Thus we cannot improve on $\hat{\mathbf{B}}$. The least squares estimate $\hat{\mathbf{B}}$ also minimizes the scalar quantities $\text{tr}(\mathbf{Y} - \mathbf{X}\hat{\mathbf{B}})'(\mathbf{Y} - \mathbf{X}\hat{\mathbf{B}})$ and $|(\mathbf{Y} - \mathbf{X}\hat{\mathbf{B}})'(\mathbf{Y} - \mathbf{X}\hat{\mathbf{B}})|$. Note that by (2.98) $\text{tr}(\mathbf{Y} - \mathbf{X}\hat{\mathbf{B}})'(\mathbf{Y} - \mathbf{X}\hat{\mathbf{B}}) = \sum_{i=1}^n \sum_{j=1}^p \hat{\varepsilon}_{ij}^2$.

We noted earlier that in the model $\mathbf{Y} = \mathbf{XB} + \boldsymbol{\Xi}$, there is a column of \mathbf{B} corresponding to each column of \mathbf{Y} ; that is, each $y_j, j = 1, 2, \dots, p$, is predicted differently by x_1, x_2, \dots, x_q . (This is illustrated in Section 10.4.1 for $p = 2$). In the estimate $\hat{\mathbf{B}} = (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{Y}$, we have a similar pattern. The matrix product $(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'$ is multiplied by each column of \mathbf{Y} [see (2.48)]. Thus the j th column of $\hat{\mathbf{B}}$ is the usual least squares estimate $\hat{\beta}$ for the j th dependent variable y_j . To give this a more precise expression, let us denote the p columns of \mathbf{Y} by $\mathbf{y}_{(1)}, \mathbf{y}_{(2)}, \dots, \mathbf{y}_{(p)}$ to distinguish them from the n rows $\mathbf{y}'_i, i = 1, 2, \dots, n$. Then

$$\begin{aligned}\hat{\mathbf{B}} &= (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{Y} = (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'(\mathbf{y}_{(1)}, \mathbf{y}_{(2)}, \dots, \mathbf{y}_{(p)}) \\ &= [(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y}_{(1)}, (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y}_{(2)}, \dots, (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y}_{(p)}] \\ &= [\hat{\beta}_{(1)}, \hat{\beta}_{(2)}, \dots, \hat{\beta}_{(p)}].\end{aligned}\quad (10.49)$$

■ EXAMPLE 10.4.2

The results of a planned experiment involving a chemical reaction are given in Table 10.1 (Box and Youle 1955).

The input (independent) variables are

$$x_1 = \text{temperature}, \quad x_2 = \text{concentration}, \quad x_3 = \text{time}.$$

The yield (dependent) variables are

$$y_1 = \text{percentage of unchanged starting material},$$

$$y_2 = \text{percentage converted to the desired product},$$

$$y_3 = \text{percentage of unwanted by-product}.$$

Using (10.48), the least squares estimator $\hat{\mathbf{B}}$ for the regression of (y_1, y_2, y_3) on (x_1, x_2, x_3) is given by

$$\begin{aligned}\hat{\mathbf{B}} &= (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{Y} \\ &= \begin{pmatrix} 332.11 & -26.04 & -164.08 \\ -1.55 & .40 & .91 \\ -1.42 & .29 & .90 \\ -2.24 & 1.03 & 1.15 \end{pmatrix}.\end{aligned}$$

Note that the first column of $\hat{\mathbf{B}}$ gives $\hat{\beta}_0, \hat{\beta}_1, \hat{\beta}_2, \hat{\beta}_3$ for regression of y_1 on x_1, x_2, x_3 ; the second column of $\hat{\mathbf{B}}$ gives $\hat{\beta}_0, \hat{\beta}_1, \hat{\beta}_2, \hat{\beta}_3$ for regression of y_2 on x_1, x_2, x_3 , and so on. \square

Table 10.1 Chemical Reaction Data

Experiment Number	Yield Variables			Input Variables		
	y_1	y_2	y_3	x_1	x_2	x_3
1	41.5	45.9	11.2	162	23	3
2	33.8	53.3	11.2	162	23	8
3	27.7	57.5	12.7	162	30	5
4	21.7	58.8	16.0	162	30	8
5	19.9	60.6	16.2	172	25	5
6	15.0	58.0	22.6	172	25	8
7	12.2	58.6	24.5	172	30	5
8	4.3	52.4	38.0	172	30	8
9	19.3	56.9	21.3	167	27.5	6.5
10	6.4	55.4	30.8	177	27.5	6.5
11	37.6	46.9	14.7	157	27.5	6.5
12	18.0	57.3	22.2	167	32.5	6.5
13	26.3	55.0	18.3	167	22.5	6.5
14	9.9	58.9	28.0	167	27.5	9.5
15	25.0	50.3	22.1	167	27.5	3.5
16	14.1	61.1	23.0	177	20	6.5
17	15.2	62.9	20.7	177	20	6.5
18	15.9	60.0	22.1	160	34	7.5
19	19.6	60.6	19.3	160	34	7.5

10.4.3 Properties of Least Squares Estimator $\hat{\mathbf{B}}$

The least squares estimator $\hat{\mathbf{B}}$ can be obtained without imposing the assumptions $E(\mathbf{y}) = \mathbf{XB}$, $\text{cov}(\mathbf{y}_i) = \Sigma$, and $\text{cov}(\mathbf{y}_i, \mathbf{y}_j) = \mathbf{O}$. However, when these assumptions hold, $\hat{\mathbf{B}}$ has the following properties:

1. The estimator $\hat{\mathbf{B}}$ is unbiased, that is, $E(\hat{\mathbf{B}}) = \mathbf{B}$. This means that if we took repeated random samples from the same population, the average value of $\hat{\mathbf{B}}$ would be \mathbf{B} .
2. All $\hat{\beta}_{jk}$'s in $\hat{\mathbf{B}}$ are correlated with each other. This is due to the correlations among the x 's and among the y 's. The $\hat{\beta}$'s within a given column of $\hat{\mathbf{B}}$ are correlated because x_1, x_2, \dots, x_q are correlated. If x_1, x_2, \dots, x_q were orthogonal to each other, the $\hat{\beta}$'s within each column of $\hat{\mathbf{B}}$ would be uncorrelated. Thus the relationship of the x 's to each other affects the relationship of the $\hat{\beta}$'s within each column to each other. On the other hand, the $\hat{\beta}$'s in each column are correlated with $\hat{\beta}$'s in other columns because y_1, y_2, \dots, y_p are correlated.

Because of the correlations among the columns of $\hat{\mathbf{B}}$, we need multivariate tests for hypotheses about \mathbf{B} . We cannot use an F -test from Section 10.2.5 on each column of \mathbf{B} , because these F -tests would not take into account the

correlations or preserve the α -level. Some appropriate multivariate tests are given in Section 10.5.

3. The least squares estimators $\hat{\beta}_{jk}$ in $\hat{\mathbf{B}}$ have minimum variance among all possible linear unbiased estimators. In other words, $\hat{\mathbf{B}}$ is the best linear unbiased estimator (BLUE) for \mathbf{B} . This result is an extension of the Gauss–Markov theorem discussed in Section 10.2.2. The restriction to unbiased estimators is necessary to exclude trivial estimators such as a constant, which has variance equal to zero, but is of no interest. This minimum variance property of least squares estimators is remarkable for its distributional generality; normality of the y 's is not required. Equally remarkable is the fact that the BLUE for \mathbf{B} is *not* a function of Σ , the common covariance matrix for the observation vector \mathbf{y}_i , $i = 1, 2, \dots, n$.

To demonstrate that the least squares estimator $\hat{\mathbf{B}}$ is the BLUE for \mathbf{B} , we consider the multivariate multiple regression model (10.46) as written in the alternative form in (10.47). The least squares estimator of β^* in (10.47) is $\hat{\beta}^* = (\mathbf{X}^{*'}\mathbf{X}^*)^{-1}\mathbf{X}^{*'}\text{vec } \mathbf{Y}$, but it is not readily apparent that $\hat{\beta}^*$ is the best linear unbiased estimator of β^* because one of the conditions for the Gauss–Markov theorem does not hold for model (10.47). Specifically, the $pn \times pn$ covariance matrix for $\text{vec } \mathbf{Y}$ is *not* equal to $\sigma^2 \mathbf{I}_{pn}$, but rather

$$\text{cov}(\text{vec } \mathbf{Y}) = \text{cov} \begin{pmatrix} \mathbf{y}_{(1)} \\ \mathbf{y}_{(2)} \\ \vdots \\ \mathbf{y}_{(p)} \end{pmatrix} = \begin{pmatrix} \sigma_{11}\mathbf{I}_n & \sigma_{12}\mathbf{I}_n & \cdots & \sigma_{1p}\mathbf{I}_n \\ \sigma_{21}\mathbf{I}_n & \sigma_{22}\mathbf{I}_n & \cdots & \sigma_{2p}\mathbf{I}_n \\ \vdots & \vdots & \ddots & \vdots \\ \sigma_{p1}\mathbf{I}_n & \sigma_{p2}\mathbf{I}_n & \cdots & \sigma_{pp}\mathbf{I}_n \end{pmatrix} = \Sigma \otimes \mathbf{I}_n = \Sigma^*.$$

The generalized least squares (GLS) estimator of β^* can be obtained by noting that there exists a nonsingular $pn \times pn$ matrix \mathbf{P} such that the positive definite covariance matrix Σ^* is equal to $\mathbf{P}\mathbf{P}'$. (One such matrix \mathbf{P} is the square-root matrix of Σ^* , as described in Section 2.11.8.) Multiplying $\text{vec } \mathbf{Y} = \mathbf{X}^*\beta^* + \varepsilon^*$ by \mathbf{P}^{-1} , we obtain the model

$$\mathbf{P}^{-1}\text{vec } \mathbf{Y} = \mathbf{P}^{-1}\mathbf{X}^*\beta^* + \mathbf{P}^{-1}\varepsilon^*. \quad (10.50)$$

Note that $E(\mathbf{P}^{-1}\varepsilon^*) = \mathbf{P}^{-1}E(\varepsilon^*) = \mathbf{0}$ and

$$\begin{aligned} \text{cov}(\mathbf{P}^{-1}\varepsilon^*) &= \mathbf{P}^{-1}\text{cov}(\varepsilon^*)(\mathbf{P}^{-1})' \\ &= \mathbf{P}^{-1}\Sigma^*(\mathbf{P}^{-1})' \\ &= \mathbf{P}^{-1}\mathbf{P}\mathbf{P}'(\mathbf{P}^{-1})' = \mathbf{I}_{pn}. \end{aligned}$$

Thus, the conditions for the Gauss–Markov theorem discussed in Section 10.2.2 are satisfied for model (10.50), and the least squares estimator

$$\hat{\beta}^* = [(\mathbf{P}^{-1}\mathbf{X}^*)'(\mathbf{P}^{-1}\mathbf{X}^*)]^{-1}(\mathbf{P}^{-1}\mathbf{X}^*)'\mathbf{P}^{-1}\text{vec } \mathbf{Y}$$

is BLUE. This estimator can be reduced to the more commonly used form

$$\hat{\beta}^* = [\mathbf{X}^{*\prime} \Sigma^{*-1} \mathbf{X}^*]^{-1} \mathbf{X}^{*\prime} \Sigma^{*-1} \text{vec } \mathbf{Y}, \quad (10.51)$$

which is called the generalized least squares (GLS) estimator of β^* . However, using properties (2.121) – (2.124), we can show that the GLS estimator (10.51) can be simplified to the least squares estimator (10.48). That is,

$$\begin{aligned}\hat{\beta}^* &= \left[(\mathbf{I}_p \otimes \mathbf{X})' (\Sigma \otimes \mathbf{I}_n)^{-1} (\mathbf{I}_p \otimes \mathbf{X}) \right]^{-1} (\mathbf{I}_p \otimes \mathbf{X})' (\Sigma \otimes \mathbf{I}_n)^{-1} \text{vec } \mathbf{Y} \\ &= [(\Sigma^{-1} \otimes \mathbf{X}') (\mathbf{I}_p \otimes \mathbf{X})]^{-1} (\mathbf{I}_p \otimes \mathbf{X}') (\Sigma^{-1} \otimes \mathbf{I}_n) \text{ vec } \mathbf{Y} \\ &= [\Sigma^{-1} \otimes (\mathbf{X}' \mathbf{X})]^{-1} (\Sigma^{-1} \otimes \mathbf{X}') \text{ vec } \mathbf{Y} \\ &= (\Sigma \otimes (\mathbf{X}' \mathbf{X})^{-1}) (\Sigma^{-1} \otimes \mathbf{X}') \text{ vec } \mathbf{Y} \\ &= (\mathbf{I}_p \otimes (\mathbf{X}' \mathbf{X})^{-1} \mathbf{X}') \text{ vec } \mathbf{Y},\end{aligned}$$

which implies that by (2.122), the GLS estimator is equivalent to the least squares estimator $\hat{\mathbf{B}} = (\mathbf{X}' \mathbf{X})^{-1} \mathbf{X}' \mathbf{Y}$ from (10.48). Thus, the ordinary least squares estimator—which is *not* a function of Σ —is in fact the BLUE for \mathbf{B} .

4. The covariances among the elements of $\hat{\mathbf{B}}$ can be characterized using the alternative form of the multivariate regression model given in (10.47). The covariance matrix for $\hat{\beta}^* = \text{vec } \hat{\mathbf{B}}$ is

$$\begin{aligned}\text{cov}(\hat{\beta}^*) &= \text{cov}((\mathbf{X}^{*\prime} \mathbf{X}^*)^{-1} \mathbf{X}^{*\prime} \text{vec } \mathbf{Y}) \\ &= \text{cov}\left([(\mathbf{I}_p \otimes \mathbf{X})' (\mathbf{I}_p \otimes \mathbf{X})]^{-1} (\mathbf{I}_p \otimes \mathbf{X})' \text{vec } \mathbf{Y}\right) \\ &= \left[\mathbf{I}_p \otimes (\mathbf{X}' \mathbf{X})^{-1} \mathbf{X}'\right] \text{cov}(\text{vec } \mathbf{Y}) \left[\mathbf{I}_p \otimes (\mathbf{X}' \mathbf{X})^{-1} \mathbf{X}'\right]' \\ &= \left[\mathbf{I}_p \otimes (\mathbf{X}' \mathbf{X})^{-1} \mathbf{X}'\right] (\Sigma \otimes \mathbf{I}_n) \left[\mathbf{I}_p \otimes \mathbf{X} (\mathbf{X}' \mathbf{X})^{-1}\right] \\ &= \Sigma \otimes \left[(\mathbf{X}' \mathbf{X})^{-1} \mathbf{X}' \mathbf{I}_n \mathbf{X} (\mathbf{X}' \mathbf{X})^{-1}\right] \\ &= \Sigma \otimes (\mathbf{X}' \mathbf{X})^{-1}.\end{aligned} \quad (10.52)$$

The variances and covariances in $\text{cov}(\hat{\beta}^*)$ can be used to calculate standard errors for individual partial slopes in $\hat{\beta}^*$ or for linear combinations of the form $\mathbf{a}' \hat{\beta}^*$.

10.4.4 An Estimator for Σ

By analogy with (10.8) and (10.9), an unbiased estimator of $\text{cov}(\mathbf{y}_i) = \Sigma$ is given by

$$\mathbf{S}_e = \frac{\mathbf{E}}{n - q - 1} = \frac{(\mathbf{Y} - \mathbf{X}\hat{\mathbf{B}})'(\mathbf{Y} - \mathbf{X}\hat{\mathbf{B}})}{n - q - 1} \quad (10.53)$$

$$= \frac{\mathbf{Y}'\mathbf{Y} - \hat{\mathbf{B}}'\mathbf{X}'\mathbf{Y}}{n - q - 1}. \quad (10.54)$$

To demonstrate the unbiasedness of \mathbf{S}_e , we first note that

$$\begin{aligned} E(\mathbf{Y} - \mathbf{X}\hat{\mathbf{B}}) &= E\{[\mathbf{I}_n - \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}']\mathbf{Y}\} \\ &= [\mathbf{I}_n - \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'] E(\mathbf{X}\mathbf{B} + \boldsymbol{\Xi}) \\ &= [\mathbf{X} - \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{X}] \mathbf{B} + [\mathbf{I}_n - \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'] E(\boldsymbol{\Xi}) \\ &= [\mathbf{I}_n - \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'] E(\boldsymbol{\Xi}) = \mathbf{O}. \end{aligned}$$

Then, denoting the columns of $\mathbf{Y} - \mathbf{X}\hat{\mathbf{B}}$ with $\hat{\varepsilon}_{(1)}, \dots, \hat{\varepsilon}_{(p)}$, we can define the expectation of the (i, j) element of $(n - q - 1)\mathbf{S}_e$ as

$$\begin{aligned} E(\hat{\varepsilon}'_{(i)} \hat{\varepsilon}_{(j)}) &= E\left(\left\{[\mathbf{I}_n - \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}']\mathbf{y}_{(i)}\right\}' \left\{[\mathbf{I}_n - \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}']\mathbf{y}_{(j)}\right\}\right) \\ &= E\left(\left\{[\mathbf{I}_n - \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}']\boldsymbol{\varepsilon}_{(i)}\right\}' \left\{[\mathbf{I}_n - \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}']\boldsymbol{\varepsilon}_{(j)}\right\}\right) \\ &= E\left(\boldsymbol{\varepsilon}'_{(i)}[\mathbf{I}_n - \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'][\mathbf{I}_n - \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}']\boldsymbol{\varepsilon}_{(j)}\right) \\ &= E\left(\boldsymbol{\varepsilon}'_{(i)}[\mathbf{I}_n - \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}']\boldsymbol{\varepsilon}_{(j)}\right) \\ &= E\left(\text{tr}\{[\mathbf{I}_n - \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}']\boldsymbol{\varepsilon}_{(j)}\boldsymbol{\varepsilon}'_{(i)}\}\right) \\ &= \text{tr}\{[\mathbf{I}_n - \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}']E(\boldsymbol{\varepsilon}_{(j)}\boldsymbol{\varepsilon}'_{(i)})\} \\ &= \sigma_{ij} \text{tr}\{[\mathbf{I}_n - \mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}']\} \\ &= \sigma_{ij}[\text{tr}\{\mathbf{I}_n\} - \text{tr}\{\mathbf{X}'\mathbf{X}(\mathbf{X}'\mathbf{X})^{-1}\}] \\ &= \sigma_{ij}[n - (q + 1)]. \end{aligned}$$

Thus,

$$E(\mathbf{S}_e) = E\left[\frac{1}{n - q - 1}(\mathbf{Y} - \mathbf{X}\hat{\mathbf{B}})'(\mathbf{Y} - \mathbf{X}\hat{\mathbf{B}})\right] = \boldsymbol{\Sigma}.$$

10.4.5 Model Corrected for Means

If the x 's are centered by subtracting their means, we have the centered \mathbf{X} matrix as in (10.13),

$$\mathbf{X}_c = \begin{pmatrix} x_{11} - \bar{x}_1 & x_{12} - \bar{x}_2 & \cdots & x_{1q} - \bar{x}_q \\ x_{21} - \bar{x}_1 & x_{22} - \bar{x}_2 & \cdots & x_{2q} - \bar{x}_q \\ \vdots & \vdots & & \vdots \\ x_{n1} - \bar{x}_1 & x_{n2} - \bar{x}_2 & \cdots & x_{nq} - \bar{x}_q \end{pmatrix}.$$

The \mathbf{B} matrix can be partitioned as

$$\mathbf{B} = \begin{pmatrix} \beta'_0 \\ \mathbf{B}_1 \end{pmatrix} = \begin{pmatrix} \beta_{01} & \beta_{02} & \cdots & \beta_{0p} \\ \beta_{11} & \beta_{12} & \cdots & \beta_{1p} \\ \vdots & \vdots & & \vdots \\ \beta_{q1} & \beta_{q2} & \cdots & \beta_{qp} \end{pmatrix}.$$

By analogy with (10.14) and (10.15), the estimates are

$$\hat{\mathbf{B}}_1 = (\mathbf{X}'_c \mathbf{X}_c)^{-1} \mathbf{X}'_c \mathbf{Y}, \quad (10.55)$$

$$\hat{\beta}'_0 = \bar{\mathbf{y}}' - \bar{\mathbf{x}}' \hat{\mathbf{B}}_1, \quad (10.56)$$

where $\bar{\mathbf{y}}' = (\bar{y}_1, \bar{y}_2, \dots, \bar{y}_p)$ and $\bar{\mathbf{x}}' = (\bar{x}_1, \bar{x}_2, \dots, \bar{x}_q)$. These estimates give the same results as $\hat{\mathbf{B}} = (\mathbf{X}' \mathbf{X})^{-1} \mathbf{X}' \mathbf{Y}$ in (10.48).

As in (10.20), the estimate $\hat{\mathbf{B}}_1$ in (10.55) can be expressed in terms of sample covariance matrices. We multiply and divide (10.55) by $n - 1$ to obtain

$$\begin{aligned} \hat{\mathbf{B}}_1 &= (n - 1)(\mathbf{X}'_c \mathbf{X}_c)^{-1} \frac{\mathbf{X}'_c \mathbf{Y}}{n - 1} = \left(\frac{\mathbf{X}'_c \mathbf{X}_c}{n - 1} \right)^{-1} \frac{\mathbf{X}'_c \mathbf{Y}}{n - 1} \\ &= \mathbf{S}_{xx}^{-1} \mathbf{S}_{xy} \end{aligned} \quad (10.57)$$

where \mathbf{S}_{xx} and \mathbf{S}_{xy} are blocks from the overall sample covariance matrix of the vector $(y_1, y_2, \dots, y_p, x_1, x_2, \dots, x_q)'$:

$$\mathbf{S} = \begin{pmatrix} \mathbf{S}_{yy} & \mathbf{S}_{yx} \\ \mathbf{S}_{xy} & \mathbf{S}_{xx} \end{pmatrix}. \quad (10.58)$$

10.4.6 Estimation in the Seemingly Unrelated Regressions (SUR) Model

In (10.47) we give an alternative version of the multivariate regression model (10.46). The alternative version can also be written

$$\begin{aligned} \text{vec } \mathbf{Y} &= \begin{pmatrix} \mathbf{y}_{(1)} \\ \mathbf{y}_{(2)} \\ \vdots \\ \mathbf{y}_{(p)} \end{pmatrix} = \begin{pmatrix} \mathbf{X} & \mathbf{0} & \cdots & \mathbf{0} \\ \mathbf{0} & \mathbf{X} & \cdots & \mathbf{0} \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{0} & \mathbf{0} & \cdots & \mathbf{X} \end{pmatrix} \begin{pmatrix} \beta_{(1)} \\ \beta_{(2)} \\ \vdots \\ \beta_{(p)} \end{pmatrix} + \begin{pmatrix} \varepsilon_{(1)} \\ \varepsilon_{(2)} \\ \vdots \\ \varepsilon_{(p)} \end{pmatrix} \\ &= \mathbf{X}^* \beta^* + \varepsilon^* \end{aligned}$$

where $\beta^* = \text{vec } \mathbf{B}$. As noted in Section 10.4.2, the estimates of the $\beta_{(i)}$ values are equivalent to those that would be obtained by running a separate regression for each $\mathbf{y}_{(i)}$, without having to adjust the estimates with the covariance matrix Σ . That is,

even though $\text{cov}(\boldsymbol{\varepsilon}^*) = \boldsymbol{\Sigma} \otimes \mathbf{I}$, the ordinary least squares estimate $\hat{\mathbf{B}} = (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}\mathbf{Y}$ is the best linear unbiased estimate (BLUE) for \mathbf{B} .

Suppose, however, that the model for each $\mathbf{y}_{(i)}$ has a potentially unique predictor variable matrix \mathbf{X}_i of size $n \times (q_i + 1)$. In this case, we have a set of p seemingly unrelated regressions, except that the responses $\mathbf{y}_i = (y_{i1}, \dots, y_{ip})'$ are measurements of the same observational unit with covariance matrix $\boldsymbol{\Sigma}$. The *seemingly unrelated regressions* (SUR) model is then:

$$\begin{aligned} \text{vec } \mathbf{Y} &= \begin{pmatrix} \mathbf{y}_{(1)} \\ \mathbf{y}_{(2)} \\ \vdots \\ \mathbf{y}_{(p)} \end{pmatrix} = \begin{pmatrix} \mathbf{X}_1 & \mathbf{0} & \cdots & \mathbf{0} \\ \mathbf{0} & \mathbf{X}_2 & \cdots & \mathbf{0} \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{0} & \mathbf{0} & \cdots & \mathbf{X}_p \end{pmatrix} \begin{pmatrix} \boldsymbol{\beta}_{(1)} \\ \boldsymbol{\beta}_{(2)} \\ \vdots \\ \boldsymbol{\beta}_{(p)} \end{pmatrix} + \begin{pmatrix} \boldsymbol{\varepsilon}_{(1)} \\ \boldsymbol{\varepsilon}_{(2)} \\ \vdots \\ \boldsymbol{\varepsilon}_{(p)} \end{pmatrix} \\ &= \mathbf{X}^{\text{SUR}} \boldsymbol{\beta}^{\text{SUR}} + \boldsymbol{\varepsilon}^*. \end{aligned}$$

In general, the ordinary least squares estimate

$$\tilde{\boldsymbol{\beta}}^{\text{SUR}} = \begin{pmatrix} (\mathbf{X}'_1 \mathbf{X}_1)^{-1} \mathbf{X}_1 \mathbf{y}_{(1)} \\ (\mathbf{X}'_2 \mathbf{X}_2)^{-1} \mathbf{X}_2 \mathbf{y}_{(2)} \\ \vdots \\ (\mathbf{X}'_p \mathbf{X}_p)^{-1} \mathbf{X}_p \mathbf{y}_{(p)} \end{pmatrix} \quad (10.59)$$

is *not* the BLUE for $\boldsymbol{\beta}^{\text{SUR}}$. Following the argument in Section 10.4.3, the generalized least squares approach can be used to obtain the BLUE for $\boldsymbol{\beta}^{\text{SUR}}$:

$$\hat{\boldsymbol{\beta}}^{\text{SUR}} = [\mathbf{X}^{\text{SUR}}' (\boldsymbol{\Sigma} \otimes \mathbf{I})^{-1} \mathbf{X}^{\text{SUR}}]^{-1} \mathbf{X}^{\text{SUR}} (\boldsymbol{\Sigma} \otimes \mathbf{I})^{-1} \text{vec } \mathbf{Y}. \quad (10.60)$$

If $\boldsymbol{\Sigma}$ is not known, an estimate $\hat{\boldsymbol{\Sigma}}$ can be calculated by first obtaining the ordinary least squares estimate for each $\mathbf{y}_{(i)}$:

$$\hat{\boldsymbol{\beta}}_{(i)} = (\mathbf{X}'_i \mathbf{X}_i)^{-1} \mathbf{X}_i \mathbf{y}_{(i)}.$$

Next, the errors associated with each of the vectors $\mathbf{y}_{(i)}$ are estimated using

$$\hat{\boldsymbol{\varepsilon}}_{(i)} = \mathbf{y}_{(i)} - \mathbf{X}_i \hat{\boldsymbol{\beta}}_{(i)}. \quad (10.61)$$

Finally, let the (i, j) element of $\hat{\boldsymbol{\Sigma}}$ equal

$$\hat{\sigma}_{ij} = \frac{1}{\sqrt{(n - q_i)(n - q_j)}} \hat{\boldsymbol{\varepsilon}}'_{(i)} \hat{\boldsymbol{\varepsilon}}_{(j)}. \quad (10.62)$$

Using $\hat{\boldsymbol{\Sigma}}$, the estimate of $\boldsymbol{\beta}$ in (10.60) can be obtained, from which a new estimate of $\hat{\boldsymbol{\Sigma}}$ in (10.62) can be obtained. The calculations associated with equations (10.60) and (10.62) can then be repeated until the estimate $\hat{\boldsymbol{\beta}}^{\text{SUR}}$ is not changing by a meaningful amount. Note that the BLUE in (10.60) reduces to the ordinary least squares estimate in (10.59) if (i) $\mathbf{X}_1 = \mathbf{X}_2 = \dots = \mathbf{X}_p$, or (ii) the off-diagonal elements of $\boldsymbol{\Sigma}$ are all equal to 0 (i.e., $\sigma_{ij} = 0$ for $i \neq j$).

10.5 MULTIVARIATE MULTIPLE REGRESSION: HYPOTHESIS TESTS

In this section we extend the two tests of Section 10.2.5 to the multivariate y case. We assume the x 's are fixed and the y 's are multivariate normal. For other tests and confidence intervals, see Rencher (1998, Chapter 7).

10.5.1 Test of Overall Regression

We first consider the hypothesis that none of the x 's predicts any of the y 's, which can be expressed as $H_0: \mathbf{B}_1 = \mathbf{0}$, where \mathbf{B}_1 includes all rows of \mathbf{B} except the first:

$$\mathbf{B} = \begin{pmatrix} \beta'_0 \\ \mathbf{B}_1 \end{pmatrix} = \begin{pmatrix} \beta_{01} & \beta_{02} & \cdots & \beta_{0p} \\ \hline \beta_{11} & \beta_{12} & \cdots & \beta_{1p} \\ \vdots & \vdots & & \vdots \\ \beta_{q1} & \beta_{q2} & \cdots & \beta_{qp} \end{pmatrix}.$$

We do not wish to include $\beta'_0 = \mathbf{0}'$ in the hypothesis, because this would restrict all y 's to have intercepts of zero. The alternative hypothesis is $H_1: \mathbf{B}_1 \neq \mathbf{0}$, which implies that we want to know if even one $\beta_{jk} \neq 0, j = 1, 2, \dots, q; k = 1, 2, \dots, p$.

The numerator of (10.54) suggests a partitioning of the total sum of squares and products matrix $\mathbf{Y}'\mathbf{Y}$,

$$\mathbf{Y}'\mathbf{Y} = (\mathbf{Y}'\mathbf{Y} - \hat{\mathbf{B}}'\mathbf{X}'\mathbf{Y}) + \hat{\mathbf{B}}'\mathbf{X}'\mathbf{Y}.$$

By analogy to (10.23), we subtract $n\bar{\mathbf{y}}\bar{\mathbf{y}}'$ from both sides to avoid inclusion of $\beta'_0 = \mathbf{0}'$:

$$\begin{aligned} \mathbf{Y}'\mathbf{Y} - n\bar{\mathbf{y}}\bar{\mathbf{y}}' &= (\mathbf{Y}'\mathbf{Y} - \hat{\mathbf{B}}'\mathbf{X}'\mathbf{Y}) + (\hat{\mathbf{B}}'\mathbf{X}'\mathbf{Y} - n\bar{\mathbf{y}}\bar{\mathbf{y}}') \\ &= \mathbf{E} + \mathbf{H}. \end{aligned} \quad (10.63)$$

The overall regression sum of squares and products matrix $\mathbf{H} = \hat{\mathbf{B}}'\mathbf{X}'\mathbf{Y} - n\bar{\mathbf{y}}\bar{\mathbf{y}}'$ can be used to test $H_0: \mathbf{B}_1 = \mathbf{0}$. The notation $\mathbf{E} = \mathbf{Y}'\mathbf{Y} - \hat{\mathbf{B}}'\mathbf{X}'\mathbf{Y}$ and $\mathbf{H} = \hat{\mathbf{B}}'\mathbf{X}'\mathbf{Y} - n\bar{\mathbf{y}}\bar{\mathbf{y}}'$ conforms with usage of \mathbf{E} and \mathbf{H} in Chapter 6.

As in Chapter 6, we can test $H_0: \mathbf{B}_1 = \mathbf{0}$ by means of

$$\Lambda = \frac{|\mathbf{E}|}{|\mathbf{E} + \mathbf{H}|} = \frac{|\mathbf{Y}'\mathbf{Y} - \hat{\mathbf{B}}'\mathbf{X}'\mathbf{Y}|}{|\mathbf{Y}'\mathbf{Y} - n\bar{\mathbf{y}}\bar{\mathbf{y}}'|}, \quad (10.64)$$

which is distributed as $\Lambda_{p,q,n-q-1}$ when $H_0: \mathbf{B}_1 = \mathbf{0}$ is true, where p is the number of y 's and q is the number of x 's. We reject H_0 if $\Lambda \leq \Lambda_{\alpha,p,q,n-q-1}$. The likelihood ratio approach leads to the same test statistic. If \mathbf{H} is "large" because of large values of the β_{jk} 's, then $|\mathbf{E} + \mathbf{H}|$ would be expected to be sufficiently greater than $|\mathbf{E}|$ so that Λ would lead to rejection. By \mathbf{H} "large," we mean that the regression sums of squares on the diagonal are large. Critical values for Λ are available in Table A.9 using $\nu_H = q$ and $\nu_E = n - q - 1$. Note that these degrees of freedom are the same

as in the univariate test for regression of y on x_1, x_2, \dots, x_q in (10.24). The F - and χ^2 -approximations for Λ in (6.15) and (6.16) can also be used, or the exact p -values described in Section 6.1.3 can be used to evaluate the significance of Λ .

There are two alternative expressions for Wilks' Λ in (10.64). We can express Λ in terms of the eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_s$ of $\mathbf{E}^{-1}\mathbf{H}$:

$$\Lambda = \prod_{i=1}^s \frac{1}{1 + \lambda_i}, \quad (10.65)$$

where $s = \min(p, q)$. Wilks' Λ can also be written in the form

$$\Lambda = \frac{|\mathbf{S}|}{|\mathbf{S}_{xx}||\mathbf{S}_{yy}|}, \quad (10.66)$$

where \mathbf{S} is partitioned as in (10.58):

$$\mathbf{S} = \begin{pmatrix} \mathbf{S}_{yy} & \mathbf{S}_{yx} \\ \mathbf{S}_{xy} & \mathbf{S}_{xx} \end{pmatrix}.$$

The form of Λ in (10.66) is the same as in the test for independence of \mathbf{y} and \mathbf{x} given in (7.30), where \mathbf{y} and \mathbf{x} are both random vectors. In the present section, the y 's are random variables and the x 's are fixed. Thus \mathbf{S}_{yy} is the sample covariance matrix of the y 's in the usual sense, while \mathbf{S}_{xx} consists of an analogous mathematical expression involving the constant x 's (see comments about \mathbf{S}_{xx} in Section 10.2.4).

By the symmetry of (10.66) in x and y , Λ is distributed as $\Lambda_{q,p,n-p-1}$ as well as $\Lambda_{p,q,n-q-1}$. This is equivalent to property 3 in Section 6.1.3. Hence, if we regressed the x 's on the y 's, we would get a different $\hat{\mathbf{B}}$ but would have the same value of Λ for the test.

The union–intersection test of $H_0: \mathbf{B}_1 = \mathbf{O}$ uses Roy's test statistic analogous to (6.20),

$$\theta = \frac{\lambda_1}{1 + \lambda_1}, \quad (10.67)$$

where λ_1 is the largest eigenvalue of $\mathbf{E}^{-1}\mathbf{H}$. Upper percentage points θ_α are given in Table A.10. The accompanying parameters are

$$s = \min(p, q), \quad m = \frac{1}{2}(|q - p| - 1), \quad N = \frac{1}{2}(n - q - p - 2).$$

The hypothesis is rejected if $\theta > \theta_\alpha$. The exact p -value for Roy's test can also be obtained using the approach discussed in Section 10.2.4.

As in Section 6.1.5, Pillai's test statistic is defined as

$$V^{(s)} = \sum_{i=1}^s \frac{\lambda_i}{1 + \lambda_i}, \quad (10.68)$$

and the Lawley–Hotelling test statistic is given by

$$U^{(s)} = \sum_{i=1}^s \lambda_i, \quad (10.69)$$

where $\lambda_1, \lambda_2, \dots, \lambda_s$ are the eigenvalues of $\mathbf{E}^{-1}\mathbf{H}$. For $V^{(s)}$, upper percentage points are found in Table A.11, indexed by s , m , and n as defined above in connection with Roy's test. Upper percentage points for $\nu_E U^{(s)} / \nu_H$ (see Section 6.1.5) are provided in Table A.12, where $\nu_H = q$ and $\nu_E = n - q - 1$. Alternatively, we can use the F -approximations or exact tests for $V^{(s)}$ and $U^{(s)}$ in Section 6.1.5.

When H_0 is true, all four test statistics have probability α of rejecting; that is, they all have the same probability of a type I error. When H_0 is false, the power ranking of the tests depends on the configuration of the population eigenvalues, as was noted in Section 6.2. (The sample eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_s$ from $\mathbf{E}^{-1}\mathbf{H}$ are estimates of the population eigenvalues.) If the population eigenvalues are equal or nearly equal, the power ranking of the tests is $V^{(s)} \geq \Lambda \geq U^{(s)} \geq \theta$. If only one population eigenvalue is nonzero, the powers are reversed: $\theta \geq U^{(s)} \geq \Lambda \geq V^{(s)}$.

In the case of a single nonzero population eigenvalue, the rank of \mathbf{B}_1 is 1. There are various ways this could occur; for example, \mathbf{B}_1 could have only one nonzero row, which would indicate that only one of the x 's predicts the y 's. On the other hand, a single nonzero column implies that only one of the y 's is predicted by the x 's. Alternatively, \mathbf{B}_1 would have rank 1 if all rows were equal or linear combinations of each other, manifesting that all x 's act alike in predicting the y 's. Similarly, all columns equal to each other or linear functions of each other would signify only one dimension in the y 's as they relate to the x 's.

■ EXAMPLE 10.5.1

For the chemical data of Table 10.1, we test the overall regression hypothesis $H_0: \mathbf{B}_1 = \mathbf{O}$. The \mathbf{E} and \mathbf{H} matrices are given by

$$\mathbf{E} = \begin{pmatrix} 80.174 & -21.704 & -65.923 \\ -21.704 & 249.462 & -179.496 \\ -65.923 & -179.496 & 231.197 \end{pmatrix},$$

$$\mathbf{H} = \begin{pmatrix} 1707.158 & -492.532 & -996.584 \\ -492.532 & 151.002 & 283.607 \\ -996.584 & 283.607 & 583.688 \end{pmatrix}.$$

The eigenvalues of $\mathbf{E}^{-1}\mathbf{H}$ are 26.3183, .1004, and .0033. The parameters for use in obtaining critical values of the four test statistics are

$$\nu_H = q = 3, \quad \nu_E = n - q - 1 = 19 - 3 - 1 = 15,$$

$$s = \min(3, 3) = 3, \quad m = \frac{1}{2}(|q - p| - 1) = -\frac{1}{2},$$

$$N = \frac{1}{2}(n - q - p - 2) = 5.5.$$

Using the eigenvalues, we obtain the test statistics

$$\begin{aligned}\Lambda &= \prod_{i=1}^3 \frac{1}{1 + \lambda_i} = \frac{1}{1 + 26.3183} \frac{1}{1 + .1004} \frac{1}{1 + .0033} \\ &= .0332 < \Lambda_{.05,3,3,15} = .309, \\ \theta &= \frac{\lambda_1}{1 + \lambda_1} = .963 > \theta(.05, 3, 0, 5) = .669, \\ V^{(s)} &= \sum_{i=1}^3 \frac{\lambda_i}{1 + \lambda_i} = 1.058 > V_{.05,3,0,5}^{(s)} = 1.040, \\ U^{(s)} &= \sum_{i=1}^3 \lambda_i = 26.422, \quad \frac{\nu_E U^{(s)}}{\nu_H} = 132.11,\end{aligned}$$

which exceeds the .05 critical value, 8.427 (interpolated), from Table A.12 (see Section 6.1.5). Thus all four tests reject H_0 . Note that the critical values given for θ and $V^{(s)}$ are conservative, since 0 was used in place of $-.5$ for m .

In this case, the first eigenvalue, 26.3183, completely dominates the other two. In Example 10.4.2, we obtained

$$\hat{\mathbf{B}}_1 = \begin{pmatrix} -1.55 & .40 & .91 \\ -1.42 & .29 & .90 \\ -2.24 & 1.03 & 1.15 \end{pmatrix}.$$

The columns are approximately proportional to each other, indicating that there is essentially only one dimension in the y 's as they are predicted by the x 's. A similar statement can be made about the rows and the dimensionality of the x 's as they predict the y 's. \square

10.5.2 Test on a Subset of the x 's

We consider the hypothesis that the y 's do not depend on the last h of the x 's, $x_{q-h+1}, x_{q-h+2}, \dots, x_q$. By this we mean that none of the p y 's is predicted by any of these h x 's. To express this hypothesis, write the \mathbf{B} matrix in partitioned form,

$$\mathbf{B} = \begin{pmatrix} \mathbf{B}_r \\ \mathbf{B}_d \end{pmatrix},$$

where, as in Section 10.2.5b, the subscript r denotes the subset of β_{jk} 's to be *retained* in the *reduced* model and d represents the subset of β_{jk} 's to be *deleted* if they are not significant predictors of the y 's. Thus \mathbf{B}_d has h rows. The hypothesis can be expressed as

$$H_0: \mathbf{B}_d = \mathbf{O}.$$

If \mathbf{X}_r contains the columns of \mathbf{X} corresponding to \mathbf{B}_r , then the reduced model is

$$\mathbf{Y} = \mathbf{X}_r \mathbf{B}_r + \boldsymbol{\Xi}. \quad (10.70)$$

To compare the fit of the full model and the reduced model, we use the difference between the regression sum of squares and products matrix for the full model, $\hat{\mathbf{B}}'\mathbf{X}'\mathbf{Y}$, and the regression sum of squares and products matrix for the reduced model, $\hat{\mathbf{B}}_r'\mathbf{X}_r'\mathbf{Y}$. By analogy to (10.26), this difference becomes our \mathbf{H} matrix:

$$\mathbf{H} = \hat{\mathbf{B}}'\mathbf{X}'\mathbf{Y} - \hat{\mathbf{B}}_r'\mathbf{X}_r'\mathbf{Y}. \quad (10.71)$$

Thus the test of $H_0: \mathbf{B}_d = \mathbf{O}$ is a full and reduced model test of the significance of $x_{q-h+1}, x_{q-h+2}, \dots, x_q$ above and beyond x_1, x_2, \dots, x_{q-h} .

To make the test, we use the \mathbf{E} matrix based on the full model, $\mathbf{E} = \mathbf{Y}'\mathbf{Y} - \hat{\mathbf{B}}'\mathbf{X}'\mathbf{Y}$. Then

$$\begin{aligned}\mathbf{E} + \mathbf{H} &= (\mathbf{Y}'\mathbf{Y} - \hat{\mathbf{B}}'\mathbf{X}'\mathbf{Y}) + (\hat{\mathbf{B}}'\mathbf{X}'\mathbf{Y} - \hat{\mathbf{B}}_r'\mathbf{X}_r'\mathbf{Y}) \\ &= \mathbf{Y}'\mathbf{Y} - \hat{\mathbf{B}}_r'\mathbf{X}_r'\mathbf{Y},\end{aligned}$$

and Wilks' Λ -statistic is given by

$$\begin{aligned}\Lambda(x_{q-h+1}, \dots, x_q | x_1, \dots, x_{q-h}) &= \frac{|\mathbf{E}|}{|\mathbf{E} + \mathbf{H}|} \\ &= \frac{|\mathbf{Y}'\mathbf{Y} - \hat{\mathbf{B}}'\mathbf{X}'\mathbf{Y}|}{|\mathbf{Y}'\mathbf{Y} - \hat{\mathbf{B}}_r'\mathbf{X}_r'\mathbf{Y}|},\end{aligned} \quad (10.72)$$

which is distributed as $\Lambda_{p,h,n-q-1}$ when $H_0: \mathbf{B}_d = \mathbf{O}$ is true. Critical values are available in Table A.9 with $\nu_H = h$ and $\nu_E = n - q - 1$. Note that these degrees of freedom for the multivariate y case are the same as for the univariate y case (multiple regression) in Section 10.2.5b. The F - and χ^2 -approximations for Λ in (6.15) and (6.16) can also be used.

As implied in the notation $\Lambda(x_{q-h+1}, \dots, x_q | x_1, \dots, x_{q-h})$, Wilks' Λ in (10.72) provides a full and reduced model test. We can express it in terms of Λ for the full model and a similar Λ for the reduced model. In the denominator of (10.72), we have $\mathbf{Y}'\mathbf{Y} - \hat{\mathbf{B}}_r'\mathbf{X}_r'\mathbf{Y}$, which is the error matrix for the reduced model $\mathbf{Y} = \mathbf{X}_r\mathbf{B}_r + \boldsymbol{\Xi}$ in (10.70). This error matrix could be used in a test for the significance of overall regression in the reduced model, as in (10.64),

$$\Lambda_r = \frac{|\mathbf{Y}'\mathbf{Y} - \hat{\mathbf{B}}_r'\mathbf{X}_r'\mathbf{Y}|}{|\mathbf{Y}'\mathbf{Y} - n\bar{\mathbf{y}}\bar{\mathbf{y}}'|}. \quad (10.73)$$

Since Λ_r in (10.73) has the same denominator as Λ in (10.64), we recognize (10.72) as the ratio of Wilks' Λ for the overall regression test in the full model to Wilks' Λ for the overall regression test in the reduced model,

$$\begin{aligned}\Lambda(x_{q-h+1}, \dots, x_q | x_1, \dots, x_{q-h}) &= \frac{|\mathbf{Y}'\mathbf{Y} - \hat{\mathbf{B}}'\mathbf{X}'\mathbf{Y}|}{|\mathbf{Y}'\mathbf{Y} - \hat{\mathbf{B}}_r'\mathbf{X}_r'\mathbf{Y}|} \\ &= \frac{|\mathbf{Y}'\mathbf{Y} - \hat{\mathbf{B}}'\mathbf{X}'\mathbf{Y}|}{|\mathbf{Y}'\mathbf{Y} - n\bar{\mathbf{y}}\bar{\mathbf{y}}'|} \\ &= \frac{|\mathbf{Y}'\mathbf{Y} - \hat{\mathbf{B}}_r'\mathbf{X}_r'\mathbf{Y}|}{|\mathbf{Y}'\mathbf{Y} - n\bar{\mathbf{y}}\bar{\mathbf{y}}'|} \\ &= \frac{\Lambda_f}{\Lambda_r},\end{aligned}\quad (10.74)$$

where Λ_f is given by (10.64). In (10.74), we have a convenient computational device. We run the overall regression test for the full model and again for the reduced model and take the ratio of the resulting Λ values.

The Wilks' Λ in (10.74), comparing the full and reduced models, is similar in appearance to (6.126). However, in (6.126), the full and reduced models involve the dependent variables y_1, y_2, \dots, y_p in MANOVA, whereas in (10.74), the reduced model is obtained by deleting a subset of the independent variables x_1, x_2, \dots, x_q in regression. The parameters of the Wilks' Λ distribution are different in the two cases. Note that in (6.126), some of the dependent variables were denoted by x_1, \dots, x_q for convenience.

Test statistics due to Roy, Pillai, and Lawley–Hotelling can be obtained from the eigenvalues of $\mathbf{E}^{-1}\mathbf{H} = (\mathbf{Y}'\mathbf{Y} - \hat{\mathbf{B}}'\mathbf{X}'\mathbf{Y})^{-1}(\hat{\mathbf{B}}'\mathbf{X}'\mathbf{Y} - \hat{\mathbf{B}}_r'\mathbf{X}_r'\mathbf{Y})$. Critical values or approximate tests for these three test statistics are based on $\nu_H = h$, $\nu_E = n - q - 1$, and

$$s = \min(p, h), \quad m = \frac{1}{2}(|h - p| - 1), \quad N = \frac{1}{2}(n - p - h - 2).$$

■ EXAMPLE 10.5.2

The chemical data in Table 10.1 originated from a response surface experiment seeking to locate optimum operating conditions. Therefore, a second-order model is of interest, and we add $x_1^2, x_2^2, x_3^2, x_1x_2, x_1x_3, x_2x_3$ to the variables x_1, x_2, x_3 considered in Example 10.5.1. There are now $q = 9$ independent variables, and we obtain an overall Wilks' Λ of

$$\Lambda = .00145 < \Lambda_{.05,3,9,9} = .024,$$

where $\nu_H = q = 9$ and $\nu_E = n - q - 1 = 19 - 9 - 1 = 9$. To see whether the six second-order variables add a significant amount to x_1, x_2, x_3 for predicting the y 's, we calculate

$$\Lambda = \frac{\Lambda_f}{\Lambda_r} = \frac{.00145}{.0332} = .0438 < \Lambda_{.05,3,6,9} = .049,$$

where $\nu_H = h = 6$ and $\nu_E = n - q - 1 = 19 - 9 - 1 = 9$. In this case, $\Lambda_r = .0332$ is from Example 10.5.1, in which we considered the model with

x_1 , x_2 , and x_3 . Thus we reject $H_0: \mathbf{B}_d = \mathbf{O}$ and conclude that the second-order terms add significant predictability to the model. \square

10.6 MULTIVARIATE MULTIPLE REGRESSION: PREDICTION

In this section we consider the problem of predicting the response variables \mathbf{y}_0 associated with a known vector of predictor variables \mathbf{x}_0 . Assuming that model (10.46) holds, we first consider a *confidence interval* for the expected value of \mathbf{y}_0 given \mathbf{x}_0 and then we discuss a *prediction interval* for a future observation \mathbf{y}_0 based on \mathbf{x}_0 .

10.6.1 Confidence Interval for $E(\mathbf{y}_0)$

Under model (10.46), the expected value of \mathbf{y}_0 given $\mathbf{x}_0 = (1, x_{01}, x_{02}, \dots, x_{0q})'$ is

$$E(\mathbf{y}_0) = E(\mathbf{B}'\mathbf{x}_0 + \boldsymbol{\varepsilon}_0) = \mathbf{B}'\mathbf{x}_0 + E(\boldsymbol{\varepsilon}_0) = \mathbf{B}'\mathbf{x}_0.$$

To construct a confidence interval for $E(\mathbf{y}_0)$, we first need an estimate of the covariance of the error associated with $\hat{\mathbf{B}}'\mathbf{x}_0$ when estimating $E(\mathbf{y}_0)$. Using (10.52) and properties (2.121) – (2.123) and (3.68),

$$\begin{aligned} \text{cov}(\hat{\mathbf{B}}'\mathbf{x}_0 - \mathbf{B}'\mathbf{x}_0) &= \text{cov}(\hat{\mathbf{B}}'\mathbf{x}_0) \\ &= \text{cov}[(\mathbf{x}'_0 \otimes \mathbf{I}_p) \text{vec } \hat{\mathbf{B}}'] \\ &= (\mathbf{x}'_0 \otimes \mathbf{I}_p)[(\mathbf{X}'\mathbf{X})^{-1} \otimes \Sigma](\mathbf{x}_0 \otimes \mathbf{I}_p) \\ &= [\mathbf{x}'_0(\mathbf{X}'\mathbf{X})^{-1}\mathbf{x}_0] \otimes \Sigma \\ &= [\mathbf{x}'_0(\mathbf{X}'\mathbf{X})^{-1}\mathbf{x}_0]\Sigma, \end{aligned}$$

and a useful estimate of $\text{cov}(\hat{\mathbf{B}}'\mathbf{x}_0)$ is

$$\widehat{\text{cov}}(\hat{\mathbf{B}}'\mathbf{x}_0) = [\mathbf{x}'_0(\mathbf{X}'\mathbf{X})^{-1}\mathbf{x}_0]\mathbf{S}_e, \quad (10.75)$$

with \mathbf{S}_e defined in (10.53).

The estimated covariance matrix in (10.75) can be used to create a p -dimensional confidence region for $E(\mathbf{y}_0)$. More commonly, interest is in individual confidence intervals for the elements of $E(\mathbf{y}_0)$. The (i, i) element of $\widehat{\text{cov}}(\hat{\mathbf{B}}'\mathbf{x}_0)$ can be used to create a $100(1 - \alpha)\%$ *confidence interval* for the i th component of $E(\mathbf{y}_0)$ with

$$\hat{\beta}'_{(i)}\mathbf{x}_0 \pm t_{\alpha/2, n-q-1} \sqrt{s_{ii}[\mathbf{x}'_0(\mathbf{X}'\mathbf{X})^{-1}\mathbf{x}_0]} \quad (10.76)$$

where $\hat{\beta}_{(i)}$ is the i th column of $\hat{\mathbf{B}}$ and s_{ii} is the i th diagonal element of \mathbf{S}_e . If $100(1 - \alpha)\%$ simultaneous confidence intervals for all p components of $E(\mathbf{y}_0)$ are desired without inflating the experimentwise Type I error rate [i.e., the probability of obtaining at least one confidence interval that does not contain $E(y_{0i})$], then the Hotelling's T^2 -based intervals can be used:

$$\hat{\beta}'_{(i)}\mathbf{x}_0 \pm \sqrt{T_{\alpha, p, n-q-1}^2 s_{ii}[\mathbf{x}'_0(\mathbf{X}'\mathbf{X})^{-1}\mathbf{x}_0]}. \quad (10.77)$$

Although the T^2 -intervals obtained using (10.77) are more conservative than the t -intervals of (10.76), the probability that at least one of the p confidence intervals does not contain $E(y_{0i})$ is bounded above by α .

10.6.2 Prediction Interval for a Future Observation \mathbf{y}_0

We are often interested in obtaining a range of reasonable values for a future observation \mathbf{y}_0 associated with \mathbf{x}_0 . Notice that we use the same point estimate for \mathbf{y}_0 that we use for $E(\mathbf{y}_0)$. However, the covariance of the error associated with $\hat{\mathbf{B}}'\mathbf{x}_0$ in estimating \mathbf{y}_0 is different from the error covariance (10.75) that is used when estimating $E(\mathbf{y}_0)$. Using (10.75) and the independence of ε_0 and \mathbf{y}_i , $i = 1, \dots, n$, the covariance of the error associated with $\hat{\mathbf{B}}'\mathbf{x}_0$ when predicting $E(\mathbf{y}_0) = \mathbf{B}'\mathbf{x}_0 + \varepsilon_0$ is

$$\begin{aligned}\text{cov}(\hat{\mathbf{B}}'\mathbf{x}_0 - [\mathbf{B}'\mathbf{x}_0 + \varepsilon_0]) &= \text{cov}(\hat{\mathbf{B}}'\mathbf{x}_0 - \varepsilon_0) \\ &= \text{cov}(\hat{\mathbf{B}}'\mathbf{x}_0) + \text{cov}(\varepsilon_0) \\ &= [\mathbf{x}_0'(\mathbf{X}'\mathbf{X})^{-1}\mathbf{x}_0]\Sigma + \Sigma,\end{aligned}$$

and a useful estimate of $\text{cov}(\hat{\mathbf{B}}'\mathbf{x}_0 - \varepsilon_0)$ is

$$\widehat{\text{cov}}(\hat{\mathbf{B}}'\mathbf{x}_0 - \varepsilon_0) = [1 + \mathbf{x}_0'(\mathbf{X}'\mathbf{X})^{-1}\mathbf{x}_0]\mathbf{S}_e, \quad (10.78)$$

with \mathbf{S}_e defined in (10.53).

The covariance matrix in (10.78) can be used to construct a p -dimensional prediction region, but similar to inference associated with $E(\mathbf{y}_0)$, the more common inferential procedure associated with the predicted observation \mathbf{y}_0 is a prediction interval for an individual element of \mathbf{y}_0 . The (i, i) element of $\widehat{\text{cov}}(\hat{\mathbf{B}}'\mathbf{x}_0 - \varepsilon_0)$ can be used to create a $100(1 - \alpha)\%$ prediction interval for the i th component of \mathbf{y}_0 with

$$\hat{\beta}_{(i)}'\mathbf{x}_0 \pm t_{\alpha/2, n-q-1} \sqrt{s_{ii}[1 + \mathbf{x}_0'(\mathbf{X}'\mathbf{X})^{-1}\mathbf{x}_0]}, \quad (10.79)$$

where $\hat{\beta}_{(i)}$ is the i th column of $\hat{\mathbf{B}}$ and s_{ii} is the i th diagonal element of \mathbf{S}_e . Similar to the simultaneous confidence intervals in (10.77), the $100(1 - \alpha)\%$ simultaneous prediction intervals for all p components of \mathbf{y}_0 are:

$$\hat{\beta}_{(i)}'\mathbf{x}_0 \pm \sqrt{T_{\alpha, p, n-q-1}^2 s_{ii}[1 + \mathbf{x}_0'(\mathbf{X}'\mathbf{X})^{-1}\mathbf{x}_0]}. \quad (10.80)$$

■ EXAMPLE 10.6.2

For the chemical data of Table 10.1, we obtain confidence intervals for the three elements of $E(\mathbf{y}_0)$ and prediction intervals for the three elements of a new observation \mathbf{y}_0 when $\mathbf{x}_0 = (1, 160, 25, 5)$. The value $\mathbf{x}_0'(\mathbf{X}'\mathbf{X})^{-1}\mathbf{x}_0$ is equal to 0.2386, $\hat{\mathbf{B}}'\mathbf{x}_0 = (37.96, 51.19, 10.40)'$, the diagonal elements of \mathbf{S}_e

in (10.53) are $(5.35, 16.63, 15.41)'$, and $t_{\alpha/2, 15} = 2.13$. So, using (10.76), the confidence intervals for $E(y_{01})$, $E(y_{02})$, and $E(y_{03})$ are

$$(35.55, 40.36), \\ (46.94, 55.43), \\ \text{and } (6.32, 14.49).$$

Using (10.79), the prediction intervals for y_{01} , y_{02} , and y_{03} are

$$(32.47, 43.44), \\ (41.51, 60.86), \\ \text{and } (1.09, 19.72).$$

□

10.7 MEASURES OF ASSOCIATION BETWEEN THE y 'S AND THE x 'S

The most widely used measures of association between two sets of variables are the *canonical correlations*, which are treated in Chapter 11. In this section, we review other measures of association that have been proposed.

In (10.34), we have $R^2 = s'_{yx} \mathbf{S}_{xx}^{-1} \mathbf{S}_{xy} / s_{yy}$ for the univariate y case. By analogy, we define an R^2 -like measure of association between y_1, y_2, \dots, y_p and x_1, x_2, \dots, x_q as

$$R_M^2 = \frac{|\mathbf{S}_{yx} \mathbf{S}_{xx}^{-1} \mathbf{S}_{xy}|}{|\mathbf{S}_{yy}|}, \quad (10.81)$$

where \mathbf{S}_{yx} , \mathbf{S}_{xy} , \mathbf{S}_{xx} , and \mathbf{S}_{yy} are defined in (10.58) and the subscript M indicates multivariate y 's.

By analogy to $r_{xy} = s_{xy} / s_x s_y$ in (3.13), Robert and Escoufier (1976) suggested

$$\text{RV} = \frac{\text{tr}(\mathbf{S}_{xy} \mathbf{S}_{yx})}{\sqrt{\text{tr}(\mathbf{S}_{xx}^2) \text{tr}(\mathbf{S}_{yy}^2)}}. \quad (10.82)$$

Kabe (1985) discussed the generalized correlation determinant

$$\text{GCD} = \frac{|\mathbf{L}' \mathbf{S}_{xy} \mathbf{M} \mathbf{M}' \mathbf{S}_{yx} \mathbf{L}|}{|\mathbf{L}' \mathbf{S}_{xx} \mathbf{L}||\mathbf{M}' \mathbf{S}_{yy} \mathbf{M}|}$$

for various choices of the transformation matrices \mathbf{L} and \mathbf{M} .

In Section 6.1.8, we introduced several measures of association that quantify the amount of relationship between the y 's and the dummy grouping variables in a MANOVA context. These are even more appropriate here in the multivariate regression setting, where both the x 's and the y 's are continuous variables. The R^2 -like indices given in (6.40), (6.42), (6.45), (6.48), and (6.50) range between 0 and 1 and

will be briefly reviewed in the remainder of this section. For more complete commentary, see Section 6.1.8.

The two measures based on Wilks' Λ are

$$\eta_{\Lambda}^2 = 1 - \Lambda,$$

$$A_{\Lambda} = 1 - \Lambda^{1/s},$$

where $s = \min(p, q)$. A measure based on Roy's θ is provided by θ itself,

$$\eta_{\theta}^2 = \frac{\lambda_1}{1 + \lambda_1} = \theta,$$

where λ_1 is the largest eigenvalue of $\mathbf{E}^{-1}\mathbf{H}$. This was identified in Section 6.1.8 as the square of the first canonical correlation (see Chapter 11) between the y 's and the grouping variables in MANOVA. In the multivariate regression setting, θ is the square of the first canonical correlation, r_1^2 , between the y 's and the x 's.

Measures of association based on the Lawley-Hotelling and Pillai statistics are given by

$$A_{LH} = \frac{U^{(s)}/s}{1 + U^{(s)}/s},$$

$$A_P = \frac{V^{(s)}}{s}. \quad (10.83)$$

By (6.47) and (6.48), A_P in (10.83) is the average of the s squared canonical correlations, $r_1^2, r_2^2, \dots, r_s^2$.

■ EXAMPLE 10.7

We use the chemical data of Table 10.1 to illustrate some measures of association. For the three dependent variables y_1, y_2 , and y_3 and the three independent variables x_1, x_2 , and x_3 , the partitioned covariance matrix is

$$\mathbf{S} = \begin{pmatrix} \mathbf{S}_{yy} & \mathbf{S}_{yx} \\ \mathbf{S}_{xy} & \mathbf{S}_{xx} \end{pmatrix}$$

$$= \left(\begin{array}{ccc|ccc} 99.30 & -28.57 & -59.03 & -41.95 & -9.49 & -7.37 \\ -28.57 & 22.25 & 5.78 & 11.85 & 1.60 & 3.03 \\ -59.03 & 5.78 & 45.27 & 24.14 & 6.43 & 3.97 \\ \hline -41.95 & 11.85 & 24.14 & 38.67 & -12.17 & -.22 \\ -9.49 & 1.60 & 6.43 & -12.17 & 17.95 & 1.22 \\ -7.37 & 3.03 & 3.97 & -.22 & 1.22 & 2.67 \end{array} \right),$$

from which we obtain R_M^2 and RV directly using (10.81) and (10.82),

$$R_M^2 = \frac{|\mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}|}{|\mathbf{S}_{yy}|} = .00029,$$

$$RV = \frac{\text{tr}(\mathbf{S}_{xy}\mathbf{S}_{yx})}{\sqrt{\text{tr}(\mathbf{S}_{xx}^2)\text{tr}(\mathbf{S}_{yy}^2)}} = .403.$$

Using the results in Example 10.5.1, we obtain

$$\begin{aligned}\eta_{\Lambda}^2 &= 1 - \Lambda = 1 - .0332 = .967, \\ A_{\Lambda} &= 1 - \Lambda^{1/s} = 1 - \Lambda^{1/3} = .679, \\ \eta_{\theta}^2 &= \frac{\lambda_1}{1 + \lambda_1} = .963, \\ A_{\text{LH}} &= \frac{U^{(s)}/s}{1 + U^{(s)}/s} = \frac{26.422/3}{1 + 26.422/3} = .898, \\ A_p &= \frac{V^{(s)}}{s} = \frac{1.058}{3} = .352.\end{aligned}$$

We have general agreement among η_{Λ}^2 , A_{Λ} , η_{θ}^2 , and A_{LH} . But R_M^2 , RV, and A_p do not appear to be measuring the same level of association, especially R_M^2 . It appears that more study is needed before one or more of these measures can be universally recommended. \square

10.8 SUBSET SELECTION

As in the univariate y case in Section 10.2.7, there may be more potential predictor variables (x 's) than are useful in a given situation. Some of the x 's may be redundant in the presence of the other x 's.

We may also be interested in deleting some of the y 's if they are not well predicted by any of the x 's. This would lead to further simplification of the model.

We present two approaches to subset selection: stepwise procedures and methods involving all possible subsets.

10.8.1 Stepwise Procedures

Subset selection among the x 's is discussed in Section 10.8.1a, followed by selection among the y 's in Section 10.8.1b.

10.8.1a Finding a Subset of the x 's

We begin with the *forward selection* procedure based on Wilks' Λ . At the first step, we test the regression of the p y 's on each x_j . There will be two rows in the $\hat{\mathbf{B}}$ matrix, a row containing the intercepts and a row corresponding to x_j :

$$\hat{\mathbf{B}}_j = \begin{pmatrix} \hat{\beta}_{01} & \hat{\beta}_{02} & \cdots & \hat{\beta}_{0p} \\ \hat{\beta}_{j1} & \hat{\beta}_{j2} & \cdots & \hat{\beta}_{jp} \end{pmatrix}.$$

We use the overall regression test statistic (10.64),

$$\Lambda(x_j) = \frac{|\mathbf{Y}'\mathbf{Y} - \hat{\mathbf{B}}_j'\mathbf{X}_j'\mathbf{Y}|}{|\mathbf{Y}'\mathbf{Y} - n\bar{\mathbf{y}}\bar{\mathbf{y}}'|},$$

which is distributed as $\Lambda_{p,1,n-2}$, since $\hat{\mathbf{B}}_j$ has two rows and \mathbf{X}_j has two columns. After calculating $\Lambda(x_j)$ for each j , we choose the variable with minimum $\Lambda(x_j)$. Note that at the first step, we are not testing each variable in the presence of the others. We search for the variable x_j that best predicts the p y 's by itself, not above and beyond the other x 's.

At the second step, we seek the variable yielding the smallest *partial* Λ for each x adjusted for the variable first entered, where the partial Λ -statistic is given by (10.74). After one variable has entered, (10.74) becomes

$$\Lambda(x_j|x_1) = \frac{\Lambda(x_1, x_j)}{\Lambda(x_1)}, \quad (10.84)$$

where x_1 denotes the variable entered at the first step. We calculate (10.84) for each $x_j \neq x_1$ and choose the variable that minimizes $\Lambda(x_j|x_1)$.

If we denote the second variable to enter by x_2 , then at the third step we seek the x_j that minimizes

$$\Lambda(x_j|x_1, x_2) = \frac{\Lambda(x_1, x_2, x_j)}{\Lambda(x_1, x_2)}. \quad (10.85)$$

By property 7 in Section 6.1.3, the partial Wilks' Λ -statistic transforms to an exact F since $\nu_H = h = 1$ at each step.

After m variables have been selected, the partial Λ would have the following form at the next step:

$$\Lambda(x_j|x_1, x_2, \dots, x_m) = \frac{\Lambda(x_1, x_2, \dots, x_m, x_j)}{\Lambda(x_1, x_2, \dots, x_m)}, \quad (10.86)$$

where the first m variables to enter are denoted x_1, x_2, \dots, x_m and x_j is a candidate variable from among the $q - m$ remaining variables. At this step, we would choose the x_j that minimizes (10.86). The partial Wilks' Λ in (10.86) is distributed as $\Lambda_{p,1,n-m-1}$ and, by Table 6.1, transforms to a partial F -statistic distributed as $F_{p,n-m-p}$. [These distributions hold for a variable x_j chosen before seeing the data but not for the x_j that minimizes (10.86) or maximizes the corresponding partial F .]

The procedure continues, bringing in the "best" variable at each step, until a step is reached at which the minimum partial Λ exceeds a predetermined threshold value or, equivalently, the associated partial F falls below a preselected value. Alternatively, the stopping rule can be cast in terms of the p -value of the partial Λ or F . If the smallest p -value at some step exceeds a predetermined value, the procedure stops.

For each x_j , there corresponds an entire row of the $\hat{\mathbf{B}}$ matrix because x_j has a coefficient for each of the p y 's. Thus if a certain x significantly predicts even one of the y 's, it should be retained.

The *stepwise* procedure is an extension of forward selection. Each time a variable enters, all the variables that have entered previously are checked by a partial Λ or F to see if the least "significant" one is now redundant and can be deleted.

The *backward elimination* procedure begins with all x 's (all rows of $\hat{\mathbf{B}}$) included in the model and deletes one at a time, using a partial Λ or F . At the first step, the

partial Λ for each x_j is

$$\Lambda(x_j|x_1, \dots, x_{j-1}, x_{j+1}, \dots, x_q) = \frac{\Lambda(x_1, \dots, x_q)}{\Lambda(x_1, \dots, x_{j-1}, x_{j+1}, \dots, x_q)}, \quad (10.87)$$

which is distributed as $\Lambda_{p,1,n-q-1}$ and can be converted to $F_{p,n-q-p}$ by Table 6.1. The variable with largest Λ or smallest F is deleted. At the second step, a partial Λ or F is calculated for each of the $q - 1$ remaining variables, and again the least important variable in the presence of the others is eliminated. This process continues until a step is reached at which the largest Λ or smallest F is “significant,” indicating that the corresponding variable is apparently not redundant in the presence of its fellows. Some preselected p -value or threshold value of Λ or F is used to determine a stopping rule.

If no automated program is available for subset selection in the multivariate case, a forward selection or backward elimination procedure could be carried out by means of a rather simple set of commands based on (10.86) or (10.87).

A sequential procedure such as stepwise selection will often fail to find the optimum subset, especially if a large pool of predictor variables is involved. However, the value of Wilks' Λ found by stepwise selection may not be far from that for the optimum subset.

The remarks in the final paragraph of Section 10.2.7b are pertinent in the multivariate context as well. True predictors of the y 's in the population may be overlooked because of inflated error variances, or, on the other hand, x 's that are not true predictors may appear to be so in the sample. The latter problem can be severe for small sample sizes (Rencher and Pun 1980).

10.8.1b Finding a Subset of the y 's

After a subset of x 's has been found, the researcher may wish to know if these x 's predict all p of the y 's. If some of the y 's do not relate to any of the x 's, they could be deleted from the model to achieve a further simplification. The y 's can be checked for redundancy in a manner analogous to the stepwise discriminant approach presented in Sections 6.11.2 and 8.9, which finds subsets of *dependent* variables, using a full and reduced model Wilks' Λ for the y 's. The partial Λ -statistic for adding or deleting a y is similar to (10.84), (10.85), or (10.86), except that dependent variables are involved rather than independent variables. Thus to add a y at the third step of a *forward selection procedure*, for example, where the first two variables already entered are denoted as y_1 and y_2 , we use (6.127) to obtain

$$\Lambda(y_j|y_1, y_2) = \frac{\Lambda(y_1, y_2, y_j)}{\Lambda(y_1, y_2)} \quad (10.88)$$

for each $y_j \neq y_1$ or y_2 , and we choose the y_j that minimizes $\Lambda(y_j|y_1, y_2)$. [In (6.127) the dependent variable of interest was denoted by x instead of y_j as here.]

Similarly, if three y 's, designated as y_1 , y_2 , and y_3 , were “in the model” and we were checking the feasibility of adding y_j , the partial Λ -statistic would be

$$\Lambda(y_j|y_1, y_2, y_3) = \frac{\Lambda(y_1, y_2, y_3, y_j)}{\Lambda(y_1, y_2, y_3)}, \quad (10.89)$$

which is distributed as $\Lambda_{1,q,n-q-4}$, where q is the number of x 's *presently* in the model and 4 is the number of y 's *presently* in the model. The two Wilks' Λ values in the numerator and denominator of the right side of (10.89), $\Lambda(y_1, y_2, y_3, y_j)$ and $\Lambda(y_1, y_2, y_3)$, are obtained from (10.64). Since $p = 1$, $\Lambda_{1,q,n-q-4}$ in (10.89) transforms to $F_{q,n-q-4}$ (see Table 6.1).

In the first step of a *backward elimination procedure*, we would delete the y_j that maximizes

$$\Lambda(y_j|y_1, \dots, y_{j-1}, y_{j+1}, \dots, y_p) = \frac{\Lambda(y_1, \dots, y_p)}{\Lambda(y_1, \dots, y_{j-1}, y_{j+1}, \dots, y_p)}, \quad (10.90)$$

which is distributed as $\Lambda_{1,\nu_H,\nu_E-p+1}$. In this case, $\nu_H = q$ and $\nu_E = n - q - 1$ so that the distribution of (10.90) is $\Lambda_{1,q,n-q-p}$, which can be transformed to an exact F . Note that q , the number of x 's, may have been reduced in a subset selection on the x 's as in Section 10.8.1a. Similarly, p is the number of y 's and will decrease in subsequent steps.

Stopping rules for either the forward or backward approach could be defined in terms of p -values or threshold values of Λ or the equivalent F . A *stepwise* procedure could be devised as a modification of the forward approach.

If a software program is available that tests the significance of one x as in (10.87), it can be adapted to test one y as in (10.90) by use of property 3 of Section 6.1.3: The distribution of Λ_{p,ν_H,ν_E} is the same as that of $\Lambda_{\nu_H,p,\nu_E+\nu_H-p}$, which can also be seen from the symmetry of Λ in (10.66),

$$\Lambda = \frac{|\mathbf{S}|}{|\mathbf{S}_{xx}||\mathbf{S}_{yy}|},$$

which is distributed as $\Lambda_{p,q,n-q-1}$ or equivalently as $\Lambda_{q,p,n-p-1}$. Thus we can reverse the y 's and x 's; we list the x 's as dependent variables in the program and the y 's as independent variables. Then the test of a single y in (10.89) or (10.90) can be carried out using (10.87) without any adjustment. The partial Λ -statistic in (10.87) is distributed as $\Lambda_{p,1,n-q-1}$. If we interchange p and q , because the y 's and x 's are interchanged as dependent and independent variables, this becomes $\Lambda_{q,1,n-p-1}$. By property 3 of Section 6.1.3 (repeated above), this is equivalent to $\Lambda_{1,q,n-p-1+1-q} = \Lambda_{1,q,n-p-q}$, which is the distribution of (10.90).

10.8.2 All Possible Subsets

In Section 10.2.7a, we discussed the criteria R_p^2 , s_p^2 , and C_p for comparing all possible subsets of x 's to predict a univariate y in multiple regression, where $p - 1$ denotes the number of x 's in a subset selected from a pool of $k - 1$ available independent variables. We now discuss some matrix analogues of these criteria for the multivariate y case, as suggested by Mallows (1973) and Sparks et al. (1983).

In this section, in order to conform with notation in the literature, we will use p for the number of columns in \mathbf{X} (and the number of rows in \mathbf{B}), rather than for the number of y 's. The number of y 's will be indicated by m .

We now extend the three criteria R_p^2 , s_p^2 , and C_p to analogous matrix expressions \mathbf{R}_p^2 , \mathbf{S}_p , and \mathbf{C}_p . These can be reduced to scalar form using trace or determinant.

1. \mathbf{R}_p^2 . In the univariate y case, R_p^2 for a $(p - 1)$ -variable subset of x 's is defined by (10.32) as

$$R_p^2 = \frac{\hat{\beta}'_p \mathbf{X}'_p \mathbf{y} - n\bar{y}^2}{\mathbf{y}' \mathbf{y} - n\bar{y}^2}.$$

A direct extension of R_p^2 for the multivariate y case is given by the matrix

$$\mathbf{R}_p^2 = (\mathbf{Y}' \mathbf{Y} - n\bar{\mathbf{y}}' \bar{\mathbf{y}})^{-1} (\hat{\mathbf{B}}'_p \mathbf{X}'_p \mathbf{Y} - n\bar{\mathbf{y}}' \bar{\mathbf{y}}'), \quad (10.91)$$

where $p - 1$ is the number of x 's selected from the $k - 1$ available x 's. To convert \mathbf{R}_p^2 to scalar form, we can use $\text{tr}(\mathbf{R}_p^2)/m$, in which we divide by m , the number of y 's, so that $0 \leq \text{tr}(\mathbf{R}_p^2)/m \leq 1$. As in the univariate case, we identify the subset that maximizes $\text{tr}(\mathbf{R}_p^2)/m$ for each value of $p = 2, 3, \dots, k$. The criterion $\text{tr}(\mathbf{R}_p^2)/m$ does not attain its maximum until p reaches k , but we look for the value of p at which further increases are deemed unimportant. We could also use $|\mathbf{R}_p^2|$ in place of $\text{tr}(\mathbf{R}_p^2)/m$.

2. \mathbf{S}_p . A direct extension of the univariate criterion $s_p^2 = \text{MSE}_p = \text{SSE}_p/(n - p)$ is provided by

$$\mathbf{S}_p = \frac{\mathbf{E}_p}{n - p}, \quad (10.92)$$

where $\mathbf{E}_p = \mathbf{Y}' \mathbf{Y} - \hat{\mathbf{B}}'_p \mathbf{X}'_p \mathbf{Y}$. To convert to a scalar, we can use $\text{tr}(\mathbf{S}_p)$ or $|\mathbf{S}_p|$, either of which will behave in an analogous fashion to s_p^2 in the univariate case. The remarks in Section 10.2.7a apply here as well; one may wish to select the subset with minimum value of $\text{tr}(\mathbf{S}_p)$ or perhaps the subset with smallest p such that $\text{tr}(\mathbf{S}_p) < \text{tr}(\mathbf{S}_k)$. A similar application could be made with $|\mathbf{S}_p|$.

3. \mathbf{C}_p . To extend the C_p criterion to the multivariate y case, we write the model under consideration as

$$\mathbf{Y} = \mathbf{X}_p \mathbf{B}_p + \Xi,$$

where $p - 1$ is the number of x 's in the subset and $k - 1$ is the number of x 's in the "full model." The predicted values of the y 's are given by

$$\hat{\mathbf{Y}} = \mathbf{X}_p \hat{\mathbf{B}}_p.$$

We are interested in predicted values of the observation vectors, $\hat{\mathbf{y}}_1, \hat{\mathbf{y}}_2, \dots, \hat{\mathbf{y}}_n$, which are given by the rows of $\hat{\mathbf{Y}}$:

$$\begin{pmatrix} \hat{\mathbf{y}}'_1 \\ \hat{\mathbf{y}}'_2 \\ \vdots \\ \hat{\mathbf{y}}'_n \end{pmatrix} = \begin{pmatrix} \mathbf{x}'_{p1} \\ \mathbf{x}'_{p2} \\ \vdots \\ \mathbf{x}'_{pn} \end{pmatrix} \hat{\mathbf{B}}_p = \begin{pmatrix} \mathbf{x}'_{p1} \hat{\mathbf{B}}_p \\ \mathbf{x}'_{p2} \hat{\mathbf{B}}_p \\ \vdots \\ \mathbf{x}'_{pn} \hat{\mathbf{B}}_p \end{pmatrix}.$$

In general, the predicted vectors $\hat{\mathbf{y}}_i$ are biased estimators of $E(\mathbf{y}_i)$ in the correct model, because we are examining many competing models for which $E(\hat{\mathbf{y}}_i) \neq$

$E(\mathbf{y}_i)$. In place of the univariate expected squared error in (10.37) and (10.38), we define a matrix of expected squares and products of errors, $E[\hat{\mathbf{y}}_i - E(\mathbf{y}_i)][\hat{\mathbf{y}}_i - E(\mathbf{y}_i)]'$. We then add and subtract $E(\hat{\mathbf{y}}_i)$ to obtain (see Problem 10.8)

$$\begin{aligned} & E[\hat{\mathbf{y}}_i - E(\mathbf{y}_i)][\hat{\mathbf{y}}_i - E(\mathbf{y}_i)]' \\ &= E[\hat{\mathbf{y}}_i - E(\hat{\mathbf{y}}_i) + E(\hat{\mathbf{y}}_i) - E(\mathbf{y}_i)][\hat{\mathbf{y}}_i - E(\hat{\mathbf{y}}_i) + E(\hat{\mathbf{y}}_i) - E(\mathbf{y}_i)]' \\ &= E[\hat{\mathbf{y}}_i - E(\hat{\mathbf{y}}_i)][\hat{\mathbf{y}}_i - E(\hat{\mathbf{y}}_i)]' + [E(\hat{\mathbf{y}}_i) - E(\mathbf{y}_i)][E(\hat{\mathbf{y}}_i) - E(\mathbf{y}_i)]' \\ &= \text{cov}(\hat{\mathbf{y}}_i) + (\text{bias in } \hat{\mathbf{y}}_i)(\text{bias in } \hat{\mathbf{y}}_i)'. \end{aligned} \quad (10.93)$$

By analogy to (10.39), our \mathbf{C}_p matrix will be an estimate of the sum of (10.93), multiplied by Σ^{-1} for standardization.

We first find an expression for $\text{cov}(\hat{\mathbf{y}}_i)$, which for convenience we write in row form,

$$\text{cov}(\hat{\mathbf{y}}'_i) = \text{cov}(\mathbf{x}'_{pi}\hat{\mathbf{B}}_p) = \text{cov}(\mathbf{x}'_{pi}\hat{\beta}_{p(1)}, \mathbf{x}'_{pi}\hat{\beta}_{p(2)}, \dots, \mathbf{x}'_{pi}\hat{\beta}_{p(m)}),$$

where $\hat{\mathbf{B}} = (\hat{\beta}_{(1)}, \hat{\beta}_{(2)}, \dots, \hat{\beta}_{(m)})$, as in (10.49). This can be written as

$$\begin{aligned} \text{cov}(\hat{\mathbf{y}}'_i) &= \begin{pmatrix} \sigma_{11}\mathbf{x}'_{pi}(\mathbf{X}'_p\mathbf{X}_p)^{-1}\mathbf{x}_{pi} & \cdots & \sigma_{1m}\mathbf{x}'_{pi}(\mathbf{X}'_p\mathbf{X}_p)^{-1}\mathbf{x}_{pi} \\ \vdots & & \vdots \\ \sigma_{m1}\mathbf{x}'_{pi}(\mathbf{X}'_p\mathbf{X}_p)^{-1}\mathbf{x}_{pi} & \cdots & \sigma_{mm}\mathbf{x}'_{pi}(\mathbf{X}'_p\mathbf{X}_p)^{-1}\mathbf{x}_{pi} \end{pmatrix} \\ &= \mathbf{x}'_{pi}(\mathbf{X}'_p\mathbf{X}_p)^{-1}\mathbf{x}_{pi}\Sigma, \end{aligned} \quad (10.94)$$

where m is the number of y 's and $\Sigma = \text{cov}(\mathbf{y}_i)$ (see Problem 10.9). As in (10.41) (see also Problem 10.5), we can sum over the n observations and use (3.65) to obtain

$$\begin{aligned} \sum_{i=1}^n \text{cov}(\hat{\mathbf{y}}'_i) &= \sum_{i=1}^n \mathbf{x}'_{pi}(\mathbf{X}'_p\mathbf{X}_p)^{-1}\mathbf{x}_{pi}\Sigma \\ &= \Sigma \sum_{i=1}^n \mathbf{x}'_{pi}(\mathbf{X}'_p\mathbf{X}_p)^{-1}\mathbf{x}_{pi} = p\Sigma. \end{aligned} \quad (10.95)$$

To sum the second term on the right of (10.93), we have, by analogy to (10.42),

$$\sum_{i=1}^n (\text{bias in } \hat{\mathbf{y}}_i)(\text{bias in } \hat{\mathbf{y}}_i)' = (n-p)E(\mathbf{S}_p - \Sigma), \quad (10.96)$$

where \mathbf{S}_p is given by (10.92).

Now by (10.95) and (10.96), we can sum (10.93) and multiply by Σ^{-1} to obtain the matrix of total expected squares and products of error standardized by Σ^{-1} ,

$$\begin{aligned} & \Sigma^{-1} \sum_{i=1}^n E[\hat{\mathbf{y}}_i - E(\hat{\mathbf{y}}_i)][\hat{\mathbf{y}}_i - E(\hat{\mathbf{y}}_i)]' \\ &= \Sigma^{-1}[p\Sigma + (n-p)E(\mathbf{S}_p - \Sigma)] \\ &= p\mathbf{I} + (n-p)\Sigma^{-1}E(\mathbf{S}_p - \Sigma). \end{aligned} \quad (10.97)$$

Using $\mathbf{S}_k = \mathbf{E}_k / (n - k)$, the sample covariance matrix based on all $k - 1$ variables, as an estimate of Σ , we can estimate (10.97) by

$$\mathbf{C}_p = p\mathbf{I} + (n - p)\mathbf{S}_k^{-1}(\mathbf{S}_p - \mathbf{S}_k) \quad (10.98)$$

$$= \mathbf{S}_k^{-1}\mathbf{E}_p + (2p - n)\mathbf{I} \quad [\text{by (10.92)}], \quad (10.99)$$

which is the form suggested by Mallows (1973). We can use $\text{tr}(\mathbf{C}_p)$ or $|\mathbf{C}_p|$ to reduce this to a scalar. But if $2p - n$ is negative, $|\mathbf{C}_p|$ can be negative, and Sparks et al. (1983) suggested a modification of $|\mathbf{C}_p|$,

$$|\mathbf{C}_p| = |\mathbf{E}_k^{-1}\mathbf{E}_p|, \quad (10.100)$$

which is always positive.

To find the optimal subset of x 's for each value of p , we could examine all possible subsets [or use a computational scheme such as that of Furnival and Wilson (1974)] and look for the "smallest" \mathbf{C}_p matrix for each p . In (10.97), we see that when the bias is \mathbf{O} , the "population \mathbf{C}_p " is equal to $p\mathbf{I}$. Thus we seek a \mathbf{C}_p that is "small" and near $p\mathbf{I}$. In terms of trace, we seek $\text{tr}(\mathbf{C}_p)$ close to pm , where m is the number of y 's in the vector of measurements; that is, $\text{tr}(\mathbf{I}) = m$.

To find a "target" value for (10.100), we write $\mathbf{E}_k^{-1}\mathbf{E}_p$ in terms of \mathbf{C}_p from (10.99),

$$\mathbf{E}_k^{-1}\mathbf{E}_p = \frac{\mathbf{C}_p + (n - 2p)\mathbf{I}}{n - k}. \quad (10.101)$$

When the bias is \mathbf{O} , we have $\mathbf{C}_p = p\mathbf{I}$, and (10.101) becomes

$$\mathbf{E}_k^{-1}\mathbf{E}_p = \frac{n - p}{n - k}\mathbf{I}, \quad (10.102)$$

whence, by (2.85),

$$|\mathbf{E}_k^{-1}\mathbf{E}_p| = \left(\frac{n - p}{n - k}\right)^m. \quad (10.103)$$

Thus we seek subsets such that

$$\text{tr}(\mathbf{C}_p) \leq pm \quad \text{or} \quad |\mathbf{E}_k^{-1}\mathbf{E}_p| \leq \left(\frac{n - p}{n - k}\right)^m.$$

In summary, when examining all possible subsets, any or all of the following criteria may be useful in finding the single best subset or the best subset for each p :

$$\text{tr}(\mathbf{R}_p^2)/m, \quad |\mathbf{R}_p^2|, \quad \text{tr}(\mathbf{S}_p), \quad |\mathbf{S}_p|, \quad \text{tr}(\mathbf{C}_p), \quad |\mathbf{E}_k^{-1}\mathbf{E}_p|.$$

10.9 MULTIVARIATE REGRESSION: RANDOM x 's

In Sections 10.4 and 10.5, it was assumed that the x 's were fixed and would have the same values in repeated sampling. In many applications, the x 's are random

variables. In such a case, the values of x_1, x_2, \dots, x_q are not under the control of the experimenter but occur randomly along with y_1, y_2, \dots, y_p . On each subject, we observe $p + q$ values in the vector $(y_1, y_2, \dots, y_p, x_1, x_2, \dots, x_q)$.

If we assume that $(y_1, y_2, \dots, y_p, x_1, x_2, \dots, x_q)$ has a multivariate normal distribution, then all estimates and tests have the same formulation as in the fixed- x case [for details, see Rencher (1998, Section 7.7)]. Thus there is no essential difference in our procedures between the fixed- x case and the random- x case.

PROBLEMS

- 10.1** Show that $\sum_{i=1}^n (y_i - \mathbf{x}'_i \hat{\beta})^2 = (\mathbf{y} - \mathbf{X}\hat{\beta})'(\mathbf{y} - \mathbf{X}\hat{\beta})$ as in (10.6).
- 10.2** Show that $\sum_{i=1}^n (y_i - \mu)^2 = \sum_{i=1}^n (y_i - \bar{y})^2 + n(\bar{y} - \mu)^2$ as in (10.10).
- 10.3** Show that $\sum_{i=1}^n (x_{i2} - \bar{x}_2)\bar{y} = 0$ as in (10.19).
- 10.4** Show that $E[\hat{y}_i - E(y_i)]^2 = E[\hat{y}_i - E(\hat{y}_i)]^2 + [E(\hat{y}_i) - E(y_i)]^2$ as in (10.37).
- 10.5** Show that $\sum_{i=1}^n \text{var}(\hat{y}_i)/\sigma^2 = \text{tr}[\mathbf{X}_p(\mathbf{X}'_p \mathbf{X}_p)^{-1} \mathbf{X}'_p]$ as in (10.40).
- 10.6** Show that the alternative form of C_p in (10.45) is equal to the original form in (10.44).
- 10.7** Show that (10.53) is the same as (10.54), that is, $(\mathbf{Y} - \mathbf{X}\hat{\mathbf{B}})'(\mathbf{Y} - \mathbf{X}\hat{\mathbf{B}}) = \mathbf{Y}'\mathbf{Y} - \hat{\mathbf{B}}'\mathbf{X}'\mathbf{Y}$.
- 10.8** Show that
- $$\begin{aligned} & E[\hat{y}_i - E(\mathbf{y}_i)][\hat{y}_i - E(\mathbf{y}_i)]' \\ &= E[\hat{\mathbf{y}}_i - E(\hat{\mathbf{y}}_i)][\hat{\mathbf{y}}_i - E(\hat{\mathbf{y}}_i)]' + [E(\hat{\mathbf{y}}_i) - E(\mathbf{y}_i)][E(\hat{\mathbf{y}}_i) - E(\mathbf{y}_i)]', \end{aligned}$$
- thus verifying (10.93).
- 10.9** Show that $\text{cov}(\hat{\mathbf{y}}'_i)$ has the form given in (10.94).
- 10.10** Show that the two forms of \mathbf{C}_p in (10.98) and (10.99) are equal.
- 10.11** Explain why $|\mathbf{E}_k^{-1} \mathbf{E}_p| > 0$, as claimed following (10.100).
- 10.12** Show that $\mathbf{E}_k^{-1} \mathbf{E}_p = [\mathbf{C}_p + (n - 2p)\mathbf{I}]/(n - k)$, as in (10.101), where \mathbf{C}_p is given in (10.98).
- 10.13** Show that if $\mathbf{C}_p = p\mathbf{I}$, then $\mathbf{E}_k^{-1} \mathbf{E}_p = [(n - p)/(n - k)]\mathbf{I}$ as in (10.102).
- 10.14** One way to show that the ordinary least squares estimate is the best linear unbiased estimator (BLUE) is to verify that the condition of Zyskind (1967) applies. The *Zyskind condition* states that for a model $\mathbf{y} = \mathbf{X}\beta + \varepsilon$ with $E(\varepsilon) = \mathbf{0}$ and $\text{cov}(\varepsilon) = \mathbf{V}$, the ordinary least squares estimate $(\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y}$ is BLUE if and only if there exists a matrix \mathbf{Q} such that $\mathbf{V}\mathbf{X} = \mathbf{X}\mathbf{Q}$.

- (a) Use the Zyskind condition to show that the ordinary least squares estimate in (10.59) is BLUE when $\mathbf{X}_1 = \mathbf{X}_2 = \dots = \mathbf{X}_p$. [Note that this is equivalent to showing that (10.48) is BLUE, because $\mathbf{X}_1 = \mathbf{X}_2 = \dots = \mathbf{X}_p$ implies that model (10.47) holds.]
- (b) Use the Zyskind condition to show that the ordinary least squares estimate (10.59) is BLUE when the off-diagonal elements of Σ are all equal to 0 (i.e., $\sigma_{ij} = 0$ for $i \neq j$).

10.15 Use the diabetes data of Table 3.5.

- (a) Find the least squares estimate $\hat{\mathbf{B}}$ for the regression of (y_1, y_2) on (x_1, x_2, x_3) .
- (b) Test the significance of overall regression using all four test statistics.
- (c) Determine what the eigenvalues of $\mathbf{E}^{-1}\mathbf{H}$ reveal about the essential rank of $\hat{\mathbf{B}}_1$ and the implications of this rank, such as the relative power of the four tests.
- (d) Test the significance of each of x_1, x_2 , and x_3 adjusted for the other two x 's.
- (e) Calculate simultaneous 95% confidence intervals for the elements of $E(y_0)$ given that $\mathbf{x}_0 = (1, 350, 200, 100)'$.
- (f) Calculate simultaneous 95% prediction intervals for the elements of y_0 given that $\mathbf{x}_0 = (1, 350, 200, 100)'$. Compare with part (e).

10.16 Use the sons data of Table 3.8.

- (a) Find the least squares estimate $\hat{\mathbf{B}}$ for the regression of (y_1, y_2) on (x_1, x_2) .
- (b) Test the significance of overall regression using all four test statistics.
- (c) Determine what the eigenvalues of $\mathbf{E}^{-1}\mathbf{H}$ reveal about the essential rank of $\hat{\mathbf{B}}_1$ and the implications of this rank, such as the relative power of the four tests.
- (d) Test the significance of x_1 adjusted for x_2 and of x_2 adjusted for x_1 .
- (e) Calculate simultaneous 95% confidence intervals for the elements of $E(y_0)$ given that $\mathbf{x}_0 = (1, 170, 140)'$.
- (f) Calculate simultaneous 95% prediction intervals for the elements of y_0 given that $\mathbf{x}_0 = (1, 170, 140)'$. Compare with part (e).

10.17 Use the glucose data of Table 3.9.

- (a) Find the least squares estimate $\hat{\mathbf{B}}$ for the regression of (y_1, y_2, y_3) on (x_1, x_2, x_3) .
- (b) Test the significance of overall regression using all four test statistics.

- (c) Determine what the eigenvalues of $\mathbf{E}^{-1}\mathbf{H}$ reveal about the essential rank of $\hat{\mathbf{B}}_1$ and the implications of this rank, such as the relative power of the four tests.
- (d) Test the significance of each of x_1, x_2 , and x_3 adjusted for the other two x 's.
- (e) Test the significance of each y adjusted for the other two by using (10.90).
- (f) Calculate simultaneous 95% confidence intervals for the elements of $E(y_0)$ given that $\mathbf{x}_0 = (1, 100, 100, 100)'$.
- (g) Calculate simultaneous 95% prediction intervals for the elements of \mathbf{y}_0 given that $\mathbf{x}_0 = (1, 100, 100, 100)'$. Compare with part (f).

10.18 Use the Seishu data of Table 7.1.

- (a) Find the least squares estimate $\hat{\mathbf{B}}$ for the regression of (y_1, y_2) on (x_1, x_2, \dots, x_8) and test for significance.
- (b) Test the significance of (x_7, x_8) adjusted for the other x 's.
- (c) Test the significance of (x_4, x_5, x_6) adjusted for the other x 's.
- (d) Test the significance of (x_1, x_2, x_3) adjusted for the other x 's.

10.19 Use the Seishu data of Table 7.1.

- (a) Do a stepwise regression to select a subset of x_1, x_2, \dots, x_8 that adequately predicts (y_1, y_2) .
- (b) After selecting a subset of x 's, use the methods of Section 10.8.1b to check if either of the y 's can be deleted.

10.20 Use the temperature data of Table 7.2.

- (a) Find the least squares estimate $\hat{\mathbf{B}}$ for the regression of (y_4, y_5, y_6) on (y_1, y_2, y_3) and test for significance.
- (b) Find the least squares estimate $\hat{\mathbf{B}}$ for the regression of (y_7, y_8, y_9) on (y_1, \dots, y_6) and test for significance.
- (c) Find the least squares estimate $\hat{\mathbf{B}}$ for the regression of (y_{10}, y_{11}) on (y_1, \dots, y_9) and test for significance.

10.21 Using the temperature data of Table 7.2, carry out a stepwise regression to select a subset of y_1, y_2, \dots, y_9 that adequately predicts (y_{10}, y_{11}) .

CHAPTER 11

CANONICAL CORRELATION

11.1 INTRODUCTION

Canonical correlation analysis is concerned with the amount of (linear) relationship between two sets of variables. We often measure two types of variables on each research unit, for example, a set of aptitude variables and a set of achievement variables, a set of personality variables and a set of ability measures, a set of price indices and a set of production indices, a set of student behaviors and a set of teacher behaviors, a set of psychological attributes and a set of physiological attributes, a set of ecological variables and a set of environmental variables, a set of academic achievement variables and a set of measures of job success, a set of closed-book exam scores and a set of open-book exam scores, and a set of personality variables of freshmen students and the same variables on the same students as seniors.

11.2 CANONICAL CORRELATIONS AND CANONICAL VARIATES

We assume that two sets of variables $\mathbf{y}' = (y_1, y_2, \dots, y_p)$ and $\mathbf{x}' = (x_1, x_2, \dots, x_q)$ are measured on the same sampling unit. We denote the two sets of variables as \mathbf{y}

and \mathbf{x} to conform to notation in Chapters 3, 7, and 10. In Section 7.4.1, we discussed the hypothesis that \mathbf{y} and \mathbf{x} were independent. In this chapter, we consider a measure of overall correlation between \mathbf{y} and \mathbf{x} .

Canonical correlation is an extension of multiple correlation, which is the correlation between one y and several x 's. Canonical correlation analysis is often a useful complement to a multivariate regression analysis.

We first review multiple correlation. The sample covariances and correlations among y, x_1, x_2, \dots, x_q can be summarized in the matrices

$$\mathbf{S} = \begin{pmatrix} s_y^2 & \mathbf{s}'_{yx} \\ \mathbf{s}_{yx} & \mathbf{S}_{xx} \end{pmatrix}, \quad (11.1)$$

$$\mathbf{R} = \begin{pmatrix} 1 & \mathbf{r}'_{yx} \\ \mathbf{r}_{yx} & \mathbf{R}_{xx} \end{pmatrix}, \quad (11.2)$$

where $\mathbf{s}'_{yx} = (s_{y1}, s_{y2}, \dots, s_{yq})$ contains the sample covariances of y with x_1, x_2, \dots, x_q and \mathbf{S}_{xx} is the sample covariance matrix of the x 's [see(10.16)]. The partitioned matrix \mathbf{R} is defined analogously; $\mathbf{r}'_{yx} = (r_{y1}, r_{y2}, \dots, r_{yq})$ contains the sample correlations of y with x_1, x_2, \dots, x_q , and \mathbf{R}_{xx} is the sample correlation matrix of the x 's [see (10.35)].

By (10.34), the squared multiple correlation between y and the x 's can be computed from the partitioned covariance matrix (11.1) or correlation matrix (11.2) as follows:

$$R^2 = \frac{\mathbf{s}'_{yx} \mathbf{S}_{xx}^{-1} \mathbf{s}_{yx}}{s_y^2} = \mathbf{r}'_{yx} \mathbf{R}_{xx}^{-1} \mathbf{r}_{yx}. \quad (11.3)$$

In R^2 , the q covariances between y and the x 's in \mathbf{s}_{yx} or the q correlations between y and the x 's in \mathbf{r}_{yx} are channeled into a single measure of linear relationship between y and the x 's. The multiple correlation R can be defined alternatively as the maximum correlation between y and a linear combination of the x 's; that is, $R = \max_{\mathbf{b}} r_{y, \mathbf{b}' \mathbf{x}}$.

We now return to the case of several y 's and several x 's. The covariance structure associated with two subvectors \mathbf{y} and \mathbf{x} was first discussed in Section 3.9.1. By (3.42), the overall sample covariance matrix for $(y_1, \dots, y_p, x_1, \dots, x_q)$ can be partitioned as

$$\mathbf{S} = \begin{pmatrix} \mathbf{S}_{yy} & \mathbf{S}_{yx} \\ \mathbf{S}_{xy} & \mathbf{S}_{xx} \end{pmatrix},$$

where \mathbf{S}_{yy} is the $p \times p$ sample covariance matrix of the y 's, \mathbf{S}_{yx} is the $p \times q$ matrix of sample covariances between the y 's and the x 's, and \mathbf{S}_{xx} is the $q \times q$ sample covariance matrix of the x 's.

In Section 10.7, we discussed several measures of association between the y 's and the x 's. The first of these is defined in (10.81) as $R_M^2 = |\mathbf{S}_{yx} \mathbf{S}_{xx}^{-1} \mathbf{S}_{xy}| / |\mathbf{S}_{yy}|$, which is analogous to $R^2 = \mathbf{s}'_{yx} \mathbf{S}_{xx}^{-1} \mathbf{s}_{yx} / s_y^2$ in (11.3). By (2.89) and (2.91), R_M^2 can be rewritten as $R_M^2 = |\mathbf{S}_{yy}^{-1} \mathbf{S}_{yx} \mathbf{S}_{xx}^{-1} \mathbf{S}_{xy}|$. By (2.108), R_M^2 can be expressed as

$$R_M^2 = |\mathbf{S}_{yy}^{-1} \mathbf{S}_{yx} \mathbf{S}_{xx}^{-1} \mathbf{S}_{xy}| = \prod_{i=1}^s r_i^2,$$

where $s = \min(p, q)$ and $r_1^2, r_2^2, \dots, r_s^2$ are the eigenvalues of $\mathbf{S}_{yy}^{-1}\mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}$. When written in this form, R_M^2 is seen to be a poor measure of association because $0 \leq r_i^2 \leq 1$ for all i , and the product will usually be too small to meaningfully reflect the amount of association. (In Example 10.7, $R_M^2 = .00029$ was a tiny fraction of the other measures of association.) The eigenvalues themselves, on the other hand, provide meaningful measures of association between the y 's and the x 's. The square roots of the eigenvalues, r_1, r_2, \dots, r_s , are called *canonical correlations*.

The best overall measure of association is the largest squared canonical correlation (maximum eigenvalue) r_1^2 of $\mathbf{S}_{yy}^{-1}\mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}$, but the other eigenvalues (squared canonical correlations) of $\mathbf{S}_{yy}^{-1}\mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}$ provide measures of supplemental dimensions of (linear) relationship between y and x . As an alternative approach, it can be shown that r_1^2 is the maximum squared correlation between a linear combination of the y 's, $u = \mathbf{a}'\mathbf{y}$, and a linear combination of the x 's, $v = \mathbf{b}'\mathbf{x}$; that is,

$$r_1 = \max_{\mathbf{a}, \mathbf{b}} r_{\mathbf{a}'\mathbf{y}, \mathbf{b}'\mathbf{x}}. \quad (11.4)$$

We denote the coefficient vectors that yield the maximum correlation as \mathbf{a}_1 and \mathbf{b}_1 . Thus r_1 (the positive square root of r_1^2) is the correlation between $u_1 = \mathbf{a}_1'\mathbf{y}$ and $v_1 = \mathbf{b}_1'\mathbf{x}$. The coefficient vectors \mathbf{a}_1 and \mathbf{b}_1 can be found as eigenvectors, see (11.7) and (11.8) below. The linear functions u_1 and v_1 are called the first *canonical variates*. There are additional canonical variates $u_i = \mathbf{a}_i'\mathbf{y}$ and $v_i = \mathbf{b}_i'\mathbf{x}$ corresponding to r_2, r_3, \dots, r_s .

It was noted in Section 2.11.5 that the (nonzero) eigenvalues of \mathbf{AB} are the same as those of \mathbf{BA} as long as \mathbf{AB} and \mathbf{BA} are square, but that the eigenvectors of \mathbf{AB} and \mathbf{BA} are not the same. If we let $\mathbf{A} = \mathbf{S}_{yy}^{-1}\mathbf{S}_{yx}$ and $\mathbf{B} = \mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}$, then $r_1^2, r_2^2, \dots, r_s^2$ can also be obtained from $\mathbf{BA} = \mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}\mathbf{S}_{yy}^{-1}\mathbf{S}_{yx}$, as well as from $\mathbf{AB} = \mathbf{S}_{yy}^{-1}\mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}$. Thus the eigenvalues can be obtained from either of the characteristic equations

$$|\mathbf{S}_{yy}^{-1}\mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy} - r^2\mathbf{I}| = 0, \quad (11.5)$$


$$|\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}\mathbf{S}_{yy}^{-1}\mathbf{S}_{yx} - r^2\mathbf{I}| = 0. \quad (11.6)$$

The coefficient vectors \mathbf{a}_i and \mathbf{b}_i in the canonical variates $u_i = \mathbf{a}_i'\mathbf{y}$ and $v_i = \mathbf{b}_i'\mathbf{x}$ are the eigenvectors of these same two matrices:

$$(\mathbf{S}_{yy}^{-1}\mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy} - r^2\mathbf{I})\mathbf{a} = \mathbf{0}, \quad (11.7)$$

$$(\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}\mathbf{S}_{yy}^{-1}\mathbf{S}_{yx} - r^2\mathbf{I})\mathbf{b} = \mathbf{0}. \quad (11.8)$$

Thus the two matrices $\mathbf{S}_{yy}^{-1}\mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}$ and $\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}\mathbf{S}_{yy}^{-1}\mathbf{S}_{yx}$ have the same (nonzero) eigenvalues, as indicated in (11.5) and (11.6), but different eigenvectors, as in (11.7) and (11.8). Since \mathbf{y} is $p \times 1$ and \mathbf{x} is $q \times 1$, the \mathbf{a}_i 's are $p \times 1$ and the \mathbf{b}_i 's are $q \times 1$. This can also be seen in the sizes of the matrices in (11.7) and (11.8); that is, $\mathbf{S}_{yy}^{-1}\mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}$ is $p \times p$ and $\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}\mathbf{S}_{yy}^{-1}\mathbf{S}_{yx}$ is $q \times q$. Since p is typically not equal to q , the matrix that is larger in size will be singular, and the smaller one will be nonsingular. We illustrate for $p < q$. In this case $\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}\mathbf{S}_{yy}^{-1}\mathbf{S}_{yx}$ has the form



With $p < q$, the rank of $\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}\mathbf{S}_{yy}^{-1}\mathbf{S}_{yx}$ is p , because \mathbf{S}_{xx}^{-1} has rank q and $\mathbf{S}_{xy}\mathbf{S}_{yy}^{-1}\mathbf{S}_{yx}$ has rank p . In this case p eigenvalues are nonzero and the remaining $q - p$ eigenvalues are equal to zero. In general, there are $s = \min(p, q)$ values of the squared canonical correlation r_i^2 with s corresponding pairs of canonical variates $u_i = \mathbf{a}'_i\mathbf{y}$ and $v_i = \mathbf{b}'_i\mathbf{x}$. For example, if $p = 3$ and $q = 7$, there will be three canonical correlations, r_1, r_2 , and r_3 .

Thus we have s canonical correlations r_1, r_2, \dots, r_s corresponding to the s pairs of canonical variates u_i and v_i :

$$\begin{array}{lll} r_1 & u_1 = \mathbf{a}'_1\mathbf{y} & v_1 = \mathbf{b}'_1\mathbf{x} \\ r_2 & u_2 = \mathbf{a}'_2\mathbf{y} & v_2 = \mathbf{b}'_2\mathbf{x} \\ \vdots & \vdots & \vdots \\ r_s & u_s = \mathbf{a}'_s\mathbf{y} & v_s = \mathbf{b}'_s\mathbf{x}. \end{array}$$

For each i , r_i is the (sample) correlation between u_i and v_i ; that is, $r_i = r_{u_i v_i}$. The pairs $(u_i, v_i), i = 1, 2, \dots, s$, provide the s dimensions of relationship. For simplicity, we would prefer only one dimension of relationship, but this occurs only when $s = 1$, that is, when $p = 1$ or $q = 1$.

The s dimensions of relationship $(u_i, v_i), i = 1, 2, \dots, s$, are nonredundant. The information each pair provides is unavailable in the other pairs because u_1, u_2, \dots, u_s are *uncorrelated*. They are not orthogonal because $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_s$ are eigenvectors of $\mathbf{S}_{yy}^{-1}\mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}$, which is nonsymmetric. Similarly, each u_i is uncorrelated with all $v_j, j \neq i$, except, of course, v_i .

We examine the elements of the coefficient vectors \mathbf{a}_i and \mathbf{b}_i for the information they provide about the contribution of the y 's and x 's to r_i . These coefficients can be standardized, as noted in the last paragraph in the present section and in Section 11.5.1.

As noted, the matrix $\mathbf{S}_{yy}^{-1}\mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}$ is not symmetric. Many algorithms for computation of eigenvalues and eigenvectors accept only symmetric matrices. Since $\mathbf{S}_{yy}^{-1}\mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}$ is the product of the two symmetric matrices \mathbf{S}_{yy}^{-1} and $\mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}$, we can proceed as in (6.23) and work with $(\mathbf{U}')^{-1}\mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}\mathbf{U}'^{-1}$, where $\mathbf{U}'\mathbf{U} = \mathbf{S}_{yy}$ is the Cholesky factorization of \mathbf{S}_{yy} (see Section 2.7). The symmetric matrix $(\mathbf{U}')^{-1}\mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}\mathbf{U}'^{-1}$ has the same eigenvalues as $\mathbf{S}_{yy}^{-1}\mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}$ but has eigenvectors $\mathbf{U}\mathbf{a}_i$, where \mathbf{a}_i is given in (11.7).

In effect, the pq covariances between the y 's and x 's in \mathbf{S}_{yx} have been replaced by $s = \min(p, q)$ canonical correlations. These succinctly summarize the relationships between \mathbf{y} and \mathbf{x} . In fact, in a typical study, we do not need all s canonical correlations. The smallest eigenvalues can be disregarded to achieve even more simplification. As in (8.13) for discriminant functions, we can judge the importance of each eigenvalue by its relative size:

$$\frac{r_i^2}{\sum_{j=1}^s r_j^2}. \quad (11.9)$$

The canonical correlations can also be obtained from the partitioned correlation matrix of the y 's and x 's,

$$\mathbf{R} = \begin{pmatrix} \mathbf{R}_{yy} & \mathbf{R}_{yx} \\ \mathbf{R}_{xy} & \mathbf{R}_{xx} \end{pmatrix},$$

where \mathbf{R}_{yy} is the $p \times p$ sample correlation matrix of the y 's, \mathbf{R}_{yx} is the $p \times q$ matrix of sample correlations between the y 's and the x 's, and \mathbf{R}_{xx} is the $q \times q$ sample correlation matrix of the x 's. The matrix $\mathbf{R}_{yy}^{-1}\mathbf{R}_{yx}\mathbf{R}_{xx}^{-1}\mathbf{R}_{xy}$ is analogous to $R^2 = \mathbf{r}'_{yx}\mathbf{R}_{xx}^{-1}\mathbf{r}_{yx}$ in the univariate y case. The characteristic equations corresponding to (11.5) and (11.6),

$$|\mathbf{R}_{yy}^{-1}\mathbf{R}_{yx}\mathbf{R}_{xx}^{-1}\mathbf{R}_{xy} - r^2\mathbf{I}| = 0, \quad (11.10)$$

$$|\mathbf{R}_{xx}^{-1}\mathbf{R}_{xy}\mathbf{R}_{yy}^{-1}\mathbf{R}_{yx} - r^2\mathbf{I}| = 0, \quad (11.11)$$

yield the same eigenvalues $r_1^2, r_2^2, \dots, r_s^2$ as (11.5) and (11.6) (the canonical correlations are scale invariant; see property 1 in Section 11.3).

If we use the partitioned correlation matrix in place of the covariance matrix in (11.7) and (11.8), we obtain the same eigenvalues (squared canonical correlations) but different eigenvectors:

$$(\mathbf{R}_{yy}^{-1}\mathbf{R}_{yx}\mathbf{R}_{xx}^{-1}\mathbf{R}_{xy} - r^2\mathbf{I})\mathbf{c} = \mathbf{0}, \quad (11.12)$$

$$(\mathbf{R}_{xx}^{-1}\mathbf{R}_{xy}\mathbf{R}_{yy}^{-1}\mathbf{R}_{yx} - r^2\mathbf{I})\mathbf{d} = \mathbf{0}. \quad (11.13)$$

The relationship between the eigenvectors \mathbf{c} and \mathbf{d} in (11.12) and (11.13) and the eigenvectors \mathbf{a} and \mathbf{b} in (11.7) and (11.8) is

$$\mathbf{c} = \mathbf{D}_y \mathbf{a} \quad \text{and} \quad \mathbf{d} = \mathbf{D}_x \mathbf{b}, \quad (11.14)$$

where $\mathbf{D}_y = \text{diag}(s_{y_1}, s_{y_2}, \dots, s_{y_p})$ and $\mathbf{D}_x = \text{diag}(s_{x_1}, s_{x_2}, \dots, s_{x_q})$.

The eigenvectors \mathbf{c} and \mathbf{d} in (11.12), (11.13), and (11.14) are *standardized coefficient vectors*. By analogy to (8.15), they would be applied to standardized variables. To show this, note that in terms of centered variables $\mathbf{y} - \bar{\mathbf{y}}$, we have

$$\begin{aligned} u &= \mathbf{a}'(\mathbf{y} - \bar{\mathbf{y}}) = \mathbf{a}'\mathbf{D}_y\mathbf{D}_y^{-1}(\mathbf{y} - \bar{\mathbf{y}}) \\ &= \mathbf{c}'\mathbf{D}_y^{-1}(\mathbf{y} - \bar{\mathbf{y}}) \quad [\text{by (11.14)}] \\ &= c_1 \frac{y_1 - \bar{y}_1}{s_{y_1}} + c_2 \frac{y_2 - \bar{y}_2}{s_{y_2}} + \cdots + c_p \frac{y_p - \bar{y}_p}{s_{y_p}}. \end{aligned} \quad (11.15)$$

Hence **c** and **d** are preferred to **a** and **b** for interpretation of the canonical variates u_i and v_i .

11.3 PROPERTIES OF CANONICAL CORRELATIONS

Two interesting properties of canonical correlations are the following [for other properties, see Rencher (1998, Section 8.3)]:

1. Canonical correlations are invariant to changes of scale on either the y 's or the x 's. For example, if the measurement scale is changed from inches to centimeters, the canonical correlations will not change (the corresponding eigenvectors will change). This property holds for simple and multiple correlations as well.
2. The first canonical correlation r_1 is the maximum correlation between linear functions of y and x . Therefore, r_1 exceeds (the absolute value of) the simple correlation between any y and any x or the multiple correlation between any y and all the x 's or between any x and all the y 's.

■ EXAMPLE 11.3

For the chemical data of Table 10.1, we obtain the canonical correlations and illustrate property 2 above. We consider the extended set of nine x 's, as in Example 10.5.2. The matrix \mathbf{R}_{yx} of correlations between the y 's and the x 's is

	x_1	x_2	x_3	x_1x_2	x_1x_3	x_2x_3	x_1^2	x_2^2	x_3^2
y_1	-.68	-.22	-.45	-.41	-.55	-.45	-.68	-.23	-.42
y_2	.40	.08	.39	.16	.44	.33	.40	.12	.33
y_3	.58	.23	.36	.40	.45	.39	.58	.22	.36

The three canonical correlations and their squares are

$$\begin{aligned} r_1 &= .9899 & r_1^2 &= .9800 \\ r_2 &= .9528 & r_2^2 &= .9078 \\ r_3 &= .4625 & r_3^2 &= .2139. \end{aligned}$$

From the relative sizes of the squared canonical correlations, we would consider only the first two to be important. A hypothesis test for the significance of each is carried out in Example 11.4.2.

To confirm that property 2 holds in this case, we compare $r_1 = .9899$ to the individual correlations and the multiple correlations. We first note that .9899 is greater than individual correlations, since (the absolute value of) the largest correlation in \mathbf{R}_{yx} is .68. The multiple correlation $R_{y_j|x}$ of each y_j with the x 's is given by

$$R_{y_1|x} = .987, \quad R_{y_2|x} = .921, \quad R_{y_3|x} = .906,$$

and for the multiple correlation of each x with the y 's we have

$$\begin{array}{lll} R_{x_1|y} = .691, & R_{x_2|y} = .237, & R_{x_3|y} = .507, \\ R_{x_1x_2|y} = .432, & R_{x_1x_3|y} = .585, & R_{x_2x_3|y} = .482, \\ R_{x_1^2|y} = .690, & R_{x_2^2|y} = .234, & R_{x_3^2|y} = .466. \end{array}$$

The first canonical correlation $r_1 = .9899$ exceeds all multiple correlations, and property 2 is satisfied. \square

11.4 TESTS OF SIGNIFICANCE

In the following two sections we discuss basic tests of significance associated with canonical correlations. For other aspects of model validation for canonical correlations and variates, see Rencher (1998, Section 8.5).

11.4.1 Tests of No Relationship Between the y 's and the x 's

In Section 7.4.1, we considered the hypothesis of independence, $H_0: \Sigma_{yx} = \mathbf{O}$. If $\Sigma_{yx} = \mathbf{O}$, the covariance of every y_i with every x_j is zero, and all corresponding correlations are likewise zero. Hence, under H_0 , there is no (linear) relationship between the y 's and the x 's, and H_0 is equivalent to the statement that all canonical correlations r_1, r_2, \dots, r_s are nonsignificant. Furthermore, H_0 is equivalent to the overall regression hypothesis in Section 10.5.1, $H_0: \mathbf{B}_1 = \mathbf{O}$, which also relates all the y 's to all the x 's. Thus by (7.30) or (10.66), the significance of r_1, r_2, \dots, r_s can be tested by

$$\Lambda_1 = \frac{|\mathbf{S}|}{|\mathbf{S}_{yy}||\mathbf{S}_{xx}|} = \frac{|\mathbf{R}|}{|\mathbf{R}_{yy}||\mathbf{R}_{xx}|}, \quad (11.16)$$

which is distributed as $\Lambda_{p,q,n-1-q}$. We reject H_0 if $\Lambda_1 \leq \Lambda_\alpha$. Critical values Λ_α are available in Table A.9 using $\nu_H = q$ and $\nu_E = n - 1 - q$. The statistic Λ_1 in (11.16) is also distributed as $\Lambda_{q,p,n-1-p}$. As in (7.31), Λ_1 is expressible in terms of the squared canonical correlations:

$$\Lambda_1 = \prod_{i=1}^s (1 - r_i^2). \quad (11.17)$$

In this form, we can see that if one or more r_i^2 is large, Λ_1 will be small. We have used the notation Λ_1 in (11.16) and (11.17) because in Section 11.4.2 we will define Λ_2, Λ_3 and so on to test the significance of r_2 and succeeding r_i 's after the first.

If the parameters exceed the range of critical values for Wilks' Λ in Table A.9, we can use the χ^2 -approximation in (6.16),

$$\chi^2 = -[n - \frac{1}{2}(p + q + 3)] \ln \Lambda_1, \quad (11.18)$$

which is approximately distributed as χ^2 with pq degrees of freedom. We reject H_0 if $\chi^2 \geq \chi_{\alpha}^2$. Alternatively, we can use the F -approximation given in (6.15):

$$F = \frac{1 - \Lambda_1^{1/t}}{\Lambda_1^{1/t}} \frac{df_2}{df_1}, \quad (11.19)$$

which has an approximate F -distribution with df_1 and df_2 degrees of freedom, where

$$df_1 = pq, \quad df_2 = wt - \frac{1}{2}pq + 1,$$

$$w = n - \frac{1}{2}(p + q + 3), \quad t = \sqrt{\frac{p^2q^2 - 4}{p^2 + q^2 - 5}}.$$

We reject H_0 if $F > F_{\alpha}$. When $pq = 2$, t is set equal to 1. If $s = \min(p, q)$ is equal to either 1 or 2, then the F -approximation in (11.19) has an exact F -distribution. For example, if one of the two sets consists of only two variables, an exact test is afforded by the F -approximation in (11.19). In contrast, the χ^2 -approximation in (11.18) does not reduce to an exact test for any parameter values.

The other three multivariate test statistics in Sections 6.1.4, 6.1.5, and 10.5.1 can also be used. Pillai's test statistic for the significance of canonical correlations is

$$V^{(s)} = \sum_{i=1}^s r_i^2. \quad (11.20)$$

Upper percentage points of $V^{(s)}$ are found in Table A.11, indexed by

$$s = \min(p, q), \quad m = \frac{1}{2}(|q - p| - 1), \quad N = \frac{1}{2}(n - q - p - 2).$$

For F -approximations for $V^{(s)}$, see Section 6.1.5.

The Lawley–Hotelling statistic for canonical correlations is

$$U^{(s)} = \sum_{i=1}^s \frac{r_i^2}{1 - r_i^2}. \quad (11.21)$$

Upper percentage points for $\nu_E U^{(s)} / \nu_H$ (see Section 6.1.5) are given in Table A.12, which is entered with p , $\nu_H = q$, and $\nu_E = n - q - 1$. For F -approximations, see Section 6.1.5.

Roy's largest root statistic is given by

$$\theta = r_1^2. \quad (11.22)$$

Upper percentage points are found in Table A.10, with s , m , and N defined as above for Pillai's test. An “upper bound” on F for Roy's test is given in (6.21). Even though this “upper bound” is routinely calculated in many software packages, it is not a valid approximation.

As noted at the beginning of this section, the following three tests are equivalent:

1. Test of $H_0: \Sigma_{yx} = \mathbf{O}$, independence of two sets of variables
2. Test of $H_0: \mathbf{B}_1 = \mathbf{O}$, significance of overall multivariate multiple regression
3. Test of significance of the canonical correlations

Even though these tests are equivalent, we have discussed them separately because each has an extension that is different from the others. The respective extensions are

1. Test of independence of three or more sets of variables (Section 7.4.2)
2. Test of full vs. reduced model in multivariate multiple regression (Section 10.5.2)
3. Test of significance of succeeding canonical correlations after the first (Section 11.4.2)

■ EXAMPLE 11.4.1

For the chemical data of Table 10.1, with the extended set of nine x 's, we obtained canonical correlations .9899, .9528, and .4625 in Example 11.3. To test the significance of these, we calculate the following four test statistics and associated approximate F 's.

Statistic	Approximate			p -Value for F
	F	df_1	df_2	
Wilks' $\Lambda = .00145$	6.537	27	21.09	< .0001
Pillai's $V^{(s)} = 2.10$	2.340	27	27	.0155
Lawley-Hotelling $U^{(s)} = 59.03$	12.388	27	17	< .0001
Roy's $\theta = .980$	48.908	9	9	< .0001

The F -approximation for Roy's test is, of course, an "upper bound." Rejection of H_0 in these tests implies that at least r_1^2 is significantly different from zero. The question of how many r_i^2 's are significant is treated in the next section. \square

11.4.2 Test of Significance of Succeeding Canonical Correlations After the First

If the test in (11.17) based on all s canonical correlations rejects H_0 , we are not sure if the canonical correlations beyond the first are significant. To test the significance of r_2, \dots, r_s , we delete r_1^2 from Λ_1 in (11.17) to obtain

$$\Lambda_2 = \prod_{i=2}^s (1 - r_i^2). \quad (11.23)$$

Table 11.1 Tests of Three Canonical Correlations of the Chemical Data

k	Λ_k	$\Lambda_{.05}$	Approximate F	df_1	df_2	p -Value for F
1	.00145	.024	6.537	27	21.1	< .0001
2	.0725	.069	2.714	16	16	.0269
3	.786	.209	.350	7	9	.91

If this test rejects the hypothesis, we conclude that at least r_2 is significantly different from zero. We can continue in this manner, testing each r_i in turn, until a test fails to reject the hypothesis. At the k th step, the test statistic is

$$\Lambda_k = \prod_{i=k}^s (1 - r_i^2), \quad (11.24)$$

which is distributed as $\Lambda_{p-k+1, q-k+1, n-k-q}$ and tests the significance of r_k, r_{k+1}, \dots, r_s . (These test statistics are analogous to those for discriminant functions in Section 8.6.2.) Note that each parameter is reduced by $k - 1$ from the parameter values p, q , and $n - 1 - q$ for Λ_1 in (11.16) or (11.17).

The usual χ^2 - and F -approximations can also be applied to Λ_k . The χ^2 -approximation analogous to (11.18) is given by

$$\chi^2 = -[n - \frac{1}{2}(p + q + 3)] \ln \Lambda_k, \quad (11.25)$$

which has $(p - k + 1)(q - k + 1)$ degrees of freedom. The F -approximation for Λ_k is a simple modification of (11.19) and the accompanying parameter definitions. In place of p, q , and n , we use $p - k + 1, q - k + 1$, and $n - k + 1$ to obtain

$$F = \frac{1 - \Lambda_k^{1/t}}{\Lambda_k^{1/t}} \frac{df_2}{df_1},$$

where

$$\begin{aligned} df_1 &= (p - k + 1)(q - k + 1), \\ df_2 &= wt - \frac{1}{2}[(p - k + 1)(q - k + 1)] + 1, \\ w &= n - \frac{1}{2}(p + q + 3), \\ t &= \sqrt{\frac{(p - k + 1)^2(q - k + 1)^2 - 4}{(p - k + 1)^2 + (q - k + 1)^2 - 5}}. \end{aligned}$$

■ EXAMPLE 11.4.2

We continue our analysis of the canonical correlations for the chemical data in Table 10.1 with three y 's and nine x 's. The tests are summarized in Table 11.1.

In the case of Λ_2 , we have a discrepancy between the exact Wilks' Λ -test and the approximate F -test. The test based on Λ is not significant, while the F -test does reach significance. This illustrates the need to check critical values for exact tests whenever p -values for approximate tests are close to the nominal value of α . From the test using Λ , we conclude that only $r_1 = .9899$ is significant. The relative sizes of the squared canonical correlations, .980, .908, and .214, would indicate two dimensions of relationship, but this is not confirmed by the Wilks' test, perhaps because of the small sample size relative to the number of variables ($p + q = 12$ and $n = 19$).

To illustrate the computations, we obtain the values in Table 11.1 for $k = 2$. Using (11.24), the computation for Λ_2 is

$$\Lambda_2 = \prod_{i=2}^3 (1 - r_i^2) = (1 - .908)(1 - .214) = .0725.$$

With $k = 2$, $p = 3$, $q = 9$, and $n = 19$, the critical value for Λ_2 is obtained from Table A.9 as

$$\Lambda_{.05,p-k+1,q-k+1,n-k-q} = \Lambda_{.05,2,8,8} = .069.$$

For the approximate F for Λ_2 , we have

$$\begin{aligned} t &= \sqrt{\frac{(3 - 2 + 1)^2(9 - 2 + 1)^2 - 4}{(3 - 2 + 1)^2 + (9 - 2 + 1)^2 - 5}} = 2, \\ w &= 19 - \frac{1}{2}(3 + 9 + 3) = 11.5, \\ df_1 &= (3 - 2 + 1)(9 - 2 + 1) = 16, \\ df_2 &= (11.5)(2) - \frac{1}{2}[(3 - 2 + 1)(9 - 2 + 1)] + 1 = 16, \\ F &= \frac{1 - (.0725)^{1/2}}{(0.725)^{1/2}} \frac{16}{16} = 2.714. \end{aligned}$$

□

11.5 INTERPRETATION

We now turn to an assessment of the information contained in the canonical correlations and canonical variates. As was done for discriminant functions in Section 8.7, a distinction can be made between interpretation of the canonical variates and assessing the contribution of each variable. In the former, the signs of the coefficients are considered; in the latter, the signs are ignored and the coefficients are ranked in order of absolute value.

In Sections 11.5.1–11.5.3, we discuss three common tools for interpretation of canonical variates: (1) standardized coefficients, (2) the correlation between each

variable and the canonical variate, and (3) rotation of the canonical variate coefficients. The second of these is the most widely recommended, but we note in Section 11.5.2 that it is the least useful. In fact, for reasons to be outlined, we recommend only the first, standardized coefficients. In Section 11.5.4, we describe redundancy analysis and discuss its shortcomings as a measure of association between two sets of variables.

11.5.1 Standardized Coefficients

The coefficients in the canonical variates $u_i = \mathbf{a}'_i \mathbf{y}$ and $v_i = \mathbf{b}'_i \mathbf{x}$ reflect differences in scaling of the variables as well as differences in contribution of the variables to canonical correlation. To remove the effect of scaling, \mathbf{a}_i and \mathbf{b}_i can be standardized by multiplying by the standard deviations of the corresponding variables as in (11.14):

$$\mathbf{c}_i = \mathbf{D}_y \mathbf{a}_i, \quad \mathbf{d}_i = \mathbf{D}_x \mathbf{b}_i,$$

where $\mathbf{D}_y = \text{diag}(s_{y_1}, s_{y_2}, \dots, s_{y_p})$ and $\mathbf{D}_x = \text{diag}(s_{x_1}, s_{x_2}, \dots, s_{x_q})$. Alternatively, \mathbf{c}_i and \mathbf{d}_i can be obtained directly from (11.12) and (11.13) as eigenvectors of $\mathbf{R}_{yy}^{-1} \mathbf{R}_{yx} \mathbf{R}_{xx}^{-1} \mathbf{R}_{xy}$ and $\mathbf{R}_{xx}^{-1} \mathbf{R}_{xy} \mathbf{R}_{yy}^{-1} \mathbf{R}_{yx}$, respectively. It was noted at the end of Section 11.2 that the coefficients in \mathbf{c}_i are applied to standardized variables [see (11.15)]. Thus the effect of differences in size or scaling of the variables is removed, and the coefficients $c_{i1}, c_{i2}, \dots, c_{ip}$ in \mathbf{c}_i reflect the relative contribution of each of y_1, y_2, \dots, y_p to u_i . A similar statement can be made about \mathbf{d}_i .

The standardized coefficients show the contribution of the variables in the presence of each other. Thus if some of the variables are deleted and others added, the coefficients will change. This is precisely the behavior we desire from the coefficients in a multivariate setting.

■ EXAMPLE 11.5.1

For the chemical data in Table 10.1 with the extended set of nine x 's, we obtain the following standardized coefficients for the three canonical variates:

	\mathbf{c}_1	\mathbf{c}_2	\mathbf{c}_3
y_1	1.5360	4.4704	5.7961
y_2	.2108	2.8291	2.2280
y_3	.4676	3.1309	5.1442
	\mathbf{d}_1	\mathbf{d}_2	\mathbf{d}_3
x_1	5.0125	-38.3053	-12.5072
x_2	5.8551	-17.7390	-24.2290
x_3	1.6500	-7.9699	-32.7392
$x_1 x_2$	-3.9209	19.2937	11.6420
$x_1 x_3$	-2.2968	6.4001	31.2189
$x_2 x_3$.5316	.8096	1.2988
x_1^2	-2.6655	32.7933	4.8454
x_2^2	-1.2346	-3.3641	10.7979
x_3^2	.5703	.8733	.9706

Thus

$$u_1 = 1.54 \frac{y_1 - \bar{y}_1}{s_{y_1}} + .21 \frac{y_2 - \bar{y}_2}{s_{y_2}} + .47 \frac{y_3 - \bar{y}_3}{s_{y_3}},$$

$$v_1 = 5.01 \frac{x_1 - \bar{x}_1}{s_{x_1}} + 5.86 \frac{x_2 - \bar{x}_2}{s_{x_2}} + \dots + .57 \frac{x_3^2 - \bar{x}_3^2}{s_{x_3^2}}.$$

The variables that contribute most to the correlation between u_1 and v_1 are y_1 and $x_1, x_2, x_1x_2, x_1x_3, x_1^2$. The correlation between u_2 and v_2 is due largely to all three y 's and x_1, x_2, x_1x_2, x_1^2 . \square

11.5.2 Correlations between Variables and Canonical Variates

Many writers recommend the additional step of converting the standardized coefficients to correlations. Thus, for example, in $\mathbf{c}'_1 = (c_{11}, c_{12}, \dots, c_{1p})$, instead of the second coefficient c_{12} we could examine $r_{y_2 u_1}$, the correlation between y_2 and the first canonical variate u_1 . Such correlations are sometimes referred to as *loadings* or *structure coefficients*, and it is widely claimed that they provide a more valid interpretation of the canonical variates. Rencher (1988; 1992b; 1998, Section 8.6.3) has shown, however, that a weighted sum of the correlations between y_j and the canonical variates u_1, u_2, \dots, u_s is equal to $R_{y_j | \mathbf{x}}^2$, the squared multiple correlation between y_j and the x 's. There is no information about how the y 's contribute jointly to canonical correlation with the x 's. Therefore, the correlations are useless in gauging the importance of a given variable in the context of the others. The researcher who uses these correlations for interpretation is unknowingly reducing the multivariate setting to a univariate one.

11.5.3 Rotation

In an attempt to improve interpretability, the canonical variate coefficients can be rotated (see Section 13.5) to increase the number of high and low coefficients and reduce the number of intermediate ones.

We do not recommend rotation of the canonical variate coefficients for two reasons [for proof and further discussion, see Rencher (1992b)]:

1. Rotation destroys the optimality of the canonical correlations. For example, the first canonical correlation is reduced and is no longer equal to $\max_{\mathbf{a}, \mathbf{b}} r_{\mathbf{a}' \mathbf{y}, \mathbf{b}' \mathbf{x}}$ as in (11.4).
2. Rotation introduces correlations among succeeding canonical variates. Thus, for example, u_1 and u_2 are correlated after rotation. Hence even though the resulting coefficients may offer a subjectively more interpretable pattern, this gain is offset by the increased complexity due to interrelationships among the canonical variates. For example, u_2 and v_2 no longer offer a new dimension of relationship uncorrelated with u_1 and v_1 . The dimensions now overlap, and some of the information in u_2 and v_2 is already available in u_1 and v_1 .

11.5.4 Redundancy Analysis

The *redundancy* is a measure of association between the y 's and the x 's based on the correlations between variables and canonical variates discussed in Section 11.5.2. Since these correlations provide only univariate information, the redundancy turns out to be a univariate rather than a multivariate measure of relationship. If the squared multiple correlation of y_j regressed on the x 's is denoted by $R_{y_j|x}^2$, then the redundancy of the y 's given the v 's is the average squared multiple correlation:

$$Rd(y|v) = \frac{\sum_{j=1}^p R_{y_j|x}^2}{p}. \quad (11.26)$$

Similarly, the redundancy of the x 's given the u 's is the average

$$Rd(x|u) = \frac{\sum_{j=1}^q R_{x_j|y}^2}{q}, \quad (11.27)$$

where $R_{x_j|y}^2$ is the squared multiple correlation of x_j regressed on the y 's. Since $Rd(y|v)$ in (11.26) is the average squared multiple correlation of each y_j regressed on the x 's, it does not take into account the correlations among the y 's. It is thus an average univariate measure of relationship between the y 's and the x 's, not a multivariate measure at all. The two redundancy measures in (11.26) and (11.27) are not symmetric; that is, $Rd(y|v) \neq Rd(x|u)$.

Thus the so-called redundancy does not really quantify the redundancy among the y 's and x 's and is, therefore, not a useful measure of association between two sets of variables. For a measure of association we recommend r_1^2 itself.

11.6 RELATIONSHIPS OF CANONICAL CORRELATION ANALYSIS TO OTHER MULTIVARIATE TECHNIQUES

In Section 11.4.1, we noted the equivalence of the test for significance of the canonical correlations and the test for significance of overall regression, $H_0: \mathbf{B}_1 = \mathbf{O}$. Additional relationships between canonical correlation and multivariate regression are developed in Section 11.6.1. The relationship of canonical correlation analysis to MANOVA and discriminant analysis is discussed in Section 11.6.2.

11.6.1 Regression

There is a direct link between canonical variate coefficients and multivariate multiple regression coefficients. The matrix of regression coefficients of the y 's regressed on the x 's (corrected for their means) is given in (10.57) as $\hat{\mathbf{B}}_1 = \mathbf{S}_{xx}^{-1} \mathbf{S}_{xy}$. This matrix can be used to relate \mathbf{a}_i and \mathbf{b}_i :

$$\mathbf{b}_i = \hat{\mathbf{B}}_1 \mathbf{a}_i. \quad (11.28)$$

[Since \mathbf{a}_i and \mathbf{b}_i are eigenvectors, (11.28) can be written as $\mathbf{b}_i = c\hat{\mathbf{B}}_1\mathbf{a}_i$, where c is an arbitrary scale factor.] By (2.67) and (11.28), the canonical variate coefficient vector \mathbf{b}_i is expressible as a linear combination of the columns of $\hat{\mathbf{B}}_1$. A similar expression for \mathbf{a}_i can be obtained from the regression of \mathbf{x} on \mathbf{y} : $\mathbf{a}_i = \mathbf{S}_{yy}^{-1}\mathbf{S}_{yx}\mathbf{b}_i$.

In Section 11.2, canonical correlation was defined as an extension of multiple correlation. Correspondingly, canonical correlation reduces to multiple correlation when one of the two sets of variables has only one variable. When $p = 1$, for example, \mathbf{R}_{yy} becomes 1, and by (11.10), the single squared canonical correlation reduces to $r^2 = \mathbf{r}'_{yx}\mathbf{R}_{xx}^{-1}\mathbf{r}_{yx}$, which we recognize from (10.34) as R^2 .

The two Wilks test statistics in multivariate regression in Sections 10.5.1 and 10.5.2, namely, the test for overall regression and the test on a subset of the x 's, can both be expressed in terms of canonical correlations. By (10.64) and (11.17), the test statistic for the overall regression hypothesis $H_0: \mathbf{B}_1 = \mathbf{O}$ can be written as

$$\Lambda_f = \frac{|\mathbf{Y}'\mathbf{Y} - \hat{\mathbf{B}}'\mathbf{X}'\mathbf{Y}|}{|\mathbf{Y}'\mathbf{Y} - n\bar{\mathbf{y}}\bar{\mathbf{y}}'|} \quad (11.29)$$

$$= \prod_{i=1}^s (1 - r_i^2), \quad (11.30)$$

where r_i^2 is the i th squared canonical correlation.

A test statistic for $H_0: \mathbf{B}_d = \mathbf{O}$, the hypothesis that the y 's do not depend on the last h of the x 's, is given by (10.74) as

$$\Lambda(x_{q-h+1}, \dots, x_q | x_1, \dots, x_{q-h}) = \frac{\Lambda_f}{\Lambda_r}, \quad (11.31)$$

where Λ_f is given in (11.29) and Λ_r is given in (10.73) as

$$\Lambda_r = \frac{|\mathbf{Y}'\mathbf{Y} - \hat{\mathbf{B}}'_r\mathbf{X}'_r\mathbf{Y}|}{|\mathbf{Y}'\mathbf{Y} - n\bar{\mathbf{y}}\bar{\mathbf{y}}'|}. \quad (11.32)$$

By analogy with (11.30), Λ_r can be expressed in terms of the squared canonical correlations $c_1^2, c_2^2, \dots, c_t^2$ between y_1, y_2, \dots, y_p and x_1, x_2, \dots, x_{q-h} :

$$\Lambda_r = \prod_{i=1}^t (1 - c_i^2), \quad (11.33)$$

where $t = \min(p, q - h)$. We have used the notation c_i^2 instead of r_i^2 to emphasize that the canonical correlations in the reduced model differ from those in the full model. By (11.30) and (11.33), the full and reduced model test of $H_0: \mathbf{B}_d = \mathbf{O}$ in (11.31) can now be expressed in terms of canonical correlations as

$$\Lambda(x_{q-h+1}, \dots, x_q | x_1, \dots, x_{q-h}) = \frac{\prod_{i=1}^s (1 - r_i^2)}{\prod_{i=1}^t (1 - c_i^2)}. \quad (11.34)$$

If $p = 1$, as in multiple regression, then $s = t = 1$, and (11.34) reduces to

$$\Lambda = \frac{1 - R_f^2}{1 - R_r^2}, \quad (11.35)$$

where R_f^2 and R_r^2 are the squared multiple correlations for the full model and for the reduced model. The distribution of Λ in (11.35) is $\Lambda_{1,h,n-q-1}$ when H_0 is true. Since $p = 1$, there is an exact F -transformation from Table 6.1,

$$F = \frac{(1 - \Lambda)(n - q - 1)}{\Lambda h},$$

which is distributed as $F_{h,n-q-1}$ when H_0 is true. Substitution of $\Lambda = (1 - R_f^2)/(1 - R_r^2)$ from (11.35) yields the F -statistic expressed in terms of R^2 ,

$$F = \frac{(R_f^2 - R_r^2)(n - q - 1)}{(1 - R_f^2)h}, \quad (11.36)$$

as given in (10.33).

Subset selection in canonical correlation analysis can be handled by the methods for multivariate regression given in Section 10.8. A subset of x 's can be found by the procedure of Section 10.8.1a. After a subset of x 's is found, the approach in Section 10.8.1b can be used to select a subset of y 's.

Muller (1982) discussed the relationship of canonical correlation analysis to multivariate regression and principal components. (Principal components are treated in Chapter 12.)

11.6.2 MANOVA and Discriminant Analysis

In Sections 6.1.8 and 8.4.2, it was noted that in a one-way MANOVA or discriminant analysis setting, $\lambda_i/(1 + \lambda_i)$ is equal to r_i^2 , where λ_i is the i th eigenvalue of $\mathbf{E}^{-1}\mathbf{H}$ and r_i^2 is the i th squared canonical correlation between the p dependent variables and the $k - 1$ grouping variables. We now give a justification of this assertion.

Let the dependent variables be denoted by y_1, y_2, \dots, y_p as usual. We represent the k groups by $k - 1$ dummy variables, x_1, x_2, \dots, x_{k-1} , defined for each member of the i th group, $i \leq k - 1$, as $x_1 = 0, \dots, x_{i-1} = 0, x_i = 1, x_{i+1} = 0, \dots, x_{k-1} = 0$. For the k th group, all x 's are zero. (See Section 6.1.8 for an introduction to dummy variables.) To illustrate with $k = 4$, the x 's are defined as follows in each group:

Group	x_1	x_2	x_3
1	1	0	0
2	0	1	0
3	0	0	1
4	0	0	0

The MANOVA model is equivalent to multivariate regression of y_1, y_2, \dots, y_p on the dummy grouping variables x_1, x_2, \dots, x_{k-1} . The MANOVA test of $H_0: \boldsymbol{\mu}_1 =$

$\mu_2 = \dots = \mu_k$ is equivalent to the multivariate regression test of $H_0: \mathbf{B}_1 = \mathbf{O}$, as given by (11.17),

$$\Lambda = \prod_{i=1}^s (1 - r_i^2). \quad (11.37)$$

When we compare this form of Λ to the MANOVA test statistic (6.14),

$$\Lambda = \prod_{i=1}^s \frac{1}{1 + \lambda_i}, \quad (11.38)$$

we obtain the relationships

$$r_i^2 = \frac{\lambda_i}{1 + \lambda_i}, \quad (11.39)$$

$$\lambda_i = \frac{r_i^2}{1 - r_i^2}. \quad (11.40)$$

To establish this relationship more formally, we write (6.22) as

$$\mathbf{H}\mathbf{a} = \lambda \mathbf{E}\mathbf{a}, \quad (11.41)$$

and (11.7) as

$$\mathbf{S}_{yy}^{-1} \mathbf{S}_{yx} \mathbf{S}_{xx}^{-1} \mathbf{S}_{xy} \mathbf{a} = r^2 \mathbf{a}. \quad (11.42)$$

We multiply (11.42) on the left by \mathbf{S}_{yy} to obtain

$$\mathbf{S}_{yx} \mathbf{S}_{xx}^{-1} \mathbf{S}_{xy} \mathbf{a} = r^2 \mathbf{S}_{yy} \mathbf{a}. \quad (11.43)$$

Using the centered matrix \mathbf{X}_c in (10.14), with an analogous definition for \mathbf{Y}_c , we can write \mathbf{B}_1 in the form [see (10.57)]

$$\hat{\mathbf{B}}_1 = \left(\frac{\mathbf{X}'_c \mathbf{X}_c}{n-1} \right)^{-1} \frac{\mathbf{X}'_c \mathbf{Y}_c}{n-1} = \mathbf{S}_{xx}^{-1} \mathbf{S}_{xy}.$$

In terms of centered matrices, $\mathbf{E} = \mathbf{Y}'\mathbf{Y} - \hat{\mathbf{B}}'\mathbf{X}'\mathbf{Y}$ in (10.54) can be written as

$$\begin{aligned} \frac{\mathbf{E}}{n-1} &= \frac{\mathbf{Y}'_c \mathbf{Y}_c}{n-1} - \hat{\mathbf{B}}'_1 \frac{\mathbf{X}'_c \mathbf{Y}_c}{n-1} \\ &= \mathbf{S}_{yy} - \mathbf{S}'_{xy} \mathbf{S}_{xx}^{-1} \mathbf{S}_{xy} = \mathbf{S}_{yy} - \mathbf{S}_{yx} \mathbf{S}_{xx}^{-1} \mathbf{S}_{xy}, \end{aligned} \quad (11.44)$$

since $\mathbf{S}'_{xy} = \mathbf{S}_{yx}$. Similarly,

$$\frac{\mathbf{H}}{n-1} = \frac{\hat{\mathbf{B}}'_1 \mathbf{X}'_c \mathbf{Y}_c}{n-1} = \mathbf{S}_{yx} \mathbf{S}_{xx}^{-1} \mathbf{S}_{xy}. \quad (11.45)$$

Since MANOVA is equivalent to multivariate regression on dummy grouping variables, we can substitute these values of \mathbf{E} and \mathbf{H} into (11.41) to obtain

$$\mathbf{S}_{yx} \mathbf{S}_{xx}^{-1} \mathbf{S}_{xy} \mathbf{a} = \lambda (\mathbf{S}_{yy} - \mathbf{S}_{yx} \mathbf{S}_{xx}^{-1} \mathbf{S}_{xy}) \mathbf{a}. \quad (11.46)$$

Subtracting $r^2 \mathbf{S}_{yx} \mathbf{S}_{xx}^{-1} \mathbf{S}_{xy} \mathbf{a}$ from both sides of (11.43) gives

$$\mathbf{S}_{yx} \mathbf{S}_{xx}^{-1} \mathbf{S}_{xy} \mathbf{a} = \frac{r^2}{1 - r^2} (\mathbf{S}_{yy} - \mathbf{S}_{yx} \mathbf{S}_{xx}^{-1} \mathbf{S}_{xy}) \mathbf{a}. \quad (11.47)$$

A comparison of (11.46) and (11.47) shows that

$$\lambda = \frac{r^2}{1 - r^2},$$

as in (11.40). Lindsey et al. (1985) discussed some advantages of using canonical correlation analysis in place of discriminant analysis in the several-group case.

PROBLEMS

- 11.1** Show that the expression for canonical correlations in (11.12) can be obtained from the analogous expression in terms of variances and covariances in (11.7).
- 11.2** Verify (11.28), $\mathbf{b}_i = \hat{\mathbf{B}}_1 \mathbf{a}_i$.
- 11.3** Verify (11.35) for Λ when $p = s = t = 1$.
- 11.4** Verify the expression in (11.36) for F in terms of R_f^2 and R_r^2 .
- 11.5** Solve (11.39), $r_i^2 = \lambda_i / (1 + \lambda_i)$, for λ_i to obtain (11.40).
- 11.6** Verify (11.46), $\mathbf{S}_{yx} \mathbf{S}_{xx}^{-1} \mathbf{S}_{xy} \mathbf{a} = \lambda (\mathbf{S}_{yy} - \mathbf{S}_{yx} \mathbf{S}_{xx}^{-1} \mathbf{S}_{xy}) \mathbf{a}$.
- 11.7** Show that (11.47) can be obtained by subtracting $r^2 \mathbf{S}_{yx} \mathbf{S}_{xx}^{-1} \mathbf{S}_{xy} \mathbf{a}$ from both sides of (11.43).
- 11.8** Use the diabetes data of Table 3.5.
 - Find the canonical correlations between (y_1, y_2) and (x_1, x_2, x_3) .
 - Find the standardized coefficients for the canonical variates.
 - Test the significance of each canonical correlation.
- 11.9** Use the sons data of Table 3.8.
 - Find the canonical correlations between (y_1, y_2) and (x_1, x_2) .
 - Find the standardized coefficients for the canonical variates.
 - Test the significance of each canonical correlation.
- 11.10** Use the glucose data of Table 3.9.
 - Find the canonical correlations between (y_1, y_2, y_3) and (x_1, x_2, x_3) .
 - Find the standardized coefficients for the canonical variates.
 - Test the significance of each canonical correlation.

11.11 Use the Seishu data of Table 7.1.

- (a) Find the canonical correlations between (y_1, y_2) and (x_1, x_2, \dots, x_8) .
- (b) Find the standardized coefficients for the canonical variates.
- (c) Test the significance of each canonical correlation.

11.12 Use canonical correlation to carry out the tests in parts (b), (c), and (d) of Problem 10.17, using the Seishu data. You will need to find the canonical correlations between (y_1, y_2) and the x 's in the indicated reduced models and use (11.34).

11.13 Using the temperature data of Table 7.2, find the canonical correlations and the standardized coefficients and carry out significance tests for the following:

- (a) (y_1, y_2, y_3) and (y_4, y_5, y_6)
- (b) (y_1, y_2, \dots, y_6) and (y_7, y_8, y_9)
- (c) (y_1, y_2, \dots, y_9) and (y_{10}, y_{11})
- (d) (y_1, y_2, \dots, y_6) and $(y_7, y_8, \dots, y_{11})$.

CHAPTER 12

PRINCIPAL COMPONENT ANALYSIS

12.1 INTRODUCTION

In principal component analysis, we seek to maximize the variance of a linear combination of the variables. For example, we might want to rank students on the basis of their scores on achievement tests in English, mathematics, reading, and so on. An average score would provide a single scale on which to compare the students, but with unequal weights we can spread the students out further on the scale and obtain a better ranking.

Essentially, principal component analysis is a one-sample technique applied to data with no groupings among the observations as in Chapters 8 and 9 and no partitioning of the variables into subsets y and x as in Chapters 10 and 11. All the linear combinations that we have considered previously were related to other variables or to the data structure. In regression, we have linear combinations of the independent variables that best predict the dependent variable(s); in canonical correlation, we have linear combinations of a subset of variables that maximally correlate with linear combinations of another subset of variables; and discriminant analysis involves linear combinations that maximally separate groups of observations. Principal components, on the other hand, are concerned only with the core structure of a single

sample of observations on p variables. None of the variables is designated as dependent, and no grouping of observations is assumed. [For a discussion of the use of principal components with data consisting of several samples or groups, see Rencher (1998, Section 9.9)].

The first principal component is the linear combination with maximal variance; we are essentially searching for a dimension along which the observations are maximally separated or spread out. The second principal component is the linear combination with maximal variance in a direction orthogonal to the first principal component, and so on. In general, the principal components define dimensions that are different from those defined by discriminant functions or canonical variates.

In some applications, the principal components are an end in themselves and may be amenable to interpretation. More often they are obtained for use as input to another analysis. For example, two situations in regression where principal components may be useful are (1) if the number of independent variables is large relative to the number of observations, a test may be ineffective or even impossible; and (2) if the independent variables are highly correlated, the estimates of regression coefficients may be unstable. In such cases, the independent variables can be reduced to a smaller number of principal components that will yield a better test or more stable estimates of the regression coefficients. For details of this application, see Rencher (1998, Section 9.8).

As another illustration, suppose that in a MANOVA application p is close to ν_E , so that a test has low power, or that $p > \nu_E$, in which case we have so many dependent variables that a test cannot be made. In such cases, we can replace the dependent variables with a smaller set of principal components and then carry out the test.

In these illustrations, principal components are used to reduce the number of dimensions. A useful dimension reduction device is to evaluate the first two principal components for each observation vector and construct a scatterplot to check for multivariate normality, outliers, and so on.

Finally, we note that in the term *principal components* we use the adjective *principal*, describing what kind of components—main, primary, fundamental, major, and so on. We do not use the noun *principle* as a modifier for *components*.

12.2 GEOMETRIC AND ALGEBRAIC BASES OF PRINCIPAL COMPONENTS

12.2.1 Geometric Approach

As noted in Section 12.1, principal components analysis deals with a single sample of n observation vectors y_1, y_2, \dots, y_n that form a swarm of points in a p -dimensional space. Principal component analysis can be applied to any distribution of y , but it will be easier to visualize geometrically if the swarm of points is ellipsoidal.

If the variables y_1, y_2, \dots, y_p in y are correlated, the ellipsoidal swarm of points is not oriented parallel to any of the axes represented by y_1, y_2, \dots, y_p . We wish to find the natural axes of the swarm of points (the axes of the ellipsoid) with origin at

$\bar{\mathbf{y}}$, the mean vector of $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n$. This is done by translating the origin to $\bar{\mathbf{y}}$ and then rotating the axes. After rotation so that the axes become the natural axes of the ellipsoid, the new variables (principal components) will be uncorrelated.

We could indicate the translation of the origin to $\bar{\mathbf{y}}$ by writing $\mathbf{y}_i - \bar{\mathbf{y}}$ but will not usually do so for economy of notation. We will write $\mathbf{y}_i - \bar{\mathbf{y}}$ when there is an explicit need; otherwise we assume that \mathbf{y}_i has been centered.

The axes can be rotated by multiplying each \mathbf{y}_i by an orthogonal matrix \mathbf{A} [see (2.101), where the orthogonal matrix was denoted by \mathbf{C}]:

$$\mathbf{z}_i = \mathbf{Ay}_i. \quad (12.1)$$

Since \mathbf{A} is orthogonal, $\mathbf{A}'\mathbf{A} = \mathbf{I}$, and the distance to the origin is unchanged:

$$\mathbf{z}_i' \mathbf{z}_i = (\mathbf{Ay}_i)'(\mathbf{Ay}_i) = \mathbf{y}_i' \mathbf{A}' \mathbf{A} \mathbf{y}_i = \mathbf{y}_i' \mathbf{y}_i$$

[see (2.103)]. Thus an orthogonal matrix transforms \mathbf{y}_i to a point \mathbf{z}_i that is the same distance from the origin, and the axes are effectively rotated.

Finding the axes of the ellipsoid is equivalent to finding the orthogonal matrix \mathbf{A} that rotates the axes to line up with the natural extensions of the swarm of points so that the new variables (principal components) z_1, z_2, \dots, z_p in $\mathbf{z} = \mathbf{Ay}$ are uncorrelated. Thus we want the sample covariance matrix of \mathbf{z} , $\mathbf{S}_z = \mathbf{ASA}'$ [see (3.64)], to be diagonal,

$$\mathbf{S}_z = \mathbf{ASA}' = \begin{pmatrix} s_{z_1}^2 & 0 & \cdots & 0 \\ 0 & s_{z_2}^2 & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & s_{z_p}^2 \end{pmatrix}, \quad (12.2)$$

where \mathbf{S} is the sample covariance matrix of $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n$. By (2.111), $\mathbf{C}'\mathbf{SC} = \mathbf{D} = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_p)$, where the λ_i 's are eigenvalues of \mathbf{S} and \mathbf{C} is an orthogonal matrix whose columns are normalized eigenvectors of \mathbf{S} . Thus the orthogonal matrix \mathbf{A} that diagonalizes \mathbf{S} is the transpose of the matrix \mathbf{C} :

$$\mathbf{A} = \mathbf{C}' = \begin{pmatrix} \mathbf{a}'_1 \\ \mathbf{a}'_2 \\ \vdots \\ \mathbf{a}'_p \end{pmatrix}, \quad (12.3)$$

where \mathbf{a}_i is the i th normalized ($\mathbf{a}'_i \mathbf{a}_i = 1$) eigenvector of \mathbf{S} . The *principal components* are the transformed variables $z_1 = \mathbf{a}'_1 \mathbf{y}$, $z_2 = \mathbf{a}'_2 \mathbf{y}, \dots, z_p = \mathbf{a}'_p \mathbf{y}$ in $\mathbf{z} = \mathbf{Ay}$. For example, $z_1 = a_{11}y_1 + a_{12}y_2 + \cdots + a_{1p}y_p$.

By (2.111), the diagonal elements of \mathbf{ASA}' on the right side of (12.2) are eigenvalues of \mathbf{S} . Hence the eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_p$ of \mathbf{S} are the (sample) variances of the principal components $z_i = \mathbf{a}'_i \mathbf{y}$:

$$s_{z_i}^2 = \lambda_i. \quad (12.4)$$

Since the rotation lines up with the natural extensions of the swarm of points, $z_1 = \mathbf{a}'_1 \mathbf{y}$ has the largest (sample) variance and $z_p = \mathbf{a}'_p \mathbf{y}$ has the smallest variance. This also follows from (12.4), because the variance of z_1 is λ_1 , the largest eigenvalue, and the variance of z_p is λ_p , the smallest eigenvalue. If some of the eigenvalues are small, we can neglect them and represent the points fairly well with fewer than p dimensions. For example, if $p = 3$ and λ_3 is small, then the swarm of points is an “elliptical pancake” and a two-dimensional representation will adequately portray the configuration of points.

Because the eigenvalues are variances of the principal components, we can speak of “the proportion of variance explained” by the first k components:

$$\begin{aligned}\text{Proportion of variance} &= \frac{\lambda_1 + \lambda_2 + \cdots + \lambda_k}{\lambda_1 + \lambda_2 + \cdots + \lambda_p} \\ &= \frac{\lambda_1 + \lambda_2 + \cdots + \lambda_k}{\sum_{j=1}^p s_{jj}},\end{aligned}\quad (12.5)$$

since $\sum_{i=1}^p \lambda_i = \text{tr}(\mathbf{S})$ by (2.107). Thus we try to represent the p -dimensional points $(y_{i1}, y_{i2}, \dots, y_{ip})$ with a few principal components $(z_{i1}, z_{i2}, \dots, z_{ik})$ that account for a large proportion of the total variance. If a few variables have relatively large variances, they will figure disproportionately in $\sum_j s_{jj}$ and in the principal components. For example, if s_{22} is strikingly larger than the other variances, then in $z_1 = a_{11}y_1 + a_{12}y_2 + \cdots + a_{1p}y_p$, the coefficient a_{12} will be large and all other a_{1j} will be small.

When a ratio analogous to (12.5) is used for discriminant functions and canonical variates [see (8.13) and (11.9)], it is frequently referred to as *percent of variance*. However, in the case of discriminant functions and canonical variates, the eigenvalues are not variances, as they are in principal components.

If the variables are highly correlated, the essential dimensionality is much smaller than p . In this case, the first few eigenvalues will be large, and (12.5) will be close to 1 for a small value of k . On the other hand, if the correlations among the variables are all small, the dimensionality is close to p and the eigenvalues will be nearly equal. In this case, no useful reduction in dimension is achieved, because the principal components essentially duplicate the variables.

Any two principal components $z_i = \mathbf{a}'_i \mathbf{y}$ and $z_j = \mathbf{a}'_j \mathbf{y}$ are orthogonal for $i \neq j$, that is, $\mathbf{a}'_i \mathbf{a}_j = 0$, because \mathbf{a}_i and \mathbf{a}_j are eigenvectors of the symmetric matrix \mathbf{S} (see Section 2.11.6). Principal components also have the secondary property of being uncorrelated in the sample [see (12.2) and (3.63)]; that is, the covariance of z_i and z_j is zero:

$$s_{z_i z_j} = \mathbf{a}'_i \mathbf{S} \mathbf{a}_j = 0 \quad \text{for } i \neq j. \quad (12.6)$$

Discriminant functions and canonical variates, on the other hand, have the weaker property of being uncorrelated but not the stronger property of orthogonality. Thus when we plot the first two discriminant functions or canonical variates on perpendicular coordinate axes, there is some distortion of their true relationship because the actual angle between their axes is not 90° .

If we change the scale on one or more of the y 's, the shape of the swarm of points will change, and we will need different components to represent the new points. Hence the principal components are not scale invariant. We therefore need to be concerned with the units in which the variables are measured. If possible, all variables should be expressed in the same units. If the variables have widely disparate variances, we could standardize them before extracting eigenvalues and eigenvectors. This is equivalent to finding principal components of the correlation matrix \mathbf{R} and is treated in Section 12.5.

If one variable has a much greater variance than the other variables, the swarm of points will be elongated and will be nearly parallel to the axis corresponding to the variable with large variance. The first principal component will largely represent that variable, and the other principal components will have negligibly small variances. Such principal components (based on \mathbf{S}) do not involve the other $p - 1$ variables, and we may prefer to analyze the correlation matrix \mathbf{R} .

■ EXAMPLE 12.2.1

To illustrate principal components as a rotation when $p = 2$, we use two variables from the sons data of Table 3.8: y_1 is head length and y_2 is head width for the first son. The mean vector and covariance matrix are

$$\bar{\mathbf{y}} = \begin{pmatrix} 185.7 \\ 151.1 \end{pmatrix}, \quad \mathbf{S} = \begin{pmatrix} 95.29 & 52.87 \\ 52.87 & 54.36 \end{pmatrix}.$$

The eigenvalues and eigenvectors of \mathbf{S} are

$$\lambda_1 = 131.52, \quad \lambda_2 = 18.14, \\ \mathbf{a}'_1 = (a_{11}, a_{12}) = (.825, .565), \quad \mathbf{a}'_2 = (a_{21}, a_{22}) = (-.565, .825).$$

The symmetric pattern in the eigenvectors is due to their orthogonality: $\mathbf{a}'_1 \mathbf{a}_2 = a_{11}a_{21} + a_{12}a_{22} = 0$.

The observations are plotted in Figure 12.1, along with the (translated and) rotated axes. The major axis is the line passing through $\bar{\mathbf{y}}' = (185.7, 151.1)$ in the direction determined by $\mathbf{a}'_1 = (.825, .565)$; the slope is $a_{12}/a_{11} = .565/.825$. Alternatively, the equation of the major axis can be obtained by setting $z_2 = 0$:

$$z_2 = 0 = a_{21}(y_1 - \bar{y}_1) + a_{22}(y_2 - \bar{y}_2) \\ = -.565(y_1 - 185.7) + .825(y_2 - 151.1).$$

Note that the line formed by the major axis can be considered to be a regression line. It is fit to the points so that the perpendicular distance of the points to the line is minimized, rather than the usual vertical distance (see Section 12.3). □




Figure 12.1 Principal component transformation for the sons data.

12.2.2 Algebraic Approach

An algebraic approach to principal components can be briefly described as follows. As noted in Section 12.1, we seek a linear combination with maximal variance. By (3.55), the sample variance of $z = \mathbf{a}'\mathbf{y}$ is $\mathbf{a}'\mathbf{S}\mathbf{a}$. Since $\mathbf{a}'\mathbf{S}\mathbf{a}$ has no maximum if \mathbf{a} is unrestricted, we seek the maximum of

$$\lambda = \frac{\mathbf{a}'\mathbf{S}\mathbf{a}}{\mathbf{a}'\mathbf{a}}. \quad (12.7)$$

By an argument similar to that used in (8.8)–(8.12), the maximum value of λ is given by the largest eigenvalue in the expression

$$(\mathbf{S} - \lambda\mathbf{I})\mathbf{a} = \mathbf{0} \quad (12.8)$$

(see Problem 12.1). The eigenvector \mathbf{a}_1 corresponding to the largest eigenvalue λ_1 is the coefficient vector in $z_1 = \mathbf{a}_1'\mathbf{y}$, the linear combination with maximum variance.

Unlike discriminant analysis or canonical correlation, there is no inverse involved before obtaining eigenvectors for principal components. Therefore, \mathbf{S} can be singular, in which case some of the eigenvalues are zero and can be ignored. A singular \mathbf{S} would arise, for example, when $n < p$, that is, when the sample size is less than the number of variables.

This tolerance of principal component analysis for a singular \mathbf{S} is important in certain research situations. For example, suppose that one has a one-way MANOVA with 10 observations in each of three groups and that $p = 50$, so that there are 50 variables in each of these 30 observation vectors. A MANOVA test involving $\mathbf{E}^{-1}\mathbf{H}$ cannot be carried out directly in this case because \mathbf{E} is singular, but we could reduce

the 50 variables to a small number of principal components and then do a MANOVA test on the components. The principal components would be based on \mathbf{S} obtained from the 30 observations (ignoring groups). For entry into the MANOVA program, we would evaluate the principal components for each observation vector. If we are retaining k components, we calculate

$$\begin{aligned} z_{1i} &= \mathbf{a}'_1 \mathbf{y}_i \\ z_{2i} &= \mathbf{a}'_2 \mathbf{y}_i \\ &\vdots \\ z_{ki} &= \mathbf{a}'_k \mathbf{y}_i \end{aligned} \tag{12.9}$$

for $i = 1, 2, \dots, 30$. These are sometimes referred to as *component scores*. In vector form, (12.9) can be rewritten as

$$\mathbf{z}_i = \mathbf{A}_k \mathbf{y}_i, \tag{12.10}$$

where

$$\mathbf{z}_i = \begin{pmatrix} z_{1i} \\ z_{2i} \\ \vdots \\ z_{ki} \end{pmatrix} \quad \text{and} \quad \mathbf{A}_k = \begin{pmatrix} \mathbf{a}'_1 \\ \mathbf{a}'_2 \\ \vdots \\ \mathbf{a}'_k \end{pmatrix}.$$

We then use $\mathbf{z}_1, \mathbf{z}_2, \dots, \mathbf{z}_{30}$ as input to the MANOVA program.

Note that in this case with $p > n$, the k components would not likely be stable; that is, they would be different in a new sample. However, this is of no concern here because we are using the components only to extract information from the sample at hand in order to compare the three groups.

■ EXAMPLE 12.2.2

Consider the football data of Table 8.3. In Example 8.8, we saw that high school football players (group 1) differed from the other two groups, college football players and college-age non-football players. Therefore, to obtain a homogeneous group of observations, we delete group 1 and use groups 2 and 3 combined. The covariance matrix is as follows:

$$\mathbf{S} = \begin{pmatrix} .370 & .602 & .149 & .044 & .107 & .209 \\ .602 & 2.629 & .801 & .666 & .103 & .377 \\ .149 & .801 & .458 & .011 & -.013 & .120 \\ .044 & .666 & .011 & 1.474 & .252 & -.054 \\ .107 & .103 & -.013 & .252 & .488 & -.036 \\ .209 & .377 & .120 & -.054 & -.036 & .324 \end{pmatrix}.$$

The total variance is

$$\sum_{j=1}^6 s_{jj} = \sum_{i=1}^6 \lambda_i = 5.743.$$

The eigenvalues of \mathbf{S} are as follows:

Eigenvalue	Proportion of Variance	Cumulative Proportion
3.323	.579	.579
1.374	.239	.818
.476	.083	.901
.325	.057	.957
.157	.027	.985
.088	.015	1.000

The first two principal components account for 81.8% of the total variance. The corresponding eigenvectors are as follows:

	a_1	a_2
WDIM	.207	-.142
CIRCUM	.873	-.219
FBEYE	.261	-.231
EYEHLD	.326	.891
EARHDL	.066	.222
JAW	.128	-.187

Thus the first two principal components are

$$\begin{aligned} z_1 &= \mathbf{a}_1' \mathbf{y} = .207y_1 + .873y_2 + .261y_3 + .326y_4 + .066y_5 + .128y_6, \\ z_2 &= \mathbf{a}_2' \mathbf{y} = -.142y_1 - .219y_2 - .231y_3 + .891y_4 + .222y_5 - .187y_6. \end{aligned}$$

Notice that the large coefficient in z_1 and the large coefficient in z_2 , .873 and .891, respectively, correspond to the two largest variances on the diagonal of \mathbf{S} . The two variables with large variances, y_2 and y_4 , have a notable influence on the first two principal components. However, z_1 and z_2 are still meaningful linear functions. If the six variances were closer in size, the six variables would enter more evenly into the first two principal components. On the other hand, if the variances of y_2 and y_4 were substantially larger, z_1 and z_2 would be essentially equal to y_2 and y_4 , respectively.

Note that y_2 and y_3 did not contribute at all when this data set was used to separate groups in Examples 8.5, 8.9, 9.3.1, and 9.6(a). However, these two variables are very useful here in the first two dimensions showing the spread of individual observations. \square

12.3 PRINCIPAL COMPONENTS AND PERPENDICULAR REGRESSION

It was noted in Section 12.2.1 that principal components constitute a rotation of axes. Another geometric property of the line formed by the first principal component is




Figure 12.2 The first principal component as a perpendicular regression line.

that it minimizes the total sum of squared perpendicular distances from the points to the line. This is easily demonstrated in the bivariate case. The first principal component line is plotted in Figure 12.2 for the first two variables of the sons data, as in Example 12.2.1. The perpendicular distance from each point to the line is simply z_2 , the second coordinate in the transformed coordinates (z_1, z_2) . Hence the sum of squares of perpendicular distances is

$$\sum_{i=1}^n z_{2i}^2 = \sum_{i=1}^n [\mathbf{a}_2'(\mathbf{y}_i - \bar{\mathbf{y}})]^2, \quad (12.11)$$

where \mathbf{a}_2 is the second eigenvector of \mathbf{S} and we use $\mathbf{y}_i - \bar{\mathbf{y}}$ because the axes have been translated to the new origin $\bar{\mathbf{y}}$. Since $\mathbf{a}_2'(\mathbf{y}_i - \bar{\mathbf{y}}) = (\mathbf{y}_i - \bar{\mathbf{y}})' \mathbf{a}_2$, we can write (12.11) in the form

$$\begin{aligned} \sum_{i=1}^n z_{2i}^2 &= \sum_{i=1}^n \mathbf{a}_2'(\mathbf{y}_i - \bar{\mathbf{y}})(\mathbf{y}_i - \bar{\mathbf{y}})' \mathbf{a}_2 \\ &= \mathbf{a}_2' \left[\sum_i (\mathbf{y}_i - \bar{\mathbf{y}})(\mathbf{y}_i - \bar{\mathbf{y}})' \right] \mathbf{a}_2 \quad [\text{by (2.44)}] \\ &= (n-1)\mathbf{a}_2' \mathbf{S} \mathbf{a}_2 = (n-1)\lambda_2 \quad [\text{by (3.27)}], \end{aligned} \quad (12.12)$$

which is a minimum [see remarks following (12.4)].

For the two variables y_1 and y_2 , as plotted in Figure 12.2, the ordinary regression line of y_2 on y_1 minimizes the sum of squares of vertical distances from the points to the line. Similarly, the regression of y_1 on y_2 minimizes the sum of squares of




Figure 12.3 Regression lines compared with first principal component (P.C.) line.

horizontal distances from the points to the line. The first principal component line represents a “perpendicular” regression line that lies between the other two. The three lines are compared in Figure 12.3 for the partial sons data. The equation of the first principal component line is easily obtained by setting $z_2 = 0$:

$$\begin{aligned} z_2 &= \mathbf{a}_2'(\mathbf{y} - \bar{\mathbf{y}}) = 0, \\ a_{21}(y_1 - \bar{y}_1) + a_{22}(y_2 - \bar{y}_2) &= 0, \\ -.565(y_1 - \bar{y}_1) + .825(y_2 - \bar{y}_2) &= 0. \end{aligned}$$

12.4 PLOTTING OF PRINCIPAL COMPONENTS

The plots in Figures 12.1 and 12.2 were illustrations of principal components as a rotation of axes when $p = 2$. When $p > 2$, we can plot the first two components as a dimension reduction device. We simply evaluate the first two components (z_1, z_2) for each observation vector and plot these n points. The plot is equivalent to a projection of the p -dimensional data swarm onto the plane that shows the greatest spread of the points.

The plot of the first two components may reveal some important features of the data set. In Example 12.4(a), we show a principal component plot that exhibits a pattern typical of a sample from a multivariate normal distribution. One of the objectives of plotting is to check for departures from normality, such as outliers or nonlinearity. In Examples 12.4(b) and 12.4(c), we illustrate principal component plots showing a nonnormal pattern characterized by the presence of outliers. Jackson

Table 12.1 Principal Components for the Ramus Bone Data of Table 3.7

Eigenvalues		First Two Eigenvectors		
Number	Value	Variable	a_1	a_2
1	25.05	AGE 8	.474	.592
2	1.74	AGE 8.5	.492	.406
3	.22	AGE 9	.515	-.304
4	.11	AGE 9.5	.517	-.627

(1980) provided a test for adequacy of representation of observation vectors in terms of principal components.

Gnanadesikan (1997, p. 308) pointed out that, in general, the first few principal components are sensitive to outliers that inflate variances or distort covariances, and the last few are sensitive to outliers that introduce artificial dimensions or mask singularities. We could examine the bivariate plots of at least the first two and the last two principal components in a search for outliers that may exert undue influence.

Devlin et al. (1981) recommended the extraction of principal components from robust estimates of S or R that reduce the influence of outliers. Campbell (1980) and Ruymgaart (1981) discussed direct robust estimation of principal components. Critchley (1985) developed methods for detection of influential observations in principal component analysis.

Another feature of the data that a plot of the first two components may reveal is a tendency of the points to cluster. The plot may reveal groupings of points; this is illustrated in Example 12.4(d).

■ EXAMPLE 12.4(a)

For the modified football data in Example 12.2.2, the first two principal components were given as follows:

$$\begin{aligned} z_1 &= \mathbf{a}'_1 \mathbf{y} = .207y_1 + .873y_2 + .261y_3 + .326y_4 + .066y_5 + .128y_6, \\ z_2 &= \mathbf{a}'_2 \mathbf{y} = -.142y_1 - .219y_2 - .231y_3 + .891y_4 + .222y_5 - .187y_6. \end{aligned}$$

These are evaluated for each observation vector and plotted in Figure 12.4. (For convenience in scaling, $\mathbf{y} - \bar{\mathbf{y}}$ was used in the computations.) The pattern is typical of that from a multivariate normal distribution. Note that the variance along the z_1 axis is greater than the variance in the z_2 direction, as expected.

□

■ EXAMPLE 12.4(b)

In Figures 4.10 and 4.11, the $Q-Q$ plot and bivariate scatterplots for the ramus bone data of Table 3.7 exhibit a nonnormal pattern. A principal component analysis using the covariance matrix is given in Table 12.1, and the first two




Figure 12.4 Plot of first two components for the modified football data.

principal components are plotted in Figure 12.5. The presence of three outliers that cause a nonnormal pattern is evident. These outliers do not appear when the four variables are examined individually. \square




Figure 12.5 First two principal components for the ramus bone data in Table 3.7.




Figure 12.6 First two principal components for economics data.

■ EXAMPLE 12.4(c)

A rather extreme example of the effect of an outlier is given by Devlin et al. (1981). The data set involved $p = 14$ economic variables for $n = 29$ chemical companies. The first two principal components are plotted in Figure 12.6. The sample correlation $r_{z_1 z_2}$ is indeed zero for all 29 points, as it must be [see (12.6)], but if the apparent outlier is excluded from the computation, then $r_{z_1 z_2} = .99$ for the remaining 28 points. If the outlier were deleted from the data set, the axes of the principal components would pass through the natural extensions of the data swarm. \square

■ EXAMPLE 12.4(d)

Jeffers (1967) applied principal component analysis to a sample of 40 alate adelges (winged aphids) on which the following 19 variables had been measured:

LENGTH	body length
WIDTH	body width
FORWING	forewing length
HINWING	hindwing length
SPIRAC	number of spiracles
ANTSEG 1	length of antennal segment I
ANTSEG 2	length of antennal segment II

ANTSEG 3	length of antennal segment III
ANTSEG 4	length of antennal segment IV
ANTSEG 5	length of antennal segment V
ANTSPIN	number of antennal spines
TARSUS 3	leg length, tarsus III
TIBIA 3	leg length, tibia III
FEMUR 3	leg length, femur III
ROSTRUM	rostrum
OVIPOS	ovipositor
OVSPIN	number of ovipositor spines
FOLD	anal fold
HOOKS	number of hindwing hooks

Table 12.2 Correlation Matrix for Winged Aphid Variables (Lower Triangle)

Table 12.3 Eigenvalues of the Correlation Matrix of the Winged Aphid Data

Component	Eigenvalue	Percent of Variance	Cumulative Percent
1	13.861	73.0	73.0
2	2.370	12.5	85.4
3	.748	3.9	89.4
4	.502	2.6	92.0
5	.278	1.4	93.5
6	.266	1.4	94.9
7	.193	1.0	95.9
8	.157	.8	96.7
9	.140	.7	97.4
10	.123	.6	98.1
11	.092	.4	98.6
12	.074	.4	99.0
13	.060	.3	99.3
14	.042	.2	99.5
15	.036	.2	99.7
16	.024	.1	99.8
17	.020	.1	99.9
18	.011	.1	100.0
19	.003	.0	100.0
			19.000

An objective in the study was to determine the number of distinct taxa present in the habitat where the sample was taken. Since adelges are difficult to identify by the usual taxonomic methods, principal component analysis was used to search for groupings among the 40 individuals in the sample.

The correlation matrix is given in Table 12.2, and the eigenvalues and first four eigenvectors are in Tables 12.3 and 12.4, respectively. The eigenvectors are scaled so that the largest value in each is 1. The first principal component is largely an index of size. The second component is associated with SPIRAC, OVIPOS, OVSPIN, and FOLD.

The first two components were computed for each of the 40 individuals and plotted in Figure 12.7. Since the first two components account for 85% of the total variance, the plot represents the data with very little distortion. There are four major groups, apparently corresponding to species. The groupings form an interesting S-shape. \square

12.5 PRINCIPAL COMPONENTS FROM THE CORRELATION MATRIX

Generally, extracting components from S rather than R remains closer to the spirit and intent of principal component analysis, especially if the components are to be

Table 12.4 Eigenvectors for the First Four Components of the Winged Aphid Data

Variable	Eigenvectors			
	1	2	3	4
LENGTH	.96	-.06	.03	-.12
WIDTH	.98	-.12	.01	-.16
FORWING	.99	-.06	-.06	-.11
HINWING	.98	-.16	.03	-.00
SPIRAC	.61	.74	-.20	1.00
ANTSEG 1	.91	.33	.04	.02
ANTSEG 2	.96	.30	.00	-.04
ANTSEG 3	.88	-.43	.06	-.18
ANTSEG 4	.90	-.08	.18	-.01
ANTSEG 5	.94	.05	.11	.03
ANTSPIN	-.49	.37	1.00	.27
TARSUS 3	.99	-.02	.03	-.29
TIBIA 3	1.00	-.05	.09	-.31
FEMUR 3	.99	-.12	.12	-.31
ROSTRUM	.96	.02	.08	-.06
OVIPOS	.76	.73	-.03	-.09
OVSPIN	.41	1.00	-.16	-.06
FOLD	-.71	.64	.04	-.80
HOOKS	.76	-.52	.06	.72

used in further computations. However, in some cases, the principal components will be more interpretable if \mathbf{R} is used. For example, if the variances differ widely or if the measurement units are not commensurate, the components of \mathbf{S} will be dominated by the variables with large variances. The other variables will contribute very little. For a more balanced representation in such cases, components of \mathbf{R} may be used (see, for example, Problem 12.9).

As with any change of scale, when the variables are standardized in transforming from \mathbf{S} to \mathbf{R} , the shape of the swarm of points will change. Note, however, that after transforming to \mathbf{R} , any further changes of scale on the variables would not affect the components because changes of scale do not change \mathbf{R} . Thus the principal components from \mathbf{R} are scale invariant.

To illustrate how the eigenvalues and eigenvectors change when converting from \mathbf{S} to \mathbf{R} , we use a simple bivariate example in which one variance is substantially larger than the other. Suppose that \mathbf{S} and the corresponding \mathbf{R} have the values

$$\mathbf{S} = \begin{pmatrix} 1 & 4 \\ 4 & 25 \end{pmatrix}, \quad \mathbf{R} = \begin{pmatrix} 1 & .8 \\ .8 & 1 \end{pmatrix}.$$

The eigenvalues and eigenvectors from \mathbf{S} are




Figure 12.7 Plotted values of the first two components for individual insects.

$$\begin{aligned}\lambda_1 &= 25.65, & \mathbf{a}_1' &= (.160, .987), \\ \lambda_2 &= .35, & \mathbf{a}_2' &= (.987, -.160).\end{aligned}$$

The patterns we see in $\lambda_1, \lambda_2, \mathbf{a}_1$, and \mathbf{a}_2 are quite predictable. The symmetry in \mathbf{a}_1 and \mathbf{a}_2 is due to their orthogonality, $\mathbf{a}_1' \mathbf{a}_2 = 0$. The large variance of y_2 in \mathbf{S} is reflected in the first principal component $z_1 = .160y_1 + .987y_2$, where y_2 is weighted heavily. Thus the first principal component z_1 essentially duplicates y_2 and does not show the mutual effect of y_1 and y_2 . As expected, z_1 accounts for virtually all of the total variance:

$$\frac{\lambda_1}{\lambda_1 + \lambda_2} = \frac{25.65}{26} = .9865.$$

The eigenvalues and eigenvectors of \mathbf{R} are

$$\begin{aligned}\lambda_1 &= 1.8, & \mathbf{a}_1' &= (.707, .707), \\ \lambda_2 &= .2, & \mathbf{a}_2' &= (.707, -.707).\end{aligned}$$

The first principal component of \mathbf{R} ,

$$z_1 = .707 \frac{y_1 - \bar{y}_1}{1} + .707 \frac{y_2 - \bar{y}_2}{5},$$

accounts for a high proportion of variance,

$$\frac{\lambda_1}{\lambda_1 + \lambda_2} = \frac{1.8}{2} = .9,$$

because the variables are fairly highly correlated ($r = .8$). But the standardized variables $(y_1 - \bar{y}_1)/1$ and $(y_2 - \bar{y}_2)/5$ are equally weighted in z_1 , due to the equality of the diagonal elements ("variances") of \mathbf{R} .

We now list some general comparisons of principal components from \mathbf{R} with those from \mathbf{S} :

1. The percent of variance in (12.5) accounted for by the components of \mathbf{R} will differ from the percent for \mathbf{S} , as illustrated above.
2. The coefficients of the principal components from \mathbf{R} differ from those obtained from \mathbf{S} , as illustrated above.
3. If we express the components from \mathbf{R} in terms of the original variables, they still will not agree with the components from \mathbf{S} . By transforming the standardized variables back to the original variables in the above illustration, the components of \mathbf{R} become

$$\begin{aligned} z_1 &= .707 \frac{y_1 - \bar{y}_1}{1} + .707 \frac{y_2 - \bar{y}_2}{5} \\ &= .707y_1 + .141y_2 + \text{const}, \\ z_2 &= .707 \frac{y_1 - \bar{y}_1}{1} - .707 \frac{y_2 - \bar{y}_2}{5} \\ &= .707y_1 - .141y_2 + \text{const}. \end{aligned}$$

As expected, these are very different from the components extracted directly from \mathbf{S} . This problem arises, of course, because of the lack of scale invariance of the components of \mathbf{S} .

4. The principal components from \mathbf{R} are scale invariant, because \mathbf{R} itself is scale invariant.
5. The components from a given matrix \mathbf{R} are not unique to that \mathbf{R} . For example, in the bivariate case, the eigenvalues of

$$\mathbf{R} = \begin{pmatrix} 1 & r \\ r & 1 \end{pmatrix}$$

are given by

$$\lambda_1 = 1 + r, \quad \lambda_2 = 1 - r, \quad (12.13)$$

and the eigenvectors are $\mathbf{a}'_1 = (.707, .707)$ and $\mathbf{a}'_2 = (.707, -.707)$, which give principal components

$$\begin{aligned} z_1 &= .707 \frac{y_1 - \bar{y}_1}{s_1} + .707 \frac{y_2 - \bar{y}_2}{s_2}, \\ z_2 &= .707 \frac{y_1 - \bar{y}_1}{s_1} - .707 \frac{y_2 - \bar{y}_2}{s_2}. \end{aligned} \quad (12.14)$$

The components in (12.14) do not depend on r . For example, they serve equally well for $r = .01$ and for $r = .99$. For $r = .01$, the proportion of variance explained by z_1 is $\lambda_1/(\lambda_1 + \lambda_2) = (1 + .01)/(1 + .01 + 1 - .01) = 1.01/2 = .505$. For $r = .99$, the ratio is $1.99/2 = .995$. Thus the statement that the first component from a correlation matrix accounts for, say, 90% of the variance is not very meaningful. In general, for $p > 2$, the components from \mathbf{R} depend only on the ratios (relative values) of the correlations, not on their actual values, and components of a given \mathbf{R} matrix will serve for other \mathbf{R} matrices [see Rencher (1998, Section 9.4)].

12.6 DECIDING HOW MANY COMPONENTS TO RETAIN

In every application, a decision must be made on how many principal components should be retained in order to effectively summarize the data. The following guidelines have been proposed:

1. Retain sufficient components to account for a specified percentage of the total variance, say 80%.
2. Retain the components whose eigenvalues are greater than the average of the eigenvalues, $\sum_{i=1}^p \lambda_i/p$. For a correlation matrix, this average is 1.
3. Use the *scree graph*, a plot of λ_i versus i , and look for a natural break between the “large” eigenvalues and the “small” eigenvalues.
4. Test the significance of the “larger” components, that is, the components corresponding to the larger eigenvalues.

We now discuss the above four criteria for choosing the components to keep. Note, however, that the smallest components may carry valuable information that should not be routinely ignored (see Section 12.7).

In method 1, the challenge lies in selecting an appropriate threshold percentage. If we aim too high, we run the risk of including components that are either *sample specific* or *variable specific*. By *sample specific* we mean that a component may not generalize to the population or to other samples. A *variable specific* component is dominated by a single variable and does not represent a composite summary of several variables.

Method 2 is widely used and is the default in many software packages. By (2.107), $\sum_i \lambda_i = \text{tr}(\mathbf{S})$, and the average eigenvalue is also the average variance of the individual variables. Thus method 2 retains those components that account for more variance than the average variance of the variables. In cases where the data can be successfully summarized in a relatively small number of dimensions, there is often a wide gap between the two eigenvalues that fall on both sides of the average. In Example 12.2.2, the average eigenvalue (of \mathbf{S}) for the football data is .957, which is




Figure 12.8 Scree graph for eigenvalues of modified football data.

amply bracketed by $\lambda_2 = 1.37$ and $\lambda_3 = .48$. In the winged aphid data in Example 12.4(d), the second and third eigenvalues (of \mathbf{R}) are 2.370 and .748, leaving a comfortable margin on both sides of 1. In some cases, one may wish to move the cutoff point slightly to accommodate a visible gap in eigenvalues.

The scree graph in method 3 is named for its similarity in appearance to a cliff with rocky debris at its bottom. The scree graph for the modified football data of Example 12.2.2 exhibits an ideal pattern, as shown in Figure 12.8. The first two eigenvalues form a steep curve, followed by a bend and then a straight-line trend with shallow slope. The recommendation is to retain those eigenvalues in the steep curve *before* the first one on the straight line. Thus in Figure 12.8, two components would be retained. In practice, the turning point between the steep curve and the straight line may not be as distinct as this or there may be more than one discernible bend. In such cases, this approach is not as conclusive. The scree graph for the winged aphid data in Example 12.4(d) is plotted in Figure 12.9. The plot would suggest that two components be retained (possibly four).

The remainder of this section is devoted to method 4, tests of significance. The tests assume multivariate normality, which is not required for estimation of principal components.

It may be useful to make a preliminary test of complete independence of the variables, as in Section 7.4.3: $H_0: \boldsymbol{\Sigma} = \text{diag}(\sigma_{11}, \sigma_{22}, \dots, \sigma_{pp})$, or equivalently, $H_0: \mathbf{P}_\rho = \mathbf{I}$. The test statistic is given in (7.37) and (7.38). If the results indicate that the variables are independent, there is no point in extracting principal components, since (except for sampling fluctuation) the variables themselves already form the principal components.




Figure 12.9 Scree graph for eigenvalues of winged aphid data.

To test the significance of the “larger” components, we test the hypothesis that the last k population eigenvalues are small and equal, $H_{0k}: \gamma_{p-k+1} = \gamma_{p-k+2} = \cdots = \gamma_p$, where $\gamma_1, \gamma_2, \dots, \gamma_p$ denote the population eigenvalues, namely, the eigenvalues of Σ . The implication is that the first sample components capture all the essential dimensions, while the last components reflect noise. If H_0 is true, the last k sample eigenvalues will tend to have the pattern shown by the straight line with small slope in the ideal scree graph, such as in Figure 12.8 or 12.9.

To test $H_{0k}: \gamma_{p-k+1} = \cdots = \gamma_p$ using a likelihood ratio approach, we calculate the average of the last k eigenvalues of \mathbf{S} ,

$$\bar{\lambda} = \sum_{i=p-k+1}^p \frac{\lambda_i}{k}$$

and use the test statistic

$$u = \left(n - \frac{2p + 11}{6} \right) \left(k \ln \bar{\lambda} - \sum_{i=p-k+1}^p \ln \lambda_i \right), \quad (12.15)$$

which has an approximate χ^2 -distribution. We reject H_0 if $u \geq \chi^2_{\alpha, \nu}$, where $\nu = \frac{1}{2}(k-1)(k+2)$.

To carry out this procedure, we could begin by testing $H_{02}: \gamma_{p-1} = \gamma_p$. If this is accepted, we could then test $H_{03}: \gamma_{p-2} = \gamma_{p-1} = \gamma_p$ and continue testing in this fashion until H_{0k} is rejected for some value of k .

In practice, when the variables are fairly highly correlated and the data can be successfully represented by a small number of principal components, the first three

methods will typically agree on the number of components to retain, and the test in method 4 will often indicate a larger number of components.

■ EXAMPLE 12.6

We apply the above four criteria to the modified football data of Example 12.2.2.

For method 1, we simply examine the eigenvalues and their proportion of variance explained, as obtained in Example 12.2.2:

Eigenvalue	Proportion of Variance	Cumulative Proportion
3.323	.579	.579
1.374	.239	.818
.476	.083	.901
.325	.057	.957
.157	.027	.985
.088	.015	1.000

To account for 82% of the variance, we would keep two components. This percent of variance is high enough for most descriptive purposes. For certain other applications, such as input to another analysis, we might wish to retain three components, which would account for 90% of the variance.

To apply method 2, we find the average eigenvalue to be

$$\bar{\lambda} = \frac{1}{6} \sum_{i=1}^6 \lambda_i = \frac{5.742824}{6} = .957.$$

Since only λ_1 and λ_2 exceed .957, we would retain two components.

For method 3, the scree graph in Figure 12.8 indicates conclusively that two components should be retained.

To implement method 4, we carry out the significance tests in (12.15). The values of the test statistic u for $k = 2, 3, \dots, 6$ are as follows:

Eigenvalue	k	u	df	$\chi^2_{.05}$
3.32341	6	245.57	20	31.41
1.37431	5	123.93	14	23.68
.47607	4	44.10	9	16.92
.32468	3	23.84	5	11.07
.15650	2	4.62	2	5.99
.08785	1			

The tests indicate that only the last two (population) eigenvalues are equal and we should retain the first four. This differs from the results of the other three criteria, which are in close agreement that two components should be retained.

□

12.7 INFORMATION IN THE LAST FEW PRINCIPAL COMPONENTS

Up to this point, we have focused on using the first few principal components to summarize and simplify the data. However, the last few components may carry useful information in some applications.

Since the eigenvalues serve as variances of the principal components, the last few principal components have smaller variances. If the variance of a component is zero or close to zero, the component represents a linear relationship among the variables that is essentially constant; that is, the relationship holds for all y_i 's in the sample. Thus if the last eigenvalue is near zero, it signifies the presence of a collinearity that may provide new information for the researcher. Suppose, for example, that there are five variables and $y_5 = \sum_{j=1}^4 y_j / 4$. Then S is singular, and barring round-off error, λ_5 will be zero. Thus $s_{z_5}^2 = 0$, and z_5 is constant. As noted early in Section 12.2, the y_i 's are centered, because the origin of the principal components is translated to \bar{y} . Hence the constant value of z_5 is its mean, which is zero:

$$z_5 = \mathbf{a}'_5 \mathbf{y} = a_{51}y_1 + a_{52}y_2 + \cdots + a_{55}y_5 = 0.$$

Since this must reflect the dependency of y_5 on y_1, y_2, y_3 , and y_4 , the eigenvector \mathbf{a}'_5 will be proportional to $(1, 1, 1, 1, -4)$.

12.8 INTERPRETATION OF PRINCIPAL COMPONENTS

In Section 12.5, we noted that principal components obtained from R are not compatible with those obtained from S . Because of this lack of scale invariance of principal components from S , the coefficients cannot be converted to standardized form, as can be done with coefficients in discriminant functions in Chapter 8 and canonical variates in Chapter 11. Hence interpretation of principal components is not as clear-cut as with previous linear functions that we have discussed. We must choose between components of S or R , knowing they will have a different interpretation. If the variables have widely disparate variances, we can use R instead of S to improve interpretation.

For certain patterns of elements in S or R , the form of the principal components can be predicted. This aid to interpretation is discussed in Section 12.8.1. As with discriminant functions and canonical variates, some writers have advocated rotation and the use of correlations between the variables and the principal components. We argue against the use of these two approaches to interpretation in Sections 12.8.2 and 12.8.3.

12.8.1 Special Patterns in S or R

In the covariance or correlation matrix, we may recognize a distinguishing pattern from which the structure of the principal components can be deduced. For example, we noted in Section 12.2 that if one variable has a much larger variance than

the other variables, this variable will dominate the first component, which will account for most of the variance. Another case in which a component will duplicate a variable occurs when the variable is uncorrelated with the other variables. We now demonstrate this by showing that if all p variables are uncorrelated, the variables themselves are the principal components. If the variables were uncorrelated (orthogonal), \mathbf{S} would have the form

$$\mathbf{S} = \begin{pmatrix} s_{11} & 0 & \cdots & 0 \\ 0 & s_{22} & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & s_{pp} \end{pmatrix}, \quad (12.16)$$

and the characteristic equation would be

$$0 = |\mathbf{S} - \lambda \mathbf{I}| = \prod_{i=1}^p (s_{ii} - \lambda) \quad [\text{by (2.83)}],$$

which has solutions

$$\lambda_i = s_{ii}, \quad i = 1, 2, \dots, p. \quad (12.17)$$

The corresponding normalized eigenvectors have a 1 in the i th position and 0's elsewhere:

$$\mathbf{a}'_i = (0, \dots, 0, 1, 0, \dots, 0). \quad (12.18)$$

Thus the i th component is

$$z_i = \mathbf{a}'_i \mathbf{y} = y_i.$$

In practice, the sample correlations (of continuous random variables) will not be zero, but if the correlations are all small, the principal components will largely duplicate the variables.

By the Perron–Frobenius theorem in Section 2.11.4, if all correlations or covariances are positive, all elements of the first eigenvector \mathbf{a}_1 are positive. Since the remaining eigenvectors $\mathbf{a}_2, \mathbf{a}_3, \dots, \mathbf{a}_p$ are orthogonal to \mathbf{a}_1 , they must have both positive and negative elements. When all elements of \mathbf{a}_1 are positive, the first component is a weighted average of the variables and is sometimes referred to as a measure of *size*. Likewise, the positive and negative coefficients in subsequent components may be regarded as defining *shape*. This pattern is often seen when the variables are various measurements of an organism.

■ EXAMPLE 12.8.1

In the modified football data of Example 12.2.2, there are a few negative covariances in \mathbf{S} , but they are small, and all elements of the first eigenvector remain positive. The second eigenvector therefore has positive and negative elements:

	First Two Eigenvectors	
	a_1	a_2
WDIM	.207	-.142
CIRCUM	.873	-.219
FBEYE	.261	-.231
EYEHD	.326	.891
EARHD	.066	.222
JAW	.128	-.187

With all positive coefficients, the first component z_1 is an overall measure of head size (z_1 increases if all six variables increase). The second component z_2 is a shape component that contrasts the vertical measurements EYEHD and EARHD with the three lateral measurements and CIRCUM (z_2 increases if EYEHD and EARHD increase and the other four variables decrease). \square

12.8.2 Rotation

The principal components are initially obtained by rotating axes in order to line up with the natural extensions of the system, whereupon the new variables become uncorrelated and reflect the directions of maximum variance. If the resulting components do not have a satisfactory interpretation, they can be further rotated, seeking dimensions in which many of the coefficients of the linear combinations are near zero to simplify interpretation.

However, the new rotated components are correlated, and they do not successively account for maximum variance. They are therefore no longer principal components in the usual sense, and their routine use is questionable. For improved interpretation, one may wish to try factor analysis (Chapter 13), in which rotation does not destroy any properties. (In factor analysis, the rotation does not involve the variables y_1, y_2, \dots, y_p , but another space, that of the factor loadings.)

12.8.3 Correlations Between Variables and Principal Components

The use of correlations between variables and principal components is widely recommended as an aid to interpretation. It was noted in Sections 8.7.3 and 11.5.2 that analogous correlations for discriminant functions and canonical variates are not useful in a multivariate context because they provide only univariate information about how each variable operates by itself, ignoring the other variables. Rencher (1992b) obtained a similar result for principal components.

We denote the correlation between the i th variable y_i and the j th principal component z_j by $r_{y_i z_j}$. Because of the orthogonality of the z_j 's, we have the simple relationship

$$r_{y_i z_1}^2 + r_{y_i z_2}^2 + \dots + r_{y_i z_k}^2 = R_{y_i | z_1, \dots, z_k}^2, \quad (12.19)$$

where k is the number of components retained and $R_{y_i | z_1, \dots, z_k}^2$ is the squared multiple correlation of y_i with the z_j 's. Thus $r_{y_i z_j}^2$ forms part of $R_{y_i | z_1, \dots, z_k}^2$, which shows how y_i relates to the z 's by itself, not what it contributes in the presence of the other

Table 12.5 Eigenvectors Obtained from \mathbf{S} , Correlations Between Variables and Principal Components, and R^2 for the First Two Principal Components

Variable	Eigenvectors from \mathbf{S}		Correlations		$R^2_{y_i z_1, z_2}$
	a_1	a_2	$r_{y_i z_1}$	$r_{y_i z_2}$	
1	.21	-.14	.62	-.27	.46
2	.87	-.22	.98	-.16	.99
3	.26	-.23	.70	-.40	.66
4	.33	.89	.49	.86	.98
5	.07	.22	.17	.37	.17
6	.13	-.19	.41	-.39	.32

y 's. The correlations are therefore not informative about the joint contribution of the y 's in a principal component.

Note that the simple partitioning of R^2 into the sum of squares of correlations in (12.19) does not happen in practice when the independent variables (x 's) are correlated. However, here the z 's are principal components and are therefore orthogonal.

Since we do not recommend rotation or correlations for interpretation, we are left with the coefficients themselves, obtained from the eigenvectors of either \mathbf{S} or \mathbf{R} .

■ EXAMPLE 12.8.3

In Example 12.8.1, the eigenvectors of \mathbf{S} from the modified football data gave a satisfactory interpretation of the first two principal components as head size and shape. We give these in Table 12.5, along with the correlations between each of the variables y_1, y_2, \dots, y_6 and the first two principal components z_1 and z_2 . For comparison we also give $R^2_{y_i|z_1, z_2}$ for each variable.

The correlations rank the variables somewhat differently in their contribution to the components, since they form part of the univariate information provided by R^2 for each variable by itself. For example, for the first component, the correlations rank the variables in the order 2, 3, 1, 4, 6, 5, whereas the coefficients (eigenvectors) from \mathbf{S} rank them in the order 2, 4, 3, 1, 6, 5. \square

12.9 SELECTION OF VARIABLES

We have previously discussed subset selection in connection with Wilks' Λ (Section 6.11.2), discriminant analysis (Section 8.9), classification analysis (Section 9.6), and regression (Sections 10.2.7 and 10.8). In each case the criterion for selection of variables was the relationship of the variables to some external factor, such as dependent variable(s), separation of groups, or correct classification rates. In the context of principal components, we have no dependent variable, as in regression, and no groupings among the observations, as in discriminant analysis. With no external in-

fluence, we simply wish to find the subset that best captures the internal variation (and covariation) of the variables.

Jolliffe (1972, 1973) discussed eight selection methods and referred to the process as *discarding variables*. The eight methods were based on three basic approaches: multiple correlation, clustering of variables, and principal components. One of the correlation methods, for example, proceeds in a stepwise fashion, deleting at each step the variable that has the largest multiple correlation with the other variables. The clustering methods partition the variables into groups or clusters and select a variable from each cluster.

We describe Jolliffe's principal component methods in the context of selecting a subset of 10 variables out of 50 variables. One of his techniques associates a variable with each of the first 10 principal components and retains these 10 variables. Another approach is to associate a variable with each of the last 40 principal components and delete the 40 variables. To associate a variable with a principal component, we choose the variable corresponding to the largest coefficient (in absolute value) in the component, providing the variable has not previously been selected. We can use components extracted from either \mathbf{S} or \mathbf{R} . For example, in the two principal components for the football data in Example 12.2.2, we would choose variables 2 and 4, which clearly have the largest coefficients in the two components. Jolliffe's methods could also be applied iteratively, with the principal components being recomputed after a variable is retained or deleted.

Jolliffe (1972) compared the eight methods using both real and simulated data and found that the methods based on principal components performed well in comparison to the regression and cluster-based methods. But he concluded that no single method was uniformly best.

McCabe (1984) suggested several criteria for selection, most of which are based on the conditional covariance matrix of the variables not selected, given those selected. He denoted the selected variables as *principal variables*. Let \mathbf{y} be partitioned as

$$\mathbf{y} = \begin{pmatrix} \mathbf{y}_1 \\ \mathbf{y}_2 \end{pmatrix},$$

where \mathbf{y}_1 contains the selected variables and \mathbf{y}_2 consists of the variables not selected. The corresponding covariance matrix is

$$\text{cov}(\mathbf{y}) = \boldsymbol{\Sigma} = \begin{pmatrix} \boldsymbol{\Sigma}_{11} & \boldsymbol{\Sigma}_{12} \\ \boldsymbol{\Sigma}_{21} & \boldsymbol{\Sigma}_{22} \end{pmatrix}.$$

By (4.8), the conditional covariance matrix is given by (assuming normality)

$$\text{cov}(\mathbf{y}_2|\mathbf{y}_1) = \boldsymbol{\Sigma}_{22} - \boldsymbol{\Sigma}_{21}\boldsymbol{\Sigma}_{11}^{-1}\boldsymbol{\Sigma}_{12},$$

which is estimated by $\mathbf{S}_{22} - \mathbf{S}_{21}\mathbf{S}_{11}^{-1}\mathbf{S}_{12}$. To find a subset \mathbf{y}_1 of size m , two of McCabe's criteria are to choose the subset \mathbf{y}_1 that

1. Minimizes $|\mathbf{S}_{22} - \mathbf{S}_{21}\mathbf{S}_{11}^{-1}\mathbf{S}_{12}|$ and

2. Maximizes $\sum_{i=1}^{m^*} r_i^2$, where $r_i, i = 1, 2, \dots, m^* = \min(m, p - m)$ are the canonical correlations between the m selected variables in \mathbf{y}_1 and the $p - m$ deleted variables in \mathbf{y}_2 .

Ideally, these criteria would be evaluated for all possible subsets so as to obtain the best subset of each size. McCabe suggested a regression approach for obtaining a percent of variance explained by a subset of variables to be compared with the percent of variance accounted for by the same number of principal components.

PROBLEMS

- 12.1** Show that the solutions to $\lambda = \mathbf{a}'\mathbf{S}\mathbf{a}/\mathbf{a}'\mathbf{a}$ in (12.7) are given by the eigenvalues and eigenvectors in (12.8), so that λ in (12.7) is maximized by the largest eigenvalue of \mathbf{S} .

- 12.2** Show that the eigenvalues of

$$\mathbf{R} = \begin{pmatrix} 1 & r \\ r & 1 \end{pmatrix}$$

are $1 \pm r$, as in (12.13), and that the eigenvectors are as given in (12.14).

- 12.3 (a)** Give a justification based on the likelihood ratio for the test statistic u in (12.15).

- (b)** Give a justification for the degrees of freedom $\nu = \frac{1}{2}(k - 1)(k + 2)$ for the test statistic in (12.15).

- 12.4** Show that when \mathbf{S} is diagonal as in (12.16), the eigenvectors have the form $\mathbf{a}'_i = (0, \dots, 0, 1, 0, \dots, 0)$, as given in (12.18).

- 12.5** Show that $r_{y_1 z_1}^2 + r_{y_1 z_2}^2 + \dots + r_{y_1 z_k}^2 = R_{y_1 | z_1, \dots, z_k}^2$, as in (12.19).

- 12.6** Carry out a principal component analysis of the diabetes data of Table 3.5. Use all five variables, including y 's and x 's. Use both \mathbf{S} and \mathbf{R} . Which do you think is more appropriate here? Show the percent of variance explained. Based on the average eigenvalue or a scree plot, decide how many components to retain. Can you interpret the components of either \mathbf{S} or \mathbf{R} ?

- 12.7** Do a principal component analysis of the probe word data of Table 3.6. Use both \mathbf{S} and \mathbf{R} . Which do you think is more appropriate here? Show the percent of variance explained. Based on the average eigenvalue or a scree plot, decide how many components to retain. Can you interpret the components of either \mathbf{S} or \mathbf{R} ?

- 12.8** Carry out a principal component analysis on all six variables of the glucose data of Table 3.9. Use both \mathbf{S} and \mathbf{R} . Which do you think is more appropriate here? Show the percent of variance explained. Based on the average

eigenvalue or a scree plot, decide how many components to retain. Can you interpret the components of either \mathbf{S} or \mathbf{R} ?

- 12.9** Carry out a principal component analysis on the hematology data of Table 4.2. Use both \mathbf{S} and \mathbf{R} . Which do you think is more appropriate here? Show the percent of variance explained. Based on the average eigenvalue or a scree plot, decide how many components to retain. Can you interpret the components of either \mathbf{S} or \mathbf{R} ? Does the large variance of y_3 affect the pattern of the components of \mathbf{S} ?
- 12.10** Carry out a principal component analysis separately for males and females in the psychological data of Table 5.1. Compare the results for the two groups. Use \mathbf{S} .
- 12.11** Carry out a principal component analysis separately for the two species in the beetle data of Table 5.5. Compare the results for the two groups. Use \mathbf{S} .
- 12.12** Carry out a principal component analysis on the engineer data of Table 5.6 as follows:
- (a) Use the pooled covariance matrix.
 - (b) Ignore groups and use a covariance matrix based on all 40 observations.
 - (c) Which of the approaches in (a) or (b) appears to be more successful?
- 12.13** Repeat the previous problem for the dystrophy data of Table 5.7.
- 12.14** Carry out a principal component analysis on all 10 variables of the Seishu data of Table 7.1. Use both \mathbf{S} and \mathbf{R} . Which do you think is more appropriate here? Show the percent of variance explained. Based on the average eigenvalue or a scree plot, decide how many components to retain. Can you interpret the components of either \mathbf{S} or \mathbf{R} ?
- 12.15** Carry out a principal component analysis on the temperature data of Table 7.2. Use both \mathbf{S} and \mathbf{R} . Which do you think is more appropriate here? Show the percent of variance explained. Based on the average eigenvalue or a scree plot, decide how many components to retain. Can you interpret the components of either \mathbf{S} or \mathbf{R} ?

CHAPTER 13

EXPLORATORY FACTOR ANALYSIS

13.1 INTRODUCTION

In factor analysis we represent the variables y_1, y_2, \dots, y_p as linear combinations of a few random variables f_1, f_2, \dots, f_m ($m < p$) called *factors*. The factors are underlying *constructs* or *latent* variables that “generate” the y ’s. Like the original variables, the factors vary from individual to individual; but unlike the variables, the factors cannot be measured or observed. The existence of these hypothetical variables is therefore open to question.

If the original variables y_1, y_2, \dots, y_p are at least moderately correlated, the basic dimensionality of the system is less than p . The goal of factor analysis is to reduce the redundancy among the variables by using a smaller number of factors.

Suppose the pattern of the high and low correlations in the correlation matrix is such that the variables in a particular subset have high correlations among themselves but low correlations with all the other variables. Then there may be a single underlying factor that gave rise to the variables in the subset. If the other variables can be similarly grouped into subsets with a like pattern of correlations, then a few factors can represent these groups of variables. In this case the pattern in the correlation ma-

trix corresponds directly to the factors. For example, suppose the correlation matrix has the form

$$\begin{pmatrix} 1 & .9 & .05 & .05 & .05 \\ .9 & 1 & .05 & .05 & .05 \\ .05 & .05 & 1 & .9 & .9 \\ .05 & .05 & .9 & 1 & .9 \\ .05 & .05 & .9 & .9 & 1 \end{pmatrix}.$$

Then variables 1 and 2 correspond to a factor and variables 3, 4, and 5 correspond to another factor. In some cases where the correlation matrix does not have such a simple pattern, factor analysis will still partition the variables into clusters.

Factor analysis is related to principal component analysis in that both seek a simpler structure in a set of variables, but they differ in many respects (see Section 13.8). For example, two differences in basic approach are as follows:

1. Principal components are defined as linear combinations of the original variables. In factor analysis, the original variables are expressed as linear combinations of the factors.
2. In principal component analysis, we explain a large part of the total variance of the variables, $\sum_i s_{ii}$. In factor analysis, we seek to account for the covariances or correlations among the variables.

Although we have thus far compared principal component analysis with “factor analysis,” we can draw further distinctions between the *exploratory factor analysis* described in this chapter and the *confirmatory factor analysis* that will be discussed in Chapter 14. Exploratory factor analysis (or EFA) is most often used to explore multivariate data to identify possible latent structure. In EFA, the number of latent factors is not determined before the analysis. In contrast, confirmatory factor analysis (or CFA) allows the researcher to hypothesize the number of latent factors and the specific nature of the latent structure in the data, and then test the hypotheses that have been formulated. Thus the objectives and methods of EFA are more closely related to those of principal component analysis, while CFA uses the more traditional statistical notions of model formulation, parameter estimation, model evaluation, and statistical inference. Because of the subjectivity and relative lack of formal statistical inference associated with EFA, it is considered by some to be controversial or of limited utility. Regardless, EFA has proven to be a useful tool in many settings, particularly as a technique for simplifying complex multivariate data and formulating hypotheses that can be later verified by using a CFA on an independently obtained data set (see Gerbing and Hamilton, 1996).

For either type of factor analysis, there will be some data sets for which the factor analysis model does not provide a satisfactory fit. Sometimes a few easily interpretable factors emerge, but for other data sets neither the number of factors nor the interpretation is clear. Some possible reasons for these failures are discussed in Section 13.7.

In this chapter, we describe the methods associated with exploratory factor analysis. Throughout this chapter, we will use “exploratory factor analysis” and “factor analysis” interchangeably.

13.2 ORTHOGONAL FACTOR MODEL

13.2.1 Model Definition and Assumptions

Factor analysis is basically a one-sample procedure [for possible applications to data with groups, see Rencher (1998, Section 10.8)]. We assume a random sample $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n$ from a homogeneous population with mean vector $\boldsymbol{\mu}$ and covariance matrix $\boldsymbol{\Sigma}$.

The factor analysis model expresses each variable as a linear combination of underlying *common factors* f_1, f_2, \dots, f_m , with an accompanying error term to account for that part of the variable that is unique (not in common with the other variables). For y_1, y_2, \dots, y_p in any observation vector \mathbf{y} , the model is as follows:

$$\begin{aligned} y_1 - \mu_1 &= \lambda_{11}f_1 + \lambda_{12}f_2 + \cdots + \lambda_{1m}f_m + \varepsilon_1 \\ y_2 - \mu_2 &= \lambda_{21}f_1 + \lambda_{22}f_2 + \cdots + \lambda_{2m}f_m + \varepsilon_2 \\ &\vdots \\ y_p - \mu_p &= \lambda_{p1}f_1 + \lambda_{p2}f_2 + \cdots + \lambda_{pm}f_m + \varepsilon_p. \end{aligned} \tag{13.1}$$

Ideally, m should be substantially smaller than p ; otherwise we have not achieved a parsimonious description of the variables as functions of a few underlying factors. We might regard the f 's in (13.1) as random variables that engender the y 's. The coefficients λ_{ij} are called *loadings* and serve as weights, showing how each y_i individually depends on the f 's. (In this chapter, we defer to common usage in the factor analysis literature and use the notation λ_{ij} for loadings rather than eigenvalues.) With appropriate assumptions, λ_{ij} indicates the importance of the j th factor f_j to the i th variable y_i and can be used in interpretation of f_j . We describe or interpret f_2 , for example, by examining its coefficients, $\lambda_{12}, \lambda_{22}, \dots, \lambda_{p2}$. The larger loadings relate f_2 to the corresponding y 's. From these y 's, we infer a meaning or description of f_2 . After estimating the λ_{ij} 's (and rotating them, see Sections 13.2.2 and 13.5), they will hopefully partition the variables into groups corresponding to factors.

The system of equations (13.1) bears a superficial resemblance to the multiple regression model (10.1), but there are fundamental differences. For example, (1) the f 's are unobserved and (2) the model in (13.1) represents only one observation vector, while (10.1) depicts all n observations.

It is assumed that for $j = 1, 2, \dots, m$, $E(f_j) = 0$, $\text{var}(f_j) = 1$, and $\text{cov}(f_j, f_k) = 0$, $j \neq k$. The assumptions for ε_i , $i = 1, 2, \dots, p$, are similar, except that we must allow each ε_i to have a different variance, since it shows the residual part of y_i that is not in common with the other variables. Thus we assume that $E(\varepsilon_i) = 0$, $\text{var}(\varepsilon_i) = \psi_i$, and $\text{cov}(\varepsilon_i, \varepsilon_k) = 0$, $i \neq k$. In addition, we assume that $\text{cov}(\varepsilon_i, f_j) = 0$ for all i and j . We refer to ψ_i as the *specific variance*.

These assumptions are natural consequences of the basic model (13.1) and the goals of factor analysis. Since $E(y_i - \mu_i) = 0$, we need $E(f_j) = 0$, $j = 1, 2, \dots, m$. The assumption $\text{cov}(f_j, f_k) = 0$ is made for parsimony in expressing the y 's as functions of as few factors as possible. The assumptions $\text{var}(f_j) = 1$, $\text{var}(\varepsilon_i) = \psi_i$, $\text{cov}(f_j, f_k) = 0$, and $\text{cov}(\varepsilon_i, f_j) = 0$ yield a simple expression for the variance of y_i ,

$$\text{var}(y_i) = \lambda_{i1}^2 + \lambda_{i2}^2 + \dots + \lambda_{im}^2 + \psi_i, \quad (13.2)$$

which plays an important role in our development. The assumption $\text{cov}(\varepsilon_i, \varepsilon_k) = 0$ implies that the factors account for all the correlations among the y 's, that is, all that the y 's have in common. Thus the emphasis in factor analysis is on modeling the covariances or correlations among the y 's.

Model (13.1) can be written in matrix notation as

$$\mathbf{y} - \boldsymbol{\mu} = \mathbf{\Lambda}\mathbf{f} + \boldsymbol{\varepsilon}, \quad (13.3)$$

where $\mathbf{y} = (y_1, y_2, \dots, y_p)'$, $\boldsymbol{\mu} = (\mu_1, \mu_2, \dots, \mu_p)'$, $\mathbf{f} = (f_1, f_2, \dots, f_m)'$, $\boldsymbol{\varepsilon} = (\varepsilon_1, \varepsilon_2, \dots, \varepsilon_p)'$, and

$$\mathbf{\Lambda} = \begin{pmatrix} \lambda_{11} & \lambda_{12} & \cdots & \lambda_{1m} \\ \lambda_{21} & \lambda_{22} & \cdots & \lambda_{2m} \\ \vdots & \vdots & & \vdots \\ \lambda_{p1} & \lambda_{p2} & \cdots & \lambda_{pm} \end{pmatrix}. \quad (13.4)$$

We illustrate the model in (13.1) and (13.3) with $p = 5$ and $m = 2$. The model for each variable in (13.1) becomes

$$\begin{aligned} y_1 - \mu_1 &= \lambda_{11}f_1 + \lambda_{12}f_2 + \varepsilon_1 \\ y_2 - \mu_2 &= \lambda_{21}f_1 + \lambda_{22}f_2 + \varepsilon_2 \\ y_3 - \mu_3 &= \lambda_{31}f_1 + \lambda_{32}f_2 + \varepsilon_3 \\ y_4 - \mu_4 &= \lambda_{41}f_1 + \lambda_{42}f_2 + \varepsilon_4 \\ y_5 - \mu_5 &= \lambda_{51}f_1 + \lambda_{52}f_2 + \varepsilon_5. \end{aligned}$$

In matrix notation as in (13.3), this becomes

$$\begin{pmatrix} y_1 - \mu_1 \\ y_2 - \mu_2 \\ y_3 - \mu_3 \\ y_4 - \mu_4 \\ y_5 - \mu_5 \end{pmatrix} = \begin{pmatrix} \lambda_{11} & \lambda_{12} \\ \lambda_{21} & \lambda_{22} \\ \lambda_{31} & \lambda_{32} \\ \lambda_{41} & \lambda_{42} \\ \lambda_{51} & \lambda_{52} \end{pmatrix} \begin{pmatrix} f_1 \\ f_2 \end{pmatrix} + \begin{pmatrix} \varepsilon_1 \\ \varepsilon_2 \\ \varepsilon_3 \\ \varepsilon_4 \\ \varepsilon_5 \end{pmatrix}, \quad (13.5)$$

or $\mathbf{y} - \boldsymbol{\mu} = \mathbf{\Lambda}\mathbf{f} + \boldsymbol{\varepsilon}$.

The assumptions described above between (13.1) and (13.2) can be expressed concisely using vector and matrix notation:

$E(f_j) = 0$, $j = 1, 2, \dots, m$, becomes

$$E(\mathbf{f}) = \mathbf{0}, \quad (13.6)$$

$\text{var}(f_j) = 1, j = 1, 2, \dots, m$, and $\text{cov}(f_j, f_k) = 0, j \neq k$, become

$$\text{cov}(\mathbf{f}) = \mathbf{I}, \quad (13.7)$$

$E(\varepsilon_i) = 0, i = 1, 2, \dots, p$, becomes

$$E(\boldsymbol{\varepsilon}) = \mathbf{0}, \quad (13.8)$$

$\text{var}(\varepsilon_i) = \psi_i, i = 1, 2, \dots, p$, and $\text{cov}(\varepsilon_i, \varepsilon_k) = 0, i \neq k$, become

$$\text{cov}(\boldsymbol{\varepsilon}) = \boldsymbol{\Psi} = \begin{pmatrix} \psi_1 & 0 & \cdots & 0 \\ 0 & \psi_2 & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & \psi_p \end{pmatrix}, \quad (13.9)$$

and $\text{cov}(\varepsilon_i, f_j) = 0$ for all i and j becomes

$$\text{cov}(\mathbf{f}, \boldsymbol{\varepsilon}) = \mathbf{O}. \quad (13.10)$$

The notation $\text{cov}(\mathbf{f}, \boldsymbol{\varepsilon})$ indicates a rectangular matrix containing the covariances of the f 's with the ε 's:

$$\text{cov}(\mathbf{f}, \boldsymbol{\varepsilon}) = \begin{pmatrix} \sigma_{f_1\varepsilon_1} & \sigma_{f_1\varepsilon_2} & \cdots & \sigma_{f_1\varepsilon_p} \\ \sigma_{f_2\varepsilon_1} & \sigma_{f_2\varepsilon_2} & \cdots & \sigma_{f_2\varepsilon_p} \\ \vdots & \vdots & & \vdots \\ \sigma_{f_m\varepsilon_1} & \sigma_{f_m\varepsilon_2} & \cdots & \sigma_{f_m\varepsilon_p} \end{pmatrix}.$$

It was noted following (13.2) that the emphasis in factor analysis is on modeling the covariances among the y 's. We wish to express the $\frac{1}{2}p(p-1)$ covariances (and the p variances) of the variables y_1, y_2, \dots, y_p in terms of a simplified structure involving the pm loadings λ_{ij} and the p specific variances ψ_i ; that is, we wish to express Σ in terms of Λ and $\boldsymbol{\Psi}$. We can do this using the model (13.3) and the assumptions (13.7), (13.9), and (13.10). Since μ does not affect variances and covariances of \mathbf{y} , we have from (13.3)

$$\boldsymbol{\Sigma} = \text{cov}(\mathbf{y}) = \text{cov}(\Lambda\mathbf{f} + \boldsymbol{\varepsilon}).$$

By (13.10), $\Lambda\mathbf{f}$ and $\boldsymbol{\varepsilon}$ are uncorrelated; therefore, the covariance matrix of their sum is the sum of their covariance matrices:

$$\begin{aligned} \boldsymbol{\Sigma} &= \text{cov}(\Lambda\mathbf{f}) + \text{cov}(\boldsymbol{\varepsilon}) \\ &= \Lambda \text{cov}(\mathbf{f}) \Lambda' + \boldsymbol{\Psi} \quad [\text{by (3.74) and (13.9)}] \\ &= \Lambda \Lambda' + \boldsymbol{\Psi} \quad [\text{by (13.7)}] \\ &= \Lambda \Lambda' + \boldsymbol{\Psi}. \end{aligned} \quad (13.11)$$

If Λ has only a few columns, say two or three, then $\boldsymbol{\Sigma} = \Lambda \Lambda' + \boldsymbol{\Psi}$ in (13.11) represents a simplified structure for $\boldsymbol{\Sigma}$, in which the covariances are modeled by the

λ_{ij} 's alone since Ψ is diagonal. For example, in the above illustration in (13.5) with $m = 2$ factors, σ_{12} would be the product of the first two rows of Λ , that is,

$$\sigma_{12} = \text{cov}(y_1, y_2) = \lambda_{11}\lambda_{21} + \lambda_{12}\lambda_{22},$$

where $(\lambda_{11}, \lambda_{12})$ is the first row of Λ and $(\lambda_{21}, \lambda_{22})$ is the second row of Λ . If y_1 and y_2 have a great deal in common, they will have similar loadings on the common factors f_1 and f_2 ; that is, $(\lambda_{11}, \lambda_{12})$ will be similar to $(\lambda_{21}, \lambda_{22})$. In this case, either $\lambda_{11}\lambda_{21}$ or $\lambda_{12}\lambda_{22}$ is likely to be high. On the other hand, if y_1 and y_2 have little in common, then their loadings λ_{11} and λ_{21} on f_1 will be different and their loadings λ_{12} and λ_{22} on f_2 will likewise differ. In this case, the products $\lambda_{11}\lambda_{21}$ and $\lambda_{12}\lambda_{22}$ will tend to be small.

We can also find the covariances of the y 's with the f 's in terms of the λ 's. Consider, for example, $\text{cov}(y_1, f_2)$. By (13.1), $y_1 - \mu_1 = \lambda_{11}f_1 + \lambda_{12}f_2 + \dots + \lambda_{1m}f_m + \varepsilon_1$. From (13.7), f_2 is uncorrelated with all other f_j 's, and by (13.10), f_2 is uncorrelated with ε_1 . Thus

$$\begin{aligned} \text{cov}(y_1, f_2) &= E[(y_1 - \mu_1)(f_2 - \mu_{f_2})] \\ &= E[(\lambda_{11}f_1 + \lambda_{12}f_2 + \dots + \lambda_{1m}f_m)f_2] \\ &= E(\lambda_{11}f_1f_2 + \lambda_{12}f_2^2 + \dots + \lambda_{1m}f_mf_2) \\ &= \lambda_{11}\text{cov}(f_1, f_2) + \lambda_{12}\text{var}(f_2) + \dots + \lambda_{1m}\text{cov}(f_m, f_2) \\ &= \lambda_{12}, \end{aligned}$$

since $\text{var}(f_2) = 1$. Hence the loadings themselves represent covariances of the variables with the factors. In general,

$$\text{cov}(y_i, f_j) = \lambda_{ij}, \quad i = 1, 2, \dots, p, \quad j = 1, 2, \dots, m. \quad (13.12)$$

Since λ_{ij} is the (ij) th element of Λ , we can write (13.12) in the form

$$\text{cov}(\mathbf{y}, \mathbf{f}) = \Lambda. \quad (13.13)$$

If standardized variables are used, (13.11) is replaced by $\mathbf{P}_\rho = \Lambda\Lambda' + \Psi$, and the loadings become correlations:

$$\text{corr}(y_i, f_j) = \lambda_{ij}. \quad (13.14)$$

In (13.2), we have a partitioning of the variance of y_i into a component due to the common factors, called the *communality*, and a component unique to y_i , called the *specific variance*:

$$\begin{aligned} \sigma_{ii} = \text{var}(y_i) &= (\lambda_{i1}^2 + \lambda_{i2}^2 + \dots + \lambda_{im}^2) + \psi_i \\ &= h_i^2 + \psi_i \\ &= \text{communality} + \text{specific variance}, \end{aligned}$$

where

$$\text{Communality} = h_i^2 = \lambda_{i1}^2 + \lambda_{i2}^2 + \cdots + \lambda_{im}^2, \quad (13.15)$$

Specific variance = ψ_i .

The communality h_i^2 is also referred to as *common variance*, and the specific variance ψ_i has been called *specificity*, *unique variance*, or *residual variance*.

Assumptions (13.6)–(13.10) lead to the simple covariance structure of (13.11), $\Sigma = \Lambda\Lambda' + \Psi$, which is an essential part of the factor analysis model. In schematic form, $\Sigma = \Lambda\Lambda' + \Psi$ has the following appearance:

$$\Sigma = \begin{array}{c} \boxed{} \\ \boxed{} \end{array} + \boxed{}$$

The diagonal elements of Σ can be easily modeled by adjusting the diagonal elements of Ψ , but $\Lambda\Lambda'$ is a simplified configuration for the off-diagonal elements. Hence the critical aspect of the model involves the covariances, and this is the major emphasis of factor analysis, as noted in Section 13.1 and in comments following (13.2) and (13.10).

It is a rare population covariance matrix Σ that can be expressed exactly as $\Sigma = \Lambda\Lambda' + \Psi$, where Ψ is diagonal and Λ is $p \times m$, with m relatively small. In practice, many sample covariance matrices do not come satisfactorily close to this ideal pattern. However, we do not relax the assumptions because the structure $\Sigma = \Lambda\Lambda' + \Psi$ is essential for estimation of Λ .

One advantage of the factor analysis model is that when it does not fit the data, the estimate of Λ clearly reflects this failure. In such cases, there are two problems in the estimates: (1) it is unclear how many factors there should be, and (2) it is unclear what the factors are. In other statistical procedures, failure of assumptions may not lead to such obvious consequences in the estimates or tests. In factor analysis, the assumptions are essentially self-checking, whereas in other procedures, we typically have to check the assumptions with residual plots, tests, and so on.

13.2.2 Nonuniqueness of Factor Loadings

The loadings in the model (13.3) can be multiplied by an orthogonal matrix without impairing their ability to reproduce the covariance matrix in $\Sigma = \Lambda\Lambda' + \Psi$. To see this, let \mathbf{T} be an arbitrary orthogonal matrix. Then by (2.102), $\mathbf{T}\mathbf{T}' = \mathbf{I}$, and we can insert $\mathbf{T}\mathbf{T}'$ into the basic model (13.3) to obtain

$$\mathbf{y} - \boldsymbol{\mu} = \Lambda\mathbf{T}\mathbf{T}'\mathbf{f} + \boldsymbol{\varepsilon}.$$

We then associate \mathbf{T} with Λ and associate \mathbf{T}' with \mathbf{f} so that the model becomes

$$\mathbf{y} - \boldsymbol{\mu} = \Lambda^* \mathbf{f}^* + \boldsymbol{\varepsilon}, \quad (13.16)$$

where

$$\Lambda^* = \Lambda \mathbf{T}, \quad (13.17)$$

$$\mathbf{f}^* = \mathbf{T}' \mathbf{f}. \quad (13.18)$$

If Λ in $\Sigma = \Lambda \Lambda' + \Psi$ is replaced by $\Lambda^* = \Lambda \mathbf{T}$, we have

$$\begin{aligned} \Sigma &= \Lambda^* \Lambda^{*''} + \Psi = \Lambda \mathbf{T} (\Lambda \mathbf{T})' + \Psi \\ &= \Lambda \mathbf{T} \mathbf{T}' \Lambda' + \Psi = \Lambda \Lambda' + \Psi, \end{aligned}$$

since $\mathbf{T} \mathbf{T}' = \mathbf{I}$. Thus the new loadings $\Lambda^* = \Lambda \mathbf{T}$ in (13.17) reproduce the covariance matrix, just as Λ does in (13.11):

$$\Sigma = \Lambda^* \Lambda^{*''} + \Psi = \Lambda \Lambda' + \Psi. \quad (13.19)$$

The new factors $\mathbf{f}^* = \mathbf{T}' \mathbf{f}$ in (13.18) satisfy the assumptions (13.6), (13.7), and (13.10); that is, $E(\mathbf{f}^*) = \mathbf{0}$, $\text{cov}(\mathbf{f}^*) = \mathbf{I}$, and $\text{cov}(\mathbf{f}^*, \boldsymbol{\varepsilon}) = \mathbf{O}$.

The communalities $h_i^2 = \lambda_{i1}^2 + \lambda_{i2}^2 + \dots + \lambda_{im}^2$, $i = 1, 2, \dots, p$, as defined in (13.15), are also unaffected by the transformation $\Lambda^* = \Lambda \mathbf{T}$. This can be seen as follows. The communality h_i^2 is the sum of squares of the i th row of Λ . If we denote the i th row of Λ by λ'_i , then the sum of squares in vector notation is $h_i^2 = \lambda'_i \lambda_i$. The i th row of $\Lambda^* = \Lambda \mathbf{T}$ is $\lambda_i^{*''} = \lambda'_i \mathbf{T}$, and the corresponding communality is

$$h_i^{*2} = \lambda_i^{*''} \lambda_i^* = \lambda'_i \mathbf{T} \mathbf{T}' \lambda_i = \lambda'_i \lambda_i = h_i^2.$$

Thus the communalities remain the same for the new loadings. Note that $h_i^2 = \lambda_{i1}^2 + \lambda_{i2}^2 + \dots + \lambda_{im}^2 = \lambda'_i \lambda_i$ is the distance from the origin to the point $\lambda'_i = (\lambda_{i1}, \lambda_{i2}, \dots, \lambda_{im})$ in the m -dimensional space of the factor loadings. Since the distance $\lambda'_i \lambda_i$ is the same as $\lambda_i^{*''} \lambda_i^*$, the points λ_i^* are rotated from the points λ_i . [This also follows because $\lambda_i^{*''} = \lambda'_i \mathbf{T}$, where \mathbf{T} is orthogonal. Multiplication of a vector by an orthogonal matrix is equivalent to a rotation of axes; see (2.103).]

The inherent potential to rotate the loadings to a new frame of reference without affecting any assumptions or properties is very useful in interpretation of the factors and will be exploited in Section 13.5.

Note that the coefficients (loadings) in (13.1) are applied to the factors, not to the variables, as they are in discriminant functions and principal components. Thus in factor analysis the observed variables are not involved in the rotation as they are in discriminant functions and principal components.

13.3 ESTIMATION OF LOADINGS AND COMMUNALITIES

In Sections 13.3.1–13.3.4, we discuss four approaches to estimation of loadings and communalities.

13.3.1 Principal Component Method

The first technique we consider is commonly called the *principal component* method. This name is perhaps unfortunate in that it adds to the confusion between factor analysis and principal component analysis. In the principal component method for estimation of loadings, we do not actually calculate any principal components. The reason for the name is given following (13.25).

From a random sample $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n$, we obtain the sample covariance matrix \mathbf{S} and then attempt to find an estimator $\hat{\Lambda}$ that will approximate the fundamental expression (13.11) with \mathbf{S} in place of Σ :

$$\mathbf{S} \cong \hat{\Lambda} \hat{\Lambda}' + \hat{\Psi}. \quad (13.20)$$

In the principal component approach, we neglect $\hat{\Psi}$ and factor \mathbf{S} into $\mathbf{S} = \hat{\Lambda} \hat{\Lambda}'$.

In order to factor \mathbf{S} , we use the spectral decomposition in (2.109),

$$\mathbf{S} = \mathbf{C} \mathbf{D} \mathbf{C}', \quad (13.21)$$

where \mathbf{C} is an orthogonal matrix constructed with normalized eigenvectors ($\mathbf{c}_i' \mathbf{c}_i = 1$) of \mathbf{S} as columns and \mathbf{D} is a diagonal matrix with the eigenvalues $\theta_1, \theta_2, \dots, \theta_p$ of \mathbf{S} on the diagonal:

$$\mathbf{D} = \begin{pmatrix} \theta_1 & 0 & \cdots & 0 \\ 0 & \theta_2 & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & \theta_p \end{pmatrix}. \quad (13.22)$$

We use the notation θ_i for eigenvalues instead of the usual λ_i in order to avoid confusion with the notation λ_{ij} used for the loadings.

To finish factoring $\mathbf{C} \mathbf{D} \mathbf{C}'$ in (13.21) into the form $\hat{\Lambda} \hat{\Lambda}'$, we observe that since the eigenvalues θ_i of the positive semidefinite matrix \mathbf{S} are all positive or zero, we can factor \mathbf{D} into

$$\mathbf{D} = \mathbf{D}^{1/2} \mathbf{D}^{1/2},$$

where

$$\mathbf{D}^{1/2} = \begin{pmatrix} \sqrt{\theta_1} & 0 & \cdots & 0 \\ 0 & \sqrt{\theta_2} & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & \sqrt{\theta_p} \end{pmatrix}.$$

With this factoring of \mathbf{D} , (13.21) becomes,

$$\begin{aligned} \mathbf{S} &= \mathbf{C} \mathbf{D} \mathbf{C}' = \mathbf{C} \mathbf{D}^{1/2} \mathbf{D}^{1/2} \mathbf{C}' \\ &= (\mathbf{C} \mathbf{D}^{1/2})(\mathbf{C} \mathbf{D}^{1/2})'. \end{aligned} \quad (13.23)$$

This is of the form $\mathbf{S} = \hat{\Lambda} \hat{\Lambda}'$, but we do not define $\hat{\Lambda}$ to be $\mathbf{C} \mathbf{D}^{1/2}$ because $\mathbf{C} \mathbf{D}^{1/2}$ is $p \times p$, and we are seeking a $\hat{\Lambda}$ that is $p \times m$ with $m < p$. We therefore define

$\mathbf{D}_1 = \text{diag}(\theta_1, \theta_2, \dots, \theta_m)$ with the m largest eigenvalues $\theta_1 > \theta_2 > \dots > \theta_m$ and $\mathbf{C}_1 = (\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_m)$ containing the corresponding eigenvectors. We then estimate Λ by the first m columns of $\mathbf{CD}_1^{1/2}$,

$$\hat{\Lambda} = \mathbf{C}_1 \mathbf{D}_1^{1/2} = (\sqrt{\theta_1} \mathbf{c}_1, \sqrt{\theta_2} \mathbf{c}_2, \dots, \sqrt{\theta_m} \mathbf{c}_m) \quad (13.24)$$

by (2.56), where $\hat{\Lambda}$ is $p \times m$, \mathbf{C}_1 is $p \times m$, and $\mathbf{D}_1^{1/2}$ is $m \times m$.

We illustrate the structure of the $\hat{\lambda}_{ij}$ in (13.24) for $p = 5$ and $m = 2$:

$$\begin{aligned} \begin{pmatrix} \hat{\lambda}_{11} & \hat{\lambda}_{12} \\ \hat{\lambda}_{21} & \hat{\lambda}_{22} \\ \hat{\lambda}_{31} & \hat{\lambda}_{32} \\ \hat{\lambda}_{41} & \hat{\lambda}_{42} \\ \hat{\lambda}_{51} & \hat{\lambda}_{52} \end{pmatrix} &= \begin{pmatrix} c_{11} & c_{12} \\ c_{21} & c_{22} \\ c_{31} & c_{32} \\ c_{41} & c_{42} \\ c_{51} & c_{52} \end{pmatrix} \begin{pmatrix} \sqrt{\theta_1} & 0 \\ 0 & \sqrt{\theta_2} \end{pmatrix} \\ &= \begin{pmatrix} \sqrt{\theta_1} c_{11} & \sqrt{\theta_2} c_{12} \\ \sqrt{\theta_1} c_{21} & \sqrt{\theta_2} c_{22} \\ \sqrt{\theta_1} c_{31} & \sqrt{\theta_2} c_{32} \\ \sqrt{\theta_1} c_{41} & \sqrt{\theta_2} c_{42} \\ \sqrt{\theta_1} c_{51} & \sqrt{\theta_2} c_{52} \end{pmatrix} \quad [\text{by (2.56)}]. \end{aligned} \quad (13.25)$$

We can see in (13.25) the source of the term *principal component* solution. The columns of $\hat{\Lambda}$ are proportional to the eigenvectors of \mathbf{S} , so that the loadings on the j th factor are proportional to coefficients in the j th principal component. The factors are thus related to the first m principal components, and it would seem that interpretation would be the same as for principal components. But after rotation of the loadings, the interpretation of the factors is usually different. The researcher will ordinarily prefer the rotated factors for reasons to be treated in Section 13.5.

By (2.52), the i th diagonal element of $\hat{\Lambda}\hat{\Lambda}'$ is the sum of squares of the i th row of $\hat{\Lambda}$, or $\hat{\lambda}_i'\hat{\lambda}_i = \sum_{j=1}^m \hat{\lambda}_{ij}^2$. Hence to complete the approximation of \mathbf{S} in (13.20), we define

$$\hat{\psi}_i = s_{ii} - \sum_{j=1}^m \hat{\lambda}_{ij}^2 \quad (13.26)$$

and write

$$\mathbf{S} \cong \hat{\Lambda}\hat{\Lambda}' + \hat{\Psi}, \quad (13.27)$$

where $\hat{\Psi} = \text{diag}(\hat{\psi}_1, \hat{\psi}_2, \dots, \hat{\psi}_p)$. Thus in (13.27) the variances on the diagonal of \mathbf{S} are modeled exactly, but the off-diagonal covariances are only approximate. Again, this is the challenge of factor analysis.

In this method of estimation, the sums of squares of the rows and columns of $\hat{\Lambda}$ are equal to communalities and eigenvalues, respectively. This is easily shown. By (13.26) and by analogy with (13.15), the i th communality is estimated by

$$\hat{h}_i^2 = \sum_{j=1}^m \hat{\lambda}_{ij}^2, \quad (13.28)$$

which is the sum of squares of the i th row of $\hat{\Lambda}$. The sum of squares of the j th column of $\hat{\Lambda}$ is the j th eigenvalue of \mathbf{S} :

$$\begin{aligned} \sum_{i=1}^p \hat{\lambda}_{ij}^2 &= \sum_{i=1}^p (\sqrt{\theta_j} c_{ij})^2 && [\text{by (13.25)}] \\ &= \theta_j \sum_{i=1}^p c_{ij}^2 \\ &= \theta_j, \end{aligned} \quad (13.29)$$

since the normalized eigenvectors (columns of \mathbf{C}) have length 1.

By (13.26) and (13.28), the variance of the i th variable is partitioned into a part due to the factors and a part due uniquely to the variable:

$$\begin{aligned} s_{ii} &= \hat{h}_i^2 + \hat{\psi}_i \\ &= \hat{\lambda}_{i1}^2 + \hat{\lambda}_{i2}^2 + \cdots + \hat{\lambda}_{im}^2 + \hat{\psi}_i. \end{aligned} \quad (13.30)$$

Thus the j th factor contributes $\hat{\lambda}_{ij}^2$ to s_{ii} . The contribution of the j th factor to the total sample variance, $\text{tr}(\mathbf{S}) = s_{11} + s_{22} + \cdots + s_{pp}$, is therefore

$$\text{variance due to } j\text{th factor} = \sum_{i=1}^p \hat{\lambda}_{ij}^2 = \hat{\lambda}_{1j}^2 + \hat{\lambda}_{2j}^2 + \cdots + \hat{\lambda}_{pj}^2, \quad (13.31)$$

which is the sum of squares of loadings in the j th column of $\hat{\Lambda}$. By (13.29), this is equal to the j th eigenvalue, θ_j . The proportion of total sample variance due to the j th factor is

$$\frac{\sum_{i=1}^p \hat{\lambda}_{ij}^2}{\text{tr}(\mathbf{S})} = \frac{\theta_j}{\text{tr}(\mathbf{S})}. \quad (13.32)$$

If the variables are not commensurate, we can use standardized variables and work with the correlation matrix \mathbf{R} . The eigenvalues and eigenvectors of \mathbf{R} are then used in place of those of \mathbf{S} in (13.24) to obtain estimates of the loadings. In practice, \mathbf{R} is used more often than \mathbf{S} and is the default in most software packages. Since the emphasis in factor analysis is on reproducing the covariances or correlations rather than the variances, use of \mathbf{R} is more appropriate in factor analysis than in principal components. In applications, \mathbf{R} often gives better results than \mathbf{S} .

If we are factoring \mathbf{R} , the proportion corresponding to (13.32) is

$$\frac{\sum_{i=1}^p \hat{\lambda}_{ij}^2}{\text{tr}(\mathbf{R})} = \frac{\theta_j}{p}, \quad (13.33)$$

where p is the number of variables.

We can assess the fit of the factor analysis model by comparing the left and right sides of (13.27). The error matrix

$$\mathbf{E} = \mathbf{S} - (\hat{\Lambda}\hat{\Lambda}' + \hat{\Psi})$$

Table 13.1 Perception Data: Ratings on Five Adjectives for Seven People

People	Kind	Intelligent	Happy	Likeable	Just
FSM1 ^a	1	5	5	1	1
SISTER	8	9	7	9	8
FSM2	9	8	9	9	8
FATHER	9	9	9	9	9
TEACHER	1	9	1	1	9
MSM ^b	9	7	7	9	9
FSM3	9	7	9	9	7

^aFemale schoolmate 1.

^bMale schoolmate.

has zeros on the diagonal but nonzero off-diagonal elements. The following inequality gives a bound on the size of the elements in $\mathbf{E} = \mathbf{S} - (\hat{\Lambda}\hat{\Lambda}' + \hat{\Psi})$:

$$\sum_{ij} e_{ij}^2 \leq \theta_{m+1}^2 + \theta_{m+2}^2 + \cdots + \theta_p^2; \quad (13.34)$$

that is, the sum of squared entries in the matrix $\mathbf{E} = \mathbf{S} - (\hat{\Lambda}\hat{\Lambda}' + \hat{\Psi})$ is at most equal to the sum of squares of the deleted eigenvalues of \mathbf{S} . If the eigenvalues are small, the residuals in the error matrix $\mathbf{S} - (\hat{\Lambda}\hat{\Lambda}' + \hat{\Psi})$ are small and the fit is good.

■ EXAMPLE 13.3.1

To illustrate the principal component method of estimation, we use a simple data set collected by Brown et al. (1984). A 12-year-old girl made five ratings on a nine-point semantic differential scale for each of seven of her acquaintances. The ratings were based on the five adjectives “kind,” “intelligent,” “happy,” “likeable,” and “just.” Her ratings are given in Table 13.1.

The correlation matrix for the five variables (adjectives) is as follows, with the larger values bolded:

$$\mathbf{R} = \begin{pmatrix} 1.000 & .296 & \mathbf{.881} & \mathbf{.995} & .545 \\ .296 & 1.000 & -.022 & .326 & \mathbf{.837} \\ \mathbf{.881} & -.022 & 1.000 & \mathbf{.867} & .130 \\ \mathbf{.995} & .326 & \mathbf{.867} & 1.000 & .544 \\ .545 & \mathbf{.837} & .130 & .544 & 1.000 \end{pmatrix}. \quad (13.35)$$

The boldface values indicate two groups of variables: {1, 3, 4} and {2, 5}. We would therefore expect that the correlations among the variables can be explained fairly well by two factors.

The eigenvalues of \mathbf{R} are 3.263, 1.538, .168, .031, and 0. Thus \mathbf{R} is singular, which is possible in a situation such as this with only seven observations on five variables recorded in a single-digit scale. The multicollinearity among

Table 13.2 Factor Loadings by the Principal Component Method for the Perception Data of Table 13.1

Variables	Loadings		Communalities, \hat{h}_i^2	Specific Variances, $\hat{\psi}_i$
	$\hat{\lambda}_{1j}$	$\hat{\lambda}_{2j}$		
Kind	.969	-.231	.993	.007
Intelligent	.519	.807	.921	.079
Happy	.785	-.587	.960	.040
Likeable	.971	-.210	.987	.013
Just	.704	.667	.940	.060
Variance accounted for	3.263	1.538	4.802	
Proportion of total variance	.653	.308	.960	
Cumulative proportion	.653	.960	.960	

the variables induced by the fifth eigenvalue, 0, could be ascertained from the corresponding eigenvector, as noted in Section 12.7 (see Problem 13.6).

By (13.33), the first two factors account for $(3.263 + 1.538)/5 = .96$ of the total sample variance. We therefore extract two factors. The first two eigenvectors are

$$\mathbf{c}_1 = \begin{pmatrix} .537 \\ .288 \\ .434 \\ .537 \\ .390 \end{pmatrix} \quad \text{and} \quad \mathbf{c}_2 = \begin{pmatrix} -.186 \\ .651 \\ -.473 \\ -.169 \\ .538 \end{pmatrix}.$$

When these are multiplied by the square roots of the respective eigenvalues 3.263 and 1.538 as in (13.25), we obtain the loadings in Table 13.2.

The communalities in Table 13.2 are obtained from the sum of squares of the rows of the loadings, as in (13.28). The first one, for example, is $(.969)^2 + (-.231)^2 = .993$. The specific variances are obtained from (13.26) as $\hat{\psi}_i = 1 - \hat{h}_i^2$ using 1 in place of s_{ii} because we are factoring \mathbf{R} rather than \mathbf{S} . The variance accounted for by each factor is the sum of squares of the corresponding column of the loadings, as in (13.31). By (13.29), the variance accounted for is also equal to the eigenvalue in each case. Note that the variance accounted for by the two factors adds to the sum of the communalities, since the latter is the sum of all squared loadings. By (13.33), the proportion of total variance for each factor is the variance accounted for divided by 5.

The two factors account for 96% of the total variance and therefore represent the five variables very well. To see how well the two-factor model reproduces the correlation matrix, we examine

$$\begin{aligned}\hat{\Lambda}\hat{\Lambda}' + \hat{\Psi} &= \begin{pmatrix} .969 & -.231 \\ .519 & .807 \\ .785 & -.587 \\ .971 & -.210 \\ .704 & .667 \end{pmatrix} \begin{pmatrix} .969 & .519 & .785 & .971 & .704 \\ -.231 & .807 & -.587 & -.210 & .667 \end{pmatrix} \\ &\quad + \begin{pmatrix} .007 & 0 & 0 & 0 & 0 \\ 0 & .079 & 0 & 0 & 0 \\ 0 & 0 & .040 & 0 & 0 \\ 0 & 0 & 0 & .013 & 0 \\ 0 & 0 & 0 & 0 & .060 \end{pmatrix} \\ &= \begin{pmatrix} 1.000 & .317 & .896 & .990 & .528 \\ .317 & 1.000 & -.066 & .335 & .904 \\ .896 & -.066 & 1.000 & .885 & .161 \\ .990 & .335 & .885 & 1.000 & .543 \\ .528 & .904 & .161 & .543 & 1.000 \end{pmatrix},\end{aligned}$$

which is very close to the original \mathbf{R} . We will not attempt to interpret the factors at this point but will wait until they have been rotated in Section 13.5.2. \square

13.3.2 Principal Factor Method

In the principal component approach to estimation of the loadings, we neglected Ψ and factored \mathbf{S} or \mathbf{R} . The *principal factor* method (also called the *principal axis* method) uses an initial estimate $\hat{\Psi}$ and factors $\mathbf{S} - \hat{\Psi}$ or $\mathbf{R} - \hat{\Psi}$ to obtain

$$\mathbf{S} - \hat{\Psi} \cong \hat{\Lambda}\hat{\Lambda}', \tag{13.36}$$

$$\mathbf{R} - \hat{\Psi} \cong \hat{\Lambda}\hat{\Lambda}', \tag{13.37}$$

where $\hat{\Lambda}$ is $p \times m$ and is calculated as in (13.24) using eigenvalues and eigenvectors of $\mathbf{S} - \hat{\Psi}$ or $\mathbf{R} - \hat{\Psi}$.

The i th diagonal element of $\mathbf{S} - \hat{\Psi}$ is given by $s_{ii} - \hat{\psi}_i$, which is the i th communality, $\hat{h}_i^2 = s_{ii} - \hat{\psi}_i$ [see (13.30)]. Likewise, the diagonal elements of $\mathbf{R} - \hat{\Psi}$ are the communalities $\hat{h}_i^2 = 1 - \hat{\psi}_i$. (Obviously, $\hat{\psi}_i$ and \hat{h}_i^2 have different values for \mathbf{S}

than for \mathbf{R} .) With these diagonal values, $\mathbf{S} - \hat{\Psi}$ and $\mathbf{R} - \hat{\Psi}$ have the form

$$\mathbf{S} - \hat{\Psi} = \begin{pmatrix} \hat{h}_1^2 & s_{12} & \cdots & s_{1p} \\ s_{21} & \hat{h}_2^2 & \cdots & s_{2p} \\ \vdots & \vdots & \ddots & \vdots \\ s_{p1} & s_{p2} & \cdots & \hat{h}_p^2 \end{pmatrix}, \quad (13.38)$$

$$\mathbf{R} - \hat{\Psi} = \begin{pmatrix} \hat{h}_1^2 & r_{12} & \cdots & r_{1p} \\ r_{21} & \hat{h}_2^2 & \cdots & r_{2p} \\ \vdots & \vdots & \ddots & \vdots \\ r_{p1} & r_{p2} & \cdots & \hat{h}_p^2 \end{pmatrix}. \quad (13.39)$$

A popular initial estimate for a communality in $\mathbf{R} - \hat{\Psi}$ is $\hat{h}_i^2 = R_i^2$, the squared multiple correlation between y_i and the other $p - 1$ variables. This can be found as

$$\hat{h}_i^2 = R_i^2 = 1 - \frac{1}{r^{ii}}, \quad (13.40)$$

where r^{ii} is the i th diagonal element of \mathbf{R}^{-1} .

For $\mathbf{S} - \hat{\Psi}$, an initial estimate of communality analogous to (13.40) is

$$\hat{h}_i^2 = s_{ii} - \frac{1}{s^{ii}}, \quad (13.41)$$

where s_{ii} is the i th diagonal element of \mathbf{S} and s^{ii} is the i th diagonal element of \mathbf{S}^{-1} . It can be shown that (13.41) is equivalent to

$$\hat{h}_i^2 = s_{ii} - \frac{1}{s^{ii}} = s_{ii}R_i^2, \quad (13.42)$$

which is a reasonable estimate of the amount of variance that y_i has in common with the other y 's.

To use (13.40) or (13.41), \mathbf{R} or \mathbf{S} must be nonsingular. If \mathbf{R} is singular we can use the absolute value or the square of the largest correlation in the i th row of \mathbf{R} as an estimate of communality.

After obtaining communality estimates, we calculate eigenvalues and eigenvectors of $\mathbf{S} - \hat{\Psi}$ or $\mathbf{R} - \hat{\Psi}$ and use (13.24) to obtain estimates of factor loadings, $\hat{\Lambda}$. Then the columns and rows of $\hat{\Lambda}$ can be used to obtain new eigenvalues (variance explained) and communalities, respectively. The sum of squares of the j th column of $\hat{\Lambda}$ is the j th eigenvalue of $\mathbf{S} - \hat{\Psi}$ or $\mathbf{R} - \hat{\Psi}$, and the sum of squares of the i th row of $\hat{\Lambda}$ is the communality of y_i . The proportion of variance explained by the j th factor is

$$\frac{\theta_j}{\text{tr}(\mathbf{S} - \hat{\Psi})} = \frac{\theta_j}{\sum_{i=1}^p \theta_i}$$

or

$$\frac{\theta_j}{\text{tr}(\mathbf{R} - \hat{\Psi})} = \frac{\theta_j}{\sum_{i=1}^p \theta_i},$$

where θ_j is the j th eigenvalue of $\mathbf{S} - \hat{\Psi}$ or $\mathbf{R} - \hat{\Psi}$. The matrices $\mathbf{S} - \hat{\Psi}$ and $\mathbf{R} - \hat{\Psi}$ are not necessarily positive semidefinite and will often have some small negative eigenvalues. In such a case, the cumulative proportion of variance will exceed 1 and then decline to 1 as the negative eigenvalues are added. [Note that loadings cannot be obtained by (13.24) for the negative eigenvalues.]

■ EXAMPLE 13.3.2

To illustrate the principal factor method, we use the perception data from Table 13.1. The correlation matrix as given in Example 13.3.1 is singular. Hence in place of multiple correlations as communality estimates, we use the (absolute value of) the largest correlation in each row of \mathbf{R} . [The multiple correlation of y with several variables is greater than the simple correlation of y with any of the individual variables; see, for example, Rencher and Schaalje (2008, p. 257).] The diagonal elements of $\mathbf{R} - \hat{\Psi}$ as given by (13.39) are therefore .995, .837, .881, .995, and .837, which are obtained from \mathbf{R} in (13.35). The eigenvalues of $\mathbf{R} - \hat{\Psi}$ are 3.202, 1.395, .030, -.0002, and -.080, whose sum is 4.546. The first two eigenvectors of $\mathbf{R} - \hat{\Psi}$ are

$$\mathbf{c}_1 = \begin{pmatrix} .548 \\ .272 \\ .431 \\ .549 \\ .373 \end{pmatrix} \quad \text{and} \quad \mathbf{c}_2 = \begin{pmatrix} -.178 \\ .656 \\ -.460 \\ -.159 \\ .549 \end{pmatrix}.$$

When these are multiplied by the square roots of the respective eigenvalues, we obtain the principal factor loadings. In Table 13.3, these are compared with the loadings obtained by the principal component method in Example 13.3.1. The two sets of loadings are very similar, as we would have expected because of the large size of the communalities. The communalities in Table 13.3 are for the principal factor loadings, as noted above. The proportion of variance in each case for the principal factor loadings is obtained by dividing the variance accounted for (eigenvalue) by the sum of the eigenvalues, 4.546; for example, $3.202/4.546 = .704$. \square

13.3.3 Iterated Principal Factor Method

The principal factor method can easily be iterated to improve the estimates of communality. After obtaining $\hat{\Lambda}$ from $\mathbf{S} - \hat{\Psi}$ or $\mathbf{R} - \hat{\Psi}$ in (13.36) or (13.37) using initial communality estimates, we can obtain new communality estimates from the loadings in $\hat{\Lambda}$ using (13.28),

$$\hat{h}_i^2 = \sum_{j=1}^m \hat{\lambda}_{ij}^2.$$

These values of \hat{h}_i^2 are substituted into the diagonal of $\mathbf{S} - \hat{\Psi}$ or $\mathbf{R} - \hat{\Psi}$, from which we obtain a new value of $\hat{\Lambda}$ using (13.24). This process is continued until the

Table 13.3 Loadings Obtained by Two Different Methods for Perception Data of Table 13.1

Variables	Principal Component Loadings		Principal Factor Loadings		Communalities
	f_1	f_2	f_1	f_2	
Kind	.969	-.231	.981	-.210	.995
Intelligent	.519	.807	.487	.774	.837
Happy	.785	-.587	.771	-.544	.881
Likeable	.971	-.210	.982	-.188	.995
Just	.704	.667	.667	.648	.837
Variance accounted for	3.263	1.538	3.202	1.395	
Proportion of total variance	.653	.308	.704	.307	
Cumulative proportion	.653	.960	.704	1.01	

communality estimates converge. (For some data sets, the iterative procedure does not converge.) Then the eigenvalues and eigenvectors of the final version of $S - \hat{\Psi}$ or $R - \hat{\Psi}$ are used in (13.24) to obtain the loadings.

The principal factor method and the iterated principal factor method will typically yield results very close to those from the principal component method when *either* of the following is true.

1. The correlations are fairly large, with a resulting small value of m .
2. The number of variables, p , is large.

A shortcoming of the iterative approach is that sometimes it leads to a communality estimate \hat{h}_i^2 exceeding 1 (when factoring R). Such a result is known as a *Heywood case* (Heywood 1931). If $\hat{h}_i^2 > 1$, then $\hat{\psi}_i < 0$ by (13.26) and (13.28), which is clearly improper, since we cannot have a negative specific variance. Thus when a communality exceeds 1, the iterative process should stop, with the program reporting that a solution cannot be reached. Some software programs have an option of continuing the iterations by setting the communality equal to 1 in all subsequent iterations. The resulting solution with $\hat{\psi}_i = 0$ is somewhat questionable because it implies exact dependence of a variable on the factors, a possible but unlikely outcome.

■ EXAMPLE 13.3.3

We illustrate the iterated principal factor method using the Seishu data in Table 7.1. The correlation matrix is given below:

$$\mathbf{R} = \begin{pmatrix} 1.00 & .56 & .22 & .10 & .20 & -.04 & .13 & .03 & -.07 & .09 \\ .56 & 1.00 & -.09 & .13 & .20 & -.17 & .17 & .24 & .16 & .06 \\ .22 & -.09 & 1.00 & .16 & .70 & -.31 & -.45 & -.34 & -.11 & .68 \\ .10 & .13 & .16 & 1.00 & .49 & -.03 & -.16 & .01 & .42 & .37 \\ .20 & .20 & .70 & .49 & 1.00 & -.32 & -.34 & -.19 & .30 & .87 \\ -.04 & -.17 & -.31 & -.03 & -.32 & 1.00 & -.42 & -.57 & -.11 & -.26 \\ .13 & .17 & -.45 & -.16 & -.34 & -.42 & 1.00 & .82 & .23 & -.30 \\ .03 & .24 & -.34 & .01 & -.19 & -.57 & .82 & 1.00 & .45 & -.17 \\ -.07 & .16 & -.11 & .42 & .30 & -.11 & .23 & .45 & 1.00 & .29 \\ .09 & .06 & .68 & .37 & .87 & -.26 & -.30 & -.17 & .29 & 1.00 \end{pmatrix}.$$

The eigenvalues of \mathbf{R} are 3.17, 2.56, 1.43, 1.28, .54, .47, .25, .12, .10, and .06. There is a notable gap between 1.28 and .54, and we therefore extract four factors (see Section 13.4). The first four eigenvalues account for a proportion

$$\frac{3.17 + 2.56 + 1.43 + 1.28}{10} = .84$$

of $\text{tr}(\mathbf{R})$.

For initial communality estimates, we use the squared multiple correlation between each variable and the other nine variables. These are given in Table 13.4, along with the final communalities after iteration. We multiply the first four eigenvectors of the final iterated version of $\mathbf{R} - \hat{\Psi}$ by the square roots of the respective eigenvalues, as in (13.24), to obtain the factor loadings given in Table 13.4. We will not attempt to interpret the factors until after they have been rotated in Example 13.5.2b(b). \square

13.3.4 Maximum Likelihood Method

If we assume that the observations $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n$ constitute a random sample from $N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, then Λ and Ψ can be estimated by the method of maximum likelihood. It can be shown that the estimates $\hat{\Lambda}$ and $\hat{\Psi}$ satisfy the following:

$$\mathbf{S}\hat{\Psi}\hat{\Lambda} = \hat{\Lambda}(\mathbf{I} + \hat{\Lambda}'\hat{\Psi}^{-1}\hat{\Lambda}), \quad (13.43)$$

$$\hat{\Psi} = \text{diag}(\mathbf{S} - \hat{\Lambda}\hat{\Lambda}'), \quad (13.44)$$

$$\hat{\Lambda}'\hat{\Psi}^{-1}\hat{\Lambda} \text{ is diagonal.} \quad (13.45)$$

These equations must be solved iteratively, and in practice the procedure may fail to converge or may yield a Heywood case (Section 13.3.3).

We note that the proportion of variance accounted for by the factors, as given by (13.32) or (13.33), will not necessarily be in descending order for maximum likelihood factors, as it is for factors obtained from the principal component or principal factor method.

Table 13.4 Iterated Principal Factor Loadings and Communalities for the Seishu Data

Variables	Loadings				Initial Communalities	Final Communalities
	f_1	f_2	f_3	f_4		
Taste	.22	.31	.92	.12	.57	1.00
Odor	.07	.40	.43	-.20	.54	.38
pH	.80	.04	.05	-.40	.78	.79
Acidity 1	.41	.22	-.11	.37	.40	.36
Acidity 2	.94	.28	-.07	.05	.88	.98
Sake meter	-.13	-.67	.10	.56	.77	.79
Reducing sugar	-.55	.66	.03	-.11	.79	.75
Total sugar	-.45	.88	-.14	-.07	.87	.99
Alcohol	.13	.54	-.37	.54	.66	.74
Formyl-nitrogen	.84	.21	-.17	-.02	.80	.78
Variance accounted for	3.00	2.37	1.25	.96	7.06	7.57

■ EXAMPLE 13.3.4

We illustrate the maximum likelihood method with the Seishu data of Table 7.1. The correlation matrix and its eigenvalues were given in Example 13.3.3. We extract four factors, as in Example 13.3.3. The iterative solution of (13.43), (13.44), and (13.45) yielded the loadings and communalities given in Table 13.5.

The pattern of the loadings is different from that obtained using the iterated principal factor method in Example 13.3.3, but we will not compare them until after rotation in Example 13.5.2b(b). Note that the four values of variance accounted for are not in descending order. \square

13.4 CHOOSING THE NUMBER OF FACTORS, m

Several criteria have been proposed for choosing m , the number of factors. We consider four criteria, which are similar to those given in Section 12.6 for choosing the number of principal components to retain.

1. Choose m equal to the number of factors necessary for the variance accounted for to achieve a predetermined percentage, say 80%, of the total variance $\text{tr}(\mathbf{S})$ or $\text{tr}(\mathbf{R})$.
2. Choose m equal to the number of eigenvalues greater than the average eigenvalue. For \mathbf{R} the average is 1; for \mathbf{S} it is $\sum_{j=1}^p \theta_j/p$.
3. Use the scree test based on a plot of the eigenvalues of \mathbf{S} or \mathbf{R} . If the graph drops sharply, followed by a straight line with much smaller slope, choose m equal to the number of eigenvalues before the straight line begins.

Table 13.5 Maximum Likelihood Loadings and Communalities for the Seishu Data

Variables	Loadings				Communalities
	f_1	f_2	f_3	f_4	
Taste	1.00	0	0	0	1.00
Odor	.45	-.05	.22	.19	.29
pH	.22	.68	-.20	-.40	.71
Acidity 1	.10	.47	.10	.37	.38
Acidity 2	.20	.98	.02	.00	1.00
Sake meter	-.04	-.31	-.68	.55	.86
Reducing sugar	.13	-.39	.76	-.02	.75
Total sugar	.03	-.22	.96	.02	.98
Alcohol	-.07	.31	.52	.60	.72
Formyl-nitrogen	.02	.79	-.05	-.10	.63
Variance accounted for	1.33	2.66	2.34	1.00	7.32

4. Test the hypothesis that m is the correct number of factors, $H_0: \Sigma = \Lambda\Lambda' + \Psi$, where Λ is $p \times m$.

Method 1 applies particularly to the principal component method. By (13.32), the proportion of total sample variance (variance accounted for) due to the j th factor from \mathbf{S} is $\sum_{i=1}^p \hat{\lambda}_{ij}^2 / \text{tr}(\mathbf{S})$. The corresponding proportion from \mathbf{R} is $\sum_{i=1}^p \hat{\lambda}_{ij}^2 / p$, as in (13.33). The contribution of all m factors to $\text{tr}(\mathbf{S})$ or p is therefore $\sum_{i=1}^p \sum_{j=1}^m \hat{\lambda}_{ij}^2$, which is the sum of squares of all elements of $\hat{\Lambda}$. For the principal component method, we see by (13.28) and (13.29) that this sum is also equal to the sum of the first m eigenvalues or to the sum of all p communalities:

$$\sum_{i=1}^p \sum_{j=1}^m \hat{\lambda}_{ij}^2 = \sum_{i=1}^p \hat{h}_i^2 = \sum_{j=1}^m \theta_j. \quad (13.46)$$

Thus we choose m sufficiently large so that the sum of the communalities or the sum of the eigenvalues (variance accounted for) constitutes a relatively large portion of $\text{tr}(\mathbf{S})$ or p .

Method 1 can be extended to the principal factor method, where prior estimates of communalities are used to form $\mathbf{S} - \hat{\Psi}$ or $\mathbf{R} - \hat{\Psi}$. However, $\mathbf{S} - \hat{\Psi}$ or $\mathbf{R} - \hat{\Psi}$ will often have some negative eigenvalues. Therefore, as values of m range from 1 to p , the cumulative proportion of eigenvalues, $\sum_{j=1}^m \theta_j / \sum_{j=1}^p \theta_j$, will exceed 1.0 and then reduce to 1.0 as the negative eigenvalues are added. Hence a percentage such as 80% will be reached for a lower value of m than would be the case for \mathbf{S} or \mathbf{R} , and a better strategy might be to choose m equal to the value for which the percentage first exceeds 100%.

In the iterated principal factor method, m is specified before iteration, and $\sum_i \hat{h}_i^2$ is obtained after iteration as $\sum_i \hat{h}_i^2 = \text{tr}(\mathbf{S} - \hat{\Psi})$. To choose m before iterating, one could use a priori considerations or the eigenvalues of \mathbf{S} or \mathbf{R} , as in the principal component method.

Method 2 is a popular criterion of long standing and is the default in many software packages. Although heuristically based, it often works well in practice. A variation to method 2 that has been suggested for use with $\mathbf{R} - \hat{\Psi}$ is to let m equal the number of positive eigenvalues. (There will typically be some negative eigenvalues of $\mathbf{R} - \hat{\Psi}$.) However, this criterion will often result in too many factors, since the sum of the positive eigenvalues will exceed the sum of the communalities.

The scree test in method 3 was named after the geological term *scree*, referring to the debris at the bottom of a rocky cliff. It also performs well in practice.

In method 4 we wish to test

$$H_0: \Sigma = \Lambda\Lambda' + \Psi \quad \text{vs.} \quad H_1: \Sigma \neq \Lambda\Lambda' + \Psi,$$

where Λ is $p \times m$. The test statistic, a function of the likelihood ratio, is

$$\left(n - \frac{2p + 4m + 11}{6} \right) \ln \left(\frac{|\hat{\Lambda}\hat{\Lambda}' + \hat{\Psi}|}{|\mathbf{S}|} \right), \quad (13.47)$$

which is approximately χ^2_ν when H_0 is true, where $\nu = \frac{1}{2}[(p - m)^2 - p - m]$ and $\hat{\Lambda}$ and $\hat{\Psi}$ are the maximum likelihood estimators. Rejection of H_0 implies that m is too small and more factors are needed.

In practice, when n is large, the test in method 4 often shows more factors to be significant than do the other three methods. We may therefore consider the value of m indicated by the test to be an upper bound on the number of factors with practical importance.

For many data sets, the choice of m will not be obvious. This indeterminacy leaves many statisticians skeptical as to the validity of factor analysis. A researcher may begin with one of the methods (say method 2) for an initial choice of m , will inspect the resulting percent of $\text{tr}(\mathbf{R})$ or $\text{tr}(\mathbf{S})$, and will then examine the rotated loadings for interpretability. If the percent of variance or interpretation does not seem satisfactory, the experimenter will try other values of m in a search for an acceptable compromise between percent of $\text{tr}(\mathbf{R})$ and interpretability of the factors. Admittedly, this is a subjective procedure, and for such data sets one could well question the outcome (see Section 13.7).

When a data set is successfully fitted by a factor analysis model, the first three methods will almost always give the same value of m , and there will be little question as to what this value should be. Thus for a “good” data set, the entire procedure becomes much more objective.

■ EXAMPLE 13.4(a)

We compare the four methods of choosing m for the perception data used in Examples 13.3.1 and 13.3.2.




Figure 13.1 Scree graph for the perception data.

Method 1 gives $m = 2$, because one eigenvalue accounts for 65% of $\text{tr}(\mathbf{R})$, while two eigenvalues account for 96%.

Method 2 gives $m = 2$, since $\lambda_2 = 1.54$ and $\lambda_3 = .17$.

For method 3, we examine the scree plot in Figure 13.1. It is clear that $m = 2$ is indicated.

Method 4 is not available for the perception data because \mathbf{R} is singular (fifth eigenvalue is zero), and the test involves $|\mathbf{R}|$.

Hence for the perception data, all three available methods agree on $m = 2$. □

■ EXAMPLE 13.4(b)

We compare the four methods of choosing m for the Seishu data used in Examples 13.3.3 and 13.3.4.

Method 1 gives $m = 4$ for the principal component method, because four eigenvalues of \mathbf{R} account for 82% of $\text{tr}(\mathbf{R})$. For the principal factor method with initial communality estimates R_i^2 , the eigenvalues of $\mathbf{R} - \hat{\Psi}$ and corresponding proportions are as follows:

Eigenvalues	2.86	2.17	.94	.88	.12	.08	.01	-.06	-.13	-.22
Proportions	.43	.33	.14	.16	.02	.01	.00	-.01	-.02	-.03
Cumulative proportions	.43	.76	.90	1.03	1.05	1.06	1.06	1.06	1.03	1.00




Figure 13.2 Scree graph for the Seishu data.

The proportions are obtained by dividing the eigenvalues by their sum, 6.63. Thus the cumulative proportion first exceeds 1.00 for $m = 4$.

Method 2 gives $m = 4$, since $\lambda_4 = 1.31$ and $\lambda_5 = .61$, where λ_4 and λ_5 are eigenvalues of \mathbf{R} .

For method 3, we examine the scree plot in Figure 13.2. There is a discernible bend in slope at the fifth eigenvalue.

For method 4, we use $m = 4$ in the approximate chi-squared statistic in (13.47) and obtain $\chi^2 = 9.039$, with degrees of freedom

$$\nu = \frac{1}{2}[(p - m)^2 - p - m] = \frac{1}{2}[(10 - 4)^2 - 10 - 4] = 11.$$

Since $9.039 < \chi^2_{.05,11} = 19.68$, we do not reject the hypothesis that four factors are adequate.

Thus for the Seishu data, all four methods agree on $m = 4$. □

13.5 ROTATION

13.5.1 Introduction

As noted in Section 13.2.2, the factor loadings (rows of Λ) in the population model are unique only up to multiplication by an orthogonal matrix that rotates the loadings. The rotated loadings preserve the essential properties of the original loadings; they reproduce the covariance matrix and satisfy all basic assumptions. The estimated loading matrix $\hat{\Lambda}$ can likewise be rotated to obtain $\hat{\Lambda}^* = \hat{\Lambda}\mathbf{T}$, where \mathbf{T} is orthogonal.

Since $\mathbf{T}\mathbf{T}' = \mathbf{I}$ by (2.102), the rotated loadings provide the same estimate of the covariance matrix as before:

$$\mathbf{S} \cong \hat{\Lambda}^* \hat{\Lambda}^{**} + \hat{\Psi} = \hat{\Lambda} \mathbf{T} \mathbf{T}' \hat{\Lambda}' + \hat{\Psi} = \hat{\Lambda} \hat{\Lambda}' + \hat{\Psi}. \quad (13.48)$$

Geometrically, the loadings in the i th row of $\hat{\Lambda}$ constitute the coordinates of a point in the loading space corresponding to y_i . Rotation of the p points gives their coordinates with respect to new axes (factors) but otherwise leaves their basic geometric configuration intact. We hope to find a new frame of reference in which the factors are more interpretable. To this end, the goal of rotation is to place the axes close to as many points as possible. If there are clusters of points (corresponding to groupings of y 's), we seek to move the axes so as to pass through or near these clusters. This would associate each group of variables with a factor (axis) and make interpretation more objective. The resulting axes then represent the natural factors.

If we can achieve a rotation in which every point is close to an axis, then each variable loads highly on the factor corresponding to the axis and has small loadings on the remaining factors. In this case, there is no ambiguity. Such a happy state of affairs is called *simple structure*, and interpretation is greatly simplified. We merely observe which variables are associated with each factor, and the factor is defined or named accordingly.

In order to identify the natural groupings of variables, we seek a rotation to an interpretable pattern for the loadings, in which the variables load highly on only one factor. The number of factors on which a variable has moderate or high loadings is called the *complexity* of the variable. In the ideal situation referred to above as simple structure, the variables all have a complexity of 1. In this case, the variables have been clearly clustered into groups corresponding to the factors.

We consider two basic types of rotation: *orthogonal* and *oblique*. The rotation in (13.48) involving an orthogonal matrix is an orthogonal rotation; the original perpendicular axes are rotated rigidly and remain perpendicular. In an orthogonal rotation, angles and distances are preserved, communalities are unchanged, and the basic configuration of the points remains the same. Only the reference axes differ. In an oblique “rotation” (transformation), the axes are not required to remain perpendicular and are thus free to pass closer to clusters of points.

In Sections 13.5.2 and 13.5.3, we discuss orthogonal and oblique rotations, followed by some guidelines for interpretation in Section 13.5.4.

13.5.2 Orthogonal Rotation

It was noted above in Section 13.5.1 that orthogonal rotations preserve communalities. This is because the rows of $\hat{\Lambda}$ are rotated, and the distance to the origin is unchanged, which, by (13.28), is the communality. However, the variance accounted for by each factor as given in (13.31) will change, as will the corresponding proportion in (13.32) or (13.33). The proportions due to the rotated loadings will not necessarily be in descending order.

In Sections 13.5.2a and 13.5.2b, we consider two approaches to orthogonal rotation.

13.5.2a Graphical Approach

If there are only two factors ($m = 2$), we can use a *graphical* rotation based on a visual inspection of a plot of factor loadings. In this case, the rows of $\hat{\Lambda}$ are pairs of loadings, $(\hat{\lambda}_{i1}, \hat{\lambda}_{i2})$, $i = 1, 2, \dots, p$, corresponding to y_1, y_2, \dots, y_p . We choose an angle ϕ through which the axes can be rotated to move them closer to groupings of points. The new rotated loadings $(\hat{\lambda}_{i1}^*, \hat{\lambda}_{i2}^*)$ can be measured directly on the graph as coordinates of the axes or calculated from $\hat{\Lambda}^* = \hat{\Lambda}\mathbf{T}$ using

$$\mathbf{T} = \begin{pmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{pmatrix}. \quad (13.49)$$

■ EXAMPLE 13.5.2a

In Example 13.3.1, the initial factor loadings for the perception data did not provide an interpretation consistent with the two groupings of variables apparent in the pattern of correlations in \mathbf{R} . The five pairs of loadings $(\hat{\lambda}_{i1}, \hat{\lambda}_{i2})$ corresponding to the five variables are plotted in Figure 13.3. An orthogonal rotation through -35° would bring the axes (factors) closer to the two clusters of points (variables) identified in Example 13.3.1. With the rotation, each cluster of variables corresponds much more closely to a factor. Using $\hat{\Lambda}$ from Example 13.3.1 and -35° in \mathbf{T} as given in (13.49), we obtain the following rotated loadings:

$$\begin{aligned} \hat{\Lambda}^* = \hat{\Lambda}\mathbf{T} &= \begin{pmatrix} .969 & -.231 \\ .519 & .807 \\ .785 & -.587 \\ .971 & -.210 \\ .704 & .667 \end{pmatrix} \begin{pmatrix} .819 & .574 \\ -.574 & .819 \end{pmatrix} \\ &= \begin{pmatrix} .927 & .367 \\ -.037 & .959 \\ .980 & -.031 \\ .916 & .385 \\ .194 & .950 \end{pmatrix}. \end{aligned}$$

In Table 13.6, we compare the rotated loadings in $\hat{\Lambda}^*$ with the original loadings in $\hat{\Lambda}$.

The interpretation of the rotated loadings is clear. As indicated by the bold-face loadings in Table 13.6, the first factor is associated with variables 1, 3, and 4: kind, happy, and likeable. The second factor is associated with the other two variables: intelligent and just. This same grouping of variables is indicated by the pattern in the correlation matrix in (13.35) and can also be seen in the two clusters of points in Figure 13.3. The first factor might be described as representing a person's perceived humanity or amiability, while the second involves more logical or rational practices.




Figure 13.3 Plot of the two loadings for each of the five variables in the perception data of Table 13.1.

Note that if the angle between the rotated axes was allowed to be less than 90° (an oblique rotation), the lower axis representing f_1^* could come closer to the points corresponding to variables 1 and 4 so that the coordinates on f_2^* , .367 and .385, could be reduced. However, the basic interpretation would not change; variables 1 and 4 would still be associated with f_1^* . \square

13.5.2b Varimax Rotation

The graphical approach to rotation is generally limited to $m = 2$. For $m > 2$, various analytical methods have been proposed. The most popular of these is the *varimax* technique, which seeks rotated loadings that maximize the variance of the squared loadings in each column of $\hat{\Lambda}^*$. If the loadings in a column were nearly equal, the variance would be close to 0. As the squared loadings approach 0 and 1 (for factoring \mathbf{R}), the variance will approach a maximum. Thus the varimax method attempts to make the loadings either large or small to facilitate interpretation.

The varimax procedure cannot guarantee that all variables will load highly on only one factor. In fact, no procedure could do this for all possible data sets. The configuration of the points in the loading space remains fixed; we merely rotate the

Table 13.6 Graphically Rotated Loadings for the Perception Data of Table 13.1

Variables	Principal Component Loadings		Graphically Rotated Loadings		Communalities, \hat{h}_i^2
	f_1	f_2	f_1	f_2	
Kind	.969	-.231	.927	.367	.993
Intelligent	.519	.807	-.037	.959	.921
Happy	.785	-.587	.980	-.031	.960
Likeable	.971	-.210	.916	.385	.987
Just	.704	.667	.194	.950	.940
Variance accounted for	3.263	1.538	2.696	2.106	4.802
Proportion of total variance	.653	.308	.539	.421	.960
Cumulative proportion	.653	.960	.539	.960	.960

axes to be as close to as many points as possible. In many cases, the points are not well clustered, and the axes simply cannot be rotated so as to be near all of them. This problem is compounded by having to choose m . If m is changed, the coordinates $(\hat{\lambda}_{i1}, \hat{\lambda}_{i2}, \dots, \hat{\lambda}_{im})$ change, and the relative position of the points is altered.

The varimax rotation is available in virtually all factor analysis software programs. The output typically includes the rotated loading matrix $\hat{\Lambda}^*$, the variance accounted for (sum of squares of each column of $\hat{\Lambda}^*$), the communalities (sum of squares of each row of $\hat{\Lambda}^*$), and the orthogonal matrix \mathbf{T} used to obtain $\hat{\Lambda}^* = \hat{\Lambda}\mathbf{T}$.

■ EXAMPLE 13.5.2b(a)

In Example 13.5.2a, a graphical rotation was devised visually to achieve interpretable loadings for the perception data of Table 13.1. As we would expect, the varimax method yields a similar result. The varimax rotated loadings are given in Table 13.7. For comparison, we have included the original unrotated loadings from Table 13.3 and the graphically rotated loadings from Table 13.6.

The orthogonal matrix \mathbf{T} for the varimax rotation is

$$\mathbf{T} = \begin{pmatrix} .859 & .512 \\ -.512 & .859 \end{pmatrix}.$$

By (13.49), $-\sin \phi = .512$, and the angle of rotation is $\phi = -\sin^{-1}(.512) = -30.8^\circ$. Thus the varimax rotation chose an angle of rotation of -30.8° as compared to the -35° we selected visually, but the results are very close and the interpretation is exactly the same. \square

Table 13.7 Varimax Rotated Factor Loadings for the Perception Data of Table 13.1

Variables	Principal Component Loadings		Graphically Rotated Loadings		Varimax Rotated Loadings		Communalities, \hat{h}_i^2
	f_1	f_2	f_1	f_2	f_1	f_2	
Kind	.969	-.231	.927	.367	.951	.298	.993
Intelligent	.519	.807	-.037	.959	.033	.959	.921
Happy	.785	-.587	.980	-.031	.975	-.103	.960
Likeable	.971	-.210	.916	.385	.941	.317	.987
Just	.704	.667	.194	.950	.263	.933	.940
Variance accounted for	3.263	1.538	2.696	2.106	2.811	1.991	4.802
Proportion of total variance	.653	.308	.539	.421	.562	.398	.960
Cumulative proportion	.653	.960	.539	.960	.562	.960	.960

■ EXAMPLE 13.5.2b(b)

In Examples 13.3.3 and 13.3.4, we obtained the iterated principal factor loadings and maximum likelihood loadings for the Seishu data. In Table 13.8, we show the varimax rotation of these two sets of loadings. The similarities in the two sets of rotated loadings are striking. The interpretation in each case is the same. The variances accounted for are virtually identical.

The rotation in each case has achieved a satisfactory simple structure, and most variables show a complexity of 1. The boldface loadings indicate the variables associated with each factor for interpretation purposes. These may be meaningful to the researcher. For example, factor 2 is associated with sake meter, reducing sugar, and total sugar, while factor 3 is aligned with taste and odor. \square

13.5.3 Oblique Rotation

The term *oblique rotation* refers to a transformation in which the axes do not remain perpendicular. Technically, the term *oblique rotation* is a misnomer, since rotation implies an orthogonal transformation that preserves distances. A more accurate characterization would be *oblique transformation*, but the term oblique rotation is well established in the literature.

Instead of the orthogonal transformation matrix \mathbf{T} used in (13.16), (13.17), and (13.18), an oblique rotation uses a general nonsingular transformation matrix \mathbf{Q} to obtain $\mathbf{f}^* = \mathbf{Q}'\mathbf{f}$, and by (3.74),

$$\text{cov}(\mathbf{f}^*) = \mathbf{Q}'\mathbf{I}\mathbf{Q} = \mathbf{Q}'\mathbf{Q} \neq \mathbf{I}. \quad (13.50)$$

Table 13.8 Varimax Rotated Loadings for the Seishu Data

Variables	Iterated Principal Factor Rotated Loadings				Maximum Likelihood Rotated Loadings			
	<i>f</i> ₁	<i>f</i> ₂	<i>f</i> ₃	<i>f</i> ₄	<i>f</i> ₁	<i>f</i> ₂	<i>f</i> ₃	<i>f</i> ₄
Taste	.16	-.01	.99	-.09	.16	-.00	.98	-.10
Odor	-.11	.14	.48	.14	-.07	.14	.49	.17
pH	.88	-.12	.02	-.13	.82	-.10	.08	-.15
Acidity 1	.26	-.09	.09	.54	.29	-.08	.11	.53
Acidity 2	.89	-.06	.10	.43	.91	-.06	.10	.39
Sake meter	-.43	-.76	.01	.07	-.46	-.80	.04	.10
Reducing sugar	-.37	.76	.18	.03	-.37	.75	.20	.08
Total sugar	-.26	.92	.10	.25	-.27	.91	.11	.26
Alcohol	-.01	.25	.00	.80	-.00	.25	.01	.81
Formyl-nitrogen	.74	-.07	-.08	.20	.76	-.07	-.08	.22
Variance accounted for	2.62	2.12	1.27	1.27	2.61	2.14	1.29	1.28

Thus the new factors are correlated. Since distances and angles are not preserved, the communalities for f^* are different from those for f . Some program packages report communalities obtained from the original loadings, rather than the oblique loadings.

When the axes are not required to be perpendicular, they can more easily pass through the major clusters of points in the loading space (assuming there are such clusters). For example, in Figure 13.4, we have plotted the varimax rotated loadings for two factors extracted from the sons data of Table 3.8 (see Example 13.5.3 at the end of this section). Oblique axes with an angle of 38° would pass much closer to the points, and the resulting loadings would be very close to 0 and 1. However, the interpretation would not change, since the same points (variables) would be associated with the oblique axes as with the orthogonal axes.

Various analytical methods for achieving oblique rotations have been proposed and are available in program packages. Typically, the output of one of these procedures includes a *pattern matrix*, a *structure matrix*, and a matrix of correlations among the oblique factors. For interpretation, we would usually prefer the pattern matrix rather than the structure matrix. The loadings in a row of the pattern matrix are the natural coordinates of the point (variable) on the oblique axes and serve as coefficients in the model relating the variable to the factors.

One use for an oblique rotation is to check on the orthogonality of the factors. The orthogonality in the original factors is imposed by the model and maintained by an orthogonal rotation. If an oblique rotation produces a correlation matrix that is nearly diagonal, we can be more confident that the factors are indeed orthogonal.




Figure 13.4 Orthogonal and oblique rotations for the sons data.

■ EXAMPLE 13.5.3

The correlation matrix for the sons data of Table 3.8 is given below:

$$\mathbf{R} = \begin{pmatrix} 1.000 & .735 & .711 & .704 \\ .735 & 1.000 & .693 & .709 \\ .711 & .693 & 1.000 & .839 \\ .704 & .709 & .839 & 1.000 \end{pmatrix}.$$

The varimax rotated loadings for two factors obtained by the principal component method are given in Table 13.9 and plotted in Figure 13.4. An analytical oblique rotation (Harris–Kaiser orthoblique method in SAS) produced oblique axes with an angle of 38° , the same as obtained by a graphical approach. The correlation between the two factors is .79 [obtained from $\mathbf{Q}'\mathbf{Q}$ in (13.50)], which is related to the angle by (3.15), $.79 = \cos 38^\circ$. The pattern loadings are given in Table 13.9.

The oblique loadings give a much cleaner simple structure than the varimax loadings, but the interpretation is essentially the same if we neglect loadings below .45 on the varimax rotation.

Table 13.9 Varimax and Orthoblique Loadings for the Sons Data

Variable	Varimax Loadings		Orthoblique Pattern Matrix	
	<i>f</i> ₁	<i>f</i> ₂	<i>f</i> ₁	<i>f</i> ₂
1	.42	.82	.03	.90
2	.40	.85	-.03	.96
3	.87	.41	.97	-.01
4	.86	.43	.95	.01

In Figure 13.4, it is evident that a single factor would be adequate since the angle between axes is less than 45°. The suggestion to let $m = 1$ is also supported by the first three criteria in Section 13.4: the eigenvalues of \mathbf{R} are 3.20, .38, .27, and .16. The first accounts for 80%, the second for an additional 9%. The large correlation, .79, between the two oblique factors constitutes additional evidence that a single factor model would suffice here. In fact, the pattern in \mathbf{R} itself indicates the presence of only one factor. The four variables form only one cluster, since all are highly correlated. There are no small correlations between groupings of variables. \square

13.5.4 Interpretation

In Sections 13.5.1, 13.5.2, and 13.5.3, we have discussed the usefulness of rotation as an aid to interpretation. Our goal is to achieve a simple structure in which each variable loads highly on only one factor, with small loadings on all other factors. In practice, we often fail to achieve this goal, but rotation usually produces loadings that are closer to the desired simple structure.

We now suggest general guidelines for interpreting the factors by examination of the matrix of rotated factor loadings. Moving horizontally from left to right across the m loadings in each row, identify the highest loading (in absolute value). If the highest loading is of a significant size (a subjective determination, see the next paragraph), circle or underline it. This is done for each of the p variables. There may be other significant loadings in a row besides the one circled. If these are considered, the interpretation is less simple. On the other hand, there may be variables with such small communalities that no significant loading appears on any factor. In this case, the researcher may wish to increase the number of factors and run the program again so that these variables might associate with a new factor.

To assess significance of factor loadings $\hat{\lambda}_{ij}$ obtained from \mathbf{R} , a threshold value of .3 has been advocated by many writers. For most successful applications, however, a critical value of .3 is too low and will result in variables of complexity greater than 1. A target value of .5 or .6 is typically more useful. The .3 criterion is loosely based on the critical value for significance of an ordinary correlation coefficient, r . However, the distribution of the sample loadings is not the same as that of r arising

from the bivariate normal. In addition, the critical value should be increased because mp values of $\hat{\lambda}_{ij}$ are being tested. On the other hand, if m is large, the critical value might possibly need to be reduced somewhat. Since $\hat{h}_i^2 = \sum_{j=1}^m \hat{\lambda}_{ij}^2$ is bounded by 1, an increase in m reduces the average squared loading in a row.

After identifying potentially significant loadings, the experimenter then attempts to discover some meaning in the factors and, ideally, to label or name them. This can readily be done if the group of variables associated with each factor makes sense to the researcher. But in many situations, the groupings are not so logical, and a revision can be tried, such as adjusting the size of loading deemed to be important, changing m , using a different method of estimating the loadings, or employing another type of rotation.

13.6 FACTOR SCORES

In many applications, the researcher wishes only to ascertain whether a factor analysis model fits the data and to identify the factors. In other applications, the experimenter wishes to obtain *factor scores* $\hat{\mathbf{f}}_i = (\hat{f}_{i1}, \hat{f}_{i2}, \dots, \hat{f}_{im})'$, $i = 1, 2, \dots, n$, which are defined as estimates of the underlying factor values for each observation. There are two potential uses for such scores: (1) the behavior of the observations in terms of the factors may be of interest and (2) we may wish to use the factor scores as input to another analysis, such as MANOVA. The latter usage resembles a similar application of principal components.

Since the f 's are not observed, we must estimate them as functions of the observed y 's. The most popular approach to estimating the factors is based on regression (Thompson 1951). We will discuss this method and also briefly describe an informal technique that can be used when \mathbf{R} (or \mathbf{S}) is singular. For other approaches see Harman (1976, Chapter 16).

Since $E(f_i) = 0$, we relate the f 's to the y 's by a centered regression model

$$\begin{aligned} f_1 &= \beta_{11}(y_1 - \bar{y}_1) + \beta_{12}(y_2 - \bar{y}_2) + \cdots + \beta_{1p}(y_p - \bar{y}_p) + \epsilon_1 \\ f_2 &= \beta_{21}(y_1 - \bar{y}_1) + \beta_{22}(y_2 - \bar{y}_2) + \cdots + \beta_{2p}(y_p - \bar{y}_p) + \epsilon_2 \\ &\vdots \\ f_m &= \beta_{m1}(y_1 - \bar{y}_1) + \beta_{m2}(y_2 - \bar{y}_2) + \cdots + \beta_{mp}(y_p - \bar{y}_p) + \epsilon_m, \end{aligned} \quad (13.51)$$

which can be written in matrix form as

$$\mathbf{f} = \mathbf{B}'_1(\mathbf{y} - \bar{\mathbf{y}}) + \boldsymbol{\epsilon}. \quad (13.52)$$

We have used the notation $\boldsymbol{\epsilon}$ to distinguish this error from $\boldsymbol{\varepsilon}$ in the original factor model $\mathbf{y} - \boldsymbol{\mu} = \boldsymbol{\Lambda}\mathbf{f} + \boldsymbol{\varepsilon}$ given in (13.3). Our approach is to estimate \mathbf{B}_1 and use the predicted value $\hat{\mathbf{f}} = \hat{\mathbf{B}}'_1(\mathbf{y} - \bar{\mathbf{y}})$ to estimate \mathbf{f} .

The model (13.52) holds for each observation:

$$f_i = \mathbf{B}'_1(\mathbf{y}_i - \bar{\mathbf{y}}) + \epsilon_i, \quad i = 1, 2, \dots, n.$$

In transposed form, the model becomes

$$\mathbf{f}'_i = (\mathbf{y}_i - \bar{\mathbf{y}})' \mathbf{B}_1 + \boldsymbol{\epsilon}'_i, \quad i = 1, 2, \dots, n,$$

and these n equations can be combined into a single model,

$$\begin{aligned} \mathbf{F} &= \begin{pmatrix} \mathbf{f}'_1 \\ \mathbf{f}'_2 \\ \vdots \\ \mathbf{f}'_n \end{pmatrix} = \begin{pmatrix} (\mathbf{y}_1 - \bar{\mathbf{y}})' \mathbf{B}_1 \\ (\mathbf{y}_2 - \bar{\mathbf{y}})' \mathbf{B}_1 \\ \vdots \\ (\mathbf{y}_n - \bar{\mathbf{y}})' \mathbf{B}_1 \end{pmatrix} + \begin{pmatrix} \boldsymbol{\epsilon}'_1 \\ \boldsymbol{\epsilon}'_2 \\ \vdots \\ \boldsymbol{\epsilon}'_n \end{pmatrix} \\ &= \begin{pmatrix} (\mathbf{y}_1 - \bar{\mathbf{y}})' \\ (\mathbf{y}_2 - \bar{\mathbf{y}})' \\ \vdots \\ (\mathbf{y}_n - \bar{\mathbf{y}})' \end{pmatrix} \mathbf{B}_1 + \boldsymbol{\Xi} \\ &= \mathbf{Y}_c \mathbf{B}_1 + \boldsymbol{\Xi} \quad [\text{by (10.11)}]. \end{aligned} \quad (13.53)$$

The model (13.53) has the appearance of a centered multivariate multiple regression model as in Section 10.4.5, with \mathbf{Y}_c in place of \mathbf{X}_c . By (10.55), the estimate for \mathbf{B}_1 would be

$$\hat{\mathbf{B}}_1 = (\mathbf{Y}'_c \mathbf{Y}_c)^{-1} \mathbf{Y}'_c \mathbf{F}. \quad (13.54)$$

However, \mathbf{F} is unobserved. To evaluate $\hat{\mathbf{B}}_1$ in spite of this, we first use (10.57) to rewrite (13.54) in terms of covariance matrices,

$$\hat{\mathbf{B}}_1 = \mathbf{S}_{yy}^{-1} \mathbf{S}_{yf}. \quad (13.55)$$

In the notation of the present chapter, \mathbf{S}_{yy} is represented by \mathbf{S} ; for \mathbf{S}_{yf} we use $\hat{\Lambda}$, since $\hat{\Lambda}$ estimates $\text{cov}(\mathbf{y}, \mathbf{f}) = \Lambda$ in (13.13). Thus, based on the assumptions in Section 13.2.1, we can write (13.55) as

$$\hat{\mathbf{B}}_1 = \mathbf{S}^{-1} \hat{\Lambda}. \quad (13.56)$$

Then from model (13.53), the estimated (predicted) \mathbf{f}_i values are given by

$$\begin{aligned} \hat{\mathbf{F}} &= \begin{pmatrix} \hat{\mathbf{f}}'_1 \\ \hat{\mathbf{f}}'_2 \\ \vdots \\ \hat{\mathbf{f}}'_n \end{pmatrix} = \mathbf{Y}_c \hat{\mathbf{B}}_1 \\ &= \mathbf{Y}_c \mathbf{S}^{-1} \hat{\Lambda}. \end{aligned} \quad (13.57)$$

If \mathbf{R} is factored instead of \mathbf{S} , (13.56) and (13.57) become

$$\hat{\mathbf{B}}_1 = \mathbf{R}^{-1} \hat{\Lambda}, \quad (13.58)$$

$$\hat{\mathbf{F}} = \mathbf{Y}_s \mathbf{R}^{-1} \hat{\Lambda}, \quad (13.59)$$

respectively, where \mathbf{Y}_s is the observed matrix of standardized variables, $(y_{ij} - \bar{y}_j)/s_j$.

We would ordinarily obtain factor scores for the rotated factors rather than the original factors. Thus $\hat{\Lambda}$ in (13.57) or (13.59) would be replaced by $\hat{\Lambda}^*$.

In order to obtain factor scores by (13.57) or (13.59), \mathbf{S} or \mathbf{R} must be nonsingular. When \mathbf{R} (or \mathbf{S}) is singular, we can obtain factor scores by a simple method based directly on the rotated loadings. We cluster the variables into groups (factors) according to the loadings and find a score for each factor by averaging the variables associated with the factor. If the variables are not commensurate, the variables should be standardized before averaging. An alternative approach would be to weight the variables by their loadings when averaging.

■ EXAMPLE 13.6

The speaking rate of four voices was artificially manipulated by means of a rate changer without altering the pitch (Brown, Strong, and Rencher 1973). There were five rates for each voice:

$$\text{FF} = 45\% \text{ faster}$$

$$\text{F} = 25\% \text{ faster}$$

$$\text{N} = \text{normal rate}$$

$$\text{S} = 22\% \text{ slower}$$

$$\text{SS} = 42\% \text{ slower.}$$

The resulting 20 voices were played to 30 judges who rated them on 15 paired-opposite adjectives (variables) with a 14-point scale between poles. The following adjectives were used: intelligent, ambitious, polite, active, confident, happy, just, likeable, kind, sincere, dependable, religious, good-looking, sociable, and strong. The results were averaged over the 30 judges to produce 20 observation vectors of 15 variables each. The averaging produced very reliable data so that even though there were only 20 observations on 15 variables, the factor analysis model fit very well. The correlation matrix is as follows:

$$\mathbf{R} = \begin{pmatrix} 1.00 & .90 & -.17 & .88 & .92 & .88 & .15 & .39 & -.02 & -.16 & .52 & -.15 & -.79 & -.78 & .73 \\ .90 & 1.00 & -.46 & .93 & .87 & .79 & -.16 & .10 & -.35 & -.42 & .25 & -.40 & .68 & -.60 & .62 \\ -.17 & -.46 & 1.00 & -.56 & -.13 & .07 & .85 & .75 & .88 & .91 & .68 & .88 & .21 & .31 & .25 \\ .88 & .93 & -.56 & 1.00 & .85 & .73 & -.25 & -.02 & -.45 & -.57 & .10 & -.53 & .58 & .84 & .50 \\ .92 & .87 & -.13 & .85 & 1.00 & .91 & .20 & .39 & -.09 & -.16 & .49 & -.10 & .85 & .80 & .81 \\ .88 & .79 & .07 & .73 & .91 & 1.00 & .27 & .53 & .12 & .06 & .66 & .08 & .90 & .85 & .78 \\ .15 & -.16 & .85 & -.25 & .20 & .27 & 1.00 & .85 & .81 & .79 & .79 & .81 & .43 & .54 & .53 \\ .39 & .10 & .75 & -.02 & .39 & .53 & .85 & 1.00 & .84 & .79 & .93 & .77 & .71 & .69 & .76 \\ -.02 & -.35 & .88 & -.45 & -.09 & .12 & .81 & .84 & 1.00 & .91 & .76 & .85 & .28 & .36 & .35 \\ -.16 & -.42 & .91 & -.57 & -.16 & .06 & .79 & .79 & .91 & 1.00 & .72 & .96 & .26 & .28 & .29 \\ .52 & .25 & .67 & .10 & .49 & .66 & .79 & .93 & .76 & .72 & 1.00 & .72 & .75 & .77 & .78 \\ -.15 & -.40 & .88 & -.53 & -.10 & .08 & .81 & .77 & .85 & .96 & .72 & 1.00 & .33 & .32 & .34 \\ .79 & .68 & .21 & .58 & .85 & .90 & .43 & .71 & .28 & .26 & .75 & .33 & 1.00 & .86 & .92 \\ .78 & .60 & .31 & .54 & .80 & .85 & .54 & .69 & .36 & .28 & .77 & .32 & .86 & 1.00 & .82 \\ .73 & .62 & .25 & .50 & .81 & .78 & .53 & .76 & .35 & .29 & .78 & .34 & .92 & .82 & 1.00 \end{pmatrix}$$




Figure 13.5 Scree graph for voice data.

The eigenvalues of \mathbf{R} are 7.91, 5.85, .31, .26, . . . , .002, with the scree plot in Figure 13.5. Clearly, by any criterion for choosing m , there are two factors.

All four major methods of factor extraction discussed in Section 13.3 produced nearly identical results (after rotation). We give the initial and rotated loadings obtained from the principal component method in Table 13.10.

The two rotated factors were labeled “competence” and “benevolence.” The same two factors emerged consistently in similar studies with different voices and different judges.

The two groupings of variables can also be seen in the correlation matrix. For example, in the first row, the large correlations correspond to the boldface rotated loadings for f_1 , while in the third row, the large correlations correspond to the boldface rotated loadings for f_2 .

The factor scores were of primary interest in this study. The goal was to ascertain the effect of the rate manipulations on the two factors, that is, to determine the perceived change in competence and benevolence when the speaking rate is increased or decreased.

The two factor scores were obtained for each of the 20 voices; these are plotted in Figure 13.6, where a consistent effect of the manipulation of speaking rate on all four voices can clearly be seen. Decreasing the speaking rate causes the speaker to be rated less competent; increasing the rate causes the speaker to be rated less benevolent. The mean vectors (centroids) are also given in Figure 13.6 for the four speakers. □

Table 13.10 Initial and Varimax Rotated Loadings for the Voice Data

Variable	Initial Loadings		Rotated Loadings		Communalities
	f_1	f_2	f_1	f_2	
Intelligent	.71	-.65	.96	-.06	.93
Ambitious	.48	-.84	.90	-.36	.94
Polite	.50	.81	-.12	.95	.92
Active	.37	-.91	.86	-.48	.97
Confident	.73	-.64	.97	-.04	.95
Happy	.83	-.47	.94	.15	.91
Just	.71	.58	.20	.89	.84
Likeable	.89	.39	.45	.87	.95
Kind	.58	.75	-.02	.95	.89
Sincere	.52	.82	-.11	.97	.95
Dependable	.93	.27	.56	.79	.94
Religious	.55	.79	-.07	.96	.92
Good-looking	.91	-.29	.89	.35	.91
Sociable	.91	-.22	.84	.40	.87
Strong	.91	-.21	.84	.41	.86
Variance accounted for	7.91	5.85	7.11	6.65	13.76
Proportion of total variance	.53	.39	.47	.44	.92
Cumulative proportion	.53	.92	.47	.92	.92

13.7 VALIDITY OF THE FACTOR ANALYSIS MODEL

For many statisticians, factor analysis is controversial and does not belong in a tool kit of legitimate multivariate techniques. The reasons for this mistrust include the following: the difficulty in choosing m , the many methods of extracting factors, the many rotation techniques, and the subjectivity in interpretation. Some statisticians also criticize factor analysis because of the indeterminacy of the factor loading matrix Λ or $\hat{\Lambda}$, first noted in Section 13.2.2. However, it is the ability to rotate that gives factor analysis its utility, if not its charm.

The basic question is whether the factors really exist. The model (13.11) for the covariance matrix is $\Sigma = \Lambda\Lambda' + \Psi$ or $\Sigma - \Psi = \Lambda\Lambda'$, where $\Lambda\Lambda'$ is of rank m . Many populations have covariance matrices that do not approach this pattern unless m is large. Thus the model will not fit data from such a population when we try to impose a small value of m . On the other hand, for a population in which Σ is reasonably close to $\Lambda\Lambda' + \Psi$ for small m , the sampling procedure leading to \mathbf{S} may obscure this pattern. The researcher may believe there are underlying factors but has difficulty collecting data that will reveal them. In many cases, the basic problem is




Figure 13.6 Factor scores of adjective rating of voices with five levels of manipulated rate.

that **S** (or **R**) contains both structure and error, and the methods of factor analysis cannot separate the two.

A statistical consultant in a university setting or elsewhere all too often sees the following scenario. A researcher designs a long questionnaire, with answers to be given in, say, a five-point semantic differential scale or Likert scale. The respondents, who vary in attitude from uninterested to resentful, hurriedly mark answers that in many cases are not even good subjective responses to the questions. Then the researcher submits the results to a handy factor analysis program. Being disappointed in the results, he or she appeals to a statistician for help. They attempt to improve the results by trying different methods of extraction, different rotations, different values of m , and so on. But it is all to no avail. The scree plot looks more like foothills than a steep cliff with gently sloping debris at the bottom. There is no clear value of m . They have to extract 10 or 12 factors to account for, say, 60% of the variance, and interpretation of this large number of factors is hopeless. If a few underlying dimensions exist, they are totally obscured by both systematic and random errors in marking the questionnaire. A factor analysis model simply does not fit such a data set, unless a large value of m is used, which gives useless results.

It is not necessarily the “discreteness” of the data that causes the problem, but the “noisiness” of the data. The specified variables are not measured accurately. In some cases, discrete variables yield satisfactory results, such as in Examples 13.3.1, 13.3.2, 13.5.2a, and 13.5.2b(a), where a 12-year-old girl, responding carefully to a semantic

differential scale, produced data leading to an unambiguous factor analysis. On the other hand, continuous variables do not guarantee good results [see Example 13.7(a) below].

In cases in which some factors are found that provide a satisfactory fit to the data, we should still be tentative in interpretation until we can independently establish the existence of the factors. If the same factors emerge in repeated sampling from the same population or a similar one, then we can have confidence that application of the model has uncovered some real factors. Thus it is good practice to repeat the experiment to check the stability of the factors. If the data set is large enough, it could be split in half and a factor analysis performed on each half. The two solutions could be compared with each other and with that for the complete set.

If there is replication in the data set, it may be helpful to average over the replications. This was done to great advantage in Example 13.6, where several judges rated the same voices. Averaging over the judges produced variables that apparently possessed very low noise. Similar experimentation with different judges always produced the same factors. Unfortunately, replication of this type is unavailable in most situations.

As with other techniques in this book, factor analysis assumes that the variables are at least approximately linearly related to each other. We could make bivariate scatterplots to check this assumption.

A basic prerequisite for a factor analysis application is that the variables not be independent. To check this requirement, we could test $H_0: \mathbf{P}_\rho = \mathbf{I}$ by using the test in Section 7.4.3.

Some writers have suggested that \mathbf{R}^{-1} should be a near-diagonal matrix in order to successfully fit a factor analysis model. To assess how close \mathbf{R}^{-1} is to a diagonal matrix, Kaiser (1970) proposed a *measure of sampling adequacy*,

$$\text{MSA} = \frac{\sum_{i \neq j} r_{ij}^2}{\sum_{i \neq j} r_{ij}^2 + \sum_{i \neq j} q_{ij}^2}, \quad (13.60)$$

where r_{ij}^2 is the square of an element from \mathbf{R} and q_{ij}^2 is the square of an element from $\mathbf{Q} = \mathbf{D}\mathbf{R}^{-1}\mathbf{D}$, with $\mathbf{D} = [(\text{diag } \mathbf{R}^{-1})^{1/2}]^{-1}$. As \mathbf{R}^{-1} approaches a diagonal matrix, MSA approaches 1. Kaiser and Rice (1974) suggest that MSA should exceed .8 for satisfactory results to be expected. We show some results for MSA in Example 13.7(b).

In summary, there are many data sets to which factor analysis should not be applied. One indication that \mathbf{R} is inappropriate for factoring is the failure of the methods in Section 13.4 to clearly and rather objectively choose a value for m . If the scree plot does not have a pronounced bend or the eigenvalues do not show a large gap around 1, then \mathbf{R} is likely to be unsuitable for factoring. In addition, the communality estimates after factoring should be fairly large.

To balance the “good” examples in this chapter, we now give an example involving a data set that cannot be successfully modeled by factor analysis. Likewise, the problems at the end of the chapter include both “good” and “bad” data sets.




Figure 13.7 Scree graph for diabetes data.

■ EXAMPLE 13.7(a)

As an illustration of an application of factor analysis that is less successful than previous examples in this chapter, we consider the diabetes data of Table 3.4. The correlation matrix for the five variables is as follows:

$$\mathbf{R} = \begin{pmatrix} 1.00 & .05 & -.13 & .07 & .21 \\ .05 & 1.00 & -0.1 & .01 & -.10 \\ -.13 & -0.1 & 1.00 & .29 & .05 \\ .07 & .01 & .29 & 1.00 & .21 \\ .21 & -.10 & .05 & .21 & 1.00 \end{pmatrix}.$$

The correlations are all small, and the variables do not appear to have much in common. The MSA value is .49. The eigenvalues are 1.40, 1.21, 1.04, .71, and .65. Three factors would be required to account for 73% of the variance and four factors to reach 87%. This is not a useful reduction in dimensionality. The eigenvalues are plotted in a scree graph in Figure 13.7. The lack of a clear value of m is apparent.

It is evident from the small correlations in \mathbf{R} that the communalities of the variables will not be large. The principal component method, which essentially estimates the initial communalities as 1, gave very different final communality estimates than did the iterated principal factor method:

	Communalities				
Principal component method	.71	.91	.71	.67	.64
Iterated principal factor method	.31	.16	.35	.37	.33

Table 13.11 Varimax Rotated Factor Loadings for Iterated Principal Factors from the Diabetes Data

Variable	Rotated Loadings			Communalities
	f_1	f_2	f_3	
1	-.08	.54	.12	.31
2	.01	.01	.40	.16
3	.57	-.15	-.03	.35
4	.57	.22	.02	.37
5	.19	.47	-.27	.33
Variance accounted for	.69	.59	.24	1.52

The communalities obtained by the iterated approach reflect more accurately the small correlations among the variables.

The varimax rotated factor loadings for three factors extracted by the iterated principal factor method are given in Table 13.11. The first factor is associated with variables 3 and 4, the second factor with variables 1 and 5, and the third with variable 2. This clustering of variables can be seen in \mathbf{R} , where variables 1 and 5 have a correlation of .21, variables 3 and 4 have a correlation of .29, and variable 2 has very low correlations with all other variables. However, these correlations (.21 and .29) are small, and in this case the collapsing of five variables to three factors is not a useful reduction in dimensionality, especially since the first three eigenvalues account for only 73% of $\text{tr}(\mathbf{R})$. The 73% is not convincingly greater than 60%, which we would expect from three original variables picked at random. This conclusion is borne out by a test of $H_0: \mathbf{P}_\rho = \mathbf{I}$. Using (7.37) and (7.38), we obtain

$$u = |\mathbf{R}| = .80276, \quad \nu = 20 - 1 = 19, \quad p = 5,$$

$$u' = -[\nu - \frac{1}{6}(2p + 5)] \ln u = -\left(19 - \frac{15}{6}\right)(-.2197) = 3.625.$$

With $\frac{1}{2}p(p - 1) = 10$ degrees of freedom, the .05 critical value for this approximate χ^2 test is 18.31, and we have no basis to question the independence of the five variables. Thus the three factors we obtained are very likely an artifact of the present sample and would not reappear in another sample from the same population. \square

■ EXAMPLE 13.7(b)

For data sets used in previous examples in this chapter, the values of MSA from (13.60) are calculated as follows:

Seishu data: MSA = .53
Sons data: MSA = .82
Voice data: MSA = .73
Diabetes data: MSA = .49.

The MSA value cannot be computed for the perception data, because \mathbf{R} is singular.

These results do not suggest great confidence in the MSA index as a sole guide to the suitability of \mathbf{R} for factoring. We see a wide disparity in the MSA values for the first three data sets. Yet all three yielded successful factor analyses. These three MSA values seem to be inversely related to the number of factors: In the sons data, there were indications that one factor would suffice; the voice data clearly had two factors; and for the Seishu data, there were four factors.

The MSA for the diabetes data is close to that of the Seishu data. Yet the diabetes data are totally unsuitable for factor analysis, while the factor analysis of the Seishu data is very convincing. \square

13.8 RELATIONSHIP OF FACTOR ANALYSIS TO PRINCIPAL COMPONENT ANALYSIS

Both factor analysis and principal component analysis have the goal of reducing dimensionality. Because the objectives are similar, many authors discuss principal component analysis as another type of factor analysis. This can be confusing, and we wish to underscore the distinguishing characteristics of the two techniques.

Two of the differences between factor analysis and principal component analysis were mentioned in Section 13.1: (1) In factor analysis, the variables are expressed as linear combinations of the factors, whereas the principal components are linear functions of the variables, and (2) in principal component analysis, the emphasis is on explaining the total variance $\sum_i s_{ii}$, as contrasted with the attempt to explain the covariances in factor analysis.

Additional differences are that (3) principal component analysis requires essentially no assumptions, while factor analysis makes several key assumptions; (4) the principal components are unique (assuming distinct eigenvalues of \mathbf{S}), whereas the factors are subject to an arbitrary rotation; and (5) if we change the number of factors, the (estimated) factors change. This does not happen in principal components.

The ability to rotate to improve interpretability is one of the advantages of factor analysis over principal components. If finding and describing some underlying factors is the goal, factor analysis may prove more useful than principal components; we would prefer factor analysis if the factor model fits the data well and we like the interpretation of the rotated factors. On the other hand, if we wish to define a smaller number of variables for input into another analysis, we would ordinarily pre-

fer principal components, although this can sometimes be accomplished with factor scores. Occasionally, principal components are interpretable, as in the size and shape components in Example 12.8.1.

PROBLEMS

- 13.1** Show that the assumptions lead to (13.2), $\text{var}(y_i) = \lambda_{i1}^2 + \lambda_{i2}^2 + \cdots + \lambda_{im}^2 + \psi_i$.
- 13.2** Verify directly that $\text{cov}(\mathbf{y}, \mathbf{f}) = \boldsymbol{\Lambda}$ as in (13.13).
- 13.3** Show that $\mathbf{f}^* = \mathbf{T}'\mathbf{f}$ in (13.18) satisfies the assumptions (13.6) and (13.7), $E(\mathbf{f}^*) = \mathbf{0}$ and $\text{cov}(\mathbf{f}^*) = \mathbf{I}$.
- 13.4** Show that $\sum_{ij} e_{ij}^2 \leq \theta_{m+1}^2 + \theta_{m+2}^2 + \cdots + \theta_p^2$ as in (13.34), where the e_{ij} 's are the elements of $\mathbf{E} = \mathbf{S} - (\hat{\boldsymbol{\Lambda}}\hat{\boldsymbol{\Lambda}}' + \hat{\boldsymbol{\Psi}})$ and the θ_i 's are eigenvalues of \mathbf{S} .
- 13.5** Show that $\sum_{i=1}^p \sum_{j=1}^m \hat{\lambda}_{ij}^2$ is equal to the sum of the first m eigenvalues and also equal to the sum of all p communalities, as in (13.46).
- 13.6** In Example 13.3.2, the correlation matrix for the perception data was shown to have an eigenvalue equal to 0. Find the multicollinearity among the five variables that this implies.
- 13.7** Use the words data of Table 5.9.
 - (a)** Obtain principal component loadings for two factors.
 - (b)** Do a graphical rotation of the two factors.
 - (c)** Do a varimax rotation and compare the results with those in part (b).
- 13.8** Use the ramus bone data of Table 3.7.
 - (a)** Extract loadings by the principal component method and do a varimax rotation. Use two factors.
 - (b)** Do all variables have a complexity of 1? Carry out an oblique rotation to improve the loadings.
 - (c)** What is the angle between the oblique axes? Would a single factor ($m = 1$) be more appropriate here?
- 13.9** Carry out a factor analysis of the rootstock data of Table 6.2. Combine the six groups into a single sample.
 - (a)** Estimate the loadings for two factors by the principal component method and do a varimax rotation.
 - (b)** Did the rotation improve the loadings?
- 13.10** Use the fish data of Table 6.17. Combine the three groups into a single sample.

- (a) Obtain loadings on two factors by the principal component method and do a varimax rotation.
- (b) Note the similarity of loadings for y_1 and y_2 . Is there any indication in the correlation matrix as to why this is so?
- (c) Compute factor scores.
- (d) Using the factor scores, carry out a MANOVA comparing the three groups.
- 13.11** Carry out a factor analysis of the flea data in Table 5.5. Combine the two groups into a single sample.
- (a) From an examination of the eigenvalues greater than 1, the scree plot, and the percentages, is there a clear choice of m ?
- (b) Extract two factors by the principal component method and carry out a varimax rotation.
- (c) Is the rotation an improvement? Try an oblique rotation.
- 13.12** Use the engineer data of Table 5.6. Combine the two groups into a single sample.
- (a) Using a scree plot, the number of eigenvalues greater than 1, and the percentages, is there a clear choice of m ?
- (b) Extract three factors by the principal component method and carry out a varimax rotation.
- (c) Extract three factors by the principal factor method and carry out a varimax rotation.
- (d) Compare the results of parts (b) and (c).
- 13.13** Use the probe word data of Table 3.6.
- (a) Obtain loadings for two factors by the principal component method and carry out a varimax rotation.
- (b) Note the near duplication of loadings for y_2 and y_4 . Is there any indication in the correlation matrix as to why this is so?
- (c) Is the rotation satisfactory? Try an oblique rotation.

CHAPTER 14

CONFIRMATORY FACTOR ANALYSIS

14.1 INTRODUCTION

In Chapter 13, we emphasized that exploratory factor analysis (EFA) is used to give a low-dimensional description of high-dimensional data, but is not based on a uniquely identifiable model. That is, no sample of data—regardless of how large—will yield a scenario in which the “true” values of the EFA model can be identified. This is due to the over-parameterized nature of the EFA model, as described in Sections 13.2.2 and 13.7. Because of the nonuniqueness of the factor loadings and other model parameters, practitioners of EFA typically find a solution to the model and then rotate the factor loadings to find a more interpretable or convenient definition of the latent factors (see Section 13.5).

As in the EFA model (13.1), the factors f_j in the *confirmatory factor analysis* (CFA) model are used to represent underlying latent variables that are of principal interest but are difficult or impossible to measure. For example, an educational researcher may be interested in a students’ attitudes about facets of their class experience, or a physician may be interested in assessing various dimensions of quality of life for patients being treated for cancer. In such cases where primary interest is in attitudes, ideas, or other hard-to-measure variables, the factor analysis approach will

be of particular utility. Thus, along with item response theory, CFA is a frequently used tool in the development and analysis of assessment tests and questionnaires. In fact, many item response theory models are special cases of confirmatory factor analysis.

What sets CFA apart from EFA (Chapter 13) is that constraints will be incorporated into the model to ensure a uniquely identifiable set of model parameters. Thus, CFA will follow the usual statistical paradigm—hypothesizing an identifiable model, fitting model parameters, assessing the goodness of model fit, and performing statistical inference for model parameters using conventional tools such as hypothesis tests and confidence intervals. CFA is often considered to be a subset or special case of structural equation modeling, wherein unobserved factors can have structural relationships. For example, a structural equation model might posit that one factor is predicted by another factor in a manner analogous to the way a response (dependent) variable y is affected by a predictor (independent) variable x in the simple linear regression model: $y = \beta_0 + \beta_1 x + \varepsilon$. In contrast, CFA allows the estimation of multiple correlated factors but does not allow the factors to have predictor/response relationships.

Throughout our discussion, we emphasize facets of CFA that are most directly useful to the practitioner. Readers interested in technical details, derivations, or extensions to structural equation modeling should refer to the excellent treatments of CFA found in Fuller (1987; Sections 1.5 and 4.3) and Bollen (1989).

14.2 MODEL SPECIFICATION AND IDENTIFICATION

14.2.1 Confirmatory Factor Analysis Model

When employing confirmatory factor analysis, we use an alternative approach for fitting the general factor analysis model that was first introduced as (13.1):

$$\begin{aligned} y_1 &= \mu_1 + \lambda_{11}f_1 + \lambda_{12}f_2 + \cdots + \lambda_{1m}f_m + \varepsilon_1 \\ y_2 &= \mu_2 + \lambda_{21}f_1 + \lambda_{22}f_2 + \cdots + \lambda_{2m}f_m + \varepsilon_2 \\ &\vdots \\ y_p &= \mu_p + \lambda_{p1}f_1 + \lambda_{p2}f_2 + \cdots + \lambda_{pm}f_m + \varepsilon_p. \end{aligned} \tag{14.1}$$

In matrix form, we write the model:

$$\mathbf{y} = \boldsymbol{\mu} + \mathbf{\Lambda}\mathbf{f} + \boldsymbol{\varepsilon}. \tag{14.2}$$

As in the EFA setting, the factor loading λ_{ij} in (14.1) quantifies the influence of the factor f_j on the observed variable y_i , and the mean and specific factor (or error) components for y_i are still defined as μ_i and ε_i , respectively. The factor vector

$\mathbf{f} = (f_1, \dots, f_m)'$ has mean $\mathbf{0}$ and covariance matrix

$$\text{cov}(\mathbf{f}) = \Phi = \begin{pmatrix} \phi_{11} & \phi_{12} & \cdots & \phi_{1m} \\ \phi_{21} & \phi_{22} & \cdots & \phi_{2m} \\ \vdots & \vdots & \ddots & \vdots \\ \phi_{m1} & \phi_{m2} & \cdots & \phi_{mm} \end{pmatrix} \quad (14.3)$$

and the error vector $\boldsymbol{\varepsilon} = (\varepsilon_1, \dots, \varepsilon_p)'$ has mean $\mathbf{0}$ and diagonal covariance matrix

$$\text{cov}(\boldsymbol{\varepsilon}) = \Psi = \begin{pmatrix} \psi_{11} & 0 & \cdots & 0 \\ 0 & \psi_{22} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \psi_{pp} \end{pmatrix}. \quad (14.4)$$

The vectors \mathbf{f} and $\boldsymbol{\varepsilon}$ are independent. The diagonality assumption for Ψ can be relaxed in more general forms of the model, but we focus here on the classical model, which assumes that all of the correlation among the observed variables is due to the factors.

Note that model (14.2) implies a specific structure for the covariance matrix for \mathbf{y} . When the model holds, we can define the population covariance matrix for \mathbf{y} to be

$$\Sigma(\boldsymbol{\theta}) = \Lambda\Phi\Lambda' + \Psi, \quad (14.5)$$

where Λ , Φ , and Ψ are defined in (14.2), (14.3), and (14.4), and $\boldsymbol{\theta}$ is the collection of the unique parameters in the model. For the unconstrained model (14.2),

$$\boldsymbol{\theta} = (\lambda', \phi', \psi')', \quad (14.6)$$

where λ contains the pm factor loadings in Λ , ϕ contains the $m(m+1)/2$ factor variances and covariances in the symmetric matrix Φ , and ψ contains the p error variances in the diagonal matrix Ψ . When model (14.2) holds, $\Sigma(\boldsymbol{\theta})$ is completely specified by the parameter vector $\boldsymbol{\theta}$.

A model formulation such as (14.1) or (14.2) is one way of describing the relationships between observed and latent variables. Equally popular is the use of the *path diagram*, wherein a latent variable model such as those described above is illustrated graphically. In a path diagram, observed variables are denoted with rectangles and latent variables (factors) are denoted with ovals. For example, Figure 14.1 illustrates the path diagram for the multiple regression model $y = \beta_1x_1 + \beta_2x_2 + \varepsilon$, with $\text{var}(\varepsilon) = \sigma^2$. Note that a one-headed arrow denotes a structural or unidirectional relationship, while a two-headed arrow denotes a covariance among variables. A two-headed arrow pointing from one variable and back to itself is used to specify the variance of the variable, which is usually an error or “disturbance” variable. As CFA models become increasingly complex, the use of path diagrams becomes increasingly attractive for efficiently communicating the hypothesized model.




Figure 14.1 Path diagram for multiple regression model $y = \beta_1x_1 + \beta_2x_2 + \varepsilon$, with $\text{var}(\varepsilon) = \sigma^2$.

14.2.2 Identified Models

As discussed in Sections 13.2.2 and 14.1, the unconstrained factor analysis model (14.1) is (in general) *underidentified*. That is, there is no unique solution for the parameter vector $\boldsymbol{\theta}$ in (14.5). More specifically, we say that a model is underidentified if it is possible to find two different vectors $\boldsymbol{\theta}_1$ and $\boldsymbol{\theta}_2$ such that $\Sigma(\boldsymbol{\theta}_1) = \Sigma(\boldsymbol{\theta}_2)$. Identification problems often occur when there are so many free parameters in $\boldsymbol{\theta}$ that there are many different values for the vector $\boldsymbol{\theta}$ that provide a solution to the equations. Every two-factor EFA model is underidentified because the factor loadings can be rotated without affecting the implied covariance matrix $\Sigma(\boldsymbol{\theta})$. In the CFA setting, our intent is to impose phenomenologically based constraints on the model in order to ensure that the model is *identified*. That is, we wish to guarantee that there exists a unique solution for the parameter vector $\boldsymbol{\theta}$.

A necessary but not sufficient condition for a model to be identified is that the number of unique statistics in the sample covariance matrix is at least as large as the number of parameters needed to specify the population covariance matrix under the hypothesized model. This is sometimes called the *order condition*. Within a $p \times p$ covariance matrix \mathbf{S} , there are $p(p+1)/2$ unique statistics—the variances and covariances. Given the unconstrained model (14.2), the parameter vector $\boldsymbol{\theta}$ described in (14.5) has pm factor loadings, $m(m+1)/2$ factor variances and covariances, and p error variances. Thus, for the unconstrained model, the order condition is met when

$$\frac{p(p+1)}{2} \geq pm + \frac{m(m+1)}{2} + p.$$

However, even when the order condition is met, say when $p = 9$ and $m = 3$ in model (14.1), we are still not guaranteed identifiability because of the way the model parameters interact. For example, the model will have equivalent goodness of fit (i) when ϕ_{11} is large and the loadings $\lambda_{11}, \dots, \lambda_{p1}$ are small and (ii) when ϕ_{11} is small and the loadings $\lambda_{11}, \dots, \lambda_{p1}$ are large. This indeterminacy associated with the scale of the latent factors is one cause of the nonidentifiability in the unconstrained model, even when the order condition is met.

There are many different ways that one can constrain model (14.1) to ensure that the parameters are uniquely identified. One widely used approach for obtaining an identified factor analysis model is to rewrite model (14.2) using the parameterization:

$$\begin{pmatrix} \mathbf{y}_0 \\ m \times 1 \\ \mathbf{y}_1 \\ (p-m) \times 1 \end{pmatrix} = \begin{pmatrix} \mathbf{0} \\ m \times 1 \\ \boldsymbol{\mu}_1 \\ (p-m) \times 1 \end{pmatrix} + \begin{pmatrix} \mathbf{I} \\ m \times m \\ \boldsymbol{\Lambda}_1 \\ (p-m) \times m \end{pmatrix} \mathbf{f} + \begin{pmatrix} \boldsymbol{\varepsilon}_0 \\ m \times 1 \\ \boldsymbol{\varepsilon}_1 \\ (p-m) \times 1 \end{pmatrix}. \quad (14.7)$$

Note that this reparameterization of (14.2) constrains the first m rows of $\boldsymbol{\Lambda}$ in (14.2) to be equal to the identity matrix, and the first m elements of $\boldsymbol{\mu}$ to be $\mathbf{0}$. This implies that each of the first m elements of \mathbf{y} in (14.2) is equal to one of the m factors plus error. Specifically, $y_i = f_i + \varepsilon_i$, $i = 1, \dots, m$. Equation (14.7) is called the *errors-in-variables* or *measurement error* parameterization because it is analogous to the errors-in-variables regression problem where the linear model of interest is $\mathbf{y}_1 = \boldsymbol{\mu}_1 + \boldsymbol{\Lambda}_1 \mathbf{f} + \boldsymbol{\varepsilon}_1$ but the predictor variables \mathbf{f} are only observable in their measurement-error-contaminated form: $\mathbf{y}_0 = \mathbf{f} + \boldsymbol{\varepsilon}_0$ (see Fuller, 1987, Section 1.3.1).

Although \mathbf{f} is related to all p of the observed variables, (14.7) effectively constrains f_1, \dots, f_m to have the same means as y_1, \dots, y_m , respectively. Further, the variances of f_1, \dots, f_m are bounded above by the variances of y_1, \dots, y_m , respectively. In practice, the modeler uses her a priori hypothesis about the nature of the multivariate structure in \mathbf{y} to select which observed variables should be the indicator variables \mathbf{y}_0 that are most closely associated with the factors \mathbf{f} . As we will show in Section 14.3, the validity of the hypothesis-driven model will be formally evaluated for goodness of fit before it is abandoned or tentatively accepted.

The order condition for model (14.7) is satisfied when

$$\frac{p(p+1)}{2} \geq (p-m)m + \frac{m(m+1)}{2} + p \quad (14.8)$$

or, equivalently (see Problem 14.1), when

$$(p-m)^2 \geq p+m.$$

For example, the errors-in-variables parameterization of the CFA model satisfies the order condition for one factor when $p \geq 3$, for two factors when $p \geq 5$, for three factors when $p \geq 6$, and so forth. However, an even more general criterion for satisfying the model's order condition [regardless of whether (14.7) is satisfied] is simply that

$$\frac{p(p+1)}{2} \geq q, \quad (14.9)$$

where q is the total number of unique parameters to be estimated in $\boldsymbol{\Lambda}$, $\boldsymbol{\Phi}$, and $\boldsymbol{\Psi}$. That is, q is the dimension of the parameter vector $\boldsymbol{\theta}$.

The indeterminacy in the unconstrained version of (14.2) is eliminated in model (14.7) by (i) satisfying the order condition in (14.8) or (14.9) and (ii) replacing an $m \times m$ submatrix of $\boldsymbol{\Lambda}$ with the rank m matrix \mathbf{I}_m . However, model (14.2) will in fact be identified as long as the order condition in (14.9) is satisfied and any m rows

of Λ are replaced by any full-rank $m \times m$ matrix of constants \mathbf{K} . Assuming that inference on μ is not of direct interest (which is usually the case), a more general model is:

$$\mathbf{y} = \boldsymbol{\mu} + \begin{pmatrix} \mathbf{K} \\ \mathbf{\Lambda}_1 \\ \vdots \\ \mathbf{0}_{(p-m) \times m} \end{pmatrix} \mathbf{f} + \boldsymbol{\varepsilon}. \quad (14.10)$$

Additional constants can be included in $\mathbf{\Lambda}_1$ and still have an identified model. Further, constraints such as setting some factor loadings equal to each other can often be accommodated in an identified model. A careful review of a variety of identifiability conditions for CFA can be found in the latent variable modeling literature (e.g., Bollen, 1987, Chapter 7). For our present purposes, it suffices to conclude that the modeler can fix the physical meaning of the factors in any desired way that satisfies the order condition along with some other identifiability-inducing constraint of the model, with the parameterizations in models (14.7) and (14.10) being examples of such acceptable constraints. As with the modeling choices associated with model (14.7), the nature of the imposed constraints utilized in model (14.10) will depend on the phenomenology under consideration and the a priori hypothesis about the nature of the covariances among the observed variables.

■ EXAMPLE 14.2.2

To illustrate the formulation of a CFA model, we use the grade data given in Table 14.1, which were obtained from a university business statistics class. As shown in Table 14.1, each of the 94 students in the class received scores for laboratory assignments (Lab), homework assignments (HW), pop quizzes (PopQuiz), midterm exam #1 (Exam1), midterm exam #2 (Exam2), and the final exam (FinalExam). The instructor might hypothesize that the 6-dimensional measure of performance in the class is being driven by an underlying 2-dimensional factor process, with the first factor associated with daily effort and the second factor associated with knowledge mastery. Consequently, we might express the model using the parameterization in (14.10):

$$\begin{aligned} \text{Lab} &= \mu_1 + \lambda_{11}f_1 + \lambda_{12}f_2 + \varepsilon_1 \\ \text{HW} &= \mu_2 + f_1 + \varepsilon_2 \\ \text{PopQuiz} &= \mu_3 + \lambda_{31}f_1 + \lambda_{32}f_2 + \varepsilon_3 \\ \text{Exam1} &= \mu_4 + \lambda_{41}f_1 + \lambda_{42}f_2 + \varepsilon_4 \\ \text{Exam2} &= \mu_5 + \lambda_{51}f_1 + \lambda_{52}f_2 + \varepsilon_5 \\ \text{FinalExam} &= \mu_6 + f_2 + \varepsilon_6 \end{aligned} \quad (14.11)$$

The order condition is satisfied for model (14.11) because the number of statistics in the sample covariance matrix \mathbf{S} [the left side of the inequality in (14.8) and (14.9)] is 21 and the number of parameters (q) in the model [the right side of the inequality in (14.8) and (14.9)] is 17. Additionally, the second and sixth rows of the model's Λ matrix form an identity matrix of rank 2. Thus,

Table 14.1 Statistics Course Grade Measurements

Lab	HW	PopQuiz	Exam1	Exam2	FinalExam
9.95	9.40	8.13	72	90	62
7.71	0.00	8.00	63	26	78
10.00	9.50	8.13	79	55	80
9.95	9.40	8.88	89	77	79
9.95	9.20	8.50	83	84	67
10.00	9.90	9.63	96	94	93
10.00	9.25	9.25	78	85	86
10.00	9.50	8.50	71	83	66
10.00	9.15	8.13	91	91	91
8.43	9.00	8.50	76	61	80
7.38	7.35	8.50	65	62	42
9.86	9.00	6.63	58	21	31
9.86	9.60	7.63	98	84	87
9.67	8.30	7.38	89	62	75
10.00	9.40	8.13	71	84	78
10.00	9.55	9.25	100	72	87
10.00	8.10	6.88	90	72	85
2.00	0.95	8.13	62	45	37
10.00	7.65	8.88	53	47	61
10.00	6.30	6.00	77	74	73
9.00	6.75	9.63	85	79	65
9.86	9.55	7.38	85	61	70
8.43	9.25	7.00	82	77	83
10.00	8.25	8.13	80	75	85
⋮	⋮	⋮	⋮	⋮	⋮
10.00	9.85	10.00	100	100	100




Figure 14.2 Path diagram for the grade data model.

model (14.11) is identified. The choice to constrain the HW and FinalExam variables as in model (14.11) was based on a priori expectations about the nature of these measures roughly corresponding to the effort and mastery factors, respectively. While choices about how to specify model structure are largely subject to the discretion of the modeler, the choice of which variables are constrained in the model does not generally affect goodness-of-fit considerations (see Problem 14.4).

In this study however, the theory we wish to test is that most variables load on only one of the two factors, with the pop quiz score being the only course component that is influenced by both daily effort *and* knowledge mastery. That is, the model we wish to test is

$$\begin{aligned}
 \text{Lab} &= \mu_1 + \lambda_{11}f_1 &+ \varepsilon_1 \\
 \text{HW} &= \mu_2 + f_1 &+ \varepsilon_2 \\
 \text{PopQuiz} &= \mu_3 + \lambda_{31}f_1 + \lambda_{32}f_2 + \varepsilon_3 \\
 \text{Exam1} &= \mu_4 &+ \lambda_{42}f_2 + \varepsilon_4 \\
 \text{Exam2} &= \mu_5 &+ \lambda_{52}f_2 + \varepsilon_5 \\
 \text{FinalExam} &= \mu_6 &+ f_2 + \varepsilon_6
 \end{aligned} \tag{14.12}$$

The path diagram associated with model (14.12) is shown in Figure 14.2. The number of statistics in the sample covariance matrix \mathbf{S} is still 21 as it was for model (14.11), but there are fewer parameters in model (14.12). There are 5 unconstrained factor loadings in Λ , $m(m+1)/2 = 3$ elements in the covariance matrix for \mathbf{f} , and $p = 6$ diagonal elements in Ψ , for a total of $q = 14$ parameters in the model. Because the order condition in (14.9) is satisfied, and because the second and sixth rows of the model's Λ matrix still form an identity matrix of rank 2, model (14.12) is also identified. \square

14.3 PARAMETER ESTIMATION AND MODEL ASSESSMENT

At an intuitive level, our approach for estimating the model parameters $\boldsymbol{\theta}$ is to find a $\boldsymbol{\theta}$ within a parameter space Θ such that the difference between \mathbf{S} and $\Sigma(\boldsymbol{\theta})$ from (14.5) is minimized. The parameter space Θ represents the collection of $\boldsymbol{\theta}$ values that yield matrices $\Sigma(\boldsymbol{\theta})$ that are nonnegative definite. We note here that a careful definition of the parameter space Θ is not merely a mathematical formality but an essential part of the parameter estimation and model assessment process. For example, proper parameter estimation will involve software that constrains factor and error variances to be nonnegative. In other settings, factor loadings or other parameters may be constrained to be equal. Such constraints will be illustrated in Example 14.3.3.

We begin by assuming that \mathbf{y} follows a multivariate normal distribution with a mean $\boldsymbol{\mu}$ and a covariance matrix $\Sigma(\boldsymbol{\theta})$. We will then apply the results of Anderson and Amemiya (1988) and Amemiya and Anderson (1990) to argue that as the sample size becomes large, the normality-based parameter estimators and associated inferential tools are valid, even for nonnormal data.

14.3.1 Maximum Likelihood Estimation

In Sections 4.3.1 and 5.4.3 we introduced the notions of maximum likelihood estimation and likelihood ratio tests, respectively. In this section, we introduce maximum likelihood estimates for the q factor analysis model parameters $\boldsymbol{\theta}$. Throughout the discussion, we use the maximum likelihood estimate of the covariance matrix for \mathbf{y} , denoted $\mathbf{S}_n = 1/n \sum_{i=1}^n (\mathbf{y}_i - \bar{\mathbf{y}})'(\mathbf{y}_i - \bar{\mathbf{y}})$.

If we assume that the factor analysis model (14.2) holds and that \mathbf{y} follows a multivariate normal distribution, then \mathbf{S}_n follows a Wishart distribution and it can be shown that the likelihood for $\Sigma(\boldsymbol{\theta})$ given \mathbf{S}_n is

$$L[\Sigma(\boldsymbol{\theta}); \mathbf{S}_n] = c \times |\Sigma(\boldsymbol{\theta})|^{-n/2} \exp\left(-\frac{1}{2} \text{tr}\left\{n\mathbf{S}_n[\Sigma(\boldsymbol{\theta})]^{-1}\right\}\right),$$

where c is a constant that is unaffected by $\boldsymbol{\theta}$ or $\Sigma(\boldsymbol{\theta})$. The value of $\boldsymbol{\theta}$ that maximizes $L(\Sigma(\boldsymbol{\theta}); \mathbf{S}_n)$ is called the maximum likelihood estimator for $\boldsymbol{\theta}$ and is denoted $\hat{\boldsymbol{\theta}}_{\text{ML}}$.

The likelihood ratio statistic associated with the hypothesized factor analysis model is

$$LR(\hat{\boldsymbol{\theta}}_{\text{ML}}) = \frac{L[\Sigma(\hat{\boldsymbol{\theta}}_{\text{ML}}); \mathbf{S}_n]}{L[\Sigma(\boldsymbol{\theta}); \mathbf{S}_n]},$$

where the numerator is the likelihood evaluated with Σ at its maximum under the hypothesized model and the denominator is the likelihood evaluated with Σ at its unrestricted maximum (i.e., $\Sigma = \mathbf{S}_n$). Taking $-2 \log LR(\boldsymbol{\theta})$, we obtain

$$\ell(\boldsymbol{\theta}; \mathbf{S}_n) = n \left(\log |\Sigma(\boldsymbol{\theta})| + \text{tr}\left\{\mathbf{S}_n [\Sigma(\boldsymbol{\theta})]^{-1}\right\} - \log |\mathbf{S}_n| - p \right). \quad (14.13)$$

It can be shown that $\hat{\boldsymbol{\theta}}_{\text{ML}}$ minimizes $\ell(\boldsymbol{\theta}; \mathbf{S})$ over the parameter space Θ and that—regardless of the actual distribution of the data—the statistic $\ell(\hat{\boldsymbol{\theta}}_{\text{ML}}; \mathbf{S}_n)$ has an

asymptotic distribution of $\chi^2_{p(p+1)/2-q}$. Further, under some fairly tractable conditions, the factor loading estimates $\hat{\lambda}_{\text{ML}}$ converge to $\boldsymbol{\lambda}$ [as defined in (14.6)] when $n \rightarrow \infty$. As we will discuss in Section 14.3.3, the statistic $\ell(\boldsymbol{\theta}; \mathbf{S}_n)$ plays a central role in the assessment of the factor analysis model's goodness of fit.

14.3.2 Least Squares Estimation

As an alternative to maximum likelihood estimation, we consider here a computational approach based on least squares estimation under the assumption that \mathbf{y} in model (14.2) is distributed normally with mean $\boldsymbol{\mu}$ and covariance matrix $\boldsymbol{\Sigma}[\boldsymbol{\theta}]$. In this scenario, $(n - 1)\mathbf{S} \sim W_p(n - 1, \boldsymbol{\Sigma}[\boldsymbol{\theta}])$, where W_p denotes the Wishart distribution introduced in Section 4.3.2. Note that we are interested in the distribution of \mathbf{S} because we wish to make inferences about the q elements of $\boldsymbol{\theta}$, which are functions of second moments. Thus, just as carrying out inferences about means (first moments) requires information about second moments (variances and covariances), carrying out inferences associated with second moments will require fourth moments for the data. Although the normality assumption is not required for least squares estimation, it is a substantial benefit to assume normality because higher-order moments will always be simple functions of first and second moments. We will later argue that normality-based approaches are asymptotically valid even when the normal assumptions are violated.

Given the assumption of normality of the data, we can show that the $p(p + 1)/2 \times p(p + 1)/2$ covariance matrix for the unique elements of \mathbf{S} is:

$$\text{cov}(\text{vech } \mathbf{S}) = \frac{2}{n - 1} \mathbf{H}_p^+ [\boldsymbol{\Sigma}(\boldsymbol{\theta}) \otimes \boldsymbol{\Sigma}(\boldsymbol{\theta})] \mathbf{H}_p^{+'}.$$

The matrices \mathbf{H}_p and \mathbf{H}_p^+ are defined as in (2.125) and (2.126), respectively, so that for any $p \times p$ matrix \mathbf{A} , $\mathbf{H}_p \text{vech } \mathbf{A} = \text{vec } \mathbf{A}$ and $\mathbf{H}_p^+ \text{vec } \mathbf{A} = \text{vech } \mathbf{A}$. Further, the normality assumption implies that if $\boldsymbol{\Sigma}(\boldsymbol{\theta})$ is positive definite, then

$$[\text{cov}(\text{vech } \mathbf{S})]^{-1} = \frac{n - 1}{2} \mathbf{H}_p' \{ [\boldsymbol{\Sigma}(\boldsymbol{\theta})]^{-1} \otimes [\boldsymbol{\Sigma}(\boldsymbol{\theta})]^{-1} \} \mathbf{H}_p. \quad (14.14)$$

In practice, we will estimate $\text{cov}(\text{vech } \mathbf{S})$ with either

$$\hat{\mathbf{V}} = \frac{2}{n - 1} \mathbf{H}_p^+ (\mathbf{S} \otimes \mathbf{S}) \mathbf{H}_p^{+'} \quad (14.15)$$

or

$$\mathbf{V}(\hat{\boldsymbol{\theta}}) = \frac{2}{n - 1} \mathbf{H}_p^+ [\boldsymbol{\Sigma}(\hat{\boldsymbol{\theta}}) \otimes \boldsymbol{\Sigma}(\hat{\boldsymbol{\theta}})] \mathbf{H}_p^{+'}. \quad (14.16)$$

The least squares estimator $\hat{\boldsymbol{\theta}}_{\text{LS}}$ is the value of $\boldsymbol{\theta}$ that minimizes the weighted sum of squares

$$q(\boldsymbol{\theta}; \mathbf{S}) = [\text{vech } \mathbf{S} - \text{vech } \boldsymbol{\Sigma}(\boldsymbol{\theta})]' \hat{\mathbf{V}}^{-1} [\text{vech } \mathbf{S} - \text{vech } \boldsymbol{\Sigma}(\boldsymbol{\theta})] \quad (14.17)$$

over the parameter space Θ , with $\hat{\mathbf{V}}^{-1}$ defined using (14.14) and (14.15). Evaluating (14.17) at the value $\hat{\boldsymbol{\theta}}_{\text{LS}}$ yields the goodness-of-fit statistic $q(\hat{\boldsymbol{\theta}}; \mathbf{S})$. Regardless of the actual distribution of the data, $q(\hat{\boldsymbol{\theta}}; \mathbf{S})$ has an asymptotic distribution of $\chi^2_{p(p+1)/2-q}$ and the factor loading estimates $\hat{\boldsymbol{\lambda}}_{\text{LS}}$ converge to $\boldsymbol{\lambda}$ as defined in (14.6).

Use of $\mathbf{V}(\hat{\boldsymbol{\theta}})$ as an estimate of $\text{cov}(\text{vech } \mathbf{S})$ implies an iteratively reweighted least squares procedure beginning with an initial estimate of $\boldsymbol{\theta}$ such as $\hat{\boldsymbol{\theta}}^{(0)} = \hat{\boldsymbol{\theta}}_{\text{LS}}$. Then, in the i th iteration of the procedure ($i = 1, 2, \dots$), $\hat{\boldsymbol{\theta}}^{(i)}$ is the value of $\boldsymbol{\theta}$ that minimizes

$$w(\boldsymbol{\theta}; \hat{\boldsymbol{\theta}}^{(i-1)}, \mathbf{S}) = [\text{vech } \mathbf{S} - \text{vech } \boldsymbol{\Sigma}(\boldsymbol{\theta})]' \left[\mathbf{V}(\hat{\boldsymbol{\theta}}^{(i-1)}) \right]^{-1} [\text{vech } \mathbf{S} - \text{vech } \boldsymbol{\Sigma}(\boldsymbol{\theta})], \quad (14.18)$$

with $\mathbf{V}(\hat{\boldsymbol{\theta}}^{(i-1)})$ defined using (14.14) and (14.16). The procedure is continued until $\hat{\boldsymbol{\theta}}^{(i)}$ has converged, with the final estimate denoted $\hat{\boldsymbol{\theta}}_{\text{IRLS}}$. Each of the estimates $\hat{\boldsymbol{\theta}}_{\text{LS}}$ and $\hat{\boldsymbol{\theta}}_{\text{IRLS}}$ has the same asymptotic distribution as $\hat{\boldsymbol{\theta}}_{\text{ML}}$, although the procedure used to obtain $\hat{\boldsymbol{\theta}}_{\text{IRLS}}$ may require a simple modification in rare instances to achieve the desired result (see Fuller, 1987, Section 4.2.2).

14.3.3 Model Assessment

As noted in Sections 14.3.1 and 14.3.2, each of the statistics $\ell(\hat{\boldsymbol{\theta}}_{\text{ML}}; \mathbf{S})$, $q(\hat{\boldsymbol{\theta}}_{\text{LS}}; \mathbf{S})$, and $w(\hat{\boldsymbol{\theta}}_{\text{IRLS}}; \hat{\boldsymbol{\theta}}_{\text{IRLS}}, \mathbf{S})$ [described in (14.13), (14.17), and (14.18), respectively] has an asymptotic $\chi^2_{p(p+1)/2-q}$ distribution. Thus, each can be used as a goodness-of-fit statistic where the degrees of freedom for the asymptotic χ^2 distribution is the difference between the number of unique statistics in \mathbf{S} [$p(p+1)/2$] and the number of parameters in $\boldsymbol{\theta}$ (q). The hypotheses being compared are

H_0 : the hypothesized model is correct.

H_1 : the hypothesized model is incorrect (more factors are needed).

For example, if maximum likelihood is used to estimate the model parameters, then a value of $\ell(\hat{\boldsymbol{\theta}}_{\text{ML}}; \mathbf{S})$ that exceeds the $100(1 - \alpha)$ percentile of the $\chi^2_{p(p+1)/2-q}$ distribution would indicate that we reject H_0 at level α and conclude that the hypothesized model is incorrect. Obtaining a statistic whose value is less than said percentile would indicate the opposite conclusion—that is, we would fail to reject H_0 and proceed with statistical inference for the model parameters under the presumption that the model is an adequate characterization of the phenomenology associated with the data. We note that the degrees of freedom for the χ^2 test will usually be increased by 1 for every parameter that is constrained at the solution. For example, if we constrain an error variance to be nonnegative and that constraint is imposed on the solution (i.e., the error variance is set to 0), then q is decreased by 1 and the degrees of freedom is increased by 1.

One of the commonly cited limitations of the χ^2 goodness-of-fit test is that the probability of rejecting H_0 (i.e., rejecting the proposed model) *increases* as the sample size gets larger. Additionally, the test may reject for slight deviations from

normality such as the presence of kurtosis. Consequently, there are many other goodness-of-fit criteria that have been proposed and evaluated (see Bollen, 1989, Chapter 7; Hu and Bentler, 1999). In this section, we discuss three additional goodness-of-fit assessment tools that have become widely used and recommended in the literature: the comparative fit index (CFI) of Bentler (1989), the root mean square error approximation (RMSEA) index of Steiger and Lind (1980), and the standardized root mean square residual (SRMR) of Jöreskog and Sörbom (1985). None of these supplementary goodness-of-fit statistics is associated with a formal statistical test [as are the χ^2 statistics in (14.13), (14.17), and (14.18)], but each can be helpful for assessing model fit when used with an associated rule of thumb.

Bentler's CFI is a measure of the scaled reduction of lack of fit when using the specified model instead of a baseline model such as the independence model:

$$\mathbf{y} = \boldsymbol{\mu} + \boldsymbol{\varepsilon}. \quad (14.19)$$

For a specified model, let χ_M^2 be one of the goodness-of-fit statistics described in (14.13), (14.17), or (14.18), with the degrees of freedom associated with the statistic denoted df_M . Further, let χ_B^2 and df_B be the corresponding statistic and degrees of freedom when fitting a baseline model such as (14.19). Then, CFI is calculated using

$$CFI = 1 - \frac{\max(\chi_M^2 - df_M, 0)}{\max(\chi_B^2 - df_B, \chi_M^2 - df_M, 0)}. \quad (14.20)$$

Because the expected value of a χ^2 random variable is equal to the degrees of freedom for the variable's distribution, the quantity $\chi_M^2 - df_M$ represents excess lack of fit for the hypothesized model. The CFI statistic has range $0 \leq CFI \leq 1$, with values near 1 indicating that the excess lack of fit associated with the hypothesized model is much smaller than that associated with the baseline model. Values of CFI near zero imply that the increased complexity of the proposed model (relative to the baseline model) yields no additional explanatory ability. Hu and Bentler (1999) recommend that CFI values greater than 0.95 indicate a good fit, although other authors argue that values greater than 0.90 are acceptable.

The RMSEA index is another measure of excess lack of fit associated with the hypothesized model. It is calculated using

$$RMSEA = \sqrt{\frac{1}{n-1} \frac{\max(\chi_M^2 - df_M, 0)}{df_M}}, \quad (14.21)$$

where n is the sample size and the other quantities are as described in (14.20). Hu and Bentler (1999) recommend that an RMSEA less than 0.06 indicates a good fit.

The final goodness-of-fit measure considered here, SRMR, is not a function of the χ^2 statistic, but rather quantifies the difference between the sample covariance matrix for the data (\mathbf{S}) and the model-constructed estimates of the covariance matrix [$\hat{\Sigma}(\hat{\theta})$]. The SRMR value is calculated using

$$SRMR = \sqrt{\frac{2}{p(p+1)} \sum_{i=1}^p \sum_{j=1}^i \frac{(s_{ij} - \sigma_{ij})^2}{s_{ii}s_{jj}}}, \quad (14.22)$$

where s_{ij} and σ_{ij} are the (i, j) elements of S and $\Sigma(\hat{\theta})$, respectively. Hu and Bentler (1999) recommend that SRMR values less than 0.08 indicate a good fit, although other authors argue that values less than 0.10 imply an adequate fit of the model.

■ EXAMPLE 14.3.3

We illustrate model fitting and model assessment using the business statistics class data given in Table 14.1 and the CFA model given in (14.12). In specifying the model, we note that the factor variances and error variances must be constrained to be nonnegative. Using maximum likelihood (“method=ML” in SAS), the parameter estimates in $\hat{\theta}_{ML}$ are:

$$\begin{pmatrix} \hat{\lambda}_{11} \\ \hat{\lambda}_{31} \\ \hat{\lambda}_{32} \\ \hat{\lambda}_{42} \\ \hat{\lambda}_{52} \end{pmatrix} = \begin{pmatrix} 0.356 \\ 0.123 \\ 0.061 \\ 0.884 \\ 1.416 \end{pmatrix}, \quad \begin{pmatrix} \hat{\phi}_{11} \\ \hat{\phi}_{21} \\ \hat{\phi}_{22} \end{pmatrix} = \begin{pmatrix} 3.162 \\ 10.267 \\ 123.947 \end{pmatrix},$$

and

$$\begin{pmatrix} \hat{\psi}_{11} \\ \hat{\psi}_{22} \\ \hat{\psi}_{33} \\ \hat{\psi}_{44} \\ \hat{\psi}_{55} \\ \hat{\psi}_{66} \end{pmatrix} = \begin{pmatrix} 0.547 \\ 0 \\ 1.667 \\ 67.006 \\ 139.496 \\ 81.423 \end{pmatrix}.$$

Note that $\hat{\psi}_{22}$ is a parameter estimate that lies at a boundary of a specified parameter space. Consequently, statistical inference procedures typically treat this scenario as if ψ_{22} were constrained to equal 0 a priori, with the number of free parameters reduced from 14 to 13 and the degrees of freedom increased from 7 to 8. We focus on $\hat{\theta}_{ML}$ in this example, but the least squares estimates $\hat{\theta}_{LS}$ found by minimizing (14.17) (using “method=GLS” in SAS) are very similar.

To assess model fit, we consider the diagnostics discussed in Section 14.3.3. The χ^2 statistic in (14.13) is equal to 12.371 with a degrees of freedom of 8, based on 21 unique elements in S with 13 unconstrained parameters fit in the model. Thus, the p -value associated with the hypothesis

$$H_0 : \text{the hypothesized model is correct}$$

is 0.135, and we determine that we have insufficient evidence to declare lack of fit. The additional diagnostic metrics yield a mixed verdict about the goodness of fit for the proposed model. Bentler’s CFI in (14.20) is 0.978 (greater than the recommended lower cutoff of 0.95) and the SRMR value in (14.22) is 0.039 (less than the recommended upper cutoff of 0.08). On the other hand, the RMSEA index in (14.21) is 0.0767 (greater than the recommended upper

cutoff of 0.06). In practice, the quality of fit of a latent variable model is determined using an ensemble of fit indexes. On balance, it appears that model (14.12) has reasonably good fit. \square

14.4 INFERENCE FOR MODEL PARAMETERS

We previously noted that normality-based methods for parameter estimation are still of practical value when data are not normally distributed. However, when conducting inference about the q -dimensional parameter vector θ using $\hat{\theta}_{\text{ML}}$, $\hat{\theta}_{\text{LS}}$, or $\hat{\theta}_{\text{IRLS}}$, we need an estimate of the covariance matrix for the vector of parameter estimates that properly accounts for the potential nonnormality of the data. Because θ is based on the second moments of the data found in the covariance matrix, inference about θ requires a fourth-moment matrix. Thus, we first define an estimated covariance matrix for $\text{vech } \mathbf{S}$.

Let \mathbf{z}_i be the vector of length $p(p+1)/2$ that is associated with the i th observation and defined by

$$\mathbf{z}_i = \text{vech} (\mathbf{y}_i - \bar{\mathbf{y}})(\mathbf{y}_i - \bar{\mathbf{y}})',$$

so that

$$\text{vech } \mathbf{S} = \frac{1}{n-1} \sum_{i=1}^n \mathbf{z}_i.$$

We can then define the covariance matrix for $\text{vech } \mathbf{S}$ as

$$\Gamma = \text{cov}(\text{vech } \mathbf{S}),$$

which can be estimated using

$$\hat{\Gamma} = \frac{1}{(n-1)^2} \sum_{i=1}^n (\mathbf{z}_i - \bar{\mathbf{z}})(\mathbf{z}_i - \bar{\mathbf{z}})', \quad (14.23)$$

where $\bar{\mathbf{z}} = \frac{1}{n} \sum_{i=1}^n \mathbf{z}_i$. When $\hat{\theta}$ is $\hat{\theta}_{\text{ML}}$, $\hat{\theta}_{\text{LS}}$, or $\hat{\theta}_{\text{IRLS}}$, the covariance matrix for the estimated model parameters is

$$\widehat{\text{cov}}(\hat{\theta}) = \{\hat{\mathbf{F}}'[\mathbf{V}(\hat{\theta})]^{-1}\hat{\mathbf{F}}\}^{-1} \hat{\Gamma} [\mathbf{V}(\hat{\theta})]^{-1} \hat{\mathbf{F}} \{\hat{\mathbf{F}}'[\mathbf{V}(\hat{\theta})]^{-1}\hat{\mathbf{F}}\}^{-1}, \quad (14.24)$$

where $\mathbf{V}(\hat{\theta})$ is defined in (14.16), \mathbf{F} is the $p(p+1)/2 \times q$ matrix of partial derivatives defined by

$$\mathbf{F} = \frac{\partial \text{vech } \Sigma(\theta)}{\partial \theta'},$$

and $\hat{\mathbf{F}}$ is the matrix \mathbf{F} when evaluated at $\theta = \hat{\theta}$.

As noted previously, Anderson and Amemiya (1988) and Amemiya and Anderson (1990) show that given general identification conditions and a large sample size, a wide array of statistical inferences about θ can be carried out, regardless of the

actual distribution of \mathbf{y} . Specifically, the parameter estimates $\hat{\boldsymbol{\theta}}_{\text{ML}}$, $\hat{\boldsymbol{\theta}}_{\text{LS}}$, and $\hat{\boldsymbol{\theta}}_{\text{IRLS}}$ are asymptotically unbiased, and the subset of $\widehat{\text{cov}}(\hat{\boldsymbol{\theta}})$ corresponding to $\widehat{\text{cov}}(\hat{\boldsymbol{\lambda}})$ (the variances and covariances among the factor loadings) are also unbiased.

The square roots of the diagonal elements of (14.24) corresponding to the estimated factor loadings ($\hat{\lambda}_{ij}$) are their associated standard errors [s.e.($\hat{\lambda}_{ij}$)]. Thus, the standard error s.e.($\hat{\lambda}_{ij}$) can be used to construct the asymptotically normal test statistic

$$z_{ij} = \frac{\hat{\lambda}_{ij}}{\text{s.e.}(\hat{\lambda}_{ij})}, \quad (14.25)$$

which is suitable for testing the hypothesis $H_0 : \lambda_{ij} = 0$. The z_{ij} statistics associated with the various factor loadings can then be used to evaluate the relative statistical evidence for relationships among observed and latent variables. Using similar logic, a $100(1 - \alpha)\%$ confidence interval for λ_{ij} can be constructed using

$$\hat{\lambda}_{ij} \pm z_{\alpha/2} \text{s.e.}(\hat{\lambda}_{ij}). \quad (14.26)$$

Because $\hat{\boldsymbol{\theta}}$ and $\widehat{\text{cov}}(\hat{\boldsymbol{\theta}})$ are valid in large samples regardless of the distribution of the data, the associated hypothesis tests and confidence intervals for $\boldsymbol{\theta}$ are valuable tools for evaluating and refining the proposed factor analysis model. A typical process for model refinement first uses the model assessment tools discussed in Section 14.3.3, then uses parameter assessment tools based on $\hat{\boldsymbol{\theta}}$ and $\widehat{\text{cov}}(\hat{\boldsymbol{\theta}})$ to fine-tune the model. For example, if a model exhibits adequate goodness of fit, we might then determine which if any of the model parameter estimates are nonsignificant, dropping any that are deemed extraneous and then refitting the model to ensure that the goodness of overall model fit has not been substantially reduced. As with any exploratory model-building exercise, there is a delicate balance between parsimony and goodness of fit. Further, such iterative model-selection procedures are subject to the overly optimistic assessments of model fit that are associated with the subset selection problem in regression analysis (see discussion in Section 10.2.7b). Thus, analyses that require several iterations of model refinement should be interpreted as exploratory in nature and used with caution.

■ EXAMPLE 14.4

In this example, we continue the analysis of the business statistics class data given in Table 14.1 with the intent of evaluating and refining the CFA model given in (14.12) and fitted in Example 14.3.3. Table 14.2 gives the test statistic z_{ij} and other pertinent inferential quantities for each of the factor loadings in the model. Note that all but λ_{31} appear to be significantly different from 0. In an effort to obtain a more parsimonious model with good fit, we consider the

Table 14.2 Factor Loading Estimates for Statistics Course Grade Data of Table 14.1 When Using Model (14.12)

Parameter	Estimate	Standard Error	z_{ij}	p-value
λ_{11}	0.356	0.043	8.254	< 0.0001
λ_{31}	0.123	0.094	1.307	0.0957
λ_{32}	0.061	0.017	3.487	0.0002
λ_{42}	0.884	0.127	6.985	< 0.0001
λ_{52}	1.416	0.197	7.185	< 0.0001

Table 14.3 Factor Loading Estimates for Statistics Course Grade Data of Table 14.1 When Using Model (14.27)

Parameter	Estimate	Standard Error	z_{ij}	p-value
λ_{11}	0.376	0.088	4.274	< 0.0001
λ_{32}	0.073	0.015	4.814	< 0.0001
λ_{42}	0.882	0.128	6.916	< 0.0001
λ_{52}	1.435	0.199	7.211	< 0.0001

revised model for the data:

$$\begin{aligned}
 \text{Lab} &= \mu_1 + \lambda_{11}f_1 &+ \varepsilon_1 \\
 \text{HW} &= \mu_2 + f_1 &+ \varepsilon_2 \\
 \text{PopQuiz} &= \mu_3 &+ \lambda_{32}f_2 + \varepsilon_3 \\
 \text{Exam1} &= \mu_4 &+ \lambda_{42}f_2 + \varepsilon_4 \\
 \text{Exam2} &= \mu_5 &+ \lambda_{52}f_2 + \varepsilon_5 \\
 \text{FinalExam} &= \mu_6 &+ f_2 + \varepsilon_6
 \end{aligned} \tag{14.27}$$

The goodness-of-fit statistics associated with the revised model given in (14.27) are: $\chi^2 = 13.945$ (degrees of freedom = 8 and p-value = 0.083), Bentler's CFI = 0.970, RMSEA = 0.0894, and SRMR = 0.039. Thus, all measures indicate an adequate fit of the model to the data except for RMSEA, which exceeds the recommended upper cutoff of 0.06. We determine that the use of the revised model in (14.27) is an improvement over the use of (14.12). Table 14.3 gives the factor loading estimates and associated inferential statistics for the revised model, which now involves only factor loadings that are statistically significant. The revised model indicates that Lab and HW load solely on the effort factor and PopQuiz, Exam1, Exam2, and FinalExam load solely on the knowledge mastery factor. \square

14.5 FACTOR SCORES

As discussed in the EFA setting in Section 13.6, it is often the case that we wish to obtain *factor scores*—estimates of the underlying factor values for each observation. In this section, we introduce two common approaches for obtaining such estimates. Factor scores are often used after fitting the factor analysis model in order to understand the underlying characteristics of the observational units.

Using the parameter estimates $\hat{\Lambda}$, $\hat{\Phi}$, and $\hat{\Psi}$, we can form factor score estimates for the i th observational unit using the so-called *regression method* formula

$$\hat{f}_i = \hat{\Phi} \hat{\Lambda}' (\hat{\Lambda} \hat{\Phi} \hat{\Lambda}' + \hat{\Psi})^{-1} (\mathbf{y}_i - \bar{\mathbf{y}}) \quad (14.28)$$

or the *weighted least squares* formula

$$\tilde{f}_i = (\hat{\Lambda}' \hat{\Psi}^{-1} \hat{\Lambda})^{-1} \hat{\Lambda}' \hat{\Psi}^{-1} (\mathbf{y}_i - \bar{\mathbf{y}}). \quad (14.29)$$

Although the two approaches generally yield very similar estimates, the regression method formula in (14.28) is perhaps the most widely used (see Bollen 1989, Chapter 7).

■ EXAMPLE 14.5

In this example, we use the regression method in (14.28) to obtain factor scores for each student in the statistics course grade data set. The data are found in Table 14.1, and the parameter estimates used are based on the revised model (14.27) fit to the data in Example 14.4. The matrices used to calculate the factor scores are:

$$\hat{\Lambda} = \begin{pmatrix} 0.376 & 0 \\ 1 & 0 \\ 0 & 0.073 \\ 0 & 0.882 \\ 0 & 1.435 \\ 0 & 1 \end{pmatrix}, \quad \hat{\Phi} = \begin{pmatrix} 2.992 & 10.542 \\ 10.542 & 122.032 \end{pmatrix},$$

$$\text{and } \hat{\Psi} = \begin{pmatrix} 0.524 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0.170 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1.664 & 0 & 0 & 0 \\ 0 & 0 & 0 & 68.982 & 0 & 0 \\ 0 & 0 & 0 & 0 & 136.699 & 0 \\ 0 & 0 & 0 & 0 & 0 & 83.338 \end{pmatrix}.$$

Values for \hat{f}_1 (estimated effort score) and \hat{f}_2 (estimated knowledge mastery score) are calculated and plotted against each other in Figure 14.3. Although we can say nothing about causal influences between the factors, it does appear that the effort score can be used to provide a rough upper bound for the knowledge mastery score. \square




Figure 14.3 Estimated effort score (\hat{f}_1) vs. estimated knowledge mastery score (\hat{f}_2) for 94 students in a business statistics course.

PROBLEMS

- 14.1** Show that the order condition given in (14.8) simplifies to $(p-m)^2 \geq p+m$.
- 14.2** Consider the confirmatory factor analysis model in (14.2). In the scenarios below, various constrained Λ matrices of dimension $p \times m$ are proposed. For each proposed model, use the rules presented in Section 14.2.2 to determine whether or not the model is identified. Assume that for each model, $\Phi = \text{cov}(\mathbf{f})$ is an $m \times m$ symmetric matrix containing $m(m+1)/2$ unique factor variances and covariances to be estimated. Further, assume that $\Psi = \text{cov}(\boldsymbol{\varepsilon})$ is a diagonal matrix with p unique error variances to be estimated. For each assessment of model identifiability, explain your reasoning.

$$(a) \quad \Lambda = \begin{bmatrix} \lambda_{11} & \lambda_{12} \\ \lambda_{21} & \lambda_{22} \\ \lambda_{31} & \lambda_{32} \\ 0.5 & 0 \\ 0 & 1 \end{bmatrix}$$

$$(b) \Lambda = \begin{bmatrix} \lambda_{11} & 0 & 0 \\ 0 & \lambda_{22} & \lambda_{23} \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$(c) \Lambda = \begin{bmatrix} \lambda_{11} & \lambda_{12} & \lambda_{13} \\ \lambda_{21} & \lambda_{22} & \lambda_{23} \\ \lambda_{31} & \lambda_{32} & \lambda_{33} \\ \lambda_{41} & \lambda_{42} & \lambda_{43} \\ 1 & 0 & 0 \\ 0 & 1 & 0.5 \\ 0 & 1 & 0.5 \end{bmatrix}$$

$$(d) \Lambda = \begin{bmatrix} \lambda_{11} & 0 \\ \lambda_{21} & \lambda_{22} \\ 1 & 0 \\ 0 & 1 \end{bmatrix}$$

14.3 Consider the 1-factor confirmatory factor analysis model in (14.2) with

$$\Lambda = \begin{bmatrix} 1 \\ \lambda_2 \\ \lambda_3 \end{bmatrix}, \text{ var}(f) = \phi, \text{ and } \text{cov}(\varepsilon) = \begin{bmatrix} \psi_{11} & 0 & 0 \\ 0 & \psi_{22} & 0 \\ 0 & 0 & \psi_{33} \end{bmatrix}.$$

- (a) Use the rules presented in Section 14.2.2 to determine whether or not the model is identified. Explain your reasoning.
- (b) How many degrees of freedom will be associated with the χ^2 goodness-of-fit statistic described in (14.13)? What does this imply about the nature of the estimate $\hat{\theta}$ —that is, how close will $\Sigma(\hat{\theta})$ be to the sample covariance matrix for the data \mathbf{S} ?
- (c) Find the model covariance matrix $\Sigma(\theta)$, with each element written as a function of λ_2 , λ_3 , ϕ , ψ_{11} , ψ_{22} , and ψ_{33} .
- (d) Let the sample covariance matrix for the data be denoted

$$\mathbf{S} = \begin{bmatrix} s_{11} & s_{12} & s_{13} \\ s_{21} & s_{22} & s_{23} \\ s_{31} & s_{32} & s_{33} \end{bmatrix}.$$

Set the six unique elements of \mathbf{S} equal to the corresponding elements of $\Sigma(\theta)$. With six equations and six parameters to estimate, find estimates of λ_2 , λ_3 , ϕ , ψ_{11} , ψ_{22} , and ψ_{33} as functions of sample variances and covariances (s_{11} , s_{12} , s_{13} , s_{22} , s_{23} , and s_{33}).

14.4 Use the probe word data of Table 3.6. Conduct a confirmatory factor analysis of the covariance matrix for the five variables. Use maximum likelihood to determine if the five word probes can be adequately modeled using a single latent factor. In formulating the model, identifiability can be ensured by letting $y_1 = f_1 + \varepsilon_1$ and letting $y_i = \lambda_i f_1 + \varepsilon_i$ ($i = 2, 3, 4, 5$).

- (a) Assess goodness of fit with the criteria discussed in Section 14.3.3.
- (b) Which variables are most significantly related to the latent factor? Which are least significant? Can the model be further simplified?
- (c) Fit the model again, this time identifying the model by setting one of the other observed variables equal to $f_1 + \varepsilon_i$. For example, set $y_2 = f_1 + \varepsilon_2$, with $y_i = \lambda_i f_1 + \varepsilon_i$ for $i = 1, 3, 4, 5$. Calculate the goodness of fit statistics discussed in Section 14.3.3 and compare with part (a). Is the goodness of fit for the one-factor model affected by which observed variable is set equal to $f_1 + \varepsilon_i$?

14.5 Use the Seishu data of Table 7.1. Conduct a confirmatory factor analysis of the covariance matrix for all 10 variables. Use maximum likelihood to fit the model and test the hypothesis that the observations are driven by four factors:

$$\begin{aligned}f_1 &= \text{"flavor"} \\f_2 &= \text{"acidity"} \\f_3 &= \text{"sweetness"} \\f_4 &= \text{"alcohol content"}\end{aligned}$$

To fit an identifiable model, define the observed variable “Taste” to be equal to f_1 plus error, define the observed variable “pH” to be equal to f_2 plus error, define the observed variable “Total Sugar” to be equal to f_3 plus error, and define the observed variable “Alcohol” to be equal to f_4 plus error. In your initial model, allow the other 6 variables to be functions of all 4 factors, for a total of 24 factor loadings to be estimated.

- (a) Assess goodness of fit with the criteria discussed in Section 14.3.3.
 - (b) Using goodness-of-fit statistics and hypothesis tests on each of the factor loadings, conduct an exploratory analysis using the iterative model-selection process discussed at the end of Section 14.5 to identify a simpler model. Specifically, determine whether a model can be specified with fewer factor loadings (i.e., simpler structure) and still exhibit a good fit. Propose a simpler model and appropriately justify your proposal using goodness-of-fit statistics and hypothesis tests.
- 14.6** Use the football data of Table 8.3, combining the three groups into a single sample. Conduct a confirmatory factor analysis of the covariance matrix using maximum likelihood to fit the model. Test the hypothesis that the observations are driven by two factors related to head size:

$$\begin{aligned}f_1 &= \text{"horizontal dimension"} \\f_2 &= \text{"vertical dimension"}\end{aligned}$$

To fit an identifiable model, define the observed variable “head circumference” to be equal to f_1 plus error, and define the observed variable “eye-to-top-of-head measurement” to be equal to f_2 plus error. In your initial model, allow the other 4 variables to be functions of both factors, for a total of 8 factor loadings to be estimated.

- (a) Assess goodness of fit with the criteria discussed in Section 14.3.3.
 - (b) For comparison, fit the 2-factor model with simple structure. That is, fit the model with head width, head circumference, front-to-back measurement at eye level, and jaw width loading only on f_1 . Similarly, let eye-to-top-of-head measurement and ear-to-top-of-head measurement load only on f_2 . Use goodness-of-fit criteria and hypothesis tests on factor loadings to compare the initial model with this simple-structure model. Which model is preferable?
- 14.7** Use the engineer data of Table 5.6, combining the two groups into a single sample. Conduct a confirmatory factor analysis of the covariance matrix using maximum likelihood. Test the hypothesis that the observations are driven by two factors related to test performance:
- $$f_1 = \text{“aptitude”}$$
- $$f_2 = \text{“effort”}$$
- To fit an identifiable model, define the observed variable “intelligence” to be equal to f_1 plus error, and define the observed variable “perseveration” to be equal to f_2 plus error. Allow the other 4 variables to be functions of both factors, for a total of 8 factor loadings to be estimated.
- (a) Assess goodness of fit with the criteria discussed in Section 14.3.3.
 - (b) For each observed y_i (performance test), quantify how much of the variability in y_i can be accounted for by the factors. This can be calculated using
- $$1 - \frac{\hat{\psi}_{ii}}{s_{ii}}.$$
- Note that if the estimated error variance ($\hat{\psi}_{ii}$) is greater than the sample variance for the associated observed variable (s_{ii}), the percent of variability explained by the factors is undefined and the quality of the model fit is compromised.
- (c) Interpret the results from parts (a) and (b). Does goodness of fit imply that the latent factors can be used as a lower-dimensional replacement for the observed variables? Would you expect that the variance explained by the factors for each y_i would improve substantially with the addition of another factor in the model?
- 14.8** Use the rootstock data of Table 6.2. Using the covariance matrix, fit a confirmatory factor analysis model using maximum likelihood to determine if the four tree measurements can be adequately modeled using a single latent factor.

To ensure model identifiability, let one of the four variables be modeled with $y_i = f_1 + \varepsilon_i$ and let $y_j = \lambda_j f_1 + \varepsilon_j$ for $j \neq i$. Assess goodness of fit with the criteria discussed in Section 14.3.3. If the model exhibits adequate fit, explain whether or not the model can be further simplified. If the model does not exhibit adequate fit, explain why you would or would not recommend fitting a 2-factor or 3-factor model.

CHAPTER 15

CLUSTER ANALYSIS

15.1 INTRODUCTION

In *cluster analysis* we search for patterns in a data set by grouping the (multivariate) observations into clusters. The goal is to find an optimal grouping for which the observations or objects within each cluster are similar but the clusters are dissimilar to each other. We hope to find the natural groupings in the data, groupings that make sense to the researcher.

Cluster analysis differs fundamentally from classification analysis (Chapter 9). In classification analysis, we allocate the observations to a known number of predefined groups or populations. In cluster analysis, neither the number of groups nor the groups themselves are known in advance.

To group the observations into clusters, many techniques begin with similarities between all pairs of observations. In many cases the similarities are based on some measure of distance. Other cluster methods use a preliminary choice for cluster centers or a comparison of within- and between-cluster variability. It is also possible to cluster the variables, in which case the similarity could be a correlation; see Section 15.7.

We can search for clusters graphically by plotting the observations. If there are only two variables ($p = 2$), this can be done in a scatterplot (see Section 3.3). For $p > 2$, we can plot the data in two dimensions using principal components (see Section 12.4) or biplots (see Section 16.3). For an example of a principal component plot, see Figure 12.7 in Section 12.4, in which four clear groupings of points can be observed. Another approach to plotting is provided by *projection pursuit*, which seeks two-dimensional projections that reveal clusters [see Friedman and Tukey (1974); Huber (1985); Sibson (1984); Jones and Sibson (1987); Yenukov (1988); Posse (1990); Nason (1995); Ripley (1996, pp. 296–303)].

Cluster analysis has also been referred to as classification, pattern recognition (specifically, unsupervised learning), and numerical taxonomy. The techniques of cluster analysis have been extensively applied to data in many fields, such as medicine, psychiatry, sociology, criminology, anthropology, archaeology, geology, geography, remote sensing, market research, economics, and engineering.

We shall concentrate largely on quantitative variables [for categorical variables, see Gordon (1999) or Everitt (1993)]. The data matrix [see (3.17)] can be written as

$$\mathbf{Y} = \begin{pmatrix} \mathbf{y}'_1 \\ \mathbf{y}'_2 \\ \vdots \\ \mathbf{y}'_n \end{pmatrix} = (\mathbf{y}_{(1)}, \mathbf{y}_{(2)}, \dots, \mathbf{y}_{(p)}), \quad (15.1)$$

where \mathbf{y}'_i is a row (observation vector) and $\mathbf{y}_{(j)}$ is a column (corresponding to a variable). We generally wish to group the n \mathbf{y}'_i 's (rows) into g clusters. We may also wish to cluster the columns $\mathbf{y}_{(j)}$, $j = 1, 2, \dots, p$ (see Section 15.7).

Two common approaches to clustering the observation vectors are hierarchical clustering and partitioning. In *hierarchical* clustering we typically start with n clusters, one for each observation, and end with a single cluster containing all n observations. At each step, an observation or a cluster of observations is absorbed into another cluster. We can also reverse this process, that is, start with a single cluster containing all n observations and end with n clusters of a single item each (see Section 15.3.10). In *partitioning*, we simply divide the observations into g clusters. This can be done by starting with an initial partitioning or with cluster centers and then reallocating the observations according to some optimality criterion. Other clustering methods that we will discuss are based on fitting mixtures of multivariate normal distributions or searching for regions of high density sometimes called modes.

There is an abundant literature on cluster analysis. Useful monographs and reviews have been given by Gordon (1999), Everitt (1993), Khattree and Naik (2000, Chapter 6), Kaufman and Rousseeuw (1990), Seber (1984, Chapter 7), Anderberg (1973), and Hartigan (1975a).

15.2 MEASURES OF SIMILARITY OR DISSIMILARITY

Since cluster analysis attempts to identify the observation vectors that are similar and group them into clusters, many techniques use an index of *similarity* or *proximity*

between each pair of observations. A convenient measure of proximity is the distance between two observations. Since distance increases as two units become further apart, distance is actually a measure of *dissimilarity*.

A common distance function is the Euclidean distance between two vectors $\mathbf{x} = (x_1, x_2, \dots, x_p)'$ and $\mathbf{y} = (y_1, y_2, \dots, y_p)'$, defined as

$$d(\mathbf{x}, \mathbf{y}) = \sqrt{(\mathbf{x} - \mathbf{y})'(\mathbf{x} - \mathbf{y})} = \sqrt{\sum_{j=1}^p (x_j - y_j)^2}. \quad (15.2)$$

To adjust for differing variances and covariances among the p variables, we could use the statistical distance

$$d(\mathbf{x}, \mathbf{y}) = \sqrt{(\mathbf{x} - \mathbf{y})' \mathbf{S}^{-1} (\mathbf{x} - \mathbf{y})} \quad (15.3)$$

[see (3.79)], where \mathbf{S} is the sample covariance matrix. After the clusters are formed, \mathbf{S} could be computed as the pooled within-cluster covariance matrix, but we do not know beforehand what the clusters will be. If we compute \mathbf{S} on the unpartitioned sample, there will be distortion of the variances and covariances because of the groups in the data (assuming there really are some natural clusters). We therefore usually use the Euclidean distance given by (15.2). In some clustering procedures, it is not necessary to take the square root in (15.2) or (15.3).

Other distance measures have been suggested, for example, the Minkowski metric

$$d(\mathbf{x}, \mathbf{y}) = \left[\sum_{j=1}^p |x_j - y_j|^r \right]^{1/r}. \quad (15.4)$$

For $r = 2$, $d(\mathbf{x}, \mathbf{y})$ in (15.4) becomes the Euclidean distance given in (15.2). For $p = 2$ and $r = 1$, (15.4) measures the “city block” distance between two observations. There are distance measures for categorical data; see Gordon (1999, Chapter 2).

For the n observation vectors $\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_n$, we can compute an $n \times n$ matrix $\mathbf{D} = (d_{ij})$ of distances (or dissimilarities) where $d_{ij} = d(\mathbf{y}_i, \mathbf{y}_j)$ is usually given by (15.2), $d(\mathbf{y}_i, \mathbf{y}_j) = \sqrt{(\mathbf{y}_i - \mathbf{y}_j)'(\mathbf{y}_i - \mathbf{y}_j)}$. We sometimes use $\mathbf{D} = (d_{ij}^2)$, where $d_{ij}^2 = d^2(\mathbf{y}_i, \mathbf{y}_j) = (\mathbf{y}_i - \mathbf{y}_j)'(\mathbf{y}_i - \mathbf{y}_j)$ is the square of (15.2). The matrix \mathbf{D} will typically be symmetric with diagonal elements equal to zero.

The scale of measurement of the variables is an important consideration when using the Euclidean distance measure in (15.2). Changing the scale can affect the relative distances among the items. For example, suppose three items have the following bivariate measurements (y_1, y_2) : (2, 5), (4, 2), (7, 9). Using d_{ij} as given by (15.2), the matrix $\mathbf{D} = (d_{ij})$ for these items is

$$\mathbf{D}_1 = \begin{pmatrix} 0 & 3.6 & 6.4 \\ 3.6 & 0 & 7.6 \\ 6.4 & 7.6 & 0 \end{pmatrix}.$$

However, if we multiply y_1 by 100 as, for example, in changing from meters to centimeters, the matrix becomes

$$\mathbf{D}_2 = \begin{pmatrix} 0 & 200 & 500 \\ 200 & 0 & 300 \\ 500 & 300 & 0 \end{pmatrix},$$

and the largest distance is now d_{13} instead of d_{23} . The distance rankings have been altered by scaling.

To counter this problem, each variable could be standardized in the usual way by subtracting the mean and dividing by the standard deviation of the variable. However, such scaling would ordinarily be based on the entire data set, that is, on all n values in each column of \mathbf{Y} in (15.1). In this case, the variables that best separate clusters might no longer do so after division by standard deviations that include between-cluster variation. If we use standardized variables, the clusters could be less well separated. The question of scaling is therefore not an easy one. However, standardization of this type is recommended by many authors.

By (15.2), the squared Euclidean distance between two observations $\mathbf{x} = (x_1, x_2, \dots, x_p)'$ and $\mathbf{y} = (y_1, y_2, \dots, y_p)'$ is $d^2(\mathbf{x}, \mathbf{y}) = \sum_{j=1}^p (x_j - y_j)^2$. This can be expressed as

$$d^2(\mathbf{x}, \mathbf{y}) = (v_x - v_y)^2 + p(\bar{x} - \bar{y})^2 + 2v_x v_y (1 - r_{xy}), \quad (15.5)$$

where $v_x^2 = \sum_{j=1}^p (x_j - \bar{x})^2$ and $\bar{x} = \sum_{j=1}^p x_j / p$, with similar expressions for v_y^2 and \bar{y} . The correlation r_{xy} in (15.5) is given by

$$r_{xy} = \frac{\sum_{j=1}^p (x_j - \bar{x})(y_j - \bar{y})}{\sqrt{\sum_{j=1}^p (x_j - \bar{x})^2 \sum_{j=1}^p (y_j - \bar{y})^2}}. \quad (15.6)$$

In Figure 15.1, we illustrate the profile (see Sections 5.9 and 6.8) for each of two observation vectors \mathbf{x} and \mathbf{y} . The squared Euclidean distance in (15.5) can be used to compare the profiles of \mathbf{x} and \mathbf{y} in terms of levels, variation, and shape, where \bar{x} and \bar{y} are the *levels* of the two profiles, v_x and v_y are the *variations* of the profiles, and the correlation r_{xy} is a measure of the closeness of the *shapes* of the two profiles. The closer r_{xy} is to 1, the greater is the similarity in shape of the two profiles. Note that \bar{x} and v_x are the mean and variation of the p variables within the observation vector \mathbf{x} , not over the n observations in the data set. A similar comment can be made about \bar{y} and v_y . Likewise, the correlation r_{xy} is between the two observation vectors \mathbf{x} and \mathbf{y} , not between two variables. The use of r_{xy} has been questioned by Jardine and Sibson (1971) and Wishart (1971), but Strauss et al. (1973) found the correlation to be superior to the Euclidean distance for finding the clusters in a particular data set.




Figure 15.1 Profiles for two observation vectors \mathbf{x} and \mathbf{y} .

15.3 HIERARCHICAL CLUSTERING

15.3.1 Introduction

Hierarchical methods and other clustering algorithms represent an attempt to find “good” clusters in the data using a computationally efficient technique. It is not generally feasible to examine all possible clustering possibilities for a data set, especially a large one. The number of ways of partitioning a set of n items into g clusters is given by

$$N(n, g) = \frac{1}{g!} \sum_{k=1}^g \binom{g}{k} (-1)^{g-k} k^n \quad (15.7)$$

[see Duran and Odell (1974, Chapter 4), Jensen (1969), Seber (1984, p. 379)]. This can be approximated by $g^n/g!$, which is large even for moderate values of n and g . For example, $N(25, 10) \cong 2.8 \times 10^{18}$. The total possible number of clusters for a set of n items is $\sum_{g=1}^n N(n, g)$, which, for $n = 25$, is greater than 10^{19} . Hence, hierarchical methods and other approaches permit us to search for a reasonable solution without having to look at all possible arrangements.

As noted in Section 15.1, hierarchical clustering algorithms involve a sequential process. In each step of the *agglomerative* hierarchical approach, an observation or a cluster of observations is merged into another cluster. In this process, the number of clusters shrinks and the clusters themselves grow larger. We start with n clusters (individual items) and end with one single cluster containing the entire data set. An alternative approach, called the *divisive* method, starts with a single cluster

containing all n items and partitions a cluster into two clusters at each step (see Section 15.3.10). The end result of the divisive approach is n clusters of one item each. Agglomerative methods are more commonly used than divisive methods. In either type of hierarchical clustering, a decision must be made as to the “optimal” number of clusters (see Section 15.5).

At each step of an agglomerative hierarchical approach, the two “closest” clusters are merged into a single new cluster. The process is therefore irreversible in the sense that any two items that are once lumped together in a cluster cannot be separated later in the procedure; any early mistakes cannot be corrected. Similarly, in a divisive hierarchical method, items cannot be moved to other clusters. An optional approach is to carry out a hierarchical procedure followed by a partitioning procedure in which items can be moved from one cluster to another (see Section 15.4.1).

Since an agglomerative hierarchical procedure combines the two “closest” clusters at each step, we must consider the question of measuring the similarity or dissimilarity of two clusters. Different approaches to measuring distance between clusters give rise to different hierarchical methods. Agglomerative techniques are discussed in Sections 15.3.2–15.3.9, and the divisive approach is considered in Section 15.3.10.

15.3.2 Single Linkage (Nearest Neighbor)

In the *single linkage* method, the distance between two clusters A and B is defined as the *minimum* distance between a point in A and a point in B :

$$D(A, B) = \min \{d(\mathbf{y}_i, \mathbf{y}_j), \text{ for } \mathbf{y}_i \text{ in } A \text{ and } \mathbf{y}_j \text{ in } B\}, \quad (15.8)$$

where $d(\mathbf{y}_i, \mathbf{y}_j)$ is the Euclidean distance in (15.2) or some other distance between the vectors \mathbf{y}_i and \mathbf{y}_j . This approach is also called the *nearest neighbor* method.

At each step in the single linkage method, the distance (15.8) is found for every pair of clusters, and we merge the two clusters with smallest distance. The number of clusters is therefore reduced by one. After two clusters are merged, the procedure is repeated for the next step: the distances between all pairs of clusters are calculated again, and the pair with minimum distance is merged into a single cluster.

The results of a hierarchical clustering procedure can be displayed graphically using a *tree diagram*, also known as a *dendrogram*, which shows all the steps in the hierarchical procedure, including the distances at which clusters are merged. Dendograms are shown in Figures 15.2 and 15.3 in Examples 15.3.2(a) and 15.3.2(b).

■ EXAMPLE 15.3.2(a)

Hartigan (1975a, p. 28) compared the crime rates per 100,000 population for various cities. The data are in Table 15.1 (taken from the 1970 US Statistical Abstract). In order to illustrate the use of the distance matrix in single linkage clustering, we use the first six observations in Table 15.1 (Atlanta through Detroit).

Table 15.1 City Crime Rates per 100,000 Population

City	Murder	Rape	Robbery	Assault	Burglary	Larceny	AutoTheft
Atlanta	16.5	24.8	106	147	1112	905	494
Boston	4.2	13.3	122	90	982	669	954
Chicago	11.6	24.7	340	242	808	609	645
Dallas	18.1	34.2	184	293	1668	901	602
Denver	6.9	41.5	173	191	1534	1368	780
Detroit	13.0	35.7	477	220	1566	1183	788
Hartford	2.5	8.8	68	103	1017	724	468
Honolulu	3.6	12.7	42	28	1457	1102	637
Houston	16.8	26.6	289	186	1509	787	697
Kansas City	10.8	43.2	255	226	1494	955	765
Los Angeles	9.7	51.8	286	355	1902	1386	862
New Orleans	10.3	39.7	266	283	1056	1036	776
New York	9.4	19.4	522	267	1674	1392	848
Portland	5.0	23.0	157	144	1530	1281	488
Tucson	5.1	22.9	85	148	1206	756	483
Washington	12.5	27.6	524	217	1496	1003	793

The distance matrix \mathbf{D} is given by

CITY	DISTANCE BETWEEN CITIES					
	Atlanta	536.6	516.4	590.2	693.6	716.2
Boston	536.6	0	447.4	833.1	915.0	881.1
Chicago	516.4	447.4	0	924.0	1073.4	971.5
Dallas	590.2	833.1	924.0	0	527.7	464.5
Denver	693.6	915.0	1073.4	527.7	0	358.7
Detroit	716.2	881.1	971.5	464.5	358.7	0

The smallest distance is 358.7 between Denver and Detroit, and therefore these two cities are joined at the first step to form $C_1 = \{\text{Denver, Detroit}\}$. In the next step, the distance matrix is calculated for Atlanta, Boston, Chicago, Dallas, and C_1 :

Atlanta	0	536.6	516.4	590.2	693.6
Boston	536.6	0	447.4	833.1	881.1
Chicago	516.4	447.4	0	924.0	971.5
Dallas	590.2	833.1	924.0	0	464.5
C_1	693.6	881.1	971.5	464.5	0

Note that all elements of this distance matrix are contained in the original distance matrix. This same pattern will hold in subsequent distance matrices below and is also characteristic of the complete linkage method [see Example 15.3.3(a)]. The smallest distance is 447.4 between Boston and Chicago. Therefore $C_2 = \{\text{Boston, Chicago}\}$. At the next step, the distance matrix is calculated for Atlanta, Dallas, C_1 , and C_2 :

Atlanta	0	516.4	590.2	693.6
C_2	516.4	0	833.1	881.1
Dallas	590.2	833.1	0	464.5
C_1	693.6	881.1	464.5	0

The smallest distance is 464.5 between Dallas and C_1 , so that $C_3 = \{\text{Dallas}, C_1\}$. The distance matrix for Atlanta, C_2 , and C_3 is given by

Atlanta	0	516.4	590.2
C_2	516.4	0	833.1
C_3	590.2	833.1	0

The smallest distance is 516.4, which defines $C_4 = \{\text{Atlanta}, C_2\}$. The distance matrix for C_3 and C_4 is

C_3	0	590.2
C_4	590.2	0

The last cluster is given by $C_5 = \{C_3, C_4\}$. The dendrogram for the steps in this example is given in Figure 15.2. The order in which the clusters were formed and the relative distances at which they formed can all be seen. Note that the distance scale runs from right to left. \square

■ EXAMPLE 15.3.2(b)

To further illustrate the single linkage method of clustering, we use the complete city crime data from Table 15.1. The dendrogram in Figure 15.3 shows the cluster groupings attained by the single linkage method. \square

15.3.3 Complete Linkage (Farthest Neighbor)

In the *complete linkage* approach, also called the *farthest neighbor* method, the distance between two clusters A and B is defined as the *maximum* distance between a point in A and a point in B :

$$D(A, B) = \max \{d(\mathbf{y}_i, \mathbf{y}_j) \text{ for } \mathbf{y}_i \text{ in } A \text{ and } \mathbf{y}_j \text{ in } B\}. \quad (15.9)$$

At each step, the distance (15.9) is found for every pair of clusters, and we merge the two clusters with the smallest distance.

■ EXAMPLE 15.3.3(a)

As in Example 15.3.2(a) for single linkage clustering, we illustrate the use of the distance matrix in complete linkage clustering with the first six observations of the city crime data in Table 15.1. The initial distance matrix is exactly the same as in Example 15.3.2(a):




Figure 15.2 Dendrogram for single linkage of the first six observations in the city crime data in Table 15.1 [See Example 15.3.2(a)].

CITY	DISTANCE BETWEEN CITIES					
	Atlanta	Boston	Chicago	Dallas	Denver	Detroit
Atlanta	0	536.6	516.4	590.2	693.6	716.2
Boston	536.6	0	447.4	833.1	915.0	881.1
Chicago	516.4	447.4	0	924.0	1073.4	971.5
Dallas	590.2	833.1	924.0	0	527.7	464.5
Denver	693.6	915.0	1073.4	527.7	0	358.7
Detroit	716.2	881.1	971.5	464.5	358.7	0

The smallest distance is 358.7 between Denver and Detroit, and these two therefore form the first cluster, $C_1 = \{\text{Denver, Detroit}\}$. Note that since the first cluster is based on the initial distance matrix, it will be the same regardless of which hierarchical clustering method is used.




Figure 15.3 Dendrogram for single linkage of the complete city crime data from Table 15.1 [see Example 15.3.2(b)].

In the next step, the distance matrix is calculated for Atlanta, Boston, Chicago, Dallas, and C_1 :

Atlanta	0	536.6	516.4	590.2	716.2
Boston	536.6	0	447.4	833.1	915.0
Chicago	516.4	447.4	0	924.0	1073.4
Dallas	590.2	833.1	924.0	0	527.7
C_1	716.2	915.0	1073.4	527.7	0

Note that this distance matrix differs from its analog for the second step in Example 15.3.2(a) only in the distances between C_1 and the other cities. All elements of this matrix and subsequent distance matrices below are contained in the original distance matrix for the six cities. The smallest distance is 447.4 between Boston and Chicago. Therefore, $C_2 = \{\text{Boston, Chicago}\}$. At the next step, distances are calculated for Atlanta, Dallas, C_1 , and C_2 :

Atlanta	0	536.6	590.2	716.2
C_2	536.6	0	924.0	833.1
Dallas	590.2	924.0	0	527.7
C_1	693.6	881.1	527.7	0

The smallest distance, 527.7, defines $C_3 = \{\text{Dallas, } C_1\}$. The distance matrix for Atlanta, C_2 , and C_3 is given by

Atlanta	0	536.6	716.2
C_2	536.6	0	1073.4
C_3	590.2	1073.4	0

The smallest distance is 536.6 between Atlanta and C_3 , so that $C_4 = \{\text{Atlanta, } C_2\}$. The distance matrix for C_3 and C_4 is

C_3	0	1073.4
C_4	1073.4	0

The last cluster is given by $C_5 = \{C_3, C_4\}$. The dendrogram in Figure 15.4 shows the steps in this example. \square

■ EXAMPLE 15.3.3(b)

To further illustrate the complete linkage method, we use the complete crime data in Table 15.1. The dendrogram in Figure 15.5 shows the clusters found for this data set by the complete linkage approach. There are some differences between these groupings and the groupings from single linkage in Figure 15.3. \square

15.3.4 Average Linkage

In the *average linkage* approach, the distance between two clusters A and B is defined as the average of the $n_A n_B$ distances between the n_A points in A and the n_B




Figure 15.4 Dendrogram for complete linkage of the first six observations in the city crime data in Table 15.1 [see Example 15.3.3(a)].




Figure 15.5 Dendrogram for complete linkage of the complete city crime data of Table 15.1 [see Example 15.3.3(b)].

points in B :

$$D(A, B) = \frac{1}{n_A n_B} \sum_{i=1}^{n_A} \sum_{j=1}^{n_B} d(\mathbf{y}_i, \mathbf{y}_j), \quad (15.10)$$

where the sum is over all \mathbf{y}_i in A and all \mathbf{y}_j in B . At each step, we join the two clusters with the smallest distance, as measured by (15.10).

■ EXAMPLE 15.3.4

Figure 15.6 shows the dendrogram resulting from the average linkage method applied to the city crime data in Table 15.1. The solution is the same as the complete linkage solution for this data set as given in Example 15.3.3(b) and Figure 15.5. \square

15.3.5 Centroid

In the *centroid* method, the distance between two clusters A and B is defined as the Euclidean distance between mean vectors (often called centroids) of the two clusters:

$$D(A, B) = d(\bar{\mathbf{y}}_A, \bar{\mathbf{y}}_B), \quad (15.11)$$

where $\bar{\mathbf{y}}_A$ and $\bar{\mathbf{y}}_B$ are the mean vectors for the observation vectors in A and the observation vectors in B , respectively, and $d(\bar{\mathbf{y}}_A, \bar{\mathbf{y}}_B)$ is defined in (15.2). We define $\bar{\mathbf{y}}_A$ and $\bar{\mathbf{y}}_B$ in the usual way, that is, $\bar{\mathbf{y}}_A = \sum_{i=1}^{n_A} \mathbf{y}_i / n_A$. The two clusters with the smallest distance between centroids are merged at each step.

After two clusters A and B are joined, the centroid of the new cluster AB is given by the weighted average

$$\bar{\mathbf{y}}_{AB} = \frac{n_A \bar{\mathbf{y}}_A + n_B \bar{\mathbf{y}}_B}{n_A + n_B}. \quad (15.12)$$

■ EXAMPLE 15.3.5

Figure 15.7 shows the dendrogram resulting from using the centroid clustering method on the complete city crime data in Table 15.1.

Note the two crossovers in the dendrogram in Figure 15.7. Boston and Chicago join at a distance of 447.4. Then that cluster joins with {Atlanta, Tucson, Hartford} at a distance of 441.1. Finally, all five join with New Orleans at a distance of 393.8. Crossovers are discussed in Section 15.3.9a. \square

15.3.6 Median

If two clusters A and B are combined using the centroid method, and if A contains a larger number of items than B , then the new centroid $\bar{\mathbf{y}}_{AB} = (n_A \bar{\mathbf{y}}_A + n_B \bar{\mathbf{y}}_B) / (n_A + n_B)$ may be much closer to $\bar{\mathbf{y}}_A$ than to $\bar{\mathbf{y}}_B$. To avoid weighting the mean vectors according to cluster size, we can use the median (midpoint) of the line




Figure 15.6 Dendrogram for average linkage clustering of the data in Table 15.1 (see Example 15.3.4).

joining A and B as the point for computing new distances to other clusters:

$$\mathbf{m}_{AB} = \frac{1}{2}(\bar{\mathbf{y}}_A + \bar{\mathbf{y}}_B). \quad (15.13)$$

The two clusters with the smallest distance between medians are merged at each step.

Note that the “median” in (15.13) is not the ordinary median in the statistical sense. The terminology arises from a median of a triangle, namely, the line from a vertex to the midpoint of the opposite side.




Figure 15.7 Dendrogram for the centroid clustering of the complete city crime data in Table 15.1 (see Example 15.3.5).

■ EXAMPLE 15.3.6

Figure 15.8 shows the dendrogram resulting from using the median distance clustering method on the complete city crime data in Table 15.1. In Figure 15.8, we see the same two crossovers as in Figure 15.7. \square




Figure 15.8 Dendrogram for the median clustering method applied to the complete city crime data in Table 15.1 (see Example 15.3.6).

15.3.7 Ward's Method

Ward's method, also called the *incremental sum of squares method*, uses the within-cluster (squared) distances and the between-cluster (squared) distances (Ward 1963, Wishart 1969a). If AB is the cluster obtained by combining clusters A and B , then

the sums of within-cluster distances (of the items from the cluster mean vectors) are

$$\text{SSE}_A = \sum_{i=1}^{n_A} (\mathbf{y}_i - \bar{\mathbf{y}}_A)'(\mathbf{y}_i - \bar{\mathbf{y}}_A), \quad (15.14)$$

$$\text{SSE}_B = \sum_{i=1}^{n_B} (\mathbf{y}_i - \bar{\mathbf{y}}_B)'(\mathbf{y}_i - \bar{\mathbf{y}}_B), \quad (15.15)$$

$$\text{SSE}_{AB} = \sum_{i=1}^{n_{AB}} (\mathbf{y}_i - \bar{\mathbf{y}}_{AB})'(\mathbf{y}_i - \bar{\mathbf{y}}_{AB}), \quad (15.16)$$

where $\bar{\mathbf{y}}_{AB} = (n_A \bar{\mathbf{y}}_A + n_B \bar{\mathbf{y}}_B) / (n_A + n_B)$ as in (15.12) and n_A, n_B , and $n_{AB} = n_A + n_B$ are the numbers of points in A , B , and AB respectively. Since these sums of distances are equivalent to within-cluster sums of squares, they are denoted by SSE_A , SSE_B , and SSE_{AB} .

Ward's method joins the two clusters A and B that minimize the increase in SSE, defined as

$$I_{AB} = \text{SSE}_{AB} - (\text{SSE}_A + \text{SSE}_B). \quad (15.17)$$

It can be shown that the increase I_{AB} in (15.17) has the following two equivalent forms:

$$I_{AB} = n_A (\bar{\mathbf{y}}_A - \bar{\mathbf{y}}_{AB})' (\bar{\mathbf{y}}_A - \bar{\mathbf{y}}_{AB}) + n_B (\bar{\mathbf{y}}_B - \bar{\mathbf{y}}_{AB})' (\bar{\mathbf{y}}_B - \bar{\mathbf{y}}_{AB}) \quad (15.18)$$

$$= \frac{n_A n_B}{n_A + n_B} (\bar{\mathbf{y}}_A - \bar{\mathbf{y}}_B)' (\bar{\mathbf{y}}_A - \bar{\mathbf{y}}_B). \quad (15.19)$$

Thus by (15.19), minimizing the increase in SSE is equivalent to minimizing the *between cluster* distances. If A consists only of \mathbf{y}_i and B consists only of \mathbf{y}_j , then SSE_A and SSE_B are zero, and (15.17) and (15.19) reduce to

$$I_{ij} = \text{SSE}_{AB} = \frac{1}{2} (\mathbf{y}_i - \mathbf{y}_j)' (\mathbf{y}_i - \mathbf{y}_j) = \frac{1}{2} d^2(\mathbf{y}_i, \mathbf{y}_j).$$

Ward's method is related to the centroid method in Section 15.3.5. If the distance $d(\bar{\mathbf{y}}_A, \bar{\mathbf{y}}_B)$ in (15.11) is squared and compared to (15.19), the only difference is the coefficient $n_A n_B / (n_A + n_B)$ for Ward's method. Thus the cluster sizes have an impact on Ward's method but not on the centroid method. Writing $n_A n_B / (n_A + n_B)$ in (15.19) as

$$\frac{n_A n_B}{n_A + n_B} = \frac{1}{1/n_A + 1/n_B},$$

we see that as n_A and n_B increase, $n_A n_B / (n_A + n_B)$ increases. Writing the coefficient as

$$\frac{n_A n_B}{n_A + n_B} = \frac{n_A}{1 + n_A/n_B},$$

we see that as n_B increases with n_A fixed, $n_A n_B / (n_A + n_B)$ increases. Therefore, compared to the centroid method, Ward's method is more likely to join smaller clusters or clusters of equal size.




Figure 15.9 Dendrogram for Ward's method applied to the complete city crime data in Table 15.1 (see Example 15.3.7).

■ EXAMPLE 15.3.7

Figure 15.9 shows the dendrogram resulting from using Ward's clustering method on the complete city crime data in Table 15.1. The vertical axis is $I_{AB} / \sum_{i=1}^n (\mathbf{y}_i - \bar{\mathbf{y}})'(\mathbf{y}_i - \bar{\mathbf{y}})$, where $\bar{\mathbf{y}}$ is the overall mean vector for the data.

□

15.3.8 Flexible Beta Method

Suppose clusters A and B have just been merged to form cluster AB . A general formula for the distance between AB and any other cluster C was given by Lance and Williams (1967):

$$\begin{aligned} D(C, AB) &= \alpha_A D(C, A) + \alpha_B D(C, B) + \beta D(A, B) \\ &\quad + \gamma |D(C, A) - D(C, B)|. \end{aligned} \quad (15.20)$$

The distances $D(C, A)$, $D(C, B)$, and $D(A, B)$ are from the distance matrix before joining A and B . The distances from AB to other clusters as given by (15.20) would be used (along with distances between other pairs of clusters) to form the next distance matrix for choosing the pair of clusters with smallest distance. This pair would then be joined at the next step.

To simplify (15.20), Lance and Williams (1967) suggested the following constraints on the parameter values:

$$\begin{aligned} \alpha_A + \alpha_B + \beta &= 1 \\ \alpha_A &= \alpha_B \\ \gamma &= 0 \\ \beta &< 1. \end{aligned}$$

With $\alpha_A = \alpha_B$ and $\gamma = 0$, we have $2\alpha_A = 1 - \beta$ or $\alpha_A = \alpha_B = (1 - \beta)/2$, and we need only choose a value of β . The resulting hierarchical clustering procedure is called the *flexible beta* method.

The choice of β determines the characteristics of the flexible beta clustering procedure. Lance and Williams (1967) suggested the use of a small negative value of β , such as $\beta = -.25$. If there are (or might be) outliers in the data, the use of a smaller value of β , such as $\beta = -.5$, may be more likely to isolate these outliers into simple clusters.

The distances defined for the agglomerative hierarchical methods in Sections 15.3.2–15.3.7 can all be expressed as special cases of (15.20). The requisite parameter values are given in Table 15.2. For the centroid, median, and Ward's methods, the distances in (15.20) must be squared distances (assuming Euclidean distances). For the other methods in Table 15.2, the distances may be either squared or unsquared.

We illustrate the choice of parameter values in Table 15.2 for the single linkage method. Using $\alpha_A = \alpha_B = \frac{1}{2}$, $\beta = 0$, and $\gamma = -\frac{1}{2}$ as in the first row of Table 15.2, (15.20) becomes

$$D(C, AB) = \frac{1}{2}D(C, A) + \frac{1}{2}D(C, B) - \frac{1}{2}|D(C, A) - D(C, B)|. \quad (15.21)$$

If $D(C, A) > D(C, B)$, then $|D(C, A) - D(C, B)| = D(C, A) - D(C, B)$, and (15.21) reduces to

$$D(C, AB) = D(C, B). \quad (15.22)$$

On the other hand, if $D(C, A) < D(C, B)$, then $|D(C, A) - D(C, B)| = D(C, B) - D(C, A)$, and (15.21) reduces to

$$D(C, AB) = D(C, A). \quad (15.23)$$

Table 15.2 Parameter Values for (15.20)

Cluster Method	α_A	α_B	β	γ
Single linkage	$\frac{1}{2}$	$\frac{1}{2}$	0	$-\frac{1}{2}$
Complete linkage	$\frac{1}{2}$	$\frac{1}{2}$	0	$\frac{1}{2}$
Average linkage	$\frac{n_A}{n_A + n_B}$	$\frac{n_B}{n_A + n_B}$	0	0
Centroid	$\frac{n_A}{n_A + n_B}$	$\frac{n_B}{n_A + n_B}$	$\frac{-n_A n_B}{(n_A + n_B)^2}$	0
Median	$\frac{1}{2}$	$\frac{1}{2}$	$-\frac{1}{4}$	0
Ward's method	$\frac{n_A + n_C}{n_A + n_B + n_C}$	$\frac{n_B + n_C}{n_A + n_B + n_C}$	$\frac{-n_C}{n_A + n_B + n_C}$	0
Flexible beta	$(1 - \beta)/2$	$(1 - \beta)/2$	$\beta (< 1)$	0

Thus, (15.21) can be written as

$$D(C, AB) = \min[D(C, A), D(C, B)], \quad (15.24)$$

which is equivalent to (15.8), the definition of distance for the single linkage method.

■ EXAMPLE 15.3.8

Figures 15.10 and 15.11 show dendograms produced when using the flexible beta clustering method on the complete city crime data in Table 15.1, with $\beta = -.25$ and $\beta = -.75$. The two results are similar. \square

15.3.9 Properties of Hierarchical Methods

15.3.9a Monotonicity

If an item or a cluster joins another cluster at a distance that is less than the distance for the previous merger of two clusters, we say that an *inversion* or a *reversal* has occurred. The reversal is represented by a *crossover* in the dendrogram. Examples of crossovers can be found in Figures 15.7 and 15.8.

A hierarchical method in which reversals cannot occur is said to be *monotonic*, because the distance at each step is greater than the distance at the previous step. A distance measure or clustering method that is monotonic is also called *ultrametric*.

We now show that the single linkage and complete linkage methods are monotonic. Let d_k be the distance at which two clusters are joined at the k th step. We can describe steps k and $k + 1$ in terms of four clusters A, B, C , and D . Suppose $D(A, B)$ is less than the distance between any other pair among these four clusters,




Figure 15.10 Dendrogram for the flexible beta method with $\beta = -.25$ applied to the complete city crime data in Table 15.1 (see Example 15.3.8).




Figure 15.11 Dendrogram for the flexible beta method with $\beta = -.75$ applied to the complete city crime data in Table 15.1 (see Example 15.3.8).

so that A and B are joined at step k to form AB . Then

$$d_k = D(A, B) < \min\{D(A, C), D(B, C), D(C, D)\}. \quad (15.25)$$

[If $D(A, B)$ is less than these three distances, it is less than the other two possible distances, $D(A, D)$ and $D(B, D)$.] Suppose at step $k + 1$ we join AB and C or we join C and D . If we merge C and D , then by (15.25), $d_k = D(A, B) < D(C, D) = d_{k+1}$. If we join AB and C , then for single linkage (15.24) gives

$$d_{k+1} = D(C, AB) = \min\{D(A, C), D(B, C)\} > d_k = D(A, B).$$

By (15.25), both of $D(A, C)$ and $D(B, C)$ exceed $D(A, B)$, and this also holds for complete linkage. Thus, the single linkage and complete linkage methods are monotonic.

For the methods in Table 15.2 other than single linkage and complete linkage, we have $\gamma = 0$ and by (15.20) and (15.25),

$$D(C, AB) > (\alpha_A + \alpha_B + \beta)D(A, B). \quad (15.26)$$

Thus we need $\alpha_A + \alpha_B + \beta \geq 1$ for monotonicity. Using this criterion, we see that all methods in Table 15.1 (beyond the first two) are monotonic except the centroid and median methods. (These two methods showed crossovers in the dendograms in Figures 15.7 and 15.8.) Because of lack of monotonicity, some authors do not recommend the centroid and median methods.

15.3.9b Contraction or Dilatation

We now consider the characteristics of the distances or proximities between the original points. As clusters form, the properties of this space of distances may be altered somewhat. A clustering method that does not alter the spatial properties is referred to by Lance and Williams (1967) as *space-conserving*. A method that is not space-conserving may either *contract* or *dilate* the space.

A method is *space-contracting* if newly formed clusters appear to move closer to individual observations, so that an individual item tends to join an existing cluster rather than join with another individual item to form a new cluster. This tendency is also called *chaining*.

A method is *space-dilating* if newly formed clusters appear to move away from individual observations, so that individual items tend to form new clusters rather than join existing clusters. In this case, clusters appear to be more distinct than they are.

Dubien and Warde (1979) described the spatial properties as follows. Suppose that the distances among three clusters satisfy

$$D(A, B) < D(A, C) < D(B, C).$$

Then a cluster method is space-conserving if

$$D(A, C) < D(AB, C) < D(B, C). \quad (15.27)$$

A method is space-contracting if the first inequality in (15.27) does not hold and space-dilating if the second inequality does not hold.

The single linkage method is very space-contracting, with marked chaining tendencies. For this reason, single linkage is not recommended by some authors. Complete linkage on the other hand, is very space-dilating, with a tendency to artificially impose cluster boundaries.

Other hierarchical methods fall in between the extremes represented by single linkage and complete linkage. The centroid and average linkage methods are largely space-conserving, while Ward's method is space-contracting. Whenever a method produces reversals for a particular data set, it can be considered to be space-contracting. Thus the centroid method is space-conserving unless it has reversals, whereupon it becomes space-contracting.

The flexible beta method is space-contracting for $\beta > 0$, space-conserving for $\beta = 0$, and space-dilating for $\beta < 0$. A small degree of dilation may help define cluster boundaries, but too much dilation may lead to too many clusters in the early stages. Thus the recommended value of $\beta = -.25$ may represent a good compromise.




Figure 15.12 Two distinct clusters with intervening individuals.




Figure 15.13 Single linkage clustering of the data in Figure 15.12.

■ EXAMPLE 15.3.9b

To illustrate chaining in the single linkage method, consider the data plotted in Figure 15.12 (see Everitt 1993, p. 68). Two distinct clusters, *A* and *C*, have points between them labeled *B* that do not belong to *A* or *C*.

In Figure 15.13, the two-cluster solution for single linkage clustering places *C*₁ and *C*₁₁ into one cluster and all other points into another cluster. The three-cluster solution has two clusters with *C*'s and a cluster with *A*'s and *B*'s.




Figure 15.14 Average linkage clustering of the data in Figure 15.12.

A dendrogram for average linkage clustering of the data in Figure 15.12 is given in Figure 15.14. For this data set, the average linkage method is more robust to chaining. The two-cluster solution separates the *C*'s from the *A*'s and *B*'s. The three-cluster solution completely separates the three groups, *A*, *B*, and *C*. \square

15.3.9c Other Properties

The single linkage method has been criticized by many authors because of its chaining tendencies and because it is sensitive to errors in distances between observations.




Figure 15.15 Clusters in a single dimension.

On the other hand, the single linkage approach is better than the other methods at identifying clusters that have curvy shapes instead of spherical or elliptical shapes, and it is somewhat robust to outliers in the data.

Ward's method and the average linkage method are also relatively insensitive to outliers. For example, in the average linkage method, outliers tend to remain isolated in the early stages and to join with other outliers rather than to join with large clusters or with less compact clusters. This is due to two properties of the average linkage method: (1) the average distance between two groups (squared Euclidean distance) increases as the points in the groups are more spread out, and (2) the average distance increases as the size of the groups increases.

These two properties of the average linkage method are illustrated in one dimension in Figure 15.15 (see Jobson 1992, pp. 524–525), where cluster A has one point at z_1 and cluster B has two points, b_1 and b_2 , located at $z_2 - h$ and $z_2 + h$. The average squared distance between A and B is

$$\begin{aligned}\overline{d^2} &= \frac{1}{2}[(z_1 - z_2 + h)^2 + (z_1 - z_2 - h)^2] \\ &= \frac{1}{2}[(z_1 - z_2)^2 + h^2 + 2h(z_1 - z_2) + (z_1 - z_2)^2 + h^2 - 2h(z_1 - z_2)] \\ &= (z_1 - z_2)^2 + h^2.\end{aligned}$$

Thus the average distance between A and B increases as the spread of b_1 and b_2 increases (that is, as h increases).

To illustrate the second property of the average linkage method, suppose cluster B in Figure 15.15 consists of a single point located at z_2 . Then, the distance between A and B is $(z_1 - z_2)^2$, and A is closer to B than it is if B consists of two points.

The centroid method is fairly robust to outliers. Complete linkage is somewhat sensitive to outliers and tends to produce clusters of the same size and shape. Ward's method tends to yield spherical clusters of the same size.

Many studies conclude that the best overall performers are Ward's method and the average linkage method. However, there seems to be an "interaction" between methods and data sets; that is, some methods work better for certain data sets, and other methods work better for other data sets.

A good strategy is to try several methods. If the results agree to some extent, the researcher may have found some natural clusters in the data.

15.3.10 Divisive Methods

In the agglomerative hierarchical methods covered in Sections 15.3.2–15.3.9, we begin with n items and end with a single cluster containing all n items. As noted in the second paragraph of Section 15.3.1, a divisive hierarchical method starts with a single cluster of n items and divides it into two groups. At each step thereafter, one of the groups is divided into two subgroups. The ultimate result of a divisive algorithm is n clusters of one item each. The results can be shown in a dendrogram.

Divisive methods suffer from the same potential drawback as the agglomerative methods, namely, that once a partition is made, an item cannot be moved into another group it does not belong to at the time of the partitioning. However, if larger clusters are of interest, then the divisive approach may sometimes be preferred over the agglomerative approach, in which the larger clusters are reached only after a large number of joinings of smaller groups.

Divisive algorithms are generally of two classes: monothetic and polythetic. In a *monothetic* approach, the division of a group into two subgroups is based on a single variable, whereas the *polythetic* approach uses all p variables to make the split.

If the variables are binary (quantitative variables can be converted to binary variables), the monothetic approach can easily be applied. Division into two groups is based on presence or absence of an attribute. The variable (attribute) is chosen which maximizes a chi-square statistic or an information statistic; see Everitt (1993, pp. 87–88) or Gordon (1999, pp. 130–134).

For a monothetic approach using a quantitative variable y , we seek to maximize the between-group sum of squares,

$$\text{SSB} = n_1(\bar{y}_1 - \bar{y})^2 + n_2(\bar{y}_2 - \bar{y})^2,$$

where n_1 and n_2 are the two group sizes (with $n_1 + n_2 = n$), \bar{y}_1 and \bar{y}_2 are the group means, and \bar{y} is the overall mean based on all n observations. The sum of squares SSB would be calculated for all possible splits into two groups of sizes n_1 and n_2 and for each of the p variables. The final division would be based on the variable that maximizes $\text{SSB}/\sum_{i=1}^n(y_i - \bar{y})^2$.

Table 15.3 Athletic Records for Eight Events in Eight Countries

Country	1	2	3	4	5	6	7	8
Australia	10.31	20.06	44.84	1.74	3.57	13.28	27.66	128.30
Belgium	10.34	20.68	45.04	1.73	3.60	13.22	27.45	129.95
Canada	10.17	20.22	45.68	1.76	3.63	13.55	28.09	130.15
GDR	10.12	20.33	44.87	1.73	3.56	13.17	27.42	129.92
GB	10.11	20.21	44.93	1.70	3.51	13.01	27.51	129.13
Kenya	10.46	20.66	44.92	1.73	3.55	13.10	27.80	129.75
USA	9.93	19.75	43.86	1.73	3.53	13.20	27.43	128.22
USSR	10.07	20.00	44.60	1.75	3.59	13.20	27.53	130.55

Event: (1) 100 m (s), (2) 200 m (s), (3) 400 m (s), (4) 800 m (min), (5) 1500 m (min),
 (6) 5000 m (min), (7) 10,000 m (min), (8) marathon (min).

Table 15.4 Average Distance from Each Country to the Other Seven

Country	Average Distance	Country	Average Distance
USA	2.068	USSR	1.513
Aust	1.643	Canada	1.594
GB	1.164	Kenya	1.156
GDR	1.083	Belgium	1.16

For a polythetic approach, we consider a technique proposed by MacNaughton-Smith et al. (1964). To divide a group, we work with a splinter group and the remainder. We seek the item in the remainder whose average distance (dissimilarity) from other items in the remainder, minus its average distance from items in the splinter group, is largest. If the largest difference is positive, the item is shifted to the splinter group. If the largest difference is negative, the procedure stops, and the division is complete. We can start the splinter group with the item that has the largest average distance from the other items in the group.

■ EXAMPLE 15.3.10

In Table 15.3 we have the track records of eight countries (Dawkins 1989). Based on the distance matrix for these eight observations, the average distance from each observation to the other seven observations is given in Table 15.4. Since USA has the greatest average distance to the other countries, USA becomes the first observation in the splinter group. Now, the average distance between each observation in the remainder to the other six observations in the remainder is calculated. Then the (average) distance between USA and each

Table 15.5 Average Distances to Remainder and Splinter Groups for Seven Countries

Country	Average Distance to Remainder (1)	Average Distance to Splinter Group (2)	Difference (1) – (2)
Australia	1.729	1.126	.603
GB	1.108	1.504	–.396
GDR	.918	2.070	–1.151
USSR	1.355	2.464	–1.111
Canada	1.392	2.808	–1.416
Kenya	.986	2.173	–1.186
Belgium	.975	2.329	–1.353

Table 15.6 Average Distances to Remainder and Splinter Groups for Six Countries

Country	Average Distance to Remainder (1)	Average Distance to Splinter Group (2)	Difference (1) – (2)
GB	1.144	1.216	–.072
GDR	.767	1.872	–1.105
USSR	1.169	2.373	–1.203
Canada	1.249	2.457	–1.208
Kenya	.865	1.884	–1.019
Belgium	.813	2.058	–1.245

item in the remainder is calculated. (This may be found using the distance matrix since there is only one observation in the splinter group.) Finally, we calculate the difference between the average distance to the remainder and the average distance to the splinter group. The results are in Table 15.5. Because Australia has a positive difference in Table 15.5, it is added to the splinter group with USA. This process is repeated for the six countries in the remainder, and the results are given in Table 15.6. Since no difference in Table 15.6 is positive, the process stops, and we have the following clusters: $C_1 = \{\text{USA}, \text{Australia}\}$, $C_2 = \{\text{GB}, \text{GDR}, \text{USSR}, \text{Canada}, \text{Kenya}, \text{Belgium}\}$. We could continue and divide C_2 into two groups in the same way. \square

15.4 NONHIERARCHICAL METHODS

In this section, we discuss three nonhierarchical techniques: partitioning, mixtures of distributions, and density estimation. Among these three methods, partitioning is the most commonly used.

15.4.1 Partitioning

In the partitioning approach, the observations are separated into g clusters without using a hierarchical approach based on a matrix of distances or similarities between all pairs of points. The methods described in this section are sometimes called *optimization methods* rather than *partitioning*.

An attractive strategy would be to examine all possible ways to partition n items into g clusters and find the optimal clustering according to some criterion. However, the number of possible partitions as given by (15.7) is prohibitively large for even moderate values of n and g . Thus we seek simpler techniques.

15.4.1a *k*-Means

We now consider an approach to partitioning that is usually called the *k-means method*. (We will continue to use the notation g rather than k for the number of clusters.) The method allows the items to be moved from one cluster to another, a reallocation that is not available in the hierarchical methods.

We first select g items to serve as “seeds.” These are later replaced by the centroids (mean vectors) of the clusters. There are various ways we can choose the seeds: select g items at random (perhaps separated by some minimum distance), choose the first g points in the data set (again subject to a minimum distance apart), select the g points that are mutually farthest apart, find the g points of maximum density, or specify g regularly spaced points in a grid-like pattern (these would not be actual data points).

For these methods of selecting seeds, the number of clusters, g , must be specified. Alternatively, a minimum distance between seeds may be specified, and then all items that satisfy this criterion are chosen as seeds.

After the seeds are chosen, each remaining point in the data set is assigned to the cluster with the nearest seed (based on Euclidean distance). As soon as a cluster has more than one member, the cluster seed is replaced by the centroid.

After all items are assigned to clusters, each item is examined to see if it is closer to the centroid of another cluster than to the centroid of its own cluster. If so, the item is moved to the new cluster and the two cluster centroids are updated. This process is continued until no further improvement is possible.

The *k*-means procedure is somewhat sensitive to the initial choice of seeds. It might be advisable to try the procedure again with a new initial choice of seeds. If different initial choices of seeds produce widely different final clusters, or if convergence is extremely slow, there may be no natural clusters in the data.

The *k*-means partitioning method can also be used as a possible improvement on hierarchical techniques. We first cluster the items using a hierarchical method and then use the centroids of these clusters as seeds for a *k*-means approach, which will allow points to be reallocated from one cluster to another.

■ EXAMPLE 15.4.1a

Protein consumption in twenty-five European countries for nine food groups is given in Table 15.7 (Hand et al. 1994, p. 298). In order to illustrate the

Table 15.7 Protein Data

Country	Red Meat	White Meat	Eggs	Milk	Fish	Cereals	Starchy Foods	Nuts	Fruit/Veg
Albania	10.1	1.4	.5	8.9	.2	42.3	.6	5.5	1.7
Austria	8.9	14.0	4.3	19.9	2.1	28.0	3.6	1.3	4.3
Belgium	13.5	9.3	4.1	17.5	4.5	26.6	5.7	2.1	4.0
Bulgaria	7.8	6.0	1.6	8.3	1.2	56.7	1.1	3.7	4.2
Czech.	9.7	11.4	2.8	12.5	2.0	34.3	5.0	1.1	4.0
Denmark	10.6	10.8	3.7	25.0	9.9	21.9	4.8	.7	2.4
E. Germany	8.4	11.6	3.7	11.1	5.4	24.6	6.5	.8	3.6
Finland	9.5	4.9	2.7	33.7	5.8	26.3	5.1	1.0	1.4
France	18.0	9.9	3.3	19.5	5.7	28.1	4.8	2.4	6.5
Greece	10.2	3.0	2.8	17.6	5.9	41.7	2.2	7.8	6.5
Hungary	5.3	12.4	2.9	9.7	.3	40.1	4.0	5.4	4.2
Ireland	13.9	10.0	4.7	25.8	2.2	24.0	6.2	1.6	2.9
Italy	9.0	5.1	2.9	13.7	3.4	36.8	2.1	4.3	6.7
Netherlands	9.5	13.6	3.6	23.4	2.5	22.4	4.2	1.8	3.7
Norway	9.4	4.7	2.7	23.3	9.7	23.0	4.6	1.6	2.7
Poland	6.9	10.2	2.7	19.3	3.0	36.1	5.9	2.0	6.6
Portugal	6.2	3.7	1.1	4.9	14.2	27.0	5.9	4.7	7.9
Romania	6.2	6.3	1.5	11.1	1.0	49.6	3.1	5.3	2.8
Spain	7.1	3.4	3.1	8.6	7.0	29.2	5.7	5.9	7.2
Sweden	9.9	7.8	3.5	24.7	7.5	19.5	3.7	1.4	2.0
Switzerland	13.1	10.1	3.1	23.8	2.3	25.6	2.8	2.4	4.9
UK	17.4	5.7	4.7	20.6	4.3	24.3	4.7	3.4	3.3
USSR	9.3	4.6	2.1	16.6	3.0	43.6	6.4	3.4	2.9
W. Germany	11.4	12.5	4.1	18.8	3.4	18.6	5.2	1.5	3.8
Yugoslavia	4.4	5.0	1.2	9.5	.6	55.9	3.0	5.7	3.2

sensitivity of the k -means clustering method to the initial choice of seeds we use the following four methods of choosing seeds:

1. Select at random g observations that are at least a distance r apart.
2. Select the first g observations that are at least a distance r apart.
3. Select the g observations that are mutually farthest apart.
4. Use the g centroids from the g -cluster solution from the average linkage (hierarchical) clustering method.

To help choose g , the number of clusters, we plot the first two principal components in Figure 15.16. It appears that there may be at least five clusters. For the first method, we select five observations at random that are at least a distance $r = 1$ from each other. The five chosen seeds are West Germany, Austria, Ireland, France, and Czechoslovakia. Using these seeds, the k -means




Figure 15.16 First two principal components z_1 and z_2 for the protein data in Table 15.7.

method produced the clusters identified in Table 15.8 along with the distance of each observation from its cluster centroid.

To view the clusters, we plot the first two discriminant functions (see Section 8.4.1) in Figure 15.17. The first two discriminant functions show good separation among clusters 3, 4 and 5. However, clusters 1 and 2 appear to overlap with the other clusters.


We now select the first five observations as cluster seeds. With these seeds, the k -means clustering method produced the clusters in Table 15.9. The first two discriminant functions are plotted in Figure 15.18. Good separation of clusters is seen except for clusters 2 and 3.

We next choose as cluster seeds the five observations that are mutually farthest apart. These seeds gave rise to the clusters in Table 15.10. The first two discriminant functions are plotted in Figure 15.19. Clusters 1, 3, and 4 seem very well separated, but clusters 2 and 5 show considerable overlap.

Finally, we obtain a five-cluster solution from average linkage and use the centroids of these clusters as the new seeds. The clusters in Table 15.11 result. The first two discriminant functions are plotted in Figure 15.20. All five clusters are well separated in the first two discriminant functions. These clusters show some resemblance to those in the principal components plot given in Figure 15.16. \square

Table 15.8 *k*-Means Cluster Solution for Seeds Chosen at Random

Country	Cluster	Distance from Centroid	Country	Cluster	Distance from Centroid
Denmark	1	1.227	Finland	3	2.261
Sweden	1	1.247	France	4	2.273
Norway	1	1.629	Greece	4	2.273
Belgium	1	1.669	Romania	5	1.514
E. Germany	1	1.987	Italy	5	1.981
Netherlands	2	0.991	Yugoslavia	5	2.040
Austria	2	1.160	Bulgaria	5	2.225
W. Germany	2	1.303	USSR	5	2.393
Czech.	2	1.433	Hungary	5	2.435
Switzerland	2	1.679	Spain	5	2.871
Poland	2	2.052	Albania	5	3.180
Ireland	3	1.334	Portugal	5	4.343
UK	3	1.821			

**Figure 15.17** First two discriminant functions z_1 and z_2 for the clusters in Table 15.8.

536 CLUSTER ANALYSIS

Table 15.9 *k*-Means Cluster Solution Using the First Five Observations as Seeds

Country	Cluster	Distance from Centroid	Country	Cluster	Distance from Centroid
Albania	1	.000	Romania	4	1.415
Netherlands	2	.648	Bulgaria	4	1.587
Austria	2	1.000	Yugoslavia	4	1.784
W. Germany	2	1.087	Italy	4	1.898
Switzerland	2	1.489	Greece	4	2.450
Belgium	3	1.368	Poland	5	1.709
Sweden	3	1.462	Czech.	5	1.956
Denmark	3	1.666	USSR	5	2.218
Ireland	3	1.832	E. Germany	5	2.285
Norway	3	1.927	Spain	5	2.344
UK	3	2.076	Hungary	5	2.558
Finland	3	2.341	Portugal	5	3.859
France	3	2.629			




Figure 15.18 First two discriminant functions z_1 and z_2 for the clusters in Table 15.9.

Table 15.10 *k*-Means Cluster Solution Using as Seeds the Five Observations Furthest Apart

Country	Cluster	Distance from Centroid	Country	Cluster	Distance from Centroid
Romania	1	.601	France	2	2.358
Yugoslavia	1	1.159	Poland	2	2.405
Bulgaria	1	1.435	UK	2	2.537
Albania	1	2.421	Greece	3	1.075
Hungary	1	2.540	Italy	3	1.075
Belgium	2	.956	Portugal	4	1.466
W. Germany	2	1.012	Spain	4	1.466
Netherlands	2	1.416	Norway	5	1.054
Austria	2	1.663	Sweden	5	1.191
Czech.	2	1.706	Finland	5	1.545
Switzerland	2	1.713	Denmark	5	1.708
Ireland	2	1.839	USSR	5	2.780
E. Germany	2	2.042			



**Figure 15.19** First two discriminant functions z_1 and z_2 for the clusters in Table 15.10.

Table 15.11 *k*-Means Cluster Solution Using Seeds from Average Linkage

Country	Cluster	Distance from Centroid	Country	Cluster	Distance from Centroid
Romania	1	.970	Norway	2	2.287
Yugoslavia	1	1.182	UK	2	2.354
Bulgaria	1	1.339	France	2	2.600
Albania	1	1.970	Finland	2	2.683
Belgium	2	1.152	Greece	3	1.075
W. Germany	2	1.245	Italy	3	1.075
Netherlands	2	1.547	Portugal	4	1.466
Sweden	2	1.604	Spain	4	1.466
Ireland	2	1.744	Czech.	5	1.337
Denmark	2	1.766	Poland	5	1.579
Switzerland	2	1.831	USSR	5	1.964
Austria	2	2.037	Hungary	5	2.023
E. Germany	2	2.251			

**Figure 15.20** First two discriminant functions z_1 and z_2 for the clusters in Table 15.11.

15.4.1b Other Partitioning Criteria

We now consider three partitioning methods that are not based directly on the distance from a point to the centroid of a cluster. These methods are based on the between-cluster and within-cluster sum of squares and products matrices \mathbf{H} and \mathbf{E} defined in (6.9) and (6.10) for one-way MANOVA. For well-defined clusters, we would like \mathbf{E} to be “small” and \mathbf{H} to be “large.”

The three criteria are as follows:

1. Minimize $\text{tr}(\mathbf{E})$
2. Minimize $|\mathbf{E}|$
3. Maximize $\text{tr}(\mathbf{E}^{-1}\mathbf{H})$.

Using criterion 1, for example, we would move an item with observation vector \mathbf{y} to the cluster for which $\text{tr}(\mathbf{E})$ is minimized after the move.

We can express the first criterion in two alternative forms. By (6.10), we have

$$\text{tr}(\mathbf{E}) = \text{tr} \left[\sum_{i=1}^g \sum_{j=1}^n (\mathbf{y}_{ij} - \bar{\mathbf{y}}_{i.})(\mathbf{y}_{ij} - \bar{\mathbf{y}}_{i.})' \right] \quad (15.28)$$

$$\begin{aligned} &= \sum_i \text{tr} \left[\sum_j (\mathbf{y}_{ij} - \bar{\mathbf{y}}_{i.})(\mathbf{y}_{ij} - \bar{\mathbf{y}}_{i.})' \right] \quad [\text{by (2.96)}] \\ &= \sum_i \text{tr}(\mathbf{E}_i), \end{aligned} \quad (15.29)$$

where $\mathbf{E}_i = \sum_{j=1}^n (\mathbf{y}_{ij} - \bar{\mathbf{y}}_{i.})(\mathbf{y}_{ij} - \bar{\mathbf{y}}_{i.})'$ is the sum of squares and products matrix of deviations of observations from the mean vector for the i th cluster. In (15.28) we are using the notation of Section 6.1.2 for a balanced design, in which n is the number of observations in each cluster.

We can write $\text{tr}(\mathbf{E}_i)$ in (15.29) in the form

$$\begin{aligned} \text{tr}(\mathbf{E}_i) &= \text{tr} \sum_j (\mathbf{y}_{ij} - \bar{\mathbf{y}}_{i.})(\mathbf{y}_{ij} - \bar{\mathbf{y}}_{i.})' \\ &= \sum_j \text{tr}(\mathbf{y}_{ij} - \bar{\mathbf{y}}_{i.})(\mathbf{y}_{ij} - \bar{\mathbf{y}}_{i.})' \quad [\text{by (2.96)}] \\ &= \sum_j (\mathbf{y}_{ij} - \bar{\mathbf{y}}_{i.})'(\mathbf{y}_{ij} - \bar{\mathbf{y}}_{i.}) \quad [\text{by (2.97)}]. \end{aligned} \quad (15.30)$$

Thus $\text{tr}(\mathbf{E}_i)$ is the sum of the (squared) Euclidean distances from the individual points to the centroid of the i th cluster.

A second form of (15.28) was given by Seber (1984, p. 277) as

$$\text{tr}(\mathbf{E}) = \frac{1}{n} \sum_i \sum_{k < m} (\mathbf{y}_{ik} - \mathbf{y}_{im})'(\mathbf{y}_{ik} - \mathbf{y}_{im}). \quad (15.31)$$

Hence minimizing $\text{tr}(\mathbf{E})$ is equivalent to minimizing the sum of squared Euclidean distances between all pairs of points in a cluster.

The second criterion, minimizing $|\mathbf{E}|$, is related to $\Lambda = |\mathbf{E}|/|\mathbf{E} + \mathbf{H}|$ in (6.13). Minimizing $|\mathbf{E}|$ is equivalent to minimizing Wilks' Λ .

Another way to look at minimizing $|\mathbf{E}|$ is to consider the effect of adding a point \mathbf{y} to a cluster with centroid $\bar{\mathbf{y}}$. Let $\mathbf{u} = \mathbf{y} - \bar{\mathbf{y}}$. By (15.28), \mathbf{E} is a sum of terms of the form $\mathbf{u}\mathbf{u}' = (\mathbf{y} - \bar{\mathbf{y}})(\mathbf{y} - \bar{\mathbf{y}})'$. Thus (ignoring the change in centroid with the added observation \mathbf{y}), the increase in $|\mathbf{E}|$ is

$$\begin{aligned} |\mathbf{E} + \mathbf{u}\mathbf{u}'| - |\mathbf{E}| &= |\mathbf{E}|(1 + \mathbf{u}'\mathbf{E}^{-1}\mathbf{u}) - |\mathbf{E}| \quad [\text{by (2.95)}] \\ &= |\mathbf{E}|\mathbf{u}'\mathbf{E}^{-1}\mathbf{u}. \end{aligned}$$

Hence, the minimum increase in $|\mathbf{E}|$ is obtained by adding \mathbf{y} to the cluster for which the standardized distance $\mathbf{u}'\mathbf{E}^{-1}\mathbf{u}$ of \mathbf{y} from $\bar{\mathbf{y}}$ is the smallest. By comparison, the $\text{tr}(\mathbf{E})$ criterion would add \mathbf{y} to the cluster for which $\mathbf{u}'\mathbf{u}$ is minimum [see (15.30)].

The third criterion, maximizing $\text{tr}(\mathbf{E}^{-1}\mathbf{H})$, is related to the Lawley–Hotelling statistic $U^{(s)} = \text{tr}(\mathbf{E}^{-1}\mathbf{H}) = \sum_{i=1}^s \lambda_i$ in (6.27), where $\lambda_1, \lambda_2, \dots, \lambda_s$ are the eigenvalues of $\mathbf{E}^{-1}\mathbf{H}$ and $s = \min(p, g - 1)$. Associated with each λ_i is the eigenvector \mathbf{a}_i and the discriminant function $z_i = \mathbf{a}_i'\mathbf{y}$ (see Section 8.4). The largest eigenvalue, λ_1 , and the accompanying first discriminant function, $z_1 = \mathbf{a}_1'\mathbf{y}$, have the greatest influence on $\text{tr}(\mathbf{E}^{-1}\mathbf{H})$. Maximizing $\text{tr}(\mathbf{E}^{-1}\mathbf{H})$ has the inclination to produce elliptical clusters of the same size. These would tend to follow a straight-line trend, especially if the first eigenvalue dominates the others. If the initial clusters or seeds are lined up in a different direction than the “true clusters,” maximizing $\text{tr}(\mathbf{E}^{-1}\mathbf{H})$ may not correct the error in subsequent iterations.

Since $\text{tr}(\mathbf{E})$ involves only the diagonal elements, the first criterion ignores the correlations and tends to yield spherical clusters. The second criterion, minimizing $|\mathbf{E}|$, takes correlations into account and tends to produce elliptical clusters. These clusters have a tendency to be of the same shape because \mathbf{E}/ν_E is a pooled estimator of the covariance matrix. A modification that may be useful is $\prod_{i=1}^g |\mathbf{E}_i|$, where \mathbf{E}_i is the error matrix for the i th cluster [see (15.29)].

Finally, we compare the three criteria in terms of invariance to nonsingular linear transformations $\mathbf{v}_{ij} = \mathbf{A}\mathbf{y}_{ij} + \mathbf{b}$, where \mathbf{A} is a constant nonsingular matrix and \mathbf{b} is a vector of constants. The first criterion, minimizing $\text{tr}(\mathbf{E})$, is not invariant to such linear transformations, while the other two criteria are invariant to these transformations. Therefore, minimizing $\text{tr}(\mathbf{E})$ will likely give different partitions for the raw data and standardized data.

15.4.2 Other Methods

We discuss mixtures of distributions in Section 15.4.2a and density estimation in Section 15.4.2b.

15.4.2a Mixtures of Distributions

In this method, we assume the existence of g distributions (usually multivariate normal), and we wish to assign each of the n items in the sample to the distribution

it most likely belongs to. Such an approach is related to classification analysis in Chapter 9. Along with partitioning in Section 15.4.1, this method has the property that points can be transferred from one cluster to another, but it requires more assumptions than partitioning.

We define the density of a mixture of g distributions as the weighted average

$$h(\mathbf{y}) = \sum_{i=1}^g \alpha_i f(\mathbf{y}, \boldsymbol{\mu}_i, \boldsymbol{\Sigma}_i), \quad (15.32)$$

where $0 \leq \alpha_i \leq 1$, $\sum_{i=1}^g \alpha_i = 1$, and $f(\mathbf{y}, \boldsymbol{\mu}_i, \boldsymbol{\Sigma}_i)$ is the multivariate normal $N_p(\boldsymbol{\mu}_i, \boldsymbol{\Sigma}_i)$ given in (4.2).

Clusters could be formed in two ways. The first approach is to assign an item with observation vector \mathbf{y} to the cluster C_i with largest value of the estimated posterior probability

$$\hat{P}(C_i|\mathbf{y}) = \frac{\hat{\alpha}_i f(\mathbf{y}, \hat{\boldsymbol{\mu}}_i, \hat{\boldsymbol{\Sigma}}_i)}{h(\mathbf{y})} \quad (15.33)$$

[see Rencher (1998, Sections 6.2.4 and 6.3.1)], where $\hat{\alpha}_i$, $\hat{\boldsymbol{\mu}}_i$, and $\hat{\boldsymbol{\Sigma}}_i$ are maximum likelihood estimates and $h(\mathbf{y})$ is given by (15.32) with estimates inserted for parameters. The posterior probability (15.33) is an estimate of the probability that an item with observation vector \mathbf{y} belongs to the i th cluster, C_i .

The second approach is to assign an item with observation vector \mathbf{y} to the cluster with largest value of

$$\ln \hat{\alpha}_i - \frac{1}{2} \ln |\hat{\boldsymbol{\Sigma}}_i| - \frac{1}{2} (\mathbf{y} - \hat{\boldsymbol{\mu}}_i)' \hat{\boldsymbol{\Sigma}}_i^{-1} (\mathbf{y} - \hat{\boldsymbol{\mu}}_i) \quad (15.34)$$

[see (9.15)]. For either of these approaches [based on (15.33) or (15.34)], we need the estimates $\hat{\alpha}_i$, $\hat{\boldsymbol{\mu}}_i$, and $\hat{\boldsymbol{\Sigma}}_i$. These estimates are obtained by maximizing the likelihood function $L = \prod_{j=1}^n h(\mathbf{y}_j)$, where $h(\mathbf{y}_j)$ is given by (15.32). The results are

$$\begin{aligned} \hat{\alpha}_i &= \frac{1}{n} \sum_{j=1}^n \hat{P}(C_i|\mathbf{y}_j), \quad i = 1, 2, \dots, g-1 \\ \hat{\boldsymbol{\mu}}_i &= \frac{1}{n\hat{\alpha}_i} \sum_{j=1}^n \mathbf{y}_j \hat{P}(C_i|\mathbf{y}_j), \quad i = 1, 2, \dots, g \\ \hat{\boldsymbol{\Sigma}}_i &= \frac{1}{n\hat{\alpha}_i} \sum_{j=1}^n (\mathbf{y}_j - \hat{\boldsymbol{\mu}}_i)(\mathbf{y}_j - \hat{\boldsymbol{\mu}}_i)' \hat{P}(C_i|\mathbf{y}_j), \quad i = 1, 2, \dots, g \end{aligned}$$

(Everitt 1993, p. 111), where $\hat{P}(C_i|\mathbf{y}_j)$ is given by (15.33). These three equations must be solved iteratively. For a given value of g , we can begin with initial estimates or guesses for the parameters and adjust them by iteration (this approach is related to the EM algorithm mentioned in Section 3.11). If g is not known, we can begin with $g = 1$, then successively try $g = 2, g = 3$, and so on, until the results are satisfactory.

The total number of parameters to be estimated is large. There are p parameters in each $\boldsymbol{\mu}_i$, $\frac{1}{2}p(p+1)$ unique parameters in each $\boldsymbol{\Sigma}_i$, and $g-1$ values of α_i (the

remaining $\hat{\alpha}_i$ is found by $\sum_{i=1}^g \hat{\alpha}_i = 1$, for a total of

$$\frac{1}{2}g(p+1)(p+2)-1 \quad (15.35)$$

parameters. If the sample size n is not sufficiently large to estimate all of these parameters, we could assume a common covariance matrix Σ , which reduces the number of parameters by $\frac{1}{2}(g-1)p(p+1)$.

The method of mixtures is invariant to full-rank linear transformations and is somewhat robust to the assumption of normality. The technique works better if the g densities are well separated or the sample sizes are large.

■ EXAMPLE 15.4.2a

To illustrate the clustering method based on mixtures of distributions, we use the protein consumption data of Table 15.7. Because of the small number of countries in the data set, there are not enough degrees of freedom to estimate a different covariance matrix for each cluster. Hence we assume equal covariance matrices and estimate a pooled covariance matrix $\hat{\Sigma}$. For illustration purposes, we choose $g = 5$ as in Example 15.4.1a.

We use the five clusters found by Ward's method to obtain initial estimates of α_i , μ_i , and Σ . Then the maximum likelihood equations are solved iteratively to find the following estimates.

$$\hat{\alpha}_1 = 0.2801, \hat{\alpha}_2 = 0.3200, \hat{\alpha}_3 = 0.1199, \hat{\alpha}_4 = 0.1600, \hat{\alpha}_5 = 0.1200$$

$$\hat{\mu}_1 = \begin{pmatrix} 8.64 \\ 6.87 \\ 2.39 \\ 14.04 \\ 2.54 \\ 39.27 \\ 3.74 \\ 4.21 \\ 4.66 \end{pmatrix}, \quad \hat{\mu}_2 = \begin{pmatrix} 13.21 \\ 10.64 \\ 3.99 \\ 21.16 \\ 3.38 \\ 24.70 \\ 4.65 \\ 2.06 \\ 4.18 \end{pmatrix}, \quad \hat{\mu}_3 = \begin{pmatrix} 6.13 \\ 5.77 \\ 1.43 \\ 9.63 \\ .93 \\ 54.07 \\ 2.40 \\ 4.90 \\ 3.40 \end{pmatrix},$$

$$\hat{\mu}_4 = \begin{pmatrix} 9.85 \\ 7.05 \\ 3.15 \\ 26.68 \\ 8.22 \\ 22.68 \\ 4.55 \\ 1.18 \\ 2.12 \end{pmatrix}, \quad \hat{\mu}_5 = \begin{pmatrix} 7.23 \\ 6.23 \\ 2.63 \\ 8.20 \\ 8.87 \\ 26.93 \\ 6.03 \\ 3.80 \\ 6.23 \end{pmatrix}$$

$$\hat{\Sigma} = \begin{pmatrix} 4.250 & -2.952 & .021 & -.047 & 1.001 & .929 & -.157 & .287 & .035 \\ -2.952 & 9.411 & .963 & .265 & -1.934 & -4.250 & 1.245 & -2.903 & -.319 \\ .021 & .963 & .471 & .552 & -.296 & -.699 & .301 & -.256 & -.008 \\ -.047 & .265 & .552 & 9.706 & -1.254 & .011 & 1.313 & -.801 & .032 \\ 1.001 & -1.934 & -.296 & -1.254 & 3.648 & .167 & .111 & .839 & 1.653 \\ .929 & -4.250 & -.699 & -.011 & .167 & 8.412 & -.777 & 1.708 & .137 \\ -.157 & 1.245 & .301 & 1.313 & .111 & -.777 & 1.634 & -.845 & -.208 \\ .287 & -2.903 & -.256 & -.801 & .839 & 1.708 & -.845 & 2.053 & .503 \\ .035 & -.319 & -.008 & .032 & 1.653 & .137 & -.208 & .503 & 1.808 \end{pmatrix}$$

Then assigning each country to the cluster for which it has the highest posterior probability of membership as in (15.33) yields the following clusters:

Cluster 1	Cluster 2	Cluster 3	Cluster 4	Cluster 5
Albania, Czech., Greece,	Austria, Belgium, France,	Bulgaria, Romania,	Denmark, Finland	E. Germany, Portugal
Hungary, Italy, Poland, USSR	Ireland, Netherlands, Switzerland, UK, W. Germany	Yugoslavia	Norway, Sweden	Spain

□

15.4.2b Density Estimation

In the method of *density estimation* or *density searching*, we seek regions of high density sometimes called *modes*. No assumption is made about the form of the density, as was done in Section 15.4.2a. We could estimate the density using a kernel function as in Section 9.7.2. Alternatively, we simply attempt to separate regions with a high concentration of points from regions with a low density.

To find regions of high density, we first choose a radius r and a value of k , the number of points in a k -nearest neighbor scheme. For each of the n points in the data, the number of points within a sphere of radius r is found. A point is called a *dense point* if at least k other points are contained in its sphere.

If a dense point is more than a distance r from all other dense points, it becomes the nucleus of a new cluster. If a dense point is within a distance r from at least one dense point that belongs to a cluster, it is added to the cluster. If the dense point is within a distance r of two or more clusters, these clusters are combined. Two clusters are also combined if the smallest distance between their dense points is less than the average of the $2k$ smallest distances between the original n points. The value of r can be gradually increased so that more points become dense. Another option is to begin with the specified value of r for each point and then gradually increase r until k observations are contained in its sphere.

■ EXAMPLE 15.4.2b

To illustrate the density estimation method, we use the protein data. For each pair of values of k and r , the value of r was allowed to increase if needed, as described above. For the following values of k and r , the number of clusters obtained are given.




Figure 15.21 First two discriminant functions for the clusters found in Example 15.4.2b.

k/r	1.6	1.7	1.8	1.9	2.0	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8
2	5	5	5	4	4	4	4	4	3	3	3	3	3
3	3	3	3	3	3	3	3	3	2	2	2	2	2
4	3	3	3	3	3	3	3	3	2	2	2	2	2

The five-cluster solution found by setting $r = 1.8$ and $k = 2$ is

Cluster 1	Cluster 2	Cluster 3	Cluster 4	Cluster 5
Austria, Belgium	Denmark	Albania,	Czech.,	Greece,
France, Ireland,	Finland,	Bulgaria,	E. Germany,	Italy,
Netherlands,	Norway,	Hungary,	Poland,	Portugal,
Switzerland,	Sweden	Romania,	USSR	Spain
UK, W. Germany		Yugoslavia		

This partitioning into five clusters is perhaps more reasonable than any of those found in Example 15.4.2a. The first two discriminant functions for these five clusters are plotted in Figure 15.21. \square

15.5 CHOOSING THE NUMBER OF CLUSTERS

In hierarchical clustering, we can select g clusters from the dendrogram by cutting across the branches at a given level of the distance measure used by one of the axes.




Figure 15.22 Cutting the dendrogram to choose the number of clusters.

This is illustrated in Figure 15.22, which is the dendrogram for the average linkage method (Section 15.3.4) applied to the city crime data in Table 15.1. Cutting the dendrogram at a level of 700 yields two clusters. Cutting it at 535 gives three clusters.

We wish to determine the value of g that provides the best fit to the data. One approach is to look for large changes in distances at which clusters are formed. For example, in Figure 15.22, the largest change in levels occurs in going from two clusters to a single cluster. The change in distance between the two-cluster solution and the three-cluster solution is 82 units-squared. The difference between the three-

cluster solution and the four-cluster solution is 73 units-squared, and the change between the four- and five-cluster solutions is only 26 units-squared. In this case we would choose two clusters.

A formalization of this procedure was proposed by Mojena (1977): Choose the number of groups given by the first stage in the dendrogram at which

$$\alpha_j > \bar{\alpha} + ks_\alpha, \quad j = 1, 2, \dots, n, \quad (15.36)$$

where $\alpha_1, \alpha_2, \dots, \alpha_n$ are the distance values for stages with $n, n-1, \dots, 1$ clusters, $\bar{\alpha}$ and s_α are the mean and standard deviation of the α 's, and k is a constant. Mojena (1977) suggested using a value of k in the range 2.75 to 3.5, but Milligan and Cooper (1985) recommended $k = 1.25$, based on a simulation study.

An index that can be used with either hierarchical or partitioning methods is

$$c = \frac{\text{tr}(\mathbf{H})/(g-1)}{\text{tr}(\mathbf{E})/(n-g)}. \quad (15.37)$$

The value of g that maximizes c is chosen. A related approach is to choose the value of g that minimizes

$$d = g^2 |\mathbf{E}|. \quad (15.38)$$

To compare two cluster solutions with g_1 and g_2 clusters where $g_2 > g_1$, we can use the test statistic

$$F = \frac{\text{tr}(\mathbf{E}_1) - \text{tr}(\mathbf{E}_2)}{\text{tr}(\mathbf{E}_2) \left[\left(\frac{n-g_1}{n-g_2} \right) \left(\frac{g_2}{g_1} \right)^{2/p} - 1 \right]}, \quad (15.39)$$

which has an approximate F -distribution with $p(g_2 - g_1)$ and $p(n - g_2)$ degrees of freedom [Beale (1969)]. The matrices \mathbf{E}_1 and \mathbf{E}_2 are within-cluster sums of squares and products matrices corresponding to g_1 and g_2 . The hypothesis is that the cluster solutions with g_1 and g_2 clusters are equally valid, and rejection implies that the cluster solution with g_2 clusters is better than the solution with g_1 clusters ($g_2 > g_1$). The F -approximation in (15.39) may not be sufficiently accurate to justify the use of p -values.

15.6 CLUSTER VALIDITY

To check the validity of a cluster solution, it may be possible to test the hypothesis that there are no clusters or groups in the population from which the sample at hand was taken. For example, the hypothesis could be that the population represents a single unimodal distribution such as the multivariate normal, or that the observations arose from a uniform distribution. Formal tests of hypotheses of this type concerning cluster validity are reviewed by Gordon (1999, Section 7.2).

A cross-validation approach can also be used to check the validity or stability of a clustering result. The data are randomly divided into two subsets, say A and B , and

the cluster analysis is carried out separately on each of A and B . The results should be similar if the clusters are valid. An alternative approach is the following (Gordon 1999, Section 7.1; Milligan 1996):

1. Use some clustering method to partition subset A into g clusters.
2. Partition subset B into g clusters in two ways:
 - (a) Assign each item in B to the cluster in A that it is closest to by using, for example, the distance to cluster centroids.
 - (b) Use the same clustering method on B that was used on A .
3. Compare the results of (a) and (b) in step 2.

15.7 CLUSTERING VARIABLES

In some cases, it may be of interest to cluster the p variables rather than the n observations. For a similarity measure between each pair of variables, we would usually use the correlation. Since most clustering methods use dissimilarities (such as distances), we need to convert the correlation matrix $\mathbf{R} = (r_{ij})$ to a dissimilarity matrix. This can conveniently be done by replacing each r_{ij} by $1 - |r_{ij}|$ or $1 - r_{ij}^2$. Using the resulting dissimilarity matrix, we can apply a clustering method such as a hierarchical technique to cluster the variables.

Clustering of variables can sometimes be done successfully with factor analysis, which groups the variables corresponding to each factor; see Sections 13.1 and 13.5.

■ EXAMPLE 15.7

We illustrate clustering of variables using the city crime data in Table 15.1. We first calculate the correlation matrix $\mathbf{R} = (r_{ij})$ and then transform \mathbf{R} to a dissimilarity matrix $\mathbf{D} = (1 - r_{ij}^2)$. The variables are then clustered using both average linkage and Ward's clustering methods, and the dendograms are given in Figures 15.23 and 15.24, respectively. Both clustering methods yield the same solution.

We next carry out a factor analysis of the data and compare the resulting groups of variables with the clusters obtained with the average linkage and Ward's methods. The factor loadings are estimated using the principal factor method (Section 13.3.2) with squared multiple correlations as initial communality estimates, and the loadings are then rotated with a varimax rotation (Section 13.5.2b). The rotated factor pattern is given in Table 15.12. The highest loading in each row is bolded. The first factor deals with crimes associated with the home. The second factor involves crimes that are violent in nature. The third factor consists of crimes of theft outside the home. Note that the three-cluster solutions found by both average linkage and Ward's methods are identical to the grouping of variables in the factor analysis solution, which is




Figure 15.23 Dendrogram for clustering the variables of Table 15.1 using average linkage (see Example 15.7).

(1) murder, rape, and assault, (2) robbery and auto theft, and (3) burglary and larceny. Since all three methods agree, we have some confidence in the validity of the solution. \square

PROBLEMS

- 15.1** Show that $d^2(\mathbf{x}, \mathbf{y}) = \sum_{j=1}^p (x_j - y_j)^2$ from (15.2) is equal to (15.5), $d^2(\mathbf{x}, \mathbf{y}) = (v_x - v_y)^2 + p(\bar{x} - \bar{y})^2 + 2v_x v_y (1 - r_{xy})$, where $v_x^2 = \sum_{j=1}^p (x_j - \bar{x})^2$, $\bar{x} = \sum_{j=1}^p x_j / p$, and r_{xy} is defined in (15.6).




Figure 15.24 Dendrogram for clustering the variables of Table 15.1 using Ward's method (see Example 15.7).

- 15.2 (a)** Show that $I_{AB} = n_A(\bar{y}_A - \bar{y}_{AB})'(\bar{y}_A - \bar{y}_{AB}) + n_B(\bar{y}_B - \bar{y}_{AB})'(\bar{y}_B - \bar{y}_{AB})$ as in (15.18).

- (b) Show that (15.18) is equal to (15.19); that is,

$$\begin{aligned} n_A(\bar{y}_A - \bar{y}_{AB})'(\bar{y}_A - \bar{y}_{AB}) + n_B(\bar{y}_B - \bar{y}_{AB})'(\bar{y}_B - \bar{y}_{AB}) \\ = \frac{n_A n_B}{n_A + n_B} (\bar{y}_A - \bar{y}_B)'(\bar{y}_A - \bar{y}_B). \end{aligned}$$

- 15.3** Using the hints provided below, show that the parameter values for (15.20) in Table 15.2 produce appropriate distances for the following cluster methods.

Table 15.12 Rotated Factor Loadings for City Crime Data

Variables	Factor 1	Factor 2	Factor 3
Murder	-.063	.734	.142
Rape	.504	.659	.160
Robbery	.133	.355	.726
Burglary	.764	.221	.181
Larceny	.847	-.014	.244
Auto theft	.240	.097	.584

- (a) Complete linkage. Use an approach analogous to that in Section 15.3.8 for the single linkage method.
- (b) Average linkage. Write (15.20) in terms of parameter values for average linkage in Table 15.2. Then use (15.9).
- (c) Centroid method. Show that

$$\begin{aligned} (\bar{\mathbf{y}}_C - \bar{\mathbf{y}}_{AB})'(\bar{\mathbf{y}}_C - \bar{\mathbf{y}}_{AB}) &= \frac{n_A}{n_A + n_B}(\bar{\mathbf{y}}_C - \bar{\mathbf{y}}_A)'(\bar{\mathbf{y}}_C - \bar{\mathbf{y}}_A) \\ &\quad + \frac{n_B}{n_A + n_B}(\bar{\mathbf{y}}_C - \bar{\mathbf{y}}_B)'(\bar{\mathbf{y}}_C - \bar{\mathbf{y}}_B) \\ &\quad - \frac{n_A n_B}{(n_A + n_B)^2}(\bar{\mathbf{y}}_A - \bar{\mathbf{y}}_B)'(\bar{\mathbf{y}}_A - \bar{\mathbf{y}}_B), \end{aligned} \quad (15.40)$$

where $\bar{\mathbf{y}}_{AB} = (n_A \bar{\mathbf{y}}_A + n_B \bar{\mathbf{y}}_B)/(n_A + n_B)$.

- (d) Median method. Use $n_A = n_B$ in (15.12) and (15.40) [see part (c)].
- (e) Ward's method. Show that

$$\begin{aligned} I_{C(AB)} &= \frac{n_A + n_C}{n_A + n_B + n_C} I_{AC} + \frac{n_B + n_C}{n_A + n_B + n_C} I_{BC} \\ &\quad - \frac{n_C}{n_A + n_B + n_C} I_{AB}, \end{aligned}$$

where I_{AB} is defined in (15.17).

- 15.4** Show that for all methods in Table 15.2 for which $\gamma = 0$, we have $D(C, AB) > (\alpha_A + \alpha_B + \beta)D(A, B)$ as in (15.26).
- 15.5** Verify the statement in the last paragraph of Section 15.4.1b, namely, that the first criterion in Section 15.4.1b is not invariant to nonsingular linear transformations $\mathbf{v}_{ij} = \mathbf{A}\mathbf{y}_{ij} + \mathbf{b}$, where \mathbf{A} is a $p \times p$ nonsingular matrix, and that the other two criteria are invariant to such transformations. Use the following approach:

- (a) Show that $\mathbf{H}_v = \mathbf{A}\mathbf{H}_y\mathbf{A}'$ and $\mathbf{E}_v = \mathbf{A}\mathbf{E}_y\mathbf{A}'$.

- (b) Show that minimizing $\text{tr}(\mathbf{E})$ is not invariant.
(c) Show that minimizing $|\mathbf{E}|$ is invariant.
(d) Show that maximizing $\text{tr}(\mathbf{E}^{-1}\mathbf{H})$ is invariant.
- 15.6** Verify the statement in Section 15.4.2a that in $\mu_i, i = 1, 2, \dots, g$; $\Sigma_i, i = 1, 2, \dots, g$; and $\alpha_i, i = 1, 2, \dots, g - 1$; the total number of parameters is given by $\frac{1}{2}g(p+1)(p+2) - 1$ as in (15.35).
- 15.7** Use the ramus bone date of Table 3.7. Carry out the following cluster methods and compare to the principal component plot in Figure 12.5.
- Find a two-cluster solution using the single linkage method.
 - Find a two-cluster solution using the average linkage method and compare to the result in (a). Which seems better?
 - Carry out a cluster analysis using the Ward's, complete linkage, centroid, and median methods.
 - Use the flexible beta method with $\beta = -0.25$, $\beta = -0.5$, and $\beta = -0.75$.
- 15.8** Use the hematology data of Table 4.2.
- Carry out a cluster analysis using the centroid method and find the distance between the centroids of the two-cluster solution.
 - Carry out a cluster analysis using the average linkage method. How many clusters are indicated in the dendrogram?
 - Using the two-cluster solution from part (b), label observations from one cluster as group 1 and the observations from the other cluster as group 2. Calculate and plot the discriminant function, as in Example 8.2. Do the two clusters overlap?
- 15.9** Use all of the variables of the Seishu data of Table 7.1.
- Find the three-cluster solution using the single linkage, complete linkage, average linkage, centroid, median, and Ward's methods. Which observation appears to be an outlier? Which cluster is the same in all six solutions?
 - Using the cluster found in part (a) to be common to all solutions as group 1 and the rest of the observations as group 2, calculate and plot the discriminant function, as in Problem 15.8(c). Do the two clusters overlap?
- 15.10** Use the first 20 observations of the temperature data of Table 7.2. Standardize the variables (columns) before doing the following:
- Carry out a k -means cluster analysis using as initial seeds the five observations that are mutually farthest apart. Plot the first two discriminant functions using the five clusters as groups.

- (b) Repeat part (a) using the first five observations as initial seeds.
- (c) Repeat part (a) using as initial seeds the centroids of the five-cluster solution found using Ward's method. Plot the dendrogram resulting from Ward's method.
- (d) Repeat part (c) using average linkage instead of Ward's method. Compare the results with those in part (c).
- (e) Plot the first and second principal components and the second and third components. Which cluster solutions found in parts (a)–(d) seem to agree most with the principal component plots?
- (f) Repeat parts (a) and (b) using three initial seeds instead of five. How do the cluster solutions compare?
- (g) Repeat part (c) using three initial seeds instead of five. How does the cluster solution compare to your answer in (f)?

15.11 Table 15.13 contains air pollution data from 41 US cities (Sokal and Rohlf 1981, p. 619). The variables are as follows:

y_1 = SO₂ content of air in micrograms per cubic meter

y_2 = Average annual temperature in °F

y_3 = Number of manufacturing enterprises employing 20 or more workers

y_4 = Population size (1970 census) in thousands

y_5 = Average annual wind speed in miles per hour

y_6 = Average annual precipitation in inches

y_7 = Average number of days with precipitation per year

Standardize each variable to mean 0 and standard deviation 1. Carry out a cluster analysis using the density estimation method with k equal to 2, 3, 4, 5, and values of r ranging from 0.2 to 2 by increments of 0.2 for each value of k . What is the maximum value of k that produces a two-cluster solution?

15.12 Table 15.14 gives the yields of winter wheat in each of the years 1970–1973 at twelve different sites in England (Hand et al. 1994, p. 31).

- (a) Carry out a cluster analysis using the density estimation method with $k = 2, 3, 4$, and $r = .2, .4, \dots, 2.0$.
- (b) Plot the first two discriminant functions from the three-cluster solution obtained with $k = 2$ and $r = 1$.
- (c) Plot the first two principal components and compare with the plot in part (b).
- (d) Repeat part (b) using a two-cluster solution obtained with $k = 3$ and $r = 1$. Which two clusters of the three-cluster solution found in part (b) merged into one cluster?

Table 15.13 Air Pollution Levels in US Cities

Cities	y_1	y_2	y_3	y_4	y_5	y_6	y_7
Phoenix	10	70.3	213	582	6.0	7.05	36
Little Rock	13	61.0	91	132	8.2	48.52	100
San Francisco	12	56.7	453	716	8.7	20.66	67
Denver	17	51.9	454	515	9.0	12.95	86
Hartford	56	49.1	412	158	9.0	43.37	127
Wilmington	36	54.0	80	80	9.0	40.25	114
Washington	29	57.3	434	757	9.3	38.89	111
Jacksonville	14	68.4	136	529	8.8	54.47	116
Miami	10	75.5	207	335	9.0	59.80	128
Atlanta	24	61.5	368	497	9.1	48.34	115
Chicago	110	50.6	3344	3369	10.4	34.44	122
Indianapolis	28	52.3	361	746	9.7	38.74	121
Des Moines	17	49.0	104	201	11.2	30.85	103
Wichita	8	56.6	125	277	12.7	30.58	82
Louisville	30	55.6	291	593	8.3	43.11	123
New Orleans	9	68.3	204	361	8.4	56.77	113
Baltimore	47	55.0	625	905	9.6	41.31	111
Detroit	35	49.9	1064	1513	10.1	30.96	129
Minneapolis-St. Paul	29	43.5	699	744	10.6	25.94	137
Kansas City	14	54.5	381	507	10.0	37.00	99
St. Louis	56	55.9	775	622	9.5	35.89	105
Omaha	14	51.5	181	347	10.9	30.18	98
Albuquerque	11	56.8	46	244	8.9	7.77	58
Albany	46	47.6	44	116	8.8	33.36	135
Buffalo	11	47.1	391	463	12.4	36.11	166
Cincinnati	23	54.0	462	453	7.1	39.04	132
Cleveland	65	49.7	1007	751	10.9	34.99	155
Columbus	26	51.5	266	540	8.6	37.01	134
Philadelphia	69	54.6	1692	1950	9.6	39.93	115
Pittsburgh	61	50.4	347	520	9.4	36.22	147
Providence	94	50.0	343	179	10.6	42.75	125
Memphis	10	61.6	337	624	9.2	49.10	105
Nashville	18	59.4	275	448	7.9	46.00	119
Dallas	9	66.2	641	844	10.9	35.94	78
Houston	10	68.9	721	1233	10.8	48.19	103
Salt Lake City	28	51.0	137	176	8.7	15.17	89
Norfolk	31	59.3	96	308	10.6	44.68	116
Richmond	26	57.8	197	299	7.6	42.59	115
Seattle	29	51.1	379	531	9.4	38.79	164
Charleston	31	55.2	35	71	6.5	40.75	148
Milwaukee	16	45.7	569	717	11.8	29.07	123

Table 15.14 Yields of Winter Wheat (kg per unit area)

Site	Year			
	1970	1971	1972	1973
Cambridge	46.81	39.40	55.64	32.61
Cockle Park	46.49	34.07	45.06	41.02
Harpers Adams	44.03	42.03	40.32	50.23
Headley Hall	52.24	36.19	47.03	34.56
Morley	36.55	43.06	38.07	43.17
Myerscough	34.88	49.72	40.86	50.08
Rosemaund	56.14	47.67	43.48	38.99
Seale-Hayne	45.67	27.30	45.48	50.32
Sparsholt	42.97	46.87	38.78	47.49
Sutton Bonington	54.44	49.34	24.48	46.94
Terrington	54.95	52.05	50.91	39.13
Wye	48.94	48.63	31.69	59.72

CHAPTER 16

GRAPHICAL PROCEDURES

In Sections 16.1, 16.2, and 16.3, we consider three graphical techniques: multidimensional scaling, correspondence analysis, and biplots. These methods are designed to reduce dimensionality and portray relationships among observations or variables.

16.1 MULTIDIMENSIONAL SCALING

16.1.1 Introduction

In the dimension reduction technique called *multidimensional scaling*, we begin with the distances δ_{ij} between each pair of items. We wish to represent the n items in a low-dimensional coordinate system, in which the distances d_{ij} between items closely match the original distances δ_{ij} , that is,

$$d_{ij} \cong \delta_{ij} \text{ for all } i, j.$$

The final distances d_{ij} are usually Euclidean. The original distances δ_{ij} may be actual measured distances between observations \mathbf{y}_i and \mathbf{y}_j in p dimensions, such as

$$\delta_{ij} = [(\mathbf{y}_i - \mathbf{y}_j)'(\mathbf{y}_i - \mathbf{y}_j)]^{1/2}. \quad (16.1)$$

On the other hand, the distances δ_{ij} may be only a proximity or similarity based on human judgment, for example, the perceived degree of similarity between all pairs of brands of a certain type of appliance (for a discussion of similarities and dissimilarities, see Section 15.2). The goal of multidimensional scaling is a plot that exhibits information about how the items relate to each other or provides some other meaningful interpretation of the data. For example, the aim may be *seriation* or ranking; if the points lie close to a curve in two dimensions, then the ordering of points along the curve is used to rank the points.

If the observation vectors $\mathbf{y}_i, i = 1, 2, \dots, n$, are available and we calculate distances using (16.1) or a similar measure, or if the original \mathbf{y}_i 's are not available but we have actual distances between items, then the process of reduction to a lower-dimensional geometric representation is called *metric multidimensional scaling*. If the original distances are only similarities based on judgment, the process is called *nonmetric multidimensional scaling*, and the final spatial representation preserves only the rank order among the similarities. We consider metric scaling in Section 16.1.2 and nonmetric scaling in Section 16.1.3. For useful discussions of various aspects of multidimensional scaling, see Davidson (1983); Gordon (1999, Sections 6.2 and 6.3); Kruskal and Wish (1978); Mardia, Kent, and Bibby (1979, Chapter 14); Seber (1984, Section 5.5); Young (1987); Jobson (1992, Section 10.3); Shepard, Romney, and Nerlove (1972); and Romney, Shepard, and Nerlove (1972).

16.1.2 Metric Multidimensional Scaling

In this section, we consider *metric multidimensional scaling*, which is also known as the *classical solution* and as *principal coordinate analysis*. We begin with an $n \times n$ distance matrix $\mathbf{D} = (\delta_{ij})$. Our goal is to find n points in k dimensions such that the interpoint distances d_{ij} in the k dimensions are approximately equal to the values of δ_{ij} in \mathbf{D} . Typically, we use $k = 2$ for plotting purposes, but $k = 1$ or 3 may also be useful.

The points are found as follows:

1. Construct the $n \times n$ matrix $\mathbf{A} = (a_{ij}) = (-\frac{1}{2} \delta_{ij}^2)$, where δ_{ij} is the ij th element of \mathbf{D} .
2. Construct the $n \times n$ matrix $\mathbf{B} = (b_{ij})$, with elements $b_{ij} = a_{ij} - \bar{a}_{i\cdot} - \bar{a}_{\cdot j} + \bar{a}_{\cdot\cdot}$, where $\bar{a}_{i\cdot} = \sum_{j=1}^n a_{ij}/n$, $\bar{a}_{\cdot j} = \sum_{i=1}^n a_{ij}/n$, $\bar{a}_{\cdot\cdot} = \sum_{ij} a_{ij}/n^2$. The matrix \mathbf{B} can be written as

$$\mathbf{B} = \left(\mathbf{I} - \frac{1}{n} \mathbf{J} \right) \mathbf{A} \left(\mathbf{I} - \frac{1}{n} \mathbf{J} \right). \quad (16.2)$$

It can be shown that there exists a q -dimensional configuration $\mathbf{z}_1, \mathbf{z}_2, \dots, \mathbf{z}_n$ with interpoint distances $d_{ij} = (\mathbf{z}_i - \mathbf{z}_j)'(\mathbf{z}_i - \mathbf{z}_j)$ such that $d_{ij} = \delta_{ij}$ if

and only if \mathbf{B} is positive semidefinite of rank q (Schoenberg 1935; Young and Householder 1938; Gower 1966; Seber 1984, p. 236).

3. Since \mathbf{B} is a symmetric matrix, we can use the spectral decomposition in (2.109) to write \mathbf{B} in the form

$$\mathbf{B} = \mathbf{V}\Lambda\mathbf{V}', \quad (16.3)$$

where \mathbf{V} is the matrix of eigenvectors of \mathbf{B} and Λ is the diagonal matrix of eigenvalues of \mathbf{B} . If \mathbf{B} is positive semidefinite of rank q , there are q positive eigenvalues, and the remaining $n - q$ eigenvalues are zero. If $\Lambda_1 = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_q)$ contains the positive eigenvalues and $\mathbf{V}_1 = (\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_q)$ contains the corresponding eigenvectors, then we can express (16.3) in the form

$$\begin{aligned} \mathbf{B} &= \mathbf{V}_1\Lambda_1\mathbf{V}'_1 \\ &= \mathbf{V}_1\Lambda_1^{1/2}\Lambda_1^{1/2}\mathbf{V}'_1 \\ &= \mathbf{Z}\mathbf{Z}', \end{aligned}$$

where

$$\mathbf{Z} = \mathbf{V}_1\Lambda_1^{1/2} = (\sqrt{\lambda_1}\mathbf{v}_1, \sqrt{\lambda_2}\mathbf{v}_2, \dots, \sqrt{\lambda_q}\mathbf{v}_q) = \begin{pmatrix} \mathbf{z}'_1 \\ \mathbf{z}'_2 \\ \vdots \\ \mathbf{z}'_n \end{pmatrix}. \quad (16.4)$$

4. The rows $\mathbf{z}'_1, \mathbf{z}'_2, \dots, \mathbf{z}'_n$ of \mathbf{Z} in (16.4) are the points whose interpoint distances $d_{ij} = (\mathbf{z}_i - \mathbf{z}_j)'(\mathbf{z}_i - \mathbf{z}_j)$ match the δ_{ij} 's in the original distance matrix \mathbf{D} , as noted following (16.2).
5. Since q in (16.4) will typically be too large to be of practical interest and we would prefer a smaller dimension k for plotting, we can use the first k eigenvalues and corresponding eigenvectors in (16.4) to obtain n points whose interpoint distances d_{ij} are approximately equal to the corresponding δ_{ij} 's.
6. If \mathbf{B} is not positive semidefinite, but its first k eigenvalues are positive and relatively large, then these eigenvalues and associated eigenvectors may be used in (16.4) to construct points that give reasonably good approximations to the δ_{ij} 's.

Note that the method used to obtain \mathbf{Z} from \mathbf{B} closely resembles principal component analysis. Note also that the solution \mathbf{Z} in (16.4) is not unique, since a shift in origin or a rotation will not change the distances d_{ij} . For example, if \mathbf{C} is a $q \times q$ orthogonal matrix producing a rotation [see (2.101)], then

$$\begin{aligned} (\mathbf{C}\mathbf{z}_i - \mathbf{C}\mathbf{z}_j)'(\mathbf{C}\mathbf{z}_i - \mathbf{C}\mathbf{z}_j) &= (\mathbf{z}_i - \mathbf{z}_j)' \mathbf{C}' \mathbf{C} (\mathbf{z}_i - \mathbf{z}_j) \\ &= (\mathbf{z}_i - \mathbf{z}_j)'(\mathbf{z}_i - \mathbf{z}_j) \quad [\text{see (2.103)}]. \end{aligned}$$

Thus the rotated points $\mathbf{C}\mathbf{z}_i$ have the same interpoint distances d_{ij} .

■ EXAMPLE 16.1.2(a)

To illustrate the first four steps of the above algorithm for metric multidimensional scaling, consider the 5×5 distance matrix

$$\mathbf{D} = (\delta_{ij}) = \begin{pmatrix} 0 & 2\sqrt{2} & 2\sqrt{2} & 2\sqrt{2} & 2\sqrt{2} \\ 2\sqrt{2} & 0 & 4 & 4\sqrt{2} & 4 \\ 2\sqrt{2} & 4 & 0 & 4 & 4\sqrt{2} \\ 2\sqrt{2} & 4\sqrt{2} & 4 & 0 & 4 \\ 2\sqrt{2} & 4 & 4\sqrt{2} & 4 & 0 \end{pmatrix}.$$

The matrix $\mathbf{A} = (-\frac{1}{2}\delta_{ij}^2)$ in step 1 is given by

$$\mathbf{A} = -\begin{pmatrix} 0 & 4 & 4 & 4 & 4 \\ 4 & 0 & 8 & 16 & 8 \\ 4 & 8 & 0 & 8 & 16 \\ 4 & 16 & 8 & 0 & 8 \\ 4 & 8 & 16 & 8 & 0 \end{pmatrix}.$$

For the means, we have $\bar{a}_{1\cdot} = \bar{a}_{\cdot 1} = -16/5$, $\bar{a}_{i\cdot} = \bar{a}_{\cdot i} = -36/5$, $i = 2, \dots, 5$, $\bar{a}_{..} = -32/5$. With $n = 5$, the matrix \mathbf{B} in step 2 is given by

$$\mathbf{B} = \left(\mathbf{I} - \frac{1}{5}\mathbf{J}\right)\mathbf{A}\left(\mathbf{I} - \frac{1}{5}\mathbf{J}\right) = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 8 & 0 & -8 & 0 \\ 0 & 0 & 8 & 0 & -8 \\ 0 & -8 & 0 & 8 & 0 \\ 0 & 0 & -8 & 0 & 8 \end{pmatrix}.$$

The rank of \mathbf{B} is clearly 2. For step 3, the (nonzero) eigenvalues and corresponding eigenvectors of \mathbf{B} are given by $\lambda_1 = 16$, $\lambda_2 = 16$,

$$\mathbf{v}_1 = \begin{pmatrix} 0 \\ \frac{1}{2}\sqrt{2} \\ 0 \\ -\frac{1}{2}\sqrt{2} \\ 0 \end{pmatrix}, \quad \mathbf{v}_2 = \begin{pmatrix} 0 \\ 0 \\ \frac{1}{2}\sqrt{2} \\ 0 \\ -\frac{1}{2}\sqrt{2} \end{pmatrix}.$$

Then for step 3 we have by (16.4)

$$\mathbf{Z} = (\sqrt{\lambda_1}\mathbf{v}_1, \sqrt{\lambda_2}\mathbf{v}_2) = \begin{pmatrix} 0 & 0 \\ 2\sqrt{2} & 0 \\ 0 & 2\sqrt{2} \\ -2\sqrt{2} & 0 \\ 0 & -2\sqrt{2} \end{pmatrix}.$$

It can be shown (step 4) that the distance matrix for these five points is \mathbf{D} . The five points constitute a square with each side of length 4 and a center point at the origin. The five points (rows of \mathbf{Z}) are plotted in Figure 16.1. \square




Figure 16.1 Plot of the five points found in Example 16.1.2(a).

■ EXAMPLE 16.1.2(b)

For another example of metric multidimensional scaling, we use airline distances between 10 US cities, as given in Table 16.1 (Kruskal and Wish 1978, pp. 7–9).

Table 16.1 Airline Distances Between Ten US Cities

City	1	2	3	4	5	6	7	8	9	10
1	0	587	1212	701	1936	604	748	2139	2182	543
2	587	0	920	940	1745	1188	713	1858	1737	597
3	1212	920	0	879	831	1726	1631	949	1021	1494
4	701	940	879	0	1374	968	1420	1645	1891	1220
5	1936	1745	831	1374	0	2339	2451	347	959	2300
6	604	1188	1726	968	2339	0	1092	2594	2734	923
7	748	713	1631	1420	2451	1092	0	2571	2408	205
8	2139	1858	949	1645	347	2594	2571	0	678	2442
9	2182	1737	1021	1891	959	2734	2408	678	0	2329
10	543	597	1494	1220	2300	923	205	2442	2329	0

Cities: (1) Atlanta, (2) Chicago, (3) Denver, (4) Houston, (5) Los Angeles, (6) Miami, (7) New York, (8) San Francisco, (9) Seattle, (10) Washington, DC.




Figure 16.2 Plot of the points found in Example 16.1.2(b).

The points given by metric multidimensional scaling are plotted in Figure 16.2. Note that in the plot, dimensions 1 and 2 roughly correspond to easting and northing, but the sign for each of these dimensions is arbitrary. That is, the eigenvectors \mathbf{v}_i in (16.4) are normalized but are subject to multiplication by -1 . \square

16.1.3 Nonmetric Multidimensional Scaling

Suppose the $m = n(n - 1)/2$ dissimilarities δ_{ij} cannot be measured as in (16.1) but can be ranked in order,

$$\delta_{r_1 s_1} < \delta_{r_2 s_2} < \cdots < \delta_{r_m s_m}, \quad (16.5)$$

where $r_1 s_1$ indicates the pair of items with the smallest dissimilarity and $r_m s_m$ represents the pair with greatest dissimilarity. In nonmetric multidimensional scaling, we seek a low-dimensional representation of the points such that the rankings of the distances

$$d_{r_1 s_1} < d_{r_2 s_2} < \cdots < d_{r_m s_m} \quad (16.6)$$

match exactly the ordering of dissimilarities in (16.5). Thus, while metric scaling uses the magnitudes of the δ_{ij} 's, nonmetric scaling is based only on the rank order of the δ_{ij} 's.

For a given set of points with distances d_{ij} , a plot of d_{ij} versus δ_{ij} may not be monotonic; that is, the ordering in (16.6) may not match exactly the ordering in (16.5). Suitable \hat{d}_{ij} 's can be estimated by *monotonic regression*, in which we seek

values of \hat{d}_{ij} to minimize the scaled sum of squared differences

$$S^2 = \frac{\sum_{i < j} (d_{ij} - \hat{d}_{ij})^2}{\sum_{i < j} d_{ij}^2} \quad (16.7)$$

subject to the constraint

$$\hat{d}_{r_1 s_1} \leq \hat{d}_{r_2 s_2} \leq \cdots \leq \hat{d}_{r_m s_m},$$

where $r_1 s_1, r_2 s_2, \dots, r_m s_m$ are defined as in (16.5) and (16.6) (Kruskal 1964a, 1964b). The minimum value of S^2 for a given dimension, k , is called the STRESS. Note that the \hat{d}_{ij} 's are *not* distances. They are merely numbers used as a reference to assess the monotonicity of the d_{ij} 's. The \hat{d}_{ij} 's are sometimes called *disparities*.

The minimum value of the STRESS over all possible configurations of points can be found using the following algorithm.

1. Rank the $m = n(n - 1)/2$ distances or dissimilarities δ_{ij} as in (16.5).
2. Choose a value of k and an initial configuration of points in k dimensions. The initial configuration could be n points chosen at random from a uniform or normal distribution, n evenly spaced points in k -dimensional space, or the metric solution obtained by treating the ordinal measurements as continuous and using the algorithm in Section 16.1.2.
3. For the initial configuration of points, find the interitem distances d_{ij} . Find the corresponding \hat{d}_{ij} 's by monotonic regression as defined above using (16.7).
4. Choose a new configuration of points whose distances d_{ij} minimize S^2 in (16.7) with respect to the \hat{d}_{ij} 's found in step 3. One approach is to use an iterative gradient technique such as the method of steepest descent or the Newton–Raphson method.
5. Using monotonic regression, find new \hat{d}_{ij} 's for the d_{ij} 's found in step 4. This gives a new value of STRESS.
6. Repeat steps 4 and 5 until STRESS converges to a minimum over all possible k -dimensional configurations of points.
7. Using the preceding six steps, calculate the minimum STRESS for values of k starting at $k = 1$ and plot these. As k increases, the curve will decrease, with occasional exceptions due to round off or numerical anomalies in the search procedure for minimum STRESS. We look for a discernible bend in the plot, following which the curve is low and relatively flat. An ideal plot is shown in Figure 16.3. The curve levels off after $k = 2$, which is convenient for plotting the resulting n points in 2 dimensions.

There is a possibility that the minimum value of STRESS found by the above seven steps for a given value of k may be a local minimum rather than the global




Figure 16.3 Ideal plot of minimum STRESS versus k .

minimum. Such an anomaly may show up in the plot of minimum STRESS versus k . The possibility of a local minimum can be checked by repeating the procedure, starting with a different initial configuration.

As was the case with metric scaling, the final configuration of points from a non-metric scaling is invariant to a rotation of axes.

■ EXAMPLE 16.1.3

The voting records for fifteen congressmen from New Jersey on nineteen environmental bills are given in Table 16.2 in the form of a dissimilarity matrix (Hand et al. 1994, p. 235). The congressmen are identified by party: R_1 for Republican 1, D_2 for Democrat 2, etc. Each entry shows how often the indicated congressman voted differently from each of the other fourteen.


Using an initial configuration of points from a multivariate normal distribution with mean vector $\mu = \mathbf{0}$ and $\Sigma = \mathbf{I}$, we find an “optimal” configuration of points for each of $k = 1, 2, \dots, 5$. A plot of the STRESS is given in Figure 16.4.

From the plot of STRESS vs. number of dimensions, we see that either two or three dimensions will be sufficient. For plotting purposes, we choose two dimensions, which has a STRESS value of 0.113. The plot of the first two dimensions is given in Figure 16.5. It is apparent that the plot separates the Republicans from the Democrats except for Republican 6, who voted much the same as the Democrats.

Table 16.2 Dissimilarity Matrix for Voting Records of 15 Congressmen

	R_1	R_2	D_1	D_2	R_3	R_4	R_5	D_3	D_4	D_5	D_6	R_6	R_7	R_8	D_7
R_1	0	8	15	15	10	9	7	15	16	14	15	16	7	11	13
R_2	8	0	17	12	13	13	12	16	17	15	16	17	13	12	16
D_1	15	17	0	9	16	12	15	5	5	6	5	4	11	10	7
D_2	15	12	9	0	14	12	13	10	8	8	8	6	15	10	7
R_3	10	13	16	14	0	8	9	13	14	12	12	12	10	11	11
R_4	9	13	12	12	8	0	7	12	11	10	9	10	6	6	10
R_5	7	12	15	13	9	7	0	17	16	15	14	15	10	11	13
D_3	15	16	5	10	13	12	17	0	4	5	5	3	12	7	6
D_4	16	17	5	8	14	11	16	4	0	3	2	1	13	7	5
D_5	14	15	6	8	12	10	15	5	3	0	1	2	11	4	6
D_6	15	16	5	8	12	9	14	5	2	1	0	1	12	5	5
R_6	16	17	4	6	12	10	15	3	1	2	1	0	12	6	4
R_7	7	13	11	15	10	6	10	12	13	11	12	12	0	9	13
R_8	11	12	10	10	11	6	11	7	7	4	5	6	9	0	9
D_7	13	16	7	7	11	10	13	6	5	6	5	4	13	9	0

We now use a different initial configuration of points drawn from a uniform distribution. The resulting plot is given in Figure 16.6. The approximate orientation of the points in the two-dimensional space is similar, with the exception of the locations of R_2 and R_5 .

**Figure 16.4** Plot of STRESS for each value of k for the voting data in Table 16.2.




Figure 16.5 Plot of points found using initial points from a multivariate normal distribution.




Figure 16.6 Plot of points found using initial points from a uniform distribution.




Figure 16.7 Plot of points found using initial points from a metric solution.

We next use a third initial configuration of points resulting from the metric solution as described in Section 16.1.2. The resulting plot is given in Figure 16.7. All three plots are very similar, indicating a good fit. \square

16.2 CORRESPONDENCE ANALYSIS

16.2.1 Introduction

Correspondence analysis is a graphical technique for representing the information in a two-way contingency table, which contains the counts (frequencies) of items for a cross-classification of two categorical variables. With correspondence analysis, we construct a plot that shows the interaction of the two categorical variables along with the relationship of the rows to each other and the columns to each other. In Sections 16.2.2–16.2.4, we consider correspondence analysis for ordinary two-way contingency tables. In Section 16.2.5 we consider multiple correspondence analysis for three-way and higher-order contingency tables. Useful treatments of correspondence analysis have been given by Greenacre (1984), Jobson (1992, Section 9.4), Khattree and Naik (1999, Chapter 7), Gower and Hand (1996, Chapters 4 and 9), and Benzécri (1992).

To test for significance of association of the two categorical variables in a contingency table, we can use a chi-square test or a log-linear model, both of which represent an asymptotic approach. Since correspondence analysis is associated with the chi-square approach, we will review it in Section 16.2.3. If a contingency table

Table 16.3 Contingency Table with a Rows and b Columns

		Columns				
		1	2	...	b	Row Total
Rows	1	n_{11}	n_{12}	...	n_{1b}	$n_{1..}$
	2	n_{21}	n_{22}	...	n_{2b}	$n_{2..}$
	:	:	:		:	:
	a	n_{a1}	n_{a2}	...	n_{ab}	$n_{a..}$
Column Total		$n_{..1}$	$n_{..2}$...	$n_{..b}$	n

has some cell frequencies that are small or zero, the chi-square approximation is not very satisfactory. In this case, some categories can be combined to increase the cell frequencies. Correspondence analysis may be useful in identifying the categories that are similar, which we may thereby wish to combine.

In correspondence analysis, we plot a point for each row and a point for each column of the contingency table. These points are in effect projections of the rows and columns of the contingency table onto a two-dimensional Euclidean space. The goal is to preserve as far as possible the relationship of the rows (or columns) to each other in a two-dimensional space. If two row points are close together, the profiles of the two rows (across the columns) are similar. Likewise, two column points that are close together represent columns with similar profiles across the rows (see Section 16.2.2 for a definition of profiles). If a row point is close to a column point, this combination of categories of the two variables occurs more frequently than would occur by chance if the two variables were independent. Another output of a correspondence analysis is the *inertia* or amount of information in each of the two dimensions in the plot (see Section 16.2.4).

16.2.2 Row and Column Profiles

A contingency table with a rows and b columns is represented in Table 16.3. The entries n_{ij} are the counts or frequencies for every two-way combination of row and column (every cell). The marginal totals are shown using the familiar “dot” notation: $n_{i..} = \sum_{j=1}^b n_{ij}$ and $n_{..j} = \sum_{i=1}^a n_{ij}$. The overall total frequency is denoted by n instead of $n_{....}$ for simplicity: $n = \sum_{ij} n_{ij}$.

The frequencies n_{ij} in a contingency table can be converted to relative frequencies p_{ij} by dividing by n : $p_{ij} = n_{ij}/n$. The matrix of relative frequencies is called the *correspondence matrix* and is denoted by \mathbf{P} :

$$\mathbf{P} = (p_{ij}) = (n_{ij}/n). \quad (16.8)$$

In Table 16.4 we show the contingency table in Table 16.3 converted to a correspondence matrix.

Table 16.4 Correspondence Matrix of Relative Frequencies

		Columns				
		1	2	...	b	Row Total
Rows	1	p_{11}	p_{12}	...	p_{1b}	$p_{1.}$
	2	p_{21}	p_{22}	...	p_{2b}	$p_{2.}$
	:	:	:		:	:
	a	p_{a1}	p_{a2}	...	p_{ab}	$p_{a.}$
Column Total		$p_{.1}$	$p_{.2}$...	$p_{.b}$	1

The last column of Table 16.4 contains the row sums $p_{i.} = \sum_{j=1}^b p_{ij}$. This column vector is denoted by \mathbf{r} and can be obtained as

$$\mathbf{r} = \mathbf{P}\mathbf{j} = (p_{1.}, p_{2.}, \dots, p_{a.})' = (n_{1.}/n, n_{2.}/n, \dots, n_{a.}/n)', \quad (16.9)$$

where \mathbf{j} is an $a \times 1$ vector of 1's. Similarly, the last row of Table 16.4 contains the column sums $p_{.j} = \sum_{i=1}^a p_{ij}$. This row vector is denoted by \mathbf{c}' and can be obtained as

$$\mathbf{c}' = \mathbf{j}'\mathbf{P} = (p_{.1}, p_{.2}, \dots, p_{.b}) = (n_{.1}/n, n_{.2}/n, \dots, n_{.b}/n), \quad (16.10)$$

where \mathbf{j}' is a $1 \times b$ vector of 1's. The elements of the vectors \mathbf{r} and \mathbf{c} are sometimes referred to as *row* and *column masses*. The correspondence matrix and marginal totals in Table 16.4 can be expressed as

$$\begin{pmatrix} \mathbf{P} & \mathbf{r} \\ \mathbf{c}' & 1 \end{pmatrix} = \left(\begin{array}{cccc|c} p_{11} & p_{12} & \cdots & p_{1b} & p_{1.} \\ p_{21} & p_{22} & \cdots & p_{2b} & p_{2.} \\ \vdots & \vdots & & \vdots & \vdots \\ p_{a1} & p_{a2} & \cdots & p_{ab} & p_{a.} \\ \hline p_{.1} & p_{.2} & \cdots & p_{.b} & 1 \end{array} \right). \quad (16.11)$$

We now convert each row and column of \mathbf{P} to a profile. The *i*th *row profile* \mathbf{r}'_i , $i = 1, 2, \dots, a$, is defined by dividing the *i*th row of either Table 16.3 or Table 16.4 by its marginal total:

$$\mathbf{r}'_i = \left(\frac{p_{i1}}{p_{i.}}, \frac{p_{i2}}{p_{i.}}, \dots, \frac{p_{ib}}{p_{i.}} \right) = \left(\frac{n_{i1}}{n_{i.}}, \frac{n_{i2}}{n_{i.}}, \dots, \frac{n_{ib}}{n_{i.}} \right). \quad (16.12)$$

The elements in each \mathbf{r}'_i are relative frequencies, and therefore they sum to 1:

$$\mathbf{r}'_i \mathbf{j} = \sum_{j=1}^b \frac{n_{ij}}{n_{i.}} = \frac{n_{i.}}{n_{i.}} = 1. \quad (16.13)$$

By defining

$$\mathbf{D}_r = \text{diag}(\mathbf{r}) = \begin{pmatrix} p_{1.} & 0 & \cdots & 0 \\ 0 & p_{2.} & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & p_{a.} \end{pmatrix} \quad (16.14)$$

and using (2.55), the matrix \mathbf{R} of row profiles can be expressed as

$$\mathbf{R} = \mathbf{D}_r^{-1} \mathbf{P} = \begin{pmatrix} \mathbf{r}'_1 \\ \mathbf{r}'_2 \\ \vdots \\ \mathbf{r}'_a \end{pmatrix} = \begin{pmatrix} \frac{p_{11}}{p_{1.}} & \frac{p_{12}}{p_{1.}} & \cdots & \frac{p_{1b}}{p_{1.}} \\ \frac{p_{21}}{p_{2.}} & \frac{p_{22}}{p_{2.}} & \cdots & \frac{p_{2b}}{p_{2.}} \\ \vdots & \vdots & & \vdots \\ \frac{p_{a1}}{p_{a.}} & \frac{p_{a2}}{p_{a.}} & \cdots & \frac{p_{ab}}{p_{a.}} \end{pmatrix}. \quad (16.15)$$

Similarly, the j th *column profile* \mathbf{c}_j , $j = 1, 2, \dots, b$, is defined by dividing the j th column of either Table 16.3 or Table 16.4 by its marginal total:

$$\mathbf{c}_j = \left(\frac{p_{1j}}{p_{.j}}, \frac{p_{2j}}{p_{.j}}, \dots, \frac{p_{aj}}{p_{.j}} \right)' = \left(\frac{n_{1j}}{n_{.j}}, \frac{n_{2j}}{n_{.j}}, \dots, \frac{n_{aj}}{n_{.j}} \right)'. \quad (16.16)$$

The elements in each \mathbf{c}_j are relative frequencies, and therefore they sum to 1:

$$\mathbf{j}' \mathbf{c}_j = \sum_{i=1}^a \frac{n_{ij}}{n_{.j}} = \frac{n_{.j}}{n_{.j}} = 1. \quad (16.17)$$

By defining

$$\mathbf{D}_c = \text{diag}(\mathbf{c}) = \begin{pmatrix} p_{.1} & 0 & \cdots & 0 \\ 0 & p_{.2} & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & p_{.b} \end{pmatrix} \quad (16.18)$$

and using (2.56), the matrix \mathbf{C} of column profiles can be expressed as

$$\mathbf{C} = \mathbf{P} \mathbf{D}_c^{-1} = (\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_b) = \begin{pmatrix} \frac{p_{11}}{p_{.1}} & \frac{p_{12}}{p_{.1}} & \cdots & \frac{p_{1a}}{p_{.1}} \\ \frac{p_{21}}{p_{.1}} & \frac{p_{22}}{p_{.1}} & \cdots & \frac{p_{2a}}{p_{.1}} \\ \vdots & \vdots & & \vdots \\ \frac{p_{a1}}{p_{.1}} & \frac{p_{a2}}{p_{.1}} & \cdots & \frac{p_{ab}}{p_{.1}} \end{pmatrix}. \quad (16.19)$$

The vector \mathbf{r} is defined in (16.9) as the column vector of row sums of \mathbf{P} . It can also be expressed as a weighted average of the column profiles:

$$\mathbf{r} = \sum_{j=1}^b p_{.j} \mathbf{c}_j. \quad (16.20)$$

Table 16.5 Piston Ring Failures

Compressor	Leg			Row Total
	North	Center	South	
1	17	17	12	46
2	11	9	13	33
3	11	8	19	38
4	14	7	28	49
Total	53	41	72	166

Similarly, \mathbf{c}' in (16.10) is the row vector of column sums of \mathbf{P} and can be expressed as a weighted average of the row profiles:

$$\mathbf{c}' = \sum_{i=1}^a p_{i\cdot} \mathbf{r}'_i. \quad (16.21)$$

Note that $\sum_{j=1}^b p_{\cdot j} = \sum_{i=1}^a p_{i\cdot} = 1$, or

$$\mathbf{j}' \mathbf{r} = \mathbf{c}' \mathbf{j} = 1, \quad (16.22)$$

where the first \mathbf{j} is $a \times 1$ and the second is $b \times 1$. Therefore the $p_{\cdot j}$'s and $p_{i\cdot}$'s serve as appropriate weights in the weighted averages (16.20) and (16.21).

■ EXAMPLE 16.2.2

In Table 16.5 (Hand et al. 1994, p. 12) we have the number of piston ring failures in each of three legs in four compressors found in the same building (the four compressors are oriented in the same direction). We obtain the correspondence matrix in Table 16.6 by dividing each element of Table 16.5 by $n = \sum_{ij} n_{ij} = 166$.

The vectors of row and column sums (marginal totals) in Table 16.6 are given by (16.9) and (16.10) as

$$\mathbf{r} = \begin{pmatrix} .277 \\ .199 \\ .229 \\ .295 \end{pmatrix},$$

$$\mathbf{c}' = (.319, .247, .434).$$

The matrix of row profiles is given by (16.15) as

$$\mathbf{R} = \mathbf{D}_r^{-1} \mathbf{P} = \begin{pmatrix} .370 & .370 & .261 \\ .333 & .273 & .394 \\ .290 & .211 & .500 \\ .286 & .143 & .571 \end{pmatrix}.$$

Table 16.6 Correspondence Matrix Obtained from Table 16.5

Compressor	Leg			Row Total
	North	Center	South	
1	.102	.102	.072	.277
2	.066	.054	.078	.199
3	.066	.048	.114	.229
4	.084	.042	.169	.295
Total	.319	.247	.434	1.000

The matrix of column profiles is given by (16.19) as

$$\mathbf{C} = \mathbf{P}\mathbf{D}_c^{-1} = \begin{pmatrix} .321 & .415 & .167 \\ .208 & .220 & .181 \\ .208 & .195 & .264 \\ .264 & .171 & .389 \end{pmatrix}.$$

□

16.2.3 Testing Independence

In Section 16.2.1, we noted that the data in a contingency table are often used to check for association of two categorical variables. If the two variables are denoted by x and y , then the assumption of independence can be expressed in terms of probabilities as

$$P(x_i y_j) = P(x_i)P(y_j), i = 1, 2, \dots, a; j = 1, 2, \dots, b, \quad (16.23)$$

where x_i and y_j correspond to the i th row and j th column of the contingency table. Using the notation in Table 16.4, we can estimate (16.23) as

$$p_{ij} = p_{i\cdot}p_{\cdot j}, i = 1, 2, \dots, a; j = 1, 2, \dots, b. \quad (16.24)$$

The usual chi-square statistic for testing independence of x and y (comparing p_{ij} with $p_{i\cdot}p_{\cdot j}$ for all i, j) is given by

$$\chi^2 = n \sum_{i=1}^a \sum_{j=1}^b \frac{(p_{ij} - p_{i\cdot}p_{\cdot j})^2}{p_{i\cdot}p_{\cdot j}}, \quad (16.25)$$

which is approximately (asymptotically) distributed as a chi-square random variable with $(a - 1)(b - 1)$ degrees of freedom. The statistic in (16.25) can also be written in terms of the frequencies n_{ij} rather than the relative frequencies p_{ij} :

$$\chi^2 = \sum_{i=1}^a \sum_{j=1}^b \frac{\left(\frac{n_{ij}}{n} - \frac{n_{i\cdot}n_{\cdot j}}{n} \right)^2}{\frac{n_{i\cdot}n_{\cdot j}}{n}}. \quad (16.26)$$

Two other alternative forms of (16.25) are

$$\chi^2 = \sum_{i=1}^a np_{i\cdot} \sum_{j=1}^b \left[\left(\frac{p_{ij}}{p_{i\cdot}} - p_{\cdot j} \right)^2 / p_{\cdot j} \right], \quad (16.27)$$

$$\chi^2 = \sum_{j=1}^b np_{\cdot j} \sum_{i=1}^a \left[\left(\frac{p_{ij}}{p_{\cdot j}} - p_{i\cdot} \right)^2 / p_{i\cdot} \right]. \quad (16.28)$$

In vector and matrix form, (16.27) and (16.28) can be written as

$$\chi^2 = \sum_{i=1}^a np_{i\cdot} (\mathbf{r}_i - \mathbf{c})' \mathbf{D}_c^{-1} (\mathbf{r}_i - \mathbf{c}), \quad (16.29)$$

$$\chi^2 = \sum_{j=1}^b np_{\cdot j} (\mathbf{c}_j - \mathbf{r})' \mathbf{D}_r^{-1} (\mathbf{c}_j - \mathbf{r}), \quad (16.30)$$

where \mathbf{r} , \mathbf{c} , \mathbf{r}_i , \mathbf{c}_j , \mathbf{D}_r , and \mathbf{D}_c are defined in (16.9), (16.10), (16.12), (16.16), (16.14), and (16.18), respectively. Thus, in (16.29) we compare \mathbf{r}_i to \mathbf{c} for each i , and in (16.30) we compare \mathbf{c}_j to \mathbf{r} for each j . Either of these is equivalent to testing independence by comparing p_{ij} to $p_{i\cdot}p_{\cdot j}$ for all i, j , since the definitions of χ^2 in (16.25)–(16.30) are all equal. Thus, the following three statements of independence are equivalent (for simplicity, we express the three statements in terms of sample quantities rather than their theoretical counterparts):

1. $p_{ij} = p_{i\cdot}p_{\cdot j}$ for all i, j (or $\mathbf{P} = \mathbf{rc}'$).
2. All rows \mathbf{r}'_i of \mathbf{R} in (16.15) are equal (and equal to their weighted average, \mathbf{c}').
3. All columns \mathbf{c}_j of \mathbf{C} in (16.19) are equal (and equal to their weighted average, \mathbf{r}).

Thus, if the variables x and y were independent, we would expect the rows of the contingency table to have similar profiles, or equivalently, the columns to have similar profiles. We can compare the row profiles to each other by comparing each row profile \mathbf{r}'_i to the weighted average \mathbf{c}' of the row profiles defined in (16.21). This comparison is made in the χ^2 statistic (16.29). Similarly, we compare column profiles in (16.30).

The chi-square statistic in (16.25) can be expressed in vector and matrix terms as

$$\chi^2 = n \operatorname{tr}[\mathbf{D}_r^{-1}(\mathbf{P} - \mathbf{rc}') \mathbf{D}_c^{-1}(\mathbf{P} - \mathbf{rc}')'] \quad (16.31)$$

$$= n \sum_{i=1}^k \lambda_i^2 \quad [\text{by (2.107)}], \quad (16.32)$$

where $\lambda_1^2, \lambda_2^2, \dots, \lambda_k^2$ are the nonzero eigenvalues of $\mathbf{D}_r^{-1}(\mathbf{P} - \mathbf{rc}') \mathbf{D}_c^{-1}(\mathbf{P} - \mathbf{rc}')'$ and

$$k = \operatorname{rank}[\mathbf{D}_r^{-1}(\mathbf{P} - \mathbf{rc}') \mathbf{D}_c^{-1}(\mathbf{P} - \mathbf{rc}')'] = \operatorname{rank}(\mathbf{P} - \mathbf{rc}'). \quad (16.33)$$

The rank of $\mathbf{P} - \mathbf{rc}'$ is ordinarily $k = \min[(a-1), (b-1)]$. It is clear that the rank is less than $\min(a, b)$ since

$$(\mathbf{P} - \mathbf{rc}')\mathbf{j} = \mathbf{P}\mathbf{j} - \mathbf{rc}'\mathbf{j} = \mathbf{r} - \mathbf{r} = \mathbf{0} \quad (16.34)$$

[see (16.9) and (16.22)].

■ EXAMPLE 16.2.3

In order to test independence of the rows (compressors) and columns (legs) of Table 16.5 in Example 16.2.2, we perform a chi-square test. Using (16.25) or (16.26), we obtain $\chi^2 = 11.722$, with 6 degrees of freedom, for which the p -value is 0.0685. There is some evidence of lack of independence between leg and compressor. \square

16.2.4 Coordinates for Plotting Row and Column Profiles

We now obtain coordinates of the row points and column points for the best two-dimensional representation of the data in a contingency table. As we will see, the metric for the row points and column points is the same, and the two sets of points can therefore be superimposed on the same plot.

In multidimensional scaling in Section 16.1, we transformed the distance matrix and then factored it by a spectral decomposition to obtain coordinates for plotting. In correspondence analysis, the matrix $\mathbf{P} - \mathbf{rc}'$ is not symmetric, and we therefore resort to a singular value decomposition to obtain coordinates.

We first scale $\mathbf{P} - \mathbf{rc}'$ to obtain

$$\mathbf{Z} = \mathbf{D}_r^{-1/2}(\mathbf{P} - \mathbf{rc}')\mathbf{D}_c^{-1/2}, \quad (16.35)$$

whose elements are

$$z_{ij} = \frac{p_{ij} - p_i \cdot p_j}{\sqrt{p_i \cdot p_j}} \quad (16.36)$$

as in (16.25). The $a \times b$ matrix \mathbf{Z} has rank $k = \min(a-1, b-1)$, the assumed rank of $\mathbf{P} - \mathbf{rc}'$. We then factor \mathbf{Z} using the singular value decomposition (2.117):

$$\mathbf{Z} = \mathbf{U}\Lambda\mathbf{V}'. \quad (16.37)$$

The columns of the $a \times k$ matrix \mathbf{U} are (normalized) eigenvectors of $\mathbf{Z}\mathbf{Z}'$; the columns of the $b \times k$ matrix \mathbf{V} are (normalized) eigenvectors of $\mathbf{Z}'\mathbf{Z}$; and $\Lambda = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_k)$, where $\lambda_1^2, \lambda_2^2, \dots, \lambda_k^2$ are the nonzero eigenvalues of $\mathbf{Z}'\mathbf{Z}$ and of $\mathbf{Z}\mathbf{Z}'$. The eigenvectors in \mathbf{U} and \mathbf{V} correspond to the eigenvalues $\lambda_1^2, \lambda_2^2, \dots, \lambda_k^2$. Since, the columns of \mathbf{U} and \mathbf{V} are orthonormal, $\mathbf{U}'\mathbf{U} = \mathbf{V}'\mathbf{V} = \mathbf{I}$. The values $\lambda_1, \lambda_2, \dots, \lambda_k$ in Λ are called the *singular values* of \mathbf{Z} . Note that, by (16.35),

$$\begin{aligned} \mathbf{Z}\mathbf{Z}' &= \mathbf{D}_r^{-1/2}(\mathbf{P} - \mathbf{rc}')\mathbf{D}_c^{-1/2}\mathbf{D}_c^{-1/2}(\mathbf{P} - \mathbf{rc})'\mathbf{D}_r^{-1/2} \\ &= \mathbf{D}_r^{-1/2}(\mathbf{P} - \mathbf{rc}')\mathbf{D}_c^{-1}(\mathbf{P} - \mathbf{rc})'\mathbf{D}_r^{-1/2}. \end{aligned} \quad (16.38)$$

The (nonzero) eigenvalues of $\mathbf{Z}\mathbf{Z}'$ in (16.38) are the same as those of

$$\mathbf{D}_r^{-1/2} \mathbf{D}_c^{-1/2} (\mathbf{P} - \mathbf{rc}') \mathbf{D}_c^{-1} (\mathbf{P} - \mathbf{rc})' \quad (16.39)$$

(see Section 2.11.5). The matrix expression in (16.39) is the same as that in (16.31). We have therefore denoted the eigenvalues as $\lambda_1^2, \lambda_2^2, \dots, \lambda_k^2$ as in (16.32).

We can obtain a decomposition of $\mathbf{P} - \mathbf{rc}'$ by equating the right-hand sides of (16.35) and (16.37) and solving for $\mathbf{P} - \mathbf{rc}'$:

$$\begin{aligned} \mathbf{D}_r^{-1/2} (\mathbf{P} - \mathbf{rc}') \mathbf{D}_c^{-1/2} &= \mathbf{U} \Lambda \mathbf{V}', \\ \mathbf{P} - \mathbf{rc}' &= \mathbf{D}_r^{1/2} \mathbf{U} \Lambda \mathbf{V}' \mathbf{D}_c^{1/2} \\ &= \mathbf{A} \Lambda \mathbf{B}' = \sum_{i=1}^k \lambda_i \mathbf{a}_i \mathbf{b}'_i, \end{aligned} \quad (16.40)$$

where $\mathbf{A} = \mathbf{D}_r^{1/2} \mathbf{U}$, $\mathbf{B} = \mathbf{D}_c^{1/2} \mathbf{V}$, \mathbf{a}_i and \mathbf{b}_i are columns of \mathbf{A} and \mathbf{B} , and $\Lambda = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_k)$.

Since $\mathbf{U}'\mathbf{U} = \mathbf{V}'\mathbf{V} = \mathbf{I}$, \mathbf{A} and \mathbf{B} in (16.40) are scaled so that $\mathbf{A}'\mathbf{D}_r^{-1}\mathbf{A} = \mathbf{B}'\mathbf{D}_c^{-1}\mathbf{B} = \mathbf{I}$. With this scaling, the decomposition in (16.40) is often called the *generalized singular value decomposition*.

In (16.40) the rows of $\mathbf{P} - \mathbf{rc}'$ are expressed as linear combinations of the rows of \mathbf{B}' , which are the columns of $\mathbf{B} = (\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_k)$. The coordinates (coefficients) for the i th row of $\mathbf{P} - \mathbf{rc}'$ are found in the i th row of $\mathbf{A}\Lambda$. In like manner, the coordinates for the columns of $\mathbf{P} - \mathbf{rc}'$ are given by the columns of $\Lambda\mathbf{B}'$, since the columns of $\Lambda\mathbf{B}'$ provide coefficients for the columns of $\mathbf{A} = (\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_k)$ in (16.40).

To find coordinates for the row deviations $\mathbf{r}'_i - \mathbf{c}'$ in $\mathbf{R} - \mathbf{jc}'$ and the column deviations $\mathbf{c}_j - \mathbf{r}$ in $\mathbf{C} - \mathbf{rj}'$, we express the two matrices as functions of $\mathbf{P} - \mathbf{rc}'$ (see Problem 16.8):

$$\mathbf{R} - \mathbf{jc}' = \mathbf{D}_r^{-1} (\mathbf{P} - \mathbf{rc}'), \quad (16.41)$$

$$\mathbf{C} - \mathbf{rj}' = \mathbf{D}_c^{-1} (\mathbf{P} - \mathbf{rc}'). \quad (16.42)$$

Thus the coordinates for the row deviations in $\mathbf{R} - \mathbf{jc}'$ with respect to the axes provided by $\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_k$ are given by the columns of

$$\mathbf{X} = \mathbf{D}_r^{-1} \mathbf{A} \Lambda. \quad (16.43)$$

Similarly, the coordinates for the column deviations in $\mathbf{C} - \mathbf{rj}'$ with respect to the axes $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_k$ are given by the columns of

$$\mathbf{Y} = \mathbf{D}_c^{-1} \mathbf{B} \Lambda. \quad (16.44)$$

Therefore, to plot the coordinates for the row profile deviations $\mathbf{r}'_i - \mathbf{c}'$, $i = 1, 2, \dots, a$, in two dimensions, we plot the rows of the first two columns of \mathbf{X} :

$$\mathbf{X}_1 = \begin{pmatrix} x_{11} & x_{12} \\ x_{21} & x_{22} \\ \vdots & \vdots \\ x_{a1} & x_{a2} \end{pmatrix}.$$

Similarly, to plot the coordinates for the column profile deviations $\mathbf{c}_j - \mathbf{r}, j = 1, 2, \dots, b$, in two dimensions, we plot the rows of the first two columns of \mathbf{Y} :

$$\mathbf{Y}_1 = \begin{pmatrix} y_{11} & y_{12} \\ y_{21} & y_{22} \\ \vdots & \vdots \\ y_{b1} & y_{b2} \end{pmatrix}.$$

Both plots can be superimposed on the same graph because \mathbf{A} and \mathbf{B} in (16.40) share the same singular values $\lambda_1, \lambda_2, \dots, \lambda_k$ in Λ . Distances between row points and distances between column points are meaningful. For example, the distance between two row points is related to the chi-square metric implicit in (16.29). The chi-square distance between two row profiles \mathbf{r}_i and \mathbf{r}_j is given by

$$d_{ij}^2 = (\mathbf{r}_i - \mathbf{r}_j)' \mathbf{D}_c^{-1} (\mathbf{r}_i - \mathbf{r}_j).$$

If two row points (or two column points) are close, the two rows (or two columns) could be combined into a single category if necessary to improve the chi-square approximation.

The distance between a row point and a column point is not meaningful, but the proximity of a row point and a column point has meaning as noted in Section 16.2.1, namely that these two categories of the two variables occur more frequently than would be expected to happen by chance if the two variables were independent.

The weighted average (weighted by p_i) of the distances $(\mathbf{r}_i - \mathbf{c})' \mathbf{D}_c^{-1} (\mathbf{r}_i - \mathbf{c})$ between the row profiles \mathbf{r}_i and their mean \mathbf{c} [see (16.21)] weighted by p_i , is called the *total inertia*. By (16.29) this can be expressed as χ^2/n :

$$\text{Total inertia} = \frac{\chi^2}{n} = \sum_{i=1}^a p_i (\mathbf{r}_i - \mathbf{c})' \mathbf{D}_c^{-1} (\mathbf{r}_i - \mathbf{c}). \quad (16.45)$$

As noted following (16.21), $\sum_i p_i = 1$, and therefore the p_i 's serve as appropriate weights.

By (16.32), we can write (16.45) as

$$\frac{\chi^2}{n} = \sum_{i=1}^k \lambda_i^2. \quad (16.46)$$

Therefore, the contribution of each of the first two dimensions (axes) of our plot to the total inertia in (16.45) is $\lambda_1^2 / \sum_i \lambda_i^2$ and $\lambda_2^2 / \sum_i \lambda_i^2$. The combined contribution of the two dimensions is

$$\frac{\lambda_1^2 + \lambda_2^2}{\sum_{i=1}^k \lambda_i^2}. \quad (16.47)$$

If (16.47) is large, then the points in the plane of the first two dimensions account for nearly all the variation in the data, including the associations. The total inertia in




Figure 16.8 Row points (1, 2, 3, 4) and column points (Center, North, South).

(16.45) and (16.46) can also be described in terms of the columns by using (16.30):

$$\text{Total inertia} = \frac{\chi^2}{n} = \sum_{j=1}^b p_{\cdot j} (\mathbf{c}_j - \mathbf{r})' \mathbf{D}_r^{-1} (\mathbf{C}_j - \mathbf{r}) = \sum_{i=1}^k \lambda_i^2. \quad (16.48)$$

Since the inertia associated with the axes for columns is the same as that for rows, the row and column points can be plotted on the same axes.

Some computer programs use a singular value decomposition of \mathbf{P} rather than of $\mathbf{P} - \mathbf{rc}'$. The results are the same if the first singular value (which is 1) is discarded along with the first column of \mathbf{A} (which is \mathbf{r}) and the first column of \mathbf{B} (which is \mathbf{c}).

■ EXAMPLE 16.2.4

We continue the analysis of the piston ring data of Table 16.5. A correspondence analysis is performed, and a plot of the row and column points is given in Figure 16.8. Row points do not lie near other row points and column points do not lie near column points. However, compressor 1 seems to be closely associated with the center leg, compressor 2 with the north leg, and compressor 4 with the south leg. These associations illustrate the lack of independence between compressor and leg position.

Singular values and inertias are given in Table 16.7. Most of the variation is due to the first dimension, and the first two dimensions explain all the variation because $\text{rank}(\mathbf{Z}) = \min(a - 1, b - 1) = \min(4 - 1, 3 - 1) = 2$, where \mathbf{Z} is defined in (16.35). \square

Table 16.7 Singular Values (λ_i), Inertia (λ_i^2), Chi-Square ($n\lambda_i^2$), and Percent ($\lambda_i^2 / \sum_j \lambda_j^2$) for the Data in Table 16.5

Singular Value	Inertia	Chi-Square	Percent
.26528	.07037	11.6819	99.66
.01560	.00024	.0404	.34
Total	.07062	11.7223	100.00

16.2.5 Multiple Correspondence Analysis

Correspondence analysis of a two-way contingency table can be extended to a three-way or higher-order multi-way table. By the method of *multiple correspondence analysis*, we obtain a two-dimensional graphical display of the information in the multi-way contingency table. The method involves a correspondence analysis of an indicator matrix G . There is a row of G for each item; thus the number of rows of G is the total number of items in the sample. The number of columns of G is the total number of categories in all variables. The elements of G are 1's and 0's. In each row, an element is 1 if the item belongs in the corresponding category of the variable; otherwise, the element is 0. Thus the number of 1's in a row of G is the number of variables; for a four-way contingency table, for example, there would be four 1's in each row of G .

We illustrate a four-way classification with the (contrived) data in Table 16.8. There are $n = 12$ items (people) and $p = 4$ categorical variables. The four variables and their categories are listed in Table 16.9. The indicator matrix G for the data in Table 16.8 is given in Table 16.10.

Table 16.8 A List of 12 People and Their Categories on Four Variables

Person	Gender	Age	Marital Status	Hair Color
1	Male	Young	Single	Brown
2	Male	Old	Single	Red
3	Female	Middle	Married	Blond
4	Male	Old	Single	Black
5	Female	Middle	Married	Black
6	Female	Middle	Single	Brown
7	Male	Young	Married	Red
8	Male	Old	Married	Blond
9	Male	Middle	Single	Brown
10	Female	Young	Married	Black
11	Female	Old	Single	Brown
12	Male	Young	Married	Blond

Table 16.9 The Categories for the Four Variables in Table 16.8

Variable	Levels
Gender	Male, female
Age	Young, middle aged, old
Marital status	Single, married
Hair color	Blond, brown, black, red

Table 16.10 Indicator Matrix \mathbf{G} for the Data in Table 16.8

Person	Gender	Age	Marital Status	Hair Color
1	1 0	1 0 0	1 0	0 1 0 0
2	1 0	0 0 1	1 0	0 0 0 1
3	0 1	0 1 0	0 1	1 0 0 0
4	1 0	0 0 1	1 0	0 0 1 0
5	0 1	0 1 0	0 1	0 0 1 0
6	0 1	0 1 0	1 0	0 1 0 0
7	1 0	1 0 0	0 1	0 0 0 1
8	1 0	0 0 1	0 1	1 0 0 0
9	1 0	0 1 0	1 0	1 0 0 0
10	0 1	1 0 0	0 1	0 0 1 0
11	0 1	0 0 1	1 0	0 1 0 0
12	1 0	1 0 0	0 1	1 0 0 0

A correspondence analysis on \mathbf{G} is equivalent to a correspondence analysis on $\mathbf{G}'\mathbf{G}$, which is called the *Burt matrix*. This equivalence can be justified as follows. In the singular value decomposition $\mathbf{G} = \mathbf{U}\Lambda\mathbf{V}'$, the matrix \mathbf{V} contains eigenvectors of $\mathbf{G}'\mathbf{G}$. The same matrix \mathbf{V} would be used in the spectral decomposition of $\mathbf{G}'\mathbf{G}$. Thus the columns of \mathbf{V} are used in plotting coordinates for the columns of \mathbf{G} or the columns of $\mathbf{G}'\mathbf{G}$. If \mathbf{G} is $n \times p$ with $p < n$, then $\mathbf{G}'\mathbf{G}$ would be smaller in size than \mathbf{G} .

The Burt matrix $\mathbf{G}'\mathbf{G}$ has a square block on the diagonal for each variable and a rectangular block off-diagonal for each pair of variables. Each diagonal block is a diagonal matrix showing the frequencies for the categories in the corresponding variable. Each off-diagonal block is a two-way contingency table for the corresponding pair of variables. In Table 16.11, we show the $\mathbf{G}'\mathbf{G}$ matrix for the \mathbf{G} matrix in Table 16.10.

A correspondence analysis of $\mathbf{G}'\mathbf{G}$ yields only column coordinates. A point is plotted for each column of \mathbf{G} (or of $\mathbf{G}'\mathbf{G}$). Thus each point represents a category (attribute) of one of the variables.

Distances between points are not as meaningful as in correspondence analysis, but points in the same quadrant or approximate vicinity indicate an association. If

Table 16.11 Burt Matrix $\mathbf{G}'\mathbf{G}$ for the Matrix \mathbf{G} in Table 16.10

Gender		Age			Marital Status		Hair Color			
7	0	3	1	3	4	3	3	1	1	2
0	5	1	3	1	2	3	1	2	2	0
3	1	4	0	0	1	3	1	1	1	1
1	3	0	4	0	2	2	2	1	1	0
3	1	0	0	4	3	1	1	1	1	1
4	2	1	2	3	6	0	1	3	1	1
3	3	3	2	1	0	6	3	0	2	1
3	1	1	2	1	1	3	4	0	0	0
1	2	1	1	1	3	0	0	3	0	0
1	2	1	1	1	1	2	0	0	3	0
2	0	1	0	1	1	1	0	0	0	2

Table 16.12 Singular Values (λ_i), Inertia (λ_i^2), Chi-Square ($n\lambda_i^2$), and Percent ($\lambda_i^2 / \sum_j \lambda_j^2$) for the Burt Matrix $\mathbf{G}'\mathbf{G}$ in Table 16.11

Singular Value	Inertia	Chi-Square	Percent
.68803	.47338	31.551	27.05
.67451	.45497	30.324	26.00
.51492	.26515	17.672	15.15
.50000	.25000	16.663	14.29
.41941	.17590	11.724	10.05
.33278	.11074	7.381	6.33
.14091	.01986	1.323	1.13
Total	1.75000	116.638	100.00

two close points represent attributes of the same variable, the two attributes may be combined into a single attribute.

Since the Burt matrix $\mathbf{G}'\mathbf{G}$ has only two-way contingency tables, three-way and higher-order interactions are not represented in the plot. The various two-way tables are analyzed simultaneously, however.

■ EXAMPLE 16.2.5(a)

We continue the illustration in this section. A correspondence analysis of the Burt matrix $\mathbf{G}'\mathbf{G}$ in Table 16.11 yielded the singular values, inertia, and chi-square values in Table 16.12. The first two singular values account for only 53.05% of the total variation. A plot of the first two dimensions for the 11




Figure 16.9 Plot of Points Representing the 11 Columns of Table 16.10 or Table 16.11.

columns in Table 16.10 or Table 16.11 is given in Figure 16.9. It appears that married and blond hair have a greater association than would be expected by chance alone. Another association is that between female and middle age. \square

■ EXAMPLE 16.2.5(b)

Table 16.13 (Edwards and Kreiner 1983) is a five-way contingency table of employed men between the ages of 18 and 67 who were asked whether they themselves carried out repair work on their home, as opposed to hiring a craftsman to do the job. The five categorical variables are as follows:

Work of respondent: Skilled, Unskilled, Office

Tenure: Rent, Own

Age: Up to 30, 31–45, over 45

Accommodation type: Apartment, House

Response to repair question: Yes, No

A multiple correspondence analysis produced the inertia and singular values in Table 16.14. The plot of the first two dimensions is given in Figure 16.10.

Unskilled employment has a high association with not doing ones own repairs. Doing ones own repairs is associated with owning a house, age between 31 and 45, and doing office work. Living in an apartment is associated with renting. \square

Table 16.13 Do-It-Yourself Data

Work	Tenure	Response	Accommodation Type					
			Apartment			House		
			Age			Age		
Skilled	Rent	Yes	18	15	6	34	10	2
		No	15	13	9	28	4	6
	Own	Yes	5	3	1	56	56	35
		No	1	1	1	12	21	8
Unskilled	Rent	Yes	17	10	15	29	3	7
		No	34	17	19	44	13	16
	Own	Yes	2	0	3	23	52	49
		No	3	2	0	9	31	51
Office	Rent	Yes	30	23	21	22	13	21
		No	25	19	40	25	16	12
	Own	Yes	8	5	1	54	191	102
		No	4	2	2	19	76	61

Table 16.14 Singular Values (λ_i), Inertia (λ_i^2), Chi-Square ($n\lambda_i^2$), and Percent ($\lambda_i^2 / \sum_j \lambda_j^2$) for the Do-It-Yourself Data in Table 16.13

Singular Value	Inertia	Chi-Square	Percent
.60707	.36853	3446.5	26.32
.49477	.24480	2289.4	17.49
.45591	.20785	1943.9	14.85
.42704	.18237	1705.5	13.03
.40516	.16415	1535.2	11.73
.39392	.15517	1451.2	11.08
.27771	.07713	721.3	5.51
Total	1.40000	13092.9	100

16.3 BIPLOTS

16.3.1 Introduction

A biplot is a two-dimensional representation of a data matrix \mathbf{Y} [see (3.17)] showing a point for each of the n observation vectors (rows of \mathbf{Y}) along with a point for each




Figure 16.10 Plot of points representing the 12 categories in Table 16.13.

of the p variables (columns of \mathbf{Y}). The prefix “bi” refers to the two kinds of points, not to the dimensionality of the plot. The method presented here could, in fact, be generalized to a three-dimensional (or higher-order) biplot. Biplots were introduced by Gabriel (1971) and have been discussed at length by Gower and Hand (1996); see also Khattree and Naik (2000), Jacoby (1998, Chapter 7), and Seber (1984, pp. 204–212).

If $p = 2$, a simple scatterplot such as in Section 3.3 has both kinds of information, namely, a point for each observation and the two axes representing the variables. We can see at a glance the placement of the points relative to each other and relative to the variables.

When $p > 2$, we can obtain a two-dimensional plot of the observations by plotting the first two principal components of \mathbf{S} as in Section 12.4. We can then add a representation of the p variables to the plot of principal components to obtain a biplot. The principal component approach is discussed in Section 16.3.2. A method based on the singular value decomposition is presented in Section 16.3.3, and other methods are reviewed in Section 16.3.5.

16.3.2 Principal Component Plots

A principal component is given by $z = \mathbf{a}'\mathbf{y}$, where \mathbf{a} is an eigenvector of \mathbf{S} , the sample covariance matrix, and \mathbf{y} is a $p \times 1$ observation vector (see Section 12.2). There are p eigenvectors $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_p$, and thus there are p principal components z_1, z_2, \dots, z_p for each observation vector $\mathbf{y}_i, i = 1, 2, \dots, n$. Hence (using the

centered form) the observation vectors are transformed to $z_{ij} = \mathbf{a}'_j(\mathbf{y}_i - \bar{\mathbf{y}}) = (\mathbf{y}_i - \bar{\mathbf{y}})' \mathbf{a}_j, i = 1, 2, \dots, n; j = 1, 2, \dots, p$. Each $p \times 1$ observation vector \mathbf{y}_i is transformed to a $p \times 1$ vector of principal components,

$$\begin{aligned}\mathbf{z}'_i &= [(\mathbf{y}_i - \bar{\mathbf{y}})' \mathbf{a}_1, (\mathbf{y}_i - \bar{\mathbf{y}})' \mathbf{a}_2, \dots, (\mathbf{y}_i - \bar{\mathbf{y}})' \mathbf{a}_p] \\ &= (\mathbf{y}_i - \bar{\mathbf{y}})' (\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_p) \\ &= (\mathbf{y}_i - \bar{\mathbf{y}})' \mathbf{A}, \quad i = 1, 2, \dots, n,\end{aligned}\tag{16.49}$$

where $\mathbf{A} = (\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_p)$ is the $p \times p$ matrix whose columns are (normalized) eigenvectors of \mathbf{S} . [Note that the matrix \mathbf{A} in (16.49) is the transpose of \mathbf{A} in (12.3)]. With \mathbf{Z} and \mathbf{Y}_c defined as

$$\mathbf{Z} = \begin{pmatrix} \mathbf{z}'_1 \\ \mathbf{z}'_2 \\ \vdots \\ \mathbf{z}'_n \end{pmatrix}, \quad \mathbf{Y}_c = \begin{pmatrix} (\mathbf{y}_1 - \bar{\mathbf{y}})' \\ (\mathbf{y}_2 - \bar{\mathbf{y}})' \\ \vdots \\ (\mathbf{y}_n - \bar{\mathbf{y}})' \end{pmatrix}\tag{16.50}$$

[see (10.13)], we can express the principal components in (16.49) as

$$\mathbf{Z} = \mathbf{Y}_c \mathbf{A}.\tag{16.51}$$

Since the eigenvectors \mathbf{a}_j of the symmetric matrix \mathbf{S} are mutually orthogonal (see Section 2.11.6), $\mathbf{A} = (\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_p)$ is an orthogonal matrix and $\mathbf{A}\mathbf{A}' = \mathbf{I}$. Multiplying (16.51) on the right by \mathbf{A}' , we obtain

$$\mathbf{Y}_c = \mathbf{Z}\mathbf{A}'.\tag{16.52}$$

The best two-dimensional representation of \mathbf{Y}_c is given by taking the first two columns of \mathbf{Z} and the first two columns of \mathbf{A} . If the resulting matrices are denoted by \mathbf{Z}_2 and \mathbf{A}_2 , we have

$$\mathbf{Y}_c \cong \mathbf{Z}_2 \mathbf{A}'_2.\tag{16.53}$$

The fit in (16.53) is best in a least squares sense. If the left side of (16.53) is represented by $\mathbf{Y}_c = \mathbf{B} = (b_{ij})$ and the right side by $\mathbf{Z}_2 \mathbf{A}'_2 = \mathbf{C} = (c_{ij})$, then $\sum_{i=1}^n \sum_{j=1}^p (b_{ij} - c_{ij})^2$ is minimized (Seber 1984, p. 206).

The coordinates for the n observations are the rows of \mathbf{Z}_2 and the coordinates for the p variables are the rows of \mathbf{A}_2 (columns of \mathbf{A}'_2). The coordinates are discussed further in Section 16.3.4.

The adequacy of the fit in (16.53) can be evaluated by examining the first two eigenvalues λ_1 and λ_2 of \mathbf{S} . Thus a large value (close to 1) of

$$\frac{\lambda_1 + \lambda_2}{\sum_{i=1}^p \lambda_i}$$

would indicate that \mathbf{Y}_c is represented well visually in the plot.

16.3.3 Singular Value Decomposition Plots

We can obtain $\mathbf{Y}_c = \mathbf{ZA}'$ in (16.52) by means of the singular value decomposition of \mathbf{Y}_c . By (2.117), we have

$$\mathbf{Y}_c = \mathbf{U}\Lambda\mathbf{V}', \quad (16.54)$$

where $\Lambda = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_p)$ is a diagonal matrix containing square roots of the nonzero eigenvalues $\lambda_1^2, \lambda_2^2, \dots, \lambda_p^2$ of $\mathbf{Y}_c'\mathbf{Y}_c$ (or of $\mathbf{Y}_c\mathbf{Y}_c'$), the columns of \mathbf{U} are the corresponding eigenvectors of $\mathbf{Y}_c\mathbf{Y}_c'$, and the columns of \mathbf{V} are the corresponding eigenvectors of $\mathbf{Y}_c'\mathbf{Y}_c$.

The product $\mathbf{U}\Lambda$ in (16.54) is equal to \mathbf{Z} , the matrix of principal component scores in (16.51). To see this we multiply (16.54) by \mathbf{V} , which is orthogonal because it contains the (normalized) eigenvectors of the symmetric matrix $\mathbf{Y}_c'\mathbf{Y}_c$ (see Section 2.11.6). This gives

$$\mathbf{Y}_c\mathbf{V} = \mathbf{U}\Lambda\mathbf{V}'\mathbf{V} = \mathbf{U}\Lambda. \quad (16.55)$$

By (10.17), $\mathbf{Y}_c'\mathbf{Y}_c$ is equal to $(n - 1)\mathbf{S}$. By (2.106), eigenvectors of $(n - 1)\mathbf{S}$ are also eigenvectors of \mathbf{S} . Thus \mathbf{V} is the same as \mathbf{A} in (16.51), which contains eigenvectors of \mathbf{S} . Hence, $\mathbf{Y}_c\mathbf{V}$ in (16.55) becomes

$$\begin{aligned} \mathbf{Y}_c\mathbf{V} &= \mathbf{Y}_c\mathbf{A} \\ &= \mathbf{Z} \quad [\text{by (16.51)}] \\ &= \mathbf{U}\Lambda \quad [\text{by (16.55)}]. \end{aligned}$$

We can therefore write (16.54) as

$$\mathbf{Y}_c = \mathbf{U}\Lambda\mathbf{V}' = \mathbf{Z}\mathbf{V}' = \mathbf{Z}\mathbf{A}'. \quad (16.56)$$

Thus the singular value decomposition of \mathbf{Y}_c gives the same factoring as the expression in (16.52) based on principal components.

16.3.4 Coordinates

In this section, we consider the coordinates for the methods of Sections 16.3.2 and 16.3.3. Let us return to (16.53), the two-dimensional representation of \mathbf{Y}_c based on principal components (which is the same representation as that based on the singular value decomposition):

$$\mathbf{Y}_c \cong \mathbf{Z}_2\mathbf{A}'_2 = \begin{pmatrix} z_{11} & z_{12} \\ z_{21} & z_{22} \\ \vdots & \vdots \\ z_{n1} & z_{n2} \end{pmatrix} \begin{pmatrix} a_{11} & a_{21} & \cdots & a_{p1} \\ a_{12} & a_{22} & \cdots & a_{p2} \end{pmatrix}. \quad (16.57)$$

The elements of (16.57) are of the form

$$y_{ij} - \bar{y}_j \cong z_{i1}a_{j1} + z_{i2}a_{j2}, \quad i = 1, 2, \dots, n; j = 1, 2, \dots, p.$$

Thus each observation is represented as a linear combination, the coordinates (coefficients) being the elements of the vector (z_{i1}, z_{i2}) and the axes being the elements of the vector (a_{j1}, a_{j2}) . We therefore plot the points $(z_{i1}, z_{i2}), i = 1, 2, \dots, n$, and the points $(a_{j1}, a_{j2}), j = 1, 2, \dots, p$. To distinguish them and to show relationship of the points to the axes, the points (a_{j1}, a_{j2}) are connected to the origin with a straight line forming an arrow. If necessary, the scale of the points (a_{j1}, a_{j2}) could be adjusted to be compatible with that of the principal components (z_{i1}, z_{i2}) .

The Euclidean distance between two points (z_{i1}, z_{i2}) and (z_{k1}, z_{k2}) is approximately equal to the distance between the corresponding points (rows) \mathbf{y}'_i and \mathbf{y}'_k in the data matrix \mathbf{Y} . If all of the principal components were used, as in (16.51) and (16.52), the distance would be the same, but with only two principal components, the distance is an approximation.

The cosine of the angle between the arrows (lines) drawn to each pair of axis points (a_{j1}, a_{j2}) and (a_{k1}, a_{k2}) shows the correlation between the two corresponding variables [see (3.14) and (3.15)]. Thus a small angle between two vectors indicates that the two variables are highly correlated, two variables whose vectors form a 90° angle are uncorrelated, and an angle greater than 90° indicates that the variables are negatively correlated.

The values of the p variables in the i th observation vector \mathbf{y}_i (corrected for means) are related to the perpendicular projection of the point (z_{1i}, z_{2i}) on the p vectors from the origin to the points (a_{j1}, a_{j2}) representing variables. The further from the origin a projection falls on an arrow, the larger the value of the observation on that variable. Hence the vectors will be oriented toward the observations that have larger values on the corresponding variables.

■ EXAMPLE 16.3.4

Using the city crime data of Table 15.1, we illustrate the principal components approach. The first two eigenvectors of the sample covariance matrix \mathbf{S} are given by

$$\mathbf{A}_2 = \begin{pmatrix} .002 & .008 \\ .017 & .014 \\ .182 & .689 \\ .104 & .221 \\ .747 & -.240 \\ .612 & -.109 \\ .153 & .638 \end{pmatrix}.$$

The matrix of the first two principal components is given by

$$\mathbf{Z}_2 = \mathbf{Y}_c \mathbf{A}_2 = \begin{pmatrix} -317.2 & -156.1 \\ -491.8 & 192.4 \\ -650.0 & 227.6 \\ 141.7 & -133.8 \\ 342 & -69.3 \\ 312.2 & 164.1 \\ -514.7 & -166.4 \\ 58.6 & -239.7 \\ -24.5 & 25.9 \\ 75.7 & 40.1 \\ 678.2 & 7.1 \\ -192.4 & 163.4 \\ 542.7 & 194.8 \\ 233.1 & -266.8 \\ -343.8 & -184.3 \\ 150.4 & 200.9 \end{pmatrix}.$$

The coordinates for the sixteen cities are found in \mathbf{Z}_2 , and the coordinates for the seven variables are found in \mathbf{A}_2 . The plot of the city and variable points could be given, but because of the different scaling for the observation points (spread out) and the variable points (clustered at the origin), the plot is not illuminating. Suitable scaling of the eigenvectors in \mathbf{A}_2 would enable the arrows representing the variables to pass through the points (see Example 16.3.5). \square

16.3.5 Other Methods

The singular value decomposition of \mathbf{Y}_c is given in (16.54) as

$$\mathbf{Y}_c = \mathbf{U}\Lambda\mathbf{V}' \quad (16.58)$$

In Section 16.3.3, it was shown that $\mathbf{U}\Lambda = \mathbf{Z}$ and $\mathbf{V} = \mathbf{A}$ [see (16.56)], so that (16.58) can be written as

$$\mathbf{Y}_c = (\mathbf{U}\Lambda)\mathbf{V}' = \mathbf{Z}\mathbf{A}',$$

which is equivalent to the principal component solution $\mathbf{Y}_c = \mathbf{Z}\mathbf{A}'$ in (16.52). Alternative factorings may be of interest. Two that have been considered are

$$\mathbf{Y}_c = \mathbf{U}(\Lambda\mathbf{V}'), \quad (16.59)$$

$$\mathbf{Y}_c = (\mathbf{U}\Lambda^{1/2})(\Lambda^{1/2}\mathbf{V}'). \quad (16.60)$$

If we denote the submatrices consisting of the first two columns of \mathbf{U} and \mathbf{V} as \mathbf{U}_2 and \mathbf{V}_2 , respectively, and define $\Lambda_2 = \text{diag}(\lambda_1, \lambda_2)$, then the two-dimensional

representations of (16.59) and (16.60) are

$$\mathbf{Y}_c \cong \mathbf{U}_2(\Lambda_2 \mathbf{V}'_2) \\ = \begin{pmatrix} u_{11} & u_{12} \\ u_{21} & u_{22} \\ \vdots & \vdots \\ u_{n1} & u_{n2} \end{pmatrix} \begin{pmatrix} \lambda_1 v_{11} & \lambda_1 v_{21} & \cdots & \lambda_1 v_{p1} \\ \lambda_2 v_{12} & \lambda_2 v_{22} & \cdots & \lambda_2 v_{p2} \end{pmatrix}, \quad (16.61)$$

$$\mathbf{Y}_c \cong (\mathbf{U}_2 \Lambda_2^{1/2})(\Lambda_2^{1/2} \mathbf{V}') \\ = \begin{pmatrix} \sqrt{\lambda_1} u_{11} & \sqrt{\lambda_2} u_{12} \\ \sqrt{\lambda_1} u_{21} & \sqrt{\lambda_2} u_{22} \\ \vdots & \vdots \\ \sqrt{\lambda_1} u_{n1} & \sqrt{\lambda_2} u_{n2} \end{pmatrix} \begin{pmatrix} \sqrt{\lambda_1} v_{11} & \sqrt{\lambda_1} v_{21} & \cdots & \sqrt{\lambda_1} v_{p1} \\ \sqrt{\lambda_2} v_{12} & \sqrt{\lambda_2} v_{22} & \cdots & \sqrt{\lambda_2} v_{p2} \end{pmatrix}. \quad (16.62)$$

For the biplot arising from (16.61), we plot the set of points (u_{i1}, u_{i2}) , $i = 1, 2, \dots, n$, and the set of points $(\lambda_1 v_{j1}, \lambda_2 v_{j2})$, $j = 1, 2, \dots, p$, with the latter points connected to the origin by an arrow to show the axes. For the biplot arising from (16.62), we plot the set of points $(\sqrt{\lambda_1} u_{i1}, \sqrt{\lambda_2} u_{i2})$, $i = 1, 2, \dots, n$, and the set of points $(\sqrt{\lambda_1} v_{j1}, \sqrt{\lambda_2} v_{j2})$, $j = 1, 2, \dots, p$, with the latter points connected to the origin with an arrow.

The presence of λ_1 and λ_2 in (16.61) and (16.62) provides scaling that is absent in (16.57). For many data sets the scaling in (16.62) will be adequate with no further adjustment.

If we write (16.59) in the form

$$\mathbf{Y}_c = \mathbf{U}(\Lambda \mathbf{V}') = \mathbf{U}(\mathbf{V} \Lambda)' = \mathbf{U} \mathbf{H}', \quad (16.63)$$

then

$$\mathbf{U} \mathbf{U}' = \left(\frac{1}{n-1} \right) \mathbf{Y}_c \mathbf{S}^{-1} \mathbf{Y}'_c, \quad (16.64)$$

$$\mathbf{H} \mathbf{H}' = (n-1) \mathbf{S} \quad (16.65)$$

(see Problem 16.10). With suitable scaling of the eigenvectors in \mathbf{U} and \mathbf{V} , we could eliminate the coefficients involving $n-1$ from (16.64) and (16.65).

By (16.64) (with scaling to eliminate $n-1$), the (Euclidean) distance $(\mathbf{u}_i - \mathbf{u}_k)'(\mathbf{u}_i - \mathbf{u}_k)$ between two points \mathbf{u}_i and \mathbf{u}_k is equal to the Mahalanobis distance $(\mathbf{y}_i - \mathbf{y}_k)' \mathbf{S}^{-1} (\mathbf{y}_i - \mathbf{y}_k)$ between the corresponding points \mathbf{y}_i and \mathbf{y}_k in the data matrix \mathbf{Y} :

$$(\mathbf{u}_i - \mathbf{u}_k)'(\mathbf{u}_i - \mathbf{u}_k) = (\mathbf{y}_i - \mathbf{y}_k)' \mathbf{S}^{-1} (\mathbf{y}_i - \mathbf{y}_k) \quad (16.66)$$

(see Problem 16.11). By (16.65), the covariance s_{jk} between the j th and k th variables (columns of \mathbf{Y}) is given by

$$s_{jk} = \mathbf{h}'_j \mathbf{h}_k, \quad (16.67)$$

where \mathbf{h}'_j and \mathbf{h}'_k are rows of \mathbf{H} . By (3.14) and (3.15), this can be converted to the correlation

$$r_{jk} = \cos \theta = \frac{\mathbf{h}'_j \mathbf{h}_k}{\sqrt{(\mathbf{h}'_j \mathbf{h}_j)(\mathbf{h}'_k \mathbf{h}_k)}}, \quad (16.68)$$

so that the angle between the two vectors \mathbf{h}_j and \mathbf{h}_k is related to the correlation between the j th and k th variables.

The two-dimensional representation of \mathbf{u}_i and \mathbf{h}_j in (16.61) has the approximate Mahalanobis distance and correlation properties discussed earlier.

■ EXAMPLE 16.3.5

Using the city crime data of Table 15.1, we illustrate the singular value decomposition method with the factorings in (16.61) and (16.62). The matrices \mathbf{U}_2 , Λ_2 , and \mathbf{V}_2 are

$$\mathbf{U}_2 = \begin{pmatrix} -.211 & -.230 \\ -.327 & .284 \\ -.432 & .335 \\ .094 & -.197 \\ .227 & -.102 \\ .208 & .242 \\ -.342 & -.245 \\ .039 & -.353 \\ -.016 & .038 \\ .050 & .059 \\ .451 & .010 \\ -.128 & .241 \\ .361 & .287 \\ .155 & -.393 \\ .229 & -.272 \\ .100 & .296 \end{pmatrix}, \quad \mathbf{V}_2 = \begin{pmatrix} .002 & .008 \\ .017 & .014 \\ .182 & .689 \\ .104 & .221 \\ .747 & -.240 \\ .612 & -.109 \\ .153 & .639 \end{pmatrix},$$

$$\Lambda_2 = \text{diag}(1503.604, 678.615).$$

By (16.61), the two-dimensional representation is given by plotting the rows of \mathbf{U}_2 and the rows of $\mathbf{V}_2 \Lambda_2$ (or the columns of $\Lambda_2 \mathbf{V}'_2$). For $\mathbf{V}_2 \Lambda_2$ we have

$$\mathbf{V}_2 \Lambda_2 = \begin{pmatrix} 3.0 & 5.2 \\ 25.4 & 9.5 \\ 273.6 & 467.5 \\ 156.3 & 150.1 \\ 1123.4 & -162.6 \\ 920.0 & -74.1 \\ 229.3 & 432.9 \end{pmatrix}.$$

As in Example 16.3.4, the plot of the observation points and variable points is not useful, again because the scale for the observation points differs greatly from the scale for the variable points.

For (16.62), we obtain

$$\mathbf{U}_2 \Lambda_2^{1/2} = \begin{pmatrix} -8.18 & -5.99 \\ -12.68 & 7.39 \\ -16.76 & 8.74 \\ 3.65 & -5.14 \\ 8.82 & -2.66 \\ 8.05 & 6.30 \\ -13.27 & -6.39 \\ 1.51 & -9.20 \\ -.63 & .99 \\ 1.95 & 1.54 \\ 17.49 & .27 \\ -4.96 & 6.27 \\ 13.99 & 7.48 \\ 6.01 & -10.24 \\ -8.87 & -7.08 \\ 3.88 & 7.71 \end{pmatrix}, \quad \mathbf{V}_2 \Lambda_2^{1/2} = \begin{pmatrix} .08 & .20 \\ .66 & .37 \\ 7.06 & 17.95 \\ 4.03 & 5.76 \\ 28.97 & -6.24 \\ 23.73 & -2.85 \\ 5.91 & 16.62 \end{pmatrix}.$$

The plot of the coordinates is given in Figure 16.11. For this data set, the factoring given by (16.62) and illustrated in Figure 16.11 is preferred because it plots both observation and variable points on the same scale. As previously noted, the factoring discussed in Example 16.3.4 and the factoring discussed at the beginning of this example would need adjustments in scaling in order to yield useful biplots. \square

PROBLEMS

- 16.1** In step 2 of the algorithm for metric scaling in Section 16.1.2, the matrix $\mathbf{B} = (b_{ij})$ is defined in terms of $\mathbf{A} = (a_{ij})$ as $b_{ij} = a_{ij} - \bar{a}_{i\cdot} - \bar{a}_{\cdot j} + \bar{a}_{\cdot\cdot}$. Show that b_{ij} in $\mathbf{B} = (\mathbf{I} - \frac{1}{n}\mathbf{J})\mathbf{A}(\mathbf{I} - \frac{1}{n}\mathbf{J})$ in (16.2) is equivalent to $b_{ij} = a_{ij} - \bar{a}_{i\cdot} - \bar{a}_{\cdot j} + \bar{a}_{\cdot\cdot}$.
- 16.2** Verify the result stated in step 2 of the algorithm in Section 16.1.2, namely, that there exists a q -dimensional configuration $\mathbf{z}_1, \mathbf{z}_2, \dots, \mathbf{z}_n$ such that $d_{ij} = \delta_{ij}$ if and only if \mathbf{B} is positive semidefinite of rank q . Use the following approach.
- Assuming the existence of $\mathbf{z}_1, \mathbf{z}_2, \dots, \mathbf{z}_n$ such that $\delta_{ij}^2 = d_{ij}^2 = (\mathbf{z}_i - \mathbf{z}_j)'(\mathbf{z}_i - \mathbf{z}_j)$, show that \mathbf{B} is positive semidefinite.
 - Assuming \mathbf{B} is positive semidefinite, show that there exist $\mathbf{z}_1, \mathbf{z}_2, \dots, \mathbf{z}_n$ such that $d_{ij}^2 = (\mathbf{z}_i - \mathbf{z}_j)'(\mathbf{z}_i - \mathbf{z}_j) = \delta_{ij}^2$.




Figure 16.11 Plot of $\mathbf{U}_2 \Lambda_2^{1/2}$ and $\mathbf{V}_2 \Lambda_2^{1/2}$ for the City Crime Data in Table 15.1.

- 16.3** (a) Show that $\mathbf{r} = \sum_{j=1}^b p_j \mathbf{c}_j$ in (16.20) is the same as $\mathbf{r} = (p_{1.}, p_{2.}, \dots, p_{a.})'$ in (16.9).
 (b) Show that $\mathbf{c}' = \sum_{i=1}^a p_i \mathbf{r}'_i$ in (16.21) is equivalent to $\mathbf{c}' = (p_{.1}, p_{.2}, \dots, p_{.b})$ in (16.10).
- 16.4** Show that $\mathbf{j}'\mathbf{r} = \mathbf{c}'\mathbf{j} = 1$ as in (16.22).
- 16.5** Show that the chi-square statistic in (16.26) is equal to that in (16.25).
- 16.6** (a) Show that the chi-square statistic in (16.27) is equal to that in (16.25).
 (b) Show that the chi-square statistic in (16.28) is equal to that in (16.25).
- 16.7** (a) Show the chi-square statistic in (16.29) is equal to that in (16.27).
 (b) Show the chi-square statistic in (16.30) is equal to that in (16.28).
- 16.8** (a) Show that $\mathbf{R} - \mathbf{j}\mathbf{c}' = \mathbf{D}_r^{-1}(\mathbf{P} - \mathbf{r}\mathbf{c}')$ as in (16.41).
 (b) Show that $\mathbf{C} - \mathbf{r}\mathbf{j}' = \mathbf{D}_c^{-1}(\mathbf{P} - \mathbf{r}\mathbf{c}')$ as in (16.42).
- 16.9** Show that if all the principal components were used, the distance between \mathbf{z}_i and \mathbf{z}_k would be the same as between \mathbf{y}_i and \mathbf{y}_k , as noted in Section 16.3.4.

- 16.10** (a) Show that $\mathbf{U}\mathbf{U}' = \mathbf{Y}_c\mathbf{S}^{-1}\mathbf{Y}'_c/(n - 1)$ as in (16.64).
 (b) Show that $\mathbf{H}\mathbf{H}' = (n - 1)\mathbf{S}$ as in (16.65).
- 16.11** Show that $(\mathbf{u}_i - \mathbf{u}_k)'(\mathbf{u}_i - \mathbf{u}_k) = (\mathbf{y}_i - \mathbf{y}_k)' \mathbf{S}^{-1} (\mathbf{y}_i - \mathbf{y}_k)$, as in (16.66).
- 16.12** In Table 16.15, we have road distances between major UK towns (Hand et al. 1994, p. 346). The towns are as follows:
 A = Aberdeen, B = Birmingham, C = Brighton, D = Bristol, E = Cardiff, F = Carlisle, G = Dover, H = Edinburgh, I = Fort William, J = Glasgow, K = Holyhead, L = Hull, M = Inverness, N = Leeds, O = Liverpool, P = London, Q = Manchester, R = Newcastle, S = Norwich, T = Nottingham, U = Penzance, V = Plymouth, W = Sheffield.
- (a) Find the matrix \mathbf{B} as in (16.2).
 - (b) Using the spectral decomposition, find the first two columns of the matrix \mathbf{Z} as in (16.4).
 - (c) Create a metric multidimensional scaling plot of the first two dimensions. What do you notice about the positions of the cities?
- 16.13** Zhang, Helander, and Drury (1996) analyzed a 43×43 similarity matrix for 43 descriptors of comfort, such as calm, tingling, restful, etc. For the similarity matrix, see the Wiley ftp site (Appendix C).
- (a) Carry out a metric multidimensional scaling analysis and plot the first two dimensions. What pattern is seen in the plot?
 - (b) For an alternative approach, carry out a cluster analysis of the configuration of points found in part (a), using Ward's method. Create a dendrogram of the cluster solution. How many clusters are indicated?
- 16.14** Use the politics data of Table 16.16 (Everitt 1987, Table 6.7). Two subjects assessed the degree of dissimilarity between World War II politicians. The data matrix represents the sum of the dissimilarities between the two subjects.
- (a) For $k = 6$, create an initial configuration of points by choosing twelve random observations taken from a multivariate normal distribution with mean vector $\mathbf{0}$ and covariance matrix \mathbf{I}_6 .
 - (b) Carry out a nonmetric multidimensional scaling analysis using the seeds found in part (a). Find the value of the STRESS statistic.
 - (c) Repeat parts (a) and (b) for $k = 1, \dots, 5$. Plot the STRESS values against the values of k . How many dimensions should be kept? Plot the final configuration of points with two dimensions.

Table 16.15 Road Distances Between Major UK Towns

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P	Q	R	S	T	U	V
B	431																					
C	611	185																				
D	515	88	170																			
E	535	108	205	47																		
F	232	198	378	282	302																	
G	595	206	78	210	245	398																
H	126	298	478	381	402	99	466															
I	159	407	587	491	511	209	608	133														
J	146	297	477	381	401	98	497	46	102													
K	461	153	333	237	205	228	353	328	438	327												
L	360	135	283	233	253	173	264	231	382	272	223											
M	106	456	636	540	560	258	657	158	65	172	488	431										
N	335	115	264	220	240	123	271	206	333	222	168	61	382									
O	358	102	282	185	205	125	302	225	335	224	105	130	383	75								
P	546	120	60	120	155	313	79	413	523	412	268	190	571	199	216							
Q	352	89	269	172	193	119	289	219	329	218	125	99	378	44	35	204						
R	237	202	350	300	320	57	352	108	241	152	268	142	268	93	175	285	144					
S	497	176	169	233	268	285	171	368	494	384	308	151	543	173	242	115	185	254				
T	402	55	196	146	166	190	217	273	400	289	177	93	449	73	109	131	70	160	120			
U	701	274	287	196	234	468	364	568	678	567	423	419	726	406	370	312	359	486	425	331		
V	630	203	216	125	162	397	292	496	607	496	352	348	655	335	300	241	287	414	354	260	78	
W	376	86	234	184	204	164	255	247	374	263	159	67	422	35	79	169	37	134	148	44	370	299

Table 16.16 Dissimilarity Matrix for World War II Politicians

Person	Mussolini	Eisenhower	Churchill	Stalin	Attlee	Franco	De Gaulle	Mao Tse	Truman	Chamberlain	Tito	
Hitler	0	5	11	15	8	17	5	10	16	17	12	16
Mussolini	5	0	14	16	13	18	3	11	18	18	14	17
Churchill	11	14	0	7	11	11	12	5	16	8	10	8
Eisenhower	15	16	7	0	16	16	14	8	17	6	7	12
Stalin	8	13	11	16	0	15	13	11	12	14	16	12
Attlee	17	18	11	16	15	0	16	12	16	12	9	13
Franco	5	3	12	14	13	16	0	9	17	16	10	12
De Gaulle	10	11	5	8	11	12	9	0	13	9	11	7
Mao Tse	16	18	16	17	12	16	17	13	0	12	17	10
Truman	17	18	8	6	14	12	16	9	12	0	9	11
Chamberlain	12	14	10	7	16	9	10	11	17	9	0	15
Tito	16	17	8	12	12	13	12	7	10	11	15	0

- (d) Repeat parts (a)–(c) using an initial configuration of points from a multivariate normal with different mean vector and covariance matrix from those in part (a). How many dimensions should be kept? Plot the final configuration of points with two dimensions. How does this solution compare to that in part (c)?
- (e) Repeat parts (a)–(c) using an initial configuration of points from a uniform distribution over $(0,1)$. How many dimensions should be kept? Plot the final configuration of points with two dimensions.
- (f) Repeat part (e) using as initial configuration of points the metric multidimensional scaling solution found by treating the ordinal measurements as continuous. How many dimensions should be kept? Plot the final configuration of points with two dimensions.

16.15 In Table 16.17 we have the months of birth and death for 1281 people (Andrews and Herzberg 1985, Table 71.2).

- (a) Find the correspondence matrix \mathbf{P} as in (16.8).
- (b) Find the matrices \mathbf{R} and \mathbf{C} , as in (16.15) and (16.19).
- (c) Perform a chi-square test for independence between birth month and death month.
- (d) Plot the row and column deviations as in Example 16.2.4.

16.16 In Table 16.18, we have a cross-classification of crimes in Norway in 1984 categorized by type and site (Clausen 1988, p. 9).

Table 16.17 Birth and Death Months of 1281 People

Death	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Total
Birth													
Jan	9	14	12	14	9	11	10	15	9	11	11	13	138
Feb	13	7	6	8	9	5	4	5	7	11	1	13	89
Mar	12	14	9	7	17	10	9	10	9	2	13	9	121
Apr	7	11	10	7	9	11	4	11	4	9	8	12	103
May	8	6	11	7	4	12	9	9	11	6	9	4	96
Jun	14	5	5	7	13	5	7	4	8	9	7	5	89
Jul	12	10	13	4	5	11	7	6	4	10	4	8	94
Aug	7	7	11	13	10	9	3	8	8	8	8	12	104
Sep	7	12	11	10	10	11	4	8	12	6	8	7	106
Oct	16	8	11	9	7	10	12	8	9	8	7	7	112
Nov	7	9	15	10	11	10	7	10	7	10	9	6	111
Dec	6	18	9	11	14	8	9	9	10	6	10	8	118
Total	118	121	123	107	118	113	85	103	98	96	95	104	1281

Table 16.18 Crimes by Type and Site

Part of Country	Burglary	Fraud	Vandalism	Total
Oslo area	395	2456	1758	4609
Mid-Norway	147	153	916	1216
North Norway	694	327	1347	2368
Total	1236	2936	4021	8193

- (a) Find the correspondence matrix \mathbf{P} as in (16.8).
- (b) Find the matrices \mathbf{R} and \mathbf{C} as in (16.15) and (16.19).
- (c) Perform a chi-square test for independence between type of crime and site.
- (d) Plot the row and column deviations as in Example 16.2.4.

16.17 In Table 16.19, we have a six-way contingency table (Andrews and Herzberg 1985, Table 34.1). Carry out a multiple correspondence analysis.

- (a) Set up an indicator matrix \mathbf{G} and find the Burt matrix $\mathbf{G}'\mathbf{G}$.
- (b) Perform a correspondence analysis on the Burt matrix found in part (a) and plot the coordinates.
- (c) What associations are present?

16.18 Use the protein consumption data of Table 15.7.

- (a) Create a biplot using the principal components approach in (16.53) or (16.57).
- (b) Create a biplot using the singular value decomposition approach with the factoring as in (16.61).
- (c) Create a biplot using the singular value decomposition approach with the factoring as in (16.62).
- (d) Which of the three biplots best represents the data?

16.19 Use the perception data of Table 13.1.

- (a) Create a biplot using the principal components approach in (16.53) or (16.57).
- (b) Create a biplot using the singular value decomposition approach with the factoring as in (16.61).
- (c) Create a biplot using the singular value decomposition approach with the factoring as in (16.62).
- (d) Which of the three biplots best represents the data?

Table 16.19 Byssinosis Data

Smoking	Gender	Years in job	Suffer Byssi- nosis	Race					
				Other			White		
				High dust	Low dust	Med dust	High dust	Low dust	Med dust
Non-smoker	Female	≤ 10	No	24	301	142	4	169	54
			Yes	1	4	3	0	2	1
		≥ 20	No	0	3	2	2	340	187
			Yes	0	0	0	0	2	3
		10–20	No	0	4	4	0	90	30
			Yes	0	0	0	0	1	0
	Male	≤ 10	No	75	122	47	16	134	35
			Yes	6	1	1	0	0	0
		≥ 20	No	15	23	1	47	182	39
			Yes	3	0	0	5	3	0
		10–20	No	9	7	0	8	58	16
			Yes	1	0	0	2	0	1
Smoker	Female	≤ 10	No	22	260	145	5	180	93
			Yes	2	3	2	0	3	1
		≥ 20	No	1	2	0	1	176	91
			Yes	0	0	0	0	3	3
		10–20	No	0	3	4	0	94	33
			Yes	0	0	0	0	2	1
	Male	≤ 10	No	139	242	88	37	258	74
			Yes	25	3	0	3	2	0
		≥ 20	No	31	45	1	77	495	141
			Yes	10	0	0	31	12	1
		10–20	No	30	33	5	21	187	50
			Yes	8	0	0	8	1	1

16.20 Use the cork data of Table 6.21.

- (a) Create a biplot using the principal components approach in (16.53) or (16.57).
- (b) Create a biplot using the singular value decomposition approach with the factoring as in (16.61).
- (c) Create a biplot using the singular value decomposition approach with the factoring as in (16.62).
- (d) Which of the three biplots best represents the data?

Appendix A

Tables

Table A.1: Upper Percentiles for $\sqrt{b_1}$

Table A.2: Coefficients for Transforming $\sqrt{b_1}$ to a Standard Normal

Table A.3: Percentiles for b_2

Table A.4: Percentiles for D'Agostino's Test for Normality

Table A.5: Upper Percentiles for $b_{1,p}$ and Upper and Lower Percentiles for $b_{2,p}$

Table A.6: Upper Percentiles for Test of Single Multivariate Normal Outlier

Table A.7: Upper Percentage Points of Hotelling's T^2 Distribution

Table A.8: Bonferroni t -Values, $t_{\alpha/2k,\nu}$, $\alpha = 0.05$

Table A.9: Lower Critical Values of Wilks' Λ , $\alpha = 0.05$

Table A.10: Upper Critical Values for Roy's Test, $\alpha = .05$

Table A.11: Upper Critical Values of Pillai's Statistic $V^{(s)}$, $\alpha = .05$

Table A.12: Upper Critical Values for the Lawley-Hotelling Statistic, $\alpha = .05$

Table A.13: Orthogonal Polynomial Contrasts

Table A.14: Test for Equal Covariance Matrices, $\alpha = 0.05$

Table A.15: Test for Independence of p Variables

*

Table A.1 Upper Percentiles for $\sqrt{b_1}$

$$\sqrt{b_1} = \frac{\sqrt{n} \sum_{i=1}^n (y_i - \bar{y})^3}{[\sum_{i=1}^n (y_i - \bar{y})^2]^{3/2}}$$

The sampling distribution of $\sqrt{b_1}$ is symmetric about zero, and the lower percentage points corresponding to negative skewness are given by the negative of the table values. Reject the hypothesis of normality if $\sqrt{b_1}$ is greater than the table value or less than the negative of the table value.

n	Upper Percentiles					
	10	5	2.5	1	0.5	0.1
4	.831	.987	1.070	1.120	1.137	1.151
5	.821	1.049	1.207	1.337	1.396	1.464
6	.795	1.042	1.239	1.429	1.531	1.671
7	.782	1.018	1.230	1.457	1.589	1.797
8	.765	.998	1.208	1.452	1.605	1.866
9	.746	.977	1.184	1.433	1.598	1.898
10	.728	.954	1.159	1.407	1.578	1.906
11	.710	.931	1.134	1.381	1.553	1.899
12	.693	.910	1.109	1.353	1.526	1.882
13	.677	.890	1.085	1.325	1.497	1.859
14	.662	.870	1.061	1.298	1.468	1.832
15	.648	.851	1.039	1.272	1.440	1.803
16	.635	.834	1.018	1.247	1.412	1.773
17	.622	.817	.997	1.222	1.385	1.744
18	.610	.801	.978	1.199	1.359	1.714
19	.599	.786	.960	1.176	1.334	1.685
20	.588	.772	.942	1.155	1.310	1.657
21	.578	.758	.925	1.134	1.287	1.628
22	.568	.746	.909	1.114	1.265	1.602
23	.559	.733	.894	1.096	1.243	1.575
24	.550	.722	.880	1.078	1.223	1.550
25	.542	.710	.866	1.060	1.203	1.526

Table A.2 Coefficients for Transforming $\sqrt{b_1}$ to a Standard Normal

n	δ	$1/\lambda$	n	δ	$1/\lambda$	n	δ	$1/\lambda$
8	5.563	0.3030	41	3.073	0.9653	98	3.913	1.1006
9	4.260	0.4080	42	3.087	0.9702	100	3.940	1.1028
10	3.734	0.4794	43	3.102	0.9750	105	4.009	1.1080
11	3.447	0.5339	44	3.117	0.9795	110	4.076	1.1128
12	3.270	0.5781	45	3.131	0.9840	115	4.142	1.1172
13	3.151	0.6153	46	3.146	0.9882	120	4.207	1.1212
14	3.069	0.6473	47	3.161	0.9923	125	4.272	1.1250
15	3.010	0.6753	48	3.176	0.9963	130	4.336	1.1285
16	2.968	0.7001	49	3.192	1.0001	135	4.398	1.1318
17	2.937	0.7224	50	3.207	1.0038	140	4.460	1.1348
18	2.915	0.7426	52	3.237	1.0108	145	4.521	1.1377
19	2.900	0.7610	54	3.268	1.0174	150	4.582	1.1403
20	2.890	0.7779	56	3.298	1.0235	155	4.641	1.1428
21	2.884	0.7934	58	3.329	1.0293	160	4.700	1.1452
22	2.882	0.8078	60	3.359	1.0348	165	4.758	1.1474
23	2.882	0.8211	62	3.389	1.0400	170	4.816	1.1496
24	2.884	0.8336	64	3.420	1.0449	175	4.873	1.1516
25	2.889	0.8452	66	3.450	1.0495	180	4.929	1.1535
26	2.895	0.8561	68	3.480	1.0540	185	1.985	1.1553
27	2.902	0.8664	70	3.510	1.0581	190	5.040	1.1570
28	2.910	0.8760	72	3.540	1.0621	195	5.094	1.1586
29	2.920	0.8851	74	3.569	1.0659	200	5.148	1.1602
30	2.930	0.8938	76	3.599	1.0695	205	5.202	1.1616
31	2.941	0.9020	78	3.628	1.0730	210	5.255	1.1631
32	2.952	0.9097	80	3.657	1.0763	215	5.307	1.1644
33	2.964	0.9171	82	3.686	1.0795	220	5.359	1.1657
34	2.977	0.9241	84	3.715	1.0825	225	5.410	1.1669
35	2.990	0.9308	86	3.744	1.0854	230	5.461	1.1681
36	3.003	0.9372	88	3.772	1.0882	235	5.511	1.1693
37	3.016	0.9433	90	3.801	1.0909	240	5.561	1.1704
38	3.030	0.9492	92	3.829	1.0934	245	5.611	1.1714
39	3.044	0.9548	94	3.857	1.0959	250	5.660	1.1724
40	3.058	0.9601	96	3.885	1.0983			

Values of δ and $1/\lambda$ are such that $g(\sqrt{b_1}) = \delta \sinh^{-1}(\sqrt{b_1}/\lambda)$ is approximately $N(0, 1)$.

Table A.3 Percentiles for b_2

Upper and lower percentiles for

$$b_2 = \frac{n \sum_{i=1}^n (y_i - \bar{y})^4}{[\sum_{i=1}^n (y_i - \bar{y})^2]^2},$$

the sample coefficient of kurtosis. Reject the hypothesis of normality if b_2 is greater than an upper percentile or less than a lower percentile.

Sample Size	Percentiles											
	1	2	2.5	5	10	20	80	90	95	97.5	98	99
7	1.25	1.30	1.34	1.41	1.53	1.70	2.78	3.20	3.55	3.85	3.93	4.23
8	1.31	1.37	1.40	1.46	1.58	1.75	2.84	3.31	3.70	4.09	4.20	4.53
9	1.35	1.42	1.45	1.53	1.63	1.80	2.98	3.43	3.86	4.28	4.41	4.82
10	1.39	1.45	1.49	1.56	1.68	1.85	3.01	3.53	3.95	4.40	4.55	5.00
12	1.46	1.52	1.56	1.64	1.76	1.93	3.06	3.55	4.05	4.56	4.73	5.20
15	1.55	1.61	1.64	1.72	1.84	2.01	3.13	3.62	4.13	4.66	4.85	5.30
20	1.65	1.71	1.74	1.82	1.95	2.13	3.21	3.68	4.17	4.68	4.87	5.36
25	1.72	1.79	1.83	1.91	2.03	2.20	3.23	3.68	4.16	4.65	4.82	5.30
30	1.79	1.86	1.90	1.98	2.10	2.26	3.25	3.68	4.11	4.59	4.75	5.21
35	1.84	1.91	1.95	2.03	2.14	2.31	3.27	3.68	4.10	4.53	4.68	5.13
40	1.89	1.96	1.98	2.07	2.19	2.34	3.28	3.67	4.06	4.46	4.61	5.04
45	1.93	2.00	2.03	2.11	2.22	2.37	3.28	3.65	4.00	4.39	4.52	4.94
50	1.95	2.03	2.06	2.15	2.25	2.41	3.28	3.62	3.99	4.33	4.45	4.88

Table A.4 Percentiles for D'Agostino's Test for Normality

Upper and lower percentiles for the statistic

$$Y = \frac{\sqrt{n}[D - (2\sqrt{\pi})^{-1}]}{.02998598},$$

where

$$D = \frac{\sum_{i=1}^n [i - \frac{1}{2}(n+1)] y_{(i)}}{\sqrt{n^3} \sum_{i=1}^n (y_i - \bar{y})^2}$$

and the observations y_1, y_2, \dots, y_n are ordered as $y_{(1)} \leq y_{(2)} \leq \dots \leq y_{(n)}$. Reject the hypothesis of normality if Y is greater than an upper percentile or less than a lower percentile.

n	Percentiles of Y									
	.5	1.0	2.5	5	10	90	95	97.5	99	99.5
10	-4.66	-4.06	-3.25	-2.62	-1.99	.149	.235	.299	.356	.385
12	-4.63	-4.02	-3.20	-2.58	-1.94	.237	.329	.381	.440	.479
14	-4.57	-3.97	-3.16	-2.53	-1.90	.308	.399	.460	.515	.555
16	-4.52	-3.92	-3.12	-2.50	-1.87	.367	.459	.526	.587	.613
18	-4.47	-3.87	-3.08	-2.47	-1.85	.417	.515	.574	.636	.667
20	-4.41	-3.83	-3.04	-2.44	-1.82	.460	.565	.628	.690	.720
22	-4.36	-3.78	-3.01	-2.41	-1.81	.497	.609	.677	.744	.775
24	-4.32	-3.75	-2.98	-2.39	-1.79	.530	.648	.720	.783	.822
26	-4.27	-3.71	-2.96	-2.37	-1.77	.559	.682	.760	.827	.867
28	-4.23	-3.68	-2.93	-2.35	-1.76	.586	.714	.797	.868	.910
30	-4.19	-3.64	-2.91	-2.33	-1.75	.610	.743	.830	.906	.941
32	-4.16	-3.61	-2.88	-2.32	-1.73	.631	.770	.862	.942	.983
34	-4.12	-3.59	-2.86	-2.30	-1.72	.651	.794	.891	.975	1.02
36	-4.09	-3.56	-2.85	-2.29	-1.71	.669	.816	.917	1.00	1.05
38	-4.06	-3.54	-2.83	-2.28	-1.70	.686	.837	.941	1.03	1.08
40	-4.03	-3.51	-2.81	-2.26	-1.70	.702	.857	.964	1.06	1.11
42	-4.00	-3.49	-2.80	-2.25	-1.69	.716	.875	.986	1.09	1.14
44	-3.98	-3.47	-2.78	-2.24	-1.68	.730	.892	1.01	1.11	1.17
46	-3.95	-3.45	-2.77	-2.23	-1.67	.742	.908	1.02	1.13	1.19
48	-3.93	-3.43	-2.75	-2.22	-1.67	.754	.923	1.04	1.15	1.22
50	-3.91	-3.41	-2.74	-2.21	-1.66	.765	.937	1.06	1.18	1.24
60	-3.81	-3.34	-2.68	-2.17	-1.64	.812	.997	1.13	1.26	1.34
70	-3.73	-3.27	-2.64	-2.14	-1.61	.849	1.05	1.19	1.33	1.42
80	-3.67	-3.22	-2.60	-2.11	-1.59	.878	1.08	1.24	1.39	1.48
90	-3.61	-3.17	-2.57	-2.09	-1.58	.902	1.12	1.28	1.44	1.54
100	-3.57	-3.14	-2.54	-2.07	-1.57	.923	1.14	1.31	1.48	1.59
150	-3.409	-3.009	-2.452	-2.004	-1.520	.990	1.233	1.423	1.623	1.746
200	-3.302	-2.922	-2.391	-1.960	-1.491	1.032	1.290	1.496	1.715	1.853
250	-3.227	-2.861	-2.348	-1.926	-1.471	1.060	1.328	1.545	1.779	1.927

Table A.5 Upper Percentiles for $b_{1,p}$ and Upper and Lower Percentiles for $b_{2,p}$

Reject the hypothesis of multivariate normality if $b_{1,p}$ is greater than table value. Reject if $b_{2,p}$ is greater than upper percentile or if $b_{2,p}$ is less than lower percentile. The statistics $b_{1,p}$ and $b_{2,p}$ are defined in Section 4.4.2.

n	Percentiles					Percentiles									
	90	92.5	95	97.5	99	99.9	n	1	2.5	5	10	90	95	97.5	99
<i>p = 2,</i>															
Upper Percentiles for $b_{1,p}$															
10	2.994	3.263	3.694	4.294	5.194	6.994	10	4.580	4.722	4.887	5.057	8.606	9.203	9.781	10.378
12	2.681	2.944	3.319	3.931	4.938	6.744	12	4.732	4.899	5.053	5.232	8.947	9.593	10.150	10.881
14	2.419	2.669	3.031	3.619	4.581	6.419	14	4.842	5.015	5.179	5.358	9.162	9.769	10.375	11.159
16	2.219	2.444	2.775	3.337	4.231	6.062	16	4.977	5.149	5.318	5.482	9.331	9.941	10.562	11.387
18	2.050	2.256	2.556	3.100	3.962	5.737	18	5.045	5.219	5.382	5.555	9.403	10.005	10.628	11.478
20	1.894	2.081	2.356	2.881	3.669	5.425	20	5.175	5.262	5.533	5.717	9.469	10.114	10.691	11.609
25	1.581	1.744	1.969	2.438	3.106	4.719	25	5.351	5.525	5.689	5.871	9.503	10.159	10.584	11.628
30	1.363	1.513	1.687	2.094	2.681	4.238	30	5.518	5.692	5.855	6.038	9.516	10.156	10.556	11.594
40	1.050	1.181	1.319	1.606	2.087	3.369	40	5.703	5.871	6.139	6.229	9.497	10.109	10.563	11.453
50	.862	.969	1.069	1.306	1.744	2.706	50	5.909	6.083	6.239	6.403	9.453	9.987	10.372	11.181
60	.731	.819	.906	1.094	1.444	2.200	60	6.015	6.189	6.335	6.505	9.401	9.889	10.250	10.994
70	.631	.725	.794	.937	1.244	1.863	70	6.139	6.290	6.437	6.602	9.356	9.781	10.106	10.753
80	.544	.637	.694	.812	1.056	1.587	80	6.223	6.372	6.539	6.683	9.309	9.694	9.981	10.537
90	.487	.569	.638	.725	.919	1.400	90	6.332	6.475	6.622	6.749	9.256	9.688	9.885	10.325
100	.438	.506	.581	.656	.831	1.231	100	6.389	6.521	6.665	6.793	9.210	9.556	9.806	10.188
150	.281	.344	.400	.444	.531	.794	150	6.615	6.749	6.858	6.972	9.027	9.300	9.475	10.253
200	.219	.269	.300	.331	.394	.569	200	6.761	6.889	6.979	7.083	8.919	9.141	9.269	9.506
300	.144	.169	.209	.225	.256	.369	300	6.949	7.052	7.142	7.245	8.776	8.916	9.031	9.219
400	.116	.129	.141	.166	.197	.275	400	7.079	7.171	7.252	7.342	8.664	8.787	8.917	9.061
600	.077	.085	.094	.110	.131	.183	600	7.232	7.295	7.369	7.464	8.547	8.647	8.749	8.874

(continued)

Table A.5 (Continued)

$p = 2$, Upper Percentiles for $b_{1,p}$										$p = 2$, Upper and Lower Percentiles for $b_{2,p}$									
n	Percentiles					n	Percentiles					n	Percentiles						
	90	92.5	95	97.5	99		1	2.5	5	10	90		95	97.5	99				
800	.058	.064	.071	.083	.099	.137	800	7.304	7.372	7.451	7.536	8.472	8.562	8.641	8.747				
1000	.046	.051	.057	.066	.079	.110	1000	7.367	7.433	7.504	7.585	8.419	8.497	8.569	8.656				
1500	.031	.034	.038	.044	.053	.074	1500	7.460	7.537	7.595	7.661	8.339	8.405	8.463	8.532				
2500	.019	.021	.023	.027	.032	.044	2000	7.535	7.599	7.649	7.707	8.293	8.351	8.401	8.461				
3000	.016	.017	.019	.022	.027	.037	2500	7.588	7.641	7.686	7.738	8.262	8.314	8.359	8.412				
4000	.012	.013	.014	.017	.020	.028	3000	7.624	7.673	7.714	7.760	8.240	8.286	8.327	8.376				
5000	.009	.010	.011	.013	.016	.022	4000	7.674	7.716	7.752	7.793	8.207	8.248	8.284	8.326				
							5000	7.709	7.746	7.778	7.714	8.186	8.222	8.254	8.291				
$p = 3$, Upper Percentiles for $b_{1,p}$										$p = 3$, Upper and Lower Percentiles for $b_{2,p}$									
n	Percentiles					n	Percentiles					n	Percentiles						
	90	92.5	95	97.5	99	99.9	1	2.5	5	10	90	95	97.5	99					
10	6.0	6.5	6.9	7.7	8.8	11.5	10	10.0	10.2	10.4	10.7	14.0	14.4	15.0	15.6				
12	5.5	5.9	6.4	7.1	8.1	10.5	12	10.2	10.4	10.7	11.0	14.7	15.2	15.9	16.4				
14	5.0	5.4	5.9	6.5	7.4	9.7	14	10.4	10.6	10.9	11.3	15.1	15.8	16.5	17.1				
16	4.6	4.9	5.4	6.1	6.8	8.9	16	10.5	10.8	11.1	11.5	15.4	16.1	16.8	17.5				
18	4.2	4.6	5.1	5.6	6.4	8.3	18	10.7	11.0	11.3	11.6	15.5	16.4	17.1	17.8				
20	3.9	4.2	4.7	5.3	6.0	7.7	20	10.8	11.1	11.4	11.8	15.7	16.5	17.2	18.0				
25	3.3	3.5	3.9	4.5	5.2	6.5	25	11.1	11.4	11.8	12.1	15.9	16.7	17.4	18.2				
30	2.8	3.0	3.3	3.9	4.4	5.6	30	11.3	11.6	12.0	12.3	16.0	16.7	17.5	18.3				
40	2.2	2.4	2.7	3.0	3.5	4.2	40	11.7	12.0	12.4	12.7	16.1	16.7	17.4	18.2				
50	1.7	1.9	2.2	2.4	2.8	3.4	50	11.9	12.3	12.6	12.9	16.1	16.7	17.3	18.0				

(continued)

Table A.5 (Continued)

$p = 3$, Upper Percentiles for $b_{1,p}$										$p = 3$, Upper and Lower Percentiles for $b_{2,p}$									
n	Percentiles					n	Percentiles					n	Percentiles						
	90	92.5	95	97.5	99		1	2.5	5	10	90		95	97.5	99				
60	1.5	1.6	1.8	2.0	2.4	2.9	60	12.1	12.5	12.8	13.1	16.1	16.6	17.2	17.9				
70	1.3	1.4	1.5	1.7	2.0	2.5	70	12.3	12.6	13.0	13.2	16.1	16.6	17.1	17.7				
80	1.13	1.2	1.3	1.5	1.7	2.2	80	12.4	12.8	13.1	13.3	16.1	16.5	17.0	17.6				
90	1.01	1.08	1.16	1.3	1.5	1.9	90	12.5	12.9	13.2	13.5	16.0	16.5	16.9	17.5				
100	.92	.97	1.05	1.18	1.3	1.7	100	12.6	13.0	13.3	13.5	16.0	16.4	16.8	17.4				
150	.62	.66	.71	.80	.90	1.15	150	13.0	13.3	13.6	13.8	15.9	16.2	16.5	17.0				
200	.47	.50	.54	.60	.68	.87	200	13.2	13.5	13.8	14.0	15.8	16.1	16.3	16.8				
300	.32	.33	.36	.40	.46	.58	300	13.6	13.8	14.0	14.2	15.7	15.9	16.1	16.5				
400	.237	.252	.272	.30	.34	.44	400	13.7	13.9	14.1	14.3	15.6	15.8	16.0	16.3				
600	.159	.168	.182	.203	.230	.294	600	13.9	14.1	14.3	14.4	15.51	15.67	15.81	15.97				
800	.119	.127	.137	.153	.173	.221	800	14.1	14.2	14.3	14.5	15.45	15.59	15.71	15.85				
1000	.095	.1010	.109	.122	.139	.177	1000	14.17	14.30	14.41	14.53	15.41	15.53	15.64	15.77				
1500	.064	.068	.073	.082	.093	.118	1500	14.33	14.43	14.52	14.62	15.34	15.44	15.53	15.63				
2000	.048	.051	.055	.061	.069	.089	2000	14.42	14.51	14.58	14.67	15.30	15.39	15.46	15.55				
3000	.032	.034	.037	.041	.046	.059	3000	14.53	14.60	14.66	14.73	15.25	15.32	15.38	15.45				
4000	.024	.025	.027	.031	.035	.044	4000	14.59	14.65	14.71	14.77	15.21	15.28	15.33	15.39				
5000	.019	.020	.022	.025	.028	.035	5000	14.63	14.69	14.74	14.80	15.19	15.25	15.30	15.35				

$p = 4$,
Upper Percentiles for $b_{1,p}$

$p = 4$,
Upper and Lower Percentiles for $b_{2,p}$

$p = 4$, Upper Percentiles for $b_{1,p}$										$p = 4$, Upper and Lower Percentiles for $b_{2,p}$									
n	Percentiles					n	Percentiles					n	Percentiles						
	90	92.5	95	97.5	99		1	2.5	5	10	90		95	97.5	99				
10	11.1	11.6	12.2	13.3	15.3	17.9	10	17.0	17.3	17.6	17.8	21.5	22.4	23.0	24.0				
12	10.1	10.6	11.2	12.2	13.9	16.2	12	17.4	17.7	18.0	18.3	22.3	23.3	24.2	25.4				

(continued)

Table A.5 (Continued)

n	Upper Percentiles for $b_{1,p}$					Upper and Lower Percentiles for $b_{2,p}$									
	90	92.5	95	97.5	99	99.9	n	1	2.5	5	10	90	95	97.5	99
$p = 4,$															
14	9.2	9.7	10.2	11.2	12.7	14.8	14	17.7	18.0	18.3	18.6	23.0	24.0	25.0	26.1
16	8.4	8.8	9.4	10.3	11.6	13.6	16	18.0	18.2	18.6	18.9	23.4	24.4	25.4	26.6
18	7.7	8.0	8.7	9.5	10.7	12.6	18	18.2	18.4	18.8	19.2	23.8	24.7	25.8	26.9
20	7.0	7.4	8.0	8.8	9.9	11.6	20	18.4	18.6	19.0	19.4	24.0	25.0	26.1	27.1
25	5.9	6.2	6.6	7.1	8.1	9.7	25	18.8	19.1	19.5	19.8	24.5	25.4	26.4	27.3
30	5.0	5.3	5.6	6.0	6.8	8.1	30	19.1	19.4	19.8	20.2	24.7	25.5	26.6	27.4
40	3.9	4.1	4.3	4.6	5.2	6.2	40	19.6	19.9	20.3	21.0	25.0	25.7	26.7	27.4
50	3.1	3.3	3.5	3.8	4.2	5.0	50	20.0	20.3	20.6	21.0	25.1	25.7	26.6	27.3
60	2.7	2.8	2.9	3.2	3.5	4.2	60	20.2	20.5	20.9	21.3	25.14	25.7	26.6	27.2
70	2.3	2.4	2.5	2.8	3.0	3.7	70	20.4	20.7	21.0	21.5	25.15	25.7	26.5	27.0
80	2.0	2.1	2.2	2.4	2.7	3.2	80	20.6	21.0	21.2	21.7	25.15	25.6	26.4	26.9
90	1.81	1.89	2.0	2.2	2.4	2.9	90	20.8	21.1	21.4	21.8	25.14	25.6	26.3	26.8
100	1.64	1.71	1.81	1.97	2.2	2.6	100	20.9	21.2	21.5	21.9	25.12	25.6	26.2	26.7
150	1.11	1.16	1.22	1.33	1.46	1.76	150	21.4	21.7	22.0	22.33	25.03	25.42	25.9	26.3
200	.84	.87	.92	1.00	1.10	1.33	200	21.7	22.0	22.2	22.57	24.95	25.29	25.6	26.0
300	.56	.59	.62	.67	.74	.89	300	22.1	22.33	22.57	22.85	24.83	25.11	25.3	25.7
400	.42	.44	.47	.51	.56	.67	400	22.3	22.56	22.77	23.02	24.75	24.99	25.20	25.46
600	.282	.295	.31	.34	.37	.45	600	22.63	22.83	23.01	23.21	24.63	24.83	25.01	25.21
800	.212	.222	.234	.255	.280	.34	800	22.82	22.99	23.15	23.32	24.56	24.74	24.89	25.06
1000	.170	.177	.188	.204	.224	.271	1000	22.94	23.10	23.24	23.40	24.51	24.67	24.80	24.96
1500	.113	.118	.125	.136	.150	.181	1500	23.14	23.27	23.38	23.51	24.42	24.55	24.66	24.79
2000	.085	.089	.094	.102	.112	.136	2000	23.26	23.37	23.47	23.58	24.37	24.48	24.58	24.69
3000	.057	.059	.063	.068	.075	.091	3000	23.40	23.49	23.57	23.66	24.31	24.40	24.48	24.57
4000	.043	.045	.047	.051	.056	.068	4000	23.48	23.56	23.63	23.71	24.27	24.35	24.42	24.50
5000	.034	.039	.038	.041	.045	.054	5000	23.54	23.61	23.67	23.74	24.24	24.31	24.37	24.45

Table A.6 Upper Percentiles for Test of Single Multivariate Normal Outlier

Upper percentage points for the test statistic

$$D_{(n)}^2 = \max_{1 \leq i \leq n} (\mathbf{y}_i - \bar{\mathbf{y}})' \mathbf{S}^{-1} (\mathbf{y}_i - \bar{\mathbf{y}}).$$

This tests for a single outlier in a sample of size n from a multivariate normal distribution. Reject and conclude that the outlier is significant if $D_{(n)}^2$ exceeds the table value.

n	$p = 2$		$p = 3$		$p = 4$		$p = 5$	
	$\alpha = .05$	$\alpha = .01$						
5	3.17	3.19						
6	4.00	4.11	4.14	4.16				
7	4.71	4.95	5.01	5.10	5.12	5.14		
8	5.32	5.70	5.77	5.97	6.01	6.09	6.11	6.12
9	5.85	6.37	6.43	6.76	6.80	6.97	7.01	7.08
10	6.32	6.97	7.01	7.47	7.50	7.79	7.82	7.98
12	7.10	8.00	7.99	8.70	8.67	9.20	9.19	9.57
14	7.74	8.84	8.78	9.71	9.61	10.37	10.29	10.90
16	8.27	9.54	9.44	10.56	10.39	11.36	11.20	12.02
18	8.73	10.15	10.00	11.28	11.06	12.20	11.96	12.98
20	9.13	10.67	10.49	11.91	11.63	12.93	12.62	13.81
25	9.94	11.73	11.48	13.18	12.78	14.40	13.94	15.47
30	10.58	12.54	12.24	14.14	13.67	15.51	14.95	16.73
35	11.10	13.20	12.85	14.92	14.37	16.40	15.75	17.73
40	11.53	13.74	13.36	15.56	14.96	17.13	16.41	18.55
45	11.90	14.20	13.80	16.10	15.46	17.74	16.97	19.24
50	12.23	14.60	14.18	16.56	15.89	18.27	17.45	19.83
100	14.22	16.95	16.45	19.26	18.43	21.30	20.26	23.17
200	15.99	18.94	18.42	21.47	20.59	23.72	22.59	25.82
500	18.12	21.22	20.75	23.95	23.06	26.37	25.21	28.62

(continued)

Table A.7 Upper Percentage Points of Hotelling's T^2 Distribution

Degrees of Freedom, ν	$p = 1$	$p = 2$	$p = 3$	$p = 4$	$p = 5$	$p = 6$	$p = 7$	$p = 8$	$p = 9$	$p = 10$
2	18.513									
3	10.128	57.000								
4	7.709	25.472	114.986							
5	6.608	17.361	46.383	192.468						
6	5.987	13.887	29.661	72.937	289.446					
7	5.591	12.001	22.720	44.718	105.157	405.920				
8	5.318	10.828	19.028	33.230	62.561	143.050	541.890			
9	5.117	10.033	16.766	27.202	45.453	83.202	186.622	697.356		
10	4.965	9.459	15.248	23.545	36.561	59.403	106.649	235.873	872.317	
11	4.844	9.026	14.163	21.108	31.205	47.123	75.088	132.903	290.806	1066.774
12	4.747	8.689	13.350	19.376	27.656	39.764	58.893	92.512	161.967	351.421
13	4.667	8.418	12.719	18.086	25.145	34.911	49.232	71.878	111.676	193.842
14	4.600	8.197	12.216	17.089	23.281	31.488	42.881	59.612	86.079	132.582
15	4.543	8.012	11.806	16.296	21.845	28.955	38.415	51.572	70.907	101.499
16	4.494	7.856	11.465	15.651	20.706	27.008	35.117	45.932	60.986	83.121
17	4.451	7.722	11.177	15.117	19.782	25.467	32.588	41.775	54.041	71.127
18	4.414	7.606	10.931	14.667	19.017	24.219	30.590	38.592	48.930	62.746
19	4.381	7.504	10.719	14.283	18.375	23.189	28.975	36.082	45.023	56.587
20	4.351	7.415	10.533	13.952	17.828	22.324	27.642	34.054	41.946	51.884
21	4.325	7.335	10.370	13.663	17.336	21.588	26.525	32.384	39.463	48.184
22	4.301	7.264	10.225	13.409	16.945	20.954	25.576	30.985	37.419	45.202
23	4.279	7.200	10.095	13.184	16.585	20.403	24.759	29.798	35.709	42.750
24	4.260	7.142	9.979	12.983	16.265	19.920	24.049	28.777	34.258	40.699
25	4.242	7.089	9.874	12.803	15.981	19.492	23.427	27.891	33.013	38.961
26	4.225	7.041	9.779	12.641	15.726	19.112	22.878	27.114	31.932	37.469

Table A.7 (Continued)

Degrees of Freedom, ν	$p = 1$	$p = 2$	$p = 3$	$p = 4$	$p = 5$	$p = 6$	$p = 7$	$p = 8$	$p = 9$	$p = 10$
27	4.210	6.997	9.692	12.493	15.496	18.770	22.388	26.428	30.985	36.176
28	4.196	6.957	9.612	12.359	15.287	18.463	21.950	25.818	30.149	35.043
29	4.183	6.919	9.539	12.236	15.097	18.184	21.555	25.272	29.407	34.044
30	4.171	6.885	9.471	12.123	14.924	17.931	21.198	24.781	28.742	33.156
35	4.121	6.744	9.200	11.674	14.240	16.944	19.823	22.913	26.252	29.881
40	4.085	6.642	9.005	11.356	13.762	16.264	18.890	21.668	24.624	27.783
45	4.057	6.564	8.859	11.118	13.409	15.767	18.217	20.781	23.477	26.326
50	4.034	6.503	8.744	10.934	13.138	15.388	17.709	20.117	22.627	25.256
55	4.016	6.454	8.652	10.787	12.923	15.090	17.311	19.600	21.972	24.437
60	4.001	6.413	8.577	10.668	12.748	14.850	16.992	19.188	21.451	23.790
70	3.978	6.350	8.460	10.484	12.482	14.485	16.510	18.571	20.676	22.834
80	3.960	6.303	8.375	10.350	12.289	14.222	16.165	18.130	20.127	22.162
90	3.947	6.267	8.309	10.248	12.142	14.022	15.905	17.801	19.718	21.663
100	3.936	6.239	8.257	10.167	12.027	13.867	15.702	17.544	19.401	21.279
110	3.927	6.216	8.215	10.102	11.934	13.741	15.540	17.340	19.149	20.973
120	3.920	6.196	8.181	10.048	11.838	13.639	15.407	17.172	18.943	20.725
150	3.904	6.155	8.105	9.931	11.693	13.417	15.121	16.814	18.504	20.196
200	3.888	6.113	8.031	9.817	11.531	13.202	14.845	16.469	18.083	19.692
400	3.865	6.052	7.922	9.650	11.297	12.890	14.447	15.975	17.484	18.976
1000	3.851	6.015	7.857	9.552	11.160	12.710	14.217	15.692	17.141	18.570
∞	3.841	5.991	7.815	9.488	11.070	12.592	14.067	15.507	16.919	18.307

(continued)

(continued)

Table A.7 (Continued)

Degrees of Freedom, ν	$p = 1$	$p = 2$	$p = 3$	$p = 4$	$p = 5$	$p = 6$	$p = 7$	$p = 8$	$p = 9$	$p = 10$
2	98.503	297.000	594.997	992.494	147.283	229.679	1489.489	329.433	2085.984	
3	34.116	82.177	147.283	229.679	1489.489	329.433	2085.984	446.571	2781.978	
4	21.198	50.652	111.839	172.908	155.219	205.293	581.106	3577.472		
5	16.258	45.000	75.125	54.890	98.703	128.067	262.076	733.045	4472.464	
6	13.745	31.857	25.491	44.838	72.882	161.015	325.576	902.392	5466.956	
7	12.246	21.821	39.118	32.598	39.454	54.150	73.676	109.441	167.499	
8	11.259	19.460	50.652	44.838	38.533	58.618	93.127	115.640	197.555	
9	10.561	17.826	28.466	20.834	28.857	36.246	48.472	65.483	90.433	
10	10.044	16.631	25.637	19.960	19.867	27.060	33.672	44.240	58.241	
11	9.646	15.722	23.588	22.041	31.171	43.745	62.114	90.907	140.429	
12	9.330	15.008	20.834	17.861	23.487	30.182	38.385	48.715	62.109	
13	9.074	14.433	19.867	13.960	19.385	22.670	28.852	36.283	45.435	
14	8.862	12.943	17.385	12.694	17.385	22.670	25.626	33.672	44.240	
15	8.683	12.185	16.015	11.958	16.973	21.972	27.734	34.546	42.779	
16	8.531	13.566	19.076	14.418	24.458	31.788	40.975	52.858	68.771	
17	8.400	13.231	18.418	17.861	23.487	30.182	38.385	48.715	62.109	
18	8.285	12.943	17.861	12.111	20.843	25.959	31.847	38.750	46.986	
19	8.185	12.694	17.385	11.958	16.015	20.381	25.244	30.779	37.188	
20	8.096	12.476	16.973	11.823	15.763	19.972	24.616	29.850	35.846	
21	8.017	12.283	16.613	11.695	15.538	19.606	24.060	29.036	34.680	
22	7.945	12.111	16.296	11.534	15.334	19.279	23.565	28.316	33.659	
23	7.881	11.958	16.015	11.581	15.334	19.279	23.565	28.316	33.659	
24	7.823	11.820	15.763	11.695	15.538	19.606	24.060	29.036	34.680	
25	7.770	11.695	15.538	11.581	15.334	19.279	23.565	28.316	33.659	
26	7.721	11.581	15.334	11.581	15.334	19.279	23.565	28.316	33.659	

Table A.7 (Continued)

Degrees of Freedom, ν	$p = 1$	$p = 2$	$p = 3$	$p = 4$	$p = 5$	$p = 6$	$p = 7$	$p = 8$	$p = 9$	$p = 10$
27	7.677	11.478	15.149	18.983	23.121	27.675	32.756	38.496	45.051	52.622
28	7.636	11.383	14.980	18.715	22.721	27.101	31.954	37.393	43.554	50.604
29	7.598	11.295	14.825	18.471	22.359	26.584	31.236	36.414	42.234	48.839
30	7.562	11.215	14.683	18.247	22.029	26.116	30.589	35.538	41.062	47.283
35	7.419	10.890	14.117	17.366	20.743	24.314	28.135	32.259	36.743	41.651
40	7.314	10.655	13.715	16.750	19.858	23.094	26.502	30.120	33.984	38.135
45	7.234	10.478	13.414	16.295	19.211	22.214	25.340	28.617	32.073	35.737
50	7.171	10.340	13.181	15.945	18.718	21.550	24.470	27.504	30.673	33.998
55	7.119	10.228	12.995	15.667	18.331	21.030	23.795	26.647	29.603	32.682
60	7.077	10.137	12.843	15.442	18.018	20.613	23.257	25.967	28.760	31.550
70	7.011	9.996	12.611	15.098	17.543	19.986	22.451	24.957	27.515	30.139
80	6.963	9.892	12.440	14.849	17.201	19.536	21.877	24.242	26.642	29.085
90	6.925	9.813	12.310	14.660	16.942	19.197	21.448	23.710	25.995	28.310
100	6.895	9.750	12.208	14.511	16.740	18.934	21.115	23.299	25.496	27.714
110	6.871	9.699	12.125	14.391	16.577	18.722	20.849	22.972	25.101	27.243
120	6.851	9.657	12.057	14.292	16.444	18.549	20.632	22.705	24.779	26.862
150	6.807	9.565	11.909	14.079	16.156	18.178	20.167	22.137	24.096	26.054
200	6.763	9.474	11.764	13.871	15.877	17.819	19.720	21.592	23.446	25.287
400	6.699	9.341	11.551	13.569	15.473	17.303	19.080	20.818	22.525	24.209
1000	6.660	9.262	11.426	13.392	15.239	17.006	18.743	20.376	22.003	23.600
∞	6.635	9.210	11.345	13.277	15.086	16.812	18.475	20.090	21.666	23.209

Note: $p = \text{number of variables}$.

Table A.8 Bonferroni *t*-Values, $t_{\alpha/2k,\nu}$, $\alpha = 0.05$

ν	k									
	1	2	3	4	5	6	7	8	9	10
	100 α/k									
	5.0000	2.5000	1.6667	1.2500	1.0000	.8333	.7143	.6250	.5556	.5000
2	4.3027	6.2053	7.6488	8.8602	9.9248	10.8859	11.7687	12.5897	13.3604	14.0890
3	3.1824	4.1765	4.8567	5.3919	5.8409	6.2315	6.5797	6.8952	7.1849	7.4533
4	2.7764	3.4954	3.9608	4.3147	4.6041	4.8510	5.0675	5.2611	5.4366	5.5976
5	2.5706	3.1634	3.5341	3.8100	4.0321	4.2193	4.3818	4.5257	4.6553	4.7733
6	2.4469	2.9687	3.2875	3.5212	3.7074	2.8630	3.9971	4.1152	4.2209	4.3168
7	2.3646	2.8412	3.1276	3.3353	3.4995	3.6358	3.7527	3.8552	3.9467	4.0293
8	2.3060	2.7515	3.0158	3.2060	3.3554	3.4789	3.5844	3.6766	3.7586	3.8325
9	2.2622	2.6850	2.9333	3.1109	3.2498	3.3642	3.4616	3.5465	3.6219	3.6897
10	2.2281	2.6338	2.8701	3.0382	3.1693	3.2768	3.3682	3.4477	3.5182	3.5814
11	2.2010	2.5931	2.8200	2.9809	3.1058	3.2081	3.2949	3.3702	3.4368	3.4966
12	2.1788	2.5600	2.7795	2.9345	3.0545	3.1527	3.2357	3.3078	3.3714	3.4284
13	2.1604	2.5326	2.7459	2.8961	3.0123	3.1070	3.1871	3.2565	3.3177	3.3725
14	2.1448	2.5096	2.7178	2.8640	2.9768	3.0688	3.1464	3.2135	3.2727	3.3257
15	2.1314	2.4899	2.6937	2.8366	2.9467	3.0363	3.1118	3.1771	3.2346	3.2860
16	2.1199	2.4729	2.6730	2.8131	2.9208	3.0083	3.0821	3.1458	3.2019	3.2520
17	2.1098	2.4581	2.6550	2.7925	2.8982	2.9840	3.0563	3.1186	3.1735	3.2224
18	2.1009	2.4450	2.6391	2.7745	2.8784	2.9627	3.0336	3.0948	3.1486	3.1966
19	2.0930	2.4334	2.6251	2.7586	2.8609	2.9439	3.0136	3.0738	3.1266	3.1737
20	2.0860	2.4231	2.6126	2.7444	2.8453	2.9271	2.9958	3.0550	3.1070	3.1534
21	2.0796	2.4138	2.6013	2.7316	2.8314	2.9121	2.9799	3.0382	3.0895	3.1352
22	2.0739	2.4055	2.5912	2.7201	2.8188	2.8985	2.9655	3.0231	3.0737	3.1188
23	2.0687	2.3979	2.5820	2.7097	2.8073	2.8863	2.9525	3.0095	3.0595	3.1040
24	2.0639	2.3909	2.5736	2.7002	2.7969	2.8751	2.9406	2.9970	3.0465	3.0905
25	2.0595	2.3846	2.5660	2.6916	2.7874	2.8649	2.9298	2.9856	3.0346	3.0782
26	2.0555	2.3788	2.5589	2.6836	2.7787	2.8555	2.9199	2.9752	3.0237	3.0669
27	2.0518	2.3734	2.5525	2.6763	2.7707	2.8469	2.9107	2.9656	3.0137	3.0565
28	2.0484	2.3685	2.5465	2.6695	2.7633	2.8389	2.9023	2.9567	3.0045	3.0469
29	2.0452	2.3638	2.5409	2.6632	2.7564	2.8316	2.8945	2.9485	2.9959	3.0380
30	2.0423	2.3596	2.5357	2.6574	2.7500	2.8247	2.8872	2.9409	2.9880	3.0298
35	2.0301	2.3420	2.5145	2.6334	2.7238	2.7966	2.8575	2.9097	2.9554	2.9960
40	2.0211	2.3289	2.4989	2.6157	2.7045	2.7759	2.8355	2.8867	2.9314	2.9712
45	2.0141	2.3189	2.4868	2.6021	2.6896	2.7599	2.8187	2.8690	2.9130	2.9521
50	2.0086	2.3109	2.4772	2.5913	2.6778	2.7473	2.8053	2.8550	2.8984	2.9370
55	2.0040	2.3044	2.4694	2.5825	2.6682	2.7370	2.7944	2.8436	2.8866	2.9247
60	2.0003	2.2990	2.4630	2.5752	2.6603	2.7286	2.7855	2.8342	2.8768	2.9146
70	1.9944	2.2906	2.4529	2.5639	2.6479	2.7153	2.7715	2.8195	2.8615	2.8987
80	1.9901	2.2844	2.4454	2.5554	2.6387	2.7054	2.7610	2.8086	2.8502	2.8870
90	1.9867	2.2795	2.4395	2.5489	2.6316	2.6978	2.7530	2.8002	2.8414	2.8779
100	1.9840	2.2757	2.4349	2.5437	2.6259	2.6918	2.7466	2.7935	2.8344	2.8707
110	1.9818	2.2725	2.4311	2.5394	2.6213	2.6868	2.7414	2.7880	2.8287	2.8648
120	1.9799	2.2699	2.4280	2.5359	2.6174	2.6827	2.7370	2.7835	2.8240	2.8599
250	1.9695	2.2550	2.4102	2.5159	2.5956	2.6594	2.7124	2.7577	2.7972	2.8322
500	1.9647	2.2482	2.4021	2.5068	2.5857	2.6488	2.7012	2.7460	2.7850	2.8195
1000	1.9623	2.2448	2.3980	2.5022	2.5808	2.6435	2.6957	2.7402	2.7790	2.8133
∞	1.9600	2.2414	2.3940	2.4977	2.5758	2.6383	2.6901	2.7344	2.7729	2.8070

(continued)

Table A.8 (Continued)

ν	k									
	11	12	13	14	15	16	17	18	19	
	$100\alpha/k$									
.4545	.4167	.3846	.3571	.3333	.3125	.2941	.2778	.2632		
2	14.7818	15.4435	16.0780	16.6883	17.2772	17.8466	18.3984	18.9341	19.4551	
3	7.7041	7.9398	8.1625	8.3738	8.5752	8.7676	8.9521	9.1294	9.3001	
4	5.7465	5.8853	6.0154	6.1380	6.2541	6.3643	6.4693	6.5697	6.6659	
5	4.8819	4.9825	5.0764	5.1644	5.2474	5.3259	5.4005	5.4715	5.5393	
6	4.4047	4.4858	4.5612	4.6317	4.6979	4.7604	4.8196	4.8759	4.9295	
7	4.1048	4.1743	4.2388	4.2989	4.3553	4.4084	4.4586	4.5062	4.5514	
8	3.8999	3.9618	4.0191	4.0724	4.1224	4.1693	4.2137	4.2556	4.2955	
9	3.7513	3.8079	3.8602	3.9088	3.9542	3.9969	4.0371	4.0752	4.1114	
10	3.6388	3.6915	3.7401	3.7852	3.8273	3.8669	3.9041	3.9394	3.9728	
11	3.5508	3.6004	3.6462	3.6887	3.7283	3.7654	3.8004	3.8335	3.8648	
12	3.4801	3.5274	3.5709	3.6112	3.6489	3.6842	3.7173	3.7487	3.7783	
13	3.4221	3.4674	3.5091	3.5478	3.5838	3.6176	3.6493	3.6793	3.7076	
14	3.3736	3.4173	3.4576	3.4949	3.5296	3.5621	3.5926	3.6214	3.6487	
15	3.3325	3.3749	3.4139	3.4501	3.4837	3.5151	3.5447	3.5725	3.5989	
16	3.2973	3.3386	3.3765	3.4116	3.4443	3.4749	3.5036	3.5306	3.5562	
17	3.2667	3.3070	3.3440	3.3783	3.4102	3.4400	3.4680	3.4944	3.5193	
18	3.2399	3.2794	3.3156	3.3492	3.3804	3.4095	3.4369	3.4626	3.4870	
19	3.2163	3.2550	3.2906	3.3235	3.3540	3.3826	3.4094	3.4347	3.4585	
20	3.1952	3.2333	3.2683	3.3006	3.3306	3.3587	3.3850	3.4098	3.4332	
21	3.1764	3.2139	3.2483	3.2802	3.3097	3.3373	3.3632	3.3876	3.4106	
22	3.1595	3.1965	3.2304	3.2618	3.2909	3.3181	3.3436	3.3676	3.3903	
23	3.1441	3.1807	3.2142	3.2451	3.2739	3.3007	3.3259	3.3495	3.3719	
24	3.1302	3.1663	3.1994	3.2300	3.2584	3.2849	3.3097	3.3331	3.3552	
25	3.1175	3.1532	3.1859	3.2162	3.2443	3.2705	3.2950	3.3181	3.3400	
26	3.1058	3.1412	3.1736	3.2035	3.2313	3.2572	3.2815	3.3044	3.3260	
27	3.0951	3.1301	3.1622	3.1919	3.2194	3.2451	3.2691	3.2918	3.3132	
28	3.0852	3.1199	3.1517	3.1811	3.2084	3.2339	3.2577	3.2801	3.3013	
29	3.0760	3.1105	3.1420	3.1712	3.1982	3.2235	3.2471	3.2694	3.2904	
30	3.0675	3.1017	3.1330	3.1620	3.1888	3.2138	3.2373	3.2594	3.2802	
35	3.0326	3.0658	3.0962	3.1242	3.1502	3.1744	3.1971	3.2185	3.2386	
40	3.0069	3.0393	3.0690	3.0964	3.1218	3.1455	3.1676	3.1884	3.2081	
45	2.9872	3.0191	3.0482	3.0751	3.1000	3.1232	3.1450	3.1654	3.1846	
50	2.9716	3.0030	3.0318	3.0582	3.0828	3.1057	3.1271	3.1472	3.1661	
55	2.9589	2.9900	3.0184	3.0446	3.0688	3.0914	3.1125	3.1324	3.1511	
60	2.9485	2.9792	3.0074	3.0333	3.0573	3.0796	3.1005	3.1202	3.1387	
70	2.9321	2.9624	2.9901	3.0156	3.0393	3.0613	3.0818	3.1012	3.1194	
80	2.9200	2.9500	2.9773	3.0026	3.0259	3.0476	3.0679	3.0870	3.1050	
90	2.9106	2.9403	2.9675	2.9924	3.0156	3.0371	3.0572	3.0761	3.0939	
100	2.9032	2.9327	2.9596	2.9844	3.0073	3.0287	3.0487	3.0674	3.0851	
110	2.8971	2.9264	2.9532	2.9778	3.0007	3.0219	3.0417	3.0604	3.0779	
120	2.8921	2.9212	2.9479	2.9724	2.9951	3.0162	3.0360	3.0545	3.0720	
250	2.8635	2.8919	2.9178	2.9416	2.9637	2.9842	3.0034	3.0213	3.0383	
500	2.8505	2.8785	2.9041	2.9276	2.9494	2.9696	2.9885	3.0063	3.0230	
1000	2.8440	2.8719	2.8973	2.9207	2.9423	2.9624	2.9812	2.9988	3.0154	
∞	2.8376	2.8653	2.8905	2.9137	2.9352	2.9552	2.9738	2.9913	3.0078	

Table A.9 Lower Critical Values of Wilks' Λ , $\alpha = 0.05$

$$\Lambda = \frac{|\mathbf{E}|}{|\mathbf{E} + \mathbf{H}|} = \prod_{i=1}^s \frac{1}{1 + \lambda_i},$$

where $\lambda_1, \lambda_2, \dots, \lambda_s$ are eigenvalues of $\mathbf{E}^{-1}\mathbf{H}$. Reject H_0 if $\Lambda \leq$ table value.

ν_E	ν_H											
	1	2	3	4	5	6	7	8	9	10	11	12
$p = 1$												
1	6.16 ^a	2.50 ^a	1.54 ^a	1.11 ^a	.868 ^a	.712 ^a	.603 ^a	.523 ^a	.462 ^a	.413 ^a	.374 ^a	.341 ^a
2	.098	.050	.034	.025	.020	.017	.015	.013	.011	.010	9.28 ^a	8.51 ^a
3	.229	.136	.097	.076	.062	.053	.046	.041	.036	.033	.030	.028
4	.342	.224	.168	.135	.113	.098	.086	.076	.069	.063	.058	.053
5	.431	.302	.236	.194	.165	.144	.128	.115	.104	.096	.088	.082
6	.501	.368	.296	.249	.215	.189	.169	.153	.140	.129	.119	.111
7	.556	.425	.349	.298	.261	.232	.209	.190	.175	.161	.150	.140
8	.601	.473	.396	.343	.303	.271	.246	.225	.208	.193	.180	.169
9	.638	.514	.437	.382	.341	.308	.281	.258	.239	.223	.209	.196
10	.668	.549	.473	.418	.376	.341	.313	.289	.269	.251	.236	.222
11	.694	.580	.505	.450	.407	.372	.343	.318	.297	.278	.262	.247
12	.717	.607	.534	.479	.436	.400	.370	.345	.323	.304	.286	.271
13	.736	.631	.560	.506	.462	.426	.396	.370	.347	.327	.310	.294
14	.753	.652	.583	.529	.486	.450	.420	.393	.370	.350	.332	.315
15	.768	.671	.603	.551	.508	.473	.442	.415	.392	.371	.352	.336
16	.781	.688	.622	.571	.529	.493	.462	.436	.412	.391	.372	.355
17	.792	.703	.639	.589	.548	.512	.482	.455	.431	.410	.390	.373
18	.803	.717	.655	.606	.565	.530	.499	.473	.449	.427	.408	.390
19	.813	.730	.669	.621	.581	.546	.516	.490	.466	.444	.425	.407
20	.821	.741	.683	.636	.596	.562	.532	.505	.482	.460	.440	.423
21	.829	.752	.695	.649	.610	.576	.547	.520	.497	.475	.455	.437
22	.836	.762	.706	.661	.623	.590	.561	.534	.511	.489	.470	.452
23	.843	.771	.717	.673	.635	.603	.574	.548	.524	.503	.483	.465
24	.849	.779	.727	.684	.647	.615	.586	.560	.537	.516	.496	.478
25	.855	.787	.736	.694	.658	.626	.598	.572	.549	.528	.508	.490
26	.860	.794	.744	.703	.668	.637	.609	.583	.560	.539	.520	.502
27	.865	.801	.752	.712	.677	.647	.619	.594	.571	.551	.531	.513
28	.870	.807	.760	.721	.686	.656	.629	.604	.582	.561	.542	.524
29	.874	.813	.767	.729	.695	.665	.638	.614	.592	.571	.552	.535
30	.878	.819	.774	.736	.703	.674	.647	.623	.601	.581	.562	.544
40	.907	.861	.824	.793	.766	.741	.718	.696	.677	.658	.641	.625
60	.938	.905	.879	.856	.835	.816	.798	.781	.766	.751	.736	.723
80	.953	.928	.907	.889	.873	.858	.843	.829	.816	.804	.792	.780
100	.962	.942	.925	.910	.897	.884	.872	.860	.849	.838	.828	.818
120	.968	.951	.937	.925	.913	.902	.891	.882	.872	.863	.854	.845
140	.973	.958	.946	.935	.925	.915	.906	.897	.889	.881	.873	.865
170	.978	.965	.955	.946	.937	.929	.922	.914	.907	.900	.893	.887
200	.981	.970	.962	.954	.947	.940	.933	.926	.920	.914	.908	.902
240	.984	.975	.968	.961	.955	.949	.944	.938	.933	.928	.923	.918
320	.988	.981	.976	.971	.966	.962	.957	.953	.949	.945	.941	.937
440	.991	.986	.982	.979	.975	.972	.969	.966	.963	.960	.957	.954
600	.994	.990	.987	.984	.982	.979	.977	.975	.972	.970	.968	.966
800	.995	.993	.990	.988	.986	.984	.983	.981	.979	.977	.976	.974
1000	.996	.994	.992	.991	.989	.988	.986	.985	.983	.982	.981	.979

^aMultiply entry by 10^{-3} .

(continued)

Table A.9 (Continued)

ν_E	ν_H											
	1	2	3	4	5	6	7	8	9	10	11	12
	$p = 2$											
1	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000
2	.250 ^a	.641 ^a	.287 ^a	.162 ^a	.104 ^a	.072 ^a	.053 ^a	.041 ^a	.032 ^a	.026 ^a	.022 ^a	.018 ^a
3	.050	.018	.953 ^a	.584 ^a	.395 ^a	.285 ^a	.215 ^a	.168 ^a	.135 ^a	.111 ^a	.928 ^a	.787 ^a
4	.136	.062	.036	.023	.017	.012	.956 ^a	.762 ^a	.621 ^a	.517 ^a	.436 ^a	.373 ^a
5	.224	.117	.074	.051	.037	.028	.023	.018	.015	.013	.011	.009
6	.302	.175	.116	.084	.063	.049	.040	.033	.027	.023	.020	.017
7	.368	.230	.160	.119	.092	.074	.060	.050	.042	.036	.032	.028
8	.4256	.280	.203	.155	.122	.099	.082	.069	.059	.051	.045	.040
9	.473	.326	.243	.190	.153	.126	.106	.090	.078	.068	.060	.053
10	.514	.367	.281	.223	.183	.152	.129	.111	.097	.085	.075	.067
11	.549	.404	.316	.255	.212	.179	.153	.133	.116	.102	.091	.082
12	.580	.437	.348	.286	.240	.204	.176	.154	.136	.120	.108	.097
13	.607	.467	.378	.314	.266	.229	.199	.175	.155	.138	.124	.112
14	.631	.495	.405	.340	.291	.252	.221	.195	.174	.156	.141	.128
15	.652	.519	.431	.365	.315	.275	.242	.215	.193	.174	.157	.143
16	.671	.542	.454	.389	.337	.296	.263	.235	.211	.191	.174	.159
17	.688	.562	.476	.410	.359	.317	.282	.254	.229	.208	.190	.174
18	.703	.581	.496	.431	.379	.337	.301	.272	.246	.225	.206	.189
19	.717	.598	.515	.450	.398	.355	.320	.289	.263	.241	.221	.204
20	.730	.614	.532	.468	.416	.373	.337	.306	.279	.256	.236	.218
21	.741	.629	.548	.485	.433	.390	.354	.322	.295	.271	.251	.232
22	.752	.643	.564	.501	.449	.406	.370	.338	.310	.286	.265	.246
23	.762	.656	.578	.516	.465	.422	.385	.353	.325	.300	.279	.259
24	.771	.668	.591	.530	.479	.436	.399	.367	.339	.314	.292	.272
25	.779	.679	.604	.544	.493	.450	.413	.381	.353	.328	.305	.285
26	.787	.689	.616	.556	.506	.464	.427	.395	.366	.341	.318	.297
27	.794	.699	.627	.568	.519	.477	.440	.407	.379	.353	.330	.309
28	.801	.708	.638	.580	.531	.489	.452	.420	.391	.365	.342	.321
29	.807	.717	.648	.591	.542	.501	.464	.432	.403	.377	.354	.332
30	.813	.725	.657	.601	.553	.512	.475	.443	.414	.388	.365	.344
40	.858	.786	.730	.682	.640	.602	.568	.537	.509	.484	.460	.439
60	.903	.853	.811	.774	.741	.710	.682	.656	.632	.609	.588	.568
80	.927	.888	.854	.825	.798	.772	.749	.727	.706	.686	.667	.649
100	.941	.909	.882	.857	.834	.813	.793	.774	.755	.738	.721	.705
120	.951	.924	.900	.879	.860	.841	.823	.807	.791	.775	.760	.746
140	.958	.934	.914	.895	.878	.862	.846	.831	.817	.803	.790	.777
170	.965	.946	.929	.913	.898	.885	.871	.859	.846	.834	.823	.812
200	.970	.954	.939	.926	.913	.901	.889	.878	.867	.857	.847	.837
240	.975	.961	.949	.938	.927	.917	.907	.897	.888	.879	.870	.862
320	.981	.971	.962	.953	.945	.937	.929	.922	.914	.907	.901	.894
440	.986	.979	.972	.965	.959	.953	.948	.942	.937	.932	.926	.921
600	.990	.984	.979	.975	.970	.966	.961	.957	.953	.949	.945	.942
800	.993	.988	.984	.981	.977	.974	.971	.968	.965	.962	.959	.956
1000	.994	.991	.987	.985	.982	.979	.977	.974	.972	.969	.967	.964

^aMultiply entry by 10⁻³.

(continued)

Table A.9 (Continued)

ν_E	ν_H											
	1	2	3	4	5	6	7	8	9	10	11	12
	$p = 3$											
1	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000
2	.000	.000	.000	.000	.000	.001 ^a	.002 ^a	.004 ^a	.005 ^a	.008 ^a	.010 ^a	.013 ^a
3	1.70 ^a	.354 ^a	.179 ^a	.127 ^a	.105 ^a	.095 ^a	.091 ^a	.090 ^a	.091 ^a	.092 ^a	.095 ^a	.098 ^a
4	.034	.010	.004	.002	.001	.001	.809 ^a	.659 ^a	.562 ^a	.496 ^a	.449 ^a	.416 ^a
5	.097	.036	.018	.010	6.36 ^a	4.37 ^a	3.20 ^a	2.46 ^a	1.97 ^a	1.64 ^a	1.40 ^a	1.22 ^a
6	.168	.074	.040	.024	.016	.011	.008	.006	.004	3.94 ^a	3.28 ^a	2.79 ^a
7	.236	.116	.068	.043	.029	.021	.016	.012	9.49 ^a	7.67 ^a	6.35 ^a	5.35 ^a
8	.296	.160	.099	.066	.046	.034	.026	.020	.016	.013	.011	9.00 ^a
9	.349	.203	.131	.091	.066	.049	.038	.030	.024	.020	.016	.014
10	.396	.243	.164	.117	.086	.066	.052	.041	.034	.028	.023	.020
11	.437	.281	.196	.143	.108	.084	.067	.054	.044	.037	.031	.026
12	.473	.316	.226	.169	.130	.103	.083	.067	.056	.047	.040	.034
13	.505	.348	.255	.194	.152	.122	.099	.082	.068	.058	.049	.042
14	.534	.378	.283	.219	.174	.141	.116	.096	.081	.069	.059	.051
15	.560	.405	.309	.243	.195	.160	.133	.111	.095	.081	.070	.061
16	.583	.431	.334	.266	.216	.179	.149	.127	.108	.093	.081	.071
17	.603	.454	.357	.288	.236	.197	.166	.142	.122	.106	.092	.081
18	.622	.476	.379	.309	.256	.215	.183	.157	.136	.118	.104	.092
19	.639	.496	.399	.329	.275	.233	.199	.172	.149	.131	.115	.102
20	.655	.515	.419	.348	.293	.250	.215	.187	.163	.144	.127	.113
21	.669	.532	.437	.366	.310	.266	.230	.201	.177	.156	.139	.124
22	.683	.548	.454	.383	.327	.282	.246	.215	.190	.169	.150	.135
23	.695	.564	.470	.399	.343	.298	.260	.229	.203	.181	.162	.146
24	.706	.578	.486	.415	.359	.313	.275	.243	.216	.193	.173	.156
25	.717	.591	.500	.430	.374	.327	.289	.256	.229	.205	.185	.167
26	.727	.604	.514	.444	.388	.341	.302	.269	.241	.217	.196	.178
27	.736	.616	.527	.458	.401	.355	.315	.282	.253	.229	.207	.188
28	.744	.627	.540	.471	.415	.368	.328	.294	.265	.240	.218	.199
29	.752	.638	.552	.483	.427	.380	.340	.306	.277	.251	.229	.209
30	.760	.648	.563	.495	.439	.392	.352	.318	.288	.262	.239	.219
40	.816	.724	.651	.591	.539	.494	.454	.419	.387	.359	.334	.311
60	.875	.808	.752	.704	.661	.623	.587	.555	.526	.498	.473	.449
80	.905	.853	.808	.769	.733	.700	.670	.641	.615	.590	.566	.544
100	.924	.881	.844	.810	.780	.751	.725	.700	.676	.654	.632	.612
120	.936	.900	.868	.839	.813	.788	.764	.742	.721	.700	.681	.663
140	.945	.913	.886	.861	.837	.815	.794	.774	.755	.736	.719	.702
170	.955	.928	.905	.884	.864	.845	.827	.809	.792	.776	.761	.746
200	.961	.939	.919	.900	.883	.866	.850	.835	.820	.806	.792	.779
240	.968	.949	.932	.916	.901	.887	.873	.860	.848	.835	.823	.811
320	.976	.961	.948	.936	.925	.914	.903	.893	.883	.873	.864	.854
440	.982	.972	.962	.953	.945	.937	.929	.921	.913	.906	.899	.891
600	.987	.979	.972	.966	.959	.953	.947	.941	.936	.930	.924	.919
800	.990	.984	.979	.974	.969	.965	.960	.956	.951	.947	.943	.939
1000	.992	.987	.983	.979	.975	.972	.968	.964	.961	.957	.954	.950

^aMultiply entry by 10⁻³.

(continued)

Table A.9 (Continued)

ν_E	ν_H											
	1	2	3	4	5	6	7	8	9	10	11	12
	$p = 4$											
1	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000
2	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000
3	.000	.000	.000	.000	.000	.001 ^a	.001 ^a	.001 ^a	.002 ^a	.002 ^a	.002 ^a	.003 ^a
4	1.38 ^a	.292 ^a	.127 ^a	.075 ^a	.052 ^a	.040 ^a	.033 ^a	.029 ^a	.026 ^a	.025 ^a	.023 ^a	.022 ^a
5	.026	6.09 ^a	2.31 ^a	1.13 ^a	.647 ^a	.416 ^a	.292 ^a	.218 ^a	.172 ^a	.141 ^a	.120 ^a	.105 ^a
6	.076	.024	.010	5.07 ^a	2.90 ^a	1.82 ^a	1.22 ^a	.872 ^a	.652 ^a	.508 ^a	.409 ^a	.338 ^a
7	.135	.051	.024	.013	7.74 ^a	4.94 ^a	3.34 ^a	2.36 ^a	1.74 ^a	1.33 ^a	1.05 ^a	.848 ^a
8	.194	.084	.043	.025	.015	.010	6.98 ^a	4.99 ^a	3.70 ^a	2.82 ^a	2.21 ^a	1.77 ^a
9	.249	.119	.066	.040	.026	.017	.012	8.91 ^a	6.66 ^a	5.11 ^a	4.01 ^a	3.21 ^a
10	.298	.155	.091	.057	.038	.027	.019	.014	.011	8.29 ^a	6.54 ^a	5.25 ^a
11	.343	.190	.117	.077	.053	.037	.027	.021	.016	.012	9.84 ^a	7.95 ^a
12	.382	.223	.143	.097	.068	.049	.037	.028	.022	.017	.014	.011
13	.418	.255	.169	.117	.085	.063	.047	.037	.029	.023	.019	.015
14	.450	.286	.194	.138	.102	.077	.059	.046	.037	.030	.024	.020
15	.479	.314	.219	.159	.119	.091	.071	.056	.045	.037	.030	.025
16	.506	.340	.243	.180	.136	.106	.083	.067	.054	.044	.037	.031
17	.529	.365	.266	.200	.154	.121	.096	.078	.064	.053	.044	.037
18	.551	.389	.288	.219	.171	.136	.109	.089	.074	.061	.051	.044
19	.571	.410	.309	.239	.188	.151	.123	.101	.084	.070	.059	.051
20	.589	.431	.329	.257	.205	.166	.136	.113	.094	.079	.068	.058
21	.606	.450	.348	.275	.221	.181	.149	.124	.105	.089	.076	.065
22	.621	.468	.366	.292	.237	.195	.162	.136	.115	.098	.085	.073
23	.636	.485	.383	.309	.253	.210	.175	.148	.126	.108	.093	.081
24	.649	.501	.399	.325	.268	.224	.188	.160	.137	.118	.102	.089
25	.661	.516	.415	.340	.283	.237	.201	.172	.148	.128	.111	.097
26	.673	.530	.430	.355	.297	.251	.214	.183	.158	.138	.120	.106
27	.684	.544	.444	.369	.311	.264	.226	.195	.169	.147	.129	.114
28	.694	.556	.458	.383	.324	.277	.238	.206	.180	.157	.138	.122
29	.703	.568	.471	.396	.337	.289	.250	.217	.190	.167	.147	.131
30	.712	.580	.483	.409	.349	.301	.261	.228	.200	.177	.157	.139
40	.779	.668	.583	.513	.455	.406	.364	.327	.295	.267	.243	.221
60	.849	.767	.700	.643	.592	.547	.507	.471	.438	.409	.382	.357
80	.885	.821	.766	.718	.675	.636	.600	.567	.536	.508	.482	.457
100	.908	.854	.809	.768	.730	.696	.664	.634	.606	.580	.555	.532
120	.923	.877	.838	.802	.770	.739	.711	.684	.658	.634	.611	.590
140	.934	.894	.860	.828	.799	.772	.746	.721	.698	.676	.655	.635
170	.945	.912	.883	.856	.831	.808	.785	.764	.743	.724	.705	.687
200	.953	.925	.900	.876	.855	.834	.814	.795	.777	.759	.742	.726
240	.961	.937	.916	.896	.877	.859	.842	.826	.810	.795	.780	.765
320	.971	.952	.936	.921	.907	.893	.879	.866	.854	.841	.829	.818
440	.979	.965	.953	.942	.931	.921	.911	.901	.891	.882	.872	.863
600	.984	.974	.966	.957	.949	.941	.934	.926	.919	.912	.905	.898
800	.988	.981	.974	.968	.961	.956	.950	.944	.938	.933	.927	.922
1000	.991	.985	.979	.974	.969	.964	.960	.955	.950	.946	.941	.937

^aMultiply entry by 10⁻³.

(continued)

Table A.9 (Continued)

ν_E	ν_H											
	1	2	3	4	5	6	7	8	9	10	11	12
	$p = 5$											
1	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000
2	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000
3	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000
4	.000	.000	.000	.000	.001 ^a							
5	1.60 ^a	.291 ^a	.105 ^a	.052 ^a	.031 ^a	.021 ^a	.015 ^a	.012 ^a	.010 ^a	.008 ^a	.007 ^a	.007 ^a
6	.021	4.39 ^a	1.48 ^a	.647 ^a	.335 ^a	.197 ^a	.126 ^a	.087 ^a	.064 ^a	.049 ^a	.039 ^a	.032 ^a
7	.063	.017	6.36 ^a	2.90 ^a	1.51 ^a	.872 ^a	.544 ^a	.361 ^a	.253 ^a	.185 ^a	.141 ^a	.110 ^a
8	.114	.037	.016	7.74 ^a	4.21 ^a	2.48 ^a	1.56 ^a	1.03 ^a	.716 ^a	.516 ^a	.385 ^a	.296 ^a
9	.165	.063	.029	.015	8.79 ^a	5.35 ^a	3.43 ^a	2.30 ^a	1.61 ^a	1.16 ^a	.861 ^a	.657 ^a
10	.215	.092	.046	.026	.015	9.64 ^a	6.34 ^a	4.34 ^a	3.06 ^a	2.22 ^a	1.66 ^a	1.27 ^a
11	.261	.122	.066	.038	.024	.015	.010	7.22 ^a	5.17 ^a	3.80 ^a	2.86 ^a	2.19 ^a
12	.303	.153	.086	.053	.034	.022	.015	.011	7.99 ^a	5.95 ^a	4.51 ^a	3.49 ^a
13	.341	.183	.108	.068	.045	.031	.022	.016	.012	8.68 ^a	6.66 ^a	5.19 ^a
14	.376	.212	.130	.085	.057	.040	.029	.021	.016	.012	9.31 ^a	7.32 ^a
15	.407	.239	.152	.102	.070	.050	.037	.027	.021	.016	.012	9.88 ^a
16	.436	.266	.174	.119	.084	.061	.045	.034	.026	.020	.016	.013
17	.462	.291	.195	.136	.098	.072	.054	.042	.032	.025	.020	.016
18	.486	.315	.216	.154	.113	.084	.064	.050	.039	.031	.025	.020
19	.508	.337	.236	.171	.127	.096	.074	.058	.046	.037	.030	.024
20	.529	.359	.256	.188	.142	.109	.085	.067	.053	.043	.035	.029
21	.548	.379	.275	.205	.156	.121	.095	.076	.061	.050	.041	.034
22	.565	.398	.293	.221	.171	.134	.106	.085	.069	.057	.047	.039
23	.581	.416	.310	.237	.185	.146	.117	.095	.077	.064	.053	.044
24	.596	.433	.327	.253	.199	.159	.128	.104	.086	.071	.060	.050
25	.610	.449	.343	.268	.213	.171	.139	.114	.094	.079	.066	.056
26	.623	.465	.359	.283	.226	.183	.150	.124	.103	.087	.073	.062
27	.635	.479	.374	.297	.239	.195	.161	.134	.112	.094	.080	.068
28	.647	.493	.388	.311	.252	.207	.172	.143	.121	.102	.087	.075
29	.658	.506	.401	.324	.265	.219	.182	.153	.130	.110	.094	.081
30	.668	.519	.415	.337	.277	.230	.193	.163	.138	.118	.102	.088
40	.744	.617	.522	.446	.384	.333	.291	.255	.224	.198	.176	.156
60	.825	.729	.652	.587	.531	.482	.438	.400	.366	.336	.308	.284
80	.867	.791	.727	.672	.623	.578	.538	.502	.469	.438	.410	.385
100	.893	.830	.776	.728	.685	.645	.609	.576	.544	.516	.489	.464
120	.910	.856	.810	.768	.730	.694	.661	.631	.602	.575	.549	.525
140	.923	.876	.835	.798	.763	.731	.701	.673	.647	.621	.598	.575
170	.936	.897	.862	.830	.801	.773	.747	.722	.698	.675	.654	.633
200	.945	.912	.882	.854	.828	.803	.780	.758	.736	.716	.696	.677
240	.954	.926	.900	.877	.855	.833	.813	.793	.775	.757	.739	.722
300	.966	.944	.925	.906	.889	.872	.856	.841	.825	.811	.797	.783
440	.975	.959	.945	.931	.918	.905	.893	.881	.870	.858	.847	.836
600	.982	.970	.959	.949	.939	.930	.920	.911	.903	.894	.885	.877
800	.986	.977	.969	.961	.954	.947	.940	.933	.926	.919	.913	.906
1000	.989	.982	.975	.969	.963	.957	.951	.946	.940	.935	.929	.924

^aMultiply entry by 10⁻³.

(continued)

Table A.9 (Continued)

ν_E	ν_H											
	1	2	3	4	5	6	7	8	9	10	11	12
	$p = 6$											
1	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000
2	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000
3	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000
4	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000
5	.007 ^a	.002 ^a	.001 ^a	.001 ^a	.001 ^a	.000	.000	.000	.000	.000	.000	.000
6	.204 ^a	.315 ^a	.095 ^a	.040 ^a	.021 ^a	.012 ^a	.008 ^a	.006 ^a	.004 ^a	.003 ^a	.003 ^a	.002 ^a
7	.019	.348 ^a	.105 ^a	.416 ^a	.197 ^a	.106 ^a	.063 ^a	.040 ^a	.027 ^a	.020 ^a	.015 ^a	.011 ^a
8	.054	.013	.437 ^a	.182 ^a	.872 ^a	.465 ^a	.270 ^a	.168 ^a	.111 ^a	.076 ^a	.055 ^a	.041 ^a
9	.098	.029	.011	.494 ^a	.248 ^a	.136 ^a	.798 ^a	.497 ^a	.325 ^a	.222 ^a	.157 ^a	.115 ^a
10	.144	.050	.021	.010	.535 ^a	.304 ^a	.183 ^a	.116 ^a	.762 ^a	.521 ^a	.369 ^a	.269 ^a
11	.189	.074	.034	.017	.964 ^a	.567 ^a	.351 ^a	.226 ^a	.151 ^a	.105 ^a	.744 ^a	.543 ^a
12	.232	.099	.049	.027	.015	.935 ^a	.594 ^a	.392 ^a	.266 ^a	.186 ^a	.134 ^a	.983 ^a
13	.271	.126	.066	.037	.022	.014	.917 ^a	.617 ^a	.427 ^a	.303 ^a	.220 ^a	.163 ^a
14	.308	.152	.084	.049	.031	.020	.013	.907 ^a	.638 ^a	.459 ^a	.337 ^a	.252 ^a
15	.341	.179	.103	.063	.040	.026	.018	.013	.90 ^a	.657 ^a	.488 ^a	.368 ^a
16	.372	.204	.122	.077	.050	.034	.024	.017	.012	.897 ^a	.674 ^a	.514 ^a
17	.400	.229	.141	.091	.061	.042	.030	.021	.016	.012	.897 ^a	.690 ^a
18	.426	.252	.160	.106	.072	.051	.037	.027	.020	.015	.012	.897 ^a
19	.450	.275	.179	.121	.084	.060	.044	.033	.025	.019	.015	.011
20	.473	.296	.197	.136	.096	.070	.052	.039	.030	.023	.018	.014
21	.493	.317	.215	.151	.109	.080	.060	.045	.035	.027	.021	.017
22	.512	.337	.233	.166	.121	.090	.068	.052	.041	.032	.025	.020
23	.530	.355	.250	.181	.134	.101	.077	.060	.047	.037	.030	.024
24	.546	.373	.266	.195	.146	.111	.086	.067	.053	.042	.034	.028
25	.562	.390	.282	.210	.159	.122	.095	.075	.060	.048	.039	.032
26	.576	.406	.298	.224	.171	.133	.104	.083	.066	.054	.044	.036
27	.590	.422	.313	.237	.183	.143	.113	.091	.073	.060	.049	.040
28	.603	.436	.327	.251	.195	.154	.123	.099	.080	.066	.054	.045
29	.615	.450	.341	.264	.207	.165	.132	.107	.088	.072	.060	.050
30	.626	.464	.355	.277	.219	.175	.142	.116	.095	.079	.066	.055
40	.711	.570	.467	.387	.324	.273	.232	.198	.170	.147	.127	.110
60	.802	.693	.608	.536	.476	.424	.379	.340	.305	.275	.249	.225
80	.849	.762	.690	.629	.574	.526	.483	.445	.410	.378	.350	.324
100	.878	.806	.745	.691	.642	.599	.559	.523	.489	.458	.430	.404
120	.898	.836	.783	.735	.692	.652	.616	.582	.551	.521	.494	.468
140	.912	.858	.811	.769	.730	.694	.660	.629	.599	.572	.546	.521
170	.927	.882	.842	.806	.772	.740	.710	.682	.656	.630	.607	.584
200	.938	.899	.864	.832	.803	.774	.748	.722	.698	.675	.653	.632
240	.948	.915	.886	.858	.833	.808	.785	.763	.741	.721	.701	.682
320	.961	.936	.913	.892	.872	.852	.834	.816	.799	.782	.766	.750
440	.972	.953	.936	.920	.905	.890	.876	.862	.849	.836	.823	.811
600	.979	.965	.953	.941	.930	.918	.908	.897	.887	.877	.867	.857
800	.984	.974	.964	.955	.947	.938	.930	.922	.914	.906	.898	.891
1000	.987	.979	.971	.964	.957	.950	.944	.937	.930	.924	.918	.912

^aMultiply entry by 10⁻³.

(continued)

Table A.9 (Continued)

ν_E	ν_H											
	1	2	3	4	5	6	7	8	9	10	11	12
	$p = 7$											
1	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000
2	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000
3	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000
4	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000
5	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000
6	.043 ^a	.006 ^a	.002 ^a	.001 ^a	.001 ^a	.000	.000	.000	.000	.000	.000	.000
7	2.62 ^a	.350 ^a	.091 ^a	.033 ^a	.015 ^a	.008 ^a	.005 ^a	.003 ^a	.002 ^a	.002 ^a	.001 ^a	.001 ^a
8	.018	2.95 ^a	.809 ^a	.292 ^a	.126 ^a	.063 ^a	.034 ^a	.020 ^a	.013 ^a	.009 ^a	.006 ^a	.005 ^a
9	.048	.010	3.20 ^a	1.22 ^a	.543 ^a	.270 ^a	.147 ^a	.086 ^a	.053 ^a	.035 ^a	.024 ^a	.017 ^a
10	.087	.023	8.07 ^a	3.34 ^a	1.56 ^a	.798 ^a	.440 ^a	.259 ^a	.160 ^a	.104 ^a	.070 ^a	.049 ^a
11	.128	.040	.016	6.97 ^a	3.43 ^a	1.83 ^a	1.04 ^a	.619 ^a	.387 ^a	.252 ^a	.170 ^a	.119 ^a
12	.170	.060	.026	.012	6.34 ^a	3.51 ^a	2.05 ^a	1.25 ^a	.796 ^a	.525 ^a	.357 ^a	.249 ^a
13	.209	.083	.038	.019	.010	5.94 ^a	3.57 ^a	2.23 ^a	1.45 ^a	.967 ^a	.665 ^a	.468 ^a
14	.246	.106	.052	.027	.015	9.17 ^a	5.67 ^a	3.63 ^a	2.40 ^a	1.62 ^a	1.13 ^a	.804 ^a
15	.281	.129	.067	.037	.022	.013	8.37 ^a	5.48 ^a	3.68 ^a	2.54 ^a	1.79 ^a	1.28 ^a
16	.313	.153	.083	.047	.029	.018	.012	7.80 ^a	5.34 ^a	3.73 ^a	2.66 ^a	1.94 ^a
17	.343	.176	.099	.059	.037	.024	.016	.011	7.38 ^a	5.24 ^a	3.78 ^a	2.78 ^a
18	.370	.199	.116	.071	.045	.030	.020	.014	9.81 ^a	7.06 ^a	5.16 ^a	3.83 ^a
19	.396	.221	.133	.083	.054	.037	.025	.018	.013	9.20 ^a	6.80 ^a	5.10 ^a
20	.420	.242	.149	.096	.064	.044	.031	.022	.016	.012	8.72 ^a	6.60 ^a
21	.442	.263	.166	.109	.074	.052	.037	.026	.019	.014	.011	8.34 ^a
22	.462	.283	.183	.123	.085	.060	.043	.031	.023	.018	.013	.010
23	.482	.301	.199	.136	.095	.068	.050	.037	.028	.021	.016	.013
24	.499	.320	.215	.149	.106	.077	.057	.042	.032	.025	.019	.015
25	.516	.337	.230	.162	.117	.086	.064	.048	.037	.029	.022	.018
26	.532	.354	.246	.175	.128	.095	.071	.055	.042	.033	.026	.020
27	.547	.370	.260	.188	.139	.104	.079	.061	.047	.037	.029	.024
28	.561	.385	.275	.201	.150	.113	.087	.068	.053	.042	.033	.027
29	.574	.399	.289	.214	.161	.123	.095	.074	.059	.047	.037	.030
30	.586	.413	.302	.226	.172	.132	.103	.081	.064	.052	.042	.034
40	.679	.526	.417	.335	.273	.224	.185	.154	.128	.108	.091	.077
60	.779	.660	.566	.490	.426	.373	.327	.288	.254	.225	.200	.178
80	.832	.735	.656	.588	.530	.479	.434	.394	.358	.326	.298	.272
100	.864	.783	.715	.656	.603	.556	.513	.475	.439	.408	.378	.352
120	.886	.817	.757	.704	.657	.613	.574	.537	.504	.473	.444	.418
140	.902	.841	.788	.741	.698	.658	.621	.587	.556	.526	.498	.472
170	.919	.868	.823	.782	.744	.709	.676	.645	.616	.589	.563	.539
200	.931	.887	.848	.812	.778	.747	.717	.689	.662	.637	.613	.590
240	.942	.905	.871	.841	.812	.784	.758	.733	.709	.687	.665	.644
320	.957	.928	.902	.878	.855	.833	.812	.792	.773	.754	.736	.719
440	.968	.947	.928	.910	.893	.876	.860	.844	.829	.814	.800	.786
600	.977	.961	.947	.933	.920	.908	.895	.883	.872	.860	.849	.838
800	.982	.971	.960	.950	.940	.930	.920	.911	.902	.893	.884	.876
1000	.986	.977	.968	.959	.951	.943	.936	.928	.921	.914	.906	.899

^aMultiply entry by 10⁻³.

(continued)

Table A.9 (Continued)

ν_E	ν_H											
	1	2	3	4	5	6	7	8	9	10	11	12
$p = 8$												
1	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000
2	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000
3	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000
4	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000
5	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000
6	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000	.000
7	.138 ^a	.015 ^a	.004 ^a	.001 ^a	.001 ^a	.000	.000	.000	.000	.000	.000	.000
8	3.30 ^a	.393 ^a	.090 ^a	.029 ^a	.012 ^a	.006 ^a	.003 ^a	.002 ^a	.001 ^a	.001 ^a	.001 ^a	.000
9	.017	2.63 ^a	.659 ^a	.218 ^a	.087 ^a	.040 ^a	.020 ^a	.011 ^a	.007 ^a	.004 ^a	.003 ^a	.002 ^a
10	.044	8.63 ^a	2.46 ^a	.872 ^a	.361 ^a	.168 ^a	.086 ^a	.047 ^a	.028 ^a	4.017 ^a	.011 ^a	.008 ^a
11	.078	.019	6.15 ^a	2.36 ^a	1.03 ^a	.497 ^a	.259 ^a	.144 ^a	.085 ^a	.052 ^a	.034 ^a	.023 ^a
12	.116	.033	.012	4.99 ^a	2.30 ^a	1.16 ^a	.619 ^a	.351 ^a	.209 ^a	.130 ^a	.084 ^a	.056 ^a
13	.154	.051	.020	8.91 ^a	4.34 ^a	2.26 ^a	1.25 ^a	.727 ^a	.441 ^a	.278 ^a	.181 ^a	.122 ^a
14	.190	.070	.030	.014	7.22 ^a	3.92 ^a	2.23 ^a	1.33 ^a	.824 ^a	.527 ^a	.347 ^a	.235 ^a
15	.225	.090	.041	.021	.011	6.17 ^a	3.63 ^a	2.22 ^a	1.40 ^a	.910 ^a	.608 ^a	.416 ^a
16	.258	.111	.054	.028	.016	9.06 ^a	5.48 ^a	3.42 ^a	2.20 ^a	1.46 ^a	.987 ^a	.683 ^a
17	.289	.133	.067	.037	.021	.013	7.80 ^a	4.98 ^a	3.27 ^a	2.20 ^a	1.51 ^a	1.06 ^a
18	.318	.154	.082	.046	.027	.017	.011	6.92 ^a	4.62 ^a	3.15 ^a	2.19 ^a	1.56 ^a
19	.345	.175	.096	.056	.034	.021	.014	9.23 ^a	6.26 ^a	4.34 ^a	3.06 ^a	2.19 ^a
20	.370	.195	.111	.067	.042	.027	.018	.012	8.22 ^a	5.77 ^a	4.12 ^a	2.99 ^a
21	.393	.215	.127	.078	.050	.033	.022	.015	.010	7.46 ^a	5.39 ^a	3.95 ^a
22	.415	.235	.142	.089	.058	.039	.026	.018	.013	9.40 ^a	6.86 ^a	5.08 ^a
23	.436	.254	.157	.101	.067	.045	.031	.022	.016	.012	8.56 ^a	6.39 ^a
24	.455	.272	.172	.113	.076	.052	.037	.026	.019	.014	.010	7.88 ^a
25	.473	.289	.187	.124	.085	.060	.042	.031	.023	.017	.013	9.56 ^a
26	.490	.306	.201	.136	.095	.067	.048	.035	.026	.020	.015	.011
27	.505	.322	.215	.148	.104	.075	.055	.040	.030	.023	.017	.013
28	.520	.338	.229	.160	.114	.083	.061	.045	.034	.026	.020	.016
29	.534	.353	.243	.172	.124	.091	.068	.051	.039	.030	.023	.018
30	.548	.367	.256	.183	.134	.099	.074	.056	.043	.034	.026	.021
40	.649	.485	.372	.290	.229	.182	.146	.118	.096	.079	.065	.054
60	.758	.627	.527	.447	.381	.327	.282	.244	.212	.184	.161	.141
80	.815	.709	.623	.551	.489	.435	.389	.348	.313	.281	.253	.229
100	.851	.761	.687	.622	.566	.516	.471	.431	.395	.362	.333	.306
120	.875	.798	.732	.675	.623	.577	.535	.496	.461	.429	.399	.372
140	.892	.825	.767	.715	.667	.625	.585	.549	.515	.484	.455	.428
170	.911	.854	.804	.759	.717	.679	.644	.610	.579	.550	.523	.497
200	.924	.875	.831	.791	.755	.720	.688	.657	.629	.602	.576	.551
240	.936	.895	.858	.823	.791	.761	.732	.705	.679	.655	.631	.609
320	.952	.920	.891	.865	.839	.815	.792	.770	.748	.728	.708	.689
440	.965	.942	.920	.900	.880	.862	.844	.827	.810	.794	.778	.762
600	.974	.957	.941	.926	.911	.897	.883	.870	.857	.844	.831	.819
800	.981	.968	.955	.944	.933	.922	.911	.901	.890	.880	.871	.861
1000	.985	.974	.964	.955	.946	.937	.928	.920	.911	.903	.895	.887

^aMultiply entry by 10⁻³.

Table A.10 Upper Critical Values for Roy's Test, $\alpha = .05$

Roy's test statistic is given by

$$\theta = \frac{\lambda_1}{1 + \lambda_1},$$

where λ_1 is the largest eigenvalue of $\mathbf{E}^{-1}\mathbf{H}$. The parameters are

$$s = \min(\nu_H, p), \quad m = \frac{|\nu_H - p| - 1}{2}, \quad N = \frac{\nu_E - p - 1}{2}.$$

Reject H_0 if $\theta >$ table value.

N	m								
	0	1	2	3	4	5	7	10	15
<i>s = 2</i>									
5	.565	.651	.706	.746	.776	.799	.834	.868	.901
10	.374	.455	.514	.561	.598	.629	.679	.732	.789
15	.278	.348	.402	.446	.483	.515	.567	.627	.696
20	.221	.281	.329	.369	.404	.434	.486	.546	.620
25	.184	.236	.278	.314	.346	.375	.424	.484	.558
30	.157	.203	.241	.274	.303	.330	.376	.433	.507
40	.122	.159	.190	.218	.243	.266	.306	.359	.428
50	.099	.130	.157	.180	.202	.222	.259	.306	.370
60	.084	.110	.133	.154	.173	.191	.223	.266	.326
80	.064	.085	.103	.119	.135	.149	.176	.211	.263
120	.043	.058	.070	.082	.093	.104	.123	.150	.190
240	.022	.030	.036	.042	.048	.054	.065	.080	.103
<i>s = 3</i>									
5	.669	.729	.770	.800	.822	.840	.867	.894	.920
10	.472	.537	.586	.625	.656	.683	.725	.770	.819
15	.362	.422	.469	.508	.541	.569	.616	.669	.730
20	.293	.346	.390	.427	.458	.486	.533	.589	.656
25	.246	.294	.333	.367	.397	.424	.470	.525	.594
30	.212	.255	.291	.322	.350	.375	.419	.473	.543
40	.166	.201	.232	.259	.283	.305	.345	.395	.462
50	.136	.167	.192	.216	.237	.257	.292	.339	.402
60	.116	.142	.164	.185	.204	.221	.254	.296	.355
80	.089	.109	.127	.144	.160	.174	.201	.237	.288
120	.061	.075	.088	.100	.111	.122	.142	.169	.209
240	.031	.039	.046	.052	.058	.064	.075	.090	.114
<i>s = 4</i>									
5	.739	.782	.813	.836	.854	.868	.889	.911	.933
10	.547	.601	.641	.674	.700	.723	.759	.798	.840
15	.431	.482	.523	.558	.587	.612	.654	.701	.756
20	.354	.402	.441	.474	.503	.529	.572	.623	.684
25	.301	.344	.380	.412	.440	.464	.507	.559	.624
30	.261	.301	.334	.364	.390	.414	.455	.507	.572
40	.207	.240	.269	.294	.318	.339	.377	.426	.490
50	.171	.199	.224	.247	.268	.287	.322	.367	.428
60	.145	.170	.193	.213	.232	.249	.280	.322	.380
80	.112	.132	.150	.167	.182	.196	.223	.259	.309
120	.077	.091	.104	.116	.127	.138	.158	.185	.226
240	.040	.047	.054	.061	.067	.073	.084	.100	.124

(continued)

Table A.10 (*Continued*)

N	<i>m</i>								
	0	1	2	3	4	5	7	10	15
<i>s</i> = 5									
5	.788	.821	.845	.863	.877	.888	.906	.924	.942
10	.607	.651	.685	.713	.735	.755	.786	.820	.857
15	.488	.533	.569	.599	.625	.648	.685	.728	.777
20	.407	.449	.485	.515	.542	.565	.604	.651	.708
25	.349	.388	.422	.451	.477	.500	.540	.588	.648
30	.305	.341	.373	.400	.425	.448	.487	.535	.597
40	.243	.275	.302	.327	.349	.370	.406	.453	.514
50	.202	.230	.254	.276	.296	.315	.348	.392	.451
60	.173	.197	.219	.238	.257	.274	.304	.345	.401
80	.134	.154	.171	.188	.203	.217	.243	.278	.329
120	.093	.107	.120	.132	.143	.154	.174	.201	.241
240	.048	.056	.063	.069	.076	.082	.093	.109	.134
<i>s</i> = 6									
5	.825	.850	.869	.883	.895	.904	.918	.934	.949
10	.655	.692	.721	.744	.764	.781	.808	.838	.871
15	.537	.576	.608	.635	.658	.678	.711	.750	.795
20	.454	.491	.523	.551	.575	.596	.632	.676	.728
25	.392	.428	.458	.485	.509	.531	.568	.613	.669
30	.345	.378	.407	.433	.457	.478	.514	.560	.618
40	.278	.307	.333	.356	.378	.397	.432	.477	.536
50	.232	.258	.281	.302	.322	.340	.372	.414	.472
60	.200	.223	.243	.262	.280	.297	.327	.366	.421
80	.156	.174	.192	.208	.222	.236	.262	.297	.346
120	.108	.122	.134	.146	.157	.168	.188	.215	.255
240	.056	.064	.071	.078	.084	.090	.101	.118	.142
<i>s</i> = 7									
5	.852	.872	.887	.899	.908	.917	.929	.941	.955
10	.695	.726	.750	.771	.788	.802	.826	.853	.882
15	.579	.613	.641	.665	.686	.704	.734	.769	.810
20	.494	.528	.557	.582	.604	.624	.657	.697	.745
25	.431	.463	.491	.516	.538	.558	.593	.635	.688
30	.381	.412	.439	.463	.485	.505	.540	.583	.638
40	.309	.337	.362	.384	.404	.423	.456	.499	.555
60	.224	.246	.266	.285	.302	.318	.347	.386	.439
80	.176	.194	.211	.226	.241	.255	.280	.314	.363
100	.145	.160	.175	.188	.200	.212	.235	.265	.310
200	.077	.085	.093	.101	.109	.116	.129	.148	.175
300	.052	.058	.064	.069	.074	.079	.089	.103	.125
500	.032	.036	.039	.042	.046	.049	.055	.064	.078
1000	.016	.018	.020	.022	.023	.025	.028	.033	.041

(continued)

Table A.10 (*Continued*)

N	<i>m</i>								
	0	1	2	3	4	5	7	10	15
<i>s</i> = 8									
5	.874	.890	.902	.912	.920	.927	.937	.948	.959
10	.728	.754	.775	.793	.808	.821	.842	.865	.892
15	.615	.645	.670	.692	.710	.727	.754	.786	.824
20	.531	.561	.587	.610	.630	.648	.679	.716	.761
25	.466	.495	.521	.544	.565	.583	.616	.655	.705
30	.414	.443	.468	.491	.511	.530	.563	.603	.655
40	.339	.365	.388	.409	.428	.446	.478	.519	.573
60	.248	.269	.288	.306	.323	.338	.367	.404	.456
80	.195	.213	.229	.244	.259	.272	.297	.330	.378
100	.161	.176	.190	.203	.216	.228	.250	.279	.323
200	.086	.094	.103	.110	.118	.125	.138	.157	.185
300	.058	.065	.070	.076	.081	.086	.096	.109	.130
500	.036	.040	.043	.047	.050	.053	.059	.068	.081
1000	.018	.020	.022	.024	.025	.027	.030	.035	.042
<i>s</i> = 9									
5	.891	.904	.914	.922	.929	.935	.944	.953	.963
10	.756	.778	.797	.812	.825	.837	.855	.876	.901
15	.647	.674	.696	.715	.732	.747	.771	.801	.835
20	.563	.591	.614	.635	.654	.670	.698	.733	.775
25	.497	.525	.549	.570	.589	.606	.636	.673	.720
30	.445	.471	.495	.516	.535	.552	.583	.622	.671
40	.366	.391	.413	.433	.451	.468	.499	.538	.590
60	.270	.291	.309	.326	.343	.358	.385	.421	.472
80	.214	.231	.247	.262	.276	.289	.313	.346	.392
100	.177	.192	.206	.219	.231	.242	.264	.293	.336
200	.095	.104	.112	.119	.127	.134	.147	.166	.194
300	.065	.071	.077	.082	.087	.092	.102	.115	.136
500	.040	.043	.047	.051	.054	.057	.063	.072	.086
1000	.020	.022	.024	.026	.028	.029	.032	.037	.044
<i>s</i> = 10									
5	.905	.916	.924	.931	.937	.941	.949	.958	.967
10	.780	.799	.815	.829	.840	.851	.867	.886	.908
15	.675	.699	.719	.736	.751	.764	.787	.814	.846
20	.592	.617	.639	.658	.675	.690	.716	.748	.787
25	.526	.551	.573	.593	.611	.627	.655	.690	.734
30	.473	.497	.519	.539	.557	.573	.603	.639	.686
40	.392	.415	.436	.455	.473	.489	.518	.555	.605
60	.292	.311	.329	.346	.361	.376	.402	.438	.487
80	.232	.249	.264	.278	.292	.305	.329	.361	.406
100	.193	.207	.220	.233	.245	.256	.278	.306	.348
200	.104	.112	.120	.128	.135	.142	.156	.174	.202
300	.071	.077	.083	.088	.093	.098	.108	.122	.143
500	.044	.047	.051	.054	.058	.061	.067	.076	.090
1000	.022	.024	.026	.028	.030	.031	.034	.039	.047

Table A.11 Upper Critical Values of Pillai's Statistic $V^{(s)}$, $\alpha = .05$

$$V^{(s)} = \sum_{i=1}^s \frac{\lambda_i}{1 + \lambda_i},$$

where $\lambda_1, \lambda_2, \dots, \lambda_s$ are eigenvalues of $E^{-1}H$. Reject H_0 if $V^{(s)}$ exceeds table value. The parameters s , m , and N are defined in Table A.10.

m	0	N										
		1	2	3	4	5	6	7	8	9	10	15
0	1.536	1.232	1.031	.890	.782	.698	.629	.573	.526	.485	.451	.333
1	1.706	1.452	1.258	.910	.991	.896	.817	.751	.694	.646	.604	.455
2	1.784	1.573	1.397	1.254	1.137	1.039	.956	.886	.825	.772	.725	.556
3	1.829	1.649	1.492	1.358	1.245	1.149	1.065	.993	.930	.875	.825	.643
4	1.859	1.703	1.560	1.436	1.329	1.235	1.153	1.081	1.018	.961	.910	.719
5	1.880	1.742	1.613	1.497	1.395	1.305	1.226	1.155	1.091	1.034	.983	.786
6	1.895	1.772	1.654	1.546	1.450	1.364	1.286	1.217	1.154	1.098	1.046	.846
7	1.907	1.796	1.687	1.586	1.495	1.413	1.338	1.270	1.209	1.153	1.102	.901
8	1.917	1.815	1.714	1.620	1.534	1.455	1.383	1.317	1.257	1.202	1.151	.950
9	1.924	1.831	1.737	1.649	1.567	1.491	1.422	1.358	1.299	1.245	1.195	.851
10	1.931	1.844	1.757	1.673	1.595	1.523	1.456	1.394	1.337	1.284	1.235	.891
15	1.951	1.888	1.822	1.758	1.695	1.636	1.580	1.527	1.477	1.430	1.386	1.036
20	1.963	1.913	1.860	1.807	1.756	1.706	1.658	1.612	1.568	1.527	1.487	.781
25	1.969	1.929	1.885	1.840	1.796	1.753	1.711	1.671	1.632	1.595	1.559	.506

(continued)

(continued)

Table A.11 (Continued)

<i>m</i>	<i>N</i>									
	0	1	2	3	4	5	6	7	8	9
0	2.037	1.710	1.473	1.294	1.153	1.040	.947	.869	.803	.746
1	2.297	1.988	1.751	1.564	1.412	1.287	1.183	1.094	1.017	.950
2	2.447	2.168	1.943	1.759	1.606	1.477	1.367	1.273	1.190	1.117
3	2.544	2.294	2.084	1.907	1.757	1.628	1.517	1.420	1.334	1.258
4	2.612	2.386	2.191	2.023	1.878	1.752	1.641	1.543	1.456	1.378
5	2.662	2.457	2.276	2.117	1.978	1.854	1.745	1.648	1.561	1.482
6	2.701	2.514	2.345	2.194	2.061	1.941	1.835	1.739	1.652	1.573
7	2.732	2.559	2.402	2.259	2.131	2.016	1.912	1.818	1.732	1.654
8	2.757	2.597	2.449	2.314	2.192	2.081	1.979	1.887	1.803	1.726
9	2.777	2.629	2.490	2.362	2.244	2.137	2.039	1.949	1.866	1.790
10	2.795	2.656	2.525	2.403	2.291	2.187	2.092	2.004	1.923	1.848
15	2.833	2.748	2.646	2.549	2.457	2.370	2.288	2.211	2.139	2.071
20	2.885	2.802	2.718	2.637	2.560	2.485	2.414	2.347	2.283	2.222
25	2.906	2.836	2.766	2.697	2.630	2.565	2.503	2.443	2.385	2.322
0	2.549	2.194	1.926	1.717	1.548	1.410	1.294	1.196	1.112	1.038
1	2.852	2.510	2.241	2.023	1.844	1.693	1.566	1.456	1.360	1.277
2	3.052	2.733	2.472	2.256	2.074	1.919	1.786	1.670	1.567	1.477
3	3.193	2.898	2.650	2.440	2.260	2.104	1.969	1.849	1.743	1.649
4	3.298	3.025	2.791	2.589	2.413	2.259	2.123	2.002	1.895	1.798
5	3.378	3.126	2.905	2.711	2.541	2.390	2.255	2.135	2.027	1.929
6	3.442	3.208	2.999	2.814	2.649	2.502	2.370	2.251	2.143	2.044
7	3.494	3.276	3.079	2.902	2.743	2.600	2.470	2.353	2.246	2.148
8	3.537	3.333	3.146	2.977	2.824	2.685	2.559	2.444	2.338	2.241
9	3.574	3.382	3.205	3.043	2.896	2.761	2.638	2.525	2.421	2.325
10	3.605	3.424	3.256	3.101	2.959	2.829	2.708	2.598	2.496	2.401
15	3.710	3.570	3.436	3.310	3.191	3.079	2.974	2.876	2.783	2.696
20	3.771	3.657	3.546	3.440	3.338	3.241	3.149			

Table A.11 (Continued)

m	N												
	0	1	2	3	4	5	6	7	8	9	10	15	20
0	3.055	2.681	2.389	2.155	1.962	1.801	1.664	1.547	1.445	1.356	1.277		
1	3.390	3.025	2.731	2.488	2.285	2.122	1.964	1.835	1.722	1.622	1.533		
2	3.628	3.281	2.993	2.751	2.545	2.367	2.213	2.077	1.957	1.850	1.754		
3	3.805	3.478	3.201	2.964	2.759	2.580	2.423	2.284	2.160	2.048	1.948		
4	3.941	3.635	3.370	3.140	2.938	2.761	2.604	2.463	2.337	2.222	2.119		
5	4.050	3.762	3.510	3.288	3.091	2.916	2.760	2.619	2.492	2.377	2.271		
6	4.138	3.868	3.627	3.414	3.223	3.052	2.897	2.758	2.630	2.514	2.408		
7	4.212	3.957	3.728	3.522	3.337	3.170	3.018	2.880					
8	4.274	4.033	3.815	3.617	3.438	3.275	3.126						
9	4.327	4.099	3.890	3.700	3.527	3.369							
10	4.372	4.156	3.957	3.774	3.607	3.45							
							$s = 6$						
0	3.559	3.171	2.859	2.604	2.390	2.209	2.053	1.918	1.799	1.694	1.601		
1	3.917	3.535	3.221	2.958	2.734	2.542	2.375	2.229	2.099	1.984	1.881		
2	4.185	3.817	3.508	3.245	3.018	2.821	2.647	2.494	2.358	2.235	2.125		
3	4.391	4.041	3.741	3.482	3.256	3.057	2.881	2.724	2.583	2.456	2.341		
4	4.556	4.223	3.934	3.681	3.458	3.260	3.084	2.925	2.782	2.652	2.534		
5	4.690	4.375	4.097	3.851	3.633	3.438	3.262	3.103	2.959	2.827	2.706		
6	4.802	4.502	4.236	3.998	3.785								
7	4.896	4.611	4.356	4.126									
8	4.976	4.706	4.461										
9	5.045	4.788	4.553										
10	5.106	4.860	4.635										

Table A.12 Upper Critical Values for the Lawley–Hotelling Statistic, $\alpha = .05$

ν_E	2	3	4	5	6	8	10	12	15	20	25	40	60
2 ^a	9.8591	10.659	11.098	11.373	11.562	11.952	11.804	12.052	12.153	12.254	12.316	12.409	12.461
3	58.428	58.915	59.161	59.308	59.407	59.531	59.606	59.655	59.705	59.755	59.785	59.830	59.855
4	23.999	23.312	22.918	22.663	22.484	22.250	22.104	22.003	21.901	21.797	21.733	21.636	21.582
5	15.639	14.864	14.422	14.135	13.934	13.670	13.504	13.391	13.275	13.156	13.083	12.972	12.909
6	12.175	11.411	10.975	10.691	10.491	10.228	10.063	9.9489	9.8320	9.7118	9.6381	9.5251	9.4610
7	10.334	9.5937	9.1694	8.8927	8.6975	8.4396	8.2765	8.16399	8.0480	7.9285	7.8549	7.7417	7.6773
8	9.2069	8.4881	8.0752	7.8054	7.6145	7.3614	7.2008	7.0896	6.9748	6.8560	6.7826	6.6694	6.6048
10	7.9095	7.2243	6.8294	6.5702	6.3860	6.1405	5.9837	5.8745	5.7612	5.6433	5.5701	5.4564	5.3910
12	7.1902	6.5284	6.1461	5.8942	5.7147	5.4744	5.3200	5.2122	5.0997	4.9820	4.9085	4.7938	4.7274
14	6.7350	6.0902	4.7168	4.5703	5.2941	5.0574	4.9048	4.7977	4.6836	4.5678	4.4939	4.3780	4.3105
16	6.4217	5.7895	5.4230	5.1804	5.0067	4.7727	4.6213	4.5147	4.4028	4.2846	4.2102	4.0930	4.0243
18	6.1932	5.5708	5.2095	4.9700	4.7982	4.5663	4.4157	4.3094	4.1976	4.0791	4.0042	3.8855	3.8158
20	6.0192	5.4046	5.0475	4.8105	4.6402	4.4099	4.2690	4.1539	4.0420	3.9231	3.8471	3.7278	3.6569
25	5.7244	5.1237	4.7741	4.5415	4.3740	4.1465	3.9977	3.8919	3.7798	3.6598	3.5832	3.4605	3.3868
30	5.5401	4.9487	4.6040	4.3743	4.2086	3.9829	3.8347	3.7291	3.6166	3.4957	3.4181	3.2926	3.2168
35	5.4140	4.8291	4.8880	4.2604	4.0959	3.8715	3.7237	3.6181	3.5054	3.3836	3.3051	3.1774	3.1000
40	5.3224	4.7424	4.4039	4.1778	4.0143	3.7908	3.6433	3.5377	3.4247	3.3022	3.2230	3.0933	3.0140
50	5.1981	4.6249	4.2900	4.0661	3.9039	3.6817	3.5346	3.4289	3.3154	3.1919	3.1115	2.9787	2.8965
60	5.1178	4.5490	4.2166	3.9941	3.8328	3.6114	3.4646	3.3588	3.2450	3.1206	3.0392	2.9041	2.8196
70	5.0616	4.4960	4.1653	3.9439	3.7831	3.5624	3.4157	3.3099	3.1957	3.0706	2.9886	2.8516	2.7652
80	5.0200	4.4569	4.1275	3.9068	3.7465	3.5262	3.3796	3.2737	3.1594	3.0338	2.9512	2.8126	2.7247
100	4.9628	4.4030	4.0754	3.8557	3.6961	3.4764	3.3200	3.2240	3.1093	2.9829	2.8994	2.7586	2.6683
200	4.8514	4.2982	3.9742	3.7567	3.5983	3.3798	3.2336	3.1275	3.0120	2.8838	2.7984	2.6520	2.5559
∞	4.7442	4.1973	3.8769	3.6614	3.5044	3.2870	3.1410	3.0346	2.9182	2.7879	2.7002	2.5470	2.4428

^aMultiply each entry in this row by 100.

(continued)

Table A.12 (Continued)

ν_E	3	4	5	6	8	10	12	15	20	25	40	60
3 ^a	25.930	26.996	27.665	28.125	28.712	29.073	29.316	29.561	29.809	29.959	30.19	30.31
4 ^a	1.1880	1.1929	1.1959	1.1978	1.2003	1.2018	1.2028	1.2038	1.2048	1.2054	1.2063	1.2068
5	42.474	41.764	1.305	40.983	40.562	40.360	40.120	39.937	39.750	39.635	39.462	39.366
6	25.456	24.715	24.235	23.899	23.458	23.182	22.992	22.799	22.600	22.479	22.294	22.190
7	18.752	18.056	17.605	17.288	16.870	16.668	16.427	16.241	16.051	15.934	15.755	15.653
8	15.308	14.657	14.233	13.934	13.540	13.290	13.118	12.941	12.758	12.646	12.473	12.375
10	11.893	11.306	10.921	10.649	10.287	10.057	9.8974	9.7320	9.5603	9.4541	9.2897	9.1955
12	10.229	9.6825	9.3234	9.0680	8.7271	8.5088	8.3566	8.1982	8.0330	7.9301	7.7700	7.6777
14	9.2550	8.7356	8.3935	8.1495	7.8225	7.6122	7.4649	7.3110	7.1497	7.0488	6.8908	6.7991
16	8.6180	8.1183	7.7884	7.5526	7.2355	7.0307	6.8868	6.7360	6.5772	6.4774	6.3204	6.2287
18	8.1701	7.6851	7.3644	7.1347	6.8251	6.6244	6.4830	6.3343	6.1771	6.0780	5.9212	5.8292
20	7.8384	7.3649	7.0513	6.8263	6.5224	6.3249	6.1853	6.0383	5.8822	5.7834	5.6266	5.5341
25	7.2943	6.8407	6.5394	6.3227	6.0287	5.8365	5.7001	5.5555	5.4010	5.3025	5.1446	5.0503
30	6.9654	6.5245	6.2311	6.0196	5.7319	5.5431	5.4085	5.2654	5.1116	5.0129	4.8555	4.7575
35	6.7453	6.3132	6.0253	5.8175	5.5341	5.3476	5.2143	5.0720	4.9185	4.8195	4.6586	4.5608
40	6.5877	6.1621	5.8783	5.6732	5.3929	5.2081	5.0757	4.9340	4.7806	4.6813	4.5189	4.4195
50	6.3773	5.9606	5.6823	5.4809	5.2050	5.0224	4.8911	4.7502	4.5967	4.4968	4.3319	4.2297
60	6.2433	5.8324	5.5577	5.3587	5.0856	4.9044	4.7739	4.6334	4.4798	4.3793	4.2123	4.1078
70	6.1504	5.7436	5.4715	5.2742	5.0031	4.8229	4.6929	4.5526	4.3988	4.2979	4.1292	4.0227
80	6.0823	5.6786	5.4084	5.2122	4.9426	4.7632	4.6336	4.4935	4.3395	4.2581	4.0680	3.9600
100	5.9891	5.5896	5.3220	5.1276	4.8601	4.6817	4.5525	4.4126	4.2583	4.1563	3.9840	3.8734
200	5.8099	5.4186	5.1562	4.9653	4.7017	4.5252	4.3970	4.2574	4.1023	3.9888	3.8212	3.7042
∞	5.6397	5.2565	4.9992	4.8116	4.5519	4.3773	4.2499	4.1104	3.9541	3.8487	3.6642	3.5384

^aMultiply each entry in this row by 100.

(continued)

Table A.12 (Continued)

ν_E	ν_H						$p = 4$					
	4	5	6	8	10	12	15	20	25	30	40	60
4 ^a	49.964	51.204	52.054	53.142	53.808	54.258	54.71	55.17	55.46	—	—	—
5 ^a	1.9964	2.0013	2.0046	2.0087	2.0112	2.0128	2.0145	2.0171	2.019	—	—	—
6	65.715	64.999	64.497	63.841	63.432	63.151	62.866	62.573	62.396	62.13	—	—
7	37.343	36.629	36.129	35.474	35.064	34.782	34.495	34.200	34.019	33.75	—	—
8	26.516	25.868	25.413	24.814	24.437	24.178	23.912	23.639	23.471	23.214	23.072	—
10	17.875	17.326	16.938	16.424	16.098	15.872	15.640	15.399	15.250	15.021	14.891	—
12	14.338	13.848	13.500	13.037	12.741	12.535	12.321	12.099	11.961	11.747	11.624	—
14	12.455	12.002	11.680	11.248	10.972	10.778	10.577	10.366	10.234	10.029	9.9103	—
16	11.295	10.868	10.563	10.154	9.8904	9.7054	9.5119	9.3085	9.1810	8.9808	8.8644	—
18	10.512	10.104	9.8121	9.4190	9.1647	8.9857	8.7978	8.5996	8.4748	8.2778	8.1626	—
20	9.9500	9.5560	9.2736	8.8926	8.6453	8.4708	8.2871	8.0926	7.9696	7.7748	7.6601	—
25	9.0585	8.6884	8.4223	8.0616	7.8261	7.6590	7.4821	7.2933	7.1750	6.9805	6.8659	—
30	8.5377	8.1825	7.9265	7.5784	7.3502	7.1876	7.0147	6.8291	6.7101	6.5181	6.4226	—
35	8.1968	7.8517	7.6026	7.2631	7.0397	6.8801	6.7099	6.5262	6.4079	6.2156	6.0989	—
40	7.9566	7.6188	7.3746	7.0413	6.8214	6.6640	6.4955	6.3131	6.1952	6.0023	5.8844	—
50	7.6404	7.3125	7.0751	6.7501	6.5350	6.3804	6.2143	6.0334	5.9157	5.7214	5.6011	—
60	7.4417	7.1202	6.8872	6.5676	6.3555	6.2027	6.0381	5.8581	5.7403	5.5446	5.4222	—
70	7.3054	6.9884	6.7584	6.4426	6.2325	6.0809	5.9173	5.7378	5.6200	5.4230	5.2987	—
80	7.2061	6.8924	6.6646	6.3515	6.1430	5.9924	5.8294	5.6503	5.5323	5.3343	5.2084	—
100	7.071	6.7619	6.5372	6.2279	6.0215	5.8721	5.7101	5.5313	5.4131	5.2133	5.0849	—
200	6.8143	6.5139	6.2952	5.9933	5.7910	5.6439	5.4836	5.3053	5.1863	4.9819	4.8471	—
∞	6.5741	6.2821	6.0692	5.7743	5.5758	5.4309	5.2721	5.0940	4.9737	4.7639	4.6190	—

^aMultiply each entry in this row by 100.

(continued)

Table A.12 (Continued)

ν_E	5	6	8	10	12	15	20	25	40	60
ν_H										
						$p = 5$				
5 ^a	81.991	83.352	85.093	86.160	86.88	—	—	—	—	—
6 ^a	3.0093	3.0142	3.0204	3.0241	3.0266	3.0291	3.032	—	—	—
7	93.762	93.042	92.102	91.515	91.113	90.705	90.29	90.04	—	—
8	51.339	50.646	49.739	49.170	48.780	48.382	47.973	47.723	47.35	—
10	27.667	27.115	26.387	25.927	25.610	25.284	24.947	24.740	24.422	—
12	20.169	19.701	19.079	18.683	18.409	18.124	17.830	17.647	17.365	17.20
14	16.643	16.224	15.666	15.309	15.059	14.800	14.530	14.361	14.100	13.95
16	14.624	14.239	13.722	13.389	13.157	12.914	12.659	12.499	12.250	12.105
18	13.326	12.963	12.476	12.161	11.939	11.708	11.463	11.310	11.068	10.928
20	12.424	12.078	11.612	11.310	11.097	10.874	10.637	10.488	10.252	10.113
25	11.046	10.728	10.297	10.016	9.8168	9.6061	9.3814	9.2386	9.0102	8.8745
30	10.270	9.9689	9.5592	9.2907	9.0995	8.8964	8.6785	8.5389	8.3141	8.1790
35	9.7739	9.4836	9.0879	8.8277	8.6419	8.4437	8.2301	8.0926	7.8693	7.7339
40	9.4292	9.1469	8.7613	8.5070	8.3250	8.1303	7.9195	7.7833	7.5607	7.4247
50	8.9825	8.7107	8.3385	8.0921	7.9150	7.7248	7.5177	7.3829	7.1605	7.0229
60	8.7057	8.4406	8.0769	7.8355	7.6615	7.4741	7.2692	7.1351	6.9124	6.7730
70	8.5174	8.2570	7.8991	7.6612	7.4894	7.3039	7.1004	6.9667	6.7434	6.6024
80	8.3811	8.1241	7.7705	7.5351	7.3648	7.1807	6.9782	6.8448	6.6208	6.4785
100	8.1969	7.9446	7.5969	7.3649	7.1968	7.0145	6.8133	6.6801	6.4550	6.3103
200	7.8505	7.6070	7.2706	7.0451	6.8811	6.7023	6.5032	6.3702	6.1416	5.9008
∞	7.5305	7.2955	6.9698	6.7505	6.5902	6.4144	6.2171	6.0838	5.8499	5.6899

^aMultiply each entry in this row by 100.

(continued)

Table A.12 (*Continued*)

ν_E	6	8	10	12	15	20	25	40	60
10	45.722	44.677	44.019	43.567	43.103	42.626	42.334	42.136	41.993
12	28.959	28.121	27.590	27.223	26.843	26.451	26.209	26.044	25.925
14	22.321	21.600	21.141	20.821	20.489	20.144	19.929	19.783	19.677
16	18.838	18.210	17.795	17.505	17.202	16.886	16.688	16.553	16.455
18	16.755	16.157	15.772	15.501	15.218	14.921	14.735	14.607	14.513
20	15.351	14.788	14.424	14.168	13.899	13.615	13.436	13.313	13.223
25	13.293	12.786	12.456	12.222	11.975	11.711	11.544	11.428	11.343
30	12.180	11.705	11.395	11.173	10.939	10.687	10.526	10.414	10.331
35	11.484	11.031	10.733	10.520	10.293	10.049	9.8921	9.7820	9.7003
40	11.009	10.571	10.282	10.075	9.8535	9.6142	9.4596	9.3508	9.2699
50	10.402	9.9832	9.7060	9.5067	9.2927	9.0598	8.9082	8.8009	8.7207
60	10.031	9.6246	9.3547	9.1602	8.9507	8.7215	8.5717	8.4651	8.3851
70	9.7813	9.3830	9.1182	8.9269	8.7204	8.4938	8.3450	8.2388	8.1589
80	9.6014	9.2093	8.9480	8.7591	8.5548	8.3300	8.1819	8.0759	7.9959
100	9.3598	8.9760	8.7197	8.5340	8.3326	8.1102	7.9629	7.8572	7.7771
200	8.9099	8.5419	8.2950	8.1153	7.9193	7.7011	7.5552	7.4494	7.3685
∞	8.4997	8.1463	7.9082	7.7340	7.5430	7.3284	7.1832	7.0768	6.9945

Table A.13 Orthogonal Polynomial Contrasts

p	Polynomial	Variable										$\mathbf{c}'_i \mathbf{c}_i$
		1	2	3	4	5	6	7	8	9	10	
3	Linear	-1	0	1								2
	Quadratic	1	-2	1								6
4	Linear	-3	-1	1	3							20
	Quadratic	1	-1	-1	1							4
	Cubic	-1	3	-3	1							20
5	Linear	-2	-1	0	1	2						10
	Quadratic	2	-1	-2	-1	2						14
	Cubic	-1	2	0	-2	1						10
	Quartic	1	-4	6	-4	1						70
6	Linear	-5	-3	-1	1	3	5					70
	Quadratic	5	-1	-4	-4	-1	5					84
	Cubic	-5	7	4	-4	-7	5					180
	Quartic	1	-3	2	2	-3	1					28
	Quintic	-1	5	-10	10	-5	1					252
7	Linear	-3	-2	-1	0	1	2	3				28
	Quadratic	5	0	-3	-4	-3	0	5				84
	Cubic	-1	1	1	0	-1	-1	1				6
	Quartic	3	-7	1	6	1	-7	3				154
	Quintic	-1	4	-5	0	5	-4	1				84
	Sextic	1	-6	15	-20	15	-6	1				924
8	Linear	-7	-5	-3	-1	1	3	5	7			168
	Quadratic	7	1	-3	-5	-5	-3	1	7			168
	Cubic	-7	5	7	3	-3	-7	-5	7			264
	Quartic	7	-13	-3	9	9	-3	-13	7			616
	Quintic	-7	23	-17	-15	15	17	-23	7			2,184
	Sextic	1	-5	9	-5	-5	9	-5	1			264
	Septic	-1	7	-21	35	-35	21	-7	1			3,432
9	Linear	-4	-3	-2	-1	0	1	2	3	4		60
	Quadratic	28	7	-8	-17	-20	-17	-8	7	28		2,772
	Cubic	-14	7	13	9	0	-9	-13	-7	14		990
	Quartic	14	-21	-11	9	18	9	-11	-21	14		2,002
	Quintic	-4	11	-4	-9	0	9	4	-11	4		468
	Sextic	4	-17	22	1	-20	1	22	-17	4		1,980
	Septic	-1	6	-14	14	0	-14	14	-6	1		858
	Octic	1	-8	28	-56	70	-56	28	-8	1		12,870
10	Linear	-9	-7	-5	-3	-1	1	3	5	7	9	330
	Quadratic	6	2	-1	-3	-4	-4	-3	-1	2	6	132
	Cubic	-42	14	35	31	12	-12	-31	-35	-14	42	8,580
	Quartic	18	-22	-17	3	18	18	3	-17	-22	18	2,860
	Quintic	-6	14	-1	-11	-6	6	11	1	-14	6	780
	Sextic	3	-11	10	6	-8	-8	6	10	11	3	660
	Septic	-9	47	-86	92	56	-56	-42	86	-47	9	29,172
	Octic	1	-7	20	-28	14	14	-28	20	-7	1	2,860
	Novic	-1	9	-36	84	-126	126	-84	36	-9	1	48,620

Note: Entries are rows \mathbf{c}'_i of the $(p - 1) \times p$ matrix \mathbf{C} illustrated in (6.90) in Section 6.10.1

Table A.14 Test for Equal Covariance Matrices, $\alpha = 0.05$

ν	$k = 2$	$k = 3$	$k = 4$	$k = 5$	$k = 6$	$k = 7$	$k = 8$	$k = 9$	$k = 10$
$p = 2$									
3	12.18	18.70	24.55	30.09	35.45	40.68	45.81	50.87	55.86
4	10.70	16.65	22.00	27.07	31.97	36.75	41.45	46.07	50.64
5	9.97	15.63	20.73	25.57	30.23	34.79	39.26	43.67	48.02
6	9.53	15.02	19.97	24.66	29.19	33.61	37.95	42.22	46.45
7	9.24	14.62	19.46	24.05	28.49	32.83	37.08	41.26	45.40
8	9.04	14.33	19.10	23.62	27.99	32.26	36.44	40.57	44.64
9	8.88	14.11	18.83	23.30	27.62	31.84	35.98	40.05	44.08
10	8.76	13.94	18.61	23.05	27.33	31.51	35.61	39.65	43.64
11	8.67	13.81	18.44	22.85	27.10	31.25	35.32	39.33	43.29
12	8.59	13.70	18.30	22.68	26.90	31.03	35.08	39.07	43.00
13	8.52	13.60	18.19	22.54	26.75	30.85	34.87	38.84	42.76
14	8.47	13.53	18.10	22.42	26.61	30.70	34.71	38.66	42.56
15	8.42	13.46	18.01	22.33	26.50	30.57	34.57	38.50	42.38
16	8.38	13.40	17.94	22.24	26.40	30.45	34.43	38.36	42.23
17	8.35	13.35	17.87	22.17	26.31	30.35	34.32	38.24	42.10
18	8.32	13.30	17.82	22.10	26.23	30.27	34.23	38.13	41.99
19	8.28	13.26	17.77	22.04	26.16	30.19	34.14	38.04	41.88
20	8.26	13.23	17.72	21.98	26.10	30.12	34.07	37.95	41.79
25	8.17	13.10	17.55	21.79	25.87	29.86	33.78	37.63	41.44
30	8.11	13.01	17.44	21.65	25.72	29.69	33.59	37.42	41.21
$p = 3$									
4	22.41	35.00	46.58	57.68	68.50	79.11	89.60	99.94	110.21
5	19.19	30.52	40.95	50.95	60.69	70.26	79.69	89.03	98.27
6	17.57	28.24	38.06	47.49	56.67	65.69	74.58	83.39	92.09
7	16.59	26.84	36.29	45.37	54.20	62.89	71.44	79.90	88.30
8	15.93	25.90	35.10	43.93	52.54	60.99	69.32	77.57	85.73
9	15.46	25.22	34.24	42.90	51.33	59.62	67.78	75.86	83.87
10	15.11	24.71	33.59	42.11	50.42	58.57	66.62	74.58	82.46
11	14.83	24.31	33.08	41.50	49.71	57.76	65.71	73.57	81.36
12	14.61	23.99	32.67	41.00	49.13	57.11	64.97	72.75	80.45
13	14.43	23.73	32.33	40.60	48.65	56.56	64.36	72.09	79.72
14	14.28	23.50	32.05	40.26	48.26	56.11	63.86	71.53	79.11
15	14.15	23.32	31.81	39.97	47.92	55.73	63.43	71.05	78.60
16	14.04	23.16	31.60	39.72	47.63	55.40	63.06	70.64	78.14
17	13.94	23.02	31.43	39.50	47.38	55.11	62.73	70.27	77.76
18	13.86	22.89	31.26	39.31	47.16	54.86	62.45	69.97	77.41
19	13.79	22.78	31.13	39.15	46.96	54.64	62.21	69.69	77.11
20	13.72	22.69	31.01	39.00	46.79	54.44	61.98	69.45	76.84
25	13.48	22.33	30.55	38.44	46.15	53.70	61.16	68.54	75.84
30	13.32	22.10	30.25	38.09	45.73	53.22	60.62	67.94	75.18

(continued)

Table A.14 (Continued)

ν	$k = 2$	$k = 3$	$k = 4$	$k = 5$	$k = 6$	$k = 7$	$k = 8$	$k = 9$	$k = 10$
$p = 4$									
5	35.39	56.10	75.36	93.97	112.17	130.11	147.81	165.39	182.80
6	30.06	48.62	65.90	82.60	98.93	115.03	130.94	146.69	162.34
7	27.31	44.69	60.89	76.56	91.88	106.98	121.90	136.71	151.39
8	25.61	42.24	57.77	72.77	87.46	101.94	116.23	130.43	144.50
9	24.45	40.57	55.62	70.17	84.42	98.46	112.32	126.08	139.74
10	23.62	39.34	54.04	68.26	82.19	95.90	109.46	122.91	136.24
11	22.98	38.41	52.84	66.81	80.48	93.95	107.27	120.46	133.57
12	22.48	37.67	51.90	65.66	79.14	92.41	105.54	118.55	131.45
13	22.08	37.08	51.13	64.73	78.04	91.15	104.12	116.98	129.74
14	21.75	36.59	50.50	63.95	77.13	90.12	102.97	115.69	128.32
15	21.47	36.17	49.97	63.30	76.37	89.26	101.99	114.59	127.14
16	21.24	35.82	49.51	62.76	75.73	88.51	101.14	113.67	126.10
17	21.03	35.52	49.12	62.28	75.16	87.87	100.42	112.87	125.22
18	20.86	35.26	48.78	61.86	74.68	87.31	99.80	112.17	124.46
19	20.70	35.02	48.47	61.50	74.25	86.82	99.25	111.56	123.79
20	20.56	34.82	48.21	61.17	73.87	86.38	98.75	111.02	123.18
25	20.06	34.06	47.23	59.98	72.47	84.78	96.95	109.01	120.99
30	19.74	33.59	46.61	59.21	71.58	83.74	95.79	107.71	119.57
$p = 5$									
6	51.11	81.99	110.92	138.98	166.54	193.71	220.66	247.37	273.88
7	43.40	71.06	97.03	122.22	146.95	171.34	195.49	219.47	243.30
8	39.29	65.15	89.45	113.03	136.18	159.04	181.65	204.14	226.48
9	36.71	61.39	84.62	107.17	129.30	151.17	172.80	194.27	215.64
10	34.93	58.78	81.25	103.06	124.48	145.64	166.56	187.37	208.02
11	33.62	56.85	78.75	100.02	120.92	141.54	161.98	182.24	202.37
12	32.62	55.37	76.83	97.68	118.15	138.38	158.38	178.23	198.03
13	31.83	54.19	75.30	95.82	115.96	135.86	155.54	175.10	194.51
14	31.19	53.23	74.05	94.29	114.16	133.80	153.21	172.49	191.68
15	30.66	52.44	73.01	93.02	112.66	132.07	151.29	170.36	189.38
16	30.22	51.76	72.14	91.94	111.41	130.61	149.66	166.53	187.32
17	29.83	51.19	71.39	91.03	110.34	129.38	148.25	166.99	185.61
18	29.51	50.69	70.74	90.23	109.39	128.29	147.03	165.65	184.10
19	29.22	50.26	70.17	89.54	108.57	127.36	145.97	164.45	182.81
20	28.97	49.88	69.67	88.93	107.85	126.52	145.02	163.38	181.65
25	28.05	48.48	67.86	86.70	105.21	123.51	141.62	159.60	177.49
30	27.48	47.61	66.71	85.29	103.56	121.60	139.47	157.22	174.87

Note: Table contains upper percentage points for

$$-2 \ln M = \nu \left(k \ln |\mathbf{S}| - \sum_{i=1}^k \ln |\mathbf{S}_i| \right)$$

for k samples, each with ν degrees of freedom. Reject $H_0 : \Sigma_1 = \Sigma_2 = \dots = \Sigma_k$ if $-2 \ln M >$ table value.

Table A.15 Test for Independence of p Variables

Upper percentage points for

$$u' = -\left(\nu - \frac{2p+5}{6}\right) \ln\left(\frac{|\mathbf{S}|}{s_{11} \cdots s_{pp}}\right) = -\left(\nu - \frac{2p+5}{6}\right) \ln|\mathbf{R}|,$$

where ν is the degrees of freedom of \mathbf{S} or \mathbf{R} . Reject independence if u' is greater than table value. The χ^2_α values are shown for comparison, since u' is approximately χ^2 distributed with $f = \frac{1}{2}p(p-1)$ degrees of freedom.

n	$p = 3$	$p = 4$	$p = 5$	$p = 6$	$p = 7$	$p = 8$	$p = 9$	$p = 10$
$\alpha = 0.05$								
4	8.020							
5	7.834	15.22						
6	7.814	13.47	24.01					
7	7.811	13.03	20.44	34.30				
8	7.811	12.85	19.45	28.75	46.05			
9	7.811	12.76	19.02	27.11	38.41	59.25		
10	7.812	12.71	18.80	26.37	36.03	49.42	73.79	
11	7.812	12.68	18.67	25.96	34.91	46.22	61.76	89.92
12	7.813	12.66	18.58	25.71	34.28	44.67	57.68	75.45
13	7.813	12.65	18.52	25.55	33.89	43.78	55.65	70.43
14	7.813	12.64	18.48	25.44	33.63	43.21	54.46	67.87
15	7.813	12.63	18.45	25.36	33.44	42.82	53.69	66.34
16	7.814	12.62	18.43	25.30	33.31	42.55	53.15	65.33
17	7.814	12.62	18.41	25.25	33.20	42.34	52.77	64.63
18	7.814	12.62	18.40	25.21	33.12	42.19	52.48	64.12
19	7.814	12.61	18.38	25.19	33.06	42.06	52.26	63.73
20	7.814	12.61	18.37	25.16	33.01	41.97	52.08	63.43
$\chi^2_{0.05}$	7.815	12.59	18.31	25.00	32.67	41.34	51.00	61.66
$\alpha = 0.01$								
4	11.79							
5	11.41	21.18						
6	11.36	18.27	32.16					
7	11.34	17.54	26.50	44.65				
8	11.34	17.24	24.95	36.09	58.61			
9	11.34	17.10	24.29	33.63	47.05	74.01		
10	11.34	17.01	23.95	32.54	43.59	59.36	90.87	
11	11.34	16.96	23.75	31.95	42.00	54.83	73.03	109.53
12	11.34	16.93	23.62	31.60	41.13	52.70	67.37	88.05
13	11.34	16.90	23.53	31.36	40.59	51.49	64.64	81.20
14	11.34	16.89	23.47	31.20	40.23	50.73	63.06	77.83
15	11.34	16.87	23.42	31.09	39.97	50.22	62.05	75.84
16	11.34	16.86	23.39	31.00	39.79	49.85	61.36	74.56
17	11.34	16.86	23.36	30.94	39.65	49.59	60.86	73.66
18	11.34	16.85	23.34	30.88	39.54	49.38	60.49	73.01
19	11.34	16.85	23.32	30.84	39.46	49.22	60.21	72.52
20	11.34	16.84	23.31	30.81	39.39	49.09	59.99	72.15
$\chi^2_{0.01}$	11.34	16.81	23.21	30.58	38.93	48.28	58.57	69.92

Appendix B

Answers and Hints to Problems

CHAPTER 2

2.1 (a) $\mathbf{A} + \mathbf{B} = \begin{pmatrix} 7 & 0 & 7 \\ 13 & 14 & 3 \end{pmatrix}$, $\mathbf{A} - \mathbf{B} = \begin{pmatrix} 1 & 4 & -1 \\ 1 & -4 & 13 \end{pmatrix}$

(b) $\mathbf{A}'\mathbf{A} = \begin{pmatrix} 65 & 43 & 68 \\ 43 & 29 & 46 \\ 68 & 46 & 73 \end{pmatrix}$, $\mathbf{A}\mathbf{A}' = \begin{pmatrix} 29 & 62 \\ 62 & 138 \end{pmatrix}$

2.2 (a) $(\mathbf{A} + \mathbf{B})' = \begin{pmatrix} 7 & 13 \\ 0 & 14 \\ 7 & 3 \end{pmatrix}$, $\mathbf{A}' + \mathbf{B}' = \begin{pmatrix} 7 & 13 \\ 0 & 14 \\ 7 & 3 \end{pmatrix}$

(b) $\mathbf{A}' = \begin{pmatrix} 4 & 7 \\ 2 & 5 \\ 3 & 8 \end{pmatrix}$, $(\mathbf{A}')' = \begin{pmatrix} 4 & 2 & 3 \\ 7 & 5 & 8 \end{pmatrix} = \mathbf{A}$

*

2.3 (a) $\mathbf{AB} = \begin{pmatrix} 5 & 15 \\ 3 & -5 \end{pmatrix}$, $\mathbf{BA} = \begin{pmatrix} 2 & 6 \\ 11 & -2 \end{pmatrix}$

(b) $|\mathbf{AB}| = -70$, $|\mathbf{A}| = -7$, $|\mathbf{B}| = 10$

2.4 (a) $\mathbf{A} + \mathbf{B} = \begin{pmatrix} 3 & 3 \\ 3 & 4 \end{pmatrix}$, $\text{tr}(\mathbf{A} + \mathbf{B}) = 7$

(b) $\text{tr}(\mathbf{A}) = 0$, $\text{tr}(\mathbf{B}) = 7$

2.5 (a) $\mathbf{AB} = \begin{pmatrix} 4 & 1 \\ 3 & -3 \end{pmatrix}$, $\mathbf{BA} = \begin{pmatrix} -1 & 8 & 7 \\ 2 & 4 & 6 \\ 1 & -3 & -2 \end{pmatrix}$

(b) $\text{tr}(\mathbf{AB}) = 1$, $\text{tr}(\mathbf{BA}) = 1$

2.6 (b) $\mathbf{x} = (1 \quad 1 \quad -1)'$

2.7 (a) $\mathbf{Bx} = (13, 6, 9)'$

(d) $\mathbf{x}'\mathbf{Ay} = 43$

(g) $\mathbf{xx}' = \begin{pmatrix} 1 & -1 & 2 \\ -1 & 1 & -2 \\ 2 & -2 & 4 \end{pmatrix}$

(i) $\mathbf{B}'\mathbf{B} = \begin{pmatrix} 62 & 7 & 22 \\ 7 & 14 & 7 \\ 22 & 7 & 41 \end{pmatrix}$

(b) $\mathbf{y}'\mathbf{B} = (25, -1, 17)$

(e) $\mathbf{x}'\mathbf{x} = 6$

(h) $\mathbf{xy}' = \begin{pmatrix} 3 & 2 & 1 \\ -3 & -2 & -1 \\ 6 & 4 & 2 \end{pmatrix}$

(c) $\mathbf{x}'\mathbf{Ax} = 10$

(f) $\mathbf{x}'\mathbf{y} = 3$

2.8 (a) $\mathbf{x} + \mathbf{y} = (4, 1, 3)'$, $\mathbf{x} - \mathbf{y} = (-2, -3, 1)'$

(b) $(\mathbf{x} - \mathbf{y})'\mathbf{A}(\mathbf{x} - \mathbf{y}) = -31$

2.9 $\mathbf{Bx} = \mathbf{b}_1x_1 + \mathbf{b}_2x_2 + \mathbf{b}_3x_3 = (1)\begin{pmatrix} 3 \\ 7 \\ 2 \end{pmatrix} + (-1)\begin{pmatrix} -2 \\ 1 \\ 3 \end{pmatrix} + (2)\begin{pmatrix} 4 \\ 0 \\ 5 \end{pmatrix} =$

$$\begin{pmatrix} 13 \\ 6 \\ 9 \end{pmatrix}$$

2.10 (a) $(\mathbf{AB})' = \begin{pmatrix} 7 & 16 \\ 8 & 4 \\ 7 & 11 \end{pmatrix}$, $\mathbf{B}'\mathbf{A}' = \begin{pmatrix} 7 & 16 \\ 8 & 4 \\ 7 & 11 \end{pmatrix}$ (c) $|\mathbf{A}| = 5$

2.11 (a) $\mathbf{a}'\mathbf{b} = 5$, $(\mathbf{a}'\mathbf{b})^2 = 25$ (b) $\mathbf{bb}' = \begin{pmatrix} 4 & 2 & 6 \\ 2 & 1 & 3 \\ 6 & 3 & 9 \end{pmatrix}$, $\mathbf{a}'(\mathbf{bb}')\mathbf{a} = 25$

2.12 $\mathbf{DA} = \begin{pmatrix} a & 2a & 3a \\ 4b & 5b & 6b \\ 7c & 8c & 9c \end{pmatrix}$, $\mathbf{AD} = \begin{pmatrix} a & 2b & 3c \\ 4a & 5b & 6c \\ 7a & 8b & 9c \end{pmatrix}$,

$$\mathbf{DAD} = \begin{pmatrix} a^2 & 2ab & 3ac \\ 4ab & 5b^2 & 6bc \\ 7ac & 8bc & 9c^2 \end{pmatrix}$$

$$2.13 \quad \mathbf{AB} = \left(\begin{array}{ccc|c} 8 & 9 & 5 & 6 \\ 7 & 5 & 5 & 4 \\ \hline 3 & 4 & 2 & 2 \end{array} \right)$$

$$2.14 \quad \mathbf{AB} = \begin{pmatrix} 3 & 5 \\ 1 & 4 \end{pmatrix}, \quad \mathbf{CB} = \begin{pmatrix} 3 & 5 \\ 1 & 4 \end{pmatrix}$$

$$2.15 \quad (\text{a}) \quad \text{tr}(\mathbf{A}) = 4, \quad \text{tr}(\mathbf{B}) = 5$$

$$(\text{b}) \quad \mathbf{A} + \mathbf{B} = \begin{pmatrix} 6 & 4 & 5 \\ 2 & -2 & 1 \\ 4 & 9 & 5 \end{pmatrix}, \quad \text{tr}(\mathbf{A} + \mathbf{B}) = 9$$

$$(\text{c}) \quad |\mathbf{A}| = 23, \quad |\mathbf{B}| = 2$$

$$(\text{d}) \quad \mathbf{AB} = \begin{pmatrix} 9 & 12 & 17 \\ 3 & -1 & 5 \\ 5 & 11 & 9 \end{pmatrix}, \quad \det(\mathbf{AB}) = 46$$

$$2.16 \quad (\text{a}) \quad |\mathbf{A}| = 36 \quad (\text{b}) \quad \mathbf{T} = \begin{pmatrix} 1.7321 & 2.3094 & 1.7321 \\ 0 & 1.6330 & 1.2247 \\ 0 & 0 & 2.1213 \end{pmatrix}$$

$$2.17 \quad (\text{a}) \quad \det(\mathbf{A}) = 1 \quad (\text{b}) \quad \mathbf{T} = \begin{pmatrix} 1.7321 & -2.8868 & -.5774 \\ 0 & 2.1602 & -.7715 \\ 0 & 0 & .2673 \end{pmatrix}$$

$$2.18 \quad (\text{a}) \quad \mathbf{C} = \begin{pmatrix} .4082 & -.5774 & .7071 \\ .8165 & .5774 & .0000 \\ .4082 & -.5774 & -.7071 \end{pmatrix}$$

$$2.19 \quad (\text{a}) \quad \text{Eigenvalues: } 2, 1, -1$$

$$\text{Eigenvectors: } \begin{pmatrix} .3015 \\ .9045 \\ .3015 \end{pmatrix}, \begin{pmatrix} .7999 \\ .5368 \\ .2684 \end{pmatrix}, \begin{pmatrix} .7071 \\ 0 \\ .7071 \end{pmatrix}$$

$$(\text{b}) \quad \text{tr}(\mathbf{A}) = 2, \quad |\mathbf{A}| = -2$$

$$2.20 \quad (\text{a}) \quad \mathbf{C} = \begin{pmatrix} .0000 & .5774 & -.8165 \\ -.7071 & -.5774 & -.4082 \\ .7071 & -.5774 & -.4082 \end{pmatrix}$$

$$(\text{b}) \quad \mathbf{C}' \mathbf{AC} = \begin{pmatrix} -2 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 4 \end{pmatrix}$$

$$(c) \mathbf{CDC}' = \begin{pmatrix} 3 & 1 & 1 \\ 1 & 0 & 2 \\ 1 & 2 & 0 \end{pmatrix} = \mathbf{A}$$

2.21 Eigenvalues: 1, 3, $\mathbf{C} = \begin{pmatrix} -.7071 & -.7071 \\ -.7071 & .7071 \end{pmatrix}$,

$$\mathbf{A}^{1/2} = \mathbf{CD}^{1/2}\mathbf{C}' = \begin{pmatrix} 1.3660 & -.3660 \\ -.3660 & 1.3660 \end{pmatrix}$$

2.22 (a) The spectral decomposition of \mathbf{A} is given by $\mathbf{A} = \mathbf{CDC}'$, where $\mathbf{C} = \begin{pmatrix} .455 & -.580 & .675 \\ .846 & .045 & -.531 \\ .278 & .813 & .511 \end{pmatrix}$ and $\mathbf{D} = \text{diag}(13.542, 3.935, -2.477)$.

(b) The spectral decomposition of \mathbf{A}^2 is given by $\mathbf{A}^2 = \mathbf{CDC}'$, where \mathbf{C} is the same as in part (a) and $\mathbf{D} = \text{diag}(183.378, 15.486, 6.135)$. Note that the diagonal elements of \mathbf{D} are the squares of the diagonal elements of \mathbf{D} in part (a).

(c) The spectral decomposition of \mathbf{A}^{-1} is given by $\mathbf{A}^{-1} = \mathbf{CDC}'$, where $\mathbf{C} = \begin{pmatrix} -.580 & .455 & .675 \\ .045 & .846 & -.531 \\ .813 & .278 & .511 \end{pmatrix}$ and $\mathbf{D} = \text{diag}(.254, .074, -.404)$.

The diagonal elements of \mathbf{D} are the reciprocals of those of \mathbf{D} in part (a). The first two columns of \mathbf{C} have been interchanged to match the interchange of the corresponding elements of \mathbf{D} ; that is, $\mathbf{D} = (1/\lambda_2, 1/\lambda_1, 1/\lambda_3)$.

2.23 $\mathbf{A} = \mathbf{UDV}'$, where $\mathbf{D} = \text{diag}(13.161, 7, 000, 3.433)$,

$$\mathbf{U} = \begin{pmatrix} .282 & -.730 & .424 \\ .591 & -.146 & .184 \\ -.225 & .404 & .886 \\ .721 & .531 & -.040 \end{pmatrix}, \quad \mathbf{V} = \begin{pmatrix} .856 & -.015 & .517 \\ -.156 & .946 & .284 \\ .494 & .324 & -.807 \end{pmatrix}.$$

2.24 (a) $\mathbf{j}'\mathbf{a} = (1)a_1 + (1)a_2 + \cdots + (1)a_n = \sum_i a_i = \mathbf{a}'\mathbf{j}$

(b) $\mathbf{j}'\mathbf{A} = [(1)a_{11} + (1)a_{21} + \cdots + (1)a_{n1}, \dots, (1)a_{1p} + (1)a_{2p} + \cdots + (1)a_{np}] = (\sum_i a_{i1}, \sum_i a_{i2}, \dots, \sum_i a_{ip})$

(c) $\mathbf{A}\mathbf{j} = \begin{pmatrix} (1)a_{11} & (1)a_{12} & \cdots & (1)a_{1p} \\ (1)a_{21} & (1)a_{22} & \cdots & (1)a_{2p} \\ \vdots & \vdots & & \vdots \\ (1)a_{n1} & (1)a_{n2} & \cdots & (1)a_{np} \end{pmatrix} = \begin{pmatrix} \sum_j a_{1j} \\ \sum_j a_{2j} \\ \vdots \\ \sum_j a_{nj} \end{pmatrix}$

2.25 $(\mathbf{x} - \mathbf{y})'(\mathbf{x} - \mathbf{y}) = (\mathbf{x}' - \mathbf{y}')(\mathbf{x} - \mathbf{y}) = \mathbf{x}'\mathbf{x} - \mathbf{x}'\mathbf{y} - \mathbf{y}'\mathbf{x} + \mathbf{y}'\mathbf{y}$
 $= \mathbf{x}'\mathbf{x} - 2\mathbf{x}'\mathbf{y} + \mathbf{y}'\mathbf{y}$

2.26 By (2.27), $(\mathbf{A}'\mathbf{A})' = \mathbf{A}'(\mathbf{A}')'$. By (2.6), $(\mathbf{A}')' = \mathbf{A}$. Thus, $(\mathbf{A}'\mathbf{A})' = \mathbf{A}'\mathbf{A}$.

2.27 (a) $\sum_i \mathbf{a}'\mathbf{x}_i = \mathbf{a}'\mathbf{x}_1 + \mathbf{a}'\mathbf{x}_2 + \cdots + \mathbf{a}'\mathbf{x}_n$
 $= \mathbf{a}'(\mathbf{x}_1 + \mathbf{x}_2 + \cdots + \mathbf{x}_n)$ [by (2.21)]
 $= \mathbf{a}' \sum_i \mathbf{x}_i$

(b) $\sum_i \mathbf{A}\mathbf{x}_i = \mathbf{A}\mathbf{x}_1 + \mathbf{A}\mathbf{x}_2 + \cdots + \mathbf{A}\mathbf{x}_n$
 $= \mathbf{A}(\mathbf{x}_1 + \mathbf{x}_2 + \cdots + \mathbf{x}_n)$ [by (2.21)]
 $= \mathbf{A} \sum_i \mathbf{x}_i$

(c) $\sum_i (\mathbf{a}'\mathbf{x}_i)^2 = \sum_i \mathbf{a}'(\mathbf{x}_i\mathbf{x}'_i)\mathbf{a}$ [by (2.40)]
 $= \mathbf{a}'(\sum_i \mathbf{x}_i\mathbf{x}'_i)\mathbf{a}$ [by (2.29)]

(d) $\sum_i \mathbf{A}\mathbf{x}_i(\mathbf{A}\mathbf{x}_i)' = \sum_i \mathbf{A}\mathbf{x}_i\mathbf{x}'_i\mathbf{A}' = \mathbf{A}(\sum_i \mathbf{x}_i\mathbf{x}'_i)\mathbf{A}'$ [by (2.29)]

2.28 (a) $\mathbf{Ax} = \begin{pmatrix} \mathbf{a}'_1 \\ \mathbf{a}'_2 \end{pmatrix}\mathbf{x} = \begin{pmatrix} \mathbf{a}'_1\mathbf{x} \\ \mathbf{a}'_2\mathbf{x} \end{pmatrix}$

(b) $\mathbf{ASA}' = \begin{pmatrix} \mathbf{a}'_1 \\ \mathbf{a}'_2 \end{pmatrix}\mathbf{S}(\mathbf{a}_1, \mathbf{a}_2) = \begin{pmatrix} \mathbf{a}'_1 \\ \mathbf{a}'_2 \end{pmatrix}(\mathbf{Sa}_1, \mathbf{Sa}_2)$ [by (2.48)]
 $= \begin{pmatrix} \mathbf{a}'_1\mathbf{Sa}_1 & \mathbf{a}'_1\mathbf{Sa}_2 \\ \mathbf{a}'_2\mathbf{Sa}_1 & \mathbf{a}'_2\mathbf{Sa}_2 \end{pmatrix}$

2.29 (a) If $\mathbf{A} = \begin{pmatrix} \mathbf{a}'_1 \\ \mathbf{a}'_2 \\ \vdots \\ \mathbf{a}'_n \end{pmatrix}$, then by (2.68), $\mathbf{A}' = (\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n)$ and

$$\mathbf{A}'\mathbf{A} = (\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_n) \begin{pmatrix} \mathbf{a}'_1 \\ \mathbf{a}'_2 \\ \vdots \\ \mathbf{a}'_n \end{pmatrix} = \mathbf{a}_1\mathbf{a}'_1 + \mathbf{a}_2\mathbf{a}'_2 + \cdots + \mathbf{a}_n\mathbf{a}'_n \quad [\text{by (2.66)}].$$

2.30 $\mathbf{A}^{-1}\mathbf{A} = \mathbf{I}$

$$(\mathbf{A}^{-1}\mathbf{A})' = \mathbf{I}' = \mathbf{I}$$

$$\mathbf{A}'(\mathbf{A}^{-1})' = \mathbf{I}$$

$$(\mathbf{A}')^{-1}\mathbf{A}'(\mathbf{A}^{-1})' = (\mathbf{A}')^{-1}\mathbf{I} = (\mathbf{A}')^{-1}$$

$$(\mathbf{A}^{-1})' = (\mathbf{A}')^{-1}$$

2.31
$$\begin{aligned} & \frac{1}{b} \begin{pmatrix} b\mathbf{A}_{11}^{-1} + \mathbf{A}_{11}^{-1}\mathbf{a}_{12}\mathbf{a}'_{12}\mathbf{A}_{11}^{-1} & -\mathbf{A}_{11}^{-1}\mathbf{a}_{12} \\ -\mathbf{a}'_{12}\mathbf{A}_{11}^{-1} & 1 \end{pmatrix} \begin{pmatrix} \mathbf{A}_{11} & \mathbf{a}_{12} \\ \mathbf{a}'_{12} & a_{22} \end{pmatrix} \\ &= \frac{1}{b} \begin{pmatrix} b\mathbf{I} + \mathbf{A}_{11}^{-1}\mathbf{a}_{12}\mathbf{a}'_{12} - \mathbf{A}_{11}^{-1}\mathbf{a}_{12}\mathbf{a}'_{12} & b\mathbf{A}_{11}^{-1}\mathbf{a}_{12} + \mathbf{A}_{11}^{-1}\mathbf{a}_{12}\mathbf{a}'_{12}\mathbf{A}_{11}^{-1}\mathbf{a}_{12} - \mathbf{A}_{11}^{-1}\mathbf{a}_{12}a_{22} \\ -\mathbf{a}'_{12} + \mathbf{a}'_{12} & -\mathbf{a}'_{12}\mathbf{A}_{11}^{-1}\mathbf{a}_{12} + a_{22} \end{pmatrix} \\ &= \frac{1}{b} \begin{pmatrix} b\mathbf{I}, & \mathbf{0} \\ \mathbf{0}', & b \end{pmatrix}, \text{ where } b = a_{22} - \mathbf{a}'_{12}\mathbf{A}_{11}^{-1}\mathbf{a}_{12} \\ &= \begin{pmatrix} \mathbf{I} & \mathbf{0} \\ \mathbf{0}' & 1 \end{pmatrix} \end{aligned}$$

2.32
$$\begin{aligned} & (\mathbf{B} + \mathbf{cc}') \left(\mathbf{B}^{-1} - \frac{\mathbf{B}^{-1}\mathbf{cc}'\mathbf{B}^{-1}}{1 + \mathbf{c}'\mathbf{B}^{-1}\mathbf{c}} \right) \\ &= \mathbf{I} - \frac{\mathbf{cc}'\mathbf{B}^{-1}}{1 + \mathbf{c}'\mathbf{B}^{-1}\mathbf{c}} + \mathbf{cc}'\mathbf{B}^{-1} - \frac{\mathbf{c}(\mathbf{c}'\mathbf{B}^{-1}\mathbf{c})\mathbf{c}'\mathbf{B}^{-1}}{1 + \mathbf{c}'\mathbf{B}^{-1}\mathbf{c}} \quad [\text{by (2.26)}] \\ &= \mathbf{I} - \mathbf{cc}'\mathbf{B}^{-1} \left(\frac{1 + \mathbf{c}'\mathbf{B}^{-1}\mathbf{c}}{1 + \mathbf{c}'\mathbf{B}^{-1}\mathbf{c}} \right) + \mathbf{cc}'\mathbf{B}^{-1} = \mathbf{I} \end{aligned}$$

2.33
$$\begin{aligned} |c\mathbf{A}| &= |c\mathbf{I}\mathbf{A}| \\ &= |c\mathbf{I}| |\mathbf{A}| \quad [\text{by (2.89)}] \\ &= c^n |\mathbf{A}| \quad [\text{by (2.84)}] \end{aligned}$$

2.34
$$\begin{aligned} \mathbf{A}\mathbf{A}^{-1} &= \mathbf{I} \\ |\mathbf{A}\mathbf{A}^{-1}| &= |\mathbf{I}| \\ |\mathbf{A}||\mathbf{A}^{-1}| &= 1 \quad [\text{by (2.83)}] \\ |\mathbf{A}^{-1}| &= \frac{1}{|\mathbf{A}|} \end{aligned}$$

2.35 In (2.93) and (2.94), let $\mathbf{A}_{11} = \mathbf{B}$, $\mathbf{A}_{12} = \mathbf{c}$, $\mathbf{A}_{21} = -\mathbf{c}'$, and $\mathbf{A}_{22} = 1$. Then equate the right-hand sides of (2.93) and (2.94) to obtain (2.95).

2.36 By (2.52), $\text{tr}(\mathbf{AA}') = \sum_{i=1}^n \mathbf{a}'_i \mathbf{a}_i = \sum_{i=1}^n (a_{i1}^2 + a_{i2}^2 + \dots + a_{in}^2)$
 $= \sum_{i=1}^n \sum_{j=1}^n a_{ij}^2$.

2.37 Show that $|\mathbf{C}| \neq 0$ by taking the determinant of both sides of $\mathbf{C}'\mathbf{C} = \mathbf{I}$. Thus \mathbf{C} is nonsingular and \mathbf{C}^{-1} exists. Multiply $\mathbf{C}'\mathbf{C} = \mathbf{I}$ on the right by \mathbf{C}^{-1} and on the left by \mathbf{C} .

2.38 Multiply $\mathbf{ABx} = \lambda\mathbf{x}$ on the left by \mathbf{B} . Then λ is an eigenvalue of \mathbf{BA} and \mathbf{Bx} is an eigenvector.

2.39 (a)
$$\begin{aligned} (\mathbf{A}^{1/2})^2 &= (\mathbf{CD}^{1/2}\mathbf{C}')^2 = \mathbf{CD}^{1/2}\mathbf{C}'\mathbf{CD}^{1/2}\mathbf{C}' \\ &= \mathbf{CDC}' \quad [\text{by (2.101)}] \\ &= \mathbf{A} \quad [\text{by (2.109)}] \end{aligned}$$

(b) By (2.114), $\mathbf{A}^{1/2}\mathbf{A}^{1/2} = \mathbf{A}$. By (2.89),

$$\begin{aligned} |\mathbf{A}^{1/2}\mathbf{A}^{1/2}| &= |\mathbf{A}| \\ |\mathbf{A}^{1/2}||\mathbf{A}^{1/2}| &= |\mathbf{A}| \\ |\mathbf{A}^{1/2}|^2 &= |\mathbf{A}| \end{aligned}$$

(c) Since \mathbf{A} is positive definite, we have, from part (b), $|\mathbf{A}^{1/2}| = |\mathbf{A}|^{1/2}$.

- 2.40** Use properties (2.123) and (2.124). Then use property (2.121) (twice). Finally, use property (2.124).

CHAPTER 3

3.1 $\bar{z} = \sum_{i=1}^n z_i/n = \sum_i a y_i/n = (ay_1 + \dots + ay_n)/n$. Now factor a out of the sum.

3.2 The numerator of s_z^2 is $\sum_{i=1}^n (z_i - \bar{z})^2 = \sum_i (ay_i - a\bar{y})^2 = \sum_i [a(y_i - \bar{y})]^2$.

3.3 $\bar{x} = 4, \bar{y} = 4$:

x	y	$x - \bar{x}$	$y - \bar{y}$	$(x - \bar{x})(y - \bar{y})$	
2	2	-2	-2	4	
2	4	-2	0	0	
2	6	-2	2	-4	
4	2	0	-2	0	
4	4	0	0	0	
4	6	0	2	0	
6	2	2	-2	-4	
6	4	2	0	0	
6	6	2	2	4	
Sum = 0					

$$\mathbf{3.4} \quad \mathbf{x} - \bar{x}\mathbf{j} = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} - \bar{x} \begin{pmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{pmatrix} = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} - \begin{pmatrix} \bar{x} \\ \bar{x} \\ \vdots \\ \bar{x} \end{pmatrix} = \begin{pmatrix} x_1 - \bar{x} \\ x_2 - \bar{x} \\ \vdots \\ x_n - \bar{x} \end{pmatrix}$$

$$\mathbf{3.5} \quad \mathbf{y}_i - \bar{\mathbf{y}} = \begin{pmatrix} y_{i1} \\ y_{i2} \\ y_{i3} \end{pmatrix} - \begin{pmatrix} \bar{y}_1 \\ \bar{y}_2 \\ \bar{y}_3 \end{pmatrix} = \begin{pmatrix} y_{i1} - \bar{y}_1 \\ y_{i2} - \bar{y}_2 \\ y_{i3} - \bar{y}_3 \end{pmatrix}$$

$$\begin{aligned} \sum_{i=1}^n (\mathbf{y}_i - \bar{\mathbf{y}})(\mathbf{y}_i - \bar{\mathbf{y}})' &= \sum_{i=1}^n \begin{pmatrix} y_{i1} - \bar{y}_1 \\ y_{i2} - \bar{y}_2 \\ y_{i3} - \bar{y}_3 \end{pmatrix} (y_{i1} - \bar{y}_1, y_{i2} - \bar{y}_2, y_{i3} - \bar{y}_3) \\ &= \sum_{i=1}^n \begin{pmatrix} (y_{i1} - \bar{y}_1)^2 & (y_{i1} - \bar{y}_1)(y_{i2} - \bar{y}_2) & (y_{i1} - \bar{y}_1)(y_{i3} - \bar{y}_3) \\ (y_{i2} - \bar{y}_2)(y_{i1} - \bar{y}_1) & (y_{i2} - \bar{y}_2)^2 & (y_{i2} - \bar{y}_2)(y_{i3} - \bar{y}_3) \\ (y_{i3} - \bar{y}_3)(y_{i1} - \bar{y}_1) & (y_{i3} - \bar{y}_3)(y_{i2} - \bar{y}_2) & (y_{i3} - \bar{y}_3)^2 \end{pmatrix} \end{aligned}$$

3.6 $\bar{z} = \sum_{i=1}^n z_i/n = \sum_i \mathbf{a}' \mathbf{y}_i/n = (\mathbf{a}' \mathbf{y}_1 + \dots + \mathbf{a}' \mathbf{y}_n)/n$. Now factor out \mathbf{a}' on the left. See also (2.42).

3.7 The numerator of s_z^2 is $\sum_{i=1}^n (z_i - \bar{z})^2 = \sum_i (\mathbf{a}'\mathbf{y}_i - \mathbf{a}'\bar{\mathbf{y}})^2 = \sum_i (\mathbf{a}'\mathbf{y}_i - \mathbf{a}'\bar{\mathbf{y}})(\mathbf{a}'\mathbf{y}_i - \mathbf{a}'\bar{\mathbf{y}})$. The scalar $\mathbf{a}'\mathbf{y}_i$ is equal to its transpose, as in (2.39). Thus $\mathbf{a}'\mathbf{y}_i = (\mathbf{a}'\mathbf{y}_i)' = \mathbf{y}_i'\mathbf{a}$, and $\sum_i (\mathbf{a}'\mathbf{y}_i - \mathbf{a}'\bar{\mathbf{y}})(\mathbf{a}'\mathbf{y}_i - \mathbf{a}'\bar{\mathbf{y}}) = \sum_i (\mathbf{a}'\mathbf{y}_i - \mathbf{a}'\bar{\mathbf{y}})(\mathbf{y}_i'\mathbf{a} - \bar{\mathbf{y}}'\mathbf{a})$. By (2.22) and (2.24), this becomes $\sum_i \mathbf{a}'(\mathbf{y}_i - \bar{\mathbf{y}})(\mathbf{y}_i - \bar{\mathbf{y}})'\mathbf{a}$. Now factor out \mathbf{a}' on the left and \mathbf{a} on the right. See also (2.44).

3.8 By (3.63) and (3.64),

$$\mathbf{A}\mathbf{S}\mathbf{A}' = \begin{pmatrix} \mathbf{a}_1'\mathbf{S}\mathbf{a}_1 & \mathbf{a}_1'\mathbf{S}\mathbf{a}_2 & \cdots & \mathbf{a}_1'\mathbf{S}\mathbf{a}_k \\ \mathbf{a}_2'\mathbf{S}\mathbf{a}_1 & \mathbf{a}_2'\mathbf{S}\mathbf{a}_2 & \cdots & \mathbf{a}_2'\mathbf{S}\mathbf{a}_k \\ \vdots & \vdots & & \vdots \\ \mathbf{a}_k'\mathbf{S}\mathbf{a}_1 & \mathbf{a}_k'\mathbf{S}\mathbf{a}_2 & \cdots & \mathbf{a}_k'\mathbf{S}\mathbf{a}_k \end{pmatrix},$$

from which the result follows immediately.

$$\begin{aligned} \mathbf{3.9} \quad \text{cov}(\mathbf{z}) &= \text{cov}[(\boldsymbol{\Sigma}^{1/2})^{-1}\bar{\mathbf{y}} - (\boldsymbol{\Sigma}^{1/2})^{-1}\boldsymbol{\mu}] \\ &= (\boldsymbol{\Sigma}^{1/2})^{-1}\text{cov}(\bar{\mathbf{y}})[(\boldsymbol{\Sigma}^{1/2})^{-1}]' \quad [\text{by (3.76)}] \\ &= (\boldsymbol{\Sigma}^{1/2})^{-1} \left(\frac{\boldsymbol{\Sigma}}{n} \right) (\boldsymbol{\Sigma}^{1/2})^{-1} \\ &= \frac{1}{n} (\boldsymbol{\Sigma}^{1/2})^{-1} \boldsymbol{\Sigma}^{1/2} \boldsymbol{\Sigma}^{1/2} (\boldsymbol{\Sigma}^{1/2})^{-1} \quad [\text{by (2.114)}] \\ &= \frac{1}{n} \mathbf{I} \end{aligned}$$

3.10 Answers are given in Examples 3.7 and 3.8.

3.11 (a) $|\mathbf{S}| = 459.956$ **(b)** $\text{tr}(\mathbf{S}) = 213.043$

3.12 (a) $|\mathbf{S}| = 27,236,586$ **(b)** $\text{tr}(\mathbf{S}) = 292.891$

$$\mathbf{3.13} \quad \mathbf{R} = \begin{pmatrix} 1.000 & .614 & .757 & .575 & .413 \\ .614 & 1.000 & .547 & .750 & .548 \\ .757 & .547 & 1.000 & .605 & .692 \\ .575 & .750 & .605 & 1.000 & .524 \\ .413 & .548 & .692 & .524 & 1.000 \end{pmatrix}$$

3.14 $\bar{z} = 83.298$, $s_z^2 = 1048.659$

3.15 $r_{zw} = -.6106$

3.16 $y_1 = (1, 0, 0)\mathbf{y} = \mathbf{a}'\mathbf{y}$, $\frac{1}{2}(y_2 + y_3) = (0, \frac{1}{2}, \frac{1}{2})\mathbf{y} = \mathbf{b}'\mathbf{y}$. Use (3.57) to obtain $r_{zw} = .4873$.

$$\mathbf{3.17 (a)} \quad \bar{\mathbf{z}} = \begin{pmatrix} 38.369 \\ 40.838 \\ -51.727 \end{pmatrix}, \quad \mathbf{S}_z = \begin{pmatrix} 323.64 & 19.25 & -460.98 \\ 19.25 & 588.67 & 104.07 \\ -460.98 & 104.07 & 686.27 \end{pmatrix}$$

$$\mathbf{(b)} \quad \mathbf{R}_z = \begin{pmatrix} 1.0000 & .0441 & -.9781 \\ .0441 & 1.0000 & .1637 \\ -.9781 & .1637 & 1.0000 \end{pmatrix}$$

3.18 (a) $\bar{\mathbf{y}} = \begin{pmatrix} 48.655 \\ 49.625 \\ 50.570 \\ 51.445 \end{pmatrix}$, $\mathbf{S} = \begin{pmatrix} 6.3300 & 6.1891 & 5.7770 & 5.5348 \\ 6.1891 & 6.4493 & 6.1534 & 5.9057 \\ 5.7770 & 6.1534 & 6.9180 & 6.9267 \\ 5.5348 & 5.9057 & 6.9267 & 7.4331 \end{pmatrix}$,

$$\mathbf{R} = \begin{pmatrix} 1.0000 & .9687 & .8730 & .8069 \\ .9687 & 1.0000 & .9212 & .8530 \\ .8730 & .9212 & 1.0000 & .9659 \\ .8069 & .8530 & .9659 & 1.0000 \end{pmatrix}$$

(b) $|\mathbf{S}| = 1.0865$, $\text{tr}(\mathbf{S}) = 27.1304$

3.19 (a) $\bar{z} = 44.1400$, $s_z^2 = 21.2309$, $\bar{w} = 103.8850$, $s_w^2 = 30.8161$

(b) $s_{zw} = 6.5359$, $r_{zw} = .2555$

3.20 $\bar{\mathbf{z}} = \begin{pmatrix} 401.40 \\ -47.55 \\ 150.48 \end{pmatrix}$, $\mathbf{S}_z = \begin{pmatrix} 398.33 & -44.35 & 148.35 \\ -44.35 & 12.36 & -16.90 \\ 148.35 & -16.90 & 59.46 \end{pmatrix}$,

$$\mathbf{R}_z = \begin{pmatrix} 1.00 & -.63 & .96 \\ -.63 & 1.00 & -.62 \\ .96 & -.62 & 1.00 \end{pmatrix}$$

3.21 (a) $\begin{pmatrix} \bar{\mathbf{y}} \\ \bar{\mathbf{x}} \end{pmatrix} = \begin{pmatrix} 185.72 \\ 151.12 \\ \hline 183.84 \\ 149.24 \end{pmatrix}$

(b) $\mathbf{S} = \left(\begin{array}{cc|cc} 95.29 & 52.87 & 69.66 & 46.11 \\ 52.87 & 54.36 & 51.31 & 35.05 \\ \hline 69.6 & 51.31 & 100.81 & 56.54 \\ 46.11 & 35.05 & 56.54 & 45.02 \end{array} \right)$

3.22 $\begin{pmatrix} \bar{\mathbf{y}} \\ \bar{\mathbf{x}} \end{pmatrix} = \begin{pmatrix} 70.08 \\ 73.54 \\ 75.10 \\ \hline 109.68 \\ 104.24 \\ 109.98 \end{pmatrix}$,

$$\mathbf{S} = \left(\begin{array}{ccc|ccc} 95.54 & 17.61 & 12.18 & 60.52 & 23.00 & 62.84 \\ 17.61 & 73.19 & 14.25 & 5.73 & 61.28 & -1.66 \\ 12.18 & 14.25 & 76.17 & 46.75 & 32.77 & 69.84 \\ \hline 60.52 & 5.73 & 46.75 & 808.63 & 320.59 & 227.36 \\ 23.00 & 61.28 & 32.77 & 320.59 & 505.86 & 167.35 \\ 62.84 & -1.66 & 69.84 & 227.36 & 167.35 & 508.71 \end{array} \right)$$

CHAPTER 4

4.1 $|\Sigma_1| = 1$, $\text{tr}(\Sigma_1) = 20$, $|\Sigma_2| = 4$, $\text{tr}(\Sigma_2) = 15$. Thus $\text{tr}(\Sigma_1) > \text{tr}(\Sigma_2)$, but $|\Sigma_1| < |\Sigma_2|$. When converted to correlations, we have

$$\mathbf{P}_{\rho_1} = \begin{pmatrix} 1 & .96 & .80 \\ .96 & 1 & .89 \\ .80 & .89 & 1 \end{pmatrix}, \quad \mathbf{P}_{\rho_2} = \begin{pmatrix} 1 & .87 & .41 \\ .87 & 1 & .71 \\ .41 & .71 & 1 \end{pmatrix}.$$

As noted at the end of Section 4.1.3, a decrease in intercorrelations or an increase in the variances will lead to a larger $|\Sigma|$. In this case, the decrease in correlations from Σ_1 to Σ_2 outweighed the increase in the variances (the increase in trace).

$$\begin{aligned} \mathbf{4.2} \quad E(\mathbf{z}) &= (\mathbf{T}')^{-1}[E(\mathbf{y}) - \boldsymbol{\mu}] \quad [\text{by (3.75)}] \\ &= (\mathbf{T}')^{-1}[\boldsymbol{\mu} - \boldsymbol{\mu}] = \mathbf{0}, \\ \text{cov}(\mathbf{z}) &= (\mathbf{T}')^{-1}\boldsymbol{\Sigma}[(\mathbf{T}')^{-1}]' \quad [\text{by (3.76)}] \\ &= (\mathbf{T}')^{-1}\mathbf{T}'\mathbf{T}\mathbf{T}'^{-1} \quad [\text{by (2.75) and (2.79)}] \\ &= \mathbf{I} \end{aligned}$$

4.3 By the last expression in Section 2.3.1,

$$\prod_{i=1}^n \frac{1}{(\sqrt{2\pi})^p |\boldsymbol{\Sigma}|^{1/2}} = \frac{1}{(\sqrt{2\pi})^{np} |\boldsymbol{\Sigma}|^{n/2}}.$$

The sum in the exponent of (4.13) follows from the basic algebra of exponents.

4.4 Since $(\mathbf{y} - \boldsymbol{\mu})'\boldsymbol{\Sigma}^{-1}(\mathbf{y} - \boldsymbol{\mu})$ is a scalar, we have $E[(\mathbf{y} - \boldsymbol{\mu})'\boldsymbol{\Sigma}^{-1}(\mathbf{y} - \boldsymbol{\mu})] = E\{\text{tr}[(\mathbf{y} - \boldsymbol{\mu})'\boldsymbol{\Sigma}^{-1}(\mathbf{y} - \boldsymbol{\mu})]\} = E\{\text{tr}[\boldsymbol{\Sigma}^{-1}(\mathbf{y} - \boldsymbol{\mu})(\mathbf{y} - \boldsymbol{\mu})']\} = \text{tr}[\boldsymbol{\Sigma}^{-1}E(\mathbf{y} - \boldsymbol{\mu})(\mathbf{y} - \boldsymbol{\mu})'] = \text{tr}(\boldsymbol{\Sigma}^{-1}\boldsymbol{\Sigma}) = \text{tr}(\mathbf{I}_p) = p$.

4.5 The other two terms are of the form $\frac{1}{2} \sum_{i=1}^n (\bar{\mathbf{y}} - \boldsymbol{\mu})'\boldsymbol{\Sigma}^{-1}(\mathbf{y}_i - \bar{\mathbf{y}})$, which is equal to $\frac{1}{2}[(\bar{\mathbf{y}} - \boldsymbol{\mu})'\boldsymbol{\Sigma}^{-1}] \sum_{i=1}^n (\bar{\mathbf{y}}_i - \bar{\mathbf{y}})$. This vanishes because $\sum_{i=1}^n (\mathbf{y}_i - \bar{\mathbf{y}}) = n\bar{\mathbf{y}} - n\bar{\mathbf{y}} = \mathbf{0}$.

- 4.6** We replace y_i in $\sqrt{b_1}$ by $z_i = ay_i + b$. By an extension of (3.3), $\bar{z} = a\bar{y} + b$. Then (4.18) becomes

$$\begin{aligned}\frac{\sqrt{n} \sum_{i=1}^n (z_i - \bar{z})^3}{[\sum_{i=1}^n (z_i - \bar{z})^2]^{3/2}} &= \frac{\sqrt{n} \sum_i (ay_i + b - a\bar{y} - b)^3}{[\sum_i (ay_i + b - a\bar{y} - b)^2]^{3/2}} \\ &= \frac{\sqrt{n} a^3 \sum_i (y_i - \bar{y})^3}{[a^2 \sum_i (y_i - \bar{y})^2]^{3/2}} = \frac{\sqrt{n} \sum_i (y_i - \bar{y})^3}{[\sum_i (y_i - \bar{y})^2]^{3/2}} = \sqrt{b_1}.\end{aligned}$$

Similarly, if (4.19) is expressed in terms of $z_i = ay_i + b$, it reduces to b_2 in terms of y_i .

- 4.7** $\beta_{2,p} = E[(\mathbf{y} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1} (\mathbf{y} - \boldsymbol{\mu})]^2$ by (4.33). But when \mathbf{y} is $N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma})$, $v = (\mathbf{y} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1} (\mathbf{y} - \boldsymbol{\mu})$ is distributed as $\chi^2(p)$ by property 3 in Section 4.2. Then $E(v^2) = \text{var}(v) + [E(v)]^2$.

- 4.8** To show that $b_{1,p}$ and $b_{2,p}$ are invariant under the transformation $\mathbf{z} = \mathbf{A}\mathbf{y}_i + \mathbf{b}$, where \mathbf{A} is nonsingular, it is sufficient to show that $g_{ij}(\mathbf{z}) = (\mathbf{y}_i - \bar{\mathbf{y}})' \hat{\boldsymbol{\Sigma}}^{-1} (\mathbf{y}_j - \bar{\mathbf{y}})$. By (3.67) and (3.68), $\bar{\mathbf{z}} = \mathbf{A}\bar{\mathbf{y}} + \mathbf{b}$ and $\hat{\boldsymbol{\Sigma}}_{\mathbf{z}} = \mathbf{A}\hat{\boldsymbol{\Sigma}}\mathbf{A}'$. Then g_{ij} for \mathbf{z} becomes

$$\begin{aligned}g_{ij}(\mathbf{z}) &= (\mathbf{z}_i - \bar{\mathbf{z}})' \hat{\boldsymbol{\Sigma}}_{\mathbf{z}}^{-1} (\mathbf{z}_j - \bar{\mathbf{z}}) \\ &= (\mathbf{A}\mathbf{y}_i + \mathbf{b} - \mathbf{A}\bar{\mathbf{y}} - \mathbf{b})' (\mathbf{A}\hat{\boldsymbol{\Sigma}}\mathbf{A}')^{-1} (\mathbf{A}\mathbf{y}_j + \mathbf{b} - \mathbf{A}\bar{\mathbf{y}} - \mathbf{b}) \\ &= (\mathbf{y}_i - \bar{\mathbf{y}})' \mathbf{A}' (\mathbf{A}')^{-1} \hat{\boldsymbol{\Sigma}}^{-1} \mathbf{A}^{-1} \mathbf{A} (\mathbf{y}_j - \bar{\mathbf{y}}) \\ &= (\mathbf{y}_i - \bar{\mathbf{y}})' \hat{\boldsymbol{\Sigma}}^{-1} (\mathbf{y}_j - \bar{\mathbf{y}}) = g_{ij}(\mathbf{y})\end{aligned}$$

- 4.9** Let $i = (n)$ in (4.47); then solve for $D_{(n)}^2$ in (4.46) and substitute into (4.47) to obtain $F_{(n)}$ in terms of w , as in (4.48).

- 4.10** (a) $\mathbf{a}' = (2, -1, 3)$, $z = \mathbf{a}'\mathbf{y}$ is $N(17, 21)$

(b) $\mathbf{A} = \begin{pmatrix} 1 & 1 & 1 \\ 1 & -1 & 2 \end{pmatrix}$, $\mathbf{z} = \mathbf{A}\mathbf{y}$ is $N_2\left[\begin{pmatrix} 8 \\ 10 \end{pmatrix}, \begin{pmatrix} 29 & -1 \\ -1 & 9 \end{pmatrix}\right]$

- (c) By property 4b in Section 4.2, y_2 is $N(1, 13)$.

(d) By property 4a in Section 4.2, $\begin{pmatrix} y_1 \\ y_3 \end{pmatrix}$ is $N_2\left[\begin{pmatrix} 3 \\ 4 \end{pmatrix}, \begin{pmatrix} 6 & -2 \\ -2 & 4 \end{pmatrix}\right]$.

(e) $\mathbf{A} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ \frac{1}{2} & \frac{1}{2} & 0 \end{pmatrix}$, $\mathbf{A}\mathbf{y}$ is $N_3\left[\begin{pmatrix} 3 \\ 4 \\ 2 \end{pmatrix}, \begin{pmatrix} 6 & -2 & 3.5 \\ -2 & 4 & 1 \\ 3.5 & 1 & 5.25 \end{pmatrix}\right]$

- 4.11** (a) $\mathbf{z} = \begin{pmatrix} .408 & 0 & 0 \\ -.047 & .279 & 0 \\ .285 & -.247 & .731 \end{pmatrix} \begin{pmatrix} y - 3 \\ y - 1 \\ y - 4 \end{pmatrix}$

(b) $\mathbf{z} = \begin{pmatrix} .465 & -.070 & .170 \\ -.070 & .326 & -.166 \\ .170 & -.166 & .692 \end{pmatrix} \begin{pmatrix} y - 3 \\ y - 1 \\ y - 4 \end{pmatrix}$

(c) By (4.6), $(\mathbf{y} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1} (\mathbf{y} - \boldsymbol{\mu})$ is distributed as χ_3^2 .

4.12 (a) $\mathbf{a}' = (4, -2, 1, -3)$, $z = \mathbf{a}' \mathbf{y}$ is $N(-30, 153)$

$$\text{(b)} \quad \mathbf{A} = \begin{pmatrix} 1 & 1 & 1 & 1 \\ -2 & 3 & 1 & -2 \end{pmatrix}, \quad \mathbf{z} = \mathbf{A} \mathbf{y} \text{ is } N_2 \left[\begin{pmatrix} 5 \\ 2 \end{pmatrix}, \begin{pmatrix} 27 & -79 \\ -79 & 361 \end{pmatrix} \right]$$

$$\text{(c)} \quad \mathbf{A} = \begin{pmatrix} 3 & 1 & -4 & -1 \\ -1 & -3 & 1 & -2 \\ 2 & 2 & 4 & -5 \end{pmatrix},$$

$$\mathbf{z} = \mathbf{A} \mathbf{y} \text{ is } N_3 \left[\begin{pmatrix} -4 \\ -18 \\ -27 \end{pmatrix}, \begin{pmatrix} 35 & -18 & -6 \\ -18 & 46 & 14 \\ -6 & 14 & 93 \end{pmatrix} \right]$$

(d) By property 4b in Section 4.2, y_3 is $N(-1, 2)$.

$$\text{(e)} \quad \text{By property 4a in Section 4.2, } \begin{pmatrix} y_2 \\ y_4 \end{pmatrix} \text{ is } N_2 \left[\begin{pmatrix} 3 \\ 5 \end{pmatrix}, \begin{pmatrix} 9 & -6 \\ -6 & 9 \end{pmatrix} \right].$$

$$\text{(f)} \quad \mathbf{A} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ \frac{1}{2} & \frac{1}{2} & 0 & 0 \\ \frac{1}{3} & \frac{1}{3} & \frac{1}{3} & 0 \\ \frac{1}{4} & \frac{1}{4} & \frac{1}{4} & \frac{1}{4} \end{pmatrix},$$

$$\mathbf{A} \mathbf{y} \text{ is } N_4 \left[\begin{pmatrix} -2 \\ 5 \\ 0 \\ 1.25 \end{pmatrix}, \begin{pmatrix} 11 & 1.5 & 2 & 3.75 \\ 1.5 & 1 & .67 & .875 \\ 2 & .67 & .67 & 1 \\ 3.75 & .875 & 1 & 1.688 \end{pmatrix} \right]$$

$$\text{4.13 (a)} \quad \mathbf{z} = \begin{pmatrix} .302 & 0 & 0 & 0 \\ .408 & .561 & 0 & 0 \\ -.087 & .261 & 1.015 & 0 \\ -.858 & -.343 & -.686 & .972 \end{pmatrix} \begin{pmatrix} y+2 \\ y-3 \\ y+1 \\ y-5 \end{pmatrix}$$

$$\text{(b)} \quad \mathbf{z} = \begin{pmatrix} .810 & .305 & .143 & -.480 \\ .305 & .582 & .249 & -.083 \\ .143 & .249 & 1.153 & -.298 \\ -.480 & -.083 & -.298 & .787 \end{pmatrix} \begin{pmatrix} y+2 \\ y-3 \\ y+1 \\ y-5 \end{pmatrix}$$

(c) $(\mathbf{y} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1} (\mathbf{y} - \boldsymbol{\mu}) = (\mathbf{y} - \boldsymbol{\mu})' \boldsymbol{\Sigma}^{-1/2} \boldsymbol{\Sigma}^{-1/2} (\mathbf{y} - \boldsymbol{\mu}) = \mathbf{z}' \mathbf{z}$, which is $\chi^2(p) = \chi^2(4)$

4.14 The variables in (b), (c), and (d) are independent.

4.15 The variables in (a), (c), (d), (f), (i), (j), and (n) are independent.

4.16 (a) $E(\mathbf{y}|\mathbf{x}) = \boldsymbol{\mu}_y + \boldsymbol{\Sigma}_{yx} \boldsymbol{\Sigma}_{xx}^{-1} (\mathbf{x} - \boldsymbol{\mu}_x)$

$$= \begin{pmatrix} 2 \\ -1 \end{pmatrix} + \begin{pmatrix} -3 & 2 \\ 0 & 4 \end{pmatrix} \begin{pmatrix} 5 & -2 \\ -2 & 4 \end{pmatrix}^{-1} \begin{pmatrix} x_1 - 3 \\ x_2 - 1 \end{pmatrix}$$

$$= \begin{pmatrix} 2 \\ -1 \end{pmatrix} + \begin{pmatrix} -.5 & .25 \\ .5 & 1.25 \end{pmatrix} \begin{pmatrix} x_1 - 3 \\ x_2 - 1 \end{pmatrix}$$

$$= \begin{pmatrix} 3.25 \\ -3.75 \end{pmatrix} + \begin{pmatrix} -.5 & .25 \\ .5 & 1.25 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$$

(b) $\text{cov}(\mathbf{y}|\mathbf{x}) = \Sigma_{yy} - \Sigma_{yx}\Sigma_{xx}^{-1}\Sigma_{xy}$

$$= \begin{pmatrix} 7 & 3 \\ 3 & 6 \end{pmatrix} - \begin{pmatrix} -3 & 2 \\ 0 & 4 \end{pmatrix} \begin{pmatrix} 5 & -2 \\ -2 & 4 \end{pmatrix}^{-1} \begin{pmatrix} -3 & 0 \\ 2 & 4 \end{pmatrix}$$

$$= \begin{pmatrix} 7 & 3 \\ 3 & 6 \end{pmatrix} - \begin{pmatrix} 2 & 1 \\ 1 & 5 \end{pmatrix} = \begin{pmatrix} 5 & 2 \\ 2 & 1 \end{pmatrix}$$

4.17 (a) $E(\mathbf{y}|\mathbf{x}) = \boldsymbol{\mu}_y + \Sigma_{yx}\Sigma_{xx}^{-1}(\mathbf{x} - \boldsymbol{\mu}_x)$

$$= \begin{pmatrix} 3 \\ -2 \end{pmatrix} + \begin{pmatrix} 15 & 0 & 3 \\ 8 & 6 & -2 \end{pmatrix} \begin{pmatrix} 50 & 8 & 5 \\ 8 & 4 & 0 \\ 5 & 0 & 1 \end{pmatrix}^{-1} \begin{pmatrix} x_1 - 4 \\ x_2 + 3 \\ x_3 - 5 \end{pmatrix}$$

$$= \begin{pmatrix} 3 \\ -2 \end{pmatrix} - \begin{pmatrix} 15 \\ -24.5 \end{pmatrix} + \begin{pmatrix} 0 & 0 & 3 \\ .67 & .167 & -5.33 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}$$

$$= \begin{pmatrix} -12 \\ 22.5 \end{pmatrix} + \begin{pmatrix} 0 & 0 & 3 \\ .67 & .167 & -5.33 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix}$$

(b) $\text{cov}(\mathbf{y}|\mathbf{x}) = \Sigma_{yy} - \Sigma_{yx}\Sigma_{xx}^{-1}\Sigma_{xy}$

$$= \begin{pmatrix} 14 & -8 \\ -8 & 18 \end{pmatrix} - \begin{pmatrix} 15 & 0 & 3 \\ 8 & 6 & -2 \end{pmatrix} \begin{pmatrix} 50 & 8 & 5 \\ 8 & 4 & 0 \\ 5 & 0 & 1 \end{pmatrix}^{-1} \begin{pmatrix} 15 & 8 \\ 0 & 6 \\ 3 & -2 \end{pmatrix}$$

$$= \begin{pmatrix} 14 & -8 \\ -8 & 18 \end{pmatrix} - \begin{pmatrix} 9 & -6 \\ -6 & 17 \end{pmatrix} = \begin{pmatrix} 5 & -2 \\ -2 & 1 \end{pmatrix}$$

4.18 (a) By the central limit theorem in Section 4.3.2, $\sqrt{n}(\bar{\mathbf{y}} - \boldsymbol{\mu})$ is approximately $N_p(\mathbf{0}, \boldsymbol{\Sigma})$.

(b) $\bar{\mathbf{y}}$ is approximately $N_p(\boldsymbol{\mu}, \boldsymbol{\Sigma}/n)$.

4.19 (a) The plots show almost no deviation from normality.

(b)	Variable	y_1	y_2	y_3	y_4
	$\sqrt{b_1}$.3069	.3111	.0645	.0637
	b_2	1.932	2.107	1.792	1.570

The values of $\sqrt{b_1}$ show a small amount of positive skewness, but none exceeds the upper 2.5% critical value for $\sqrt{b_1}$ given in Table A.1 as .942. The values of b_2 show negative kurtosis. For y_4 , the kurtosis is significant, since $b_2 < 1.74$, the lower 2.5 percentile in Table A.3.

(c)	Variable	y_1	y_2	y_3	y_4
	D	.2848	.2841	.2866	.2851
	Y	.4021	.2934	.6730	.4491

From Table A.4, the lower 2.5 percentile for Y is -3.04 and the upper 97.5 percentile is $.628$. We reject the hypothesis of normality only for y_3 .

- (d) z defined in (4.24) is approximately $N(0, 3/n)$. To obtain a $N(0,1)$ statistic, we calculate $z^* = z/\sqrt{3/n}$.

	Variable	y_1	y_2	y_3	y_4
	z^*	-.3366	-.3095	-.0737	-.0856

4.20	(a)	i	1	2	3	4	5	6	7	8	9	10
		D_i^2	1.06	1.60	7.54	3.54	4.61	.63	.81	2.47	.95	3.78

- (b) The .05 critical value from Table A.6 is 7.01 . $D_{(10)}^2 = 7.54 > 7.01$.

(c)	i	1	2	3	4	5	6	7	8	9	10
	$u_{(i)}$.08	.10	.12	.13	.20	.30	.44	.47	.57	.93
	v_i	.07	.13	.18	.23	.28	.34	.40	.47	.55	.68

The plot of $(v_i, u_{(i)})$ shows some evidence of nonlinearity and an outlier.

- (d) $b_{1,p} = 7.255$, $b_{2,p} = 14.406$. Both (barely) exceed upper .05 critical values in Table A.5.

4.21	(b)	Variable	y_1	y_2	y_3	y_4	y_5
		$\sqrt{b_1}$.2176	.5857	.7461	-.3327	-.1772
		b_2	2.079	1.681	2.583	1.774	2.456

None of the values of $\sqrt{b_1}$ exceeds 1.134 (from Table A.1) or is less than -1.134 . None of the values of b_2 is less than 1.53 (from Table A.3). Thus there is no significant departure from normality.

(c)	Variable	y_1	y_2	y_3	y_4	y_5
	D	.279	.269	.275	.281	.276
	Y	-.305	-1.399	-.805	-.114	-.669

- (d) $z^* = z/\sqrt{3/n}$, where z is defined in (4.24).

	Variable	y_1	y_2	y_3	y_4	y_5
	z^*	-.4848	-1.7183	-1.3627	.8091	.3686

4.22	(a)	i	1	2	3	4	5	6	7	8	9	10	11
		D_i^2	5.20	2.15	7.63	5.34	5.54	1.73	5.21	5.90	2.72	6.02	2.56
	(c)	i	1	2	3	4	5	6	7	8	9	10	11
		$u_{(i)}$.19	.24	.28	.30	.57	.57	.59	.61	.65	.66	.84
		v_i	.18	.27	.34	.39	.45	.50	.55	.61	.66	.73	.82

The plot shows a sharp break from the fourth to the fifth points.

(d) $b_{1,p} = 12.985, b_{2,p} = 29.072$

- 4.23** (a) The $Q - Q$ plots for y_1 and y_5 show little departure from normality. The $Q - Q$ plots for y_2 and y_3 show some evidence of heavier tails than the normal. The $Q - Q$ plots for y_4 and y_6 show some evidence of positive skewness.

(b)	Variable	y_1	y_2	y_3	y_4	y_5	y_6
	$\sqrt{b_1}$.5521	.0302	.7827	1.4627	.2219	.9974
	b_2	3.160	3.275	2.772	6.675	2.176	4.528
(c)	Variable	y_1	y_2	y_3	y_4	y_5	y_6
	D	.276	.274	.275	.260	.286	.271
	Y	-1.469	-1.845	-1.675	-5.249	.889	-2.741
(d)	Variable	y_1	y_2	y_3	y_4	y_5	y_6
	z^*	-1.640	-.062	-2.803	-2.961	-.870	-2.456

- 4.24** (a) $D_i^2 = 7.816, 3.640, 5.730, \dots, 6.433$
- (b) $D_{(51)}^2 = 25.628$. By extrapolation in Table A.6, the .05 critical value for $p = 6$ is approximately 19. Thus we reject the hypothesis of multivariate normality.
- (c) $(v_i, u_{(i)}) = (.021, .024), (.029, .028), \dots, (.306, .523)$. The plot shows nonlinearity for the last four points.
- (d) $b_{1,p} = 16.287, b_{2,p} = 58.337$. By extrapolation to $p = 6$ in Table A.5, both appear to exceed their critical values.

- 4.25** (a) $\lambda_1 = 0.5, \lambda_2 = -0.3, \lambda_3 = 1.1$

(b) $\lambda = (0.7, 0, 1.1)'$

- 4.26** (a) $\lambda_1 = -0.1, \lambda_2 = -0.4, \lambda_3 = 0.4$

(b) $\lambda = (-0.2, -0.5, 0.5)'$

CHAPTER 5

- 5.1** By (5.6), we have

$$\begin{aligned} (\bar{\mathbf{y}} - \boldsymbol{\mu}_0)' \left(\frac{\mathbf{S}}{n} \right)^{-1} (\bar{\mathbf{y}} - \boldsymbol{\mu}_0) &= (\bar{\mathbf{y}} - \boldsymbol{\mu}_0)' \left(\frac{1}{n} \right)^{-1} \mathbf{S}^{-1} (\bar{\mathbf{y}} - \boldsymbol{\mu}_0) \\ &= n(\bar{\mathbf{y}} - \boldsymbol{\mu}_0)' \mathbf{S}^{-1} (\bar{\mathbf{y}} - \boldsymbol{\mu}_0). \end{aligned}$$

5.2 From (5.9), we have

$$\begin{aligned} & \frac{n_1 n_2}{n_1 + n_2} (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{\text{pl}}^{-1} (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2) \\ &= (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \left(\frac{n_1 + n_2}{n_1 n_2} \right)^{-1} \mathbf{S}_{\text{pl}}^{-1} (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2) \\ &= (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \left(\frac{n_1 + n_2}{n_1 n_2} \mathbf{S}_{\text{pl}} \right)^{-1} (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2) \\ &= (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \left[\left(\frac{1}{n_1} + \frac{1}{n_2} \right) \mathbf{S}_{\text{pl}} \right]^{-1} (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2). \end{aligned}$$

5.3 By (5.13) and (5.14),

$$t^2(\mathbf{a}) = \frac{[\mathbf{a}'(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)]^2}{[(n_1 + n_2)/n_1 n_2] \mathbf{a}' \mathbf{S}_{\text{pl}} \mathbf{a}} = \frac{n_1 n_2}{n_1 + n_2} \frac{[(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{\text{pl}}^{-1} (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)]^2}{(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{\text{pl}}^{-1} \mathbf{S}_{\text{pl}} \mathbf{S}_{\text{pl}}^{-1} (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)}.$$

5.4 It is assumed that y and x have a bivariate normal distribution. Let $\mathbf{y}_i = \begin{pmatrix} y_i \\ x_i \end{pmatrix}$. Then d_i can be expressed as $d_i = y_i - x_i = \mathbf{a}' \mathbf{y}_i$, where $\mathbf{a}' = (1, -1)$. By property 1a in Section 4.2, d_i is $N(\mathbf{a}'\boldsymbol{\mu}, \mathbf{a}'\boldsymbol{\Sigma}\mathbf{a})$. Show that $\mathbf{a}'\bar{\mathbf{y}} = \bar{y} - \bar{x}$, $\mathbf{a}'\mathbf{S}\mathbf{a} = s_y^2 - 2s_{yx} + s_x^2 = s_d^2$, and that $T^2 = n(\mathbf{a}'\bar{\mathbf{y}})'(\mathbf{a}'\mathbf{S}\mathbf{a})^{-1}(\mathbf{a}'\bar{\mathbf{y}})$ is the square of $t = \bar{d}/(s_d/\sqrt{n})$.

$$\begin{aligned} \mathbf{5.5} \quad \bar{d} &= \frac{1}{n} \sum_{i=1}^n d_i = \frac{1}{n} \sum_{i=1}^n (y_i - x_i) = \frac{1}{n} \sum_i y_i - \frac{1}{n} \sum_i x_i = \bar{y} - \bar{x}, \\ s_d^2 &= \frac{1}{n-1} \sum_{i=1}^n (d_i - \bar{d})^2 = \frac{1}{n-1} \sum_i (y_i - x_i - \bar{y} + \bar{x})^2 \\ &= \frac{1}{n-1} \sum_i [(y_i - \bar{y}) - (x_i - \bar{x})]^2 \end{aligned}$$

When this is expanded, we obtain $s_d^2 = s_y^2 + s_x^2 - 2s_{yx}$.

5.6 Similar to Problem 5.1.

5.7 By (5.7), $[(\nu - p + 1)/\nu_p] T_{p,\nu}^2 = F_{p,\nu-p+1}$. By (5.29), $(\nu - q)(T_{p+q}^2 - T_p^2)/(\nu + T^2)$ is $T_{q,\nu-p}^2$. Replacing p by q and ν by $\nu - p$ in (5.7), we obtain $\frac{(\nu-p)-q+1}{(\nu-p)q} (\nu - p) \frac{T_{p+q}^2 - T_p^2}{\nu + T_p^2}$ is $F_{q,(\nu-p)-q+1}$.

5.9 Under H_{03} , we have $\mathbf{C}\boldsymbol{\mu}_1 = 0$ and $\mathbf{C}\boldsymbol{\mu}_2 = \mathbf{0}$. Then

$$E(\mathbf{C}\bar{\mathbf{y}}) = \mathbf{C}E(\bar{\mathbf{y}}) = \mathbf{C}E\left(\frac{n_1 \bar{\mathbf{y}}_1 + n_2 \bar{\mathbf{y}}_2}{n_1 + n_2}\right) = \frac{n_1 \mathbf{C}\boldsymbol{\mu}_1 + n_2 \mathbf{C}\boldsymbol{\mu}_2}{n_1 + n_2} = \mathbf{0}.$$

Since $\bar{\mathbf{y}}_1$ and $\bar{\mathbf{y}}_2$ are independent,

$$\begin{aligned} \text{cov}(\bar{\mathbf{y}}) &= \text{cov}\left(\frac{n_1 \bar{\mathbf{y}}_1 + n_2 \bar{\mathbf{y}}_2}{n_1 + n_2}\right) = \frac{n_1^2 \boldsymbol{\Sigma}/n_1 + n_2^2 \boldsymbol{\Sigma}/n_2}{(n_1 + n_2)^2} \\ &= \frac{(n_1 + n_2)\boldsymbol{\Sigma}}{(n_1 + n_2)^2}. \end{aligned}$$

- 5.10** $\mathbf{C}\mathbf{S}_{\mathbf{p}1}\mathbf{C}'/(n_1+n_2)$ is the sample covariance matrix of $\mathbf{C}\bar{\mathbf{y}}$. Hence the equation immediately above (5.39) exhibits the characteristic form of the T^2 -distribution.
- 5.11** $T^2 = .061$
- 5.12** (a) $T^2 = 85.3327$
 (b) $t_1 = 2.5039, t_2 = .2665, t_3 = -2.5157, t_4 = .9510, t_5 = .3161$
- 5.13** $T^2 = 30.2860$
- 5.14** (a) $T^2 = 1.8198$
 (b) $t_1 = 1.1643, t_2 = 1.1006, t_3 = .9692, t_4 = .7299$. None of these is significant. In fact, ordinarily they would not have been examined because the T^2 -test in part (a) did not reject H_0 .
- 5.15** $T^2 = 79.5510$
- 5.16** (a) $T^2 = 133.4873$
 (b) $t_1 = 3.8879, t_2 = -3.8652, t_3 = -5.6911, t_4 = -5.0426$
 (c) $\mathbf{a}' = (.345, -.130, -.106, -.143)$
 (d) $t^2(\mathbf{a}) = 133.4873$
 (e) $R^2 = .782975, T^2 = 133.4873$
 (f) By (5.32), $t^2(y_1|y_2, y_3, y_4) = 35.9336, t^2(y_2|y_1, y_3, y_4) = 5.7994, t^2(y_3|y_1, y_2, y_4) = 1.7749, t^2(y_4|y_1, y_2, y_3) = 8.2592$
 (g) By (5.29), $T^2(y_3, y_4|y_1, y_2) = 12.5206, F(y_3, y_4|y_1, y_2) = 6.0814$
- 5.17** By (5.34), the test for parallelism gives $T^2 = 132.6863$. The discriminant function coefficient vector is given by (5.35) as $\mathbf{a}' = (-.362, -.223, -.137)$.
- 5.18** (a) $T^2 = 66.6604$
 (b) $t_1 = -.6556, t_2 = 2.6139, t_3 = -3.2884, t_4 = -4.6315, t_5 = 1.8873, t_6 = -3.2205$
 (c) By (5.32),

$$\begin{aligned} t^2(y_1|y_2, y_3, y_4, y_5, y_6) &= .0758, t^2(y_2|y_1, y_3, y_4, y_5, y_6) = 6.4513, \\ t^2(y_3|y_1, y_2, y_4, y_5, y_6) &= 6.9518, t^2(y_4|y_1, y_2, y_3, y_5, y_6) = 6.0309, \\ t^2(y_5|y_1, y_2, y_3, y_4, y_6) &= 3.7052, t^2(y_6|y_1, y_2, y_3, y_4, y_5) = 6.2619 \end{aligned}$$

 (d) By (5.29), $T^2(y_4, y_5, y_6|y_1, y_2, y_3) = 27.547$
- 5.19** (a) $T^2 = 70.5679$
 (b) $T^2(y_5, y_6|y_3, y_4) = 13.1517$
 (c) $T^2(y_1, y_2|y_3, y_4, y_5, y_6) = 8.5162$

5.20 (a) $T^2 = 18.4625$

(b) $\mathbf{a}' = (-.057, -.010, -.242, -.071)$

(c) By (5.32),

$$\begin{aligned}t^2(y_1|y_2, y_3, y_4) &= 3.3315, t^2(y_2|y_1, y_3, y_4) = .0102, \\t^2(y_3|y_1, y_2, y_4) &= 1.4823, t^2(y_4|y_1, y_2, y_3) = .0013\end{aligned}$$

5.21 (a) $T^2 = 15.1912$

(b) $\mathbf{a}' = (-.036, .048)$

(c) $t_1 = -3.8371, t_2 = -2.4362$

5.22 $T^2 = 22.3238$

5.23 (a) $T^2 = 206.1188$

(b) $t^2(d_1|d_2, d_3) = 59.0020, t^2(d_2|d_1, d_3) = 53.4507, t^2(d_3|d_1, d_2) = 80.9349$

CHAPTER 6

6.1 (a) Using $\bar{y}_{i\cdot} = y_{i\cdot}/n$, we have

$$\begin{aligned}\sum_{i=1}^k \sum_{j=1}^n (y_{ij} - \bar{y}_{i\cdot})^2 &= \sum_{ij} (y_{ij}^2 - 2y_{ij}\bar{y}_{i\cdot} + \bar{y}_{i\cdot}^2) \\&= \sum_{ij} y_{ij}^2 - \sum_i \bar{y}_{i\cdot} \sum_j y_{ij} + n \sum_i \bar{y}_{i\cdot}^2 \\&= \sum_{ij} y_{ij}^2 - 2 \sum_i \frac{y_{i\cdot}}{n} y_{i\cdot} + n \sum_i \left(\frac{y_{i\cdot}}{n}\right)^2 \\&= \sum_{ij} y_{ij}^2 - 2 \sum_i \frac{y_{i\cdot}^2}{n} + \sum_i \frac{y_{i\cdot}^2}{n}.\end{aligned}$$

6.2 $\frac{|\mathbf{E}^{-1}| |\mathbf{E}|}{|\mathbf{E}^{-1}| |\mathbf{E} + \mathbf{H}|} = \frac{|\mathbf{E}^{-1}\mathbf{E}|}{|\mathbf{E}^{-1}(\mathbf{E} + \mathbf{H})|} = \frac{|\mathbf{I}|}{|\mathbf{I} + \mathbf{E}^{-1}\mathbf{H}|} = \frac{1}{\prod_{i=1}^s (1 + \lambda_i)};$
see Section 2.11.2.

6.3 $(\mathbf{E}^{-1}\mathbf{H} - \lambda\mathbf{I})\mathbf{a} = \mathbf{0}$

$[(\mathbf{E}^{1/2}\mathbf{E}^{1/2})^{-1}\mathbf{H} - \lambda\mathbf{I}]\mathbf{a} = \mathbf{0}$

$[(\mathbf{E}^{1/2})^{-1}(\mathbf{E}^{1/2})^{-1}\mathbf{H} - \lambda\mathbf{I}]\mathbf{a} = \mathbf{0}$

$[(\mathbf{E}^{1/2})^{-1}\mathbf{H} - \lambda\mathbf{E}^{1/2}]\mathbf{a} = \mathbf{0}$

$[(\mathbf{E}^{1/2})^{-1}\mathbf{H} - \lambda\mathbf{E}^{1/2}](\mathbf{E}^{1/2})^{-1}\mathbf{E}^{1/2}\mathbf{a} = \mathbf{0}$

$[(\mathbf{E}^{1/2})^{-1}\mathbf{H}(\mathbf{E}^{1/2})^{-1} - \lambda\mathbf{I}]\mathbf{E}^{1/2}\mathbf{a} = \mathbf{0}$

6.4 We need to show that $(2N + s + 1)/(2m + s + 1) = (\nu_E - p + s)/d$. Using the definitions $N = \frac{1}{2}(\nu_E - p - 1)$, $m = \frac{1}{2}(|\nu_H - p| - 1)$, $d = \max(p, \nu_H)$, and $s = \min(p, \nu_H)$, we have $2N + s + 1 = 2(\frac{1}{2})(\nu_E - p - 1) + s + 1 = \nu_E - p - 1 + s + 1 = \nu_E - p + s$. For the denominator, we have $2m + s + 1 = 2(\frac{1}{2})(|\nu_H - p| - 1) + s + 1 = |\nu_H - p| + s$. Suppose $\nu_H > p$. Then $|\nu_H - p| + s = \nu_H - p + p = \nu_H = d$. On the other hand, if $\nu_H < p$, then $|\nu_H - p| + s = p - \nu_H + \nu_H = p = d$.

6.5 If $p \leq \nu_H$, we have $s = p$ and $|\nu_H - p| = \nu_H - p$. Then (6.29) becomes

$$\begin{aligned}\frac{2(sN + 1)U^{(s)}}{s^2(2m + s + 1)} &= \frac{2[p(\frac{1}{2})(\nu_E - p - 1) + 1]U^{(s)}}{p^2[2(\frac{1}{2})(\nu_H - p - 1) + p + 1]} \\ &= \frac{[p(\nu_E - p - 1) + 2]U^{(s)}}{p^2(\nu_H - p - 1 + p + 1)} \\ &= \frac{[p(\nu_E - p - 1) + 2]U^{(s)}}{p^2\nu_H},\end{aligned}$$

which is the same as (6.30) because $p = s$. If $\nu_H \leq p$, then $s = \nu_H$, $|\nu_H - p| = p - \nu_H$, and (6.29) can be shown to equal (6.30) in a similar manner.

6.6 When $s = 1$, we have $V^{(1)} = \lambda_1/(1 + \lambda_1)$, $U^{(1)} = \lambda_1$, $\Lambda = 1/(1 + \lambda_1)$, and $\theta = \lambda_1/(1 + \lambda_1)$. Solving the last of these for λ_1 gives $\lambda_1 = \theta/(1 - \theta)$, and the results in (6.33), (6.34), and (6.35) follow immediately.

6.7 With $T^2 = (n_1 + n_2 - 2)U^{(1)}$ and $U^{(1)} = \theta/(1 - \theta)$, we obtain (5.19). We obtain (5.18) from (5.19) by $V^{(1)} = \theta$. A similar argument leads to (5.16).

6.8 (a) With $\bar{\mathbf{y}}_{i\cdot} = \mathbf{y}_{i\cdot}/n_i$ and $\bar{\mathbf{y}}_{..} = \mathbf{y}_{..}/N$, we obtain

$$\begin{aligned}\mathbf{H} &= \sum_{i=1}^k n_i(\bar{\mathbf{y}}_{i\cdot} - \bar{\mathbf{y}}_{..})(\bar{\mathbf{y}}_{i\cdot} - \bar{\mathbf{y}}_{..})' \\ &= \sum_i n_i(\bar{\mathbf{y}}_{i\cdot}\bar{\mathbf{y}}_{i\cdot}' - \bar{\mathbf{y}}_{i\cdot}\bar{\mathbf{y}}_{..}' - \bar{\mathbf{y}}_{..}\bar{\mathbf{y}}_{i\cdot}' + \bar{\mathbf{y}}_{..}\bar{\mathbf{y}}_{..}') \\ &= \sum_i n_i\bar{\mathbf{y}}_{i\cdot}\bar{\mathbf{y}}_{i\cdot}' - (\sum_i n_i\bar{\mathbf{y}}_{i\cdot})\bar{\mathbf{y}}_{..}' - \bar{\mathbf{y}}_{..}\sum_i n_i\bar{\mathbf{y}}_{i\cdot}' + \bar{\mathbf{y}}_{..}\bar{\mathbf{y}}_{..}'\sum_i n_i \\ &= \sum_i n_i \frac{\mathbf{y}_{i\cdot}\mathbf{y}_{i\cdot}'}{n_i^2} - \frac{(\sum_i \mathbf{y}_{i\cdot})\mathbf{y}_{..}'}{N} - \frac{\mathbf{y}_{..}}{N}\sum_i \mathbf{y}_{i\cdot}' + \frac{N\mathbf{y}_{..}\mathbf{y}_{..}'}{N^2} \\ &= \sum_i \frac{\mathbf{y}_{i\cdot}\mathbf{y}_{i\cdot}'}{n_i} - \frac{\mathbf{y}_{..}\mathbf{y}_{..}'}{N} - \frac{\mathbf{y}_{..}\mathbf{y}_{..}'}{N} + \frac{\mathbf{y}_{..}\mathbf{y}_{..}'}{N}.\end{aligned}$$

6.9 $\bar{\mathbf{y}}_{1\cdot} - \bar{\mathbf{y}}_{..}$ becomes

$$\bar{\mathbf{y}}_{1\cdot} - \frac{n_1\bar{\mathbf{y}}_{1\cdot} + n_2\bar{\mathbf{y}}_{2\cdot}}{n_1 + n_2} = \frac{n_1\bar{\mathbf{y}}_{1\cdot} + n_2\bar{\mathbf{y}}_{1\cdot} - n_1\bar{\mathbf{y}}_{1\cdot} - n_2\bar{\mathbf{y}}_{2\cdot}}{n_1 + n_2} = \frac{n_2(\bar{\mathbf{y}}_{1\cdot} - \bar{\mathbf{y}}_{2\cdot})}{n_1 + n_2}.$$

The first term in the sum is

$$\frac{n_1 n_2^2}{(n_1 + n_2)^2}(\bar{\mathbf{y}}_{1\cdot} - \bar{\mathbf{y}}_{2\cdot})(\bar{\mathbf{y}}_{1\cdot} - \bar{\mathbf{y}}_{2\cdot})'.$$

The second term in the sum is

$$\frac{n_1^2 n_2}{(n_1 + n_2)^2} (\bar{\mathbf{y}}_{1..} - \bar{\mathbf{y}}_{2..})(\bar{\mathbf{y}}_{1..} - \bar{\mathbf{y}}_{2..})'.$$

6.10 $\theta = \frac{\lambda_1}{1 + \lambda_1} = \frac{\text{SSH}(z)/\text{SSE}(z)}{1 + \text{SSH}(z)/\text{SSE}(z)} = \frac{\text{SSH}(z)}{\text{SSE}(z) + \text{SSH}(z)}$

6.11 From $r_i^2 = \lambda_i/(1+\lambda_i)$, obtain $\lambda_i = r_i^2/(1-r_i^2)$. Substitute this into $1/(1+\lambda_i)$ to obtain the result.

6.12 Substitute $A_p = V^{(s)}/s$ into (6.49) to obtain (6.26).

6.13 When $s = 1$, (6.50) becomes

$$A_{\text{LH}} = \frac{U^{(1)}}{1 + U^{(1)}}.$$

By (6.33), $U^{(1)} = \lambda_1$.

6.14 Substitute $A_{\text{LH}} = U^{(s)}/(s + U^{(s)})$ from (6.50) into (6.51) to obtain F_3 in (6.30).

6.15 To show $\text{cov}(c_i \bar{\mathbf{y}}_{i..}) = c_i^2 \Sigma / n$, use (3.74), $\text{cov}(\mathbf{Ay}) = \mathbf{A}\Sigma\mathbf{A}'$, with $\mathbf{A} = c_i \mathbf{I}$.

6.16 By (6.9),

$$\begin{aligned} \mathbf{H}_z &= n \sum_{i=1}^k (\bar{\mathbf{z}}_{i..} - \bar{\mathbf{z}}_{..})(\bar{\mathbf{z}}_{i..} - \bar{\mathbf{z}}_{..})' \\ &= n \sum_i (\mathbf{C}\bar{\mathbf{y}}_{i..} - \mathbf{C}\bar{\mathbf{y}}_{..})(\mathbf{C}\bar{\mathbf{y}}_{i..} - \mathbf{C}\bar{\mathbf{y}}_{..})' \\ &= n \sum_i [\mathbf{C}(\bar{\mathbf{y}}_{i..} - \bar{\mathbf{y}}_{..})][\mathbf{C}(\bar{\mathbf{y}}_{i..} - \bar{\mathbf{y}}_{..})]' \\ &= n\mathbf{C}[\sum_i (\bar{\mathbf{y}}_{i..} - \bar{\mathbf{y}}_{..})(\bar{\mathbf{y}}_{i..} - \bar{\mathbf{y}}_{..})']\mathbf{C}' \quad [\text{by (2.45)}] \end{aligned}$$

6.17 \mathbf{C} is not square.

6.18 $E(\mathbf{C}\bar{\mathbf{y}}_{..}) = \mathbf{C}E(\bar{\mathbf{y}}_{..}) = \mathbf{C}E(\sum_{i=1}^k \bar{\mathbf{y}}_{i..}/k)$
 $= \mathbf{C} \sum_i E(\bar{\mathbf{y}}_{i..})/k = \mathbf{C} \sum_i \boldsymbol{\mu}_i/k$
 $= \mathbf{0} \quad [\text{by } H_{03} \text{ in (6.82)}]$

$\text{cov}(\mathbf{C}\bar{\mathbf{y}}_{..}) = \mathbf{C}\Sigma\mathbf{C}'/kn$ if there are no differences in the group means, $\mathbf{C}\boldsymbol{\mu}_1, \mathbf{C}\boldsymbol{\mu}_2, \dots$. This condition is assured by H_{01} in (6.77).

6.19 For our purposes, it will suffice to show that T^2 has the characteristic form of the T^2 -distribution in (5.6).

6.20 If $\boldsymbol{\Sigma} = \sigma^2 \mathbf{I}$, (6.88) becomes

$$\varepsilon = \frac{[\text{tr}(\sigma^2 \mathbf{I} - \mathbf{J}\sigma^2 \mathbf{I}/p)]^2}{(p-1)\text{tr}(\sigma^2 \mathbf{I} - \mathbf{J}\sigma^2 \mathbf{I}/p)^2} = \frac{[\sigma^2 \text{tr}(\mathbf{I} - \mathbf{J}/p)]^2}{\sigma^4(p-1)\text{tr}(\mathbf{I} - \mathbf{J}/p)^2}.$$

Show that $(\mathbf{I} - \mathbf{J}/p)^2 = \mathbf{I} - \mathbf{J}/p$. Then

$$\varepsilon = \frac{\sigma^4(p - p/p)^2}{\sigma^4(p - 1)(p - p/p)} = \frac{(p - 1)^2}{(p - 1)^2} = 1.$$

- 6.21** The (univariate) expected mean square corresponding to $\bar{\mu}_i$ in a one-way ANOVA is $\sigma^2 + N\mu_i^2$. Thus the mean square for $\bar{\mu}_i$ is tested with MSE. The corresponding multivariate test therefore uses \mathbf{H}^* and \mathbf{E} .

- 6.22** From (6.104) we have

$$\Lambda = \frac{|\mathbf{A}\mathbf{E}\mathbf{A}'|}{|\mathbf{A}(\mathbf{E} + \mathbf{H}^*)\mathbf{A}'|} = \frac{|\mathbf{A}\mathbf{E}\mathbf{A}'|}{|\mathbf{A}\mathbf{E}\mathbf{A}' + \mathbf{A}\mathbf{H}^*\mathbf{A}'|}.$$

Substitute $\mathbf{H}^* = kn\bar{\mathbf{y}}.. \bar{\mathbf{y}}'..$ to obtain

$$\Lambda = \frac{|\mathbf{A}\mathbf{E}\mathbf{A}'|}{|\mathbf{A}\mathbf{E}\mathbf{A}' + \sqrt{kn}\mathbf{A}\bar{\mathbf{y}}..(\sqrt{kn}\mathbf{A}\bar{\mathbf{y}}..)'|}.$$

Now use (2.95) with $\mathbf{B} = \mathbf{A}\mathbf{E}\mathbf{A}'$ and $\mathbf{c} = \sqrt{kn}\mathbf{A}\bar{\mathbf{y}}..$ to obtain

$$\Lambda = \frac{1}{1 + kn(\mathbf{A}\bar{\mathbf{y}}..)'(\mathbf{A}\mathbf{E}\mathbf{A}')^{-1}(\mathbf{A}\bar{\mathbf{y}}..)}.$$

Multiply and divide by ν_E and use (6.100) to obtain (6.105).

- 6.23** Solve for T^2 in (6.105).

- 6.24** In $\mathbf{C}_1\mathbf{B}'$ the rows of \mathbf{C}_1 are multiplied by the rows of \mathbf{B} . Show that $\mathbf{C}_1\mathbf{B}' = \mathbf{O}$.

- 6.25** As noted, the function $(\bar{\mathbf{y}} - \mathbf{A}\hat{\beta})'\mathbf{S}^{-1}(\bar{\mathbf{y}} - \mathbf{A}\hat{\beta})$ is similar to $\text{SSE} = (\mathbf{y} - \mathbf{X}\hat{\beta})'(\mathbf{y} - \mathbf{X}\hat{\beta})$ in (10.4) and (10.6). By an argument similar to that used in Section 10.2.2 to obtain $\hat{\beta} = (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{y}$, it follows that $\hat{\beta} = (\mathbf{A}'\mathbf{S}^{-1}\mathbf{A})^{-1}\mathbf{A}'\mathbf{S}^{-1}\bar{\mathbf{y}}$. An alternative approach (for those familiar with differentiation with respect to a vector) is to expand $(\bar{\mathbf{y}} - \mathbf{A}\hat{\beta})'\mathbf{S}^{-1}(\bar{\mathbf{y}} - \mathbf{A}\hat{\beta})$ to four terms, differentiate with respect to $\hat{\beta}$, and set the result equal to $\mathbf{0}$.

- 6.26** Expand $n(\bar{\mathbf{y}} - \mathbf{A}\hat{\beta})'\mathbf{S}^{-1}(\bar{\mathbf{y}} - \mathbf{A}\hat{\beta})$ to four terms and substitute

$$\hat{\beta} = (\mathbf{A}'\mathbf{S}^{-1}\mathbf{A})^{-1}\mathbf{A}'\mathbf{S}^{-1}\bar{\mathbf{y}}$$

into the last one.

- 6.27 (a)** $\mathbf{E} = \begin{pmatrix} 13.41 & 7.72 & 8.68 & 5.86 \\ 7.72 & 8.48 & 7.53 & 6.21 \\ 8.68 & 7.53 & 11.61 & 7.04 \\ 5.86 & 6.21 & 7.04 & 10.57 \end{pmatrix}$

$$\mathbf{H} = \begin{pmatrix} 1.05 & 2.17 & -1.38 & -.76 \\ 2.17 & 4.88 & -2.37 & -1.26 \\ -1.38 & -2.37 & 2.38 & 1.38 \\ -.76 & -1.26 & 1.38 & .81 \end{pmatrix}$$

$\Lambda = .224, V^{(s)} = .860, U^{(s)} = 3.08$, and $\theta = .747$ All four are significant.

- (b) $\eta_{\Lambda}^2 = 1 - \Lambda = .776, \eta_{\theta}^2 = \theta = .747, A_{\Lambda} = 1 - \Lambda^{1/s} = .526, A_{\text{LH}} = .606, A_p = V^{(s)}/s = .430$
- (c) The eigenvalues of $\mathbf{E}^{-1}\mathbf{H}$ are 2.9515 and .1273. The essential dimensionality of the space of the mean vectors is 1.
- (d) For 1, 2 vs. 3 we have $\Lambda = .270, V^{(s)} = .730, U^{(s)} = 2.702$, and $\theta = .730$. All four are significant. For 1 vs. 2 we obtain $\Lambda = .726, V^{(s)} = .274, U^{(s)} = .377$, and $\theta = .274$. All four are significant.

(e)	Variable	y_1	y_2	y_3	y_4
	F	1.29	9.50	3.39	1.27

The F 's for y_2 and y_3 are significant. For the discriminant function $z = \mathbf{a}'\mathbf{y}$, where \mathbf{a} is the first eigenvector of $\mathbf{E}^{-1}\mathbf{H}$, we have $\mathbf{a}' = (-.032, -.820, .533, .208)$. Again y_2 and y_3 contribute most to separation of groups.

- (f) By (6.126), $\Lambda(y_3, y_4 | y_1, y_2) = \Lambda(y_1, y_2, y_3, y_4) / \Lambda(y_1, y_2) = .224/.568 = .395 < \Lambda_{.05} = .725$
- (g) By (6.127),

$$\begin{aligned} \Lambda(y_1 | y_2, y_3, y_4) &= \Lambda(y_1, y_2, y_3, y_4) / \Lambda(y_2, y_3, y_4) \\ &= .224/.240 = .934 > \Lambda_{.05} = .819 \\ \Lambda(y_2 | y_1, y_3, y_4) &= .224/.538 = .417 < .819 \\ \Lambda(y_3 | y_1, y_2, y_4) &= .224/.369 = .609 < .819 \\ \Lambda(y_4 | y_1, y_2, y_3) &= .224/.243 = .924 > .819 \end{aligned}$$

- 6.28** (a) S effect: $\Lambda = .00065, V^{(s)} = 2.357, U^{(s)} = 142.304, \theta = .993$. All are significant.

V effect: $\Lambda = .065, V^{(s)} = 1.107, U^{(s)} = 11.675, \theta = .920$. All are significant.

SV interaction: $\Lambda = .138, V^{(s)} = 1.321, U^{(s)} = 3.450, \theta = .726$. All are significant.

- (b) Contrast on V comparing 2 vs. 1, 3: $\Lambda = .0804, V^{(s)} = .920, U^{(s)} = 11.445, \theta = .920$. All are significant.
- (c) Linear contrast for S : $\Lambda = .0073, V^{(s)} = .993, U^{(s)} = 135.273, \theta = .993$. All are significant.

Quadratic contrast for S : $\Lambda = .168, V^{(s)} = .832, U^{(s)} = 4.956, \theta = .832$. All are significant.

Cubic contrast for S : $\Lambda = .325, V^{(s)} = .675, U^{(s)} = 2.076, \theta = .675$. All are significant.

(d) The ANOVA F 's for each variable are as follows:

Source	y_1	y_2	y_3	y_4
S	980.21	214.24	876.13	73.91
V	251.22	9.47	14.77	27.12
SV	20.37	2.84	3.44	2.08

All F 's are significant except the last one, 2.08.

(e) Test of significance of y_3 and y_4 adjusted for y_1 and y_2 :

	S	V	SV
$\Lambda(y_3, y_4 y_1, y_2)$.1226	.9336	.6402

(f) Test of significance of each variable adjusted for the other three:

	S	V	SV
$\Lambda(y_1 y_2, y_3, y_4)$.1158	.2099	.3082
$\Lambda(y_2 y_1, y_3, y_4)$.5586	.8134	.7967
$\Lambda(y_3 y_1, y_2, y_4)$.2271	.9627	.7604
$\Lambda(y_4 y_1, y_2, y_3)$.6692	.9795	.8683

6.29 V = velocity (fixed), L = lubricant (random)

V effect (using \mathbf{H}_{VL} for error matrix): $\Lambda = .0492, V^{(s)} = .951, U^{(s)} = 19.315, \theta = .951$. With $p = 2, \nu_H = 1$, and $\nu_E = 3, \Lambda_{.05} = .050, V_{.05}^{(s)} = .950, U_{.05}^{(s)} = T_{.05}^2 / \nu_E = 19.00, \theta_{.05} = .950$. Thus all four test statistics are significant.

L effect (using \mathbf{E} for error matrix): $\Lambda = .692, V^{(s)} = .314, U^{(s)} = .438, \theta = .295$. None is significant.

VL interaction (using \mathbf{E} for error matrix): $\Lambda = .932, V^{(s)} = .069, U^{(s)} = .073, \theta = .061$. None is significant.

6.30	Source	Λ	$V^{(s)}$	$U^{(s)}$	θ	Significant?
	(a) Reagent	.0993	1.126	6.911	.868	Yes
	(b) Contrast 1 vs. 2, 3, 4	.146	.854	5.871	.854	Yes
	Subjects	.00000082	2.847	1091.127	.999	Yes

6.31 P = proportion of filler, T = surface treatment, F = filler:

Source	Λ	$V^{(s)}$	$U^{(s)}$	θ	Significant?
P	.138	.977	5.441	.841	Yes
T	.080	.920	11.503	.920	Yes
PT	.712	.295	.396	.271	No
F	.019	.980	51.180	.981	Yes
PF	.179	.958	3.835	.784	Yes
TF	.355	.645	1.815	.645	Yes
PTF	.752	.264	.309	.172	No

6.32 A = period; P, T , and F are defined in Problem 6.31:

Source	Λ	$V^{(s)}$	$U^{(s)}$	θ	Significant?
A	.021	.979	47.099	.979	Yes
AP	.475	.545	1.063	.505	No
AT	.142	.858	6.049	.858	Yes
APT	.777	.228	.282	.208	No
AF	.095	.905	9.486	.905	Yes
APF	.622	.387	.594	.363	No
ATF	.387	.613	1.586	.613	Yes
$APTF$.781	.229	.267	.169	No

For the between-subject factors and interactions, we have

Source	df	F	p-Value
P	2	21.79	< .0001
T	1	78.34	< .0001
PT	2	1.28	.3143
F	1	345.04	< .0001
PF	2	15.79	.0004
TF	1	5.36	.0392
PTF	2	.48	.6294
Error	12		

6.33 For parallelism, we use (6.78) to obtain $\Lambda = .2397$. For levels, we use (6.80) and (6.81) to obtain $\Lambda = .9651$ and $F = .597$. For flatness we use (6.83) to obtain $T^2 = 110.521$.

6.34 (a) By (6.89), $T^2 = 20.7420$. By (6.104) or (6.105), $\Lambda = .5655$.

(b) For each row \mathbf{c}_i' of \mathbf{C} , we use $T_i^2 = n(\mathbf{c}_i'\bar{\mathbf{y}})'(\mathbf{c}_i'\mathbf{S}\mathbf{c}_i)^{-1}\mathbf{c}_i'\bar{\mathbf{y}}$, as in Example 6.9.2: $T_1^2 = 17.0648$, $T_2^2 = .3238$, $T_3^2 = .2714$. This can also be done by Wilks' Λ using $\Lambda_i = \mathbf{c}_i'\mathbf{E}\mathbf{c}_i/\mathbf{c}_i'(\mathbf{E} + \mathbf{H}^*)\mathbf{c}_i$: $\Lambda_1 = .6127$, $\Lambda_2 = .9882$, $\Lambda_3 = .9900$.

6.35 The six variables represent two within-subjects factors: y_1 is A_1B_1 , y_2 is A_1B_2 , y_3 is A_1B_3 , x_1 is A_2B_1 , x_2 is A_2B_2 , and x_3 is A_2B_2 . Using linear and quadratic effects (other orthogonal contrasts could be used), the matrices

A, B, and G in (6.97), (6.98), and (6.99) become

$$\mathbf{A} = \begin{pmatrix} 1 & 1 & 1 & -1 & -1 & -1 \end{pmatrix}$$

$$\mathbf{B} = \begin{pmatrix} 1 & 0 & -1 & 1 & 0 & -1 \\ 1 & -2 & 1 & 1 & -2 & 1 \end{pmatrix}$$

$$\mathbf{G} = \begin{pmatrix} 1 & 0 & -1 & -1 & 0 & 1 \\ 1 & -2 & 1 & -1 & 2 & -1 \end{pmatrix}.$$

Using these in T^2 as given by (6.100), (6.101), and (6.102), we obtain $T_A^2 = 193.0901$, $T_B^2 = 2.8000$, and $T_{AB}^2 = 6.8676$. Using MANOVA tests for the same within-subjects factors, we obtain

Source	Λ	$V^{(s)}$	$U^{(s)}$	θ	Significant?
A	.202	.798	3.941	.798	Yes
B	.946	.054	.057	.054	No
AB	.877	.123	.140	.123	Yes

- 6.36** MANOVA tests for the within-subjects effect T (time), and interactions of time with the between-subjects effects C (cancer) and G (gender):

Source	Λ	$V^{(s)}$	$U^{(s)}$	θ
T	.258	.742	2.874	.742
TC	.363	.809	1.299	.444
TG	.929	.071	.077	.071
TCG	.809	.201	.225	.130

ANOVA F -tests for between-subjects factors and interactions:

Source	df	F	p -Value
C	5	4.16	.003
G	1	2.69	.107
CG	5	.37	.869

- 6.37 (a)** $T^2 = 79.551$

(b) Using $t_i = \mathbf{c}'_i \bar{\mathbf{y}} / \sqrt{\mathbf{c}'_i \mathbf{S} \mathbf{c}_i / n}$, where \mathbf{c}'_i is the i th row of \mathbf{C} , we obtain $t_1 = 7.155$, $t_2 = -.445$, $t_3 = -.105$.

- 6.38 (a)** $T^2 = 1712.2201$

(b) Using $t_i = \mathbf{c}'_i \bar{\mathbf{y}} / \sqrt{\mathbf{c}'_i \mathbf{S} \mathbf{c}_i / n}$, we obtain $t_1 = 332.358$, $t_2 = 54.589$, $t_3 = .056$, $t_4 = 7.637$, $t_5 = 4.344$, $t_6 = 1.968$.

- 6.39 (a)** Using $T^2 = N(\mathbf{C}\bar{\mathbf{y}}_{..})'(\mathbf{C}\mathbf{S}_{p1}\mathbf{C}')^{-1}(\mathbf{C}\bar{\mathbf{y}}_{..})$ in (6.121), we obtain $T^2 = 17.582 < T_{.05,3,9}^2 = 27.202$.

(b) $t_1 = .951$, $t_2 = 1.606$, $t_3 = .127$ [Since the T^2 -test in part (a) did not reject H_0 , these would ordinarily not be calculated.]

(c) Using $\Lambda = |\mathbf{CEC}'|/|\mathbf{C}(\mathbf{E} + \mathbf{H})\mathbf{C}'|$ in (6.123), we obtain $\Lambda = .3107 > \Lambda_{.05,3,2,9} = .203$.

- (d) To compare groups using each row of \mathbf{C} , we use $\Lambda_i = \mathbf{c}_i' \mathbf{E} \mathbf{c}_i / \mathbf{c}_i' (\mathbf{E} + \mathbf{H}) \mathbf{c}_i$ to obtain $\Lambda_1 = .833$, $\Lambda_2 = .988$, $\Lambda_3 = .650$. [Since the Λ -test in part (c) did not reject H_0 , we would ordinarily not have calculated these.]
- 6.40** (a) Using $T^2 = N(\bar{\mathbf{y}}_{..})'(\mathbf{C}\mathbf{S}_{p1}\mathbf{C}')^{-1}(\bar{\mathbf{y}}_{..})$ in (6.121), we obtain $T^2 = 33.802 > T^2_{.05,4,24} = 12.983$.
- (b) Using $t_i^2 = N(\mathbf{c}_i' \bar{\mathbf{y}})^2 / \mathbf{c}_i' \mathbf{S}_{p1} \mathbf{c}_i$, we obtain $t_1^2 = .675$, $t_2^2 = .393$, $t_3^2 = 32.626$. Only the cubic effect is significant.
- (c) For an overall test comparing groups, we use (6.123),

$$\Lambda = |\mathbf{C}\mathbf{E}\mathbf{C}'| / |\mathbf{C}(\mathbf{E} + \mathbf{H})\mathbf{C}'| = .4361.$$

- (d) To compare groups using each row of \mathbf{C} , we use $\Lambda_i = \mathbf{c}_i' \mathbf{E} \mathbf{c}_i / \mathbf{c}_i' (\mathbf{E} + \mathbf{H}) \mathbf{c}_i$: $\Lambda_1 = .534$, $\Lambda_2 = .764$, $\Lambda_3 = .941$.
- 6.41** (a) Using $T^2 = N(\bar{\mathbf{y}}_{..})'(\mathbf{C}\mathbf{S}_{p1}\mathbf{C}')^{-1}(\bar{\mathbf{y}}_{..})$ in (6.121), we obtain $T^2 = 45.500$.
- (b) Using $t_i^2 = N(\mathbf{c}_i' \bar{\mathbf{y}})^2 / \mathbf{c}_i' \mathbf{S}_{p1} \mathbf{c}_i$, we obtain $t_1^2 = 18.410$, $t_2^2 = 8.385$, $t_3^2 = 3.446$, $t_4^2 = .011$, $t_5^2 = .098$, $t_6^2 = 2.900$.
- (c) For an overall test comparing groups, we use (6.123),

$$\Lambda = |\mathbf{C}\mathbf{E}\mathbf{C}'| / |\mathbf{C}(\mathbf{E} + \mathbf{H})\mathbf{C}'| = .304.$$

- (d) To compare groups using each row of \mathbf{C} , we use $\Lambda_i = \mathbf{c}_i' \mathbf{E} \mathbf{c}_i / \mathbf{c}_i' (\mathbf{E} + \mathbf{H}) \mathbf{c}_i$: $\Lambda_1 = .695$, $\Lambda_2 = .925$, $\Lambda_3 = .731$, $\Lambda_4 = .814$, $\Lambda_5 = .950$, $\Lambda_6 = .894$.
- 6.42** (a) Combined groups (pooled covariance matrix). Using $t = \text{number of minutes} - 30$, we obtain, by (6.114),

$$\hat{\beta}' = (98.1, .981, .0418, -.00101, -.000048)$$

By (6.115), we obtain $T^2 = .216$. By (6.117), we have

$$\hat{\mu}' = (95.5, 96.7, 95.6, 93.8, 98.1, 99.2)$$

- (b) Group 1: $\hat{\beta}'_1 = (100.7, .819, .040, -.00085, -.000038)$, $T^2 = .0113$, $\hat{\mu}'_1 = (105.2, 104.4, 101.5, 98.6, 100.6, 108.1)$
- (c) Groups 2–4: $\hat{\beta}'_2 = (97.4, 1.010, .0403, -.00103, -.000049)$, $T^2 = .2554$, $\hat{\mu}'_2 = (92.6, 94.4, 93.8, 92.4, 97.4, 96.6)$
- 6.43** (a) For the control group, the overall test is

$$T^2 = n_1(\bar{\mathbf{y}}_{1..})'(\mathbf{C}\mathbf{S}_1\mathbf{C}')^{-1}(\bar{\mathbf{y}}_{1..}) = 554.749.$$

For each row of \mathbf{C} (linear, quadratic, etc.), we have

$$t_i^2 = n_1(\mathbf{c}_i' \bar{\mathbf{y}}_{1..})^2 / \mathbf{c}_i' \mathbf{S}_1 \mathbf{c}_i:$$

$$t_1^2 = 5.714, t_2^2 = 50.111, t_3^2 = 50.767, t_4^2 = 8.011, t_5^2 = .508.$$

- (b) For the obese group, we obtain $T^2 = n_2(\mathbf{C}\bar{\mathbf{y}}_{2..})'(\mathbf{C}\mathbf{S}_2\mathbf{C}')^{-1}(\mathbf{C}\bar{\mathbf{y}}_{2..}) = 128.552$. For the five rows of \mathbf{C} , we obtain $t_1^2 = 4.978, t_2^2 = 107.129, t_3^2 = 5.225, t_4^2 = 10.750, t_5^2 = 3.572$.
- (c) For the combined groups (\mathbf{S}_{pl} = pooled covariance matrix), we use $T^2 = N(\bar{\mathbf{y}}_{..})'(\mathbf{C}\mathbf{S}_{pl}\mathbf{C}')^{-1}(\bar{\mathbf{y}}_{..})$ in (6.121) to obtain $T^2 = 247.0079$. We test for linear, quadratic, etc., trends using the rows of \mathbf{C} in $t_i^2 = N(\mathbf{c}_i'\bar{\mathbf{y}}_{..})^2/\mathbf{c}_i'\mathbf{S}\mathbf{c}_i$; $t_1^2 = 1.162, t_2^2 = 155.017, t_3^2 = 30.540, t_4^2 = 1.319, t_5^2 = .506$. To compare groups, we use $\Lambda = |\mathbf{C}\mathbf{E}\mathbf{C}'|/|\mathbf{C}(\mathbf{E} + \mathbf{H})\mathbf{C}'|$ in (6.123) and $\Lambda_i = \mathbf{c}_i'\mathbf{E}\mathbf{c}_i/\mathbf{c}_i'(\mathbf{E} + \mathbf{H})\mathbf{c}_i$: $\Lambda = .4902, \Lambda_1 = .7947, \Lambda_2 = .9940, \Lambda_3 = .7987, \Lambda_4 = .6228, \Lambda_5 = .9172$.

6.44 Control group: By (6.114), $\hat{\beta}'_1 = (3.129, .656, -.283, -.334, .192, .037, -0.20)$. By (6.115), $T^2 = .7633$. By (6.117), $\hat{\mu}'_1 = (\hat{\mu}_{11}, \hat{\mu}_{12}, \dots, \hat{\mu}_{18}) = (4.11, 3.29, 2.71, 2.71, 3.04, 3.39, 3.54, 3.95)$.

Obese group: $\hat{\beta}'_2 = (3.207, -.187, .463, .056, -.102, -.010, .010), T^2 = .3943, \hat{\mu}'_2 = (4.51, 4.12, 3.81, 3.48, 3.24, 3.37, 3.70, 4.02)$

Combined groups (pooled covariance matrix): $\hat{\beta}' = (3.15, .162, .183, -.115, .012, .010, -.002), T^2 = .0158, \hat{\mu}' = (4.36, 3.80, 3.36, 3.15, 3.13, 3.37, 3.63, 3.98)$

6.45 A = activator, T = time, C = group. In (6.100), (6.101), and (6.102), we use

$$\mathbf{A} = \begin{pmatrix} 2 & 2 & 2 & -1 & -1 & -1 & -1 & -1 & -1 \\ 0 & 0 & 0 & 1 & 1 & 1 & -1 & -1 & -1 \end{pmatrix}$$

$$\mathbf{T} = \begin{pmatrix} -1 & 0 & 1 & -1 & 0 & 1 & -1 & 0 & 1 \\ 1 & -2 & 1 & 1 & -2 & 1 & 1 & -2 & 1 \end{pmatrix}$$

$$\mathbf{G} = \begin{pmatrix} -2 & 0 & 2 & 1 & 0 & -1 & 1 & 0 & -1 \\ 2 & -4 & 2 & -1 & 2 & -1 & -1 & 2 & -1 \\ 0 & 0 & 0 & 1 & 0 & -1 & -1 & 0 & 1 \\ 0 & 0 & 0 & 1 & -2 & 1 & -1 & 2 & -1 \end{pmatrix}$$

$T_A^2 = 5072.579, T_T^2 = 268.185, T_{AT}^2 = 143.491$. The same within-sample factors and interaction can be tested with Wilks' Λ using (6.104) and the other three MANOVA tests:

Source	Λ	$V^{(s)}$	$U^{(s)}$	θ	Significant?
A	.003	.997	317.04	.997	Yes
T	.056	.944	16.76	.944	Yes
AT	.100	.900	8.97	.900	Yes

The interactions of the within factors with the between factor G are tested with Wilks' Λ (Section 6.9.5) and with the other three MANOVA tests:

Source	Λ	$V^{(s)}$	$U^{(s)}$	θ	Significant?
AC	.884	.116	.131	.116	No
TC	.889	.111	.125	.111	No
ATC	.795	.205	.258	.205	No

The between-subjects factor C is tested with an ANOVA F -test: $F = .47$, p -value = .504.

CHAPTER 7

7.1 If Σ_0 is substituted for \mathbf{S} in (7.1), we have

$$u = \nu[\ln |\Sigma_0| - \ln |\Sigma_0| + \text{tr}(\mathbf{I}) - p] = \nu[0 + p - p] = 0$$

$$\begin{aligned} \mathbf{7.2} \quad \ln |\Sigma_0| - \ln |\mathbf{S}| &= -\ln |\Sigma_0|^{-1} - \ln |\mathbf{S}| \\ &= -\ln |\Sigma_0^{-1}| - \ln |\mathbf{S}| \quad [\text{by (2.91)}] \\ &= -(\ln |\mathbf{S}| + \ln |\Sigma_0^{-1}|) \\ &= -\ln |\mathbf{S}\Sigma_0^{-1}| \quad [\text{by (2.89)}] \end{aligned}$$

$$\mathbf{7.3} \quad -\ln (\prod_{i=1}^p \lambda_i) + \sum_{i=1}^p \lambda_i = -\sum_{i=1}^p \ln \lambda_i + \sum_{i=1}^p \lambda_i$$

7.4 As noted in Section 7.1, the likelihood ratio in this case involves the ratio of the determinants of the sample covariance matrices under H_0 and H_1 . Under H_1 , which is essentially unrestricted, the maximum likelihood estimate of Σ (corrected for bias) is given by (4.12) as \mathbf{S} . Under H_0 it is assumed that each of the p y_i 's in \mathbf{y} has variance σ^2 and that all y_i 's are independent. Thus we estimate σ^2 (unbiasedly) in each of the p columns of the \mathbf{Y} matrix [see (3.17) and (3.23)] and pool the p estimates to obtain

$$\hat{\sigma}^2 = \sum_{i=1}^n \sum_{j=1}^p \frac{(y_{ij} - \bar{y}_j)^2}{(n-1)p}.$$

Show that by (3.22) and (3.23) this is equal to

$$\hat{\sigma}^2 = \sum_{j=1}^p \frac{s_{jj}}{p} = \frac{\text{tr}(\mathbf{S})}{p}.$$

Thus the likelihood ratio is

$$\text{LR} = \left(\frac{|\mathbf{S}|}{|\hat{\sigma}^2 \mathbf{I}|} \right)^{n/2} = \left(\frac{|\mathbf{S}|}{|\mathbf{I} \text{tr}(\mathbf{S})/p|} \right)^{n/2}.$$

Show that by (2.85) this becomes

$$\text{LR} = \left(\frac{|\mathbf{S}|}{(\text{tr}\mathbf{S}/p)^p} \right)^{n/2}.$$

7.5 If $\lambda_1 = \lambda_2 = \dots = \lambda_p = \lambda$, say, then by (7.5),

$$u = \frac{p^p \prod_{i=1}^p \lambda_i}{\left(\sum_{i=1}^p \lambda_i \right)^p} = \frac{p^p \lambda^p}{(p\lambda)^p} = 1.$$

$$\begin{aligned} \mathbf{7.6} \quad & [(1-\rho)\mathbf{I} + \rho\mathbf{J}] = \begin{pmatrix} 1-\rho & 0 & \dots & 0 \\ 0 & 1-\rho & \dots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & 1-\rho \end{pmatrix} + \begin{pmatrix} \rho & \rho & \dots & \rho \\ \rho & \rho & \dots & \rho \\ \vdots & \vdots & & \vdots \\ \rho & \rho & \dots & \rho \end{pmatrix} \\ &= \begin{pmatrix} 1 & \rho & \dots & \rho \\ \rho & 1 & \dots & \rho \\ \vdots & \vdots & & \vdots \\ \rho & \rho & \dots & 1 \end{pmatrix} \end{aligned}$$

7.7 (a) Substitute $\mathbf{J} = \mathbf{j}\mathbf{j}'$ and $\mathbf{x} = \mathbf{j}$ into $\mathbf{J}\mathbf{x} = \lambda\mathbf{x}$ to obtain $\mathbf{j}\mathbf{j}'\mathbf{j} = \lambda\mathbf{j}$ which gives $p\mathbf{j} = \lambda\mathbf{j}$.

$$\mathbf{7.7 (b)} \quad \mathbf{S}_0 = s^2[(1-r)\mathbf{I} + r\mathbf{J}] = s^2(1-r)\left(\mathbf{I} + \frac{r}{1-r}\mathbf{J}\right)$$

(c) By (2.85) and (2.108), we have

$$\begin{aligned} |\mathbf{S}_0| &= \left| s^2(1-r)\left(\mathbf{I} + \frac{r}{1-r}\mathbf{J}\right) \right| = (s^2)^p(1-r)^p \left| \mathbf{I} + \frac{r}{1-r}\mathbf{J} \right| \\ &= (s^2)^p(1-r)^p \prod_{i=1}^p (1+\lambda_i) = (s^2)^p(1-r)^p \left(1 + \frac{rp}{1-r} \right) \\ &= (s^2)^p(1-r)^{p-1}(1-r+rp) = (s^2)^p(1-r)^{p-1}[1+(p-1)r] \end{aligned}$$

$$\mathbf{7.8} \quad M = \frac{|\mathbf{S}_1|^{\nu_1/2} |\mathbf{S}_2|^{\nu_2/2} \dots |\mathbf{S}_k|^{\nu_k/2}}{|\mathbf{S}|^{\sum_i \nu_i/2}} = \frac{|\mathbf{S}_1|^{\nu_1/2} |\mathbf{S}_2|^{\nu_2/2} \dots |\mathbf{S}_k|^{\nu_k/2}}{|\mathbf{S}|^{\nu_1/2} |\mathbf{S}|^{\nu_2/2} \dots |\mathbf{S}|^{\nu_k/2}}$$

$$\mathbf{7.9 (a)} \quad M = .7015 \quad \mathbf{7.9 (b)} \quad M = .0797$$

$$\begin{aligned} \mathbf{7.10} \quad \Lambda &= \frac{|\mathbf{S}|}{|\mathbf{S}_{yy}| |\mathbf{S}_{xx}|} = \frac{|\mathbf{S}_{xx}| |\mathbf{S}_{yy} - \mathbf{S}_{yx} \mathbf{S}_{xx}^{-1} \mathbf{S}_{xy}|}{|\mathbf{S}_{yy}| |\mathbf{S}_{xx}|} \\ &= |\mathbf{S}_{yy}^{-1}| |\mathbf{S}_{yy} - \mathbf{S}_{yx} \mathbf{S}_{xx}^{-1} \mathbf{S}_{xy}| \quad [\text{by (2.91)}] \\ &= |\mathbf{S}_{yy}^{-1} (\mathbf{S}_{yy} - \mathbf{S}_{yx} \mathbf{S}_{xx}^{-1} \mathbf{S}_{xy})| \quad [\text{by (2.89)}] \\ &= |\mathbf{I} - \mathbf{S}_{yy}^{-1} \mathbf{S}_{yx} \mathbf{S}_{xx}^{-1} \mathbf{S}_{xy}| \end{aligned}$$

$$= \prod_{i=1}^s (1 - r_i^2) \quad [\text{by (2.108)}],$$

where the r_i^2 's are the nonzero eigenvalues of $\mathbf{S}_{yy}^{-1}\mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}$. It was shown in Section 2.11.2 that $1 - \lambda_i$ is an eigenvalue of $\mathbf{I} - \mathbf{A}$, where λ_i is an eigenvalue of \mathbf{A} .

7.11 When all $p_i = 1$, we have $k = p$, and the submatrices in the denominators of (7.33) and (7.34) reduce to $\mathbf{S}_{jj} = s_{jj}$, $j = 1, 2, \dots, p$, and $\mathbf{R}_{jj} = 1$, $j = 1, 2, \dots, p$.

7.12 When all $p_i = 1$, we have $k = p$ and

$$\begin{aligned} a_2 &= p^2 - \sum_{i=1}^p p_i^2 = p^2 - p, \quad a_3 = p^3 - p, \\ c &= 1 - \frac{1}{12f\nu}(2a_3 + 3a_2) \\ &= 1 - \frac{1}{6(p^2 - p)\nu}[2(p^3 - p) + 3(p^2 - p)] \\ &= 1 - \frac{1}{6(p-1)\nu}[2(p^2 - 1) + 3(p-1)] \\ &= 1 - \frac{1}{6(p-1)\nu}[2(p-1)(p+1) + 3(p-1)] \\ &= 1 - \frac{1}{6\nu}[2p + 5]. \end{aligned}$$

7.13 As noted below (7.6), the degrees of freedom for the χ^2 -approximation is the total number of parameters minus the number estimated under H_0 . The number of distinct parameters in Σ is $p + \binom{p}{2} = \frac{1}{2}p(p+1)$. The number of parameters estimated under H_0 is p . The difference is $\frac{1}{2}p(p+1) - p = \frac{1}{2}p(p-1)$.

7.14 By (7.1) and (7.2), $u = 11.094$ and $u' = 10.668$.

7.15 By (7.7), $u = .0000594$. By (7.9), $u' = 23.519$. For H_0 : $\mathbf{C}\Sigma\mathbf{C}' = \sigma^2\mathbf{I}$, $u = .471$ and $u' = 2.050$.

7.16 For H_0 : $\Sigma = \sigma^2\mathbf{I}$, $u = .00513$ and $u' = 131.922$. For H_0 : $\mathbf{C}\Sigma\mathbf{C}' = \sigma^2\mathbf{I}$, $u = .129$ and $u' = 36.278$.

7.17 For H_0 : $\Sigma = \sigma^2\mathbf{I}$, $u = .00471$ and $u' = 136.190$. For H_0 : $\mathbf{C}\Sigma\mathbf{C}' = \sigma^2\mathbf{I}$, $u = .747$ and $u' = 7.486$.

7.18 By (7.16), $u' = 6.3323$ with 13 degrees of freedom. The F -approximation is $F = .4802$ with 13 and 1147 degrees of freedom.

7.19 $u' = 21.488$, $F = 2.511$ with 8 and 217 degrees of freedom

7.20 $u' = 35.795$, $F = 4.466$ with 8 and 4905 degrees of freedom

7.21 $u = 8.7457$, $F = .8730$ with 10 and 6502 degrees of freedom

- 7.22** $|\mathbf{S}_1| = 2.620 \times 10^{14}$, $|\mathbf{S}_2| = 2.410 \times 10^{14}$, $|\mathbf{S}_{p1}| = 4.368 \times 10^{14}$, $u = 17.502$, $F = .829$
- 7.23** $\ln M = -85.965$, $u = 156.434$, $a_1 = 21$, $a_2 = 17,797$, $F = 7.4396$
- 7.24** $\ln M = -7.082$, $u = 10.565$, $a_1 = 10$, $a_2 = 1340$, $F = 1.046$
- 7.25** $\ln M = -8.6062$, $u = 14.222$, $a_1 = 20$, $a_2 = 3909$, $F = .707$
- 7.26** $\ln M = -28.917$, $u = 44.018$, $a_1 = 50$, $a_2 = 3238$, $F = .8625$
- 7.27** $\ln M = -142.435$, $u = 174.285$, $a_1 = 110$, $a_2 = 2084$, $F = 1.448$
- 7.28** $|\mathbf{S}| = 1,207,109.5$, $|\mathbf{S}_{yy}| = 2385.1$, $|\mathbf{S}_{xx}| = 1341.9$, $\Lambda = .3772$
- 7.29** $|\mathbf{S}| = 4.237 \times 10^{13}$, $|\mathbf{S}_{yy}| = 484,926.6$, $|\mathbf{S}_{xx}| = 131,406,938$, $\Lambda = .6650$
- 7.30** $|\mathbf{S}| = 9.676 \times 10^{-8}$, $|\mathbf{S}_{yy}| = .02097$, $|\mathbf{S}_{xx}| = 9.94 \times 10^{-6}$, $\Lambda = .4642$
- 7.31** $|\mathbf{S}| = 1.7148 \times 10^{16}$, $|\mathbf{S}_{11}| = 11,284.967$, $|\mathbf{S}_{22}| = 11,891.15$, $|\mathbf{S}_{33}| = 25,951.605$, $s_{44} = 22,227.158$, $s_{55} = 214.06$, $u = .00103$, $u' = 274.787$, $\nu = 46$
- 7.32** $|\mathbf{S}| = 459.96$, $s_{11} = 140.54$, $s_{22} = 72.25$, $s_{33} = .250$, $u = .1811$, $u' = 12.246$, $f = 3$
- 7.33** $u = .0001379$, $u' = 16.297$
- 7.34** $u = .0005176$, $u' = 127.367$
- 7.35** $u = .005071$, $u' = 131.226$

CHAPTER 8

8.1 Using $\mathbf{a} = \mathbf{S}_{p1}^{-1}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)$, we obtain

$$\begin{aligned}\frac{[\mathbf{a}'(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)]^2}{\mathbf{a}'\mathbf{S}_{p1}\mathbf{a}} &= \frac{[(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{p1}^{-1}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)]^2}{(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{p1}^{-1} \mathbf{S}_{p1} \mathbf{S}_{p1}^{-1} (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)} \\ &= \frac{[(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{p1}^{-1}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)]^2}{(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{p1}^{-1} (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)}.\end{aligned}$$

8.2 You may wish to use the following steps:

- (i) In Section 5.6.2 the grouping variable w is defined as $n_2/(n_1 + n_2)$ for each observation in group 1 and $-n_1/(n_1 + n_2)$ for group 2. Show that with this formulation, $\bar{w} = 0$.

(ii) Because $\bar{w} = 0$, there is no intercept and the fitted model becomes

$$\hat{w}_i = b_1(y_{i1} - \bar{y}_1) + b_2(y_{i2} - \bar{y}_2) + \cdots + b_p(y_{ip} - \bar{y}_p), \\ i = 1, 2, \dots, n_1 + n_2.$$

Denote the resulting matrix of y -values corrected for their means as \mathbf{Y}_c and the vector of w 's as \mathbf{w} . Then the least squares estimate $\mathbf{b} = (b_1, b_2, \dots, b_p)'$ is obtained as

$$\mathbf{b} = (\mathbf{Y}'_c \mathbf{Y}_c)^{-1} \mathbf{Y}'_c \mathbf{w}.$$

Using (2.51), show that

$$\begin{aligned} \mathbf{Y}'_c \mathbf{Y}_c &= \sum_{i=1}^2 \sum_{j=1}^{n_i} (\mathbf{y}_{ij} - \bar{\mathbf{y}})(\mathbf{y}_{ij} - \bar{\mathbf{y}})' \\ &= \sum_{i=1}^2 \sum_{j=1}^{n_i} (\mathbf{y}_{ij} - \bar{\mathbf{y}}_i)(\mathbf{y}_{ij} - \bar{\mathbf{y}}_i)' + \frac{n_1 n_2}{n_1 + n_2} (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)', \end{aligned}$$

where $\bar{\mathbf{y}} = (n_1 \bar{\mathbf{y}}_1 + n_2 \bar{\mathbf{y}}_2)/(n_1 + n_2)$. It will be helpful to write the first sum above as

$$\sum_{j=1}^{n_1} (\mathbf{y}_{1j} - \bar{\mathbf{y}})(\mathbf{y}_{1j} - \bar{\mathbf{y}})' + \sum_{j=1}^{n_2} (\mathbf{y}_{2j} - \bar{\mathbf{y}})(\mathbf{y}_{2j} - \bar{\mathbf{y}})'$$

and add and subtract $\bar{\mathbf{y}}_1$ in the first term and $\bar{\mathbf{y}}_2$ in the second.

(iii) Show that

$$\mathbf{Y}'_c \mathbf{w} = \sum_{i=1}^2 \sum_{j=1}^{n_i} (\mathbf{y}_{ij} - \bar{\mathbf{y}}) w_{ij} = \frac{n_1 n_2}{n_1 + n_2} (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2).$$

Again it will be helpful to sum separately over the two groups.

(iv) From (ii) and (iii) we have

$$\mathbf{b} = (\nu \mathbf{S} + k \bar{\mathbf{d}} \bar{\mathbf{d}}')^{-1} k \bar{\mathbf{d}},$$

where $\mathbf{S} = \sum_{ij} (\mathbf{y}_{ij} - \bar{\mathbf{y}}_i)(\mathbf{y}_{ij} - \bar{\mathbf{y}}_i)' / (n_1 + n_2 - 2)$, $\nu = n_1 + n_2 - 2$, $k = n_1 n_2 / (n_1 + n_2)$, and $\bar{\mathbf{d}} = \bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2$. Use (2.77) for the inverse of a patterned matrix of the type $\nu \mathbf{S} + k \bar{\mathbf{d}} \bar{\mathbf{d}}'$ to obtain (8.4).

8.3 You may want to use the following steps:

(i) R^2 is defined as [see (10.30)]

$$R^2 = \frac{\mathbf{b}' \mathbf{Y}'_c \mathbf{w} - n \bar{w}^2}{\mathbf{w}' \mathbf{w} - n \bar{w}^2}.$$

In this case the expression simplifies because $\bar{w} = 0$. Using $\mathbf{Y}'_c \mathbf{w}$ in Problem 8.2(iii) above, show that $R^2 = \mathbf{b}'(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)$.

- (ii) Show that

$$\mathbf{b}'(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2) = \frac{T^2}{n_1 + n_2 - 2 + T^2}.$$

8.4 $[\mathbf{a}'(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)]^2 = \mathbf{a}'(\bar{\mathbf{y}} - \bar{\mathbf{y}}_2)\mathbf{a}'(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2) = \mathbf{a}'(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{a}$

8.5 $\mathbf{H}\mathbf{a} - \lambda \mathbf{E}\mathbf{a} = \mathbf{0}$

$$\mathbf{E}^{-1}(\mathbf{H}\mathbf{a} - \lambda \mathbf{E}\mathbf{a}) = \mathbf{E}^{-1}\mathbf{0}$$

$$\mathbf{E}^{-1}\mathbf{H}\mathbf{a} - \lambda \mathbf{E}^{-1}\mathbf{E}\mathbf{a} = \mathbf{0}$$

$$(\mathbf{E}^{-1}\mathbf{H} - \lambda \mathbf{I})\mathbf{a} = \mathbf{0}$$

8.6 Substituting $a_r^* = s_r a_r$, $r = 1, 2, \dots, p$, into (8.15), we obtain

$$\begin{aligned} z_{1i} &= s_1 a_1 \frac{y_{1i1} - \bar{y}_{11}}{s_1} + s_2 a_2 \frac{y_{1i2} - \bar{y}_{12}}{s_2} + \cdots + s_p a_p \frac{y_{1ip} - \bar{y}_{1p}}{s_p} \\ &= a_1 y_{1i1} + a_2 y_{1i2} + \cdots + a_p y_{1ip} - a_1 \bar{y}_{11} - a_2 \bar{y}_{12} - \cdots - a_p \bar{y}_{1p} \\ &= a_1 y_{1i1} + a_2 y_{1i2} + \cdots + a_p y_{1ip} - \mathbf{a}' \bar{\mathbf{y}}_1 \end{aligned}$$

8.7 (a) $\mathbf{a}^* = (1.366, -.810, 2.525, -1.463)$

(b) $t_1 = 5.417, t_2 = 2.007, t_3 = 7.775, t_4 = .688$

(c) The standardized coefficients rank the variables in the order y_3, y_4, y_1, y_2 .
The t -tests rank them in the order y_3, y_1, y_2, y_4 .

(d) The partial F 's calculated by (8.26) are $F(y_1|y_2, y_3, y_4) = 7.844$,
 $F(y_2|y_1, y_3, y_4) = 2.612$, $F(y_3|y_1, y_2, y_4) = 40.513$, and
 $F(y_4|y_1, y_2, y_3) = 9.938$.

8.8 (a) $\mathbf{a}' = (.345, -.130, -.106, -.143)$

(b) $\mathbf{a}^* = (4.137, -2.501, -1.158, -2.068)$

(c) $t_1 = 3.888, t_2 = -3.865, t_3 = -5.691, t_4 = -5.043$

(e) $F(y_1|y_2, y_3, y_4) = 35.934, F(y_2|y_1, y_3, y_4) = 5.799$
 $F(y_3|y_1, y_2, y_4) = 1.775, F(y_4|y_1, y_2, y_3) = 8.259$

8.9 (a) $\mathbf{a}' = (-.145, .052, -.005, -.089, -.007, -.022)$

(b) $\mathbf{a}^* = (-1.016, .147, -.542, -1.035, -.107, -1.200)$

(c) $t_1 = -4.655, t_2 = .592, t_3 = -4.354, t_4 = -5.257, t_5 = -4.032, t_6 = -6.439$

(e) $F(y_1|y_2, y_3, y_4, y_5, y_6) = 8.081, F(y_2|y_1, y_3, y_4, y_5, y_6) = .150$,
 $F(y_3|y_1, y_2, y_4, y_5, y_6) = .835, F(y_4|y_1, y_2, y_3, y_5, y_6) = 8.503$,
 $F(y_5|y_1, y_2, y_3, y_4, y_6) = .028, F(y_6|y_1, y_2, y_3, y_4, y_5) = 9.192$

8.10 (a) $\mathbf{a}' = (.057, .010, .242, .071)$

(b) $\mathbf{a}^* = (1.390, .083, 1.025, .032)$

(c) $t_1 = -3.713, t_2 = .549, t_3 = -3.262, t_4 = -.724$

(e) $F(y_1|y_2, y_3, y_4) = 3.332, F(y_2|y_1, y_3, y_4) = .010$
 $F(y_3|y_1, y_2, y_4) = 1.482, F(y_4|y_1, y_2, y_3) = .001$

8.11 (a) $\mathbf{a}_1' = (.021, .533, -.347, -.135), \mathbf{a}_2' = (-.317, .298, .243, -.026)$

(b) $\lambda_1/(\lambda_1 + \lambda_2) = .958, \lambda_2/(\lambda_1 + \lambda_2) = .042$. Using the methods of Section 8.6.2, we have two tests, the first for significance of λ_1 and λ_2 and the second for significance of λ_2 :

Test	Λ	F	p -Value for F
1	.2245	8.3294	<.0001
2	.8871	1.3157	.2869

(c) $\mathbf{a}_1^* = (.076, 1.553, -1.182, -.439), \mathbf{a}_2^* = (-1.162, .869, .828, -.085)$

(d) $F(y_1|y_2, y_3, y_4) = 1.067, F(y_2|y_1, y_3, y_4) = 20.975$
 $F(y_3|y_1, y_2, y_4) = 9.630, F(y_4|y_1, y_2, y_3) = 1.228$

(e) In the plot, the first discriminant function separates groups 1 and 2 from group 3, but the second is ineffective in separating group 1 from group 2.

8.12 (a) $\lambda_i \quad \lambda_i / \sum_{j=1}^4 \lambda_j \quad \text{Eigenvector}$

1.8757	.6421	$\mathbf{a}_1' = (.470, -.263, .653, -.074)$
.7907	.2707	$\mathbf{a}_2' = (.176, .188, -1.058, 1.778)$
.2290	.0784	$\mathbf{a}_3' = (-.155, .258, .470, -.850)$
.0260	.0089	$\mathbf{a}_4' = (-3.614, .475, .310, -.479)$

(b) Test of significance of each eigenvalue and those that follow it:

Test	Λ	Approximate F	p -Value for F
------	-----------	-----------------	--------------------

1	.1540	4.937	<.0001
2	.4429	3.188	.0006
3	.7931	1.680	.1363
4	.9747	.545	.5839

(c) $\mathbf{a}_1^* = (.266, -.915, 1.353, -.097), \mathbf{a}_2^* = (.100, .654, -2.291, 2.333),$
 $\mathbf{a}_3^* = (-.087, .899, .973, -1.115), \mathbf{a}_4^* = (-2.044, 1.654, .643, -.628)$

(d) $F(y_1|y_2, y_3, y_4) = .299, F(y_2|y_1, y_3, y_4) = 1.931$
 $F(y_3|y_1, y_2, y_4) = 6.085, F(y_4|y_1, y_2, y_3) = 4.659$

(e) In the plot, the first discriminant function separates groups 1, 4, and 6 from groups 2, 3, and 5. The second function achieves some separation of group 6 from groups 1 and 4 and some separation of group 3 from groups 2 and 5.

8.13 Three variables entered the model in the stepwise selection. The summary table is as follows:

Step	Variable Entered	Overall Λ	p-Value	Partial Λ	Partial F	p-Value
1	y_4	.4086	<.0001	.4086	12.158	<.0001
2	y_3	.2655	<.0001	.6499	4.418	.0026
3	y_2	.1599	<.0001	6022	5.284	.0008

8.14 Summary table:

Step	Variable Entered	Overall Λ	p-Value	Partial Λ	Partial F	p-Value
1	y_4	.6392	<.0001	.6392	21.451	<.0001
2	y_3	.5430	<.0001	.8495	6.554	.0147
3	y_6	.4594	<.0001	.8461	6.549	.0148
4	y_2	.4063	<.0001	.8843	4.578	.0394
5	y_5	.3639	<.0001	.8957	3.959	.0547

In this case, the fifth variable to enter, y_5 , would not ordinarily be included in the subset. The p -value of .0547 is large in this setting, where several tests are run at each step and the variable with smallest p -value is selected.

8.15 Summary table:

Step	Variable Entered	Overall Λ	p-Value	Partial Λ	Partial F	p-Value
1	y_2	.6347	.0006	.6347	9.495	.0006
2	y_3	.2606	<.0001	.4106	22.975	<.0001

CHAPTER 9

$$\mathbf{9.1} \quad \bar{z}_1 - \bar{z}_2 = \mathbf{a}'\bar{\mathbf{y}}_1 - \mathbf{a}'\bar{\mathbf{y}}_2 = \mathbf{a}'(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2) = (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{pl}^{-1} (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)$$

$$\mathbf{9.2} \quad \frac{1}{2}(\bar{z}_1 + \bar{z}_2) = \frac{1}{2}(\mathbf{a}'\bar{\mathbf{y}}_1 + \mathbf{a}'\bar{\mathbf{y}}_2) = \frac{1}{2}\mathbf{a}'(\bar{\mathbf{y}}_1 + \bar{\mathbf{y}}_2) \\ = \frac{1}{2}(\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{pl}^{-1} (\bar{\mathbf{y}}_1 + \bar{\mathbf{y}}_2)$$

9.3 Write (9.8) in the form

$$\frac{f(\mathbf{y}|G_1)}{f(\mathbf{y}|G_2)} > \frac{p_2}{p_1}$$

and substitute $f(\mathbf{y}|G_i) = N_p(\boldsymbol{\mu}_i, \boldsymbol{\Sigma})$ from (4.2) to obtain

$$\frac{f(\mathbf{y}|G_1)}{f(\mathbf{y}|G_2)} = e^{(\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2)' \boldsymbol{\Sigma}^{-1} \mathbf{y} - (\boldsymbol{\mu}_1 - \boldsymbol{\mu}_2)' \boldsymbol{\Sigma}^{-1} (\boldsymbol{\mu}_1 + \boldsymbol{\mu}_2)/2} > \frac{p_2}{p_1}.$$

Substitute estimates for $\boldsymbol{\mu}_1$, $\boldsymbol{\mu}_2$, and $\boldsymbol{\Sigma}$, and take the logarithm of both sides to obtain (9.9). Note that if $a > b$, then $\ln a > \ln b$.

9.4 Maximizing $p_i f(\mathbf{y}, G_i)$ is equivalent to maximizing $\ln[p_i f(\mathbf{y}|G_i)]$. Use $f(\mathbf{y}|G_i) = N_p(\boldsymbol{\mu}_i, \boldsymbol{\Sigma})$ from (4.2) and take the logarithm to obtain

$$\ln[p_i f(\mathbf{y}|G_i)] = \ln p_i - \frac{1}{2}p \ln(2\pi) - \frac{1}{2}|\boldsymbol{\Sigma}| - \frac{1}{2}(\mathbf{y} - \boldsymbol{\mu}_i)' \boldsymbol{\Sigma}^{-1} (\mathbf{y} - \boldsymbol{\mu}_i).$$

Expand the last term, delete terms common to all groups (terms that do not involve i), and substitute estimators of μ_i and Σ to obtain (9.13).

- 9.5** Use $f(\mathbf{y}|G_i) = N_p(\boldsymbol{\mu}_i, \boldsymbol{\Sigma}_i)$ in $\ln[p_i f(\mathbf{y}|G_i)]$, delete $-(p/2) \ln(2\pi)$, and substitute $\bar{\mathbf{y}}_i$ and \mathbf{S}_i for $\boldsymbol{\mu}_i$ and $\boldsymbol{\Sigma}_i$.

- 9.6** Substitute the constant c for each occurrence of p_i/n_i to obtain the result.

- 9.7 (a)** $\mathbf{a}' = (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{pl}^{-1} = (.345, -.130, -.106, -.143)$,
 $\frac{1}{2}(\bar{z}_1 + \bar{z}_2) = -15.8054$

Actual Group	Number of Observations	Predicted Group	
		1	2
1	19	19	0
2	20	1	19

$$\text{Error rate} = \frac{1}{39} = .0256$$

- (c)** Using the k nearest neighbor method with $k = 5$, we obtain the same classification table as in part (b). With $k = 4$, two observations are misclassified, and the error rate becomes $2/39 = .0513$.

- 9.8 (a)** $\mathbf{a}' = (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{pl}^{-1} = (-.145, .052, -.005, -.089, -.007, -.022)$,
 $\frac{1}{2}(\bar{z}_1 + \bar{z}_2) = -17.045$

Actual Group	Number of Observations	Predicted Group	
		1	2
1	39	37	20
2	34	8	26

$$\text{Error rate} = (2 + 8)/73 = .1370$$

- (c)** p_1 and p_2 Proportional to Sample Sizes

Actual Group	Number of Observations	Predicted Group	
		1	2
1	39	37	2
2	34	8	26

$$\text{Error rate} = (2 + 8)/73 = .1370$$

- 9.9 (a)** $\mathbf{a}' = (\bar{\mathbf{y}}_1 - \bar{\mathbf{y}}_2)' \mathbf{S}_{pl}^{-1} = (-.057, -.010, -.242, -.071)$,
 $\frac{1}{2}(\bar{z}_1 + \bar{z}_2) = -7.9686$

(b)

Linear Classification

Actual Group	Number of Observations	Predicted Group	
		1	2
1	9	8	1
2	10	1	9

$$\text{Error rate} = \frac{2}{19} = .1053$$

(c)

Holdout Method

Actual Group	Number of Observations	Predicted Group	
		1	2
1	9	6	3
2	10	3	7

$$\text{Error rate} = (3 + 3)/19 = .3158$$

(d)

Kernel Density Estimator with $h = 2$

Actual Group	Number of Observations	Predicted Group	
		1	2
1	9	9	0
2	10	1	9

$$\text{Error rate} = \frac{1}{19} = .0526$$

9.10 (a)

Actual Group	Number of Observations	Predicted Group	
		1	2
1	20	18	2
2	20	2	18

$$\text{Error rate} = (2 + 2)/40 = .100.$$

- (b) Four variables were selected by the stepwise discriminant analysis: y_2 , y_3 , y_4 , and y_6 (see Problem 8.14). With these four variables we obtain the classification table in part (c).

(c)

Actual Group	Number of Observations	Predicted Group	
		1	2
1	20	18	2
2	20	2	18

Error rate = $(2 + 2)/40 = .100$. The four variables classified the sample as well as did all six variables in part (a).

- 9.11 (a)** By (9.11), $L_i(\mathbf{y}) = \bar{\mathbf{y}}_i' \mathbf{S}_{\text{pl}}^{-1} \mathbf{y} - \frac{1}{2} \bar{\mathbf{y}}_i' \mathbf{S}_{\text{pl}}^{-1} \bar{\mathbf{y}}_i = \mathbf{c}_i' \mathbf{y} + c_{0i}$. The vectors $\begin{pmatrix} c_{0i} \\ \mathbf{c}_i \end{pmatrix}, i = 1, 2, 3$, are

Group 1	Group 2	Group 3
-72.77	-65.18	-68.57
.81	2.12	.68
15.15	10.11	2.79
-1.03	-.24	6.54
10.02	11.06	13.09

- (b)** Linear Classification

Actual Group	Number of Observations	Predicted Group		
		1	2	3
1	12	9	3	0
2	12	3	7	2
3	12	0	1	11

$$\text{Error rate} = (3 + 3 + 2 + 1)/36 = .250$$

- (c)** Quadratic Classification

Actual Group	Number of Observations	Predicted Group		
		1	2	3
1	12	10	2	0
2	12	2	8	2
3	12	0	1	11

$$\text{Error rate} = (2 + 2 + 2 + 1)/36 = .194$$

- (d)** Linear Classification–Holdout Method

Actual Group	Number of Observations	Predicted Group		
		1	2	3
1	12	7	5	0
2	12	4	5	3
3	12	0	1	11

$$\text{Error rate} = (5 + 4 + 3 + 1)/12 = .361$$

(e) k Nearest Neighbor with $k = 5$

Actual Group	Number of Observations	Predicted Group		
		1	2	3
1	11	9	2	0
2	11	2	7	2
3	12	0	1	11

$$\text{Error rate} = (2 + 2 + 2 + 1)/34 = .206$$

- 9.12 (a)** By (9.11), $L_i(\mathbf{y}) = \bar{\mathbf{y}}'_i \mathbf{S}_{\text{pl}}^{-1} \mathbf{y} - \frac{1}{2} \bar{\mathbf{y}}'_i \mathbf{S}_{\text{pl}}^{-1} \bar{\mathbf{y}}_i = \mathbf{c}'_i \mathbf{y} + c_{0i}$. The vectors $\begin{pmatrix} c_{0i} \\ \mathbf{c}_i \end{pmatrix}, i = 1, 2, \dots, 6$, are

Group 1	Group 2	Group 3	Group 4	Group 5	Group 6
-300.0	-353.2	-328.5	-291.8	-347.5	-315.8
314.6	317.1	324.6	307.3	316.8	311.3
-59.4	-64.0	-65.2	-59.4	-65.8	-63.1
149.6	168.2	154.9	147.7	168.2	160.6
-161.2	-172.6	-150.4	-153.4	-172.9	-175.5

(b) Linear Classification

Actual Group	Number of Observations	Predicted Group					
		1	2	3	4	5	6
1	8	5	0	0	1	0	2
2	8	0	3	2	1	2	0
3	8	0	0	6	1	1	0
4	8	3	0	1	4	0	0
5	8	0	3	1	0	3	1
6	8	2	0	0	0	2	4

$$\text{Correct classification rate} = (5 + 3 + 6 + 4 + 3 + 4)/48 = .521$$

$$\text{Error rate} = 1 - .521 = .479$$

(c) Quadratic Classification

Actual Group	Number of Observations	Predicted Group					
		1	2	3	4	5	6
1	8	8	0	0	0	0	0
2	8	0	7	0	1	0	0
3	8	1	0	6	0	1	0
4	8	0	0	1	7	0	0
5	8	0	3	0	0	4	1
6	8	2	0	0	0	1	5

$$\text{Correct classification rate} = (8 + 7 + 6 + 7 + 4 + 5)/48 = .771$$

$$\text{Error rate} = 1 - .771 = .229$$

(d)

k Nearest Neighbor with $k = 3$

Actual Group	Number of Observations	Predicted Group						
		1	2	3	4	5	6	Ties
1	8	5	0	0	2	0	0	1
2	8	0	4	0	0	1	0	3
3	8	1	0	6	0	1	0	0
4	8	0	0	0	5	0	0	3
5	8	0	1	0	0	6	1	0
6	8	2	0	0	0	0	5	1

Correct classification rate = $(5 + 4 + 6 + 5 + 6 + 5)/40 = .775$ Error rate = $1 - .775 = .225$

(e)

Normal Kernel with $h = 1$ (For this data set, larger values of h do much worse.)

Actual Group	Number of Observations	Predicted Group					
		1	2	3	4	5	6
1	8	8	0	0	0	0	0
2	8	0	8	0	0	0	0
3	8	1	0	6	0	1	0
4	8	1	0	0	7	0	0
5	8	0	0	0	0	7	1
6	8	2	0	0	0	0	6

Correct classification rate = $(8 + 8 + 6 + 7 + 7 + 6)/48 = .875$ Error rate = $1 - .875 = .125$

CHAPTER 10

$$10.1 \quad \mathbf{y} - \mathbf{X}\hat{\beta} = \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{pmatrix} - \begin{pmatrix} \mathbf{x}'_1 \\ \mathbf{x}'_2 \\ \vdots \\ \mathbf{x}'_n \end{pmatrix} \hat{\beta} = \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{pmatrix} - \begin{pmatrix} \mathbf{x}'_1 \hat{\beta} \\ \mathbf{x}'_2 \hat{\beta} \\ \vdots \\ \mathbf{x}'_n \hat{\beta} \end{pmatrix} = \begin{pmatrix} y_1 - \mathbf{x}'_1 \hat{\beta} \\ y_2 - \mathbf{x}'_2 \hat{\beta} \\ \vdots \\ y_n - \mathbf{x}'_n \hat{\beta} \end{pmatrix}$$

By (2.33), $\sum_{i=1}^n (y_i - \mathbf{x}'_i \hat{\beta})^2 = (\mathbf{y} - \mathbf{X}\hat{\beta})'(\mathbf{y} - \mathbf{X}\hat{\beta})$.

$$\begin{aligned} 10.2 \quad \sum_{i=1}^n (y_i - \mu)^2 &= \sum_i (y_i - \bar{y} + \bar{y} - \mu)^2 \\ &= \sum_i (y_i - \bar{y})^2 + 2 \sum_i (y_i - \bar{y})(\bar{y} - \mu) + \sum_i (\bar{y} - \mu)^2 \\ &= \sum_i (y_i - \bar{y})^2 + (\bar{y} - \mu) \sum_i (y_i - \bar{y}) + n(\bar{y} - \mu)^2 \\ &= \sum_i (y_i - \bar{y})^2 + n(\bar{y} - \mu)^2 \quad [\text{since } \sum_i (y_i - \bar{y}) = 0] \end{aligned}$$

$$10.3 \quad \sum_{i=1}^n (x_{i2} - \bar{x}_2)\bar{y} = \bar{y} \sum_i (x_{i2} - \bar{x}_2) = \bar{y}(\sum_i x_{i2} - n\bar{x}_2) = \bar{y}(n\bar{x}_2 - n\bar{x}_2)$$

$$\begin{aligned} \mathbf{10.4} \quad E[\hat{y}_i - E(y_i)]^2 &= E[\hat{y}_i - E(\hat{y}_i) + E(\hat{y}_i) - E(y_i)]^2 \\ &= E[\hat{y}_i - E(\hat{y}_i)]^2 + 2E[\hat{y}_i - E(\hat{y}_i)][E(\hat{y}_i) - E(y_i)] \\ &\quad + E[E(\hat{y}_i) - E(y_i)]^2 \end{aligned}$$

The second term on the right vanishes because $[E(\hat{y}_i) - E(y_i)]$ is constant and $E[\hat{y}_i - E(\hat{y}_i)] = E(\hat{y}_i) - E(\hat{y}_i) = 0$. For the third term, we have $E[E(\hat{y}_i) - E(y_i)]^2 = [E(\hat{y}_i) - E(y_i)]^2$, because $[E(\hat{y}_i) - E(y_i)]^2$ is constant.

10.5 First show that $\text{cov}(\hat{\beta}_p) = \sigma^2(\mathbf{X}'_p \mathbf{X}_p)^{-1}$. This can be done by noting that $\hat{\beta}_p = (\mathbf{X}'_p \mathbf{X}_p)^{-1} \mathbf{X}'_p \mathbf{y} = \mathbf{A}\mathbf{y}$, say. Then, by (3.74), $\text{cov}(\mathbf{A}\mathbf{y}) = \mathbf{A}\text{cov}(\mathbf{y})\mathbf{A}' = \mathbf{A}(\sigma^2 \mathbf{I})\mathbf{A}' = \sigma^2 \mathbf{A}\mathbf{A}'$. By substituting $\mathbf{A} = (\mathbf{X}'_p \mathbf{X}_p)^{-1} \mathbf{X}'_p$, this becomes $\text{cov}(\hat{\beta}_p) = \sigma^2(\mathbf{X}'_p \mathbf{X}_p)^{-1}$. Then by (3.70), $\text{var}(\mathbf{x}'_{pi} \hat{\beta}_p) = \mathbf{x}'_{pi} \text{cov}(\hat{\beta}_p) \mathbf{x}_{pi}$ and the remaining steps follow as indicated.

10.6 By (10.36), $s_p^2 = \text{SSE}_p/(n-p)$. Then by (10.44),

$$\begin{aligned} C_p &= p + (n-p) \frac{s_p^2 - s_k^2}{s_k^2} = p + (n-p) \left(\frac{s_p^2}{s_k^2} - 1 \right) \\ &= p + (n-p) \frac{s_p^2}{s_k^2} - (n-p) = (n-p) \frac{\text{SSE}_p/s_k^2}{n-p} - n + 2p \\ &= \frac{\text{SSE}_p}{s_k^2} - (n-2p). \end{aligned}$$

10.7 $(\mathbf{Y} - \mathbf{X}\hat{\mathbf{B}})'(\mathbf{Y} - \mathbf{X}\hat{\mathbf{B}}) = \mathbf{Y}'\mathbf{Y} - \mathbf{Y}'\mathbf{X}\hat{\mathbf{B}} - \hat{\mathbf{B}}'\mathbf{X}'\mathbf{Y} - \hat{\mathbf{B}}'\mathbf{X}'\mathbf{X}\hat{\mathbf{B}}$. Transpose $\hat{\mathbf{B}} = (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\mathbf{Y}$ from (10.48) and substitute into $\hat{\mathbf{B}}'\mathbf{X}'\mathbf{X}\hat{\mathbf{B}}$.

$$\begin{aligned} \mathbf{10.8} \quad E[\hat{\mathbf{y}}_i - E(\mathbf{y}_i)][\hat{\mathbf{y}}_i - E(\mathbf{y}_i)]' &= E[\hat{\mathbf{y}}_i - E(\hat{\mathbf{y}}_i) + E(\hat{\mathbf{y}}_i) - E(\mathbf{y}_i)][\hat{\mathbf{y}}_i - E(\hat{\mathbf{y}}_i)] \\ &\quad + E(\hat{\mathbf{y}}_i) - E(\mathbf{y}_i)]' \\ &= E[\hat{\mathbf{y}}_i - E(\hat{\mathbf{y}}_i)][\hat{\mathbf{y}}_i - E(\hat{\mathbf{y}}_i)]' \\ &\quad + E[\hat{\mathbf{y}}_i - E(\hat{\mathbf{y}}_i)][E(\hat{\mathbf{y}}_i) - E(\mathbf{y}_i)]' \\ &\quad + E[E(\hat{\mathbf{y}}_i) - E(\mathbf{y}_i)][\hat{\mathbf{y}}_i - E(\hat{\mathbf{y}}_i)]' \\ &\quad + E[E(\hat{\mathbf{y}}_i) - E(\mathbf{y}_i)][E(\hat{\mathbf{y}}_i) - E(\mathbf{y}_i)]' \end{aligned}$$

The second and third terms are equal to \mathbf{O} because $[E(\hat{\mathbf{y}}_i) - E(\mathbf{y}_i)]$ is a constant vector and $E[\hat{\mathbf{y}}_i - E(\hat{\mathbf{y}}_i)] = E(\hat{\mathbf{y}}_i) - E(\hat{\mathbf{y}}_i) = \mathbf{0}$. The fourth term is a constant matrix and the first E can be deleted.

10.9 As in Problem 10.5, we have $\text{cov}(\hat{\beta}_{p(j)}) = \sigma_{jj}(\mathbf{X}'_p \mathbf{X}_p)^{-1}$, where $\sigma_{jj} = \text{var}(y_j)$ is the j th diagonal element of $\Sigma = \text{cov}(\mathbf{y})$. Similarly, $\text{cov}(\hat{\beta}_{p(k)}) = \sigma_{kk}(\mathbf{X}'_p \mathbf{X}_p)^{-1}$, where $\sigma_{jk} = \text{cov}(y_j, y_k)$ is the (jk) th element of Σ . The notation $\text{cov}(\hat{\beta}_{p(j)}, \hat{\beta}_{p(k)})$ indicates a matrix containing the covariance of each element of $\hat{\beta}_{p(j)}$ and each element of $\hat{\beta}_{p(k)}$. Now for the covariance matrix, $\text{cov}(\hat{\mathbf{y}}'_i) = \text{cov}(\mathbf{x}'_{pi} \hat{\beta}_{p(1)}, \dots, \mathbf{x}'_{pi} \hat{\beta}_{p(m)})$, we need the variance of each of the m random variables and the covariance of each pair. By Problem 10.5

and (3.70), $\text{var}(\mathbf{x}'_{pi}\hat{\beta}_{p(1)}) = \mathbf{x}'_{pi}\text{cov}(\hat{\beta}_{p(1)})\mathbf{x}_{pi} = \sigma_{11}\mathbf{x}'_{pi}(\mathbf{X}'_p\mathbf{X}_p)^{-1}\mathbf{x}_{pi}$. Similarly, $\text{cov}(\mathbf{x}'_{pi}\hat{\beta}_{p(1)}, \mathbf{x}'_{pi}\hat{\beta}_{p(2)}) = \sigma_{12}\mathbf{x}'_{pi}(\mathbf{X}'_p\mathbf{X}_p)^{-1}\mathbf{x}_{pi}$. The other variances and covariances can be obtained in an analogous manner.

- 10.10** By (10.92), $\mathbf{S}_p = \mathbf{E}_p/(n - p)$. Then by (10.98),

$$\begin{aligned}\mathbf{C}_p &= p\mathbf{I} + (n - p)\mathbf{S}_k^{-1}(\mathbf{S}_p - \mathbf{S}_k) \\ &= p\mathbf{I} + (n - p)\mathbf{S}_k^{-1}\frac{\mathbf{E}_p}{n - p} - (n - p)\mathbf{I} \\ &= \mathbf{S}_k^{-1}\mathbf{E}_p + (2p - n)\mathbf{I}.\end{aligned}$$

- 10.11** $|\mathbf{E}_k^{-1}\mathbf{E}_p| = |\mathbf{E}_k^{-1}| |\mathbf{E}_p| > 0$, because both \mathbf{E}_k^{-1} and \mathbf{E}_p are positive definite.

- 10.12** By (10.99), $\mathbf{C}_p = \mathbf{S}_k^{-1}\mathbf{E}_p + (2p - n)\mathbf{I}$. Using $\mathbf{S}_k = \mathbf{E}_k/(n - k)$, we obtain

$$\begin{aligned}\left(\frac{\mathbf{E}_k}{n - k}\right)^{-1}\mathbf{E}_p &= \mathbf{C}_p - (2p - n)\mathbf{I} \\ (n - k)\mathbf{E}_k^{-1}\mathbf{E}_p &= \mathbf{C}_p + (n - 2p)\mathbf{I}.\end{aligned}$$

- 10.13** If \mathbf{C}_p is replaced by $p\mathbf{I}$ in (10.101), we obtain

$$\mathbf{E}_k^{-1}\mathbf{E}_p = \frac{\mathbf{C}_p + (n - 2p)\mathbf{I}}{n - k} = \frac{p\mathbf{I} + n\mathbf{I} - 2p\mathbf{I}}{n - k} = \frac{(n - p)\mathbf{I}}{n - k}.$$

- 10.14** (a) For this scenario, the Zyskind condition holds if $(\Sigma \otimes \mathbf{I})\mathbf{X}^{\text{SUR}} = \mathbf{X}^{\text{SUR}}\mathbf{Q}$. Use property (2.121) to find the matrix \mathbf{Q} .
(b) For this scenario, the Zyskind condition holds if $(\Sigma \otimes \mathbf{I})\mathbf{X}^{\text{SUR}} = \mathbf{X}^{\text{SUR}}\mathbf{Q}$. Use the diagonal property of Σ to find the matrix \mathbf{Q} .

10.15 (a) $\hat{\mathbf{B}} = \begin{pmatrix} .6264 & 83.243 \\ .0009 & .029 \\ -.0010 & -.013 \\ .0015 & -.004 \end{pmatrix}$

- (b) $\Lambda = .724, V^{(s)} = .280, U^{(s)} = .375, \theta = .264$

- (c) $\lambda_1 = .3594, \lambda_2 = .0160$. The essential rank of $\hat{\mathbf{B}}_1$ is 1, and the power ranking is $\theta > U^{(s)} > \Lambda > V^{(s)}$.

- (d) The Wilks' Λ test of x_2 adjusted for x_1 and x_3 , for example, is given by (10.74) as

$$\Lambda(x_2|x_1, x_3) = \frac{\Lambda(x_1, x_2, x_3)}{\Lambda(x_1, x_3)},$$

which is distributed as $\Lambda_{p,1,n-4}$ and has an exact F -transformation. The tests for x_1 and x_3 are similar. For the three tests we obtain the following:

	Λ	F	p -Value
$x_1 x_2, x_3$.931	1.519	.231
$x_2 x_1, x_3$.887	2.606	.086
$x_3 x_1, x_2$.762	6.417	.004

- (e) Confidence intervals for the two elements of $E(\mathbf{y}_0)$ are: (0.86, 0.94) and (87.20, 93.39).
- (f) Prediction intervals for the two elements of \mathbf{y}_0 are: (0.67, 1.13) and (72.63, 107.98).

10.16 (a) $\hat{\mathbf{B}} = \begin{pmatrix} 34.282 & 35.802 \\ .394 & .245 \\ .529 & .471 \end{pmatrix}$

(b) $\Lambda = .377, V^{(s)} = .625, U^{(s)} = 1.647, \theta = .622$

(c) $\lambda_1 = 1.644, \lambda_2 = .0029$. The essential rank of $\hat{\mathbf{B}}_1$ is 1, and the power ranking is $\theta > U^{(s)} > \Lambda > V^{(s)}$.

	Λ	F	p -Value
$x_1 x_2$.888	1.327	.287
$x_2 x_1$.875	1.506	.245

- (e) Confidence intervals for the two elements of $E(\mathbf{y}_0)$ are: (170.31, 180.44) and (139.51, 147.25).

- (f) Prediction intervals for the two elements of \mathbf{y}_0 are: (160.23, 190.52) and (131.82, 154.94).

10.17 (a) $\hat{\mathbf{B}} = \begin{pmatrix} 54.870 & 65.679 & 58.106 \\ .054 & -.048 & .018 \\ -.024 & .163 & .012 \\ .107 & -.036 & .125 \end{pmatrix}$

(b) $\Lambda = .665, V^{(s)} = .365, U^{(s)} = .458, \theta = .240$

(c) $\lambda_1 = .3159, \lambda_2 = .1385, \lambda_3 = .0037$. The essential rank of $\hat{\mathbf{B}}_1$ is 2, and the power ranking is $V^{(s)} > \Lambda > U^{(s)} > \theta$.

	Λ	F	p -Value
$x_1 x_2, x_3$.942	.903	.447
$x_2 x_1, x_3$.847	2.653	.060
$x_3 x_1, x_2$.829	3.020	.040

	Λ	F	p -Value
$y_1 y_2, y_3$.890	1.804	.160
$y_2 y_1, y_3$.833	2.932	.044
$y_3 y_1, y_2$.872	2.159	.106

- (f) Confidence intervals for the three elements of $E(\mathbf{y}_0)$ are: (65.55, 71.62), (71.06, 76.27), and (70.96, 76.29).

- (g) Prediction intervals for the three elements of \mathbf{y}_0 are: (49.07, 88.10), (56.92, 90.41), and (56.51, 90.74).

10.18 (a) $\hat{\mathbf{B}} = \begin{pmatrix} -4.140 & 4.935 \\ 1.103 & -.955 \\ .231 & -.222 \\ 1.171 & 1.773 \\ .111 & .048 \\ .617 & -.058 \\ .267 & .485 \\ -.263 & -.209 \\ -.004 & -.004 \end{pmatrix}$

Test of overall regression of (y_1, y_2) on (x_1, x_2, \dots, x_8) : $\Lambda = .4642$ (with $p = 2$, exact $F = 1.169$, p -value = .332). Tests on subsets (the F 's are exact because $p = 2$):

	Λ	F	p -Value
(b) $x_7, x_8 x_1, x_2, \dots, x_6$.856	.808	.527
(c) $x_4, x_5, x_6 x_1, x_2, x_3, x_7, x_8$.674	1.457	.218
(d) $x_1, x_2, x_3 x_4, x_5, \dots, x_8$.569	2.170	.066

- 10.19 (a)** The overall test of (y_1, y_2) on (x_1, x_2, \dots, x_8) gives $\Lambda = .4642$, with (exact) $F = 1.169$ (p -value = .332). Even though this test result is not significant, we give the results of a backward elimination for illustrative purposes:

Step	Partial Λ -Test on Each x_i Using (10.87)							
	x_1	x_2	x_3	x_4	x_5	x_6	x_7	x_8
1	.723	.969	.817	.859	.821	.945	.924	.943
2	.741		.801	.851	.839	.948	.927	.940
3	.737		.837	.798	.757		.949	.938
4	.675		.852	.821	.794			.925
5	.680		.861	.835	.817			
6	.701			.805	.806			
7	.855				.930			
8	.891							

At each step, the variable deleted was not significant. In fact, the variable remaining at the last step, x_1 , is not a significant predictor of y_1 and y_2 .

- (b)** There were no significant x 's, but to illustrate, we will use the three x 's at step 6 above and test for each y :

	Λ	F	p -Value
$y_1 y_2$.701	3.548	.029
$y_2 y_1$.808	1.984	.142

10.20 (a) $\hat{\mathbf{B}} = \begin{pmatrix} 43.703 & 46.793 & 187.923 \\ .019 & -.098 & 1.016 \\ .139 & .185 & -4.953 \\ .204 & .107 & 1.606 \end{pmatrix}$

$\Lambda = .167, V^{(s)} = .883, U^{(s)} = 4.709, \theta = .823$

(b) $\hat{\mathbf{B}} = \begin{pmatrix} 99.817 & -29.120 & 121.595 \\ -.008 & -.224 & -.027 \\ .097 & 1.252 & 5.775 \\ -.049 & -.442 & -1.768 \\ -.022 & -.631 & -.488 \\ -.159 & 2.128 & 4.387 \\ .054 & -.037 & -.476 \end{pmatrix}$

$\Lambda = .110, V^{(s)} = 1.350, U^{(s)} = 4.319, \theta = .769$

(c) $\hat{\mathbf{B}} = \begin{pmatrix} 710.236 & 123.403 \\ -.1625 & .055 \\ 24.648 & .094 \\ -.8622 & -.334 \\ -.8224 & .462 \\ 23.626 & -.110 \\ 2.862 & .427 \\ -.16.186 & -.267 \\ -.268 & .014 \\ -.1160 & -.336 \end{pmatrix}$

$\Lambda = .102, V^{(s)} = 1.236, U^{(s)} = 5.475, \theta = .827$

10.21 Using a backward elimination based on (10.87), we obtain the following partial Λ -values:

Step	x_1	x_2	x_3	x_4	x_5	x_6	x_7	x_8	x_9
1	.993	.962	.916	.958	.919	.879	.981	.999	.797
2	.994	.962	.916	.956	.909	.874	.980		.626
3		.951	.883	.954	.912	.873	.981		.626
4		.948	.884	.955	.861	.867			.561
5		.953	.862		.840	.803			.561
6		.830			.781	.783			.535

At step 6, we stop and retain all four x 's because each Λ has a p -value less than .05.

CHAPTER 11

- 11.1** By (3.38), $\mathbf{S}_{yy} = \mathbf{D}_y \mathbf{R}_{yy} \mathbf{D}_y$ and $\mathbf{S}_{xx} = \mathbf{D}_x \mathbf{R}_{xx} \mathbf{D}_x$, where \mathbf{D}_y and \mathbf{D}_x are defined below (11.14). Similarly, $\mathbf{S}_{yx} = \mathbf{D}_y \mathbf{R}_{yx} \mathbf{D}_x$ and $\mathbf{S}_{xy} = \mathbf{D}_x \mathbf{R}_{xy} \mathbf{D}_y$. Substitute these into (11.7), replace \mathbf{I} by $\mathbf{D}_y^{-1} \mathbf{D}_y$, and factor out \mathbf{D}_y on the right.

11.2 Multiply (11.7) by $\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}$ on the left to obtain $(\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}\mathbf{S}_{yy}^{-1}\mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy} - r^2\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy})\mathbf{a} = \mathbf{0}$. Factor out $\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}$ on the right to write this in the form $(\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}\mathbf{S}_{yy}^{-1}\mathbf{S}_{yx} - r^2\mathbf{I})\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}\mathbf{a} = \mathbf{0}$. Upon comparing this to (11.8), we see that $\mathbf{b} = \mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}\mathbf{a}$.

11.3 When $p = 1, s$ is also 1, and there is only one canonical correlation, which is equal to R^2 from multiple regression [see comments between (11.28) and (11.29)]. Thus

$$\Lambda = \frac{1 - r_1^2}{1 - c_1^2} = \frac{1 - R_f^2}{1 - R_r^2}.$$

$$\begin{aligned} \mathbf{11.4} \quad F &= \frac{(1 - \Lambda)(n - q - 1)}{\Lambda h} = \frac{[1 - (1 - R_f^2)/(1 - R_r^2)](n - q - 1)}{[(1 - R_f^2)/(1 - R_r^2)]h} \\ &= \frac{[1 - R_r^2 - (1 - R_f^2)](n - q - 1)}{(1 - R_f^2)h} \\ &= \frac{(R_f^2 - R_r^2)(n - q - 1)}{(1 - R_f^2)h} \end{aligned}$$

11.5 By (11.39),

$$r_i^2 = \frac{\lambda_i}{1 + \lambda_i}, \quad r_i^2 + r_i^2 \lambda_i = \lambda_i, \quad \lambda_i(1 - r_i^2) = r_i^2.$$

11.6 Substitute $\mathbf{E} = (n - 1)(\mathbf{S}_{yy} - \mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy})$ and $\mathbf{H} = (n - 1)\mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}$ from (11.44) and (11.45) into (11.37):

$$\begin{aligned} \mathbf{H}\mathbf{a} &= \lambda\mathbf{E}\mathbf{a}, \\ (n - 1)\mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}\mathbf{a} &= (n - 1)\lambda(\mathbf{S}_{yy} - \mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy})\mathbf{a}, \\ \mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}\mathbf{a} &= \lambda(\mathbf{S}_{yy} - \mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy})\mathbf{a}. \end{aligned}$$

11.7 By (11.42), $\mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}\mathbf{a} = r^2\mathbf{S}_{yy}\mathbf{a}$. Subtracting $r^2\mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}\mathbf{a}$ from both sides gives

$$\begin{aligned} \mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}\mathbf{a} - r^2\mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}\mathbf{a} &= r^2\mathbf{S}_{yy}\mathbf{a} - r^2\mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}\mathbf{a}, \\ (1 - r^2)\mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy}\mathbf{a} &= r^2(\mathbf{S}_{yy} - \mathbf{S}_{yx}\mathbf{S}_{xx}^{-1}\mathbf{S}_{xy})\mathbf{a}. \end{aligned}$$

11.8 (a) $r_1 = .5142, r_2 = .1255$

(b)	c_1	c_2	d_1	d_2
y_1	1.020	-.048	.436	.823
y_2	-.160	1.009	-.704	-.455
			x_3	1.081
				-.401

(c)	k	Λ	Approximate F	p -Value
	1	.7240	2.395	.035
	2	.9843	.336	.716

11.9 (a) $r_1 = .7885$, $r_2 = .0537$

(b)	\mathbf{c}_1		\mathbf{d}_1		\mathbf{d}_2
y_1	.5522	-1.3664	x_1	.5044	-1.7686
y_2	.5215	1.3784	x_2	.5383	1.7586

(c)	k	Λ	Approximate F	p -Value
	1	.3772	6.5972	.0003
	2	.9971	0.0637	.8031

11.10 (a) $r_1 = .4900$, $r_2 = .3488$, $r_3 = .0609$

(b)	\mathbf{c}_1			\mathbf{d}_1			\mathbf{d}_2	\mathbf{d}_3
y_1	.633	.091	.806	x_1	.482	-.262	1.054	
y_2	-.624	816	.147	x_2	-.578	1.024	-.059	
y_3	.643	.400	-.690	x_3	.865	.216	-.626	

(c)	k	Λ	Approximate F	p -Value
	1	.665	2.175	.029
	2	.875	1.552	.194
	3	.996	.171	.681

11.11 (a) $r_1 = .6251$, $r_2 = .4135$

(b)	\mathbf{c}_1		\mathbf{d}_1		\mathbf{d}_2
y_1	1.120	-.007	x_1	1.091	-.794
y_2	-.498	1.003	x_2	.184	-.288
			x_3	.842	1.807
			x_4	.944	.641
			x_5	1.040	-.154
			x_6	.215	1.256
			x_7	-.603	-.528
			x_8	-.641	-.588

(c)	k	Λ	Approximate F	p -Value
	1	.4642	1.1692	.3321
	2	.7553	.9718	.4766

11.12 (b) By (11.34),

$$\Lambda(x_7, x_8 | x_1, x_2, \dots, x_6) = \frac{\prod_{i=1}^2 (1 - r_i^2)}{\prod_{i=1}^2 (1 - c_i^2)},$$

where r_1^2 and r_2^2 are the squared canonical correlations from the full model and c_1^2 and c_2^2 are the squared canonical correlations from the reduced model:

$$\Lambda(x_7, x_8 | x_1, x_2, \dots, x_6) = \frac{(1 - .6208^2)(1 - .4947^2)}{(1 - .2650^2)(1 - .0886^2)} = \frac{.4643}{.9225} = .5033$$

$$\begin{aligned}
 \text{(c)} \quad \Lambda(x_4, x_5, x_6 | x_1, x_2, x_3, x_7, x_8) &= \frac{(1 - .6208^2)(1 - .4947^2)}{(1 - .3301^2)(1 - .1707^2)} \\
 &= \frac{.4643}{.8651} = .5367 \\
 \text{(d)} \quad \Lambda(x_1, x_2, x_3 | x_4, x_5, \dots, x_8) &= \frac{(1 - .6208^2)(1 - .4947^2)}{(1 - .4831^2)(1 - .2185^2)} \\
 &= \frac{.4643}{.7300} = .6359
 \end{aligned}$$

11.13 (a) $r_1 = .9279, r_2 = .5622, r_3 = .1660,$

k	Λ	Approximate F	p -Value
1	.0925	17.9776	<.0001
2	.6651	4.6366	.0020
3	.9725	1.1898	.2816

(b) $r_1 = .8770, r_2 = .6776, r_3 = .3488,$

k	Λ	Approximate F	p -Value
1	.1097	6.919	<.0001
2	.4751	3.427	.001
3	.8783	1.351	.269

(c) $r_1 = .9095, r_2 = .6395,$

k	Λ	Approximate F	p -Value
1	.1022	8.2757	<.0001
2	.5911	3.1129	.0089

(d) $r_1 = .9029, r_2 = .7797, r_3 = .3597, r_4 = .3233, r_5 = .0794,$

k	Λ	Approximate F	p -Value
1	.0561	4.992	<.0001
2	.3037	2.601	.0007
3	.7747	.829	.6210
4	.8898	.761	.6030
5	.9937	.124	.8840

CHAPTER 12

12.1 From $\lambda = \mathbf{a}'\mathbf{S}\mathbf{a}/\mathbf{a}'\mathbf{a}$ in (12.7), we obtain $\lambda\mathbf{a}'\mathbf{a} = \mathbf{a}'\mathbf{S}\mathbf{a}$, which can be factored as $\mathbf{a}'(\mathbf{S}\mathbf{a} - \lambda\mathbf{a}) = 0$. Since $\mathbf{a} = \mathbf{0}$ is not a solution to $\lambda = \mathbf{a}'\mathbf{S}\mathbf{a}/\mathbf{a}'\mathbf{a}$, we have $\mathbf{S}\mathbf{a} - \lambda\mathbf{a} = \mathbf{0}$.

$$\text{12.2 } |\mathbf{R} - \lambda\mathbf{I}| = 0, \quad \begin{vmatrix} 1 - \lambda & r \\ r & 1 - \lambda \end{vmatrix} = (1 - \lambda)^2 - r^2 = 0,$$

$$(1 - \lambda + r)(1 - \lambda - r) = 0, \quad \lambda = 1 \pm r$$

With $\lambda_1 = 1 + r$ in $(\mathbf{R} - \lambda_1\mathbf{I})\mathbf{a}_1 = \mathbf{0}$, we obtain

$$\begin{pmatrix} -r & r \\ r & -r \end{pmatrix} \begin{pmatrix} a_{11} \\ a_{12} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

which gives $a_{11} = a_{12}$ for any r . Normalizing to $\mathbf{a}'_1 \mathbf{a}_1 = 1$, yields $a_{11} = 1/\sqrt{2}$.

- 12.3 (a)** By (4.14) and a comment in Section 7.1, the likelihood ratio is given by $\text{LR} = (|\mathbf{S}|/|\mathbf{S}_0|)^{n/2}$, where \mathbf{S}_0 is the estimate of Σ under H_0 . By (2.108) and (7.6), the test statistic is

$$\begin{aligned}-2 \ln \text{LR} &= -2 \ln \left(\frac{|\mathbf{S}|}{|\mathbf{S}_0|} \right)^{n/2} = -2 \left(\frac{n}{2} \right) \ln \frac{\prod_{i=1}^{p-k} \lambda_i \prod_{i=p-k+1}^p \lambda_i}{\prod_{i=1}^{p-k} \lambda_i \prod_{i=p-k+1}^p \bar{\lambda}} \\ &= -n \ln \frac{\prod_{i=p-k+1}^p \lambda_i}{\bar{\lambda}^k} = n \left(k \ln \bar{\lambda} - \sum_{i=p-k+1}^p \lambda_i \right).\end{aligned}$$

In (12.15), the coefficient n is modified to give an improved chi-square approximation.

- 12.4** If \mathbf{S} is diagonal, then $\lambda_i = s_{ii}$, as in (12.17). Thus

$$\begin{aligned}\mathbf{S} \mathbf{a}_i &= \lambda_i \mathbf{a}_i = s_{ii} \mathbf{a}_i \\ \begin{pmatrix} s_{11} & 0 & \cdots & 0 \\ 0 & s_{22} & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \cdots & s_{pp} \end{pmatrix} \begin{pmatrix} a_{i1} \\ a_{i2} \\ \vdots \\ a_{ip} \end{pmatrix} &= \begin{pmatrix} s_{11} a_{i1} \\ s_{22} a_{i2} \\ \vdots \\ s_{pp} a_{ip} \end{pmatrix} = \begin{pmatrix} s_{ii} a_{i1} \\ s_{ii} a_{i2} \\ \vdots \\ s_{ii} a_{ip} \end{pmatrix}\end{aligned}$$

From the first element, we obtain $s_{11} a_{i1} = s_{ii} a_{i1}$ or $(s_{11} - s_{ii}) a_{i1} = 0$. Since $s_{11} - s_{ii} \neq 0$ (unless $i = 1$), we must have $a_{i1} = 0$. Thus, $\mathbf{a}_i = (0, \dots, 0, a_{ii}, 0, \dots, 0)'$, and normalizing \mathbf{a}_i leads to $a_{ii} = 1$.

- 12.5** By (10.34) and (12.2),

$$\begin{aligned}R_{y_i|z_1, \dots, z_k}^2 &= \frac{\mathbf{s}'_{y_i z} \mathbf{S}_{zz}^{-1} \mathbf{s}_{y_i z}}{s_{y_i}^2} \\ &= (s_{y_i z_1}, s_{y_i z_2}, \dots, s_{y_i z_k}) \begin{pmatrix} s_{z_1}^2 & 0 & \cdots & 0 \\ 0 & s_{z_2}^2 & \cdots & 0 \\ \vdots & & & \vdots \\ 0 & \cdots & & s_{z_k}^2 \end{pmatrix}^{-1} \begin{pmatrix} s_{y_i z_1} \\ s_{y_i z_2} \\ \vdots \\ s_{y_i z_k} \end{pmatrix} / s_{y_i}^2.\end{aligned}$$

Show that this is equal to

$$R_{y_i|z_1, \dots, z_k}^2 = \sum_{j=1}^k \frac{s_{y_i z_j}^2}{s_{z_j}^2 s_{y_i}^2} = \sum_{j=1}^k r_{y_i z_j}^2.$$

- 12.6** The variances of y_1, y_2, x_1, x_2 , and x_3 on the diagonal of \mathbf{S} are .016, 70.6, 1106.4, 2381.9, and 2136.4. The eigenvalues of \mathbf{S} and \mathbf{R} are as follows:

S			R		
λ_i	$\lambda_i / \sum_j \lambda_j$	Cumulative	λ_i	$\lambda_i / \sum_j \lambda_j$	Cumulative
3466.18	.608607	.60861	1.72	.34	.34
1264.47	.222021	.83063	1.23	.25	.59
895.27	.157195	.98782	.96	.19	.78
69.34	.012174	.99999	.79	.16	.94
.01	.000002	1.00000	.30	.06	1.00

Two principal components of \mathbf{S} account for 83% of the variance, but it requires three principal components of \mathbf{R} to reach 78%. For most purposes we would use two components of \mathbf{S} , although with three we could account for 99% of the variance. However, we show all five eigenvectors below because of the interesting pattern they exhibit. The first principal component is largely a weighted average of the last two variables, x_2 and x_3 , which have the largest variances. The second and third components represent contrasts in the last three variables and could be described as “shape” components. The fourth and fifth components are associated uniquely with y_2 and y_1 , respectively. These components are “variable specific,” as described in the discussion of method 1 in Section 12.6.

As expected, the principal components of \mathbf{R} show an entirely different pattern. All five variables contribute to the first three components of \mathbf{R} , whereas in \mathbf{S} , y_1 and y_2 have small variances and contribute almost nothing to the first three components. The eigenvectors of \mathbf{S} and \mathbf{R} are as follows:

S					R					
	a₁	a₂	a₃	a₄	a₅	a₁	a₂	a₃	a₄	a₅
y_1	.0004	-.0008	.0018	.0029	.9999	.42	.53	-.42	-.40	.46
y_2	-.0080	.0166	.0286	.9994	-.0029	.07	.68	.16	.70	-.10
x_1	.1547	.6382	.7535	-.0309	-.0008	.36	.20	.76	-.44	-.24
x_2	.7430	.4279	-.5145	.0136	.0009	.54	-.43	.25	.39	.56
x_3	.6511	-.6397	.4083	.0042	-.0015	.63	-.18	-.40	.10	-.64

$$12.7 \quad \mathbf{S} = \begin{pmatrix} 65.1 & 33.6 & 47.6 & 36.8 & 25.4 \\ 33.6 & 46.1 & 28.9 & 40.3 & 28.4 \\ 47.6 & 28.9 & 60.7 & 37.4 & 41.1 \\ 36.8 & 40.3 & 37.4 & 62.8 & 31.7 \\ 25.4 & 28.4 & 41.1 & 31.7 & 58.2 \end{pmatrix}$$

$$\mathbf{R} = \begin{pmatrix} 1.00 & .61 & .76 & .58 & .41 \\ .61 & 1.00 & .55 & .75 & .55 \\ .76 & .55 & 1.00 & .61 & .69 \\ .58 & .75 & .61 & 1.00 & .52 \\ .41 & .55 & .69 & .52 & 1.00 \end{pmatrix}$$

The eigenvalues of \mathbf{S} and \mathbf{R} are as follows:

S			R		
λ_i	$\lambda_i / \sum_j \lambda_j$	Cumulative	λ_i	$\lambda_i / \sum_j \lambda_i$	Cumulative
200.4	.684	.684	3.42	.683	.683
36.1	.123	.807	.61	.123	.806
34.1	.116	.924	.57	.114	.921
15.0	.051	.975	.27	.054	.975
7.4	.025	1.000	.13	.025	1.000

The first three eigenvectors of \mathbf{S} and \mathbf{R} are as follows:

S			R		
\mathbf{a}_1	\mathbf{a}_2	\mathbf{a}_3	\mathbf{a}_1	\mathbf{a}_2	\mathbf{a}_3
.47	-.58	-.42	.44	-.20	-.68
.39	-.11	.45	.45	-.43	.35
.49	.10	-.48	.47	.37	-.38
.47	-.12	.62	.45	-.39	.33
.41	.80	-.09	.41	.70	.41

The variances in \mathbf{S} are nearly identical, and the covariances are likewise similar in magnitude. Consequently, the percent of variance explained by the eigenvalues of \mathbf{S} and \mathbf{R} are indistinguishable. The interpretation of the second principal component from \mathbf{S} is slightly different from that of the second one from \mathbf{R} , but otherwise there is little to choose between them.

- 12.8** The variances on the diagonal of \mathbf{S} are 95.5, 73.2, 76.2, 808.6, 505.9, and 508.7. The eigenvalues of \mathbf{S} and \mathbf{R} are as follows:

S			R		
λ_i	$\lambda_i / \sum_j \lambda_j$	Cumulative	λ_i	$\lambda_i / \sum_j \lambda_j$	Cumulative
1152.0	.557	.557	2.17	.363	.363
394.1	.191	.748	1.08	.180	.543
310.8	.150	.898	.98	.163	.706
97.8	.047	.945	.87	.144	.850
68.8	.033	.978	.55	.092	.942
44.6	.022	1.000	.35	.058	1.000

We could keep either two or three components from \mathbf{S} . The first three components of \mathbf{S} account for a larger percent of variance than do those from \mathbf{R} . The first three eigenvectors of \mathbf{S} and \mathbf{R} are as follows:

S			R		
a₁	a₂	a₃	a₁	a₂	a₃
.080	.092	-.069	.336	.176	.497
.034	-.018	.202	.258	.843	-.093
.076	.122	-.011	.370	.049	.466
.758	-.446	-.469	.475	-.329	-.358
.493	-.081	.844	.486	.079	-.567
.412	.878	-.147	.471	-.376	.278

As expected, the first three principal components from **S** are heavily influenced by the last three variables because of their relatively large variances.

- 12.9** The variances on the diagonal of **S** are .69; 5.4; 2,006, 682.4; 90.3; 56.4; 18.1. With the large variance of y_3 , we would expect the first principal component from **S** to account for most of the variance, and y_3 would essentially constitute that single component. This is indeed the pattern that emerges in the eigenvalues and eigenvectors of **S**. The principal components from **R**, on the other hand, are not dominated by y_3 . The eigenvalues of **S** and **R** are as follows:

S		R			
λ_i	$\lambda_i / \sum_j \lambda_j$	λ_i	$\lambda_i / \sum_j \lambda_j$	Cumulative	
2,006,760	.999954	2.42	.404	.404	
65	.000033	1.40	.234	.638	
18	.000009	1.03	.171	.809	
7	.000003	.92	.153	.963	
3	.000001	.20	.033	.996	
0	.000000	.02	.004	1.000	

Most of the correlations in **R** are small (only three exceed .3), and its first three principal components account for only 72% of the variance. The first three eigenvectors of **S** and **R** are as follows:

S			R		
a₁	a₂	a₃	a₁	a₂	a₃
.00016	.005	-.0136	.424	-.561	-.150
.00051	.017	.0787	.446	-.528	.087
.99998	-.001	-.0002	.563	.387	-.051
.00529	.698	.0174	.454	.267	.166
.00322	-.716	.0195	.303	.425	-.296
.00020	.025	.9965	.073	.069	.923

- 12.10** Covariance matrix for males:

$$\mathbf{S}_M = \begin{pmatrix} 5.19 & 4.55 & 6.52 & 5.25 \\ 4.55 & 13.18 & 6.76 & 6.27 \\ 6.52 & 6.76 & 28.67 & 14.47 \\ 5.25 & 6.27 & 14.47 & 16.65 \end{pmatrix}$$

Covariance matrix for females:

$$\mathbf{S}_F = \begin{pmatrix} 9.14 & 7.55 & 4.86 & 4.15 \\ 7.55 & 18.60 & 10.22 & 5.45 \\ 4.86 & 10.22 & 30.04 & 13.49 \\ 4.15 & 5.45 & 13.49 & 28.00 \end{pmatrix}$$

The eigenvalues are as follows:

Males			Females		
λ_i	$\lambda_i / \sum_j \lambda_j$	Cumulative	λ_i	$\lambda_i / \sum_j \lambda_j$	Cumulative
43.56	.684	.684	48.96	.571	.571
11.14	.175	.858	18.46	.215	.786
6.47	.102	.960	13.54	.158	.944
2.52	.040	1.000	4.82	.056	1.000

The first two eigenvectors are as follows:

Males		Females	
\mathbf{a}_1	\mathbf{a}_2	\mathbf{a}_1	\mathbf{a}_2
.24	.21	.22	.27
.31	.85	.39	.62
.76	-.48	.68	.17
.52	.09	.58	-.72

The variances in \mathbf{S}_M have a slightly wider range (5.19 to 28.67) than those in \mathbf{S}_F (9.14 to 30.04), and this is reflected in the eigenvalues. The first two components account for 86% of the variance from \mathbf{S}_M , whereas the first two account for 79% from \mathbf{S}_F .

12.11 Covariance matrix for species 1:

$$\mathbf{S}_1 = \begin{pmatrix} 187.6 & 176.9 & 48.4 & 113.6 \\ 176.9 & 345.4 & 76.0 & 118.8 \\ 48.4 & 76.0 & 66.4 & 16.2 \\ 113.6 & 118.8 & 16.2 & 239.9 \end{pmatrix}$$

Covariance matrix for species 2:

$$\mathbf{S}_2 = \begin{pmatrix} 101.8 & 128.1 & 37.0 & 32.6 \\ 128.1 & 389.0 & 165.4 & 94.4 \\ 37.0 & 165.4 & 167.5 & 66.5 \\ 32.6 & 94.4 & 66.5 & 177.9 \end{pmatrix}$$

The eigenvalues are as follows:

Species 1			Species 2		
λ_i	$\lambda_i / \sum_j \lambda_j$	Cumulative	λ_i	$\lambda_i / \sum_j \lambda_j$	Cumulative
561.3	.669	.669	555.7	.664	.664
169.0	.201	.870	145.4	.174	.838
65.3	.078	.948	93.5	.112	.950
43.7	.057	1.000	41.7	.050	1.000

The first two eigenvectors are as follows:

Species 1		Species 2	
a_1	a_2	a_1	a_2
.50	.01	.28	-.20
.72	-.48	.81	-.34
.17	-.22	.42	.14
.45	.85	.30	.91

The variances in S_1 have a wider range than those in S_2 , and the first two components of S_1 account for a higher percent of variance.

12.12 The variances on the diagonal of S in each case are:

- (a) Pooled: 536.0, 59.9, 116.0, 896.4, 248.1, 862.0
- (b) Unpooled: 528.2, 68.9, 145.2, 1366.4, 264.4, 1069.1

The eigenvalues are as follows:

Pooled			Unpooled		
λ_i	$\lambda_i / \sum_j \lambda_j$	Cumulative	λ_i	$\lambda_i / \sum_j \lambda_j$	Cumulative
1050.6	.386	.386	1722.0	.500	.500
858.3	.316	.702	878.4	.255	.755
398.9	.147	.849	401.4	.117	.872
259.2	.095	.944	261.1	.076	.948
108.1	.040	.984	128.9	.037	.985
43.4	.016	1.000	50.4	.015	1.000

The first three eigenvectors are as follows:

Pooled			Unpooled		
a_1	a_2	a_3	a_1	a_2	a_3
.441	-.190	.864	.212	.389	.888
.041	-.038	.082	-.039	.064	.096
-.039	.031	.143	.080	-.066	.081
.450	.892	-.033	.776	-.608	.081
-.019	-.001	-.054	-.096	.010	.015
.774	-.407	-.471	.580	.686	-.434

- (c) The pattern in both eigenvalues and eigenvectors is similar for the pooled and unpooled cases. The first three principal components account for 87.2% of the variance in the unpooled case compared to 84.9% for the pooled case.

12.13 The variances on the diagonal of \mathbf{S} in each case are:

- (a) Pooled: 49.1, 8.1, 12140.8, 136.2, 210.8, 2983.9
 (b) Unpooled: 63.2, 8.0, 15168.9, 186.6, 255.4, 4660.7

The eigenvalues are as follows:

Pooled			Unpooled		
λ_i	$\lambda_i / \sum_j \lambda_j$	Cumulative	λ_i	$\lambda_i / \sum_j \lambda_j$	Cumulative
12,809.0	.8249	.8249	17,087.0	.8400	.8400
2,455.9	.1582	.9830	2,958.0	.1454	.9854
137.1	.0088	.9918	168.6	.0083	.9937
77.2	.0050	.9968	77.1	.0038	.9974
42.2	.0027	.9995	44.7	.0022	.9996
7.4	.0005	1.0000	7.3	.0004	1.0000

The eigenvectors are as follows:

Pooled		Unpooled	
a_1	a_2	a_1	a_2
-.004	-.000	.013	.027
-.005	.004	-.004	.004
.968	-.233	.931	-.355
-.002	.023	.028	.069
.103	.041	.103	.021
.228	.971	.350	.932

12.14 The variances on the diagonal of \mathbf{S} are all less than 1 except $s_{x_4}^2 = 5.02$ and $s_{x_8}^2 = 1541.08$. We therefore expect the last variable, x_8 , to dominate the principal components of \mathbf{S} . This is the case for \mathbf{S} but not for \mathbf{R} . The eigenvalues of \mathbf{S} and \mathbf{R} are as follows:

S		R		
λ_i	$\lambda_i / \sum_j \lambda_j$	λ_i	$\lambda_i / \sum_j \lambda_j$	Cumulative
1541.55	.996273	3.174	.317	.317
4.83	.003123	2.565	.256	.574
.44	.000286	1.432	.143	.717
.27	.000174	1.277	.128	.845
.10	.000066	.542	.054	.899
.07	.000043	.473	.047	.946
.02	.000014	.251	.025	.971
.02	.000011	.118	.012	.983
.01	.000005	.104	.010	.994
.00	.000003	.064	.006	1.000

The eigenvectors of **S** and **R** are as follows:

S		R			
a_1	a_2	a_1	a_2	a_3	a_4
.0009	-.005	.12	.19	.69	.10
.0007	-.034	.06	.32	.54	.26
.0029	-.007	.46	-.06	.07	-.38
.0014	.004	.29	.17	-.18	.49
.0059	-.009	.52	.14	-.04	-.01
-.0150	.982	-.09	-.42	.07	.55
-.0028	-.092	-.31	.45	-.01	-.14
-.0022	-.158	-.23	.54	-.14	-.10
.0044	-.011	.09	.36	-.38	.44
.9998	.014	.50	.11	-.13	-.09

- 12.15** The variances in the diagonal of **S** are: 55.7, 10.9, 402.7, 25.7, 13.4, 438.3, 1.5, 106.2, 885.6, 22227.2, 214.1

The eigenvalues of **S** and **R** are as follows:

S			R		
λ_i	$\lambda_i / \sum_j \lambda_j$	Cumulative	λ_i	$\lambda_i / \sum_j \lambda_j$	Cumulative
22,303.5	.91479	.91479	6.020	.54730	.54730
1590.7	.06524	.98003	2.119	.19267	.73996
358.0	.01469	.99471	1.130	.10275	.84272
63.4	.00260	.99731	.760	.06909	.91181
29.3	.00120	.99852	.355	.03231	.94411
17.1	.00070	.99922	.259	.02358	.96769
12.7	.00052	.99974	.122	.01110	.97879
2.8	.00012	.99986	.110	.01004	.98883
1.9	.00008	.99994	.060	.00544	.99427
.9	.00004	.99997	.042	.00384	.99810
.7	.00003	1.00000	.021	.00190	1.00000

The eigenvectors of \mathbf{S} and \mathbf{R} are as follows:

S		R				
	a₁	a₂	a₁	a₂	a₃	a₄
y_1	-.0097	.1331	.3304	-.0787	.0880	-.2807
y_2	.0006	.0608	.3542	.1928	.1071	-.2301
y_3	-.0141	.4397	.3923	.0518	.1105	-.1413
y_4	-.0033	.1078	.3820	.0474	.1334	-.0104
y_5	.0101	.0398	.2323	.5303	.0154	-.0710
y_6	.0167	.4290	.3621	.2361	.1198	.1350
y_7	-.0012	-.0072	-.0884	.0213	.7946	.5414
y_8	.0275	-.1844	-.2501	.5023	.0826	-.1506
y_9	.0456	-.6657	-.3111	.3595	.2136	-.2278
y_{10}	.9982	.0346	-.0243	.4685	-.4669	.5001
y_{11}	.0034	.3311	.3357	-.1153	-.1853	.4550

For most purposes, one or two principal components would suffice for \mathbf{S} , with 91% or 98% of the variance explained. For \mathbf{R} , on the other hand, three components are required to explain 84% of the variance, and seven components are necessary to reach 98%. The reduction to one or two components for \mathbf{S} is due in part to the relatively large variances of y_3, y_6, y_9 , and y_{10} . In the eigenvectors of \mathbf{S} , we see that these four variables figure prominently in the first two principal components.

CHAPTER 13

$$\begin{aligned} \text{13.1 } \text{var}(y_i) &= \text{var}(y_i - \mu_i) = \text{var}(\lambda_{i1}f_1 + \lambda_{i2}f_2 + \cdots + \lambda_{im}f_m + \varepsilon_i) \\ &= \sum_{j=1}^m \lambda_{ij}^2 \text{var}(f_j) + \text{var}(\varepsilon_i) + \sum_{j \neq k} \lambda_{ij}\lambda_{ik} \text{cov}(f_j, f_k) \\ &\quad + \sum_{j=1}^m \lambda_{ij} \text{cov}(f_j, \varepsilon_i) \\ &= \sum_{j=1}^m \lambda_{ij}^2 + \psi_i. \end{aligned}$$

The last equality follows by the assumptions $\text{var}(f_j) = 1, \text{var}(\varepsilon_i) = \psi_i, \text{cov}(f_j, f_k) = 0$, and $\text{cov}(f_j, \varepsilon_i) = 0$.

$$\begin{aligned} \text{13.2 } \text{cov}(\mathbf{y}, \mathbf{f}) &= \text{cov}(\Lambda\mathbf{f} + \boldsymbol{\varepsilon}, \mathbf{f}) && [\text{by (13.3)}] \\ &= \text{cov}(\Lambda\mathbf{f}, \mathbf{f}) && [\text{by (13.10)}] \\ &= E[\Lambda\mathbf{f} - E(\Lambda\mathbf{f})][\mathbf{f} - E(\mathbf{f})]' && [\text{by analogy to (3.31)}] \\ &= E[\Lambda\mathbf{f} - \Lambda E(\mathbf{f})][\mathbf{f} - E(\mathbf{f})]' \\ &= \Lambda E[\mathbf{f} - E(\mathbf{f})][\mathbf{f} - E(\mathbf{f})]' \\ &= \Lambda \text{cov}(\mathbf{f}) = \Lambda && [\text{by (13.7)}] \end{aligned}$$

$$\begin{aligned} \text{13.3 } E(\mathbf{f}^*) &= E(\mathbf{T}'\mathbf{f}) = \mathbf{T}'E(\mathbf{f}) = \mathbf{T}'\mathbf{0} = \mathbf{0}, \\ \text{cov}(\mathbf{f}^*) &= \text{cov}(\mathbf{T}'\mathbf{f}) = \mathbf{T}'\text{cov}(\mathbf{f})\mathbf{T} = \mathbf{T}'\mathbf{I}\mathbf{T} = \mathbf{I} \end{aligned}$$

$$\text{13.4 Let } \mathbf{E} = \mathbf{S} - (\hat{\Lambda}\hat{\Lambda}' + \hat{\Psi}). \text{ Then by (2.98), } \text{tr}(\mathbf{E}'\mathbf{E}) = \sum_{ij} e_{ij}^2. \text{ By (13.26), } \hat{\Psi} = \text{diag}(\mathbf{S} - \hat{\Lambda}\hat{\Lambda}'), \text{ and } \mathbf{E} \text{ has zeros on the diagonal. This gives the inequality}$$

$$\sum_{ij} e_{ij}^2 \leq \text{sum of squared elements of } \mathbf{S} - \hat{\mathbf{\Lambda}}\hat{\mathbf{\Lambda}}'.$$

By (2.98),

$$\text{sum of squared elements of } \mathbf{S} - \hat{\mathbf{\Lambda}}\hat{\mathbf{\Lambda}}' = \text{tr}(\mathbf{S} - \hat{\mathbf{\Lambda}}\hat{\mathbf{\Lambda}}')'(\mathbf{S} - \hat{\mathbf{\Lambda}}\hat{\mathbf{\Lambda}}').$$

Since $\mathbf{S} - \hat{\mathbf{\Lambda}}\hat{\mathbf{\Lambda}}'$ is symmetric, we have by (13.20), (13.23), and (13.24),

$$\begin{aligned}\mathbf{S} - \hat{\mathbf{\Lambda}}\hat{\mathbf{\Lambda}}' &= \mathbf{C}\mathbf{D}\mathbf{C}' - \mathbf{C}_1\mathbf{D}_1^{1/2}\mathbf{D}_1^{1/2}\mathbf{C}_1' \\ &= \mathbf{C}\mathbf{D}\mathbf{C}' - \mathbf{C}_1\mathbf{D}_1\mathbf{C}_1',\end{aligned}$$

where $\mathbf{C} = (\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_p)$ contains normalized eigenvectors of \mathbf{S} , $\mathbf{D} = \text{diag}(\theta_1, \theta_2, \dots, \theta_p)$ contains eigenvalues of \mathbf{S} , $\mathbf{C}_1 = (\mathbf{c}_1, \mathbf{c}_2, \dots, \mathbf{c}_m)$, and $\mathbf{D}_1 = \text{diag}(\theta_1, \theta_2, \dots, \theta_m)$.

Using the partitioned forms $\mathbf{C} = (\mathbf{C}_1, \mathbf{C}_2)$ and $\mathbf{D} = \begin{pmatrix} \mathbf{D}_1 & \mathbf{O} \\ \mathbf{O} & \mathbf{D}_2 \end{pmatrix}$, show that $\mathbf{C}_1'\mathbf{C}_1 = \mathbf{I}_m$, $\mathbf{C}_1'\mathbf{C}_2 = \mathbf{O}$, $\mathbf{C}'\mathbf{C}_1 = \begin{pmatrix} \mathbf{I}_m \\ \mathbf{O} \end{pmatrix}$, $\mathbf{D}\begin{pmatrix} \mathbf{I}_m \\ \mathbf{O} \end{pmatrix} = \begin{pmatrix} \mathbf{D}_1 \\ \mathbf{O} \end{pmatrix}$, $\mathbf{C}\begin{pmatrix} \mathbf{D}_1 \\ \mathbf{O} \end{pmatrix} = \mathbf{C}_1\mathbf{D}_1$, and $\mathbf{C}\mathbf{D}\mathbf{C}'\mathbf{C}_1\mathbf{D}_1\mathbf{C}_1' = \mathbf{C}_1\mathbf{D}_1^2\mathbf{C}_1'$. Show similarly that $\mathbf{C}_1\mathbf{D}_1\mathbf{C}_1'\mathbf{C}\mathbf{D}\mathbf{C}' = \mathbf{C}_1\mathbf{D}_1^2\mathbf{C}_1'$ and $\mathbf{C}_1\mathbf{D}_1\mathbf{C}_1'\mathbf{C}_1\mathbf{D}_1\mathbf{C}_1' = \mathbf{C}_1\mathbf{D}_1^2\mathbf{C}_1'$. Now by (2.97) $\text{tr}(\mathbf{CD}^2\mathbf{C}') = \text{tr}(\mathbf{C}'\mathbf{CD}^2) = \text{tr}(\mathbf{D}^2) = \sum_{i=1}^p \theta_i^2$. Similarly, $\text{tr}(\mathbf{C}_1\mathbf{D}_1^2\mathbf{C}_1') = \sum_{i=1}^m \theta_i^2$. Then

$$\begin{aligned}\text{tr}(\mathbf{S} - \hat{\mathbf{\Lambda}}\hat{\mathbf{\Lambda}}')'(\mathbf{S} - \hat{\mathbf{\Lambda}}\hat{\mathbf{\Lambda}}') &= \text{tr}(\mathbf{C}\mathbf{D}\mathbf{C}' - \mathbf{C}_1\mathbf{D}_1\mathbf{C}_1')(\mathbf{C}\mathbf{D}\mathbf{C}' - \mathbf{C}_1\mathbf{D}_1\mathbf{C}_1') \\ &= \text{tr}(\mathbf{C}\mathbf{D}\mathbf{C}'\mathbf{C}\mathbf{D}\mathbf{C}' - \mathbf{C}\mathbf{D}\mathbf{C}'\mathbf{C}_1\mathbf{D}_1\mathbf{C}_1' \\ &\quad - \mathbf{C}_1\mathbf{D}_1\mathbf{C}_1'\mathbf{C}\mathbf{D}\mathbf{C}' + \mathbf{C}_1\mathbf{D}_1\mathbf{C}_1'\mathbf{C}_1\mathbf{D}_1\mathbf{C}_1') \\ &= \sum_{i=1}^p \theta_i^2 - \sum_{i=1}^m \theta_i^2 - \sum_{i=1}^m \theta_i^2 + \sum_{i=1}^m \theta_i^2 \\ &= \sum_{i=m+1}^p \theta_i^2.\end{aligned}$$

$$13.5 \sum_{i=1}^p \sum_{j=1}^m \hat{\lambda}_{ij}^2 = \sum_{i=1}^p \left[\sum_{j=1}^m \hat{\lambda}_{ij}^2 \right] = \sum_{i=1}^p \hat{\lambda}_i^2 \quad [\text{by (13.28)}]$$

By interchanging the order of summation, we have

$$\sum_{i=1}^p \sum_{j=1}^m \hat{\lambda}_{ij}^2 = \sum_{j=1}^m \sum_{i=1}^p \hat{\lambda}_{ij}^2 = \sum_{j=1}^m \theta_j \quad [\text{by (13.29)}].$$

13.6 We use the covariance matrix to avoid working with standardized variables. The eigenvalues of \mathbf{S} are 39.16, 8.78, .66, .30, and 0. The eigenvector corresponding to $\lambda_5 = 0$ is

$$\mathbf{a}_5' = (-.75, -.25, .25, .50, .25).$$

As noted in Section 12.7, $s_{z_5}^2 = 0$ implies $z_5 = 0$. Thus

$$\begin{aligned}z_5 &= \mathbf{a}_5' \mathbf{y} = -.75y_1 - .25y_2 + .25y_3 + .50y_4 + .25y_5 = 0, \\ 3y_1 + y_2 &= y_3 + 2y_4 + y_5.\end{aligned}$$

13.7 Words data of Table 5.9:

	Principal Component Loadings		Varimax Rotated Loadings		Communalities, \hat{h}_i^2
	f_1	f_2	f_1	f_2	
Variables					
Informal words	.802	-.535	.956	.129	.930
Informal verbs	.856	-.326	.858	.321	.839
Formal words	.883	.270	.484	.786	.853
Formal verbs	.714	.658	.101	.966	.943
Variance	2.666	.899	1.894	1.671	3.565
Proportion	.666	.225	.474	.418	.891

The orthogonal matrix T for the varimax rotation as given by (13.49) is

$$\mathbf{T} = \begin{pmatrix} .750 & .661 \\ -.661 & .750 \end{pmatrix}.$$

Thus $\sin \phi = -0.661$ and $\phi = -41.4^\circ$. A graphical rotation of -40° would produce results very close to the varimax rotation.

13.8 Ramus bone data of Table 3.7:

	Principal Component Loadings		Varimax Rotated Loadings		Orthoblique Pattern Loadings
	f_1	f_2	f_1	f_2	
Variables					
8 years	.949	-.295	.884	.455	.988
8 $\frac{1}{2}$ years	.974	-.193	.830	.545	.986
9 years	.978	.171	.578	.808	.986
9 $\frac{1}{2}$ years	.943	.319	.449	.888	.991
Variance	3.695	.255	2.005	1.946	3.951
Proportion	.924	.064	.501	.486	.988

The Harris–Kaiser orthoblique rotation produced loadings for which the variables have a complexity of 1. These oblique loadings provide a much cleaner simple structure than the varimax loadings. For interpretation, we see that one factor represents variables 1 and 2, and the other factor represents variables 3 and 4. This same clustering of variables can be deduced from the varimax loadings if we simply use the larger of the two loadings for each variable.

The correlation between the two oblique factors is .87. The angle between the oblique axes is $\cos^{-1}(0.87) = 29.5^\circ$. With such a small angle between the axes and a large correlation between the factors, it is clear that a single factor

would better represent the variables. This is also borne out by the eigenvalues of the correlation matrix: 3.695, .255, .033, and .017. The first accounts for 92% of the variance and the second for only 6%.

13.9 Rootstock data of Table 6.2:

	Principal Component Loadings		Varimax Rotated Loadings		Communalities, \hat{h}_i^2
	f_1	f_2	f_1	f_2	
Variables					
Trunk 4 years	.787	.575	.167	.960	.949
Extension 4 years	.849	.467	.287	.925	.939
Trunk 15 years	.875	-.455	.946	.280	.973
Weight 15 years	.824	-.547	.973	.179	.978
Variance	2.785	1.054	1.951	1.888	3.839
Proportion	.696	.264	.488	.472	.960

The rotation was successful in producing variables with a complexity of 1, that is, partitioning the variables into two groups, each with two variables.

13.10 Fish data of Table 6.17:

	Principal Component Loadings		Varimax Rotated Loadings		Communalities, \hat{h}_i^2
	f_1	f_2	f_1	f_2	
Variables					
y_1	.830	-.403	.874	.294	.851
y_2	.783	-.504	.911	.189	.866
y_3	.803	.432	.270	.871	.831
y_4	.769	.497	.200	.893	.838
Variance	2.537	.850	1.709	1.678	3.386
Proportion	.634	.213	.427	.420	.847

- (b) The loadings for y_1 and y_2 are similar. In R below we see some indication of the reason for this; y_1 and y_2 are more highly correlated than any other pair of variables and their correlations with y_3 and y_4 are similar:

$$\mathbf{R} = \begin{pmatrix} 1.00 & .71 & .51 & .40 \\ .71 & 1.00 & .38 & .40 \\ .51 & .38 & 1.00 & .67 \\ .40 & .40 & .67 & 1.00 \end{pmatrix}.$$

- (c) By (13.58), the factor score coefficient matrix is

$$\hat{\mathbf{B}}_1 = \mathbf{R}^{-1}\hat{\Lambda} = \begin{pmatrix} .566 & -.109 \\ .636 & -.207 \\ -.130 & .584 \\ -.194 & .630 \end{pmatrix},$$

where $\hat{\Lambda}$ is the matrix of rotated factor loadings given above in part (a). The factor scores are given by (13.59) as follows:

Method 1		Method 2		Method 3	
\hat{f}_1	\hat{f}_2	\hat{f}_1	\hat{f}_2	\hat{f}_1	\hat{f}_2
.544	1.151	-.254	.309	-1.156	2.104
1.250	-.254	-.309	-1.534	-.321	.878
1.017	1.120	-1.865	-1.558	-.671	.947
-.147	-1.583	-.999	-.690	.067	1.130
.219	-.103	.520	-.343	-1.610	-.458
1.007	.679	.919	-.111	.557	.491
1.413	-.186	-.443	-.018	-.454	1.157
-.666	-2.279	-.265	.676	-.961	.063
1.057	-1.870	1.449	-.295	-.230	1.721
.388	-.440	1.371	.295	-1.309	.054
1.328	-.298	1.260	-.027	-1.766	-.111
.694	-.033	-.000	-1.452	-1.636	-.048

- (d) A one-way MANOVA on the two factor scores comparing the three methods yielded the following values for \mathbf{E} and \mathbf{H} :

$$\mathbf{E} = \begin{pmatrix} 21.8606 & 10.3073 \\ 10.3073 & 25.2081 \end{pmatrix}, \quad \mathbf{H} = \begin{pmatrix} 13.1394 & -10.3073 \\ -10.3073 & 9.7919 \end{pmatrix}$$

The four MANOVA test statistics are $\Lambda = .3631$, $V^{(s)} = .6552$, $U^{(s)} = 1.7035$, and $\theta = .6259$. All are highly significant.

- 13.11 (a)** For the flea data of Table 5.5, the eigenvalues of \mathbf{R} are 2.273, 1.081, .450, and .196. There is a noticeable gap between 1.081 and .450, and the first two factors account for 83.9% of the variance. Thus $m = 2$ factors seem to be indicated for this set of data.

(b)

	Principal Component Loadings		Varimax Rotated Loadings		Communalities, \hat{h}_i^2	Orthoblique Pattern Loadings	
	f_1	f_2	f_1	f_2		f_1	f_2
Variables							
y_1	-.038	.989	-.025	.990	.980	-.003	.990
y_2	.889	.269	.892	.256	.862	.898	.253
y_3	.893	-.157	.891	-.170	.823	.887	-.173
y_4	.827	-.073	.823	-.084	.689	.824	-.087
Variance	2.273	1.081	2.273	1.081	3.354		
Proportion	.568	.270	.568	.270	.839		

(The variance explained by the varimax rotated factors remains the same as for the initial factors when rounded to three decimal places.)

- (c) In this case, neither of the rotations changes the initial loadings appreciably. The reason for this unusual outcome can be seen in the correlation matrix:

$$\mathbf{R} = \begin{pmatrix} 1.00 & .18 & -.17 & -.07 \\ .18 & 1.00 & .73 & .59 \\ -.17 & .73 & 1.00 & .59 \\ -.07 & .59 & .59 & 1.00 \end{pmatrix}.$$

There are clearly two clusters of variables: $\{y_1\}$ and $\{y_2, y_3, y_4\}$. We would expect two factors corresponding to these groupings to emerge after rotation. That the same pattern surfaces in the initial factor loadings (based on eigenvectors) is due to their affiliation with principal components. As noted in Section 12.8.1, if a variable has small correlations with all other variables, the variable itself will essentially constitute a principal component. In this case, y_1 has this property and makes up most of the second principal component. The first component is comprised of the other three variables.

- 13.12 (a)** For the engineer data of Table 5.6, the number of eigenvalues greater than 1 is three, but the three account for only 70% of the variance. It requires four eigenvalues to reach 84%. The scree plot also indicates four eigenvalues.

(b)

	Principal Component Loadings			Varimax Rotated Loadings			Communalities, \hat{h}_i^2
	f_1	f_2	f_3	f_1	f_2	f_3	
Variables							
y_1	.536	.461	.478	-.063	.834	.170	.729
y_2	-.129	.870	-.182	-.357	.100	.818	.806
y_3	.514	-.254	-.448	.724	-.026	.068	.529
y_4	.724	-.366	-.110	.739	.295	-.193	.670
y_5	-.416	-.414	.649	-.484	-.013	-.729	.766
y_6	.715	.124	.420	.239	.800	-.069	.702
Variance	1.775	1.354	1.073	1.493	1.435	1.275	4.202
Proportion	.296	.226	.179	.249	.239	.212	.700

- (c) The initial communality estimates for the six variables are given by (13.36) as .215, .225, .113, .255, .161, .248. With these substituted for the diagonal of \mathbf{R} , the eigenvalues of $\mathbf{R} - \hat{\Psi}$ are

Eigenvalue	.994	.569	.255	-.025	-.237	-.339
Proportion	.816	.468	.209	-.020	-.195	-.278
Cumulative	.816	1.284	1.493	1.473	1.278	1.000

The principal factor loadings and varimax rotation are as follows:

	Principal Component Loadings			Varimax Rotated Loadings			Communalities, \hat{h}_i^2
	f_1	f_2	f_3	f_1	f_2	f_3	
Variables							
y_1	.403	.312	.227	.030	.536	.151	.311
y_2	-.106	.569	-.100	-.288	.083	.505	.345
y_3	.343	-.139	-.197	.413	.060	.037	.176
y_4	.559	-.247	-.090	.564	.233	-.094	.381
y_5	-.286	-.246	.328	-.262	-.088	-.417	.250
y_6	.556	.089	.197	.258	.537	.003	.356

- (d) The pattern of loadings is similar in parts (b) and (c), and the interpretation of the three factors would be the same.

13.13 Probe word data of Table 3.6:

	Principal Component Loadings		Varimax Rotated Loadings		Communalities, \hat{h}_i^2	Orthoblique Pattern Loadings	
	f_1	f_2	f_1	f_2		f_1	f_2
Variables							
y_1	.817	-.157	.732	.395	.692	.737	.131
y_2	.838	-.336	.861	.271	.815	.963	-.092
y_3	.874	.288	.494	.776	.847	.248	.734
y_4	.838	-.308	.844	.292	.798	.931	-.057
y_5	.762	.547	.244	.905	.879	-.134	1.023
Variance	3.416	.614	2.294	1.736	4.031		
Proportion	.683	.123	.459	.347	.806		

The loadings for y_2 are similar to those for y_4 in all three sets of loadings. The reason for this can be seen in the correlation matrix

$$\mathbf{R} = \begin{pmatrix} 1.00 & .61 & .76 & .58 & .41 \\ .61 & 1.00 & .55 & .75 & .55 \\ .76 & .55 & 1.00 & .61 & .69 \\ .58 & .75 & .61 & 1.00 & .52 \\ .41 & .55 & .69 & .52 & 1.00 \end{pmatrix}.$$

The correlations of y_2 with y_1, y_3 , and y_5 are very similar to the correlations of y_4 with y_1, y_3 , and y_5 .

CHAPTER 14**14.1** Multiply both sides of the inequality by 2 to obtain:

$$\begin{aligned} p^2 + p &\geq 2pm - 2m^2 + m^2 + m + 2p \\ \Rightarrow p^2 - 2pm + m^2 &\geq p + m \\ \Rightarrow (p - m)^2 &\geq p + m \end{aligned}$$

- 14.2** (a) Identified
 (b) Identified
 (c) Not identified
 (d) Identified

- 14.3** (a) Identified
 (b) 0

- (c) Find $\text{var}(y_1)$, $\text{var}(y_2)$, $\text{var}(y_3)$, $\text{cov}(y_1, y_2)$, $\text{cov}(y_1, y_3)$, and $\text{cov}(y_2, y_3)$.
For example, $\text{var}(y_1) = \phi + \psi_1$ and $\text{cov}(y_1, y_2) = \lambda_2\phi$.

- (d) $\hat{\lambda}_2 = s_{23}/s_{13}$, $\hat{\lambda}_3 = s_{23}/s_{12}$, $\hat{\phi} = s_{12}s_{13}/s_{23}$, $\hat{\psi}_{11} = s_{11} - s_{12}s_{13}/s_{23}$,
 $\hat{\psi}_{22} = s_{22} - s_{12}s_{23}/s_{13}$, and $\hat{\psi}_{33} = s_{33} - s_{13}s_{23}/s_{12}$.

14.4 (a) $\chi^2 = 6.5811$, df = 5, p-value = 0.2537

Bentler's CFI = 0.9280

RMSEA = 0.1778

SRMR = 0.0711

- (b) Using (14.25), $z_{21} = 2.5483$, $z_{31} = 2.8623$, $z_{41} = 2.5851$, and $z_{51} = 2.2674$. Since all exceed $z_{\alpha/2} = 1.96$ in absolute value, all loadings are significant and the model cannot be simplified.

- (c) Same as part (a).

14.5 (a) $\chi^2 = 14.8718$, df = 12 (due to one error variance being constrained to equal zero), p-value = 0.2485

Bentler's CFI = 0.9824

RMSEA = 0.0908

SRMR = 0.0308

14.6 (a) $\chi^2 = 17.7127$, df = 5 (due to one error variance being constrained to equal zero), p-value = 0.0033

Bentler's CFI = 0.9401

RMSEA = 0.1690

SRMR = 0.0630

- (b) $\chi^2 = 34.3303$, df = 9 (due to one error variance being constrained to equal zero), p-value < 0.0001

Bentler's CFI = 0.8806

RMSEA = 0.1778

SRMR = 0.0790

14.7 (a) $\chi^2 = 4.1224$, df = 4, p-value = 0.3897

Bentler's CFI = 0.9895

RMSEA = 0.0280

SRMR = 0.0608

(b) Variable	Variance explained by the factors
Intelligence	20.5%
Form relations	undefined
Dynamometer	4.4%
Dotting	18.8%
Sensory motor coordination	2.3%
Perseveration	53.8%

14.8 $\chi^2 = 101.3717$, df = 2, p-value < 0.0001

Bentler's CFI = 0.9895

$$\begin{aligned} \text{RMSEA} &= 1.0282 \\ \text{SRMR} &= 0.2237 \end{aligned}$$

CHAPTER 15

15.1 Adding and subtracting \bar{x} and \bar{y} in (15.2) (squared), we obtain

$$\begin{aligned} d^2(\mathbf{x}, \mathbf{y}) &= \sum_{j=1}^p [(x_j - \bar{x}) - (y_j - \bar{y}) + (\bar{x} - \bar{y})]^2 \\ &= \sum_{j=1}^p (x_j - \bar{x})^2 + \sum_{j=1}^p (y_j - \bar{y})^2 + p(\bar{x} - \bar{y})^2 \\ &\quad - 2 \sum_{j=1}^p (x_j - \bar{x})(y_j - \bar{y}). \end{aligned}$$

The other two terms vanish because $\sum_j (x_j - \bar{x}) = \sum_j (y_j - \bar{y}) = 0$. Substituting $v_x^2 = \sum_{j=1}^p (x_j - \bar{x})^2$ and $v_y^2 = \sum_{j=1}^p (y_j - \bar{y})^2$ and adding and subtracting $-2\sqrt{v_x^2 v_y^2} = -2v_x v_y$, we obtain

$$\begin{aligned} d^2(\mathbf{x}, \mathbf{y}) &= v_x^2 + v_y^2 - 2\sqrt{v_x^2 v_y^2} + p(\bar{x} - \bar{y})^2 + 2v_x v_y \\ &\quad - 2\sqrt{v_x^2 v_y^2} \frac{\sum_{j=1}^p (x_j - \bar{x})(y_j - \bar{y})}{\sqrt{v_x^2 v_y^2}} \\ &= (v_x - v_y)^2 + p(\bar{x} - \bar{y})^2 + 2v_x v_y (1 - r_{xy}) \end{aligned}$$

15.2 (a) Since $\bar{\mathbf{y}}_{AB} = \sum_{i=1}^{n_{AB}} \mathbf{y}_i / n_{AB}$, we have by (15.16),

$$\begin{aligned} \text{SSE}_{AB} &= \sum_{i=1}^{n_{AB}} (\mathbf{y}_i - \bar{\mathbf{y}}_{AB})' (\mathbf{y}_i - \bar{\mathbf{y}}_{AB}) \\ &= \sum_{i=1}^{n_{AB}} \mathbf{y}_i' \mathbf{y}_i - \sum_{i=1}^{n_{AB}} \mathbf{y}_i' \bar{\mathbf{y}}_{AB} - \sum_{i=1}^{n_{AB}} \bar{\mathbf{y}}_{AB}' \mathbf{y}_i \\ &\quad + \sum_{i=1}^{n_{AB}} \bar{\mathbf{y}}_{AB}' \bar{\mathbf{y}}_{AB} \\ &= \sum_{i=1}^{n_{AB}} \mathbf{y}_i' \mathbf{y}_i - n_{AB} \bar{\mathbf{y}}_{AB}' \bar{\mathbf{y}}_{AB} - n_{AB} \bar{\mathbf{y}}_{AB}' \bar{\mathbf{y}}_{AB} \\ &\quad + n_{AB} \bar{\mathbf{y}}_{AB}' \bar{\mathbf{y}}_{AB} \\ &= \sum_{i=1}^{n_{AB}} \mathbf{y}_i' \mathbf{y}_i - n_{AB} \bar{\mathbf{y}}_{AB}' \bar{\mathbf{y}}_{AB}. \end{aligned}$$

Similarly, $\text{SSE}_A = \sum_{i=1}^{n_A} \mathbf{y}'_i \mathbf{y}_i - n_A \bar{\mathbf{y}}'_A \bar{\mathbf{y}}_A$ and $\text{SSE}_B = \sum_{i=1}^{n_B} \mathbf{y}'_i \mathbf{y}_i - n_B \bar{\mathbf{y}}'_B \bar{\mathbf{y}}_B$. Now

$$\begin{aligned} n_{AB} \bar{\mathbf{y}}'_{AB} \bar{\mathbf{y}}_{AB} &= (n_A + n_B) \frac{(n_A \bar{\mathbf{y}}_A + n_B \bar{\mathbf{y}}_B)'}{n_A + n_B} \frac{(n_A \bar{\mathbf{y}}_A + n_B \bar{\mathbf{y}}_B)}{n_A + n_B} \\ &= \frac{n_A^2 \bar{\mathbf{y}}'_A \bar{\mathbf{y}}_A + n_A n_B \bar{\mathbf{y}}'_A \bar{\mathbf{y}}_B + n_A n_B \bar{\mathbf{y}}'_B \bar{\mathbf{y}}_B + n_B^2 \bar{\mathbf{y}}'_B \bar{\mathbf{y}}_B}{n_A + n_B}. \end{aligned}$$

Thus

$$\begin{aligned} \text{SSE}_{AB} - (\text{SSE}_A + \text{SSE}_B) &= \sum_{i=1}^{n_{AB}} \mathbf{y}'_i \mathbf{y}_i - \sum_{i=1}^{n_A} \mathbf{y}'_i \mathbf{y}_i - \sum_{i=1}^{n_B} \mathbf{y}'_i \mathbf{y}_i \\ &\quad + n_A \bar{\mathbf{y}}'_A \bar{\mathbf{y}}_A + n_B \bar{\mathbf{y}}'_B \bar{\mathbf{y}}_B - n_{AB} \bar{\mathbf{y}}'_{AB} \bar{\mathbf{y}}_{AB} \\ &= n_A \bar{\mathbf{y}}'_A \bar{\mathbf{y}}_A + n_B \bar{\mathbf{y}}'_B \bar{\mathbf{y}}_B - n_{AB} \bar{\mathbf{y}}'_{AB} \bar{\mathbf{y}}_{AB}. \end{aligned}$$

Show that when the right side of (15.16) is expanded, it reduces to this same expression [see Problem 15.3(b) below].

- (b)** Multiplying out the right side of (15.16), we have

$$\begin{aligned} &n_A \bar{\mathbf{y}}'_A \bar{\mathbf{y}}_A - n_A \bar{\mathbf{y}}'_A \bar{\mathbf{y}}_{AB} - n_A \bar{\mathbf{y}}'_{AB} \bar{\mathbf{y}}_A + n_A \bar{\mathbf{y}}'_{AB} \bar{\mathbf{y}}_{AB} + n_B \bar{\mathbf{y}}'_B \bar{\mathbf{y}}_B \\ &\quad - n_B \bar{\mathbf{y}}'_B \bar{\mathbf{y}}_{AB} - n_B \bar{\mathbf{y}}'_{AB} \bar{\mathbf{y}}_B + n_B \bar{\mathbf{y}}'_{AB} \bar{\mathbf{y}}_A \\ &= n_A \bar{\mathbf{y}}'_A \bar{\mathbf{y}}_A + n_B \bar{\mathbf{y}}'_B \bar{\mathbf{y}}_B - 2(n_A \bar{\mathbf{y}}'_A + n_B \bar{\mathbf{y}}'_B) \bar{\mathbf{y}}_{AB} + (n_A + n_B) \bar{\mathbf{y}}'_{AB} \bar{\mathbf{y}}_{AB} \\ &= n_A \bar{\mathbf{y}}'_A \bar{\mathbf{y}}_A + n_B \bar{\mathbf{y}}'_B \bar{\mathbf{y}}_B - 2(n_A + n_B) \bar{\mathbf{y}}'_{AB} \bar{\mathbf{y}}_{AB} + (n_A + n_B) \bar{\mathbf{y}}'_{AB} \bar{\mathbf{y}}_{AB} \\ &= n_A \bar{\mathbf{y}}'_A \bar{\mathbf{y}}_A + n_B \bar{\mathbf{y}}'_B \bar{\mathbf{y}}_B - (n_A + n_B) \bar{\mathbf{y}}'_{AB} \bar{\mathbf{y}}_{AB}. \end{aligned}$$

Substitute $\bar{\mathbf{y}}_{AB} = (n_A \bar{\mathbf{y}}_A + n_B \bar{\mathbf{y}}_B)/(n_A + n_B)$.

15.3 (a) Complete linkage.

From Table 15.2, we have

$$D(C, AB) = \frac{1}{2} D(C, A) + \frac{1}{2} D(C, B) + \frac{1}{2} |D(C, A) - D(C, B)| \quad (1)$$

If $D(C, A) > D(C, B)$, then $|D(C, A) - D(C, B)| = D(C, A) - D(C, B)$, and (1) becomes $D(C, AB) = D(C, A)$. If $D(C, A) > D(C, B)$, then $|D(C, A) - D(C, B)| = D(C, B) - D(C, A)$ and equation (1) becomes $D(C, AB) = D(C, B)$. Thus equation (1) can be written as $D(C, AB) = \max[D(C, A), D(C, B)]$, which is equivalent to (15.9), the definition of distance for the complete linkage method.

- (b)** Average linkage. From Table 15.2, we have

$$D(C, AB) = \frac{n_A}{n_A + n_B} D(C, A) + \frac{n_B}{n_A + n_B} D(C, B). \quad (2)$$

By (15.10) equation (2) can be written as

$$\begin{aligned}
 D(C, AB) &= \frac{n_A}{n_A + n_B} \cdot \frac{1}{n_C n_A} \sum_{i=1}^{n_C} \sum_{j=1}^{n_A} d(\mathbf{y}_i, \mathbf{y}_j) \\
 &\quad + \frac{n_B}{n_A + n_B} \cdot \frac{1}{n_C n_B} \sum_{i=1}^{n_C} \sum_{j=1}^{n_B} d(\mathbf{y}_i, \mathbf{y}_j) \\
 &= \frac{1}{n_C(n_A + n_B)} \sum_{i=1}^{n_C} \left[\sum_{j=1}^{n_A} d(\mathbf{y}_i, \mathbf{y}_j) + \sum_{j=1}^{n_B} d(\mathbf{y}_i, \mathbf{y}_j) \right] \\
 &= \frac{1}{n_C n_{AB}} \sum_{i=1}^{n_C} \sum_{j=1}^{n_{AB}} d(\mathbf{y}_i, \mathbf{y}_j),
 \end{aligned}$$

which, by (15.10), is the definition of distance for the average linkage method.

- (c) Substitute $\bar{\mathbf{y}}_{AB} = (n_A \bar{\mathbf{y}}_A + n_B \bar{\mathbf{y}}_B)/(n_A + n_B)$ in the left side of (15.40) in the statement of Problem 15.3(c) and multiply to obtain

$$\begin{aligned}
 \bar{\mathbf{y}}'_C \bar{\mathbf{y}}_C - \frac{2n_A \bar{\mathbf{y}}'_A \bar{\mathbf{y}}_C}{n_A + n_B} + \frac{2n_A n_B \bar{\mathbf{y}}'_A \bar{\mathbf{y}}_B}{(n_A + n_B)^2} - \frac{2n_B \bar{\mathbf{y}}'_B \bar{\mathbf{y}}_C}{n_A + n_B} \\
 + \frac{n_A^2 \bar{\mathbf{y}}'_A \bar{\mathbf{y}}_A}{(n_A + n_B)^2} + \frac{n_B^2 \bar{\mathbf{y}}'_B \bar{\mathbf{y}}_B}{(n_A + n_B)^2}.
 \end{aligned}$$

Similarly, multiply on the right side of (15.40) to obtain the same result.

- (d) Using $n_A = n_B$ in $\bar{\mathbf{y}}_{AB} = (n_A \bar{\mathbf{y}}_A + n_B \bar{\mathbf{y}}_B)/(n_A + n_B)$ in (15.12), we obtain $\mathbf{m}_{AB} = \frac{1}{2}(\bar{\mathbf{y}}_A + \bar{\mathbf{y}}_B)$ in (15.13). Then (15.40) [see part (c)] becomes

$$\begin{aligned}
 (\bar{\mathbf{y}}_C - \mathbf{m}_{AB})'(\bar{\mathbf{y}}_C - \mathbf{m}_{AB}) &= \frac{1}{2}(\bar{\mathbf{y}}_C - \bar{\mathbf{y}}_A)'(\bar{\mathbf{y}}_C - \bar{\mathbf{y}}_A) \\
 &\quad + \frac{1}{2}(\bar{\mathbf{y}}_C - \bar{\mathbf{y}}_B)'(\bar{\mathbf{y}}_C - \bar{\mathbf{y}}_B) - \frac{1}{4}(\bar{\mathbf{y}}_A - \bar{\mathbf{y}}_B)'(\bar{\mathbf{y}}_A - \bar{\mathbf{y}}_B),
 \end{aligned}$$

which matches the parameter values for the median method in Table 15.2.

- (e) By (15.19),

$$(\bar{\mathbf{y}}_A - \bar{\mathbf{y}}_B)'(\bar{\mathbf{y}}_A - \bar{\mathbf{y}}_B) = \frac{n_A + n_B}{n_A n_B} I_{AB},$$

and we have analogous expressions for $(\bar{\mathbf{y}}_C - \bar{\mathbf{y}}_{AB})'(\bar{\mathbf{y}}_C - \bar{\mathbf{y}}_{AB})$, $(\bar{\mathbf{y}}_C - \bar{\mathbf{y}}_A)'(\bar{\mathbf{y}}_C - \bar{\mathbf{y}}_A)$, and $(\bar{\mathbf{y}}_C - \bar{\mathbf{y}}_B)'(\bar{\mathbf{y}}_C - \bar{\mathbf{y}}_B)$. Then (15.40) in part (c) becomes

$$\begin{aligned}
 \frac{n_C + n_{AB}}{n_C n_{AB}} I_{C(AB)} &= \left(\frac{n_A}{n_A + n_B} \right) \left(\frac{n_C + n_A}{n_C n_A} \right) I_{CA} \\
 &\quad + \left(\frac{n_B}{n_A + n_B} \right) \left(\frac{n_C + n_B}{n_C n_B} \right) I_{CB} \\
 &\quad - \left[\frac{n_A n_B}{(n_A + n_B)^2} \right] \left(\frac{n_A + n_B}{n_A n_B} \right) I_{AB} \\
 &= \frac{n_A + n_B}{n_C n_{AB}} I_{AC} + \frac{n_B + n_C}{n_C n_{AB}} I_{CB} - \frac{1}{n_{AB}} I_{AB}.
 \end{aligned}$$

Solve for $I_{C(AB)}$.

15.4 If $\gamma = 0$, then (15.20) becomes

$$D(C, AB) = \alpha_A D(C, A) + \alpha_B D(C, B) + \beta D(A, B). \quad (1)$$

By (15.25), we have $D(A, C) > D(A, B)$ and $D(B, C) > D(A, B)$. Thus, replacing $D(C, A)$ and $D(C, B)$ in equation (1) by $D(A, B)$, we obtain

$$D(C, AB) > \alpha_A D(A, B) + \alpha_B D(A, B) + \beta D(A, B),$$

which is equivalent to (15.26).

$$\begin{aligned}
 \mathbf{15.5} \quad (\mathbf{a}) \quad \bar{\mathbf{v}}_{..} &= \frac{1}{gn} \sum_{i=1}^g \sum_{j=1}^n \mathbf{v}_{ij} = \frac{1}{gn} \sum_{i=1}^g \sum_{j=1}^n (\mathbf{A}\mathbf{y}_{ij} + \mathbf{b}) \\
 &= \frac{1}{gn} \left(\mathbf{A} \sum_{ij} \mathbf{y}_{ij} + gn\mathbf{b} \right) = \mathbf{A} \left(\frac{1}{gn} \sum_{ij} \mathbf{y}_{ij} \right) + \mathbf{b} = \mathbf{A}\bar{\mathbf{y}}_{..} + \mathbf{b}
 \end{aligned}$$

Show similarly that $\bar{\mathbf{v}}_{i..} = \mathbf{A}\bar{\mathbf{y}}_{i..} + \mathbf{b}$. Then by (6.9), we have

$$\begin{aligned}
 \mathbf{H}_v &= n \sum_{i=1}^g (\bar{\mathbf{v}}_{i..} - \bar{\mathbf{v}}_{..})(\bar{\mathbf{v}}_{i..} - \bar{\mathbf{v}}_{..})' \\
 &= n \sum_i [\mathbf{A}\bar{\mathbf{y}}_{i..} + \mathbf{b} - (\mathbf{A}\bar{\mathbf{y}}_{..} + \mathbf{b})][\mathbf{A}\bar{\mathbf{y}}_{i..} + \mathbf{b} - (\mathbf{A}\bar{\mathbf{y}}_{..} + \mathbf{b})]' \\
 &= n \sum_i (\mathbf{A}\bar{\mathbf{y}}_{i..} - \mathbf{A}\bar{\mathbf{y}}_{..})(\mathbf{A}\bar{\mathbf{y}}_{i..} - \mathbf{A}\bar{\mathbf{y}}_{..})' \\
 &= n \sum_i \mathbf{A}(\bar{\mathbf{y}}_{i..} - \bar{\mathbf{y}}_{..})(\bar{\mathbf{y}}_{i..} - \bar{\mathbf{y}}_{..})' \mathbf{A}' \quad [\text{by (2.27)}] \\
 &= n \mathbf{A} \left[\sum_i (\bar{\mathbf{y}}_{i..} - \bar{\mathbf{y}}_{..})(\bar{\mathbf{y}}_{i..} - \bar{\mathbf{y}}_{..})' \right] \mathbf{A}' \quad [\text{by (2.45)}] \\
 &= \mathbf{A}\mathbf{H}_y\mathbf{A}'.
 \end{aligned}$$

Show similarly that $\mathbf{E}_v = \mathbf{A}\mathbf{E}_y\mathbf{A}'$.

- (b) $\text{tr}(\mathbf{E}_v) = \text{tr}(\mathbf{A}\mathbf{E}_y\mathbf{A}') = \text{tr}(\mathbf{A}'\mathbf{A}\mathbf{E}_y) \neq \text{tr}(\mathbf{E}_y)$
- (c) $|\mathbf{E}_v| = |\mathbf{A}\mathbf{E}_y\mathbf{A}'| = |\mathbf{A}||\mathbf{E}_y||\mathbf{A}'| = |\mathbf{A}|^2|\mathbf{E}_y| = c|\mathbf{E}_y|$, where $c > 0$.
Thus minimizing $|\mathbf{E}_v|$ is equivalent to minimizing $|\mathbf{E}_y|$.
- (d)
$$\begin{aligned}\text{tr}(\mathbf{E}_v^{-1}\mathbf{H}_v) &= \text{tr}[(\mathbf{A}\mathbf{E}_y\mathbf{A}')^{-1}(\mathbf{A}\mathbf{H}_y\mathbf{A}')] \\ &= \text{tr}[(\mathbf{A}')^{-1}\mathbf{E}_y^{-1}\mathbf{A}^{-1}\mathbf{A}\mathbf{H}_y\mathbf{A}'] \\ &= \text{tr}[(\mathbf{A}')^{-1}\mathbf{E}_y^{-1}\mathbf{H}_y\mathbf{A}'] \\ &= \text{tr}[\mathbf{A}'(\mathbf{A}')^{-1}\mathbf{E}_y^{-1}\mathbf{H}_y] \\ &= \text{tr}(\mathbf{E}_y^{-1}\mathbf{H}_y)\end{aligned}$$

- 15.6** There are p parameters in each μ_i , $\frac{1}{2}p(p+1)$ unique parameters in each Σ_i , and $g-1$ unique parameters α_i . Thus the total number is

$$\begin{aligned}gp + g[\frac{1}{2}p(p+1)] + g - 1 &= g[p + \frac{1}{2}p(p+1) + 1] - 1 \\ &= \frac{1}{2}g[2p + p^2 + p + 2] - 1 \\ &= \frac{1}{2}g(3p + p^2 + 2) - 1 \\ &= \frac{1}{2}g(p+1)(p+2) - 1.\end{aligned}$$

- 15.7** (a) The two-cluster solution from single linkage puts boy number 20 in one cluster and the other 19 boys in the other cluster.
- (b)-(d) Based on the change in distance, average linkage and the other cluster solutions in parts (c) and (d) clearly indicate two clusters. These solutions generally agree and also correspond to a division into two groups seen in the first principal component in Figure 12.5. The separation of the three apparent outliers from the other 17 observations is less pronounced in the cluster analyses than in Figure 12.5. Note that the scale of the second component in Figure 12.5 is much larger than that of the first component, so the separation of points 9, 12, and 20 from the rest is not as large as it appears in the figure. Of the methods in parts (b), (c), and (d), only flexible beta with $\beta = -.50$ and $-.75$ place points 9, 12, and 20 together in one cluster. All others place 9 and 12 in one of the clusters and 20 in the other.
- 15.8** (a) The distance between centroids of the two clusters is $\sqrt{2994.9} = 54.7$.
- (b) From the dendrogram produced by the average linkage method, the largest change in distance corresponds to a two-cluster solution.
- (c) The discriminant function completely separates the two clusters, with no overlap.
- 15.9** (a) Observation 22 seems to be an outlier, because it forms its own cluster in both the single linkage and average linkage methods. The cluster consisting of observations 2, 21, 24, 26 and 30 is the same in all six methods.

- (b) The discriminant function completely separates the two clusters, with no overlap.

- 15.10** (a) The following five clusters were found using as seeds the five observations that are mutually farthest apart.

Cluster	1	2	3	4	5
Observation(s)	9,15,16, 18,19	1,2,3, 4,5,17	6,7, 8,20	10,11, 12,13	14

In the plot of the first two discriminant functions, observation 14 is relatively far removed from the rest. Clusters 1, 2, and 3 are somewhat closer to each other.

- (b) The following five clusters were found using as seeds the first five observations.

Cluster	1	2	3	4	5
Observation(s)	1,3,4	2	5,17, 18,19	6,7,8, 15,16,20	9,10,11, 12,13,14

The plot of the first two discriminant functions shows a pattern different from that in part (a).

- (c) The following five clusters were found using as seeds the centroids of the five-cluster solution resulting from Ward's method.

Cluster	1	2	3	4	5
Observation(s)	6,7,8,15, 16,20	5,9,17, 18,19	10,11, 12,13	1,2, 3,4	14

The plot of the first two discriminant functions shows a pattern similar to that found in part (a), with observation 14 isolated. The dendrogram shows that Ward's method gives the same five-cluster solution as the *k*-means result.

- (d) The following five clusters were found using the *k*-means method with seeds equal to the centroids of the five clusters from average linkage.

Cluster	1	2	3	4	5
Observation(s)	6,7,8,15, 16,20	1,2,3,4,5, 17,18,19	10,11, 12,13	9	14

The plot of the first two discriminant functions shows a pattern somewhat similar to that in part (a). In the dendrogram for average linkage, observations 9 and 14 are isolated clusters in the five-cluster solution, which is identical to the five-cluster solution using *k*-means clustering with these seeds.

- (e) Observation 14 does not appear as an outlier in the plot of the first two principal components, but it does show up as an outlier in the plot of the

second and third components. The solutions found in parts (a) and (c) seem to agree most with the principal component plots. This suggests that a different number of initial cluster seeds be used.

- (f) The two clustering solutions are identical. The results are given below.

Cluster	1	2	3
Observation(s)	6,7,8,15,16,20	9,10,11,12,13,14	1,2,3,4,5,17,18,19

- (g) The clustering solution is identical to that found in part (e), which indicates that the three-cluster solution is appropriate.

- 15.11** The numbers of clusters obtained from the indicated combinations of k and r are shown in the table below. Note that for each pair of values of k and r , the value of r was increased if necessary for each point until k points were included in the sphere.

k/r	.2	.4	.6	.8	1.0	1.2	1.4	1.6	1.8	2.0
2	10	10	10	10	8	6	4	3	3	2
3	5	5	5	5	5	3	2	2	2	2
4	2	2	2	2	2	2	2	2	2	2
5	1	1	1	1	1	1	1	1	1	1

The maximum value of k that yields a two-cluster solution is 4.

- 15.12** (a) The numbers of clusters obtained from the initial combinations of k and r are shown in the table below. The value of r was variable, as noted in Problem 15.11 above.

k/r	.2	.4	.6	.8	1.0	1.2	1.4	1.6	1.8	2.0
2	3	3	3	3	3	3	3	3	3	3
3	2	2	2	2	2	2	2	2	2	2
4	1	1	1	1	1	1	1	1	1	1

- (b) The plot of the first two discriminant functions for $k = 2$ and $r = 1$ shows the three clusters to be well separated.
- (c) The plot of the first two principal components shows the same groupings as in the plot in part (b).
- (d) The plot of the discriminant function shows wide separation of the two clusters. The clusters do not overlap. The three-cluster solution found in part (b) is given below

Cluster 1	Cluster 2	Cluster 3
Harpers	Rosemaund	Cambridge
Morley	Terrington	Cockle Park
Myerscough	Headley	
Sparsholt	Seale-Hayne	
Sutton Bonington		
Wye		

The two-cluster solution found in part (d) merges clusters two and three of part (b).

CHAPTER 16

$$\begin{aligned} \mathbf{16.1} \quad \mathbf{B} &= \left(\mathbf{I} - \frac{1}{n} \mathbf{J} \right) \mathbf{A} \left(\mathbf{I} - \frac{1}{n} \mathbf{J} \right) \\ &= \mathbf{A} - \frac{1}{n} \mathbf{AJ} - \frac{1}{n} \mathbf{JA} + \frac{1}{n^2} \mathbf{JAJ} \end{aligned} \quad (1)$$

By (2.38),

$$\frac{1}{n} \mathbf{AJ} = \frac{1}{n} \begin{pmatrix} \sum_j a_{1j} \\ \sum_j a_{2j} \\ \vdots \\ \sum_j a_{nj} \end{pmatrix} = \begin{pmatrix} \bar{a}_{1.} \\ \bar{a}_{2.} \\ \vdots \\ \bar{a}_{n.} \end{pmatrix}. \quad (2)$$

Hence,

$$\begin{aligned} \frac{1}{n} \mathbf{AJ} &= \frac{1}{n} \mathbf{A}(\mathbf{j}, \mathbf{j}, \dots, \mathbf{j}) = \left(\frac{1}{n} \mathbf{Aj}, \dots, \frac{1}{n} \mathbf{Aj} \right) \\ &= \begin{pmatrix} \bar{a}_{1.} & \cdots & \bar{a}_{1.} \\ \bar{a}_{2.} & \cdots & \bar{a}_{2.} \\ \vdots & & \vdots \\ \bar{a}_{n.} & \cdots & \bar{a}_{n.} \end{pmatrix}. \end{aligned}$$

Show that

$$\frac{1}{n} \mathbf{JA} = \begin{pmatrix} \bar{a}_{.1} & \bar{a}_{.2} & \cdots & \bar{a}_{.n} \\ \bar{a}_{.1} & \bar{a}_{.2} & \cdots & \bar{a}_{.n} \\ \vdots & \vdots & & \vdots \\ \bar{a}_{.1} & \bar{a}_{.2} & \cdots & \bar{a}_{.n} \end{pmatrix}.$$

Using equation (2), we obtain

$$\frac{1}{n^2} \mathbf{j}' \mathbf{AJ} = \frac{1}{n^2} (1, 1, \dots, 1) \begin{pmatrix} \sum_j a_{1j} \\ \sum_j a_{2j} \\ \vdots \\ \sum_j a_{nj} \end{pmatrix} = \frac{1}{n^2} \sum_{ij} a_{ij} = \bar{a} \dots$$

By (3.63),

$$\frac{1}{n^2} \mathbf{J} \mathbf{A} \mathbf{J} = \frac{1}{n^2} \begin{pmatrix} \mathbf{j}' \mathbf{A} \mathbf{j} & \cdots & \mathbf{j}' \mathbf{A} \mathbf{j} \\ \vdots & & \vdots \\ \mathbf{j}' \mathbf{A} \mathbf{j} & \cdots & \mathbf{j}' \mathbf{A} \mathbf{j} \end{pmatrix} = \begin{pmatrix} \bar{a}_{..} & \cdots & \bar{a}_{..} \\ \vdots & & \vdots \\ \bar{a}_{..} & \cdots & \bar{a}_{..} \end{pmatrix}.$$

Hence the ij th element of equation (1) is $b_{ij} = a_{ij} - \bar{a}_{i\cdot} - \bar{a}_{\cdot j} + \bar{a}_{..}$.

- 16.2 (a)** (Seber 1984, pp. 236-237) The elements of $\mathbf{B} = (b_{ij})$ are defined as $b_{ij} = a_{ij} - \bar{a}_{i\cdot} - \bar{a}_{\cdot j} + \bar{a}_{..}$, where $a_{ij} = -\frac{1}{2}\delta_{ij}^2$. Thus

$$\begin{aligned} -2a_{ij} &= \delta_{ij}^2 = (\mathbf{z}_i - \mathbf{z}_j)'(\mathbf{z}_i - \mathbf{z}_j) \\ &= \mathbf{z}'_i \mathbf{z}_i + \mathbf{z}'_j \mathbf{z}_j - 2\mathbf{z}'_i \mathbf{z}_j. \end{aligned}$$

Then

$$\begin{aligned} -2\bar{a}_{i\cdot} &= \frac{1}{n} \sum_{j=1}^n (-2a_{ij}) = \frac{1}{n} \sum_j (\mathbf{z}'_i \mathbf{z}_i + \mathbf{z}'_j \mathbf{z}_j - 2\mathbf{z}'_i \mathbf{z}_j) \\ &= \mathbf{z}'_i \mathbf{z}_i + \frac{1}{n} \sum_j \mathbf{z}'_j \mathbf{z}_j - \frac{2}{n} \mathbf{z}'_i \sum_j \mathbf{z}_j \\ &= \mathbf{z}'_i \mathbf{z}_i + \frac{1}{n} \sum_j \mathbf{z}'_j \mathbf{z}_j - 2\mathbf{z}'_i \bar{\mathbf{z}}. \end{aligned}$$

Similarly, show that

$$\begin{aligned} -2\bar{a}_{\cdot j} &= \mathbf{z}'_j \mathbf{z}_j + \frac{1}{n} \sum_i \mathbf{z}'_i \mathbf{z}_i - 2\bar{\mathbf{z}}' \mathbf{z}_j, \\ -2\bar{a}_{..} &= \frac{2}{n} \sum_i \mathbf{z}'_i \mathbf{z}_i - 2\bar{\mathbf{z}}' \bar{\mathbf{z}}. \end{aligned}$$

Solve for a_{ij} , $\bar{a}_{i\cdot}$, $\bar{a}_{\cdot j}$, and $\bar{a}_{..}$ and substitute into $b_{ij} = a_{ij} - \bar{a}_{i\cdot} - \bar{a}_{\cdot j} + \bar{a}_{..}$ to obtain $b_{ij} = \mathbf{z}'_i \mathbf{z}_j - \mathbf{z}'_i \bar{\mathbf{z}} - \bar{\mathbf{z}}' \mathbf{z}_j + \bar{\mathbf{z}}' \bar{\mathbf{z}}$, which can be factored as $b_{ij} = (\mathbf{z}_i - \bar{\mathbf{z}})'(\mathbf{z}_j - \bar{\mathbf{z}})$. Hence

$$\begin{aligned} \mathbf{B} &= \begin{pmatrix} (\mathbf{z}_1 - \bar{\mathbf{z}})'(\mathbf{z}_1 - \bar{\mathbf{z}}) & \cdots & (\mathbf{z}_1 - \bar{\mathbf{z}})'(\mathbf{z}_n - \bar{\mathbf{z}}) \\ \vdots & & \vdots \\ (\mathbf{z}_n - \bar{\mathbf{z}})'(\mathbf{z}_1 - \bar{\mathbf{z}}) & \cdots & (\mathbf{z}_n - \bar{\mathbf{z}})'(\mathbf{z}_n - \bar{\mathbf{z}}) \end{pmatrix} \\ &= \begin{pmatrix} (\mathbf{z}_1 - \bar{\mathbf{z}})' \\ \vdots \\ (\mathbf{z}_n - \bar{\mathbf{z}})' \end{pmatrix} (\mathbf{z}_1 - \bar{\mathbf{z}}, \dots, \mathbf{z}_n - \bar{\mathbf{z}}) \\ &= \mathbf{Z}_c \mathbf{Z}'_c \quad [\text{see (10.13)}]. \end{aligned}$$

Thus \mathbf{B} is positive semidefinite (see Section 2.7).

- (b) If \mathbf{B} is positive semidefinite of rank q , then by (2.109) and Section 2.11.4, \mathbf{B} can be expressed in the form $\mathbf{B} = \mathbf{V}\Lambda\mathbf{V}'$, where $\mathbf{V} = (\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n)$ is an orthogonal matrix of eigenvectors of \mathbf{B} , and Λ is a diagonal matrix of eigenvalues, q of which are positive, with the rest equal to zero. Letting Λ_1 be the $q \times q$ upper left hand block of Λ with positive eigenvalues and $\mathbf{V}_1 = (\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_q)$ be the $n \times q$ matrix with the corresponding eigenvectors, we can write $\mathbf{B} = \mathbf{V}\Lambda\mathbf{V}'$ as

$$\begin{aligned}\mathbf{B} &= (\mathbf{V}_1, \mathbf{V}_2) \begin{pmatrix} \Lambda_1 & \mathbf{O} \\ \mathbf{O} & \mathbf{O} \end{pmatrix} \begin{pmatrix} \mathbf{V}_1' \\ \mathbf{V}_2' \end{pmatrix} \\ &= \mathbf{V}_1 \Lambda_1 \mathbf{V}_1' = \mathbf{V}_1 \Lambda_1^{1/2} \Lambda_1^{1/2} \mathbf{V}_1' \\ &= \mathbf{Z} \mathbf{Z}',\end{aligned}\quad (1)$$

where the $n \times q$ matrix \mathbf{Z} is

$$\begin{aligned}\mathbf{Z} &= \mathbf{V}_1 \Lambda_1^{1/2} = (\sqrt{\lambda_1} \mathbf{v}_1, \sqrt{\lambda_2} \mathbf{v}_2, \dots, \sqrt{\lambda_q} \mathbf{v}_q) \\ &= \begin{pmatrix} \mathbf{z}_1' \\ \mathbf{z}_2' \\ \vdots \\ \mathbf{z}_n' \end{pmatrix}.\end{aligned}$$

To show that $(\mathbf{z}_i - \mathbf{z}_j)'(\mathbf{z}_i - \mathbf{z}_j)$ is equal to δ_{ij}^2 , we can proceed as follows:

$$(\mathbf{z}_i - \mathbf{z}_j)'(\mathbf{z}_i - \mathbf{z}_j) = \mathbf{z}_i' \mathbf{z}_i + \mathbf{z}_j' \mathbf{z}_j - 2\mathbf{z}_i' \mathbf{z}_j. \quad (2)$$

By equation (1), we have

$$\begin{aligned}\mathbf{B} &= \mathbf{Z} \mathbf{Z}' = \begin{pmatrix} \mathbf{z}_1' \\ \mathbf{z}_2' \\ \vdots \\ \mathbf{z}_n' \end{pmatrix} (\mathbf{z}_1, \mathbf{z}_2, \dots, \mathbf{z}_n) \\ &= \begin{pmatrix} \mathbf{z}_1' \mathbf{z}_1 & \mathbf{z}_1' \mathbf{z}_2 & \cdots & \mathbf{z}_1' \mathbf{z}_n \\ \mathbf{z}_2' \mathbf{z}_1 & \mathbf{z}_2' \mathbf{z}_2 & \cdots & \mathbf{z}_2' \mathbf{z}_n \\ \vdots & \vdots & & \vdots \\ \mathbf{z}_n' \mathbf{z}_1 & \mathbf{z}_n' \mathbf{z}_2 & \cdots & \mathbf{z}_n' \mathbf{z}_n \end{pmatrix}.\end{aligned}$$

Hence equation (2) becomes

$$\begin{aligned}(\mathbf{z}_i - \mathbf{z}_j)'(\mathbf{z}_i - \mathbf{z}_j) &= \mathbf{z}_i' \mathbf{z}_i + \mathbf{z}_j' \mathbf{z}_j - 2\mathbf{z}_i' \mathbf{z}_j \\ &= b_{ii} + b_{jj} - 2b_{ij}.\end{aligned}\quad (3)$$

Show that substituting $b_{ij} = a_{ij} - \bar{a}_{i\cdot} - \bar{a}_{\cdot j} + \bar{a}_{\cdot\cdot}$ into equation (3) leads to

$$(\mathbf{z}_i - \mathbf{z}_j)'(\mathbf{z}_i - \mathbf{z}_j) = a_{ii} + a_{jj} - 2a_{ij} + \bar{a}_{i\cdot} - \bar{a}_{\cdot j} + \bar{a}_{\cdot\cdot}.$$

Show that the symmetry of \mathbf{A} implies $\bar{a}_{i.} = \bar{a}_{.i}$ and $\bar{a}_{.j} - \bar{a}_{j.}$ Hence,

$$(\mathbf{z}_i - \mathbf{z}_j)'(\mathbf{z}_i - \mathbf{z}_j) = a_{ii} + a_{jj} - 2a_{ij} = -2a_{ij} = \delta_{ij}^2,$$

since $a_{ii} = -\frac{1}{2}\delta_{ii}^2 = 0$ and $-2a_{ij} = \delta_{ij}^2$.

$$\begin{aligned} \mathbf{16.3 (a)} \quad \mathbf{r} &= \sum_{j=1}^b p_{.j} \mathbf{c}_j = \sum_{j=1}^b p_{.j} \left(\frac{p_{1j}}{p_{.j}}, \frac{p_{2j}}{p_{.j}}, \dots, \frac{p_{aj}}{p_{.j}} \right)' \\ &= \sum_{j=1}^b (p_{1j}, p_{2j}, \dots, p_{aj})' \quad [\text{by (2.61)}] \\ &= (\sum_j p_{1j}, \sum_j p_{2j}, \dots, \sum_j p_{aj})' \\ &= (p_{.1}, p_{.2}, \dots, p_{.a})' \end{aligned}$$

$$\begin{aligned} \mathbf{(b)} \quad \mathbf{c}' &= \sum_{i=1}^a p_{i.} \mathbf{r}'_i = \sum_{i=1}^a p_{i.} \left(\frac{p_{i1}}{p_{i.}}, \frac{p_{i2}}{p_{i.}}, \dots, \frac{p_{ib}}{p_{i.}} \right)' \\ &= \sum_{i=1}^a (p_{i1}, p_{i2}, \dots, p_{ib}) \quad [\text{by (2.61)}] \\ &= (\sum_i p_{i1}, \sum_i p_{i2}, \dots, \sum_i p_{ib}) \\ &= (p_{.1}, p_{.2}, \dots, p_{.b}) \end{aligned}$$

$$\begin{aligned} \mathbf{16.4} \quad \mathbf{j}'\mathbf{r} &= \sum_{i=1}^a p_{i.} = \sum_{i=1}^a \sum_{j=1}^b p_{ij} = \sum_{ij} n_{ij}/n = n/n = 1, \\ \mathbf{c}'\mathbf{j} &= \sum_{j=1}^b p_{.j} = \sum_j n_{.j}/n = \sum_j \sum_i n_{ij}/n = n/n \end{aligned}$$

16.5 By (16.8), (16.9), and (16.10), $p_{ij} = n_{ij}/n$, $p_{i.} = n_{i.}/n$, and $p_{.j} = n_{.j}/n$. Substituting these into (16.25), we obtain

$$\begin{aligned} \chi^2 &= \sum_{ij} \frac{n \left(\frac{n_{ij}}{n} - \frac{n_{i.} n_{.j}}{n^2} \right)^2}{\frac{n_{i.} n_{.j}}{n^2}} = \sum_{ij} \frac{n \left[\frac{1}{n} (n_{ij} - \frac{n_{i.} n_{.j}}{n}) \right]^2}{\frac{n_{i.} n_{.j}}{n^2}} \\ &= \sum_{ij} \frac{\frac{n}{n^2} (n_{ij} - \frac{n_{i.} n_{.j}}{n})^2}{\frac{n_{i.} n_{.j}}{n^2}} = \sum_{ij} \frac{(n_{ij} - \frac{n_{i.} n_{.j}}{n})^2}{\frac{n_{i.} n_{.j}}{n}}. \end{aligned}$$

16.6 (a) Multiplying numerator and denominator of (16.25) by $p_{i.}$, we obtain

$$\begin{aligned} \chi^2 &= \sum_i n \sum_j \frac{p_{i.}}{p_{i.}^2 p_{.j}} (p_{ij} - p_{i.} p_{.j})^2 \\ &= \sum_i n p_{i.} \sum_j \frac{1}{p_{.j}} \left[\frac{1}{p_{i.}} (p_{ij} - p_{i.} p_{.j}) \right]^2 \\ &= \sum_i n p_{i.} \sum_j \left(\frac{p_{ij}}{p_{i.}} - p_{.j} \right)^2 / p_{.j}. \end{aligned}$$

16.7 (a) By (16.29), (16.10), (16.12), and (16.18), we obtain

$$\begin{aligned}\chi^2 &= \sum_i n p_i (\mathbf{r}_i - \mathbf{c})' \mathbf{D}_c^{-1} (\mathbf{r}_i - \mathbf{c}) \\&= \sum_i n p_i \left(\frac{p_{i1}}{p_{i.}} - p_{.1}, \dots, \frac{p_{ib}}{p_{i.}} - p_{.b} \right) \begin{pmatrix} p_{.1} & \cdots & 0 \\ \vdots & & \vdots \\ 0 & \cdots & p_{.b} \end{pmatrix}^{-1} \begin{pmatrix} \frac{p_{i1}}{p_{i.}} - p_{.1} \\ \vdots \\ \frac{p_{ib}}{p_{i.}} - p_{.b} \end{pmatrix} \\&= \sum_i n p_i \left(\frac{\frac{p_{i1}}{p_{i.}} - p_{.1}}{p_{.1}}, \dots, \frac{\frac{p_{ib}}{p_{i.}} - p_{.b}}{p_{.b}} \right) \begin{pmatrix} \frac{p_{i1}}{p_{i.}} - p_{.1} \\ \vdots \\ \frac{p_{ib}}{p_{i.}} - p_{.b} \end{pmatrix}.\end{aligned}$$

16.8 (a) By (16.9) $\mathbf{r} = \mathbf{Pj}$. Then $\mathbf{D}_r^{-1}\mathbf{r} = \mathbf{D}_r^{-1}\mathbf{Pj} = \mathbf{Rj}$ by (16.15). By (16.13) $\mathbf{r}'_i \mathbf{j} = 1$, and therefore $\mathbf{Rj} = \mathbf{j}$. Now

$$\mathbf{D}_r^{-1}(\mathbf{P} - \mathbf{rc}') = \mathbf{D}_r^{-1}\mathbf{P} - \mathbf{D}_r^{-1}\mathbf{rc}' = \mathbf{R} - \mathbf{Rjc}' = \mathbf{R} - \mathbf{jc}'.$$

16.9 By (16.49), $\mathbf{z}'_i = \mathbf{y}'_i \mathbf{A}$ (ignoring the centering on \mathbf{y}_i). Thus the squared Euclidean distance can be written as

$$\begin{aligned}(\mathbf{z}_i - \mathbf{z}_k)'(\mathbf{z}_i - \mathbf{z}_k) &= (\mathbf{z}'_i - \mathbf{z}'_k)(\mathbf{z}_i - \mathbf{z}_k) \\&= (\mathbf{y}'_i \mathbf{A} - \mathbf{y}'_k \mathbf{A})(\mathbf{A}' \mathbf{y}_i - \mathbf{A}' \mathbf{y}_k) \\&= (\mathbf{y}'_i - \mathbf{y}'_k) \mathbf{A} \mathbf{A}' (\mathbf{y}_i - \mathbf{y}_k) \\&= (\mathbf{y}_i - \mathbf{y}_k)'(\mathbf{y}_i - \mathbf{y}_k),\end{aligned}$$

since \mathbf{A} is orthogonal.

16.10 (a) From $\mathbf{Y}_c \mathbf{V} = \mathbf{U} \Lambda$ in (16.55), we have $\mathbf{Y}_c \mathbf{V} \Lambda^{-1} = \mathbf{U}$. Then

$$\begin{aligned}\mathbf{U} \mathbf{U}' &= \mathbf{Y}_c \mathbf{V} \Lambda^{-1} \Lambda^{-1} \mathbf{V}' \mathbf{Y}'_c \\&= \mathbf{Y}_c \mathbf{V} (\Lambda^{-1})^2 \mathbf{V}' \mathbf{Y}'_c.\end{aligned}\tag{1}$$

Since $(\Lambda^{-1})^2 = \text{diag}(1/\lambda_1^2, 1/\lambda_2^2, \dots, 1/\lambda_p^2)$, where the λ_i^2 's are eigenvalues of $\mathbf{Y}'_c \mathbf{Y}_c$, the matrix $(\Lambda^{-1})^2$ contains eigenvalues of $(\mathbf{Y}'_c \mathbf{Y}_c)^{-1} = [(n-1)\mathbf{S}]^{-1} = \mathbf{S}^{-1}/(n-1)$ [see (2.115) and (2.116)]. The matrix \mathbf{V} contains eigenvectors of $\mathbf{Y}'_c \mathbf{Y}_c$ and thereby of $(\mathbf{Y}'_c \mathbf{Y}_c)^{-1}$ (see Section 2.11.9). Hence we recognize $\mathbf{V}(\Lambda^{-1})^2 \mathbf{V}'$ as the spectral decomposition of $(\mathbf{Y}'_c \mathbf{Y}_c)^{-1}$ [see (2.109), (2.115), and (2.116)]. Therefore, equation (1) can be written as

$$\begin{aligned}\mathbf{U} \mathbf{U}' &= \mathbf{Y}_c \mathbf{V} (\Lambda^{-1})^2 \mathbf{V}' \mathbf{Y}'_c = \mathbf{Y}_c (\mathbf{Y}'_c \mathbf{Y}_c)^{-1} \mathbf{Y}'_c \\&= \mathbf{Y}_c \mathbf{S}^{-1} \mathbf{Y}'_c / (n-1).\end{aligned}$$

- (b) If $\mathbf{H} = \mathbf{V}\Lambda$, then $\mathbf{HH}' = \mathbf{V}\Lambda\Lambda'\mathbf{V}' = \mathbf{V}\Lambda^2\mathbf{V}'$. The diagonal matrix Λ^2 contains the eigenvalues λ_i^2 of the matrix $\mathbf{Y}_c'\mathbf{Y}_c$. Thus by (2.115), $\mathbf{V}\Lambda^2\mathbf{V}'$ is the spectral decomposition of $\mathbf{Y}_c'\mathbf{Y}_c$, and

$$\mathbf{HH}' = \mathbf{V}\Lambda^2\mathbf{V}' = \mathbf{Y}_c'\mathbf{Y}_c = (n - 1)\mathbf{S}.$$

- 16.11** By (16.64), (3.63), and (3.64) (ignoring $n - 1$ and assuming the \mathbf{y}_i 's are centered),

$$\begin{aligned} (\mathbf{u}_i - \mathbf{u}_k)'(\mathbf{u}_i - \mathbf{u}_k) &= \mathbf{u}_i' \mathbf{u}_i + \mathbf{u}_k' \mathbf{u}_k - 2\mathbf{u}_i' \mathbf{u}_k \\ &= \mathbf{y}_i' \mathbf{S}^{-1} \mathbf{y}_i + \mathbf{y}_k' \mathbf{S}^{-1} \mathbf{y}_k - 2\mathbf{y}_i' \mathbf{S}^{-1} \mathbf{y}_k \\ &= (\mathbf{y}_i - \mathbf{y}_k)' \mathbf{S}^{-1} (\mathbf{y}_i - \mathbf{y}_k). \end{aligned}$$

- 16.12 (a)** The first ten rows and columns of the matrix \mathbf{B} are as follows:

129849	-26801	-88750	-53847	-59118	43583	-73877	81571	112101	80909
-26801	2310	17029	11125	14394	-11076	18149	-18662	-21852	-16306
-88750	17029	65973	32378	31044	-31085	68156	-56671	-79481	-54135
-53847	11125	32378	27683	31808	-18550	30003	-34154	-46882	-32096
-59118	14394	31044	31808	38141	-19161	27269	-37147	-51673	-34687
45383	-11076	-31085	-18550	-19161	14741	-33620	27054	45347	29211
-73877	18149	68156	30003	27269	-33620	76423	-45782	-86804	-58650
81571	-18662	-56671	-34154	-37147	27054	-45782	49169	75557	50169
112101	-21852	-79481	-46882	-51673	45347	-86804	75557	119634	81258
80909	-16306	-54135	-32096	-34687	29211	-58650	50169	81258	53286

- (b) The first two columns of the matrix \mathbf{Z} are given by:

City	z_1	z_2	City	z_1	z_2
A	354.1	-10.2	M	391.6	47.5
B	-77.1	25.0	N	21.0	-44.7
C	-238.2	-75.7	O	9.8	30.9
D	-154.9	65.9	P	-173.8	-78.5
E	-163.2	72.2	Q	6.3	17.1
F	126.0	24.9	R	117.0	-48.0
G	-228.8	-149.4	S	-102.3	-170.2
H	223.9	1.5	T	-53.2	-27.3
I	337.7	44.8	U	-315.2	190.9
J	226.7	34.7	V	-255.7	140.2
K	-33.4	22.3	W	-19.3	-34.3
L	1.1	-79.4			

- (c) The metric multidimensional scaling plot shows the relative positions of the cities.

- 16.13 (a)** The multidimensional scaling plot shows two clusters, one for positive values of the first dimension, and one for negative values. The two clusters can be interpreted as comfort (positive values) and discomfort (negative values). Hence, the axis of the first dimension can be interpreted as the level of comfort.
- (b) The dendrogram for Ward's method clearly shows two clusters, the same as in part (a).

- 16.14 (a)** The initial configuration of points will vary. One example is as follows:

y_1	y_2	y_3	y_4	y_5	y_6
1.458	.769	-1.350	.456	-1.610	1.827
-.598	-1.069	-2.667	.458	.416	1.094
-1.777	-.409	.369	.655	-.058	1.177
.071	.361	1.157	-.154	.343	-.417
-.060	1.361	.743	1.436	.332	-.894
-.757	-.432	-.545	.233	.646	-.102
-1.971	-.492	-.461	.078	1.441	.039
-1.560	-.173	.657	-.528	1.001	1.030
-.597	.814	-.898	.283	-.355	-1.115
1.449	-.942	.867	-.922	.833	1.196
-1.809	-.093	-1.762	-.533	-1.136	-.226
1.067	.199	.978	.884	-1.060	-.800

- (b)** Answers will vary. For the seeds given in part (a), STRESS = 0.0266.
- (c)** Answers will vary. The plot of STRESS versus k for one solution showed that two dimensions should be retained. The nonmetric MDS plot showed that Franco, Mussolini, and Hitler lie closely together, as well as Churchill and De Gaulle, and Eisenhower and Truman.
- (d)** Answers will vary. One solution gave results similar to part (c).
- (e)** Answers will vary. One solution showed three dimensions. A plot of two dimensions showed Mussolini and Franco together in the center with the others forming a circle around them almost equally spaced.
- (f)** Answers will vary. One solution was similar to that in part (c).

- 16.15 (a)** The correspondence matrix \mathbf{P} is found by dividing each element of Table 16.16 by $n = 1281$ to obtain the following:

Death Birth	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec	Total
Jan	.007	.011	.009	.011	.007	.009	.008	.012	.007	.009	.009	.010	.108
Feb	.010	.005	.005	.006	.007	.004	.003	.004	.005	.009	.001	.010	.069
Mar	.009	.011	.007	.005	.013	.008	.007	.008	.007	.002	.010	.007	.094
Apr	.005	.009	.008	.005	.007	.009	.003	.009	.003	.007	.006	.009	.080
May	.006	.005	.009	.005	.003	.009	.007	.007	.009	.005	.007	.003	.075
Jun	.011	.004	.004	.005	.010	.004	.005	.003	.006	.007	.005	.004	.069
Jul	.009	.008	.010	.003	.004	.009	.005	.005	.003	.008	.003	.006	.073
Aug	.005	.005	.009	.010	.008	.007	.002	.006	.006	.006	.006	.009	.081
Sep	.005	.009	.009	.008	.008	.009	.003	.006	.009	.005	.006	.005	.083
Oct	.012	.006	.009	.007	.005	.008	.009	.006	.007	.006	.005	.005	.087
Nov	.005	.007	.012	.008	.009	.008	.005	.008	.005	.008	.007	.005	.087
Dec	.005	.014	.007	.009	.011	.006	.007	.007	.008	.005	.008	.006	.092
Total	.092	.094	.096	.084	.092	.088	.066	.080	.077	.075	.074	.081	1.000

(b) The \mathbf{R} matrix is given by

$$\mathbf{R} = \begin{pmatrix} .07 & .11 & .12 & .11 & .07 & .04 & .11 & .10 & .09 & .08 & .09 & .04 \\ .13 & .08 & .12 & .07 & .07 & .03 & .09 & .11 & .10 & .08 & .08 & .08 \\ .09 & .08 & .07 & .15 & .05 & .08 & .07 & .08 & .12 & .08 & .05 & .08 \\ .09 & .06 & .15 & .08 & .15 & .04 & .06 & .07 & .10 & .01 & .12 & .08 \\ .10 & .11 & .09 & .10 & .07 & .07 & .08 & .09 & .07 & .08 & .08 & .07 \\ .04 & .06 & .09 & .11 & .13 & .07 & .12 & .14 & .05 & .04 & .11 & .04 \\ .08 & .04 & .06 & .06 & .16 & .08 & .06 & .06 & .15 & .08 & .10 & .09 \\ .06 & .08 & .07 & .12 & .10 & .07 & .08 & .07 & .14 & .11 & .02 & .07 \\ .07 & .09 & .04 & .06 & .08 & .09 & .13 & .11 & .04 & .09 & .06 & .11 \\ .09 & .09 & .05 & .08 & .06 & .06 & .09 & .14 & .10 & .08 & .09 & .06 \\ .08 & .07 & .06 & .07 & .14 & .11 & .09 & .10 & .06 & .06 & .07 & .08 \\ .09 & .08 & .07 & .11 & .07 & .04 & .10 & .10 & .09 & .08 & .06 & .11 \end{pmatrix},$$

and the \mathbf{C} matrix is given by

$$\mathbf{C} = \begin{pmatrix} .07 & .11 & .12 & .09 & .06 & .05 & .10 & .08 & .08 & .08 & .09 & .04 \\ .12 & .08 & .12 & .06 & .04 & .08 & .09 & .08 & .08 & .08 & .08 & .08 \\ .10 & .09 & .08 & .15 & .05 & .11 & .07 & .07 & .12 & .11 & .06 & .10 \\ .07 & .05 & .13 & .06 & .11 & .05 & .04 & .05 & .08 & .01 & .11 & .07 \\ .13 & .15 & .13 & .12 & .08 & .12 & .10 & .10 & .08 & .12 & .11 & .09 \\ .04 & .06 & .08 & .08 & .10 & .08 & .10 & .11 & .04 & .04 & .10 & .04 \\ .07 & .04 & .05 & .04 & .12 & .08 & .04 & .04 & .11 & .07 & .09 & .08 \\ .07 & .10 & .09 & .12 & .10 & .11 & .09 & .07 & .14 & .14 & .02 & .09 \\ .07 & .09 & .04 & .05 & .07 & .11 & .11 & .09 & .03 & .09 & .10 & .07 \\ .08 & .08 & .07 & .07 & .14 & .14 & .09 & .09 & .06 & .07 & .08 & .09 \\ .09 & .08 & .07 & .10 & .06 & .05 & .10 & .09 & .08 & .08 & .06 & .12 \end{pmatrix}.$$

- (c)** The chi-square statistic is 117.7742 with 121 degrees of freedom, which gives a p -value of .5660. The two variables appear to be independent.
- (d)** In the correspondence plot, the following associations are seen:
- {November births, June deaths}, {March deaths, April deaths, January births}, {September births, February deaths}, {August births, April births}, {May deaths, September deaths, May births}.

- 16.16 (a)** The correspondence matrix \mathbf{P} is found by dividing each element of Table 16.17 by 8193 to obtain the following:

Part of Country	Burglary	Fraud	Vandalism	Total
Oslo area	.048	.300	.215	.563
Mid-Norway	.018	.019	.112	.148
North Norway	.085	.040	.164	.289
Total	.151	.358	.491	1.000

- (b)** The \mathbf{R} matrix is given by

Part of Country	Burglary	Fraud	Vandalism
Oslo area	.086	.533	.381
Mid-Norway	.121	.126	.753
North Norway	.293	.138	.569

and the \mathbf{C} matrix is given by

Part of Country	Burglary	Fraud	Vandalism
Oslo area	.320	.837	.437
Mid-Norway	.119	.052	.228
North Norway	.561	.111	.335

- (c)** The chi-square statistic is 1662.6 with 4 degrees of freedom, which gives a p -value less than .0001. The two variables are dependent.
- (d)** In the correspondence plot, North Norway is associated with burglaries, Oslo is associated with fraud, and mid-Norway is associated with vandalism.

16.17 (a) The Burt matrix is given below.

No	5254	0	564	3408	1282	1830	3424	2466	2788	2190	3064	686	2666	1902
Yes	0	165	105	42	18	73	92	37	128	4	125	26	63	76
High dust	564	105	669	0	0	402	267	62	607	218	451	87	359	223
Low dust	3408	42	0	3450	0	1056	2394	1642	1808	1446	2004	480	1684	1286
Medium dust	1282	18	0	0	1300	445	855	799	501	566	734	145	686	469
Race—other	1830	73	402	1056	445	1930	0	932	971	799	1104	108	1658	137
White	3424	92	267	2394	855	0	3516	1571	1945	1431	2085	604	1071	1841
Female	2466	37	62	1642	799	932	1571	2503	0	1373	1130	266	1421	816
Male	2788	128	607	1808	501	971	1945	0	2916	857	2059	446	1308	1162
Nonsmoker	2190	40	218	1446	566	799	1431	1373	857	2230	0	231	1142	857
Smoker	3064	125	451	2004	734	1104	2085	1130	2059	0	3189	481	1587	1121
10–20	686	26	87	480	145	108	604	266	446	231	481	712	0	0
≤ 10	2666	63	359	1684	686	1658	1071	1421	1308	1142	1587	0	2729	0
≥ 20	1902	76	223	1286	469	137	1841	816	1162	857	1121	0	0	1978

(b) The column coordinates for the plot are given by

Variables	y_1	y_2
No	-.032	-.087
Yes	1.013	2.761
High dust	1.072	1.648
Low dust	-.209	-.107
Medium dust	.003	-.564
Race—other	1.184	-.153
White	-.641	.083
Female	.007	-.791
Male	-.006	.679
Nonsmoker	-.036	-.592
Smoker	.025	.414
10–20	-.605	.535
≤ 10	.789	-.300
≥ 20	-.871	.221

- (c)** Some associations seen in the plot are {byssinosis-yes, high dust}, {female, nonsmoker, medium dust}, {smoker, male}.

- 16.18 (a)** The two-dimensional coordinates of the observation points and variable points are given below.

Observation Points

Name	Coordinate 1	Coordinate 2
Albania	14.102	-1.322
Austria	-5.461	1.548
Belgium	-6.077	-1.479
Bulgaria	26.116	3.319
Czech.	3.317	-2.092
Denmark	-13.861	1.374
E. Germany	-4.902	-8.360
Finland	-12.262	11.290
France	-6.345	.672
Greece	9.036	3.033
Hungary	10.805	-2.363
Ireland	-11.857	5.312
Italy	6.309	-1.314
Netherlands	-11.809	2.133
Norway	-11.005	-.077
Poland	2.526	2.999
Portugal	.784	-16.753
Romania	19.067	2.591
Spain	1.923	-10.483
Sweden	-14.842	.726
Switzerland	-9.068	4.000
UK	-9.311	.698
USSR	10.586	4.355
W. Germany	-13.514	-3.353
Yugoslavia	25.742	3.548

Variable Points

Name	Coordinate 1	Coordinate 2
R_MEAT	-.151	.133
W_MEAT	-.129	.043
EGGS	-.067	.021
MILK	-.425	.831
FISH	-.127	-.292
CEREALS	.861	.406
STARCHY	-.067	-.076
NUTS	.114	-.070
FRUIT_VEG	.020	-.169

In the biplot, the arrows for variables are too short to pass through the points for observations.

- (b) The two-dimensional coordinates of the observation points and variable points are given below.

Observation Points

Name	Coordinate 1	Coordinate 2
Albania	.231	-.049
Austria	-.089	.057
Belgium	-.100	-.055
Bulgaria	.428	.122
Czech.	.054	-.077
Denmark	-.227	.051
E. Germany	-.080	-.308
Finland	-.201	.416
France	-.104	.025
Greece	.148	.112
Hungary	.177	-.087
Ireland	-.194	.196
Italy	.103	-.048
Netherlands	-.193	.079
Norway	-.180	-.003
Poland	.041	.110
Portugal	.013	-.617
Romania	.312	.095
Spain	.032	-.386
Sweden	-.243	.027
Switzerland	-.149	.147
UK	-.153	.026
USSR	.173	.160
W. Germany	-.221	-.124
Yugoslavia	.422	.131

Variable Points

Name	Coordinate 1	Coordinate 2
R_MEAT	-9.196	3.602
W_MEAT	-7.904	1.179
EGGS	-4.106	.569
MILK	-25.964	22.552
FISH	-7.750	-7.934
CEREALS	52.545	11.025
STARCHY	-4.080	-2.064
NUTS	6.953	-1.902
FRUIT_VEG	1.235	-4.593

In the biplot, the observation points are tightly clustered around the point (0,0), making them difficult to distinguish, whereas variable points are easily discerned. Red meats, white meats, and milk are highly positively

correlated. These three variables are negatively correlated with nuts and fruit_veg.

- (c) The two-dimensional coordinates of the observation points and variable points are given below.

Observation Points

Name	Coordinate 1	Coordinate 2
Albania	1.805	-.254
Austria	-.699	.297
Belgium	-.778	-.284
Bulgaria	3.343	.637
Czech.	.425	-.402
Denmark	-1.774	.264
E. Germany	-.627	-1.605
Finland	-1.570	2.167
France	-.812	.129
Greece	1.157	.582
Hungary	1.383	-.454
Ireland	-1.518	1.020
Italy	.808	-.252
Netherlands	-1.511	.409
Norway	-1.409	-.015
Poland	.323	.576
Portugal	.100	-3.216
Romania	2.441	.497
Spain	.246	-2.012
Sweden	-1.900	.139
Switzerland	-1.161	.768
UK	-1.192	.134
USSR	1.355	.836
W. Germany	-1.730	-.644
Yugoslavia	3.295	.681

Variable Points

Name	Coordinate 1	Coordinate 2
R_MEAT	-1.177	.691
W_MEAT	-1.012	.226
EGGS	-.526	.109
MILK	-3.323	4.329
FISH	-.992	-1.523
CEREALS	6.726	2.116
STARCHY	-.522	-.396
NUTS	.890	-.365
FRUIT_VEG	.158	-.882

In the biplot, the variable points and observation points are both well spaced. Finland scored high on the milk variable. Yugoslavia and Bulgaria scored high on the cereal variable. Spain and Portugal scored highest on the fish and frut_ veg variables.

- (d)** The biplot from part (c) seems better because the scales on the variables and points are more evenly matched.

- 16.19 (a)** The two-dimensional coordinates of the observation points and variable points are as follows.

Observation Points

Name	Coordinate 1	Coordinate 2
FSM1	-9.535	-4.752
Sister	2.705	.796
FSM2	4.043	-.584
Father	4.392	.614
Teacher	-8.708	5.008
MSM	3.409	.701
FSM3	3.694	-1.782

Variable Points

Name	Coordinate 1	Coordinate 2
KIND	.610	-.054
INTEL	.085	.413
HAPPY	.407	-.456
LIKE	.621	-.039
JUST	.264	.785

In the biplot, the arrows for the variables are too short to pass through the points for observations.

- (b)** The two-dimensional coordinates of the observation points and variable points are given below.

Observation Points

Name	Coordinate 1	Coordinate 2
FSM1	-.622	-.655
Sister	.176	.110
FSM2	.264	-.080
Father	.287	.085
Teacher	-.568	.690
MSM	.222	.097
FSM3	.241	-.246

Variable Points

Name	Coordinate 1	Coordinate 2
KIND	9.345	-.391
INTEL	1.298	2.997
HAPPY	6.235	-3.313
LIKE	9.521	-.282
JUST	4.054	5.700

In the biplot, the observation points are tightly clustered around the point (0,0) making them difficult to distinguish, whereas variable points are well spaced. Just and intelligent are highly positively correlated, as are kind and likeable.

- (c) The two-dimensional coordinates of the observation points and variable points are given below.

Observation Points

Name	Coordinate 1	Coordinate 2
FSM1	-2.435	-1.764
Sister	.691	.295
FSM2	1.033	-.217
Father	1.122	.228
Teacher	-2.224	1.859
MSM	.871	.260
FSM3	.943	-.662

Variable Points

Name	Coordinate 1	Coordinate 2
KIND	2.387	-.145
INTEL	.331	1.113
HAPPY	1.593	-1.230
LIKE	2.432	-.105
JUST	1.036	2.116

In the biplot, the variable points and observation points are both well spaced. Father, sister, FSM2 and FSM3 all scored high on the kind, likeable, and happy variables, whereas teacher and FSM1 scored negatively on those variables.

- (d) The biplot from part (c) seems better because the scales on the variables and points are more evenly matched.

- 16.20 (a)** The two-dimensional coordinates of the observation points and variable points are given below.

Observation Points

Name	Coordinate 1	Coordinate 2
1	49.410	-5.832
2	25.407	-7.658
3	21.600	-2.340
4	-23.545	-6.367
5	-28.477	-4.773
6	-33.341	2.315
7	-28.176	7.992
8	-25.786	12.655
9	-29.703	9.275
10	-33.868	-3.776
11	-33.529	-1.977
12	28.186	-16.031
13	10.804	-6.608
14	.566	3.021
15	77.970	.109
16	12.859	16.294
17	41.960	5.103
18	46.930	19.064
19	34.958	-1.018
20	-16.477	1.148
21	-23.634	-1.055
22	-34.036	-2.424
23	20.632	-5.882
24	-15.873	-6.731
25	-23.023	.745
26	-15.183	-1.942
27	-11.903	-6.917
28	5.273	3.610

Variable Points

Name	Coordinate 1	Coordinate 2
North	.526	.225
East	.429	.752
South	.579	-.379
West	.452	-.490

In the biplot, the variable points are tightly grouped and the corresponding arrows do not pass through the points for observations.

- (b) The two-dimensional coordinates of the observation points and variable points are given below.

Observation Points

Name	Coordinate 1	Coordinate 2
1	.303	-.145
2	.156	-.191
3	.132	-.058
4	-.144	-.158
5	-.175	-.119
6	-.205	.058
7	-.173	.199
8	-.158	.315
9	-.182	.231
10	-.208	-.094
11	-.206	-.049
12	.173	-.399
13	.066	-.164
14	.003	.075
15	.478	.003
16	.079	.406
17	.257	.127
18	.288	.474
19	.214	-.025
20	-.101	.029
21	-.145	-.026
22	-.209	-.060
23	.127	-.146
24	-.097	-.168
25	-.141	.019
26	-.093	-.048
27	-.073	-.172
28	.032	.090

Variable Points

Name	Coordinate 1	Coordinate 2
North	85.779	9.026
East	69.899	30.223
South	94.377	-15.213
West	73.682	-19.694

In the biplot, the observation points are tightly clustered around the point (0,0), making them difficult to distinguish, whereas variable points are well spaced. All the variables are positively correlated, with south and west showing the closest relationship.

- (c) The two-dimensional coordinates of the observation points and variable points are given below.

Observation Points

Name	Coordinate 1	Coordinate 2
1	3.870	-.920
2	1.990	-1.208
3	1.692	-.369
4	-1.844	-1.004
5	-2.230	-.753
6	-2.611	.365
7	-2.207	1.261
8	-2.020	1.996
9	-2.326	1.463
10	-2.652	-.596
11	-2.626	-.312
12	2.207	-2.529
13	.846	-1.042
14	.044	.477
15	6.106	.017
16	1.007	2.571
17	3.286	.805
18	3.675	3.008
19	2.738	-.161
20	-1.290	.181
21	-1.851	-.166
22	-2.666	-.382
23	1.616	-.928
24	-1.243	-1.062
25	-1.803	.118
26	-1.189	-.306
27	-.932	-1.091
28	.413	.569

Variable Points

Name	Coordinate 1	Coordinate 2
North	6.718	1.424
East	5.474	4.768
South	7.391	-.2400
West	5.771	-.3107

In the biplot, the variable points and observation points are both well spaced. Tree 18 is associated with east, 17 with north, 1 and 3 with south, and 2 and 23 with west.

- (d) The biplot from part (c) seems better because the scales on the variables and points are more evenly matched.

Appendix C

Data Sets and SAS Files

Two sets of files are located on the ftp server of John Wiley & Sons STM (Scientific, Technical, and Medical) Division.

All data sets in the book

SAS command files for nearly all numerical examples

DOWNLOADING DATA SET FILES FROM FTP SERVER

The new FTP site for the third edition can be found at:

ftp://ftp.wiley.com/public/sci_tech_med/multivariate_analysis_3e

The zipped files at this site can be downloaded and extracted. Additional information about the files can be found in the file named README.TXT.

*

References

- Amemiya, Y., and Anderson, T. W. (1990), "Asymptotic Chi-Square Tests for a Large Class of Factor Analysis Models," *The Annals of Statistics*, **18**, 1453–1463.
- Anderberg, M. R. (1973), *Cluster Analysis for Applications*, New York: Academic Press.
- Anderson, E. (1960), "A Semi-Graphical Method for Analysis of Complex Problems," *Technometrics*, **2**, 287–292.
- Anderson, T. W., and Amemiya, Y. (1988), "The Asymptotic Normal Distribution of Estimators in Factor Analysis Under General Conditions," *The Annals of Statistics*, **16**, 759–771.
- Andrews, D. F., Gnanadesikan, R., and Warner, J. L. (1971), "Transformations of Multivariate Data," *Biometrics*, **27**, 825–840.
- Andrews, D. F., and Herzberg, A. M. (1985), *Data*, New York: Springer-Verlag.
- Asimov, D. (1985), "The Grand Tour: A Tool for Viewing Multidimensional Data," *SIAM Journal of Scientific Statistical Computing*, **6**, 128–143.
- Bailey, B. J. R. (1977), "Tables of the Bonferroni *t* Statistic," *Journal of the American Statistical Association*, **72**, 469–479.
- Barcikowski, R. S. (1981), "Statistical Power with Group Means as the Unit of Analysis," *Journal of Educational Statistics*, **6**, 267–285.
- Barnett, V., and Lewis, T. (1978), *Outliers in Statistical Data*, New York: Wiley.

- Bartlett, M. S. (1937), "Properties of Sufficiency and Statistical Tests," *Proceedings of the Royal Society of London, Ser. A*, **160**, 268–282.
- Baten, W. D., Tack, P. I., and Baeder, H. A. (1958), "Testing for Differences between Methods of Preparing Fish by Use of Discriminant Function," *Industrial Quality Control*, **14**, 6–10.
- Bayne, C. K., Beauchamp, J. J., Kane, V. E., and McCabe, G. P. (1983), "Assessment of Fisher and Logistic Linear and Quadratic Discrimination Models," *Computational Statistics and Data Analysis*, **1**, 257–273.
- Beale, E. M. L. (1969), "Euclidean Cluster Analysis," *Bulletin of the International Statistical Institute*, **43** (2), 92–4.
- Beall, G. (1945), "Approximate Methods in Calculating Discriminant Functions," *Psychometrika*, **10**(3), 205–217.
- Beauchamp, J. J., and Hoel, D. G. (1973), "Some Investigation and Simulation Studies of Canonical Analysis," *Journal of Statistical Computation and Simulation*, **2**, 197–209.
- Beckman, R. J., and Cook, R. D. (1983), "Outliers" (with comments), *Technometrics*, **25**, 119–163.
- Bentler, P. M. (1989), *EQS, Structural Equations, Program Manual*, Program Version 3.0, Los Angeles: BMDP Statistical Software, Inc.
- Benzécri, J.-P. (1992), *Correspondence Analysis Handbook*, New York: Marcel Dekker.
- Bock, R. D. (1963), "Multivariate Analysis of Variance of Repeated Measurements," in *Problems of Measuring Change*, C. W. Harris (ed.), Madison, WI: University of Wisconsin Press, pp. 85–103.
- Bock, R. D. (1975), *Multivariate Statistical Methods in Behavioral Research*, New York: McGraw-Hill.
- Boik, R. J. (1981), "A Priori Tests in Repeated Measures Designs: Effects of Non-sphericity," *Psychometrika*, **46**, 241–255.
- Bollen, K. A. (1989), *Structural Equations with Latent Variables*, New York: John Wiley & Sons.
- Bonferroni, C. E. (1936), "Il Calcolo delle Assicurazioni su Gruppi di Teste," in *Studii in Onore del Profesor S. O. Carboni Roma*.
- Box, G. E. P. (1949), "A General Distribution Theory for a Class of Likelihood Criteria," *Biometrika*, **36**, 317–346.
- Box, G. E. P. (1950), "Problems in the Analysis of Growth and Linear Curves," *Biometrics*, **6**, 362–389.
- Box, G. E. P. (1954), "Some Theorems on Quadratic Forms Applied in the Study of Analysis of Variance Problems: II. The Effect of Inequality of Variance and of Correlation between Errors in the Two-Way Classification," *Annals of Mathematical Statistics*, **25**, 484–498.
- Box, G. E. P., and Cox, D. R. (1964), "An Analysis of Transformations," *Journal of the Royal Statistical Society (B)*, **26**, 211–252.
- Box, G. E. P., and Youle, P. V. (1955), "The Exploration of Response Surfaces: An Example of the Link Between the Fitted Surface and the Basic Mechanism of the System," *Biometrics*, **11**, 287–323.

- Breiman, L., Friedman, H., Olshen, R. A., and Stone, C. J. (1984), *Classification and Regression Trees*, Belmont, CA: Wadsworth International.
- Brown, B. L., Strong, W. J., and Rencher, A. C. (1973), "Perceptions of Personality from Speech: Effects of Manipulations of Acoustical Parameters," *The Journal of the Acoustical Society of America*, **54**, 29–35.
- Brown, B. L., Williams, R. N., and Barlow, C. D. (1984), "PRIFAC: A Pascal Factor Analysis Program," *Journal of Pascal, Ada, and Modula*, **2**, 18–24.
- Brown, T. A., and Koplowitz, J. (1979), "The Weighted Nearest Neighbor Rule for Class Dependent Sample Sizes," *IEEE Transactions on Information Theory*, **IT-25**, 617–619.
- Bryce, G. R. (1980), "Some Observations on the Analysis of Growth Curves," Paper No. SD-025-R, Brigham Young University, Department of Statistics.
- Buck, S. F. A. (1960), "A Method of Estimation of Missing Values in Multivariate Data Suitable for Use with an Electronic Computer," *Journal of the Royal Statistical Society, Series B*, **22**, 302–307.
- Buja, A., Cook, D., Asimov, D., and Hurley, C. (2005), "Computational Methods for High-Dimensional Rotations in Data Visualization," in *Handbook of Statistics: Data Mining and Visualization*, Vol. 24, C. R. Rao, E. J. Wegman, and J. L. Solka (eds.), New York: Elsevier/North Holland, pp. 391–413.
- Burdick, R. K. (1979), "On the Use of Canonical Analysis to Test MANOVA Hypotheses," Presented at the Annual Meeting of the American Statistical Association, Washington, DC, August 1979.
- Cacoullos, T. (1966), "Estimation of a Multivariate Density," *Annals of the Institute of Statistical Mathematics*, **18**, 179–189.
- Cameron, E., and Pauling, L. (1978), "Supplemental Ascorbate in the Supportive Treatment of Cancer: Reevaluation of Prolongation of Survival Times in Terminal Human Cancer," *Proceedings of the National Academy of Sciences, U.S.A.*, **75**, 4538–4552.
- Campbell, N. A. (1980), "Robust Procedures in Multivariate Analysis I. Robust Covariance Estimation," *Applied Statistics*, **29**, 231–237.
- Chambers, J. M., and Kleiner, B. (1982), "Graphical Techniques for Multivariate Data and for Clustering," in *Handbook of Statistics*, Vol. 2, P. R. Krishnaiah and L. N. Kanal (eds.), New York: North-Holland, pp. 209–244.
- Chernoff, H. (1973), "The Use of Faces to Represent Points in k -Dimensional Space Graphically," *Journal of the American Statistical Association*, **68**, 361–368.
- Chidananda Gowda, K., and Krishna, G. (1979), "The Condensed Nearest Neighbor Rule Using the Concept of Mutual Nearest Neighborhood," *IEEE Transactions on Information Theory*, **IT-25**, 488–490.
- Clausen, Sten-Erik (1998), *Applied Correspondence Analysis*, Thousand Oaks, CA: Sage Publications.
- Cochran, W. G., and Cox, G. M. (1957), *Experimental Designs*, 2nd ed. New York: Wiley.
- Collier, R. O., Baker, F. B., Mandeville, G. K., and Hayes, T. F. (1967), "Estimates of Test Size for Several Procedures Based on Conventional Ratios in the Repeated Measure Design," *Psychometrika*, **32**, 339–353.

- Colvin, T. S., Jaynes, D. B., Karlen, D. L., Laird, D. A., and Ambuel, J. R. (1997), "Yield Variability Within a Central Iowa Field," *Transactions of the ASAE*, **40**, 883–889.
- Cook, D., and Swayne, D. F. (2007), *Interactive and Dynamic Graphics for Data Analysis, With R and GGobi*, New York: Springer.
- Cramer, E. M., and Nicewander, W. A. (1979), "Some Symmetric, Invariant Measures of Multivariate Association," *Psychometrika*, **44**, 43–54.
- Crepeau, H., Koziol, J., Reid, N., and York, Y. S. (1985), "Analysis of Incomplete Multivariate Data from Repeated Measurement Experiments," *Biometrics*, **41**, 505–514.
- Critchley, F. (1985), "Influence in Principal Components Analysis," *Biometrika*, **72**, 627–636.
- Crowder, M. J., and Hand, D. J. (1990), *Analysis of Repeated Measures*, New York: Chapman & Hall.
- D'Agostino, R. B. (1971), "An Omnibus Test of Normality for Moderate and Large Size Samples," *Biometrika*, **58**, 341–348.
- D'Agostino, R. B. (1972), "Small Sample Probability Points for the *D* Test of Normality," *Biometrika*, **59**, 219–221.
- D'Agostino, R. B., and Pearson, E. S. (1973), "Tests for Departure from Normality. Empirical Results for the Distributions of b_2 and $\sqrt{b_1}$," *Biometrika*, **60**, 613–622; correction, **61**, 647.
- D'Agostino, R. B., and Tietjen, G. L. (1971), "Simulated Probability Points of b_2 for Small Samples," *Biometrika*, **58**, 669–672.
- Davidson, M. L. (1972), "Univariate versus Multivariate Tests in Repeated Measures Experiments," *Psychological Bulletin*, **77**, 446–452.
- Davidson, M. L. (1983), *Multidimensional Scaling*, New York: Wiley.
- Davis, A. W. (1970a), "Exact Distributions of Hotelling's Generalized T_0^2 -Test," *Biometrika*, **57**, 187–191.
- Davis, A. W. (1970b), "Further Applications of a Differential Equation for Hotelling's Generalized T_0^2 -Test," *Annals of the Institute of Statistical Mathematics*, **22**, 77–87.
- Davis, A. W. (1970c), "Differential Equation of Hotellings Generalized," *Annals of Statistics*, **39**, 815–832.
- Davis, A. W. (1972), "On the Marginal Distributions of the Latent Roots of the Multivariate Beta Matrix," *Biometrika*, **43**, 1664–1670.
- Davis, A. W. (1979), "On the Differential Equation for Meijer $C_{p,p}^{0,0}$ Function, and Further Wilks's Likelihood Ratio Criterion," *Biometrika*, **66**, 519–531.
- Davis, A. W. (1980), "Further Tabulation of Hotelling's Generalized T_0^2 -Test," *Communications in Statistics—Part B Simulation and Computation*, **9**, 321–336.
- Dawkins, B. (1989), "Multivariate Analysis of National Track Records," *American Statistician*, **43**, 110–115.
- Dempster, A. P., Laird, N. M., and Rubin, D. B. (1977), "Maximum Likelihood from Incomplete Data via the EM Algorithm," *Journal of the Royal Statistical Society, Series B*, **39**, 1–38.

- Devlin, S. J., Gnanadesikan, R., and Kettenring, J. R. (1981), "Robust Estimation of Dispersion Matrices and Principal Components," *Journal of the American Statistical Association*, **76**, 354–362.
- Dubien, J. L. and Warde, W. D. (1979), "A Mathematical Comparison of the Members of an Infinite Family of Agglomerative Clustering Algorithms," *Canadian Journal of Statistics*, **7**, 29–38.
- Duran, B. S. and Odell, P. L. (1974), "Cluster Analysis: A Survey," *Lecture Notes in Economics and Mathematical Systems*, New York: Springer-Verlag.
- Edwards, D. and Kreiner, S. (1983), "The Analysis of Contingency Tables by Graphical Models," *Biometrika*, **70** (3), pp. 553–65.
- Ehrenberg, A. S. C. (1977), "Rudiments of Numeracy," *Journal of the Royal Statistical Society, Series A*, **140**, 277–297.
- Elston, R. C., and Grizzle, J. E. (1962), "Estimation of Time-Response Curves and Their Confidence Bands," *Biometrics*, **18**, 148–159.
- Everitt, B. S. (1987), *Introduction to Optimization Methods and Their Application in Statistics*, London: Chapman & Hall.
- Everitt, B. S. (1993), *Cluster Analysis*, 3rd ed., London: Edward Arnold.
- Fearn, T. (1975), "A Bayesian Approach to Growth Curves," *Biometrika*, **62**, 89–100.
- Fearn, T. (1977), "A Two-Stage Model for Growth Curves Which Leads to Rao's Covariance Adjusted Estimators," *Biometrika*, **64**, 141–143.
- Federer, W. T. (1986), "On Planning Repeated Measures Experiments," Unit Report BU-909-M, Cornell University, Biometrics Unit.
- Fehlberg, W. T. (1980), "Repeated Measures: The Effect of a Preliminary Test for the Structure of the Covariance Matrix upon the Alpha Levels of the Tests for Experimental Design Factors," M. A. Thesis, Brigham Young University, Department of Statistics.
- Ferguson, T. S. (1961), "On the Rejection of Outliers," *Proceedings of the Fourth Berkeley Symposium on Mathematical Statistics and Probability*, Vol. 1, Los Angeles, CA: University of California Press, pp. 253–287.
- Fisher, R. A. (1936), "The Use of Multiple Measurement in Taxonomic Problems," *Annals of Eugenics*, **7**, 179–188.
- Fix, E., and Hodges, J. L. (1951), "Discriminatory Analysis, Nonparametric Discrimination: Consistency Properties," Report No. 4, Project No. 21-49-004, Brooks Air Force Base, USAF School of Aviation Medicine.
- Flack, V. F., and Chang, P. C. (1987), "Frequency of Selecting Noise Variables in Subset Regression Analysis: A Simulation Study," *American Statistician*, **41**, 84–86.
- Flury, B., and Riedwyl, H. (1981), "Graphical Representation of Multivariate Data by Means of Asymmetric Faces," *Journal of the American Statistical Association*, **76**, 757–765.
- Flury, B., and Riedwyl, H. (1985), " T^2 Tests, the Linear Two-Group Discriminant Function, and Their Computation by Linear Regression," *American Statistician*, **39**, 20–25.
- Frets, G. P. (1921), "Heredity of Head Form in Man," *Genetica*, **3**, 193–384.

- Friedman, J. H. and Tukey, J. W. (1974), "A Projection Pursuit Algorithm for Exploratory Data Analysis," *IEEE Transactions on Component Parts*, **9**, 881–890.
- Fuller, W. A. (1987), *Measurement Error Models*, New York: John Wiley & Sons.
- Furnival, G. M., and Wilson, R. W. (1974), "Regression by Leaps and Bounds," *Technometrics*, **16**, 499–511.
- Gabriel, K. R. (1968), "Simultaneous Test Procedures in Multivariate Analysis of Variance," *Biometrika*, **55**, 489–504.
- Gabriel, K. R. (1969), "Simultaneous Test Procedures—Some Theory of Multiple Comparisons," *Annals of Mathematics and Statistics*, **40**, 224–250.
- Gabriel, K. R. (1971), "The Biplot Graphical Display of Matrices with Application to Principal Component Analysis," *Biometrika*, **58**, 453–467.
- Gates, G. W. (1972), "The Reduced Nearest Neighbor Rule," *IEEE Transactions on Information Theory*, **IT-18**, 431.
- Geisser, S. (1980), "Growth Curve Analysis," In *Handbook of Statistics*, Vol. 1, P. R. Krishnaiah (ed.), Amsterdam: North-Holland, pp. 89–115.
- Gerbing, D. W. and Hamilton, J. G. (1996), "Viability of Exploratory Factor Analysis as a Precursor to Confirmatory Factor Analysis," *Structural Equation Modeling*, **3**, 62–72.
- Gilbert, E. S. (1968), "On Discrimination Using Qualitative Variables," *Journal of the American Statistical Association*, **63**, 1399–1412.
- Gnanadesikan, R. (1997), *Methods for Statistical Data Analysis of Multivariate Observations*, 2nd ed., New York: Wiley.
- Gnanadesikan, R., and Kettenring, J. R. (1972), "Robust Estimates, Residuals, and Outlier Detection with Multiresponse Data," *Biometrics*, **28**, 81–124.
- Gordon, A. D. (1999), *Classification*, 2nd ed., London: Chapman & Hall/CRC.
- Gower, J. C. (1966), "Some Distance Properties of Latent Root and Vector Methods used in Multivariate Analysis," *Biometrika*, **53**, 325–338.
- Gower, J. C. and Hand, D. J. (1996), *Biplots*, London: Chapman & Hall.
- Graybill, F. A. (1969), *Introduction to Matrices with Applications in Statistics*, Belmont, CA: Wadsworth.
- Greenacre, M. J. (1984), *Theory and Applications of Correspondence Analysis*, London: Academic Press.
- Greenhouse, S. W., and Geisser, S. (1959), "On Methods in the Analysis of Profile Data," *Psychometrika*, **24**, 95–112.
- Grizzle, J. E., and Allen, D. M. (1969), "Analysis of Growth and Dose Response Curves," *Biometrics*, **25**, 357–381.
- Guttman, I. (1982), *Linear Models: An Introduction*, New York: Wiley.
- Habbema, J. D. F., Hermans, J., and Remme, J. (1978), "Variable Kernel Density Estimation in Discriminant Analysis," In *Compstat 1978: Proceedings in Computational Statistics*, G. Bruckman (ed.), Vienna: Physica-Verlag, pp. 178–185.
- Habbema, J. D. F., Hermans, J., and van den Broek, K. (1974), "A Stepwise Discriminant Analysis Program Using Density Estimation," In *Comstat*, G. Bruckmann, F. Ferschl, and L. Schmetterer (eds.), Vienna: Physica-Verlag, pp. 101–110.

- Hammond, J., Smith, D., and Gill, D. S. (1981), "Selection of Adequate Subsets in Multivariate Multiple Regression," Presented at the Annual Meeting of the American Statistical Association, August 10–13, Detroit, MI.
- Hand, D. J., and Batchelor, B. G. (1978), "An Edited Condensed Nearest Neighbor Rule," *Information Sciences*, **14**, 171–180.
- Hand, D. J., Daly, F., Lunn, A. D., McConway, K. J., and Ostrowski, E. (eds.) (1994), *A Handbook of Small Data Sets*, London: Chapman & Hall.
- Harman, H. H. (1976), *Modern Factor Analysis*, 3rd rev. ed., Chicago: University of Chicago Press.
- Hart, P. E. (1968), "The Condensed Nearest Neighbor Rule," *IEEE Transactions on Information Theory*, **IT-14**, 515–516.
- Hartigan, J. A. (1975a), *Clustering Algorithms*, New York: John Wiley & Sons.
- Hartigan, J. A. (1975b), "Printer Graphics for Clustering," *Journal of Statistical Computation and Simulation*, **4**, 187–213.
- Harville, D. A. (1997), *Matrix Algebra from a Statistician's Perspective*, New York: Springer-Verlag.
- Hastie, T., Tibshirani, R., and Friedman, J. (2009), *The Elements of Statistical Learning: Data Mining, Inference, and Prediction (Second Edition)*, New York: Springer.
- Hawkins, D. M. (1980), *Identification of Outliers*, London: Chapman & Hall.
- Healy, M. J. R. (1969), "Rao's Paradox Concerning Multivariate Tests of Significance," *Biometrics*, **25**, 411–413.
- Heywood, H. B. (1931), "On Finite Sequences of Real Numbers," *Proceedings of the Royal Society, Series A*, **134**, 486–501.
- Hilton, D. K. (1983), "A Simulation and Comparison of Three Criterion Functions Used for Subset Selection in Linear Regression," M.A. Thesis, Brigham Young University, Department of Statistics.
- Hocking, R. R. (1976), "The Analysis and Selection of Variables in Linear Regression," *Biometrics*, **32**, 1–51.
- Hotelling, H. (1931), "The Generalization of Student's Ratio," *Annals of Mathematical Statistics*, **2**, 360–378.
- Hotelling, H. (1951), "A Generalized *T* Test and Measure of Multivariate Dispersion," *Proceedings of the Second Berkeley Symposium on Mathematical Statistics and Probability*, **1**, 23–41.
- Hsu, P. L. (1938), "Notes on Hotelling's Generalized T^2 ," *Annals of Mathematical Statistics*, **9**, 231–243.
- Hu, L.-T., and Bentler, P. M. (1999), "Cutoff Criteria for Fit Indexes in Covariance Structure Analysis: Conventional Criteria Versus New Alternatives," *Structural Equation Modeling: A Multidisciplinary Journal*, **6**, 1–55.
- Huber, P. J. (1985), "Projection Pursuit," *Annals of Statistics*, **13**, 435–525.
- Huberty, C. J. (1975), "The Stability of Three Indices of Relative Variable Contribution in Discriminant Analysis," *Journal of Experimental Education*, **44**, 59–64.
- Hummel, T. J., and Sligo, J. (1971), "Empirical Comparison of Univariate and Multivariate Analysis of Variance Procedures," *Psychological Bulletin*, **76**, 49–57.

- Huynh, H. (1978), "Some Approximate Tests in Repeated Measures Designs," *Psychometrika*, **43**, 1582–1589.
- Huynh, H., and Feldt, L. S. (1970), "Conditions Under Which Mean Square Ratios in Repeated Measurement Designs Have Exact *F* Distributions," *Journal of the American Statistical Association*, **65**, 1582–1589.
- Huynh, H., and Feldt, L. S. (1976), "Estimation of the Box Correction for Degrees of Freedom from Sample Data in Randomized Block and Split-Plot Designs," *Journal of Educational Statistics*, **1**(1), 69–82.
- Inselberg, A. (1985), "The Plane with Parallel Coordinates," *The Visual Computer*, **1**, 69–91.
- Izenman, A. J., and Williams, J. S. (1989), "A Class of Linear Spectral Models and Analyses for the Study of Longitudinal Data," *Biometrics*, **45**, 831–849.
- Jackson, D., and Bryce, G. R. (1981), "A Univariate-Like Method of Analyzing Growth Curve Data with Nonuniform Variance-Covariance Matrices," Paper No. SD-028-R, Brigham Young University, Department of Statistics.
- Jackson, J. E. (1980), "Principal Components and Factor Analysis: Part I—Principal Components," *Journal of Quality Technology*, **12**, 201–213.
- Jacoby, W. G. (1998), *Statistical Graphics for Visualizing Multivariate Data*, Thousand Oaks, CA: Sage Publications.
- Jardine, N. and Sibson, R. (1968), "The Construction of Hierarchic and Non-hierarchic Classifications," *Computer Journal*, **11**, 117–184.
- Jardine, N. and Sibson, R. (1971), *Mathematical Taxonomy*, London: Wiley.
- Jeffers, J. N. R. (1967), "Two Case Studies in the Application of Principal Component Analysis," *Applied Statistics*, **16**, 225–236.
- Jensen, D. R. (1982), "Efficiency and Robustness in the Use of Repeated Measurements," *Biometrics*, **38**, 813–825.
- Jensen, R. E. (1969), "A Dynamic Programming Algorithm for Cluster Analysis," *Operations Research*, **12**, 1034–1057.
- Jobson, J. D. (1992), *Applied Multivariate Data Analysis, Volume II: Categorical and Multivariate Methods*, New York: Springer-Verlag.
- Jolliffe, I. T. (1972), "Discarding Variables in a Principal Component Analysis, I: Artificial Data," *Applied Statistics*, **21**, 160–173.
- Jolliffe, I. T. (1973), "Discarding Variables in a Principal Component Analysis, II: Real Data," *Applied Statistics*, **22**, 21–31.
- Jones, M. C. and Sibson, R. (1987), "What is Projection Pursuit?" *Journal of the Royal Statistical Society*, **150**, 1–36.
- Joreskog, K. G. and Sörbom, D. (1985), *LISREL VI; Analysis of Linear Structural Relationships by Maximum Likelihood, Instrumental Variables, and Least Squares*, Uppsala, Sweden: University of Uppsala.
- Kabe, D. G. (1985), "On Some Multivariate Statistical Methodology with Applications to Statistics, Psychology, and Mathematical Programming," *Journal of the Industrial Mathematics Society*, **35**, 1–18.
- Kaiser, H. F. (1970), "A Second Generation Little Jiffy," *Psychometrika*, **35**, 401–415.

- Kaiser, H. F., and Rice, J. (1974), "Little Jiffy, Mark IV," *Educational and Psychological Measurement*, **34**, 111–117.
- Kaufman, L., and Rousseeuw, P. J. (1990), *Finding Groups in Data: An Introduction to Cluster Analysis*, New York: Wiley.
- Keuls, M., Martakis, G. F. P., and Magid, A. H. A. (1984), "The Relationship between Pod Yield and Specific Leaf Area in Snapbeans; an Example of Stepwise Multivariate Analysis of Variance," *Scientia Horticulturae*, **23**, 231–246.
- Khattree, R. and Naik, D. N. (1999), *Applied Multivariate Statistics with SAS Software*, 2nd ed, New York: Wiley.
- Khattree, R. and Naik, D. N. (2000), *Multivariate Data Reduction and Discrimination with SAS Software*, New York: Wiley.
- Khuri, A. I., Mathew, T., and Nel, D. G. (1994), "A Test to Determine Closeness of Multivariate Satterthwaite's Approximation," *Journal of Multivariate Analysis*, **51**, 201–209.
- Kleinbaum, D. G., Kupper, L. L., and Muller, K. E. (1988), *Applied Regression Analysis and Other Multivariable Methods*, Boston: PWS-KENT.
- Kleiner, B., and Hartigan, J. A. (1981), "Representing Points in Many Dimensions by Trees and Castles," *Journal of the American Statistical Association*, **76**, 260–269.
- Kramer, C. Y. (1972), *A First Course in Multivariate Analysis*, Published by the author, Blacksburg, VA.
- Kramer, C. Y., and Jensen, D. R. (1969a), "Fundamentals of Multivariate Analysis, Part I. Inference about Means," *Journal of Quality Technology*, **1**(2), 120–133.
- Kramer, C. Y., and Jensen, D. R. (1969b), "Fundamentals of Multivariate Analysis, Part II. Inference about Two Treatments," *Journal of Quality Technology*, **1**(3), 189–204.
- Kramer, C. Y., and Jensen, D. R. (1970), "Fundamentals of Multivariate Analysis, Part IV. Analysis of Variance for Balanced Experiments," *Journal of Quality Technology*, **2**, 32–40.
- Kruskal, J. B. (1964a), "Multidimensional Scaling by Optimizing Goodness of Fit to a Nonmetric Hypothesis," *Psychometrika*, **29**, 1–27.
- Kruskal, J. B. (1964b), "Nonmetric Multidimensional Scaling: A Numerical Method," *Psychometrika*, **29**, 115–129.
- Kruskal, J. B. and Wish, M. (1978), *Multidimensional Scaling*, Beverly Hills, CA: Sage Publications.
- Krzanowski, W. J. (1975), "Discrimination and Classification Using Both Binary and Continuous Variables," *Journal of the American Statistical Association*, **70**, 782–790.
- Krzanowski, W. J. (1976), "Canonical Representation of the Location Model for Discrimination or Classification," *Journal of the American Statistical Association*, **71**, 845–848.
- Krzanowski, W. J. (1977), "The Performance of Fisher's Linear Discriminant Function Under Non-Optimal Conditions," *Technometrics*, **19**, 191–200.
- Krzanowski, W. J. (1979), "Some Linear Transformations for Mixtures of Binary and Continuous Variables, with Particular Reference to Linear Discriminant Analysis," *Biometrika*, **66**, 33–39.

- Krzanowski, W. J. (1980), "Mixtures of Continuous and Categorical Variables in Discriminant Analysis," *Biometrics*, **36**, 493–499.
- Lachenbruch, P. A. (1975), *Discriminant Analysis*, New York: Hafner.
- Lachenbruch, P. A., and Goldstein, M. (1979), "Discriminant Analysis," *Biometrics*, **35**, 69–85.
- Lance, G. N. and Williams, W. T. (1966), "Computer Programs for Hierarchical Polythetic Classification," *Computer Journal*, **9**, 60–64.
- Lance, G. N. and Williams, W. T. (1967), "A General Theory of Classificatory Sorting Strategies: I. Hierarchical Systems," *Computer Journal*, **9**, 373–380.
- Lavine, B., and Carlson, D. (1987), "European Bee or Africanized Bee?: Species Identification through Chemical Analysis," *Analytical Chemistry*, **59**(6), 468–470.
- Lawley, D. N. (1938), "A Generalization of Fisher's Z-Test," *Biometrika*, **30**, 180–187.
- Lee, J. C. (1988), "Prediction and Estimation of Growth Curves with Special Covariance Structures," *Journal of the American Statistical Association*, **83**, 432–440.
- Lee, J. C., Chang, T. C., and Krishnaiah, P. R. (1977), "Approximations to the Distributions of the Likelihood Ratio Statistics for Testing Certain Structures on the Covariance Matrices of Real Multivariate Normal Populations," In *Multivariate Analysis*, Vol. 4, P. R. Krishnaiah (ed.), Amsterdam: North-Holland, pp. 105–118.
- Lee, Y. (1972), "Some Results on the Distribution of Wilks's Likelihood Ratio Criterion," *Biometrika*, **95**, 649–664.
- Lin, C. C., and Mudholkar, G. S. (1980), "A Simple Test for Normality Against Asymmetric Alternatives," *Biometrika*, **67**, 455–461.
- Lindsey, H., Webster, J. T., and Halpern, H. (1985), "Canonical Correlation as a Discriminant Tool in a Periodontal Problem," *Biometrika*, **72**, 257–264.
- Loftsgaarden, D. O., and Quesenberry, C. P. (1965), "A Nonparametric Estimate of a Multivariate Density Function," *Annals of Mathematical Statistics*, **36**, 1049–1051.
- Lubischew, A. A. (1962), "On the Use of Discriminant Functions in Taxonomy," *Biometrics*, **18**, 455–477.
- MacNaughton-Smith, P., Williams, W. T., Dale, M. B. and Mockett, L. G. (1964), "Dissimilarity Analysis," *Nature*, **202**, 1034–1035.
- Mahalanobis, P. C. (1936), "On the Generalized Distance in Statistics," *Proceedings of the National Institute of Sciences of India*, **12**, 49–55.
- Mallows, C. L. (1964), "Choosing Variables in a Linear Regression: A Graphical Aid," Paper presented at the Central Regional Meeting of the Institute of Mathematical Statistics, Manhattan, KS.
- Mallows, C. L. (1973), "Some Comments on C_p ," *Technometrics*, **15**, 661–675.
- Mardia, K. V. (1970), "Measures of Multivariate Skewness and Kurtosis with Applications," *Biometrika*, **57**, 519–530.
- Mardia, K. V. (1974), "Applications of Some Measures of Multivariate Skewness and Kurtosis for Testing Normality and Robustness Studies," *Sankhya, Series B*, **36**, 115–128.

- Mardia, K. V., Kent, J. T. and Bibby, J. M. (1979), *Multivariate Analysis*, New York: Academic Press.
- Marks, R. G., and Rao, P. V. (1978), "A Modified Tiao-Guttman Rule for Multiple Outliers," *Communications in Statistics*, **A7**, 113–126.
- Mathai, A. M., and Katiyar, R. S. (1979), "Exact Percentage Points for Testing Independence," *Biometrika*, **66**, 353–356.
- Mauchly, J. W. (1940), "Significance Test for Sphericity of a Normal N -Variate Distribution," *Annals of Mathematical Statistics*, **11**, 204–209.
- Maxwell, S. E., and Avery, R. D. (1982), "Small Sample Profile Analysis with Many Variables," *Psychological Bulletin*, **92**, 778–785.
- McCabe, G. P. (1984), "Principal Variables," *Technometrics*, **26**, 137–144.
- McKay, R. J. (1977), "Variable Selection in Multivariate Regression: An Application of Simultaneous Test Procedures," *Journal of the Royal Statistical Society, Series B*, **39**, 371–380.
- McKay, R. J. (1979), "The Adequacy of Variable Subsets in Multivariate Regression," *Technometrics*, **21**, 475–480.
- Milligan, G. W. (1996), "Clustering Validation: Results and Implications for Applied Analyses," *Clustering and Classification* (eds P. Arabie, L. J. Hubert and G. De Soete), World Scientific, Singapore, 341–375.
- Milligan, G. W. and Cooper, M. C. (1985), "An Examination of Procedures for Determining the Number of Clusters in a Data Set," *Psychometrika*, **50**, 159–179.
- Mojena, R. (1977), "Hierarchical Grouping Methods and Stopping Rules: An Evaluation," *Computer Journal*, **20**, 359–363.
- Morrison, D. F. (1972), "The Analysis of a Single Sample of Repeated Measurements," *Biometrics*, **28**, 55–71.
- Morrison, D. F. (1983), *Applied Linear Statistical Methods*, Englewood Cliffs, NJ: Prentice-Hall.
- Morrison, D. F. (1990), *Multivariate Statistical Methods*, 3rd ed, New York: McGraw-Hill.
- Mosteller, F., and Wallace, D. L. (1984), *Applied Bayesian and Classical Inference: The Case of the Federalist Papers*, New York: Springer.
- Mulholland, H. P. (1977), "On the Null Distribution of $\sqrt{b_1}$ for Samples of Size at Most 25, with Tables," *Biometrika*, **64**, 401–409.
- Muller, K. E. (1982), "Understanding Canonical Correlation through the General Linear Model and Principal Components," *American Statistician*, **36**, 342–354.
- Muller, K. E. (1998), "A New F Approximation for the Pillai-Bartlett Trace Under H_0 ," *Journal of Computational and Graphical Statistics*, **7**, 131–137.
- Muller, K. E., and Peterson, B. L. (1984), "Practical Methods for Computing Power in Testing the Multivariate General Linear Hypothesis," *Computational Statistics and Data Analysis*, **2**, 143–158.
- Myers, R. H. (1990), *Classical and Modern Regression with Applications*, 2nd ed, Boston: PWS-KENT.
- Nagarsenker, B. N., and Pillai, K. C. S. (1973), "The Distribution of the Sphericity Test Criterion," *Journal of Multivariate Analysis*, **3**, 226–235.

- Nason, G. (1995), "Three-Dimensional Projection Pursuit," *Journal of Applied Statistics*, **44**(4), 411–430.
- Olson, C. L. (1974), "Comparative Robustness of Six Tests in Multivariate Analysis of Variance," *Journal of the American Statistical Association*, **69**, 894–908.
- O'Sullivan, J. B., and Mahan, C. M. (1966), "Glucose Tolerance Test: Variability in Pregnant and Non-pregnant Women," *American Journal of Clinical Nutrition*, **19**, 345–351.
- Parzen, E. (1962), "On Estimation of a Probability Density Function and Mode," *Annals of Mathematical Statistics*, **33**, 1065–1076.
- Patel, H. I. (1986), "Analysis of Repeated Measures Designs with Changing Covariates in Clinical Trials," *Biometrika*, **73**, 707–715.
- Pearson, E. S., and Hartley, H. O. (eds.) (1972), *Biometrika Tables for Statisticians*, Volume 2, Cambridge: Cambridge University Press.
- Pfeiffer, K. P. (1985), "Stepwise Variable Selection and Maximum Likelihood Estimation of Smoothing Factors of Kernel Functions for Nonparametric Discriminant Functions Evaluated by Different Criteria," *Computers and Biomedical Research*, **18**, 46–61.
- Pillai, K. C. S. (1964), "On the Distribution of the Largest of Seven Roots of a Matrix in Multivariate Analysis," *Biometrika*, **51**, 270–275.
- Pillai, K. C. S. (1965), "On the Distribution of the Largest Characteristic Root of a Matrix in Multivariate Analysis," *Biometrika*, **52**, 405–414.
- Pillai, K. C. S. and Flury, B. N. (1984), "Percentage Points of the Largest Characteristic Root of the Multivariate Beta Matrix," *Communications in Statistics, Part A*, **13**, 2199–2237.
- Posse, C. (1990), "An Effective Two-Dimensional Projection Pursuit Algorithm," *Communications in Statistics: Simulation and Computation*, **19**, 1143–1164.
- Posten, H. O. (1962), "Analysis of Variance and Analysis of Regression with More than One Response," *Proceedings of a Symposium on Applications of Statistics and Computers to Fuel and Lubricant Problems*, 91–109.
- Potthoff, R. F., and Roy, S. N. (1964), "A Generalized Multivariate Analysis of Variance Model Useful Especially for Growth Curve Problems," *Biometrika*, **51**, 313–326.
- R Development Core Team (2010), *R: A Language and Environment for Statistical Computing*, Vienna: R Foundation for Statistical Computing.
- Rao, C. R. (1948), "Tests of Significance in Multivariate Analysis," *Biometrika*, **35**, 58–79.
- Rao, C. R. (1959), "Some Problems Involving Linear Hypotheses in Multivariate Analysis," *Biometrika*, **46**, 49–58.
- Rao, C. R. (1966), "Covariance Adjustment and Related Problems in Multivariate Analysis," In *Multivariate Analysis*, P. Krishnaiah (ed.), New York: Academic Press, pp. 87–103.
- Rao, C. R. (1973), *Linear Statistical Inference and Its Applications*, 2nd ed, New York: Wiley.
- Rao, C. R. (1984), "Prediction of Future Observations in Polynomial Growth Curve Models," in *Proceedings of the Indian Statistical Institute Golden Jubilee Inter-*

- national Conference on Statistics: Applications and New Directions*, Calcutta: Indian Statistical Institute, pp. 512–520.
- Rao, C. R. (1987), “Prediction of Future Observations in Growth Curve Models,” *Statistical Science*, **2**, 434–471.
- Reaven, G. M., and Miller, R. G. (1979), “An Attempt to Define the Nature of Chemical Diabetes Using a Multidimensional Analysis,” *Diabetologia*, **16**, 17–24.
- Reinsel, G. (1982), “Multivariate Repeated-Measurement or Growth Curve Models with Random-Effects Covariance Structure,” *Journal of the American Statistical Association*, **77**, 190–195.
- Remme, J., Habbema, J. D. F., and Hermans, J. (1980), “A Simulative Comparison of Linear, Quadratic, and Kernel Discrimination,” *Journal of Statistical Computation and Simulation*, **11**, 87–106.
- Rencher, A. C. (1988), “On the Use of Correlations to Interpret Canonical Functions,” *Biometrika*, **75**, 363–365.
- Rencher, A. C. (1992a), “Bias in Apparent Classification Rates in Stepwise Discriminant Analysis,” *Communications in Statistics—Simulation and Computation*, **21**, 373–389.
- Rencher, A. C. (1992b), “Interpretation of Canonical Discriminant Functions, Canonical Variates, and Principal Components,” *American Statistician*, **46**, 217–225.
- Rencher, A. C. (1993), “The Contribution of Individual Variables to Hotelling’s T^2 , Wilks’ Λ , and R^2 ,” *Biometrics*, **49**, 479–489.
- Rencher, A. C. (1998), *Multivariate Statistical Inference and Applications*, New York: John Wiley & Sons.
- Rencher, A. C., and Larson, S. F. (1980), “Bias in Wilks’ Λ in Stepwise Discriminant Analysis,” *Technometrics*, **22**, 349–356.
- Rencher, A. C., and Pun, F. C. (1980), “Inflation of R^2 in Best Subset Regression,” *Technometrics*, **22**, 49–53.
- Rencher, A. C., and Schaalje, G. B. (2008), *Linear Models in Statistics*, 2nd ed., New York: John Wiley & Sons.
- Rencher, A. C., and Scott, D. T. (1990), “Assessing the Contribution of Individual Variables Following Rejection of a Multivariate Hypothesis,” *Communications in Statistics: Simulation and Computation*, **19**(2), 535–553.
- Rencher, A. C., Wadham, R. A., and Young, J. R. (1978), “A Discriminant Analysis of Four Levels of Teacher Competence,” *Journal of Experimental Education*, **46**, 46–51.
- Ripley, B. D. (1996), *Pattern Recognition and Neural Networks*, Cambridge: Cambridge University Press.
- Robert, P., and Escoufier, Y. (1976), “A Unifying Tool for Linear Multivariate Analysis: The RV-Coefficient,” *Journal of the Royal Statistical Society, Series C*, **25**, 257–265.
- Roebruck, P. (1982), “Canonical Forms and Tests of Hypotheses; Part II: Multivariate Mixed Linear Models,” *Statistica Neerlandica*, **36**, 75–80.
- Rogan, J. C., Keselman, H. J., and Mendoza, J. L. (1979), “Analysis of Repeated Measurements,” *British Journal of Mathematical and Statistical Psychology*, **32**, 269–286.

- Rogers, W. H., and Wagner, T. J. (1978), "A Finite Sample Distribution Free Performance Bound for Local Discrimination Rules," *Annals of Statistics*, **6**, 506–514.
- Romney, A. K., Shepard, R. N., and Nerlove, S. B. (eds.) (1972), *Multidimensional Scaling: Theory and Applications in the Behavioral Sciences, Volume II: Applications*, New York: Seminar Press.
- Rosenblatt, M. (1956), "Remarks on Some Nonparametric Estimates of a Density Function," *Annals of Mathematical Statistics*, **27**, 832–837.
- Royston, J. P. (1983), "Some Techniques for Assessing Multivariate Normality Based on the Shapiro-Wilk W ," *Applied Statistics*, **32**, 121–133.
- Ruyymgaart, F. H. (1981), "A Robust Principal Component Analysis," *Journal of Multivariate Analysis*, **11**, 485–497.
- SAS Institute Inc. (2009), "SAS OnlineDoc® 9.2," Cary, NC: SAS Institute Inc.
- Satterthwaite, F. E. (1941), "Synthesis of Variances," *Psychometrika*, **6**, 309–316.
- Schoenberg, I. J. (1935), "Remarks to Maurice Fréchet's article 'Sur la definition axiomatique d'une classe d'espace distanciés vectoriellement applicable sur l'espace de Hilbert,'" *Annals of Mathematics*, **36**, 724–732.
- Schott, J. R., and Saw, J. G. (1984), "A Multivariate One-Way Classification Model with Random Effects," *Journal of Multivariate Analysis*, **15**, 1–12.
- Schuurmann, F. J., Krishnaiah, P. R., and Chattopadhyay, A. K. (1975), "Exact Percentage Points of the Distribution of the Trace of a Multivariate Beta Matrix," *Journal of Statistical Computation and Simulation*, **3**, 331–343.
- Schwager, S. J., and Margolin, B. H. (1982), "Detection of Multivariate Normal Outliers," *Annals of Statistics*, **10**, 943–954.
- Scott, D. W. (1992), *Multivariate Density Estimation Theory, Practice and Visualization*, New York: John Wiley & Sons.
- Searle, S. R., (1982), *Matrix Algebra Useful for Statistics*, New York: Wiley.
- Seber, G. A. F. (1984), *Multivariate Observations*, New York: Wiley.
- Shapiro, S. S., and Wilk, M. B. (1965), "An Analysis of Variance Test for Normality (Complete Samples)," *Biometrika*, **52**, 591–611.
- Shepard, R. N., Romney, A. K., and Nelove, S. B. (eds.) (1972), *Multidimensional Scaling: Theory and Applications in the Behavioral Sciences, Volume I: Theory*, New York: Seminar Press.
- Sibson, R. (1984), "Present Position and Potential Developments: Some Personal Views. Multivariate Analysis (with discussion)," *Journal of the Royal Statistical Society, Series A*, **147**, 198–207.
- Silverman, B. W. (1986), *Density Estimation for Statistics and Data Analysis*, New York: Chapman & Hall.
- Sinha, B. K. (1984), "Detection of Multivariate Outliers in Elliptically Symmetric Distributions," *Annals of Statistics*, **12**, 1558–1565.
- Siotani, M., Hayakawa, T., and Fujikoshi, Y. (1985), *Modern Multivariate Statistical Analysis*, Columbus, OH: American Sciences.
- Siotani, M., Yoshida, K., Kawakami, H., Nojiro, K., Kawashima, K., and Sato, M. (1963), "Statistical Research on the Taste Judgement: Analysis of the Preliminary Experiment on Sensory and Chemical Characters of SEISHU," *Proceedings of the Institute of Statistical Mathematics*, **10**, 99–118.

- Small, N. J. H. (1978), "Plotting Squared Radii," *Biometrika*, **65**, 657–658.
- Smith, D. W., Gill, D. S., and Hammond, J. J. (1985), "Variable Selection in Multivariate Multiple Regression," *Journal of Statistical Computation and Simulation*, **22**, 217–227.
- Snee, R. D. (1972), "On the Analysis of Response Curve Data," *Technometrics*, **14**, 47–62.
- Snee, R. D., Acuff, S. J., and Gibson, J. R. (1979), "A Useful Method for the Analysis of Growth Studies," *Biometrics*, **35**, 835–848.
- Sokal, R. R. and Rohlf, F. J. (1981), *Biometry: The Principles and Practices of Statistics in Biological Research*, 2nd ed., San Francisco, CA: W. H. Freeman and Co.
- Sparks, R. S., Coutsourides, D., and Troskie, L. (1983), "The Multivariate C_p ," *Communications in Statistics—Part A Theory and Methods*, **12**(15), 1775–1793.
- Steiger, J. H. and Lind, J. C. (1980), "Statistically Based Tests for the Number of Common Factors," paper presented at the annual meeting of the Psychometric Society, Iowa City, IA.
- Strauss, J. S., Bartko, J. J., and Carpenter, W. T. (1973), "The Use of Clustering Techniques for the Classification of Psychiatric Patients," *British Journal of Psychiatry*, **122**, 531–540.
- Swayne, D. F., Lang, D. T., Buja, A., Cook, D. (2003), "GGobi: Evolving from XGobi into an Extensible Framework for Interactive Data Visualization," *Computational Statistics & Data Analysis*, **43**, 423–444.
- Therneau, T. M. and Atkinson, B. (1997), *An Introduction to Recursive Partitioning Using the RPART Routines*, Technical Report 61, Section of Biostatistics, Mayo Clinic, Rochester, MN, URL: <http://www.mayo.edu/hsr/tech rpt/61.pdf>.
- Thomas, D. R. (1983), "Univariate Repeated Measures Techniques Applied to Multivariate Data," *Psychometrika*, **48**, 451–464.
- Thomson, G. H. (1951), *The Factorial Analysis of Human Ability*, London: London University Press.
- Timm, N. H. (1975), *Multivariate Analysis: With Applications in Education and Psychology*, Monterey, CA: Brooks/Cole.
- Timm, N. H. (1980), "Multivariate Analysis of Variance of Repeated Measures," In *Handbook of Statistics*, Vol. 1, P. R. Krishnaiah (ed.), Amsterdam: North-Holland, pp. 41–87.
- Tintner, G. (1946), "Some Applications of Multivariate Analysis to Economic Data," *Journal of the American Statistical Association*, **41**, 472–500.
- Titterington, D. M., Murray, G. D., Murray, L. S., Speigelhalter, D. J., Skene, A. M., Habbema, J. D. F., and Gelpke, D. J. (1981), "Comparison of Discrimination Techniques Applied to a Complex Data Set of Head Injured Patients," *Journal of the Royal Statistical Society, Series A*, **144**, 145–175.
- Travers, R. M. W. (1939), "The Use of a Discriminant Function in the Treatment of Psychological Group Differences," *Psychometrika*, **4**(1), 25–32.
- Tu, C. T., and Han, C. P. (1982), "Discriminant Analysis Based on Binary and Continuous Variables," *Journal of the American Statistical Association*, **77**, 447–454.

- Venables, W. N. (1976), "Some Implications of the Union Intersection Principle for Test of Sphericity," *Journal of Multivariate Analysis*, **6**, 185–190.
- Vonesh, E. F. (1986), "Sample Sizes in the Multivariate Analysis of Repeated Measurements," *Biometrics*, **42**, 601–610.
- Wainer, H. (1981), "Comments on a Paper by Kleiner and Hartigan," *Journal of the American Statistical Association*, **76**, 270–272.
- Wall, F. J. (1967), *The Generalized Variance Ratio or U-Statistic*, Albuquerque, NM: The Dikewood Corporation.
- Wang, C. M. (1983), "On the Analysis of Multivariate Repeated Measures Designs," *Communications in Statistics—Part A Theory and Methods*, **12**, 1647–1659.
- Ward, J. H. (1963), "Hierarchical Grouping to Optimize an Objective Function," *Journal of the American Statistical Association*, **58**, 236–244.
- Wegman, E. J. (1972), "Nonparametric Probability Density Estimation I: A Summary of Available Methods," *Technometrics*, **14**, 533–546.
- Wegman, E. J. (1990), "Hyperdimensional Data Analysis Using Parallel Coordinates," *Journal of the American Statistical Association*, **85**, 664–675.
- Welch, B. L. (1939), "Note on Discriminant Functions," *Biometrika*, **31**, 218–220.
- Wickham, H., Cook, D., Hofmann, H., Buja, A. (2011), "Tourr: An R Package for Exploring Multivariate Data with Projections," *Journal of Statistical Software*, **40**, 1–18.
- Wilks, S. S. (1932), "Certain Generalizations in the Analysis of Variance," *Biometrika*, **24**, 471–494.
- Wilks, S. S. (1946), "Sample Criteria for Testing Equality of Means, Equality of Variances and Equality of Covariances in a Normal Multivariate Distribution," *Annals of Mathematical Statistics*, **17**, 257–281.
- Wilks, S. S. (1963), "Multivariate Statistical Outliers," *Sankhya, Series A*, **25**, 407–426.
- Williams, E. J. (1970), "Comparing Means of Correlated Variates," *Biometrika*, **57**, 459–461.
- Williams, J. S., and Izenman, A. J. (1981), "A Class of Linear Spectral Models and Analysis for the Study of Longitudinal Data," Technical Report, Colorado State University, Department of Statistics.
- Wishart, D. (1969a), "An Algorithm for Hierarchical Classifications," *Biometrics*, **25**, 165–170.
- Wishart, D. (1969b), "Numerical Classification Method for Deriving Natural Classes," *Nature*, **221**, 97–98.
- Wishart, D. (1971), *A Generalised Approach to Cluster Analysis*. Part of Ph.D. Thesis, University of St. Andrews.
- Yang, S. S., and Lee, Y. (1987). "Identification of a Multivariate Outlier," Presented at the Annual Meeting of the American Statistical Association, San Francisco, CA, August 1987.
- Yenyukov, I. S. (1988), "Detecting Structures by Means of Projection Pursuit," *Compstat*, **88**, 47–58.
- Young, F. W. (1987), *Multidimensional Scaling: History, Theory and Applications*, R. M. Hammer (ed.), Hillsdale, NJ: Lawrence Erlbaum.

- Young, G. and Householder, A. S. (1938), "Discussion of a Set of Points in Terms of Their Mutual Distances," *Psychometrika*, **3**, 19–22.
- Zellner, A. (1962), "An Efficient Method of Estimating Seemingly Unrelated Regressions and Tests for Aggregation Bias," *Journal of the American Statistical Association*, **57**, 54–78.
- Zerbe, G. O. (1979a), "Randomization Analysis of the Completely Randomized Design Extended to Growth Curves," *Journal of the American Statistical Association*, **74**, 215–221.
- Zerbe, G. O. (1979b), "Randomization Analysis of Randomized Block Design Extended to Growth and Response Curves," *Communications in Statistics—Part A Theory and Methods*, **8**, 191–205.
- Zhang, L., Helander, M. G. and Drury, C. G. (1996), "Identifying Factors of Comfort and Discomfort in Sitting," *Human Factors*, **38**(3), 377–380.
- Zyskind, G. (1967), "On Canonical Forms, Non-Negative Covariance Matrices and Best and Simple Least Square Linear Estimator in Linear Models," *Annals of Mathematical Statistics*, **38**, 1092–1109.

INDEX

- Additional information, test for, 149–152, 241–243
- Air pollution data, 553
- Airline distance data, 559
- Algebra, matrix, *see* Matrix, algebra
- Analysis of variance, multivariate (MANOVA):
 additional information, test for, 241–243
 association, measures of, 186–188
 assumptions, checking on, 210–211
 and canonical correlation, 400–402
 contrasts, 192–195
 orthogonal, 194
 discriminant function, 178, 197–198, 203
 growth curves, 232–241, *see also* Growth curves; Repeated measures designs
 signs
H and **E** matrices, 173–174
 higher-order models, 207–208
 individual variables, tests on, 177–178, 195–198
 discriminant function, 178, 197–198, 203
 experimentwise error rate, 195–197
 protected tests, 196
 Lawley–Hotelling test, 180–181
- table of critical values, 627–636
- likelihood ratio test, 178
- mixed models, 208–210
 expected mean squares, 208
- multivariate association, measures of, 186–188
- one-way, 171–174
 contrasts, 192–195
 model, 172
 unbalanced, 181
- Pillai's test, 179
- table of critical values, 624–626
- profile analysis, 211–213
- repeated measures, 215–232, *see also* Repeated measures designs; Growth curves
- Roy's test (union-intersection), 178–179
- table of critical values, 621–623
- stepwise discriminant analysis, 243–244
- stepwise selection of variables, 243–244
- test for additional information, 241–243
- test statistics, 174–186
 comparison of, 182–183, 189–190
 and eigenvalues, 182
 power of, 189–190
 and T^2 , 182

- tests on a subvector, 241–243
 tests on individual variables, *see* individual variables, tests on
 two-way, 201–207
 contrasts, 202–203
 discriminant function, 203
 interactions, 201–202
 main effects, 201–202
 model, 201
 test statistics, 202
 tests on individual variables, 203
 unbalanced one-way, 181
 union-intersection test, 178–179
 Wilks' Λ (likelihood ratio) test, 174–178
 F approximation, 176
 chi-square approximation, 177
 partial Λ statistic, 242
 properties of, 175–178
 table of critical values, 613–636
 transformation to an exact F , 176
- Analysis of variance, univariate (ANOVA):**
 one-way, 169–171
 contrasts, 191–192
 SSH, SSE, F -statistic, 171
 two-way, 198–200
 F -test, 200
 contrasts, 200
 interactions, 199
 main effects, 199–200
 model, 198
- ANOVA, *see* Analysis of variance, univariate (ANOVA)**
- Association, measures of**, 186–188
Association, measures of, 372–374
Athletic record data, 530
- Baker corn field data**, 61
Bar steel data, 204
Beetles data, 162
Bilinear form, 21
Biplots, 580–588
 coordinates of points, 583–585
 correlation, 584
 cosine, 584, 587
 points for observations, 580–585
 points for variables, 580–585
 principal component approach, 581–582, 585
 singular value decomposition, 583, 585
Birth and death data, 593
Bivariate normal distribution, 51, 93, 98–99, 146
Blood data, 248
- Blood pressure data**, 255
Bonferroni critical values, 140
 table, 611
Box's M -test, 267–269
 table of exact critical values, 633–636
Box–Cox transformations, 109–111
Bronchus data, 167
Burt matrix, 577–579
Byssinosis data, 594, 595
- Calcium data**, 66
Calculator speed data, 222
Canonical correlation(s), 187, 271, 385–402
 canonical variates, *see* Canonical variates
 definition of, 387
 and discriminant analysis, 400–402
 and eigenvalues, 387, 400–402
 and MANOVA, 400–402
 dummy variables, 400
 and measures of association, 386, 398
 and multiple correlation, 386, 390, 400
 properties of, 390–391
 redundancy analysis, 398
 and regression, 393, 398–400
 subset selection, 400
 tests of significance, 391–395
 subset of canonical correlations, 393–395
 subset selection, 400
 test of a subset in regression, 399–400
 and test of independence, 391–392
 and test of overall regression, 391–392
 with all canonical correlations, 391–393
 with grouping variables, 187
 with test for independence of two subvectors, 271, 391–392
- Canonical variates:**
 correlations among, 388
 definition of, 387
 interpretation, 395–398
 by correlations (structure coefficients), 397
 by rotation, 397
 by standardized coefficients, 396–397
 redundancy analysis, 398
 and regression, 398–400
 standardized coefficients, 389, 396–397
- Categorical variables**, *see* Dummy variables
Central limit theorem (multivariate), 100, 109
CFA, *see* Confirmatory factor analysis

- Characteristic form:
- of T^2 -statistic, 131, 136
 - of t -statistic, 131, 135
- Characteristic roots, *see* Eigenvalues
- Chemical data, 358
- Chi-square distribution, 96, 101, 127
- Cholesky decomposition, 27
- City crime data, 507
- Classification analysis, 309–337
- assigning a sampling unit to a group, 309
 - asymptotic optimality, 313
 - classification trees, 331–335
 - impurity, 332
 - prior probabilities, 333
 - correct classification rates, 318–320
 - error rates, 318–326, *see also* Error rate(s)
 - as a stopping rule, 323–326
 - k -nearest neighbor rule, 330–331
 - nonparametric classification procedures, 312, 326–335
 - classification trees, 331–335
 - density estimators (kernel), 327–330
 - multinomial data (categorical variables), 326
 - nearest neighbor rule, 330–331
 - several groups, 314–318
 - linear classification functions, 315
 - optimal classification rule (Welch), 315
 - quadratic classification functions, 317
 - subset selection, 322–326
 - stepwise discriminant analysis, 323
 - two groups, 310–314
 - Fisher's classification function, 311
 - linear classification function, 311
 - misclassification costs, 312
 - optimal classification rule (Welch), 313
 - prior probabilities, 312
- Cluster analysis, 501–552
- average linkage method, 511
 - centroid method, 514
 - choosing the number of clusters, 544
 - and classification, 501
 - clustering observations, 501–547
 - clustering variables, 501, 547
 - comparison of methods, 527
 - complete linkage method, 508
 - definition, 501
 - dendrogram, 506
 - crossover, 521
 - examples of, 509, 510, 512, 513, 515–517, 519, 522, 523, 526, 527
 - inversion, 521
 - reversal, 521
 - dissimilarity, 503
 - distance, 502–504
 - distance matrix, 503
 - Euclidean distance, 503
 - Minkowski metric, 503
 - profile of observation vector, 504
 - scale of measurement, 503
 - statistical distance, 503
 - farthest neighbor method, *see* complete linkage method
 - flexible beta method, 520
 - hierarchical clustering, 502, 505–531
 - agglomerative method, 506–529
 - comparison of methods, 527
 - dendrogram, 506
 - divisive method, 505, 529
 - properties, 521–529
 - incremental sum of squares method, *see* Ward's method
 - median method, 514
 - nearest neighbor method, *see* single linkage method
 - nonhierarchical methods, 531–544
 - density estimation, 543
 - mixtures of distributions, 540
 - partitioning, 532–540
 - optimization methods, *see* nonhierarchical methods, partitioning
 - partitioning, 502, 532–540
 - plotting of clusters:
 - discriminant functions, 535–538, 544
 - principal components, 502, 534
 - projection pursuit, 502
 - profile of observation vector, 504
 - similarity, 502
 - single linkage method, 506
 - tree diagram, *see* dendrogram
 - validity of a cluster solution, 546
 - cross-validation, 546
 - hypothesis test, 546
 - variables, clustering of, 501, 547
 - correlations, 547
 - and factor analysis, 547
 - Ward's method, 517
- Coated pipe data, 148
- Coefficient of determination, *see* R^2
- Commensurate variables, *see* Variables, commensurate
- Commonality, *see* Confirmatory factor analysis; Exploratory factor analysis

- Confidence interval (reference), 132, 140
 Confirmatory factor analysis, 479–500
 covariance matrix under the model, 481
 errors-in-variables parameterization, 483
 estimation, 487–489
 least squares, 488–489
 likelihood ratio tests, 487–488
 maximum likelihood, 487–488
 and exploratory factor analysis, 479–480, 482
 factor scores, 495
 regression method formula, 495
 weighted least squares formula, 495
 goodness-of-fit criteria, 489–492
 Bentler's comparative fit index (CFI), 490
 χ^2 test, 489–490
 root mean square error approximation (RMSEA), 490
 standardized root mean square residual (SRMR), 490
 hypotheses, 484, 489
 identified model, 479, 482–486, 496
 indeterminacy, 482, 483
 order condition, 482–484
 underidentified, 482
 indicator variable, 483
 inference, 492–494
 confidence intervals, 493
 hypothesis tests, 493
 model selection and refinement, 493
 and item response theory, 480
 measurement error parameterization, 483
 model, 480–481
 normality assumption, 487
 parameter space, 487
 parameters, 481
 path diagram, 481, 486
 and regression, 481, 493
 and structural equation modeling, 480
 Contingency table:
 graphical analysis of, *see* Correspondence analysis
 higher-way table, 576
 two-way table, 565, 570, 572
 Contour plots, 93
 Contrasts:
 contrast matrices in growth curves, 233–236, 239
 contrast matrices in repeated measures, 218–232
 one-sample profile analysis, 152
 one-way ANOVA, 191
 one-way MANOVA, 192
 orthonormal, 218
 two-sample profile analysis, 154
 two-way ANOVA, 200
 two-way MANOVA, 202
 Cork data, 249
 Corn field data, 61
 Correct classification rate, 318–320
 Correlation matrix:
 and covariance matrix, 69
 factor analysis on, 445
 partitioned, 389
 population correlation matrix, 70
 principal components from, 409, 419–423
 sample correlation matrix, 69
 Correlation:
 canonical, *see* Canonical correlation(s)
 and cosine of angle between two vectors, 53
 intraclass correlation, 210
 and law of cosines, 53
 multiple, *see* Multiple correlation
 and orthogonality of two vectors, 54
 population correlation (ρ), 53
 sample correlation (r), 53
 of two linear combinations, 76, 81
 Correspondence analysis, 565–579
 contingency table, *see* Contingency table
 coordinates for row and column points, 572–575
 distances between column points, 574
 distances between row points, 574
 singular value decomposition, 572
 correspondence matrix, 566
 definition of (graph of contingency table), 565
 independence of rows and columns, testing, 570
 chi-square, 565, 570
 inertia, 566, 574
 multiple correspondence analysis, 576–579
 Burt matrix, 577–579
 indicator matrix, 576
 profiles of rows and columns, 566–570
 rows and columns, 565–575
 inertia, 566, 574
 interaction, 565
 points for plotting, 572–575
 profiles, 566–570
 singular value decomposition, 572, 575
 generalized singular value decomposition, 573

- Covariance:
- and independence, 50
 - and orthogonality, 52
 - population covariance (σ_{xy}), 49
 - sample covariance (s_{xy}), 51
 - expected value of, 51
 - and linear relationships, 51
 - of two linear combinations, 76, 80
- Covariance matrix:
- compound symmetry, 217
 - and correlation matrix, 69
 - of linear combinations of variables, 78–81
 - partitioned, 71–75, 386
 - dependence of \mathbf{y} and \mathbf{x} and $\text{cov}(\mathbf{y}, \mathbf{x})$, 72
 - difference between $\text{cov}(\mathbf{y})$ and $\text{cov}(\mathbf{y}, \mathbf{x})$, 72
 - three or more subsets, 73
 - pooled covariance matrix, 136
 - population covariance matrix (Σ), 68
 - sample covariance matrix (S), 66
 - from data matrix, 67
 - distribution of, 100
 - from observations, 67
 - positive definiteness of, 76
 - and sample mean vector, independence of, 101
 - sphericity, 218, 261
 - tests on, 259–280, *see also* Tests of hypothesis, covariance matrices
 - unbiasedness of, 69
 - uniformity, 217, 263
- Cross-validation, 322
- Cyclical data, 166
- Data matrix (\mathbf{Y}), 64
- Data sets:
- air pollution data, 553
 - airline distance data, 559
 - athletic record data, 530
 - Baker corn field, 61
 - bar steel data, 204
 - beetles data, 162
 - birth and death data, 593
 - blood data, 248
 - blood pressure data, 255
 - bronchus data, 167
 - byssinosis data, 594, 595
 - calcium data, 66
 - calculator speed data, 222
 - chemical data, 358
 - city crime data, 507
 - coated pipe data, 148
 - cork data, 249
 - cyclical data, 166
 - dental data, 238
 - diabetes data, 74
 - do-it-yourself data, 580
 - dogs data, 254
 - dystrophy data, 165
 - engineer data, 163
 - fabric wear data, 249
 - fish data, 246
 - football data, 291–292
 - glucose data, 90
 - guinea pig data, 213
 - height-weight data, 49
 - hematology data, 121, 122
 - mandible data, 257
 - mice data, 252
 - Norway crime data, 594
 - people data, 576
 - perception data, 446
 - piston ring data, 569
 - plasma data, 256
 - politics data, 592
 - probe word data, 79
 - protein data, 533
 - psychological data, 138
 - ramus bone data, 87
 - repeated data, 229
 - Republican vote data, 58
 - road distance data, 591
 - rootstock data, 184
 - Seishu data, 274
 - snap bean data, 247
 - sons data, 88
 - statistics course grade data, 485
 - steel data, 285
 - survival data, 250, 251
 - temperature data, 279
 - trout data, 252
 - voting data, 563
 - weight gain data, 253
 - wheat data, 554
 - words data, 167
- Data, types of, 4, *see also* Multivariate data
- Density function, 47
- Dental data, 238
- Descriptive statistics, 2
- Determinant, 28–30
 - definition of, 28
 - of diagonal matrix, 29
 - of inverse, 30
 - of nonsingular matrix, 29
 - of partitioned matrix, 30
 - of positive definite matrix, 29

- as product of eigenvalues, 34
of product, 29
of scalar multiple of a matrix, 29
of singular matrix, 29
- Diabetes data, 74
- Diagonal matrix, 10
- Discriminant analysis (descriptive), 281–306
and canonical correlation, 292, 400–402
and classification analysis, 282
discriminant functions:
for several groups, 178, 197–198, 203, 288–292
for two groups, 139–145, 282–286
and eigenvalues, 289–290
interpretation of discriminant functions, 298–301
correlations (structure coefficients), 300
partial F -values, 299
rotation, 301
standardized coefficients, 298
purposes of, 288
scatterplots, 301–302
selection of variables, 243, 303–306
several groups, 288–290
standardized discriminant functions, 292–294
stepwise discriminant analysis, 243, 303–306
tests of significance, 294–298
two groups, 282–286
and multiple regression, 144–145, 286–288
- Discriminant analysis (predictive), *see* Classification analysis
- Dispersion matrix, *see* Covariance matrix
- Distance between vectors, 84–85, 93, 128, 131, 136, 282
- Distribution:
beta, 107
bivariate normal, 49, 93, 98–99
chi-square, 96
elliptically symmetric, 115
 F , 132, 151, 171, 176, 192, 265–266
multivariate normal, *see* Multivariate normal distribution; Multivariate normality, tests for
univariate normal, 92, 95
tests for, *see* Univariate normality, tests for
Wishart, 100–101
- Do-it-yourself data, 580
- Dogs data, 254
- Dummy variables, 186, 292, 327, 400
- Dynamic graphics, 58–60
grand tour, 58, 107
linked brushing, 59
parallel coordinates plot, 59
- Dystrophy data, 165
- E matrix, 173, 356, 364–366
- EFA, *see* Exploratory factor analysis
- Eigenvalues, 32–37, 182, 387–390, 407–409, 423, 443
- Eigenvectors, 32–37, 387–390, 407–409, 444
- Elliptically symmetric distribution, 115
- EM algorithm, 83, 541
- Engineer data, 163
- Error rate(s), 318–322
actual error rate, 319
apparent error rate, 319
bias in, 319, 320
classification table, 319
cross-validation, 322
experimentwise error rate, 3, 141, 196
holdout method, 322, 330
leaving-one-out method, 322, 330
partitioning the sample, 321
resubstitution, 319
- Expected value:
of random matrix, 68
of random vector [$E(\mathbf{y})$], 65
of sample covariance matrix [$E(\mathbf{S})$], 69
of sample mean [$E(\bar{\mathbf{y}})$], 48
of sample mean vector [$E(\bar{\mathbf{y}})$], 65
of sample variance [$E(s^2)$], 48
of sum or product of random variables, 50
of univariate random variable [$E(y)$], 48
- Experimental units, 1
- Exploratory factor analysis, 435–477, 479
assumptions, 437–439
failure of assumptions, consequences of, 441, 470
common factors, 437
communalities, 440, 444, 448–449, 454
estimation of, 444, 449, 450, 454
and confirmatory factor analysis, 479–480, 482
eigenvalues, 443–446, 448–450, 453, 469, 473
eigenvectors, 443–445, 447, 449
factor scores, 466–469
averaging method, 468
regression method, 466–468
- factors, 441
common, 437

- definition of, 435
- interpretation of, 436, 465
- number of, 453–457
- Heywood case, 451
- loadings:
 - definition of, 437
 - estimation of, 442–453
- model, 437–442
- modeling covariances or correlations, 435, 438, 440, 441, 444
- number of factors to retain, 453–457
 - average eigenvalue, 453–455
 - comparison of methods, 455–457
 - hypothesis test, 454–455
 - indeterminacy for certain data sets, 455
 - scree plot, 453–455
 - variance accounted for, 453–455
- orthogonal factors, 437–442, 458–462
- and principal components, 436, 475–476
- and regression, 437, 466–468
- rotation, 441–442, 444, 457–466
 - complexity of the variables, 458
 - interpretation of factors, 436, 465
 - oblique rotation, 458, 462–465
 - orthogonal rotation, 457–462
 - simple structure, 458
 - scree plot, 453–455
 - simple structure, 458
- singular matrix and, 449
- specific variance, 437, 444
- total variance, 445, 453
- validity of factor analysis model, 470–475
 - how well the model fits the data, 445, 470
 - measure of sampling adequacy, 472
- variance due to a factor, 445
- F*-test(s):
 - ANOVA, 171, 200
 - between-subjects tests in repeated measures, 223, 227, 231
 - comparing two variances, 265
 - contrasts, 192
 - equivalent to T^2 , 132, 137, 150
 - in multiple regression, 150, 347
 - partial *F*-test, 140, 151, 242, 303
 - stepwise selection, 243, 303–306, 352–354
 - test for additional information, 149
 - test for individual variables in MANOVA, 195–198
- Wilks' Λ :
 - exact *F* transformation for, 176
 - F* approximation for, 176
- Fabric wear data, 249
- Factor analysis, *see* Exploratory factor analysis, Confirmatory factor analysis
- Fish data, 246
- Fisher's classification function, 311
- Football data, 291–292
- Gauss–Markov theorem, 343, 359
- Generalized population variance, 93, 118
- Generalized sample variance, 81
 - total sample variance, 82, 408, 436, 445, 453
- Generalized singular value decomposition, 573
- Geometric mean, 187
- Glucose data, 90
- Grade data, 485
- Grand tour, 58, 107
- Graphical display of multivariate data, 56–58
- Graphical procedures, 555–596
 - biplots, *see* Biplots
 - correspondence analysis, *see* Correspondence analysis
 - multidimensional scaling, *see* Multidimensional scaling
- Growth curves, 232–241
 - contrast matrices, 233–236, 238–241
 - for one sample, 232–239
 - orthogonal polynomials, 232–236
 - polynomial function of t , 236–237
 - several samples, 239–241
 - unequally spaced time points, 236–237
- Guinea pig data, 213
- H matrix, 173, 364
- Height-weight data, 49
- Hematology data, 121, 122
- Hierarchical clustering, *see* Cluster analysis, hierarchical clustering
- Hotelling's T^2 -statistic, *see* T^2 -statistic
- Hotelling-Lawley test statistic, *see* Lawley-Hotelling test statistic
- Hyperellipsoid, 82
- Hypothesis tests, *see* Tests of hypotheses
- Identity matrix, 10
- Imputation, 82–84
- Independence of variables, test for, 275–276
 - table of exact critical values, 635
- Indicator variables, *see* Dummy variables
- Inferential statistics, 2
- Intraclass correlation, 210

- Kernel density estimators, 327–330
 Kronecker and vec notation, 37–39, 45, 356
 Kurtosis, 103, 107, 115
- Largest root test, *see* Roy's test statistic
 Latent roots, *see* Eigenvalues
 Lawley-Hotelling test statistic:
 definition of, 188
 table of critical values, 627–631
 Length of vector, 16
 Likelihood function, 99
 Likelihood ratio tests:
 in confirmatory factor analysis, 487–488
 for covariance matrices, 260, 261, 264,
 267, 270, 272, 275
 in exploratory factor analysis, 455
 for mean vectors, 139, 178
- Linear classification functions, 311–317
 Linear combination of matrices, 21
 Linear combination of vectors, 21
 Linear combination(s) of variables, 2, 75–81,
 126
 correlation matrix for several linear combinations, 80
 correlation of two linear combinations, 76, 80
 covariance matrix for several linear combinations, 78, 80
 covariance of two linear combinations, 76, 80
 distribution of, 94
 mean of a single linear combination, 76,
 79
 mean vector for several linear combinations, 77
 variance of a single linear combination, 76, 79
- Linear hypotheses, 154–155, 211–213, 219–241
 Linked brushing, 59
- Mahalanobis distance, 84–85, 93
 Mandible data, 257
 MANOVA, 143, 171, *see also* Analysis of variance, multivariate (MANOVA)
- Matrix (matrices):
 algebra of, 7–37
 bilinear form, 21
 Burt matrix, 577–579
 Cholesky decomposition, 27
 conformable, 13
 covariance matrix, 66–69
 definition, 8
 determinant, 28–30, *see also* Determinant
 diagonal, 10
 eigenvalues, 32–37, *see also* Eigenvalues
 characteristic equation, 33
 and determinant, 34
 of $\mathbf{I} + \mathbf{A}$, 34
 of inverse matrix, 36
 of positive definite matrix, 35
 of product, 35
 singular value decomposition, 37
 spectral decomposition, 35
 of square matrix, 36
 of square root matrix, 36
 of symmetric matrix, 35
 and trace, 34
 eigenvectors, 32–37, *see also* Eigenvectors
 equality, 9
 identity, 10
 indicator matrix, 576
 inverse, 25–26
 of partitioned matrix, 26
 of product, 25
 of transpose, 26
 J matrix, 11
 j vector, 11
 linear combination of, 21
 nonsingular matrix, 25
 notation for matrix and vector, 8
 O (zero matrix), 11
 operations with, 11–21
 distributive law, 14
 factoring, 17
 product, 13–21
 sum, 12
 orthogonal, 31
 rotation of axes, 32
 partitioned matrices, 22–23
 determinant of, 30
 inverse of, 26
 product of, 22
 transpose of, 23
 Perron–Frobenius theorem, 35, 428
 positive definite, 26, 35
 positive semidefinite, 26, 35
 quadratic form, 21
 rank, 23
 full rank, 24
 scalar, 8
 product of scalar and matrix, 21
 singular matrix, 25
 singular value decomposition, 37

- size of a matrix, 8
 spectral decomposition, 35
 square root matrix, 36
 sum of products in vector notation, 16
 sum of squares in vector notation, 16
 symmetric, 10, 35
 trace, 31, 34, 78
 and eigenvalues, 34
 of product, 31
 of sum, 31
 transpose, 9
 of product, 14
 of sum, 12
 triangular, 11
 vectors, *see* Vector(s)
 zero matrix ($\mathbf{0}$) and zero vector ($\mathbf{0}$), 11
- Maximum likelihood estimation, 99–100
 and confirmatory factor analysis, 487–488
 of correlation matrix, 100
 of covariance matrix, 99
 likelihood function, 99
 of mean vector, 99
 multivariate normal, 99
- Mean vector, 63–66, 92, 99
 notation, 64
 population mean vector (μ), 65
 sample mean vector ($\bar{\mathbf{y}}$), 64–66
 from data matrix, 65
 distribution of, 100
 and sample covariance matrix, independence of, 101
- Mean:
 geometric, 187
 of linear function, 76
 population mean (μ), 48
 of product, 50
 sample mean ($\bar{\mathbf{y}}$), 48
 of sum, 50
- Measurement scale, 3
 interval scale, 3
 ordinal scale, 3
 ratio scale, 3
- Mice data, 252
- Misclassification rates, *see* Error rate(s)
- Missing values, 82–84
- Multicollinearity, 82, 93
- Multidimensional scaling, 555–565
 classical solution, *see* metric multidimensional scaling
 definition, 555–556
 distances, 555
 seriation (ranking), 556
- metric multidimensional scaling, 556–560
 algorithm for finding the points, 556
 and principal components analysis, 557
- nonmetric multidimensional scaling, 560–565
 monotonic regression, 560
 ranked dissimilarities, 560
 STRESS, 561–562
- principal coordinate analysis, *see* metric multidimensional scaling
 spectral decomposition, 557
- Multiple correlation, 349, 386, 450, *see also R²*
- Multiple correspondence analysis, 576–579
 Burt matrix, 577–579
 column coordinates, 577
 indicator matrix, 576
- Multiple regression, *see* Regression, multiple
- Multivariate analysis, 1
 descriptive statistics, 2
 inferential statistics, 2
- Multivariate analysis of variance (MANOVA), *see* Analysis of variance, multivariate (MANOVA)
- Multivariate data:
 basic types of, 4
 plotting of, 56
 sparseness of, 106
- Multivariate inference, 2
- Multivariate normal distribution, 91–117
 applicability of, 93
 conditional distribution, 97
 contour plots, 93
 density function, 92
 distribution of $\bar{\mathbf{y}}$ and \mathbf{S} , 100
 features of, 91
 independence of $\bar{\mathbf{y}}$ and \mathbf{S} , 101
 linear combinations of, 94
 marginal distribution, 96
 maximum likelihood estimates, 99–100,
 see also Maximum likelihood estimation
 properties of, 94–98
 quadratic form and chi-square distribution, 96
 standardized variables, 95
 zero covariance matrix implies independence of subvectors, 97
- Multivariate normality, tests for, 101, 106–108
 D_i^2 , 106, 114
 and chi-square, 107
 table of critical values, 606

- scatterplots, 107
skewness and kurtosis, multivariate, 107,
115
table of critical values, 602–605
- Multivariate normality, transformations to, 110–
111
- Multivariate regression, *see* Regression, multivariate
- Nonsingular matrix, 25
- Normal distribution:
bivariate normal, 51, 93, 98–99, 146
multivariate normal, *see* Multivariate normal distribution
univariate normal, 92, 95
- Normality, tests for, *see* Multivariate normality; Univariate normality
- Normality, transformations to, *see* Multivariate normality; Univariate normality
- Norway crime data, 594
- Numerical taxonomy, *see* Cluster analysis
- Objectives of this book, 3
- Observations, 1
- One-sample test for a mean vector, 130–134
- Orthogonal matrix, 31
- Orthogonal polynomials, 232–236
table of, 632
- Orthogonal vectors, 54
- Outliers, multivariate
kurtosis, 115–116
elliptically symmetric distributions, 115
and principal components, 415
slippage in mean, variance, and correlation, 113
Wilks' statistic, 114–115
- Outliers, univariate
accommodation, 112
block test, 113
identification, 112
masking, 113
maximum studentized residual, 112
skewness and kurtosis, 113
slippage in mean and variance, 112
swamping, 113
- Overall probability, 81–82
- Paired observation test, 145–149
- Parallel coordinates plot, 59
- Partial *F*-tests, 140, 151, 242, 303–306
- Partitioned matrices, *see* Matrix (matrices), partitioned matrices
- Partitioning, *see* Cluster analysis, partitioning
- Pattern recognition, *see* Cluster analysis
- People data, 576
- Perception data, 446
- Perron–Frobenius theorem, 35, 428
- Pillai's test statistic:
definition of, 179
table of critical values, 624–626
- Piston ring data, 569
- Plasma data, 256
- Plotting multivariate data, 56–58
- Politics data, 592
- Positive definite matrix, 26
positive definite sample covariance matrix, 76
- Prerequisites for this book, 3
- Principal components, 405–433
algebra of, 410–412
and biplots, 581–582
and cluster analysis, 415, 417–419, 421, 533, 534
component scores, 411
definition of, 405, 407
dimension reduction, 406–412, 414
eigenvalues and eigenvectors, 407–409, 423
major axis, 409, 414
and factor analysis, 429, 436, 475–476
geometry of, 406–409
interpretation of, 427–430
correlations, 429–430
rotation, 429
special patterns in **S** or **R**, 427–429
large variance of a variable, effect of, 409, 428
last few principal components, 408, 415, 427
maximum variance, 405, 410
minimum perpendicular distances to line, 412–414
number of components to retain, 423–426
orthogonality of, 406, 408
percent of variance, 408, 423
and perpendicular regression, 409, 412–414
plotting of, 414–419
assessing normality, 414
detection of outliers, 415
properties of, 406–411
proportion of variance, 408
robust, 415
as rotation of axes, 406–409
from **S** or **R**, 408–409, 419–423

- nonuniqueness of components from **R**, 422
- sample specific components, 423
- scale invariance, lack of, 408
- scree graph, 423–424
- selection of variables, 430–432
- singular matrix and, 410–411
- size and shape, 428
- smaller principal components, 408, 415, 427
- tests of significance for, 423–426
- variable specific components, 423
- variances of, 408
- Probe word data, 79
- Product notation (\prod), 12
- Profile, 152
 - profile of observation vector, 504
- Profile analysis, 152–160, 211–215
 - and contrasts, 155
 - one-sample, 152–154
 - and one-way ANOVA, 152
 - profile, definition of, 152
 - and repeated measures, 152
 - several-sample, 211–215
 - two-sample, 154–160
 - and two-way ANOVA, 156
- Projection pursuit, 502
- Protein data, 533
- Psychological data, 138
- Q-Q* plot, 101–103
- Quadratic classification functions, 317–318
- Quadratic form, 21
- Quantiles, 101–103, 107
- R^2 (squared multiple correlation), 349–350, 353, 372, 378, 386, 389, 399, 400, 449
- Ramus bone data, 87
- Random variable(s):
 - bivariate, 49
 - bivariate normal distribution, 51, 93, 98–99
 - correlation of, 53–55
 - covariance of, 49–52
 - independent, 50
 - orthogonal, 52
 - scatterplot, 55–56
 - linear combinations, *see* Linear combination(s) of variables
 - univariate, 47
 - expected value of, 48
 - mean of, 48
 - variance of, 48
- vector, 63–66
- Random vector(s), 63–66
 - distance between, 84–85, 93, 128, 131, 136, 282
 - linear functions of, 75–81, *see also* Linear combination(s) of variables
 - mean of, 64–66
 - partitioned random vector, 71–75
 - standardized, 95
 - subvectors, 71–75
- Rank of a matrix, 23
- Rao's paradox, 130
- Redundancy analysis, 398
- Regression, monotonic, 560
- Regression, multiple (one y and several x 's), 144–145, 340–354, *see also* Regression, multivariate
 - centered x 's, 344–346
 - estimation of β :
 - centered x 's, 344–345
 - covariances, 345–346
 - least squares, 342–343
 - estimation of σ^2 , 343–344
 - fixed x 's, 340–354
 - Gauss–Markov theorem, 343
 - model, 340–341
 - assumptions, 340–341
 - corrected for means (centered), 344–346
 - multiple correlation, 349
 - R^2 , *see* R^2 (squared multiple correlation)
 - random x 's, 340, 354
 - regression coefficients, 340
 - SSE, 342–343, 347–348, 350–353
 - SSR, 347–348
 - subset selection, 350–354
 - all possible subsets, 350–352
 - stepwise selection, 352–354
 - tests of hypotheses, 346–349
 - full and reduced model, 347–349
 - overall regression test, 347
 - partial F -test, 348–349
 - subset of the β 's, 347–349
 - variables:
 - dependent (y), 339
 - independent (x), 339
 - predictor (x), 339
 - response (y), 339
- Regression, multivariate (several y 's and several x 's), 339–340, 354–381
 - association, measures of, 372–374
 - centered x 's, 361–362
 - estimation of Σ , 360

- estimation of **B** (matrix of coefficients):
 centered x 's, 361–362
 covariances, 362
 generalized least squares, 359–360,
 363
 least squares, 356–357
 properties of estimators, 358–360
 for seemingly unrelated regressions,
 362–363
 fixed x 's, 354–370
Gauss–Markov theorem, 359
model, 354–356
 assumptions, 355
 corrected for means (centered), 361–
 362
prediction, 370–372
 confidence interval for $E(y_0)$, 370–
 371
 prediction interval for y_0 , 371–372
random x 's, 380
regression coefficients, matrix of (**B**),
 97, 355
 seemingly unrelated regressions, 362–
 363
subset selection, 374–380
 all possible subsets, 377–380
 stepwise procedures, 374–377
tests of hypotheses, 364–370
E matrix, 364–365
 full and reduced model, 367–370,
 376–377, 399–400
H matrix, 364–365
 overall regression test, 364–367
 subset of the x 's, 374–376, 399, 400
 subset of the y 's, 376–377
Zyskind condition, 381
- Repeated data set**, 229
- Repeated measures designs**, 230–232, *see also*
 Growth curves
 assumptions, 215–219
 computation of test statistics, 224
 contrast matrices, 218–232
 doubly multivariate data, 232
 higher-order designs, 224–232
 multivariate approach, advantages of,
 215–217
 one sample, 219–222
 likelihood ratio test, 221
 and randomized block designs, 219
 and profile analysis, 152
 several samples, 222–223
 univariate approach, 215–219
- Republican vote data**, 58
- Research units**, 1
- Road distance data, 591
- Rootstock data, 184
- Rotation**, *see* Confirmatory factor analysis;
 Exploratory factor analysis
- Roy's test statistic**:
 definition of, 178
 table of critical values, 621–623
- Sampling units**, 1
- Scalar**, 9
- Scale of measurement**, 2
- Scatterplot**, 55–56, 107, 117
- Seishu data**, 274
- Selection of variables**, 243, 350–354, 374–380
- Singular value decomposition**, 37, 572, 575,
 583
 generalized singular value decomposi-
 tion, 573
- Size and shape**, 428
- Skewness**, 103, 107, 116
- Snap bean data**, 247
- Sons data**, 88
- Specific variance**, *see* Confirmatory factor anal-
 ysis; Exploratory factor analysis
- Spectral decomposition**, 35, 407, 443, 557
- Squared multiple correlation**, *see* R^2
- Standard deviation**, 48
- Standardized vector**, 95
- Statistics course grade data**, 485
- Steel data**, 285
- Stepwise selection of variables**, 243, 352–354,
 374–377
- STRESS**, 561
- Subvectors**, 71–75
 conditional distribution of, 97
 covariance matrix of, 71–75
 distribution of sum of, 97
 independence of, 72, 97
 mean vector, 71, 75
 tests of, 149–152, 241–243, 367–370,
 376–377
- Summation notation** (\sum), 11
- Survival data**, 250, 251
- t-tests**:
 characteristic form, 131, 135
 contrasts, 192
 equal levels in profile analysis, 157
 growth curves, 235, 238
 matched pairs, 145–146
 one sample, 130
 paired observations, 145–146
 repeated measures, 222
 two samples, 134, 140

- T^2 -statistic:
- additional information, test for, 149–152
 - assumptions for, 135
 - characteristic form, 131, 136
 - chi-square approximation for, 133
 - computation of, 143–145
 - by MANOVA, 143
 - by regression, 144–145
 - and F -distribution, 132, 137, 150
 - full and reduced model test, 149
 - likelihood ratio test, 139
 - matched pairs, 147–149
 - one-sample, 131–134
 - paired observations, 147–149
 - and profile analysis, 152–160
 - one sample, 152–154
 - two samples, 154–160
 - properties of, 132–133, 136–137
 - for a subvector, 149–152
 - table of critical values for T^2 , 607–610
 - two-sample, 135–139
- Taxonomy, numerical, *see* Cluster analysis
- Temperature data, 279
- Tests of hypotheses:
- accepting H_0 , 131
 - for additional information, 149–152, 241–243, 367–370, 376–380
 - partial F -tests, 140, 150, 242
 - covariance matrices, 259–280
 - one covariance matrix, 259–265
 - several covariance matrices, 265–269
 - on individual variables, 139–143
 - Bonferroni critical values for, 140
 - discriminant functions, 139–145
 - experimentwise error rate, 141
 - partial F -tests, 140, 242
 - protected tests, 141
 - likelihood ratio test, 139, *see also* Likelihood ratio tests
 - for linear combinations:
 - one sample ($H_0: \mathbf{C}\boldsymbol{\mu} = \mathbf{0}$), 130, 153, 219–222
 - two samples ($H_0: \mathbf{C}\boldsymbol{\mu}_1 = \mathbf{C}\boldsymbol{\mu}_2$), 155
 - mean vectors:
 - likelihood ratio tests, 139
 - one sample, Σ known, 127–130
 - one sample, Σ unknown, 131–134
 - several samples, 171–186
 - two-sample T^2 -test, 135–139
 - multivariate vs. univariate testing, 2–3, 125–130, 140–143
 - paired observations (matched pairs), 145–149
 - multivariate, 147–149
 - univariate, 145–146
 - partial F -tests, 140, 150, 242
 - power of a test, 126
 - protected tests, 141
 - on regression coefficients, 346–349, 364–370
 - on a subvector, 149–152, 241–243, 367–370, 376–381
 - univariate tests:
 - ANOVA F -test, 169–171, 198–200
 - one-sample test on a mean, σ known, 126
 - one-sample test on a mean, σ unknown, 130
 - paired observation test, 145–146
 - tests on variance, 265–266
 - two-sample t -test, 134–135, 140
 - variances, equality of, 265–266
 - Total sample variance, 82, 408, 436, 445, 453
 - Trace of a matrix, 31, 34, 78
 - Trout data, 252
 - Two-sample test for equal mean vectors, 135–139
 - Union-intersection test, 178–179
 - Unit:
 - experimental, 1
 - research, 1
 - sampling, 1
 - Univariate normal distribution, 92, 95
 - Univariate normality, tests for, 101–106
 - D'Agostino's D -statistic, 105
 - table of critical values, 601
 - goodness-of-fit test, 106
 - normal probability paper, 103
 - $Q-Q$ plot, 101–103
 - quantiles, 101–103
 - skewness and kurtosis, 103–104
 - tables of critical values, 598–636
 - transformation of correlation, 105
 - Univariate normality, transformations to, 109–110
 - Variables, 1, *see also* Random variable(s)
 - commensurate, 2
 - dummy variables, 186, 292, 327, 400
 - linear combinations of, 75–81
 - Variance matrix, *see* Covariance matrix
 - Variance-covariance matrix, *see* Covariance matrix
 - Variance:
 - generalized sample variance, 81
 - pooled variance, 134

- population variance (σ^2), 48
- sample variance (s^2), 48
- total sample variance, 82
- Varimax rotation, 460–462
- Vector(s):
 - $\mathbf{0}$ vector, 11
 - definition of, 8
 - distance:
 - Mahalanobis, 85
 - from origin to a point, 16
 - between two vectors, 85
 - geometry of, 9
 - j vector, 11
 - length of, 16
 - linear combinations of, 21
 - linear independence and dependence of, 23
 - normalized, 32
 - notation for vector, 8
 - observation vector, 64
 - orthogonal, 31, 54
 - perpendicular, 54
 - product of, 16
 - rows and columns of a matrix, 17
 - standardized, 95
 - subvectors, 71–75
 - sum of products, 16
 - sum of squares, 16
 - transpose of, 8
 - zero vector, 11
- Voting data, 563
- Weight gain data, 253
- Wheat data, 554
- Wilks' Λ test statistic:
 - definition of, 174–178
 - partial Λ statistic, 242
 - table of critical values, 613–636
- Wishart distribution, 100–101
- Words data, 167

WILEY SERIES IN PROBABILITY AND STATISTICS
 ESTABLISHED BY WALTER A. SHEWHART AND SAMUEL S. WILKS

Editors: *David J. Balding, Noel A. C. Cressie, Garrett M. Fitzmaurice, Harvey Goldstein, Iain M. Johnstone, Geert Molenberghs, David W. Scott, Adrian F. M. Smith, Ruey S. Tsay, Sanford Weisberg*
 Editors Emeriti: *Vic Barnett, J. Stuart Hunter, Joseph B. Kadane, Jozef L. Teugels*

The *Wiley Series in Probability and Statistics* is well established and authoritative. It covers many topics of current research interest in both pure and applied statistics and probability theory. Written by leading statisticians and institutions, the titles span both state-of-the-art developments in the field and classical methods.

Reflecting the wide range of current research in statistics, the series encompasses applied, methodological and theoretical statistics, ranging from applications and new techniques made possible by advances in computerized practice to rigorous treatment of theoretical approaches.

This series provides essential and invaluable reading for all statisticians, whether in academia, industry, government, or research.

- † ABRAHAM and LEDOLTER · Statistical Methods for Forecasting
 AGRESTI · Analysis of Ordinal Categorical Data, *Second Edition*
 AGRESTI · An Introduction to Categorical Data Analysis, *Second Edition*
 AGRESTI · Categorical Data Analysis, *Second Edition*
 ALTMAN, GILL, and McDONALD · Numerical Issues in Statistical Computing for the Social Scientist
 AMARATUNGA and CABRERA · Exploration and Analysis of DNA Microarray and Protein Array Data
 ANDĚL · Mathematics of Chance
 ANDERSON · An Introduction to Multivariate Statistical Analysis, *Third Edition*
 * ANDERSON · The Statistical Analysis of Time Series
 ANDERSON, AUQUIER, HAUCK, OAKES, VANDAELE, and WEISBERG · Statistical Methods for Comparative Studies
 ANDERSON and LOYNES · The Teaching of Practical Statistics
 ARMITAGE and DAVID (editors) · Advances in Biometry
 ARNOLD, BALAKRISHNAN, and NAGARAJA · Records
 * ARTHANARI and DODGE · Mathematical Programming in Statistics
 * BAILEY · The Elements of Stochastic Processes with Applications to the Natural Sciences
 BAJORSKI · Statistics for Imaging, Optics, and Photonics
 BALAKRISHNAN and KOUTRAS · Runs and Scans with Applications
 BALAKRISHNAN and NG · Precedence-Type Tests and Applications
 BARNETT · Comparative Statistical Inference, *Third Edition*
 BARNETT · Environmental Statistics
 BARNETT and LEWIS · Outliers in Statistical Data, *Third Edition*
 BARTHOLOMEW, KNOTT, and MOUSTAKI · Latent Variable Models and Factor Analysis: A Unified Approach, *Third Edition*
 BARTOSZYNSKI and NIEWIADOMSKA-BUGAJ · Probability and Statistical Inference, *Second Edition*
 BASILEVSKY · Statistical Factor Analysis and Related Methods: Theory and Applications
 BATES and WATTS · Nonlinear Regression Analysis and Its Applications
 BECHHOFER, SANTNER, and GOLDSMAN · Design and Analysis of Experiments for Statistical Selection, Screening, and Multiple Comparisons

*Now available in a lower priced paperback edition in the Wiley Classics Library.

†Now available in a lower priced paperback edition in the Wiley-Interscience Paperback Series.

- BEIRLANT, GOEGEBEUR, SEGERS, TEUGELS, and DE WAAL · Statistics of Extremes: Theory and Applications
- BELSLEY · Conditioning Diagnostics: Collinearity and Weak Data in Regression
- † BELSLEY, KUH, and WELSCH · Regression Diagnostics: Identifying Influential Data and Sources of Collinearity
- BENDAT and PIERSOL · Random Data: Analysis and Measurement Procedures, *Fourth Edition*
- BERNARDO and SMITH · Bayesian Theory
- BHAT and MILLER · Elements of Applied Stochastic Processes, *Third Edition*
- BHATTACHARYA and WAYMIRE · Stochastic Processes with Applications
- BIEMER, GROVES, LYBERG, MATHIOWETZ, and SUDMAN · Measurement Errors in Surveys
- BILLINGSLEY · Convergence of Probability Measures, *Second Edition*
- BILLINGSLEY · Probability and Measure, *Anniversary Edition*
- BIRKES and DODGE · Alternative Methods of Regression
- BISGAARD and KULAHCI · Time Series Analysis and Forecasting by Example
- BISWAS, DATTA, FINE, and SEGAL · Statistical Advances in the Biomedical Sciences: Clinical Trials, Epidemiology, Survival Analysis, and Bioinformatics
- BLISCHKE and MURTHY (editors) · Case Studies in Reliability and Maintenance
- BLISCHKE and MURTHY · Reliability: Modeling, Prediction, and Optimization
- BLOOMFIELD · Fourier Analysis of Time Series: An Introduction, *Second Edition*
- BOLLEN · Structural Equations with Latent Variables
- BOLLEN and CURRAN · Latent Curve Models: A Structural Equation Perspective
- BOROVKOV · Ergodicity and Stability of Stochastic Processes
- BOSQ and BLANKE · Inference and Prediction in Large Dimensions
- BOULEAU · Numerical Methods for Stochastic Processes
- * BOX and TIAO · Bayesian Inference in Statistical Analysis
- BOX · Improving Almost Anything, *Revised Edition*
- * BOX and DRAPER · Evolutionary Operation: A Statistical Method for Process Improvement
- BOX and DRAPER · Response Surfaces, Mixtures, and Ridge Analyses, *Second Edition*
- BOX, HUNTER, and HUNTER · Statistics for Experimenters: Design, Innovation, and Discovery, *Second Edition*
- BOX, JENKINS, and REINSEL · Time Series Analysis: Forecasting and Control, *Fourth Edition*
- BOX, LUCEÑO, and PANIAGUA-QUIÑONES · Statistical Control by Monitoring and Adjustment, *Second Edition*
- * BROWN and HOLLANDER · Statistics: A Biomedical Introduction
- CAIROLI and DALANG · Sequential Stochastic Optimization
- CASTILLO, HADI, BALAKRISHNAN, and SARABIA · Extreme Value and Related Models with Applications in Engineering and Science
- CHAN · Time Series: Applications to Finance with R and S-Plus®, *Second Edition*
- CHARALAMBIDES · Combinatorial Methods in Discrete Distributions
- CHATTERJEE and HADI · Regression Analysis by Example, *Fourth Edition*
- CHATTERJEE and HADI · Sensitivity Analysis in Linear Regression
- CHERNICK · Bootstrap Methods: A Guide for Practitioners and Researchers, *Second Edition*
- CHERNICK and FRIIS · Introductory Biostatistics for the Health Sciences
- CHILÈS and DELFINER · Geostatistics: Modeling Spatial Uncertainty, *Second Edition*
- CHOW and LIU · Design and Analysis of Clinical Trials: Concepts and Methodologies, *Second Edition*
- CLARKE · Linear Models: The Theory and Application of Analysis of Variance
- CLARKE and DISNEY · Probability and Random Processes: A First Course with Applications, *Second Edition*

*Now available in a lower priced paperback edition in the Wiley Classics Library.

†Now available in a lower priced paperback edition in the Wiley-Interscience Paperback Series.

- * COCHRAN and COX · Experimental Designs, *Second Edition*
- COLLINS and LANZA · Latent Class and Latent Transition Analysis: With Applications in the Social, Behavioral, and Health Sciences
- CONGDON · Applied Bayesian Modelling
- CONGDON · Bayesian Models for Categorical Data
- CONGDON · Bayesian Statistical Modelling, *Second Edition*
- CONOVER · Practical Nonparametric Statistics, *Third Edition*
- COOK · Regression Graphics
- COOK and WEISBERG · An Introduction to Regression Graphics
- COOK and WEISBERG · Applied Regression Including Computing and Graphics
- CORNELL · A Primer on Experiments with Mixtures
- CORNELL · Experiments with Mixtures, Designs, Models, and the Analysis of Mixture Data, *Third Edition*
- COX · A Handbook of Introductory Statistical Methods
- CRESSIE · Statistics for Spatial Data, *Revised Edition*
- CRESSIE and WIKLE · Statistics for Spatio-Temporal Data
- CSÖRGÖ and HORVÁTH · Limit Theorems in Change Point Analysis
- DAGPUNAR · Simulation and Monte Carlo: With Applications in Finance and MCMC
- DANIEL · Applications of Statistics to Industrial Experimentation
- DANIEL · Biostatistics: A Foundation for Analysis in the Health Sciences, *Eighth Edition*
- * DANIEL · Fitting Equations to Data: Computer Analysis of Multifactor Data, *Second Edition*
- DASU and JOHNSON · Exploratory Data Mining and Data Cleaning
- DAVID and NAGARAJA · Order Statistics, *Third Edition*
- * DEGROOT, FIENBERG, and KADANE · Statistics and the Law
- DEL CASTILLO · Statistical Process Adjustment for Quality Control
- DEMARIS · Regression with Social Data: Modeling Continuous and Limited Response Variables
- DEMIDENKO · Mixed Models: Theory and Applications
- DENISON, HOLMES, MALLICK and SMITH · Bayesian Methods for Nonlinear Classification and Regression
- DETTE and STUDDEN · The Theory of Canonical Moments with Applications in Statistics, Probability, and Analysis
- DEY and MUKERJEE · Fractional Factorial Plans
- DILLON and GOLDSTEIN · Multivariate Analysis: Methods and Applications
- * DODGE and ROMIG · Sampling Inspection Tables, *Second Edition*
- * DOOB · Stochastic Processes
- DOWDY, WEARDEN, and CHILKO · Statistics for Research, *Third Edition*
- DRAPER and SMITH · Applied Regression Analysis, *Third Edition*
- DRYDEN and MARDIA · Statistical Shape Analysis
- DUDEWICZ and MISHRA · Modern Mathematical Statistics
- DUNN and CLARK · Basic Statistics: A Primer for the Biomedical Sciences, *Fourth Edition*
- DUPUIS and ELLIS · A Weak Convergence Approach to the Theory of Large Deviations
- EDLER and KITSOS · Recent Advances in Quantitative Methods in Cancer and Human Health Risk Assessment
- * ELANDT-JOHNSON and JOHNSON · Survival Models and Data Analysis
- ENDERS · Applied Econometric Time Series, *Third Edition*
- † ETHIER and KURTZ · Markov Processes: Characterization and Convergence
- EVANS, HASTINGS, and PEACOCK · Statistical Distributions, *Third Edition*
- EVERITT, LANDAU, LEESE, and STAHL · Cluster Analysis, *Fifth Edition*
- FEDERER and KING · Variations on Split Plot and Split Block Experiment Designs

*Now available in a lower priced paperback edition in the Wiley Classics Library.

†Now available in a lower priced paperback edition in the Wiley-Interscience Paperback Series.

- FELLER · An Introduction to Probability Theory and Its Applications, Volume I,
Third Edition, Revised; Volume II, *Second Edition*
- FITZMAURICE, LAIRD, and WARE · Applied Longitudinal Analysis, *Second Edition*
- * FLEISS · The Design and Analysis of Clinical Experiments
- FLEISS · Statistical Methods for Rates and Proportions, *Third Edition*
- † FLEMING and HARRINGTON · Counting Processes and Survival Analysis
- FUJIKOSHI, ULYANOV, and SHIMIZU · Multivariate Statistics: High-Dimensional and Large-Sample Approximations
- FULLER · Introduction to Statistical Time Series, *Second Edition*
- † FULLER · Measurement Error Models
- GALLANT · Nonlinear Statistical Models
- GEISSEY · Modes of Parametric Statistical Inference
- GELMAN and MENG · Applied Bayesian Modeling and Causal Inference from Incomplete-Data Perspectives
- GEWEKE · Contemporary Bayesian Econometrics and Statistics
- GHOSH, MUKHOPADHYAY, and SEN · Sequential Estimation
- GIESBRECHT and GUMPERTZ · Planning, Construction, and Statistical Analysis of Comparative Experiments
- GIFI · Nonlinear Multivariate Analysis
- GIVENS and HOETING · Computational Statistics
- GLASSERMAN and YAO · Monotone Structure in Discrete-Event Systems
- GNANADESIKAN · Methods for Statistical Data Analysis of Multivariate Observations, *Second Edition*
- GOLDSTEIN · Multilevel Statistical Models, *Fourth Edition*
- GOLDSTEIN and LEWIS · Assessment: Problems, Development, and Statistical Issues
- GOLDSTEIN and WOOFF · Bayes Linear Statistics
- GREENWOOD and NIKULIN · A Guide to Chi-Squared Testing
- GROSS, SHORTLE, THOMPSON, and HARRIS · Fundamentals of Queueing Theory, *Fourth Edition*
- GROSS, SHORTLE, THOMPSON, and HARRIS · Solutions Manual to Accompany Fundamentals of Queueing Theory, *Fourth Edition*
- * HAHN and SHAPIRO · Statistical Models in Engineering
- HAHN and MEEKER · Statistical Intervals: A Guide for Practitioners
- HALD · A History of Probability and Statistics and their Applications Before 1750
- † HAMEL · Robust Statistics: The Approach Based on Influence Functions
- HARTUNG, KNAPP, and SINHA · Statistical Meta-Analysis with Applications
- HEIBERGER · Computation for the Analysis of Designed Experiments
- HEDAYAT and SINHA · Design and Inference in Finite Population Sampling
- HEDEKER and GIBBONS · Longitudinal Data Analysis
- HELLER · MACSYMA for Statisticians
- HERITIER, CANTONI, COPT, and VICTORIA-FESER · Robust Methods in Biostatistics
- HINKELMANN and KEMPTHORNE · Design and Analysis of Experiments, Volume 1: Introduction to Experimental Design, *Second Edition*
- HINKELMANN and KEMPTHORNE · Design and Analysis of Experiments, Volume 2: Advanced Experimental Design
- HINKELMANN (editor) · Design and Analysis of Experiments, Volume 3: Special Designs and Applications
- HOAGLIN, MOSTELLER, and TUKEY · Fundamentals of Exploratory Analysis of Variance
- * HOAGLIN, MOSTELLER, and TUKEY · Exploring Data Tables, Trends and Shapes
- * HOAGLIN, MOSTELLER, and TUKEY · Understanding Robust and Exploratory Data Analysis

*Now available in a lower priced paperback edition in the Wiley Classics Library.

†Now available in a lower priced paperback edition in the Wiley-Interscience Paperback Series.

- HOCHBERG and TAMHANE · Multiple Comparison Procedures
- HOCKING · Methods and Applications of Linear Models: Regression and the Analysis of Variance, *Second Edition*
- HOEL · Introduction to Mathematical Statistics, *Fifth Edition*
- HOGG and KLUGMAN · Loss Distributions
- HOLLANDER and WOLFE · Nonparametric Statistical Methods, *Second Edition*
- HOSMER and LEMESHOW · Applied Logistic Regression, *Second Edition*
- HOSMER, LEMESHOW, and MAY · Applied Survival Analysis: Regression Modeling of Time-to-Event Data, *Second Edition*
- HUBER · Data Analysis: What Can Be Learned From the Past 50 Years
- HUBER · Robust Statistics
- † HUBER and RONCHETTI · Robust Statistics, *Second Edition*
- HUBERTY · Applied Discriminant Analysis, *Second Edition*
- HUBERTY and OLEJNIK · Applied MANOVA and Discriminant Analysis, *Second Edition*
- HUITEMA · The Analysis of Covariance and Alternatives: Statistical Methods for Experiments, Quasi-Experiments, and Single-Case Studies, *Second Edition*
- HUNT and KENNEDY · Financial Derivatives in Theory and Practice, *Revised Edition*
- HURD and MIAMEE · Periodically Correlated Random Sequences: Spectral Theory and Practice
- HUSKOVA, BERAN, and DUPAC · Collected Works of Jaroslav Hajek— with Commentary
- HUZURBAZAR · Flowgraph Models for Multistate Time-to-Event Data
- JACKMAN · Bayesian Analysis for the Social Sciences
- † JACKSON · A User's Guide to Principle Components
- JOHN · Statistical Methods in Engineering and Quality Assurance
- JOHNSON · Multivariate Statistical Simulation
- JOHNSON and BALAKRISHNAN · Advances in the Theory and Practice of Statistics: A Volume in Honor of Samuel Kotz
- JOHNSON, KEMP, and KOTZ · Univariate Discrete Distributions, *Third Edition*
- JOHNSON and KOTZ (editors) · Leading Personalities in Statistical Sciences: From the Seventeenth Century to the Present
- JOHNSON, KOTZ, and BALAKRISHNAN · Continuous Univariate Distributions, Volume 1, *Second Edition*
- JOHNSON, KOTZ, and BALAKRISHNAN · Continuous Univariate Distributions, Volume 2, *Second Edition*
- JOHNSON, KOTZ, and BALAKRISHNAN · Discrete Multivariate Distributions
- JUDGE, GRIFFITHS, HILL, LÜTKEPOHL, and LEE · The Theory and Practice of Econometrics, *Second Edition*
- JUREK and MASON · Operator-Limit Distributions in Probability Theory
- KADANE · Bayesian Methods and Ethics in a Clinical Trial Design
- KADANE AND SCHUM · A Probabilistic Analysis of the Sacco and Vanzetti Evidence
- KALBFLEISCH and PRENTICE · The Statistical Analysis of Failure Time Data, *Second Edition*
- KARIYA and KURATA · Generalized Least Squares
- KASS and VOS · Geometrical Foundations of Asymptotic Inference
- † KAUFMAN and ROUSSEEUW · Finding Groups in Data: An Introduction to Cluster Analysis
- KEDEM and FOKIANOS · Regression Models for Time Series Analysis
- KENDALL, BARDEN, CARNE, and LE · Shape and Shape Theory
- KHURI · Advanced Calculus with Applications in Statistics, *Second Edition*
- KHURI, MATHEW, and SINHA · Statistical Tests for Mixed Linear Models
- * KISH · Statistical Design for Research

*Now available in a lower priced paperback edition in the Wiley Classics Library.

†Now available in a lower priced paperback edition in the Wiley-Interscience Paperback Series.

- KLEIBER and KOTZ · Statistical Size Distributions in Economics and Actuarial Sciences
KLEMELÄ · Smoothing of Multivariate Data: Density Estimation and Visualization
KLUGMAN, PANJER, and WILLMOT · Loss Models: From Data to Decisions,
Third Edition
KLUGMAN, PANJER, and WILLMOT · Solutions Manual to Accompany Loss Models:
From Data to Decisions, Third Edition
KOSKI and NOBLE · Bayesian Networks: An Introduction
KOTZ, BALAKRISHNAN, and JOHNSON · Continuous Multivariate Distributions,
Volume 1, *Second Edition*
KOTZ and JOHNSON (editors) · Encyclopedia of Statistical Sciences: Volumes 1 to 9
with Index
KOTZ and JOHNSON (editors) · Encyclopedia of Statistical Sciences: Supplement
Volume
KOTZ, READ, and BANKS (editors) · Encyclopedia of Statistical Sciences: Update
Volume 1
KOTZ, READ, and BANKS (editors) · Encyclopedia of Statistical Sciences: Update
Volume 2
KOWALSKI and TU · Modern Applied U-Statistics
KRISHNAMOORTHY and MATHEW · Statistical Tolerance Regions: Theory,
Applications, and Computation
KROESE, TAIMRE, and BOTEV · Handbook of Monte Carlo Methods
KROONENBERG · Applied Multiway Data Analysis
KULINSKAYA, MORGENTHALER, and STAUDTE · Meta Analysis: A Guide to
Calibrating and Combining Statistical Evidence
KULKARNI and HARMAN · An Elementary Introduction to Statistical Learning Theory
KUROWICKA and COOKE · Uncertainty Analysis with High Dimensional Dependence
Modelling
KVAM and VIDAKOVIC · Nonparametric Statistics with Applications to Science
and Engineering
LACHIN · Biostatistical Methods: The Assessment of Relative Risks, *Second Edition*
LAD · Operational Subjective Statistical Methods: A Mathematical, Philosophical, and
Historical Introduction
LAMPERTI · Probability: A Survey of the Mathematical Theory, *Second Edition*
LAWLESS · Statistical Models and Methods for Lifetime Data, *Second Edition*
LAWSON · Statistical Methods in Spatial Epidemiology, *Second Edition*
LE · Applied Categorical Data Analysis, *Second Edition*
LE · Applied Survival Analysis
LEE · Structural Equation Modeling: A Bayesian Approach
LEE and WANG · Statistical Methods for Survival Data Analysis, *Third Edition*
LEPAGE and BILLARD · Exploring the Limits of Bootstrap
LESSLER and KALSBECK · Nonsampling Errors in Surveys
LEYLAND and GOLDSTEIN (editors) · Multilevel Modelling of Health Statistics
LIAO · Statistical Group Comparison
LIN · Introductory Stochastic Analysis for Finance and Insurance
LITTLE and RUBIN · Statistical Analysis with Missing Data, *Second Edition*
LLOYD · The Statistical Analysis of Categorical Data
LOWEN and TEICH · Fractal-Based Point Processes
MAGNUS and NEUDECKER · Matrix Differential Calculus with Applications in
Statistics and Econometrics, *Revised Edition*
MALLER and ZHOU · Survival Analysis with Long Term Survivors
MARCHETTE · Random Graphs for Statistical Pattern Recognition
MARDIA and JUPP · Directional Statistics
MARKOVICH · Nonparametric Analysis of Univariate Heavy-Tailed Data: Research and
Practice

*Now available in a lower priced paperback edition in the Wiley Classics Library.

†Now available in a lower priced paperback edition in the Wiley-Interscience Paperback Series.

- MARONNA, MARTIN and YOHAI · Robust Statistics: Theory and Methods
 MASON, GUNST, and HESS · Statistical Design and Analysis of Experiments with Applications to Engineering and Science, *Second Edition*
 McCULLOCH, SEARLE, and NEUHAUS · Generalized, Linear, and Mixed Models, *Second Edition*
 McFADDEN · Management of Data in Clinical Trials, *Second Edition*
 * McLACHLAN · Discriminant Analysis and Statistical Pattern Recognition
 McLACHLAN, DO, and AMBROISE · Analyzing Microarray Gene Expression Data
 McLACHLAN and KRISHNAN · The EM Algorithm and Extensions, *Second Edition*
 McLACHLAN and PEEL · Finite Mixture Models
 McNEIL · Epidemiological Research Methods
 MEEKER and ESCOBAR · Statistical Methods for Reliability Data
 MEERSCHAERT and SCHEFFLER · Limit Distributions for Sums of Independent Random Vectors: Heavy Tails in Theory and Practice
 MENGERSEN, ROBERT, and TITTERINGTON · Mixtures: Estimation and Applications
 MICKEY, DUNN, and CLARK · Applied Statistics: Analysis of Variance and Regression, *Third Edition*
 * MILLER · Survival Analysis, *Second Edition*
 MONTGOMERY, JENNINGS, and KULAHCI · Introduction to Time Series Analysis and Forecasting
 MONTGOMERY, PECK, and VINING · Introduction to Linear Regression Analysis, *Fifth Edition*
 MORGENTHALER and TUKEY · Configural Polysampling: A Route to Practical Robustness
 MUIRHEAD · Aspects of Multivariate Statistical Theory
 MULLER and STOYAN · Comparison Methods for Stochastic Models and Risks
 MURTHY, XIE, and JIANG · Weibull Models
 MYERS, MONTGOMERY, and ANDERSON-COOK · Response Surface Methodology: Process and Product Optimization Using Designed Experiments, *Third Edition*
 MYERS, MONTGOMERY, VINING, and ROBINSON · Generalized Linear Models. With Applications in Engineering and the Sciences, *Second Edition*
 NATVIG · Multistate Systems Reliability Theory With Applications
 † NELSON · Accelerated Testing, Statistical Models, Test Plans, and Data Analyses
 † NELSON · Applied Life Data Analysis
 NEWMAN · Biostatistical Methods in Epidemiology
 NG, TAIN, and TANG · Dirichlet Theory: Theory, Methods and Applications
 OKABE, BOOTS, SUGIHARA, and CHIU · Spatial Tesselations: Concepts and Applications of Voronoi Diagrams, *Second Edition*
 OLIVER and SMITH · Influence Diagrams, Belief Nets and Decision Analysis
 PALTA · Quantitative Methods in Population Health: Extensions of Ordinary Regressions
 PANJER · Operational Risk: Modeling and Analytics
 PANKRATZ · Forecasting with Dynamic Regression Models
 PANKRATZ · Forecasting with Univariate Box-Jenkins Models: Concepts and Cases
 PARDOUX · Markov Processes and Applications: Algorithms, Networks, Genome and Finance
 PARMIGIANI and INOUE · Decision Theory: Principles and Approaches
 * PARZEN · Modern Probability Theory and Its Applications
 PEÑA, TIAO, and TSAY · A Course in Time Series Analysis
 PESARIN and SALMASO · Permutation Tests for Complex Data: Applications and Software
 PIANTADOSI · Clinical Trials: A Methodologic Perspective, *Second Edition*
 POURAHMADI · Foundations of Time Series Analysis and Prediction Theory

*Now available in a lower priced paperback edition in the Wiley Classics Library.

†Now available in a lower priced paperback edition in the Wiley-Interscience Paperback Series.

- POWELL · Approximate Dynamic Programming: Solving the Curses of Dimensionality, *Second Edition*
- POWELL and RYZHOV · Optimal Learning
- PRESS · Subjective and Objective Bayesian Statistics, *Second Edition*
- PRESS and TANUR · The Subjectivity of Scientists and the Bayesian Approach
- PURI, VILAPLANA, and WERTZ · New Perspectives in Theoretical and Applied Statistics
- † PUTERMAN · Markov Decision Processes: Discrete Stochastic Dynamic Programming
- QIU · Image Processing and Jump Regression Analysis
- * RAO · Linear Statistical Inference and Its Applications, *Second Edition*
- RAO · Statistical Inference for Fractional Diffusion Processes
- RAUSAND and HØYLAND · System Reliability Theory: Models, Statistical Methods, and Applications, *Second Edition*
- RAYNER, THAS, and BEST · Smooth Tests of Goodness of Fit: Using R, *Second Edition*
- RENCHER and SCHAALJE · Linear Models in Statistics, *Second Edition*
- RENCHER and CHRISTENSEN · Methods of Multivariate Analysis, *Third Edition*
- RENCHER · Multivariate Statistical Inference with Applications
- RIGDON and BASU · Statistical Methods for the Reliability of Repairable Systems
- * RIPLEY · Spatial Statistics
- * RIPLEY · Stochastic Simulation
- ROHATGI and SALEH · An Introduction to Probability and Statistics, *Second Edition*
- ROLSKI, SCHMIDL, SCHMIDT, and TEUGELS · Stochastic Processes for Insurance and Finance
- ROSENBERGER and LACHIN · Randomization in Clinical Trials: Theory and Practice
- ROSSI, ALLENBY, and McCULLOCH · Bayesian Statistics and Marketing
- † ROUSSEEUW and LEROY · Robust Regression and Outlier Detection
- ROYSTON and SAUERBREI · Multivariate Model Building: A Pragmatic Approach to Regression Analysis Based on Fractional Polynomials for Modeling Continuous Variables
- * RUBIN · Multiple Imputation for Nonresponse in Surveys
- RUBINSTEIN and KROESE · Simulation and the Monte Carlo Method, *Second Edition*
- RUBINSTEIN and MELAMED · Modern Simulation and Modeling
- RYAN · Modern Engineering Statistics
- RYAN · Modern Experimental Design
- RYAN · Modern Regression Methods, *Second Edition*
- RYAN · Statistical Methods for Quality Improvement, *Third Edition*
- SALEH · Theory of Preliminary Test and Stein-Type Estimation with Applications
- SALTELLI, CHAN, and SCOTT (editors) · Sensitivity Analysis
- SCHERER · Batch Effects and Noise in Microarray Experiments: Sources and Solutions
- * SCHEFFE · The Analysis of Variance
- SCHIMEK · Smoothing and Regression: Approaches, Computation, and Application
- SCHOTT · Matrix Analysis for Statistics, *Second Edition*
- SCHOUTENS · Levy Processes in Finance: Pricing Financial Derivatives
- SCOTT · Multivariate Density Estimation: Theory, Practice, and Visualization
- * SEARLE · Linear Models
- † SEARLE · Linear Models for Unbalanced Data
- † SEARLE · Matrix Algebra Useful for Statistics
- † SEARLE, CASELLA, and McCULLOCH · Variance Components
- SEARLE and WILLETT · Matrix Algebra for Applied Economics
- SEBER · A Matrix Handbook For Statisticians
- † SEBER · Multivariate Observations
- SEBER and LEE · Linear Regression Analysis, *Second Edition*
- † SEBER and WILD · Nonlinear Regression
- SENNOTT · Stochastic Dynamic Programming and the Control of Queueing Systems

*Now available in a lower priced paperback edition in the Wiley Classics Library.

†Now available in a lower priced paperback edition in the Wiley-Interscience Paperback Series.

- * SERFLING · Approximation Theorems of Mathematical Statistics
- SHAFER and VOVK · Probability and Finance: It's Only a Game!
- SHERMAN · Spatial Statistics and Spatio-Temporal Data: Covariance Functions and Directional Properties
- SILVAPULLE and SEN · Constrained Statistical Inference: Inequality, Order, and Shape Restrictions
- SINGPURWALLA · Reliability and Risk: A Bayesian Perspective
- SMALL and MCLEISH · Hilbert Space Methods in Probability and Statistical Inference
- SRIVASTAVA · Methods of Multivariate Statistics
- STAPLETON · Linear Statistical Models, *Second Edition*
- STAPLETON · Models for Probability and Statistical Inference: Theory and Applications
- STAUDTE and SHEATHER · Robust Estimation and Testing
- STOYAN · Counterexamples in Probability, *Second Edition*
- STOYAN, KENDALL, and MECKE · Stochastic Geometry and Its Applications, *Second Edition*
- STOYAN and STOYAN · Fractals, Random Shapes and Point Fields: Methods of Geometrical Statistics
- STREET and BURGESS · The Construction of Optimal Stated Choice Experiments: Theory and Methods
- STYAN · The Collected Papers of T. W. Anderson: 1943–1985
- SUTTON, ABRAMS, JONES, SHELDON, and SONG · Methods for Meta-Analysis in Medical Research
- TAKEZAWA · Introduction to Nonparametric Regression
- TAMHANE · Statistical Analysis of Designed Experiments: Theory and Applications
- TANAKA · Time Series Analysis: Nonstationary and Noninvertible Distribution Theory
- THOMPSON · Empirical Model Building: Data, Models, and Reality, *Second Edition*
- THOMPSON · Sampling, *Third Edition*
- THOMPSON · Simulation: A Modeler's Approach
- THOMPSON and SEBER · Adaptive Sampling
- THOMPSON, WILLIAMS, and FINDLAY · Models for Investors in Real World Markets
- TIERNEY · LISP-STAT: An Object-Oriented Environment for Statistical Computing and Dynamic Graphics
- TSAY · Analysis of Financial Time Series, *Third Edition*
- UPTON and FINGLETON · Spatial Data Analysis by Example, Volume II: Categorical and Directional Data
- † VAN BELLE · Statistical Rules of Thumb, *Second Edition*
- VAN BELLE, FISHER, HEAGERTY, and LUMLEY · Biostatistics: A Methodology for the Health Sciences, *Second Edition*
- VESTRUP · The Theory of Measures and Integration
- VIDAKOVIC · Statistical Modeling by Wavelets
- VIERTL · Statistical Methods for Fuzzy Data
- VINOD and REAGLE · Preparing for the Worst: Incorporating Downside Risk in Stock Market Investments
- WALLER and GOTWAY · Applied Spatial Statistics for Public Health Data
- WEISBERG · Applied Linear Regression, *Third Edition*
- WEISBERG · Bias and Causation: Models and Judgment for Valid Comparisons
- WELSH · Aspects of Statistical Inference
- WESTFALL and YOUNG · Resampling-Based Multiple Testing: Examples and Methods for *p*-Value Adjustment
- * WHITTAKER · Graphical Models in Applied Multivariate Statistics
- WINKER · Optimization Heuristics in Economics: Applications of Threshold Accepting
- WOODWORTH · Biostatistics: A Bayesian Introduction
- WOOLSON and CLARKE · Statistical Methods for the Analysis of Biomedical Data, *Second Edition*

*Now available in a lower priced paperback edition in the Wiley Classics Library.

†Now available in a lower priced paperback edition in the Wiley-Interscience Paperback Series.

- WU and HAMADA · Experiments: Planning, Analysis, and Parameter Design Optimization, *Second Edition*
- WU and ZHANG · Nonparametric Regression Methods for Longitudinal Data Analysis
- YIN · Clinical Trial Design: Bayesian and Frequentist Adaptive Methods
- YOUNG, VALERO-MORA, and FRIENDLY · Visual Statistics: Seeing Data with Dynamic Interactive Graphics
- ZACKS · Stage-Wise Adaptive Designs
- * ZELLNER · An Introduction to Bayesian Inference in Econometrics
- ZELTERMAN · Discrete Distributions—Applications in the Health Sciences
- ZHOU, OBUCHOWSKI, and McCLISH · Statistical Methods in Diagnostic Medicine, *Second Edition*

*Now available in a lower priced paperback edition in the Wiley Classics Library.

†Now available in a lower priced paperback edition in the Wiley–Interscience Paperback Series.