


APPROXIMATIONS TO THE LOOP SPACE OF FLAG MANIFOLDS



PROGRESS REPORT

SIMON STOLTZE

JANUARY 6, 2015

ADVISOR: MARCEL BÖKSTEDT

INSTITUT FOR MATEMATIK
AARHUS UNIVERSITET

Preface

The following document is part of the midway exam for Ph.D. students at Aarhus University. It describes the work I have done on my project so far, and ends with a small presentation of ideas for future work. Due to the limited number of pages, the report is meant more as an overview and will occasionally only sketch a proof or argument instead of giving all the details.

I would like to thank Thomas Schmidt for proofreading an earlier version of this report, and my advisor Marcel Bökstedt for his help and inspiration during the past two years.

Conventions and notation

What follows is a brief description of the notation used in the report. For two topological spaces X and Y , we will write $X \cong Y$ to denote that X and Y are homeomorphic and $X \simeq Y$ to denote that they are homotopy equivalent. Likewise, $f \simeq g$ means that the maps f and g are homotopic. For subsets, we will use $A \subset X$ when A is any subset of X , and we will generally not make a distinction between points of X and one-point subsets of X . The set difference of X and A will be written as $X - A$. With a distinguished basepoint x_0 in X , the set of based loops in X will be denoted $\Omega(X) = \{\gamma : S^1 \rightarrow X \mid \gamma(1) = x_0\}$, and this comes equipped with the compact-open topology. The one-point compactification of X is $X^+ = X \cup \{\infty\}$.

We will be working exclusively with topology in the complex numbers, so the disc D is the unit disc in the complex plane \mathbb{C} and D_p is the unit disc in \mathbb{C}^p . The non-zero elements in \mathbb{C} will be written as $\mathbb{C}^* = \mathbb{C} - 0$. GL_m will denote the group of linear automorphisms of \mathbb{C}^m and the standard basis of \mathbb{C}^m is e_1, \dots, e_m . Linear maps will be written as matrices with respect to the standard basis.

Since we are mainly concerned with algebraic topology, a space will occasionally be replaced with a homotopy equivalent one without mention. For example, T^n will denote both the torus $(\mathbb{C}^*)^n \subset \mathrm{GL}_n$ and $(S^1)^n \subset \mathbb{C}^n$ depending on context, and the punctured disc $D_p - 0$ will sometimes be identified with the sphere $S^{2p-1} \subset \mathbb{C}^p$. Likewise, we will not distinguish between the different models of the flag manifold of \mathbb{C}^m , given as $\mathrm{GL}_m/B_m \cong \mathrm{U}_m/T^m \cong \mathrm{SU}_m/T^{m-1}$ with $B_m \subset \mathrm{GL}_m$ the upper (or lower) triangular matrices in GL_m , $T^m \subset \mathrm{U}_m$ the diagonal matrices in the unitary group and $T^{m-1} \subset \mathrm{SU}_m$ the diagonal matrices in the special unitary group.

Homology and cohomology groups will always be singular groups with integer coefficients. If a theorem is mentioned by name without a reference, it can be found in [Hat02].

Simon Stoltze
Aarhus University
January, 2015

Contents

Preface	i
Contents	ii
Introduction	iii
1 The space $X_{m,n}$	1
1.1 Stabilization	2
1.2 Symmetries	3
1.3 The quotient space	4
2 Cohomology calculations for $X_{3,3}$	5
2.1 A filtration on $X_{m,n}$	5
2.2 The spectral sequence of $X_{3,3}$	7
2.3 Differentials on E_1	13
2.4 Turning the page	19
3 Loop spaces	21
3.1 The map $Y_{m,n} \rightarrow \Omega$	21
3.2 Homotopy equivalence	23
3.3 Future work	26
Bibliography	27

Introduction

This document gives a description of a family of spaces $X_{m,n} \subset \mathbb{C}^{mn}$, indexed by two natural numbers m and n , and a quotient of these spaces, $Y_{m,n} = X_{m,n}/T^n$. The spaces were originally defined as the zero-set of a section on a vector bundle of a Grassmann manifold, and were studied to understand the properties of families of Calabi-Yau manifolds. However, the spaces turned out to have other properties that made them interesting in their own right. The most important property is that they can be used to form a space that is homotopy equivalent to a space that appears in physics, namely the loop space $\Omega(\mathrm{SU}_m/T^{m-1})$. This hints at a way of extracting information about the loop space by working with the simpler spaces $Y_{m,n}$.

The spaces have been studied by Marcel Bökstedt, and some of his results are gathered in [Bök13]. In particular it includes the following theorem that covers the special case where $m = 2$:

Theorem 1. *The homology of $Y_{2,n}$ is*

$$H_q(Y_{2,n}) = \begin{cases} \mathbb{Z} & 0 \leq q \leq n \\ 0 & \text{otherwise} \end{cases}$$

There is a stabilization map $s : Y_{2,n} \rightarrow Y_{2,n+1}$ that induces an isomorphism

$$s_* : H_q(Y_{2,n}) \rightarrow H_q(Y_{2,n+1})$$

for $q \leq n$. There is also a map $f : Y_{2,n} \rightarrow \Omega(\mathrm{SU}_2/T^1)$ that induces an isomorphism in homology in degree $q \leq n$.

The aim of the following is to try and extend these results to the general case of $Y_{m,n}$, with $m \geq 2$.

This progress report is divided into three main parts. The first chapter gives an introduction to the spaces we are interested in and lists some of the properties that will be used in the other chapters. The second chapter is mainly concerned with an explicit calculation of the cohomology of the spaces $X_{3,3}$ and $Y_{3,3}$. This is done by applying the spectral sequence coming from a filtration of $X_{3,3}$, computing the various relative cohomology groups needed, and finding the differentials of the sequence. The third and final chapter concerns the relation to the loop space described above. Here it is shown that by taking a limit of the spaces $Y_{m,n}$, we get a space homotopy equivalent to the loop space. The chapter concludes with a brief overview of plans for future studies.

Chapter 1

The space $X_{m,n}$

This chapter will introduce the spaces to be studied, along with a brief listing of various useful structure.

Definition 2. For natural numbers m and n , the space $X_{m,n} \subset \mathbb{C}^{mn}$ is defined as

$$X_{m,n} = \left\{ (a_1, \dots, a_n) \in (\mathbb{C}^m)^n \left| \begin{array}{l} \text{Any } m \text{ subsequent vectors in} \\ (e_1, \dots, e_m, a_1, \dots, a_n, e_1, \dots, e_m) \\ \text{are linearly independent.} \end{array} \right. \right\}$$

For any two elements $X = [x_1, \dots, x_m]$ and $Y = [y_1, \dots, y_m]$ in GL_m , define the space $X_{m,n}(X, Y)$ as

$$X_{m,n}(X, Y) = \left\{ (a_1, \dots, a_n) \in (\mathbb{C}^m)^n \left| \begin{array}{l} \text{Any } m \text{ subsequent vectors in} \\ (x_1, \dots, x_m, a_1, \dots, a_n, y_1, \dots, y_m) \\ \text{are linearly independent.} \end{array} \right. \right\}$$

The special case $X = \text{Id}$ will be denoted $X_{m,n}(Y)$. Elements $(a_1, \dots, a_n) \in X_{m,n}$ will be identified with the $m \times n$ matrix $A = [a_1, \dots, a_n]$ without mention.

Since the space $X_{m,n}(X, Y)$ only depends on linear independence of subsequent vectors, it can be described by giving the two flags in \mathbb{C}^m defined from the columns of the matrices $X = [x_1, \dots, x_m]$ and $Y = [y_1, \dots, y_m]$:

$$\begin{aligned} \text{Fl}_R(X) &= \left(\text{span}(x_m) \subset \text{span}(x_{m-1}, x_m) \subset \dots \subset \mathbb{C}^m \right) \\ \text{Fl}_L(Y) &= \left(\text{span}(y_1) \subset \text{span}(y_1, y_2) \subset \dots \subset \mathbb{C}^m \right) \end{aligned}$$

Multiplying with X^{-1} on each vector in $X_{m,n}(X, Y)$ defines a homeomorphism

$$X_{m,n}(X, Y) \cong X_{m,n}(\text{Id}, X^{-1}Y) = X_{m,n}(X^{-1}Y)$$

Hence we only need to consider these spaces.

Note that we can also define the space $X_{m,n}$ as the preimage of $(\mathbb{C}^*)^{m+n-1}$ under the map $\text{Det} : (\mathbb{C}^m)^n \rightarrow \mathbb{C}^{m+n-1}$, given by

$$\begin{aligned} \text{Det}(a_1, \dots, a_n) &= \left(\det(e_2, \dots, e_m, a_1), \right. \\ &\quad \det(e_3, \dots, e_m, a_1, a_2), \\ &\quad \dots, \\ &\quad \left. \det(a_n, e_1, \dots, e_{m-1}) \right) \end{aligned}$$

where \det is the determinant map. There is a similar description of $X_{m,n}(Y)$ as the preimage under a map Det_Y . Since the map Det_Y is continuous, this shows that the space $X_{m,n}(Y)$ is open in \mathbb{C}^{mn} .

1.1 Stabilization

We would like to relate the spaces $X_{m,n}$ and $X_{m,n+1}$. To do this, consider a lower-triangular $m \times m$ invertible matrix L . This preserves the right flag $\text{Fl}_R(\text{Id})$, so it defines a homeomorphism $\tilde{L} : X_{m,n}(Y) \rightarrow X_{m,n}(LY)$ by

$$\tilde{L}(a_1, \dots, a_n) = (La_1, \dots, La_n)$$

By using the Bruhat decomposition of the general linear group, any matrix Y can be written uniquely as a product $L\sigma U$, where L is an invertible lower triangular matrix, U is an invertible upper triangular matrix, and σ is a permutation matrix, see [BB05, Example 1.2.11] or [Hil82, Proposition 4.5]. Since the flag $\text{Fl}_L(\sigma U)$ is preserved by right multiplication with upper triangular matrices, this only depends on the permutation σ and not on U . All of this together shows the following lemma:

Lemma 3. *For $Y \in \text{GL}_m$ with $Y = L\sigma U$, the space $X_{m,n}(Y)$ is homeomorphic to $X_{m,n}(\sigma)$.*

Using this lemma, we only need to understand the spaces $X_{m,n}(\sigma)$ to understand all of the spaces $X_{m,n}(Y)$. More importantly, it turns out that the space $X_{m,n+1}(\sigma)$ is given as a union of spaces homeomorphic to a torus times $X_{m,n}(\tau)$, for various permutations τ . This is summarized in the following lemma.

Lemma 4. *Any choice of indices $I = (i_1 < \dots < i_k) \subset \{1, \dots, m\}$ gives a subspace of $X_{m,n+1}(\sigma)$:*

$$X_{m,n+1}^I(\sigma) = \left\{ (a_1, \dots, a_{n+1}) \in X_{m,n+1}(\sigma) \mid \begin{array}{l} (a_{n+1})_{i_j} \neq 0 \ \forall i_j \in I, \\ (a_{n+1})_j = 0 \ \forall j \notin I \end{array} \right\}$$

This space is empty if I does not contain the number $\sigma(m)$, so we will assume that $\sigma(m) \in I$ but otherwise leave it out of the notation.

These subspaces cover $X_{m,n+1}(\sigma)$. There is a map φ , defined for an indexing set I and a permutation σ , which gives a new permutation such that

$$X_{m,n+1}^I(\sigma) \cong X_{m,n}(\varphi(I, \sigma)) \times (\mathbb{C}^*)^{|I|}$$

If we define the permutation $\hat{\sigma} = \sigma \cdot (m \ m-1 \ \dots \ 1)$ and the indexing set is $I = (i_1 < \dots < i_k)$, then φ is given by

$$\varphi(I, \sigma) = \left(\prod_{j \in J} (i_j \ \sigma(m)) \right) \hat{\sigma}$$

where the product is over the set

$$J = \{j \in \{1, \dots, k\} \mid i_j < \sigma(m), \ \hat{\sigma}^{-1}(i_j) > \hat{\sigma}^{-1}(i_r) \ \forall r < j\}$$

The proof of this lemma relies on taking $X_{m,n+1}^I(\sigma)$ and multiplying with a lower triangular matrix to reduce it to something in $X_{m,n}(\tau)$. By considering the matrix used, the formula for $\tau = \varphi(I, \sigma)$ given in the lemma can be found. In practice, it is often easier to do the reduction by hand rather than using the lemma, but the formula can be useful. For example, it is clear that $\varphi(I, \sigma)$ is given as a product of transpositions and $\varphi((\sigma(m)), \sigma)$. By considering exactly which transpositions, we can show that for all I we have

$$\varphi(I, \sigma) \leq \varphi((\sigma(m)), \sigma)$$

in the Bruhat ordering on the symmetric group. The Bruhat order is defined in [BB05]. It also follows that for all σ ,

$$\varphi((1 < 2 < \dots < m), \sigma) = \text{Id}$$

With this we can identify $X_{m,n}$ with the subspace of $X_{m,n+1}$ where the last column has a 1 on each entry:

$$X_{m,n} \cong \{(a_1, \dots, a_{n+1}) \in X_{m,n+1} \mid a_{n+1} = (1, \dots, 1)\}$$

The identification is given by the map $s : X_{m,n} \rightarrow X_{m,n+1}$,

$$s(a_1, \dots, a_n) = (La_1, \dots, La_n, Le_1)$$

where L is the lower triangular matrix with all entries on or below the diagonal equal to one:

$$L = \begin{pmatrix} 1 & 0 & \dots & 0 \\ \vdots & \ddots & \ddots & \vdots \\ \vdots & & \ddots & 0 \\ 1 & \dots & \dots & 1 \end{pmatrix}$$

These maps give a directed system of spaces,

$$X_{m,1} \xrightarrow{s} X_{m,2} \xrightarrow{s} \dots \xrightarrow{s} X_{m,n} \xrightarrow{s} X_{m,n+1} \xrightarrow{s} \dots$$

and we can take the direct limit of this system,

$$X_{m,\infty} = \varinjlim_n X_{m,n}$$

The direct limit is the disjoint union of all the spaces $X_{m,n}$, with the identification $A \sim B$ if there are j, k such that $s^k(A) = s^j(B)$. The topology is the finest such that the maps $X_{m,n} \rightarrow X_{m,\infty}$ are all continuous.

1.2 Symmetries

There are various symmetries and group actions worth keeping in mind when working with these spaces. Some of these will be described here.

Since we are interested in linear independence of vectors in \mathbb{C}^m , the group of complex units \mathbb{C}^* acts on $X_{m,n}(\sigma)$ in various ways. For any index i , we can define a group action l_i by scaling the i^{th} vector:

$$\begin{aligned} l_i : \mathbb{C}^* \times X_{m,n}(\sigma) &\rightarrow X_{m,n}(\sigma) \\ (\lambda, (a_1, \dots, a_n)) &\mapsto (a_1, \dots, a_{i-1}, \lambda a_i, a_{i+1}, \dots, a_n) \end{aligned}$$

These actions commute with each other, so we get an action l of $(\mathbb{C}^*)^n$ on $X_{m,n}(\sigma)$, given by

$$l((\lambda_1, \dots, \lambda_n), (a_1, \dots, a_n)) = (\lambda_1 a_1, \dots, \lambda_n a_n)$$

Likewise, we can also multiply the rows of $A \in X_{m,n}(\sigma)$ by non-zero complex numbers. This defines an action l' of $(\mathbb{C}^*)^m$ on $X_{m,n}(\sigma)$. The two actions together satisfy the relation

$$l((\lambda, \dots, \lambda), l'((\lambda^{-1}, \dots, \lambda^{-1}), A)) = A$$

since multiplying all rows by λ^{-1} while multiplying all columns by λ is the same as doing nothing.

For the special case of $X_{m,n} = X_{m,n}(\text{Id})$, there is an additional symmetry that can be useful to know. By thinking of $A \in X_{m,n}$ as a matrix, it is possible to transpose it. Checking the various determinants used in the definition of $X_{m,n}$ show that they do not change after transposition. Hence transposing defines a homeomorphism $T : X_{m,n} \rightarrow X_{n,m}$, allowing us to switch the order of m and n .

1.3 The quotient space

While the above definition of $X_{m,n}$ gives an open subset of \mathbb{C}^{mn} , it has some issues that makes the space hard to work with. As we shall see shortly, the fundamental group and the cohomology ring of $X_{m,n}$ both become larger as n grows. This complicates things slightly, but it can be fixed by taking the quotient of a group action.

Definition 5. The space $Y_{m,n}$ is the quotient space

$$Y_{m,n} = X_{m,n}/T^n$$

where T^n acts on $X_{m,n}$ by scaling the columns, as defined above:

$$(\lambda_1, \dots, \lambda_n) \cdot (a_1, \dots, a_n) = (\lambda_1 a_1, \dots, \lambda_n a_n)$$

These spaces retain some of the structure described above. For example, the stabilization map s is linear in A , which means it respects the group action.

$$s((\lambda_1, \dots, \lambda_n) \cdot A) = (\lambda_1, \dots, \lambda_n, 1) \cdot s(A)$$

Hence s descends to a stabilization map $s : Y_{m,n} \rightarrow Y_{m,n+1}$ and there is a direct limit,

$$Y_{m,\infty} = \varinjlim_n Y_{m,n}$$

which can be thought of as $X_{m,\infty}/T^\infty$.

The space $Y_{m,n}$ is homeomorphic to the subspace of $X_{m,n}$ consisting of elements with the last n coordinates of the determinant map Det equal to 1:

$$Y_{m,n} \cong \{A \in X_{m,n} \mid \text{Det}(A) \in (\mathbb{C}^*)^{m-1} \times \{1\}^n\}$$

The group action gives an identification,

$$X_{m,n} \cong Y_{m,n} \times (\mathbb{C}^*)^n$$

This shows the claim about the fundamental group of $X_{m,n}$, but it can also be used to get information about $Y_{m,n}$ from $X_{m,n}$ or vice versa. For example, it allows us to calculate the cohomology of one when we know the cohomology of the other, by using the Künneth formula to add or remove the cohomology of $(\mathbb{C}^*)^n$.

Chapter 2

Cohomology calculations for $X_{3,3}$

In this section we will calculate the cohomology of the space $X_{3,3}$, which is the set:

$$X_{3,3} = \left\{ \begin{pmatrix} a & b & g \\ c & d & h \\ e & f & i \end{pmatrix} \in \mathbb{C}^9 \mid \begin{array}{l} a \neq 0, ad - bc \neq 0, \\ adi + beh + cfg - deg - bci - afh \neq 0, \\ di - fh \neq 0, i \neq 0 \end{array} \right\}$$

The way we will calculate this is by using the spectral sequence for a filtration of $X_{3,3}$. Some of the calculations have been done in Sage and the relevant source code can be found at <http://home.math.au.dk/sstoltze/del-a.html>.

2.1 A filtration on $X_{m,n}$

Before we start the calculation, we need some general setup. To work with the space $X_{m,n}$ and compute the cohomology, it will be beneficial to build it up from simpler subspaces. This is done by considering a filtration.

For $0 \leq p \leq m-1$, let F_p be the subset

$$F_p = \{(a_1, \dots, a_n) \in X_{m,n} \mid a_n \text{ has at most } p \text{ zeroes.}\}$$

F_p is open in $X_{m,n}$ since we stay inside the subspace when we vary the coordinates of a point $A \in F_p$, as long as we do not introduce new zeroes in the last column. Together, the subspaces form an increasing sequence:

$$F_0 \subset F_1 \subset \dots \subset F_{m-1} = X_{m,n}$$

The general method of computing the homology and cohomology of $X_{m,n}$ is by using the spectral sequence of this filtration, as described in [Hat04] and [McC01]. Here we will focus on cohomology, but similar arguments could of course be applied to homology.

The spectral sequence has E_1 -page given by

$$E_1^{p,q} = H^{p+q}(F_p, F_{p-1})$$

with differentials given by the boundary map in the long exact sequence of the triple (F_{p+1}, F_p, F_{p-1}) :

$$d_1 : H^{p+q}(F_p, F_{p-1}) = E_1^{p,q} \rightarrow E_1^{p+1,q} = H^{p+q+1}(F_{p+1}, F_p)$$

To use the spectral sequence, we need to compute the cohomology of the pair (F_p, F_{p-1}) . To calculate this we will use excision, so we would like to find an open subset of F_p that contains the complement of F_{p-1} . For any choice of exactly p different indices $\{v_1, \dots, v_p\} \subset \{1, \dots, m-1\}$, define $X_{\{v_1, \dots, v_p\}}$ to be the points $(a_1, \dots, a_n) \in F_p$ which has $(a_n)_{v_i} = 0$ for all i . Since $X_{m,n}$ is open, we can get an open set in F_p by inserting a small disc on these zero-entries. More precisely, the set we get is

$$X_{\{v_1, \dots, v_p\}} \times D_p \cong \left\{ A \in F_p \left| \begin{array}{l} |((a_n)_{v_1}, \dots, (a_n)_{v_p})| < \min_{i \notin \{v_1, \dots, v_p\}} |(a_n)_i|, \\ (a_1, \dots, a_{n-1}, a_n - t \sum_i (a_n)_{v_i} e_{v_i}) \in F_p \forall t \in [0, 1] \end{array} \right. \right\}$$

This is an open set in F_p , with $X_\emptyset = F_0$, and it contains all points $(a_1, \dots, a_n) \in F_p$ where the entries of a_n indicated by $\{v_1, \dots, v_p\}$ are zero. For a different choice of indices, the condition that the radius of the disc is smaller than the norm of the non-zero entries in a_n ensures disjointness, so the intersection is empty:

$$(X_{\{v_1, \dots, v_p\}} \times D_p) \cap (X_{\{u_1, \dots, u_p\}} \times D_p) = \emptyset \quad \text{if } \{v_1, \dots, v_p\} \neq \{u_1, \dots, u_p\}$$

The intersection of $X_{m,n} \times D_p$ and F_{p-1} corresponds to removing 0 from the disc, since one of the entries inserted on the last column has to be non-zero.

$$(X_{\{v_1, \dots, v_p\}} \times D_p) \cap F_{p-1} = X_{\{v_1, \dots, v_p\}} \times (D_p - 0)$$

This means that the disjoint union of all of these form an open set in F_p with

$$F_p = F_{p-1} \cup \left(\coprod_{\{v_1, \dots, v_p\}} X_{\{v_1, \dots, v_p\}} \times D_p \right)$$

and by excision, the inclusion of pairs

$$\coprod_{\{v_1, \dots, v_p\}} X_{\{v_1, \dots, v_p\}} \times (D_p, D_p - 0) \hookrightarrow (F_p, F_{p-1})$$

induces an isomorphism of cohomology groups,

$$\begin{aligned} E_1^{p,q} &= H^{p+q}(F_p, F_{p-1}) \\ &\cong H^{p+q} \left(\coprod_{\{v_1, \dots, v_p\}} X_{\{v_1, \dots, v_p\}} \times (D_p, D_p - 0) \right) \\ &\cong \bigoplus_{\{v_1, \dots, v_p\}} H^{p+q}(X_{\{v_1, \dots, v_p\}} \times (D_p, D_p - 0)) \\ &\cong \bigoplus_{\{v_1, \dots, v_p\}} H^{q-p}(X_{\{v_1, \dots, v_p\}}) \otimes H^{2p}(D_p, D_p - 0) \end{aligned}$$

The last isomorphism is given by the Künneth formula, and the fact that the pair $(D_p, D_p - 0)$ only has non-zero cohomology in degree $2p$:

$$H^q(D_p, D_p - 0) \cong \tilde{H}^{q-1}(D_p - 0) = \tilde{H}^{q-1}(S^{2p-1}) = \begin{cases} \mathbb{Z} & q = 2p \\ 0 & \text{otherwise} \end{cases}$$

2.2 The spectral sequence of $X_{3,3}$

We can apply the above filtration to get a spectral sequence of $X = X_{3,3}$. The groups we need to compute for the first page are:

$$E_1^{p,q} \cong \begin{cases} H^q(X_\emptyset) & p = 0 \\ H^{q+1}(X_{\{1\}} \times (D, D-0)) \oplus H^{q+1}(X_{\{2\}} \times (D, D-0)) & p = 1 \\ H^{q+2}(X_{\{1,2\}} \times (D_2, D_2-0)) & p = 2 \\ 0 & \text{otherwise} \end{cases}$$

The four spaces that appear will be handled separately.

The cohomology of $X_{\{1,2\}}$

Consider the space $X_{\{1,2\}}$. This consists of the elements of $X_{3,3}$ that have two zeroes in the last column:

$$X_{\{1,2\}} = \left\{ \begin{pmatrix} a & b & 0 \\ c & d & 0 \\ e & f & i \end{pmatrix} \mid \begin{array}{l} a \neq 0, ad - bc \neq 0 \\ adi - bci \neq 0 \\ di \neq 0, i \neq 0 \end{array} \right\}$$

The space is homeomorphic to $(\mathbb{C}^*)^3 \times \mathbb{C}^2 \times \{(b, c) \in \mathbb{C}^2 \mid bc \neq 1\}$, with the homeomorphism given by

$$\begin{aligned} X_{\{1,2\}} &\rightarrow (\mathbb{C}^*)^3 \times \mathbb{C}^2 \times \{(b, c) \in \mathbb{C}^2 \mid bc \neq 1\} \\ \begin{pmatrix} a & b & 0 \\ c & d & 0 \\ e & f & i \end{pmatrix} &\mapsto \left(a, d, i, \frac{e}{i}, \frac{f}{i}, \frac{b}{a}, \frac{c}{d} \right) \end{aligned}$$

The space $\{(b, c) \mid bc \neq 1\}$ will be denoted Y . To get the cohomology of this space, consider it as a subspace of the one-point compactification $S^4 = \mathbb{C}^2 \cup \{\infty\}$. Then the complement is:

$$S^4 - Y = \{(b, c) \in \mathbb{C}^2 \mid bc = 1\} \cup \{\infty\}$$

This can be identified with the wedge sum $S^1 \vee S^2$ by considering the subspace

$$\mathbb{C}^* \cong \left\{ \left(b, \frac{1}{b} \right) \mid b \in \mathbb{C}^* \right\} = (S^4 - Y) - \{\infty\}$$

The point at infinity is glued to this space when either b or c is large, which for our copy of \mathbb{C}^* corresponds to b large or b near 0. Hence what we get is $S^2 = \mathbb{C} \cup \{\infty\}$ with the identification $0 \sim \infty$. This is homotopy equivalent to $S^1 \vee S^2$.

Now we can apply Alexander duality to this space to compute

$$\tilde{H}_q(Y) \cong \tilde{H}^{3-q}(S^4 - Y) = \begin{cases} \mathbb{Z} & q \in \{1, 2\} \\ 0 & \text{otherwise} \end{cases}$$

and hence

$$H^q(Y) \cong \begin{cases} \mathbb{Z} & q \in \{0, 1, 2\} \\ 0 & \text{otherwise} \end{cases}$$

From all of this, we can calculate the cohomology of $X_{\{1,2\}}$

$$H^q(X_{\{1,2\}}) = \begin{cases} \mathbb{Z} & q = 0 \\ \mathbb{Z}^4 & q = 1 \\ \mathbb{Z}^7 & q = 2 \\ \mathbb{Z}^7 & q = 3 \\ \mathbb{Z}^4 & q = 4 \\ \mathbb{Z} & q = 5 \\ 0 & \text{otherwise} \end{cases}$$

Since we know the product structure from the Künneth formula, we can name the generators and get the ring structure,

$$H^*(X_{\{1,2\}}) = \bigwedge^* [a_{00}, d_{00}, i_{00}] \otimes H^*(Y)$$

The two generators of Y will be denoted $y_{00,1}$ for the degree one generator and $y_{00,2}$ for the degree two generator. All other generators have degree one. The ring structure is the usual one on a tensor product, with the added relation $y_{00,1} \cup y_{00,2} = 0$. The element $y_{00,1} \in H^1(Y)$ is given by the map

$$Y \ni (b, c) \mapsto 1 - bc \in \mathbb{C}^*$$

in cohomology, since there is another map

$$\mathbb{C}^* \ni \lambda \mapsto (1, 1 - \lambda) \in Y$$

and the composition is the identity on \mathbb{C}^* . Applying H^1 shows that the two maps give isomorphisms in degree 1.

The cohomology of $X_{\{1\}}$

In $X_{\{1\}}$ we have the first entry of the last column equal to zero, which is the space

$$X_{\{1\}} = \left\{ \begin{pmatrix} a & b & 0 \\ c & d & h \\ e & f & i \end{pmatrix} \mid \begin{array}{l} a \neq 0, ad - bc \neq 0 \\ adi + beh - bci - afh \neq 0 \\ di - fh \neq 0, h \neq 0, i \neq 0 \end{array} \right\}$$

This space is homeomorphic to $(\mathbb{C}^*)^5 \times \mathbb{C} \times Y$:

$$\begin{pmatrix} a & b & 0 \\ c & d & h \\ e & f & i \end{pmatrix} \mapsto \left(a, i, h, \frac{d}{h} - \frac{f}{i}, \frac{b}{h} - \frac{bc}{ah}, \frac{e}{i}, \frac{bhi}{adi - afh}, \frac{c}{h} - \frac{e}{i} \right)$$

Since the cohomology of Y is already known, applying the Künneth formula gives

$$H^q(X_{\{1\}}) = \begin{cases} \mathbb{Z} & q = 0 \\ \mathbb{Z}^6 & q = 1 \\ \mathbb{Z}^{16} & q = 2 \\ \mathbb{Z}^{25} & q = 3 \\ \mathbb{Z}^{25} & q = 4 \\ \mathbb{Z}^{16} & q = 5 \\ \mathbb{Z}^6 & q = 6 \\ \mathbb{Z} & q = 7 \\ 0 & \text{otherwise} \end{cases}$$

As before, the ring structure is given by

$$H^*(X_{\{1\}}) = \bigwedge^* [a_{01}, d_{01}, f_{01}, h_{01}, i_{01}] \otimes H^*(Y)$$

The cohomology of $X_{\{2\}}$

The space is

$$X_{\{2\}} = \left\{ \begin{pmatrix} a & b & g \\ c & d & 0 \\ e & f & i \end{pmatrix} \mid \begin{array}{l} a \neq 0, ad - bc \neq 0 \\ adi + cfg - bci - deg \neq 0 \\ di \neq 0, g \neq 0, i \neq 0 \end{array} \right\}$$

which can also be identified with $(\mathbb{C}^*)^5 \times \mathbb{C} \times Y$ using the homeomorphism

$$\begin{pmatrix} a & b & g \\ c & d & 0 \\ e & f & i \end{pmatrix} \mapsto \left(\frac{a}{g}, d, i, g, 1 + \frac{cfg}{adi} - \frac{eg}{ai} - \frac{bc}{ad}, \frac{f}{i}, \frac{b}{a}, \frac{c}{d} \right)$$

Since we already know the cohomology of this space, we know

$$H^q(X_{\{2\}}) = \left(\bigwedge^* [a_{10}, d_{10}, f_{10}, g_{10}, i_{10}] \otimes H^*(Y) \right)^q = \begin{cases} \mathbb{Z} & q = 0 \\ \mathbb{Z}^6 & q = 1 \\ \mathbb{Z}^{16} & q = 2 \\ \mathbb{Z}^{25} & q = 3 \\ \mathbb{Z}^{25} & q = 4 \\ \mathbb{Z}^{16} & q = 5 \\ \mathbb{Z}^6 & q = 6 \\ \mathbb{Z} & q = 7 \\ 0 & \text{otherwise} \end{cases}$$

The cohomology of X_\emptyset

This space is slightly more complicated than the others. The space consists of the elements in $X_{3,3}$ where all entries of the last vector are non-zero.

$$X_\emptyset = \left\{ \begin{pmatrix} a & b & g \\ c & d & h \\ e & f & i \end{pmatrix} \mid \begin{array}{l} a \neq 0, ad - bc \neq 0, di - fh \neq 0, \\ adi + beh + cfg - deg - bci - afh \neq 0, \\ g \neq 0, h \neq 0, i \neq 0 \end{array} \right\}$$

The space is homeomorphic to $(\mathbb{C}^*)^7 \times \{(x, y, z) \in \mathbb{C}^3 \mid y - z - xyz = 1\}$, with the homeomorphism given by

$$\begin{pmatrix} a & b & g \\ c & d & h \\ e & f & i \end{pmatrix} \mapsto \left(g, h, i, \frac{a}{g}, \frac{f}{i} - \frac{d}{h}, \right.$$

$$D = \frac{ad - bc}{ah},$$

$$C = \frac{c}{h} - \frac{a}{g} + \frac{cdi - deh}{fh^2 - dhi} + \frac{beh - bci}{fgh - dgi},$$

$$\left. \frac{bC}{aD}, \frac{c}{hC} - \frac{a}{gC}, \frac{D}{C} \frac{eh - ci}{fh - di} \right)$$

To get the cohomology, we need to compute the cohomology of the space:

$$Z = \{(x, y, z) \in \mathbb{C}^3 \mid y - z - xyz = 1\}$$

This is the same as the cohomology of S^2 , but this is hard to see directly. Instead consider the space

$$\tilde{Z} = \{(b, c, d, e) \in \mathbb{C}^4 \mid d \neq bc, c \neq de\}$$

This space is homeomorphic to $Z \times (\mathbb{C}^*)^2$:

$$(b, c, d, e) \mapsto \left(\frac{b(c-de)}{d-bc}, \frac{c}{c-de}, \frac{e(d-bc)}{c-de}, d-bc, c-de \right)$$

We will use Alexander duality to compute the cohomology as we did for Y . Consider the complement in S^8 ,

$$S^8 - \tilde{Z} = \{(b, c, d, e) \mid d = bc\} \cup \{(b, c, d, e) \mid c = de\} \cup \{\infty\}$$

and denote the two sets as W_1 and W_2 . Then

$$W_1 \cap W_2 = \{(b, c, d, e) \mid d = bc, c = de\} \cup \{\infty\}$$

We glue the point at infinity to this space when c is large, but this is the same as either d or e being large. Hence there is a homeomorphism,

$$\begin{aligned} W_1 \cap W_2 &\cong \{(b, d, e) \mid d = bde\} \cup \{\infty\} \\ &= \{(b, d, e) \mid 0 = (be - 1)d\} \cup \{\infty\} \\ &= \{(b, d, e) \mid d = 0\} \cup \{(b, d, e) \mid be = 1\} \cup \{\infty\} \end{aligned}$$

These new spaces have computable homology, since they can be identified with spaces built from spheres. The first is relatively easy,

$$Z_1 = \{(b, d, e) \mid d = 0\} \cup \{\infty\} = \{(b, 0, 0, d)\} \cup \{\infty\} \cong S^4$$

The second is slightly harder.

$$Z_2 = \{(b, d, e) \mid be = 1\} \cup \{\infty\} = \left\{ \left(b, \frac{d}{b}, d, \frac{1}{b} \right) \right\} \cup \{\infty\}$$

This set is homeomorphic to the one-point compactification of $\{(b, d) \in \mathbb{C}^* \times \mathbb{C}\}$. By [Dup68, Chapter 2.3], this is given by the smash product of the one-point compactifications,

$$Z_2 \simeq (\mathbb{C}^* \times \mathbb{C})^+ \cong (\mathbb{C}^*)^+ \wedge \mathbb{C}^+ \cong (S^1 \vee S^2) \wedge S^2 = S^3 \vee S^4$$

The intersection of these spaces is

$$Z_1 \cap Z_2 = \left\{ \left(b, 0, 0, \frac{1}{b} \right) \right\} \cup \{\infty\} \simeq S^1 \vee S^2$$

From all of this, we get a commutative diagram of inclusions,

$$\begin{array}{ccc} S^1 \vee S^2 & \longrightarrow & S^3 \vee S^4 \\ \downarrow & & \downarrow \\ S^4 & \longrightarrow & W_1 \cap W_2 \end{array}$$

The two maps from $S^1 \vee S^2$ are zero in cohomology (when the degree is greater than zero) since in each degree either the domain or the codomain is zero. By applying Mayer-Vietoris, we get

$$H^q(W_1 \cap W_2) = \begin{cases} \mathbb{Z} & q = 0 \\ 0 & q = 1 \\ \mathbb{Z} & q = 2 \\ \mathbb{Z}^2 & q = 3 \\ \mathbb{Z}^2 & q = 4 \\ 0 & \text{otherwise} \end{cases}$$

where the case $q = 0$ follows from $W_1 \cap W_2$ being connected and $q = 1$ follows from writing out the relevant part of the long exact sequence,

$$H^0(S^3 \vee S^4) \oplus H^0(S^4) \twoheadrightarrow H^0(S^1 \vee S^4) \longrightarrow H^1(W_1 \cap W_2) \longrightarrow 0$$

W_1 and W_2 are both a copy of S^6 since we can get one of the coordinates from two of the others. Hence we get another diagram,

$$\begin{array}{ccc} W_1 \cap W_2 & \longrightarrow & S^6 \\ \downarrow & & \downarrow \\ S^6 & \longrightarrow & S^8 - \tilde{Z} \end{array}$$

and again the map from the intersection is zero. Repeating the above, we can compute:

$$H^q(S^8 - \tilde{Z}) = \begin{cases} \mathbb{Z} & q = 0 \\ 0 & q = 1 \\ 0 & q = 2 \\ \mathbb{Z} & q = 3 \\ \mathbb{Z}^2 & q = 4 \\ \mathbb{Z}^2 & q = 5 \\ \mathbb{Z}^2 & q = 6 \\ 0 & \text{otherwise} \end{cases}$$

Applying Alexander duality then gives

$$H^q(\tilde{Z}) = \begin{cases} \mathbb{Z} & q = 0 \\ \mathbb{Z}^2 & q = 1 \\ \mathbb{Z}^2 & q = 2 \\ \mathbb{Z}^2 & q = 3 \\ \mathbb{Z} & q = 4 \\ 0 & \text{otherwise} \end{cases}$$

from which we get that the cohomology of Z is the same as the cohomology of S^2 by factoring out the torus. The generator of $H^2(Z)$ will be called z_{11} with $z_{11}^2 = 0$.

All in all, this shows

$$H^q(X_\emptyset) = \begin{cases} \mathbb{Z} & q = 0 \\ \mathbb{Z}^7 & q = 1 \\ \mathbb{Z}^{22} & q = 2 \\ \mathbb{Z}^{42} & q = 3 \\ \mathbb{Z}^{56} & q = 4 \\ \mathbb{Z}^{56} & q = 5 \\ \mathbb{Z}^{42} & q = 6 \\ \mathbb{Z}^{22} & q = 7 \\ \mathbb{Z}^7 & q = 8 \\ \mathbb{Z} & q = 9 \\ 0 & \text{otherwise} \end{cases}$$

and the generators are

$$H^*(X_\emptyset) = \bigwedge^* [a_{11}, d_{11}, g_{11}, h_{11}, i_{11}, C_{11}, D_{11}] \otimes H^*(Z)$$

A note on the cohomology of Z

The space Z defined above is a manifold, which we will prove by applying the implicit function theorem. Consider the formula defining Z ,

$$(x, y, z) \mapsto y - z - xyz - 1$$

Taking partial derivatives gives the three functions

$$\begin{aligned} (x, y, z) &\mapsto yz \\ (x, y, z) &\mapsto 1 - xz \\ (x, y, z) &\mapsto -1 - xy \end{aligned}$$

If the first function is zero, then at least one of the two others must be non-zero. By the implicit function theorem, the coordinate that has non-zero derivative can be written as a smooth function of the other coordinates. This shows that Z is locally the graph of a smooth function, so it is a smooth manifold.

Later on it will be necessary to have a description of $H^2(Z)$. This is done by considering the subspace

$$R = \left\{ (x, y, z) \in \mathbb{R}^3 \mid x > 0, y < 0, xy < -1, z = \frac{y-1}{1+xy} > 0 \right\} \subset Z$$

This space is closed in Z since any sequence in R that converges in Z must also converge in R . Consider a subbundle $NR \cong R \times D$ of the normal bundle of R in the tangent bundle TZ , such that the exponential map is a diffeomorphism on NR . This gives us a diagram

$$H^2(NR, NR_0) \xleftarrow{\cong} H^2(Z, Z - R) \xrightarrow{j^*} H^2(Z)$$

where NR_0 is NR without the zero section, $NR_0 \cong R \times (D - 0)$. The isomorphism is given by excision, since $Z = NR \cup (Z - R)$. The space R is homeomorphic to \mathbb{R}^2 , so we have

$$H^2(Z, Z - R) \cong H^2(NR, NR_0) \cong H^2(R \times (D, D - 0)) \cong \mathbb{Z}$$

From the diagram, $j^*(1)$ gives us an element of $H^2(Z)$. To see that this is a generator, consider the torus

$$\hat{T} = \{(x, y, z) \in Z \mid |x| = |y| = 2\} \subset Z$$

This intersects R in exactly one point, namely $p = (2, -2, 1)$. By taking the intersection with \hat{T} we can extend the previous diagram,

$$\begin{array}{ccccc} H^2(NR, NR_0) & \xleftarrow{\cong} & H^2(Z, Z - R) & \xrightarrow{j^*} & H^2(Z) \\ \downarrow i^* & & \downarrow i^* & & \downarrow i^* \\ H^2(p \times (D, D - 0)) & \xleftarrow{\cong} & H^2(\hat{T}, \hat{T} - p) & \xrightarrow{\cong} & H^2(\hat{T}) \end{array}$$

The left i^* is an isomorphism since the cohomology in degree two is given by the pair $(D, D - 0)$, which does not change under the inclusion. This shows that the inclusion $\hat{T} \rightarrow Z$ is an isomorphism in cohomology, so $j^*(1)$ is a generator of $H^2(Z)$. Taking the same diagram with cohomology replaced with homology reverses all arrows but is otherwise exactly the same. This shows that to calculate the homology class of a cycle σ in $H_2(Z)$, we can count the number of points of intersection between σ and R with orientation, since each of these will contribute either 1 or -1 , depending on orientation, to the homology class in $(Z, Z - R)$.

There is a similar argument with the space Y from before and the subspace $R_Y = \{(b, c) \in \mathbb{R}^2 \mid b > 0, c > 0, bc > 1\}$. In particular, the generator of $H_2(Y)$ can be taken as $i_*([\hat{T}])$, where $[\hat{T}]$ is a generator of $H_2(\hat{T})$ for the torus $\hat{T} = \{(b, c) \mid |b| = |c| = 2\}$.

The first page

With all of the above, we can draw the E_1 -page of the spectral sequence. Remember that this is

$$E_1^{p,q} \cong \begin{cases} H^q(X_\emptyset) & p = 0 \\ H^{q+1}(X_{\{1\}} \times (D, D - 0)) \oplus H^{q+1}(X_{\{2\}} \times (D, D - 0)) & p = 1 \\ H^{q+2}(X_{\{1,2\}} \times (D_2, D_2 - 0)) & p = 2 \\ 0 & \text{otherwise} \end{cases}$$


By inserting the groups that were calculated above into this, we get Figure 2.1 with the non-zero differentials moving directly to the right. To get the E_2 -page of the spectral sequence, we need to compute the homology of the differentials.

2.3 Differentials on E_1

By definition, the differentials are the boundary maps of the long exact sequence of the triple (F_{p+1}, F_p, F_{p-1}) ,

$$d_1 : E_1^{p,q} = H^{p+q}(F_p, F_{p-1}) \rightarrow H^{p+q+1}(F_{p+1}, F_p) = E_1^{p+1,q}$$

We will consider the two cases $p = 0$ and $p = 1$ separately.

Figure 2.1: The E_1 -page with differentials.

First column

For $p = 0$, we are looking at the boundary map of the pair (F_1, F_0) . Consider the inclusion of pairs,

$$i : X_{\{1\}} \times (D, D - 0) \sqcup X_{\{2\}} \times (D, D - 0) \rightarrow (F_1, F_0)$$

This is an isomorphism in cohomology and gives a map between the long exact sequences that makes the following diagram commute:

$$\begin{array}{ccc}
 H^{p+q}(F_0) & \xrightarrow{d_1} & H^{p+q+1}(F_1, F_0) \\
 \downarrow i^* & & \downarrow \cong i^* \\
 H^{p+q}((X_{\{1\}} \sqcup X_{\{2\}}) \times (D - 0)) & \xrightarrow{\delta} & H^{p+q+1}((X_{\{1\}} \sqcup X_{\{2\}}) \times (D, D - 0))
 \end{array}$$

The bottom map is the boundary map from the long exact sequence of the pair $(D, D - 0)$. This map is well-known and will be treated at the end of this section, so the only missing part is to calculate the inclusion i^* . Since we are working with a disjoint union, we only need to calculate the inclusion on each factor.

The inclusion $X_{\{1\}} \times (D - 0) \hookrightarrow X_\emptyset$

We can write the inclusion out in coordinates as

$$\begin{aligned} (\mathbb{C}^*)^5 \times Y \times (D - 0) &\longrightarrow (\mathbb{C}^*)^7 \times Z \\ (a, d, f, h, i, b, c, g) &\mapsto \left(g, h, i, \frac{a}{g}, -f, d, \right. \\ &\quad \left. C = c - \frac{a}{g} - \frac{cd}{f} - bc^2 + \frac{abc}{g}, \right. \\ &\quad \left. \frac{bfC}{d}, \frac{c}{C} - \frac{a}{gC}, -\frac{cd}{fC} \right) \end{aligned}$$

On the generators, this is given by

$$\begin{aligned} i^* : H^*(X_\emptyset) &\rightarrow H^*(X_{\{1\}} \times (D - 0)) \\ a_{11} &\mapsto a_{01} - g_{01} \\ d_{11} &\mapsto f_{01} \\ g_{11} &\mapsto g_{01} \\ h_{11} &\mapsto h_{01} \\ i_{11} &\mapsto i_{01} \\ D_{11} &\mapsto d_{01} \\ C_{11} &\mapsto a_{01} - g_{01} + y_{01,1} \\ z_{11} &\mapsto y_{01,2} + (a_{01} - d_{01} + f_{01} - g_{01})y_{01,1} \end{aligned}$$

where g_{01} is the generator of $H^1(D - 0)$.

In degree 1, this is calculated by dualizing to homology, writing out the map on generators and working out what they map to. For z_{11} it is slightly harder. To calculate this, we use the submanifold $R \subset Z$ defined previously as

$$R = \left\{ (x, y, z) \in \mathbb{R}^3 \mid x > 0, y < 0, z = \frac{y-1}{1+xy} > 0 \right\}$$

by choosing a generator $\alpha = [\sigma]$ of $H_2(X_{\{1\}} \times (D - 0))$ and calculating the intersection of the image of $f = \text{pr}_Z \circ i \circ \sigma$ and R . Note that if the coordinates of Y are fixed at $(0, 1)$, then the image of

$$f(a, d, f, h, i, 0, 1, g) = \left(0, \frac{1}{C} - \frac{a}{gC}, -\frac{d}{fC} \right)$$

never intersects R since $x = 0$. Likewise, the coordinates i and h do not appear in the projection to Z and can not influence the result. Hence we only have to check the duals of the following generators in degree 2:

$$a_{01}y_{01,1}, d_{01}y_{01,1}, f_{01}y_{01,1}, y_{01,1}g_{01}, y_{01,2}$$

These will be handled individually. All non-relevant coordinates will be set to 1, except g which will be denoted ε to make it clear that this coordinate is small. If $y_{01,1}$ is involved, we choose the generator

$$\sigma(\lambda) = (1 - \lambda, 1) \quad \lambda \in S^1$$

as this gives

$$y_{01,1} = [t \mapsto 1 - (1 - \lambda) \cdot 1] = [t \mapsto \lambda]$$

a₀₁y_{01,1}: The relevant map is

$$(a, \lambda) \mapsto \left((1 - \lambda)C, \frac{1}{C} - \frac{a}{\varepsilon C}, \frac{1}{C} \right)$$

where $C = -1 - \frac{a\lambda}{\varepsilon}$. If this intersects R then C has to be real and positive by looking at the last coordinate, which forces λ real and negative by the first coordinate and a real and positive by the second. Hence there is at most one intersection, at $(1, -1)$:

$$f(1, -1) = \left(\frac{2 - 2\varepsilon}{\varepsilon}, -1, \frac{\varepsilon}{1 - \varepsilon} \right)$$

By inspection, this satisfies the conditions and hence belongs to R , so we conclude that the image intersects once. This gives the following formula:

$$\langle f^*(z_{11}), a_{01}y_{01,1} \rangle = \langle z_{11}, f_*(a_{01}y_{01,1}) \rangle = \pm 1$$

To check the sign, we need to check if this map preserves the orientation of the torus in R . This is done by differentiating the map to get a map of tangent spaces

$$Tf : TT^2 \rightarrow TR$$

and checking that the determinant of this map is positive. This has been done in Sage.

d₀₁y_{01,1}: In this case the map becomes

$$(d, \lambda) \mapsto \left(\frac{(1 - \lambda)C}{d}, \frac{\varepsilon - 1}{\varepsilon C}, \frac{d}{C} \right)$$

and by a similar analysis, this intersects in exactly the point $f(1, -1)$, which is the same as before. However, this time the orientation is reversed.

f₀₁y_{01,1}: This has the same point of intersection and preserves the orientation.

y_{01,1}g₀₁: This also intersects and preserves the orientation.

y_{01,2}: This is calculated by the intersection points of the fundamental class of the torus in Y , which was mentioned earlier:

$$\hat{T} = \{(b, c) \in \mathbb{C}^2 \mid |b| = |c| = 2\} \subset Y$$

The map becomes

$$(b, c) \mapsto \left(bC, \frac{c}{C} - \frac{1}{\varepsilon C}, \frac{c}{C} \right)$$

with $C = -bc^2 - \frac{1-bc}{\varepsilon}$. We will write x for the first coordinate, y for the second and z for the third. An analysis as before gives $bc = xz > 0$ and $c = \frac{1}{\varepsilon(z-y)} \in \mathbb{R}$. So both b and c must be real and c should be positive. Then $xy = bc - b\varepsilon^{-1} < 0$ shows that b must be positive since bc is positive. The only possible point of intersection is

$$f(2, 2) = \left(\frac{6 - 16\varepsilon}{\varepsilon}, \frac{2\varepsilon - 1}{3 - 8\varepsilon}, \frac{2\varepsilon}{3 - 8\varepsilon} \right)$$

Since ε is very small, this intersects R and the orientation is preserved.

The inclusion $X_{\{2\}} \times (D - 0) \hookrightarrow X_\emptyset$

The inclusion in coordinates is

$$\begin{aligned} (\mathbb{C}^*)^5 \times Y \times (D - 0) &\longrightarrow (\mathbb{C}^*)^7 \times Z \\ (a, d, f, g, i, b, c, h) &\mapsto \left(g, h, i, a, -\frac{d}{h}, \right. \\ &\quad D = \frac{d}{h}(1 - bc), \\ &\quad C = -af - \frac{a^2bh}{d} + \frac{a^2bhf}{d} + \frac{a^2b^2ch}{d}, \\ &\quad \left. \frac{bC}{D}, \frac{cd}{hC} - \frac{a}{C}, \frac{cD}{C} - \frac{ahD}{dC} + \frac{afhD}{dC} + \frac{abchD}{dC} \right) \end{aligned}$$

We calculate as before and get the following on generators:

$$\begin{aligned} i^* : H^*(X_\emptyset) &\rightarrow H^*(X_{\{2\}} \times (D - 0)) \\ a_{11} &\mapsto a_{10} \\ d_{11} &\mapsto d_{10} - h_{10} \\ g_{11} &\mapsto g_{10} \\ h_{11} &\mapsto h_{10} \\ i_{11} &\mapsto i_{10} \\ D_{11} &\mapsto d_{10} + y_{10,1} - h_{10} \\ C_{11} &\mapsto f_{10} + a_{10} \\ z_{11} &\mapsto y_{10,2} + (a_{10} - d_{10} + f_{10} + h_{10})y_{10,1} \end{aligned}$$

where h_{10} is the generator of $H^1(D - 0)$.

The calculations have been left out, but are very similar to the previous ones.

Second column

The differential on the second column is the boundary map

$$\delta : H^q((X_{\{1\}} \sqcup X_{\{2\}}) \times (D, D - 0)) \cong H^q(F_1, F_0) \rightarrow H^{q+1}(F_2, F_1)$$

To calculate this map, consider the inclusion of triples

$$X_{\{1,2\}} \times (D_2, D_2 - 0, (D - 0) \times (D - 0)) \hookrightarrow (F_2, F_1, F_0)$$

given by inserting a small disc, sphere or torus on the zero-entries of $X_{\{1,2\}}$. There are homeomorphisms of pairs,

$$\begin{aligned} (D_2 - 0, (D - 0) \times (D - 0)) &\cong (D \times (D - 0) \sqcup (D - 0) \times D, (D - 0) \times (D - 0)) \\ (D_2, D_2 - 0) &\cong (D \times D, D \times (D - 0) \cup (D - 0) \times D) = (D, D - 0) \times (D, D - 0) \end{aligned}$$

These gives a commuting diagram,

$$\begin{array}{ccc} H^q(F_1, F_0) & \xrightarrow{\quad\quad\quad} & H^{q+1}(F_2, F_1) \\ \downarrow & & \downarrow \cong \\ H^q(X_{\{1,2\}} \times (D_2 - 0, (D - 0) \times (D - 0))) & \longrightarrow & H^{q+1}(X_{\{1,2\}} \times (D_2, D_2 - 0)) \end{array}$$

The marked isomorphism follows from excision and the left map is induced by an inclusion. If we apply the homeomorphism of pairs to the bottom of the diagram, the boundary map at the bottom becomes the sum of the two boundary maps,

$$\begin{array}{ccc} H^q(X_{\{1,2\}} \times (D, D-0) \times (D-0)) & \xrightarrow{\delta} & \\ & \searrow & \\ & & H^{q+1}(X_{\{1,2\}} \times (D, D-0) \times (D, D-0)) \\ & \nearrow & \\ H^q(X_{\{1,2\}} \times (D-0) \times (D, D-0)) & \xrightarrow{\delta} & \end{array}$$

The boundary maps are given by the long exact sequence of the pair $(D, D-0)$, so we only need to consider the inclusion on the left. By choosing the discs sufficiently small and using the identification of $H^q(F_1, F_0)$ defined in the first section, we see that the inclusion maps the space $X_{\{1,2\}} \times (D, D-0) \times (D-0)$ to $X_{\{1\}} \times (D, D-0)$ and $X_{\{1,2\}} \times (D-0) \times (D, D-0)$ to $X_{\{2\}} \times (D, D-0)$. To get the differential, we must calculate the sum of these two maps:

$$\begin{aligned} i_1^* : H^q(X_{\{1\}} \times (D, D-0)) &\rightarrow H^q(X_{\{1,2\}} \times (D, D-0) \times (D-0)) \\ i_2^* : H^q(X_{\{2\}} \times (D, D-0)) &\rightarrow H^q(X_{\{1,2\}} \times (D-0) \times (D, D-0)) \end{aligned}$$

The inclusion $X_{\{1,2\}} \times (D, D-0) \times (D-0) \rightarrow X_{\{1\}} \times (D, D-0)$

In coordinates, the map becomes

$$\begin{aligned} (\mathbb{C}^*)^3 \times Y \times (D, D-0) \times (D-0) &\rightarrow (\mathbb{C}^*)^5 \times Y \times (D, D-0) \\ (a, d, i, b, c, g, h) &\mapsto \left(a, i, h, \frac{d}{h}, \frac{d}{h}(1-bc), \frac{bh}{d}, \frac{dc}{h}, g \right) \end{aligned}$$

Again, we calculate the map on cohomology by dualizing to homology and working with the dual generators. To check what happens to the elements of $H^q(Y)$, we count intersections with the subspace $R_Y = \{(b, c) \in \mathbb{R}^2 \mid b > 0, c > 0, bc > 1\}$ of Y , exactly as for Z above.

$$\begin{aligned} i_1^* : H^q(X_{\{1\}} \times (D, D-0)) &\rightarrow H^q(X_{\{1,2\}} \times (D, D-0) \times (D-0)) \\ a_{01} &\mapsto a_{00} \\ d_{01} &\mapsto d_{00} - h_{00} + y_{00,1} \\ f_{01} &\mapsto d_{00} - h_{00} \\ g_{01} &\mapsto g_{00} \\ h_{01} &\mapsto h_{00} \\ i_{01} &\mapsto i_{00} \\ y_{01,1} &\mapsto y_{00,1} \\ y_{01,2} &\mapsto y_{00,2} - d_{00}y_{00,1} + h_{00}y_{00,1} \end{aligned}$$

We denote the generator of $H^2(D, D-0)$ in $H^*(X_{\{1\}} \times (D, D-0))$ by g_{01} and for $H^*(X_\emptyset \times (D, D-0) \times (D-0))$ we consider $g_{00} \in H^2(D, D-0)$ and $h_{00} \in H^1(D-0)$.

The inclusion $X_{\{1,2\}} \times (D-0) \times (D, D-0) \rightarrow X_{\{2\}} \times (D, D-0)$

This is exactly as above for the map

$$\begin{aligned} (\mathbb{C}^*)^3 \times Y \times (D-0) \times (D, D-0) &\rightarrow (\mathbb{C}^*)^5 \times Y \times (D, D-0) \\ (a, d, i, b, c, g, h) &\mapsto \left(\frac{a}{g}, d, i, g, 1-bc, b, c, h \right) \end{aligned}$$

In cohomology, this becomes

$$\begin{aligned} i_2^* : H^q(X_{\{2\}} \times (D, D-0)) &\rightarrow H^q(X_{\{1,2\}} \times (D-0) \times (D, D-0)) \\ a_{10} &\mapsto a_{00} - g_{00} \\ d_{10} &\mapsto d_{00} \\ f_{10} &\mapsto y_{00,1} \\ g_{10} &\mapsto g_{00} \\ h_{10} &\mapsto h_{00} \\ i_{10} &\mapsto i_{00} \\ y_{10,1} &\mapsto y_{00,1} \\ y_{10,2} &\mapsto y_{00,2} \end{aligned}$$

with $g_{01} \in H^2(D, D-0)$ inside $H^*(X_{\{2\}} \times (D, D-0))$ and $g_{00} \in H^2(D, D-0)$, $h_{00} \in H^1(D-0)$ inside $H^*(X_\emptyset \times (D-0) \times (D, D-0))$.

Boundary maps

We have now reduced everything to calculating some well-known boundary maps for the pair $(D, D-0)$. By writing out the long exact sequence, the boundary map

$$\delta : H^*(D-0) \rightarrow H^*(D, D-0)$$

is an isomorphism in degree 1,

$$H^1(D-0) \cong H^2(D, D-0)$$

and zero in all other degrees. The boundary map for the pair $X \times (D, D-0)$ is $\text{Id} \otimes \delta$ in tensor-notation. So the boundary map in the cases above picks out the “new” coordinate added to $X \times (D-0)$ and throws away everything that does not depend on it. To be more precise, the map is given on generators by:

$$\begin{aligned} H^*(X) \otimes H^*(D-0) &\rightarrow H^*(X) \otimes H^*(D, D-0) \\ x \otimes d &\mapsto \begin{cases} x \otimes \delta d & d \in H^1(D-0) \\ 0 & \text{otherwise} \end{cases} \end{aligned}$$

2.4 Turning the page

We can now use the above to calculate the E_2 -page. We have calculated the groups appearing on the first page and the differential. To get the second page, we need to compute the homology groups of d_1 . This reduces to linear algebra over the integers, and is done using Sage. The result can be found in Figure 2.2.

9	\mathbb{Z}	0	0
8	\mathbb{Z}^5	0	0
7	\mathbb{Z}^{11}	0	0
6	\mathbb{Z}^{15}	\mathbb{Z}	0
5	\mathbb{Z}^{16}	\mathbb{Z}^3	0
4	\mathbb{Z}^{16}	\mathbb{Z}^3	0
3	\mathbb{Z}^{15}	\mathbb{Z}	0
2	\mathbb{Z}^{11}	0	0
1	\mathbb{Z}^5	0	0
0	\mathbb{Z}	0	0
q/p	0	1	2

Figure 2.2: The E_2 -page.

The differentials from E_2 onwards are given as maps:

$$d_r : E_r^{p,q} \rightarrow E_r^{p+r, q+1-r}$$

For $r \geq 2$, this increases p by at least 2. This means that the domain or codomain of d_r will always be zero, so taking homology does not change anything. Hence $E_2 = E_\infty$, the spectral sequence collapses at the 2nd term, and there are no extension problems since all the groups are free and abelian. This allows us to read off the cohomology of $X_{3,3}$. Since we are also interested in factoring out the T^3 torus, we note this as well.

$$H^q(X_{3,3}) = E_2^{0,q} \oplus E_2^{1,q-1} = \begin{cases} \mathbb{Z} & q = 0 \\ \mathbb{Z}^5 & q = 1 \\ \mathbb{Z}^{11} & q = 2 \\ \mathbb{Z}^{15} & q = 3 \\ \mathbb{Z}^{17} & q = 4 \\ \mathbb{Z}^{19} & q = 5 \\ \mathbb{Z}^{18} & q = 6 \\ \mathbb{Z}^{12} & q = 7 \\ \mathbb{Z}^5 & q = 8 \\ \mathbb{Z} & q = 9 \\ 0 & \text{otherwise} \end{cases} \quad H^q(Y_{3,3}) = \begin{cases} \mathbb{Z} & q = 0 \\ \mathbb{Z}^2 & q = 1 \\ \mathbb{Z}^2 & q = 2 \\ \mathbb{Z}^2 & q = 3 \\ \mathbb{Z}^3 & q = 4 \\ \mathbb{Z}^2 & q = 5 \\ \mathbb{Z} & q = 6 \\ 0 & \text{otherwise} \end{cases}$$

We can also read off the ring structure from the spectral sequence, as it is induced from the spaces we start with. Since we do not need it, it will not be included here.

Chapter 3

Loop spaces

We now return to the general case. We want to show that the space $Y_{m,\infty}$ is equivalent to the space of loops in the flag manifold GL_m/B_m .

Definition 6. The loop space $\Omega = \Omega(\mathrm{GL}_m/B_m)$ is the space

$$\Omega = \{\gamma : [0, 1] \rightarrow \mathrm{GL}_m/B_m \mid \gamma(0) = \gamma(1) = [\mathrm{Id}]\}$$

with the compact-open topology.

Remark 7. Following [Mil63, Chapter 17], the compact-open topology is also the topology given by the metric

$$d(\gamma, \gamma') = \max_{t \in [0, 1]} \tilde{d}(\gamma(t), \gamma'(t))$$

where \tilde{d} is a metric on GL_m/B_m induced from a Riemannian metric. The space Ω is homotopy equivalent to the space of piecewise smooth loops with a slightly modified metric, and we will generally choose the most convenient model for our arguments.

3.1 The map $Y_{m,n} \rightarrow \Omega$

For $A = [a_1, \dots, a_n] \in X_{m,n}$ we can define a path γ_A in GL_m by starting with $\gamma_A(0) = \mathrm{Id}$. Since the matrix $[e_2, \dots, e_m, a_1]$ is invertible by the definition of $X_{m,n}$, we know that a_1 is linearly independent of e_2, \dots, e_m . Hence it can be written as

$$a_1 = \sum_{i=1}^m c_i e_i$$

with $c_1 \neq 0$. Rewriting the expression leads to the formula

$$\frac{1}{c_1} a_1 = e_1 + \frac{1}{c_1} \left(\sum_{i=2}^m c_i e_i \right)$$

This gives us a path in GL_m from the identity to $\left[\frac{1}{c_1} a_1, e_2, \dots, e_m \right]$:

$$[0, 1] \ni t \mapsto \left[e_1 + t \left(\sum_{i=2}^m \frac{c_i}{c_1} e_i \right), e_2, \dots, e_m \right] \in \mathrm{GL}_m$$

This process can be continued, since we can write

$$\frac{1}{d_2}a_2 = e_2 + \frac{1}{d_2} \left(\sum_{i=3}^m d_i e_i \right) + \frac{d_1}{d_2 c_1} a_1$$

and we get a path

$$t \mapsto \left[\frac{1}{c_1} a_1, e_2 + t \left(\sum_{i=3}^m \frac{d_i}{d_2} e_i + \frac{d_1}{d_2 c_1} a_1 \right), e_3, \dots, e_m \right]$$

that starts where the other path stopped. If we continue through the vectors in A , we will eventually end at a point

$$[\lambda_i e_i, \dots, \lambda_m e_m, \lambda_1 e_1, \dots, \lambda_{i-1} e_{i-1}] \in \mathrm{GL}_m$$

which is diagonal, except the columns may have been permuted if n is not divisible by m . Choosing a path in GL_m from $[e_i, \dots, e_{i-1}]$ to Id , scaling the columns and appending it to all the other paths then gives a path γ_A with

$$\gamma_A(0) = \mathrm{Id}, \quad \gamma_A(1) = [\lambda_1 e_1, \dots, \lambda_m e_m]$$

If we consider this as a path in GL_m/B_m we get a loop, since $\gamma_A(1)$ is a diagonal matrix. The assignment

$$f(A) = \gamma_A \in \Omega(\mathrm{GL}_m/B_m)$$

defines a continuous function from $X_{m,n}$ to the loop space. Note that this function descends to the quotient space $Y_{m,n} = X_{m,n}/T^n$, since elements of $X_{m,n}$ that are equivalent under the T^n -action gives paths in GL_m that are equivalent under the B_m -action.

The goal of this chapter is to prove the following theorem:

Theorem 8. *The map $f : Y_{m,n} \rightarrow \Omega$ induces a map*

$$f : Y_{m,\infty} \rightarrow \Omega$$

This is a (weak) homotopy equivalence.

Since the space Ω is relatively well-studied, this gives information about $Y_{m,\infty}$. In particular, it allows us to calculate the homology and cohomology of this space.

We want to show that the stabilization map

$$s : Y_{m,n} \rightarrow Y_{m,n+1}$$

$$A = (a_1, \dots, a_n) \mapsto \left(La_1, \dots, La_n, \begin{pmatrix} 1 \\ \vdots \\ 1 \end{pmatrix} \right) = (LA, Le_1)$$

does not affect the map f , or more precisely that f induces a well-defined map on $Y_{m,\infty}$. To do this we will work with the subsequence $Y_{m,nm}$ of the sequence $Y_{m,n}$, as the two limits are isomorphic.

We will show that the diagram

$$\begin{array}{ccc} Y_{m,nm} & \xrightarrow{s^m} & Y_{m,(n+1)m} \\ & \searrow f & \swarrow f \\ & \Omega & \end{array}$$

commutes up to homotopy. This is done by showing that the map s^m is homotopic to the map \tilde{s} that inserts the identity on the last m columns of $A \in Y_{m,nm}$:

$$\tilde{s}(a_1, \dots, a_{nm}) = (a_1, \dots, a_{nm}, e_1, \dots, e_m)$$

If we connect Id and L^{-m} by a path γ in the space of invertible lower triangular matrices, we get a homotopy

$$s_t(A) = (\gamma(t)L^m A, \gamma(t)L^m e_1, \gamma(t)L^{m-1} e_1, \dots, \gamma(t)L e_1)$$

which goes from $s_0 = s^m$ to the map

$$s_1 : A \mapsto (A, e_1, L^{-1}e_1, \dots, L^{-m+1}e_1)$$

The matrix $T = [e_1, L^{-1}e_1, \dots, L^{-m+1}e_1]$ is upper triangular with entries given by the formula:

$$T_{ij} = (L^{-j+1}e_1)_i = (-1)^{i-1} \binom{j-1}{i-1} \quad \text{when } i \leq j$$

This is proven by considering negative powers of L and showing the identity:

$$(L^{-j+1})_{ik} = (-1)^{i-k} \binom{j-1}{i-k}$$

But since T is an upper triangular matrix, it can be moved to the identity without affecting the linear independence. Hence the maps s^m and \tilde{s} are homotopic. The composition $f \circ \tilde{s}$ is the same as f , except that the loop $f \circ \tilde{s}$ is stationary at the identity for a while before it terminates. Putting it all together shows the following:

Lemma 9. *The diagram*

$$\begin{array}{ccc} Y_{m,nm} & \xrightarrow{s^m} & Y_{m,(n+1)m} \\ & \searrow f & \swarrow f \\ & \Omega & \end{array}$$

commutes up to homotopy and f induces a map:

$$f : Y_{m,\infty} \rightarrow \Omega$$

3.2 Homotopy equivalence

To show that the map $f : Y_{m,\infty} \rightarrow \Omega$ gives an isomorphism of homotopy groups, we will again follow [Mil63]. For each element U of SU_m/T^{m-1} , we can find a geodesically convex open ball centered on U . Since SU_m/T^{m-1} is compact, there is some $\varepsilon > 0$ such that any ball of radius ε or less is geodesically convex. By the same argument, we can also choose ε so small that for any two points $A = [a_1, \dots, a_m]$ and $B = [b_1, \dots, b_m]$, with the distance from A to B less than ε , any m subsequent vectors in the sequence $(a_1, \dots, a_m, b_1, \dots, b_m)$ are linearly independent. With this, we get an increasing sequence of open sets in Ω , defined by requiring the loops to be piecewise contained in a ball of radius ε :

$$\Omega_k = \left\{ \gamma \in \Omega \mid \gamma|_{\left[\frac{j-1}{2^k}, \frac{j}{2^k}\right]} \text{ is contained in an } \varepsilon\text{-ball } B_\varepsilon \right\}$$

By the choice of ε made above, we can define a map

$$\begin{aligned}\varphi_k : \Omega_k &\rightarrow Y_{m, (2^k - 1)m} \\ \gamma &\mapsto \left[\gamma\left(\frac{1}{2^k}\right), \gamma\left(\frac{2}{2^k}\right), \dots, \gamma\left(\frac{2^k - 1}{2^k}\right) \right]\end{aligned}$$

Note that $\gamma(t)$ is an element of SU_m/T^{m-1} , so the notation $[\gamma(t), \dots]$ is shorthand for $[\gamma(t)e_1, \dots, \gamma(t)e_m, \dots]$. The map is continuous since two paths being close in Ω_k means that the maximum distance between them is small. But this translates to the vectors of the image being close together. The maps φ_k and φ_{k+1} are related in the following fashion:

Lemma 10. *The following diagram commutes up to homotopy:*

$$\begin{array}{ccc}\Omega_k & \xhookrightarrow{\quad} & \Omega_{k+1} \\ \downarrow \varphi_k & & \downarrow \varphi_{k+1} \\ Y_{m, (2^k - 1)m} & \xrightarrow{s} & Y_{m, (2^{k+1} - 1)m}\end{array}$$

Proof. The proof is by direct computation.

$$\begin{aligned}\varphi_{k+1} &= \left(\gamma \mapsto \left[\gamma\left(\frac{1}{2^{k+1}}\right), \gamma\left(\frac{2}{2^{k+1}}\right), \dots, \gamma\left(\frac{2^{k+1} - 1}{2^{k+1}}\right) \right] \right) \\ &\simeq \left(\gamma \mapsto \left[\gamma\left(\frac{2}{2^{k+1}}\right), \gamma\left(\frac{2}{2^{k+1}}\right), \gamma\left(\frac{4}{2^{k+1}}\right), \dots, \gamma\left(\frac{2^{k+1}}{2^{k+1}}\right) \right] \right) \\ &\simeq \left(\gamma \mapsto \left[\gamma\left(\frac{1}{2^k}\right), \gamma\left(\frac{2}{2^k}\right), \dots, \gamma\left(\frac{2^k - 1}{2^k}\right), \mathrm{Id}, \dots, \mathrm{Id} \right] \right) \\ &= s \circ \varphi_k\end{aligned}$$

The first homotopy is given by sliding along the minimal geodesic between the points and gives an allowed path by our choice of ε . The second homotopy is somewhat similar. If (a_1, \dots, a_m) and (b_1, \dots, b_m) are two elements of SU_m/T^{m-1} and we have an element of $Y_{m,n}$ of the form

$$[\dots, a_1, \dots, a_m, a_1, \dots, a_m, b_1, \dots, b_m, \dots]$$

then we can change the middle columns one by one, as we did in the construction of f , and get a path in $Y_{m,n}$ that changes the middle from $[a_1, \dots, a_m]$ to $[b_1, \dots, b_m]$:

$$\begin{aligned}[a_1, \dots, a_m, a_1, a_2, \dots, a_m, b_1, \dots, b_m] &\rightsquigarrow [a_1, \dots, a_m, b_1, a_2, \dots, a_m, b_1, \dots, b_m] \\ &\rightsquigarrow [a_1, \dots, a_m, b_1, b_2, \dots, a_m, b_1, \dots, b_m] \\ &\rightsquigarrow [a_1, \dots, a_m, b_1, b_2, \dots, b_m, b_1, \dots, b_m]\end{aligned}$$

This allows us to take all the repetitions $[\dots, \gamma(t), \gamma(t), \dots]$ and move them to the end as diagonal matrices, which results in the stabilization map s . \square

The above lemma shows that the maps φ_k fit together to define a map from Ω to $Y_{m,\infty}$. The idea is to use the maps φ_k as homotopy inverses of the loop space map f . We are now ready to prove the result.

Theorem 11. *The map $f : Y_{m,\infty} \rightarrow \Omega$ is a weak homotopy equivalence, i.e. the map*

$$f_* : \pi_i(Y_{m,\infty}) \rightarrow \pi_i(\Omega)$$

is an isomorphism for all i .

Proof. We will start with surjectivity. Let

$$g : W \rightarrow \Omega$$

be a finite cell complex in Ω . We want to show that we can construct a map \tilde{g} from W to $Y_{m,\infty}$ such that the composition $f \circ \tilde{g}$ is homotopic to g .

The image of g is compact and hence contained in Ω_k for some k , since $\{\Omega_k\}$ is an open cover of Ω . Consider the map

$$\tilde{g} = \varphi_k \circ g : W \rightarrow Y_{m,(2^k-1)m}$$

Evaluating at a point $w \in W$ gives

$$\tilde{g}(w) = \varphi_k(\gamma_w) = \left[\gamma_w \left(\frac{1}{2^k} \right), \dots, \gamma_w \left(\frac{2^k-1}{2^k} \right) \right]$$

Evaluating f on this gives a path connecting the points $\gamma_w \left(\frac{j}{2^k} \right)$. But by our choice of ε , any such path is homotopic to the path connecting these points with minimal geodesics, so $f \circ \tilde{g} \simeq g$.

To show injectivity, let

$$g : W \rightarrow Y_{m,\infty}$$

be a finite cell complex with $f \circ g$ null-homotopic. We want to show that g is null-homotopic in $Y_{m,\infty}$. In the direct limit, a compact set is contained in the image of $Y_{m,n}$ for some n (see [Hat02, Proposition A.1] or [May99, Chapter 9.4]), so g is represented by a map:

$$g : W \rightarrow Y_{m,n}$$

We will work with this representative. The map $f \circ g$ is null-homotopic in Ω and the image is compact, so it is contained in Ω_k for some k . By applying φ_k , we get a function

$$\varphi_k \circ f \circ g : W \rightarrow Y_{m,(2^k-1)m}$$

Since $f \circ g$ is null-homotopic, this function is null-homotopic. But by the same argument that was used in Lemma 10, the map

$$\varphi_k(f(g(w))) = \left[\gamma_w \left(\frac{1}{2^k} \right), \dots, \gamma_w \left(\frac{2^k-1}{2^k} \right) \right]$$

is homotopic to the stabilization map

$$s(g(w)) = [g(w), \text{Id}, \dots, \text{Id}]$$

with the homotopy given by sliding along the path $\gamma_w = f(g(w))$. So the map g can be stabilized to a null-homotopic map, so it must be null-homotopic in $Y_{m,\infty}$. This shows that f_* is injective on homotopy groups. \square

Since f is a weak homotopy equivalence, it follows that f_* is an isomorphism on homology and cohomology groups. The Pontrjagin homology ring of the loop space, with product given by loop concatenation, has been calculated in [GT10, Theorem 4.1] as

$$H_*(\Omega(\mathrm{SU}_m/T^{m-1})) \cong T(x_1, \dots, x_{m-1}) \otimes \mathbb{Z}[y_1, \dots, y_{m-1}]$$

where $T(x_1, \dots, x_{m-1})$ is the tensor algebra generated by x_1, \dots, x_{m-1} . The product has the relation $x_k^2 = x_p x_q + x_q x_p = 2y_1$ for $1 \leq k, p, q \leq m-1$ and $p \neq q$. The degree of x_i is 1 and the degree of y_i is $2i$. By the above result, this is also the homology of $Y_{m,\infty}$. The groups are finitely generated and torsion free, so by the Universal Coefficient Theorem the homology and cohomology groups are isomorphic as abelian groups.

3.3 Future work

In the above, we showed that the space

$$Y_{m,\infty} = \varinjlim_n X_{m,n}/T^n$$

is equivalent to the loop space $\Omega(\mathrm{SU}_m/T^{m-1})$ and used this to calculate the homology and cohomology groups of the limit. The current aim of the project is to work with the spaces $Y_{m,n}$ directly and see what we can prove about these, in particular by using the stabilization map

$$s : Y_{m,n} \rightarrow Y_{m,n+1}$$

The conjecture at the moment is the following:

Conjecture. *The induced maps on cohomology*

$$s^* : H^*(Y_{m,n+1}) \rightarrow H^*(Y_{m,n})$$

are always surjective and the induced maps on homology

$$s_* : H_*(Y_{m,n}) \rightarrow H_*(Y_{m,n+1})$$

are always injective.

The idea for a proof would be to use a spectral sequence argument like the one applied to $X_{3,3}$ by defining a filtration that is easier to work with and then proceeding by induction, using that the maps s^* and s_* induce maps of the spectral sequences.

Another thing to consider is modifying the space $X_{m,n}$. One possible change could be to replace $X_{m,n}$ by a union:

$$X'_{m,n} = \bigcup_{A \in \mathrm{GL}_m} X_{m,n}(A, A)$$

This would allow all sequences of vectors that start and end at the same matrix, and one would expect the space $Y'_{m,\infty}$ to be equivalent to the free loop space, consisting of all maps from the circle S^1 to SU_m/T^{m-1} .

There is also the option of replacing the complex numbers \mathbb{C} with some other space, like the reals or quaternions. The fact that \mathbb{R}^* is not connected would most likely make the space $X_{m,n}^{\mathbb{R}}$ quite different, but possibly the space $X_{m,n}^{\mathbb{H}}$ would behave in a similar manner to $X_{m,n}$, with some shifts in dimension.

Bibliography

- [BB05] A. Bjorner and F. Brenti.
Combinatorics of Coxeter Groups.
Graduate Texts in Mathematics.
Springer, 2005.
ISBN: 9783540442387.
URL: <http://books.google.dk/books?id=0saeruFjCRsC>.
- [Bök13] M. Bökstedt.
“bokstedt.pdf”.
2013.
URL: <http://home.math.au.dk/sstoltze/del-a/bokstedt.pdf>.
- [Dup68] J.L. Dupont.
K-theory.
Lecture notes series nr. 11.
Aarhus universitet, Matematisk institut, 1968.
- [GT10] J. Grbić and S Terzić.
“The integral Pontrjagin homology of the based loop space on a flag manifold”.
In: *Osaka J. Math.* 47.2 (June 2010), pp. 439–460.
URL: <http://projecteuclid.org/euclid.ojm/1277298912>.
- [Hat02] A. Hatcher.
Algebraic topology.
Cambridge University Press, 2002.
ISBN: 9780521795401.
URL: <http://books.google.com/books?id=BjKs86kosqgC>.
- [Hat04] A. Hatcher.
Spectral Sequences in Algebraic Topology.
2004.
URL: <http://www.math.cornell.edu/~hatcher/SSAT/SSATpage.html>.
- [Hil82] H. Hiller.
Geometry of Coxeter groups.
Research notes in mathematics.
Pitman Pub., 1982.
ISBN: 9780273085171.
URL: <https://books.google.dk/books?id=7jzvAAAAAAAJ>.
- [May99] J.P. May.
A Concise Course in Algebraic Topology.
Chicago Lectures in Mathematics.

University of Chicago Press, 1999.
ISBN: 9780226511832.
URL: <http://www.math.uchicago.edu/~may/CONCISE/ConciseRevised.pdf>.

- [McC01] J. McCleary.
A User's Guide to Spectral Sequences.
Cambridge Studies in Advanced Mathematics.
Cambridge University Press, 2001.
ISBN: 9780521567596.
URL: <https://books.google.dk/books?id=B5XPPXiezT0C>.
- [Mil63] J.W. Milnor.
Morse Theory.
Annals of mathematics studies.
Princeton University Press, 1963.
ISBN: 9780691080086.
URL: http://books.google.dk/books?id=A9QZZ3S_QxwC.