

Solution 2 (cont'd):

The improved electromagnetic energy-momentum tensor reads: $\hat{T}^{\mu\nu} = T^{\mu\nu} + \partial_\lambda(F^{\mu\lambda}A^\nu)$.

First we note that the expression for the energy-momentum tensor for a scalar field,

$$T^{\mu\nu} = \frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi)} \partial^\nu \phi - \mathcal{L} g^{\mu\nu},$$

is generalized to

$$T^{\mu\nu} = \frac{\partial \mathcal{L}}{\partial(\partial_\mu A_\sigma)} \partial^\nu A_\sigma - \mathcal{L} g^{\mu\nu}$$

for a vector field. From last week's part of this exercise we know that $\frac{\partial \mathcal{L}}{\partial(\partial_\mu A_\sigma)} = -F^{\mu\sigma}$, resulting in

$$T^{\mu\nu} = -F^{\mu\sigma} \partial^\nu A_\sigma + \frac{1}{4} F_{\alpha\beta} F^{\alpha\beta} g^{\mu\nu}.$$

With the help of

$$\partial_\lambda(F^{\mu\lambda}A^\nu) = (\partial_\lambda F^{\mu\lambda})A^\nu + F^{\mu\lambda}(\partial_\lambda A^\nu) \stackrel{\text{week 1}}{=} 0 + F^{\mu\lambda}(\partial_\lambda A^\nu)$$

and

$$-F^{\mu\sigma} \partial^\nu A_\sigma + F^{\mu\lambda}(\partial_\lambda A^\nu) = F^{\mu\lambda}(\partial_\lambda A^\nu - \partial^\nu A_\lambda) = F^{\mu\lambda} F_\lambda{}^\nu$$

one eventually obtains

$$\hat{T}^{\mu\nu} = F^{\mu\lambda} F_\lambda{}^\nu + \frac{1}{4} F_{\alpha\beta} F^{\alpha\beta} g^{\mu\nu},$$

which is indeed symmetric under $\mu \leftrightarrow \nu$. Therefore, summing over repeated indices we get

$$\begin{aligned} \mathcal{E} &= \hat{T}^{00} = F^{0\lambda} F_\lambda{}^0 + \frac{1}{4} F_{\alpha\beta} F^{\alpha\beta} g^{00} = F^{0i} F^{0i} - \frac{1}{2} F^{0i} F^{0i} g^{00} + \frac{1}{4} F^{ij} F^{ij} g^{00} \\ &= E^i E^i - \frac{1}{2} E^i E^i + \frac{1}{4} \epsilon^{ijk} B^k \epsilon^{ijl} B^l \stackrel{\text{hint week 1}}{=} \vec{E}^2 - \frac{1}{2} (\vec{E}^2 - \vec{B}^2) = \frac{1}{2} (\vec{E}^2 + \vec{B}^2) \end{aligned}$$

for the energy density carried by the electromagnetic field, and

$$S^i = \hat{T}^{0i} = \hat{T}^{i0} = F^{0\lambda} F_\lambda{}^i + 0 = F^{0j} F_j{}^i = -F^{0j} F^{ji} = E^j (-\epsilon^{jik} B^k) = \epsilon^{ijk} E^j B^k = (\vec{E} \times \vec{B})^i$$

for the momentum density (which is also known as the Poynting vector).

Solution 3:

$$\mathcal{L} = (\partial_\mu \phi_1^*)(\partial^\mu \phi_1) + (\partial_\mu \phi_2^*)(\partial^\mu \phi_2) - m^2(\phi_1^* \phi_1 + \phi_2^* \phi_2)$$

(a) The equations of motion for $\phi_{1,2}(x)$ and $\phi_{1,2}^*(x)$ are of the Klein-Gordon type:

$$\partial_\mu \frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi_{1,2})} - \frac{\partial \mathcal{L}}{\partial \phi_{1,2}} = 0 \implies \partial_\mu \partial^\mu \phi_{1,2}^* + m^2 \phi_{1,2}^* = 0,$$

and

$$\partial_\mu \frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi_{1,2}^*)} - \frac{\partial \mathcal{L}}{\partial \phi_{1,2}^*} = 0 \implies \partial_\mu \partial^\mu \phi_{1,2} + m^2 \phi_{1,2} = 0.$$

(b) One obtains now four conjugate momentum fields $\Pi = \frac{\partial \mathcal{L}}{\partial \dot{\phi}} = \frac{\partial \mathcal{L}}{\partial(\partial_0 \phi)}$:

$$\Pi_{\phi_{1,2}} = \frac{\partial \mathcal{L}}{\partial(\partial_0 \phi_{1,2})} = \partial^0 \phi_{1,2}^* \equiv \Pi_{1,2} \quad \text{and} \quad \Pi_{\phi_{1,2}^*} = \frac{\partial \mathcal{L}}{\partial(\partial_0 \phi_{1,2}^*)} = \partial^0 \phi_{1,2} \equiv \Pi_{1,2}^*.$$

(c) The Hamiltonian is the integral over the Hamiltonian density, i.e. $H = \int d^3x \mathcal{H}$, where

$$\mathcal{H} = \Pi_1 \partial_0 \phi_1 + \Pi_2 \partial_0 \phi_2 + \Pi_1^* \partial_0 \phi_1^* + \Pi_2^* \partial_0 \phi_2^* - \mathcal{L} = 2\Pi_1^* \Pi_1 + 2\Pi_2^* \Pi_2 - \mathcal{L}.$$

Since the Lagrangian density \mathcal{L} is given by

$$\mathcal{L} = \Pi_1^* \Pi_1 + \Pi_2^* \Pi_2 - (\vec{\nabla} \phi_1) \cdot (\vec{\nabla} \phi_1^*) - (\vec{\nabla} \phi_2) \cdot (\vec{\nabla} \phi_2^*) - m^2(\phi_1^* \phi_1 + \phi_2^* \phi_2)$$

one has

$$\mathcal{H} = \Pi_1^* \Pi_1 + \Pi_2^* \Pi_2 + (\vec{\nabla} \phi_1) \cdot (\vec{\nabla} \phi_1^*) + (\vec{\nabla} \phi_2) \cdot (\vec{\nabla} \phi_2^*) + m^2(\phi_1^* \phi_1 + \phi_2^* \phi_2),$$

consisting of kinetic terms, elastic terms and rest mass terms.

(d) Now we introduce the vector (doublet) notation $\vec{\Phi} \equiv \begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix}$ for the two complex scalar fields and write compactly $\mathcal{L} = (\partial_\mu \vec{\Phi}^\dagger)(\partial^\mu \vec{\Phi}) - m^2 \vec{\Phi}^\dagger \vec{\Phi}$. Under the continuous global $U(1)$ transformation

$$\Phi(x) \rightarrow e^{i\alpha} \Phi(x) \quad \text{and} \quad \Phi^\dagger(x) \rightarrow e^{-i\alpha} \Phi^\dagger(x) \quad (\alpha \in \mathbb{R})$$

the Lagrangian is invariant:

$$\mathcal{L} \rightarrow (\partial_\mu \vec{\Phi}^\dagger) e^{-i\alpha} e^{i\alpha} (\partial^\mu \vec{\Phi}) - m^2 \vec{\Phi}^\dagger e^{-i\alpha} e^{i\alpha} \vec{\Phi} = \mathcal{L}.$$

Keeping a close eye on the order of the vectors, the corresponding Noether current is given by

$$j^\mu = \frac{\partial \mathcal{L}}{\partial(\partial_\mu \vec{\Phi})} \Delta \vec{\Phi} + \Delta \vec{\Phi}^\dagger \frac{\partial \mathcal{L}}{\partial(\partial_\mu \vec{\Phi}^\dagger)} = (\partial^\mu \vec{\Phi}^\dagger) i \vec{\Phi} + (-i \vec{\Phi}^\dagger) (\partial^\mu \vec{\Phi}) = i \left[(\partial^\mu \vec{\Phi}^\dagger) \vec{\Phi} - \vec{\Phi}^\dagger (\partial^\mu \vec{\Phi}) \right]$$

using the first-order variations¹ $\Delta \vec{\Phi} = i \vec{\Phi}$ and $\Delta \vec{\Phi}^\dagger = -i \vec{\Phi}^\dagger$. The corresponding Noether charge is

$$\int d^3x j^0 = i \int d^3x \left[(\partial^0 \vec{\Phi}^\dagger) \vec{\Phi} - \vec{\Phi}^\dagger (\partial^0 \vec{\Phi}) \right] \stackrel{(b)}{=} i \sum_{a=1}^2 \int d^3x [\Pi_a \phi_a - \phi_a^* \Pi_a^*].$$

(e) Under the continuous global $SU(2)$ transformation

$$\Phi(x) \rightarrow e^{i\alpha^k \sigma^k} \Phi(x) \quad \text{and} \quad \Phi^\dagger(x) \rightarrow \Phi^\dagger(x) e^{-i\alpha^k \sigma^k} \quad (\alpha^k \in \mathbb{R} \text{ for } k = 1, 2, 3)$$

the Lagrangian is also invariant, which can be shown by an analogous calculation as above. Here the Pauli matrices $\sigma^{1,2,3}$ are the generators of $SU(2)$. With

$$(\Delta \vec{\Phi})^k = i \sigma^k \vec{\Phi} \quad \text{and} \quad (\Delta \vec{\Phi}^\dagger)^k = -i \vec{\Phi}^\dagger \sigma^k,$$

the corresponding three conserved Noether currents and charges read

$$(j^\mu)^k = (\partial^\mu \vec{\Phi}^\dagger) i \sigma^k \vec{\Phi} - i \vec{\Phi}^\dagger \sigma^k (\partial^\mu \vec{\Phi}),$$

$$Q^k \equiv \int d^3x (j^0)^k = i \int d^3x \left[(\partial^0 \vec{\Phi}^\dagger) \sigma^k \vec{\Phi} - \vec{\Phi}^\dagger \sigma^k (\partial^0 \vec{\Phi}) \right] = i \sum_{a,b=1}^2 \int d^3x [\Pi_a (\sigma^k)_{ab} \phi_b - \phi_a^* (\sigma^k)_{ab} \Pi_b^*].$$

¹By definition one has $\vec{\Phi}' = \vec{\Phi} + \alpha \Delta \vec{\Phi} + \mathcal{O}(\alpha^2)$. With $e^{i\alpha} = 1 + i\alpha + \mathcal{O}(\alpha^2)$ the given expressions for $\Delta \vec{\Phi}$ and $\Delta \vec{\Phi}^\dagger$ follow.