

BROWN UNIVERSITY
PROBLEM SET 5
INSTRUCTOR: SAMUEL S. WATSON
DUE: 13 OCTOBER 2017

Print out these pages, including the additional space at the end, and complete the problems by hand. Then use Gradescope to scan and upload the entire packet by 18:00 on the due date.

Problem 1

- (a) Find the quadratic Maclaurin polynomial for the function $f(x, y) = e^{3x+y}$ by calculating all the relevant partial derivatives.
- (b) Find the quadratic Maclaurin polynomial for e^t and substitute $t = 3x + y$.

Solution

- (a) The relevant partial derivatives of f at the origin are

$$\begin{aligned}f(0, 0) &= 1 \\f_x(0, 0) &= 3 \\f_y(0, 0) &= 1 \\f_{xx}(0, 0) &= 9 \\f_{yy}(0, 0) &= 1 \\f_{xy}(0, 0) &= 3.\end{aligned}$$

So the quadratic Maclaurin polynomial for f is

$$1 + 3x + y + \frac{9}{2}x^2 + 3xy + \frac{1}{2}y^2.$$

- (b) The Maclaurin polynomial for e^t is $1 + t + t^2/2$. If we substitute $t = 3x + y$, we get $1 + 3x + y + (3x + y)^2/2 = 1 + 3x + y + \frac{9}{2}x^2 + 3xy + \frac{1}{2}y^2$, the same as in (a).

Problem 2

Consider the function $f(x, y) = \frac{e^{xy}}{e(1+x^2)}$.

- (a) Use a quadratic Taylor polynomial centered at $(1, 1)$ to approximate $f(0.99, 0.98)$. Compare your answer to Example 4.3.3 in the book.
- (b) Use a quadratic Taylor polynomial centered at $(0, 0)$ to approximate $f(0.99, 0.98)$.
- (c) The following code can be copy-pasted at sagecell.sagemath.org (or [click here](#)) to calculate the degree-50 Maclaurin polynomial of f and evaluate it at $(0.99, 0.98)$.

```
var("x y") # declares x and y to be symbolic variables
f(x,y) = exp(x*y-1)/(1+x^2) # defines f
taylor(f(x,y), (x,0), (y,0), 50).subs(x=0.99, y=0.98).n()
```

How good is this estimate compared to the ones in (a) above and in Example 4.3.3 in the text?

- (d) Repeat (c) but with the function $g(x, y) = \frac{e^{xy}}{e(9+x^2)}$. Does the degree-50 Maclaurin polynomial approximate the value of $g(0.99, 0.98)$ well?

Solution

(a) The Taylor polynomial centered at $(1, 1)$ is $\frac{1}{2}y + \frac{1}{2}(x-1)(y-1) + \frac{1}{4}(y-1)^2$. Substituting $(0.99, 0.98)$ gives exactly 0.4902. The linear approximation yielded 0.49, and the actual value is 0.4901972..., so the quadratic approximation is much more accurate.

(b) We get an approximation of $(x^2 + xy + 1)/e = 0.364237$, which is not nearly as good as the approximations centered at $(1, 1)$.

(c) We get 0.4492..., which is also not very good.

(d) We run the code

```
var('x y') # declares x and y to be symbolic variables
f(x,y) = exp(x*y-1)/(9+x^2) # defines f
taylor(f(x,y), (x,0), (y,0), 50).subs(x=0.99, y=0.98).n(), f(0.99, 0.98)
```

which returns

```
(0.0972575066447653, 0.0972575066447653)
```

Remarkably, this estimate is so good that all the digits displayed by the system are the same.

Here's an amazing fact: the Maclaurin polynomial of the function $1/(1+x^2)$ only approximates it well up to a certain distance from the origin. This critical distance is determined by the least distance from the origin to a *complex* root of the polynomial $1+x^2$ in the denominator. This is one sense in which the study of complex numbers can be relevant even when looking at real-valued functions. This idea is explored further in my course notes for Math 19 [[link](#)].

Problem 3

Consider the function $f(x, y) = \frac{1}{xy}$. Show that f has no maximum or minimum value on the open unit square $S = (0, 1) \times (0, 1)$. In other words, show that for any $(x, y) \in S$, there exists $(x', y') \in S$ with $f(x', y') > f(x, y)$ (and similarly for the minimum).

Solution

Given any $(x, y) \in S$, we can choose x' to be any number strictly between 0 and x , and we can set $y' = y$. Then $f(x', y') = 1/(x'y') > 1/(xy) = f(x, y)$. Thus f has no maximum value on S .

Similarly, if $(x, y) \in S$, then we can choose x' to be strictly between x and 1 and set $y' = y$. Then $f(x', y') = 1/(x'y') < 1/(xy) = f(x, y)$. Thus f has no minimum value on S .

This example does not contradict the extreme value theorem because the set S is not **closed**.

Problem 4

Let D be the closed unit disk $\{(x, y) \in \mathbb{R}^2 : x^2 + y^2 \leq 1\}$. Come up with a function $f : D \rightarrow \mathbb{R}$ with the property that f does not have a maximum value on D . Explain why your function does indeed have this property.

Solution

There are many such functions, but let us set $f(x, y) = 1/(1 + x^2 + y^2)$ for $(x, y) \neq (0, 0)$ and $f(0, 0) = 0$. Then from any $(x, y) \neq (0, 0)$, we can increase the value of f by moving a bit closer to the origin, and $f(0, 0)$ is actually the function's *minimum* value.

This function does not contradict the extreme value theorem because the function is **discontinuous**.

Problem 5

Find the maximum and minimum values of $f(x, y) = x^4 + y^4 - 4xy$ on the rectangle $[0, 3] \times [0, 2]$.

Solution

The system of equations $\partial_x f = \partial_y f = 0$ tells us that $y = x^3$ and $x = y^3$. Substituting, we find $x = x^9$, which can be rearranged and factored (by repeated applications of difference of squares) to get

$$x(x-1)(x+1)(x^2+1)(x^4+1) = 0.$$

The only value of x in the interior of the rectangle which satisfies this equation is $x = 1$. Substituting into $y = x^3$, we find the critical point $(1, 1)$ in the interior of the rectangle. The value of the function this point is -2 .

Along the bottom edge of the rectangle, the value of the function at $(x, 0)$ is x^4 which ranges monotonically from 0 to 81. Along the top edge, we have $f(x, 2) = x^4 - 8x + 16$. This has a critical point at $x = \sqrt[3]{2}$.

Along the left edge, the function is equal to y^4 , which has no interior critical points. And finally, along the right edge, we have $f(3, y) = y^4 - 12y + 81$, which has a critical point at $\sqrt[3]{3}$.

So altogether, checking critical points and corners gives

(x, y)	$f(x, y)$
$(1, 1)$	-2
$(0, 0)$	0
$(3, 0)$	81
$(0, 2)$	16
$(3, 2)$	73
$(3, \sqrt[3]{3})$	$81 - 9\sqrt[3]{3} \approx 68.02$
$(\sqrt[3]{2}, 2)$	$16 - 6\sqrt[3]{2} \approx 8.44$

So the absolute maximum is $\boxed{81}$ and the absolute minimum is $\boxed{-2}$.

