

**A Novel Weighted Mean Approach to Estimate the Distribution Function  
of Potential Outcomes in Observational Studies**

Asymptotic Analysis

**Ioan Scheffel**

A thesis presented for the degree of  
Master of Science Mathematics

supervised by PD Dr. Jürgen Dippon  
Institute for Stochastics and Applications  
Faculty 8: Mathematics and Physics  
University of Stuttgart  
submitted at April 17, 2023

### **Eigenständigkeitserklärung**

Ich erkläre mit meiner Unterschrift, dass ich diese Arbeit selbstständig verfasst habe und keine anderen als die angegebenen Quellen benutzt habe. Alle Stellen dieser Arbeit, die dem Wortlaut, dem Sinn oder der Argumentation nach anderen Werken entnommen sind (einschließlich des World Wide Web und anderer elektronischer Text- und Datensammlungen), habe ich unter Angabe der Quellen vollständig kenntlich gemacht.

## **Abstract**

english

In this thesis I extend the balancing weights framework of [WZ19] to estimate the distribution function of potential outcomes. I also suggest to balance basis functions of non-parametric partitioning estimates. This greatly simplifies the proofs and allows for rigorous mathematical treatment of the method. The asymptotic analysis shows convergence of the error to a Gaussian process. My findings allow to apply the functional delta method to plug-in estimators. This makes classical statistical methods such as quantile estimation, hypothesis testing or survival analysis accessible to causal inference in observational studies. While the theoretical results are promising, this novel approach waits for testing in practice.

## **Abstract**

german

In dieser Abschlussarbeit erweitere ich den Balacing Weights Ansatz von [WZ19] um die Verteilungsfunktion von Potential Outcomes zu schätzen. Ich schlage vor als Basis Funktionen die Basis nicht-parametrischer Partitioning Estimates zu wählen. Diese Wahl vereinfacht die Beweise deutlich und erlaubt es, die Methode mathematisch rigoros zu behandeln. Die asymptotische Analyse zeigt Konvergenz des Fehlers gegen einen Gaußschen Prozess. Meine Ergebnisse erlauben es direkt die Funktionale Delta-Methode auf Substitutions-Schätzer anzuwenden. Dies erschließt der kausalen Inferenz in Beobachtungsstudien klassische Methoden der Statistik, wie Quantil-Schätzung, Hypothesen-Tests oder Überlebenszeit-Analyse. Die theoretischen Eigenschaften der von mir vorgeschlagenen Erweiterung sind vielversprechend - der Praxis-Test steht allerdings noch aus.



# Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>The Optimization Problem behind the Weights</b>	<b>7</b>
2.1	Introduction . . . . .	7
2.2	Objective Function . . . . .	10
2.3	Dual Problem . . . . .	12
<b>3</b>	<b>Constructing the Weights Process</b>	<b>19</b>
3.1	Argmax Measurability Theorem . . . . .	19
3.2	Measurable Dual Solution . . . . .	20
3.3	Basis Functions . . . . .	21
3.4	Weights Process . . . . .	26
<b>4</b>	<b>Consistency of the Weights Process</b>	<b>29</b>
4.1	Propensity Score Function . . . . .	29
4.2	Consistency of the Dual Solution . . . . .	29
4.3	Main Result . . . . .	36
<b>5</b>	<b>Convergence of the Weighted Mean</b>	<b>39</b>
5.1	Tools . . . . .	39
5.1.1	Empirical Processes - Definition . . . . .	39
5.1.2	Bracketing Numbers and Integral . . . . .	40
5.1.3	Maximal Inequality . . . . .	47
5.1.4	Donsker's Theorem . . . . .	48
5.1.5	Propensity Score Weights . . . . .	49
5.2	Main Result . . . . .	50
5.3	Error Decomposition . . . . .	50
5.4	Analysis of the Error Terms . . . . .	53
5.4.1	Analysis of $R_1$ . . . . .	53
5.4.2	Analysis of $R_2$ . . . . .	54
5.4.3	Analysis of $R_3$ . . . . .	55

5.4.4	Analysis of $R_4$ . . . . .	56
5.4.5	Proof of Theorem 5.3 . . . . .	58
<b>6</b>	<b>Discussion and Outlook</b>	<b>59</b>
6.1	Discussion . . . . .	59
6.1.1	Summary of Assumptions . . . . .	59
6.2	Outlook . . . . .	59
6.2.1	Matching . . . . .	59
6.2.2	Application of the Functional Delta Method . . . . .	60
6.2.3	Bootstrapping . . . . .	60
6.2.4	Non-binary Treatment . . . . .	61
6.2.5	Different Basis Functions . . . . .	63
<b>7</b>	<b>Convex Analysis</b>	<b>65</b>
7.1	A Convex Analysis Primer . . . . .	65
7.2	Duality of Optimal Solutions . . . . .	72
	<b>References</b>	<b>79</b>
	<b>Index</b>	<b>83</b>

# 1 Introduction

How does action change an outcome? How should I guide my actions towards a better outcome? The first question is about causality, the second about ethics.

How do causality and ethics reflect on statistics? If you have not spent much time thinking about study design, this is a good way to start: As an analyst, ask yourself “Who acted? Who assigned treatment?” As a researcher – plan your study accurately. You can ask yourself “How do we act? How do we assign treatment? Can we act?”

Let’s say, you gather a sample from a study population, assign treatment (but forget how you did it). Some units get the drug, others don’t. Then the statistical analysis shows a strong correlation of treatment and outcome. You hurry to your supervisor. “How was treatment assigned”, asks she. “I forgot”, says you. “How do you know your analysis is correct then?” You show her the data and together you find out, that all units that received treatment were significantly taller than the rest of the sample. After all, is the drug or the height responsible for the change in outcome? You realise that the data is worthless for answering this question. But you are lucky: It is just grass and fertiliser you were studying.

You get a second chance. A new medication needs testing before it enters the market. A company shall recruit participants, but the board requires you to write an outline for the study. You carefully explain steps to minimize risks for participants. You include plans to meet other requirements of human research. Then you have to decide how to assign treatment. No hand waving this time. You talk to your supervisor. “Last time, too many tall grass blades received fertiliser. The distribution of treatment was not really random...” You decide to determine treatment status by the flip of a fair coin. You call the procedure ‘randomization’.

Would you smoke if a coin tells you to? If you say yes - you likely smoke anyway. The point is that forcing someone to smoke is unethical. But so is not studying the risks of smoking.

A professor is curious if the smoking habits of his students affect their grades. He observes the smoking area through his field glasses. His assistant gets to know his plans. He warns him. “Many students attend parties the night before exams. Maybe they are also more likely to smoke.” “I shall see this for myself...” says the professor. He puts



## 1 Introduction

away the field glasses. After a while, he visits the local club. He talks to a few of his students. Some smoke, some don't. The chats are enjoyable. He thinks: "Some of my best students celebrate before the exams."

I hope, by now it's clear that we should focus on treatment assignment. The propensity score [RR83], that is, the probability of treatment given (observed) individual characteristics, helps with that.

**Theorem.** [RR83, Theorem 1] *Observed individual characteristics are independent of treatment assignment given the propensity score.*

In the second example, where you flip a fair coin to assign treatment, the propensity score is  $1/2$ , despite variation across individual characteristics. The coin ignores everything. What is the propensity score in the other examples? I admit, I don't know. It varies, but we can see trends. In the first example, tall grass blades had a large propensity score. In the third example, the assistant thinks that students attending parties have a larger propensity score. This is not true, after all, but somehow the best students have a large propensity to celebrate before exams.

The propensity score is a simple concept that works well with potential outcomes. They are potential, because they exist (or we assume they exist) independent of our observation. They live in parallel universes. If we have a binary treatment, that is, you either treat or don't, there are two potential outcomes. One under treatment and one under no treatment. Ideally we would like to compare (for one unit) those two potential outcomes. But that is impossible. Instead people keep asking: "Had it been better if (20 years ago) I made a different decision?" You know what happened but don't know what would have happened. On a high-level: If you act, you can't observe at the same time the effect of no action. Thus, one of the potential outcomes always remains potential. Of course there are tricks. You can wait for the effect of an action to vanish and then observe the outcome (under similar conditions) again. This works well when the effect of an action is short term.

If the propensity score is known we actually observe one of the potential outcomes. This is because treatment assignment carries no more information [RR83, Theorem 1]. But we saw that assignment often carries more information, especially if the assignment mechanism is unknown. This is typical for observational studies. Somehow grass blades that received fertiliser were also taller. Or students attending parties before the exams had better results. It is not clear, if the effect on the outcome stems from observed or unobserved individual characteristics or the received treatment. Then we observe neither of the two potential outcomes, but a biased version. Why then bother?

Since the introduction of the propensity score in 1983 [RR83], statisticians developed

different ideas how to incorporate it in their analysis. There are two important branches of application - matching and weighting. In matching the idea is to pair two or more units with different treatment status but similar propensity score and compare their outcome. The assumption is that the propensity score eliminates imbalances and makes the paired units comparable. In weighting - the method we will focus on - the idea is to re-weight the population, ideally with the inverse propensity score, that is, 1 divided by the propensity score. Both methods share the goal to minimize imbalances in the population and make the two groups, that is, treatment and control (no treatment) group more comparable. Let's introduce some notation to be more precise.

Let  $T \in \{0, 1\}$  be the **indicator of treatment**. Let  $X \in \mathcal{X}$  be a vector with individual characteristics. We call this the **covariate vector**.

Furthermore, let  $(Y(0), Y(1))$  be the **potential outcomes**, that is,  $Y(0)$  is the potential outcome under no treatment and  $Y(1)$  the potential outcome under treatment. All the quantities we introduce are random variables.

We define the propensity score  $\pi$  with individual characteristics  $x \in \mathcal{X}$  to be

$$\pi(x) := \mathbf{P}[T = 1 | X = x]$$

We observe

$$\text{either } Y(0) | T = 0 \quad \text{or} \quad Y(1) | T = 1.$$

We show in Lemma 5.10, that if treatment assignment is **strongly ignorable** [RR83, (1.3)]

$$(Y(0), Y(1)) \perp T | X \quad \text{and} \quad 0 < \pi(X) < 1,$$

that is, potential outcomes are independent of treatment given covariates and every possible set of characteristic has a chance to receive treatment, we get

$$\mathbf{E} \left[ \frac{T}{\pi(X)} Y(T) \right] = \mathbf{E} [Y(1)]. \quad (1.1)$$

That is, by weighting the observed outcome under treatment with the inverse propensity score we recover (in expectation) the potential outcome under treatment. This is relevant, because  $Y(t) | T = t$  does not have the same distribution as  $Y(t)$  for  $t \in \{0, 1\}$ .

In observational studies - independent of which method is applied, that is, matching, weighting or other methods - the propensity score is unknown. This is because we can't assign treatment. We only observe the treated or untreated. Who assigned treatment is a philosophical question - at least to some extent.

## 1 Introduction

It used to be very popular to use estimates of the propensity score - either for matching or to create weights or for some other purpose. In weighting, the hope is to recover (1.1) from the estimate. In practice, however, estimating the propensity score is a difficult task. Researchers often compare estimates from different models and check for covariate balance. This is not surprising, because that's what the weights are designed for - minimizing imbalances in the population. The poor performance of classical propensity score estimates, such as logistic regression - and the insight that the task is to eliminate imbalances - led to the development of methods, such as the Covariate Balancing Propensity Score [IR14] that tries to estimate the propensity score and balance covariates simultaneously (the name is indicative).

Recently, new balancing frameworks were developed that do not rely on estimates of the propensity score [Hai12, Zub15]. They generate weights with a constrained convex optimization problem that explicitly bounds imbalances by the constraints. The constraints responsible are of the form

$$\sum_{i=1}^N T_i \cdot w_i \cdot B_k(X_i) = \sum_{i=1}^N B_k(X_i) \quad \text{for all } k, \quad (1.2)$$

where  $B_k$  are basis functions of the covariates. The aim is to balance the weighted group (here the treatment group) against the whole sample. The basis functions can be moments of the covariates - a natural aim. But it is a non-trivial question, which basis to choose in practice to obtain best performance. How strictly to enforce covariate balance is another question. It is relevant, because very strict assumptions can render the problem infeasible, whereas loosening can result in bias of the estimator. In [Hai12] the authors choose the (known) moments of the covariate as basis functions and enforce strict balance, that is, (1.2). In [WZ19] they consider the regression basis of sieve estimators [New97], where the number of basis functions grows with the sample size. Also they loosen the strict constraints on the covariate balance as to vanish only for  $N \rightarrow \infty$ , that is,

$$\left| \frac{1}{N} \left( \sum_{i=1}^n w_i B_k(X_i) - \sum_{i=1}^N B_k(X_i) \right) \right| \leq \delta_k \quad \text{for all } k,$$

with  $\delta_k > 0$  and  $\delta_k \rightarrow 0$  for  $N \rightarrow \infty$ .

What attracted my attention was that the paper [WZ19] also contains theoretical analysis. The authors reveal a surprising connection to propensity score estimation. They show that with the regression basis of sieve estimators [New97], their method (implicitly) models the inverse propensity score. They use this to obtain asymptotic normality of a weighted mean estimate of the expectation of potential outcomes.

One novelty introduced in this thesis is to balance basis functions of partitioning estimates [GKKW02, §4]. I show that this simplifies the proofs of [WZ19]. Furthermore, I extend the framework to estimate the distribution function of potential outcomes.

I show, that (under mild assumptions) with the regression basis of partitioning estimates, the weighted mean is asymptotically well behaved in estimating distribution functions. By the functional delta method [vdV00, §20] results of this thesis immediately open access to a large class of plug-in estimators. Therefore, with this thesis I contribute to one of the main purposes in causal inference – reinforcing classical methods of statistical analysis for use in observational studies.



## 2 The Optimization Problem behind the Weights

There are different ways to generate weights for covariate balance. We discussed this in the introduction. Now, we introduce the balancing weights framework of [WZ19]. It is a (generic) convex optimization problem that enforces covariate balance by constraints on the search space. Similar to classical propensity score estimates, it only extracts from the data information about treatment status and individual characteristics. It ignores the outcome. This gives the additional option to balance covariates before observing outcomes.

The primary optimization task is to minimize an objective function over a predefined search space. From a practical point of view, the objective function instils additional goodness in the weights, for example, low sample variance [Zub15, Introduction]. More important, however, are the constraints that enforce covariate balance. Both objective function and design of the constraints distinguish the method.

From a mathematical point of view, we have slightly different requirements. The proofs should be as clear and short as possible. The mathematical objects involved should help with that (or at least not prevent it). Therefore, we focus on theoretical properties of the method. As a by-product, we create new ideas that wait for testing in practice.

The notion of what to balance is defined by the basis functions of the covariates. It is common that they form a regression basis. We shall use this flexibility to introduce basis functions with random design to this framework. This makes the proofs easier.

### 2.1 Introduction

Let  $(T_1, X_1), \dots, (T_N, X_N)$  be independent and identically-distributed copies of  $T$  and  $X$  (see Introduction page). We gather them in the (random) data set

$$D_N := \{ (T_i, X_i) : i \in \{1, \dots, N\} \} .$$

## 2 The Optimization Problem behind the Weights

Furthermore, let

$$n := \# \{i \in \{1, \dots, N\} : T_i = 1\}$$

be the number of treated units. This is a random variable. We assume the order  $T_i = 1$  for all  $i \leq n$ . For a

$$\text{(proper) convex function} \quad \varphi : \mathbb{R} \rightarrow \mathbb{R} \cup \{+\infty\},$$

a vector of  $N$  basis functions of the covariates

$$B := [B_1, \dots, B_N]^\top \quad \text{with} \quad B_k : \mathbb{R}^d \rightarrow \mathbb{R} \quad \text{for all } k \in \{1, \dots, N\},$$

and a (random) constraints vector

$$\delta := [\delta_1, \dots, \delta_N]^\top \quad \text{with} \quad \delta_k : (\Omega, \sigma(D_N), \mathbf{P}) \rightarrow \mathbb{R} \quad \text{for all } k \in \{1, \dots, N\},$$

we consider the (random) convex optimization problem

**Problem 1.**

$$\begin{aligned}
& \underset{w_1, \dots, w_n \in \mathbb{R}}{\text{minimize}} && \sum_{i=1}^n \varphi(w_i) \\
& \text{subject to} && w_i \geq 0 \quad \text{for all } i \in \{1, \dots, n\}, \\
& && \frac{1}{N} \sum_{i=1}^n w_i = 1 \\
& && \left| \frac{1}{N} \left( \sum_{i=1}^n w_i \cdot B_k(X_i) - \sum_{i=1}^N B_k(X_i) \right) \right| \leq \delta_k \quad \text{for all } k \in \{1, \dots, N\}.
\end{aligned}$$

What is random in Problem 1? First, the dimension of the search space ( $w \in \mathbb{R}^n$ ) depends on the random variable  $n$ . Thus, we only compute weights for the treated units (the ones with  $T_i = 1$ ). Next, consider the **objective function**

$$w \mapsto \sum_{i=1}^n \varphi(w_i).$$

The number of summands is random (again  $n$ ). Note, that sometimes we use the equivalent notation

$$w \mapsto \sum_{i=1}^N T_i \cdot \varphi(w_i),$$

where we set the weights of the untreated (the ones with  $T_i = 0$ ) to some arbitrary value in the domain of  $\varphi$ . Let's consider the **constraints**. There is no randomness in the first two constraints.

$$w_i \geq 0 \quad \text{for all } i \in \{1, \dots, n\} \quad \text{and} \quad \frac{1}{N} \sum_{i=1}^n w_i = 1.$$

They only make sure, that the weights (divided by  $N$ ) form a convex combination. If, for example, the outcome space  $\mathcal{Y}$  is convex we make sure that a weighted-mean-estimate of  $\mathbf{E}[Y(1)]$  satisfies

$$\hat{Y}(1) := \frac{1}{N} \sum_{i=1}^n w_i \cdot Y_i \in \mathcal{Y}$$

or that a weighted-mean-estimate of the distribution function of  $Y(1)$  satisfies

$$\hat{F}_{Y(1)} := \frac{1}{N} \sum_{i=1}^n w_i \cdot \mathbf{1}\{Y_i \leq z\} \in [0, 1].$$



## 2 The Optimization Problem behind the Weights

We talked about the covariate balancing constraint in the introduction (we shall call them the **box constraints**, because of the absolute value).

$$\left| \frac{1}{N} \left( \sum_{i=1}^n w_i \cdot B_k(X_i) - \sum_{i=1}^N B_k(X_i) \right) \right| \leq \delta_k \quad \text{for all } k \in \{1, \dots, N\}.$$

They are crucial - we shall discuss their implications as the analysis unfolds. For now, note that the number of summands in

$$\sum_{i=1}^n w_i \cdot B_k(X_i)$$

is random again, and sometimes we switch to

$$\sum_{i=1}^N T_i \cdot w_i \cdot B_k(X_i).$$

In section?, we shall specify the vector of basis functions  $B$ . Instead of sieve estimators as in [WZ19], where the number of basis functions grows slower than  $N$  to  $\infty$  and the basis functions have fixed design, we shall choose the basis of partitioning estimates as in [GKKW02, §4], which depends on the whole data set  $D_N$  and therefore has random design. We shall see that this choice greatly simplifies the consistency proofs. Finally, note that [WZ19, Algorithm 1 on page 11] is a (random) algorithm to specify  $\delta$  based on  $D_N$ .

## 2.2 Objective Function

The formulation of Problem 1 allows for great flexibility. To obtain clear and short proofs, however, we have to restrict it.

**Definition 2.1.** We define  $\varphi$  in Problem 1 by

$$\varphi : \mathbb{R} \rightarrow [0, \infty), \quad x \mapsto (x - 1)^2.$$

**Remark.** If we plug this choice in Problem 1, we observe

$$\sum_{i=1}^n \varphi(w_i) = \sum_{i=1}^N T_i (T_i \cdot w_i - 1)^2 = \sum_{i=1}^N T_i \left( T_i \cdot w_i - \frac{1}{N} \sum_{i=1}^N T_i \cdot w_i \right)^2.$$

Thus Problem 1 minimizes the sample variance of the weights  $(T_i \cdot w_i)$ . This is in line with the objective function in [Zub15].  $\diamond$

Next, we derive theoretical properties of  $\varphi$  that we will use in the subsequent analysis.

**Lemma 2.1.** *The function  $\varphi$  of Definition 2.1 satisfies*

- (i)  $\varphi$  is strictly convex and continuously differentiable on  $\mathbb{R}$ , with derivative  $\varphi'$
- (ii) The inverse of the derivative  $(\varphi')^{-1}$  exists and is continuously differentiable
- (iii) Both  $\varphi'$  and  $(\varphi')^{-1}$  are uniformly continuous

**Proof.** The proof is easy. We omit the details. □

The next lemma prepares a link to the assumptions of Theorem 7.4.

**Lemma 2.2.** *The convex conjugate of  $\varphi$  (see (7.12)) is*

$$\varphi^*: \mathbb{R} \rightarrow \mathbb{R}, \quad x^* \mapsto x^* \cdot (\varphi')^{-1}(x^*) - \varphi\left((\varphi')^{-1}(x^*)\right).$$

*Furthermore,  $\varphi^*$  is strictly convex and continuously differentiable on  $\mathbb{R}$ .*

**Proof.** We define

$$\phi: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}, \quad (x, x^*) \mapsto x \cdot x^* - \varphi(x).$$

Let  $x^* \in \mathbb{R}$ . By Lemma 2.1.(i),  $\varphi$  is continuously differentiable on  $\mathbb{R}$  with derivative  $\varphi'$ . The same holds for  $\phi(\cdot, x^*)$  with derivative satisfying

$$\frac{\partial}{\partial x} \phi(x, x^*) = x^* - \varphi'(x) \quad \text{for all } x \in \mathbb{R}.$$

By Lemma 2.1.(ii), it holds that

$$z := (\varphi')^{-1}(x^*)$$

is an extreme point of  $\phi(\cdot, x^*)$ . Since  $\varphi$  is strictly convex by Lemma 2.1.(i),  $\phi(\cdot, x^*)$  is strictly concave. Thus,  $z$  is the unique maximum point of  $\phi(\cdot, x^*)$  on  $\mathbb{R}$ . Thus

$$\begin{aligned} \varphi^*(x^*) &= \sup_{x \in \mathbb{R}} x \cdot x^* - \varphi(x) = \sup_{x \in \mathbb{R}} \phi(x, x^*) \\ &= \phi(z, x^*) \\ &= x^* \cdot (\varphi')^{-1}(x^*) - \varphi\left((\varphi')^{-1}(x^*)\right) \quad \text{for all } x^* \in \mathbb{R}. \end{aligned}$$

## 2 The Optimization Problem behind the Weights

Now we proof the second statement. Since  $(\varphi')^{-1}$  is continuously differentiable by Lemma 2.1.(ii), it holds

$$\begin{aligned} \frac{\partial}{\partial x^*} \varphi^*(x^*) &= (\varphi')^{-1}(x^*) + x^* \cdot \frac{\partial}{\partial x^*} (\varphi')^{-1}(x^*) - \varphi' \left( (\varphi')^{-1}(x^*) \right) \cdot \frac{\partial}{\partial x^*} (\varphi')^{-1}(x^*) \\ &= (\varphi')^{-1}(x^*) + x^* \cdot \frac{\partial}{\partial x^*} (\varphi')^{-1}(x^*) - x^* \cdot \frac{\partial}{\partial x^*} (\varphi')^{-1}(x^*) \\ &= (\varphi')^{-1}(x^*) \quad \text{for all } x^* \in \mathbb{R}. \end{aligned} \tag{2.1}$$

Since  $\varphi$  is strictly convex and continuously differentiable,  $\varphi'$  is continuous and strictly non-decreasing. Thus  $(\varphi')^{-1}$  is continuous and strictly non-decreasing. It follows from (2.1) that  $\varphi^*$  is strictly convex and continuously differentiable.  $\square$

With Lemma 2.2 we are ready to complete the link.

**Lemma 2.3.** *The function*

$$\Phi : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}, \quad [w_1, \dots, w_n]^\top \mapsto \sum_{i=1}^n \varphi(w_i),$$

*satisfies Assumption 2.*

**Proof.** By Example 7.1 the convex conjugate of  $\Phi$  is

$$\Phi^* : \mathbb{R}^n \rightarrow \mathbb{R}, \quad [\lambda_1, \dots, \lambda_n]^\top \mapsto \sum_{i=1}^n \varphi^*(\lambda_i),$$

where  $\varphi^*$  is the convex conjugate of  $\varphi$ . By Lemma 2.1,  $\varphi$  is strictly convex. Thus,  $\Phi$  is strictly convex. By Lemma 2.2,  $\varphi^*$  continuously differentiable on  $\mathbb{R}$ . Thus,  $\Phi$  is continuously differentiable on  $\mathbb{R}^n$ . It follows the statement of Assumption 2 for  $\Phi$ .  $\square$

**Takeaways** The choice of Definition 2.1 introduces the sample variance to Problem 1. It has good practical and theoretical properties. Among the theoretical are strict convexity that allows linking Problem 1 to the theory of convex analysis.

## 2.3 Dual Problem

In the previous section we have expounded our choice of  $\varphi$  - and with it the objective function of Problem 1. Now, we want to apply Theorem 7.4 to Problem 1. To this end, we provide its proper formulation.

**Lemma 2.4.** *A matrix formulation of Problem 1 is*

$$\begin{aligned} & \underset{w \in \mathbb{R}^n}{\text{minimize}} && \Phi(w) \\ & \text{subject to} && \mathbf{U}w \geq d, \\ & && \mathbf{A}w = a, \end{aligned} \tag{2.2}$$

*with objective function*

$$\Phi : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}, \quad [w_1, \dots, w_n]^\top \mapsto \sum_{i=1}^n \varphi(w_i),$$

*inequality matrix and vector*

$$\mathbf{U} := \begin{bmatrix} \mathbf{I}_n \\ \pm \mathbf{B}(\mathbf{X}) \end{bmatrix} \in \mathbb{R}^{(n+2N) \times n} \quad d := \begin{bmatrix} 0_n \\ -N \cdot \delta \pm \sum_{i=1}^N B(X_i) \end{bmatrix} \in \mathbb{R}^{n+2N},$$

*and equality matrix and vector*

$$\mathbf{A} := \mathbf{1}_n^\top \in \mathbb{R}^{1 \times n} \quad a := N \in \mathbb{N}.$$

**Proof.** Recall that the box constraints of Problem 1 are

$$\left| \frac{1}{N} \left( \sum_{i=1}^n w_i B_k(X_i) - \sum_{i=1}^N B_k(X_i) \right) \right| \leq \delta_k \quad \text{for all } k \in \{1, \dots, N\}.$$

Put differently, it holds both

$$-\sum_{i=1}^n w_i B_k(X_i) \geq -N\delta_k - \sum_{i=1}^N B_k(X_i) \quad \text{and} \quad \sum_{i=1}^n w_i B_k(X_i) \geq -N\delta_k + \sum_{i=1}^N B_k(X_i)$$

for all  $k \in \{1, \dots, N\}$ . In matrix notation this is

$$\pm \mathbf{B}(\mathbf{X})w \geq [d_{n+1}, \dots, d_{n+2N}]^\top.$$

Proving the rest of the statements is straightforward. We omit the details.  $\square$

**Remark.** The inequality constraints of Lemma 2.4 differ from its counterpart [WZ19, Proof of Lemma 1]. We don't transform the variable  $w$ , but shift to  $d$  what prevents us from keeping  $w$ . Note, that the choice of [WZ19, Proof of Lemma 1] leads to a mistake on page 21. The mistake is most obvious in the second display, where the first implication follows from dividing by 0. I discussed this with the authors and proposed a version of Lemma 2.4 to solve the problem. I think it's best not to transform variables,

## 2 The Optimization Problem behind the Weights

because the mistake comes from (wrongly) calculating the convex conjugate of the (more complicated) transformed version of the objective function. The subsequent analysis even simplifies with my version.

I was surprised to find the (exact) same mistake in the earlier paper [CYZ16, page 35 second display]. There is no reference in [WZ19, Proof of Lemma 1] to [CYZ16]. Yet the formulation and the mistake are the same. Did the authors of [WZ19] (inadvertently?) plagiarize the mathematical analysis of [CYZ16] ?  $\diamond$

In the next lemma we apply Theorem 7.4 to Problem 1.

**Lemma 2.5.** *Consider the optimization problem*

$$\begin{aligned} \underset{\substack{\rho, \lambda^+, \lambda^- \geq 0 \\ \lambda_0 \in \mathbb{R}}}{\text{maximize}} \quad & - \sum_{i=1}^n \varphi^*(\rho_i + \lambda_0 + \langle B(X_i), \lambda^+ - \lambda^- \rangle) \\ & + \sum_{i=1}^N (\lambda_0 + \langle B(X_i), \lambda^+ - \lambda^- \rangle) - \langle \delta, \lambda^+ + \lambda^- \rangle. \end{aligned} \quad (2.3)$$

*If there exists the optimal solution  $(\rho^\dagger, \lambda_0^\dagger, \lambda^{+, \dagger}, \lambda^{-, \dagger})$  then the unique optimal solutions to Problem 1 are*

$$w_i^\dagger := (\varphi')^{-1} \left( \rho_i^\dagger + \lambda_0^\dagger + \langle B(X_i), \lambda^{+, \dagger} - \lambda^{-, \dagger} \rangle \right) \quad \text{for all } i \in \{1, \dots, n\}.$$

**Proof.** First, note that by the strict convexity of  $\varphi^*$  (see Lemma 2.2), a solution to Problem (2.3) is unique (if it exists). By Lemma 2.4, Problem 1 has the form required in Theorem 7.4. By Lemma 2.3, the objective function  $\Phi$  of Problem 1 satisfies Assumption 2. Thus we can apply Theorem 7.4 to Problem 1. Calculations yield the result.  $\square$

With the next theorem we merge  $\lambda^+, \lambda^- \geq 0$  to  $\lambda = \lambda^+ - \lambda^- \in \mathbb{R}$ . Let  $[x]^+ := 0 \wedge x$  be the positive part of  $x \in \mathbb{R}$ .

**Theorem 2.1.** Consider the optimization problem

$$\begin{aligned}
& \underset{\substack{\rho \in \mathbb{R}^N \\ \lambda_0 \in \mathbb{R} \\ \lambda \in \mathbb{R}^N}}{\text{minimize}} & \frac{1}{N} \sum_{i=1}^N \left[ T_i \cdot \varphi^*(\rho_i + \lambda_0 + \langle B(X_i), \lambda \rangle) - \lambda_0 - \langle B(X_i), \lambda \rangle \right] + \langle \delta, |\lambda| \rangle, \\
& \text{subject to} & \rho_i \geq 0 \quad \text{for all } i \leq n \\
& \text{and} & \rho_i = [\varphi^{-1}(0) - (\lambda_0 + \langle B(X_i), \lambda \rangle)]^+ \quad \text{for all } i > n.
\end{aligned} \tag{2.4}$$

If there exists the optimal solution  $(\rho^\dagger, \lambda_0^\dagger, \lambda^\dagger)$  then the unique optimal solutions to Problem 1 are

$$w_i^\dagger := (\varphi')^{-1} \left( \rho_i^\dagger + \lambda_0^\dagger + \langle B(X_i), \lambda^\dagger \rangle \right) \quad \text{for all } i \in \{1, \dots, n\}.$$

**Proof.** Assume that  $(\rho^\dagger, \lambda_0^\dagger, \lambda^{+, \dagger}, \lambda^{-, \dagger})$  is an optimal solution to Problem 2.3. We write

$$\begin{aligned}
G(\rho, \lambda_0, \lambda^+, \lambda^-) &:= - \sum_{i=1}^n \varphi^*(\rho_i + \lambda_0 + \langle B(X_i), \lambda^+ - \lambda^- \rangle) \\
&\quad + \sum_{i=1}^N (\lambda_0 + \langle B(X_i), \lambda^+ - \lambda^- \rangle) - \langle \delta, \lambda^+ + \lambda^- \rangle.
\end{aligned}$$

To eliminate the remaining constraints, we paraphrase [WZ19, pages 19-20]. We show for all  $i \in \{1, \dots, N\}$

$$\begin{aligned}
& \text{either} & \lambda_i^{+, \dagger} > 0 \\
& \text{or} & \lambda_i^{-, \dagger} > 0.
\end{aligned} \tag{2.5}$$

Assume towards a contradiction that

$$\text{there exists } i \in \{1, \dots, N\} \text{ such that } \lambda_i^{+, \dagger} > 0 \quad \text{and} \quad \lambda_i^{-, \dagger} > 0. \tag{2.6}$$

Consider

$$\tilde{\lambda}^{+, \dagger} := \left[ \lambda_1^{+, \dagger}, \dots, \lambda_i^{+, \dagger} - (\lambda_i^{+, \dagger} \wedge \lambda_i^{-, \dagger}), \dots, \lambda_N^{+, \dagger} \right]^\top$$

and

$$\tilde{\lambda}^{-, \dagger} := \left[ \lambda_1^{-, \dagger}, \dots, \lambda_i^{-, \dagger} - (\lambda_i^{+, \dagger} \wedge \lambda_i^{-, \dagger}), \dots, \lambda_N^{-, \dagger} \right]^\top.$$

Since

$$\lambda_i^{\pm, \dagger} - (\lambda_i^{+, \dagger} \wedge \lambda_i^{-, \dagger}) \geq 0,$$

## 2 The Optimization Problem behind the Weights

the perturbed vectors  $\tilde{\lambda}^{\pm, \dagger}$  are in the domain of the optimization problem. By Assumption (2.6) and  $\delta > 0$  it follows

$$G\left(\rho^\dagger, \lambda_0^\dagger, \tilde{\lambda}^{+, \dagger}, \tilde{\lambda}^{-, \dagger}\right) - G\left(\rho^\dagger, \lambda_0^\dagger, \lambda^{+, \dagger}, \lambda^{-, \dagger}\right) = 2 \cdot \delta_i \cdot (\lambda_i^{+, \dagger} \wedge \lambda_i^{-, \dagger}) > 0,$$

which contradicts the optimality of  $(\rho^\dagger, \lambda^{+, \dagger}, \lambda^{-, \dagger}, \lambda_0^\dagger)$  (it is supposed to be a maximum in the domain of the optimization problem). It follows (2.5). But then  $\lambda_i^{\pm, \dagger} \geq 0$  collapses to  $\lambda_i^\dagger \in \mathbb{R}$  for all  $i \in \{0, \dots, N\}$ , that is, we set

$$\lambda_i^\dagger = \lambda_i^{+, \dagger} - \lambda_i^{-, \dagger} \quad \text{and} \quad |\lambda_i^\dagger| = \lambda_i^{+, \dagger} + \lambda_i^{-, \dagger}.$$

Thus, we can extend the domain of Problem 2.3 to  $\lambda \in \mathbb{R}^N$  and update the objective function in the following way (without changing the optimal solution).

$$\begin{aligned} G(\rho, \lambda_0, \lambda) &:= - \sum_{i=1}^n \varphi^*(\rho_i + \lambda_0 + \langle B(X_i), \lambda \rangle) \\ &\quad + \sum_{i=1}^N (\lambda_0 + \langle B(X_i), \lambda \rangle) - \langle \delta, |\lambda| \rangle. \end{aligned}$$

Multiplying  $G$  with  $-1/N$  doesn't change the solution either (if we search instead for a minimum). To finish the proof, we choose the notation with  $T_i$  instead of  $n$ . This extends the domain of  $\rho$  to  $\mathbb{R}_{\geq 0}^N$ , but the new  $\rho_i$  are not effective because of  $T_i = 0$  for all  $i > n$ . Thus we may set them to an arbitrary value.  $\square$

**Remark.** This is the final form of the dual of Problem 1. Since the constraints in the dual problem are elementary, a result such as Lemma 4.1 keeps the initiative going. The dual variables  $(\rho, \lambda_0, \lambda)$  are connected to the constraints of Problem 1, that is,  $\rho \in \mathbb{R}_{\geq 0}^N$  to  $T_i \cdot w_i \geq 0$  for all  $i \in \{1, \dots, N\}$ ,  $\lambda_0 \in \mathbb{R}$  to  $\frac{1}{N} \sum_{i=1}^N T_i \cdot w_i - 1 = 0$ , and  $\lambda \in \mathbb{R}^N$  to the  $N$  box constraints.  $\diamond$

As an example for the interested reader and as transition to the next section, we show how  $\rho$  is linked to the first constraint of Problem 1.

**Lemma 2.6.** *Let  $(\rho^\dagger, \lambda_0^\dagger, \lambda^\dagger)$  be the optimal solution to Problem 2.4. It holds*

$$\rho_i^\dagger = \left[ \varphi'(0) - \left( \lambda_0^\dagger + \langle B(X_i), \lambda^\dagger \rangle \right) \right]^+ \quad \text{for all } i \in \{1, \dots, N\}.$$

*Furthermore, the unique optimal solutions to Problem 1 are*

$$w_i^\dagger := \left[ (\varphi')^{-1} \left( \lambda_0^\dagger + \langle B(X_i), \lambda^\dagger \rangle \right) \right]^+ \quad \text{for all } i \in \{1, \dots, n\}.$$

**Proof.** For  $i > n$  this is clear. Let  $i \leq n$ , that is,  $T_i = 1$ . By the complementary slackness (see the proof of Theorem 7.4), if  $\rho_i^\dagger > 0$  it holds

$$w_i^\dagger = (\varphi')^{-1} \left( \rho_i^\dagger + \lambda_0^\dagger + \langle B(X_i), \lambda^\dagger \rangle \right) = 0.$$

By Lemma 2.1 it follows

$$\rho_i = \varphi'(0) - \left( \lambda_0^\dagger + \langle B(X_i), \lambda^\dagger \rangle \right).$$

If on the other hand  $\rho_i = 0$ , then by the complementary slackness it follows

$$w_i^\dagger = (\varphi')^{-1} \left( \lambda_0^\dagger + \langle B(X_i), \lambda^\dagger \rangle \right) \geq 0,$$

and

$$0 \geq \varphi'(0) - \left( \lambda_0^\dagger + \langle B(X_i), \lambda^\dagger \rangle \right).$$

□

**Takeaways** We derive a dual formulation of Problem 1 that is easier to analyse. Theorem 2.1 provides a functional relationship of optimal dual solutions and optimal weights. The dual variables are connected to the constraints of the primal problem.





## 3 Constructing the Weights Process

In the formulation of Theorem 2.4 we encounter "If there exists the optimal solution  $(\rho^\dagger, \lambda_0^\dagger, \lambda) \dots$ ". To be able to study asymptotic properties of the weights, we shall assume that Problem 2.4 is feasible, construct a measurable dual solution, and plug it in  $(\varphi')^{-1}$ . Before we formulate concrete assumptions, we provide tools from functional analysis to obtain measurability. Afterwards, we tailor the feasibility assumptions to the capability of this tools. Then, we interpose a section on basis functions before we construct the weights process - the theoretical analogy of optimal weights.

### 3.1 Argmax Measurability Theorem

We follow [AB07]. A **correspondence**  $\psi$  from a set  $S_1$  to a set  $S_2$  assigns to each  $s_1 \in S_1$  a subset  $\psi(s_1) \subset S_2$ . To clarify that we map  $s_1$  to a set, we use the double arrow, that is,  $\psi: S_1 \rightrightarrows S_2$ . Let  $(\mathcal{Z}, \Sigma_{\mathcal{Z}})$  be a measurable space and  $\mathcal{S}$  a topological space. We say, that a correspondence  $\psi: \mathcal{Z} \rightrightarrows \mathcal{S}$  is **weakly measurable**, if

$$\{z \in \mathcal{Z} \mid \psi(z) \cap O \neq \emptyset\} \in \Sigma_{\mathcal{Z}} \quad \text{for all open subsets } O \subset \mathcal{S}.$$

A **selector** from a correspondence  $\psi: \mathcal{Z} \rightrightarrows \mathcal{S}$  is a function  $s: \mathcal{Z} \rightarrow \mathcal{S}$  that satisfies

$$s(z) \in \psi(z) \quad \text{for all } z \in \mathcal{Z}.$$

**Definition 3.1.** Let  $(\mathcal{Z}, \Sigma_{\mathcal{Z}})$  be a measurable space, and let  $\mathcal{S}_1$  and  $\mathcal{S}_2$  be topological space. A function  $f: \mathcal{Z} \times \mathcal{S}_1 \rightarrow \mathcal{S}_2$  is a **Caratheodory function** if

$$f(\cdot, s_1): \mathcal{Z} \rightarrow \mathcal{S}_2 \quad \text{is } (\Sigma_{\mathcal{Z}}, \mathcal{B}(\mathcal{S}_2)) - \text{measurable for all } s_1 \in \mathcal{S}_1,$$

and

$$f(z, \cdot): \mathcal{S}_1 \rightarrow \mathcal{S}_2 \quad \text{is continuous for all } z \in \mathcal{Z}.$$

**Theorem 3.1.** *Let  $\mathcal{S}$  be a separable metrizable space and  $(\mathcal{Z}, \Sigma_{\mathcal{Z}})$  a measurable space. Let  $\psi: \mathcal{Z} \rightrightarrows \mathcal{S}$  be a weakly measurable correspondence with non-empty compact values, and suppose  $f: \mathcal{Z} \times \mathcal{S} \rightarrow \mathbb{R}$  is a Caratheodory function. Define the value function  $m: \mathcal{Z} \rightarrow \mathbb{R}$  by*

$$m(z) := \max_{s \in \psi(z)} f(z, s),$$

*and the correspondence  $\mu: \mathcal{Z} \rightrightarrows \mathcal{S}$  of maximizers by*

$$\mu(z) := \{s \in \psi(z) | f(z, s) = m(z)\}.$$

*Then the value function  $m$  is measurable, the argmax correspondence  $\mu$  has non-empty and compact values, is measurable and admits a measurable selector.*

**Proof.** [AB07, Theorem 18.19] □

### 3.2 Measurable Dual Solution

Next, we formulate the feasibility assumption. Note that we assume compactness to be able to apply Theorem 3.1.

**Assumption 1.** *For all  $N \in \mathbb{N}$  there exists a compact and deterministic parameter space  $\Theta_N \subset \mathbb{R}_{\geq 0}^N \times \mathbb{R} \times \mathbb{R}^N$  such that the optimal solution  $(\rho^\dagger, \lambda_0^\dagger, \lambda^\dagger)$  of Problem 2.4 are contained in  $\Theta_N$ .*

Based on this assumption it is easy to derive measurability for the dual solutions  $(\rho^\dagger, \lambda_0^\dagger, \lambda^\dagger)$ . To this end, we take a closer look on the objective function.

**Definition 3.2.** We define the (random) objective function of (the maximize version of) Problem 2.4 by

$$G : (\Omega, \sigma(D_N)) \times (\mathbb{R}_{\geq 0}^N \times \mathbb{R} \times \mathbb{R}^N) \rightarrow \overline{\mathbb{R}}$$

with

$$G(\omega, (\rho, \lambda_0, \lambda)) = \infty \quad \text{if } \rho_i \neq 0 \text{ for some } i > n,$$

and else

$$\begin{aligned} G(\omega, (\rho, \lambda_0, \lambda)) &= \frac{1}{N} \sum_{i=1}^N \left[ T_i(\omega) \cdot \varphi^*(\rho_i + \lambda_0 + \langle B(X_i)(\omega), \lambda \rangle) - \lambda_0 - \langle B(X_i)(\omega), \lambda \rangle \right] \\ &\quad + \langle \delta(\omega), |\lambda| \rangle. \end{aligned}$$

**Lemma 3.1.** *The function  $G$  of Definition 3.2 is Caratheodory.*

**Proof.** This follows from continuity of  $\varphi^*$  and the measurability of all random variables included.  $\square$

In the proof of the lemma gathers the arguments and applies Theorem 3.1.

**Lemma 3.2.** *For all  $N \in \mathbb{N}$  the dual solution*

$$(\rho^\dagger, \lambda_0^\dagger, \lambda^\dagger) : \Omega \rightarrow \mathbb{R}_{\geq 0}^N \times \mathbb{R} \times \mathbb{R}^N$$

*to Problem 2.4 is*

$$\left( \sigma \left( (T_i, X_i)_{i \in \{1, \dots, N\}} \right), \mathcal{B}(\mathbb{R}_{\geq 0}^N \times \mathbb{R} \times \mathbb{R}^N) \right) - \text{measurable}.$$

**Proof.** Since  $\Theta_N$  is deterministic (by Assumption 1) we can define the (constant) correspondence  $\omega \mapsto \Theta_N$ . Clearly, this is weakly-measurable. Next, we consider the (random) objective function of (the maximize version of) Problem 2.4, that is,  $-G$  (see Definition 3.2). By Lemme 3.1,  $-G$  is a Caratheodory function. Since  $-G$  is also strictly concave, it has a unique argmax in  $\Theta_N$ . By Assumption 1 this is  $(\rho^\dagger, \lambda_0^\dagger, \lambda^\dagger)$ . By Theorem 3.1 this is

$$(\sigma(D_N), \mathcal{B}(\mathbb{R}_{\geq 0}^N \times \mathbb{R} \times \mathbb{R}^N)) - \text{measurable}.$$

$\square$

### 3.3 Basis Functions

Going back to the functional relationship of optimal dual solution and optimal weights (see Theorem 2.1), we see that the basis vector of the covariates plays an important

### 3 Constructing the Weights Process

role. Now, we present our choice. To the best of our knowledge, this is a novelty in the framework of balancing weights.

Let  $(\mathcal{P}_N)$  denote a sequence of countable,  $\mathcal{B}$ -measurable partitions

$$\mathcal{P}_N = \{A_{N,1}, A_{N,2}, \dots\} \subset \mathcal{B}(\mathbb{R}^d)$$

of  $\mathbb{R}^d$ , that is,

$$A_{N,i} \cap A_{N,j} = \emptyset \quad \text{if } i \neq j \quad \text{and} \quad \bigcap_{i \in \mathbb{N}} A_{N,i} = \mathbb{R}^d.$$

We define  $A_N(x)$  to be the cell of  $\mathcal{P}_N$  containing  $x$ , that is,

$$A_N: \mathbb{R}^d \rightarrow \mathbb{R}^d, \quad x \mapsto A_N(x),$$

where  $A_N(x)$  is the only cell containing  $x$ .

**Lemma 3.3.** *The relation*

$$x \sim y \quad :\Leftrightarrow \quad x \in A_N(y)$$

*is an equivalence relation.*

**Proof.** The proof is simple. We omit it. □

Before we define the basis vector, we assume uniform partition width such that

$$\lambda(A_N) =: h_N^d \rightarrow 0 \quad \text{for } N \rightarrow \infty.$$

Next, we define the (empirical) basis functions vector

$$B: \mathbb{R}^d \times \mathbb{R}^{d \cdot N} \rightarrow \mathbb{R}, \quad (x, (x_1, \dots, x_N)) \mapsto \frac{[\mathbf{1}_{A_N(x)}(x_k)]_{k \in \{1, \dots, N\}}}{\sum_{j=1}^N \mathbf{1}_{A_N(x)}(x_j)}, \quad (3.1)$$

where we keep to the convention "0/0 = 0". We shall extend  $B$  to depend on the random vectors  $X, X_1, \dots, X_N$ . The next lemma studies the measurability of the extensions.

**Lemma 3.4.**

- (i)  $B(\cdot, (X_1, \dots, X_N))(\omega)$  is  $(\mathcal{B}(\mathbb{R}^d), \mathcal{B}(\mathbb{R}^N))$ -measurable and constant on each cell  $A_N \in \mathcal{P}_N$  for all  $\omega \in \Omega$ .
- (ii)  $B(X, (X_1, \dots, X_N))$  is  $(\sigma(X, D_N), \mathcal{B}(\mathbb{R}^N))$ -measurable.

**Proof.** Consider for  $k \in \{1, \dots, N\}$  and  $\omega \in \Omega$  the indicator function

$$\mathbf{1}_{A_N(X_k(\omega))}: \mathbb{R}^d \rightarrow \{0, 1\} . \quad (3.2)$$

Since  $A_N(X_k(\omega)) \in \mathcal{B}(\mathbb{R}^d)$  this is a  $(\mathcal{B}(\mathbb{R}^d), \mathcal{B}(\mathbb{R}))$ -measurable function. From the definition of  $B$  (3.1) it follows the first part of (i). Since the indicator function in (3.2) is 1 if  $x \in A_N(X_k(\omega))$  and 0 else, it is also constant on each cell  $A_N \in \mathcal{P}_N$ . It follows (i). To prove (ii), note that

$$\mathbf{1}_{A_N(X_k(\omega))}(X(\omega)) = \mathbf{1} \bigcup_{i \in \mathbb{N}} \{X, X_k \in A_{N,i}\}(\omega) \quad \text{for all } \omega \in \Omega ,$$

and  $\bigcup_{i \in \mathbb{N}} \{X, X_k \in A_{N,i}\} \in \sigma(X, D_N)$ . □

Now we gather some useful properties of the (empirical) basis vector.

**Lemma 3.5.** Let  $(x, x_1, \dots, x_N) \in \mathbb{R}^{d(N+1)}$ .

- (i)  $\sum_{k=1}^N B_k(x, x_1, \dots, x_N) \in \{0, 1\}$ . In particular,  $x_1, \dots, x_N \notin A_N(x)$  is equivalent to  $\sum_{k=1}^N B_k(x, x_1, \dots, x_N) = 0$
- (ii)  $\sum_{k=1}^N B_k(x_i, x_1, \dots, x_N) = 1$  for all  $i \in \{1, \dots, N\}$ .
- (iii)  $\|B(x, x_1, \dots, x_N)\|_2 \leq 1$
- (iv)  $B_k(x_i, x_1, \dots, x_N) = B_i(x_k, x_1, \dots, x_N)$  for all  $i, k \in \{1, \dots, N\}$

**Proof.** Let  $(x, x_1, \dots, x_N) \in \mathbb{R}^{d(N+1)}$ . We prove (i). Then (ii) is a direct consequence of (i). If  $x_1, \dots, x_N \notin A_N(x)$ , then

$$B_k(x, x_1, \dots, x_N) = \frac{\mathbf{1}_{A_N(x)}(x_k)}{\sum_{j=1}^N \mathbf{1}_{A_N(x)}(x_j)} = 0 \quad \text{for all } k \in \{1, \dots, N\} .$$

On the other hand, if the sum is 0 it holds

$$\mathbf{1}_{A_N(x)}(x_k) = 0 \quad \text{for all } k \in \{1, \dots, N\} .$$

It follows the desired equivalence. If

$$\mathbf{1}_{A_N(x)}(x_k) = 1 \quad \text{for some } k \in \{1, \dots, N\} ,$$

then  $\sum_{j=1}^N \mathbf{1}_{A_N(x)}(x_j) \geq 1$  and thus "0/0" doesn't occur. It follows

$$\sum_{k=1}^N B_k(x, x_1, \dots, x_N) = \frac{\sum_{k=1}^N \mathbf{1}_{A_N(x)}(x_k)}{\sum_{j=1}^N \mathbf{1}_{A_N(x)}(x_j)} = 1 .$$

### 3 Constructing the Weights Process

To prove (iii), note that by (i)

$$\|B(x, x_1, \dots, x_N)\|_2^2 = \sum_{k=1}^N B_k(x, x_1, \dots, x_N)^2 \leq \sum_{k=1}^N B_k(x, x_1, \dots, x_N) \leq 1.$$

To prove (iv), note that by Lemma 3.3 and by symmetry and transitivity of the equivalence relation  $x \in A_N(y)$  it holds

$$\begin{aligned} B_k(x_i, x_1, \dots, x_N) &= \frac{\mathbf{1}\{x_k \in A_N(x_i)\}}{\sum_{j=1}^N \mathbf{1}\{x_j, x_k \in A_N(x_i)\}} = \frac{\mathbf{1}\{x_i \in A_N(x_k)\}}{\sum_{j=1}^N \mathbf{1}\{x_j \in A_N(x_k)\}} \\ &= B_i(x_k, x_1, \dots, x_N). \end{aligned}$$

□

Now we show that the basis vector plays well with uniformly continuous functions. The result seems simple, yet the consequence are great. It allows us later on to specify an oracle parameter instead of assuming its existence (see [WZ19, Assumption 1.6]). This greatly clarifies the proofs.

**Lemma 3.6.** *Let  $(x, x_1, \dots, x_N) \in \mathbb{R}^{d(N+1)}$ . For all uniformly continuous functions  $f: \mathbb{R}^d \rightarrow \mathbb{R}$  it holds*

$$\left| \sum_{k=1}^N B_k(x_i, x_1, \dots, x_N) \cdot f(x_k) - f(x_i) \right| \leq \omega\left(f, h_N^d\right) \quad \text{for all } i \in \{1, \dots, N\},$$

where  $\omega(f, \cdot)$  is the uniform modulus of continuity of  $f$ .

**Proof.** It follows from Lemma 3.5(ii)

$$\begin{aligned} &\left| \sum_{k=1}^N B_k(x_i, x_1, \dots, x_N) \cdot f(x_k) - f(x_i) \right| \\ &\leq \left| \sum_{k=1}^N B_k(x_i, x_1, \dots, x_N) (f(x_k) - f(x_i)) \right| \\ &\leq \sum_{k=1}^N B_k(x_i, x_1, \dots, x_N) \cdot \mathbf{1}\{x_k \in A_N(x_i)\} |f(x_k) - f(x_i)| \\ &\leq \omega\left(f, h_N^d\right). \end{aligned}$$

□

Next, we bring forward the applications of Lemma 3.6 that we need. On a high-level it says that the basis functions estimate both treatment (i) and outcome model (ii) well. This feature is connected to double robustness discussed in [ZP17].

In the following, let  $F_{Y(1)}(\cdot|x)$  denote the distribution function of  $Y(1)$  conditional on  $X = x \in \mathcal{X}$  (see (5.1)).

**Lemma 3.7.** *Let  $(x, x_1, \dots, x_N) \in \mathcal{X}^{N+1}$ . It holds for  $N \rightarrow \infty$*

(i)

$$\frac{1}{N} \sum_{i,k=1}^N \left| B_k(x_i, x_1, \dots, x_N) \cdot \varphi' \left( \frac{1}{\pi(x_k)} \right) - \varphi' \left( \frac{1}{\pi(x_i)} \right) \right| \rightarrow 0,$$

(ii) *If it holds  $\sqrt{N} \sup_{z \in \mathbb{R}} \omega(F_{Y(1)}(z|\cdot), h_N^d) \rightarrow 0$  for  $N \rightarrow \infty$ , then*

$$\sqrt{N} \sup_{z \in \mathbb{R}} \max_{i \in \{1, \dots, N\}} \sum_{k=1}^N |B_k(x_i, x_1, \dots, x_N) \cdot F_{Y(1)}(z|x_k) - F_{Y(1)}(z|x_i)| \rightarrow 0.$$

**Proof.** By Lemma 3.6, the uniform continuity of  $\varphi'$  it holds

$$\frac{1}{N} \sum_{i,k=1}^N \left| B_k(x_i, x_1, \dots, x_N) \cdot \varphi' \left( \frac{1}{\pi(x_k)} \right) - \varphi' \left( \frac{1}{\pi(x_i)} \right) \right| \leq \omega(\varphi', h_N^d) \rightarrow 0$$

for  $N \rightarrow \infty$ . Likewise

$$\begin{aligned} & \sqrt{N} \sup_{z \in \mathbb{R}} \max_{i \in \{1, \dots, N\}} \sum_{k=1}^N |B_k(x_i, x_1, \dots, x_N) \cdot F_{Y(1)}(z|x_k) - F_{Y(1)}(z|x_i)| \\ & \leq \sqrt{N} \sup_{z \in \mathbb{R}} \omega(F_{Y(1)}(z|\cdot), h_N^d) \rightarrow 0 \quad \text{for } N \rightarrow \infty. \end{aligned}$$

□

**Remark.** We want to comment on the assumption

$$\sqrt{N} \sup_{z \in \mathbb{R}} \omega(F_{Y(1)}(z|\cdot), h_N^d) \rightarrow 0 \quad \text{for } N \rightarrow \infty,$$

I decided to keep this more general (and abstract) assumption, although there are many (more concrete, yet stronger) assumptions on the regularity of  $F_{Y(1)}(z|\cdot)$  and the convergence speed of  $h_N$ . If for example  $F_{Y(1)}(z|\cdot)$  is  $\alpha$ -Hölder continuous with  $\alpha \in (0, 1]$  for all  $z \in \mathbb{R}$ , it suffices  $\sqrt{N} h_N^{\alpha \cdot d} \rightarrow 0$ .

◇

**Takeaways** Basis functions of non-parametric partitioning estimates are new to the framework of balancing weights. They play well with uniformly continuous functions and promise to simplify the analysis. This choice of basis functions waits to be tested in practice.



### 3.4 Weights Process

Based on Theorem 2.1 we want to use the dual solution  $(\rho^\dagger, \lambda_0^\dagger, \lambda^\dagger)$  to construct weights. To this end, we define the (empirical) weights function

$$w : (\mathbb{R}^d \times \mathbb{R}^{d \cdot N}) \times (\mathbb{R}_{\geq 0}^N \times \mathbb{R} \times \mathbb{R}^N) \rightarrow \mathbb{R}^N$$

$$((x, x_1, \dots, x_N), (\rho, \lambda_0, \lambda)) \mapsto \left[ (\varphi')^{-1}(\rho_i + \lambda_0 + \langle B(x, x_1, \dots, x_N), \lambda \rangle) \right]_{i \in \{1, \dots, N\}}.$$

**Definition 3.3.** Let  $(\rho^\dagger, \lambda_0^\dagger, \lambda^\dagger)$  be the dual solution of Lemma 3.2. We define the weights process  $\{w^\dagger(x) | x \in \mathbb{R}^d\}$  by

$$w^\dagger(x) := w\left((x, X_1, \dots, X_N), (\rho^\dagger, \lambda_0^\dagger, \lambda^\dagger)\right) \quad \text{for all } x \in \mathbb{R}^d.$$

**Lemma 3.8.**

- (i)  $w^\dagger(\cdot)(\omega)$  is  $(\mathcal{B}(\mathbb{R}^d), \mathcal{B}(\mathbb{R}^N))$ -measurable and constant on each cell  $A_N \in \mathcal{P}_N$  for all  $\omega \in \Omega$ .
- (ii)  $w^\dagger(X)$  is  $(\sigma(X, D_N), \mathcal{B}(\mathbb{R}^N))$ -measurable.

**Proof.** This is a direct consequence of Lemme 3.4, Lemma 3.2 and the (assumed) continuity of  $(\varphi')^{-1}$ .  $\square$

**Lemma 3.9.** It holds  $w_i^\dagger(X) \in L^\infty(\mathbf{P})$  for all  $i \in \{1, \dots, N\}$ .

**Proof.** By Lemma 3.5.(iii) it holds

$$\left| \rho_i^\dagger + \lambda_0^\dagger + \langle B(x, x_1, \dots, x_N), \lambda^\dagger \rangle \right| \lesssim \left\| (\rho^\dagger, \lambda_0^\dagger, \lambda^\dagger) \right\|_2 \quad \text{for all } i \in \{1, \dots, N\},$$

where  $\lesssim$  denotes the lesser-or-equal-up-to-a-uniform-constant order, that is, we choose  $C > 1$  independent of  $N$  large enough, such that  $a \lesssim b$  means  $a \leq C \cdot b$ . Since  $(\rho^\dagger, \lambda_0^\dagger, \lambda^\dagger)$  is contained in the deterministic and compact parameter space  $\Theta_N$ , it holds

$$\left\| (\rho^\dagger, \lambda_0^\dagger, \lambda^\dagger) \right\|_2 \in L^\infty(\mathbf{P}).$$

By the (assumed) uniform continuity of  $(\varphi')^{-1}$  on  $\mathbb{R}$ , it follows  $w_i^\dagger(X) \in L^\infty(\mathbf{P})$  for all  $i \in \{1, \dots, N\}$ .  $\square$

Next, we want to simplify the weights process in the spirit of Lemma 2.6. In other words, we want to become independent of the index  $i$  in  $w_i^\dagger$ . This will be helpful in the subsequent analysis. To this end, we define the (empirical) simplified weights function

$$w_0 : (\mathbb{R}^d \times \mathbb{R}^{d \cdot N}) \times (\mathbb{R} \times \mathbb{R}^N) \rightarrow [0, \infty)$$

$$((x, x_1, \dots, x_N), (\lambda_0, \lambda)) \mapsto \left[ (\varphi')^{-1} (\lambda_0 + \langle B(x, x_1, \dots, x_N), \lambda \rangle) \right]^+.$$

**Definition 3.4.** Let  $(\rho^\dagger, \lambda_0^\dagger, \lambda^\dagger)$  be the dual solution of Lemma 3.2. We define the simplified weights process  $\{w_0^\dagger(x) \mid x \in \mathbb{R}^d\}$  by

$$w_0^\dagger(x) := w_0((x, X_1, \dots, X_N), (\lambda_0^\dagger, \lambda^\dagger)) \quad \text{for all } x \in \mathbb{R}^d.$$

**Lemma 3.10.**

- (i)  $w_0^\dagger(\cdot)(\omega)$  is  $(\mathcal{B}(\mathbb{R}^d), \mathcal{B}(\mathbb{R}^N))$ -measurable and constant on each cell  $A_N \in \mathcal{P}_N$  for all  $\omega \in \Omega$ .
- (ii)  $w_0^\dagger(X)$  is  $(\sigma(X, D_N), \mathcal{B}(\mathbb{R}^N))$ -measurable.

**Proof.** The proof is as that of Lemma 3.8. □

**Lemma 3.11.** It holds  $w_0^\dagger(X) \in L^\infty(\mathbf{P})$ .

**Proof.** By Lemma 3.9, the monotonicity of  $(\varphi')^{-1}$  and  $\rho_i \geq 0$  for  $i \leq n$ , it holds

$$\begin{aligned} w_0^\dagger(X) &\leq \left[ (\varphi')^{-1} (\lambda_0^\dagger + \langle B(X), \lambda^\dagger \rangle) \right]^+ \\ &\leq \left[ (\varphi')^{-1} (\rho_i^\dagger + \lambda_0^\dagger + \langle B(X), \lambda^\dagger \rangle) \right]^+ \leq |w_i^\dagger(X)| \in L^\infty(\mathbf{P}) \end{aligned}$$

□

**Lemma 3.12.** Let  $Z \in L^1(\mathbf{P})$  be a random variable that is independent of  $D_N =$

### 3 Constructing the Weights Process

$(T_i, X_i)_{i \in \{1, \dots, N\}}$  with  $\mathbf{E}[Z | X] = 0$  almost surely. It holds

$$\mathbf{E} \left[ w_0^\dagger(X) \cdot Z \right] = 0.$$

**Proof.** By Lemma 3.11 it holds

$$\left\| w_0^\dagger(X) \cdot Z \right\|_{L^1(\mathbf{P})} \leq \left\| w_0^\dagger(X) \right\|_{L^\infty(\mathbf{P})} \|Z\|_{L^1(\mathbf{P})} < \infty. \quad (3.3)$$

By (3.3),  $Z \perp D_N$  and  $\mathbf{E}[Z | X] = 0$  almost surely it holds

$$\begin{aligned} \mathbf{E} \left[ w_0^\dagger(X) \cdot Z \mid D_N, X \right] &= w_0^\dagger(X) \cdot \mathbf{E}[Z \mid D_N, X] \\ &= w_0^\dagger(X) \cdot \mathbf{E}[Z \mid X] = 0 \quad \text{almost surely.} \end{aligned}$$

Note, that  $w_0^\dagger(X)$  is  $(\sigma(D_N, X), \mathcal{B}(\mathbb{R}))$ -measurable. Thus

$$\mathbf{E} \left[ w_0^\dagger(X) \cdot Z \right] = \mathbf{E} \left[ \mathbf{E} \left[ w_0^\dagger(X) \cdot Z \mid D_N, X \right] \right] = 0.$$

□

**Theorem 3.2.** *The simplified weights process satisfies the constraints of Problem 1, that is,*

$$(i) \quad T_i \cdot w_0^\dagger(X_i) \geq 0 \quad \text{for all } i \in \{1, \dots, N\}$$

$$(ii) \quad \frac{1}{N} \sum_{i=1}^N T_i \cdot w_0^\dagger(X_i) = 1$$

(iii) For all  $k \in \{1, \dots, N\}$  it holds

$$\left| \frac{1}{N} \left( \sum_{i=1}^N T_i \cdot w_0^\dagger(X_i) \cdot B_k(X_i, X_1, \dots, X_N) - \sum_{i=1}^N B_k(X_i, X_1, \dots, X_N) \right) \right| \leq \delta_k$$

**Proof.** This follows from Theorem 2.1, Lemma 2.6 and the construction of the simplified weights process. □

To avoid notational overload, from now on we write

$$B(x) := B(x, X_1, \dots, X_N) \quad \text{for all } x \in \mathbb{R}^d.$$

## 4 Consistency of the Weights Process

The goal of this section is to establish consistency of the weights process for the inverse propensity score. To this end, we first show that asymptotically there exists an optimal solution  $(\rho^\dagger, \lambda^\dagger, \lambda_0^\dagger)$  to Problem 2.4 that converges to the oracle parameter

$$(0_N, 0, \lambda^*) \quad \text{where} \quad \lambda^* := \left[ \varphi' \left( \frac{1}{\pi(X_k)} \right) \right]_{k \in \{1, \dots, N\}}$$

in probability (see Theorem). This result justifies Assumption 1. Furthermore, we will identify the dual solution from Lemma 3.2 with the consistent dual solution to derive consistency of the weights process for the inverse propensity score. Before we start the analysis we interpose a section on the propensity score function.

### 4.1 Propensitiy Score Function

Here is the Propensitiy Score.

### 4.2 Consistency of the Dual Solution

We get a grip by the following lemma. The high-level idea is that the existence of the optimal dual solution and its proximity to the oracle parameter can be analysed by the objective function.

**Lemma 4.1.** *Let  $m, N \in \mathbb{N}$  and let  $g : \mathbb{R}_{\geq 0}^N \times \mathbb{R}^m \rightarrow \overline{\mathbb{R}}$  be a continuous and proper convex function. Consider*

$$\tilde{S}(\varepsilon) := \{(\Delta_\rho, \Delta) \in \mathbb{R}_{\geq 0}^N \times \mathbb{R}^m : \|(\Delta_\rho, \Delta)\|_2 = \varepsilon\} \quad \text{for } \varepsilon > 0.$$

*Then for all  $y \in \mathbb{R}^m$  and  $\varepsilon > 0$*

$$\inf \left\{ g(\Delta_\rho, y + \Delta) - g(0, y) : (\Delta_\rho, \Delta) \in \tilde{S}(\varepsilon) \right\} \geq 0 \quad (4.1)$$

*implies the existence of a global minimum*

$$(y_\rho^*, y^*) \in \mathbb{R}_{\geq 0}^N \times \mathbb{R}^m \quad \text{of } g \text{ such that} \quad \|(y_\rho^*, y^*) - (0, y)\|_2 \leq \varepsilon.$$

**Proof.** We start by defining the convex set

$$\tilde{B}(\varepsilon) := \{(\Delta_\rho, \Delta) \in \mathbb{R}_{\geq 0}^N \times \mathbb{R}^m : \|(\Delta_\rho, \Delta)\|_2 \leq \varepsilon\} \quad \text{for } \varepsilon > 0.$$

Then the translation  $(0, y) + \tilde{B}(\varepsilon)$  is also convex. Assume towards a contradiction that it holds (4.1) and that there exists

$$(x_\rho^*, x^*) \in \mathbb{R}_{\geq 0}^N \times \mathbb{R}^m \setminus \left((0, y) + \tilde{B}(\varepsilon)\right) \quad \text{such that} \quad g(x_\rho^*, x^*) < g(0, y). \quad (4.2)$$

Since  $(0, y) + \tilde{B}(\varepsilon)$  is bounded, the line segment between  $(x_\rho^*, x^*)$  and  $(0, y)$  crosses its boundary. The boundary consists of two disjoint sets

$$S_0(\varepsilon) := \{(0, y + \Delta) : \Delta \in \mathbb{R}^m \text{ and } \|\Delta\|_2 < \varepsilon\} \quad \text{and} \quad \tilde{S}(\varepsilon).$$

Clearly, if the line segment does not cross  $\tilde{S}(\varepsilon)$  it leaves  $\mathbb{R}_{\geq 0}^N \times \mathbb{R}^m$ . But this is not possible. Thus, there exists  $(\Delta_\rho, \Delta) \in \tilde{S}(\varepsilon)$  and  $\theta \in (0, 1)$  such that

$$\theta \cdot (x_\rho^*, x^*) + (1 - \theta) \cdot (0, y) = (\Delta_\rho, y + \Delta). \quad (4.3)$$

It follows

$$\begin{aligned} g(0, y) &\leq g(\Delta_\rho, y + \Delta) = g(\theta \cdot (x_\rho^*, x^*) + (1 - \theta) \cdot (0, y)) \\ &\leq \theta \cdot g(x_\rho^*, x^*) + (1 - \theta) \cdot g(0, y) < g(0, y), \end{aligned}$$

which is a contradiction. The first inequality is due to (4.1), the equality is due to (4.3), the second inequality is due to the convexity of  $g$ , and the strict inequality is due to assumption (4.2). Thus, all values outside  $(0, y) + \tilde{B}(\varepsilon)$  are greater or equal  $g(0, y)$ . Since  $(0, y) + \tilde{B}(\varepsilon)$  is also compact, the continuous function  $g$  has a local minimum

$$(y_\rho^*, y^*) \in (0, y) + \tilde{B}(\varepsilon).$$

But then it holds

$$g(y_\rho^*, y^*) \leq g(0, y) \leq g(x_\rho, x) \quad \text{for all} \quad (x_\rho, x) \in \mathbb{R}_{\geq 0}^N \times \mathbb{R}^m \setminus \left((0, y) + \tilde{B}(\varepsilon)\right)$$

and

$$g(y_\rho^*, y^*) \leq g(z_\rho, z) \quad \text{for all} \quad (z_\rho, z) \in (0, y) + \tilde{B}(\varepsilon).$$

Thus,  $(y_\rho^*, y^*)$  is also a global minimum in  $\mathbb{R}_{\geq 0}^N \times \mathbb{R}^m$ . Since  $(y_\rho^*, y^*) \in (0, y) + \tilde{B}(\varepsilon)$  there exists  $(\Delta_\rho, \Delta) \in \tilde{B}(\varepsilon)$  such that

$$(y_\rho^*, y^*) = (\Delta_\rho, y + \Delta) \quad \text{for some } (\Delta_\rho, \Delta) \in \tilde{B}(\varepsilon).$$

Thus

$$\|(y_\rho^*, y^*) - (0, y)\|_2 = \|(\Delta_\rho, \Delta)\|_2 \leq \varepsilon.$$

This finish the proof.  $\square$

**Remark.** I learned of the high-level idea from [WZ19, page 22]. I adapted it to the needs of the subsequent analysis and provided the details by myself. Note, that the hint in [WZ19, page 22] uses strict inequality in the statement. I found out that this can be relaxed. It is crucial to my further approach that this holds (only) with inequality, because I use measurability properties to obtain convergence.  $\diamond$

On the basis of the (random) objective function  $G$  of Problem 2.4 (see Definition 3.2) we define, for  $\varepsilon > 0$ , an auxiliary function

$$\begin{aligned} \underline{\Delta G}_\varepsilon^*: (\Omega, \sigma(D_N), \mathbf{P}) &\rightarrow \overline{\mathbb{R}} \\ \omega &\mapsto \inf \{ G(\omega, (\Delta_\rho, \Delta_0, \lambda^*(\omega) + \Delta)) - G(\omega, (0_N, 0, \lambda^*(\omega))) : \| \Delta_\rho, \Delta_0, \Delta \|_2 = \varepsilon \} \end{aligned}$$

**Lemma 4.2.** For all  $\varepsilon > 0$  the function  $\underline{\Delta G}_\varepsilon^*$  is  $(\sigma(D_N), \mathcal{B}(\overline{\mathbb{R}}))$ -measurable.

**Proof.** Let  $\varepsilon > 0$ . By Lemma 3.1, the function

$$\begin{aligned} \Delta G_\varepsilon: \Omega \times (\mathbb{R}^N \times (\mathbb{R}_{\geq 0}^N \times \mathbb{R} \times \mathbb{R}^N)) &\rightarrow \overline{\mathbb{R}} \\ (\omega, (\lambda, (\Delta_\rho \Delta_0 \Delta))) &\mapsto G(\omega, (\Delta_\rho, \Delta_0, \lambda + \Delta)) - G(\omega, (0_N, 0, \lambda)) \end{aligned}$$

is Caratheodory. Since  $\{ \| \Delta_\rho \Delta_0 \Delta \|_2 = \varepsilon \}$  is compact in  $\mathbb{R}_{\geq 0}^N \times \mathbb{R} \times \mathbb{R}^N$ , the function

$$\begin{aligned} \underline{\Delta G}_\varepsilon: \Omega \times \mathbb{R}^N &\rightarrow \overline{\mathbb{R}} \\ (\omega, \lambda) &\mapsto \inf \{ G(\omega, (\Delta_\rho, \Delta_0, \lambda + \Delta)) - G(\omega, (0_N, 0, \lambda)) : \| \Delta_\rho \Delta_0 \Delta \|_2 = \varepsilon \} \end{aligned}$$

is Caratheodory. Since  $\lambda^*$  is  $(\sigma(D_N), \mathcal{B}(\mathbb{R}^N))$ -measurable it follows the statement.  $\square$

**Lemma 4.3.** *It holds for all  $\varepsilon > 0$*

$$\mathbf{P} \left[ \underline{\Delta G_{\varepsilon}^*} \geq 0 \right] \rightarrow 1 \quad \text{for } N \rightarrow \infty.$$

**Proof.** Let  $\varepsilon > 0$  and  $\|\Delta_{\rho}, \Delta_0, \Delta\|_2 = \varepsilon$ . We show

$$\mathbf{P} \left[ \underline{\Delta G_{\varepsilon}^*} \geq -\tilde{\varepsilon} \right] \rightarrow 1 \quad \text{for } N \rightarrow \infty \text{ for all } \tilde{\varepsilon} > 0.$$

Then the result follows from the measurability of  $\underline{\Delta G_{\varepsilon}^*}$  (see Lemma 4.2). To this end, note, that

$$G(\rho, \lambda_0, \lambda) = g(\rho, \lambda_0, \lambda) + \langle \delta, |\lambda| \rangle \quad \text{for all } (\rho, \lambda_0, \lambda) \in \mathbb{R}_{\geq 0}^N \times \mathbb{R} \times \mathbb{R}^N,$$

with

$$g := (\rho, \lambda_0, \lambda) \mapsto \frac{1}{N} \left( \sum_{i=1}^N T_i \cdot \varphi^*(\rho_i + \lambda_0 + \langle B(X_i), \lambda \rangle) - \lambda_0 - \langle B(X_i), \lambda \rangle \right).$$

Since  $\varphi^*$  is continuously differentiable by Lemma 2.2 (it is always convex),  $g$  is a continuously differentiable convex function with gradient

$$\begin{aligned} & (\rho, \lambda_0, \lambda) \\ & \mapsto \frac{1}{N} \left( \sum_{i=1}^N T_i \cdot (\varphi')^{-1}(\rho_i + \lambda_0 + \langle B(X_i), \lambda \rangle) \left[ e_i^\top, 1, B(X_i)^\top \right]^\top - \left[ 0_N^\top, 1, B(X_i)^\top \right]^\top \right). \end{aligned}$$

Thus, by (7.9), it holds

$$\begin{aligned}
 & G(\Delta_\rho, \Delta_0, \lambda^* + \Delta) - G(0_N, 0, \lambda^*) \\
 & \geq \frac{1}{N} \left( \sum_{i=1}^N T_i \cdot (\varphi')^{-1}(\langle B(X_i), \lambda^* \rangle) \begin{bmatrix} e_i^\top, 1, B(X_i)^\top \end{bmatrix} - \begin{bmatrix} 0_N^\top, 1, B(X_i)^\top \end{bmatrix} \right) \begin{bmatrix} \Delta_\rho \\ \Delta_0 \\ \Delta \end{bmatrix} \\
 & \quad + \langle \delta, |\lambda^* + \Delta| - |\lambda^*| \rangle \\
 & \geq \frac{1}{N} \sum_{i=1}^N \left( T_i \cdot (\varphi')^{-1}(\langle B(X_i), \lambda^* \rangle) - 1 \right) \begin{bmatrix} e_i^\top, 1, B(X_i)^\top \end{bmatrix} \cdot \begin{bmatrix} \Delta_\rho \\ \Delta_0 \\ \Delta \end{bmatrix} + \langle e_i, \Delta_\rho \rangle \\
 & \quad + \langle \delta, |\lambda^* + \Delta| - |\lambda^*| \rangle \\
 & \geq -\frac{1}{N} \sum_{i=1}^N \left| \left( T_i \cdot (\varphi')^{-1}(\langle B(X_i), \lambda^* \rangle) - 1 \right) \begin{bmatrix} e_i^\top, 1, B(X_i)^\top \end{bmatrix} \cdot \begin{bmatrix} \Delta_\rho \\ \Delta_0 \\ \Delta \end{bmatrix} \right| \\
 & \quad - \langle \delta, |\Delta| \rangle \\
 & =: -I_1 \\
 & \quad - I_2.
 \end{aligned} \tag{4.4}$$

Note, that  $\Delta_\rho \in \mathbb{R}_{\geq 0}^N$ , and thus  $\langle e_i, \Delta_\rho \rangle \geq 0$  for all  $i \in \{1, \dots, N\}$ , where  $e_i$  is the  $i$ -the unit vector.

### Analysis of $I_1$

By the Cauchy-Schwarz inequality and Lemma 3.5.(iii) it holds

$$\left| \begin{bmatrix} e_i^\top, 1, B(X_i)^\top \end{bmatrix} \cdot \begin{bmatrix} \Delta_\rho \\ \Delta_0 \\ \Delta \end{bmatrix} \right| \leq \|\Delta_\rho, \Delta_0, \Delta\|_2 \leq \varepsilon.$$



Furthermore,

$$\begin{aligned}
 & \frac{1}{N} \sum_{i=1}^N \left| \left( T_i \cdot (\varphi')^{-1} (\langle B(X_i), \lambda^* \rangle) - 1 \right) \right| \\
 & \leq \frac{1}{N} \sum_{i=1}^N \left| 1 - \frac{T_i}{\pi(X_i)} \right| \\
 & \quad + \frac{1}{N} \sum_{i=1}^N \omega \left( (\varphi')^{-1}, \left| \sum_{k=1}^N B_k(X_i) \cdot \varphi' \left( \frac{1}{\pi(X_k)} \right) - \varphi' \left( \frac{1}{\pi(X_i)} \right) \right| \right) \\
 & =: J_1 \\
 & \quad + J_2
 \end{aligned}$$

### Analysis of $J_1$

By the properties of conditional expectation it holds

$$\mathbf{E} \left[ \frac{T}{\pi(X)} \right] = \mathbf{E} \left[ \frac{\mathbf{E}[T|X]}{\pi(X)} \right] = 1.$$

Also

$$\mathbf{E} \left[ \left| 1 - \frac{T}{\pi(X)} \right| \right] \leq 1 + \mathbf{E} \left[ \frac{T}{\pi(X)} \right] = 2. \quad (4.5)$$

Thus Etemadi's ( $\mathcal{L}_1$  version) strong law of large numbers (cf. [Kle20, Theorem 5.17]) applies to  $J_1$ , that is,  $J_1 \xrightarrow{\mathbf{P}} 0$ .

### Analysis of $J_2$

By Lemma 3.7.(i) and the uniform continuity of  $(\varphi')^{-1}$  it holds

$$\begin{aligned}
 \omega \left( (\varphi')^{-1}, \left| \sum_{k=1}^N B_k(X_i) \cdot \varphi' \left( \frac{1}{\pi(X_k)} \right) - \varphi' \left( \frac{1}{\pi(X_i)} \right) \right| \right) & \leq \omega \left( (\varphi')^{-1}, \omega \left( \varphi', h_N^d \right) \right) \\
 & \rightarrow 0.
 \end{aligned}$$

Thus  $J_2 \rightarrow 0$ .

### Conclusion $I_1$

It follows from the analysis of  $J_1$  and  $J_2$

$$\mathbf{P} [I_1 \leq \tilde{\varepsilon}] \rightarrow 1 \quad \text{for all } \tilde{\varepsilon} > 0.$$

Note, that this holds independently of the specific choice of  $\varepsilon > 0$  in the beginning of the proof.

**Analysis of  $I_2$** 

Since  $\delta > 0$ , we get

$$\langle \delta, |\Delta| \rangle \leq \|\delta\|_1 \|\Delta\|_\infty \leq \|\delta\|_1 \varepsilon,$$

Since  $\|\delta\|_1$  converges to 0 in probability, we get

$$\mathbf{P}[I_2 \leq \tilde{\varepsilon}] \rightarrow 1 \quad \text{for all } \tilde{\varepsilon} > 0.$$

**Conclusion**

By (4.4), and the analysis of  $I_1$  and  $I_2$ , we get

$$\mathbf{P}[G(\Delta_\rho, \Delta_0, \lambda^* + \Delta) - G(0_N, 0, \lambda^*) \geq -\tilde{\varepsilon}] \rightarrow 1 \quad \text{for all } \tilde{\varepsilon} > 0.$$

This holds uniformly for all  $\|\Delta_\rho, \Delta_0, \Delta\|_2 = \varepsilon$ . Thus

$$\mathbf{P}[\underline{\Delta G_\varepsilon^*} \geq -\tilde{\varepsilon}] \rightarrow 1 \quad \text{for all } \tilde{\varepsilon} > 0.$$

From the measurability of  $\underline{\Delta G_\varepsilon^*}$  (see Lemma 4.2) it follows

$$\mathbf{P}[\underline{\Delta G_\varepsilon^*} \geq 0] \rightarrow 1.$$

But this holds independently of the choice  $\varepsilon > 0$ . □

**Remark.** The last proof is a simplification of the similar [WZ19, Proof of Lemma 2]. There, the authors claim to derive concrete learning rates. Their proof is very obscure. The shortcomings range from missing assumptions on the eigenvalues of a Hessian matrix (needed to bound the quadratic term (first display of page 23) away from 0), to mixing up the terms in the conclusion (see [WZ19, page 25]). The application of matrix concentration inequalities additionally clouds the proof. I decided, to aim for less, that is, only consistency instead of concrete learning rates. With this choice, a linear bound like (7.9) suffices. Then I get rid of the quadratic Taylor expansion, assumptions on eigenvalues of a Hessian matrix and the application of matrix concentration inequalities. Later, I show that the lack of concrete learning rates is compensated by good approximation properties of the basis functions (see the remark of the section *Analysis of  $R_2$* ). ◇

**Theorem 4.1.** *With probability going to 1 Problem 2.4 is feasible. Furthermore, if the solution  $(\rho^\dagger, \lambda^\dagger, \lambda_0^\dagger)$  exists, it converges in probability to  $(0_N, 0, \lambda^*)$ .*

**Proof.** By Lemma 4.1 and Lemma 4.3 it holds for all  $\varepsilon > 0$

$$\begin{aligned} & \mathbf{P} \left[ \text{Problem 2.4 is feasible and } \left\| (\rho^\dagger, \lambda^\dagger, \lambda_0^\dagger) - (0_N, 0, \lambda^*) \right\|_2 \leq \varepsilon \right] \\ & \geq \mathbf{P} \left[ \underline{\Delta G_\varepsilon^*} \geq 0 \right] \rightarrow 1 \quad \text{for } N \rightarrow \infty. \end{aligned}$$

□

### 4.3 Main Result

**Theorem 4.2.** *If Problem 2.4 is feasible it holds*

$$\max_{i \in \{1, \dots, n\}} \left| w_0^\dagger(X_i) - 1/\pi(X_i) \right| \xrightarrow{\mathbf{P}} 0.$$

*Furthermore, there exists a decreasing sequence  $(\varepsilon_N) \subset (0, 1]$  such that  $\varepsilon_N \rightarrow 0$  and*

$$\mathbf{P} \left[ \max_{i \in \{1, \dots, n\}} \left| w_0^\dagger(X_i) - 1/\pi(X_i) \right| \leq \varepsilon_N \right] \rightarrow 0 \quad \text{for } N \rightarrow \infty.$$

**Proof.** Let  $i \in \{1, \dots, n\}$ . By Lemma 2.6 it holds

$$w_0^\dagger(X_i) = (\varphi')^{-1} \left( \rho_i^\dagger + \lambda_0^\dagger + \langle B(X_i), \lambda^\dagger \rangle \right).$$

Thus

$$\left| w_0^\dagger(X_i) - \frac{1}{\pi(X_i)} \right| \leq \omega \left( (\varphi')^{-1}, \left| \rho_i^\dagger + \lambda_0^\dagger + \sum_{k=1}^N B_k(X_i) \cdot \lambda_k^\dagger - \varphi' \left( \frac{1}{\pi(X_i)} \right) \right| \right). \quad (4.6)$$

With

$$\begin{aligned} & \left| \rho_i^\dagger + \lambda_0^\dagger + \sum_{k=1}^N B_k(X_i) \cdot \lambda_k^\dagger - \varphi' \left( \frac{1}{\pi(X_i)} \right) \right| \\ & \leq \left\| (\rho^\dagger, \lambda_0^\dagger, \lambda^\dagger) - (0_N, 0, \lambda^*) \right\|_2 + \left| \sum_{k=1}^N B_k(X_i) \cdot \varphi' \left( \frac{1}{\pi(X_k)} \right) - \varphi' \left( \frac{1}{\pi(X_i)} \right) \right| \\ & \leq \left\| (\rho^\dagger, \lambda_0^\dagger, \lambda^\dagger) - (0_N, 0, \lambda^*) \right\|_2 + \omega \left( \varphi', h_N^d \right) \end{aligned}$$

we get an upper bound that is independent of  $i$ . Thus,

$$\max_{i \in \{1, \dots, n\}} \left| w_0^\dagger(X_i) - 1/\pi(X_i) \right| \leq \left\| (\rho^\dagger, \lambda_0^\dagger, \lambda^\dagger) - (0_N, 0, \lambda^*) \right\|_2 + \omega \left( \varphi', h_N^d \right)$$

Since

$$\left\| (\rho^\dagger, \lambda_0^\dagger, \lambda^\dagger) - (0_N, 0, \lambda^*) \right\|_2 \xrightarrow{\mathbf{P}} 0 \quad \text{for } N \rightarrow \infty \quad \text{by Theorem 4.1 ,}$$

and  $\omega(\varphi', h_N^{d'}) \rightarrow 0$  by the uniform continuity of  $\varphi'$  and  $h_N \rightarrow 0$ , it follows the first statement. The second statement follows from the selection lemma [SC08, A.1.4.].  $\square$



## 5 Convergence of the Weighted Mean

Throughout this section we use the following notation. Let  $F_{Y(1)}$  denote the distribution function of  $Y(1)$ , that is,

$$F_{Y(1)} : \mathbb{R} \rightarrow [0, 1], \quad z \mapsto \mathbf{P}[Y(1) \leq z].$$

Let  $F_{Y(1)}(\cdot|x)$  denote the distribution function of  $Y(1)$  conditional on  $X = x \in \mathcal{X}$ , that is,

$$F_{Y(1)}(z|x) = \mathbf{P}[Y(1) \leq z | X = x] \quad \text{for all } (z, x) \in \mathbb{R} \times \mathcal{X}. \quad (5.1)$$

We illustrate the flexibility of the weighted mean estimator by extending the method of [WZ19] to estimates of the distribution function of  $Y(1)$ , that is,  $F_{Y(1)}$ . For the asymptotic analysis of estimating the mean  $\mathbf{E}[Y(1)]$  see [WZ19, Proof of Theorem 3]. To make this extension, the central observation is, that we can adapt the error decomposition in [WZ19, page 27] to estimates of the distribution function  $F_{Y(1)}$  of  $Y(1)$ . We do this in Lemma 5.11. With this modification, we aim at proving the convergence of

$$\sqrt{N} \left( \frac{1}{N} \sum_{i=1}^n w_0^\dagger(X_i) \mathbf{1}\{Y_i(T_i) \leq z\} - F_{Y(1)}(z) \right)_{z \in \mathbb{R}} \quad (5.2)$$

in  $l^\infty(\mathbb{R})$  to a Gaussian process with mean 0 and covariance specified in Theorem ???. We show, that this is possible under mild assumptions. We discuss this in the next section.

### 5.1 Tools

For the subsequent analysis we need the theory of empirical processes. For an introduction to empirical processes see [vdV00, §19]. For a thorough treatment see [vdVW13, §2].

#### 5.1.1 Empirical Processes - Definition

Let  $(\Omega, \mathcal{A}, \mathbf{P})$  be a probability space,  $(\mathcal{Z}, \Sigma)$  a measurable space, and

$$\xi_1, \dots, \xi_N : (\Omega, \mathcal{A}, \mathbf{P}) \rightarrow (\mathcal{Z}, \Sigma) \quad \text{independent and identically-distributed}$$

## 5 Convergence of the Weighted Mean

random variables with probability distribution  $\mathbf{P}_\xi$ . Let  $\mathcal{F}$  be a class of measurable functions  $f : (\mathcal{Z}, \Sigma) \rightarrow (\mathbb{R}, \mathcal{B}(\mathbb{R}))$ , where  $\mathcal{B}(\mathbb{R})$  is the Borel- $\sigma$ -algebra on  $\mathbb{R}$ . Then  $\mathcal{F}$  induces a stochastic process by

$$f \mapsto \mathbb{G}_N f := \frac{1}{\sqrt{n}} \sum_{i=1}^N (f(\xi_i) - \mathbf{E}_\xi[f]), \quad (5.3)$$

where  $\mathbf{E}_\xi[f] := \int_{\mathcal{Z}} f d\mathbf{P}_\xi$ . We call  $\mathbb{G}_N$  the **empirical process** indexed by  $\mathcal{F}$ . The purpose of this construction is, to study the behaviour of a centered, scaled arithmetic mean uniformly over  $\mathcal{F}$ . To this end, we define the (random) norm

$$\|\mathbb{G}_n\|_{\mathcal{F}} := \sup_{f \in \mathcal{F}} |\mathbb{G}_N f|. \quad (5.4)$$

We stress that  $\|\mathbb{G}_n\|_{\mathcal{F}}$  often ceases to be measurable, even in simple situations [vdVW13, page 3]. To deal with this, we introduce the notion of **outer expectation**  $\mathbf{E}^*$  (see [vdVW13, page 6])

$$\mathbf{E}^*[Z] := \inf \{ \mathbf{E}[U] \mid U \geq Z, U : (\Omega, \mathcal{A}, \mathbf{P}) \rightarrow (\overline{\mathbb{R}}, \mathcal{B}(\overline{\mathbb{R}})) \text{ measurable and } \mathbf{E}[U] < \infty \}.$$

In our application the technical difficulties halt at this point, because we only consider  $Z$  with  $\mathbf{E}^*[Z] < \infty$ . Then there exists a smallest measurable function  $Z^*$  dominating  $Z$  with  $\mathbf{E}^*[Z] = \mathbf{E}[Z^*]$  (see [vdVW13, Lemma 1.2.1]).

An **envelope function**  $F$  of a class  $\mathcal{F}$  satisfies

$$|f(z)| \leq F(z) < \infty \quad \text{for all } f \in \mathcal{F} \text{ and all } z \in \mathcal{Z}.$$

### 5.1.2 Bracketing Numbers and Integral

To control empirical processes - apart from strong theorems - we need the notion of bracketing number and integral (see [vdV00, page 270]). Given two functions  $\underline{f} \leq \overline{f}$ ,

$$\text{the bracket } [\underline{f}, \overline{f}] \text{ is the set of all functions } f \text{ with } \underline{f} \leq f \leq \overline{f}.$$

For  $\varepsilon > 0$  we define a

$$(\varepsilon, L^r(\mathbf{P}))\text{-bracket to be a bracket } [\underline{f}, \overline{f}] \text{ with } \|\overline{f} - \underline{f}\|_{L^r(\mathbf{P})} < \varepsilon.$$

The **bracketing number**  $N_{[]}(\varepsilon, \mathcal{F}, L^r(\mathbf{P}))$  is the minimum number of  $(\varepsilon, L^r(\mathbf{P}))$ -brackets needed to cover  $\mathcal{F}$ .

For most classes  $\mathcal{F}$  the bracketing number grows to infinity for  $\varepsilon \rightarrow 0$ . To measure the speed of growth we introduce for  $\delta > 0$  the **bracketing integral**

$$J_{[]}(\delta, \mathcal{F}, L^r(\mathbf{P})) = \int_0^\delta \sqrt{\log N_{[]}(\varepsilon, \mathcal{F}, L^r(\mathbf{P}))} d\varepsilon.$$

Next we give a technical lemma to bound the bracketing numbers of products of two function classes, that is,

$$\mathcal{F} \cdot \mathcal{G} := \{f \cdot g : f \in \mathcal{F}, g \in \mathcal{G}\}.$$

**Lemma 5.1.** *Let  $\mathcal{F}$  and  $\mathcal{G}$  be two function classes with envelope functions  $F$  and  $G$  satisfying  $\|F\|_\infty, \|G\|_\infty \leq 1$ . For all  $\varepsilon > 0$  and all  $r \in [1, \infty)$  it holds*

$$N_{[]} (2\varepsilon, \mathcal{F} \cdot \mathcal{G}, L_r(\mathbf{P})) \leq N_{[]} (\varepsilon, \mathcal{F}, L_r(\mathbf{P})) \cdot N_{[]} (\varepsilon, \mathcal{G}, L_r(\mathbf{P})).$$

**Proof.** The proof is simple. We omit the details.  $\square$

The following has the advantage of being both example (for the interested reader) and helpful for the subsequent analysis.

For  $z \in \mathbb{R}$  we define the function

$$\begin{aligned} f_z &: \{0, 1\} \times \mathcal{X} \times \mathcal{Y} \rightarrow \mathbb{R} \\ (t, x, y) &\mapsto t (\mathbf{1}_{\{y \leq z\}} - F_{Y(1)}(z|x)), \end{aligned}$$

Next we define the function classes

$$\begin{aligned} \mathcal{F} &:= \{f_z \mid z \in \mathbb{R}\} \\ \mathcal{G} &:= \left\{ \frac{f_z}{\pi(\cdot)} + F_{Y(1)}(z|\cdot) - F_{Y(1)}(z) : z \in \mathbb{R} \right\}. \end{aligned} \tag{5.5}$$

Next, we provide bracketing numbers for these classes.

**Lemma 5.2.** *The function class  $\mathcal{F}$  and  $\mathcal{G}$  defined in (5.5) are measurable. Furthermore,*

$$N_{[]} (\varepsilon, \mathcal{F}, L^2(\mathbf{P})) \lesssim \left( \frac{1}{\varepsilon} \right)^2 \quad \text{for all } \varepsilon > 0.$$

*If  $1/\pi(X) \in L^2(\mathbf{P})$ , it also holds*

$$N_{[]} (\varepsilon, \mathcal{G}, L^2(\mathbf{P})) \lesssim \left( \frac{1 + \|1/\pi(X)\|_{L^2(\mathbf{P})}}{\varepsilon} \right)^4 \quad \text{for all } \varepsilon > 0.$$

**Proof.** As in [vdV00, Example 19.6] we choose for  $\varepsilon > 0$  and  $m \in \mathbb{N}$

$$-\infty = z_0 < z_1 < \cdots < z_{m-1} < z_m = \infty$$



## 5 Convergence of the Weighted Mean

such that

$$\mathbf{P}[Y(1) \in [z_{l-1}, z_l]] \leq \varepsilon \quad \text{for all } l \in \{1, \dots, m\} \quad (5.6)$$

and  $m \leq 2/\varepsilon$ . Next, we define  $m$  brackets by

$$\begin{aligned} \overline{f}_l(t, x, y) &:= t(\mathbf{1}_{\{y \leq z_l\}} - F_{Y(1)}(z_{l-1}|x)) , \\ \underline{f}_l(t, x, y) &:= t(\mathbf{1}_{\{y \leq z_{l-1}\}} - F_{Y(1)}(z_l|x)) , \end{aligned}$$

for  $l \in \{1, \dots, m\}$ . These brackets cover  $\mathcal{F}$ . Indeed,

$$\text{for all } z \in \mathbb{R} \text{ there exists } l \in \{1, \dots, m\} \quad \text{such that} \quad z_{l-1} \leq z \leq z_l .$$

By the monotonicity of  $\mathbf{1}_{\{y \leq (\cdot)\}}$  and  $F_{Y(1)}(\cdot|x)$  and the non-negativity of  $T$  it follows

$$\text{for all } z \in \mathbb{R} \text{ there exists } l \in \{1, \dots, m\} \quad \text{such that} \quad \underline{f}_l \leq f_z \leq \overline{f}_l .$$

Thus, the  $m$  brackets  $[\underline{f}_l, \overline{f}_l]$  cover  $\mathcal{F}$ .

Let's calculate the size of the brackets. It holds

$$\begin{aligned} &\mathbf{E} \left[ T \cdot (\mathbf{1}_{\{Y(T) \leq z_l\}} - F_{Y(1)}(z_{l-1}|X) - \mathbf{1}_{\{Y(T) \leq z_{l-1}\}} + F_{Y(1)}(z_l|X)) \right] \\ &= \mathbf{E} \left[ T \cdot (\mathbf{1}_{\{Y(T) \in [z_{l-1}, z_l]\}} + \mathbf{P}[Y(1) \in [z_{l-1}, z_l] | X]) \right] \\ &\leq \mathbf{E} [\pi(X) \cdot \mathbf{P}[Y(1) \in [z_{l-1}, z_l] | X]] + \varepsilon \\ &\leq 2\varepsilon . \end{aligned}$$

We used (5.6),  $0 \leq T, \pi(X) \leq 1$  and Lemma 5.10. It follows

$$\begin{aligned} &\|(\overline{f}_l - \underline{f}_l)(T, X, Y(T))\|_{L^2(\mathbf{P})} \\ &\lesssim \mathbf{E} \left[ T \cdot (\mathbf{1}_{\{Y(T) \in [z_{l-1}, z_l]\}} + \mathbf{P}[Y(1) \in [z_{l-1}, z_l] | X]) \right]^{1/2} \lesssim \varepsilon^{1/2} . \end{aligned}$$

Since  $m \leq 2/\varepsilon$  it holds

$$N_{[]}(\varepsilon^{1/2}, \mathcal{F}, L^2(\mathbf{P})) \lesssim \frac{1}{\varepsilon}$$

and thus

$$N_{[]}(\varepsilon, \mathcal{F}, L^2(\mathbf{P})) \lesssim \left(\frac{1}{\varepsilon}\right)^2 .$$

Next, we look at  $\mathcal{G}$ . To this end, we define  $m$  brackets by

$$\begin{aligned} \overline{g}_l(t, x, y) &:= \frac{t}{\pi(x)} (\mathbf{1}_{\{y \leq z_l\}} - F_{Y(1)}(z_{l-1}|x)) + F_{Y(1)}(z_l|x) - F_{Y(1)}(z_{l-1}) , \\ \underline{g}_l(t, x, y) &:= \frac{t}{\pi(x)} (\mathbf{1}_{\{y \leq z_{l-1}\}} - F_{Y(1)}(z_l|x)) + F_{Y(1)}(z_{l-1}|x) - F_{Y(1)}(z_l) , \end{aligned}$$

for  $l \in \{1, \dots, m\}$ . With the same arguments as before, we see that these brackets cover  $\mathcal{G}$ . Let's calculate the size. It holds

$$\begin{aligned}
& \left\| \frac{T}{\pi(X)} (\mathbf{1}\{Y(T) \in [z_{l-1}, z_l]\} + \mathbf{P}[Y(1) \in [z_{l-1}, z_l] | X]) \right\|_{L^2(\mathbf{P})} \\
& \lesssim \left( \mathbf{E} \left[ \frac{1}{\pi(X)} \frac{T}{\pi(X)} (\mathbf{1}\{Y(T) \in [z_{l-1}, z_l]\} + \mathbf{P}[Y(1) \in [z_{l-1}, z_l] | X]) \right] \right)^{1/2} \\
& \lesssim \left( \mathbf{E} \left[ \frac{1}{\pi(X)} \mathbf{P}[Y(1) \in [z_{l-1}, z_l] | X] \right] \right)^{1/2} \\
& \lesssim \left( \|1/\pi(X)\|_{L^2(\mathbf{P})} \sqrt{\varepsilon} \right)^{1/2} = \varepsilon^{1/4} \|1/\pi(X)\|_{L^2(\mathbf{P})}^{1/2}
\end{aligned}$$

and

$$\|\mathbf{P}[Y(1) \in [z_{l-1}, z_l] | X] + \mathbf{P}[Y(1) \in [z_{l-1}, z_l]]\|_{L^2(\mathbf{P})} \lesssim \varepsilon^{1/2}.$$

Thus

$$\begin{aligned}
\|(\bar{g}_l - \underline{g}_l)(T, X, Y(T))\|_{L^2(\mathbf{P})} & \lesssim \varepsilon^{1/4} \left( 1 + \|1/\pi(X)\|_{L^2(\mathbf{P})}^{1/2} \right) \\
& \lesssim \varepsilon^{1/4} \left( 1 + \|1/\pi(X)\|_{L^2(\mathbf{P})} \right).
\end{aligned}$$

As before, it follows

$$N_{[]}(\varepsilon, \mathcal{G}, L^2(\mathbf{P})) \lesssim \left( \frac{1 + \|1/\pi(X)\|_{L^2(\mathbf{P})}}{\varepsilon} \right)^4.$$

□

Before we give another example, we fix some useful properties of  $f_z$ .

**Lemma 5.3.** *It holds  $f_z(T, X, Y(T)) \in L^1(\mathbf{P})$  and  $f_z(T, X, Y(T)) \perp D_N$  for all  $z \in \mathbb{R}$ . If also Assumption ?? holds, then for all  $z \in \mathbb{R}$*

$$\mathbf{E}[f_z(T, X, Y(T)) | X] = 0 \quad \text{almost surely.}$$

**Proof.** Since  $f_z$  is bounded by 1, it holds  $f_z(T, X, Y(T)) \in L^1(\mathbf{P})$ . Since

$$(T, X, Y(T)) \perp D_N = (T_i, X_i)_{i \in \{1, \dots, N\}}$$

it holds  $f_z(T, X, Y(T)) \perp D_N$  for all  $z \in \mathbb{R}$ . For the third statement, note that

$$\begin{aligned}
\mathbf{E}[f_z(T, X, Y(T)) | X] &= \mathbf{E}[T(\mathbf{1}_{\{Y(T) \leq z\}} - F_{Y(1)}(z|X)) | X] \\
&= \mathbf{E}[\mathbf{1}_{\{Y(1) \leq z\}} - F_{Y(1)}(z|X) | X, T = 1] \pi(X) \\
&= (\mathbf{E}[\mathbf{1}_{\{Y(1) \leq z\}} | X] - F_{Y(1)}(z|X)) \pi(X) \\
&= 0 \quad \text{almost surely.}
\end{aligned}$$

The third equality is due to Assumption ??.

□

## 5 Convergence of the Weighted Mean

Next, consider the stochastic process (indexed over  $x \in \mathbb{R}^d$ )

$$\mathbf{1} \left\{ \sup_{y \in A_N(x)} \left| w_0^\dagger(y) - \frac{1}{\pi(y)} \right| \leq \varepsilon_N \right\} \left( w_0^\dagger(x) - \frac{1}{\pi(x)} \right) \cdot \mathbf{1} \bigcup_{k=1}^n \{x = X_k\}, \quad (5.7)$$

where  $(\varepsilon_N)$  is the learning rate of Theorem 4.2. We show, that under mild regularity conditions on the inverse propensity score function all paths of (5.7) are contained in shrinking function classes  $(\mathcal{F}_N)$  - and provide bracketing numbers. To be more precise, we need theory from [vdVW13, §2.7.1].

Let for any vector  $k \in \mathbb{N}_0^d$  ( $d \in \mathbb{N}$ )

$$D^k := \frac{\partial^{\|k\|_1}}{\partial^{k_1} x_1 \cdots \partial^{k_d} x_d},$$

and let  $\lfloor a \rfloor$  be the greatest integer smaller than  $a > 0$ . For  $\alpha > 0$ , a bounded set  $\mathcal{Z} \subset \mathbb{R}^d$  ( $d \in \mathbb{N}$ ) and  $M > 0$ , we define  $C_M^\alpha(\mathcal{Z})$  to be the space of all continuous functions  $f: \mathcal{Z} \rightarrow \mathbb{R}$  with

$$\max_{\|k\|_1 \leq \alpha} \sup_{x \in \mathcal{Z}} |D^k f(x)| + \max_{\|k\|_1 = \lfloor \alpha \rfloor} \sup_{x, y} \frac{|D^k f(x) - D^k f(y)|}{\|x - y\|_2^{\alpha - \lfloor \alpha \rfloor}} \leq M.$$

where the suprema in the second term are taken over all  $x, y$  in the interior of  $\mathcal{Z}$  with  $x \neq y$ . Furthermore, let

$$\mathcal{Z}^1 := \left\{ y \in \mathbb{R}^d: \|x - y\|_2 < 1 \text{ for some } x \in \mathcal{Z} \right\}.$$

**Lemma 5.4.** *Let  $\mathcal{P} = \{A_1, A_2, \dots\}$  be a partition of  $\mathbb{R}^d$  into bounded, convex sets with non-empty interior, and let  $\mathcal{F}$  be a class of functions  $f: \mathbb{R}^d \rightarrow \mathbb{R}$  such that the restrictions  $\mathcal{F}|_{A_j}$  belong to  $C_{M_j}^\alpha(A_j)$  for all  $j \in \mathbb{N}$ . Then there exists a constant  $K$ , depending only on  $\alpha$ ,  $V$ ,  $r$  and  $d$  such that*

$$\log N_{[]}(\varepsilon, \mathcal{F}, L^r(\mathbf{Q})) \leq K \left( \frac{1}{\varepsilon} \right)^V \left( \sum_{j=1}^{\infty} \lambda(A_j^1)^{r/(V+r)} M_j^{Vr/(V+r)} \mathbf{Q}(A_j)^{V/(V+r)} \right)^{(V+r)/r} \quad (5.8)$$

for every  $\varepsilon > 0$ ,  $V \geq d/\alpha$ , and probability measure  $\mathbf{Q}$ .

**Proof.** [vdVW13, Corollary 2.7.4] □

The next lemma gives sufficient conditions on the regularity of the inverse propensity score function.

**Lemma 5.5.** *Let  $(\mathcal{P}_N)$  denote a sequence of cubic partitions  $\mathcal{P}_N = \{A_{N,1}, A_{N,2}, \dots\}$  of  $\mathbb{R}^d$  with decreasing width  $(h_N) \subset (0, 1]$  such that  $h_N \rightarrow 0$  for  $N \rightarrow \infty$ . Furthermore, assume that there exists  $\alpha > d/2$ , where  $\mathcal{X} \subseteq \mathbb{R}^d$ , such that for  $V := d/\alpha$  and for all  $(j, N) \in \mathbb{N}^2$  there exists  $M_{N,j} \geq 1$  such that*

$$\frac{1}{\pi(\cdot)} \in C_{M_{N,j}}^\alpha(A_{N,j}) \quad \text{and} \quad \sum_{j=1}^{\infty} M_{N,j}^{2V/(V+2)} \mathbf{P}[X \in A_{N,j}]^{V/(V+2)} \lesssim 1. \quad (5.9)$$

*Then for any decreasing sequence  $(\varepsilon_N)$  with  $\varepsilon_N \rightarrow 0$  for  $N \rightarrow \infty$ , there exists a sequence of (measurable) function classes  $(\mathcal{F}_N)$  with envelope functions  $(F_N)$ , satisfying for some  $k < 2$*

$$\|F_N\|_{L^2(\mathbf{P})} \leq \varepsilon_N \quad \text{and} \quad \log N_{[]}(\varepsilon, \mathcal{F}_N, L_2(\mathbf{P}_X)) \lesssim \left(\frac{1}{\varepsilon}\right)^k \quad \text{for all } N \in \mathbb{N},$$

*such that for all  $N \in \mathbb{N}$  the paths of the stochastic process*

$$\mathbf{1} \left\{ \sup_{y \in A_N(x)} \left| w_0^\dagger(y) - \frac{1}{\pi(y)} \right| \leq \varepsilon_N \right\} \left( w_0^\dagger(x) - \frac{1}{\pi(x)} \right) \cdot \mathbf{1} \bigcup_{k=1}^N \{x = X_k\}. \quad (5.10)$$

*are contained in  $\mathcal{F}_N$ .*

**Proof.** We want to employ Lemma 5.4. To do this, the crucial observation is that by Lemma 3.8.(ii)

the paths  $w^\dagger(\cdot)(\omega)$  are constant on each cell  $A_N \in \mathcal{P}_N$  for all  $\omega \in \Omega$ .

Thus, the regularity of a path of (5.10) on each cell  $A_N \in \mathcal{P}_N$  is decided by  $1/\pi(\cdot)$ . Indeed, a path of (5.10) is either 0 if the threshold of  $\varepsilon_N$  is exceeded somewhere in the cell, or has the form constant-minus-smooth-function. In any case, it is continuous and bounded by  $\varepsilon_N$ . All its derivatives are 0 (if the threshold is exceeded) or are governed by  $1/\pi(\cdot)$ . Thus, it follows from (5.9)

$$(5.10)(\cdot)(\omega) \in C_{M_{N,j}}^\alpha(A_{N,j}) \quad \text{and} \quad \sum_{j=1}^{\infty} M_{N,j}^{2V/(V+2)} \mathbf{P}[X \in A_{N,j}]^{V/(V+2)} \lesssim 1. \quad (5.11)$$

To bound the right-hand-side in (5.8) we note that  $\lambda(A_{N,j}) = h_N^d$  and thus  $\lambda(A_{N,j}^1) \lesssim 1$  for all  $(j, N) \in \mathbb{N}^2$ . Thus

$$\sum_{j=1}^{\infty} \lambda(A_{N,j}^1)^{2/(V+2)} M_{N,j}^{2V/(V+2)} \mathbf{P}[X \in A_{N,j}]^{V/(V+2)} \lesssim 1.$$

## 5 Convergence of the Weighted Mean

(5.10)( $\cdot$ )( $\omega$ )  $\in \mathcal{F}_N$ , where  $\mathcal{F}_N$  restricted to  $A_{N,j}$  is  $C_{M_{N,j}}^\alpha(A_{N,j})$  and satisfies the requirements of Lemma 5.4. Since  $V = d/\alpha \in (0, 2)$  by  $\alpha > d/2$ , applying Lemma 5.4 finishes the proof.  $\square$

**Remark.** Note, that we only get  $L^2(\mathbf{P}_X)$  bracketing numbers in this way. If we assume, that all functions in  $\mathcal{F}_N$  are independent of  $(T, Y)$  we readily obtain  $L^2(\mathbf{P})$  bracketing numbers. Note, that  $w^\dagger(X)$  and  $1/\pi(X)$  are independent of  $(T, Y)$ .  $\diamond$

Next, we show, that a finite covariate space always meets the requirements of Lemma 5.5 - and that a continuous distribution of  $X$  never does so.

**Lemma 5.6.** *Consider the covariate space  $\mathcal{X}$ .*

(i) *If  $\#\mathcal{X} < \infty$ , that is,  $X$  can take only finitely many values with positive probability, then*

$$\sum_{j=1}^{\infty} M_{N,j}^{2V/(V+2)} \mathbf{P}[X \in A_{N,j}]^{V/(V+2)} \lesssim 1.$$

(ii) *If  $X$  is continuously distributed, then*

$$\sum_{j=1}^{\infty} \mathbf{P}[X \in A_{N,j}]^{V/(V+2)} \rightarrow \infty \quad \text{for } N \rightarrow \infty.$$

**Proof.** Assume  $\#\mathcal{X} < \infty$ , that is,  $X$  can take only finitely many values with positive probability. We write

$$J_N := \{j \in \mathbb{N} : \mathbf{P}[X \in A_{N,j}] > 0\}.$$

It holds  $\#J_N \leq \#\mathcal{X} < \infty$ . Thus, the following maximum is attained

$$\max_{j \in J_N} M_{N,j} =: M_N^*.$$

But the partitions increasingly better fit the support of  $X$ . Thus  $M_N^*$  is decreasing in  $N$ , that is,  $\infty > M_1^* \geq M_N^*$ . It follows

$$\sum_{j=1}^{\infty} M_{N,j}^{2V/(V+2)} \mathbf{P}[X \in A_{N,j}]^{V/(V+2)} \leq (M_1^*)^{2V/(V+2)} \cdot \#J_N \lesssim 1.$$

Now let  $f_X$  be the probability density of  $X$ . Then there exists a compact set  $K \subset \mathcal{X} \subset \mathbb{R}^d$ , such that  $\inf_{x \in K} f_X(x) > 0$ . Since  $\mathcal{P}_N$  are cubic partitions, it holds for

$$I_N := \{i \in \mathbb{N} : A_{N,i} \subset K\} \quad \text{that} \quad \bigcup_{i \in I_N} A_{N,i} \nearrow K.$$

Thus

$$\begin{aligned}
\sum_{i=1}^{\infty} \mathbf{P}[X \in A_{N,i}]^{V/(V+2)} &\geq \sum_{i \in I_N} \mathbf{P}[X \in A_{N,i}]^{V/(V+2)} \\
&\geq \inf_{x \in K} f_X(x)^{V/(V+2)} \cdot h_N^{d \cdot (V/(V+2)-1)} \sum_{i \in I_N} \lambda(A_{N,i}) \\
&\rightarrow \infty.
\end{aligned}$$

This follows from  $\sum_{i \in I_N} \lambda(A_{N,i}) \rightarrow \lambda(K) > 0$ ,  $\inf_{x \in K} f_X(x) > 0$ ,  $V/(V+2) - 1 < 0$  and  $h_N \rightarrow 0$ .  $\square$

### 5.1.3 Maximal Inequality

In our application we need concentration inequalities for  $\|\mathbb{G}_n\|_{\mathcal{F}}^*$ . One easy way to obtain this is, to use a maximal inequality (see Theorem 5.1) to control the expectation, together with Markov's inequality. There are also Bernstein-like inequalities for empirical processes (see [vdVW13, §2.14.2]).

**Theorem 5.1.** (Maximal inequality) *For any class  $\mathcal{F}$  of measurable functions with envelope function  $F$ ,*

$$\mathbf{E}^* \|\mathbb{G}_n\|_{\mathcal{F}} \lesssim J_{[]}(\|F\|_{L^2(\mathbf{P})}, \mathcal{F}, L^2(\mathbf{P})).$$

**Proof.** [vdV00, Corollary 19.35]  $\square$

**Lemma 5.7.** *Let  $(\mathcal{H}_N)$  be a sequence of measurable function classes with envelope functions  $(H_N)$ . If*

$$J_{[]}(\|H_N\|_{L^2(\mathbf{P})}, \mathcal{H}_N, L^2(\mathbf{P})) \rightarrow 0 \quad \text{for } N \rightarrow \infty,$$

*it holds  $\|\mathbb{G}_N\|_{\mathcal{H}_N}^* \xrightarrow{\mathbf{P}} 0$ .*

**Proof.** By Markov's inequality and Theorem 5.1 it holds for all  $\varepsilon > 0$

$$\begin{aligned}
\mathbf{P}[\|\mathbb{G}_N\|_{\mathcal{H}_N}^* \geq \varepsilon] &\leq \varepsilon^{-1} \mathbf{E}[\|\mathbb{G}_N\|_{\mathcal{H}_N}^*] = \varepsilon^{-1} \mathbf{E}^*[\|\mathbb{G}_N\|_{\mathcal{H}_N}] \\
&\lesssim \varepsilon^{-1} J_{[]}(\|H_N\|_{L^2(\mathbf{P})}, \mathcal{H}_N, L^2(\mathbf{P})) \\
&\rightarrow 0 \quad \text{for } N \rightarrow \infty.
\end{aligned}$$

$\square$

**Lemma 5.8.** *Let  $(\varepsilon_N) \subset (0, 1]$  be a decreasing sequence with  $\varepsilon_N \rightarrow 0$  for  $N \rightarrow \infty$  and  $(\mathcal{F}_N)$  a sequence of (measurable) function classes with envelope functions  $(F_N)$ , satisfying for some  $k < 2$*

$$\|F_N\|_{L^2(\mathbf{P})} \leq \varepsilon_N \quad \text{and} \quad \log N_{[]}(\varepsilon, \mathcal{F}_N, L_2(\mathbf{P}_X)) \lesssim \left(\frac{1}{\varepsilon}\right)^k \quad \text{for all } N \in \mathbb{N}.$$

*Then*

$$J_{[]}(\|F_N\|_{L^2(\mathbf{P})}, \mathcal{F}_N \cdot \mathcal{F}, L_2(\mathbf{P})) \rightarrow 0 \quad \text{and} \quad \|\mathbb{G}_N\|_{\mathcal{F}_N \cdot \mathcal{F}}^* \xrightarrow{\mathbf{P}} 0 \quad \text{for } N \rightarrow \infty,$$

*where  $\mathcal{F}$  is defined in (5.5).*

**Proof.** By assumption and Lemma 5.2 it holds for some  $k < 2$

$$\|F_N\|_{L^2(\mathbf{P})} \leq \varepsilon_N \quad \text{and} \quad \log N_{[]}(\varepsilon, \mathcal{F}_N, L_2(\mathbf{P})) \lesssim \left(\frac{1}{\varepsilon}\right)^k \quad \text{for all } N \in \mathbb{N},$$

and

$$N_{[]}(\varepsilon, \mathcal{F}, L_2(\mathbf{P})) \lesssim \left(\frac{1}{\varepsilon}\right)^2 \quad \text{for all } \varepsilon > 0.$$

Since  $\mathcal{F}_N$  and  $\mathcal{F}$  have envelope functions smaller 1, we can apply Lemma 5.1 to get

$$\log N_{[]}(\varepsilon, \mathcal{F}_N \cdot \mathcal{F}, L_2(\mathbf{P})) \lesssim \left(\frac{1}{\varepsilon}\right)^k + \log(1/\varepsilon) \lesssim \left(\frac{1}{\varepsilon}\right)^k \quad \text{for all } \varepsilon > 0.$$

Since  $k/2 \in (0, 1)$  it holds

$$\begin{aligned} J_{[]}(\|F_N\|_{L^2(\mathbf{P})}, \mathcal{F}_N \cdot \mathcal{F}, L_2(\mathbf{P})) &= \int_0^{\|F_N\|_{L^2(\mathbf{P})}} \sqrt{\log N_{[]}(\varepsilon, \mathcal{F}_N \cdot \mathcal{F}, L_2(\mathbf{P}))} d\varepsilon \\ &\lesssim \int_0^{\varepsilon_N} \left(\frac{1}{\varepsilon}\right)^{k/2} d\varepsilon \\ &= \frac{\varepsilon_N^{1-k/2}}{1-k/2} \rightarrow 0 \quad \text{for } N \rightarrow \infty. \end{aligned}$$

The second statement follows from Lemma 5.7 for  $\mathcal{H}_N := \mathcal{F}_N \cdot \mathcal{F}$  and  $H_N := F_N$ .  $\square$

#### 5.1.4 Donsker's Theorem

There is a powerful theorem - a central limit theorem for  $\mathbb{G}_N$  uniform in  $\mathcal{F}$  - that we now introduce.

**Definition 5.1.** We call a class  $\mathcal{F}$  of measurable functions  $\mathbf{P}$ -Donsker if the sequence of processes  $\{\mathbb{G}_N f : f \in \mathcal{F}\}$  converges in  $l^\infty(\mathcal{F})$  to a tight limit process.

**Theorem 5.2.** Every class  $\mathcal{F}$  of measurable functions with

$$J_{[]} (1, \mathcal{F}, L_2(\mathbf{P})) < \infty$$

is  $\mathbf{P}$ -Donsker. Furthermore, the sequence of processes  $\{\mathbb{G}_N f : f \in \mathcal{F}\}$  converges in  $l^\infty(\mathcal{F})$  to a Gaussian process with mean 0 and covariance function given by

$$\mathbf{Cov}(f, g) := \mathbf{E}[fg] - \mathbf{E}[f]\mathbf{E}[g].$$

**Proof.** [vdV00, Theorem 19.5] □

**Lemma 5.9.** The function class  $\mathcal{F}$  defined in (5.5) is  $\mathbf{P}$ -Donsker.

**Proof.** By Theorem 5.2 it suffices to show that the bracketing integral is finite. By Lemma 5.2 it holds

$$N_{[]}(\varepsilon, \mathcal{F}, L^2(\mathbf{P})) \lesssim \left(\frac{1}{\varepsilon}\right)^2 \quad \text{for all } \varepsilon > 0.$$

Thus

$$J_{[]} (1, \mathcal{F}, L_2(\mathbf{P})) \lesssim \int_0^1 \sqrt{-\log(\varepsilon)} d\varepsilon \leq 1 + \int_0^1 -\log(\varepsilon) d\varepsilon = 2.$$

□

### 5.1.5 Propensity Score Weights

The next lemma shows what effect the **propensity score weights**  $T/\pi(X)$  have on other functions.

**Lemma 5.10.** Let  $g_1 : \mathcal{X} \rightarrow \mathbb{R}$  and  $g_2 : \mathcal{Y} \rightarrow \mathbb{R}$  be measurable functions.

(i) It holds

$$\mathbf{E} \left[ \frac{T}{\pi(X)} g_1(X) \right] = \mathbf{E} [g_1(X)].$$



(ii) If Assumption ?? holds true, then

$$\mathbf{E} \left[ \frac{T}{\pi(X)} g_2(Y(T)) \right] = \mathbf{E} [f(Y(1))] .$$

## 5.2 Main Result

**Theorem 5.3.** *The stochastic process*

$$\sqrt{N} \left( \frac{1}{N} \sum_{i=1}^N T_i \cdot w_0^\dagger(X_i) \cdot \mathbf{1} \{Y_i \leq z\} - F_{Y(1)}(z) \right)_{z \in \mathbb{R}} \quad (5.12)$$

*converges in  $l^\infty(\mathbb{R})$  to a Gaussian process with mean 0 and covariance function satisfying for all  $z_1, z_2 \in \mathbb{R}$*

$$\begin{aligned} & \mathbf{Cov}(z_1, z_2) \\ &= \mathbf{E} \left[ \frac{F_{Y(1)}(z_1 \wedge z_2 | X)}{\pi(X)} - \frac{1 - \pi(X)}{\pi(X)} F_{Y(1)}(z_1 | X) \cdot F_{Y(1)}(z_2 | X) \right] \\ & \quad - F_{Y(1)}(z_1) \cdot F_{Y(1)}(z_2) . \end{aligned} \quad (5.13)$$

## 5.3 Error Decomposition

**Lemma 5.11.** *It holds*

$$\sqrt{N} \left( \frac{1}{N} \sum_{i=1}^N T_i \cdot w_0^\dagger(X_i) \cdot \mathbf{1} \{Y_i \leq z\} - F_{Y(1)}(z) \right)_{z \in \mathbb{R}} = R_1 + R_2 + R_3 + R_4 \quad (5.14)$$

with

$$\begin{aligned}
R_1 &:= \sqrt{N} \sum_{k=1}^N \left[ \frac{1}{N} \left( \sum_{i=1}^N T_i \cdot w_0^\dagger(X_i) \cdot B_k(X_i) - \sum_{i=1}^N B_k(X_i) \right) \cdot F_{Y(1)}(z|X_k) \right]_{z \in \mathbb{R}}, \\
R_2 &:= \sqrt{N} \sum_{i=1}^N \frac{1}{N} \left[ \left( T_i \cdot w_0^\dagger(X_i) - 1 \right) \left( F_{Y(1)}(z|X_i) - \sum_{k=1}^N B_k(X_i) \cdot F_{Y(1)}(z|X_k) \right) \right]_{z \in \mathbb{R}}, \\
R_3 &:= \sqrt{N} \left( \frac{1}{N} \sum_{i=1}^N \left[ T_i \cdot \left( w_0^\dagger(X_i) - \frac{1}{\pi(X_i)} \right) \cdot (\mathbf{1}\{Y_i \leq z\} - F_{Y(1)}(z|X_i)) \right] \right)_{z \in \mathbb{R}}, \\
R_4 &:= \sqrt{N} \left( \frac{1}{N} \sum_{i=1}^N \frac{T_i}{\pi(X_i)} (\mathbf{1}\{Y_i \leq z\} - F_{Y(1)}(z|X_i)) + (F_{Y(1)}(z|X_i) - F_{Y(1)}(z)) \right)_{z \in \mathbb{R}}.
\end{aligned}$$

**Proof.** We fix  $z \in \mathbb{R}$ . It holds

$$\begin{aligned}
 & \frac{1}{N} \sum_{i=1}^N w_0^\dagger(X_i) \cdot T_i \cdot \mathbf{1}\{Y_i \leq z\} \\
 &= \frac{1}{N} \sum_{i=1}^N \left( w_0^\dagger(X_i) - \frac{1}{\pi(X_i)} \right) T_i \cdot \mathbf{1}\{Y_i \leq z\} \\
 &+ \frac{1}{N} \sum_{i=1}^N \frac{T_i}{\pi(X_i)} \mathbf{1}\{Y_i \leq z\} \\
 &= \frac{1}{N} \sum_{i=1}^N \left( w_0^\dagger(X_i) - \frac{1}{\pi(X_i)} \right) T_i (\mathbf{1}\{Y_i \leq z\} - F_{Y(1)}(z|X_i)) \\
 &+ \frac{1}{N} \sum_{i=1}^N \frac{T_i}{\pi(X_i)} (\mathbf{1}\{Y_i \leq z\} - F_{Y(1)}(z|X_i)) \\
 &+ \frac{1}{N} \sum_{i=1}^N w_0^\dagger(X_i) \cdot T_i \cdot F_{Y(1)}(z|X_i) \\
 &= R_3(z)/\sqrt{N} \\
 &+ \frac{1}{N} \sum_{i=1}^N \frac{T_i}{\pi(X_i)} (\mathbf{1}\{Y_i \leq z\} - F_{Y(1)}(z|X_i)) + (F_{Y(1)}(z|X_i) - F_{Y(1)}(z)) \\
 &+ \frac{1}{N} \sum_{i=1}^N \left( w_0^\dagger(X_i) \cdot T_i - 1 \right) F_{Y(1)}(z|X_i) \\
 &+ F_{Y(1)}(z) \\
 &= R_3(z)/\sqrt{N} \\
 &+ R_4(z)/\sqrt{N} \\
 &+ \frac{1}{N} \sum_{i=1}^N \left( w_0^\dagger(X_i) \cdot T_i - 1 \right) \left( F_{Y(1)}(z|X_i) - \sum_{k=1}^N B_k(X_i) \cdot F_{Y(1)}(z|X_k) \right) \\
 &+ \frac{1}{N} \sum_{i=1}^N \left( w_0^\dagger(X_i) \cdot T_i - 1 \right) \sum_{k=1}^N B_k(X_i) \cdot F_{Y(1)}(z|X_k) \\
 &+ F_{Y(1)}(z) \\
 &= R_3(z)/\sqrt{N} \\
 &+ R_4(z)/\sqrt{N} \\
 &+ R_2(z)/\sqrt{N} \\
 &+ \sum_{k=1}^N \frac{1}{N} \sum_{i=1}^N \left( w_0^\dagger(X_i) \cdot T_i B_k(X_i) - B_k(X_i) \right) \cdot F_{Y(1)}(z|X_k) \\
 &+ F_{Y(1)}(z) \\
 &= (R_3(z) + R_4(z) + R_2(z) + R_1(z))/\sqrt{N} + F_{Y(1)}(z).
 \end{aligned}$$

This holds for all  $z \in \mathbb{R}$ . Multiplying with  $\sqrt{N}$  yields the result.  $\square$

## 5.4 Analysis of the Error Terms

### 5.4.1 Analysis of $R_1$

**Lemma 5.12.** *Let  $\sqrt{N} \|\delta\|_1 \xrightarrow{\mathbf{P}} 0$ . Then it holds  $\sup_{z \in \mathbb{R}} |R_1(z)| \xrightarrow{\mathbf{P}} 0$ .*

**Proof.** By Theorem 3.2  $(w_i^\dagger(X_i))$  satisfy the box constraints of Problem 1 (in the form with the  $T_i$  instead of  $n$ ). Thus

$$\begin{aligned} \sup_{z \in \mathbb{R}} |R_1(z)| &= \sqrt{N} \sup_{z \in \mathbb{R}} \sum_{k=1}^N \left[ \frac{1}{N} \left( \sum_{i=1}^N T_i \cdot w_i^\dagger(X_i) \cdot B_k(X_i) - \sum_{i=1}^N B_k(X_i) \right) \cdot F_{Y(1)}(z|X_k) \right] \\ &\leq \sqrt{N} \sum_{k=1}^N \left| \frac{1}{N} \left( \sum_{i=1}^N T_i \cdot w_i^\dagger(X_i) \cdot B_k(X_i) - \sum_{i=1}^N B_k(X_i) \right) \right| \cdot \sup_{z \in \mathbb{R}} F_{Y(1)}(z|X_k) \\ &\leq \sqrt{N} \|\delta\|_1 \end{aligned} \tag{5.15}$$

The last inequality is due to  $F_{Y(1)} \in [0, 1]$ . Since we assume  $\sqrt{N} \|\delta\|_1 \xrightarrow{\mathbf{P}} 0$  it holds  $\sup_{z \in \mathbb{R}} |R_1(z)| \xrightarrow{\mathbf{P}} 0$ .  $\square$

**Remark.** We want to comment on the box constraints of Problem 1, that is,

$$\left| \frac{1}{N} \left( \sum_{i=1}^n w_i^\dagger(X_i) B_k(X_i) - \sum_{i=1}^N B_k(X_i) \right) \right| \leq \delta_k \quad \text{for all } k \in \{1, \dots, N\}.$$

Note, that the first sum goes over  $\{1, \dots, n\}$  while the second sum goes over  $\{1, \dots, N\}$ . A second, equivalent version of the constraints is

$$\left| \frac{1}{N} \left( \sum_{i=1}^N T_i w_i^\dagger(X_i) B_k(X_i) - \sum_{i=1}^N B_k(X_i) \right) \right| \leq \delta_k \quad \text{for all } k \in \{1, \dots, N\}.$$

Now both sums go over  $\{1, \dots, N\}$  and the indicator of treatment  $T_i$  takes care that in the first sum only the terms with  $i \leq n$  are effective. Having this flexibility with the versions helps. I regard the first version as suitable for non-probabilistic computations, although  $n$  is of course a random variable. On the other hand, the second version is more honest, exactly telling the dependence on the indicator of treatment. This version is useful in probabilistic computations.

## 5 Convergence of the Weighted Mean

Also we want to comment on the assumption on  $\|\delta\|$ . Playing around with norm equivalences we discover that  $\sqrt{N}\|\delta\|_1 \xrightarrow{\mathbf{P}} 0$  for  $N \rightarrow \infty$  is the weakest (natural) assumption to control  $R_1$ . Indeed, other ways to continue the second row in (5.15) are

$$(\dots) \leq \sqrt{N}\|\delta\|_2 \left( \sum_{k=1}^N \left( \sup_{z \in \mathbb{R}} F_{Y(1)}(z|X_k) \right)^2 \right)^{1/2} \leq N\|\delta\|_2,$$

by the Cauchy-Schwarz inequality and  $F_{Y(1)} \in [0, 1]$ , or

$$(\dots) \leq \sqrt{N}\|\delta\|_\infty \sum_{k=1}^N \sup_{z \in \mathbb{R}} F_{Y(1)}(z|X_k) \leq N^{3/2}\|\delta\|_\infty.$$

Since  $\delta \in \mathbb{R}^N$ , however, it holds

$$\sqrt{N}\|\delta\|_1 \leq N\|\delta\|_2 \leq N^{3/2}\|\delta\|_\infty.$$

With hindsight, the assumption  $\sqrt{N}\|\delta\|_1 \xrightarrow{\mathbf{P}} 0$  for  $N \rightarrow \infty$  also suffices to control the second (or first) occurrence of a term, that we control by assumptions on  $\|\delta\|$ . This is the term  $I_2$  in (4.4), where we estimate

$$\langle \delta, |\Delta| \rangle = \sum_{k=1}^N \delta_k |\Delta_k| \leq \|\delta\|_1 \|\Delta\|_\infty \leq \|\delta\|_1 \|\Delta\|_2 \leq \|\delta\|_1 \varepsilon \xrightarrow{\mathbf{P}} 0 \quad \text{for } N \rightarrow \infty.$$

◇

### 5.4.2 Analysis of $R_2$

**Lemma 5.13.** *Assume*

$$\sqrt{N} \sup_{z \in \mathbb{R}} \omega \left( F_{Y(1)}(z|\cdot), h_N^d \right) \rightarrow 0 \quad \text{for } N \rightarrow \infty.$$

*Then  $\sup_{z \in \mathbb{R}} |R_2(z)| \xrightarrow{\mathbf{P}} 0$ .*

**Proof.**

$$\begin{aligned} \sup_{z \in \mathbb{R}} |R_2(z)| &\leq \sqrt{N} \sup_{z \in \mathbb{R}} \max_{i \in \{1, \dots, N\}} \sum_{k=1}^N |B_k(X_i, X_1, \dots, X_N) \cdot F_{Y(1)}(z|X_k) - F_{Y(1)}(z|X_i)| \\ &\quad \cdot \frac{1}{N} \sum_{i=1}^N |T_i \cdot w_i^\dagger(X_i) - 1| \end{aligned}$$

Note, that by Theorem 3.2.(i)-(ii) it holds

$$\frac{1}{N} \sum_{i=1}^N |T_i \cdot w_i^\dagger(X_i) - 1| \leq 1 + \frac{1}{N} \sum_{i=1}^N T_i \cdot w_i^\dagger(X_i) = 2.$$

The statement follows from Lemma 3.7.(ii) □

**Remark.** In the original paper [WZ19] the authors derive concrete learning rates for the weights and employ them in bounding this term. They obtain a multiplied learning rate, which is sufficiently fast. Their approach, however, calls for concrete learning rates of the weights. Arguably, the process of deriving such rates is the most complicated part of the paper. I found out, that we don't need concrete rates for the weights. Consistency of the weights is enough and gives us an (arbitrarily slow but sufficient) learning rate to establish the results. We don't even need rates for the weights to control  $R_2$ . They only play a role in bounding  $R_3$ .  $\diamond$

### 5.4.3 Analysis of $R_3$

**Lemma 5.14.** *It holds  $\sup_{z \in \mathbb{R}} |R_3(z)| \xrightarrow{\mathbf{P}} 0$ .*

**Proof.** Let  $z \in \mathbb{R}$ . By Lemma ?? it holds

$$\begin{aligned} f_z(T, X, Y(T)) &\in L^1(\mathbf{P}), \\ f_z(T, X, Y(T)) &\perp D_N, \\ \mathbf{E}[f_z(T, X, Y(T))|X] &= 0. \end{aligned}$$

Thus, it follows from Lemma 3.12

$$\mathbf{E} \left[ w_0^\dagger(X) \cdot f_z(T, X, Y(T)) \right] = 0.$$

Since

$$\mathbf{E} \left[ \frac{T}{\pi(X)} f_z(T, X, Y(T)) \right] = \mathbf{E} \left[ \frac{T}{\pi(X)} (\mathbf{1}_{\{Y(T) \leq z\}} - F_{Y(1)}(z|X)) \right] = 0$$

by Lemma 5.10, it follows

$$\mathbf{E} \left[ \left( w_0^\dagger(X) - \frac{1}{\pi(X)} \right) \cdot f_z(T, X, Y(T)) \right] = 0.$$

But then

$$R_3(z) = \frac{1}{\sqrt{N}} \sum_{i=1}^N \left[ \left( w_0^\dagger(X_i) - \frac{1}{\pi(X_i)} \right) T_i (\mathbf{1}_{\{Y_i \leq z\}} - F_{Y(1)}(z|X_i)) \right] = \mathbb{G}_N \left( \left( w_0^\dagger - \frac{1}{\pi} \right) \cdot f_z \right).$$

Let  $g^\dagger$  denote the stochastic process (5.7), that is,

$$g^\dagger(x) := \mathbf{1} \left\{ \sup_{y \in A_N(x)} \left| w_0^\dagger(y) - \frac{1}{\pi(y)} \right| \leq \varepsilon_N \right\} \left( w_0^\dagger(x) - \frac{1}{\pi(x)} \right) \cdot \mathbf{1} \bigcup_{k=1}^n \{x = X_k\}$$

## 5 Convergence of the Weighted Mean

for all  $x \in \mathbb{R}^d$ . If

$$\left| w_0^\dagger(X_i) - \frac{1}{\pi(X_i)} \right| \leq \varepsilon_N \quad \text{for all } i \in \{1, \dots, n\} \quad (5.16)$$

it holds for all  $i \in \{1, \dots, N\}$

$$g^\dagger(X_i) \cdot f_z(T_i, X_i, Y_i(T_i)) = \left( w_0^\dagger(X_i) - \frac{1}{\pi(X_i)} \right) f_z(T_i, X_i, Y_i(T_i)).$$

Thus, if (5.16) holds, it follows

$$R_3(z) = \mathbb{G}_N(g^\dagger \cdot f_z).$$

It follows

$$\begin{aligned} \mathbf{P} \left[ \sup_{z \in \mathbb{R}} |R_3(z)| \geq \varepsilon \right] &\leq \mathbf{P} \left[ \sup_{z \in \mathbb{R}} |R_3(z)| \geq \varepsilon \text{ and } |w_0^\dagger(X_i) - 1/\pi(X_i)| \leq \varepsilon_N \text{ for all } i \in \{1, \dots, n\} \right] \\ &\quad + \mathbf{P} \left[ |w_0^\dagger(X_i) - 1/\pi(X_i)| > \varepsilon_N \text{ for some } i \in \{1, \dots, n\} \right] \\ &\leq \mathbf{P} \left[ \|\mathbb{G}_N\|_{\mathcal{F}_N, \mathcal{F}}^* \geq \varepsilon \right] + \mathbf{P} \left[ \max_{i \in \{1, \dots, n\}} |w_0^\dagger(X_i) - 1/\pi(X_i)| > \varepsilon_N \right] \\ &\rightarrow 0. \end{aligned}$$

The convergence of the first term follows from Lemma 5.8. The convergence of the second term follows from Theorem 4.2. □

### 5.4.4 Analysis of $R_4$

**Lemma 5.15.** *Let  $1/\pi(X) \in L^2(\mathbf{P})$ .  $R_4$  converges in  $l^\infty(\mathbb{R})$  to a Gaussian process with mean 0 and covariance*

$$\begin{aligned} &\text{Cov}(z_1, z_2) \\ &= \mathbf{E} \left[ \frac{F_{Y(1)}(z_1 \wedge z_2 | X)}{\pi(X)} - \frac{1 - \pi(X)}{\pi(X)} F_{Y(1)}(z_1 | X) \cdot F_{Y(1)}(z_2 | X) \right] - F_{Y(1)}(z_1) \cdot F_{Y(1)}(z_2) \end{aligned}$$

**Proof.** By Lemma ?? it follows

$$\mathbf{E} \left[ \frac{f_z(T, X, Y(T))}{\pi(X)} + F_{Y(1)}(z | X) - F_{Y(1)}(z) \right] = \mathbf{E} \left[ \frac{1}{\pi(X)} \mathbf{E} [f_z(T, X, Y(T)) | X] \right] = 0.$$

Thus

$$\begin{aligned}
R_4(z) &= \frac{1}{\sqrt{N}} \sum_{i=1}^N \frac{T_i}{\pi(X_i)} (\mathbf{1}\{Y_i \leq z\} - F_{Y(1)}(z|X_i)) + (F_{Y(1)}(z|X_i) - F_{Y(1)}(z)) \\
&= \frac{1}{\sqrt{N}} \sum_{i=1}^N \frac{f_z(T_i, X_i, Y_i)}{\pi(X_i)} + (F_{Y(1)}(z|X_i) - F_{Y(1)}(z)) \\
&= \mathbb{G}_N \left( \frac{f_z}{\pi(\cdot)} + F_{Y(1)}(z|\cdot) - F_{Y(1)}(z) \right).
\end{aligned}$$

By Lemma 5.2 it holds

$$\begin{aligned}
&\log N_{[]}(\varepsilon, \mathcal{G}, L^2(\mathbf{P})) \\
&\lesssim \log \left( \frac{1 + \|1/\pi(X)\|_{L^2(\mathbf{P})}}{\varepsilon} \right) \lesssim \frac{1 + \|1/\pi(X)\|_{L^2(\mathbf{P})}}{\varepsilon} \quad \text{for all } \varepsilon \in (0, 1).
\end{aligned}$$

Thus

$$J_{[]} (1, \mathcal{G}, L^2(\mathbf{P})) \lesssim \int_0^1 \sqrt{\frac{1 + \|1/\pi(X)\|_{L^2(\mathbf{P})}}{\varepsilon}} d\varepsilon \lesssim 1 + \|1/\pi(X)\|_{L^2(\mathbf{P})} < \infty.$$

But then  $\mathcal{G}$  is  $\mathbf{P}$ -Donsker. By the Donsker Theorem [vdV00, Theorem 19.5] the process  $R_4$  converges in  $l^\infty(\mathbb{R})$  to a Gaussian process, called  $\mathbf{P}$ -Brownian bridge, with mean 0. We now calculate the covariance of the limiting process.

### Covariance

$$\begin{aligned}
&\mathbf{E} \left[ \left( f_{1/\pi}^{z_1} + F_{Y(1)}(z_1|X) - F_{Y(1)}(z_1) \right) \left( f_{1/\pi}^{z_2} + F_{Y(1)}(z_2|X) - F_{Y(1)}(z_2) \right) \right] \\
&= \mathbf{E} \left[ f_{1/\pi}^{z_1} \cdot f_{1/\pi}^{z_2} \right] \\
&\quad + \mathbf{E} \left[ f_{1/\pi}^{z_1} (F_{Y(1)}(z_2|X) - F_{Y(1)}(z_2)) \right] + \mathbf{E} \left[ f_{1/\pi}^{z_2} (F_{Y(1)}(z_1|X) - F_{Y(1)}(z_1)) \right] \\
&\quad + \mathbf{E} \left[ (F_{Y(1)}(z_1|X) - F_{Y(1)}(z_1)) (F_{Y(1)}(z_2|X) - F_{Y(1)}(z_2)) \right] \\
&=: C_0 + C_1 + C_2 + C_3.
\end{aligned}$$

It holds

$$\begin{aligned}
C_0 &= \mathbf{E} \left[ f_{1/\pi}^{z_1} \cdot f_{1/\pi}^{z_2} \right] \\
&= \mathbf{E} \left[ \frac{1}{\pi(X)} \frac{T}{\pi(X)} (\mathbf{1}\{Y(T) \leq z_1\} - F_{Y(1)}(z_1|X)) (\mathbf{1}\{Y(T) \leq z_2\} - F_{Y(1)}(z_2|X)) \right] \\
&= \mathbf{E} \left[ \frac{1}{\pi(X)} (\mathbf{1}\{Y(1) \leq z_1\} - F_{Y(1)}(z_1|X)) (\mathbf{1}\{Y(1) \leq z_2\} - F_{Y(1)}(z_2|X)) \right] \\
&= \mathbf{E} \left[ \frac{1}{\pi(X)} (F_{Y(1)}(z_1 \wedge z_2|X) - F_{Y(1)}(z_1|X) \cdot F_{Y(1)}(z_2|X)) \right].
\end{aligned}$$



$$\begin{aligned}
C_1 &= \mathbf{E} \left[ f_{1/\pi}^{z_1} (F_{Y(1)}(z_2|X) - F_{Y(1)}(z_2)) \right] \\
&= \mathbf{E} \left[ \frac{T}{\pi(X)} (\mathbf{1}\{Y(T) \leq z_1\} - F_{Y(1)}(z_1|X)) (F_{Y(1)}(z_2|X) - F_{Y(1)}(z_2)) \right] \\
&= \mathbf{E} \left[ (\mathbf{1}\{Y(1) \leq z_1\} - F_{Y(1)}(z_1|X)) (F_{Y(1)}(z_2|X) - F_{Y(1)}(z_2)) \right] \\
&= 0.
\end{aligned}$$

In the same way we see  $C_2 = 0$ .

$$\begin{aligned}
C_3 &= \mathbf{E} \left[ (F_{Y(1)}(z_1|X) - F_{Y(1)}(z_1)) (F_{Y(1)}(z_2|X) - F_{Y(1)}(z_2)) \right] \\
&= \mathbf{E} [F_{Y(1)}(z_1|X) \cdot F_{Y(1)}(z_2|X)] - F_{Y(1)}(z_1) \cdot F_{Y(1)}(z_2).
\end{aligned}$$

Adding up the results gives us (5.13).  $\square$

### 5.4.5 Proof of Theorem 5.3

We have gathered all the results to prove Theorem 5.3.

**Proof.** (*Theorem 5.3*) We connect the statement of the theorem to the error decomposition by Lemma 5.11. By Lemma 5.12, Lemma ??, Lemma 5.14 it follows  $\sup_{z \in \mathbb{R}} |R_i(z)| \xrightarrow{\mathbf{P}} 0$  for  $i = 1, 2, 3$ . Thus, by Slutsky's theorem (cf. [Kle20, Theorem 13.18]) the behaviour of the limiting process is the one of Lemma 5.15.  $\square$

# 6 Discussion and Outlook

## 6.1 Discussion

### 6.1.1 Summary of Assumptions

Here are Assumptions.

## 6.2 Outlook

### 6.2.1 Matching

#### Motivation

The papers [WZ19, WZ23] are closely related. In [WZ19] - the paper this thesis is based on - the authors study weighting methods. In [WZ23] the authors propose (in similar style) a matching framework based a constrained convex optimization.

#### Conjecture

The extensions proposed in this thesis (or parts) can be applied to the matching framework of [WZ23].

#### Ideas/Brainstorming

The constraints in the problem [WZ23, (2.1)] are more complicated. Nevertheless, they rely on the notion of basis function. While in [WZ19] the estimation objective is the expectation of potential outcomes, in [WZ23] it is the average treatment effect. The structure of the proofs is similar - first reveal connection to the inverse propensity score and then employ it in the error analysis of the estimator.

#### Organisation

Get familiar with [WZ19, WZ23]. Point out the differences and similarities. Make the mathematical analysis of [WZ23] rigorous. You can use ideas from this thesis. Try to extend the matching framework - either with ides from this thesis or your own ideas.

### Next Step

Read the introductions of [WZ19, WZ23].

## 6.2.2 Application of the Functional Delta Method

### Motivation

Theorem ?? immediately allows to apply the functional delta method [vdVW13, §3.9], [vdV00, §20]. This readily generates theoretic results for a large class of plug-in estimators. The plug-in estimators have not been tested before in practice.

### Conjecture

A large class of plug-in estimators converges in distribution to a nice limiting process. The estimators work well in practice.

### Ideas/Brainstorming

A plethora of applications of the delta method to estimates of the distribution function are to be found in [vdV00] and [vdVW13]. This includes Quantile estimation [vdV00, §21] [vdVW13, §3.9.21/24], survival analysis via Nelson-Aalen and Kaplan-Meier estimator [vdVW13, §3.9.19/31], Wilcoxon Test [vdVW13, §3.9.4.1], and much more.

### Organisation

Understand the functional delta method. Determine your interests. Pick an example and compute the limiting process. If this is fun and successful try another example - or come up with a new plug-in estimator that works with the functional delta method. Do a simulation study and evaluate the performance of the method in practice. Start with a plain estimate of the distribution function. If this works well, plug-in estimators might as well. Apply the method to real world data.

### Next Step

Read [vdV00, §20].

## 6.2.3 Bootstrapping

### Motivation

A very natural idea is to bootstrap from the weighted distribution  $(w_i \cdot X_i)$ . I discussed this with Jose Zubizarreta, one of the authors of [WZ19, WZ23]. Jose told me that

testing in practice showed promising results. To the best of his and my knowledge the theoretical properties of this particular weighted bootstrap wait to be studied.

### Conjecture

Results similar to [vdV00, Theorem 23.5] holds for the weighted bootstrap.

### Ideas/Brainstorming

A good starting point to become familiar with the asymptotic theory of bootstrap is [vdVW13, §3.6] and [vdV00, §23]. For more details, a good starting point could be [BB95]. The project seems to be challenging - maybe at PHD level.

### Organisation

Understand the mathematical theory of bootstrap. Talk to Jose Zubizarreta about practical results and possible collaboration. Develop ideas based on the existing literature.

### Next Step

Get acquainted with the method of bootstrap by reading the (non-mathematical) introduction [ET94].

## 6.2.4 Non-binary Treatment

### Motivation

In practice, there often exists multiple treatments. For example,  $T \in \{0, 1, 2\}$ ,  $T \in I \subset \mathbb{N}$  or even  $T \in \mathbb{R}$ . There exists a general notion of propensity score [HI05]. There is a need for methods covering this scenarios.

### Conjecture

The framework of [WZ19] can be extended to for non-binary treatment.

### Ideas/Brainstorming

There are already ideas [Tüb20, VGC<sup>+</sup>20]. They try to estimate one set of weights to cover all possible treatments. Jose Zubizarreta, one of the authors of [WZ19], told me,

that he works on a similar (practical) project. I think it's better to compute weights for one treatment at a time. My idea is: For fixed  $t \in \mathcal{T}$  this could be

$$\underset{w_1, \dots, w_n \in \mathbb{R}}{\text{minimize}} \quad \sum_{i=1}^n d_n(t, T_i) \varphi(w_i)$$

subject to the constraints

$$\left| \frac{1}{N} \left( \sum_{i=1}^n w_i \cdot d_n(t, T_i) B_k(X_i) - \sum_{i=1}^N B_k(X_i) \right) \right| \leq \delta_k, \quad k = 1, \dots, N,$$

where

$$d_n(t, s) := \frac{\mathbf{1}_{s \in N_n(t)}}{\lambda[N_n(t)]} \quad (6.1)$$

where  $N_n(t)$  is a neighborhood of  $t$  with  $\lambda[N_n(t)] \rightarrow 0$  for  $n \rightarrow \infty$ . In a consistency proof, such as that of Lemma 4.3, we have to control a term like

$$\frac{1}{N} \sum_{i=1}^N \left| 1 - \frac{d_n(t, T_i)}{h_{T|X}(t, X_i)} \right|,$$

where  $h_{T|X}$  is the generalized propensity score of [HI05]. We need a result such as

$$\mathbf{E} \left[ \frac{d_n(t, T_i)}{h_{T|X}(t, X_i)} \right] = \mathbf{E} \left[ \frac{\mathbf{P}[T_i \in N_n(t) | X_i]}{\lambda[N_n(t)]} \cdot \frac{1}{h_{T|X}(t, X_i)} \right] \rightarrow \mathbf{E} \left[ \frac{h_{T|X}(t, X_i)}{h_{T|X}(t, X_i)} \right] = 1.$$

A possible error decomposition could be derived from

$$\begin{aligned} & \left| \frac{1}{n} \sum_{i=1}^n d_n(t, T_i) w_i Y_i - \mathbf{E}[Y(t)] \right| \\ & \leq \left| \frac{1}{n} \sum_{i=1}^n (w_i d_n(t, T_i) - 1) \langle B(X_i), \mathbf{Y}(t) \rangle \right| \\ & \quad + \left| \frac{1}{n} \sum_{i=1}^n (w_i d_n(t, T_i) - 1) (\mathbf{E}[Y(t) | X_i] - \langle B(X_i), \mathbf{Y}(t) \rangle) \right| \\ & \quad + \left| \frac{1}{n} \sum_{i=1}^n d_n(t, T_i) \cdot (w_i - 1/h_{T|X}(t, X_i)) (Z_i - \mathbf{E}[Y(t) | X_i]) \right| \\ & \quad + \left| \frac{1}{n} \sum_{i=1}^n h_T(t)/h_{T|X}(t, X_i) (Z_i - \mathbf{E}[Y(t) | X_i]) \right| \\ & \quad + \left| \frac{1}{n} \sum_{i=1}^n (h_T(t) - d_n(t, T_i)) / h_{T|X}(t, X_i) (Z_i - \mathbf{E}[Y(t) | X_i]) \right| \\ & \quad + \left| \frac{1}{n} \sum_{i=1}^n (\mathbf{E}[Y(t) | X_i] - \mathbf{E}[Y(t)]) \right| \\ & \quad + \left| \frac{1}{n} \sum_{i=1}^n w_i d_n(t, T_i) (Y_i - Z_i) \right|, \end{aligned}$$

where  $Z_i \sim Y(t)|T_i$ .

### Organisation

Get familiar with generalized propensity score [HI05]. Try to adapt ideas to make the proofs work. Talk to Jose Zubizarreta, one of the authors of [WZ19].

### Next Step

Read [HI05].

## 6.2.5 Different Basis Functions

### Motivation

The introduction of partitioning estimates [GKKW02, §4] - as done in this thesis - was successful. Thus the implementation of other local averaging regression techniques, such as kernel estimates [GKKW02, §5] is promising.

### Conjecture

Similar results as of this thesis hold for basis functions of (boxed) kernel estimates [GKKW02, §5]. They have good practical performance.

### Ideas/Brainstorming

For boxed kernels it is likely easy to prove a lemma similar to Lemma 3.7. For kernels with unbounded support, such as gaussian kernels, this might be more difficult. Generally, the basis functions should approximate treatment and outcome model well (see [WZ19, Assumptions 1.6 & 2.3]). Partitioning estimates work well in this thesis, because we can define concrete oracle parameters. If concrete oracle parameters are not readily available, there might be theoretic results to rely on.

### Organisation

Get familiar with the notion of (boxed) kernel [GKKW02]. Find a result similar to Lemma 3.7. Try other basis functions, maybe with unbounded support. Rely on theoretic results to derive results such as Lemma 3.7 for a more abstract oracle parameter.

### Next Step

Look up the definition of boxed kernel [GKKW02, Theorem 5.1, Figure 5.7].



## 7 Convex Analysis

In our application we want to analyse a convex optimization problem by its dual problem. In particular we want to obtain primal optimal solutions from dual solutions. To accomplish the task we need technical tools from convex analysis, mainly conjugate calculus and some Karush-Kuhn-Tucker related results.

Our starting point is the support function intersection rule [MMN22, Theorem 4.23]. We give the details in the case of finite dimensions and refer for the rest of the proof to the book. The support function intersection rule is applied to give first conjugate sum and then chain rule, which are vital to calculating convex conjugates. The proofs are omitted, since the book is thorough enough. The material we present is very well known. As an introduction, we recommend the recent book [MMN22] and the classical reference [Roc70]. We finish the chapter with ideas from [TB91]. They provide the high-level ideas to obtain for strictly convex functions a dual relationship between optimal solutions. We will deliver the details that are omitted in the paper.

### 7.1 A Convex Analysis Primer

#### My Contribution

I present the relevant facts from Convex analysis. I prove some results that I did not find in the literature, but likely are folklore.

Throughout this section let  $n \in \mathbb{N}$ .

#### Sets

A subset  $C \subseteq \mathbb{R}^n$  is called **convex set**, if for all  $x, y \in C$  and all  $\theta \in [0, 1]$ , we have  $\theta x + (1 - \theta)y \in C$ . Many set operations preserve convexity. Among them forming the **Cartesian product** of two convex sets, **intersection** of a collection of convex sets and taking the **inverse image under linear functions**.

The classical theory evolves around the question if convex sets can be separated.

**Definition.** Let  $C_1$  and  $C_2$  be two non-empty convex sets in  $\mathbb{R}^n$ . A hyperplane  $H$  is said to **separate**  $C_1$  and  $C_2$  if  $C_1$  is contained in one of the closed half-spaces



associated with  $H$  and  $C_2$  lies in the opposite closed half-space. It is said to separate  $C_1$  and  $C_2$  **properly** if  $C_1$  and  $C_2$  are not both contained in  $H$ .

We need a refined concept of interiors, since some convex sets have empty interior. To this end, we call a set  $A \subseteq \mathbb{R}^n$  **affine set**, if  $\alpha x + (1 - \alpha)y \in A$  for all  $x, y \in A$  and all  $\alpha \in \mathbb{R}$ . The **affine hull**  $\text{aff}(\Omega)$  of a set  $\Omega \subseteq \mathbb{R}^n$  is the smallest affine set that includes  $\Omega$ . We define the **relative interior**  $\text{ri}\Omega$  of a set  $\Omega \subseteq \mathbb{R}^n$  to be the interior relative to the affine hull, that is,

$$\text{ri}(\Omega) := \{x \in \Omega \mid \exists \varepsilon > 0 : (x + \varepsilon B_{\mathbb{R}^n}) \cap \text{aff}(\Omega) \subset \Omega\}. \quad (7.1)$$

**Theorem 7.1.** (Convex separation in finite dimension) *Let  $C_1$  and  $C_2$  be two non-empty convex sets in  $\mathbb{R}^n$ . Then  $C_1$  and  $C_2$  can be properly separated if and only if  $\text{ri}(C_1) \cap \text{ri}(C_2) = \emptyset$ .*

**Proof.** [Roc70, Theorem 11.3] □

We collect some useful properties of relative interiors before we get on to convex functions.

**Proposition 7.1.** *Let  $C$  be a non-empty convex set in  $\mathbb{R}^n$ . The following holds:*

- (i)  $\text{ri}(C) \neq \emptyset$  if and only if  $C \neq \emptyset$
- (ii)  $\text{cl}(\text{ri} C) = \text{cl} C$  and  $\text{ri}(\text{cl} C) = \text{ri}(C)$
- (iii)  $\text{ri}(C) = \{z \in C : \text{for all } x \in C \text{ there exists } t > 0 \text{ such that } z + t(z - x) \in C\}$
- (iv) Suppose  $\bigcap_{i \in I} C_i \neq \emptyset$  for a finite index set  $I$ . Then  $\text{ri}(\bigcap_{i \in I} C_i) = \bigcap_{i \in I} \text{ri}(C_i)$ .
- (v) Let  $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear function. Then  $\text{ri} L(C) = L(\text{ri} C)$ . If it also holds  $L^{-1}(\text{ri} C) \neq \emptyset$ , we have  $\text{ri} L^{-1}(C) = L^{-1}(\text{ri} C)$ .
- (vi)  $\text{ri}(C_1 \times C_2) = \text{ri} C_1 \times \text{ri} C_2$

**Proof.** For a proof of (i)-(v) we refer to [Roc70, Theorem 6.2 - 6.7].

To prove (vi) we use (iii). Let  $(z_1, z_2) \in \text{ri}(C_1 \times C_2)$ . Then for all  $(x_1, x_2) \in C_1 \times C_2$  there exists  $t > 0$  such that

$$z_i + t(z_i - x_i) \in C_i \quad \text{for all } i \in \{1, 2\}. \quad (7.2)$$

Using (iii) again, we get  $\text{ri}(C_1 \times C_2) \subseteq \text{ri } C_1 \times \text{ri } C_2$ . Suppose  $(z_1, z_2) \in \text{ri } C_1 \times \text{ri } C_2$ . By (iii), for all  $(x_1, x_2) \in C_1 \times C_2$  there exist  $(t_1, t_2) > 0$  such that

$$z_i + t_i(z_i - x_i) \in C_i \quad \text{for all } i \in \{1, 2\}. \quad (7.3)$$

If  $t_1 = t_2$  we recover (7.2) from (7.3). By (iii) it holds  $(z_1, z_2) \in \text{ri}(C_1 \times C_2)$ . If  $t_1 < t_2$  we define  $\theta := \frac{t_1}{t_2} \in (0, 1)$ . Consider (7.3) with  $i = 2$ , together with  $z_2 \in C_2$  and the convexity of  $C_2$ . It follows

$$z_2 + t_1(z_2 - x_2) = \theta \cdot (z_2 + t_2(z_2 - x_2)) + (1 - \theta) \cdot z_2 \in C_2. \quad (7.4)$$

Now we consider (7.4) and (7.3) with  $i = 1$ . This gives (7.2) with  $t = t_1$ . As before, it follows  $(z_1, z_2) \in \text{ri}(C_1 \times C_2)$ . If  $t_1 > t_2$  similar arguments lead to the same result. We have proven  $\text{ri}(C_1 \times C_2) \supseteq \text{ri } C_1 \times \text{ri } C_2$  and equality.  $\square$

## Functions

A function  $f: \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$  is called **convex function**, if the area above its graph, that is, its epigraph(cf. [MMN22, §2.4.1]), is convex. We shall often use an equivalent definition. To this end, a function  $f$  is convex if and only if

$$f(\theta x + (1 - \theta)y) \leq \theta f(x) + (1 - \theta)f(y) \quad \text{for all } x, y \in \mathbb{R}^n \text{ and all } \theta \in [0, 1]. \quad (7.5)$$

This definition extends to convex combinations  $\theta_1, \dots, \theta_m \in [0, 1]$  with  $\sum_{i=1}^m \theta_i = 1$ , that is, a function  $f$  is convex if and only if

$$f\left(\sum_{i=1}^m \theta_i x_i\right) \leq \sum_{i=1}^m \theta_i f(x_i) \quad \text{for all } x_1, \dots, x_m \in \mathbb{R}^n. \quad (7.6)$$

We call a function **strictly convex** if the inequality in (7.5) is strict.

We define the **domain**  $\text{dom } f$  of a convex function  $f$  to be the set where  $f$  is finite, that is,

$$\text{dom } f := \{x \in \mathbb{R}^n : f(x) < \infty\}. \quad (7.7)$$

The domain of a convex function is convex. We say that  $f$  is a **proper function** if  $\text{dom } f \neq \emptyset$ .

For any  $\bar{x} \in \text{dom } f$  we call  $x^* \in \mathbb{R}^n$  a **subgradient** of  $f$  at  $\bar{x}$  if for all  $x \in \mathbb{R}^n$  it holds

$$\langle x^*, x - \bar{x} \rangle \leq f(x) - f(\bar{x}). \quad (7.8)$$

We denote the collection of all subgradients at  $\bar{x}$ , that is, the **subdifferential** of  $f$  at  $\bar{x}$ , as  $\partial f(\bar{x})$ . If  $f$  is differentiable at  $\bar{x}$  it holds  $\partial f(\bar{x}) = \{\nabla f(\bar{x})\}$  and thus

$$\langle \nabla f(\bar{x}), x - \bar{x} \rangle \leq f(x) - f(\bar{x}). \quad (7.9)$$

**Definition 7.1.** Given a nonempty subset  $\Omega \subseteq \mathbb{R}^n$ , we define the **support function** of  $\Omega$  to be

$$\sigma_\Omega : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}, \quad x^* \mapsto \sup_{x \in \Omega} \langle x^*, x \rangle.$$

**Definition 7.2.** Given functions  $f_i : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$  for  $i = 1, \dots, m$ , we define the **infimal convolution** of these functions to be

$$f_1 \square \dots \square f_m : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}, \quad x \mapsto \inf \left\{ \sum_{i=1}^m f_i(x_i) : x_i \in \mathbb{R}^n \text{ and } \sum_{i=1}^m x_i = x \right\}.$$

The next result establishes a connection between the support function of the intersection of two convex sets and the infimal convolution of the support functions of the sets taken by themselves. The proof translates the geometric concept of convex separation to the world of convex functions.

**Lemma 7.1.** Let  $C_1$  and  $C_2$  be two non-empty convex sets in  $\mathbb{R}^n$ . For any  $x^* \in \text{dom } \sigma_{C_1 \cap C_2}$  the sets

$$\begin{aligned} \Theta_1 &:= C_1 \times [0, \infty), \\ \Theta_2(x^*) &:= \{(x, \lambda) \in \mathbb{R}^n : x \in C_2 \text{ and } \lambda \leq \langle x^*, x \rangle - \sigma_{C_1 \cap C_2}(x^*)\} \end{aligned}$$

can be properly separated.

**Proof.** We fix  $x^* \in \text{dom } \sigma_{C_1 \cap C_2}$  and write  $\alpha := \sigma_{C_1 \cap C_2}(x^*)$ . In order to apply convex separation in finite dimension (Theorem 7.1) to the sets  $\Theta_1$  and  $\Theta_2(x^*)$ , it suffices to show their convexity and  $\text{ri } \Theta_1 \cap \text{ri } \Theta_2(x^*) = \emptyset$ .

**Convexity of  $\Theta_1$  and  $\Theta_2(x^*)$**

Clearly,  $\Theta_1$  is convex by the convexity of  $C_1$  and  $[0, \infty)$ . To see that  $\Theta_2(x^*)$  is convex consider the linear function

$$L : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}, \quad (x, \lambda) \mapsto \langle x^*, x \rangle - \lambda.$$

From the definitions of  $L$  and  $\Theta_2(x^*)$  we get

$$\Theta_2(x^*) = (C_2 \times \mathbb{R}) \cap L^{-1}[\alpha, \infty).$$

Thus, by Proposition 7.1 (v) and the convexity of  $C_2$  we get the convexity of  $L^{-1}[\alpha, \infty)$  and with it that of  $\Theta_2(x^*)$ .

### Relative interiors of $\Theta_1$ and $\Theta_2(x^*)$ are disjoint

We start by calculating the relative interiors. It holds

$$\begin{aligned} \text{ri } \Theta_1 &= \text{ri}(C_1 \times [0, \infty)) = \text{ri } C_1 \times \text{ri } [0, \infty) = \text{ri } C_1 \times (0, \infty), \\ \text{ri } \Theta_2(x^*) &= \text{ri}(L^{-1}[\alpha, \infty)) = L^{-1}(\text{ri } [\alpha, \infty)) = L^{-1}(\alpha, \infty). \end{aligned}$$

Suppose there exists  $(\lambda, x) \in \text{ri } \Theta_1 \cap \text{ri } \Theta_2(x^*)$ . Then it holds  $x \in C_1 \times C_2$  and  $\lambda > 0$ . We also note, that

$$\alpha = \sigma_{C_1 \cap C_2}(x^*) = \sup_{z \in C_1 \cap C_2} \langle x^*, z \rangle \geq \langle x^*, x \rangle.$$

Then it follows

$$\alpha < \langle x^*, x \rangle - \lambda \leq \alpha,$$

a contradiction. Thus, the relative interiors of  $\Theta_1$  and  $\Theta_2(x^*)$  are disjoint.

Applying Theorem 7.1 finishes the proof.  $\square$

**Theorem.** Let  $C_1$  and  $C_2$  be two non-empty convex sets in  $\mathbb{R}^n$  with  $\text{ri } C_1 \cap \text{ri } C_2 \neq \emptyset$ . Then the support function of the intersection  $C_1 \cap C_2$  is represented as

$$(\sigma_{C_1 \cap C_2})(x^*) = (\sigma_{C_1} \square \sigma_{C_2})(x^*) \quad \text{for all } x^* \in \mathbb{R}^n. \quad (7.10)$$

Furthermore, for any  $x^* \in \text{dom}(\sigma_{C_1 \cap C_2})$  there exist dual elements  $x_1^*, x_2^* \in \mathbb{R}^n$  such that  $x^* = x_1^* + x_2^*$ . and

$$(\sigma_{C_1 \cap C_2})(x^*) = \sigma_{C_1}(x_1^*) + \sigma_{C_2}(x_2^*). \quad (7.11)$$

**Proof.** Using Lemma 7.1 the rest of the proof is as that of [MMN22, Theorem 4.23(b)].  $\square$

**Takeaways** The support function intersection rule connects the geometric property of convex separation to an identity of support functions. This result is central to the analysis of convex conjugates.

One important application of convex functions is in optimization. There we often analyse a dual problem instead, which relies on the notion of **convex conjugate**  $f^* : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$  of  $f$  defined by

$$f^*(x^*) := \sup_{x \in \mathbb{R}^n} \langle x^*, x \rangle - f(x). \quad (7.12)$$

Even for arbitrary functions, the convex conjugate is convex(cf. [MMN22, Proposition 4.2] ). Like in differential calculus, there exist sum and chain rule for computing the convex conjugate.

**Theorem 7.2.** *Let  $f, g : \mathbb{R}^n \rightarrow (-\infty, \infty]$  be proper convex functions and*

$$ri(dom(f)) \cap ri(dom(g)) \neq \emptyset.$$

*Then we have the **conjugate sum rule***

$$(f + g)^*(x^*) = (f^* \square g^*)(x^*) \quad (7.13)$$

*for all  $x^* \in \mathbb{R}^n$ . Moreover, the infimum in  $(f^* \square g^*)(x^*)$  is attained, i.e., for any  $x^* \in dom(f + g)^*$  there exists vectors  $x_1^*, x_2^*$  for which*

$$(f + g)^*(x^*) = f^*(x_1^*) + g^*(x_2^*), \quad x^* = x_1^* + x_2^*. \quad (7.14)$$

**Proof.** [MMN22, Theorem 4.27(c)] □

**Theorem 7.3.** *Let  $A : \mathbb{R}^m \rightarrow \mathbb{R}^n$  be a linear map (matrix) and  $g : \mathbb{R}^n \rightarrow (-\infty, \infty]$  a proper convex function. If  $Im(A) \cap ri(dom(g)) \neq \emptyset$  it follows the **conjugate chain rule***

$$(g \circ A)^*(x^*) = \inf_{y^* \in (A^*)^{-1}(x^*)} g^*(y^*). \quad (7.15)$$

*Furthermore, for any  $x^* \in dom(g \circ A)^*$  there exists  $y^* \in (A^*)^{-1}(x^*)$  such that  $(g \circ A)^*(x^*) = g^*(y^*)$ .*

**Proof.** [MMN22, Theorem 4.28(c)] □

**Example 7.1.** Let  $f : \mathbb{R} \rightarrow \overline{\mathbb{R}}$  be a proper convex function, that is,  $dom f \neq \emptyset$  and  $f$  is convex. In steps we apply the conjugate chain and sum rule, together with

mathematical induction, to prove the conjugate relationship

$$\begin{aligned} S_{f,n} : \mathbb{R}^n &\rightarrow \overline{\mathbb{R}}, & (x_1, \dots, x_n) &\mapsto \sum_{i=1}^n f(x_i), \\ S_{f,n}^* : \mathbb{R}^n &\rightarrow \overline{\mathbb{R}}, & (x_1^*, \dots, x_n^*) &\mapsto \sum_{i=1}^n f^*(x_i^*). \end{aligned}$$

This relationship is very natural and the ensuing calculations serve to confirm our intuition.

First, we work in the projections on the coordinates. For the  $i$ -th coordinate, where  $i = 1, \dots, n$ , this is

$$p_i : \mathbb{R}^n \rightarrow \mathbb{R}, \quad (x_1, \dots, x_n) \mapsto x_i. \quad (7.16)$$

All projections  $p_i$  are linear function with matrix representation  $e_i^\top$ , where  $e_i$  is  $i$ -the coordinate vector. The adjoint of  $p_i$  is therefore

$$p_i^* : \mathbb{R} \rightarrow \mathbb{R}^n, \quad x \mapsto e_i \cdot x. \quad (7.17)$$

For the inverse image of the adjoint of  $p_i$  it holds

$$(p_i^*)^{-1} \{(x_1^*, \dots, x_n^*)\} = \begin{cases} \{x_i^*\}, & \text{if } x_j^* = 0 \text{ for all } j \neq i, \\ \emptyset & \text{else.} \end{cases} \quad (7.18)$$

Throughout this example we use the asterisk character  $*$  somewhat inconsistently. Note that  $f^*$  is the convex conjugate of the function  $f$  and  $p_i^*$  is the adjoint linear function of the projection on the  $i$ -th coordinate. Likewise, we denote dual variables, that is, the arguments of convex conjugates, as  $x^*$ .

Next, we employ the conjugate chain rule to establish the conjugate relationship

$$\begin{aligned} f_i : \mathbb{R}^n &\rightarrow \overline{\mathbb{R}}, & (x_1, \dots, x_n) &\mapsto x_i \mapsto f(x_i), \\ f_i^* : \mathbb{R} &\rightarrow \overline{\mathbb{R}}, & x_i^* &\mapsto \begin{cases} f^*(x_i^*), & \text{if } x_j^* = 0 \text{ for all } j \neq i, \\ \infty & \text{else.} \end{cases} \end{aligned}$$

Note, that  $f_i = (f \circ p_i)$  and  $f_i^* = (f \circ p_i)^*$ . Since  $\text{Im } p_i = \mathbb{R}$  and  $\text{dom } f \neq \emptyset$ , it holds  $\text{Im } p_i \cap \text{ri}(\text{dom } f) \neq \emptyset$ . Then  $f$  and  $p_i$  conform with the demands of the conjugate chain rule. It follows

$$\begin{aligned} f_i^*(x_1^*, \dots, x_n^*) &= (f \circ p_i)^*(x_1^*, \dots, x_n^*) = \inf \{f^*(y) \mid y \in (p_i^*)^{-1} \{(x_1^*, \dots, x_n^*)\}\} \\ &= \begin{cases} f^*(x_i^*), & \text{if } x_j^* = 0 \text{ for all } j \neq i, \\ \infty & \text{else,} \end{cases} \end{aligned}$$

where we keep to the convention  $\inf \emptyset = \infty$ . In the same way it follows

$$(S_{f,n} \circ p_{\{1,\dots,n\}})^*(x_1^*, \dots, x_{n+1}^*) = \begin{cases} S_{f,n}^*(x_1^*, \dots, x_n^*) & \text{if } x_{n+1}^* = 0, \\ \infty & \text{else,} \end{cases} \quad (7.19)$$

Next, note that for  $n = 1$  we arrive at the result. Thus, for some  $n \in \mathbb{N}$  it holds  $(S_{f,n})^* = S_{f,n}^*$ . In order to apply the conjugate sum rule to  $S_{f,n}$  and  $f_{n+1}$  we note that

$$\begin{aligned} \text{dom } f_i &= \{(x_1, \dots, x_{n+1}) \in \mathbb{R}^{n+1} : x_i \in \text{dom } f\} \neq \emptyset \quad \text{for all } i = 1, \dots, n+1, \\ \bigcap_{i=1}^{n+1} \text{dom } f_i &= \{(x_1, \dots, x_{n+1}) \in \mathbb{R}^{n+1} : x_i \in \text{dom } f \text{ for all } i = 1, \dots, n+1\} \neq \emptyset, \end{aligned}$$

and

$$\begin{aligned} \text{ri}(\text{dom}(S_{f,n} \circ p_{\{1,\dots,n\}})) \cap \text{ri}(\text{dom } f_{n+1}) \\ = \text{ri}(\text{dom}(S_{f,n} \circ p_{\{1,\dots,n\}}) \cap \text{dom } f_{n+1}) = \text{ri}\left(\bigcap_{i=1}^{n+1} \text{dom } f_i\right) \neq \emptyset. \end{aligned}$$

By the conjugate sum rule it follows

$$\begin{aligned} (S_{f,n+1})^* &= (S_{f,n} \circ p_{\{1,\dots,n\}} + f_{n+1})^* = (S_{f,n} \circ p_{\{1,\dots,n\}})^* \square f_{n+1}^* \\ &= S_{f,n}^* \circ p_{\{1,\dots,n\}} + f_{n+1}^* = S_{f,n+1}^*. \end{aligned}$$

◇

## 7.2 Duality of Optimal Solutions

### My Contribution

I adapt ideas from [TB91] to take also equality constraints. For this, I had to understand the connection to my version of the primal optimization problem. I filled in many details that were omitted in the paper: I derived the Karush-Kuhn-Tucker conditions for the problem from the general result [Roc70, Theorem 28.3]. I prove in detail, that they hold for the adapted problem.

We consider a general convex optimization problem with matrix equality and inequality constraints. For this problem there exists a related problem, which we call its dual. With ideas from [TB91] we establish a functional relationship between the optimal solution of the original problem and optimal solutions of the dual. The main assumption is that in the original problem we have a strictly convex objective function with continuously differentiable convex conjugate.

**Assumption 2.** The objective function  $f: \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$  is strictly convex and its convex conjugate  $f^*$  is continuously differentiable.

**Theorem 7.4.** Consider the optimization problem

$$\begin{aligned} & \underset{w \in \mathbb{R}^n}{\text{minimize}} && f(w) \\ & \text{subject to} && \mathbf{U}w \geq d, \\ & && \mathbf{A}w = a, \end{aligned} \tag{7.20}$$

and its dual problem

$$\begin{aligned} & \underset{\lambda_d \in \mathbb{R}^r, \lambda_a \in \mathbb{R}^s}{\text{maximize}} && \langle \lambda_d, d \rangle + \langle \lambda_a, a \rangle - f^*(\mathbf{U}^\top \lambda_d + \mathbf{A}^\top \lambda_a) \\ & \text{subject to} && \lambda_d \geq 0. \end{aligned} \tag{7.21}$$

Let  $(\lambda_d^\dagger, \lambda_a^\dagger)$  be an optimal solution to (7.21). If the objective function  $f$  of (7.20) is strictly convex and its convex conjugate  $f^*$  is continuously differentiable, then the unique optimal solution to (7.20) is given by

$$w^\dagger = \nabla f^*(\mathbf{U}^\top \lambda_d^\dagger + \mathbf{A}^\top \lambda_a^\dagger). \tag{7.22}$$

### Plan of Proof

We show that  $w^\dagger$  and  $(\lambda_d^\dagger, \lambda_a^\dagger)$  meet the Karush-Kuhn-Tucker conditions for 7.20, that is, **complementary slackness**

$$\langle \lambda_d^\dagger, d - \mathbf{U}w^\dagger \rangle = 0, \tag{7.23}$$

**primal and dual feasibility**

$$\mathbf{U}w^\dagger \geq d, \tag{7.24}$$

$$\mathbf{A}w^\dagger = a,$$

$$\lambda_d^\dagger \geq 0, \tag{7.25}$$

and **stationarity**

$$0_n \in [\partial f(w^\dagger) + \partial(w \mapsto d - \mathbf{U}w)(w^\dagger) \cdot \lambda_d^\dagger + \partial(w \mapsto a - \mathbf{A}w)(w^\dagger) \cdot \lambda_a^\dagger]. \tag{7.26}$$

Applying the well know result [Roc70, Theorem 28.3] finishes the proof. Apart from elementary calculations, our main tools are the strict convexity of  $f$ , the smoothness of  $f^*$  and



**Proposition 7.2.** [Roc70, Theorem 23.5(a)-(b)]. *For any proper convex function  $g$  and any vector  $w$ , it holds  $t \in \partial f(w)$  if and only if  $x \mapsto \langle x, t \rangle - f(x)$  achieves its supremum at  $w$ .*

**Proof.** Let  $(\lambda_d^\dagger, \lambda_a^\dagger)$  be an optimal solution to (7.21).

### Complementary Slackness

We fix  $\lambda_a^\dagger$  and work with the objective function  $G$  of the dual problem, that is,

$$G(\lambda_d) := \langle \lambda_d, d \rangle + \langle \lambda_a^\dagger, a \rangle - f^* \left( \mathbf{U}^\top \lambda_d + \mathbf{A}^\top \lambda_a^\dagger \right).$$

Since  $f^*$  is continuously differentiable, so is  $G$ . Thus

$$\nabla G(\lambda_d^\dagger) := d - \mathbf{U} \cdot \nabla f^* \left( \mathbf{U}^\top \lambda_d^\dagger + \mathbf{A}^\top \lambda_a^\dagger \right) = d - \mathbf{U} w^\dagger.$$

Let  $\lambda_{d,i}^\dagger$  be the  $i$ -th coordinate of  $\lambda_d^\dagger$  and  $\nabla G_i(\lambda_d^\dagger)$  be the  $i$ -th coordinate of  $\nabla G(\lambda_d^\dagger)$ . To establish (7.23) we will show for all coordinates

$$\begin{aligned} \text{either} \quad & \lambda_{d,i}^\dagger = 0 \quad \text{and} \quad \nabla G_i(\lambda_d^\dagger) \leq 0 \\ \text{or} \quad & \lambda_{d,i}^\dagger > 0 \quad \text{and} \quad \nabla G_i(\lambda_d^\dagger) = 0. \end{aligned}$$

It is well known that a concave function  $g$  satisfies

$$g(x) - g(y) \geq \nabla g(x)^\top (x - y) \quad \text{for all } x, y. \quad (7.27)$$

But  $G$  is concave by the convexity of  $f^*$ .

First, we show

$$\nabla G_i(\lambda_d^\dagger) \leq 0 \quad \text{for all } i \in \{1, \dots, s\}. \quad (7.28)$$

Assume towards a contradiction that  $\nabla G_i(\lambda_d^\dagger) > 0$  for some  $i \in \{1, \dots, s\}$ . By the continuity of  $\nabla G$  there exists  $\varepsilon > 0$  such that  $\nabla G_i(\lambda_d^\dagger + e_i \cdot \varepsilon) > 0$ . It follows from (7.27)

$$G(\lambda_d^\dagger + e_i \cdot \varepsilon) - G(\lambda_d^\dagger) \geq \nabla G_i(\lambda_d^\dagger + e_i \cdot \varepsilon) \cdot \varepsilon > 0,$$

which contradicts the optimality of  $\lambda_d^\dagger$  for (7.21). It follows (7.28).

Next, we assume that  $\lambda_{d,i}^\dagger > 0$  and  $\nabla G_i(\lambda_d^\dagger) < 0$  for some  $i \in \{1, \dots, s\}$ . Again, by the continuity of  $\nabla G$  there exists  $\varepsilon > 0$  such that  $\nabla G_i(\lambda_d^\dagger - e_i \cdot \varepsilon) < 0$  and  $\varepsilon - \lambda_{d,i}^\dagger < 0$ . Thus

$$G(\lambda_d^\dagger - e_i \cdot \varepsilon) - G(\lambda_d^\dagger) \geq \nabla G_i(\lambda_d^\dagger - e_i \cdot \varepsilon) \cdot (-\varepsilon) > 0,$$

which contradicts the optimality of  $\lambda_d^\dagger$ . It follows (7.23), that is, we proved complementary slackness.

**Primal Feasibility**

Since  $f^*$  is continuously differentiable it holds

$$\nabla G(\lambda_d^\dagger) = d - \mathbf{U} \cdot \nabla f^* \left( \mathbf{U}^\top \lambda_d^\dagger + \mathbf{A}^\top \lambda_a^\dagger \right) = d - \mathbf{U} w^\dagger.$$

Thus, by (7.28),  $w^\dagger$  satisfies the inequality constraints in (7.20). To prove this for the equality constraints, we view  $G$  from a different angel. Let for fixed  $\lambda_d^\dagger$

$$G(\lambda_a) := \langle \lambda_a, a \rangle - \left( f^* \left( \mathbf{U}^\top \lambda_d^\dagger + \mathbf{A}^\top \lambda_a \right) - \langle \lambda_d^\dagger, d \rangle \right) =: \langle \lambda_a, a \rangle - g(\lambda_a).$$

The function  $g$  inherits convexity and differentiability from  $f^*$ . From the optimality of  $\lambda_a^\dagger$  we know that  $G$  takes its maximum there. But then by Proposition 7.2 and the differentiability of  $g$  it holds

$$a \in \partial g(\lambda_a^\dagger) = \left\{ \mathbf{A} \cdot \nabla f^* \left( \mathbf{U}^\top \lambda_d^\dagger + \mathbf{A}^\top \lambda_a^\dagger \right) \right\} = \left\{ \mathbf{A} w^\dagger \right\}. \quad (7.29)$$

Thus  $a = \mathbf{A} w^\dagger$ . But then  $w^\dagger$  satisfies also the equality constraints. We proved (7.24).

**Stationarity**

First we show

$$\mathbf{U}^\top \lambda_d^\dagger + \mathbf{A}^\top \lambda_a^\dagger \in \partial f(w^\dagger). \quad (7.30)$$

By Proposition 7.2 it suffices to show that

$$w \mapsto \langle w, \mathbf{U}^\top \lambda_d^\dagger + \mathbf{A}^\top \lambda_a^\dagger \rangle - f(w)$$

achieves its supremum at  $w^\dagger$ . Since  $f$  is strictly convex there exists a unique vector  $x^\dagger$  where the above expression achieves its maximum. Since  $f^*$  is differentiable it holds

$$w^\dagger = \nabla f^* \left( \mathbf{U}^\top \lambda_d^\dagger + \mathbf{A}^\top \lambda_a^\dagger \right) = \nabla \left( \lambda \mapsto \langle x^\dagger, \lambda \rangle - f(x^\dagger) \right) \left( \mathbf{U}^\top \lambda_d^\dagger + \mathbf{A}^\top \lambda_a^\dagger \right) = x^\dagger.$$

It follows (7.30). Next we show

$$-\mathbf{U}^\top \in \partial (w \mapsto d - \mathbf{U}w)(w^\dagger) \quad \text{and} \quad -\mathbf{A}^\top \in \partial (w \mapsto d - \mathbf{A}w)(w^\dagger). \quad (7.31)$$

To this end, note that

$$\langle -\mathbf{U}^\top e_i, w - w^\dagger \rangle = (d - \mathbf{U}w)_i - (d - \mathbf{U}w^\dagger)_i \quad \text{for all } i \in \{1, \dots, r\}.$$

Thus  $-\mathbf{U}^\top \in \partial (w \mapsto d - \mathbf{U}w)(w^\dagger)$ . In the same way it follows  $-\mathbf{A}^\top \in \partial (w \mapsto d - \mathbf{A}w)(w^\dagger)$ .

From (7.30) and (7.31) we conclude

$$\begin{aligned} 0_n &= \left( \mathbf{U}^\top \lambda_d^\dagger + \mathbf{A}^\top \lambda_a^\dagger \right) - \mathbf{U}^\top \lambda_d^\dagger - \mathbf{A}^\top \lambda_a^\dagger \\ &\in [\partial f(w^\dagger) + \partial (w \mapsto d - \mathbf{U}w)(w^\dagger) \cdot \lambda_d^\dagger + \partial (w \mapsto d - \mathbf{A}w)(w^\dagger) \cdot \lambda_a^\dagger]. \end{aligned}$$

We have proved (7.26), that is, stationarity.

**Dual Feasibility and Conclusion**

Dual feasibility (7.25) follows immediately from the optimality of  $\lambda_d^\dagger$  for (7.21). Thus,  $(\lambda_d^\dagger, \lambda_a^\dagger)$  and  $w^\dagger$  satisfy the Karush-Kuhn-Tucker conditions for (7.20). Applying [Roc70, Theorem 28.3] finishes the proof.  $\square$

**Takeaways** For strictly convexity objective functions with continuously differentiable convex conjugate we get a functional relationship of primal and dual solutions via the Karush-Kuhn-Tucker conditions.

# Notation Index

$\#A$  cardinality of the set  $A$

$\mathbf{E}[X|Y]$  conditional expectation of the random variable  $X$  with respect to  $\sigma(Y)$

$\mathbf{E}[X]$  expectation of the random variable  $X$

$\mathbf{Var}[X]$  variance of the random variable  $X$

$\overline{\mathbb{R}} = \mathbb{R} \cup \{+\infty\}$  extension of the real numbers

$\xrightarrow{\mathcal{D}}$  convergence of distributions

$\mathbf{P}$  generic probability measure

$\mathbf{P}_X = \mathbf{P} \circ X^{-1}$  distribution of the random variable  $X$

$\mathbb{R}$  set of real numbers

$x \vee y, x \wedge y, x^+, x^-$  maximum, minimum, positive part, negative part of real numbers

$X \sim \mu$  the random variable has distribution  $\mu$



## References

- [AB07] Charalambos D. Aliprantis and Kim C. Border. *Infinite Dimensional Analysis: A Hitchhiker's Guide*. Springer Science & Business Media, May 2007.
- [BB95] Philippe Barbe and Patrice Bertail. *The Weighted Bootstrap*, volume 98 of *Lecture Notes in Statistics*. Springer, New York, NY, 1995.
- [CYZ16] Kwun Chuen Gary Chan, Sheung Chi Phillip Yam, and Zheng Zhang. Globally Efficient Non-Parametric Inference of Average Treatment Effects by Empirical Balancing Calibration Weighting. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 78(3):673–700, June 2016.
- [ET94] Bradley Efron and R. J. Tibshirani. *An Introduction to the Bootstrap*. CRC Press, May 1994.
- [GKKW02] László Györfi, Michael Kohler, Adam Krzyżak, and Harro Walk. *A Distribution-Free Theory of Nonparametric Regression*. Springer Series in Statistics. Springer, New York, NY, 2002.
- [Hai12] Jens Hainmueller. Entropy Balancing for Causal Effects: A Multivariate Reweighting Method to Produce Balanced Samples in Observational Studies. *Political Analysis*, 20(1):25–46, 2012.
- [HI05] Keisuke Hirano and Guido W. Imbens. The Propensity Score with Continuous Treatments. In Andrew Gelman and Xiao-Li Meng, editors, *Wiley Series in Probability and Statistics*, pages 73–84. John Wiley & Sons, Ltd, Chichester, UK, July 2005.
- [IR14] Kosuke Imai and Marc Ratkovic. Covariate balancing propensity score. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 76:243–263, 2014.
- [Kle20] Achim Klenke. *Probability Theory: A Comprehensive Course*. Universitext. Springer International Publishing, Cham, 2020.

- [MMN22] Boris S. Mordukhovich and Nguyen Mau Nam. *Convex Analysis and Beyond: Volume I: Basic Theory*. Springer Series in Operations Research and Financial Engineering. Springer International Publishing, Cham, 2022.
- [New97] Whitney K. Newey. Convergence rates and asymptotic normality for series estimators. *Journal of Econometrics*, 79(1):147–168, July 1997.
- [Roc70] R. Tyrrell Rockafellar. *Convex Analysis*. Princeton University Press, 1970.
- [RR83] Paul R. Rosenbaum and Donald B. Rubin. The Central Role of the Propensity Score in Observational Studies for Causal Effects. *Biometrika*, 70(1):41–55, 1983.
- [SC08] Ingo Steinwart and Andreas Christmann. *Support Vector Machines*. Springer Science & Business Media, September 2008.
- [TB91] Paul Tseng and Dimitri P. Bertsekas. Relaxation Methods for Problems with Strictly Convex Costs and Linear Constraints. *Mathematics of Operations Research*, 16(3):462–481, 1991.
- [Tüb20] Stefan Tübbicke. Entropy Balancing for Continuous Treatments, May 2020.
- [vdV00] Aad van der Vaart. *Asymptotic Statistics*. Cambridge University Press, June 2000.
- [vdVW13] Aad van der Vaart and Jon Wellner. *Weak Convergence and Empirical Processes: With Applications to Statistics*. Springer Science & Business Media, March 2013.
- [VGC<sup>+</sup>20] Brian G. Vegetabile, Beth Ann Griffin, Donna L. Coffman, Matthew Cefalu, and Daniel F. McCaffrey. Nonparametric Estimation of Population Average Dose-Response Curves using Entropy Balancing Weights for Continuous Exposures, March 2020.
- [WZ19] Yixin Wang and José R. Zubizarreta. Minimal Dispersion Approximately Balancing Weights: Asymptotic Properties and Practical Considerations. *Biometrika*, page asz050, October 2019.
- [WZ23] Yixin Wang and José R. Zubizarreta. Large Sample Properties of Matching for Balance. *Statistica Sinica*, 2023.
- [ZP17] Qingyuan Zhao and Daniel Percival. Entropy balancing is doubly robust. *Journal of Causal Inference*, 5(1):20160010, September 2017.

- [Zub15] José R. Zubizarreta. Stable Weights that Balance Covariates for Estimation With Incomplete Outcome Data. *Journal of the American Statistical Association*, 110(511):910–922, July 2015.





# Index

- $B$  , vector of basis functions of the covariates, 8, 21
- $D_N$  , (random) data set without observed outcome, 7
- $F_{Y(1)}$  , distribution function of the potential outcome under treatment, 39
- $F_{Y(1)}(\cdot, x)$  , conditional distribution function of the potential outcome under treatment39
- $[x]^+$  , positive part of  $x \in \mathbb{R}$  , 14
- $\delta$  , (random) constraints vector , 8
- $\lesssim$  , lesser-or-equal-up-to-a-uniform-constant order , 26
- $\text{aff}(\cdot)$  , affine hull, 66
- $\text{dom } f$  , domain of a convex function, 67
- $\text{ri}(\cdot)$  , relative interior, 66
- $\sigma_\omega$  , support function, 68
- $\varphi$  , objective function of Problem 1, 9
- $e_i$  ,  $i$ -the unit vector, 33
- $f \square g$  , infimal convolution of  $f$  and  $g$  , 68
- $f^*$  , convex conjugate of  $f$ , 70
- $n$  , (random) number of treated units, 8
- $\partial f(x)$  , subdifferential of  $f$  at  $x$  , 68
- affine set, 66
- conjugate chain rule, 70
- conjugate sum rule, 70
- convex function, 67
- convex set, 65
- proper (convex) function, 67
- strictly convex, 67
- subgradient, 67