

Robust Weighting and Matching Techniques for Causal Inference in Observational Studies with Continuous Treatment

Universität Stuttgart



Universität Stuttgart

Ioan Scheffel

October 19, 2022

Contents

1	Chapter One Title	2
2	Convex Analysis	3
3	Random Matrix Inequality	4

Chapter 1

Chapter One Title

hello \mathbb{R}

Chapter 2

Convex Analysis

We begin by defining convex sets

Definition 1 A subset $\Omega \subseteq \mathbb{R}^n$ is called *CONVEX* if we have $\lambda x + (1 - \lambda)y \in \Omega$ for all $x, y \in \Omega$ and $\lambda \in (0, 1)$.

Clearly, the line segment $[a, b] := \{\lambda a + (1 - \lambda)b \mid \lambda \in [0, 1]\}$ is contained in Ω for all $a, b \in \Omega$ if and only if Ω is a convex set.

Next we define convex functions.

The concept of convex functions is closely related to convex sets.

The line segment between two points on the graph of a convex function lies on or above and does not intersect the graph.

In other words: The area above the graph of a convex function f is a convex set, i.e. the *epigraph* $\text{epi}(f) := \{(x, \alpha) \in \mathbb{R}^n \times \mathbb{R} \mid f(x) \leq \alpha\}$ is a convex set in \mathbb{R}^{n+1} .

Often an equivalent characterisation of convex functions is more useful.

Theorem 1 The convexity of a function $f : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$ on \mathbb{R}^n is equivalent to the following statement:

For all $x, y \in \mathbb{R}^n$ and $\lambda \in (0, 1)$ we have

$$f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y). \quad (2.1)$$

Chapter 3

Random Matrix Inequality

Theorem 2 *Let $(A_k)_{1 \leq k \leq n} \subseteq \mathbb{R}^{d_1 \times d_2}$ be a finite sequence of independent, random matrices. Assume that*

$$\mathbb{E}(A_k) = 0 \quad \text{and} \quad \|A_k\| \leq L \quad \text{for each } k \in \{1, \dots, n\}. \quad (3.1)$$

Introduce the random matrix

$$S := \sum_{k=1}^n A_k. \quad (3.2)$$

Let $v(S)$ be the matrix variance statistic of the sum:

$$v(S) := \max \left\{ \|\mathbb{E}(SS^T)\|, \|\mathbb{E}(S^T S)\| \right\} \quad (3.3)$$

$$= \max \left\{ \left\| \sum_{k=1}^n \mathbb{E}(A_k A_k^T) \right\|, \left\| \sum_{k=1}^n \mathbb{E}(A_k^T A_k) \right\| \right\}. \quad (3.4)$$

Then

$$\mathbb{E} \|S\| \leq \sqrt{2v(S) \log(d_1 + d_2)} + \frac{1}{3} L \log(d_1 + d_2). \quad (3.5)$$

Furthermore, for all $t \geq 0$,

$$\mathbb{P}(\|S\| \geq t) \leq (d_1 + d_2) \exp \left(\frac{-t^2/2}{v(S) + Lt/3} \right). \quad (3.6)$$