

# Alcohol Consumption in Schools

Hypothesis Heroes - Drew Davison, Lisa Zhang, Ellie Culman, Austin Chang

2023-11-14

## Introduction and data


All around the world, underage drinking among students is a significant public health issue and takes a huge toll on the quality of students' lives and education. There is plenty of existing documentation on the effects of socioeconomic status on risky drinking behavior amongst college students. For example, a paper by Susan E. Collins, PhD, professor and licensed clinical psychologist, found that individuals with lower socioeconomic status as well as people of racial and ethnic minorities and homelessness experience greater alcohol-related consequences. Additionally, studies from the National Institutes of Health have found that “aspects of college life—such as unstructured time, widespread availability of alcohol, inconsistent enforcement of underage drinking laws, and limited interactions with parents and other adults” drives up rates of underage drinking, and “college students have higher binge-drinking rates and a higher incidence of driving under the influence of alcohol than their noncollege peers.” Since most existing literature on underage student drinking focuses on college students, we wanted to examine the factors contributing to drinking amongst secondary school students. Our research question is as follows:

How do social indicators affect student alcohol consumption in secondary schools?

We hypothesize that factors like gender, familial status, family and school support, and other social and economic indicators will strongly influence the rates that secondary school students consume alcohol and that increased alcohol consumption is correlated with their school performance.

This data is from a Kaggle public dataset, originally sourced from the UC Irvine Machine Learning Repository. The data consists of information collected in 2008 on secondary school students from two schools in Portugal: Gabriel Pereira and Mousinho da Silveira. There are 649 observations, each one being a student, and 33 variables which cover a range of characteristics about each student's family, education, social situation, alcohol consumption, and grades in their Portuguese language class. Some key variables we will be examining are sex, male or female, Pstatus, whether their parents are together or apart, schoolsup, whether or not their school provides them extra academic support, famsup, whether or not their family provides

them extra academic support, Mjob and Fjob, the categories that their parents' jobs fall under, freetime and studytime, the amount of free time after school and amount of time spent studying on a scale of 1-5, number of absences, and amount of class failures, on a scale of 1-4. Our response variable is Walc, which is the students' average weekend alcohol consumption, on a scale of 1-5, with 5 being very high.



## Methodology

We are using a multivariable linear regression to examine the effects of the predictor variables sex, Pstatus, schoolsup, famsup, Mjob, Fjob, freetime, studytime, absences, and failures on the response variable Walc, or weekend alcohol consumption. We are using a linear model rather than logistic since Walc is not binary. In our model we are also including interactions between freetime and studytime, failures and studytime, failures and absences, as well as famsup and Mjob and famsup and Fjob. In our model we used `step_dummy()` to create dummy variables for all categorical variables, `step_interact()` to create all of our interaction variables, and `step_zv()` to remove all variables with only one value.

## Results

Linear Regression Model Specification (regression)

Computational engine: lm

### StepAIC Model

Start: AIC=184.95

Walc ~ sex + Pstatus + schoolsup + famsup + Mjob + Fjob + freetime +  
studytime + absences + failures

	Df	Sum of Sq	RSS	AIC
- Mjob	4	6.830	669.85	181.93
- failures	1	0.027	663.05	182.97
- schoolsup	1	0.874	663.90	183.59
- famsup	1	1.341	664.37	183.94
- Pstatus	1	1.646	664.67	184.16
- freetime	1	1.923	664.95	184.36
<none>			663.02	184.95
- Fjob	4	13.202	676.23	186.53
- absences	1	19.232	682.26	196.85
- studytime	1	21.440	684.46	198.42
- sex	1	41.513	704.54	212.47

Step: AIC=181.93

Walc ~ sex + Pstatus + schoolsup + famsup + Fjob + freetime +  
studytime + absences + failures

	Df	Sum of Sq	RSS	AIC
- failures	1	0.032	669.89	179.96
- schoolsup	1	0.558	670.41	180.34
- famsup	1	0.832	670.69	180.54
- Pstatus	1	1.544	671.40	181.05
- freetime	1	2.191	672.04	181.52
<none>			669.85	181.93
- Fjob	4	13.307	683.16	183.49
- absences	1	19.376	689.23	193.79
- studytime	1	20.916	690.77	194.88
- sex	1	45.151	715.00	211.63

Step: AIC=179.96

Walc ~ sex + Pstatus + schoolsup + famsup + Fjob + freetime +  
studytime + absences

	Df	Sum of Sq	RSS	AIC
- schoolsup	1	0.553	670.44	178.36
- famsup	1	0.837	670.72	178.56
- Pstatus	1	1.564	671.45	179.09
- freetime	1	2.159	672.04	179.52
<none>			669.89	179.96
- Fjob	4	13.279	683.16	181.50
- absences	1	19.355	689.24	191.80
- studytime	1	20.974	690.86	192.94
- sex	1	45.126	715.01	209.64

Step: AIC=178.36

Walc ~ sex + Pstatus + famsup + Fjob + freetime + studytime +  
absences

	Df	Sum of Sq	RSS	AIC
- famsup	1	0.747	671.19	176.90
- Pstatus	1	1.575	672.01	177.50
- freetime	1	2.243	672.68	177.98
<none>			670.44	178.36
- Fjob	4	13.004	683.44	179.69
- absences	1	19.067	689.51	189.99
- studytime	1	20.552	690.99	191.03
- sex	1	44.635	715.07	207.68

Step: AIC=176.9

Walc ~ sex + Pstatus + Fjob + freetime + studytime + absences

	Df	Sum of Sq	RSS	AIC
- Pstatus	1	1.464	672.65	175.96
- freetime	1	2.156	673.34	176.46
<none>			671.19	176.90
- Fjob	4	13.377	684.56	178.49
- absences	1	18.660	689.85	188.23
- studytime	1	21.734	692.92	190.39
- sex	1	45.597	716.78	206.84

Step: AIC=175.96

Walc ~ sex + Fjob + freetime + studytime + absences

	Df	Sum of Sq	RSS	AIC
- freetime	1	2.193	674.84	175.54
<none>			672.65	175.96
- Fjob	4	13.916	686.57	177.91
- absences	1	17.605	690.25	186.51
- studytime	1	21.518	694.17	189.26
- sex	1	48.428	721.08	207.75

Step: AIC=175.54

Walc ~ sex + Fjob + studytime + absences

	Df	Sum of Sq	RSS	AIC
<none>			674.84	175.54
- Fjob	4	13.363	688.20	177.07
- absences	1	17.354	692.20	185.88
- studytime	1	23.041	697.88	189.86
- sex	1	51.864	726.71	209.52

# A tibble: 8 x 5

term	estimate	std.error	statistic	p.value
<chr>	<dbl>	<dbl>	<dbl>	<dbl>
1 (Intercept)	2.15	0.249	8.65	7.67e-17
2 sexM	0.688	0.114	6.06	2.75e- 9
3 Fjobhealth	0.0785	0.338	0.232	8.17e- 1
4 Fjobother	0.287	0.211	1.36	1.76e- 1
5 Fjobservices	0.473	0.224	2.11	3.50e- 2
6 Fjobteacher	-0.164	0.314	-0.520	6.03e- 1
7 studytime	-0.274	0.0678	-4.04	6.23e- 5
8 absences	0.0400	0.0114	3.51	4.98e- 4

## Model

term	estimate	std.error	statistic	p.value
(Intercept)	2.211	0.265	8.354	0.000
studytime	-0.307	0.085	-3.622	0.000
absences	0.022	0.030	0.732	0.464
sex_M	0.695	0.114	6.092	0.000
Fjob_health	0.066	0.339	0.194	0.847
Fjob_other	0.285	0.212	1.347	0.179
Fjob_services	0.468	0.224	2.089	0.037
Fjob_teacher	-0.153	0.315	-0.484	0.629
studytime_x_absences	0.011	0.017	0.649	0.517

! Important

Before you submit, make sure your code chunks are turned off with `echo: false` and there are no warnings or messages with `warning: false` and `message: false` in the YAML.