

# Prediction and hypothesis testing

# Data

**Question:** What is the relationship between age and contraceptive use for women in Indonesia?

**Data:** 1473 Indonesian couples, with variables

- +  $Y_i$  = contraceptive method used (1 = no use, 2 = long-term, 3 = short-term)
- +  $X_i$  = Wife's age (numeric)

## Last time: Fitted model

$$\log\left(\frac{\hat{\pi}_{i(Short)}}{\hat{\pi}_{i(None)}}\right) = -8.234 + 0.456Age_i - 0.0065Age_i^2$$

$$\log\left(\frac{\hat{\pi}_{i(Long)}}{\hat{\pi}_{i(None)}}\right) = -5.083 + 0.366Age_i - 0.00628Age_i^2$$

## Last time: Predictions

##		Actual		
##	Prediction	None	Short	Long
##	None	342	166	189
##	Short	0	0	0
##	Long	287	167	322

How good are our predictions?

## Last time: Predictions

##		Actual		
##	Prediction	None	Short	Long
##	None	342	166	189
##	Short	0	0	0
##	Long	287	167	322

We can also assess our predictions by comparing to random guessing.

What are our predicted probabilities for each observation from random guessing?

# Random guessing

- + If we don't have any data, our estimated probability would be  $1/3$  for each level
- + If we have data but we don't use age, our estimated probability for each level is just the proportion of observations in that group:

```
table(cmc_data$Choice)/nrow(cmc_data)
```

```
##  
##      None      Short      Long  
## 0.4270197 0.2260692 0.3469111
```

# Class activity

[https://sta214-f22.github.io/class\\_activities/ca\\_lecture\\_16.html](https://sta214-f22.github.io/class_activities/ca_lecture_16.html)

## Class activity

What would our confusion matrix look like if our predictions randomly assigned each person to one of the three categories, with a  $1/3$  chance for each category?



## Class activity

What would our confusion matrix look like if our predictions randomly assigned each person to one of the three categories, with a  $1/3$  chance for each category?

Something like

		Actual		
		None	Short	Long
Predicted	None	210	111	170
	Short	210	111	170
	Long	209	111	171

## Class activity

		Actual		
		None	Short	Long
Predicted	None	210	111	170
	Short	210	111	170
	Long	209	111	171

What is the accuracy of our predictions in this confusion matrix?

## Class activity

What would our confusion matrix look like if for every individual, we just predicted the most common contraception choice in the data?

## Class activity

What would our confusion matrix look like if for every individual, we just predicted the most common contraception choice in the data?

The most common choice is None, so

		Actual		
		None	Short	Long
Predicted	None	629	333	511
	Short	0	0	0
	Long	0	0	0

## Class activity

		Actual		
		None	Short	Long
Predicted	None	629	333	511
	Short	0	0	0
	Long	0	0	0

What is the accuracy of our predictions in this confusion matrix?

## Class activity

Do we do better than random guessing?

# Moral

- + By itself, accuracy isn't particularly useful for summarizing prediction performance
- + It is helpful to interpret accuracy in relation to simple random guessing. Our model isn't very good if we can't beat a random guess
- + We also need to look at predictive ability for each class

# Hypothesis testing

**Research question:** Is there a relationship between age and contraceptive choice?

What are my steps to answer this question with a hypothesis test?



# Specify hypotheses

**Research question:** Is there a relationship between age and contraceptive choice?

$$\log\left(\frac{\pi_{i(Long)}}{\pi_{i(None)}}\right) = \beta_{0(Long)} + \beta_{1(Long)}Age_i + \beta_{2(Long)}Age_i^2$$

$$\log\left(\frac{\pi_{i(Short)}}{\pi_{i(None)}}\right) = \beta_{0(Short)} + \beta_{1(Short)}Age_i + \beta_{2(Short)}Age_i^2$$

What should our null and alternative hypotheses be?

## Specify hypotheses

$$\log\left(\frac{\pi_{i(Long)}}{\pi_{i(None)}}\right) = \beta_{0(Long)} + \beta_{1(Long)}Age_i + \beta_{2(Long)}Age_i^2$$

$$\log\left(\frac{\pi_{i(Short)}}{\pi_{i(None)}}\right) = \beta_{0(Short)} + \beta_{1(Short)}Age_i + \beta_{2(Short)}Age_i^2$$

$$H_0 : \beta_{1(Short)} = \beta_{2(Short)} = \beta_{1(Long)} = \beta_{2(Long)} = 0$$

$$H_A : \text{at least one of } \beta_{1(Short)}, \beta_{2(Short)}, \beta_{1(Long)}, \beta_{2(Long)} \neq 0$$

What are the full and reduced models?

# Test statistic

What test can I use to compare nested models?

## Drop in deviance

```
m1 <- multinom(Choice ~ WifeAge + I(WifeAge^2),  
               data = cmc_data)
```

```
summary(m1)
```

```
...
```

```
##
```

```
## Residual Deviance: 3015.821
```

```
...
```

Deviance for full model: 3015.821

How would we fit the reduced model in R?

## Drop in deviance

```
m0 <- multinom(Choice ~ 1,  
                data = cmc_data)
```

```
summary(m0)
```

```
...  
##  
## Residual Deviance: 3142.726  
...
```

Reduced model deviance: 3142.726

Drop in deviance:  $G = 3142.726 - 3015.821 = 126.905$

What distribution do we use to calculate the p-value?

## Calculating a p-value

Under  $H_0$ ,  $G \sim \chi_q^2$ , where  $q$  is the number of parameters tested.

Here  $q = 4$  (2 parameters for each log relative risk model)

```
pchisq(126.905, df=4, lower.tail=F)
```

```
## [1] 1.787184e-26
```

So we have very strong evidence that there is a relationship between age and contraceptive choice.