

Logistic regression interpretation

Ciaran Evans

Logistic regression model

Y_i = dengue status (0 = negative, 1 = positive)

Age_i = age (in years)

Random component: $Y_i \sim \text{Bernoulli}(\pi_i)$

Systematic component: $\log\left(\frac{\pi_i}{1 - \pi_i}\right) = \beta_0 + \beta_1 Age_i$

Where did the ε term go?

Linear regression, written two ways:

Option 1:

$$Y_i = \beta_0 + \beta_1 X_i + \varepsilon_i \quad \varepsilon_i \sim N(0, \sigma^2)$$

Option 2:

$$Y_i \sim N(\mu_i, \sigma^2)$$

$$\mu_i = \beta_0 + \beta_1 X_i$$

Fitting the logistic regression model

$$Y_i \sim \text{Bernoulli}(\pi_i)$$

$$\log\left(\frac{\pi_i}{1 - \pi_i}\right) = \beta_0 + \beta_1 \text{Age}_i$$

```
m1 <- glm(Dengue ~ Age, data = dengue,  
          family = binomial)  
summary(m1)
```

Fitting the logistic regression model

$$Y_i \sim \text{Bernoulli}(\pi_i)$$

$$\log\left(\frac{\pi_i}{1 - \pi_i}\right) = \beta_0 + \beta_1 \text{Age}_i$$

```
m1 <- glm(Dengue ~ Age, data = dengue,  
          family = binomial)  
summary(m1)
```

...

Coefficients:

##		Estimate	Std. Error	z value	Pr(> z)	
##	(Intercept)	-2.454345	0.075068	-32.70	<2e-16	***
##	Age	0.217312	0.008826	24.62	<2e-16	***

...

Activity 1

Work on the activity (handout) with a neighbor, then we will discuss as a class.

Activity

$$\log\left(\frac{\hat{\pi}_i}{1 - \hat{\pi}_i}\right) = -2.45 + 0.22 \text{ Age}_i$$

Does the probability of dengue increase or decrease with age?

Activity

$$\log\left(\frac{\hat{\pi}_i}{1 - \hat{\pi}_i}\right) = -2.45 + 0.22 \text{ Age}_i$$

What are the predicted odds of dengue for a 5 year old patient?

Activity

$$\log\left(\frac{\hat{\pi}_i}{1 - \hat{\pi}_i}\right) = -2.45 + 0.22 \text{ Age}_i$$

For a 5 year old patient: is the predicted *probability* of dengue > 0.5 , < 0.5 , or $= 0.5$?

Activity 2

Work on the activity (handout) with a neighbor, then we will discuss as a class.

Activity

$$\log\left(\frac{\hat{\pi}_i}{1 - \hat{\pi}_i}\right) = -2.45 + 0.22 \text{ Age}_i$$

What is the predicted probability of dengue for a 10 year old patient?

Activity

$$\log\left(\frac{\hat{\pi}_i}{1 - \hat{\pi}_i}\right) = -2.45 + 0.22 \text{ Age}_i$$

Suppose we want to identify patients for whom the predicted probability of dengue is at least 0.5. What age range should we focus on?

Activity

$$\log\left(\frac{\hat{\pi}_i}{1 - \hat{\pi}_i}\right) = -2.45 + 0.22 \text{ Age}_i$$

Compare the odds of dengue for a 12 year old patient to the odds of dengue for an 11 year old patient. What do you notice?

Interpretation

$$\log\left(\frac{\hat{\pi}_i}{1 - \hat{\pi}_i}\right) = -2.45 + 0.22 \text{ Age}_i$$

Recap: logistic regression

$$Y_i \sim \textit{Bernoulli}(\pi_i)$$

$$\log\left(\frac{\pi_i}{1 - \pi_i}\right) = \beta_0 + \beta_1 \textit{Age}_i$$

Adding more variables

Now let's add WBC as a variable to the model:

```
m2 <- glm(Dengue ~ Age + WBC, data = dengue,  
          family = binomial)  
summary(m2)
```

$$\log\left(\frac{\hat{\pi}_i}{1 - \hat{\pi}_i}\right) = 0.34 + 0.15 \text{ Age}_i - 0.31 \text{ WBC}_i$$

How should I interpret each coefficient in the fitted model?

Adding more variables

What about a *categorical* predictor?

```
m3 <- glm(Dengue ~ Age + Abdominal, data = dengue,  
          family = binomial)  
summary(m3)
```

$$\log\left(\frac{\hat{\pi}_i}{1 - \hat{\pi}_i}\right) = -0.25 + 0.22 \text{ Age}_i + 0.21 \text{ Abdominal}_i$$

How should I interpret the coefficient on abdominal pain?

For next week

Reading: in the textbook, read

+ sections 6.7.1 - 6.7.3

+ sections 2.2 - 2.4

Class on Monday: lab day (most of the class period to work on HW 1)