


Model diagnostics

Ciaran Evans

Model assessment

Essentially, all models are wrong, but some are useful. (Box and Draper, 1987)




Question: Is our model useful?

Limitations of empirical logit plots


- ▶ Does not assess a fitted model (performed *before* fitting the model)
- ▶ One point per *bin*, not one point per *observation*
- ▶ Can only plot one or two explanatory variables at a time

Why residuals in linear regression are nice


Dataset 1




Dataset 2



Residuals 1



Residuals 2




Randomized quantile residuals

Simulated data: $\log\left(\frac{p_i}{1-p_i}\right) = -1 + 2X_i$

Model fit: $\log\left(\frac{p_i}{1-p_i}\right) = \beta_0 + \beta_1 X_i$

Correctly specified model




Randomized quantile residuals

Simulated data: $\log\left(\frac{p_i}{1-p_i}\right) = -1 + 2X_i^2$


Model fit: $\log\left(\frac{p_i}{1-p_i}\right) = \beta_0 + \beta_1 X_i$

Mis-specified model




Example: dengue data

Model: $\log\left(\frac{p_i}{1-p_i}\right) = \beta_0 + \beta_1 \text{WBC}_i$



Example: dengue data

Model: $\log\left(\frac{p_i}{1-p_i}\right) = \beta_0 + \beta_1 \text{WBC}_i$




Question: How can we decide whether a plot is consistent with a correctly-specified model?

Parametric bootstrap


The *parametric bootstrap* generates data from the *fitted* model:

Visual inference

```
## decrypt("QUg2 qFyF Rx 8tLRyRtx Ko")
```



Visual inference



```
decrypt("QUg2 qFyF Rx 8tLRyRtx Ko")
```

```
## [1] "True data in position 4"
```