

# STA286 Lecture 11

Neil Montgomery

Last edited: 2017-02-02 13:00

reminder “theorem”

$$E(g(X)) = \begin{cases} \sum_x g(x)p(x) & : \text{discrete} \\ \int_{-\infty}^{\infty} g(x)f(x) dx & : \text{continuous} \end{cases}$$

## “constant” random variables

From last time:  $E(a + bX) = a + bE(X)$ . The left hand side is a bit of an odd expression.

## “constant” random variables

From last time:  $E(a + bX) = a + bE(X)$ . The left hand side is a bit of an odd expression.

For convenience we can treat constants as random variables. Formally, we can consider a random variable  $X$  that is always, say, some real  $a$  no matter what. The pmf of  $X$  is:

$$P(X = a) = 1$$

and 0 otherwise.

## “constant” random variables

From last time:  $E(a + bX) = a + bE(X)$ . The left hand side is a bit of an odd expression.

For convenience we can treat constants as random variables. Formally, we can consider a random variable  $X$  that is always, say, some real  $a$  no matter what. The pmf of  $X$  is:

$$P(X = a) = 1$$

and 0 otherwise.

Informally, for convenience, we dispense with the  $X$  notation and just treat the constant  $a$  as a random variable, allowing for statements like:

$$E(a) = a$$

## expected values and joint distributions

Given  $X$  and  $Y$  with joint density  $f(x, y)$ , and given a function  $g : \mathbb{R}^2 \rightarrow \mathbb{R}$ , a sophisticated application of the theorem from last time is:

$$E(g(X, Y)) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(x, y) f(x, y) dx dy$$

(discrete version is the same, with sums)

## first key example

Consider  $g(x, y) = x + y$ .

$$E(g(X, Y)) = E(X + Y) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (x + y)f(x, y) dx dy$$

## first key example

Consider  $g(x, y) = x + y$ .

$$\begin{aligned} E(g(X, Y)) &= E(X + Y) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (x + y) f(x, y) \, dx \, dy \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} x f(x, y) \, dy \, dx + \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} y f(x, y) \, dx \, dy \end{aligned}$$



## first key example

Consider  $g(x, y) = x + y$ .

$$\begin{aligned} E(g(X, Y)) &= E(X + Y) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (x + y) f(x, y) dx dy \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} x f(x, y) dy dx + \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} y f(x, y) dx dy \\ &= \int_{-\infty}^{\infty} x \int_{-\infty}^{\infty} f(x, y) dy dx + \int_{-\infty}^{\infty} y \int_{-\infty}^{\infty} f(x, y) dx dy \end{aligned}$$

## first key example

Consider  $g(x, y) = x + y$ .

$$\begin{aligned} E(g(X, Y)) &= E(X + Y) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (x + y) f(x, y) dx dy \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} x f(x, y) dy dx + \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} y f(x, y) dx dy \\ &= \int_{-\infty}^{\infty} x \int_{-\infty}^{\infty} f(x, y) dy dx + \int_{-\infty}^{\infty} y \int_{-\infty}^{\infty} f(x, y) dx dy \\ &= \int_{-\infty}^{\infty} x f_X(x) dx + \int_{-\infty}^{\infty} y f_Y(y) dy \end{aligned}$$

## first key example

Consider  $g(x, y) = x + y$ .

$$\begin{aligned} E(g(X, Y)) &= E(X + Y) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (x + y) f(x, y) dx dy \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} x f(x, y) dy dx + \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} y f(x, y) dx dy \\ &= \int_{-\infty}^{\infty} x \int_{-\infty}^{\infty} f(x, y) dy dx + \int_{-\infty}^{\infty} y \int_{-\infty}^{\infty} f(x, y) dx dy \\ &= \int_{-\infty}^{\infty} x f_X(x) dx + \int_{-\infty}^{\infty} y f_Y(y) dy \\ &= E(X) + E(Y) \end{aligned}$$

## second key example

Suppose  $X$  and  $Y$  are independent. Consider  $g(x, y) = xy$ .

$$E(g(X, Y)) = E(XY) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy f(x, y) dx dy$$

## second key example

Suppose  $X$  and  $Y$  are independent. Consider  $g(x, y) = xy$ .

$$\begin{aligned} E(g(X, Y)) &= E(XY) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy f(x, y) dx dy \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy f_X(x) f_Y(y) dx dy \end{aligned}$$

## second key example

Suppose  $X$  and  $Y$  are independent. Consider  $g(x, y) = xy$ .

$$\begin{aligned} E(g(X, Y)) &= E(XY) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy f(x, y) dx dy \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy f_X(x) f_Y(y) dx dy \\ &= \int_{-\infty}^{\infty} x f_X(x) dx \int_{-\infty}^{\infty} y f_Y(y) dy \end{aligned}$$

## second key example

Suppose  $X$  and  $Y$  are independent. Consider  $g(x, y) = xy$ .

$$\begin{aligned} E(g(X, Y)) &= E(XY) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy f(x, y) dx dy \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} xy f_X(x) f_Y(y) dx dy \\ &= \int_{-\infty}^{\infty} x f_X(x) dx \int_{-\infty}^{\infty} y f_Y(y) dy \\ &= E(X)E(Y) \end{aligned}$$

## notation

It is common to use  $\mu$  as a shorter stand-in for  $E(X)$ .

I'm not so convinced this is a good idea.



measuring variation

## BIG MONEY versus BIG MONEY SCHNAPPS VERSION

$$E(X) = E(Y) = 0$$

	Roll	1, 2	3	4, 5	6
BIG MONEY Outcome X \$		-2	0	1	2
BIG MONEY SCHNAPPS Outcome Y \$		-200	0	100	200
Probability		2/6	1/6	2/6	1/6

## BIG MONEY versus BIG MONEY SCHNAPPS VERSION

$$E(X) = E(Y) = 0$$

	Roll	1, 2	3	4, 5	6
BIG MONEY Outcome $X$ \$		-2	0	1	2
BIG MONEY SCHNAPPS Outcome $Y$ \$		-200	0	100	200
Probability		2/6	1/6	2/6	1/6

But  $Y$  is clearly a riskier game. The distribution is more spread out, by a factor of 100.

## BIG MONEY versus BIG MONEY SCHNAPPS VERSION

$$E(X) = E(Y) = 0$$

	Roll	1, 2	3	4, 5	6
BIG MONEY Outcome $X$ \$		-2	0	1	2
BIG MONEY SCHNAPPS Outcome $Y$ \$		-200	0	100	200
Probability		2/6	1/6	2/6	1/6

But  $Y$  is clearly a riskier game. The distribution is more spread out, by a factor of 100. The question is—how to measure this difference in variation?

## variance

Recall the sample variance, expressed a little differently:

$$s^2 = \sum_{i=1}^n (x_i - \bar{x})^2 \frac{1}{n-1}$$

This is a weighted sum of squared deviations with weights  $w_i = 1/(n-1)$

## variance

Recall the sample variance, expressed a little differently:

$$s^2 = \sum_{i=1}^n (x_i - \bar{x})^2 \frac{1}{n-1}$$

This is a weighted sum of squared deviations with weights  $w_i = 1/(n-1)$

The theoretical analogue is called the *variance*:

$$\text{Var}(X) = E \left( (X - E(X))^2 \right) = E \left( (X - \mu)^2 \right)$$

provided the expectations all exist.

## variance

Recall the sample variance, expressed a little differently:

$$s^2 = \sum_{i=1}^n (x_i - \bar{x})^2 \frac{1}{n-1}$$

This is a weighted sum of squared deviations with weights  $w_i = 1/(n-1)$

The theoretical analogue is called the *variance*:

$$\text{Var}(X) = E\left((X - E(X))^2\right) = E\left((X - \mu)^2\right)$$

provided the expectations all exist.

It is also possible to view this as an application of the  $E(g(X))$  theorem with  $g(x) = (x - \mu)^2$ .

## examples - BIG MONEY

$$\text{Var}(X) = (-2 - 0)^2 \frac{2}{6} + (0 - 0)^2 \frac{1}{6} + (1 - 0)^2 \frac{2}{6} + (2 - 0)^2 \frac{1}{6} = \frac{14}{6}$$



## examples - BIG MONEY

$$\text{Var}(X) = (-2 - 0)^2 \frac{2}{6} + (0 - 0)^2 \frac{1}{6} + (1 - 0)^2 \frac{2}{6} + (2 - 0)^2 \frac{1}{6} = \frac{14}{6}$$

Schnapps version:

$$\text{Var}(Y) = (-200 - 0)^2 \frac{2}{6} + (0 - 0)^2 \frac{1}{6} + (100 - 0)^2 \frac{2}{6} + (200 - 0)^2 \frac{1}{6} = \frac{140000}{6}$$

## how to actually calculate variance

Using the two rules  $E(a + bX) = a + bE(X)$  and  $E(X + Y) = E(X) + E(Y)$  we can derive the better way to calculate variance:

$$\begin{aligned}\text{Var}(X) &= E\left((X - \mu)^2\right) \\ &= E\left(X^2 - 2X\mu + \mu^2\right)\end{aligned}$$

## how to actually calculate variance

Using the two rules  $E(a + bX) = a + bE(X)$  and  $E(X + Y) = E(X) + E(Y)$  we can derive the better way to calculate variance:

$$\begin{aligned}\text{Var}(X) &= E\left((X - \mu)^2\right) \\ &= E\left(X^2 - 2X\mu + \mu^2\right) \\ &= E\left(X^2\right) - 2E(X)\mu + \mu^2\end{aligned}$$

## how to actually calculate variance

Using the two rules  $E(a + bX) = a + bE(X)$  and  $E(X + Y) = E(X) + E(Y)$  we can derive the better way to calculate variance:

$$\begin{aligned}\text{Var}(X) &= E\left((X - \mu)^2\right) \\ &= E\left(X^2 - 2X\mu + \mu^2\right) \\ &= E\left(X^2\right) - 2E(X)\mu + \mu^2 \\ &= E\left(X^2\right) - \mu^2\end{aligned}$$

## how to actually calculate variance

Using the two rules  $E(a + bX) = a + bE(X)$  and  $E(X + Y) = E(X) + E(Y)$  we can derive the better way to calculate variance:

$$\begin{aligned}\text{Var}(X) &= E\left((X - \mu)^2\right) \\ &= E\left(X^2 - 2X\mu + \mu^2\right) \\ &= E\left(X^2\right) - 2E(X)\mu + \mu^2 \\ &= E\left(X^2\right) - \mu^2\end{aligned}$$

Or as I prefer:  $\text{Var}(X) = E(X^2) - E(X)^2$

## another variance example

Gas pipes revisited:

Y	X			Marginal
	1	1.5	1.75	
0.5	0.075	0.100	0.150	0.325
1	0.110	0.080	0.090	0.280
2	0.160	0.140	0.095	0.395
Marginal	0.345	0.320	0.335	1.000

## another variance example

Gas pipes revisited:

Y	X			Marginal
	1	1.5	1.75	
0.5	0.075	0.100	0.150	0.325
1	0.110	0.080	0.090	0.280
2	0.160	0.140	0.095	0.395
Marginal	0.345	0.320	0.335	1.000

$$E(X) = 1 \cdot 0.345 + 1.5 \cdot 0.320 + 1.75 \cdot 0.335 = 1.41125$$

## another variance example

Gas pipes revisited:

Y	X			Marginal
	1	1.5	1.75	
0.5	0.075	0.100	0.150	0.325
1	0.110	0.080	0.090	0.280
2	0.160	0.140	0.095	0.395
Marginal	0.345	0.320	0.335	1.000

$$E(X) = 1 \cdot 0.345 + 1.5 \cdot 0.320 + 1.75 \cdot 0.335 = 1.41125$$

$$E(X^2) = 1^2 \cdot 0.345 + 1.5^2 \cdot 0.320 + 1.75^2 \cdot 0.335 = 2.0909375$$



## another variance example

Gas pipes revisited:

Y	X			Marginal
	1	1.5	1.75	
0.5	0.075	0.100	0.150	0.325
1	0.110	0.080	0.090	0.280
2	0.160	0.140	0.095	0.395
Marginal	0.345	0.320	0.335	1.000

$$E(X) = 1 \cdot 0.345 + 1.5 \cdot 0.320 + 1.75 \cdot 0.335 = 1.41125$$

$$E(X^2) = 1^2 \cdot 0.345 + 1.5^2 \cdot 0.320 + 1.75^2 \cdot 0.335 = 2.0909375$$

$$\text{Var}(X) = 0.0993109$$

## variance non-example

Reconsider  $X$  with density  $f(x) = 2x^{-3}$  for  $x > 1$ , and 0 otherwise. Last time we found  $E(X) = 2$ .

But  $X$  does not have a variance, because  $E(X^2)$  does not exist.

notation; standard deviation

It is common to use  $\sigma^2$  as a shorter stand-in for  $\text{Var}(X)$ .

## notation; standard deviation

It is common to use  $\sigma^2$  as a shorter stand-in for  $\text{Var}(X)$ .

I'm not so convinced this is a good idea.

## notation; standard deviation

It is common to use  $\sigma^2$  as a shorter stand-in for  $\text{Var}(X)$ .

I'm not so convinced this is a good idea.

Recall the “unit” problem with the sample variance  $s^2$ ; hence the use of the sample standard deviation  $s = \sqrt{s^2}$ .

## notation; standard deviation

It is common to use  $\sigma^2$  as a shorter stand-in for  $\text{Var}(X)$ .

I'm not so convinced this is a good idea.

Recall the “unit” problem with the sample variance  $s^2$ ; hence the use of the sample standard deviation  $s = \sqrt{s^2}$ .

The theoretical analogue is the *standard deviation*, defined as:

$$\text{SD}(X) = \sqrt{\text{Var}(X)}$$

## notation; standard deviation

It is common to use  $\sigma^2$  as a shorter stand-in for  $\text{Var}(X)$ .

I'm not so convinced this is a good idea.

Recall the “unit” problem with the sample variance  $s^2$ ; hence the use of the sample standard deviation  $s = \sqrt{s^2}$ .

The theoretical analogue is the *standard deviation*, defined as:

$$\text{SD}(X) = \sqrt{\text{Var}(X)}$$

It is common to use  $\sigma$  as a shorter stand-in for  $\text{SD}(X)$ .

## notation; standard deviation

It is common to use  $\sigma^2$  as a shorter stand-in for  $\text{Var}(X)$ .

I'm not so convinced this is a good idea.

Recall the “unit” problem with the sample variance  $s^2$ ; hence the use of the sample standard deviation  $s = \sqrt{s^2}$ .

The theoretical analogue is the *standard deviation*, defined as:

$$\text{SD}(X) = \sqrt{\text{Var}(X)}$$

It is common to use  $\sigma$  as a shorter stand-in for  $\text{SD}(X)$ .

I will leave to you to guess as to whether or not I am convinced this is a good idea.



## variance rules

$$\text{Var}(a + bX) = b^2 \text{Var}(X)$$

## variance rules

$$\text{Var}(a + bX) = b^2 \text{Var}(X)$$

What about  $\text{Var}(X + Y)$ ?