

STA303: M3 Team Up! activity

I grow rows and rows of roses

Instructions

To participate in this activity you will need have to two windows readily available to you:

- 1) Your Zoom window
- 2) The Team Up! activity linked from Quercus in a browser window for voting.

I would recommend that one member of the team shares their screen with the rest of the team and shows this activity where you can see the question and options.

Note 1: In the Team Up! activity you will just see the letters for the questions, not the options themselves.

Note 2: There are hints for some questions, but if you get really stuck, please use the ‘Ask for Help’ option in Zoom, our join the Zoom main room.

Data



Checkmate



Devotion



Hearts



Wanted

1. Pipe this changed `roses` data directly into a `ggplot`.
2. Put `greenhouse` on your x-axis, `profit` on your y, and colour by `variety`.
3. Use points as your geometry. (There is probably a bit of overprinting, but we won't worry about that here.)
4. You can add `theme_minimal()` if you like, but use the default colour palette.

Which greenhouse and variety combination had the lowest profit blue dot on your graph?

A. Devotion variety in Central greenhouse. B. Wanted variety in the West greenhouse. C. Hearts variety in the South greenhouse. D. Impossible to say.

Question 2

Considering the `roses` data described in the Data section and the plot you made, which ONE of the following statements best describes an appropriate analysis for this data?

- A. We can approach this analysis with linear regression (all fixed effects) as all our assumptions appear to be valid.
- B. We should make `variety` a fixed effect and `greenhouse` as a random effect.
- C. We should make `greenhouse` a random effect and `variety` a fixed effect and consider including a `greenhouse-variety` interaction (also a random effect).
- D. We should make everything a random effect.

Question 3

The below code aggregates the data to get the average profit by greenhouse and variety.

```
## # A tibble: 20 x 3
##   variety    greenhouse profit_avg
##   <chr>      <chr>      <dbl>
## 1 Checkmate central      23.8
## 2 Checkmate east        22.1
## 3 Checkmate north       29.5
## 4 Checkmate south       22.3
## 5 Checkmate west        18.1
## 6 Devotion  central      21.4
## 7 Devotion  east        22.2
## 8 Devotion  north       24.8
## 9 Devotion  south       22.4
## 10 Devotion west        19.1
## 11 Hearts   central      26.8
## 12 Hearts   east        27.7
## 13 Hearts   north       35.3
## 14 Hearts   south       22.3
## 15 Hearts   west        26.3
## 16 Wanted   central      16.0
## 17 Wanted   east        15.7
## 18 Wanted   north       24.3
## 19 Wanted   south       15.9
## 20 Wanted   west        11.0
```

How many values are averaged to get each observation of profit_avg in agg_int?

- A. 4
- B. 5
- C. 8
- D. 32

Question 4

Suppose an appropriate model for this data is:

$$y_{ijk} = \mu + \alpha_i + b_j + (\alpha b)_{ij} + \epsilon_{ijk}$$

where:

- μ is the grand mean profit of strawberries, in thousands of dollars, across all the data.
- α_i is the fixed effect of variety.
- b_j is the random effect of greenhouse, $N(0, \sigma_b^2)$.
- $(\alpha b)_{ij}$ is in interaction between variety and greenhouse, $N(0, \sigma_{\alpha b}^2)$.
- ϵ_{ijk} is the error term, $N(0, \sigma^2)$.

Which ONE of the following statements is FALSE? (Notice, FALSE *not* TRUE!)

- A. α_i and $(\alpha b)_{ij}$ are the only fixed effects in this model.
- B. μ is the same for all values of i, j and k.
- C. There are three random quantities being estimated.
- D. $b_j \sim N(0, \sigma_b^2)$ in this model.

Question 5

This question takes you through the ‘teaching and learning’ world version of thinking about sources of variance with random effects.

Which of the following correctly describes what each of the values being calculated under A, B and C below is matches with in the model specification? (Scroll!)

- A. A is $\hat{\sigma}^2$, B is $\hat{\sigma}_b^2$, C is $\hat{\sigma}_{\alpha b}^2$.
- B. A is $\hat{\sigma}_b^2$, B is $\hat{\sigma}^2$, C is $\hat{\sigma}_{\alpha b}^2$.
- C. A is $\hat{\sigma}_{\alpha b}^2$, B is $\hat{\sigma}_b^2$, C is $\hat{\sigma}^2$.
- D. A is $\hat{\sigma}_{\alpha b}^2$, B is not actually one of the variances for our random effects, C is $\hat{\sigma}^2$.

Question 6

Suppose our data was presented as follows. We are still interested in making the same plot and doing the same models as before.

observation	Checkmate	Devotion	Hearts	Wanted
north, plot group 1	29.49	23.44	32.77	24.64
north, plot group 2	27.60	20.42	36.41	27.38

observation	Checkmate	Devotion	Hearts	Wanted
north, plot group 3	26.67	23.86	34.05	25.21
north, plot group 4	30.53	28.74	35.06	27.29
north, plot group 5	28.60	29.10	38.84	21.98
north, plot group 6	30.01	24.73	35.12	23.56
north, plot group 7	33.30	23.86	33.98	19.85
north, plot group 8	29.52	24.11	35.94	24.88
east, plot group 1	20.67	26.56	32.15	14.46
east, plot group 2	25.45	22.04	26.92	20.31
east, plot group 3	20.08	20.43	28.88	15.07
east, plot group 4	23.47	20.78	27.10	14.06
east, plot group 5	18.79	20.16	28.30	16.04
east, plot group 6	23.34	23.60	25.80	13.80
east, plot group 7	21.17	23.59	25.60	17.74
east, plot group 8	23.82	20.48	26.93	13.91
south, plot group 1	25.01	25.27	22.00	16.59
south, plot group 2	20.34	21.86	24.00	15.70
south, plot group 3	20.33	20.25	19.61	19.11
south, plot group 4	21.08	22.72	25.24	15.12
south, plot group 5	20.58	18.89	24.08	13.81
south, plot group 6	21.11	21.63	20.57	15.47
south, plot group 7	24.35	25.45	23.21	16.38
south, plot group 8	25.81	23.30	19.32	14.93
west, plot group 1	15.34	18.76	28.10	11.77
west, plot group 2	17.62	20.19	29.68	9.60
west, plot group 3	20.22	19.09	24.92	11.26
west, plot group 4	16.87	20.10	22.88	8.88
west, plot group 5	15.04	19.14	27.65	10.18
west, plot group 6	21.45	19.56	24.28	11.33
west, plot group 7	22.27	16.08	25.32	15.54
west, plot group 8	15.75	19.88	27.74	9.70
central, plot group 1	23.03	22.76	25.75	15.02
central, plot group 2	27.94	22.54	28.18	16.20
central, plot group 3	22.52	23.61	26.80	15.21
central, plot group 4	23.64	17.95	28.30	17.06
central, plot group 5	26.36	22.99	29.59	16.72
central, plot group 6	21.95	18.82	23.02	15.56
central, plot group 7	22.46	22.55	26.93	13.51
central, plot group 8	22.68	19.67	26.09	18.35

Which ONE of the following statements best describes our situation with this new version of the dataset?

- A. This dataset is also tidy for our purposes.
- B. This dataset is NOT tidy for our purposes because each observational unit has too many observations.
- C. This dataset is NOT tidy for our purposes because it is not the case that each value is in one and only one cell.
- D. This dataset is NOT tidy for our purposes because it is not the case that each variable is in one and only one column.