

## Activity: Motivating Gaussian quadrature

**Instructions:** Submit your work as a single PDF. You may choose to either hand-write your work and submit a PDF scan, or type your work using LaTeX and submit the resulting PDF. See the course website for a homework template file and instructions on getting started with LaTeX and Overleaf.

### Part 1

Suppose we observe  $n$  points  $(x_1, y_1), \dots, (x_n, y_n)$ , and let

$$L_{n,i}(x) = \prod_{k:k \neq i} \frac{(x - x_k)}{(x_i - x_k)}$$

This function  $L_{n,i}(x)$  is a polynomial, and it turns out that  $L_{n,i}(x)$  plays an important role in deriving Gaussian quadrature. To begin, let's explore some properties of  $L_{n,i}(x)$ .

1. Show that  $L_{n,i}(x_i) = 1$
2. Show that  $L_{n,i}(x_k) = 0$  for all  $k \neq i$

Now let

$$q(x) = \sum_{i=1}^n y_i L_{n,i}(x)$$

$q(x)$  is also a polynomial.

3. Using the results from questions 1 and 2, calculate  $q(x_1), \dots, q(x_n)$ .

## Plotting $q(x)$

The following code provides a function 'q' to plot  $q(x)$  between -1 and 1:

```
# calculate q at a single point
# x: point to evaluate q(x)
# xi: the points x1,...,xn
# yi: the points y1,...,yn
q_helper <- function(x, xi, yi){
  lp <- sapply(1:length(xi),
               function(i){prod((x - xi[-i])/(xi[i] - xi[-i]))})
  sum(yi*lp)
}

# calculate q at a vector of new points
# x: point to evaluate q(x)
# xi: the points x1,...,xn
# yi: the points y1,...,yn
q <- function(x, xi, yi){
  sapply(x, function(t){q_helper(t, xi, yi)})
}

xi <- seq(-1, 1, length.out = 5)
yi <- xi^3
plot(xi, yi, pch=16)

x <- seq(-1, 1, 0.01)
lines(x, q(x, xi, yi))
```

4. Run the code to add  $q(x)$  to the plot with the five points  $(x_1, y_1), \dots, (x_n, y_n)$ . What do you notice about  $q(x)$ ?
  
5. To your plot from question 4, add the curve  $y = x^3$  (the original function from which the  $(x_i, y_i)$  were sampled). Comment on  $q(x)$  vs.  $x^3$ .

## Another example

The following code samples  $n = 4$  points  $(x_1, y_1), \dots, (x_n, y_n)$  from the 7th degree polynomial

$$f(x) = 10(x^7 - 1.6225x^5 + 0.79875x^3 - 0.113906x)$$

and plots both the true polynomial  $f(x)$  (in red) and the polynomial  $q(x)$  (in black):

```
f <- function(x){  
  10*(x^7 -1.6225*x^5 +0.79875*x^3 - 0.113906*x)  
}  
  
n <- 4  
xi <- seq(-1, 1, length.out=n)  
yi <- f(xi)  
  
plot(xi, yi, pch=16, xlab="x", ylab="y")  
  
x <- seq(-1, 1, 0.01)  
lines(x, q(x, xi, yi))  
lines(x, f(x), col="red")
```

6. Comment on  $q(x)$  vs.  $f(x)$ .

7. Now rerun the code with  $n = 5, 6, 7$ , and 8 nodes. For each  $n$ , compare  $q(x)$  to  $f(x)$ .

## Key points

8. What does the function  $q(x)$  do?

9. Why is the number of points  $n$  important?

## Part 2

Previously in class, we found that the "best" two-point rule to approximate the integral of  $f$  was

$$\int_{-1}^1 f(x)dx \approx w_1 f(x_1) + w_2 f(x_2)$$

with  $x_1 = -1/\sqrt{3}$ ,  $x_2 = 1/\sqrt{3}$ , and  $w_1 = w_2 = 1$ .

Where do these weights come from? By using the polynomial interpolation  $q(x) = \sum_{i=1}^n f(x_i)L_{n,i}(x)$ , we argued that

$$w_i = \int_{-1}^1 L_{n,i}(x)dx$$

10. For the two-point rule, we have points  $L_{2,1}(x) = \frac{x - x_2}{x_1 - x_2}$  with  $x_1 = -1/\sqrt{3}$ ,  $x_2 = 1/\sqrt{3}$ . Show that

$$\int_{-1}^1 L_{2,1}(x)dx = 1$$