

# STA 711 Exam 1 Make-up

**Due:** Friday, March 3, 12:00pm (noon) on Canvas.

**Instructions:** Submit your work as a single PDF. For this assignment, you may include written work by scanning it and incorporating it into the PDF. Include all R code needed to reproduce your results in your submission.

**Rules:** This is a closed-book, closed-notes exam. You may:

- Email me, or come to office hours, with specific questions (I may be somewhat less helpful than for regular assignments)
- Use the internet for general R help and debugging
- Use R for question 7

You may *not*:

- Use any resources from the course (the textbook, the course website, class notes, previous assignments, etc.)
- Use the internet to look up any questions on the exam
- Discuss the exam with anyone else

## Maximum likelihood and Fisher information

1. Let  $Y_1, \dots, Y_n$  be iid random variables with pdf

$$f(y|\mu, \sigma^2) = \frac{1}{y\sqrt{2\pi\sigma^2}} \exp \left\{ -\frac{1}{2\sigma^2} (\log(y) - \mu)^2 \right\}$$

where  $\sigma^2 > 0$ ,  $y > 0$ , and  $\mu \in \mathbb{R}$ . Find the maximum likelihood estimators  $\hat{\mu}$  and  $\hat{\sigma}^2$ .

2. Let  $Y_1, \dots, Y_n \stackrel{iid}{\sim} \text{Gamma}(\alpha, \beta)$ . Recall that the gamma distribution has pdf

$$f(y|\alpha, \beta) = \frac{\beta^\alpha}{\Gamma(\alpha)} y^{\alpha-1} e^{-\beta y}$$

where  $y > 0$ ,  $\alpha > 0$ , and  $\beta > 0$ . Find the Fisher information matrix  $\mathcal{I}(\alpha, \beta)$ . (You may assume that desired regularity conditions hold for the gamma distribution). Note: the derivative of the gamma function  $\Gamma(x)$  has no nice closed form. You may use  $\psi(x)$  to represent  $\frac{d}{dx} \log \Gamma(x)$  (the digamma function), and  $\psi_1(x)$  to represent  $\frac{d}{dx} \psi(x)$  (the trigamma function).

## Modeling an Exponential response

In this section, you will model a response variable  $Y_i \sim \text{Exponential}(\lambda_i)$ .

3. Recall that if  $Y \sim \text{Exponential}(\lambda)$ , then the pdf of  $Y$  is given by

$$f(Y; \lambda) = \frac{1}{\lambda} e^{-Y/\lambda}$$

Show that  $\mathbb{E}[Y] = \lambda$  and  $\text{Var}(Y) = \lambda^2$ .

4. Suppose that  $Y_i$  is a response variable of interest, and we use the following model for  $Y_i$ :

$$Y_i \sim \text{Exponential}(\lambda_i)$$
$$\frac{1}{\lambda_i} = \beta^T X_i,$$

where  $X_i = (1, X_{i,1}, \dots, X_{i,k})^T \in \mathbb{R}^{k+1}$ , and  $\beta = (\beta_0, \dots, \beta_k) \in \mathbb{R}^{k+1}$  is the vector of regression coefficients. We observe  $n$  independent observations  $(X_1, Y_1), \dots, (X_n, Y_n)$ .

- (a) Derive a matrix form for the score  $U(\beta)$ .
- (b) Derive a matrix form for the Fisher information  $\mathcal{I}(\beta)$ . You may assume that desired regularity conditions hold.
5. Now we apply the model from Question 4 to real data. A factory is interested in the relationship between the amount of stress applied to a piece of steel, and the time it takes until that steel breaks. We use the following model:

$$time_i \sim \text{Exponential}(\lambda_i)$$
$$\frac{1}{\lambda_i} = \beta_0 + \beta_1 stress_i$$

The raw data contains  $n = 40$  observations  $(stress_1, time_1), \dots, (stress_{40}, time_{40})$ .

You can load the data into R by

```
steel <- read.csv("https://sta711-s23.github.io/exams/steel.csv")
```

In this question, we want to begin Fisher scoring to estimate  $\beta_0$  and  $\beta_1$  using the observed data.

- (a) In R, write functions `U` and `I` to calculate  $U(\beta)$  and  $\mathcal{I}(\beta)$  using the observed data.
- (b) Use your functions to estimate  $\beta$  using Fisher scoring with the observed data. Iterate the Fisher scoring algorithm until convergence, beginning with  $\beta^{(0)} = (1, 2)^T$ . For the purpose of this question, stop when

$$\max\{|\beta_0^{(r+1)} - \beta_0^{(r)}|, |\beta_1^{(r+1)} - \beta_1^{(r)}|\} < 0.0001$$