

# STA 711 HW 1 Solutions

Ciaran Evans

1. Suppose that  $X \sim \text{Poisson}(\lambda)$ .

(a) We use the fact that  $e^\lambda = \sum_{x=0}^{\infty} \frac{\lambda^x}{x!}$ .

$$\begin{aligned}\mathbb{E}[X] &= \sum_{x=0}^{\infty} x \frac{\lambda^x e^{-\lambda}}{x!} = \sum_{x=1}^{\infty} x \frac{\lambda^x e^{-\lambda}}{x!} \\ &= e^{-\lambda} \lambda \sum_{x=1}^{\infty} \frac{\lambda^{x-1}}{(x-1)!} = e^{-\lambda} \lambda \sum_{x=0}^{\infty} \frac{\lambda^x}{x!} \\ &= e^{-\lambda} \lambda e^\lambda \\ &= \lambda\end{aligned}$$

$$\begin{aligned}\mathbb{E}[X^2] &= \sum_{x=0}^{\infty} x^2 \frac{\lambda^x e^{-\lambda}}{x!} = \sum_{x=1}^{\infty} x^2 \frac{\lambda^x e^{-\lambda}}{x!} \\ &= \lambda e^{-\lambda} \sum_{x=1}^{\infty} x \frac{\lambda^{x-1}}{(x-1)!} \\ &= \lambda e^{-\lambda} \sum_{x=1}^{\infty} x \frac{\lambda^{x-1}}{(x-1)!} \\ &= \lambda e^{-\lambda} \left( \sum_{x=1}^{\infty} (x-1) \frac{\lambda^{x-1}}{(x-1)!} + \sum_{x=1}^{\infty} \frac{\lambda^{x-1}}{(x-1)!} \right) \\ &= \lambda e^{-\lambda} (\lambda e^\lambda + e^\lambda) \\ &= \lambda^2 + \lambda\end{aligned}$$

Thus,  $\text{Var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2 = \lambda$ .

(b)

$$\mathbb{E}[e^{tX}] = \sum_{x=0}^{\infty} e^{tx} \frac{\lambda^x e^{-\lambda}}{x!} = e^{-\lambda} \sum_{x=0}^{\infty} \frac{(\lambda e^t)^x}{x!} = e^{-\lambda} e^{\lambda e^t} = \exp\{\lambda(e^t - 1)\}$$

(c) From (b),  $M_X(t) = \exp\{\lambda(e^t - 1)\}$ , so

$$M'_X(t) = \exp\{\lambda(e^t - 1)\} \lambda e^t$$

and

$$M''_X(t) = \exp\{\lambda(e^t - 1)\} \lambda^2 e^{2t} + \exp\{\lambda(e^t - 1)\} \lambda$$

At  $t = 0$ , we get

$$\mathbb{E}[X] = M'_X(0) = \lambda$$

$$\mathbb{E}[X^2] = M''_X(0) = \lambda^2 + \lambda$$

and thus  $\text{Var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2 = \lambda$ .

2. Suppose that  $X \sim \text{Poisson}(\lambda)$  and  $Y \sim \text{Poisson}(\mu)$  are independent. Let  $Z = X + Y$ .

(a) We use the fact that the MGF for the sum of independent random variables is the product of their individual MGFs:

$$\begin{aligned} M_Z(t) &= M_X(t)M_Y(t) = \exp\{\lambda(e^t - 1)\} \exp\{\mu(e^t - 1)\} \\ &= \exp\{(\lambda + \mu)(e^t - 1)\} \end{aligned}$$

This is the MGF of a  $\text{Poisson}(\lambda + \mu)$  distribution, and so by the uniqueness of MGFs we conclude  $X + Y \sim \text{Poisson}(\lambda + \mu)$ .

(b)

$$\begin{aligned} P(X = x|Z = n) &= \frac{P(X = x, Z = n)}{P(Z = n)} = \frac{P(X = x, Y = n - x)}{P(Z = n)} \\ &= \frac{\frac{\lambda^x e^{-\lambda}}{x!} \cdot \frac{\mu^{n-x} e^{-\mu}}{(n-x)!}}{\frac{(\lambda + \mu)^n e^{-(\lambda + \mu)}}{n!}} \\ &= \frac{n!}{x!(n-x)!} \left(\frac{\lambda}{\lambda + \mu}\right)^x \left(1 - \frac{\lambda}{\lambda + \mu}\right)^{n-x} \end{aligned}$$

This is the pmf of a  $\text{Binomial}\left(n, \frac{\lambda}{\lambda + \mu}\right)$  distribution.

3. The joint pdf is

$$f(x, y) = \begin{cases} c(x + 2y) & 0 < y < 1, 0 < x < 2 \\ 0 & \text{else} \end{cases}$$

(a) To find  $c$ , we note that  $f(x, y)$  must integrate to 1.

$$\begin{aligned} c \int_0^1 \int_0^2 (x + 2y) dx dy &= c \int_0^1 \int_0^2 x dx dy + c \int_0^1 \int_0^2 2y dx dy \\ &= 2c + 2c \end{aligned}$$

so we conclude that  $c = 1/4$ .

(b)

$$f(x) = \int_0^1 f(x, y) dy = \frac{1}{4} \int_0^1 (x + 2y) dy = \frac{x + 1}{4}$$

(c)

$$\begin{aligned} P(Z < t) &= P(9 \leq (1 + X)^2 t) = P\left(\frac{3}{\sqrt{t}} - 1 \leq X\right) = \int_{3/\sqrt{t}-1}^1 (x + 1)/4 dx \\ &= \frac{1}{2} - \frac{9}{8t} \end{aligned}$$

Then, differentiating,

$$f_Z(t) = \frac{9}{8t^2}$$

for  $t \in [9/4, 9]$

4.

5.

6. (a) Since  $F_X$  is a monotonic, continuous cdf, then  $F_X$  and  $F_X^{-1}$  are both strictly increasing, and

$$P(X \leq t) = P(F_X^{-1}(U) \leq t) = P(U \leq F_X(t)) = F_X(t),$$

where the final step follows from the cdf of a *Uniform*(0, 1) distribution.

- (b) See the R part below.

7. (a)

$$\begin{aligned} \text{Cov}\left(\sum_{i=1}^m a_i X_i, \sum_{j=1}^n b_j Y_j\right) &= \mathbb{E}\left[\left(\sum_{i=1}^m a_i X_i - \sum_{i=1}^m a_i \mathbb{E}[X_i]\right)\left(\sum_{j=1}^n b_j Y_j - \sum_{j=1}^n b_j \mathbb{E}[Y_j]\right)\right] \\ &= \mathbb{E}\left[\left(\sum_{i=1}^m a_i (X_i - \mathbb{E}[X_i])\right)\left(\sum_{j=1}^n b_j (Y_j - \mathbb{E}[Y_j])\right)\right] \\ &= \mathbb{E}\left[\sum_{i=1}^m \sum_{j=1}^n a_i b_j (X_i - \mathbb{E}[X_i])(Y_j - \mathbb{E}[Y_j])\right] \\ &= \sum_{i=1}^m \sum_{j=1}^n a_i b_j \mathbb{E}[(X_i - \mathbb{E}[X_i])(Y_j - \mathbb{E}[Y_j])] \\ &= \sum_{i=1}^m \sum_{j=1}^n a_i b_j \text{Cov}(X_i, Y_j) \end{aligned}$$

- (b)

$$\begin{aligned} \text{Var}\left(\sum_{i=1}^m X_i\right) &= \text{Cov}\left(\sum_{i=1}^m X_i, \sum_{i=1}^m X_i\right) \\ &= \sum_{i=1}^m \sum_{j=1}^m \text{Cov}(X_i, X_j) \\ &= \sum_{i=1}^m \text{Var}(X_i) + \sum_{i=1}^m \sum_{j \neq i}^m \text{Cov}(X_i, X_j) \\ &= \sum_{i=1}^m \text{Var}(X_i) + 2 \sum_{i < j}^m \text{Cov}(X_i, X_j) \end{aligned}$$

8.  $X_1, \dots, X_n \stackrel{iid}{\sim} \text{Uniform}(0, 1)$ , and  $Y = \max\{X_1, \dots, X_n\}$

(a) For  $t \in [0, 1]$ :

$$\begin{aligned} P(Y \leq t) &= P(X_1, \dots, X_n \leq t) = \prod_{i=1}^n P(X_i \leq t) \text{ (independence)} \\ &= t^n \end{aligned}$$

Then, the pdf is

$$f_Y(t) = nt^{n-1}$$