

Convergence of random variables

Ciaran Evans

Recap: convergence

Definition: A sequence of random variables X_1, X_2, \dots *converges in probability* to a random variable X if, for every $\varepsilon > 0$,

$$\lim_{n \rightarrow \infty} P(|X_n - X| \geq \varepsilon) = 0$$

We write $X_n \xrightarrow{p} X$.

Definition: A sequence of random variables X_1, X_2, \dots *converges in distribution* to a random variable X if

$$\lim_{n \rightarrow \infty} F_{X_n}(x) = F_X(x)$$

at all points where $F_X(x)$ is continuous. We write $X_n \xrightarrow{d} X$.

Central Limit Theorem

Let X_1, X_2, \dots be iid random variables, with $\mu = \mathbb{E}[X_i]$ and $0 < \sigma^2 = \text{Var}(X_i) < \infty$. Then

$$\frac{\sqrt{n}(\bar{X}_n - \mu)}{\sigma} \xrightarrow{d} Z$$

where $Z \sim N(0, 1)$.

Intuition:

$$\bar{X}_n \xrightarrow{P} \mu \quad (\text{WLLN})$$

$$\bar{X}_n - \mu \xrightarrow{P} 0$$

$$\text{var}(\bar{X}_n) = \frac{\sigma^2}{n} \rightarrow 0$$

$$\text{var}\left(\frac{\sqrt{n}}{\sigma}(\bar{X}_n - \mu)\right) = 1$$

Other key results

Continuous mapping theorem: Let X_1, X_2, \dots be a sequence of random variables, and g a continuous function.

$$\text{if } X_n \xrightarrow{d} X, \text{ then } g(X_n) \xrightarrow{d} g(X)$$

$$\text{if } X_n \xrightarrow{p} X, \text{ then } g(X_n) \xrightarrow{p} g(X) \leftarrow \text{proof: HW}$$

Slutsky's theorem: Let $\{X_n\}, \{Y_n\}$ be sequences of random variables, and suppose that $X_n \xrightarrow{d} X$ and $Y_n \xrightarrow{p} c$, where c is a constant. Then:

$$X_n + Y_n \xrightarrow{d} X + c$$

$$X_n Y_n \xrightarrow{d} cX$$

$$\frac{X_n}{Y_n} \xrightarrow{d} \frac{X}{c}, \text{ if } c \neq 0$$

Asymptotic normality with sample standard deviation

Let X_1, X_2, \dots be iid random variables, with $\mu = \mathbb{E}[X_i]$ and $0 < \sigma^2 = \text{Var}(X_i) < \infty$. Let $\hat{\sigma}^2$ be an estimate of σ^2 such that $\hat{\sigma}^2 \xrightarrow{P} \sigma^2$. Then

$$\frac{\sqrt{n}(\bar{X}_n - \mu)}{\hat{\sigma}} \xrightarrow{d} Z \sim N(0, 1)$$

$$\textcircled{1} \quad \hat{\sigma}^2 \xrightarrow{P} \sigma^2 \quad \hat{\sigma} = \sqrt{\hat{\sigma}^2} \xrightarrow{P} \sigma \quad (\text{continuous mapping})$$
$$\Rightarrow \frac{\sigma}{\hat{\sigma}} \xrightarrow{P} 1$$

$$\textcircled{2} \quad \text{By CLT, } \frac{\sqrt{n}(\bar{X}_n - \mu)}{\sigma} \xrightarrow{d} Z$$


$$\textcircled{3} \quad \frac{\sqrt{n}(\bar{X}_n - \mu)}{\hat{\sigma}} = \frac{\sqrt{n}(\bar{X}_n - \mu)}{\sigma} \cdot \frac{\sigma}{\hat{\sigma}} \xrightarrow{d} Z \cdot 1 = Z \quad (\text{by Slutsky's theorem})$$

Relationships between types of convergence

(a) If $X_n \xrightarrow{d} c$, where c is a constant, then $X_n \xrightarrow{P} c$
 $P(c=c) = 1$

(b) If $X_n \xrightarrow{P} X$, then $X_n \xrightarrow{d} X$

(c) point mass at c :



$$pmf: f_c(x) = \begin{cases} 1 & x=c \\ 0 & x \neq c \end{cases}$$

$$cdf: F_c(x) = \begin{cases} 0 & x < c \\ 1 & x \geq c \end{cases}$$

Suppose $X_n \xrightarrow{d} c$, and let $\varepsilon > 0$.

$$\begin{aligned} P(|X_n - c| \geq \varepsilon) &= 1 - P(|X_n - c| < \varepsilon) = 1 - P(c - \varepsilon < X_n < c + \varepsilon) \\ &= 1 - (F_{X_n}(c + \varepsilon) - F_{X_n}(c - \varepsilon)) \end{aligned}$$

Since $X_n \xrightarrow{d} c$, $F_{X_n}(t) \rightarrow F_c(t) \quad \forall t \neq c$

$$F_{X_n}(c + \varepsilon) \rightarrow F_c(c + \varepsilon) = 1$$

$$F_{X_n}(c - \varepsilon) \rightarrow F_c(c - \varepsilon) = 0$$

$$\begin{aligned} \Rightarrow \lim_{n \rightarrow \infty} P(|X_n - c| \geq \varepsilon) &= 1 - \lim_{n \rightarrow \infty} (F_{X_n}(c + \varepsilon) - F_{X_n}(c - \varepsilon)) \\ &= 1 - (1 - 0) = 0 \end{aligned}$$

//

Relationships between types of convergence

(a) If $X_n \xrightarrow{d} c$, where c is a constant, then $X_n \xrightarrow{P} c$

(b) If $X_n \xrightarrow{P} X$, then $X_n \xrightarrow{d} X$

(b) WTS: $F_{X_n}(t) \rightarrow F_X(t) \quad \forall t$ where F_X is continuous

If F_X is continuous at t : $\lim_{\varepsilon \rightarrow 0^+} F_X(t-\varepsilon) = F_X(t) = \lim_{\varepsilon \rightarrow 0^+} F_X(t+\varepsilon)$

It suffices to show that $\forall \varepsilon > 0, F_X(t-\varepsilon) \leq \lim_{n \rightarrow \infty} F_{X_n}(t) \leq F_X(t+\varepsilon)$

Let $\varepsilon > 0$, and let t be a continuity point of F_X

$$F_{X_n}(t) = P(X_n \leq t) = \underbrace{P(X_n \leq t, X \leq t+\varepsilon)}_{\leq P(X \leq t+\varepsilon)} + \underbrace{P(X_n \leq t, X > t+\varepsilon)}_{\leq P(|X_n - X| \geq \varepsilon)}$$

$$\Rightarrow F_{X_n}(t) \leq F_X(t+\varepsilon) + P(|X_n - X| \geq \varepsilon)$$

$$\Rightarrow \lim_{n \rightarrow \infty} F_{X_n}(t) \leq F_X(t+\varepsilon) + \underbrace{\lim_{n \rightarrow \infty} P(|X_n - X| \geq \varepsilon)}_{=0 \text{ b/c } X_n \xrightarrow{P} X}$$

$$\leq F_X(t+\varepsilon)$$

Relationships between types of convergence

(a) If $X_n \xrightarrow{d} c$, where c is a constant, then $X_n \xrightarrow{p} c$

(b) If $X_n \xrightarrow{p} X$, then $X_n \xrightarrow{d} X$

(b) continued...

$$\text{So: } \lim_{n \rightarrow \infty} F_{X_n}(t) \leq F_X(t + \varepsilon)$$

$$\text{Similarly, } F_X(t - \varepsilon) \leq \lim_{n \rightarrow \infty} F_{X_n}(t)$$

$$\Rightarrow F_X(t - \varepsilon) \leq \lim_{n \rightarrow \infty} F_{X_n}(t) \leq F_X(t + \varepsilon) \quad \forall \varepsilon > 0$$

$$\Rightarrow \lim_{\varepsilon \rightarrow 0^+} F_X(t - \varepsilon) \leq \lim_{n \rightarrow \infty} F_{X_n}(t) \leq \lim_{\varepsilon \rightarrow 0^+} F_X(t + \varepsilon)$$

$$\Rightarrow F_X(t) \leq \lim_{n \rightarrow \infty} F_{X_n}(t) \leq F_X(t)$$

$$\Rightarrow \lim_{n \rightarrow \infty} F_{X_n}(t) = F_X(t)$$

$$\text{So, } X_n \xrightarrow{d} X$$