

# Identifiability, Gauss Markov & Predictive Distributions

Merlise Clyde

STA721 Linear Models

Duke University

September 16, 2019

# Outline

## Topics

- ▶ Gauss-Markov Theorem
- ▶ Estimability and Prediction

Readings: Christensen Chapter 2, Chapter 6.3, ( Appendix A, and Appendix B as needed)

# Non-Identifiable

Recall the One-way ANOVA model

$$\mu_{ij} = \mu + \tau_j \quad \boldsymbol{\mu} = (\mu_{11}, \dots, \mu_{n_1 1}, \mu_{12}, \dots, \mu_{n_2 2}, \dots, \mu_{1J}, \dots, \mu_{n_J J})^T$$

- ▶ Let  $\boldsymbol{\beta}_1 = (\mu, \tau_1, \dots, \tau_J)^T$
- ▶ Let  $\boldsymbol{\beta}_2 = (\mu - 42, \tau_1 + 42, \dots, \tau_J + 42)^T$
- ▶ Then  $\boldsymbol{\mu}_1 = \boldsymbol{\mu}_2$  even though  $\boldsymbol{\beta}_1 \neq \boldsymbol{\beta}_2$
- ▶  $\boldsymbol{\beta}$  is not identifiable
- ▶ yet  $\boldsymbol{\mu}$  is identifiable, where  $\boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$  (a linear combination of  $\boldsymbol{\beta}$ )

# Identifiability and Estimability

## Theorem

*A function  $g(\beta)$  is identifiable if and only if  $g(\beta)$  is a function of  $\mu(\beta)$*

# Identifiability and Estimability

## Theorem

*A function  $g(\beta)$  is identifiable if and only if  $g(\beta)$  is a function of  $\mu(\beta)$*

In linear models, focus on linear functions. Identifiable linear functions are called *estimable* functions historically

# Identifiability and Estimability

## Theorem

*A function  $g(\beta)$  is identifiable if and only if  $g(\beta)$  is a function of  $\mu(\beta)$*

In linear models, focus on linear functions. Identifiable linear functions are called *estimable* functions historically

## Definition

A scalar function  $\lambda^T \beta$  is *estimable* if  $\lambda^T \beta = \mathbf{a}^T \mathbf{X} \beta$  for some vector  $\mathbf{a} \in \mathbb{R}^n$

# Identifiability and Estimability

## Theorem

A function  $g(\beta)$  is identifiable if and only if  $g(\beta)$  is a function of  $\mu(\beta)$

In linear models, focus on linear functions. Identifiable linear functions are called *estimable* functions historically

## Definition

A scalar function  $\lambda^T \beta$  is *estimable* if  $\lambda^T \beta = \mathbf{a}^T \mathbf{X} \beta$  for some vector  $\mathbf{a} \in \mathbb{R}^n$

Equivalently

## Definition

A function  $\lambda^T \beta$  is *estimable* if it has an unbiased linear estimator, i.e. there exists an  $\mathbf{a}$  such that  $E(\mathbf{a}^T \mathbf{Y}) = \lambda^T \beta$  for all  $\beta$

# Estimability

## Theorem

*The function  $\psi = \boldsymbol{\lambda}^T \boldsymbol{\beta}$  is estimable if and only if  $\boldsymbol{\lambda}^T$  is a linear combination of the rows of  $\mathbf{X}$ . i.e. there exists  $\mathbf{a}^T$  such that  $\boldsymbol{\lambda}^T = \mathbf{a}^T \mathbf{X}$*



# Estimability

## Theorem

*The function  $\psi = \boldsymbol{\lambda}^T \boldsymbol{\beta}$  is estimable if and only if  $\boldsymbol{\lambda}^T$  is a linear combination of the rows of  $\mathbf{X}$ . i.e. there exists  $\mathbf{a}^T$  such that  $\boldsymbol{\lambda}^T = \mathbf{a}^T \mathbf{X}$*

## Proof.

The function  $\psi = \boldsymbol{\lambda}^T \boldsymbol{\beta}$  is estimable if there exists an  $\mathbf{a}^T$  such that  $E[\mathbf{a}^T \mathbf{Y}] = \boldsymbol{\lambda}^T \boldsymbol{\beta}$

# Estimability

## Theorem

*The function  $\psi = \boldsymbol{\lambda}^T \boldsymbol{\beta}$  is estimable if and only if  $\boldsymbol{\lambda}^T$  is a linear combination of the rows of  $\mathbf{X}$ . i.e. there exists  $\mathbf{a}^T$  such that  $\boldsymbol{\lambda}^T = \mathbf{a}^T \mathbf{X}$*

## Proof.

The function  $\psi = \boldsymbol{\lambda}^T \boldsymbol{\beta}$  is estimable if there exists an  $\mathbf{a}^T$  such that  $E[\mathbf{a}^T \mathbf{Y}] = \boldsymbol{\lambda}^T \boldsymbol{\beta}$

$$E[\mathbf{a}^T \mathbf{Y}] = \mathbf{a}^T E[\mathbf{Y}]$$

# Estimability

## Theorem

*The function  $\psi = \boldsymbol{\lambda}^T \boldsymbol{\beta}$  is estimable if and only if  $\boldsymbol{\lambda}^T$  is a linear combination of the rows of  $\mathbf{X}$ . i.e. there exists  $\mathbf{a}^T$  such that  $\boldsymbol{\lambda}^T = \mathbf{a}^T \mathbf{X}$*

## Proof.

The function  $\psi = \boldsymbol{\lambda}^T \boldsymbol{\beta}$  is estimable if there exists an  $\mathbf{a}^T$  such that  $E[\mathbf{a}^T \mathbf{Y}] = \boldsymbol{\lambda}^T \boldsymbol{\beta}$

$$\begin{aligned} E[\mathbf{a}^T \mathbf{Y}] &= \mathbf{a}^T E[\mathbf{Y}] \\ &= \mathbf{a}^T \mathbf{X} \boldsymbol{\beta} \end{aligned}$$

# Estimability

## Theorem

*The function  $\psi = \boldsymbol{\lambda}^T \boldsymbol{\beta}$  is estimable if and only if  $\boldsymbol{\lambda}^T$  is a linear combination of the rows of  $\mathbf{X}$ . i.e. there exists  $\mathbf{a}^T$  such that  $\boldsymbol{\lambda}^T = \mathbf{a}^T \mathbf{X}$*

## Proof.

The function  $\psi = \boldsymbol{\lambda}^T \boldsymbol{\beta}$  is estimable if there exists an  $\mathbf{a}^T$  such that  $E[\mathbf{a}^T \mathbf{Y}] = \boldsymbol{\lambda}^T \boldsymbol{\beta}$

$$\begin{aligned} E[\mathbf{a}^T \mathbf{Y}] &= \mathbf{a}^T E[\mathbf{Y}] \\ &= \mathbf{a}^T \mathbf{X} \boldsymbol{\beta} \\ &= \boldsymbol{\lambda}^T \boldsymbol{\beta} \end{aligned}$$

# Estimability

## Theorem

*The function  $\psi = \boldsymbol{\lambda}^T \boldsymbol{\beta}$  is estimable if and only if  $\boldsymbol{\lambda}^T$  is a linear combination of the rows of  $\mathbf{X}$ . i.e. there exists  $\mathbf{a}^T$  such that  $\boldsymbol{\lambda}^T = \mathbf{a}^T \mathbf{X}$*

## Proof.

The function  $\psi = \boldsymbol{\lambda}^T \boldsymbol{\beta}$  is estimable if there exists an  $\mathbf{a}^T$  such that  $E[\mathbf{a}^T \mathbf{Y}] = \boldsymbol{\lambda}^T \boldsymbol{\beta}$

$$\begin{aligned} E[\mathbf{a}^T \mathbf{Y}] &= \mathbf{a}^T E[\mathbf{Y}] \\ &= \mathbf{a}^T \mathbf{X} \boldsymbol{\beta} \\ &= \boldsymbol{\lambda}^T \boldsymbol{\beta} \end{aligned}$$

if and only if  $\boldsymbol{\lambda}^T = \mathbf{a}^T \mathbf{X}$  for all  $\boldsymbol{\beta}$



# Estimability of Individual $\beta_j$

## Proposition

For

$$\mu = \mathbf{X}\beta = \sum_j \mathbf{X}_j \beta_j$$

$\beta_j$  is not identifiable if and only if there exists  $\alpha_j$  such that

$$\mathbf{X}_j = \sum_{i \neq j} \mathbf{X}_i \alpha_i$$

# Estimability of Individual $\beta_j$

## Proposition

For

$$\mu = \mathbf{X}\beta = \sum_j \mathbf{X}_j \beta_j$$

$\beta_j$  is not identifiable if and only if there exists  $\alpha_j$  such that

$$\mathbf{X}_j = \sum_{i \neq j} \mathbf{X}_i \alpha_i$$

One-way Anova Model:

$$Y_{ij} = \mu + \tau_j + \epsilon_{ij}$$

$$\mu = \begin{bmatrix} \mathbf{1}_{n_1} & \mathbf{1}_{n_1} & \mathbf{0}_{n_1} & \cdots & \mathbf{0}_{n_1} \\ \mathbf{1}_{n_2} & \mathbf{0}_{n_2} & \mathbf{1}_{n_2} & \cdots & \mathbf{0}_{n_2} \\ \vdots & \vdots & \ddots & \vdots & \\ \mathbf{1}_{n_J} & \mathbf{0}_{n_J} & \mathbf{0}_{n_J} & \cdots & \mathbf{1}_{n_J} \end{bmatrix} \begin{pmatrix} \mu \\ \tau_1 \\ \tau_2 \\ \vdots \\ \tau_J \end{pmatrix}$$

# Estimability of Individual $\beta_j$

## Proposition

For

$$\mu = \mathbf{X}\beta = \sum_j \mathbf{X}_j \beta_j$$

$\beta_j$  is not identifiable if and only if there exists  $\alpha_j$  such that

$$\mathbf{X}_j = \sum_{i \neq j} \mathbf{X}_i \alpha_i$$

One-way Anova Model:

$$Y_{ij} = \mu + \tau_j + \epsilon_{ij}$$

$$\mu = \begin{bmatrix} \mathbf{1}_{n_1} & \mathbf{1}_{n_1} & \mathbf{0}_{n_1} & \cdots & \mathbf{0}_{n_1} \\ \mathbf{1}_{n_2} & \mathbf{0}_{n_2} & \mathbf{1}_{n_2} & \cdots & \mathbf{0}_{n_2} \\ \vdots & \vdots & \ddots & \vdots & \\ \mathbf{1}_{n_J} & \mathbf{0}_{n_J} & \mathbf{0}_{n_J} & \cdots & \mathbf{1}_{n_J} \end{bmatrix} \begin{pmatrix} \mu \\ \tau_1 \\ \tau_2 \\ \vdots \\ \tau_J \end{pmatrix}$$

Are any parameters  $\mu$  or  $\tau_j$  identifiable?



# Gauss-Markov Theorem

## Theorem

*Under the assumptions:*

$$E[\mathbf{Y}] = \boldsymbol{\mu}$$

# Gauss-Markov Theorem

## Theorem

*Under the assumptions:*

$$\begin{aligned} E[\mathbf{Y}] &= \boldsymbol{\mu} \\ \text{Cov}(\mathbf{Y}) &= \sigma^2 \mathbf{I}_n \end{aligned}$$

# Gauss-Markov Theorem

## Theorem

*Under the assumptions:*

$$\begin{aligned}E[\mathbf{Y}] &= \boldsymbol{\mu} \\ \text{Cov}(\mathbf{Y}) &= \sigma^2 \mathbf{I}_n\end{aligned}$$

*every estimable function  $\psi = \boldsymbol{\lambda}^T \boldsymbol{\beta}$  has a unique unbiased linear estimator  $\hat{\psi}$  which has minimum variance in the class of all unbiased linear estimators.*

# Gauss-Markov Theorem

## Theorem

*Under the assumptions:*

$$\begin{aligned}E[\mathbf{Y}] &= \boldsymbol{\mu} \\ \text{Cov}(\mathbf{Y}) &= \sigma^2 \mathbf{I}_n\end{aligned}$$

*every estimable function  $\psi = \boldsymbol{\lambda}^T \boldsymbol{\beta}$  has a unique unbiased linear estimator  $\hat{\psi}$  which has minimum variance in the class of all unbiased linear estimators.  $\hat{\psi} = \boldsymbol{\lambda}^T \hat{\boldsymbol{\beta}}$  where  $\hat{\boldsymbol{\beta}}$  is any set of ordinary least squares estimators.*

# Unique Unbiased Estimator

## Lemma

- ▶ If  $\psi = \boldsymbol{\lambda}^T \boldsymbol{\beta}$  is estimable, there exists a unique linear unbiased estimator of  $\psi = \mathbf{a}^{*T} \mathbf{Y}$  with  $\mathbf{a}^* \in C(\mathbf{X})$ .

# Unique Unbiased Estimator

## Lemma

- ▶ If  $\psi = \boldsymbol{\lambda}^T \boldsymbol{\beta}$  is estimable, there exists a unique linear unbiased estimator of  $\psi = \mathbf{a}^{*T} \mathbf{Y}$  with  $\mathbf{a}^* \in C(\mathbf{X})$ .
- ▶ If  $\mathbf{a}^T \mathbf{Y}$  is any unbiased linear estimator of  $\psi$  then  $\mathbf{a}^*$  is the projection of  $\mathbf{a}$  onto  $C(\mathbf{X})$ , i.e.  $\mathbf{a}^* = \mathbf{P}_X \mathbf{a}$ .

# Unique Unbiased Estimator

## Proof

- ▶ Since  $\psi$  is estimable, there exists an  $\mathbf{a} \in \mathbb{R}^n$  for which  $E[\mathbf{a}^T \mathbf{Y}] = \boldsymbol{\lambda}^T \boldsymbol{\beta} = \psi$  with  $\boldsymbol{\lambda}^T = \mathbf{a}^T \mathbf{X}$

# Unique Unbiased Estimator

## Proof

- ▶ Since  $\psi$  is estimable, there exists an  $\mathbf{a} \in \mathbb{R}^n$  for which  $E[\mathbf{a}^T \mathbf{Y}] = \boldsymbol{\lambda}^T \boldsymbol{\beta} = \psi$  with  $\boldsymbol{\lambda}^T = \mathbf{a}^T \mathbf{X}$
- ▶ Let  $\mathbf{a} = \mathbf{a}^* + \mathbf{u}$  where  $\mathbf{a}^* \in C(\mathbf{X})$  and  $\mathbf{u} \in C(\mathbf{X})^\perp$



# Unique Unbiased Estimator

## Proof

- ▶ Since  $\psi$  is estimable, there exists an  $\mathbf{a} \in \mathbb{R}^n$  for which  $E[\mathbf{a}^T \mathbf{Y}] = \boldsymbol{\lambda}^T \boldsymbol{\beta} = \psi$  with  $\boldsymbol{\lambda}^T = \mathbf{a}^T \mathbf{X}$
- ▶ Let  $\mathbf{a} = \mathbf{a}^* + \mathbf{u}$  where  $\mathbf{a}^* \in C(\mathbf{X})$  and  $\mathbf{u} \in C(\mathbf{X})^\perp$
- ▶ Then

$$\psi = E[\mathbf{a}^T \mathbf{Y}] = E[\mathbf{a}^{*T} \mathbf{Y}] + E[\mathbf{u}^T \mathbf{Y}]$$

# Unique Unbiased Estimator

## Proof

- ▶ Since  $\psi$  is estimable, there exists an  $\mathbf{a} \in \mathbb{R}^n$  for which  $E[\mathbf{a}^T \mathbf{Y}] = \boldsymbol{\lambda}^T \boldsymbol{\beta} = \psi$  with  $\boldsymbol{\lambda}^T = \mathbf{a}^T \mathbf{X}$
- ▶ Let  $\mathbf{a} = \mathbf{a}^* + \mathbf{u}$  where  $\mathbf{a}^* \in C(\mathbf{X})$  and  $\mathbf{u} \in C(\mathbf{X})^\perp$
- ▶ Then

$$\begin{aligned}\psi = E[\mathbf{a}^T \mathbf{Y}] &= E[\mathbf{a}^{*T} \mathbf{Y}] + E[\mathbf{u}^T \mathbf{Y}] \\ &= E[\mathbf{a}^{*T} \mathbf{Y}] + 0\end{aligned}$$

$$E[\mathbf{u}^T \mathbf{Y}] = \mathbf{u}^T \mathbf{X} \boldsymbol{\beta}$$

since  $\mathbf{u} \perp C(\mathbf{X})$  (i.e.  $\mathbf{u} \in C(\mathbf{X})^\perp$ )  $E[\mathbf{u}^T \mathbf{Y}] = 0$

# Unique Unbiased Estimator

## Proof

- ▶ Since  $\psi$  is estimable, there exists an  $\mathbf{a} \in \mathbb{R}^n$  for which  $E[\mathbf{a}^T \mathbf{Y}] = \boldsymbol{\lambda}^T \boldsymbol{\beta} = \psi$  with  $\boldsymbol{\lambda}^T = \mathbf{a}^T \mathbf{X}$
- ▶ Let  $\mathbf{a} = \mathbf{a}^* + \mathbf{u}$  where  $\mathbf{a}^* \in C(\mathbf{X})$  and  $\mathbf{u} \in C(\mathbf{X})^\perp$
- ▶ Then

$$\begin{aligned}\psi = E[\mathbf{a}^T \mathbf{Y}] &= E[\mathbf{a}^{*T} \mathbf{Y}] + E[\mathbf{u}^T \mathbf{Y}] \\ &= E[\mathbf{a}^{*T} \mathbf{Y}] + 0\end{aligned}$$

$$E[\mathbf{u}^T \mathbf{Y}] = \mathbf{u}^T \mathbf{X} \boldsymbol{\beta}$$

since  $\mathbf{u} \perp C(\mathbf{X})$  (i.e.  $\mathbf{u} \in C(\mathbf{X})^\perp$ )  $E[\mathbf{u}^T \mathbf{Y}] = 0$

- ▶ Thus  $\mathbf{a}^{*T} \mathbf{Y}$  is also an unbiased linear estimator of  $\psi$  with  $\mathbf{a}^* \in C(\mathbf{X})$

# Uniqueness

Proof.

Suppose that there is another  $\mathbf{v} \in C(\mathbf{X})$  such that  $E[\mathbf{v}^T \mathbf{Y}] = \psi$ .

Then for all  $\beta$

# Uniqueness

Proof.

Suppose that there is another  $\mathbf{v} \in C(\mathbf{X})$  such that  $E[\mathbf{v}^T \mathbf{Y}] = \psi$ .

Then for all  $\beta$

$$0 = E[\mathbf{a}^{*T} \mathbf{Y}] - E[\mathbf{v}^T \mathbf{Y}]$$

# Uniqueness

## Proof.

Suppose that there is another  $\mathbf{v} \in C(\mathbf{X})$  such that  $E[\mathbf{v}^T \mathbf{Y}] = \psi$ .

Then for all  $\beta$

$$\begin{aligned} 0 &= E[\mathbf{a}^{*T} \mathbf{Y}] - E[\mathbf{v}^T \mathbf{Y}] \\ &= (\mathbf{a}^* - \mathbf{v})^T \mathbf{X} \beta \end{aligned}$$

# Uniqueness

## Proof.

Suppose that there is another  $\mathbf{v} \in C(\mathbf{X})$  such that  $E[\mathbf{v}^T \mathbf{Y}] = \psi$ .

Then for all  $\beta$

$$\begin{aligned} 0 &= E[\mathbf{a}^{*T} \mathbf{Y}] - E[\mathbf{v}^T \mathbf{Y}] \\ &= (\mathbf{a}^* - \mathbf{v})^T \mathbf{X} \beta \end{aligned}$$

$$\text{So } (\mathbf{a}^* - \mathbf{v})^T \mathbf{X} = 0 \quad \text{for all } \beta$$

# Uniqueness

## Proof.

Suppose that there is another  $\mathbf{v} \in C(\mathbf{X})$  such that  $E[\mathbf{v}^T \mathbf{Y}] = \psi$ .  
Then for all  $\beta$

$$\begin{aligned} 0 &= E[\mathbf{a}^{*T} \mathbf{Y}] - E[\mathbf{v}^T \mathbf{Y}] \\ &= (\mathbf{a}^* - \mathbf{v})^T \mathbf{X} \beta \\ \text{So } (\mathbf{a}^* - \mathbf{v})^T \mathbf{X} &= 0 \quad \text{for all } \beta \end{aligned}$$

► Implies  $(\mathbf{a}^* - \mathbf{v}) \in C(\mathbf{X})^\perp$



# Uniqueness

## Proof.

Suppose that there is another  $\mathbf{v} \in C(\mathbf{X})$  such that  $E[\mathbf{v}^T \mathbf{Y}] = \psi$ .  
Then for all  $\beta$

$$\begin{aligned} 0 &= E[\mathbf{a}^{*T} \mathbf{Y}] - E[\mathbf{v}^T \mathbf{Y}] \\ &= (\mathbf{a}^* - \mathbf{v})^T \mathbf{X} \beta \\ \text{So } (\mathbf{a}^* - \mathbf{v})^T \mathbf{X} &= 0 \quad \text{for all } \beta \end{aligned}$$

- Implies  $(\mathbf{a}^* - \mathbf{v}) \in C(\mathbf{X})^\perp$
- but by assumption  $(\mathbf{a}^* - \mathbf{v}) \in C(\mathbf{X})$

# Uniqueness

## Proof.

Suppose that there is another  $\mathbf{v} \in C(\mathbf{X})$  such that  $E[\mathbf{v}^T \mathbf{Y}] = \psi$ .

Then for all  $\beta$

$$\begin{aligned} 0 &= E[\mathbf{a}^{*T} \mathbf{Y}] - E[\mathbf{v}^T \mathbf{Y}] \\ &= (\mathbf{a}^* - \mathbf{v})^T \mathbf{X} \beta \end{aligned}$$

$$\text{So } (\mathbf{a}^* - \mathbf{v})^T \mathbf{X} = 0 \quad \text{for all } \beta$$

- Implies  $(\mathbf{a}^* - \mathbf{v}) \in C(\mathbf{X})^\perp$
- but by assumption  $(\mathbf{a}^* - \mathbf{v}) \in C(\mathbf{X})$  ( $C(\mathbf{X})$  is a vector space)

# Uniqueness

## Proof.

Suppose that there is another  $\mathbf{v} \in C(\mathbf{X})$  such that  $E[\mathbf{v}^T \mathbf{Y}] = \psi$ .  
Then for all  $\beta$

$$\begin{aligned} 0 &= E[\mathbf{a}^{*T} \mathbf{Y}] - E[\mathbf{v}^T \mathbf{Y}] \\ &= (\mathbf{a}^* - \mathbf{v})^T \mathbf{X} \beta \\ \text{So } (\mathbf{a}^* - \mathbf{v})^T \mathbf{X} &= 0 \quad \text{for all } \beta \end{aligned}$$

- Implies  $(\mathbf{a}^* - \mathbf{v}) \in C(\mathbf{X})^\perp$
- but by assumption  $(\mathbf{a}^* - \mathbf{v}) \in C(\mathbf{X})$  ( $C(\mathbf{X})$  is a vector space)
- the only vector in BOTH is  $\mathbf{0}$ , so  $\mathbf{a}^* = \mathbf{v}$

# Uniqueness

## Proof.

Suppose that there is another  $\mathbf{v} \in C(\mathbf{X})$  such that  $E[\mathbf{v}^T \mathbf{Y}] = \psi$ .

Then for all  $\beta$

$$\begin{aligned} 0 &= E[\mathbf{a}^{*T} \mathbf{Y}] - E[\mathbf{v}^T \mathbf{Y}] \\ &= (\mathbf{a}^* - \mathbf{v})^T \mathbf{X} \beta \end{aligned}$$

$$\text{So } (\mathbf{a}^* - \mathbf{v})^T \mathbf{X} = 0 \quad \text{for all } \beta$$

- ▶ Implies  $(\mathbf{a}^* - \mathbf{v}) \in C(\mathbf{X})^\perp$
- ▶ but by assumption  $(\mathbf{a}^* - \mathbf{v}) \in C(\mathbf{X})$  ( $C(\mathbf{X})$  is a vector space)
- ▶ the only vector in BOTH is  $\mathbf{0}$ , so  $\mathbf{a}^* = \mathbf{v}$

Therefore  $\mathbf{a}^{*T} \mathbf{Y}$  is the unique linear unbiased estimator of  $\psi$  with  $\mathbf{a}^* \in C(\mathbf{X})$ . □

# Proof of Minimum Variance (G-M)

- ▶ Let  $\mathbf{a}^{*T}\mathbf{Y}$  be the unique unbiased linear estimator of  $\psi$  with  $\mathbf{a}^* \in C(\mathbf{X})$ .

# Proof of Minimum Variance (G-M)

- ▶ Let  $\mathbf{a}^{*T}\mathbf{Y}$  be the unique unbiased linear estimator of  $\psi$  with  $\mathbf{a}^* \in C(\mathbf{X})$ .
- ▶ Let  $\mathbf{a}^T\mathbf{Y}$  be any unbiased estimate of  $\psi$ ;  $\mathbf{a} = \mathbf{a}^* + \mathbf{u}$  with  $\mathbf{a}^* \in C(\mathbf{X})$  and  $\mathbf{u} \in C(\mathbf{X})^\perp$

# Proof of Minimum Variance (G-M)

- ▶ Let  $\mathbf{a}^{*T}\mathbf{Y}$  be the unique unbiased linear estimator of  $\psi$  with  $\mathbf{a}^* \in C(\mathbf{X})$ .
- ▶ Let  $\mathbf{a}^T\mathbf{Y}$  be any unbiased estimate of  $\psi$ ;  $\mathbf{a} = \mathbf{a}^* + \mathbf{u}$  with  $\mathbf{a}^* \in C(\mathbf{X})$  and  $\mathbf{u} \in C(\mathbf{X})^\perp$

$$\text{Var}(\mathbf{a}^T\mathbf{Y}) = \mathbf{a}^T\text{Cov}(\mathbf{Y})\mathbf{a}$$

# Proof of Minimum Variance (G-M)

- ▶ Let  $\mathbf{a}^{*T}\mathbf{Y}$  be the unique unbiased linear estimator of  $\psi$  with  $\mathbf{a}^* \in C(\mathbf{X})$ .
- ▶ Let  $\mathbf{a}^T\mathbf{Y}$  be any unbiased estimate of  $\psi$ ;  $\mathbf{a} = \mathbf{a}^* + \mathbf{u}$  with  $\mathbf{a}^* \in C(\mathbf{X})$  and  $\mathbf{u} \in C(\mathbf{X})^\perp$

$$\begin{aligned}\text{Var}(\mathbf{a}^T\mathbf{Y}) &= \mathbf{a}^T\text{Cov}(\mathbf{Y})\mathbf{a} \\ &= \sigma^2\|\mathbf{a}\|^2\end{aligned}$$



# Proof of Minimum Variance (G-M)

- ▶ Let  $\mathbf{a}^{*T}\mathbf{Y}$  be the unique unbiased linear estimator of  $\psi$  with  $\mathbf{a}^* \in C(\mathbf{X})$ .
- ▶ Let  $\mathbf{a}^T\mathbf{Y}$  be any unbiased estimate of  $\psi$ ;  $\mathbf{a} = \mathbf{a}^* + \mathbf{u}$  with  $\mathbf{a}^* \in C(\mathbf{X})$  and  $\mathbf{u} \in C(\mathbf{X})^\perp$

$$\begin{aligned}\text{Var}(\mathbf{a}^T\mathbf{Y}) &= \mathbf{a}^T\text{Cov}(\mathbf{Y})\mathbf{a} \\ &= \sigma^2\|\mathbf{a}\|^2 \\ &= \sigma^2(\|\mathbf{a}^*\|^2 + \|\mathbf{u}\|^2 + 2\mathbf{a}^{*T}\mathbf{u})\end{aligned}$$

# Proof of Minimum Variance (G-M)

- ▶ Let  $\mathbf{a}^{*T}\mathbf{Y}$  be the unique unbiased linear estimator of  $\psi$  with  $\mathbf{a}^* \in C(\mathbf{X})$ .
- ▶ Let  $\mathbf{a}^T\mathbf{Y}$  be any unbiased estimate of  $\psi$ ;  $\mathbf{a} = \mathbf{a}^* + \mathbf{u}$  with  $\mathbf{a}^* \in C(\mathbf{X})$  and  $\mathbf{u} \in C(\mathbf{X})^\perp$

$$\begin{aligned}\text{Var}(\mathbf{a}^T\mathbf{Y}) &= \mathbf{a}^T\text{Cov}(\mathbf{Y})\mathbf{a} \\ &= \sigma^2\|\mathbf{a}\|^2 \\ &= \sigma^2(\|\mathbf{a}^*\|^2 + \|\mathbf{u}\|^2 + 2\mathbf{a}^{*T}\mathbf{u}) \\ &= \sigma^2(\|\mathbf{a}^*\|^2 + \|\mathbf{u}\|^2) + 0\end{aligned}$$

# Proof of Minimum Variance (G-M)

- ▶ Let  $\mathbf{a}^{*T}\mathbf{Y}$  be the unique unbiased linear estimator of  $\psi$  with  $\mathbf{a}^* \in C(\mathbf{X})$ .
- ▶ Let  $\mathbf{a}^T\mathbf{Y}$  be any unbiased estimate of  $\psi$ ;  $\mathbf{a} = \mathbf{a}^* + \mathbf{u}$  with  $\mathbf{a}^* \in C(\mathbf{X})$  and  $\mathbf{u} \in C(\mathbf{X})^\perp$

$$\begin{aligned}\text{Var}(\mathbf{a}^T\mathbf{Y}) &= \mathbf{a}^T\text{Cov}(\mathbf{Y})\mathbf{a} \\ &= \sigma^2\|\mathbf{a}\|^2 \\ &= \sigma^2(\|\mathbf{a}^*\|^2 + \|\mathbf{u}\|^2 + 2\mathbf{a}^{*T}\mathbf{u}) \\ &= \sigma^2(\|\mathbf{a}^*\|^2 + \|\mathbf{u}\|^2) + 0 \\ &= \text{Var}(\mathbf{a}^{*T}\mathbf{Y}) + \sigma^2\|\mathbf{u}\|^2\end{aligned}$$

# Proof of Minimum Variance (G-M)

- ▶ Let  $\mathbf{a}^{*T}\mathbf{Y}$  be the unique unbiased linear estimator of  $\psi$  with  $\mathbf{a}^* \in C(\mathbf{X})$ .
- ▶ Let  $\mathbf{a}^T\mathbf{Y}$  be any unbiased estimate of  $\psi$ ;  $\mathbf{a} = \mathbf{a}^* + \mathbf{u}$  with  $\mathbf{a}^* \in C(\mathbf{X})$  and  $\mathbf{u} \in C(\mathbf{X})^\perp$

$$\begin{aligned}\text{Var}(\mathbf{a}^T\mathbf{Y}) &= \mathbf{a}^T\text{Cov}(\mathbf{Y})\mathbf{a} \\ &= \sigma^2\|\mathbf{a}\|^2 \\ &= \sigma^2(\|\mathbf{a}^*\|^2 + \|\mathbf{u}\|^2 + 2\mathbf{a}^{*T}\mathbf{u}) \\ &= \sigma^2(\|\mathbf{a}^*\|^2 + \|\mathbf{u}\|^2) + 0 \\ &= \text{Var}(\mathbf{a}^{*T}\mathbf{Y}) + \sigma^2\|\mathbf{u}\|^2 \\ &\geq \text{Var}(\mathbf{a}^{*T}\mathbf{Y})\end{aligned}$$

# Proof of Minimum Variance (G-M)

- ▶ Let  $\mathbf{a}^{*T}\mathbf{Y}$  be the unique unbiased linear estimator of  $\psi$  with  $\mathbf{a}^* \in C(\mathbf{X})$ .
- ▶ Let  $\mathbf{a}^T\mathbf{Y}$  be any unbiased estimate of  $\psi$ ;  $\mathbf{a} = \mathbf{a}^* + \mathbf{u}$  with  $\mathbf{a}^* \in C(\mathbf{X})$  and  $\mathbf{u} \in C(\mathbf{X})^\perp$

$$\begin{aligned}\text{Var}(\mathbf{a}^T\mathbf{Y}) &= \mathbf{a}^T\text{Cov}(\mathbf{Y})\mathbf{a} \\ &= \sigma^2\|\mathbf{a}\|^2 \\ &= \sigma^2(\|\mathbf{a}^*\|^2 + \|\mathbf{u}\|^2 + 2\mathbf{a}^{*T}\mathbf{u}) \\ &= \sigma^2(\|\mathbf{a}^*\|^2 + \|\mathbf{u}\|^2) + 0 \\ &= \text{Var}(\mathbf{a}^{*T}\mathbf{Y}) + \sigma^2\|\mathbf{u}\|^2 \\ &\geq \text{Var}(\mathbf{a}^{*T}\mathbf{Y})\end{aligned}$$

with equality if and only if  $\mathbf{a} = \mathbf{a}^*$

# Proof of Minimum Variance (G-M)

- ▶ Let  $\mathbf{a}^{*T}\mathbf{Y}$  be the unique unbiased linear estimator of  $\psi$  with  $\mathbf{a}^* \in C(\mathbf{X})$ .
- ▶ Let  $\mathbf{a}^T\mathbf{Y}$  be any unbiased estimate of  $\psi$ ;  $\mathbf{a} = \mathbf{a}^* + \mathbf{u}$  with  $\mathbf{a}^* \in C(\mathbf{X})$  and  $\mathbf{u} \in C(\mathbf{X})^\perp$

$$\begin{aligned}\text{Var}(\mathbf{a}^T\mathbf{Y}) &= \mathbf{a}^T\text{Cov}(\mathbf{Y})\mathbf{a} \\ &= \sigma^2\|\mathbf{a}\|^2 \\ &= \sigma^2(\|\mathbf{a}^*\|^2 + \|\mathbf{u}\|^2 + 2\mathbf{a}^{*T}\mathbf{u}) \\ &= \sigma^2(\|\mathbf{a}^*\|^2 + \|\mathbf{u}\|^2) + 0 \\ &= \text{Var}(\mathbf{a}^{*T}\mathbf{Y}) + \sigma^2\|\mathbf{u}\|^2 \\ &\geq \text{Var}(\mathbf{a}^{*T}\mathbf{Y})\end{aligned}$$

with equality if and only if  $\mathbf{a} = \mathbf{a}^*$

Hence  $\mathbf{a}^{*T}\mathbf{Y}$  is the unique linear unbiased estimator of  $\psi$  with minimum variance

# Proof of Minimum Variance (G-M)

- ▶ Let  $\mathbf{a}^{*T}\mathbf{Y}$  be the unique unbiased linear estimator of  $\psi$  with  $\mathbf{a}^* \in C(\mathbf{X})$ .
- ▶ Let  $\mathbf{a}^T\mathbf{Y}$  be any unbiased estimate of  $\psi$ ;  $\mathbf{a} = \mathbf{a}^* + \mathbf{u}$  with  $\mathbf{a}^* \in C(\mathbf{X})$  and  $\mathbf{u} \in C(\mathbf{X})^\perp$

$$\begin{aligned}\text{Var}(\mathbf{a}^T\mathbf{Y}) &= \mathbf{a}^T\text{Cov}(\mathbf{Y})\mathbf{a} \\ &= \sigma^2\|\mathbf{a}\|^2 \\ &= \sigma^2(\|\mathbf{a}^*\|^2 + \|\mathbf{u}\|^2 + 2\mathbf{a}^{*T}\mathbf{u}) \\ &= \sigma^2(\|\mathbf{a}^*\|^2 + \|\mathbf{u}\|^2) + 0 \\ &= \text{Var}(\mathbf{a}^{*T}\mathbf{Y}) + \sigma^2\|\mathbf{u}\|^2 \\ &\geq \text{Var}(\mathbf{a}^{*T}\mathbf{Y})\end{aligned}$$

with equality if and only if  $\mathbf{a} = \mathbf{a}^*$

Hence  $\mathbf{a}^{*T}\mathbf{Y}$  is the unique linear unbiased estimator of  $\psi$  with minimum variance "BLUE" = Best Linear Unbiased Estimator

# Continued

Proof.

Show that  $\hat{\psi} = \mathbf{a}^{*T} \mathbf{Y} = \boldsymbol{\lambda}^T \hat{\boldsymbol{\beta}}$



# Continued

Proof.

Show that  $\hat{\psi} = \mathbf{a}^{*T} \mathbf{Y} = \boldsymbol{\lambda}^T \hat{\boldsymbol{\beta}}$

Since  $\mathbf{a}^* \in C(\mathbf{X})$  we have  $\mathbf{a}^* = \mathbf{P}_X \mathbf{a}^*$

## Continued

Proof.

Show that  $\hat{\psi} = \mathbf{a}^{*T} \mathbf{Y} = \boldsymbol{\lambda}^T \hat{\boldsymbol{\beta}}$

Since  $\mathbf{a}^* \in C(\mathbf{X})$  we have  $\mathbf{a}^* = \mathbf{P}_X \mathbf{a}^*$

$$\mathbf{a}^{*T} \mathbf{Y} = \mathbf{a}^{*T} \mathbf{P}_X^T \mathbf{Y}$$

# Continued

Proof.

Show that  $\hat{\psi} = \mathbf{a}^{*T} \mathbf{Y} = \boldsymbol{\lambda}^T \hat{\boldsymbol{\beta}}$

Since  $\mathbf{a}^* \in C(\mathbf{X})$  we have  $\mathbf{a}^* = \mathbf{P}_X \mathbf{a}^*$

$$\begin{aligned} \mathbf{a}^{*T} \mathbf{Y} &= \mathbf{a}^{*T} \mathbf{P}_X^T \mathbf{Y} \\ &= \mathbf{a}^{*T} \mathbf{P}_X \mathbf{Y} \end{aligned}$$

## Continued

Proof.

Show that  $\hat{\psi} = \mathbf{a}^{*T} \mathbf{Y} = \boldsymbol{\lambda}^T \hat{\boldsymbol{\beta}}$

Since  $\mathbf{a}^* \in C(\mathbf{X})$  we have  $\mathbf{a}^* = \mathbf{P}_X \mathbf{a}^*$

$$\begin{aligned} \mathbf{a}^{*T} \mathbf{Y} &= \mathbf{a}^{*T} \mathbf{P}_X^T \mathbf{Y} \\ &= \mathbf{a}^{*T} \mathbf{P}_X \mathbf{Y} \\ &= \mathbf{a}^{*T} \mathbf{X} \hat{\boldsymbol{\beta}} \end{aligned}$$

## Continued

Proof.

Show that  $\hat{\psi} = \mathbf{a}^{*T} \mathbf{Y} = \boldsymbol{\lambda}^T \hat{\boldsymbol{\beta}}$

Since  $\mathbf{a}^* \in C(\mathbf{X})$  we have  $\mathbf{a}^* = \mathbf{P}_X \mathbf{a}^*$

$$\begin{aligned} \mathbf{a}^{*T} \mathbf{Y} &= \mathbf{a}^{*T} \mathbf{P}_X^T \mathbf{Y} \\ &= \mathbf{a}^{*T} \mathbf{P}_X \mathbf{Y} \\ &= \mathbf{a}^{*T} \mathbf{X} \hat{\boldsymbol{\beta}} \\ &= \boldsymbol{\lambda}^T \hat{\boldsymbol{\beta}} \end{aligned}$$

## Continued

### Proof.

Show that  $\hat{\psi} = \mathbf{a}^{*T} \mathbf{Y} = \boldsymbol{\lambda}^T \hat{\boldsymbol{\beta}}$

Since  $\mathbf{a}^* \in C(\mathbf{X})$  we have  $\mathbf{a}^* = \mathbf{P}_X \mathbf{a}^*$

$$\begin{aligned} \mathbf{a}^{*T} \mathbf{Y} &= \mathbf{a}^{*T} \mathbf{P}_X^T \mathbf{Y} \\ &= \mathbf{a}^{*T} \mathbf{P}_X \mathbf{Y} \\ &= \mathbf{a}^{*T} \mathbf{X} \hat{\boldsymbol{\beta}} \\ &= \boldsymbol{\lambda}^T \hat{\boldsymbol{\beta}} \end{aligned}$$

for  $\boldsymbol{\lambda}^T = \mathbf{a}^{*T} \mathbf{X}$  or  $\boldsymbol{\lambda} = \mathbf{X}^T \mathbf{a}^*$



- ▶ Gauss-Markov Theorem says that OLS has minimum variance in the class of all Linear Unbiased estimators

# MVUE

- ▶ Gauss-Markov Theorem says that OLS has minimum variance in the class of all Linear Unbiased estimators
- ▶ Requires just first and second moments



# MVUE

- ▶ Gauss-Markov Theorem says that OLS has minimum variance in the class of all Linear Unbiased estimators
- ▶ Requires just first and second moments
- ▶ Additional assumption of normality, OLS = MLEs have minimum variance out of **ALL** unbiased estimators (MVUE); not just linear estimators

# MVUE

- ▶ Gauss-Markov Theorem says that OLS has minimum variance in the class of all Linear Unbiased estimators
- ▶ Requires just first and second moments
- ▶ Additional assumption of normality, OLS = MLEs have minimum variance out of ALL unbiased estimators (MVUE); not just linear estimators (requires Completeness and Rao-Blackwell Theorem - next semester)

# MVUE

- ▶ Gauss-Markov Theorem says that OLS has minimum variance in the class of all Linear Unbiased estimators
- ▶ Requires just first and second moments
- ▶ Additional assumption of normality, OLS = MLEs have minimum variance out of ALL unbiased estimators (MVUE); not just linear estimators (requires Completeness and Rao-Blackwell Theorem - next semester)

# Prediction

- For predicting at new  $\mathbf{x}_*$  is there always a unique unbiased estimator of  $E[\mathbf{Y} \mid \mathbf{x}_*]$ ?

# Prediction

- ▶ For predicting at new  $\mathbf{x}_*$  is there always a unique unbiased estimator of  $E[\mathbf{Y} \mid \mathbf{x}_*]$ ?
- ▶ If one does exist, how do we know that if we are given  $\lambda$ ?

# Existence

- ▶  $\mathbf{x}_*^T \boldsymbol{\beta}$  has a unique unbiased estimator if  $\mathbf{x}_* \equiv \boldsymbol{\lambda} = \mathbf{X}^T \mathbf{a}$

# Existence

- ▶  $\mathbf{x}_*^T \boldsymbol{\beta}$  has a unique unbiased estimator if  $\mathbf{x}_* \equiv \boldsymbol{\lambda} = \mathbf{X}^T \mathbf{a}$
- ▶ Clearly if  $\mathbf{x}_* = \mathbf{x}_i$  ( $i$ th row of observed data) then it is estimable with  $\mathbf{a}$  equal to the vector with a 1 in the  $i$ th position even if  $\mathbf{X}$  is not full rank!

# Existence

- ▶  $\mathbf{x}_*^T \boldsymbol{\beta}$  has a unique unbiased estimator if  $\mathbf{x}_* \equiv \boldsymbol{\lambda} = \mathbf{X}^T \mathbf{a}$
- ▶ Clearly if  $\mathbf{x}_* = \mathbf{x}_i$  ( $i$ th row of observed data) then it is estimable with  $\mathbf{a}$  equal to the vector with a 1 in the  $i$ th position even if  $\mathbf{X}$  is not full rank!
- ▶ What about out of sample prediction?



# Existence

- ▶  $\mathbf{x}_*^T \boldsymbol{\beta}$  has a unique unbiased estimator if  $\mathbf{x}_* \equiv \boldsymbol{\lambda} = \mathbf{X}^T \mathbf{a}$
- ▶ Clearly if  $\mathbf{x}_* = \mathbf{x}_i$  ( $i$ th row of observed data) then it is estimable with  $\mathbf{a}$  equal to the vector with a 1 in the  $i$ th position even if  $\mathbf{X}$  is not full rank!
- ▶ What about out of sample prediction?

# Example

```
x1 = -4:4
x2 = c(-2, 1, -1, 2, 0, 2, -1, 1, -2)
x3 = 3*x1 - 2*x2
x4 = x2 - x1 + 4
Y = 1+x1+x2+x3+x4 + c(-.5,.5,.5,-.5,0,.5,-.5,-.5,.5)
dev.set = data.frame(Y, x1, x2, x3, x4)
lm1234 = lm(Y ~ x1 + x2 + x3 + x4, data=dev.set)
round(coefficients(lm1234), 4)
```

## (Intercept)	x1	x2	x3	x4
## 5	3	0	NA	NA

```
lm3412 = lm(Y ~ x3 + x4 + x1 + x2, data = dev.set)
round(coefficients(lm3412), 4)
```

## (Intercept)	x3	x4	x1	x2
## -19	3	6	NA	NA

# In Sample Predictions

```
cbind(dev.set, predict(lm1234), predict(lm3412))
```

##		Y	x1	x2	x3	x4	predict(lm1234)	predict(lm3412)
## 1		-7.5	-4	-2	-8	6	-7	-7
## 2		-3.5	-3	1	-11	8	-4	-4
## 3		-0.5	-2	-1	-4	5	-1	-1
## 4		1.5	-1	2	-7	7	2	2
## 5		5.0	0	0	0	4	5	5
## 6		8.5	1	2	-1	5	8	8
## 7		10.5	2	-1	8	1	11	11
## 8		13.5	3	1	7	2	14	14
## 9		17.5	4	-2	16	-2	17	17

Both models agree for estimating the mean at the observed **X** points!

# Out of Sample

```
out = data.frame(test.set,  
  Y1234=predict(lm1234, new=test.set),  
  Y3412=predict(lm3412, new=test.set))
```

out

##	x1	x2	x3	x4	Y1234	Y3412
## 1	3	1	7	2	14	14
## 2	6	2	14	4	23	47
## 3	6	2	14	0	23	23
## 4	0	0	0	4	5	5
## 5	0	0	0	0	5	-19
## 6	1	2	3	4	8	14

# Out of Sample

```
out = data.frame(test.set,  
  Y1234=predict(lm1234, new=test.set),  
  Y3412=predict(lm3412, new=test.set))
```

out

##	x1	x2	x3	x4	Y1234	Y3412
## 1	3	1	7	2	14	14
## 2	6	2	14	4	23	47
## 3	6	2	14	0	23	23
## 4	0	0	0	4	5	5
## 5	0	0	0	0	5	-19
## 6	1	2	3	4	8	14

Agreement for cases 1, 3, and 4 only! Can we determine that without finding the predictions and comparing?

# Determining Estimable $\lambda$

- ▶ Estimable means that  $\lambda^T = \mathbf{a}^T \mathbf{X}$  for  $\mathbf{a} \in C(\mathbf{X})$

## Determining Estimable $\lambda$

- ▶ Estimable means that  $\lambda^T = \mathbf{a}^T \mathbf{X}$  for  $\mathbf{a} \in C(\mathbf{X})$
- ▶ Transpose:  $\lambda = \mathbf{X}^T \mathbf{a}$  for  $\mathbf{a} \in C(\mathbf{X})$

# Determining Estimable $\lambda$

- ▶ Estimable means that  $\lambda^T = \mathbf{a}^T \mathbf{X}$  for  $\mathbf{a} \in C(\mathbf{X})$
- ▶ Transpose:  $\lambda = \mathbf{X}^T \mathbf{a}$  for  $\mathbf{a} \in C(\mathbf{X})$
- ▶  $\lambda \in C(\mathbf{X}^T)$  ( $\lambda \in R(\mathbf{X})$ )



# Determining Estimable $\lambda$

- ▶ Estimable means that  $\lambda^T = \mathbf{a}^T \mathbf{X}$  for  $\mathbf{a} \in C(\mathbf{X})$
- ▶ Transpose:  $\lambda = \mathbf{X}^T \mathbf{a}$  for  $\mathbf{a} \in C(\mathbf{X})$
- ▶  $\lambda \in C(\mathbf{X}^T)$  ( $\lambda \in R(\mathbf{X})$ )
- ▶  $\lambda \perp C(\mathbf{X}^T)^\perp$

# Determining Estimable $\lambda$

- ▶ Estimable means that  $\lambda^T = \mathbf{a}^T \mathbf{X}$  for  $\mathbf{a} \in C(\mathbf{X})$
- ▶ Transpose:  $\lambda = \mathbf{X}^T \mathbf{a}$  for  $\mathbf{a} \in C(\mathbf{X})$
- ▶  $\lambda \in C(\mathbf{X}^T)$  ( $\lambda \in R(\mathbf{X})$ )
- ▶  $\lambda \perp C(\mathbf{X}^T)^\perp$
- ▶  $C(\mathbf{X}^T)^\perp$  is the null space of  $\mathbf{X}$

# Determining Estimable $\lambda$

- ▶ Estimable means that  $\lambda^T = \mathbf{a}^T \mathbf{X}$  for  $\mathbf{a} \in C(\mathbf{X})$
- ▶ Transpose:  $\lambda = \mathbf{X}^T \mathbf{a}$  for  $\mathbf{a} \in C(\mathbf{X})$
- ▶  $\lambda \in C(\mathbf{X}^T)$  ( $\lambda \in R(\mathbf{X})$ )
- ▶  $\lambda \perp C(\mathbf{X}^T)^\perp$
- ▶  $C(\mathbf{X}^T)^\perp$  is the null space of  $\mathbf{X}$

$$\mathbf{v} \perp C(\mathbf{X}^T) : \mathbf{X}\mathbf{v} = 0 \Leftrightarrow \mathbf{v} \in N(\mathbf{X})$$

# Determining Estimable $\lambda$

- ▶ Estimable means that  $\lambda^T = \mathbf{a}^T \mathbf{X}$  for  $\mathbf{a} \in C(\mathbf{X})$
- ▶ Transpose:  $\lambda = \mathbf{X}^T \mathbf{a}$  for  $\mathbf{a} \in C(\mathbf{X})$
- ▶  $\lambda \in C(\mathbf{X}^T)$  ( $\lambda \in R(\mathbf{X})$ )
- ▶  $\lambda \perp C(\mathbf{X}^T)^\perp$
- ▶  $C(\mathbf{X}^T)^\perp$  is the null space of  $\mathbf{X}$

$$\mathbf{v} \perp C(\mathbf{X}^T) : \mathbf{X}\mathbf{v} = 0 \Leftrightarrow \mathbf{v} \in N(\mathbf{X})$$

- ▶  $\lambda \perp N(\mathbf{X})$

# Determining Estimable $\lambda$

- ▶ Estimable means that  $\lambda^T = \mathbf{a}^T \mathbf{X}$  for  $\mathbf{a} \in C(\mathbf{X})$
- ▶ Transpose:  $\lambda = \mathbf{X}^T \mathbf{a}$  for  $\mathbf{a} \in C(\mathbf{X})$
- ▶  $\lambda \in C(\mathbf{X}^T)$  ( $\lambda \in R(\mathbf{X})$ )
- ▶  $\lambda \perp C(\mathbf{X}^T)^\perp$
- ▶  $C(\mathbf{X}^T)^\perp$  is the null space of  $\mathbf{X}$

$$\mathbf{v} \perp C(\mathbf{X}^T) : \mathbf{X}\mathbf{v} = 0 \Leftrightarrow \mathbf{v} \in N(\mathbf{X})$$

- ▶  $\lambda \perp N(\mathbf{X})$
- ▶ if  $\mathbf{P}$  is a projection onto  $C(\mathbf{X}^T)$  then  $\mathbf{I} - \mathbf{P}$  is a projection onto  $N(\mathbf{X})$  and therefore  $(\mathbf{I} - \mathbf{P})\lambda = \mathbf{0}$  if  $\lambda$  is estimable

## Determining Estimable $\lambda$

- ▶ Estimable means that  $\lambda^T = \mathbf{a}^T \mathbf{X}$  for  $\mathbf{a} \in C(\mathbf{X})$
- ▶ Transpose:  $\lambda = \mathbf{X}^T \mathbf{a}$  for  $\mathbf{a} \in C(\mathbf{X})$
- ▶  $\lambda \in C(\mathbf{X}^T)$  ( $\lambda \in R(\mathbf{X})$ )
- ▶  $\lambda \perp C(\mathbf{X}^T)^\perp$
- ▶  $C(\mathbf{X}^T)^\perp$  is the null space of  $\mathbf{X}$

$$\mathbf{v} \perp C(\mathbf{X}^T) : \mathbf{X}\mathbf{v} = 0 \Leftrightarrow \mathbf{v} \in N(\mathbf{X})$$

- ▶  $\lambda \perp N(\mathbf{X})$
- ▶ if  $\mathbf{P}$  is a projection onto  $C(\mathbf{X}^T)$  then  $\mathbf{I} - \mathbf{P}$  is a projection onto  $N(\mathbf{X})$  and therefore  $(\mathbf{I} - \mathbf{P})\lambda = \mathbf{0}$  if  $\lambda$  is estimable

Take  $\mathbf{P}_{\mathbf{X}^T} = (\mathbf{X}^T \mathbf{X})(\mathbf{X}^T \mathbf{X})^-$  as a projection onto  $C(\mathbf{X}^T)$  and show  $(\mathbf{I} - \mathbf{P}_{\mathbf{X}^T})\lambda = \mathbf{0}_p$

# Example

```
library("estimability" )  
cbind(epredict(lm1234, test.set), epredict(lm3412, test.set))  
  
##      [,1] [,2]  
## 1      14  14  
## 2      NA  NA  
## 3      23  23  
## 4       5   5  
## 5      NA  NA  
## 6      NA  NA
```

Rows 2, 5, and 6 are not estimable! No linear unbiased estimator

# Summary

- ▶ When BLUEs exist, under normality they are MVUE (ditto for prediction - BLUP)



# Summary

- ▶ When BLUEs exist, under normality they are MVUE (ditto for prediction - BLUP)
- ▶ BLUE/BLUP do not always exist for estimation/prediction if  $\mathbf{X}$  is not full rank

# Summary

- ▶ When BLUEs exist, under normality they are MVUE (ditto for prediction - BLUP)
- ▶ BLUE/BLUP do not always exist for estimation/prediction if  $\mathbf{X}$  is not full rank
- ▶ may occur with redundancies for modest  $p < n$  and of course  $p > n$

# Summary

- ▶ When BLUEs exist, under normality they are MVUE (ditto for prediction - BLUP)
- ▶ BLUE/BLUP do not always exist for estimation/prediction if  $\mathbf{X}$  is not full rank
- ▶ may occur with redundancies for modest  $p < n$  and of course  $p > n$
- ▶ Eliminate redundancies by removing variables (variable selection)

# Summary

- ▶ When BLUEs exist, under normality they are MVUE (ditto for prediction - BLUP)
- ▶ BLUE/BLUP do not always exist for estimation/prediction if  $\mathbf{X}$  is not full rank
- ▶ may occur with redundancies for modest  $p < n$  and of course  $p > n$
- ▶ Eliminate redundancies by removing variables (variable selection)
- ▶ Consider alternative estimators (Bayes and related)

# Other Estimators

What about some estimator  $g(\mathbf{Y})$  that is not unbiased?

# Other Estimators

What about some estimator  $g(\mathbf{Y})$  that is not unbiased?

- ▶ Mean Squared Error for estimator  $g(\mathbf{Y})$  of  $\boldsymbol{\lambda}^T \boldsymbol{\beta}$  is

$$E[g(\mathbf{Y}) - \boldsymbol{\lambda}^T \boldsymbol{\beta}]^2 = \text{Var}(g(\mathbf{Y})) + \text{Bias}^2(g(\mathbf{Y}))$$

where  $\text{Bias} = E[g(\mathbf{Y})] - \boldsymbol{\lambda}^T \boldsymbol{\beta}$

# Other Estimators

What about some estimator  $g(\mathbf{Y})$  that is not unbiased?

- ▶ Mean Squared Error for estimator  $g(\mathbf{Y})$  of  $\boldsymbol{\lambda}^T \boldsymbol{\beta}$  is

$$E[g(\mathbf{Y}) - \boldsymbol{\lambda}^T \boldsymbol{\beta}]^2 = \text{Var}(g(\mathbf{Y})) + \text{Bias}^2(g(\mathbf{Y}))$$

where  $\text{Bias} = E[g(\mathbf{Y})] - \boldsymbol{\lambda}^T \boldsymbol{\beta}$

- ▶ Bias vs Variance tradeoff

# Other Estimators

What about some estimator  $g(\mathbf{Y})$  that is not unbiased?

- ▶ Mean Squared Error for estimator  $g(\mathbf{Y})$  of  $\boldsymbol{\lambda}^T \boldsymbol{\beta}$  is

$$E[g(\mathbf{Y}) - \boldsymbol{\lambda}^T \boldsymbol{\beta}]^2 = \text{Var}(g(\mathbf{Y})) + \text{Bias}^2(g(\mathbf{Y}))$$

where  $\text{Bias} = E[g(\mathbf{Y})] - \boldsymbol{\lambda}^T \boldsymbol{\beta}$

- ▶ Bias vs Variance tradeoff
- ▶ Can have smaller MSE if we allow some Bias!



# Bayes

- ▶ Next Class Bayes Theorem & Conjugate Normal-Gamma Prior/Posterior distributions
- ▶ Read Chapter 2 in Christensen or Wakefield 5.7
- ▶ Review Multivariate Normal and Gamma distributions