

# Multivariate Normal Theory

STA721 Linear Models Duke University

Merlise Clyde

September 5, 2017

# Outline

- ▶ Multivariate Normal Distribution
- ▶ Linear Transformations
- ▶ Distribution of estimates under normality

## Properties of MLE's Recap

- ▶  $\hat{\mathbf{Y}} = \hat{\boldsymbol{\mu}} = \mathbf{P}_{\mathbf{X}}\mathbf{Y}$  is an unbiased estimate of  $\boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$

# Properties of MLE's Recap

- ▶  $\hat{\mathbf{Y}} = \hat{\boldsymbol{\mu}} = \mathbf{P}_{\mathbf{X}}\mathbf{Y}$  is an unbiased estimate of  $\boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$
- ▶  $E[\mathbf{e}] = \mathbf{0}$  if  $\boldsymbol{\mu} \in C(\mathbf{X})$

## Properties of MLE's Recap

- ▶  $\hat{\mathbf{Y}} = \hat{\boldsymbol{\mu}} = \mathbf{P}_{\mathbf{X}}\mathbf{Y}$  is an unbiased estimate of  $\boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$
- ▶  $E[\mathbf{e}] = \mathbf{0}$  if  $\boldsymbol{\mu} \in C(\mathbf{X})$

$$E[\mathbf{e}] = E[(\mathbf{I} - \mathbf{P}_{\mathbf{X}})\mathbf{Y}]$$

# Properties of MLE's Recap

- ▶  $\hat{\mathbf{Y}} = \hat{\boldsymbol{\mu}} = \mathbf{P}_\mathbf{X}\mathbf{Y}$  is an unbiased estimate of  $\boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$
- ▶  $E[\mathbf{e}] = \mathbf{0}$  if  $\boldsymbol{\mu} \in C(\mathbf{X})$

$$E[\mathbf{e}] = E[(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}]$$

- ▶ MLE of  $\sigma^2$ :

$$\hat{\sigma}^2 = \frac{\mathbf{e}^T \mathbf{e}}{n} = \frac{\mathbf{Y}^T (\mathbf{I} - \mathbf{P}_\mathbf{X}) \mathbf{Y}}{n}$$

# Properties of MLE's Recap

- ▶  $\hat{\mathbf{Y}} = \hat{\boldsymbol{\mu}} = \mathbf{P}_\mathbf{X}\mathbf{Y}$  is an unbiased estimate of  $\boldsymbol{\mu} = \mathbf{X}\boldsymbol{\beta}$
- ▶  $E[\mathbf{e}] = \mathbf{0}$  if  $\boldsymbol{\mu} \in C(\mathbf{X})$

$$E[\mathbf{e}] = E[(\mathbf{I} - \mathbf{P}_\mathbf{X})\mathbf{Y}]$$

- ▶ MLE of  $\sigma^2$ :

$$\hat{\sigma}^2 = \frac{\mathbf{e}^T \mathbf{e}}{n} = \frac{\mathbf{Y}^T (\mathbf{I} - \mathbf{P}_\mathbf{X}) \mathbf{Y}}{n}$$

Is not an unbiased estimate of  $\sigma^2$ , but

$$\hat{\sigma}^2 \equiv \frac{\mathbf{e}^T \mathbf{e}}{n - p} = \frac{\mathbf{Y}^T (\mathbf{I} - \mathbf{P}_\mathbf{X}) \mathbf{Y}}{n - p}$$

where  $p$  equals the rank of  $\mathbf{X}$  is an unbiased estimate.

# Sampling Distributions

- ▶ Distribution of  $\hat{\beta}$
- ▶ Distribution of  $\mathbf{P}_X \mathbf{Y}$
- ▶ Distribution of  $\mathbf{e}$



# Univariate Normal

## Definition

We say that  $Z$  has a standard Normal distribution

$$Z \sim N(0, 1)$$

with mean 0 and variance 1 if it has density

# Univariate Normal

## Definition

We say that  $Z$  has a standard Normal distribution

$$Z \sim N(0, 1)$$

with mean 0 and variance 1 if it has density

$$f_Z(z) = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}z^2}$$

# Univariate Normal

## Definition

We say that  $Z$  has a standard Normal distribution

$$Z \sim N(0, 1)$$

with mean 0 and variance 1 if it has density

$$f_Z(z) = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}z^2}$$

If  $Y = \mu + \sigma Z$  then  $Y \sim N(\mu, \sigma^2)$  with mean  $\mu$  and variance  $\sigma^2$

# Univariate Normal

## Definition

We say that  $Z$  has a standard Normal distribution

$$Z \sim N(0, 1)$$

with mean 0 and variance 1 if it has density

$$f_Z(z) = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}z^2}$$

If  $Y = \mu + \sigma Z$  then  $Y \sim N(\mu, \sigma^2)$  with mean  $\mu$  and variance  $\sigma^2$

$$f_Y(y) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{1}{2}\left(\frac{y-\mu}{\sigma}\right)^2}$$

# Standard Multivariate Normal

Let  $z_i \stackrel{\text{iid}}{\sim} \mathcal{N}(0, 1)$  for  $i = 1, \dots, d$  and define

$$\mathbf{z} \equiv \begin{bmatrix} z_1 \\ z_2 \\ \vdots \\ z_d \end{bmatrix}$$

# Standard Multivariate Normal

Let  $z_i \stackrel{\text{iid}}{\sim} \mathcal{N}(0, 1)$  for  $i = 1, \dots, d$  and define

$$\mathbf{Z} \equiv \begin{bmatrix} z_1 \\ z_2 \\ \vdots \\ z_d \end{bmatrix}$$

- Density of  $Z$ :

# Standard Multivariate Normal

Let  $z_i \stackrel{\text{iid}}{\sim} \mathcal{N}(0, 1)$  for  $i = 1, \dots, d$  and define

$$\mathbf{z} \equiv \begin{bmatrix} z_1 \\ z_2 \\ \vdots \\ z_d \end{bmatrix}$$

► Density of  $Z$ :

$$f_{\mathbf{z}}(\mathbf{z}) = \prod_{j=1}^d \frac{1}{\sqrt{2\pi}} e^{-z_i^2/2}$$

# Standard Multivariate Normal

Let  $z_i \stackrel{\text{iid}}{\sim} \mathcal{N}(0, 1)$  for  $i = 1, \dots, d$  and define

$$\mathbf{z} \equiv \begin{bmatrix} z_1 \\ z_2 \\ \vdots \\ z_d \end{bmatrix}$$

► Density of  $\mathbf{Z}$ :

$$\begin{aligned} f_{\mathbf{Z}}(\mathbf{z}) &= \prod_{j=1}^d \frac{1}{\sqrt{2\pi}} e^{-z_j^2/2} \\ &= (2\pi)^{-d/2} e^{-\frac{1}{2}(\mathbf{z}^T \mathbf{z})} \end{aligned}$$



# Standard Multivariate Normal

Let  $z_i \stackrel{\text{iid}}{\sim} \mathcal{N}(0, 1)$  for  $i = 1, \dots, d$  and define

$$\mathbf{Z} \equiv \begin{bmatrix} z_1 \\ z_2 \\ \vdots \\ z_d \end{bmatrix}$$

- Density of  $\mathbf{Z}$ :

$$\begin{aligned} f_{\mathbf{Z}}(\mathbf{z}) &= \prod_{j=1}^d \frac{1}{\sqrt{2\pi}} e^{-z_j^2/2} \\ &= (2\pi)^{-d/2} e^{-\frac{1}{2}(\mathbf{Z}^T \mathbf{Z})} \end{aligned}$$

- $E[\mathbf{Z}] = \mathbf{0}$  and  $\text{Cov}[\mathbf{Z}] = \mathbf{I}_d$

# Standard Multivariate Normal

Let  $z_i \stackrel{\text{iid}}{\sim} \mathcal{N}(0, 1)$  for  $i = 1, \dots, d$  and define

$$\mathbf{Z} \equiv \begin{bmatrix} z_1 \\ z_2 \\ \vdots \\ z_d \end{bmatrix}$$

- Density of  $\mathbf{Z}$ :

$$\begin{aligned} f_{\mathbf{Z}}(\mathbf{z}) &= \prod_{j=1}^d \frac{1}{\sqrt{2\pi}} e^{-z_j^2/2} \\ &= (2\pi)^{-d/2} e^{-\frac{1}{2}(\mathbf{Z}^T \mathbf{Z})} \end{aligned}$$

- $\mathbb{E}[\mathbf{Z}] = \mathbf{0}$  and  $\text{Cov}[\mathbf{Z}] = \mathbf{I}_d$
- $\mathbf{Z} \sim \mathcal{N}(\mathbf{0}_d, \mathbf{I}_d)$

# Standard Multivariate Normal

Let  $z_i \stackrel{\text{iid}}{\sim} \mathcal{N}(0, 1)$  for  $i = 1, \dots, d$  and define

$$\mathbf{Z} \equiv \begin{bmatrix} z_1 \\ z_2 \\ \vdots \\ z_d \end{bmatrix}$$

- Density of  $\mathbf{Z}$ :

$$\begin{aligned} f_{\mathbf{Z}}(\mathbf{z}) &= \prod_{j=1}^d \frac{1}{\sqrt{2\pi}} e^{-z_j^2/2} \\ &= (2\pi)^{-d/2} e^{-\frac{1}{2}(\mathbf{Z}^T \mathbf{Z})} \end{aligned}$$

- $\mathbb{E}[\mathbf{Z}] = \mathbf{0}$  and  $\text{Cov}[\mathbf{Z}] = \mathbf{I}_d$
- $\mathbf{Z} \sim \mathcal{N}(\mathbf{0}_d, \mathbf{I}_d)$

# Multivariate Normal

For a  $d$  dimensional multivariate normal random vector, we write  
 $\mathbf{Y} \sim N_d(\boldsymbol{\mu}, \boldsymbol{\Sigma})$

# Multivariate Normal

For a  $d$  dimensional multivariate normal random vector, we write

$$\mathbf{Y} \sim N_d(\boldsymbol{\mu}, \boldsymbol{\Sigma})$$

- ▶  $E[\mathbf{Y}] = \boldsymbol{\mu}$ :  $d$  dimensional vector with means  $E[Y_j]$

# Multivariate Normal

For a  $d$  dimensional multivariate normal random vector, we write  $\mathbf{Y} \sim N_d(\boldsymbol{\mu}, \boldsymbol{\Sigma})$

- ▶  $E[\mathbf{Y}] = \boldsymbol{\mu}$ :  $d$  dimensional vector with means  $E[Y_j]$
- ▶  $\text{Cov}[\mathbf{Y}] = \boldsymbol{\Sigma}$ :  $d \times d$  matrix with diagonal elements that are the variances of  $Y_j$  and off diagonal elements that are the covariances  $E[(Y_j - \mu_j)(Y_k - \mu_k)]$

# Multivariate Normal

For a  $d$  dimensional multivariate normal random vector, we write  $\mathbf{Y} \sim N_d(\boldsymbol{\mu}, \boldsymbol{\Sigma})$

- ▶  $E[\mathbf{Y}] = \boldsymbol{\mu}$ :  $d$  dimensional vector with means  $E[Y_j]$
- ▶  $\text{Cov}[\mathbf{Y}] = \boldsymbol{\Sigma}$ :  $d \times d$  matrix with diagonal elements that are the variances of  $Y_j$  and off diagonal elements that are the covariances  $E[(Y_j - \mu_j)(Y_k - \mu_k)]$

## Density

If  $\boldsymbol{\Sigma}$  is positive definite ( $\mathbf{x}'\boldsymbol{\Sigma}\mathbf{x} > 0$  for any  $\mathbf{x} \neq 0$  in  $\mathbb{R}^d$ ) then  $\mathbf{Y}$  has a density <sup>1</sup>

---

<sup>1</sup>with respect to Lebesgue measure on  $\mathbb{R}^d$

# Multivariate Normal

For a  $d$  dimensional multivariate normal random vector, we write  $\mathbf{Y} \sim N_d(\boldsymbol{\mu}, \boldsymbol{\Sigma})$

- ▶  $E[\mathbf{Y}] = \boldsymbol{\mu}$ :  $d$  dimensional vector with means  $E[Y_j]$
- ▶  $\text{Cov}[\mathbf{Y}] = \boldsymbol{\Sigma}$ :  $d \times d$  matrix with diagonal elements that are the variances of  $Y_j$  and off diagonal elements that are the covariances  $E[(Y_j - \mu_j)(Y_k - \mu_k)]$



# Spectral Theorem

## Theorem

*If  $\mathbf{A}$  ( $n \times n$ ) is a symmetric real matrix then there exists a  $\mathbf{U}$  ( $n \times n$ ) such that  $\mathbf{U}^T \mathbf{U} = \mathbf{U} \mathbf{U}^T = \mathbf{I}_n$  and a diagonal matrix  $\mathbf{\Lambda}$  with elements  $\lambda_i$  such that  $\mathbf{A} = \mathbf{U} \mathbf{\Lambda} \mathbf{U}^T$*

# Spectral Theorem

## Theorem

If  $\mathbf{A}$  ( $n \times n$ ) is a symmetric real matrix then there exists a  $\mathbf{U}$  ( $n \times n$ ) such that  $\mathbf{U}^T \mathbf{U} = \mathbf{U} \mathbf{U}^T = \mathbf{I}_n$  and a diagonal matrix  $\mathbf{\Lambda}$  with elements  $\lambda_i$  such that  $\mathbf{A} = \mathbf{U} \mathbf{\Lambda} \mathbf{U}^T$

- ▶  $\mathbf{U}$  is an orthogonal matrix;  $\mathbf{U}^{-1} = \mathbf{U}^T$

# Spectral Theorem

## Theorem

If  $\mathbf{A}$  ( $n \times n$ ) is a symmetric real matrix then there exists a  $\mathbf{U}$  ( $n \times n$ ) such that  $\mathbf{U}^T \mathbf{U} = \mathbf{U} \mathbf{U}^T = \mathbf{I}_n$  and a diagonal matrix  $\mathbf{\Lambda}$  with elements  $\lambda_i$  such that  $\mathbf{A} = \mathbf{U} \mathbf{\Lambda} \mathbf{U}^T$

- ▶  $\mathbf{U}$  is an orthogonal matrix;  $\mathbf{U}^{-1} = \mathbf{U}^T$
- ▶ The columns of  $\mathbf{U}$  form an Orthonormal Basis for  $\mathbb{R}^n$

# Spectral Theorem

## Theorem

If  $\mathbf{A}$  ( $n \times n$ ) is a symmetric real matrix then there exists a  $\mathbf{U}$  ( $n \times n$ ) such that  $\mathbf{U}^T \mathbf{U} = \mathbf{U} \mathbf{U}^T = \mathbf{I}_n$  and a diagonal matrix  $\mathbf{\Lambda}$  with elements  $\lambda_i$  such that  $\mathbf{A} = \mathbf{U} \mathbf{\Lambda} \mathbf{U}^T$

- ▶  $\mathbf{U}$  is an orthogonal matrix;  $\mathbf{U}^{-1} = \mathbf{U}^T$
- ▶ The columns of  $\mathbf{U}$  form an Orthonormal Basis for  $\mathbb{R}^n$
- ▶ rank of  $\mathbf{A}$  equals the number of non-zero eigenvalues  $\lambda_i$

# Spectral Theorem

## Theorem

If  $\mathbf{A}$  ( $n \times n$ ) is a symmetric real matrix then there exists a  $\mathbf{U}$  ( $n \times n$ ) such that  $\mathbf{U}^T \mathbf{U} = \mathbf{U} \mathbf{U}^T = \mathbf{I}_n$  and a diagonal matrix  $\mathbf{\Lambda}$  with elements  $\lambda_i$  such that  $\mathbf{A} = \mathbf{U} \mathbf{\Lambda} \mathbf{U}^T$

- ▶  $\mathbf{U}$  is an orthogonal matrix;  $\mathbf{U}^{-1} = \mathbf{U}^T$
- ▶ The columns of  $\mathbf{U}$  form an Orthonormal Basis for  $\mathbb{R}^n$
- ▶ rank of  $\mathbf{A}$  equals the number of non-zero eigenvalues  $\lambda_i$
- ▶ Columns of  $\mathbf{U}$  associated with non-zero eigenvalues form an ONB for  $C(\mathbf{A})$  (eigenvectors of  $\mathbf{A}$ )

# Spectral Theorem

## Theorem

If  $\mathbf{A}$  ( $n \times n$ ) is a symmetric real matrix then there exists a  $\mathbf{U}$  ( $n \times n$ ) such that  $\mathbf{U}^T \mathbf{U} = \mathbf{U} \mathbf{U}^T = \mathbf{I}_n$  and a diagonal matrix  $\mathbf{\Lambda}$  with elements  $\lambda_i$  such that  $\mathbf{A} = \mathbf{U} \mathbf{\Lambda} \mathbf{U}^T$

- ▶  $\mathbf{U}$  is an orthogonal matrix;  $\mathbf{U}^{-1} = \mathbf{U}^T$
- ▶ The columns of  $\mathbf{U}$  form an Orthonormal Basis for  $\mathbb{R}^n$
- ▶ rank of  $\mathbf{A}$  equals the number of non-zero eigenvalues  $\lambda_i$
- ▶ Columns of  $\mathbf{U}$  associated with non-zero eigenvalues form an ONB for  $C(\mathbf{A})$  (eigenvectors of  $\mathbf{A}$ )
- ▶  $\mathbf{A}^p = \mathbf{U} \mathbf{\Lambda}^p \mathbf{U}^T$  (matrix powers)

# Spectral Theorem

## Theorem

If  $\mathbf{A}$  ( $n \times n$ ) is a symmetric real matrix then there exists a  $\mathbf{U}$  ( $n \times n$ ) such that  $\mathbf{U}^T \mathbf{U} = \mathbf{U} \mathbf{U}^T = \mathbf{I}_n$  and a diagonal matrix  $\mathbf{\Lambda}$  with elements  $\lambda_i$  such that  $\mathbf{A} = \mathbf{U} \mathbf{\Lambda} \mathbf{U}^T$

- ▶  $\mathbf{U}$  is an orthogonal matrix;  $\mathbf{U}^{-1} = \mathbf{U}^T$
- ▶ The columns of  $\mathbf{U}$  form an Orthonormal Basis for  $\mathbb{R}^n$
- ▶ rank of  $\mathbf{A}$  equals the number of non-zero eigenvalues  $\lambda_i$
- ▶ Columns of  $\mathbf{U}$  associated with non-zero eigenvalues form an ONB for  $C(\mathbf{A})$  (eigenvectors of  $\mathbf{A}$ )
- ▶  $\mathbf{A}^p = \mathbf{U} \mathbf{\Lambda}^p \mathbf{U}^T$  (matrix powers)
- ▶ a (symmetric) square root of  $\mathbf{A} > 0$  is  $\mathbf{U} \mathbf{\Lambda}^{1/2} \mathbf{U}^T$

# Spectral Theorem

## Theorem

If  $\mathbf{A}$  ( $n \times n$ ) is a symmetric real matrix then there exists a  $\mathbf{U}$  ( $n \times n$ ) such that  $\mathbf{U}^T \mathbf{U} = \mathbf{U} \mathbf{U}^T = \mathbf{I}_n$  and a diagonal matrix  $\mathbf{\Lambda}$  with elements  $\lambda_i$  such that  $\mathbf{A} = \mathbf{U} \mathbf{\Lambda} \mathbf{U}^T$

- ▶  $\mathbf{U}$  is an orthogonal matrix;  $\mathbf{U}^{-1} = \mathbf{U}^T$
- ▶ The columns of  $\mathbf{U}$  form an Orthonormal Basis for  $\mathbb{R}^n$
- ▶ rank of  $\mathbf{A}$  equals the number of non-zero eigenvalues  $\lambda_i$
- ▶ Columns of  $\mathbf{U}$  associated with non-zero eigenvalues form an ONB for  $C(\mathbf{A})$  (eigenvectors of  $\mathbf{A}$ )
- ▶  $\mathbf{A}^p = \mathbf{U} \mathbf{\Lambda}^p \mathbf{U}^T$  (matrix powers)
- ▶ a (symmetric) square root of  $\mathbf{A} > 0$  is  $\mathbf{U} \mathbf{\Lambda}^{1/2} \mathbf{U}^T$



# Multivariate Normal Density

- Density of  $Z \sim N(\mathbf{0}, \mathbf{I}_d)$ :

# Multivariate Normal Density

- Density of  $Z \sim N(\mathbf{0}, \mathbf{I}_d)$ :

$$f_{\mathbf{Z}}(\mathbf{z}) = \prod_{j=1}^d \frac{1}{\sqrt{2\pi}} e^{-z_j^2/2}$$

# Multivariate Normal Density

- Density of  $Z \sim N(\mathbf{0}, \mathbf{I}_d)$ :

$$\begin{aligned} f_{\mathbf{Z}}(\mathbf{z}) &= \prod_{j=1}^d \frac{1}{\sqrt{2\pi}} e^{-z_j^2/2} \\ &= (2\pi)^{-d/2} e^{-\frac{1}{2}(\mathbf{Z}^T \mathbf{Z})} \end{aligned}$$

# Multivariate Normal Density

- Density of  $Z \sim N(\mathbf{0}, \mathbf{I}_d)$ :

$$\begin{aligned} f_{\mathbf{Z}}(\mathbf{z}) &= \prod_{j=1}^d \frac{1}{\sqrt{2\pi}} e^{-z_j^2/2} \\ &= (2\pi)^{-d/2} e^{-\frac{1}{2}(\mathbf{Z}^T \mathbf{Z})} \end{aligned}$$

- Write  $\mathbf{Y} = \boldsymbol{\mu} + \mathbf{AZ}$

# Multivariate Normal Density

- Density of  $\mathbf{Z} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_d)$ :

$$\begin{aligned} f_{\mathbf{Z}}(\mathbf{z}) &= \prod_{j=1}^d \frac{1}{\sqrt{2\pi}} e^{-z_j^2/2} \\ &= (2\pi)^{-d/2} e^{-\frac{1}{2}(\mathbf{Z}^T \mathbf{Z})} \end{aligned}$$

- Write  $\mathbf{Y} = \boldsymbol{\mu} + \mathbf{AZ}$
- Solve for  $\mathbf{Z} = g(\mathbf{Y})$

# Multivariate Normal Density

- Density of  $\mathbf{Z} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_d)$ :

$$\begin{aligned} f_{\mathbf{Z}}(\mathbf{z}) &= \prod_{j=1}^d \frac{1}{\sqrt{2\pi}} e^{-z_j^2/2} \\ &= (2\pi)^{-d/2} e^{-\frac{1}{2}(\mathbf{Z}^T \mathbf{Z})} \end{aligned}$$

- Write  $\mathbf{Y} = \boldsymbol{\mu} + \mathbf{A}\mathbf{Z}$
- Solve for  $\mathbf{Z} = \mathbf{g}(\mathbf{Y})$
- Jacobian of the transformation  $J(\mathbf{Z} \rightarrow \mathbf{Y}) = \left| \frac{\partial \mathbf{g}}{\partial \mathbf{Y}} \right|$

# Multivariate Normal Density

- Density of  $Z \sim N(\mathbf{0}, \mathbf{I}_d)$ :

$$\begin{aligned}f_{\mathbf{Z}}(\mathbf{z}) &= \prod_{j=1}^d \frac{1}{\sqrt{2\pi}} e^{-z_j^2/2} \\&= (2\pi)^{-d/2} e^{-\frac{1}{2}(\mathbf{Z}^T \mathbf{Z})}\end{aligned}$$

- Write  $\mathbf{Y} = \boldsymbol{\mu} + \mathbf{A}\mathbf{Z}$
- Solve for  $\mathbf{Z} = g(\mathbf{Y})$
- Jacobian of the transformation  $J(\mathbf{Z} \rightarrow \mathbf{Y}) = \left| \frac{\partial g}{\partial \mathbf{Y}} \right|$
- substitute  $g(\mathbf{Y})$  for  $\mathbf{Z}$  in density and multiply by Jacobian

$$f_{\mathbf{Y}}(\mathbf{y}) = f_{\mathbf{Z}}(\mathbf{z})J(\mathbf{Z} \rightarrow \mathbf{Y})$$

# Multivariate Normal Density

$$\mathbf{Y} = \boldsymbol{\mu} + \mathbf{AZ} \quad \text{for } \mathbf{Z} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_d) \quad (1)$$

Proof.



# Multivariate Normal Density

$$\mathbf{Y} = \boldsymbol{\mu} + \mathbf{AZ} \quad \text{for } \mathbf{Z} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_d) \quad (1)$$

Proof.

- ▶ since  $\boldsymbol{\Sigma} > 0$ ,  $\exists$  by the spectral theorem an  $\mathbf{A}$  ( $d \times d$ ) such that  $\mathbf{A} > 0$  and  $\mathbf{AA}^T = \boldsymbol{\Sigma}$

# Multivariate Normal Density

$$\mathbf{Y} = \boldsymbol{\mu} + \mathbf{AZ} \quad \text{for } \mathbf{Z} \sim N(\mathbf{0}, \mathbf{I}_d) \quad (1)$$

Proof.

- ▶ since  $\boldsymbol{\Sigma} > 0$ ,  $\exists$  by the spectral theorem an  $\mathbf{A}$  ( $d \times d$ ) such that  $\mathbf{A} > 0$  and  $\mathbf{AA}^T = \boldsymbol{\Sigma}$
- ▶  $\mathbf{A} > 0 \Rightarrow \mathbf{A}^{-1}$  exists

# Multivariate Normal Density

$$\mathbf{Y} = \boldsymbol{\mu} + \mathbf{A}\mathbf{Z} \quad \text{for } \mathbf{Z} \sim N(\mathbf{0}, \mathbf{I}_d) \quad (1)$$

Proof.

- ▶ since  $\boldsymbol{\Sigma} > 0$ ,  $\exists$  by the spectral theorem an  $\mathbf{A}$  ( $d \times d$ ) such that  $\mathbf{A} > 0$  and  $\mathbf{A}\mathbf{A}^T = \boldsymbol{\Sigma}$
- ▶  $\mathbf{A} > 0 \Rightarrow \mathbf{A}^{-1}$  exists
- ▶ Multiply both sides (1) by  $\mathbf{A}^{-1}$ :

$$\mathbf{A}^{-1}\mathbf{Y} = \mathbf{A}^{-1}\boldsymbol{\mu} + \mathbf{A}^{-1}\mathbf{A}\mathbf{Z}$$

# Multivariate Normal Density

$$\mathbf{Y} = \boldsymbol{\mu} + \mathbf{AZ} \quad \text{for } \mathbf{Z} \sim N(\mathbf{0}, \mathbf{I}_d) \quad (1)$$

Proof.

- ▶ since  $\boldsymbol{\Sigma} > 0$ ,  $\exists$  by the spectral theorem an  $\mathbf{A}$  ( $d \times d$ ) such that  $\mathbf{A} > 0$  and  $\mathbf{AA}^T = \boldsymbol{\Sigma}$
- ▶  $\mathbf{A} > 0 \Rightarrow \mathbf{A}^{-1}$  exists
- ▶ Multiply both sides (1) by  $\mathbf{A}^{-1}$ :

$$\mathbf{A}^{-1}\mathbf{Y} = \mathbf{A}^{-1}\boldsymbol{\mu} + \mathbf{A}^{-1}\mathbf{AZ}$$

- ▶ Rearrange  $\mathbf{A}^{-1}(\mathbf{Y} - \boldsymbol{\mu}) = \mathbf{Z}$

# Multivariate Normal Density

$$\mathbf{Y} = \boldsymbol{\mu} + \mathbf{A}\mathbf{Z} \quad \text{for } \mathbf{Z} \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_d) \quad (1)$$

Proof.

- ▶ since  $\boldsymbol{\Sigma} > 0$ ,  $\exists$  by the spectral theorem an  $\mathbf{A}$  ( $d \times d$ ) such that  $\mathbf{A} > 0$  and  $\mathbf{A}\mathbf{A}^T = \boldsymbol{\Sigma}$
- ▶  $\mathbf{A} > 0 \Rightarrow \mathbf{A}^{-1}$  exists
- ▶ Multiply both sides (1) by  $\mathbf{A}^{-1}$ :

$$\mathbf{A}^{-1}\mathbf{Y} = \mathbf{A}^{-1}\boldsymbol{\mu} + \mathbf{A}^{-1}\mathbf{A}\mathbf{Z}$$

- ▶ Rearrange  $\mathbf{A}^{-1}(\mathbf{Y} - \boldsymbol{\mu}) = \mathbf{Z}$
- ▶ Jacobian of transformation  $d\mathbf{Z} = |\mathbf{A}^{-1}|d\mathbf{Y}$

# Multivariate Normal Density

$$\mathbf{Y} = \boldsymbol{\mu} + \mathbf{A}\mathbf{Z} \quad \text{for } \mathbf{Z} \sim N(\mathbf{0}, \mathbf{I}_d) \quad (1)$$

Proof.

- ▶ since  $\boldsymbol{\Sigma} > 0$ ,  $\exists$  by the spectral theorem an  $\mathbf{A}$  ( $d \times d$ ) such that  $\mathbf{A} > 0$  and  $\mathbf{A}\mathbf{A}^T = \boldsymbol{\Sigma}$
- ▶  $\mathbf{A} > 0 \Rightarrow \mathbf{A}^{-1}$  exists
- ▶ Multiply both sides (1) by  $\mathbf{A}^{-1}$ :

$$\mathbf{A}^{-1}\mathbf{Y} = \mathbf{A}^{-1}\boldsymbol{\mu} + \mathbf{A}^{-1}\mathbf{A}\mathbf{Z}$$

- ▶ Rearrange  $\mathbf{A}^{-1}(\mathbf{Y} - \boldsymbol{\mu}) = \mathbf{Z}$
- ▶ Jacobian of transformation  $d\mathbf{Z} = |\mathbf{A}^{-1}|d\mathbf{Y}$
- ▶ Substitute and simplify algebra

$$f(\mathbf{Y}) = (2\pi)^{-d/2} |\boldsymbol{\Sigma}|^{-1/2} \exp\left(-\frac{1}{2}(\mathbf{Y} - \boldsymbol{\mu})^T \boldsymbol{\Sigma}^{-1}(\mathbf{Y} - \boldsymbol{\mu})\right)$$

## Singular Case

$$\mathbf{Y} = \boldsymbol{\mu} + \mathbf{AZ} \text{ with } \mathbf{Z} \in \mathbb{R}^d \text{ and } \mathbf{A} \text{ is } n \times d$$

# Singular Case

$\mathbf{Y} = \boldsymbol{\mu} + \mathbf{AZ}$  with  $\mathbf{Z} \in \mathbb{R}^d$  and  $\mathbf{A}$  is  $n \times d$

►  $E[\mathbf{Y}] = \boldsymbol{\mu}$



# Singular Case

$\mathbf{Y} = \boldsymbol{\mu} + \mathbf{AZ}$  with  $\mathbf{Z} \in \mathbb{R}^d$  and  $\mathbf{A}$  is  $n \times d$

- ▶  $E[\mathbf{Y}] = \boldsymbol{\mu}$
- ▶  $\text{Cov}(\mathbf{Y}) = \mathbf{AA}^T \geq 0$

## Singular Case

$\mathbf{Y} = \boldsymbol{\mu} + \mathbf{A}\mathbf{Z}$  with  $\mathbf{Z} \in \mathbb{R}^d$  and  $\mathbf{A}$  is  $n \times d$

- ▶  $E[\mathbf{Y}] = \boldsymbol{\mu}$
- ▶  $\text{Cov}(\mathbf{Y}) = \mathbf{A}\mathbf{A}^T \geq 0$
- ▶  $\mathbf{Y} \sim N(\boldsymbol{\mu}, \boldsymbol{\Sigma})$  where  $\boldsymbol{\Sigma} = \mathbf{A}\mathbf{A}^T$

If  $\boldsymbol{\Sigma}$  is singular then there is no density (on  $\mathbb{R}^n$ ), but claim that  $\mathbf{Y}$  still has a multivariate normal distribution!

# Singular Case

$\mathbf{Y} = \boldsymbol{\mu} + \mathbf{AZ}$  with  $\mathbf{Z} \in \mathbb{R}^d$  and  $\mathbf{A}$  is  $n \times d$

- ▶  $E[\mathbf{Y}] = \boldsymbol{\mu}$
- ▶  $\text{Cov}(\mathbf{Y}) = \mathbf{AA}^T \geq 0$
- ▶  $\mathbf{Y} \sim N(\boldsymbol{\mu}, \boldsymbol{\Sigma})$  where  $\boldsymbol{\Sigma} = \mathbf{AA}^T$

If  $\boldsymbol{\Sigma}$  is singular then there is no density (on  $\mathbb{R}^n$ ), but claim that  $\mathbf{Y}$  still has a multivariate normal distribution!

## Definition

$\mathbf{Y} \in \mathbb{R}^n$  has a multivariate normal distribution  $N(\boldsymbol{\mu}, \boldsymbol{\Sigma})$  if for any  $\mathbf{v} \in \mathbb{R}^n$   $\mathbf{v}^T \mathbf{Y}$  has a normal distribution with mean  $\mathbf{v}^T \boldsymbol{\mu}$  and variance  $\mathbf{v}^T \boldsymbol{\Sigma} \mathbf{v}$

# Singular Case

$\mathbf{Y} = \boldsymbol{\mu} + \mathbf{AZ}$  with  $\mathbf{Z} \in \mathbb{R}^d$  and  $\mathbf{A}$  is  $n \times d$

- ▶  $E[\mathbf{Y}] = \boldsymbol{\mu}$
- ▶  $\text{Cov}(\mathbf{Y}) = \mathbf{AA}^T \geq 0$
- ▶  $\mathbf{Y} \sim N(\boldsymbol{\mu}, \boldsymbol{\Sigma})$  where  $\boldsymbol{\Sigma} = \mathbf{AA}^T$

If  $\boldsymbol{\Sigma}$  is singular then there is no density (on  $\mathbb{R}^n$ ), but claim that  $\mathbf{Y}$  still has a multivariate normal distribution!

## Definition

$\mathbf{Y} \in \mathbb{R}^n$  has a multivariate normal distribution  $N(\boldsymbol{\mu}, \boldsymbol{\Sigma})$  if for any  $\mathbf{v} \in \mathbb{R}^n$   $\mathbf{v}^T \mathbf{Y}$  has a normal distribution with mean  $\mathbf{v}^T \boldsymbol{\mu}$  and variance  $\mathbf{v}^T \boldsymbol{\Sigma} \mathbf{v}$

see linked videos using characteristic functions:

$$Y \sim N(\mu, \sigma^2) \Leftrightarrow \varphi_Y(t) \equiv E[e^{itY}] = e^{it\mu - t^2\sigma^2/2}$$

# Linear Transformations are Normal

If  $\mathbf{Y} \sim N_n(\boldsymbol{\mu}, \boldsymbol{\Sigma})$

# Linear Transformations are Normal

If  $\mathbf{Y} \sim N_n(\boldsymbol{\mu}, \boldsymbol{\Sigma})$  then for  $\mathbf{A} \ m \times n$

$$\mathbf{AY} \sim N_m(\mathbf{A}\boldsymbol{\mu}, \mathbf{A}\boldsymbol{\Sigma}\mathbf{A}^T)$$

# Linear Transformations are Normal

If  $\mathbf{Y} \sim N_n(\boldsymbol{\mu}, \boldsymbol{\Sigma})$  then for  $\mathbf{A} \ m \times n$

$$\mathbf{AY} \sim N_m(\mathbf{A}\boldsymbol{\mu}, \mathbf{A}\boldsymbol{\Sigma}\mathbf{A}^T)$$

$\mathbf{A}\boldsymbol{\Sigma}\mathbf{A}^T$  does not have to be positive definite!

# Linear Transformations are Normal

If  $\mathbf{Y} \sim N_n(\boldsymbol{\mu}, \boldsymbol{\Sigma})$  then for  $\mathbf{A} \ m \times n$

$$\mathbf{AY} \sim N_m(\mathbf{A}\boldsymbol{\mu}, \mathbf{A}\boldsymbol{\Sigma}\mathbf{A}^T)$$

$\mathbf{A}\boldsymbol{\Sigma}\mathbf{A}^T$  does not have to be positive definite!  
(Proof in book or linked video)



# Distribution of $\hat{\mathbf{Y}}$ and $\mathbf{e}$ (marginally)

# Equal in Distribution

Multiple ways to define the same normal:

# Equal in Distribution

Multiple ways to define the same normal:

- ▶  $\mathbf{Z}_1 \sim N(\mathbf{0}, \mathbf{I}_n)$ ,  $\mathbf{Z}_1 \in \mathbb{R}^n$  and take  $\mathbf{A} \ d \times n$

# Equal in Distribution

Multiple ways to define the same normal:

- ▶  $\mathbf{Z}_1 \sim N(\mathbf{0}, \mathbf{I}_n)$ ,  $\mathbf{Z}_1 \in \mathbb{R}^n$  and take  $\mathbf{A} \ d \times n$
- ▶  $\mathbf{Z}_2 \sim N(\mathbf{0}, \mathbf{I}_p)$ ,  $\mathbf{Z}_2 \in \mathbb{R}^p$  and take  $\mathbf{B} \ d \times p$

# Equal in Distribution

Multiple ways to define the same normal:

- ▶  $\mathbf{Z}_1 \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_n)$ ,  $\mathbf{Z}_1 \in \mathbb{R}^n$  and take  $\mathbf{A} \ d \times n$
- ▶  $\mathbf{Z}_2 \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$ ,  $\mathbf{Z}_2 \in \mathbb{R}^p$  and take  $\mathbf{B} \ d \times p$
- ▶ Define  $\mathbf{Y} = \boldsymbol{\mu} + \mathbf{AZ}_1$

# Equal in Distribution

Multiple ways to define the same normal:

- ▶  $\mathbf{Z}_1 \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_n)$ ,  $\mathbf{Z}_1 \in \mathbb{R}^n$  and take  $\mathbf{A} \ d \times n$
- ▶  $\mathbf{Z}_2 \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$ ,  $\mathbf{Z}_2 \in \mathbb{R}^p$  and take  $\mathbf{B} \ d \times p$
- ▶ Define  $\mathbf{Y} = \boldsymbol{\mu} + \mathbf{A}\mathbf{Z}_1$
- ▶ Define  $\mathbf{W} = \boldsymbol{\mu} + \mathbf{B}\mathbf{Z}_2$

# Equal in Distribution

Multiple ways to define the same normal:

- ▶  $\mathbf{Z}_1 \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_n)$ ,  $\mathbf{Z}_1 \in \mathbb{R}^n$  and take  $\mathbf{A} \ d \times n$
- ▶  $\mathbf{Z}_2 \sim \mathcal{N}(\mathbf{0}, \mathbf{I}_p)$ ,  $\mathbf{Z}_2 \in \mathbb{R}^p$  and take  $\mathbf{B} \ d \times p$
- ▶ Define  $\mathbf{Y} = \boldsymbol{\mu} + \mathbf{AZ}_1$
- ▶ Define  $\mathbf{W} = \boldsymbol{\mu} + \mathbf{BZ}_2$

## Theorem

If  $\mathbf{Y} = \boldsymbol{\mu} + \mathbf{AZ}_1$  and  $\mathbf{W} = \boldsymbol{\mu} + \mathbf{BZ}_2$  then  $\mathbf{Y} \stackrel{\text{D}}{=} \mathbf{W}$  if and only if  $\mathbf{AA}^T = \mathbf{BB}^T = \boldsymbol{\Sigma}$

# Equal in Distribution

Multiple ways to define the same normal:

- ▶  $\mathbf{Z}_1 \sim N(\mathbf{0}, \mathbf{I}_n)$ ,  $\mathbf{Z}_1 \in \mathbb{R}^n$  and take  $\mathbf{A} \ d \times n$
- ▶  $\mathbf{Z}_2 \sim N(\mathbf{0}, \mathbf{I}_p)$ ,  $\mathbf{Z}_2 \in \mathbb{R}^p$  and take  $\mathbf{B} \ d \times p$
- ▶ Define  $\mathbf{Y} = \boldsymbol{\mu} + \mathbf{AZ}_1$
- ▶ Define  $\mathbf{W} = \boldsymbol{\mu} + \mathbf{BZ}_2$

## Theorem

If  $\mathbf{Y} = \boldsymbol{\mu} + \mathbf{AZ}_1$  and  $\mathbf{W} = \boldsymbol{\mu} + \mathbf{BZ}_2$  then  $\mathbf{Y} \stackrel{D}{=} \mathbf{W}$  if and only if  $\mathbf{AA}^T = \mathbf{BB}^T = \boldsymbol{\Sigma}$

see linked video



# Zero Correlation and Independence

## Theorem

For a random vector  $\mathbf{Y} \sim N(\boldsymbol{\mu}, \boldsymbol{\Sigma})$  partitioned as

$$\mathbf{Y} = \begin{bmatrix} \mathbf{Y}_1 \\ \mathbf{Y}_2 \end{bmatrix} \sim N\left(\begin{bmatrix} \boldsymbol{\mu}_1 \\ \boldsymbol{\mu}_2 \end{bmatrix}, \begin{bmatrix} \boldsymbol{\Sigma}_{11} & \boldsymbol{\Sigma}_{12} \\ \boldsymbol{\Sigma}_{21} & \boldsymbol{\Sigma}_{22} \end{bmatrix}\right)$$

# Zero Correlation and Independence

## Theorem

For a random vector  $\mathbf{Y} \sim N(\boldsymbol{\mu}, \boldsymbol{\Sigma})$  partitioned as

$$\mathbf{Y} = \begin{bmatrix} \mathbf{Y}_1 \\ \mathbf{Y}_2 \end{bmatrix} \sim N\left(\begin{bmatrix} \boldsymbol{\mu}_1 \\ \boldsymbol{\mu}_2 \end{bmatrix}, \begin{bmatrix} \boldsymbol{\Sigma}_{11} & \boldsymbol{\Sigma}_{12} \\ \boldsymbol{\Sigma}_{21} & \boldsymbol{\Sigma}_{22} \end{bmatrix}\right)$$

then  $\text{Cov}(\mathbf{Y}_1, \mathbf{Y}_2) = \boldsymbol{\Sigma}_{12} = \boldsymbol{\Sigma}_{21}^T = \mathbf{0}$  if and only if  $\mathbf{Y}_1$  and  $\mathbf{Y}_2$  are independent.

# Independence Implies Zero Covariance

Proof.

$$\text{Cov}(\mathbf{Y}_1, \mathbf{Y}_2) = \mathbb{E}[(\mathbf{Y}_1 - \boldsymbol{\mu}_1)(\mathbf{Y}_2 - \boldsymbol{\mu}_2)^T]$$

# Independence Implies Zero Covariance

Proof.

$$\text{Cov}(\mathbf{Y}_1, \mathbf{Y}_2) = E[(\mathbf{Y}_1 - \boldsymbol{\mu}_1)(\mathbf{Y}_2 - \boldsymbol{\mu}_2)^T]$$

If  $\mathbf{Y}_1$  and  $\mathbf{Y}_2$  are independent

# Independence Implies Zero Covariance

Proof.

$$\text{Cov}(\mathbf{Y}_1, \mathbf{Y}_2) = E[(\mathbf{Y}_1 - \boldsymbol{\mu}_1)(\mathbf{Y}_2 - \boldsymbol{\mu}_2)^T]$$

If  $\mathbf{Y}_1$  and  $\mathbf{Y}_2$  are independent

$$E[(\mathbf{Y}_1 - \boldsymbol{\mu}_1)(\mathbf{Y}_2 - \boldsymbol{\mu}_2)^T] = E[(\mathbf{Y}_1 - \boldsymbol{\mu}_1)E(\mathbf{Y}_2 - \boldsymbol{\mu}_2)^T] = \mathbf{0}\mathbf{0}^T = \mathbf{0}$$

# Independence Implies Zero Covariance

Proof.

$$\text{Cov}(\mathbf{Y}_1, \mathbf{Y}_2) = E[(\mathbf{Y}_1 - \boldsymbol{\mu}_1)(\mathbf{Y}_2 - \boldsymbol{\mu}_2)^T]$$

If  $\mathbf{Y}_1$  and  $\mathbf{Y}_2$  are independent

$$E[(\mathbf{Y}_1 - \boldsymbol{\mu}_1)(\mathbf{Y}_2 - \boldsymbol{\mu}_2)^T] = E[(\mathbf{Y}_1 - \boldsymbol{\mu}_1)E(\mathbf{Y}_2 - \boldsymbol{\mu}_2)^T] = \mathbf{0}\mathbf{0}^T = \mathbf{0}$$

therefore  $\boldsymbol{\Sigma}_{12} = \mathbf{0}$



# Zero Covariance Implies Independence

Assume  $\Sigma_{12} = \mathbf{0}$

Proof

► Choose an

$$\mathbf{A} = \begin{bmatrix} \mathbf{A}_1 & \mathbf{0} \\ \mathbf{0} & \mathbf{A}_2 \end{bmatrix}$$

such that  $\mathbf{A}_1 \mathbf{A}_1^T = \Sigma_{11}$ ,  $\mathbf{A}_2 \mathbf{A}_2^T = \Sigma_{22}$

# Zero Covariance Implies Independence

Assume  $\Sigma_{12} = \mathbf{0}$

## Proof

- Choose an

$$\mathbf{A} = \begin{bmatrix} \mathbf{A}_1 & \mathbf{0} \\ \mathbf{0} & \mathbf{A}_2 \end{bmatrix}$$

such that  $\mathbf{A}_1 \mathbf{A}_1^T = \Sigma_{11}$ ,  $\mathbf{A}_2 \mathbf{A}_2^T = \Sigma_{22}$

- Partition

$$\mathbf{z} = \begin{bmatrix} \mathbf{z}_1 \\ \mathbf{z}_2 \end{bmatrix} \sim \mathcal{N} \left( \begin{bmatrix} \mathbf{0}_1 \\ \mathbf{0}_2 \end{bmatrix}, \begin{bmatrix} \mathbf{I}_1 & \mathbf{0} \\ \mathbf{0} & \mathbf{I}_2 \end{bmatrix} \right) \text{ and } \boldsymbol{\mu} = \begin{bmatrix} \mu_1 \\ \mu_2 \end{bmatrix}$$



# Zero Covariance Implies Independence

Assume  $\Sigma_{12} = \mathbf{0}$

## Proof

- ▶ Choose an

$$\mathbf{A} = \begin{bmatrix} \mathbf{A}_1 & \mathbf{0} \\ \mathbf{0} & \mathbf{A}_2 \end{bmatrix}$$

such that  $\mathbf{A}_1 \mathbf{A}_1^T = \Sigma_{11}$ ,  $\mathbf{A}_2 \mathbf{A}_2^T = \Sigma_{22}$

- ▶ Partition

$$\mathbf{Z} = \begin{bmatrix} \mathbf{Z}_1 \\ \mathbf{Z}_2 \end{bmatrix} \sim N \left( \begin{bmatrix} \mathbf{0}_1 \\ \mathbf{0}_2 \end{bmatrix}, \begin{bmatrix} \mathbf{I}_1 & \mathbf{0} \\ \mathbf{0} & \mathbf{I}_2 \end{bmatrix} \right) \text{ and } \boldsymbol{\mu} = \begin{bmatrix} \mu_1 \\ \mu_2 \end{bmatrix}$$

- ▶ then  $\mathbf{Y} \stackrel{D}{=} \mathbf{AZ} + \boldsymbol{\mu} \sim N(\boldsymbol{\mu}, \Sigma)$

# Continued

Proof.



$$\begin{bmatrix} \mathbf{Y}_1 \\ \mathbf{Y}_2 \end{bmatrix} \stackrel{\text{D}}{=} \begin{bmatrix} \mathbf{A}_1 \mathbf{Z}_1 + \boldsymbol{\mu}_1 \\ \mathbf{A}_2 \mathbf{Z}_2 + \boldsymbol{\mu}_2 \end{bmatrix}$$

## Continued

Proof.



$$\begin{bmatrix} \mathbf{Y}_1 \\ \mathbf{Y}_2 \end{bmatrix} \stackrel{D}{=} \begin{bmatrix} \mathbf{A}_1 \mathbf{Z}_1 + \boldsymbol{\mu}_1 \\ \mathbf{A}_2 \mathbf{Z}_2 + \boldsymbol{\mu}_2 \end{bmatrix}$$

- ▶ But  $\mathbf{Z}_1$  and  $\mathbf{Z}_2$  are independent

# Continued

Proof.



$$\begin{bmatrix} \mathbf{Y}_1 \\ \mathbf{Y}_2 \end{bmatrix} \stackrel{D}{=} \begin{bmatrix} \mathbf{A}_1 \mathbf{Z}_1 + \boldsymbol{\mu}_1 \\ \mathbf{A}_2 \mathbf{Z}_2 + \boldsymbol{\mu}_2 \end{bmatrix}$$

- ▶ But  $\mathbf{Z}_1$  and  $\mathbf{Z}_2$  are independent
- ▶ Functions of  $\mathbf{Z}_1$  and  $\mathbf{Z}_2$  are independent

# Continued

Proof.



$$\begin{bmatrix} \mathbf{Y}_1 \\ \mathbf{Y}_2 \end{bmatrix} \stackrel{D}{=} \begin{bmatrix} \mathbf{A}_1 \mathbf{Z}_1 + \boldsymbol{\mu}_1 \\ \mathbf{A}_2 \mathbf{Z}_2 + \boldsymbol{\mu}_2 \end{bmatrix}$$

- ▶ But  $\mathbf{Z}_1$  and  $\mathbf{Z}_2$  are independent
- ▶ Functions of  $\mathbf{Z}_1$  and  $\mathbf{Z}_2$  are independent
- ▶ Therefore  $\mathbf{Y}_1$  and  $\mathbf{Y}_2$  are independent

## Continued

Proof.



$$\begin{bmatrix} \mathbf{Y}_1 \\ \mathbf{Y}_2 \end{bmatrix} \stackrel{D}{=} \begin{bmatrix} \mathbf{A}_1 \mathbf{Z}_1 + \boldsymbol{\mu}_1 \\ \mathbf{A}_2 \mathbf{Z}_2 + \boldsymbol{\mu}_2 \end{bmatrix}$$

- ▶ But  $\mathbf{Z}_1$  and  $\mathbf{Z}_2$  are independent
- ▶ Functions of  $\mathbf{Z}_1$  and  $\mathbf{Z}_2$  are independent
- ▶ Therefore  $\mathbf{Y}_1$  and  $\mathbf{Y}_2$  are independent



For Multivariate Normal Zero Covariance implies independence

# Corollary

## Corollary

*If  $\mathbf{Y} \sim N(\boldsymbol{\mu}, \sigma^2 \mathbf{I}_n)$  and  $\mathbf{AB}^T = \mathbf{0}$  then  $\mathbf{AY}$  and  $\mathbf{BY}$  are independent.*

# Corollary

## Corollary

If  $\mathbf{Y} \sim N(\boldsymbol{\mu}, \sigma^2 \mathbf{I}_n)$  and  $\mathbf{AB}^T = \mathbf{0}$  then  $\mathbf{AY}$  and  $\mathbf{BY}$  are independent.

## Proof.



$$\begin{bmatrix} \mathbf{W}_1 \\ \mathbf{W}_2 \end{bmatrix} = \begin{bmatrix} \mathbf{A} \\ \mathbf{B} \end{bmatrix} \mathbf{Y} = \begin{bmatrix} \mathbf{AY} \\ \mathbf{BY} \end{bmatrix}$$



# Corollary

## Corollary

If  $\mathbf{Y} \sim N(\boldsymbol{\mu}, \sigma^2 \mathbf{I}_n)$  and  $\mathbf{AB}^T = \mathbf{0}$  then  $\mathbf{AY}$  and  $\mathbf{BY}$  are independent.

## Proof.



$$\begin{bmatrix} \mathbf{W}_1 \\ \mathbf{W}_2 \end{bmatrix} = \begin{bmatrix} \mathbf{A} \\ \mathbf{B} \end{bmatrix} \mathbf{Y} = \begin{bmatrix} \mathbf{AY} \\ \mathbf{BY} \end{bmatrix}$$

►  $\text{Cov}(\mathbf{W}_1, \mathbf{W}_2) = \text{Cov}(\mathbf{AY}, \mathbf{BY}) = \sigma^2 \mathbf{AB}^T$

# Corollary

## Corollary

If  $\mathbf{Y} \sim N(\boldsymbol{\mu}, \sigma^2 \mathbf{I}_n)$  and  $\mathbf{AB}^T = \mathbf{0}$  then  $\mathbf{AY}$  and  $\mathbf{BY}$  are independent.

## Proof.



$$\begin{bmatrix} \mathbf{W}_1 \\ \mathbf{W}_2 \end{bmatrix} = \begin{bmatrix} \mathbf{A} \\ \mathbf{B} \end{bmatrix} \mathbf{Y} = \begin{bmatrix} \mathbf{AY} \\ \mathbf{BY} \end{bmatrix}$$

- ▶  $\text{Cov}(\mathbf{W}_1, \mathbf{W}_2) = \text{Cov}(\mathbf{AY}, \mathbf{BY}) = \sigma^2 \mathbf{AB}^T$
- ▶  $\mathbf{AY}$  and  $\mathbf{BY}$  are independent if  $\mathbf{AB}^T = \mathbf{0}$



# Joint Distribution of $\hat{\mathbf{Y}}$ and $\mathbf{e}$

# More Distribution Theory

Distributions unconditional on  $\sigma^2$

- ▶  $\chi^2$  distributions ( $\hat{\sigma}^2$ )
- ▶  $t$  distribution ( $\hat{\mathbf{Y}}$ ,  $\mathbf{e}$ ,  $\hat{\beta}$ )