

Choice of Prior Distributions

STA721 Linear Models Duke University

Merlise Clyde

September 23, 2019

Bayesian Estimation

Model

$$\mathbf{Y} \sim \mathcal{N}(\mathbf{X}\boldsymbol{\beta}, \mathbf{I}_n/\phi)$$

precision $\phi = 1/\sigma^2$

Bayesian Estimation

Model

$$\mathbf{Y} \sim N(\mathbf{X}\boldsymbol{\beta}, \mathbf{I}_n/\phi)$$

precision $\phi = 1/\sigma^2$

Normal-Gamma Conjugate prior $NG(\mathbf{b}_0, \Phi_0, \nu_0, SS_0)$

$$\Phi_n = \mathbf{X}^T \mathbf{X} + \Phi_0$$

$$\mathbf{b}_n = \Phi_n^{-1}(\mathbf{X}^T \mathbf{X} \hat{\boldsymbol{\beta}} + \Phi_0 \mathbf{b}_0)$$

$$\nu_n = \nu_0 + n$$

$$SS_n = SSE + SS_0 + \hat{\boldsymbol{\beta}}^T \mathbf{X}^T \mathbf{X} \hat{\boldsymbol{\beta}} + \mathbf{b}_0^T \Phi_0 \mathbf{b}_0 - \mathbf{b}_n^T \Phi_n \mathbf{b}_n$$

$$\hat{\sigma}_n^2 \equiv SS_n / \nu_n$$

Bayesian Estimation

Model

$$\mathbf{Y} \sim N(\mathbf{X}\beta, \mathbf{I}_n/\phi)$$

precision $\phi = 1/\sigma^2$

Normal-Gamma Conjugate prior $NG(\mathbf{b}_0, \Phi_0, \nu_0, SS_0)$

$$\Phi_n = \mathbf{X}^T \mathbf{X} + \Phi_0$$

$$\mathbf{b}_n = \Phi_n^{-1}(\mathbf{X}^T \mathbf{X} \hat{\beta} + \Phi_0 \mathbf{b}_0)$$

$$\nu_n = \nu_0 + n$$

$$SS_n = SSE + SS_0 + \hat{\beta}^T \mathbf{X}^T \mathbf{X} \hat{\beta} + \mathbf{b}_0^T \Phi_0 \mathbf{b}_0 - \mathbf{b}_n^T \Phi_n \mathbf{b}_n$$

$$\hat{\sigma}_n^2 \equiv SS_n / \nu_n$$

Posterior Distribution

$$\beta \mid \phi, \mathbf{Y} \sim N(\mathbf{b}_n, (\phi \Phi_n)^{-1})$$

Bayesian Estimation

Model

$$\mathbf{Y} \sim \mathcal{N}(\mathbf{X}\boldsymbol{\beta}, \mathbf{I}_n/\phi)$$

precision $\phi = 1/\sigma^2$

Normal-Gamma Conjugate prior $\text{NG}(\mathbf{b}_0, \Phi_0, \nu_0, \text{SS}_0)$

$$\Phi_n = \mathbf{X}^T \mathbf{X} + \Phi_0$$

$$\mathbf{b}_n = \Phi_n^{-1}(\mathbf{X}^T \mathbf{X} \hat{\boldsymbol{\beta}} + \Phi_0 \mathbf{b}_0)$$

$$\nu_n = \nu_0 + n$$

$$\text{SS}_n = \text{SSE} + \text{SS}_0 + \hat{\boldsymbol{\beta}}^T \mathbf{X}^T \mathbf{X} \hat{\boldsymbol{\beta}} + \mathbf{b}_0^T \Phi_0 \mathbf{b}_0 - \mathbf{b}_n^T \Phi_n \mathbf{b}_n$$

$$\hat{\sigma}_n^2 \equiv \text{SS}_n / \nu_n$$

Posterior Distribution

$$\boldsymbol{\beta} \mid \phi, \mathbf{Y} \sim \mathcal{N}(\mathbf{b}_n, (\phi \Phi_n)^{-1})$$

$$\phi \mid \mathbf{Y} \sim \mathbf{G}\left(\frac{\nu_n}{2}, \frac{\nu_n \hat{\sigma}_n^2}{2}\right)$$

Marginal Distribution from Normal–Gamma

Theorem

Let $\boldsymbol{\theta} \mid \phi \sim N(m, \frac{1}{\phi}\Sigma)$ and $\phi \sim \mathbf{G}(\nu/2, \nu\hat{\sigma}^2/2)$. Then \mathbf{t} ($p \times 1$) has a p dimensional multivariate t distribution

$$\boldsymbol{\theta} \sim t_{\nu}(m, \hat{\sigma}^2\Sigma)$$

with density

$$p(\boldsymbol{\theta}) \propto \left[1 + \frac{1}{\nu} \frac{(\boldsymbol{\theta} - m)^T \Sigma^{-1} (\boldsymbol{\theta} - m)}{\hat{\sigma}^2} \right]^{-\frac{p+\nu}{2}}$$

Derivation

Marginal density $p(\boldsymbol{\theta}) = \int p(\boldsymbol{\theta} \mid \phi)p(\phi) d\phi$

Derivation

Marginal density $p(\boldsymbol{\theta}) = \int p(\boldsymbol{\theta} \mid \phi) p(\phi) d\phi$

$$p(\boldsymbol{\theta}) \propto \int |\Sigma/\phi|^{-1/2} e^{-\frac{\phi}{2}(\boldsymbol{\theta}-\mathbf{m})^T \Sigma^{-1}(\boldsymbol{\theta}-\mathbf{m})} \phi^{\nu/2-1} e^{-\phi \frac{\nu \hat{\sigma}^2}{2}} d\phi$$

Derivation

Marginal density $p(\boldsymbol{\theta}) = \int p(\boldsymbol{\theta} \mid \phi) p(\phi) d\phi$

$$\begin{aligned} p(\boldsymbol{\theta}) &\propto \int |\Sigma/\phi|^{-1/2} e^{-\frac{\phi}{2}(\boldsymbol{\theta}-\mathbf{m})^T \Sigma^{-1}(\boldsymbol{\theta}-\mathbf{m})} \phi^{\nu/2-1} e^{-\phi \frac{\nu \hat{\sigma}^2}{2}} d\phi \\ &\propto \int \phi^{p/2} \phi^{\nu/2-1} e^{-\phi \frac{(\boldsymbol{\theta}-\mathbf{m})^T \Sigma^{-1}(\boldsymbol{\theta}-\mathbf{m}) + \nu \hat{\sigma}^2}{2}} d\phi \end{aligned}$$

Derivation

Marginal density $p(\boldsymbol{\theta}) = \int p(\boldsymbol{\theta} \mid \phi) p(\phi) d\phi$

$$\begin{aligned} p(\boldsymbol{\theta}) &\propto \int |\Sigma/\phi|^{-1/2} e^{-\frac{\phi}{2}(\boldsymbol{\theta}-\mathbf{m})^T \Sigma^{-1}(\boldsymbol{\theta}-\mathbf{m})} \phi^{\nu/2-1} e^{-\phi \frac{\nu \hat{\sigma}^2}{2}} d\phi \\ &\propto \int \phi^{p/2} \phi^{\nu/2-1} e^{-\phi \frac{(\boldsymbol{\theta}-\mathbf{m})^T \Sigma^{-1}(\boldsymbol{\theta}-\mathbf{m}) + \nu \hat{\sigma}^2}{2}} d\phi \\ &\propto \int \phi^{\frac{p+\nu}{2}-1} e^{-\phi \frac{(\boldsymbol{\theta}-\mathbf{m})^T \Sigma^{-1}(\boldsymbol{\theta}-\mathbf{m}) + \nu \hat{\sigma}^2}{2}} d\phi \end{aligned}$$

Derivation

Marginal density $p(\boldsymbol{\theta}) = \int p(\boldsymbol{\theta} \mid \phi)p(\phi) d\phi$

$$\begin{aligned} p(\boldsymbol{\theta}) &\propto \int |\Sigma/\phi|^{-1/2} e^{-\frac{\phi}{2}(\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m)} \phi^{\nu/2-1} e^{-\phi \frac{\nu \hat{\sigma}^2}{2}} d\phi \\ &\propto \int \phi^{p/2} \phi^{\nu/2-1} e^{-\phi \frac{(\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m) + \nu \hat{\sigma}^2}{2}} d\phi \\ &\propto \int \phi^{\frac{p+\nu}{2}-1} e^{-\phi \frac{(\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m) + \nu \hat{\sigma}^2}{2}} d\phi \\ &= \Gamma((p+\nu)/2) \left(\frac{(\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m) + \nu \hat{\sigma}^2}{2} \right)^{-\frac{p+\nu}{2}} \end{aligned}$$

Derivation

Marginal density $p(\boldsymbol{\theta}) = \int p(\boldsymbol{\theta} \mid \phi)p(\phi) d\phi$

$$\begin{aligned} p(\boldsymbol{\theta}) &\propto \int |\Sigma/\phi|^{-1/2} e^{-\frac{\phi}{2}(\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m)} \phi^{\nu/2-1} e^{-\phi \frac{\nu \hat{\sigma}^2}{2}} d\phi \\ &\propto \int \phi^{p/2} \phi^{\nu/2-1} e^{-\phi \frac{(\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m) + \nu \hat{\sigma}^2}{2}} d\phi \\ &\propto \int \phi^{\frac{p+\nu}{2}-1} e^{-\phi \frac{(\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m) + \nu \hat{\sigma}^2}{2}} d\phi \\ &= \Gamma((p+\nu)/2) \left(\frac{(\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m) + \nu \hat{\sigma}^2}{2} \right)^{-\frac{p+\nu}{2}} \\ &\propto \left((\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m) + \nu \hat{\sigma}^2 \right)^{-\frac{p+\nu}{2}} \end{aligned}$$

Derivation

Marginal density $p(\boldsymbol{\theta}) = \int p(\boldsymbol{\theta} \mid \phi)p(\phi) d\phi$

$$\begin{aligned} p(\boldsymbol{\theta}) &\propto \int |\Sigma/\phi|^{-1/2} e^{-\frac{\phi}{2}(\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m)} \phi^{\nu/2-1} e^{-\phi \frac{\nu \hat{\sigma}^2}{2}} d\phi \\ &\propto \int \phi^{p/2} \phi^{\nu/2-1} e^{-\phi \frac{(\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m) + \nu \hat{\sigma}^2}{2}} d\phi \\ &\propto \int \phi^{\frac{p+\nu}{2}-1} e^{-\phi \frac{(\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m) + \nu \hat{\sigma}^2}{2}} d\phi \\ &= \Gamma((p+\nu)/2) \left(\frac{(\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m) + \nu \hat{\sigma}^2}{2} \right)^{-\frac{p+\nu}{2}} \\ &\propto \left((\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m) + \nu \hat{\sigma}^2 \right)^{-\frac{p+\nu}{2}} \\ &\propto \left(1 + \frac{1}{\nu} \frac{(\boldsymbol{\theta}-m)^T \Sigma^{-1}(\boldsymbol{\theta}-m)}{\hat{\sigma}^2} \right)^{-\frac{p+\nu}{2}} \end{aligned}$$

Marginal Posterior Distribution of β

$$\beta \mid \phi, \mathbf{Y} \sim \mathbf{N}(\mathbf{b}_n, \phi^{-1} \Phi_n^{-1})$$

Marginal Posterior Distribution of β

$$\beta \mid \phi, \mathbf{Y} \sim \mathbf{N}(\mathbf{b}_n, \phi^{-1} \Phi_n^{-1})$$

$$\phi \mid \mathbf{Y} \sim \mathbf{G}\left(\frac{\nu_n}{2}, \frac{SS_n}{2}\right)$$

Marginal Posterior Distribution of β

$$\beta \mid \phi, \mathbf{Y} \sim \mathbf{N}(\mathbf{b}_n, \phi^{-1} \Phi_n^{-1})$$

$$\phi \mid \mathbf{Y} \sim \mathbf{G}\left(\frac{\nu_n}{2}, \frac{SS_n}{2}\right)$$

Let $\hat{\sigma}^2 = SS_n/\nu_n$ (Bayesian MSE)

Marginal Posterior Distribution of β

$$\beta \mid \phi, \mathbf{Y} \sim \mathbf{N}(\mathbf{b}_n, \phi^{-1} \Phi_n^{-1})$$

$$\phi \mid \mathbf{Y} \sim \mathbf{G}\left(\frac{\nu_n}{2}, \frac{SS_n}{2}\right)$$

Let $\hat{\sigma}^2 = SS_n/\nu_n$ (Bayesian MSE)

Then the marginal posterior distribution of β is

$$\beta \mid \mathbf{Y} \sim t_{\nu_n}(\mathbf{b}_n, \hat{\sigma}^2 \Phi_n^{-1})$$

Marginal Posterior Distribution of β

$$\begin{aligned}\beta \mid \phi, \mathbf{Y} &\sim \mathbf{N}(\mathbf{b}_n, \phi^{-1} \Phi_n^{-1}) \\ \phi \mid \mathbf{Y} &\sim \mathbf{G}\left(\frac{\nu_n}{2}, \frac{SS_n}{2}\right)\end{aligned}$$

Let $\hat{\sigma}^2 = SS_n/\nu_n$ (Bayesian MSE)

Then the marginal posterior distribution of β is

$$\beta \mid \mathbf{Y} \sim t_{\nu_n}(\mathbf{b}_n, \hat{\sigma}^2 \Phi_n^{-1})$$

Any linear combination $\lambda^T \beta$

$$\lambda^T \beta \mid \mathbf{Y} \sim t_{\nu_n}(\lambda^T \mathbf{b}_n, \hat{\sigma}^2 \lambda^T \Phi_n^{-1} \lambda)$$

has a univariate t distribution with ν_n degrees of freedom

Predictive Distribution

Suppose $\mathbf{Y}^* \mid \beta, \phi \sim \mathcal{N}(\mathbf{X}^*\beta, \mathbf{I}/\phi)$ and is conditionally independent of \mathbf{Y} given β and ϕ

Predictive Distribution

Suppose $\mathbf{Y}^* \mid \boldsymbol{\beta}, \phi \sim \mathcal{N}(\mathbf{X}^* \boldsymbol{\beta}, \mathbf{I}/\phi)$ and is conditionally independent of \mathbf{Y} given $\boldsymbol{\beta}$ and ϕ

What is the predictive distribution of $\mathbf{Y}^* \mid \mathbf{Y}$?

Predictive Distribution

Suppose $\mathbf{Y}^* \mid \boldsymbol{\beta}, \phi \sim \mathcal{N}(\mathbf{X}^*\boldsymbol{\beta}, \mathbf{I}/\phi)$ and is conditionally independent of \mathbf{Y} given $\boldsymbol{\beta}$ and ϕ

What is the predictive distribution of $\mathbf{Y}^* \mid \mathbf{Y}$?

$\mathbf{Y}^* = \mathbf{X}^*\boldsymbol{\beta} + \boldsymbol{\epsilon}^*$ and $\boldsymbol{\epsilon}^*$ is independent of \mathbf{Y} given ϕ

Predictive Distribution

Suppose $\mathbf{Y}^* \mid \boldsymbol{\beta}, \phi \sim \mathcal{N}(\mathbf{X}^*\boldsymbol{\beta}, \mathbf{I}/\phi)$ and is conditionally independent of \mathbf{Y} given $\boldsymbol{\beta}$ and ϕ

What is the predictive distribution of $\mathbf{Y}^* \mid \mathbf{Y}$?

$\mathbf{Y}^* = \mathbf{X}^*\boldsymbol{\beta} + \boldsymbol{\epsilon}^*$ and $\boldsymbol{\epsilon}^*$ is independent of \mathbf{Y} given ϕ

$$\mathbf{X}^*\boldsymbol{\beta} + \boldsymbol{\epsilon}^* \mid \phi, \mathbf{Y} \sim \mathcal{N}(\mathbf{X}^*\mathbf{b}_n, (\mathbf{X}^*\boldsymbol{\Phi}_n^{-1}\mathbf{X}^{*T} + \mathbf{I})/\phi)$$

Predictive Distribution

Suppose $\mathbf{Y}^* \mid \beta, \phi \sim \mathcal{N}(\mathbf{X}^*\beta, \mathbf{I}/\phi)$ and is conditionally independent of \mathbf{Y} given β and ϕ

What is the predictive distribution of $\mathbf{Y}^* \mid \mathbf{Y}$?

$\mathbf{Y}^* = \mathbf{X}^*\beta + \epsilon^*$ and ϵ^* is independent of \mathbf{Y} given ϕ

$$\mathbf{X}^*\beta + \epsilon^* \mid \phi, \mathbf{Y} \sim \mathcal{N}(\mathbf{X}^*\mathbf{b}_n, (\mathbf{X}^*\Phi_n^{-1}\mathbf{X}^{*T} + \mathbf{I})/\phi)$$

$$\mathbf{Y}^* \mid \phi, \mathbf{Y} \sim \mathcal{N}(\mathbf{X}^*\mathbf{b}_n, (\mathbf{X}^*\Phi_n^{-1}\mathbf{X}^{*T} + \mathbf{I})/\phi)$$

Predictive Distribution

Suppose $\mathbf{Y}^* \mid \boldsymbol{\beta}, \phi \sim \mathcal{N}(\mathbf{X}^* \boldsymbol{\beta}, \mathbf{I}/\phi)$ and is conditionally independent of \mathbf{Y} given $\boldsymbol{\beta}$ and ϕ

What is the predictive distribution of $\mathbf{Y}^* \mid \mathbf{Y}$?

$\mathbf{Y}^* = \mathbf{X}^* \boldsymbol{\beta} + \boldsymbol{\epsilon}^*$ and $\boldsymbol{\epsilon}^*$ is independent of \mathbf{Y} given ϕ

$$\mathbf{X}^* \boldsymbol{\beta} + \boldsymbol{\epsilon}^* \mid \phi, \mathbf{Y} \sim \mathcal{N}(\mathbf{X}^* \mathbf{b}_n, (\mathbf{X}^* \boldsymbol{\Phi}_n^{-1} \mathbf{X}^{*T} + \mathbf{I})/\phi)$$

$$\mathbf{Y}^* \mid \phi, \mathbf{Y} \sim \mathcal{N}(\mathbf{X}^* \mathbf{b}_n, (\mathbf{X}^* \boldsymbol{\Phi}_n^{-1} \mathbf{X}^{*T} + \mathbf{I})/\phi)$$

$$\phi \mid \mathbf{Y} \sim \mathbf{G}\left(\frac{\nu_n}{2}, \frac{\hat{\sigma}^2 \nu_n}{2}\right)$$

Predictive Distribution

Suppose $\mathbf{Y}^* \mid \boldsymbol{\beta}, \phi \sim \mathcal{N}(\mathbf{X}^* \boldsymbol{\beta}, \mathbf{I}/\phi)$ and is conditionally independent of \mathbf{Y} given $\boldsymbol{\beta}$ and ϕ

What is the predictive distribution of $\mathbf{Y}^* \mid \mathbf{Y}$?

$\mathbf{Y}^* = \mathbf{X}^* \boldsymbol{\beta} + \boldsymbol{\epsilon}^*$ and $\boldsymbol{\epsilon}^*$ is independent of \mathbf{Y} given ϕ

$$\mathbf{X}^* \boldsymbol{\beta} + \boldsymbol{\epsilon}^* \mid \phi, \mathbf{Y} \sim \mathcal{N}(\mathbf{X}^* \mathbf{b}_n, (\mathbf{X}^* \boldsymbol{\Phi}_n^{-1} \mathbf{X}^{*T} + \mathbf{I})/\phi)$$

$$\mathbf{Y}^* \mid \phi, \mathbf{Y} \sim \mathcal{N}(\mathbf{X}^* \mathbf{b}_n, (\mathbf{X}^* \boldsymbol{\Phi}_n^{-1} \mathbf{X}^{*T} + \mathbf{I})/\phi)$$

$$\phi \mid \mathbf{Y} \sim \mathbf{G}\left(\frac{\nu_n}{2}, \frac{\hat{\sigma}^2 \nu_n}{2}\right)$$

$$\mathbf{Y}^* \mid \mathbf{Y} \sim t_{\nu_n}(\mathbf{X}^* \mathbf{b}_n, \hat{\sigma}^2 (\mathbf{I} + \mathbf{X}^* \boldsymbol{\Phi}_n^{-1} \mathbf{X}^T))$$

Conjugate Priors

Definition

A class of prior distributions \mathcal{P} for θ is conjugate for a sampling model $p(y \mid \theta)$ if for every $p(\theta) \in \mathcal{P}$, $p(\theta \mid \mathbf{Y}) \in \mathcal{P}$.

Conjugate Priors

Definition

A class of prior distributions \mathcal{P} for θ is conjugate for a sampling model $p(y \mid \theta)$ if for every $p(\theta) \in \mathcal{P}$, $p(\theta \mid \mathbf{Y}) \in \mathcal{P}$.

Advantages:

Conjugate Priors

Definition

A class of prior distributions \mathcal{P} for θ is conjugate for a sampling model $p(y \mid \theta)$ if for every $p(\theta) \in \mathcal{P}$, $p(\theta \mid \mathbf{Y}) \in \mathcal{P}$.

Advantages:

- ▶ Closed form distributions for most quantities; bypass MCMC for calculations

Conjugate Priors

Definition

A class of prior distributions \mathcal{P} for θ is conjugate for a sampling model $p(y \mid \theta)$ if for every $p(\theta) \in \mathcal{P}$, $p(\theta \mid \mathbf{Y}) \in \mathcal{P}$.

Advantages:

- ▶ Closed form distributions for most quantities; bypass MCMC for calculations
- ▶ Simple updating in terms of sufficient statistics “weighted average”

Conjugate Priors

Definition

A class of prior distributions \mathcal{P} for θ is conjugate for a sampling model $p(y \mid \theta)$ if for every $p(\theta) \in \mathcal{P}$, $p(\theta \mid \mathbf{Y}) \in \mathcal{P}$.

Advantages:

- ▶ Closed form distributions for most quantities; bypass MCMC for calculations
- ▶ Simple updating in terms of sufficient statistics “weighted average”
- ▶ Interpretation as prior samples - prior sample size

Conjugate Priors

Definition

A class of prior distributions \mathcal{P} for θ is conjugate for a sampling model $p(y \mid \theta)$ if for every $p(\theta) \in \mathcal{P}$, $p(\theta \mid \mathbf{Y}) \in \mathcal{P}$.

Advantages:

- ▶ Closed form distributions for most quantities; bypass MCMC for calculations
- ▶ Simple updating in terms of sufficient statistics “weighted average”
- ▶ Interpretation as prior samples - prior sample size
- ▶ Elicitation of prior through imaginary or historical data

Conjugate Priors

Definition

A class of prior distributions \mathcal{P} for θ is conjugate for a sampling model $p(y \mid \theta)$ if for every $p(\theta) \in \mathcal{P}$, $p(\theta \mid \mathbf{Y}) \in \mathcal{P}$.

Advantages:

- ▶ Closed form distributions for most quantities; bypass MCMC for calculations
- ▶ Simple updating in terms of sufficient statistics “weighted average”
- ▶ Interpretation as prior samples - prior sample size
- ▶ Elicitation of prior through imaginary or historical data
- ▶ limiting “non-proper” form recovers MLEs

Conjugate Priors

Definition

A class of prior distributions \mathcal{P} for θ is conjugate for a sampling model $p(y \mid \theta)$ if for every $p(\theta) \in \mathcal{P}$, $p(\theta \mid \mathbf{Y}) \in \mathcal{P}$.

Advantages:

- ▶ Closed form distributions for most quantities; bypass MCMC for calculations
- ▶ Simple updating in terms of sufficient statistics “weighted average”
- ▶ Interpretation as prior samples - prior sample size
- ▶ Elicitation of prior through imaginary or historical data
- ▶ limiting “non-proper” form recovers MLEs

Choice of conjugate prior?

Unit Information Prior

Unit information prior $\beta \mid \phi \sim \mathbf{N}(\hat{\beta}, n(\mathbf{X}^T \mathbf{X})^{-1} / \phi)$

Unit Information Prior

Unit information prior $\beta \mid \phi \sim N(\hat{\beta}, n(\mathbf{X}^T \mathbf{X})^{-1} / \phi)$

- Fisher Information is $\phi \mathbf{X}^T \mathbf{X}$ based on a sample of n observations

Unit Information Prior

Unit information prior $\beta \mid \phi \sim N(\hat{\beta}, n(\mathbf{X}^T \mathbf{X})^{-1} / \phi)$

- ▶ Fisher Information is $\phi \mathbf{X}^T \mathbf{X}$ based on a sample of n observations
- ▶ Inverse Fisher information is covariance matrix of MLE

Unit Information Prior

Unit information prior $\beta \mid \phi \sim N(\hat{\beta}, n(\mathbf{X}^T \mathbf{X})^{-1} / \phi)$

- ▶ Fisher Information is $\phi \mathbf{X}^T \mathbf{X}$ based on a sample of n observations
- ▶ Inverse Fisher information is covariance matrix of MLE
- ▶ “average information” in one observation is $\phi \mathbf{X}^T \mathbf{X} / n$

Unit Information Prior

Unit information prior $\beta \mid \phi \sim N(\hat{\beta}, n(\mathbf{X}^T \mathbf{X})^{-1} / \phi)$

- ▶ Fisher Information is $\phi \mathbf{X}^T \mathbf{X}$ based on a sample of n observations
- ▶ Inverse Fisher information is covariance matrix of MLE
- ▶ “average information” in one observation is $\phi \mathbf{X}^T \mathbf{X} / n$
- ▶ center prior at MLE and base covariance on the information in “1” observation

Unit Information Prior

Unit information prior $\beta \mid \phi \sim N(\hat{\beta}, n(\mathbf{X}^T \mathbf{X})^{-1} / \phi)$

- ▶ Fisher Information is $\phi \mathbf{X}^T \mathbf{X}$ based on a sample of n observations
- ▶ Inverse Fisher information is covariance matrix of MLE
- ▶ “average information” in one observation is $\phi \mathbf{X}^T \mathbf{X} / n$
- ▶ center prior at MLE and base covariance on the information in “1” observation
- ▶ Posterior mean

$$\frac{n}{1+n} \hat{\beta} + \frac{1}{1+n} \hat{\beta} = \hat{\beta}$$

Unit Information Prior

Unit information prior $\beta \mid \phi \sim N(\hat{\beta}, n(\mathbf{X}^T \mathbf{X})^{-1}/\phi)$

- ▶ Fisher Information is $\phi \mathbf{X}^T \mathbf{X}$ based on a sample of n observations
- ▶ Inverse Fisher information is covariance matrix of MLE
- ▶ “average information” in one observation is $\phi \mathbf{X}^T \mathbf{X}/n$
- ▶ center prior at MLE and base covariance on the information in “1” observation
- ▶ Posterior mean

$$\frac{n}{1+n} \hat{\beta} + \frac{1}{1+n} \hat{\beta} = \hat{\beta}$$

- ▶ Posterior Distribution

$$\beta \mid \mathbf{Y}, \phi \sim N\left(\hat{\beta}, \frac{n}{1+n}(\mathbf{X}^T \mathbf{X})^{-1} \phi^{-1}\right)$$

Unit Information Prior

Unit information prior $\beta \mid \phi \sim N(\hat{\beta}, n(\mathbf{X}^T \mathbf{X})^{-1}/\phi)$

- ▶ Fisher Information is $\phi \mathbf{X}^T \mathbf{X}$ based on a sample of n observations
- ▶ Inverse Fisher information is covariance matrix of MLE
- ▶ “average information” in one observation is $\phi \mathbf{X}^T \mathbf{X}/n$
- ▶ center prior at MLE and base covariance on the information in “1” observation
- ▶ Posterior mean

$$\frac{n}{1+n} \hat{\beta} + \frac{1}{1+n} \hat{\beta} = \hat{\beta}$$

- ▶ Posterior Distribution

$$\beta \mid \mathbf{Y}, \phi \sim N \left(\hat{\beta}, \frac{n}{1+n} (\mathbf{X}^T \mathbf{X})^{-1} \phi^{-1} \right)$$

Cannot represent real prior beliefs; double use of data but has the “right” behaviour.

Zellner's g -prior

$$\text{Zellner's } g\text{-prior(s) } \beta \mid \phi \sim N(\mathbf{b}_0, g(\mathbf{X}^T \mathbf{X})^{-1} / \phi)$$

Zellner's g -prior

Zellner's g -prior(s) $\beta \mid \phi \sim N(\mathbf{b}_0, g(\mathbf{X}^T \mathbf{X})^{-1} / \phi)$

$$\beta \mid \mathbf{Y}, \phi \sim N \left(\frac{g}{1+g} \hat{\beta} + \frac{1}{1+g} \mathbf{b}_0, \frac{g}{1+g} (\mathbf{X}^T \mathbf{X})^{-1} \phi^{-1} \right)$$

Zellner's g -prior

Zellner's g -prior(s) $\beta \mid \phi \sim N(\mathbf{b}_0, g(\mathbf{X}^T \mathbf{X})^{-1} / \phi)$

$$\beta \mid \mathbf{Y}, \phi \sim N \left(\frac{g}{1+g} \hat{\beta} + \frac{1}{1+g} \mathbf{b}_0, \frac{g}{1+g} (\mathbf{X}^T \mathbf{X})^{-1} \phi^{-1} \right)$$

- Invariance: Require posterior of $\mathbf{X}\beta$ equal the posterior of $\mathbf{X}\mathbf{H}\alpha$

Zellner's g -prior

Zellner's g -prior(s) $\beta \mid \phi \sim N(\mathbf{b}_0, g(\mathbf{X}^T \mathbf{X})^{-1} / \phi)$

$$\beta \mid \mathbf{Y}, \phi \sim N \left(\frac{g}{1+g} \hat{\beta} + \frac{1}{1+g} \mathbf{b}_0, \frac{g}{1+g} (\mathbf{X}^T \mathbf{X})^{-1} \phi^{-1} \right)$$

- ▶ Invariance: Require posterior of $\mathbf{X}\beta$ equal the posterior of $\mathbf{X}\mathbf{H}\alpha$ ($\mathbf{a}_0 = \mathbf{H}^{-1}\mathbf{b}_0$) (take $\mathbf{b}_0 = \mathbf{0}$)
- ▶ Choice of g ?

Zellner's g -prior

Zellner's g -prior(s) $\beta \mid \phi \sim N(\mathbf{b}_0, g(\mathbf{X}^T \mathbf{X})^{-1} / \phi)$

$$\beta \mid \mathbf{Y}, \phi \sim N \left(\frac{g}{1+g} \hat{\beta} + \frac{1}{1+g} \mathbf{b}_0, \frac{g}{1+g} (\mathbf{X}^T \mathbf{X})^{-1} \phi^{-1} \right)$$

- ▶ Invariance: Require posterior of $\mathbf{X}\beta$ equal the posterior of $\mathbf{X}\mathbf{H}\alpha$ ($\mathbf{a}_0 = \mathbf{H}^{-1}\mathbf{b}_0$) (take $\mathbf{b}_0 = \mathbf{0}$)
- ▶ Choice of g ?
- ▶ $\frac{g}{1+g}$ weight given to the data

Zellner's g -prior


Zellner's g -prior(s) $\beta \mid \phi \sim N(\mathbf{b}_0, g(\mathbf{X}^T \mathbf{X})^{-1} / \phi)$

$$\beta \mid \mathbf{Y}, \phi \sim N \left(\frac{g}{1+g} \hat{\beta} + \frac{1}{1+g} \mathbf{b}_0, \frac{g}{1+g} (\mathbf{X}^T \mathbf{X})^{-1} \phi^{-1} \right)$$

- ▶ Invariance: Require posterior of $\mathbf{X}\beta$ equal the posterior of $\mathbf{X}\mathbf{H}\alpha$ ($\mathbf{a}_0 = \mathbf{H}^{-1}\mathbf{b}_0$) (take $\mathbf{b}_0 = \mathbf{0}$)
- ▶ Choice of g ?
- ▶ $\frac{g}{1+g}$ weight given to the data
- ▶ Fixed g effect does not vanish as $n \rightarrow \infty$
- ▶ Use $g = n$ or place a prior distribution on g

Shrinkage

Posterior mean under g -prior with $\mathbf{b}_0 = 0$ $\frac{g}{1+g}\hat{\beta}$



Jeffreys Prior

Jeffreys proposed a default procedure so that resulting prior would be invariant to model parameterization

Jeffreys Prior

Jeffreys proposed a default procedure so that resulting prior would be invariant to model parameterization

$$p(\boldsymbol{\theta}) \propto |\mathcal{J}(\boldsymbol{\theta})|^{1/2}$$

Jeffreys Prior

Jeffreys proposed a default procedure so that resulting prior would be invariant to model parameterization

$$p(\boldsymbol{\theta}) \propto |\mathcal{I}(\boldsymbol{\theta})|^{1/2}$$

where $\mathcal{I}(\boldsymbol{\theta})$ is the Expected Fisher Information matrix

Jeffreys Prior

Jeffreys proposed a default procedure so that resulting prior would be invariant to model parameterization

$$p(\boldsymbol{\theta}) \propto |\mathcal{I}(\boldsymbol{\theta})|^{1/2}$$

where $\mathcal{I}(\boldsymbol{\theta})$ is the Expected Fisher Information matrix

$$\mathcal{I}(\boldsymbol{\theta}) = -\mathbb{E}\left[\frac{\partial^2 \log(\mathcal{L}(\boldsymbol{\theta}))}{\partial \theta_i \partial \theta_j}\right]$$

Fisher Information Matrix

Log Likelihood

$$\log(\mathcal{L}(\boldsymbol{\beta}, \phi)) = \frac{n}{2} \log(\phi) - \frac{\phi}{2} \|(\mathbf{I} - \mathbf{P}_{\mathbf{x}})\mathbf{Y}\|^2 - \frac{\phi}{2} (\boldsymbol{\beta} - \hat{\boldsymbol{\beta}})^T (\mathbf{X}^T \mathbf{X}) (\boldsymbol{\beta} - \hat{\boldsymbol{\beta}})$$

Fisher Information Matrix

Log Likelihood

$$\log(\mathcal{L}(\boldsymbol{\beta}, \phi)) = \frac{n}{2} \log(\phi) - \frac{\phi}{2} \|(\mathbf{I} - \mathbf{P}_{\mathbf{X}}) \mathbf{Y}\|^2 - \frac{\phi}{2} (\boldsymbol{\beta} - \hat{\boldsymbol{\beta}})^T (\mathbf{X}^T \mathbf{X}) (\boldsymbol{\beta} - \hat{\boldsymbol{\beta}})$$

$$\frac{\partial^2 \log \mathcal{L}}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T} = \begin{bmatrix} -\phi(\mathbf{X}^T \mathbf{X}) & -(\mathbf{X}^T \mathbf{X})(\boldsymbol{\beta} - \hat{\boldsymbol{\beta}}) \\ -(\boldsymbol{\beta} - \hat{\boldsymbol{\beta}})^T (\mathbf{X}^T \mathbf{X}) & -\frac{n}{2} \frac{1}{\phi^2} \end{bmatrix}$$

Fisher Information Matrix

Log Likelihood

$$\log(\mathcal{L}(\boldsymbol{\beta}, \phi)) = \frac{n}{2} \log(\phi) - \frac{\phi}{2} \|(\mathbf{I} - \mathbf{P}_{\mathbf{x}})\mathbf{Y}\|^2 - \frac{\phi}{2} (\boldsymbol{\beta} - \hat{\boldsymbol{\beta}})^T (\mathbf{X}^T \mathbf{X}) (\boldsymbol{\beta} - \hat{\boldsymbol{\beta}})$$

$$\begin{aligned} \frac{\partial^2 \log \mathcal{L}}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T} &= \begin{bmatrix} -\phi(\mathbf{X}^T \mathbf{X}) & -(\mathbf{X}^T \mathbf{X})(\boldsymbol{\beta} - \hat{\boldsymbol{\beta}}) \\ -(\boldsymbol{\beta} - \hat{\boldsymbol{\beta}})^T (\mathbf{X}^T \mathbf{X}) & -\frac{n}{2} \frac{1}{\phi^2} \end{bmatrix} \\ \mathbb{E}\left[\frac{\partial^2 \log \mathcal{L}}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T}\right] &= \begin{bmatrix} -\phi(\mathbf{X}^T \mathbf{X}) & \mathbf{0}_p \\ \mathbf{0}_p^T & -\frac{n}{2} \frac{1}{\phi^2} \end{bmatrix} \end{aligned}$$

Fisher Information Matrix

Log Likelihood

$$\log(\mathcal{L}(\boldsymbol{\beta}, \phi)) = \frac{n}{2} \log(\phi) - \frac{\phi}{2} \|(\mathbf{I} - \mathbf{P}_{\mathbf{x}})\mathbf{Y}\|^2 - \frac{\phi}{2} (\boldsymbol{\beta} - \hat{\boldsymbol{\beta}})^T (\mathbf{X}^T \mathbf{X}) (\boldsymbol{\beta} - \hat{\boldsymbol{\beta}})$$

$$\frac{\partial^2 \log \mathcal{L}}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T} = \begin{bmatrix} -\phi(\mathbf{X}^T \mathbf{X}) & -(\mathbf{X}^T \mathbf{X})(\boldsymbol{\beta} - \hat{\boldsymbol{\beta}}) \\ -(\boldsymbol{\beta} - \hat{\boldsymbol{\beta}})^T (\mathbf{X}^T \mathbf{X}) & -\frac{n}{2} \frac{1}{\phi^2} \end{bmatrix}$$

$$\mathbb{E}\left[\frac{\partial^2 \log \mathcal{L}}{\partial \boldsymbol{\theta} \partial \boldsymbol{\theta}^T}\right] = \begin{bmatrix} -\phi(\mathbf{X}^T \mathbf{X}) & \mathbf{0}_p \\ \mathbf{0}_p^T & -\frac{n}{2} \frac{1}{\phi^2} \end{bmatrix}$$

$$\mathcal{I}((\boldsymbol{\beta}, \phi)^T) = \begin{bmatrix} \phi(\mathbf{X}^T \mathbf{X}) & \mathbf{0}_p \\ \mathbf{0}_p^T & \frac{n}{2} \frac{1}{\phi^2} \end{bmatrix}$$

Jeffreys Prior

Jeffreys Prior

$$p_J(\boldsymbol{\beta}, \phi) \propto |\mathcal{I}((\boldsymbol{\beta}, \phi)^T)|^{1/2}$$

Jeffreys Prior

Jeffreys Prior

$$\begin{aligned} p_J(\boldsymbol{\beta}, \phi) &\propto |\mathcal{I}((\boldsymbol{\beta}, \phi)^T)|^{1/2} \\ &= |\phi(\mathbf{X}^T \mathbf{X})|^{1/2} \left(\frac{n}{2} \frac{1}{\phi^2} \right)^{1/2} \end{aligned}$$

Jeffreys Prior

Jeffreys Prior

$$\begin{aligned} p_J(\boldsymbol{\beta}, \phi) &\propto |\mathcal{I}((\boldsymbol{\beta}, \phi)^T)|^{1/2} \\ &= |\phi(\mathbf{X}^T \mathbf{X})|^{1/2} \left(\frac{n}{2} \frac{1}{\phi^2} \right)^{1/2} \\ &\propto \phi^{p/2-1} |\mathbf{X}^T \mathbf{X}|^{1/2} \end{aligned}$$

Jeffreys Prior

Jeffreys Prior

$$\begin{aligned} p_J(\boldsymbol{\beta}, \phi) &\propto |\mathcal{I}((\boldsymbol{\beta}, \phi)^T)|^{1/2} \\ &= |\phi(\mathbf{X}^T \mathbf{X})|^{1/2} \left(\frac{n}{2} \frac{1}{\phi^2} \right)^{1/2} \\ &\propto \phi^{p/2-1} |\mathbf{X}^T \mathbf{X}|^{1/2} \\ &\propto \phi^{p/2-1} \end{aligned}$$

Jeffreys Prior

Jeffreys Prior

$$\begin{aligned} p_J(\boldsymbol{\beta}, \phi) &\propto |\mathcal{I}((\boldsymbol{\beta}, \phi)^T)|^{1/2} \\ &= |\phi(\mathbf{X}^T \mathbf{X})|^{1/2} \left(\frac{n}{2} \frac{1}{\phi^2} \right)^{1/2} \\ &\propto \phi^{p/2-1} |\mathbf{X}^T \mathbf{X}|^{1/2} \\ &\propto \phi^{p/2-1} \end{aligned}$$

Improper prior $\iint p_J(\boldsymbol{\beta}, \phi) d\boldsymbol{\beta} d\phi$ not finite

Formal Bayes Posterior

$$p(\beta, \phi \mid \mathbf{Y}) \propto p(\mathbf{Y} \mid \beta, \phi) \phi^{p/2-1}$$

Formal Bayes Posterior

$$p(\beta, \phi \mid \mathbf{Y}) \propto p(\mathbf{Y} \mid \beta, \phi) \phi^{p/2-1}$$

if this is integrable, then renormalize to obtain formal posterior distribution

Formal Bayes Posterior

$$p(\boldsymbol{\beta}, \phi \mid \mathbf{Y}) \propto p(\mathbf{Y} \mid \boldsymbol{\beta}, \phi) \phi^{p/2-1}$$

if this is integrable, then renormalize to obtain formal posterior distribution

$$\begin{aligned}\boldsymbol{\beta} \mid \phi, \mathbf{Y} &\sim \mathbf{N}(\hat{\boldsymbol{\beta}}, (\mathbf{X}^T \mathbf{X})^{-1} \phi^{-1}) \\ \phi \mid \mathbf{Y} &\sim \mathbf{G}(n/2, \|\mathbf{Y} - \mathbf{X}\hat{\boldsymbol{\beta}}\|^2/2)\end{aligned}$$

Formal Bayes Posterior

$$p(\boldsymbol{\beta}, \phi \mid \mathbf{Y}) \propto p(\mathbf{Y} \mid \boldsymbol{\beta}, \phi) \phi^{p/2-1}$$

if this is integrable, then renormalize to obtain formal posterior distribution

$$\begin{aligned}\boldsymbol{\beta} \mid \phi, \mathbf{Y} &\sim \mathbf{N}(\hat{\boldsymbol{\beta}}, (\mathbf{X}^T \mathbf{X})^{-1} \phi^{-1}) \\ \phi \mid \mathbf{Y} &\sim \mathbf{G}(n/2, \|\mathbf{Y} - \mathbf{X}\hat{\boldsymbol{\beta}}\|^2/2)\end{aligned}$$

Limiting case of Conjugate prior with $\mathbf{b}_0 = \mathbf{0}$, $\Phi = \mathbf{0}$, $\nu_0 = 0$ and $SS_0 = 0$

Formal Bayes Posterior

$$p(\boldsymbol{\beta}, \phi \mid \mathbf{Y}) \propto p(\mathbf{Y} \mid \boldsymbol{\beta}, \phi) \phi^{p/2-1}$$

if this is integrable, then renormalize to obtain formal posterior distribution

$$\begin{aligned}\boldsymbol{\beta} \mid \phi, \mathbf{Y} &\sim \mathbf{N}(\hat{\boldsymbol{\beta}}, (\mathbf{X}^T \mathbf{X})^{-1} \phi^{-1}) \\ \phi \mid \mathbf{Y} &\sim \mathbf{G}(n/2, \|\mathbf{Y} - \mathbf{X}\hat{\boldsymbol{\beta}}\|^2/2)\end{aligned}$$

Limiting case of Conjugate prior with $\mathbf{b}_0 = \mathbf{0}$, $\Phi = \mathbf{0}$, $\nu_0 = 0$ and $SS_0 = 0$

Posterior does not depend on dimension p ;

Formal Bayes Posterior

$$p(\boldsymbol{\beta}, \phi \mid \mathbf{Y}) \propto p(\mathbf{Y} \mid \boldsymbol{\beta}, \phi) \phi^{p/2-1}$$

if this is integrable, then renormalize to obtain formal posterior distribution

$$\begin{aligned}\boldsymbol{\beta} \mid \phi, \mathbf{Y} &\sim \mathbf{N}(\hat{\boldsymbol{\beta}}, (\mathbf{X}^T \mathbf{X})^{-1} \phi^{-1}) \\ \phi \mid \mathbf{Y} &\sim \mathbf{G}(n/2, \|\mathbf{Y} - \mathbf{X}\hat{\boldsymbol{\beta}}\|^2/2)\end{aligned}$$

Limiting case of Conjugate prior with $\mathbf{b}_0 = \mathbf{0}$, $\Phi = \mathbf{0}$, $\nu_0 = 0$ and $SS_0 = 0$

Posterior does not depend on dimension p ;

Jeffreys did not recommend using this

Independent Jeffreys Prior

- ▶ Treat β and ϕ separately (“orthogonal parameterization”)

Independent Jeffreys Prior

- ▶ Treat β and ϕ separately (“orthogonal parameterization”)
- ▶ $p_{IJ}(\beta) \propto |\mathcal{I}(\beta)|^{1/2}$

Independent Jeffreys Prior

- ▶ Treat β and ϕ separately (“orthogonal parameterization”)
- ▶ $p_{IJ}(\beta) \propto |\mathcal{I}(\beta)|^{1/2}$
- ▶ $p_{IJ}(\phi) \propto |\mathcal{I}(\phi)|^{1/2}$

Independent Jeffreys Prior

- ▶ Treat β and ϕ separately (“orthogonal parameterization”)
- ▶ $p_{IJ}(\beta) \propto |\mathcal{I}(\beta)|^{1/2}$
- ▶ $p_{IJ}(\phi) \propto |\mathcal{I}(\phi)|^{1/2}$

$$\mathcal{I}((\beta, \phi)^T) = \begin{bmatrix} \phi(\mathbf{X}^T \mathbf{X}) & \mathbf{0}_p \\ \mathbf{0}_p^T & \frac{n}{2} \frac{1}{\phi^2} \end{bmatrix}$$

Independent Jeffreys Prior

- ▶ Treat β and ϕ separately (“orthogonal parameterization”)
- ▶ $p_{IJ}(\beta) \propto |\mathcal{I}(\beta)|^{1/2}$
- ▶ $p_{IJ}(\phi) \propto |\mathcal{I}(\phi)|^{1/2}$

$$\mathcal{I}((\beta, \phi)^T) = \begin{bmatrix} \phi(\mathbf{X}^T \mathbf{X}) & \mathbf{0}_p \\ \mathbf{0}_p^T & \frac{n}{2} \frac{1}{\phi^2} \end{bmatrix}$$

$$p_{IJ}(\beta) \propto |\phi \mathbf{X}^T \mathbf{X}|^{1/2} \propto 1$$

Independent Jeffreys Prior

- ▶ Treat β and ϕ separately (“orthogonal parameterization”)
- ▶ $p_{IJ}(\beta) \propto |\mathcal{I}(\beta)|^{1/2}$
- ▶ $p_{IJ}(\phi) \propto |\mathcal{I}(\phi)|^{1/2}$

$$\mathcal{I}((\beta, \phi)^T) = \begin{bmatrix} \phi(\mathbf{X}^T \mathbf{X}) & \mathbf{0}_p \\ \mathbf{0}_p^T & \frac{n}{2} \frac{1}{\phi^2} \end{bmatrix}$$

$$p_{IJ}(\beta) \propto |\phi \mathbf{X}^T \mathbf{X}|^{1/2} \propto 1$$

$$p_{IJ}(\phi) \propto \phi^{-1}$$

Independent Jeffreys Prior

- ▶ Treat β and ϕ separately (“orthogonal parameterization”)
- ▶ $p_{IJ}(\beta) \propto |\mathcal{I}(\beta)|^{1/2}$
- ▶ $p_{IJ}(\phi) \propto |\mathcal{I}(\phi)|^{1/2}$

$$\mathcal{I}((\beta, \phi)^T) = \begin{bmatrix} \phi(\mathbf{X}^T \mathbf{X}) & \mathbf{0}_p \\ \mathbf{0}_p^T & \frac{n}{2} \frac{1}{\phi^2} \end{bmatrix}$$

$$p_{IJ}(\beta) \propto |\phi \mathbf{X}^T \mathbf{X}|^{1/2} \propto 1$$

$$p_{IJ}(\phi) \propto \phi^{-1}$$

Independent Jeffreys Prior is

$$p_{IJ}(\beta, \phi) \propto p_{IJ}(\beta) p_{IJ}(\phi) = \phi^{-1}$$

Formal Posterior Distribution

With Independent Jeffreys Prior

$$p_{IJ}(\beta, \phi) \propto p_{IJ}(\beta) p_{IJ}(\phi) = \phi^{-1}$$

Formal Posterior Distribution

With Independent Jeffreys Prior

$$p_{IJ}(\beta, \phi) \propto p_{IJ}(\beta)p_{IJ}(\phi) = \phi^{-1}$$

Formal Posterior Distribution

Formal Posterior Distribution

With Independent Jeffreys Prior

$$p_{IJ}(\beta, \phi) \propto p_{IJ}(\beta) p_{IJ}(\phi) = \phi^{-1}$$

Formal Posterior Distribution

$$\beta \mid \phi, \mathbf{Y} \sim \mathcal{N}(\hat{\beta}, (\mathbf{X}^T \mathbf{X})^{-1} \phi^{-1})$$

Formal Posterior Distribution

With Independent Jeffreys Prior

$$p_{IJ}(\beta, \phi) \propto p_{IJ}(\beta)p_{IJ}(\phi) = \phi^{-1}$$

Formal Posterior Distribution

$$\begin{aligned}\beta \mid \phi, \mathbf{Y} &\sim \mathbf{N}(\hat{\beta}, (\mathbf{X}^T \mathbf{X})^{-1} \phi^{-1}) \\ \phi \mid \mathbf{Y} &\sim \mathbf{G}((n-p)/2, \|\mathbf{Y} - \mathbf{X}\hat{\beta}\|^2/2)\end{aligned}$$

Formal Posterior Distribution

With Independent Jeffreys Prior

$$p_{IJ}(\beta, \phi) \propto p_{IJ}(\beta)p_{IJ}(\phi) = \phi^{-1}$$

Formal Posterior Distribution

$$\begin{aligned}\beta \mid \phi, \mathbf{Y} &\sim \mathbf{N}(\hat{\beta}, (\mathbf{X}^T \mathbf{X})^{-1} \phi^{-1}) \\ \phi \mid \mathbf{Y} &\sim \mathbf{G}((n-p)/2, \|\mathbf{Y} - \mathbf{X}\hat{\beta}\|^2/2) \\ \beta \mid \mathbf{Y} &\sim t_{n-p}(\hat{\beta}, \hat{\sigma}^2(\mathbf{X}^T \mathbf{X})^{-1})\end{aligned}$$

Formal Posterior Distribution

With Independent Jeffreys Prior

$$p_{IJ}(\beta, \phi) \propto p_{IJ}(\beta)p_{IJ}(\phi) = \phi^{-1}$$

Formal Posterior Distribution

$$\begin{aligned}\beta \mid \phi, \mathbf{Y} &\sim \mathbf{N}(\hat{\beta}, (\mathbf{X}^T \mathbf{X})^{-1} \phi^{-1}) \\ \phi \mid \mathbf{Y} &\sim \mathbf{G}((n-p)/2, \|\mathbf{Y} - \mathbf{X}\hat{\beta}\|^2/2) \\ \beta \mid \mathbf{Y} &\sim t_{n-p}(\hat{\beta}, \hat{\sigma}^2(\mathbf{X}^T \mathbf{X})^{-1})\end{aligned}$$

Bayesian Credible Sets $p(\beta \in C_\alpha) = 1 - \alpha$ correspond to frequentist Confidence Regions

$$\frac{\lambda^T \beta - \lambda^T \hat{\beta}}{\sqrt{\hat{\sigma}^2 \lambda^T (\mathbf{X}^T \mathbf{X})^{-1} \lambda}} \sim t_{n-p}$$

Disadvantages of Conjugate Priors

Disadvantages:

Disadvantages of Conjugate Priors

Disadvantages:

- ▶ Results may have be sensitive to prior “outliers” due to linear updating

Disadvantages of Conjugate Priors

Disadvantages:

- ▶ Results may have be sensitive to prior “outliers” due to linear updating
- ▶ Cannot capture all possible prior beliefs

Disadvantages of Conjugate Priors

Disadvantages:

- ▶ Results may have be sensitive to prior “outliers” due to linear updating
- ▶ Cannot capture all possible prior beliefs
- ▶ Mixtures of Conjugate Priors