

G-Priors and Mixture Distributions

STA721 Linear Models Duke University

Merlise Clyde

September 30, 2019

Bayesian Estimation

Model

$$\mathbf{Y} \sim \mathcal{N}(\mathbf{X}\boldsymbol{\beta}, \mathbf{I}_n/\phi)$$

with precision $\phi = 1/\sigma^2$.

Bayesian Estimation

Model

$$\mathbf{Y} \sim N(\mathbf{X}\beta, \mathbf{I}_n/\phi)$$

with precision $\phi = 1/\sigma^2$.

Default Prior Choices for β and ϕ :

- ▶ “Non-Informative Priors”: Independent Jeffreys’ Priors (improper)
- ▶ g-prior $N(0, \frac{g}{\phi}(\mathbf{X}^T\mathbf{X})^{-1})$
- ▶ Partitioned g-priors
- ▶ Zellner-Siow Cauchy Prior, mixtures and MCMC (if time)

Readings: Hoff Chapter 9

Zellner's g -prior

$$\text{Zellner's } g\text{-prior(s) } \beta \mid \phi \sim N(\mathbf{b}_0, g(\mathbf{X}^T \mathbf{X})^{-1} / \phi)$$

Zellner's g -prior

Zellner's g -prior(s) $\beta \mid \phi \sim N(\mathbf{b}_0, g(\mathbf{X}^T \mathbf{X})^{-1} / \phi)$

$$\beta \mid \mathbf{Y}, \phi \sim N \left(\frac{g}{1+g} \hat{\beta} + \frac{1}{1+g} \mathbf{b}_0, \frac{g}{1+g} (\mathbf{X}^T \mathbf{X})^{-1} \phi^{-1} \right)$$

Zellner's g -prior

Zellner's g -prior(s) $\beta \mid \phi \sim N(\mathbf{b}_0, g(\mathbf{X}^T \mathbf{X})^{-1} / \phi)$

$$\beta \mid \mathbf{Y}, \phi \sim N \left(\frac{g}{1+g} \hat{\beta} + \frac{1}{1+g} \mathbf{b}_0, \frac{g}{1+g} (\mathbf{X}^T \mathbf{X})^{-1} \phi^{-1} \right)$$

Invariance: Require posterior of $\mathbf{X}\beta$ equal the posterior of $\mathbf{X}\mathbf{H}\alpha$

Zellner's g -prior


Zellner's g -prior(s) $\beta \mid \phi \sim N(\mathbf{b}_0, g(\mathbf{X}^T \mathbf{X})^{-1} / \phi)$

$$\beta \mid \mathbf{Y}, \phi \sim N \left(\frac{g}{1+g} \hat{\beta} + \frac{1}{1+g} \mathbf{b}_0, \frac{g}{1+g} (\mathbf{X}^T \mathbf{X})^{-1} \phi^{-1} \right)$$

Invariance: Require posterior of $\mathbf{X}\beta$ equal the posterior of $\mathbf{X}\mathbf{H}\alpha$
($\mathbf{a}_0 = \mathbf{H}^{-1}\mathbf{b}_0$)

Shrinkage

Posterior mean under g -prior with $\mathbf{b}_0 = 0$ $\frac{g}{1+g}\hat{\beta}$



Choice of g

- ▶ $\frac{g}{1+g}$ weight given to the data

Choice of g

- ▶ $\frac{g}{1+g}$ weight given to the data
- ▶ Fixed g effect does not vanish as $n \rightarrow \infty$ (asymptotic bias)

Choice of g

- ▶ $\frac{g}{1+g}$ weight given to the data
- ▶ Fixed g effect does not vanish as $n \rightarrow \infty$ (asymptotic bias)
- ▶ Use $g = n$ or place a prior distribution on g

Partitioned Zellner's g -prior

Zellner recognized that some parameters might have less information

$$\mathbf{Y} = \mathbf{X}_0\boldsymbol{\beta}_0 + \mathbf{X}_1\boldsymbol{\beta}_1 + \boldsymbol{\epsilon}$$

- ▶ $\mathbf{X}_0^T\mathbf{X}_1 = \mathbf{0}$ (orthogonal columns)

Partitioned Zellner's g -prior

Zellner recognized that some parameters might have less information

$$\mathbf{Y} = \mathbf{X}_0\boldsymbol{\beta}_0 + \mathbf{X}_1\boldsymbol{\beta}_1 + \boldsymbol{\epsilon}$$

- ▶ $\mathbf{X}_0^T \mathbf{X}_1 = \mathbf{0}$ (orthogonal columns)
- ▶ Fisher information block diagonal

Partitioned Zellner's g -prior

Zellner recognized that some parameters might have less information

$$\mathbf{Y} = \mathbf{X}_0\boldsymbol{\beta}_0 + \mathbf{X}_1\boldsymbol{\beta}_1 + \boldsymbol{\epsilon}$$

- ▶ $\mathbf{X}_0^T\mathbf{X}_1 = \mathbf{0}$ (orthogonal columns)
- ▶ Fisher information block diagonal
- ▶ $\boldsymbol{\beta}_0 \sim N(\mathbf{b}_0, g_0(\mathbf{X}_0^T\mathbf{X}_0)^{-1}/\phi)$

Partitioned Zellner's g -prior

Zellner recognized that some parameters might have less information

$$\mathbf{Y} = \mathbf{X}_0\boldsymbol{\beta}_0 + \mathbf{X}_1\boldsymbol{\beta}_1 + \boldsymbol{\epsilon}$$

- ▶ $\mathbf{X}_0^T\mathbf{X}_1 = \mathbf{0}$ (orthogonal columns)
- ▶ Fisher information block diagonal
- ▶ $\boldsymbol{\beta}_0 \sim N(\mathbf{b}_0, g_0(\mathbf{X}_0^T\mathbf{X}_0)^{-1}/\phi)$
- ▶ $\boldsymbol{\beta}_1 \sim N(\mathbf{b}_1, g_1(\mathbf{X}_1^T\mathbf{X}_1)^{-1}/\phi)$

Partitioned Zellner's g -prior

Zellner recognized that some parameters might have less information

$$\mathbf{Y} = \mathbf{X}_0\boldsymbol{\beta}_0 + \mathbf{X}_1\boldsymbol{\beta}_1 + \boldsymbol{\epsilon}$$

- ▶ $\mathbf{X}_0^T\mathbf{X}_1 = \mathbf{0}$ (orthogonal columns)
- ▶ Fisher information block diagonal
- ▶ $\boldsymbol{\beta}_0 \sim N(\mathbf{b}_0, g_0(\mathbf{X}_0^T\mathbf{X}_0)^{-1}/\phi)$
- ▶ $\boldsymbol{\beta}_1 \sim N(\mathbf{b}_1, g_1(\mathbf{X}_1^T\mathbf{X}_1)^{-1}/\phi)$
- ▶ $p(\phi) \propto 1/\phi$

Special case $\mathbf{X}_0 = \mathbf{1}_n$ and let $g_0 \rightarrow \infty$

$$p(\boldsymbol{\beta}_0, \phi) \propto 1/\phi$$

Bayesian Estimation with g prior

$$\mathbf{Y} = \mathbf{1}\alpha_0 + \mathbf{X}_1\boldsymbol{\beta} + \boldsymbol{\epsilon}$$

$$p(\alpha_0, \phi) \propto 1$$

$$\boldsymbol{\beta} \mid \phi \sim \text{N}(\mathbf{0}, \frac{g}{\phi}(\mathbf{X}^T(\mathbf{I}_n - \mathbf{P}_1)\mathbf{X})^{-1})$$

Bayesian Estimation with g prior

$$\begin{aligned}\mathbf{Y} &= \mathbf{1}\alpha_0 + \mathbf{X}_1\boldsymbol{\beta} + \boldsymbol{\epsilon} \\ p(\alpha_0, \phi) &\propto 1 \\ \boldsymbol{\beta} \mid \phi &\sim \text{N}(\mathbf{0}, \frac{g}{\phi}(\mathbf{X}^T(\mathbf{I}_n - \mathbf{P}_1)\mathbf{X})^{-1})\end{aligned}$$

Equivalent to

$$\begin{aligned}\mathbf{Y} &= \mathbf{1}\beta_0 + (\mathbf{I}_n - \mathbf{P}_1)\mathbf{X}_1\boldsymbol{\beta} + \boldsymbol{\epsilon} \\ \beta_0 &= \alpha + \bar{\mathbf{x}}^T\boldsymbol{\beta} \\ p(\beta_0, \phi) &\propto 1 \\ \boldsymbol{\beta} \mid \phi &\sim \text{N}(\mathbf{0}, \frac{g}{\phi}(\mathbf{X}^T(\mathbf{I}_n - \mathbf{P}_1)\mathbf{X})^{-1})\end{aligned}$$

Prior Data

Note

$$(\mathbf{X}^T(\mathbf{I}_n - \mathbf{P}_1)\mathbf{X}) = (\mathbf{X}^T(\mathbf{I}_n - \mathbf{P}_1)^T(\mathbf{I}_n - \mathbf{P}_1)\mathbf{X}) = (\mathbf{X} - \mathbf{1}_n\bar{\mathbf{X}}^T)^T(\mathbf{X} - \mathbf{1}_n\bar{\mathbf{X}})$$

Prior Data

Note

$$(\mathbf{X}^T(\mathbf{I}_n - \mathbf{P}_1)\mathbf{X}) = (\mathbf{X}^T(\mathbf{I}_n - \mathbf{P}_1)^T(\mathbf{I}_n - \mathbf{P}_1)\mathbf{X}) = (\mathbf{X} - \mathbf{1}_n\bar{\mathbf{X}}^T)^T(\mathbf{X} - \mathbf{1}_n\bar{\mathbf{X}})$$

$$\text{Let } (\mathbf{X} - \mathbf{1}_n\bar{\mathbf{X}}^T)^T(\mathbf{X} - \mathbf{1}_n\bar{\mathbf{X}}) = SS_{\mathbf{X}} = \mathbf{U}^T\mathbf{U}$$

Prior Data

Note

$$(\mathbf{X}^T(\mathbf{I}_n - \mathbf{P}_1)\mathbf{X}) = (\mathbf{X}^T(\mathbf{I}_n - \mathbf{P}_1)^T(\mathbf{I}_n - \mathbf{P}_1)\mathbf{X}) = (\mathbf{X} - \mathbf{1}_n\bar{\mathbf{X}}^T)^T(\mathbf{X} - \mathbf{1}_n\bar{\mathbf{X}})$$

$$\text{Let } (\mathbf{X} - \mathbf{1}_n\bar{\mathbf{X}}^T)^T(\mathbf{X} - \mathbf{1}_n\bar{\mathbf{X}}) = SS_{\mathbf{X}} = \mathbf{U}^T\mathbf{U}$$

Quadratic contribution to the log likelihood from prior after integrating out β_0

$$(\mathbf{Y}_c - \mathbf{X}_c\beta)^T(\mathbf{Y}_c - \mathbf{X}_c\beta) + (\beta^T \frac{\mathbf{U}^T\mathbf{U}}{g} \beta)$$

Prior Data

Note

$$(\mathbf{X}^T(\mathbf{I}_n - \mathbf{P}_1)\mathbf{X}) = (\mathbf{X}^T(\mathbf{I}_n - \mathbf{P}_1)^T(\mathbf{I}_n - \mathbf{P}_1)\mathbf{X}) = (\mathbf{X} - \mathbf{1}_n\bar{\mathbf{X}}^T)^T(\mathbf{X} - \mathbf{1}_n\bar{\mathbf{X}})$$

$$\text{Let } (\mathbf{X} - \mathbf{1}_n\bar{\mathbf{X}}^T)^T(\mathbf{X} - \mathbf{1}_n\bar{\mathbf{X}}) = SS_{\mathbf{X}} = \mathbf{U}^T\mathbf{U}$$

Quadratic contribution to the log likelihood from prior after integrating out β_0

$$(\mathbf{Y}_c - \mathbf{X}_c\beta)^T(\mathbf{Y}_c - \mathbf{X}_c\beta) + (\beta^T \frac{\mathbf{U}^T\mathbf{U}}{g}\beta)$$

$$(\mathbf{Y}_c - \mathbf{X}_c\beta)^T(\mathbf{Y}_c - \mathbf{X}_c\beta) + (\mathbf{0}_p - \frac{\mathbf{U}}{\sqrt{g}}\beta)^T(\mathbf{0}_p - \frac{\mathbf{U}}{\sqrt{g}}\beta)$$

Prior observations with $Y_c = 0$.

Example: $g=5$, $n=30$


In SLR it is like an extra $Y_0 = 0$ at $\mathbf{X}_o = \sqrt{\frac{SS_x}{g}}$:

$$(\mathbf{Y}_c - \mathbf{X}_c\beta)^T(\mathbf{Y}_c - \mathbf{X}_c\beta) + (0 - \sqrt{\frac{SS_x}{g}}\beta)^T(0 - \sqrt{\frac{SS_x}{g}}\beta)$$

Example: $g=5$, $n=30$

In SLR it is like an extra $Y_0 = 0$ at $\mathbf{X}_0 = \sqrt{\frac{SS_x}{g}}$:

$$(\mathbf{Y}_c - \mathbf{X}_c\beta)^T(\mathbf{Y}_c - \mathbf{X}_c\beta) + (0 - \sqrt{\frac{SS_x}{g}}\beta)^T(0 - \sqrt{\frac{SS_x}{g}}\beta)$$



Disadvantages of Conjugate Priors

Disadvantages:

Disadvantages of Conjugate Priors

Disadvantages:

- ▶ Results may have be sensitive to prior “outliers” due to linear updating

Disadvantages of Conjugate Priors

Disadvantages:

- ▶ Results may have be sensitive to prior “outliers” due to linear updating
- ▶ Problem potentially with all Normal priors, not just the g -prior.

Disadvantages of Conjugate Priors

Disadvantages:

- ▶ Results may have be sensitive to prior “outliers” due to linear updating
- ▶ Problem potentially with all Normal priors, not just the g -prior.
- ▶ Cannot capture all possible prior beliefs

Disadvantages of Conjugate Priors

Disadvantages:

- ▶ Results may have be sensitive to prior “outliers” due to linear updating
- ▶ Problem potentially with all Normal priors, not just the g -prior.
- ▶ Cannot capture all possible prior beliefs
- ▶ Mixtures of Conjugate Priors

Mixtures of Conjugate Priors

Theorem (Diaconis & Ylvisaker 1985)

Given a sampling model $p(y \mid \theta)$ from an exponential family, any prior distribution can be expressed as a mixture of conjugate prior distributions

► Prior $p(\theta) = \int p(\theta \mid \omega) p(\omega) d\omega$

Mixtures of Conjugate Priors

Theorem (Diaconis & Ylvisaker 1985)

Given a sampling model $p(y \mid \theta)$ from an exponential family, any prior distribution can be expressed as a mixture of conjugate prior distributions

- ▶ Prior $p(\theta) = \int p(\theta \mid \omega) p(\omega) d\omega$
- ▶ Posterior

Mixtures of Conjugate Priors

Theorem (Diaconis & Ylvisaker 1985)

Given a sampling model $p(y \mid \theta)$ from an exponential family, any prior distribution can be expressed as a mixture of conjugate prior distributions

- ▶ Prior $p(\theta) = \int p(\theta \mid \omega)p(\omega) d\omega$
- ▶ Posterior

$$p(\theta \mid \mathbf{Y}) \propto \int p(\mathbf{Y} \mid \theta)p(\theta \mid \omega)p(\omega) d\omega$$

Mixtures of Conjugate Priors

Theorem (Diaconis & Ylvisaker 1985)

Given a sampling model $p(y \mid \theta)$ from an exponential family, any prior distribution can be expressed as a mixture of conjugate prior distributions

- ▶ Prior $p(\theta) = \int p(\theta \mid \omega)p(\omega) d\omega$
- ▶ Posterior

$$\begin{aligned} p(\theta \mid \mathbf{Y}) &\propto \int p(\mathbf{Y} \mid \theta)p(\theta \mid \omega)p(\omega) d\omega \\ &\propto \int \frac{p(\mathbf{Y} \mid \theta)p(\theta \mid \omega)}{p(\mathbf{Y} \mid \omega)}p(\mathbf{Y} \mid \omega)p(\omega) d\omega \end{aligned}$$

Mixtures of Conjugate Priors

Theorem (Diaconis & Ylvisaker 1985)

Given a sampling model $p(y \mid \theta)$ from an exponential family, any prior distribution can be expressed as a mixture of conjugate prior distributions

- ▶ Prior $p(\theta) = \int p(\theta \mid \omega)p(\omega) d\omega$
- ▶ Posterior

$$\begin{aligned} p(\theta \mid \mathbf{Y}) &\propto \int p(\mathbf{Y} \mid \theta)p(\theta \mid \omega)p(\omega) d\omega \\ &\propto \int \frac{p(\mathbf{Y} \mid \theta)p(\theta \mid \omega)}{p(\mathbf{Y} \mid \omega)} p(\mathbf{Y} \mid \omega)p(\omega) d\omega \\ &\propto \int p(\theta \mid \mathbf{Y}, \omega)p(\mathbf{Y} \mid \omega)p(\omega) d\omega \end{aligned}$$

Mixtures of Conjugate Priors

Theorem (Diaconis & Ylvisaker 1985)

Given a sampling model $p(y \mid \theta)$ from an exponential family, any prior distribution can be expressed as a mixture of conjugate prior distributions

- ▶ Prior $p(\theta) = \int p(\theta \mid \omega)p(\omega) d\omega$
- ▶ Posterior

$$\begin{aligned} p(\theta \mid \mathbf{Y}) &\propto \int p(\mathbf{Y} \mid \theta)p(\theta \mid \omega)p(\omega) d\omega \\ &\propto \int \frac{p(\mathbf{Y} \mid \theta)p(\theta \mid \omega)}{p(\mathbf{Y} \mid \omega)} p(\mathbf{Y} \mid \omega)p(\omega) d\omega \\ &\propto \int p(\theta \mid \mathbf{Y}, \omega)p(\mathbf{Y} \mid \omega)p(\omega) d\omega \\ p(\theta \mid \mathbf{Y}) &= \frac{\int p(\theta \mid \mathbf{Y}, \omega)p(\mathbf{Y} \mid \omega)p(\omega) d\omega}{\int p(\mathbf{Y} \mid \omega)p(\omega) d\omega} \end{aligned}$$

Zellner-Siow prior (assume \mathbf{X} is centered)

Zellner's g-prior $\beta \mid \phi \sim N(\mathbf{0}_p, g(\mathbf{X}^T \mathbf{X})^{-1} / \phi)$

Zellner-Siow prior (assume \mathbf{X} is centered)

Zellner's g-prior $\beta \mid \phi \sim \mathcal{N}(\mathbf{0}_p, g(\mathbf{X}^T \mathbf{X})^{-1} / \phi)$

- Choice of g ?

Zellner-Siow prior (assume \mathbf{X} is centered)

Zellner's g-prior $\beta \mid \phi \sim \mathcal{N}(\mathbf{0}_p, g(\mathbf{X}^T \mathbf{X})^{-1} / \phi)$

- ▶ Choice of g ?
- ▶ $\frac{g}{1+g}$ weight given to the data

Zellner-Siow prior (assume \mathbf{X} is centered)

Zellner's g-prior $\beta \mid \phi \sim N(\mathbf{0}_p, g(\mathbf{X}^T \mathbf{X})^{-1} / \phi)$

- ▶ Choice of g ?
- ▶ $\frac{g}{1+g}$ weight given to the data
- ▶ Let $\tau = 1/g$ assign $\tau \sim G(1/2, n/2)$

Zellner-Siow prior (assume \mathbf{X} is centered)

Zellner's g-prior $\beta \mid \phi \sim N(\mathbf{0}_p, g(\mathbf{X}^T \mathbf{X})^{-1} / \phi)$

- ▶ Choice of g ?
- ▶ $\frac{g}{1+g}$ weight given to the data
- ▶ Let $\tau = 1/g$ assign $\tau \sim G(1/2, n/2)$
- ▶ Marginal prior on $\beta \sim C(0, \phi^{-1} \mathbf{X}^T \mathbf{X})$
- ▶ Can express posterior as a mixture of g-priors

$$p(\tau \mid \mathbf{Y}) = \frac{p(\mathbf{Y} \mid \tau) p(\tau)}{\int p(\mathbf{Y} \mid \tau) p(\tau) d\tau}$$

Zellner-Siow prior (assume \mathbf{X} is centered)

Zellner's g-prior $\beta \mid \phi \sim N(\mathbf{0}_p, g(\mathbf{X}^T \mathbf{X})^{-1} / \phi)$

- ▶ Choice of g ?
- ▶ $\frac{g}{1+g}$ weight given to the data
- ▶ Let $\tau = 1/g$ assign $\tau \sim G(1/2, n/2)$
- ▶ Marginal prior on $\beta \sim C(0, \phi^{-1} \mathbf{X}^T \mathbf{X})$
- ▶ Can express posterior as a mixture of g-priors

$$p(\tau \mid \mathbf{Y}) = \frac{p(\mathbf{Y} \mid \tau) p(\tau)}{\int p(\mathbf{Y} \mid \tau) p(\tau) d\tau}$$

- ▶ Problem: no analytic expression for integral

Zellner-Siow prior (assume \mathbf{X} is centered)

Zellner's g-prior $\beta \mid \phi \sim N(\mathbf{0}_p, g(\mathbf{X}^T \mathbf{X})^{-1} / \phi)$

- ▶ Choice of g ?
- ▶ $\frac{g}{1+g}$ weight given to the data
- ▶ Let $\tau = 1/g$ assign $\tau \sim G(1/2, n/2)$
- ▶ Marginal prior on $\beta \sim C(0, \phi^{-1} \mathbf{X}^T \mathbf{X})$
- ▶ Can express posterior as a mixture of g-priors

$$p(\tau \mid \mathbf{Y}) = \frac{p(\mathbf{Y} \mid \tau) p(\tau)}{\int p(\mathbf{Y} \mid \tau) p(\tau) d\tau}$$

- ▶ Problem: no analytic expression for integral
- ▶ Need 2 one dimensional integrals to obtain posterior.

Zellner-Siow prior (assume \mathbf{X} is centered)

Zellner's g-prior $\beta \mid \phi \sim N(\mathbf{0}_p, g(\mathbf{X}^T \mathbf{X})^{-1} / \phi)$

- ▶ Choice of g ?
- ▶ $\frac{g}{1+g}$ weight given to the data
- ▶ Let $\tau = 1/g$ assign $\tau \sim G(1/2, n/2)$
- ▶ Marginal prior on $\beta \sim C(0, \phi^{-1} \mathbf{X}^T \mathbf{X})$
- ▶ Can express posterior as a mixture of g-priors

$$p(\tau \mid \mathbf{Y}) = \frac{p(\mathbf{Y} \mid \tau) p(\tau)}{\int p(\mathbf{Y} \mid \tau) p(\tau) d\tau}$$

- ▶ Problem: no analytic expression for integral
- ▶ Need 2 one dimensional integrals to obtain posterior.
- ▶ What about credible intervals?

Markov Chain Monte Carlo

- ▶ We know that $\beta_0, \beta, \phi \mid \mathbf{Y}, g = 1/\tau$ has a Normal-Gamma distribution

Markov Chain Monte Carlo

- ▶ We know that $\beta_0, \beta, \phi \mid \mathbf{Y}, g = 1/\tau$ has a Normal-Gamma distribution
- ▶ We can show that $\tau \mid \beta_0, \beta, \phi, \mathbf{Y}$ has a Gamma distribution

Markov Chain Monte Carlo

- ▶ We know that $\beta_0, \beta, \phi \mid \mathbf{Y}, g = 1/\tau$ has a Normal-Gamma distribution
- ▶ We can show that $\tau \mid \beta_0, \beta, \phi, \mathbf{Y}$ has a Gamma distribution

$$p(\tau \mid \beta, \phi, \mathbf{Y}) \propto \mathcal{L}(\beta_0, \beta, \phi) \tau^{p/2} e^{(-\tau \frac{\phi}{2} \beta^T (\mathbf{X}^T \mathbf{X}) \beta)} \tau^{1/2-1} e^{-\tau n/2}$$

Markov Chain Monte Carlo

- ▶ We know that $\beta_0, \beta, \phi \mid \mathbf{Y}, g = 1/\tau$ has a Normal-Gamma distribution
- ▶ We can show that $\tau \mid \beta_0, \beta, \phi, \mathbf{Y}$ has a Gamma distribution

$$p(\tau \mid \beta, \phi, \mathbf{Y}) \propto \mathcal{L}(\beta_0, \beta, \phi) \tau^{p/2} e^{(-\tau \frac{\phi}{2} \beta^T (\mathbf{X}^T \mathbf{X}) \beta)} \tau^{1/2-1} e^{-\tau n/2}$$

- ▶ alternate sampling from full conditional distributions given current values of other parameters. (STA 601)

Markov Chain Monte Carlo

- ▶ We know that $\beta_0, \beta, \phi \mid \mathbf{Y}, g = 1/\tau$ has a Normal-Gamma distribution
- ▶ We can show that $\tau \mid \beta_0, \beta, \phi, \mathbf{Y}$ has a Gamma distribution

$$p(\tau \mid \beta, \phi, \mathbf{Y}) \propto \mathcal{L}(\beta_0, \beta, \phi) \tau^{p/2} e^{(-\tau \frac{\phi}{2} \beta^T (\mathbf{X}^T \mathbf{X}) \beta)} \tau^{1/2-1} e^{-\tau n/2}$$

- ▶ alternate sampling from full conditional distributions given current values of other parameters. (STA 601)
- ▶ JAGS or STAN

JAGS Code: library(R2jags)

```
model = function(){  
  for (i in 1:n) {  
    Y[i] ~ dnorm(beta0+ (X[i] -Xbar)*beta, phi)  
  }  
  beta0 ~ dnorm(0, .000001*phi) #precision is 2nd arg  
  beta ~ dnorm(0, phi*tau*SSX) #precision is 2nd arg  
  phi ~ dgamma(.001, .001)  
  tau ~ dgamma(.5, .5*n)  
  g <- 1/tau  
  sigma <- pow(phi, -.5)  
}  
data = list(Y=Y, X=X, n=length(Y), SSX=sum(Xc^2),  
            Xbar=mean(X))  
ZSout = jags(data, inits=NULL,  
             parameters.to.save=c("beta0", "beta", "g",  
                                  "sigma"),  
             model=model, n.iter=10000)
```

HPD intervals


```
confint(lm(Y ~ Xc))
```

```
##                2.5 %    97.5 %  
## (Intercept) -0.3985359 0.2048303  
## Xc           2.7945824 3.4555162
```

```
HPDinterval(as.mcmc(ZSout$BUGSoutput$sims.matrix))
```

```
##                lower      upper  
## beta           2.7823047    3.4453690  
## beta0          -0.3764027    0.2095465  
## deviance       70.2043917    78.4813041  
## g              19.4503373 3782.7134974  
## sigma          0.6171029    1.0504892  
## attr(,"Probability")  
## [1] 0.95
```

Compare



ZSout

```
## Inference for Bugs model at "/var/folders/n4/nj1122xj6bn5_xgbptv7bml40000gp/T//Rtmpf0Ltcz/modeld51989a
## 3 chains, each with 10000 iterations (first 5000 discarded), n.thin = 5
## n.sims = 3000 iterations saved
##           mu.vect   sd.vect   2.5%    25%    50%    75%    97.5%  Rhat
## beta          3.112     0.170   2.782   2.997   3.115   3.225   3.445  1.001
## beta0         -0.099     0.152  -0.384  -0.204  -0.099   0.001   0.204  1.002
## g            2263.147 38967.029 48.273 146.129 282.298 697.063 9018.709 1.001
## sigma          0.827     0.114   0.636   0.747   0.816   0.896   1.079  1.001
## deviance       73.347     2.563  70.390  71.458  72.680  74.500  79.882  1.002
##           n.eff
## beta          3000
## beta0         1200
## g             3000
## sigma         3000
## deviance      1600
##
## For each parameter, n.eff is a crude measure of effective sample size,
## and Rhat is the potential scale reduction factor (at convergence, Rhat=1).
##
## DIC info (using the rule, pD = var(deviance)/2)
## pD = 3.3 and DIC = 76.6
## DIC is an estimate of expected predictive error (lower deviance is better).
```

Cauchy Summary

- ▶ Cauchy rejects prior mean if it is an "outlier"
- ▶ robustness related to "bounded" influence (more later)
- ▶ numerical integration or Monte Carlo sampling (MCMC)