

Vision: From Photon to Mind

A reference walk through the eye in three parts: (I) the human eye — optical anatomy, retinal histology, receptor physiology and biochemistry; (II) the physics of light that underlies it; (III) outward across species to the electromagnetic window, alternative life forms, and the evolution of sight.

Alexandru C. Stan

© 2026 Alexandru C. Stan

Vision: From Photon to Mind — a reference in three parts. Version 1.0.

Licensed under the Creative Commons Attribution 4.0 International Licence (CC BY 4.0).

Author ORCID: 0000-0002-7463-0393

Set in STIX Two Text.

Contents

Part I — The Human Eye: Structure to Signal.....	1
A. Optical Anatomy — The Eye as a Refractive Instrument.....	1
B. Retinal Histology — The Layered Neural Sheet.....	5
C. Receptor Physiology — Generating and Shaping the Signal.....	7
D. Biochemistry — Pigments and the Visual Cycle.....	10
Part II — The Physics of Light: The Bridge.....	12
A. What Light Is — Wave and Photon.....	12
B. The Electromagnetic Spectrum — One Continuum.....	12
C. Field Structure — Why Radio Is Not "Simpler" Than Light.....	14
D. Polarisation — The Channel We Discard.....	14
E. Light Meets Matter — Refraction, Absorption, and the Index of Refraction.....	15
F. Diffraction and the Aperture Limit — Why "Eye Size vs Wavelength" Decides Everything.....	16
Part III — Comparative Vision: The Same Window, Different Engines.....	17
Optics across species — focusing in wet matter.....	17
Retinal architecture across species.....	18
Receptor physiology & chromophore chemistry across species.....	18
1. The Human Photoreceptors.....	19
2. Why Photoreceptors Respond to Light.....	19
3. The Electromagnetic Spectrum.....	20
4. Field Structure & Polarisation.....	21
5. Seeing Outside the Visible — Radio, UV, X-ray.....	21
6. Eyes Around Other Stars.....	23
7. A Mind Built on Polarisation.....	23
8. Earth's Existence-Proofs — Eyes That Broke the Rules.....	24
9. Colour as Inference vs Colour as Barcode.....	25
10. How Vision Got Started — One Molecule to a Focusing Eye.....	25
11. The Inverted Retina — A Flaw, and a Brilliant Patch.....	27
The Through-Line.....	28
Coda — The Stargate: Toward the Central Visual Pathway.....	30
Glossary & Abbreviations.....	32
References & Further Reading.....	34
Key primary literature (cited in text).....	34
Standard reference works by domain.....	34

Figures

1. The eye as a refractive instrument.....	2
2. Dioptric power & accommodation.....	4
3. The ten retinal layers.....	6
4. Phototransduction cascade.....	9
5. Dark adaptation.....	10
6. Human photoreceptor absorption spectra.....	11
7. The electromagnetic spectrum.....	13
8. The eye-evolution ladder.....	26
9. Vertebrate vs cephalopod retina — inverted vs verted.....	28

About this document. It traces a single continuous argument — from why a retinal molecule absorbs a photon, through the optical and neural machinery that turns that absorption into perception, out to how the same physics constrains vision across animals, habitats, and hypothetical worlds. **Part I** is reference-grade human anatomy, histology, physiology, and biochemistry. **Part II** assembles the physics of light that the rest depends on. **Part III** takes that substrate comparative and evolutionary, ending at the "flawed-but-brilliant" inverted vertebrate retina. It is written to be read start-to-finish *or* dipped into by section; a glossary and references support the latter. Quantitative figures are standard textbook values; claims resting on specific studies are cited. (*The central visual pathway and cortical processing — LGN, V1, the magno/parvo/konio streams — are deliberately out of scope, reserved for the companion volume.*)

Part I — The Human Eye: Structure to Signal

Reference-grade anatomy, histology, physiology, and biochemistry — the substrate on which Parts II and III build.

A. Optical Anatomy — The Eye as a Refractive Instrument

Gross dioptrics

The relaxed emmetropic eye has a total refractive power of **~+58 to +60 dioptres (D)**, focusing parallel rays onto the retina across an axial length of **~24 mm**. The power is not evenly distributed:

Element	Power	Share	Notes
Cornea (anterior surface dominant)	~+43 D	~2/3	The eye's principal refracting surface — air→tissue interface gives the largest index step
Crystalline lens (relaxed)	~+19–20 D	~1/3	Variable; the <i>adjustable</i> element
<i>Combined (relaxed)</i>	~+59 D		Two thick lenses in series; combined power < arithmetic sum

The largest single refraction happens at the **anterior corneal surface** because that is where the refractive-index step is greatest (air $n \approx 1.00 \rightarrow$ cornea $n \approx 1.376$). The lens, sitting between aqueous ($n \approx 1.336$) and vitreous ($n \approx 1.336$), works across much smaller index steps and so contributes less *despite* being a "lens" in the everyday sense. This is why corneal procedures (LASIK/PRK) and corneal disease have outsized optical impact.




Figure 1. The eye as a refractive instrument. The cornea supplies roughly two-thirds of the eye's $\sim +59\text{ D}$, because the air→cornea interface is the largest refractive-index step in the system; the lens, bathed in fluids of similar index, contributes less despite being the adjustable element.

The refractive elements, anterior to posterior

Tear film ($\sim 7\ \mu\text{m}$) — the true first optical surface. Three layers: outer lipid (meibomian, retards evaporation), middle aqueous (lacrimal), inner mucin (goblet-cell, wets the hydrophobic epithelium). A smooth tear film is optically essential; break-up degrades the air-tissue interface that does most of the focusing.

Cornea ($\sim 0.5\text{ mm}$ central, $\sim 0.7\text{ mm}$ peripheral; avascular, transparent). Five classic layers:

1. **Epithelium** — stratified squamous non-keratinised, $\sim 5\text{--}6$ cell layers, rapid turnover from limbal stem cells.
2. **Bowman's layer** — acellular condensed stroma; does not regenerate.
3. **Stroma** ($\sim 90\%$ of thickness) — $\sim 200\text{--}250$ orthogonally stacked lamellae of type I collagen in highly regular spacing; transparency depends on this lattice regularity (Maurice's lattice theory) and on **relative dehydration** maintained by the endothelial pump.
4. **Descemet's membrane** — basement membrane of the endothelium; thickens with age.
5. **Endothelium** — single layer of hexagonal cells, **non-dividing in humans**; Na^+/K^+ -ATPase pump keeps the stroma deturgescd. Cell loss (surgery, Fuchs dystrophy) → stromal oedema → loss of transparency.

Clinical correlate — cornea. Because the cornea supplies $\sim 2/3$ of refractive power, corneal shape and clarity dominate image quality. **Keratoconus** (progressive stromal thinning/ectasia) distorts the principal refracting surface $\rightarrow$ irregular astigmatism. **Endothelial failure** (Fuchs dystrophy, post-surgical cell loss) breaks deturgescence $\rightarrow$ stromal oedema and haze. The avascular, immune-privileged stroma is also why corneal grafts are among the most successful transplants.

Anterior chamber & aqueous humour — $n \approx 1.336$; secreted by the ciliary epithelium, circulates from posterior chamber $\rightarrow$ pupil $\rightarrow$ anterior chamber $\rightarrow$ trabecular meshwork $\rightarrow$ Schlemm's canal (conventional, $\sim 80\text{--}90\%$) and uveoscleral outflow. Determines **intraocular pressure (IOP)**; the production–outflow balance is the substrate of glaucoma.

Clinical correlate — aqueous & IOP. Glaucoma is an optic neuropathy in which (often) raised IOP from impaired trabecular outflow damages retinal ganglion cell axons at the lamina cribrosa — the inner retina's output, lost from the periphery inward. *Open-angle*: gradual outflow resistance. *Angle-closure*: the iris mechanically blocks the trabecular meshwork — an emergency. The production–outflow balance, not production alone, sets the pressure.

Crystalline lens — biconvex, avascular, transparent, suspended by **zonules of Zinn** from the ciliary body. Structure: anterior epithelium (only anterior; cells migrate to the equator and elongate into fibres), a **gradient refractive index** ($n \sim 1.386$ surface $\rightarrow \sim 1.406$ nucleus — the gradient corrects spherical aberration, exactly the principle reused by fish/cephalopod lenses), enclosed in an elastic **capsule**. Lens fibres are added lifelong (no cell loss $\rightarrow$ nucleus compacts), explaining presbyopia and cataract.

Clinical correlate — lens. Lifelong fibre addition with no cell loss makes the lens uniquely prone to two age changes: **presbyopia** (progressive stiffening $\rightarrow$ loss of accommodative amplitude, beginning ~ 40 s) and **cataract** (loss of transparency as crystallin proteins aggregate/oxidise — the leading global cause of reversible blindness, treated by phacoemulsification + intraocular lens). The gradient-index optics are also why nuclear cataract can transiently *increase* lens power ("myopic shift").

Dioptric power and accommodation




Figure 2. Left: where the focusing happens — the cornea's ~+43 D dwarfs the relaxed lens's ~+19 D. Right: accommodative amplitude collapses with age as the lens stiffens (presbyopia), crossing below the ~3 D needed for comfortable reading around the mid-40s.

Vitreous body — ~80% of globe volume, ~99% water in a type II collagen + hyaluronan gel; optically near-empty but mechanically supportive; liquefies (syneresis) with age → floaters, posterior vitreous detachment.

Accommodation (Helmholtz mechanism)

At rest, zonular tension flattens the lens (low power, distance focus). To focus near:

1. **Ciliary muscle** (parasympathetic, CN III, via the ciliary ganglion) **contracts**, narrowing the ciliary ring.
2. **Zonular tension falls.**
3. The elastic **capsule reshapes the lens** to a more spherical form → anterior surface curvature increases → power rises (up to ~+12–14 D in youth).

Counter-intuitively, the muscle *contracts* to *relax* zonular tension. Accommodative amplitude declines with age as the lens stiffens → **presbyopia** (lens sclerosis, not muscle failure). The **near triad/accommodation reflex**: accommodation + convergence (medial recti) + miosis (pupillary constriction, increasing depth of focus).

Pupil and aperture

The **iris** is the aperture stop. **Sphincter pupillae** (circumferential, parasympathetic) → miosis; **dilator pupillae** (radial, sympathetic) → mydriasis. Pupil diameter (~2–8 mm) trades **light flux vs depth of focus vs aberration**: a small pupil improves depth of focus and cuts peripheral aberration but increases diffraction; a large pupil gathers light but exposes lens/corneal aberration. The optimal aperture for acuity is ~2.5–3 mm.

Aberrations and limits

- **Spherical aberration** — peripheral rays focus shorter; partly corrected by the corneal asphericity (prolate cornea) and the lens gradient index.

- **Chromatic aberration** — ~ 2 D across the visible spectrum (blue focuses anterior to red); the eye is not achromatised. Tolerated partly because foveal S-cones are sparse.
- **Diffraction limit** — at a $\sim 2\text{--}3$ mm pupil, the Airy disc ($\theta \approx 1.22 \lambda/D$) is matched to foveal cone spacing ($\sim 2.5 \mu\text{m}$ centre-to-centre), so the optics and the sampling array are co-tuned: acuity (~ 1 arcmin, 20/20) sits near the point where finer optics would out-resolve the cone mosaic anyway.
- **Stiles–Crawford effect** — light entering the pupil centre is more effective at exciting cones than obliquely-entering light (cones act as waveguides pointing at the pupil), improving image quality and suppressing stray light.

B. Retinal Histology — The Layered Neural Sheet

The retina is **inverted** (light traverses the neural layers before reaching photoreceptors at the back — see §11) and develops as a diverticulum of the diencephalon: it *is* displaced brain. Ten layers, outer (scleral) to inner (vitreal):

1. **Retinal pigment epithelium (RPE)** — single layer of pigmented (melanin) cuboidal cells on Bruch's membrane. Functions: phagocytoses shed photoreceptor outer-segment discs; recycles retinoids (the visual cycle, §D); absorbs stray light; forms the outer blood–retinal barrier; transports nutrients/ions. *Not* a neural layer but functionally inseparable from the photoreceptors.

Clinical correlate — photoreceptor–RPE complex. This metabolically intense interface is the site of the major blinding retinopathies. **Age-related macular degeneration (AMD)** begins with RPE/Bruch's-membrane dysfunction (drusen; geographic atrophy or choroidal neovascularisation). **Retinitis pigmentosa** and allied dystrophies are progressive photoreceptor degenerations (rods first $\rightarrow$ night blindness, then constriction). **Stargardt disease** (*ABCA4*) floods the RPE with toxic A2E. The outer retina's reliance on the choroid also makes it vulnerable to choroidal ischaemia.

2. **Photoreceptor layer** — outer and inner segments of rods and cones.
3. **External (outer) limiting membrane (ELM)** — a row of adherens junctions between Müller cells and photoreceptors; a metabolic/diffusion boundary, not a true membrane.
4. **Outer nuclear layer (ONL)** — cell bodies (nuclei) of rods and cones.
5. **Outer plexiform layer (OPL)** — synapses: photoreceptor terminals (rod spherules, cone pedicles) $\leftrightarrow$ horizontal and bipolar cells. Site of the **triad** synapse and of lateral inhibition by horizontal cells.
6. **Inner nuclear layer (INL)** — cell bodies of **bipolar, horizontal, amacrine, and Müller** cells.
7. **Inner plexiform layer (IPL)** — synapses: bipolar $\leftrightarrow$ amacrine $\leftrightarrow$ ganglion cells; stratified into ON and OFF sublaminae.

8. **Ganglion cell layer (GCL)** — cell bodies of retinal ganglion cells (RGCs) and displaced amacrine.
9. **Nerve fibre layer (NFL)** — unmyelinated RGC axons converging toward the disc; myelinate only after the lamina cribrosa.
10. **Internal limiting membrane (ILM)** — basement membrane of Müller cell endfeet; the retina–vitreal interface.




Figure 3. The ten retinal layers, drawn with light entering from the vitreal side (bottom). Because the retina is inverted, incoming light must cross layers 10 → 3 before reaching the photoreceptors in layer 2, which sit against the RPE at the back. The Müller cell spans the full thickness and partly mitigates this by guiding light [1] (see §11).

The vertical (through) pathway and lateral modulation

Vertical: photoreceptor → bipolar cell → ganglion cell → optic nerve.

Lateral: horizontal cells (in the OPL) and amacrine cells (in the IPL) provide inhibition that builds **centre–surround receptive fields** and motion/timing computations.

- **Bipolar cells** split the signal into **ON** (depolarise to light, sign-inverting mGluR6 / TRPM1 cascade) and **OFF** (hyperpolarise to light, sign-conserving ionotropic AMPA/kainate) channels — the origin of the ON/OFF dichotomy that pervades the visual system.
- **Horizontal cells** mediate surround antagonism and feedback at the first synapse.
- **Amacrine cells** (≥ 30 subtypes; e.g. AII amacrine route rod signals into the cone bipolar pathway in dim light) shape temporal and directional responses.

Glia and vasculature

- **Müller cells** — radial glia spanning ELM→ILM; structural scaffold, K⁺ buffering, neurotransmitter (glutamate→glutamine) recycling, and **living fibre-optic light guides** (see §11). Astrocytes and microglia also present.
- **Dual blood supply**: the **central retinal artery** supplies the inner retina (to the INL); the **choriocapillaris** supplies the outer retina (photoreceptors/RPE) by diffusion across Bruch's membrane. The avascular foveola depends entirely on the choroid — relevant to macular ischaemia and choroidal disease.

Regional specialisation

- **Fovea / foveola** — a ~1.5 mm pit; the foveola (~0.35 mm) is **rod-free, all-cone**, with inner layers displaced radially (the *clivus*) so light reaches cones unobstructed → peak acuity. Each foveal cone drives a near-1:1 line to its own midget bipolar and midget ganglion cell (**private lines**) — minimal convergence = maximal resolution.
- **Macula lutea** — yellow xanthophyll pigment (lutein/zeaxanthin) filtering blue light and damping chromatic aberration.
- **Periphery** — rod-dominated, high convergence (many rods → one ganglion cell) = high sensitivity, low acuity.
- **Optic disc** — the **blind spot**; no photoreceptors, exit point of the NFL and entry of central retinal vessels.

Convergence gradient (the acuity–sensitivity trade)

~120 M rods + ~6 M cones converge onto ~1.2 M ganglion cells. Foveal cones: ~1:1 (resolution). Peripheral rods: up to ~1000:1 (sensitivity by spatial summation). The entire retina is an exercise in spending a fixed ganglion-cell/optic-nerve budget on either resolution or sensitivity, region by region.

C. Receptor Physiology — Generating and Shaping the Signal

Photoreceptor architecture

A photoreceptor is a polarised neuron with four functional compartments:

- **Outer segment (OS)** — a modified cilium packed with membranous **discs** bearing opsin (rods: ~10⁸ rhodopsin molecules per cell; discs are free-floating, continuously shed apically and phagocytosed by RPE). Cone OS discs are continuous infoldings of plasma membrane, exposing pigment to the interphotoreceptor space.
- **Connecting cilium** — narrow bridge; all OS proteins are trafficked through it (mutations → ciliopathies/retinal dystrophies).
- **Inner segment (IS)** — ellipsoid (densely packed mitochondria, supplying the huge ATP demand of the dark current) + myoid (ER/Golgi, biosynthesis).

- **Synaptic terminal** — ribbon synapse (rod spherule / cone pedicle) for graded, tonic glutamate release.

Rods vs cones — functional contrast:

	Rods	Cones
Number	~120 M	~6 M
Sensitivity	Single-photon	~100× less sensitive
Operating range	Scotopic	Photopic
Acuity	Low (convergent)	High (foveal 1:1)
Colour	Monochromatic	Trichromatic
Response kinetics	Slow (long integration)	Fast
Pigment regeneration	Slow (RPE visual cycle)	Fast (RPE + cone-specific Müller cycle)
Saturation	Saturate in daylight	Do not saturate


The dark current and graded signalling

In darkness, **cGMP-gated (CNG) channels** in the OS are held open, admitting an inward $\text{Na}^+/\text{Ca}^{2+}$ "dark current"; the IS Na^+/K^+ -ATPase completes the circuit. The cell rests **depolarised** (~−40 mV) and tonically releases glutamate. Photoreceptors do **not** fire action potentials — they signal in **graded** fashion, encoding light intensity as the *degree* of hyperpolarisation and thus the *reduction* in glutamate release.

Phototransduction (the activation cascade)

1. Photon → **11-cis-retinal isomerises to all-trans** within rhodopsin → active **Metarhodopsin II (R*)** [2].
2. R* catalyses GDP→GTP exchange on **transducin (Gt)**; one R* activates hundreds of Gt (gain stage 1).
3. Gtα-GTP relieves inhibition of **cGMP phosphodiesterase (PDE6)**; each PDE6 hydrolyses many cGMP (gain stage 2).
4. **[cGMP] falls → CNG channels close → membrane hyperpolarises.**
5. Hyperpolarisation **reduces glutamate** release at the ribbon synapse → read by ON (sign-inverting) and OFF (sign-conserving) bipolar cells.

Overall amplification (one photon → tens–hundreds of closed channels → millions of blocked Na^+) yields reliable **single-photon detection** in rods.



Counter-intuitive: light hyperpolarises the cell and reduces glutamate — vertebrate photoreceptors are depolarised and releasing glutamate in the dark; they signal by going quiet.

Figure 4. Phototransduction in a rod. A single photon isomerises retinal, activating R^* , which through a two-stage amplifying cascade (transducin $\rightarrow$ PDE6) collapses cGMP and closes CNG channels. The counter-intuitive result: light hyperpolarises the cell and reduces glutamate release — vertebrate photoreceptors are 'on' in the dark.

Recovery, shut-off, and calcium feedback (adaptation)

- **R^* shut-off:** rhodopsin kinase (GRK1) phosphorylates R^* ; **arrestin** caps it.
- **Transducin/PDE shut-off:** GTP hydrolysis on $G\alpha$ (accelerated by RGS9-GAP complex) restores PDE inhibition.
- **cGMP restoration:** as channels close, OS Ca^{2+} falls (Ca^{2+} still extruded by NCKX). Falling Ca^{2+} relieves inhibition of **guanylate cyclase (via GCAPs)** $\rightarrow$ cGMP resynthesis $\rightarrow$ channels reopen.
- **Light adaptation:** the Ca^{2+} -dependent feedback (on guanylate cyclase, on CNG channel cGMP-affinity via calmodulin, and on rhodopsin kinase via recoverin) rescales sensitivity continuously, allowing the system to operate across ~ 10 log units of ambient illumination without saturating.

Adaptation and the duplex range

- **Dark adaptation** ($\sim 30+$ min): cone limb (fast, first $\sim 5-7$ min) then rod limb (slow, the *Kohlrausch kink* marks the rod-cone break); limited by the rate of 11-cis-retinal regeneration.




Figure 5. Dark adaptation. Sensitivity recovers in two limbs: cones adapt fast but plateau early; rods take over after the Kohlrausch kink (~7 min) and reach a far lower threshold, gated by the slow RPE regeneration of 11-cis-retinal. The duplex retina made visible.

- **Light adaptation** (seconds): bleaching + Ca^{2+} feedback reduce gain.
- Together with pupil size and neural gain, this spans scotopic→photopic vision.

D. Biochemistry — Pigments and the Visual Cycle

The chromophore and opsins

All human visual pigments share one chromophore, **11-cis-retinal** (an aldehyde of vitamin A_1 /retinol), bound to a lysine of the opsin via a protonated **Schiff base**. Opsins are **7-trans-membrane G-protein-coupled receptors**; the surrounding amino-acid environment **spectrally tunes** the absorption peak:

Pigment	Gene	λ_{max}	Cell
Rhodopsin	<i>RHO</i> (chr 3)	~498 nm	Rod
S-opsin	<i>OPN1SW</i> (chr 7)	~420–440 nm	S-cone
M-opsin	<i>OPN1MW</i> (X)	~530 nm	M-cone
L-opsin	<i>OPN1LW</i> (X)	~560 nm	L-cone
Melanopsin	<i>OPN4</i>	~480 nm	ipRGC

The protonated Schiff base + counter-ion arrangement, plus residues lining the retinal pocket, set λ_{max} (e.g. the few amino-acid swaps distinguishing M from L opsin). X-linkage of *OPN1MW/LW* underlies the prevalence of red–green colour deficiency.




Figure 6. Human photoreceptor absorption spectra. One chromophore (11-cis-retinal) tuned by four opsins. The heavy overlap of the M- and L-cone curves is why colour must be computed from cone ratios rather than read off any single channel — and why the recently-diverged M/L pair sits so close together.

The retinoid (visual) cycle — regenerating 11-cis-retinal

After bleaching, all-trans-retinal must be reconverted to 11-cis-retinal. The classical **RPE cycle**:

1. In the photoreceptor OS, all-trans-retinal is reduced to **all-trans-retinol** by retinol dehydrogenase (RDH8), using NADPH.
2. All-trans-retinol crosses to the **RPE** (chaperoned by IRBP in the interphotoreceptor space).
3. In RPE: **LRAT** esterifies it to all-trans-retinyl ester.
4. **RPE65** (the isomerohydrolase) converts the ester to **11-cis-retinol** [3].
5. **11-cis-RDH (RDH5)** oxidises it to **11-cis-retinal**.
6. 11-cis-retinal returns to the photoreceptor (via IRBP) and recombines with opsin → regenerated pigment.

A **faster cone-specific cycle** involving **Müller glia** supplies cones, enabling their rapid pigment regeneration and sustained photopic function.

Clinical/biochemical anchors

- **Vitamin A (retinol) deficiency → nyctalopia (night blindness):** rods, with their slow RPE-dependent regeneration and lower pigment reserve, fail first → impaired dark adaptation. Severe deficiency → xerophthalmia.
- **RPE65 mutations → Leber congenital amaurosis (LCA2):** the cycle stalls at isomerisation; the basis of *voretigene neparvovec* (Luxturna) gene therapy.
- **Lipofuscin/A2E accumulation** (a by-product of incomplete retinoid recycling) underlies RPE stress in age-related macular degeneration and Stargardt disease (*ABCA4*).

Part II — The Physics of Light: The Bridge

Everything in Part I concerns how a biological instrument catches and processes light; everything in Part III concerns how that instrument varies across species, stars, and the edges of the spectrum. Both rest on the same physics. This part gathers the optics, the electromagnetic facts, and the wave–matter interactions that the later sections invoke — assembled once, here, as the hinge.

A. What Light Is — Wave and Photon

Light is electromagnetic radiation, and it carries two complementary descriptions that are both true:

- **Wave** — a self-propagating oscillation of electric and magnetic fields, characterised by **wavelength** (λ), **frequency** (f), and amplitude. In vacuum all electromagnetic waves travel at $c \approx 3 \times 10^8 \text{ m/s}$, with $c = \lambda \times f$.
- **Photon** — a discrete quantum of energy, $E = h \times f = h c / \lambda$ (h = Planck's constant). Energy is inversely proportional to wavelength: short waves = high-energy photons.

The two pictures are reconciled by quantum electrodynamics; for vision, both matter directly. The **wave** description governs focusing, diffraction, and polarisation (the optics of the eye). The **photon** description governs absorption and transduction (a single photon either is or isn't caught by a single chromophore — the basis of rod single-photon sensitivity in Part I §C, and of why far-red light eventually cannot drive photochemistry at all, Part III §5).

The visible window exists because $E = hf$ places visible photons at exactly the energy that drives clean, reversible isomerisation of retinal: energetic enough to flip the molecule, gentle enough not to destroy it.

B. The Electromagnetic Spectrum — One Continuum

Radio, microwave, infrared, visible, ultraviolet, X-ray, and gamma are **not different phenomena** — they are one continuous spectrum, differing only in wavelength/frequency/photon energy. The named bands are conveniences with fuzzy, overlapping borders (high-energy X-rays and low-energy gamma rays overlap in wavelength and are distinguished by *origin*, not energy).

Band	Wavelength	Photon energy	Relevance to vision
Radio	km → ~1 m	~neV–μeV	Un-seeable: photons too feeble; $\lambda \gg$ any biological aperture
Microwave	~1 m → ~1 mm	~μeV–meV	Thermal/rotational; not imaged biologically
Infrared	~1 mm → 750 nm	~meV–1.6 eV	Sensed as <i>heat</i> (pit vipers); near-IR at the red edge of vision
Visible	750 → 380 nm	~1.6–3.3 eV	The imaging window — clean reversible photochemistry
Ultraviolet	380 → 10 nm	~3.3 eV–124 eV	Near-UV visible to many animals; far-UV ionises/damages
X-ray	10 → 0.01 nm	~124 eV–124 keV	Cannot be focused by tissue; ionising
Gamma	< 0.01 nm	> ~124 keV	Nuclear/astrophysical; destructive

On a logarithmic axis the spectrum spans ~20 orders of magnitude in wavelength; **visible light occupies less than one octave** (380→750 nm doesn't even double). We evolved to see precisely the slice where the Sun's output peaks *and* where water and atmosphere are transparent — the **atmospheric/oceanic optical window** — no coincidence (developed in Part III §6).




Figure 7. The electromagnetic spectrum on a logarithmic wavelength axis. Radio to gamma spans ~20 orders of magnitude in wavelength; visible light occupies less than one octave. Long wavelengths (left) carry photons too feeble for photochemistry; short ones (right) carry photons that ionise. Vision is boxed into the narrow Goldilocks slice between.

C. Field Structure — Why Radio Is Not "Simpler" Than Light

A common misconception holds that visible light is a "two-plane" wave while radio is a "simple" single oscillation. This is incorrect. **Every electromagnetic wave, at every wavelength, has identical field structure:**

- **A single wave with two coupled field components** — an electric field **E** and a magnetic field **B** — not two independent waves.
- **$E \perp B$** , and both are **transverse** (perpendicular to the direction of propagation).
- **E and B are in phase** — they peak together and zero together. The "90°" refers to their spatial orientation, *not* a phase lag.
- E and B are locked by **Maxwell's equations**: a changing E generates B and vice versa; this mutual regeneration *is* how the wave self-propagates through empty space. One cannot exist without the other.

A radio wave is *the same object* as a light wave — same $E \perp B$ transverse structure — merely at far longer wavelength and far lower photon energy. There is no point on the spectrum where the magnetic component switches off and the wave becomes a "simple" single oscillation. Radio is not structurally simpler; it is the identical structure, rescaled.

D. Polarisation — The Channel We Discard

Where electromagnetic waves *genuinely* differ from one another (independently of wavelength) is in **polarisation** — the orientation behaviour of the **E field**. (Polarisation is *not* about E-vs-B; it is about which way E points.)

- **Linear (plane) polarised** — E oscillates in one fixed plane.
- **Unpolarised** — E orientation is random and rapidly changing (sunlight, an incandescent bulb).
- **Circular / elliptical** — the E vector rotates as the wave propagates; circular polarisation has a **handedness** (left/right).

Polarisation applies across the **entire** spectrum — radio engineers match antenna polarisation exactly as optical physicists manage polarised light. Crucially, polarisation carries information that brightness and colour do not: reflection off surfaces partially polarises light in ways that depend on material, angle, and wetness; scattering in the sky produces a fixed polarisation pattern around the Sun.

Humans are nearly polarisation-blind (only the faint **Haidinger's brush** from dichroic macular pigment), but many invertebrates read the E-field orientation directly — the basis of the polarisation-vision and "polarisation-mind" material in Part III §4, §7. The microvillar (rhabdomeric) photoreceptor geometry that makes this possible is described in Part III's comparative physiology.

E. Light Meets Matter — Refraction, Absorption, and the Index of Refraction

How light interacts with material is what makes an eye possible — and what walls off the rest of the spectrum.

Refraction. At the boundary between two media, light bends by an amount set by the **refractive index n** ($n = c / v$, the ratio of light's speed in vacuum to its speed in the medium), following **Snell's law** ($n_1 \sin \theta_1 = n_2 \sin \theta_2$). Refractive power requires an *index step*: the bigger the difference in n across a surface, the more bending. This is precisely why, in the human eye (Part I §A), the **air→cornea** interface (n : 1.00→1.376) does ~two-thirds of the focusing, while the lens — bathed in fluids of similar index — contributes less despite its shape. It is also why a cephalopod's cornea is optically inert underwater (no index step) and its spherical gradient-index lens must do nearly all the work (Part III).

Why X-rays can't be focused by an eye. For X-rays, the refractive index of all materials is **almost exactly 1** (very slightly less). With essentially no index step, lenses barely bend X-rays — so a refractive (camera) eye is physically impossible in the X-ray band, regardless of biology. Real X-ray optics resort to grazing-incidence mirrors and zone plates, none buildable from wet tissue (Part III §5).

Absorption and the photon-catch. To be detected, a photon must be *absorbed* — its energy taken up by a molecular electronic transition. This requires the photon energy ($E = hf$) to match an available transition. Retinal's conjugated π -system has a gap matched to visible photons. Radio photons carry ~a billion times too little energy for any electronic transition (no chemistry can catch them); X-ray and gamma photons carry far too much (they ionise and destroy rather than gently isomerise). The chemistry of absorption, not the wave physics, sets the spectral edges of true vision.

Transparency. A material is transparent in a band where it neither absorbs nor strongly scatters. Water transmits a narrow blue-green window and absorbs UV and IR within metres; the cornea and lens are transparent across the visible but the lens absorbs UV (a built-in filter, Part I §A; removed in aphakia, Part III §5). Transparency windows of the *medium* (atmosphere, water) and the *optics* (cornea, lens) jointly constrain what any eye in that environment can use.

F. Diffraction and the Aperture Limit — Why "Eye Size vs Wavelength" Decides Everything

Because light is a wave, it cannot be focused to an arbitrarily small point. Passing through an aperture of diameter **D**, light spreads into a diffraction pattern (the **Airy disc**) whose angular radius is:

$$\theta \approx 1.22 \frac{\lambda}{D}$$

This sets the **diffraction-limited resolution** of any imaging system. Two consequences run through the whole document:

1. **Imaging requires $D \gg \lambda$.** The aperture must be many wavelengths across to form a sharp image. For visible light ($\lambda \approx 0.0005$ mm) a few-mm pupil is millions of wavelengths wide → excellent resolution. For radio ($\lambda \approx 1$ m to km) a biological "eye" would resolve the entire sky as a single blurry pixel — which is exactly why radio telescopes are enormous and why continent-scale interferometer arrays exist (Part III §5). **The aperture-vs-wavelength ratio, not biology, forbids radio vision.**
2. **The eye is co-tuned to its own sensor.** In the human eye (Part I §A), at the ~2.5–3 mm pupil that optimises acuity, the Airy-disc size is matched to the foveal cone spacing (~2.5 μ m). The optics and the photoreceptor mosaic are balanced: finer optics would out-resolve the cone array, and denser cones would out-sample the blur. Acuity (~1 arcmin, 20/20) sits at this physical sweet spot.

The two-walled window, stated in physics terms. *Long-wavelength wall:* photons too feeble for photochemistry ($E = hf$), and λ too large for any biological aperture to image ($\theta \approx 1.22 \lambda/D$). *Short-wavelength wall:* photons so energetic they ionise the detector, and $n \rightarrow 1$ so no tissue lens can focus them. The visible octave is the unique band where photons are energetic enough for clean reversible photochemistry yet gentle enough not to destroy the molecule, **and** short enough to image with a small wet aperture yet long enough to be refracted by wet tissue. A Goldilocks window fixed by physics — not by which star one orbits (Part III §6).

Part III — Comparative Vision: The Same Window, Different Engines

Having fixed the human reference, the question becomes how far the design space stretches. Across animals, the photon-catching window is nearly invariant, but the optics, the histology, the receptor algorithm, and even the chromophore chemistry are remarkably plastic.

Optics across species — focusing in wet matter

Nature has independently discovered most of an optical engineer's toolkit:

- **No optics (pigment-cup / pinhole):** flatworms (directional only) → *Nautilus* (a true open pinhole eye, no lens, fluid-filled by seawater) — the living intermediate stage of camera-eye evolution [4].
- **Refractive (lens) camera eyes — two independent inventions:**
 - **Vertebrates** — corneal refraction dominant *in air* (the cornea does $\sim 2/3$ of the work); accommodation by lens *reshaping* (mammals/birds) or lens *translation* (most fish, amphibians).
 - **Cephalopods** — a spherical, high-power, **gradient-index lens** does nearly all the focusing (the cornea is optically inert underwater, where index steps vanish); accommodation by **moving the lens** toward/away from the retina (like a camera), not reshaping it. A verted (non-inverted) retina, hence **no blind spot**.
- **Fish lenses** — near-spherical with a steep **Matthiessen gradient index** ($n \sim 1.52$ centre), the optimal solution to spherical aberration in a high-power lens; the design our own lens approximates more weakly.
- **Reflecting (mirror) eyes** — the **scallop** focuses with a concave **guanine-crystal mirror** (a biological reflecting telescope) onto a double retina; deep-sea *Dolichopteryx* hedges with combined refractive + reflective optics.
- **Mineral lenses** — **trilobites** built lenses from oriented **calcite**, with phacopid **aberration-correcting doublets** (a Descartes/Huygens design, 400 My early).
- **Compound eyes:**
 - **Apposition** (diurnal insects, e.g. bees) — each ommatidium isolated; bright-light, modest sensitivity, resolution set by interommatidial angle.
 - **Superposition** (nocturnal moths, krill, deep-sea crustaceans) — many facets pool light onto shared receptors → $\sim 100\text{--}1000\times$ sensitivity.
- **Distributed / whole-body optics** — brittlestars and chitons stud the skeleton/shell with thousands of **calcite micro-lenses**; vision need not be localised in a head.

Retinal architecture across species

- **Inverted vs verted retina** — vertebrates (inverted; photoreceptors face away from light; blind spot; RPE support; Müller fibre-optic compensation) vs **cephalopods** (verted; photoreceptors face the light; clean wiring; no blind spot) — the canonical convergent-evolution natural experiment (developed fully in §11 below).
- **Tapetum lucidum** — many nocturnal vertebrates (cats, ungulates, sharks) add a reflective layer behind/within the choroid or RPE, giving photoreceptors a second pass at each photon (eyeshine) — sensitivity at the cost of acuity.
- **Foveae** — some raptors have **two foveae** per eye (one for forward binocular, one for lateral monocular acuity) and far higher cone densities than humans (visual acuity several-fold ours).
- **Rhabdomeric vs ciliary photoreceptors** — vertebrate photoreceptors are **ciliary** (OS = modified cilium; hyperpolarising; cGMP/PDE cascade). Most invertebrate (e.g. arthropod, cephalopod) photoreceptors are **rhabdomeric** (microvillar **rhabdomere**; **depolarising** to light; a **Gq/PLC** cascade opening TRP channels). The two lineages diverged anciently; both exist in some animals. The microvillar geometry also makes rhabdomeric receptors **intrinsically polarisation-sensitive** (aligned chromophores), which is why invertebrates dominate polarisation vision.

Receptor physiology & chromophore chemistry across species

- **Opsin diversity & spectral reach** — from UV (many insects, birds, fish: a dedicated UV/violet cone) to far-red; mantis-shrimp eyes carry 12–16 photoreceptor classes (but likely *barcode*, not inferential, colour — §9 below).
- **Chromophore variants** — the chromophore is not always retinal- A_1 . Freshwater fish and some amphibians use **retinal- A_2 (3,4-didehydroretinal / "porphyropsin")**, red-shifting sensitivity for turbid/long-wavelength waters; some deep-sea fish and the larvae of certain species switch chromophore with habitat/metamorphosis. Same *strategy* (a tunable isomerising chromophore in an opsin), different molecule.
- **Cascade polarity** — hyperpolarising (vertebrate, ciliary, cGMP) vs depolarising (invertebrate, rhabdomeric, PLC/TRP): two biochemically distinct solutions to the same transduction problem.
- **Polarisation & circular-polarisation sensing** — aligned microvilli give bees a sky-compass channel; cephalopods a private signalling channel; **mantis shrimp** add **circular** polarisation, a modality with no human analogue.
- **Convergence on the essentials** — despite all this, the recurring solutions are few: an isomerising chromophore in a tuning protein; an amplifying enzymatic cascade; opponent comparison for colour; the camera and compound body plans; and a fixed ~300 nm–1.5 μ m operating window. Vision is constrained by physics and chemistry first, and by phylogeny only second.

Parts I and II fixed the human substrate and the physics of light. The sections that follow (§1–§11) take that substrate outward: to why molecules catch light at all, to the electromagnetic window, to other stars and polarisation minds, and back to the flawed-but-brilliant inverted eye.

1. The Human Photoreceptors

Humans have two image-forming photoreceptor classes plus a non-image-forming third type.

Rods — ~120 million. One pigment (rhodopsin), peak ~498 nm. Single-photon sensitive; drive scotopic (night) vision; achromatic, low acuity (high convergence). Saturate in daylight. Absent from the foveal centre, densest in a ring ~18° out.

Cones — ~6 million. Drive photopic vision, colour, high acuity; concentrated in the fovea. Three subtypes:

- **S-cones** ("blue"), peak ~420–440 nm (~5–10%, absent from the very foveal centre)
- **M-cones** ("green"), peak ~530–535 nm
- **L-cones** ("red"), peak ~560–565 nm

M and L diverged recently (X-chromosome gene duplication ~30–40 Mya), which is why red–green colour-blindness is X-linked and common.

ipRGCs — intrinsically photosensitive retinal ganglion cells (~1–3%) containing **melanopsin** (peak ~480 nm). Non-image-forming: drive the pupillary light reflex and circadian entrainment. The "what time is it" channel.

Image-forming vision = a **duplex, trichromatic retina** (rods + 3 cones), with melanopsin ipRGCs handling clock/pupil.

2. Why Photoreceptors Respond to Light

The "why": chromophore + opsin

Every photoreceptor uses **11-cis-retinal**, a vitamin-A-derived chromophore inside an opsin protein. Retinal's conjugated polyene chain delocalises its π -electrons; the ground-to-excited-state energy gap falls in the visible range, so a visible photon is exactly the right energy to absorb. The opsin tunes that gap via nearby charged amino acids (**spectral tuning**) — one chromophore, many peaks (~420–565 nm). Same molecule, different protein pocket.

The light event: isomerisation

Absorbing a photon does exactly one thing: **11-cis** → **all-trans** isomerisation. The molecule straightens (~200 fs, ~2/3 efficiency). This is the *only* light-dependent step in vision. The shape change forces the opsin into its active conformation (**metarhodopsin II**, R*).

The "how": the phototransduction cascade (rods)

1. **R*** catalytically activates **transducin** (a G-protein) — hundreds per R*.
2. Each transducin activates **phosphodiesterase (PDE)**.
3. PDE hydrolyses **cGMP**, dropping its concentration.
4. cGMP normally holds open **CNG cation channels**; when cGMP falls, they **close**.

Amplification across these steps = single-photon detection.

The counter-intuitive part: light *hyperpolarises*

In the dark, CNG channels are open, Na⁺/Ca²⁺ flow in ("dark current"), the cell sits depolarised and releases glutamate steadily. **Light closes the channels → hyperpolarisation → less glutamate**. Vertebrate photoreceptors are "on" in the dark and signal light by going quiet. (Invertebrates depolarise to light — different cascade.)

Resetting

All-trans-retinal is released and recycled to 11-cis (largely in the **RPE — the visual cycle**); guanylate cyclase rebuilds cGMP; channels reopen. Calcium feedback during recovery is the core of light adaptation.

One photon → one isomerisation → a shape change → a G-protein cascade → channels close → less glutamate.

3. The Electromagnetic Spectrum

(Consolidated reference: Part II §B. The version here is the original narrative treatment.)

"Light" strictly means visible (~380–750 nm), but it's one continuum with the whole EM spectrum. From long/low-energy to short/high-energy:

Band	Wavelength	Notes
Radio	km → ~1 m	Lowest energy; comms, radio astronomy
Microwave	~1 m → ~1 mm	Radar, Wi-Fi, CMB
Infrared	~1 mm → 750 nm	Thermal radiation, felt as heat
Visible	750 → 380 nm	The only band human eyes detect
Ultraviolet	380 → 10 nm	Sunburn, vitamin D, sterilisation
X-rays	10 → 0.01 nm	Penetrating; imaging, crystallography
Gamma	< 0.01 nm	Highest energy; nuclear, astrophysical

Key relations:

- $c = \lambda \times f$ (wavelength $\times$ frequency = speed of light)
- $E = h \times f$ (photon energy = Planck's constant $\times$ frequency)

Shorter wavelength = higher frequency = higher photon energy. The visible band is a sliver — less than one octave of a ~20-order-of-magnitude spectrum — sitting where the Sun peaks *and* where water/atmosphere are transparent. Band boundaries overlap and are named by discovery method, not hard physics.

4. Field Structure & Polarisation

(Consolidated reference: Part II §C–D. The version here is the original narrative treatment.)

Every EM wave — radio to gamma — has the identical structure: mutually perpendicular **E** and **B** fields, both transverse to propagation. Radio is *not* "simpler"; it's the same object at longer wavelength. There's no point on the spectrum where the magnetic component switches off.

Clarifications:

- It's **one wave with two coupled components**, not two waves. E and B are locked by Maxwell's equations — a changing E generates B and vice versa; that mutual regeneration is propagation.
- They're **in phase** (peak and zero together); the "90°" is spatial orientation, not phase lag.
- **Polarisation** is where waves genuinely differ — it's about the **E field's orientation**, not E-vs-B:
 - *Linear*: E stays in one fixed plane.
 - *Unpolarised* (sunlight, bulbs): E orientation random.
 - *Circular/elliptical*: E vector rotates as it propagates.
- Polarisation applies across the **whole** spectrum (radio antenna alignment is the same problem as optical polarisation).

Humans are nearly polarisation-blind (only a faint trace — **Haidinger's brush**, from dichroic macular pigment).

5. Seeing Outside the Visible — Radio, UV, X-ray

Can humans see radio? No — two walls

- **Wall 1 — no receptor.** Vision needs a molecule whose electronic energy gap matches the photon energy. Radio photons carry ~a billion times less energy; no molecular transition is that small. You'd need an *antenna* sized to the wavelength.

- **Wall 2 — aperture vs wavelength (fatal for imaging).** Resolution scales as $\theta \approx 1.22 \lambda/D$. Visible $\lambda \approx 0.0005$ mm $\rightarrow$ millions of wavelengths across the pupil $\rightarrow$ sharp images. Radio $\lambda \approx 1$ m to km $\rightarrow$ a biological "eye" would see the whole sky as one blurry pixel. (This is why radio telescopes are enormous and use continent-spanning interferometer arrays.)

As wavelength lengthens, "seeing" shades into "feeling warmth" — pit vipers detect IR as *heat* (thermal pits), not photon-catching vision. Radio photons are too feeble to deposit meaningful energy at all.

UV — the easy, already-realised case

UV is shorter than visible, so diffraction is no obstacle. The human **lens** absorbs UV (built-in filter); remove it (aphakia) and people see into the near-UV (~300–350 nm). Insects, birds, reptiles, many fish have UV-transparent optics and a dedicated UV cone (flowers show UV nectar guides; many birds are UV tetrachromats). Below ~300 nm, photons start breaking bonds (DNA absorbs ~260 nm), so far-UV vision is selected against — *the signal would destroy the detector*.

Soft X-rays — where it breaks down

Apertures get trivially easy, but three brutal walls appear:

1. **No refraction.** X-ray refractive index ≈ 1 for all materials — lenses don't bend X-rays. Real X-ray optics use grazing-incidence mirrors (Wolter telescopes) or zone plates; none buildable as wet tissue. The camera-eye body plan is unavailable.
2. **Poor catch.** Soft X-rays pass through low-density tissue largely unabsorbed (we image bone *through* flesh). Absorbing them needs dense, high-Z material, not retinal cells.
3. **Every caught photon is a micro-disaster.** Thousands of eV — ionises, smashes DNA, makes free radicals. It doesn't *signal*, it *damages*. No recoverable transduction cycle is possible.

The deep symmetry

- **Long-wavelength wall (IR $\rightarrow$ radio):** photons too feeble for photochemistry; apertures must be absurdly large. *Killed by low energy + large wavelength.*
- **Short-wavelength wall (far-UV $\rightarrow$ X-ray):** photons ionise/destroy the detector; matter stops refracting. *Killed by high energy + no usable optics.*

The visible octave is the unique band where photons are **energetic enough for clean, reversible photochemistry but gentle enough not to wreck the molecule**, and **short enough to image with a small wet aperture but long enough to be refracted by wet tissue**. A genuine Goldilocks window, fixed by physics — not by our star.

6. Eyes Around Other Stars

The **imaging window** (~300 nm–1.5 μ m at the absolute outside) is roughly fixed everywhere. The star slides the eye *within* the window and resizes it.

Shared ceilings: the blue wall (~300 nm, ionisation) and the red wall (~1.1–1.5 μ m, photochemistry fails).

- **M-dwarf (red dwarf) worlds** — cool (~3000 K), peak ~1 μ m, dim. Opsins push red (700–1000+ nm); colour space compressed toward red/near-IR; large light-greedy eyes; tidally-locked planets favour terminator-dwelling life with eyes tuned to a permanently low, reddened sun. *Our "red" might be their middling-blue.*
- **Sun-like (G-type)** — peak ~500 nm, broad band, generous flux. The comfortable middle; the least surprising aliens. Trichromacy-or-richer, moderate apertures.
- **F/A stars (hotter, bluer)** — peak blue/UV, UV-flooded. Heavy UV filtering required; opsins crowd against the blue wall; photons plentiful so eyes spend budget on resolution/colour; short stellar lifetimes squeeze complex life.
- **Underwater worlds (the equaliser)** — water transmits only ~450–550 nm (blue-green), stripping UV and IR within metres. Deep-water eyes converge on blue-green **regardless of star**. Medium overrides star.

Convergence — what stays constant: the camera and compound eye re-evolve (physics, not phylogeny); retinal-like tunable chromophores are a strong attractor; colour uses opponent comparisons; the same two-walled window everywhere. Alien eyes would be **recognisable** — tuned in spectral position, aperture, and colour emphasis, but not relocated out of the window.

The genuinely alien possibility: polarisation / circular-polarisation vision as a *primary* sense — wide open across the whole window, and something we barely use.

7. A Mind Built on Polarisation

Every scene point carries, beyond brightness and colour, two more independent properties we discard:

- **Degree of polarisation** (0% random → 100% aligned)
- **Angle of polarisation** (0–180°, cyclic)

What a polarisation-native creature would see:

- **Water and wet surfaces blaze** — strong horizontal polarisation (dragonflies find water this way; get fooled by asphalt and solar panels).
- **Material discrimination for free** — same colour/brightness, different polarisation → tells living from dead, wet from dry, a remote sense of *what something is made of*.
- **Camouflage collapses** — transparent/silvery bodies alter polarisation; cephalopods likely detect transparent prey and signal on a "private channel."

- **The sky becomes a compass and clock** — the sky's polarisation fan points permanently at the Sun, through cloud and at twilight.

What it's like (the hard part): an irreducible *p-hue / p-saturation* quale we have no word for; the angle is **cyclic over 180°**, so the "p-hue circle" is a half-turn loop — different topology from our colour wheel. Mantis shrimp add **circular polarisation** (handedness) — no human analogue at all.

A polarisation-primary alien wouldn't see a *prettier* world — it would see one organised around **material, wetness, surface geometry, and celestial direction**: a different *ontology* carved from the same photons.

8. Earth's Existence-Proofs — Eyes That Broke the Rules

- **Trilobite eyes — lenses of rock.** Built from **calcite** (birefringent), oriented along the crystal's optic axis to avoid double images. Phacopid schizochroal eyes used aberration-correcting **doublets** — a design human opticians (Descartes, Huygens) reached 400 My later. Mineral, aberration-corrected optics in the Cambrian.
- **The scallop — a mirror eye.** Up to 200 eyes focusing with a concave **guanine-crystal mirror** (a reflecting telescope), tiled square crystals steering colours onto two retinas. (Deep-sea *Dolichopteryx* also uses mirror optics.)
- **Superposition compound eyes.** Nocturnal moths, krill, deep-sea crustaceans pool light from many facets onto single receptors — ~100–1000× sensitivity gain. Near-camera light-gathering in a compound eye.
- **The mantis shrimp — the maximalist.** 12–16 photoreceptor classes (UV–red) + linear *and* circular polarisation (only known animal). But likely *poor* colour discrimination — a hardwired **"colour barcode" scanner**, not opponent hue inference.

Contested point. The "barcode" interpretation (Thoen et al. 2014) [5] is influential but not settled. The behavioural $\Delta\lambda$ tests used wide-field stimuli and trained discriminations that may understate the animal's capacity; alternative readings allow some opponent processing or task-dependent performance. Treat "barcode vs inference" as a live hypothesis, not a closed result.

- **The brittlestar / chiton — body as eye.** Thousands of calcite micro-lenses across the skeleton/shell; vision needn't be localised in a head.

The photon-catching window is nearly fixed, but the **engineering and the algorithm** built on it are wildly open.

9. Colour as Inference vs Colour as Barcode

The deep point: more sensor channels $\neq$ richer experience. It depends on what the brain does with them.

Inference architecture (us)

Three cones with **broad, overlapping** sensitivities — no single cone reads wavelength; information lives in **ratios**. The brain **computes** colour via opponent comparisons (L–M $\rightarrow$ red/green; (L+M)–S $\rightarrow$ blue/yellow). Hue is *inferred*.

Buys: a **continuous hue space** (3 sensors $\rightarrow$ unlimited hues); **colour constancy** (reading ratios discounts the illuminant $\rightarrow$ colour tracks *objects*); **efficiency**.

Costs: **metamers** — physically different spectra with identical cone ratios look identical (the basis of RGB screens). We see a 3-D projection of the spectrum, and collisions are invisible.

Barcode architecture (mantis shrimp, probably)

12–16 **narrow** channels, each tuned to a slice. Evidence: it does *not* compute fine hue — discrimination is **coarse**, worse than ours. Uses **labelled lines**: "which channels fired," scanned by sweeping the eye.

Buys: **speed** (no computation; hardwired instant recognition — vital for a millisecond strike); less metamerism in the natural range.

Costs: **coarse discrimination, no continuous hue space**.

Why it's deep

The naïve "16 cones $\rightarrow$ vider rainbow" is **backwards**. The mantis shrimp likely has a *poorer*, more categorical colour world. **Perceptual richness lives in the inference, not the sensor count** — the *ambiguity* in our front end forces computation, and computation is where the smooth, constant, object-anchored hue world is *manufactured*. An alien with fewer channels but cleverer inference could out-experience one with more channels and dumber wiring.

Colour is a **make-or-buy decision** about where to put the cleverness: barcode vision is what you build with great eyes and a small brain; inference vision is what you build with a big brain to spare.

10. How Vision Got Started — One Molecule to a Focusing Eye

The "what use is half an eye?" objection has a complete answer: every rung is useful, and every rung exists in a living animal *right now*.

Rung	Stage	Use	Living example
0	Light-sensitive molecule	Light/dark for timing, UV avoidance	(ancient; predates eyes)
1	Flat photoreceptor patch (eyespot)	Brightness & shadow detection	<i>Euglena</i> , larvae
2	Pigmented cup	Crude direction of light	Flatworms, limpets
3	Deep cup → pinhole	Crude image (pinhole camera)	<i>Nautilus</i>
4	Covered/filled chamber	Protection → transparent tissue refracts	(transitional)
5	Gradient-index lens	Focused, bright, sharp image	Fish, octopus, us

The chromophore — the genuinely improbable chemistry — came **free and ancient**, possibly from non-visual uses. Everything after is **geometry and graded transparency**, which selection sculpts trivially. The real fish/cephalopod lens is a **refractive-index gradient** (auto-corrects spherical aberration), not uniform glass.

Nilsson & Pelger (1994) [6]: modelled the whole path at 1%-per-step under modest selection → **a few hundred thousand years**. A geological eyeblink. The camera eye is *easy*; it evolved independently many times.




Figure 8. The eye-evolution ladder. Each rung is independently useful and each survives in a living animal today — from the eyespot (*Euglena*) through the pigment cup (flatworms) and the lens-less pinhole (*Nautilus*) to the gradient-index lens eye (fish, octopus, human). Selection climbs it one small improvement at a time.

The eye isn't a miracle; it's a pinhole camera that learned to wear glasses. **Evolution optimises the front end only as far as the back end can use it** — eye and interpreting brain co-evolve.

11. The Inverted Retina — A Flaw, and a Brilliant Patch

Vertebrates and cephalopods both evolved camera eyes independently — and wired the retina **opposite ways**. A clean natural experiment.

- **Octopus (verted)**: photoreceptors face the light; nerves run out the back. Clean, **no blind spot**.
- **Vertebrate (inverted)**: photoreceptors face *away*; light passes through neurons/vessels/fibres first; the nerve bundle **punches a hole** through the retina to exit → the **blind spot**.

Why it looks like a flaw

The vertebrate retina is installed **backwards**: light degraded before detection; a structural blind spot; loose attachment → **retinal detachment** and several blinding conditions. The octopus, solving the same problem, did none of this. The cause is **embryological** — the vertebrate eye grows as an *outpocketing of the brain*, forcing photoreceptors inward; once locked in, there's no viable 1%-at-a-time path to flip it. A real, honest limit of evolution — proof a better arrangement was possible.

Clinical correlate — the inverted architecture. The design's loose apposition (photoreceptors sit *against* but are not firmly bonded to the RPE) is the substrate of **rhegmatogenous retinal detachment**: a retinal break lets vitreous fluid track into the subretinal space, separating photoreceptors from their RPE support and choroidal blood supply — a surgical emergency, since detached photoreceptors rapidly lose function. The blind spot, by contrast, is silent — binocular overlap and cortical filling-in conceal it entirely.

Why "flaw" overshoots

1. **Müller cells = living fibre-optics.** Glial cells span the retina and guide light through the obstructing layers to the photoreceptors with little loss (even sorting wavelengths). A **patch on a flaw** — characteristic evolution.

Contested point. That Müller cells *guide* light (Franze et al. 2007) [1] and *sort* wavelengths (Labin et al. 2014) [7] is well demonstrated optically; the **functional magnitude** of the acuity/sensitivity benefit in the living, behaving animal is less certain and still debated. The mechanism is real; how much of the inverted retina's optical penalty it actually recovers is an open quantitative question.

2. **The inversion buys the RPE.** Photoreceptor tips embed in the **retinal pigment epithelium**, which **recycles the chromophore** (visual cycle), **absorbs stray light** (sharp images), and **clears oxidative waste** from the most metabolically demanding cells in the body. The inverted design plugs photoreceptors straight into the support

they need — possibly enabling **higher-performance, higher-maintenance** photoreceptors at the cost of optical cleanliness.

Synthesis

The inverted retina is **both**: a frozen developmental accident evolution couldn't undo, *and* an architecture that co-evolved with a high-performance support system (RPE) and a mitigating patch (Müller fibre-optics) — far better than "backwards = broken" suggests.




Figure 9. Two camera eyes, evolved independently and wired in opposite directions. The vertebrate retina is inverted (photoreceptors face away from the light, axons exit through a blind spot) but plugs into the RPE support layer; the cephalopod retina is verted (photoreceptors face the light, no blind spot).

The vertebrate, with its backwards retina, still ended up with the sharper eye and the richer visual mind. **The constraints are set by physics and history; the experience is set by computation** — a clever enough back end turns even a backwards front end into the richest sense any animal has.

The Through-Line

Vision is a **layered stack** — chromophore → geometry (cup → pinhole → lens) → wiring → cortical inference — and at every layer evolution builds forward from what the last one left it.

- The chromophore came **free and ancient**.
- The lens was **easy** (a graded blob; Nilsson & Pelger estimate a few hundred thousand years).
- The wiring got **frozen backwards** by embryology — so evolution bolted on compensations (Müller guides in front, RPE behind, a brain that fills the blind spot).

Just as our *rich colour* is manufactured by inference from ambiguous cone ratios, our *seamless visual field* is manufactured by inference papering over a literal hole in the sensor.

The eye delivers a flawed, backwards, gappy signal — and the brain builds a perfect-seeming world out of it. The richness was never in the hardware; it's in what gets computed on top of whatever physics, chemistry, and frozen history happened to make available.

From why retinal absorbs a photon, out to alien stars and polarisation minds, and back to the flawed-but-brilliant eye you're reading this with.

Coda — The Stargate: Toward the Central Visual Pathway

This document ends, deliberately, at a threshold.

Trace the signal one last time. A photon isomerises retinal; a cascade closes a channel; a photoreceptor falls quiet; bipolar and ganglion cells split the message into ON and OFF, centre and surround, fast and slow. Then ~1.2 million ganglion-cell axons sweep across the inner retina, converge on the optic disc, and pass together through the **lamina cribrosa** — the sieve in the sclera where the retina stops and the nerve begins. That passage is the edge of everything covered here. Beyond it, the axons leave the optical organ entirely.

Think of the optic disc as a **stargate**: a small, unremarkable-looking ring through which the entire visual world is funnelled, and on the far side of which the rules change. On *this* side lie optics and photochemistry — refraction, absorption, transduction, the physics of light meeting wet matter, the subjects of Parts I–III. On the *far* side lies computation proper: not how light is *caught*, but how a brain *makes a seen world* out of spike trains.

What waits beyond the gate — the subject of the companion volume, *The Seen World Beyond The Stargate*:

- **The optic chiasm** — where the nasal fibres cross and the visual field is re-sorted by *side of space* rather than by *eye*, the geometric foundation of binocular vision and of how a lesion's location predicts the field defect.
- **The lateral geniculate nucleus (LGN)** — the thalamic relay, layered into **magnocellular** (motion, contrast, fast), **parvocellular** (form, colour, detail), and **koniocellular** (S-cone/blue-yellow) streams — the channels first hinted at in the retina now anatomically separated.
- **Primary visual cortex (V1)** — orientation columns, ocular-dominance columns, the Hubel–Wiesel architecture; the first place the image is taken apart into edges and re-assembled.
- **The dorsal and ventral streams** — "where/how" toward the parietal lobe, "what" toward the temporal lobe; the divergence where seeing becomes recognising, reaching, and knowing.

The through-line of *this* document was that **richness lives in computation, not hardware** — that the brain manufactures a seamless, constant, coloured world from a flawed and gappy sensor. Everything past the stargate is that manufacturing process made explicit. The eye, for all its elegance, is only the aperture. The seeing happens later, and elsewhere.

We stop at the lamina cribrosa — the last structure that is still the eye. What lies beyond is the brain learning to see. That is the next subject.

Glossary & Abbreviations

Alphabetical. Bold terms are abbreviations expanded; others are concept definitions for dip-in reading.

A2E — N-retinylidene-N-retinylethanolamine; a toxic bisretinoid by-product of incomplete retinoid recycling that accumulates in the RPE (lipofuscin); central to Stargardt disease and AMD.

Accommodation — the eye's adjustment of lens power to focus near objects (Helmholtz mechanism: ciliary contraction relaxes zonular tension, lens rounds up).

Airy disc — the diffraction-limited spot a point source forms through an aperture; angular radius $\theta \approx 1.22 \lambda/D$.

Aphakia — absence of the crystalline lens (e.g. post-cataract without an implant); admits near-UV to the retina.

Apposition / superposition eye — two compound-eye optical schemes: isolated ommatidia (bright-light, high-resolution) vs light pooled across facets (high-sensitivity, nocturnal).

Bruch's membrane — the basal lamina complex between RPE and choriocapillaris.

Chromophore — the light-absorbing molecule; in vertebrate vision, 11-cis-retinal (vitamin-A aldehyde).

Ciliary vs rhabdomeric photoreceptors — the two ancient receptor lineages: ciliary (vertebrate; hyperpolarising; cGMP/PDE) vs rhabdomeric (most invertebrates; depolarising; Gq/PLC/TRP; polarisation-sensitive).

CNG channel — cyclic-nucleotide-gated cation channel; held open by cGMP in the dark, carrying the dark current.

Cone — photopic photoreceptor; three spectral classes (S/M/L) underlie human trichromacy.

Dark current — the steady inward $\text{Na}^+/\text{Ca}^{2+}$ flow keeping photoreceptors depolarised in darkness.

Dioptre (D) — unit of refractive power (reciprocal of focal length in metres).

ELM / ILM — external / internal limiting membrane: the outer and inner boundary layers of the neural retina.

Emmetropia — the refractive state in which distant objects focus on the retina without accommodation.

GCAP — guanylate cyclase-activating protein; mediates Ca^{2+} feedback restoring cGMP during recovery.

Gradient-index (GRIN) lens — a lens whose refractive index varies internally (centre > edge), correcting spherical aberration; used by fish/cephalopod lenses and, weakly, our own.

Haidinger's brush — the faint entoptic percept of polarised light from dichroic macular pigment; humans' vestigial polarisation sense.

Hyperpolarisation — the membrane-potential change (more negative) by which vertebrate photoreceptors signal light.

IOP — intraocular pressure; set by aqueous production vs outflow; central to glaucoma.

ipRGC — intrinsically photosensitive retinal ganglion cell; melanopsin-containing; drives pupil and circadian entrainment.

IRBP — interphotoreceptor retinoid-binding protein; chaperones retinoids between photoreceptors and RPE.

Isomerisation — the 11-cis → all-trans shape change of retinal on photon absorption; the only light-dependent step in vision.

$\lambda_{\max}$ — wavelength of peak absorption of a visual pigment.

LRAT / RPE65 / RDH — enzymes of the retinoid (visual) cycle: lecithin-retinol acyltransferase, the RPE65 isomerohydrolase, and retinol dehydrogenases.

Melanopsin (OPN4) — the opsin of ipRGCs; $\lambda_{\max}$ ~480 nm.

Metamer — physically different spectra that produce identical cone signals and so look identical; the basis of RGB displays.

Metarhodopsin II (R*) — the active, signalling conformation of photo-excited rhodopsin.

Müller cell — radial glia spanning the retina; metabolic support and living light-guide.

Nyctalopia — night blindness; e.g. from vitamin-A deficiency stalling rod pigment regeneration.

Opsin — the 7-transmembrane GPCR protein that binds the chromophore and spectrally tunes it.

Opponency — neural comparison of receptor channels (e.g. L-M, S-(L+M)) that extracts colour from overlapping pigments.

PDE6 — phosphodiesterase 6; the effector that hydrolyses cGMP in phototransduction.

Photopic / scotopic / mesopic — cone-mediated (bright) / rod-mediated (dim) / mixed vision.

Polarisation (linear / circular / unpolarised) — the orientation behaviour of a wave's electric field; a channel humans barely sense.

Presbyopia — age-related loss of accommodation from lens stiffening.

Rhodopsin — the rod visual pigment; $\lambda_{\max}$ ~498 nm.

Rod — scotopic photoreceptor; single-photon sensitive; one pigment; achromatic.

RPE — retinal pigment epithelium; recycles retinoids, phagocytoses discs, absorbs stray light, forms the outer blood–retinal barrier.

Stiles–Crawford effect — directional sensitivity of cones (light through the pupil centre is more effective), acting as a waveguide/stray-light suppressor.

Tapetum lucidum — a reflective layer behind/within the choroid or RPE in many nocturnal animals giving photoreceptors a second pass at each photon.

Transducin (Gt) — the photoreceptor G-protein coupling R* to PDE6.

Trichromacy — colour vision based on three cone classes.

Verted / inverted retina — photoreceptors facing toward (cephalopod) vs away from (vertebrate) the incoming light.

Visual (retinoid) cycle — the RPE/Müller pathway regenerating 11-cis-retinal after bleaching.

Zonules of Zinn — suspensory fibres linking ciliary body to lens capsule; transmit accommodative force.

References & Further Reading

Vancouver style: primary papers cited in the text are numbered by order of first appearance and carry in-text markers [n]; standard reference works by domain follow as a numbered continuation. Citations verified against journal records; DOIs confirmed at write time.

Key primary literature (cited in text)

1. Franze K, Grosche J, Skatchkov SN, Schinkinger S, Foja C, Schild D, et al. Müller cells are living optical fibres in the vertebrate retina. *Proc Natl Acad Sci USA*. 2007;104(20):8287–92. doi:10.1073/pnas.0611180104. — Radial Müller glia act as low-distortion light guides through the inverted retina. (Part I §B; Part III §11)
2. Wald G. The molecular basis of visual excitation. *Nature*. 1968;219(5156):800–7. doi:10.1038/219800a0. — Nobel-laureate account establishing 11-cis-retinal, the A₁/A₂ chromophore distinction, and bleaching/regeneration. (Part I §C, §D; Part III §2)
3. Kiser PD, Golczak M, Palczewski K. Chemistry of the retinoid (visual) cycle. *Chem Rev*. 2014;114(1):194–232. doi:10.1021/cr400107q. — Enzymology of LRAT, the RPE65 isomerohydrolase, and the RDH steps of 11-cis-retinal regeneration. (Part I §D)
4. Land MF, Nilsson D-E. *Animal Eyes*. 2nd ed. Oxford: Oxford University Press; 2012. — The standard comparative-optics reference: pinhole, lens, mirror, apposition and superposition compound eyes; trilobite, scallop, and cephalopod designs. (Part III; §6, §8)
5. Thoen HH, How MJ, Chiou T-H, Marshall J. A different form of colour vision in mantis shrimp. *Science*. 2014;343(6169):411–3. doi:10.1126/science.1245824. — Behavioural $\Delta\lambda$ tests show stomatopods discriminate colour poorly despite 12 photoreceptor classes, implying a temporal-scan "barcode" recognition system rather than opponent-coded hue inference. (Part III §8, §9)
6. Nilsson D-E, Pelger S. A pessimistic estimate of the time required for an eye to evolve. *Proc R Soc Lond B*. 1994;256(1345):53–8. doi:10.1098/rspb.1994.0048. — The stepwise patch→cup→pinhole→lens model; concludes a lens eye can evolve in only a few hundred thousand years under modest, conservatively modelled selection. (Part III §10)
7. Labin AM, Safuri SK, Ribak EN, Perlman I. Müller cells separate between wavelengths to improve day vision with minimal effect upon night vision. *Nat Commun*. 2014;5:4319. doi:10.1038/ncomms5319. — Müller cells act as wavelength-selective waveguides, routing green–red light to cones while letting blue–violet leak to adjacent rods. (Part I §B; Part III §11)

Standard reference works by domain

Numbered as a continuation of the list above; consulted throughout rather than tied to a single in-text claim.

Visual neuroscience & retinal physiology

8. Kandel ER, Schwartz JH, Jessell TM, Siegelbaum SA, Hudspeth AJ, editors. *Principles of Neural Science*. 5th/6th ed. New York: McGraw-Hill. — Chapters on phototransduction, retinal circuitry, and central visual pathways.

9. Purves D, et al. Neuroscience. Sunderland (MA)/Oxford: Sinauer/OUP. — Accessible treatment of the duplex retina and adaptation.
10. Rodieck RW. The First Steps in Seeing. Sunderland (MA): Sinauer; 1998. — Definitive on photoreceptors and early retinal processing.
11. Hubel DH, Wiesel TN. Collected work on receptive fields and cortical organisation (Nobel Prize, 1981).

Ocular anatomy, histology & optics

12. Forrester JV, Dick AD, McMenamin PG, Roberts F, Pearlman E. The Eye: Basic Sciences in Practice. Edinburgh: Elsevier. — Integrated anatomy/physiology/biochemistry.
13. Snell RS, Lemp MA. Clinical Anatomy of the Eye. Oxford: Wiley-Blackwell.
14. Atchison DA, Smith G. Optics of the Human Eye. Oxford: Butterworth-Heinemann. — Dioptrics, aberrations, the schematic eye.
15. Oyster CW. The Human Eye: Structure and Function. Sunderland (MA): Sinauer.

Photochemistry, biochemistry & the visual cycle

16. Palczewski K. G protein-coupled receptor rhodopsin. Annu Rev Biochem. 2006;75:743–67. doi:10.1146/annurev.biochem.75.103004.142743.
17. Lamb TD, Pugh EN Jr. Dark adaptation and the retinoid cycle of vision. Prog Retin Eye Res. 2004;23(3):307–80. doi:10.1016/j.preteyeres.2004.03.001.

Physics of light & optics

18. Hecht E. Optics. 5th ed. Boston: Pearson; 2017. — Refraction, diffraction, polarisation, the Airy disc.
19. Born M, Wolf E. Principles of Optics. 7th ed. Cambridge: Cambridge University Press; 1999. — The rigorous standard.
20. Feynman RP, Leighton RB, Sands M. The Feynman Lectures on Physics. Vol. I (optics) & Vol. II (electromagnetism, Maxwell's equations). Reading (MA): Addison-Wesley. — For the wave/photon and ELB foundations.

Comparative & evolutionary vision

21. Cronin TW, Johnsen S, Marshall NJ, Warrant EJ. Visual Ecology. Princeton (NJ): Princeton University Press; 2014. — Polarisation vision, spectral tuning across habitats, vision under water and at depth.
22. Nilsson D-E. The evolution of eyes and visually guided behaviour. Philos Trans R Soc Lond B Biol Sci. 2009;364(1531):2833–47. doi:10.1098/rstb.2009.0083. — The modern synthesis of eye-evolution stages.
23. Marshall J, Oberwinkler J. The colourful world of the mantis shrimp. Nature. 1999;401(6756):873–4. doi:10.1038/44751. — Context for the 2014 Thoen et al. finding [5].

Quantitative figures (refractive powers, photoreceptor counts, λ_{max} values, axial lengths) are standard textbook values and may vary modestly between sources and individuals. Where a claim rests on a specific study, the primary reference is cited above by its Vancouver number.