Secondary Motor Cortex Dynamics Are Consistent with Langevin Diffusion

and Somatosensory Cortex Shows a Weak Correlation with Plan Certainty in Mice

Stamelos Loutsos
stamoutsos@gmail.com (Independent Researcher)

April 2026

Abstract

The prefrontal cortex is thought to represent action possibilities during motor planning, but the underlying neural dynamics remain unclear. Here we test the hypothesis that mouse secondary motor cortex (M2, a prefrontal cortical proxy) performs Langevin sampling – a stochastic diffusion process – to represent a distribution over actions, and that somatosensory cortex (S1) signals the certainty of this distribution. Using electrophysiological data from 11 mouse sessions (N=2041 trials) and human vmPFC recordings (N=378 trials), we show: (1) In both species, the mean squared displacement (MSD) of the population trajectory scales linearly with time (mouse: $\Delta\text{AIC} = -30.97$; human: $\Delta\text{AIC} = -30.12$), consistent with Langevin diffusion rather than drift. An Ornstein-Uhlenbeck model was rejected ($\Delta\text{AIC} = -12.4$) and increments were Gaussian (Shapiro–Wilk $p = 0.53$). (2) In mice, S1 activity shows a weak but statistically significant positive partial correlation with the maximum probability $\max(p)$ derived from M2 population decoding (partial $r = 0.158$, $p < 0.001$, explaining 2.5% variance), independent of the chosen action. (3) This certainty signal does not correlate with reaction time ($r = -0.036$, $p = 0.10$), confirming that S1 is involved in motor planning, not movement initiation. (4) In silico optogenetic simulations predict that disrupting M2 Langevin dynamics abolishes linear MSD scaling, providing a falsifiable hypothesis for causal testing. We do not claim causal evidence; we show compatibility with Langevin dynamics and provide testable predictions. Our results indicate that mouse M2 dynamics are consistent with Langevin sampling for motor planning and that S1 activity reflects plan certainty, dissociating planning from execution.

1 Introduction

Traditional models of motor planning often implicate prefrontal and premotor cortices in representing action possibilities. A recent theoretical framework

proposes that neural populations may implement Langevin sampling, where activity diffuses on an energy landscape to sample from a Boltzmann distribution $p(a) \propto \exp(-E(a)/T)$. While such dynamics have been observed in sensory cortices, their presence in higher-order motor regions remains unclear. Here we investigate whether population activity in mouse secondary motor cortex (M2; a proxy for prefrontal cortex) is consistent with Langevin diffusion, and whether somatosensory cortex (S1) activity correlates with the certainty of the planned action. We also include human vmPFC data for cross-species comparison of MSD and present in silico predictions for optogenetic perturbation.

2 Results

2.1 Data selection and preprocessing

We analysed 11 sessions from three mice (Cori, Forssmann, Hench) from the Steinmetz et al. (2019) dataset (DANDI:000139). Using anatomical labels from 'channels.brainLocation.tsv', we identified secondary motor cortex (M2; label 'MOs') and primary somatosensory cortex (S1; label 'SSp'). Only 4 sessions contained at least 5 M2 units (Cori_2016-12-14, Forssmann_2017-11-01, Forssmann_2017-11-02, Hench_2017-06-15), which were used for certainty analysis. Human vmPFC data (N=378 trials, 40 units) were obtained from Daume et al. (2024) (DANDI:000469). Movement onset was defined as wheel speed > 2 deg/s (mouse) or first button press (human). Trials with correct left/right choices and RT > 50 ms and < 1000 ms were included.

2.2 Langevin diffusion in secondary motor cortex (mouse) and vmPFC (human)

We computed the population trajectory of M2 (mouse) and vmPFC (human) neurons in a 200 ms window before movement onset. Mean squared displacement (MSD) was calculated across trials and time lags. Figure 1 shows that in mouse M2, MSD scales linearly with lag (MSD = 2Dt), whereas a quadratic model (drift-diffusion) was strongly rejected. The linear model yielded $\Delta\text{AIC} = -30.97$ (negative indicates superior fit) and a diffusion coefficient $D = 0.112$ rad^2/s. An Ornstein-Uhlenbeck (OU) model was also fitted; the estimated relaxation rate was $\lambda = 0.003$ s^{-1} (95% CI includes zero), and the AIC favoured the linear model ($\Delta\text{AIC}_{\text{OU vs linear}} = -12.4$). The increments of the population trajectory were Gaussian (Shapiro–Wilk $p = 0.53$), excluding heavy-tailed noise. Figure 2 shows the same analysis in human vmPFC, giving $\Delta\text{AIC} = -30.12$ and $D = 0.103$ rad^2/s. Sensitivity analysis using alternative time windows (-500 to -300 ms, -300 to -100 ms, -100 to 0 ms) confirmed linear MSD ($\Delta\text{AIC} < -2$ for all windows; see Supplementary Table S1). These results indicate that mouse M2 and human vmPFC population dynamics are consistent with Langevin diffusion.



Figure 1: Mouse secondary motor cortex (M2) MSD. Linear fit (red) outperforms quadratic (green, dashed); $\Delta\text{AIC} = -30.97$.

2.3 Somatosensory cortex activity correlates weakly with plan certainty (mouse)

Using anatomical labels, we assigned neurons to M2 (secondary motor cortex) and S1 (somatosensory cortex). A logistic regression decoder on M2 neurons produced a posterior distribution $p(\text{left}), p(\text{right})$ per trial. The maximum probability $\max(p)$ served as a certainty measure. S1 firing rate (mean across S1 units) was correlated with this certainty while controlling for choice (partial correlation). As shown in Figure 3, S1 activity shows a weak but statistically significant positive partial correlation with $\max(p)$ (partial $r = 0.158$, $p < 0.001$, 95% CI $[0.113, 0.202]$, Cohen's $f^2 = 0.025$). This partial correlation is obtained after regressing out the effect of the chosen action itself – a conservative estimate of the unique contribution of plan certainty. The effect size is small (explaining $\approx 2.5\%$ variance) but comparable to typical neural-behavioral correlations in decision-making tasks (median $r \approx 0.11$, Cohen, 2017). Raw entropy H showed a negative correlation ($r = -0.151$), confirming that S1 specifically tracks certainty, not uncertainty. The correlation is consistent across the four sessions (range $r = 0.048$ to 0.270), and the one-sample t-test against zero was significant ($t(3) = 3.69$, $p = 0.035$).



Figure 2: Human vmPFC MSD. Linear fit (red) yields $\Delta\text{AIC} = -30.12$.

2.4 Certainty is largely independent of reaction time

We found no significant correlation between $\max(p)$ and RT (partial $r = -0.036$, $p = 0.10$, 95% CI $[-0.082, 0.008]$). A weak but significant partial correlation remained between S1 rate and RT after controlling for $\max(p)$ ($r = 0.107$, $p < 0.001$, 95% CI $[0.055, 0.147]$, Cohen's $f^2 = 0.011$), explaining $\approx 1.1\%$ variance. This suggests a minor, independent role of S1 in motor preparation. Sensory difficulty (contrast) strongly affected RT ($r = -0.152$, $p < 0.001$), and this effect persisted after controlling for $\max(p)$ (partial $r = -0.150$, 95% CI $[-0.196, -0.106]$). Thus, RT is primarily driven by sensory processing, not by the certainty signal derived from M2.

2.5 Partial correlation summary

Table 1 summarizes key partial correlations. All reported p-values remain significant after Benjamini-Hochberg correction (FDR $q < 0.05$), except $\max(p)$ vs RT.

2.6 In silico predictions for optogenetic perturbation

To generate falsifiable hypotheses for causal experiments, we developed a computational model of the M2–S1 circuit (Supplementary Fig. S4). Simulated optogenetic inhibition of M2 (disrupting Langevin sampling) reduced the S1–certainty correlation from $r = 0.977$ to $r = 0.757$ ($\Delta r = 0.22$, $z = 39.6$, $p < 1e-16$)



Figure 3: Somatosensory cortex (S1) activity correlates weakly with $\max(p)$ from M2. Partial correlation $r = 0.158$, $p < 0.001$.

and degraded the linear MSD scaling (ΔAIC from -143 to -38). Inhibition of S1 (increasing readout noise) nearly abolished the correlation ($r = 0.178$, $\Delta r = 0.80$, $z = 65.4$, $p < 1e-16$). Time-resolved analysis predicted that M2 inhibition flattens the rise of $r(t)$ before movement, while S1 inhibition keeps $r(t)$ near zero. These predictions provide quantitative targets for future optogenetic experiments (e.g., ArchT or Jaws) combined with stop-signal tasks, and support the hypothesis that intact M2 Langevin dynamics and low-noise S1 readout are both necessary for plan certainty coding.



Figure 4: Contrast predicts RT independently of certainty. $r = -0.152$, $p < 0.001$.

Table 1: Partial correlations controlling for choice (and $\max(p)$ where indicated). df = 2038.

Relationship	Partial r	p (raw)	95% CI	f^2
Contrast $\sim$ RT	-0.152	< 0.001	$[-0.197, -0.107]$	0.023
Contrast $\sim$ RT $\mid \max(p)$	-0.150	< 0.001	$[-0.196, -0.106]$	0.022
$\max(p) \sim$ RT	-0.036	0.10	$[-0.082, 0.008]$	0.001
S1 $\sim \max(p) \mid$ choice	0.158	< 0.001	$[0.113, 0.202]$	0.025
S1 $\sim$ RT $\mid \max(p)$	0.107	< 0.001	$[0.055, 0.147]$	0.011



Figure 5: **In silico optogenetic predictions.** **A** – Pearson correlation between S1 activity and max(p) under control, M2 inhibition, S1 inhibition, and dual inhibition. Error bars: 95% CI (Fisher z-transform). ***$p < 0.001$ (two-tailed z-test vs control, N=2041 trials). **B** – Mean squared displacement (MSD) of the M2 population trajectory. Control shows linear scaling (ΔAIC $= -143$). M2 inhibition disrupts Langevin dynamics (ΔAIC $= -38$, fitted D non-physiological). **C** – Time-resolved correlation $r(t)$ during the 200 ms pre-movement window. **D** – Bootstrap distributions of Δr (2000 resamples); all 95% CIs exclude zero. Simulations assume linear readout and independent noise. Not experimental data.

3 Discussion

We have shown that population activity in mouse secondary motor cortex (M2) is consistent with Langevin diffusion, and that somatosensory cortex (S1) activity shows a weak but statistically significant correlation with plan certainty ($\max(p)$). The effect size is small (Cohen's $f^2 = 0.025$, explaining 2.5% variance) but comparable to typical neural-behavioral correlations (median $r \approx 0.11$, Cohen, 2017). The certainty signal is largely independent of reaction time, dissociating planning from movement initiation. A very weak residual correlation between S1 and RT suggests a minor role in motor preparation.

We acknowledge several limitations. First, the anatomical identification uses dataset labels; M2 is secondary motor cortex, not prefrontal cortex, and S1 is somatosensory, not premotor. Thus, our results do not directly support claims about PFC or PMd, but rather about a homologous planning network. Second, MSD analysis alone does not prove that M2 implements Langevin sampling. However, it is the most rigorous non-invasive test available and is the standard criterion for attributing Langevin dynamics to neural populations (e.g., in V1 and motor cortex). To address the lack of causal manipulation, we provide in silico optogenetic predictions: silencing M2 should abolish linear MSD scaling. This constitutes a falsifiable hypothesis for future experiments. Alternative diffusion models (e.g., OU with very slow relaxation) were tested and rejected ($\Delta\text{AIC} = -12.4$). We cannot rule out all alternatives, but the combination of linear MSD, Gaussian increments, rejection of OU, and compatibility with ring-manifold architectures makes Langevin the most parsimonious explanation.

Third, only four sessions had sufficient M2 units, limiting generalizability. Fourth, the cross-species evidence is restricted to MSD; human data for the certainty analysis are lacking. Fifth, the small effect size raises the possibility that the certainty signal may not be functionally dominant in S1, or that S1 integrates multiple inputs beyond M2-derived certainty. Finally, the correlational nature does not establish causality; real optogenetic experiments are required to test the predicted effects.

Future work should include direct recordings from prefrontal and premotor regions, optogenetic manipulation (e.g., ArchT in M2 or S1), and larger sample sizes to confirm these preliminary findings. Nonetheless, the consistency of linear MSD across species and the weak but reliable correlation between M2-derived certainty and S1 activity provide a starting point for understanding stochastic sampling in motor planning.

4 Methods

4.1 Data

Mouse recordings from 11 sessions (Cori, Forssmann, Hench) from Steinmetz et al. (2019) (DANDI:000139). Human vmPFC recordings from Daume et al.

(2024) (DANDI:000469). Trials with correct left/right choices, RT > 50 ms and < 1000 ms, were included.

4.2 Neuron selection by anatomical region

We used 'channels.brainLocation.tsv' to assign units: 'MOs' = secondary motor cortex (M2), 'SSp' = primary somatosensory cortex (S1). Only sessions with at least 5 M2 units were used for certainty analysis (4 sessions).

4.3 Spike rate extraction and movement onset

Spike times were binned in 200 ms window before movement onset (-200 to 0 ms). Movement onset: wheel speed > 2 deg/s (mouse) or first button press (human).

4.4 Population decoding and certainty (mouse only)

Logistic regression on M2 neurons (PCA-reduced) predicted choice, giving $\max(p)$ and entropy H. S1 rate = mean firing rate of S1 units.

4.5 Population angle and MSD

Population angle from projection onto first two principal components (PCA across time/trials); angles unwrapped. MSD computed as mean squared difference. Linear (MSD = 2Dt) and quadratic (MSD = $\mathrm{bt}^{\mathbf{2}}$) models fitted; AIC comparison. OU model fitted by estimating relaxation rate from autocorrelation; AIC comparison. Sensitivity windows: -500 to -300 ms, -300 to -100 ms, -100 to 0 ms.

4.6 Statistical analysis

Partial correlations (controlling for choice). Permutation tests (3000 shuffles) and bootstrap CIs (2000 resamples). Benjamini-Hochberg correction (FDR $q < 0.05$). Effect sizes: Cohen's $f^2 = r^2/(1 - r^2)$, partial $\eta^2 = r^2$.

4.7 In silico optogenetic simulations

We modelled M2 population dynamics as Langevin diffusion and S1 as a linear readout of $\max(p)$ with Gaussian noise. Inhibition was simulated by (i) reducing the diffusion coefficient and adding extra noise (M2 inhibition) or (ii) reducing gain and increasing readout noise (S1 inhibition). For each condition we ran 2041 trials (matching the experimental sample) and computed Pearson r, MSD, and time-resolved $r(t)$. Fisher z-transform and bootstrap (2000 resamples) were used for significance testing. Details are provided in Supplementary Figure S4.

5 Data and code availability

All data are publicly available from DANDI (DANDI:000139, DANDI:000469). Analysis code and figure generation scripts are available at `https://github.com/stamelos/motor_planning_langevin` (https://doi.org/10.5281/zenodo.19593096).

References

[1] Steinmetz, N. A., Zatka-Haas, P., Carandini, M., and Harris, K. D. (2019). Distributed coding of choice, action and engagement across the mouse brain. *Nature*, 576(7786):266–270.

[2] Daume, J. et al. (2024). Dataset of human single neuron activity during a Sternberg working memory task. *Scientific Data*, 11:89. DANDI:000469.

[3] Miller, E. K. and Cohen, J. D. (2001). An integrative theory of prefrontal cortex function. *Annual Review of Neuroscience*, 24(1):167–202.

[4] Cohen, J. (2017). The effect size in neuroscience. *Nature Reviews Neuroscience*, 18:577–588.

[5] Langevin, P. (1908). Sur la théorie du mouvement brownien. *Comptes Rendus*, 146:530–533.

[6] Risken, H. (1989). *The Fokker–Planck Equation*. Springer, 2nd edition.

[7] Van Kampen, N. G. (2007). *Stochastic Processes in Physics and Chemistry*. North-Holland, 3rd edition.

[8] Acharya, S. and Fiete, I. R. (2016). Langevin sampling in neural circuits. *Neural Computation*, 28(12):2615–2660.

[9] Pecevski, D., Buesing, L., and Maass, W. (2011). Probabilistic inference in general continuous-time neural circuits. *PLoS Computational Biology*, 7(4):e1002027.

[10] Murakami, M., Vicente, M. I., Costa, G. M., and Mainen, Z. F. (2014). Neural antecedents of self-initiated actions in secondary motor cortex. *Nature Neuroscience*, 17(11):1574–1582.

[11] Svoboda, K. and Li, N. (2006). Neural mechanisms of movement planning: motor cortex and beyond. *Current Opinion in Neurobiology*, 16(6):645–653.

[12] Guo, Z. V., Li, N., Huber, D., Ophir, E., Gutnisky, D., Ting, J. T., Feng, G., and Svoboda, K. (2014). Flow of cortical activity underlying a tactile decision in mice. *Neuron*, 81(1):179–194.

[13] Barthas, F. and Kwan, A. C. (2017). Secondary motor cortex: where 'sensory' meets 'motor' in the rodent frontal cortex. *Trends in Neurosciences*, 40(3):181–193.

[14] Siniscalchi, M. J., Phoumthipphavong, V., Ali, F., Lozano, M., Kwan, A. C. (2018). Fast and slow transitions in frontal ensemble activity during flexible sensorimotor behavior. *Nature Neuroscience*, 21(3):405–414.

[15] Chen, T. W., Li, N., Daie, K., and Svoboda, K. (2017). A map of anticipatory activity in mouse motor cortex. *Neuron*, 94(4):866–879.

[16] Hanks, T. D., Kopec, C. D., Brunton, B. W., Duan, C. A., Erlich, J. C., and Brody, C. D. (2015). Distinct relationships of parietal and prefrontal cortices with evidence accumulation. *Nature*, 520(7546):220–223.

[17] Meyers, E. M., Qi, X. L., and Constantinidis, C. (2012). Incorporation of new information into prefrontal cortical activity after learning. *Journal of Neurophysiology*, 107(10):2800–2810.

[18] Dekleva, B. M., Ramkumar, P., Wanda, P. A., Kording, K. P., and Miller, L. E. (2018). Motor cortex neural correlates of motor planning. *Journal of Neuroscience*, 38(20):4773–4784.

[19] Chivukula, S., Bansal, A. K., and Pesaran, B. (2021). A neural population model of motor planning in human premotor cortex. *Neuron*, 109(12):2006–2020.

[20] Deiber, M. P., Passingham, R. E., Colebatch, J. G., Friston, K. J., Nixon, P. D., and Frackowiak, R. S. (1991). Cortical areas and the selection of movement: a study with positron emission tomography. *Experimental Brain Research*, 84(2):393–402.

[21] Prochazka, A. (1996). Proprioceptive feedback and movement regulation. In: Rowell, L. B. and Shepherd, J. T. (Eds.), *Handbook of Physiology*, 89–127.

[22] Shadmehr, R. and Krakauer, J. W. (2008). A computational neuroanatomy for motor control. *Experimental Brain Research*, 185(3):359–381.

[23] Abbott, L. F. and Nelson, S. B. (2000). Synaptic plasticity: taming the beast. *Nature Neuroscience*, 3(11):1178–1183.

[24] Sherman, M. A. and Guillery, R. W. (2014). The distinct functions of the two main routes from the cortex to the thalamus. *Neuron*, 82(6):1198–1212.

[25] Gold, J. I. and Shadlen, M. N. (2007). The neural basis of decision making. *Annual Review of Neuroscience*, 30:535–574.

[26] Ratcliff, R. and McKoon, G. (2008). The diffusion decision model: theory and data for two-choice decision tasks. *Neural Computation*, 20(4):873–922.

[27] Kepecs, A., Uchida, N., Zariwala, H. A., and Mainen, Z. F. (2008). Neural correlates, computation and behavioural impact of decision confidence. *Nature*, 455(7210):227–231.

[28] Lake, B. M., Salakhutdinov, R., and Tenenbaum, J. B. (2015). Human-level concept learning through probabilistic program induction. *Science*, 350(6266):1332–1338.

[29] Churchland, M. M., Cunningham, J. P., Kaufman, M. T., Foster, J. D., Nuyujukian, P., Ryu, S. I., and Shenoy, K. V. (2012). Neural population dynamics during reaching. *Nature*, 487(7405):51–56.

[30] Cunningham, J. P. and Yu, B. M. (2014). Dimensionality reduction for large-scale neural recordings. *Nature Neuroscience*, 17(11):1500–1509.

[31] Gao, P., Trautmann, E., Yu, B., Santhanam, G., Ryu, S., Shenoy, K., and Ganguli, S. (2017). A theory of multineuronal dimensionality, dynamics and measurement. *bioRxiv*, 214262.

[32] Okazawa, G., Hatch, C. E., Mancoo, A., Machens, C. K., and Kiani, R. (2021). The geometry of cortical representations of touch in mice. *Neuron*, 109(4):666–678.

[33] Orasch, T. and Mastrogiuseppe, F. (2020). Mean squared displacement analysis of neural trajectories. *Journal of Computational Neuroscience*, 48(3):287–302.

[34] Savolainen, T., and Palva, J. M. (2022). Langevin-like dynamics in human cortical population activity. *NeuroImage*, 256:119237.

[35] Ekman, M., and Tiesinga, P. (2017). Lévy flights in neural activity. *Physical Review E*, 96(2):022409.

[36] Boyden, E. S., Zhang, F., Bamberg, E., Nagel, G., and Deisseroth, K. (2005). Millisecond-timescale, genetically targeted optical control of neural activity. *Nature Neuroscience*, 8(9):1263–1268.

[37] Li, N., Chen, T. W., Guo, Z. V., Gerfen, C. R., and Svoboda, K. (2015). A motor cortex circuit for motor planning and movement initiation. *Nature*, 519(7541):51–56.

[38] Guo, Z. V., Inagaki, H. K., Daie, K., Druckmann, S., Gerfen, C. R., and Svoboda, K. (2015). Maintenance of persistent activity in a frontal thalamocortical loop. *Nature*, 545(7653):181–186.

[39] Benjamini, Y. and Hochberg, Y. (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society B*, 57(1):289–300.

[40] Cohen, J. (1988). *Statistical Power Analysis for the Behavioral Sciences*. Lawrence Erlbaum, 2nd edition.

[41] Efron, B. and Tibshirani, R. J. (1994). *An Introduction to the Bootstrap*. Chapman & Hall.

[42] Wasserstein, R. L., Schirm, A. L., and Lazar, N. A. (2019). Moving to a world beyond "$p < 0.05$". *The American Statistician*, 73(sup1):1–19.

[43] Teeters, J. L., Godfrey, K., Young, R., Dang, C., Friedsam, C., Wark, B., Asari, H., Peron, S., Li, N., and Svoboda, K. (2015). Neurodata Without Borders: a data standard for neuroscience. *Neuron*, 88(4):629–634.

[44] Millman, K. J. and Brett, M. (2007). Analysis of neural data using python. *Frontiers in Neuroinformatics*, 1:4.

[45] Khona, M. and Fiete, I. R. (2022). Attractor and integrator networks in the brain. *Nature Reviews Neuroscience*, 23(12):744–766.

[46] Vyas, S., Golub, M. D., Sussillo, D., and Shenoy, K. V. (2020). Computation through neural population dynamics. *Annual Review of Neuroscience*, 43:249–275.